

A \mathbb{Z}_2 -index of symmetry protected topological phases with reflection symmetry for quantum spin chains

Yoshiko Ogata *

April 8, 2024

Abstract

For the classification of SPT phases, defining an index is a central problem. In the famous paper [PTBO1], Pollmann, Tuner, Berg, and Oshikawa introduced \mathbb{Z}_2 -indices for injective matrix products states (MPS) which have either $\mathbb{Z}_2 \times \mathbb{Z}_2$ dihedral group (of π -rotations about x , y , and z -axes) symmetry, time-reversal symmetry, or reflection symmetry. The first two are on-site symmetries. In [O4], an index for on-site symmetries, which generalizes the index in [PTBO1], was introduced for general unique gapped ground state phases in quantum spin chains. It was proved that the index is an invariant of the C^1 -classification of SPT phases. The index for the reflection symmetry, which is not an on-site symmetry, was left as an open question. In this paper, we introduce a \mathbb{Z}_2 -index for the reflection symmetric unique gapped ground state phases, and complete the generalization problem of index by Pollmann et.al. We also show that the index is an invariant of the C^1 -classification.

1 Introduction

Classification of unique gapped ground states in quantum many-body systems is an important problem in modern condensed matter physics and quantum information science. In one dimension, it is believed that all unique gapped ground states belong to a single phase, in the sense that any two such ground states can be smoothly connected with each other thorough a series of models with unique gapped ground states. This conjecture was verified for frustration free models with uniformly bounded degeneracy [O3]. Motivated by the study of the Haldane phenomena in antiferromagnetic quantum spin chains, Gu and Wen [GW] proposed a finer classification based on the notion of symmetry protected topological (SPT) phase. Instead of considering the whole family of Hamiltonians, we fix some symmetry and consider the set of all Hamiltonians with a unique gapped ground state in the bulk, satisfying the symmetry. We then say such two Hamiltonians are equivalent if they can be connected to each other via a continuous path of symmetric Hamiltonians with unique gapped ground state. It can be possible that two symmetric Hamiltonians which can be connected via a path of *non-symmetric* gapped Hamiltonians fails to be connected via a path of *symmetric* gapped Hamiltonians. A Hamiltonian which can not be connected to trivial Hamiltonians (i.e, Hamiltonians with on-site interactions) via a symmetry preserving path belongs to the SPT phase. The question is how to show some Hamiltonian is in the SPT phase. One way should be defining some index which is stable along the path of symmetric gapped Hamiltonians. If some Hamiltonian has an index which is different from that of trivial phases, the Hamiltonian should be in a SPT phase. Finding such an index is a non-trivial important question for the classification problem of SPT phases.

In the famous paper [PTBO1], Pollmann, Tuner, Berg, and Oshikawa introduced \mathbb{Z}_2 -indices for injective matrix products states (MPS) which have either $\mathbb{Z}_2 \times \mathbb{Z}_2$ dihedral group (of π -rotations

*Graduate School of Mathematical Sciences The University of Tokyo, Komaba, Tokyo, 153-8914, Japan Supported in part by the Grants-in-Aid for Scientific Research, JSPS.

about x , y , and z -axes) symmetry, time-reversal symmetry, or reflection symmetry. The first two are on-site symmetry, and the index is the cohomology class of some projective representation associated to the symmetric injective MPS. It was claimed there, that as the index takes discrete values, it should be stable under the continuous path of gapped Hamiltonians.

The \mathbb{Z}_2 -index beyond the framework of matrix product state was recently introduced by Tasaki for systems satisfying on-site $U(1)$ -symmetry together with one of $\mathbb{Z}_2 \times \mathbb{Z}_2$ -onsite symmetry/reflection symmetry/time reversal symmetry [Tas1]. He showed that these are actually invariant of the classification.

In [O4], we extended the index of Pollmann et.al. for *on-site* symmetry with full generality (without asking $U(1)$ -symmetry). We also proved that our index is an invariant of the C^1 -classification of SPT phases. The index for the reflection symmetry, which is not an on-site symmetry, was left as an open question. In this paper, we introduce a \mathbb{Z}_2 -index for the reflection symmetric unique gapped ground state phase, and complete the generalization problem of index by Pollmann et.al.

Now let us state our result more in details. For a Hilbert space \mathcal{H} , $B(\mathcal{H})$ denotes the set of all bounded operators on \mathcal{H} . If $V : \mathcal{H}_1 \rightarrow \mathcal{H}_2$ is a linear/anti-linear map from a Hilbert space \mathcal{H}_1 to another Hilbert space \mathcal{H}_2 , then $\text{Ad}(V) : B(\mathcal{H}_1) \rightarrow B(\mathcal{H}_2)$ denotes the map $\text{Ad}(V)(x) := VxV^*$, $x \in B(\mathcal{H}_1)$.

We start by summarizing standard setup of quantum spin chains on the infinite chain [BR1, BR2]. Throughout this paper, we fix some $2 \leq d \in \mathbb{N}$. We denote the algebra of $d \times d$ matrices by M_d . We denote the standard basis of \mathbb{C}^d by $\{\psi_\mu\}_{\mu=1,\dots,d}$, and set $e_{\mu,\nu} = |\psi_\mu\rangle\langle\psi_\nu|$ for each $\mu, \nu = 1, \dots, d$.

We denote the set of all finite subsets in \mathbb{Z} by $\mathfrak{S}_{\mathbb{Z}}$. For each $n \in \mathbb{N}$, we set $\Lambda_n := [-n, n] \cap \mathbb{Z}$. For each $z \in \mathbb{Z}$, let $\mathcal{A}_{\{z\}}$ be an isomorphic copy of M_d , and for any finite subset $\Lambda \subset \mathbb{Z}$, let $\mathcal{A}_\Lambda = \otimes_{z \in \Lambda} \mathcal{A}_{\{z\}}$, which is the local algebra of observables in Λ . For finite Λ , the algebra \mathcal{A}_Λ can be regarded as the set of all bounded operators acting on the Hilbert space $\otimes_{z \in \Lambda} \mathbb{C}^d$. We use this identification freely. If $\Lambda_1 \subset \Lambda_2$, the algebra \mathcal{A}_{Λ_1} is naturally embedded in \mathcal{A}_{Λ_2} by tensoring its elements with the identity. The algebra \mathcal{A}_R (resp. \mathcal{A}_L) representing the half-infinite chain is given as the inductive limit of the algebras \mathcal{A}_Λ with $\Lambda \in \mathfrak{S}_{\mathbb{Z}}$, $\Lambda \subset [0, \infty)$ (resp. $\Lambda \subset (-\infty, -1]$). The algebra \mathcal{A} , representing the two sided infinite chain is given as the inductive limit of the algebras \mathcal{A}_Λ with $\Lambda \in \mathfrak{S}_{\mathbb{Z}}$. Note that \mathcal{A}_Λ for $\Lambda \in \mathfrak{S}_{\mathbb{Z}}$, \mathcal{A}_L , and \mathcal{A}_R can be regarded naturally as subalgebras of \mathcal{A} . We denote the set of local observables by $\mathcal{A}_{\text{loc}} = \bigcup_{\Lambda \in \mathfrak{S}_{\mathbb{Z}}} \mathcal{A}_\Lambda$. We denote by β_x the automorphisms on \mathcal{A} representing the space translation by $x \in \mathbb{Z}$. By $Q^{(j)}$, $j \in \mathbb{Z}$, we denote the element of \mathcal{A} with $Q \in M_d$ in the j -th component of the tensor product of \mathcal{A} and the unit in any other component. The reflection γ is the unique $*$ -automorphism on \mathcal{A} which satisfies

$$\gamma(Q^{(j)}) = Q^{(-j-1)}, \quad \text{for all } Q \in M_d \text{ and } j \in \mathbb{Z}. \quad (1)$$

From γ , we define $*$ -isomorphisms $\gamma_{R \rightarrow L} : \mathcal{A}_R \rightarrow \mathcal{A}_L$ and $\gamma_{L \rightarrow R} : \mathcal{A}_L \rightarrow \mathcal{A}_R$ by

$$\gamma(\mathbb{I}_{\mathcal{A}_L} \otimes A) = \gamma_{R \rightarrow L}(A) \otimes \mathbb{I}_{\mathcal{A}_R}, \quad A \in \mathcal{A}_R, \quad (2)$$

and

$$\gamma(B \otimes \mathbb{I}_{\mathcal{A}_R}) = \mathbb{I}_{\mathcal{A}_L} \otimes \gamma_{L \rightarrow R}(B), \quad B \in \mathcal{A}_L. \quad (3)$$

We introduce the \mathbb{Z}_2 -index for reflection invariant pure states satisfying the split property (see Definition 2.1) in Section 2 Definition 2.7. Since a unique gapped ground state of a reflection invariant Hamiltonians satisfies these properties, this defines an index for such systems. (See Section 3.) The definition of the index is simple. Let ω be a reflection invariant pure state which satisfies the split property with respect to \mathcal{A}_L and \mathcal{A}_R . We then can find its GNS triple of the form $(\mathcal{H}_\omega \otimes \mathcal{H}_\omega, \pi_\omega \circ \gamma_{L \rightarrow R} \otimes \pi_\omega, \Omega_\omega)$ (where π_ω is an irreducible representation of \mathcal{A}_R) and a unitary operator Γ_ω on $\mathcal{H}_\omega \otimes \mathcal{H}_\omega$ implementing γ . (Lemma 2.5.) From the structure, we either

have $\Gamma_\omega(\xi \otimes \eta) = \eta \otimes \xi$ for all $\xi, \eta \in \mathcal{H}_\omega$ or $\Gamma_\omega(\xi \otimes \eta) = -\eta \otimes \xi$ for all $\xi, \eta \in \mathcal{H}_\omega$. (Theorem 2.6.) This sign $\sigma_\omega = \pm 1$ is our \mathbb{Z}_2 -index. The same index can be obtained from the Tomita-Takesaki modular conjugation.: For the above GNS triple of ω , let $\mathbb{I} \otimes s_\omega$ be the support projection of Ω_ω in $\mathbb{I} \otimes B(\mathcal{H}_\omega)$. Then we can consider modular conjugation J_ω associated to $s_\omega \otimes B(s_\omega \mathcal{H}_\omega)$ and Ω_ω . (Lemma 4.2.) There exists an anti-unitary $\theta : s_\omega \mathcal{H}_\omega \rightarrow s_\omega \mathcal{H}_\omega$ such that

$$J_\omega(s_\omega \otimes x) J_\omega^* = \theta x \theta^* \otimes s_\omega, \quad J_\omega(x \otimes s_\omega) J_\omega^* = s_\omega \otimes \theta x \theta^*, \quad (4)$$

for all $x \in B(s_\omega \mathcal{H}_\omega)$, (Proposition 4.3.) This θ satisfies $\theta^2 = \kappa_\omega s_\omega$ with some $\kappa_\omega \in \{-1, 1\}$, because of $J_\omega^2 = s_\omega \otimes s_\omega$. It turns out that κ_ω coincides with our \mathbb{Z}_2 -index σ_ω . (Theorem 4.4.) This θ is related to the Schmidt decomposition of Ω_ω . (Lemma 4.2.) Therefore, considering the Schmidt decomposition can be one way to calculate the index σ_ω . (Remark 4.5.)

As stated above, for reflection invariant injective matrix product states, a \mathbb{Z}_2 -index was introduced in [PTBO1]. It turns out that our \mathbb{Z}_2 -index restricted to such states coincides with that of [PTBO1]. This is proven in Section 5 using the relation $\kappa_\omega = \sigma_\omega$.

The \mathbb{Z}_2 -index σ_ω is invariant under automorphic equivalence via an automorphism which allows a reflection invariant decomposition.:

Definition. We say an automorphism α of \mathcal{A} allows a reflection invariant decomposition if there is an automorphisms α_R on \mathcal{A}_R , and a unitary W in \mathcal{A} such that

$$\tilde{\alpha}^{-1} \circ \alpha = \text{Ad}(W), \quad \gamma(W) = W, \quad (5)$$

where

$$\tilde{\alpha} := (\gamma_{R \rightarrow L} \circ \alpha_R \circ \gamma_{L \rightarrow R}) \otimes \alpha_R. \quad (6)$$

From the definition, we can show the following:

Theorem. (See Theorem 2.9 for details.) Let ω_0, ω_1 be reflection invariant pure states satisfying the split property. Suppose that ω_0 and ω_1 are automorphic equivalent via an automorphism which allows a reflection invariant decomposition. Then the \mathbb{Z}_2 -indices $\sigma_{\omega_0}, \sigma_{\omega_1}$ associated to ω_0, ω_1 are equal.

Recalling that a unique gapped ground state is pure and satisfies the split property (see Theorem 3.2), our \mathbb{Z}_2 -index can be understood as an index of reflection invariant Hamiltonians with unique gapped ground states. It turns out that this \mathbb{Z}_2 -index is an invariant of the C^1 -classification.:

Corollary. (See Theorem 3.6 for more precise statement) Let us consider a C^1 -path of interactions, in the reflection invariant unique gapped ground state phase. Suppose that if we associate some suitable boundary conditions along the path, they give local Hamiltonians which are gapped for an increasing sequence of finite boxes. (See Definition 3.4.) Then the \mathbb{Z}_2 -index σ_ω does not change along the path.

This can be shown from the fact that ground states along the C^1 -path are mutually automorphic equivalent via an automorphism which allows a reflection invariant decomposition. The boundary conditions in the Corollary can be arbitrary, as long as they guarantee the gap. We may take it as periodic boundary condition, for example. Furthermore, the boundary condition itself does not need to be reflection invariant.

Our theorem, along with results in [PTOB1,PTOB2,CGW,Tas2] about matrix product states, shows that AKLT interaction and trivial interaction belong to different reflection symmetric unique gapped ground state phases. In other words, AKLT interaction and trivial interaction can never be connected by a C^1 -path of reflection invariant interactions without closing the gap.

2 The \mathbb{Z}_2 -index associated to the reflection symmetric split states

We introduce \mathbb{Z}_2 -index for reflection invariant pure state satisfying the split property. Let us first recall the definition of the split property. Here we give the following definition, which is most suitable for our purpose. It corresponds to the standard definition [DL] in our setting (see [M3]).

Definition 2.1. Let φ be a pure state on \mathcal{A} . Let φ_R be the restriction of φ to \mathcal{A}_R , and $(\mathcal{H}_{\varphi_R}, \pi_{\varphi_R}, \Omega_{\varphi_R})$ be the GNS triple of φ_R . We say φ satisfies the split property with respect to \mathcal{A}_L and \mathcal{A}_R , if the von Neumann algebra $\pi_{\varphi_R}(\mathcal{A}_R)''$ is a type I factor.

Recall that a type I factor is isomorphic to $B(\mathcal{K})$, the set of all bounded operators on a Hilbert space \mathcal{K} . See [T1]. We consider following type of GNS-triple for a reflection invariant pure state which satisfies the split property.

Definition 2.2. Let ω be a reflection invariant pure state on \mathcal{A} which satisfies the split property with respect to \mathcal{A}_R and \mathcal{A}_L . We say $(\mathcal{H}_\omega, \pi_\omega, \Omega_\omega, \Gamma_\omega)$ is a reflection-split representation associated to ω if setting $\hat{\mathcal{H}}_\omega := \mathcal{H}_\omega \otimes \mathcal{H}_\omega$ and $\hat{\pi}_\omega := (\pi_\omega \circ \gamma_{L \rightarrow R}) \otimes \pi_\omega$, following hold:

1. $(\mathcal{H}_\omega, \pi_\omega)$ is an irreducible representation of \mathcal{A}_R ,
2. Ω_ω is a unit vector of $\hat{\mathcal{H}}_\omega$,
3. $(\hat{\mathcal{H}}_\omega, \hat{\pi}_\omega, \Omega_\omega)$ is a GNS triple of ω ,
4. Γ_ω is the unique unitary operator on $\hat{\mathcal{H}}_\omega$ such that

$$\Gamma_\omega \hat{\pi}_\omega(A) \Omega_\omega = \hat{\pi}_\omega \circ \gamma(A) \Omega_\omega, \quad A \in \mathcal{A}. \quad (7)$$

Remark 2.3. Because of $\gamma^2 = \text{id}$, from the definition of Γ_ω (7), we have $\Gamma_\omega^2 = \mathbb{I}_{\mathcal{H}_\omega}$.

Remark 2.4. For the rest of this paper, for any reflection-split representation $(\mathcal{H}_\omega, \pi_\omega, \Omega_\omega, \Gamma_\omega)$ we use the notation $\hat{\mathcal{H}}_\omega := \mathcal{H}_\omega \otimes \mathcal{H}_\omega$ and $\hat{\pi}_\omega := (\pi_\omega \circ \gamma_{L \rightarrow R}) \otimes \pi_\omega$.

Lemma 2.5. *For any reflection invariant pure state ω on \mathcal{A} which satisfies the split property with respect to \mathcal{A}_R and \mathcal{A}_L , there exists a reflection-split representation $(\mathcal{H}_\omega, \pi_\omega, \Omega_\omega, \Gamma_\omega)$ associated to ω . Furthermore, if $(\mathcal{H}_\omega, \bar{\pi}_\omega, \bar{\Omega}_\omega, \bar{\Gamma}_\omega)$ is another reflection-split representation of ω , there exists a unitary $V : \mathcal{H}_\omega \rightarrow \mathcal{H}_\omega$ such that*

$$\text{Ad}(V) \circ \pi_\omega(A) = \bar{\pi}_\omega(A), \quad A \in \mathcal{A}_R, \quad (8)$$

$$\bar{\Gamma}_\omega = \text{Ad}(V \otimes V)(\Gamma_\omega), \quad (9)$$

$$(V \otimes V) \Omega_\omega = \bar{\Omega}_\omega. \quad (10)$$

Proof. Let $(\mathcal{H}_R, \pi_R, \Omega_R)$ be a GNS triple of $\omega|_{\mathcal{A}_R}$. Note that from the reflection invariance of ω , $(\mathcal{H}_R, \pi_R \circ \gamma_{L \rightarrow R}, \Omega_R)$ is a GNS triple of $\omega|_{\mathcal{A}_L}$. Therefore, $(\mathcal{H}_R \otimes \mathcal{H}_R, \pi_R \circ \gamma_{L \rightarrow R} \otimes \pi_R, \Omega_R \otimes \Omega_R)$ is a GNS triple of $\omega|_{\mathcal{A}_L \otimes \mathcal{A}_R}$.

Since ω satisfies the split property, there exists a Hilbert space \mathcal{H}_ω and a $*$ -isomorphism $\iota : \pi_R(\mathcal{A}_R)'' \rightarrow B(\mathcal{H}_\omega)$. We introduce a representation $\pi_\omega := \iota \circ \pi_R$ of \mathcal{A}_R on \mathcal{H}_ω . Since ι is a $*$ -isomorphism, $(\mathcal{H}_\omega, \pi_\omega)$ is an irreducible representation of \mathcal{A}_R .

Set $\hat{\mathcal{H}}_\omega := \mathcal{H}_\omega \otimes \mathcal{H}_\omega$ and let $\hat{\pi}_\omega := (\pi_\omega \circ \gamma_{L \rightarrow R}) \otimes \pi_\omega$ be the representation of \mathcal{A} on $\hat{\mathcal{H}}_\omega$. Now we would like to show the existence of a unit vector $\Omega_\omega \in \hat{\mathcal{H}}_\omega$ such that $(\hat{\mathcal{H}}_\omega, \hat{\pi}_\omega, \Omega_\omega)$ is a GNS triple of ω . Since there is a $*$ -isomorphism

$$\iota \otimes \iota : (\pi_R \circ \gamma_{L \rightarrow R}(\mathcal{A}_L))'' \otimes \pi_R(\mathcal{A}_R)'' = (\pi_R(\mathcal{A}_R))'' \otimes \pi_R(\mathcal{A}_R)'' \rightarrow B(\mathcal{H}_\omega \otimes \mathcal{H}_\omega)$$

(Theorem 5.2 IV of [T1]), the representation $\hat{\pi}_\omega = (\iota \otimes \iota) \circ (\pi_R \circ \gamma_{L \rightarrow R} \otimes \pi_R)$ is quasi-equivalent to $\pi_R \circ \gamma_{L \rightarrow R} \otimes \pi_R$, the GNS representation of $\omega|_{\mathcal{A}_L \otimes \mathcal{A}_R}$. (Theorem 2.4.26 [BR1].) On the

other hand, as ω satisfies the split property, by the proof of Proposition 2.2 of [M2], $\omega|_{\mathcal{A}_L} \otimes \omega|_{\mathcal{A}_R}$ is quasi-equivalent to ω . (In Proposition 2.2 of [M2], it is assumed that the state is translationally invariant because of the first equivalent condition (i). However, the proof for the equivalence between (ii) and (iii) does not require translation invariance.) Hence $\hat{\pi}_\omega$ is quasi-equivalent to the GNS representation of ω . (See section 2.4 of [BR1].) Therefore, there is a density matrix ρ on $\hat{\mathcal{H}}_\omega$ such that

$$\omega(A) = \text{Tr } \rho (\hat{\pi}_\omega(A)), \quad A \in \mathcal{A}. \quad (11)$$

Since ω is pure and $\hat{\pi}_\omega(\mathcal{A})'' = B(\hat{\mathcal{H}}_\omega)$, this ρ should be a one rank operator onto a one dimensional space $\mathbb{C}\Omega_\omega$, with some unit vector $\Omega_\omega \in \hat{\mathcal{H}}_\omega$. Because of $\hat{\pi}_\omega(\mathcal{A})'' = B(\hat{\mathcal{H}}_\omega)$, Ω_ω is cyclic for $\hat{\pi}_\omega(\mathcal{A})$, and $(\hat{\mathcal{H}}_\omega, \hat{\pi}_\omega, \Omega_\omega)$ is a GNS triple of ω . From the γ -invariance of ω , there exists Γ_ω satisfying (7). (Corollary 2.3.17 of [BR1].)

Now let us show the uniqueness. Let $(\bar{\mathcal{H}}_\omega, \bar{\pi}_\omega, \bar{\Omega}_\omega, \bar{\Gamma}_\omega)$ be another reflection-spilt representation of ω . Since both of $(\mathcal{H}_\omega \otimes \mathcal{H}_\omega, (\pi_\omega \circ \gamma_{L \rightarrow R}) \otimes \pi_\omega, \Omega_\omega)$ and $(\bar{\mathcal{H}}_\omega \otimes \bar{\mathcal{H}}_\omega, (\bar{\pi}_\omega \circ \gamma_{L \rightarrow R}) \otimes \bar{\pi}_\omega, \bar{\Omega}_\omega)$ are GNS triple of ω , there is a unitary $U : \mathcal{H}_\omega \rightarrow \bar{\mathcal{H}}_\omega$ such that

$$U(\pi_\omega \circ \gamma_{L \rightarrow R} \otimes \pi_\omega)(A)U^* = (\bar{\pi}_\omega \circ \gamma_{L \rightarrow R} \otimes \bar{\pi}_\omega)(A), \quad A \in \mathcal{A}, \text{ and } U\Omega_\omega = \bar{\Omega}_\omega. \quad (12)$$

(Theorem 2.3.16 of [BR1].) Restricting the first equation to \mathcal{A}_R , we have

$$U(\mathbb{I}_{\mathcal{H}_\omega} \otimes \pi_\omega(A))U^* = \mathbb{I}_{\bar{\mathcal{H}}_\omega} \otimes \bar{\pi}_\omega(A), \quad A \in \mathcal{A}_R. \quad (13)$$

From this we obtain a $*$ -isomorphism τ from $B(\mathcal{H}_\omega) = \pi_\omega(\mathcal{A}_R)''$ to $B(\bar{\mathcal{H}}_\omega) = \bar{\pi}_\omega(\mathcal{A}_R)''$ such that

$$U(\mathbb{I}_{\mathcal{H}_\omega} \otimes x)U^* = \mathbb{I}_{\bar{\mathcal{H}}_\omega} \otimes \tau(x), \quad x \in B(\mathcal{H}_\omega), \quad (14)$$

which satisfies

$$\tau \circ \pi_\omega(A) = \bar{\pi}_\omega(A), \quad A \in \mathcal{A}_R. \quad (15)$$

Applying Wigner's theorem to τ , there exists a unitary $\tilde{V} : \mathcal{H}_\omega \rightarrow \bar{\mathcal{H}}_\omega$ such that

$$\tau(x) = \tilde{V}x\tilde{V}^*, \quad x \in B(\mathcal{H}_\omega). \quad (16)$$

In particular, we have

$$\text{Ad}(\tilde{V})(\pi_\omega(A)) = \bar{\pi}_\omega(A), \quad A \in \mathcal{A}_R. \quad (17)$$

From this and (12), we have

$$\begin{aligned} \text{Ad}(U)((\pi_\omega \circ \gamma_{L \rightarrow R} \otimes \pi_\omega)(A)) &= (\bar{\pi}_\omega \circ \gamma_{L \rightarrow R} \otimes \bar{\pi}_\omega)(A) \\ &= \text{Ad}(\tilde{V} \otimes \tilde{V})((\pi_\omega \circ \gamma_{L \rightarrow R} \otimes \pi_\omega)(A)), \quad A \in \mathcal{A}. \end{aligned} \quad (18)$$

Since $((\pi_\omega \circ \gamma_{L \rightarrow R} \otimes \pi_\omega)(\mathcal{A}))'' = B(\mathcal{H}_\omega \otimes \mathcal{H}_\omega)$, $(\tilde{V} \otimes \tilde{V})^*U = c\mathbb{I}_{\mathcal{H}_\omega}$, for some $c \in \mathbb{T}$. Choosing one $c_1 \in \mathbb{T}$ such that $c_1^2 = c$, we set $V := c_1\tilde{V}$. Then we have $U = V \otimes V$, and from (12), (10) holds. The property (9) holds from (12) and (10). By (17), we obtain (8). \square

For a reflection invariant pure state which satisfies the split property with respect to \mathcal{A}_R and \mathcal{A}_L we can define an index via the reflection-split representation of ω .

Theorem 2.6. *Let ω be a reflection invariant pure state which satisfies the split property with respect to \mathcal{A}_R and \mathcal{A}_L . Let $(\mathcal{H}_\omega, \pi_\omega, \Omega_\omega, \Gamma_\omega)$ be a reflection-spilt representation associated to ω . Then we have*

$$\Gamma_\omega(\mathbb{I}_{\mathcal{H}_\omega} \otimes x)\Gamma_\omega^* = x \otimes \mathbb{I}_{\mathcal{H}_\omega}, \quad x \in B(\mathcal{H}_\omega). \quad (19)$$

Furthermore, there exists a constant $\sigma_\omega = \pm 1$ such that

$$\Gamma_\omega(\xi \otimes \eta) = \sigma_\omega \eta \otimes \xi, \quad \text{for all } \xi, \eta \in \mathcal{H}_\omega. \quad (20)$$

This σ_ω is independent of the choice of the reflection-spilt representation $(\mathcal{H}_\omega, \pi_\omega, \Omega_\omega, \Gamma_\omega)$.

Definition 2.7. From Theorem 2.6, for each reflection invariant pure state ω which satisfies the split property with respect to \mathcal{A}_R and \mathcal{A}_L , we can define a \mathbb{Z}_2 -index σ_ω .

Proof. We first prove (19). For any $A \in \mathcal{A}_R$, we have

$$\begin{aligned}\Gamma_\omega(\mathbb{I}_{\mathcal{H}_\omega} \otimes \pi_\omega(A)) \Gamma_\omega^* &= \Gamma_\omega(\hat{\pi}_\omega(\mathbb{I}_{\mathcal{A}_L} \otimes A)) \Gamma_\omega^* = \hat{\pi}_\omega \circ \gamma(\mathbb{I}_{\mathcal{A}_L} \otimes A) \\ &= \hat{\pi}_\omega(\gamma_{R \rightarrow L}(A) \otimes \mathbb{I}_{\mathcal{H}_\omega}) = \pi_\omega \circ \gamma_{L \rightarrow R}(\gamma_{R \rightarrow L}(A)) \otimes \mathbb{I}_{\mathcal{A}_R} = \pi_\omega(A) \otimes \mathbb{I}_{\mathcal{H}_\omega}.\end{aligned}\quad (21)$$

Since both sides are σ -weak continuous and $\pi_\omega(\mathcal{A}_R)'' = B(\mathcal{H}_\omega)$, we obtain (19).

From (19), we derive (20).: For any nonzero $\xi, \eta \in \mathcal{H}_\omega$, there exists $\sigma_{\xi, \eta} \in \mathbb{T}$ such that

$$\Gamma_\omega(\xi \otimes \eta) = \sigma_{\xi, \eta}(\eta \otimes \xi), \quad (22)$$

because

$$\Gamma_\omega|\xi \otimes \eta\rangle \langle \xi \otimes \eta| \Gamma_\omega^* = |\eta \otimes \xi\rangle \langle \eta \otimes \xi|, \quad (23)$$

from (19). Considering the case $\xi = \eta \neq 0$ in (22), we have

$$\xi \otimes \xi = \Gamma_\omega^2(\xi \otimes \xi) = \sigma_{\xi, \xi} \Gamma_\omega(\xi \otimes \xi) = \sigma_{\xi, \xi}^2 \xi \otimes \xi. \quad (24)$$

The first equality is from Remark 2.3. From this, we obtain $\sigma_{\xi, \xi} = \pm 1$. Again by (19), for nonzero $\xi, \eta \in \mathcal{H}_\omega$, we obtain

$$\Gamma_\omega|\xi \otimes \xi\rangle \langle \eta \otimes \eta| \Gamma_\omega^* = |\xi \otimes \xi\rangle \langle \eta \otimes \eta|. \quad (25)$$

On the other hand, from the above argument, we have

$$\Gamma_\omega|\xi \otimes \xi\rangle \langle \eta \otimes \eta| \Gamma_\omega^* = \sigma_{\xi, \xi} \sigma_{\eta, \eta} |\xi \otimes \xi\rangle \langle \eta \otimes \eta|. \quad (26)$$

Since η, ξ are not zero, we obtain $\sigma_{\xi, \xi} \sigma_{\eta, \eta} = 1$. Recalling that $\sigma_{\xi, \xi}, \sigma_{\eta, \eta}$ take values in ± 1 , we obtain $\sigma_{\xi, \xi} = \sigma_{\eta, \eta}$. Therefore, we set $\sigma_\omega := \sigma_{\xi, \xi}$, which is independent of the choice of nonzero $\xi \in \mathcal{H}_\omega$. To prove (20), we use (19) again and for any nonzero $\xi, \eta \in \mathcal{H}_\omega$, we have

$$|\eta \otimes \xi\rangle \langle \xi \otimes \xi| = \Gamma_\omega|\xi \otimes \eta\rangle \langle \xi \otimes \xi| \Gamma_\omega^* = \sigma_{\xi, \eta} \sigma_\omega |\eta \otimes \xi\rangle \langle \xi \otimes \xi|. \quad (27)$$

The first equality follows from (19) and the second one is the definition of $\sigma_{\xi, \eta}$ and σ_ω . From this and $\sigma_\omega = \pm 1$, we obtain $\sigma_{\xi, \eta} = \sigma_\omega$, completing the proof of (20).

To show that the sign σ_ω is independent of the choice of the reflection-split representation, let $(\bar{\mathcal{H}}_\omega, \bar{\pi}_\omega, \bar{\Omega}_\omega, \bar{\Gamma}_\omega)$ be another reflection-split representations associated to ω . Let $V : \mathcal{H}_\omega \rightarrow \bar{\mathcal{H}}_\omega$ be the unitary given in Lemma 2.5. We have

$$\bar{\Gamma}_\omega(\xi \otimes \eta) = (V \otimes V) \Gamma_\omega(V^* \xi \otimes V^* \eta) = (V \otimes V) \sigma_\omega(V^* \eta \otimes V^* \xi) = \sigma_\omega(\eta \otimes \xi), \quad (28)$$

for any $\xi, \eta \in \bar{\mathcal{H}}_\omega$, proving the claim. \square

Now we prove that the \mathbb{Z}_2 -index is invariant under automorphic equivalence via an automorphism which allows a reflection invariant decomposition. Let us recall the definition:

Definition 2.8. We say an automorphism α of \mathcal{A} allows a reflection invariant decomposition if there is an automorphisms α_R on \mathcal{A}_R , and a unitary W in \mathcal{A} such that

$$\tilde{\alpha}^{-1} \circ \alpha = \text{Ad}(W), \quad \gamma(W) = W, \quad (29)$$

where

$$\tilde{\alpha} := (\gamma_{R \rightarrow L} \circ \alpha_R \circ \gamma_{L \rightarrow R}) \otimes \alpha_R. \quad (30)$$

We call (α_R, W) , a reflection invariant decomposition of α .

We prove the following theorem:

Theorem 2.9. *Let ω_0, ω_1 be reflection invariant pure states satisfying the split property with respect to \mathcal{A}_R and \mathcal{A}_L . Suppose that ω_0 and ω_1 are automorphic equivalent via an automorphism α , i.e., $\omega_1 = \omega_0 \circ \alpha$, which allows a reflection invariant decomposition (α_R, W) . Then the \mathbb{Z}_2 -indices $\sigma_{\omega_0}, \sigma_{\omega_1}$ associated to ω_0, ω_1 are equal.*

Proof. Let $(\mathcal{H}_{\omega_0}, \pi_{\omega_0}, \Omega_{\omega_0}, \Gamma_{\omega_0})$ be a reflection-split representation associated to ω_0 . Set $\hat{\mathcal{H}}_{\omega_0} := \mathcal{H}_{\omega_0} \otimes \mathcal{H}_{\omega_0}$, $\hat{\pi}_{\omega_0} := (\pi_{\omega_0} \circ \gamma_{L \rightarrow R}) \otimes \pi_{\omega_0}$. We also set $\tilde{\alpha} := (\gamma_{R \rightarrow L} \circ \alpha_R \circ \gamma_{L \rightarrow R}) \otimes \alpha_R$ and

$$\hat{\pi}_{\omega_1} := (\pi_{\omega_0} \circ \alpha_R \circ \gamma_{L \rightarrow R}) \otimes (\pi_{\omega_0} \circ \alpha_R) = \hat{\pi}_{\omega_0} \circ \tilde{\alpha}.$$

We claim that $(\mathcal{H}_{\omega_0}, \pi_{\omega_0} \circ \alpha_R, \hat{\pi}_{\omega_1}(W^*)\Omega_{\omega_0}, \Gamma_{\omega_0})$ is a reflection-split representation of ω_1 . From this, we obtain the statement of the Theorem i.e., $\sigma_{\omega_0} = \sigma_{\omega_1}$.

The first condition of Definition 2.2 is from $\pi_{\omega_0} \circ \alpha_R(\mathcal{A}_R)'' = \pi_{\omega_0}(\mathcal{A}_R)'' = B(\mathcal{H}_{\omega_0})$. The second one is trivial because W is unitary. To prove the third one, note that $(\hat{\mathcal{H}}_{\omega_0}, \hat{\pi}_{\omega_1}, \Omega_{\omega_0})$ is a GNS triple of $\omega_0 \circ \tilde{\alpha}$. From $\tilde{\alpha}^{-1} \circ \alpha = \text{Ad}(W)$ and $\omega_1 = \omega_0 \circ \alpha$, we have

$$\omega_1 = \omega_0 \circ \alpha = \omega_0 \circ \tilde{\alpha} \circ \text{Ad}(W). \quad (31)$$

Combining these two facts, we have

$$\omega_1(A) = \langle \hat{\pi}_{\omega_1}(W^*)\Omega_{\omega_0}, \hat{\pi}_{\omega_1}(A)\hat{\pi}_{\omega_1}(W^*)\Omega_{\omega_0} \rangle, \quad A \in \mathcal{A}. \quad (32)$$

Since $\hat{\pi}_{\omega_1}(\mathcal{A})'' = B(\hat{\mathcal{H}}_{\omega_0})$, $\hat{\pi}_{\omega_1}(W^*)\Omega_{\omega_0}$ is cyclic for $\hat{\pi}_{\omega_1}(\mathcal{A})$. Hence $(\hat{\mathcal{H}}_{\omega_0}, \hat{\pi}_{\omega_1}, \hat{\pi}_{\omega_1}(W^*)\Omega_{\omega_0})$ is a GNS triple of ω_1 . Finally for the fourth condition of Definition 2.2, note that γ and $\tilde{\alpha}$ commute because

$$\begin{aligned} \gamma \circ \tilde{\alpha}(A \otimes B) &= \gamma \circ ((\gamma_{R \rightarrow L} \circ \alpha_R \circ \gamma_{L \rightarrow R}(A)) \otimes \alpha_R(B)) = (\gamma_{R \rightarrow L} \circ \alpha_R(B)) \otimes (\alpha_R \circ \gamma_{L \rightarrow R}(A)) \\ &= (\gamma_{R \rightarrow L} \circ \alpha_R \circ \gamma_{L \rightarrow R} \circ \gamma_{R \rightarrow L}(B)) \otimes \alpha_R \circ \gamma_{L \rightarrow R}(A) = \tilde{\alpha} \circ \gamma(A \otimes B), \end{aligned} \quad (33)$$

for any $A \in \mathcal{A}_L$ and $B \in \mathcal{A}_R$. From this fact, we obtain

$$\begin{aligned} \Gamma_{\omega_0} \hat{\pi}_{\omega_1}(A) \hat{\pi}_{\omega_1}(W^*)\Omega_{\omega_0} &= \Gamma_{\omega_0} \hat{\pi}_{\omega_0} \circ \tilde{\alpha}(AW^*)\Omega_{\omega_0} = (\hat{\pi}_{\omega_0} \circ \gamma \circ \tilde{\alpha}(AW^*))\Omega_{\omega_0} \\ &= (\hat{\pi}_{\omega_0} \circ \tilde{\alpha} \circ \gamma(AW^*))\Omega_{\omega_0} = (\hat{\pi}_{\omega_0} \circ \tilde{\alpha} \circ \gamma(A))(\hat{\pi}_{\omega_0} \circ \tilde{\alpha}(W^*))\Omega_{\omega_0} = (\hat{\pi}_{\omega_1} \circ \gamma(A))(\hat{\pi}_{\omega_1}(W^*))\Omega_{\omega_0}, \end{aligned} \quad (34)$$

for all $A \in \mathcal{A}$. For the fourth equality, we used $\gamma(W) = W$. This completes the proof. \square

3 C^1 -classification of gapped Hamiltonians with the reflection symmetry.

Let us now apply the result in Section 2 to the C^1 -classification of gapped Hamiltonians preserving the reflection symmetry.

A mathematical model of a quantum spin chain is fully specified by its interaction Φ . An interaction is a map Φ from $\mathfrak{S}_{\mathbb{Z}}$ into \mathcal{A}_{loc} such that $\Phi(X) \in \mathcal{A}_X$ and $\Phi(X) = \Phi(X)^*$ for each $X \in \mathfrak{S}_{\mathbb{Z}}$. Let $R : \mathbb{Z} \rightarrow \mathbb{Z}$ be the reflection : $R(i) := -i - 1$, $i \in \mathbb{Z}$. An interaction Φ is reflection invariant if $\gamma(\Phi(X)) = \Phi(R(X))$ for all $X \in \mathfrak{S}_{\mathbb{Z}}$. An interaction Φ is of finite range if there exists an $m \in \mathbb{N}$ such that $\Phi(X) = 0$ for X with diameter larger than m . We denote by \mathcal{B}_f , the set of all finite range interactions Φ which satisfy

$$a_{\Phi} := \sup_{X \in \mathfrak{S}_{\mathbb{Z}}} \|\Phi(X)\| < \infty. \quad (35)$$

We may define addition on \mathcal{B}_f : for $\Phi, \Psi \in \mathcal{B}_f$, $\Phi + \Psi$ denotes the interaction given by $(\Phi + \Psi)(X) = \Phi(X) + \Psi(X)$ for each $X \in \mathfrak{S}_{\mathbb{Z}}$.

For an interaction Φ and a finite set $\Lambda \in \mathfrak{S}_{\mathbb{Z}}$, we define the local Hamiltonian on Λ by

$$(H_{\Phi})_{\Lambda} := \sum_{X \subset \Lambda} \Phi(X). \quad (36)$$

The dynamics given by this local Hamiltonian is denoted by

$$\tau_t^{\Phi, \Lambda}(A) := e^{it(H_{\Phi})_{\Lambda}} A e^{-it(H_{\Phi})_{\Lambda}}, \quad t \in \mathbb{R}, \quad A \in \mathcal{A}. \quad (37)$$

If Φ belongs to \mathcal{B}_f , the limit

$$\tau_t^{\Phi}(A) = \lim_{\Lambda \rightarrow \mathbb{Z}} \tau_t^{\Phi, \Lambda}(A) \quad (38)$$

exists for each $A \in \mathcal{A}$ and $t \in \mathbb{R}$, and defines a strongly continuous one parameter group of automorphisms τ^{Φ} on \mathcal{A} . (See [BR2].) We denote the generator of the C^* -dynamics τ^{Φ} by δ_{Φ} .

For $\Phi \in \mathcal{B}_f$, a state φ on \mathcal{A} is called a τ^{Φ} -ground state if the inequality $-i\varphi(A^*\delta_{\Phi}(A)) \geq 0$ holds for any element A in the domain $\mathcal{D}(\delta_{\Phi})$ of δ_{Φ} . Let φ be a τ^{Φ} -ground state, with the GNS triple $(\mathcal{H}_{\varphi}, \pi_{\varphi}, \Omega_{\varphi})$. Then there exists a unique positive operator $H_{\varphi, \Phi}$ on \mathcal{H}_{φ} such that $e^{itH_{\varphi, \Phi}} \pi_{\varphi}(A) \Omega_{\varphi} = \pi_{\varphi}(\tau_t^{\Phi}(A)) \Omega_{\varphi}$, for all $A \in \mathcal{A}$ and $t \in \mathbb{R}$. We call this $H_{\varphi, \Phi}$ the bulk Hamiltonian associated with φ . Note that Ω_{φ} is an eigenvector of $H_{\varphi, \Phi}$ with eigenvalue 0. See [BR2] for the general theory.

The following definition clarifies what we mean by a model with a unique gapped ground state.

Definition 3.1. We say that a model with an interaction $\Phi \in \mathcal{B}_f$ has a unique gapped ground state if (i) the τ^{Φ} -ground state, which we denote as φ , is unique, and (ii) there exists a $g > 0$ such that $\sigma(H_{\varphi, \Phi}) \setminus \{0\} \subset [g, \infty)$, where $\sigma(H_{\varphi, \Phi})$ is the spectrum of $H_{\varphi, \Phi}$.

Note that the uniqueness of φ implies that 0 is a non-degenerate eigenvalue of $H_{\varphi, \Phi}$.

If φ is a τ^{Φ} -ground state of reflection invariant interaction $\Phi \in \mathcal{B}_f$, then its reflection $\varphi \circ \gamma$ is also a τ^{Φ} -ground state. In particular, if φ is a unique τ^{Φ} -ground state, it is pure and reflection invariant.

In [M3], T.Matsui showed that the spectral gap implies the split property.

Theorem 3.2 (Theorem 1.5, Lemma 4.1, and Proposition 4.2 of [M3]). *Let φ be a pure τ^{Φ} -ground state of $\Phi \in \mathcal{B}_f$, and denote by $H_{\varphi, \Phi}$ the corresponding bulk Hamiltonian. Assume that 0 is a non-degenerate eigenvalue of $H_{\varphi, \Phi}$ and there exists $g > 0$ such that $\sigma(H_{\varphi, \Phi}) \setminus \{0\} \subset [g, \infty)$. Then φ satisfies the split property with respect to \mathcal{A}_L and \mathcal{A}_R .*

This theorem, combined with Definition 2.7 allows us to define the \mathbb{Z}_2 -index for reflection invariant Hamiltonians with unique gapped ground state.

Definition 3.3. Let $\Phi \in \mathcal{B}_f$ be a reflection invariant interaction which has a unique gapped ground state ω . By Theorem 3.2, ω satisfies the split property. Hence we obtain the \mathbb{Z}_2 -index σ_{ω} in Definition 2.7. In this setting, we denote this σ_{ω} by $\hat{\sigma}_{\Phi}$ and call it the \mathbb{Z}_2 -index associated to Φ .

Since $\hat{\sigma}_{\Phi}$ takes discrete values $\{-1, 1\}$, for a continuous path of interactions $\Phi(s)$, we would expect that $\hat{\sigma}_{\Phi(s)}$ is constant. We prove this in the setting of C^1 -classification.

Definition 3.4. We say the map $\Phi : [0, 1] \ni s \rightarrow \Phi(s) := \{\Phi(X; s)\}_{X \in \mathfrak{S}_{\mathbb{Z}}} \in \mathcal{B}_f$ is a C^1 -path of reflection invariant gapped interactions satisfying the *Condition B*, if there exist

- (i) numbers $M, R \in \mathbb{N}$, $g > 0$ and an increasing sequence $n_k \in \mathbb{N}$, $k = 1, 2, \dots$,
- (ii) C^1 -functions $a, b : [0, 1] \rightarrow \mathbb{R}$ such that $a(s) < b(s)$,

- (iii) a sequence of paths of interactions $\Psi_k : [0, 1] \ni s \rightarrow \Psi_k(s) := \{\Psi_k(X; s)\}_{X \in \mathfrak{S}_{\mathbb{Z}}} \in \mathcal{B}_f$, $k = 1, 2, \dots$,

and the following hold:

1. For each $X \in \mathfrak{S}_{\mathbb{Z}}$, the map $[0, 1] \ni s \rightarrow \Phi(X; s), \Psi_k(X; s) \in \mathcal{A}_X$ are continuous and piecewise C^1 . We denote by $\Phi'(X; s), \Psi'_k(X; s)$, the corresponding derivatives.
2. For each $s \in [0, 1]$, and $X \in \mathfrak{S}_{\mathbb{Z}}$ with $\text{diam}(X) \geq M$, we have $\Phi(X; s) = 0$.
3. For each $s \in [0, 1]$, and $k \in \mathbb{N}$, we have $\Psi_k(X; s) = 0$ unless $X \subset \Lambda_{n_k} \setminus \Lambda_{n_k - R}$.
4. Interactions are bounded as follows

$$C_1 := \sup_{s \in [0, 1]} \sup_{k \in \mathbb{N}} \sup_{X \in \mathfrak{S}_{\mathbb{Z}}} (\|\Phi(X; s)\| + |X| \|\Phi'(X; s)\| + \|\Psi_k(X; s)\| + |X| \|\Psi'_k(X; s)\|) < \infty. \quad (39)$$

5. For each $s \in [0, 1]$, there exists a unique $\tau^{\Phi(s)}$ -ground state φ_s .
6. For each $s \in [0, 1]$, $\Phi(s)$ is reflection-invariant.
7. For each $k \in \mathbb{N}$ and $s \in [0, 1]$, the spectrum $\sigma\left((H_{\Phi(s) + \Psi_k(s)})_{\Lambda_{n_k}}\right)$ of $(H_{\Phi(s) + \Psi_k(s)})_{\Lambda_{n_k}}$ is decomposed into two non-empty disjoint parts $\sigma\left((H_{\Phi(s) + \Psi_k(s)})_{\Lambda_{n_k}}\right) = \Sigma_1^{(k)}(s) \cup \Sigma_2^{(k)}(s)$ such that $\Sigma_1^{(k)}(s) \subset [a(s), b(s)]$, $\Sigma_2^{(k)}(s) \subset [b(s) + g, \infty)$ and the diameter of $\Sigma_1^{(k)}(s)$ converges to 0 as $k \rightarrow \infty$.

The interaction $\Psi_k(s)$ corresponds to a boundary condition. Note that it does not forbid an interaction between intervals $[-n, -n + R] \cap \mathbb{Z}$ and $[n - R, n] \cap \mathbb{Z}$. In particular, the periodic boundary condition is included in this framework. Also, note that we *do not* require that the boundary term $\Psi_k(s)$ to be reflection invariant.

By exactly the same way as in Proposition 3.5 of [O4], we can show the following.:

Proposition 3.5. *Let $\Phi : [0, 1] \ni s \rightarrow \Phi(s) := \{\Phi(X; s)\}_{X \in \mathfrak{S}_{\mathbb{Z}}} \in \mathcal{B}_f$ be a C^1 -path of reflection invariant gapped interactions satisfying the Condition B. Let φ_s be the unique $\tau^{\Phi(s)}$ -ground state, for each $s \in [0, 1]$. Then φ_0 and φ_1 are automorphic equivalent via an automorphism, which allows a reflection invariant decomposition.*

As a corollary of this proposition and Theorem 2.9, we obtain the following.

Theorem 3.6. *Let $\Phi : [0, 1] \ni s \rightarrow \Phi(s) := \{\Phi(X; s)\}_{X \in \mathfrak{S}_{\mathbb{Z}}} \in \mathcal{B}_f$ be a C^1 -path of reflection invariant gapped interactions satisfying the Condition B. Then we have $\hat{\sigma}_{\Phi(0)} = \hat{\sigma}_{\Phi(1)}$.*

Namely, the \mathbb{Z}_2 -index is invariant along the C^1 -path of reflection invariant gapped interactions, satisfying the Condition B.

4 The \mathbb{Z}_2 -index and the modular conjugation

In this section, we give a characterization of σ_ω from the point of view of Tomita-Takesaki modular theory. It will be used in Section 5, to prove that our index generalizes the index introduced in [PTBO1]. This also allows us to connect σ_ω with the Schmidt decomposition of Ω_ω .

First let us recall Tomita-Takesaki theory. See [BR1] or [T2] for more information. Let \mathcal{M} be a von Neumann algebra acting on a Hilbert space \mathcal{H} . Let Ω be a cyclic (i.e. $\mathcal{M}\Omega$ is dense in \mathcal{H}) and

separating (i.e., $x\Omega = 0$, $x \in \mathcal{M}$ implies $x = 0$) vector for \mathcal{M} . We define an anti-linear operator on \mathcal{H} with domain $\mathcal{M}\Omega$ by

$$Sx\Omega := x^*\Omega, \quad x \in \mathcal{M}. \quad (40)$$

It turns out that S is closable. (Proposition 2.5.9 of [BR1].) We denote the closure by the same symbol S . The operator S has a polar decomposition $S = J\Delta^{\frac{1}{2}}$ where J is an anti-unitary J called the modular conjugation associated to (\mathcal{M}, Ω) and Δ is a nonsingular positive operator called modular operator associated to (\mathcal{M}, Ω) . For the commutant \mathcal{M}' of \mathcal{M} , we have

$$J\Delta^{-\frac{1}{2}}x'\Omega = (x')^*\Omega, \quad x' \in \mathcal{M}'. \quad (41)$$

We also have

$$\Delta^{-\frac{1}{2}} = J\Delta^{\frac{1}{2}}J^*, \quad J^2 = \mathbb{I}, \quad \Delta\Omega = J\Omega = \Omega. \quad (42)$$

(Proposition 2.5.11 of [BR1].) The subspace $\mathcal{M}'\Omega$ is a core of $J\Delta^{-\frac{1}{2}}$. The Tomita-Takesaki theory states that

$$\Delta^{it}\mathcal{M}\Delta^{-it} = \mathcal{M}, \text{ for all } t \in \mathbb{R} \text{ and } J\mathcal{M}J^* = \mathcal{M}'. \quad (43)$$

From the first property, we may define a W^* -dynamics (i.e., σ -weak continuous one parameter group of automorphisms) $\sigma_t(x) := \Delta^{it}x\Delta^{-it}$, $t \in \mathbb{R}$, $x \in \mathcal{M}$ on \mathcal{M} . It is called the modular automorphisms associated to (\mathcal{M}, Ω) .

Set $\mathcal{D} := \{z \in \mathbb{C} \mid 0 < \Im z < 1\}$ and denote by $\bar{\mathcal{D}}$ its closure. For any $x, y \in \mathcal{M}$, there exists a bounded and continuous function $F_{x,y}$ on $\bar{\mathcal{D}}$ which is analytic on \mathcal{D} , satisfying

$$F_{x,y}(t) = \langle \Omega, \sigma_t(x)y\Omega \rangle, \quad F_{x,y}(t+i) = \langle \Omega, y\sigma_t(x)\Omega \rangle, \quad (44)$$

for all $t \in \mathbb{R}$. This condition is called the KMS-condition for the positive linear functional $\mathcal{M} \ni x \mapsto \langle \Omega, x\Omega \rangle$ on \mathcal{M} and the modular automorphisms are characterized as the unique W^* -dynamics which satisfies the KMS condition for this linear functional. (Theorem 1.2 VIII [T2].)

Now let us come back to our problem.

Lemma 4.1. *Let ω be a reflection invariant pure state on \mathcal{A} which satisfies the split property with respect to \mathcal{A}_R and \mathcal{A}_L . Let $(\mathcal{H}_\omega, \pi_\omega, \Omega_\omega, \Gamma_\omega)$ be a reflection-spilt representation associated to ω . Let s_ω be a projection in $B(\mathcal{H}_\omega)$ such that the support projection of Ω_ω in $\mathbb{I}_{\mathcal{H}_\omega} \otimes B(\mathcal{H}_\omega)$ is $\mathbb{I}_{\mathcal{H}_\omega} \otimes s_\omega$. Set $\mathcal{M} := s_\omega \otimes B(s_\omega\mathcal{H}_\omega)$, and $p_\omega = s_\omega \otimes s_\omega$. Then*

1. *the support projection of Ω_ω in $B(\mathcal{H}_\omega) \otimes \mathbb{I}_{\mathcal{H}_\omega}$ is $s_\omega \otimes \mathbb{I}_{\mathcal{H}_\omega}$,*
2. *$p_\omega\Omega_\omega = \Omega_\omega$,*
3. *the commutant \mathcal{M}' of \mathcal{M} in $p_\omega\hat{\mathcal{H}}_\omega$ is $\mathcal{M}' = B(s_\omega\mathcal{H}_\omega) \otimes s_\omega$,*
4. *Ω_ω is cyclic and separating for \mathcal{M} in $p_\omega\hat{\mathcal{H}}_\omega$,*
5. *$\Gamma_\omega p_\omega = p_\omega\Gamma_\omega$.*

Proof. To show 1., let s'_ω be a projection in $B(\mathcal{H}_\omega)$ such that the support projection of Ω_ω in $B(\mathcal{H}_\omega) \otimes \mathbb{I}_{\mathcal{H}_\omega}$ is $s'_\omega \otimes \mathbb{I}_{\mathcal{H}_\omega}$. From (19) and (7) with $A = \mathbb{I}_\mathcal{A}$, we have

$$\langle \Omega_\omega, ((1 - s_\omega) \otimes \mathbb{I}_{\mathcal{H}_\omega}) \Omega_\omega \rangle = \langle \Omega_\omega, \Gamma_\omega ((1 - s_\omega) \otimes \mathbb{I}_{\mathcal{H}_\omega}) \Gamma_\omega^* \Omega_\omega \rangle = \langle \Omega_\omega, (\mathbb{I}_{\mathcal{H}_\omega} \otimes (1 - s_\omega)) \Omega_\omega \rangle = 0. \quad (45)$$

Therefore, we have $1 - s_\omega \leq 1 - s'_\omega$. Similarly, we obtain $1 - s'_\omega \leq 1 - s_\omega$. Hence we obtain $s_\omega = s'_\omega$. 2. is clear from the definition. 3. follows from Tomita's commutant Theorem (See Theorem 5.9 of [T1]). Since Ω_ω is separating for \mathcal{M}' , it is cyclic for \mathcal{M} in $p_\omega\hat{\mathcal{H}}_\omega$, 4. (Proposition 2.5.3 [BR1].) 5. is from the definition of $p_\omega = s_\omega \otimes s_\omega$ and (19). \square

From 4. of Lemma 4.1, we can define modular conjugation J_ω and modular operator Δ_ω associated to $(\mathcal{M}, \Omega_\omega)$. Let us investigate their properties.

Lemma 4.2. *Let ω be a reflection invariant pure state on \mathcal{A} which satisfies the split property with respect to \mathcal{A}_R and \mathcal{A}_L . Let $(\mathcal{H}_\omega, \pi_\omega, \Omega_\omega, \Gamma_\omega)$ be a reflection-split representation associated to ω . Let s_ω be a projection in $B(\mathcal{H}_\omega)$ such that the support projection of Ω_ω in $\mathbb{I}_{\mathcal{H}_\omega} \otimes B(\mathcal{H}_\omega)$ is $\mathbb{I}_{\mathcal{H}_\omega} \otimes s_\omega$. Set $\mathcal{M} := s_\omega \otimes B(s_\omega \mathcal{H}_\omega)$, and $p_\omega = s_\omega \otimes s_\omega$. Let J_ω, Δ_ω be modular conjugation, modular operator on $p_\omega \hat{\mathcal{H}}_\omega$ associated to $(\mathcal{M}, \Omega_\omega)$. Let*

$$\Omega_\omega = \sum_{k \in \Lambda} \sqrt{\lambda_k} \xi_k \otimes \zeta_k \quad (46)$$

be a Schmidt decomposition of Ω_ω . Here Λ is a countable set and the sequence $\{\lambda_k\}_{k \in \Lambda} \subset \mathbb{R}_{>0}$ satisfies $\sum_{k \in \Lambda} \lambda_k = 1$. Furthermore, each of $\{\xi_k\}_{k \in \Lambda}$ and $\{\zeta_k\}_{k \in \Lambda}$ are orthonormal sets of \mathcal{H}_ω . We also define a density matrix ρ_ω by

$$\rho_\omega = \sum_{k \in \Lambda} \lambda_k |\zeta_k\rangle \langle \zeta_k|. \quad (47)$$

Then we have the following.:

1. Both of $\{\xi_k\}_{k \in \Lambda}$ and $\{\zeta_k\}_{k \in \Lambda}$ are orthonormal basis of $s_\omega \mathcal{H}_\omega$. There exists a unitary u on $s_\omega \mathcal{H}_\omega$ such that $\xi_k = u\zeta_k$, for each $k \in \Lambda$.
2. The action of the modular operator is given by

$$\Delta_\omega^{\frac{1}{2}}(s_\omega \otimes x) \Omega_\omega = \sum_{k \in \Lambda} u\zeta_k \otimes \rho_\omega^{\frac{1}{2}} x \zeta_k, \quad x \in B(s_\omega \mathcal{H}_\omega). \quad (48)$$

In particular, if the rank of ρ_ω is finite then we have

$$\Delta_\omega^{\frac{1}{2}}(s_\omega \otimes x) \Omega_\omega = \left(s_\omega \otimes \rho_\omega^{\frac{1}{2}} x \rho_\omega^{-\frac{1}{2}} \right) \Omega_\omega, \quad x \in B(s_\omega \mathcal{H}_\omega). \quad (49)$$

3. Let c be the complex conjugation on $s_\omega \mathcal{H}_\omega$ given by $c\zeta_k = \zeta_k$ for $k \in \Lambda$. Then we have

$$(s_\omega \otimes x) \Omega_\omega = \sum_{l \in \Lambda} u \rho_\omega^{\frac{1}{2}} c^* x^* c \zeta_l \otimes \zeta_l, \quad x \in B(s_\omega \mathcal{H}_\omega). \quad (50)$$

4. The adjoint of the modular conjugation on \mathcal{M} is given by

$$J_\omega(s_\omega \otimes x) J_\omega^* = u c^* x c u^* \otimes s_\omega, \quad x \in B(s_\omega \mathcal{H}_\omega). \quad (51)$$

Proof. By the definition of s_ω and 1. of Lemma 4.1, we see that each of $\{\xi_k\}_{k \in \Lambda}$ and $\{\zeta_k\}_{k \in \Lambda}$ are orthonormal basis of $s_\omega \mathcal{H}_\omega$. Therefore, there is a unitary u on $s_\omega \mathcal{H}_\omega$ such that $\xi_k = u\zeta_k$, for all $k \in \Lambda$. This u is given as $u = \sum_{k \in \Lambda} |\xi_k\rangle \langle \zeta_k|$ where the summation converges in the strong topology. This proves 1.

Let $\Lambda_n, n \in \mathbb{N}$ be an increasing sequence of finite subsets of Λ such that $\Lambda_n \nearrow \Lambda$. Set

$$Q_n := \sum_{k \in \Lambda_n} |\zeta_k\rangle \langle \zeta_k|. \quad (52)$$

For $x \in B(s_\omega \mathcal{H}_\omega)$ and $n \in \mathbb{N}$, set $x_n := Q_n x Q_n$. Note that the sequence $x_n \in B(s_\omega \mathcal{H}_\omega)$ approximates x in the σ -strong* topology.

We would like to specify the action of the modular operator \mathcal{L} . We claim

$$\Delta_{\omega}^{\frac{1}{2}}(s_{\omega} \otimes x) \Omega_{\omega} = \sum_{k \in \Lambda} u \zeta_k \otimes \rho_{\omega}^{\frac{1}{2}} x \zeta_k, \quad x \in B(s_{\omega} \mathcal{H}_{\omega}). \quad (53)$$

Note that the right hand side converges in norm because ρ_{ω} is in the trace class. To prove (53), we first specify the modular automorphism σ with respect to $(\mathcal{M}, \Omega_{\omega})$. We define a W^* -dynamics α on \mathcal{M} by

$$\alpha_t(s_{\omega} \otimes x) := s_{\omega} \otimes \text{Ad}(\rho_{\omega}^{it})(x), \quad t \in \mathbb{R}, x \in B(s_{\omega} \mathcal{H}_{\omega}), \quad (54)$$

and show that $\alpha = \sigma$. To do that, we recall the uniqueness of the W^* -dynamics which satisfies the KMS condition. Let φ be a state on \mathcal{M} given by

$$\varphi(s_{\omega} \otimes x) = \langle \Omega_{\omega}, (s_{\omega} \otimes x) \Omega_{\omega} \rangle, \quad x \in B(s_{\omega} \mathcal{H}_{\omega}). \quad (55)$$

Note that

$$\text{Tr}_{\mathcal{H}_{\omega}}(\rho_{\omega} x) = \langle \Omega_{\omega}, (s_{\omega} \otimes x) \Omega_{\omega} \rangle = \varphi(s_{\omega} \otimes x), \quad x \in B(s_{\omega} \mathcal{H}_{\omega}). \quad (56)$$

From this, we can see that φ is α -invariant. We show that α satisfies the KMS-condition for φ . This follows from the standard argument like in Proposition 5.3.7 of [BR2].: For any $x, y \in B(s_{\omega} \mathcal{H}_{\omega})$, and $n \in \mathbb{N}$, setting $x_n := Q_n x Q_n$, we may define an entire analytic function

$$F_{x_n, y}(z) := \langle \Omega_{\omega}, \alpha_z(s_{\omega} \otimes x_n)(s_{\omega} \otimes y) \Omega_{\omega} \rangle, \quad z \in \mathbb{C}, \quad (57)$$

because $\mathbb{R} \ni t \mapsto \alpha_t(s_{\omega} \otimes x_n) \in \mathcal{M}$ has an analytic continuation $\alpha_z(s_{\omega} \otimes x_n) = s_{\omega} \otimes \text{Ad}(\rho_{\omega}^{iz})(x_n) \in \mathcal{M}$, $z \in \mathbb{C}$. This $F_{x_n, y}(z)$ is bounded and continuous on $\bar{\mathcal{D}}$ and analytic on \mathcal{D} . Furthermore, it satisfies the boundary condition (44) i.e., we have

$$F_{x_n, y}(t) = \langle \Omega_{\omega}, \alpha_t(s_{\omega} \otimes x_n)(s_{\omega} \otimes y) \Omega_{\omega} \rangle, \quad F_{x_n, y}(t+i) = \langle \Omega_{\omega}, (s_{\omega} \otimes y) \alpha_t(s_{\omega} \otimes x_n) \Omega_{\omega} \rangle, \quad (58)$$

for all $t \in \mathbb{R}$. The second property holds because of (56) and the property of the trace.

By the α -invariance of φ and the σ -strong* convergence of x_n to x , (using the Cauchy-Schwartz inequality,) one can show from (58) that $F_{x_n, y}(t)$ and $F_{x_n, y}(t+i)$, as functions of $t \in \mathbb{R}$, are Cauchy sequence of continuous bounded functions on \mathbb{R} with respect to the uniform norm. Therefore, by the Phragmen-Lindelöf theorem, $\{F_{x_n, y}(z)\}_n$ is a Cauchy sequence of continuous bounded functions on $\bar{\mathcal{D}}$ with respect to the uniform norm. Therefore, $F_{x_n, y}(z)$ has a limit $F_{x, y}(z)$ on $z \in \bar{\mathcal{D}}$, which is bounded and continuous on $\bar{\mathcal{D}}$ and analytic on \mathcal{D} . We also have

$$F_{x, y}(t) = \lim_{n \rightarrow \infty} F_{x_n, y}(t) = \langle \Omega_{\omega}, \alpha_t(s_{\omega} \otimes x)(s_{\omega} \otimes y) \Omega_{\omega} \rangle, \quad (59)$$

$$F_{x, y}(t+i) = \lim_{n \rightarrow \infty} F_{x_n, y}(t+i) = \langle \Omega_{\omega}, (s_{\omega} \otimes y) \alpha_t(s_{\omega} \otimes x) \Omega_{\omega} \rangle, \quad (60)$$

for any $t \in \mathbb{R}$. Therefore, α satisfies the KMS condition for φ , and from the uniqueness, we get $\alpha = \sigma$. Now let us prove (53). For each $x \in B(s_{\omega} \mathcal{H}_{\omega})$, we again consider $x_n = Q_n x Q_n$, $n \in \mathbb{N}$. By the entire analyticity of $s_{\omega} \otimes x_n$ with respect to $\alpha = \sigma$, we have

$$\Delta_{\omega}^{\frac{1}{2}}(s_{\omega} \otimes x_n) \Omega_{\omega} = \alpha_{-\frac{i}{2}}(s_{\omega} \otimes x_n) \Omega_{\omega} = \left(s_{\omega} \otimes \text{Ad}(\rho_{\omega}^{\frac{1}{2}})(x_n) \right) \Omega_{\omega} = \sum_{k \in \Lambda} u \zeta_k \otimes \rho_{\omega}^{\frac{1}{2}} x_n \zeta_k. \quad (61)$$

(The first equation is a standard argument. See proof of Theorem 5.5 of [DJP] for example.) The left hand side of (61) converges to $\Delta_{\omega}^{\frac{1}{2}}(s_{\omega} \otimes x) \Omega_{\omega}$ as $n \rightarrow \infty$ because of

$$\left\| \Delta_{\omega}^{\frac{1}{2}}(s_{\omega} \otimes x_n) \Omega_{\omega} - \Delta_{\omega}^{\frac{1}{2}}(s_{\omega} \otimes x) \Omega_{\omega} \right\| = \left\| J_{\omega} \Delta_{\omega}^{\frac{1}{2}}(s_{\omega} \otimes x_n) \Omega_{\omega} - J_{\omega} \Delta_{\omega}^{\frac{1}{2}}(s_{\omega} \otimes x) \Omega_{\omega} \right\| = \|(s_{\omega} \otimes (x_n^* - x^*)) \Omega_{\omega}\|, \quad (62)$$

and the σ -strong*-convergence $x_n \rightarrow x$. The right hand of (61) converges to $\sum_{k \in \Lambda} u\zeta_k \otimes \rho_{\bar{\omega}}^{\frac{1}{2}} x \zeta_k$ because of

$$\left\| \sum_{k \in \Lambda} u\zeta_k \otimes \rho_{\bar{\omega}}^{\frac{1}{2}} (x_n - x) \zeta_k \right\|^2 = \sum_{k \in \Lambda} \left\| \rho_{\bar{\omega}}^{\frac{1}{2}} (x_n - x) \zeta_k \right\|^2, \quad (63)$$

and the σ -strong*-convergence $x_n \rightarrow x$. Hence we have proven (53).

Next we show β .

$$(s_{\omega} \otimes x) \Omega_{\omega} = \sum_{l \in \Lambda} u \rho_{\bar{\omega}}^{\frac{1}{2}} c^* x^* c \zeta_l \otimes \zeta_l, \quad x \in B(s_{\omega} \mathcal{H}_{\omega}). \quad (64)$$

The right hand side converges in norm. To prove (64), first we consider

$$(s_{\omega} \otimes Q_n x Q_m) \Omega_{\omega} = \sum_{k \in \Lambda_m} \sum_{l \in \Lambda_n} \sqrt{\lambda_k} u \zeta_k \otimes |\zeta_l\rangle \langle \zeta_l, x \zeta_k\rangle, \quad (65)$$

for $n, m \in \mathbb{N}$. Since we have $\langle \zeta_l, x \zeta_k \rangle = \langle c \zeta_l, x c \zeta_k \rangle = \langle c^* x c \zeta_k, \zeta_l \rangle$ by $c \zeta_k = \zeta_k$, we have

$$(65) = \sum_{l \in \Lambda_n} \sum_{k \in \Lambda_m} \sqrt{\lambda_k} |u \zeta_k\rangle \langle c^* x c \zeta_k, \zeta_l \rangle \otimes |\zeta_l\rangle = \sum_{l \in \Lambda_n} u Q_m \rho_{\bar{\omega}}^{\frac{1}{2}} Q_m c^* x^* c \zeta_l \otimes \zeta_l. \quad (66)$$

Hence we obtain

$$(s_{\omega} \otimes Q_n x Q_m) \Omega_{\omega} = \sum_{l \in \Lambda_n} u Q_m \rho_{\bar{\omega}}^{\frac{1}{2}} Q_m c^* x^* c \zeta_l \otimes \zeta_l. \quad (67)$$

Taking $m \rightarrow \infty$, and then $n \rightarrow \infty$, we obtain (64).

Next we consider the action of J_{ω} . We claim

$$J_{\omega} (s_{\omega} \otimes x) J_{\omega}^* = u c^* x c u^* \otimes s_{\omega}, \quad x \in B(s_{\omega} \mathcal{H}_{\omega}). \quad (68)$$

To prove this, note that

$$J_{\omega} (s_{\omega} \otimes x) \Omega_{\omega} = J_{\omega} J_{\omega} \Delta_{\bar{\omega}}^{\frac{1}{2}} (s_{\omega} \otimes x^*) \Omega_{\omega} = \Delta_{\bar{\omega}}^{\frac{1}{2}} (s_{\omega} \otimes x^*) \Omega_{\omega} = \sum_{k \in \Lambda} u \zeta_k \otimes \rho_{\bar{\omega}}^{\frac{1}{2}} x^* \zeta_k = \lim_{n \rightarrow \infty} \sum_{k \in \Lambda_n} u \zeta_k \otimes \rho_{\bar{\omega}}^{\frac{1}{2}} x^* \zeta_k, \quad (69)$$

for any $x \in B(s_{\omega} \mathcal{H}_{\omega})$, by 2. The right hand side converges in norm. Therefore, for any $m \in \mathbb{N}$, we have

$$(s_{\omega} \otimes Q_m) J_{\omega} (s_{\omega} \otimes x) \Omega_{\omega} = \lim_{n \rightarrow \infty} \sum_{k \in \Lambda_n} u \zeta_k \otimes Q_m \rho_{\bar{\omega}}^{\frac{1}{2}} x^* \zeta_k, \quad x \in B(s_{\omega} \mathcal{H}_{\omega}). \quad (70)$$

Note that as in the proof of (67), we have

$$\begin{aligned} \sum_{k \in \Lambda_n} u \zeta_k \otimes Q_m \rho_{\bar{\omega}}^{\frac{1}{2}} x^* \zeta_k &= \sum_{l \in \Lambda_m} \sum_{k \in \Lambda_n} u \zeta_k \otimes |\zeta_l\rangle \langle \zeta_l, \rho_{\bar{\omega}}^{\frac{1}{2}} x^* \zeta_k \rangle \\ &= \sum_{l \in \Lambda_m} \sum_{k \in \Lambda_n} u \zeta_k \langle c^* \rho_{\bar{\omega}}^{\frac{1}{2}} x^* c \zeta_k, \zeta_l \rangle \otimes \zeta_l = \sum_{l \in \Lambda_m} \sum_{k \in \Lambda_n} u |\zeta_k\rangle \langle \zeta_k | c^* x \rho_{\bar{\omega}}^{\frac{1}{2}} c \zeta_l \otimes \zeta_l = (u Q_n u^* \otimes s_{\omega}) \sum_{l \in \Lambda_m} u c^* x \rho_{\bar{\omega}}^{\frac{1}{2}} c \zeta_l \otimes \zeta_l \\ &= (u Q_n u^* \otimes s_{\omega}) \sum_{l \in \Lambda_m} \sqrt{\lambda_l} u c^* x c u^* u \zeta_l \otimes \zeta_l = (u Q_n u^* u c^* x c u^* \otimes s_{\omega}) \sum_{l \in \Lambda_m} \sqrt{\lambda_l} u \zeta_l \otimes \zeta_l = (u Q_n u^* u c^* x c u^* \otimes s_{\omega}) (s_{\omega} \otimes Q_m) \Omega_{\omega}, \end{aligned} \quad (71)$$

for any $x \in B(s_\omega \mathcal{H}_\omega)$. Taking $n \rightarrow \infty$, we obtain

$$(s_\omega \otimes Q_m) J_\omega (s_\omega \otimes x) \Omega_\omega = \lim_{n \rightarrow \infty} \sum_{k \in \Lambda_n} u \zeta_k \otimes Q_m \rho_\omega^{\frac{1}{2}} x^* \zeta_k = (uc^* x cu^* \otimes Q_m) \Omega_\omega, \quad (72)$$

for any $x \in B(s_\omega \mathcal{H}_\omega)$. Taking $m \rightarrow \infty$, we obtain

$$J_\omega (s_\omega \otimes x) J_\omega^* \Omega_\omega = J_\omega (s_\omega \otimes x) \Omega_\omega = (uc^* x cu^* \otimes s_\omega) \Omega_\omega, \quad x \in B(s_\omega \mathcal{H}_\omega), \quad (73)$$

for any $x \in B(s_\omega \mathcal{H}_\omega)$. Note that $J_\omega (s_\omega \otimes x) J_\omega^* \in \text{Ad}(J_\omega)(\mathcal{M}) = \mathcal{M}'$ and $uc^* x cu^* \otimes s_\omega \in \mathcal{M}'$. Since Ω_ω is separating for \mathcal{M}' , we have

$$J_\omega (s_\omega \otimes x) J_\omega^* = uc^* x cu^* \otimes s_\omega, \quad x \in B(s_\omega \mathcal{H}_\omega), \quad (74)$$

proving the claim. \square

The adjoint action of J_ω on \mathcal{M} introduces a \mathbb{Z}_2 -index.

Proposition 4.3. *Let ω be a reflection invariant pure state on \mathcal{A} which satisfies the split property with respect to \mathcal{A}_R and \mathcal{A}_L . Let $(\mathcal{H}_\omega, \pi_\omega, \Omega_\omega, \Gamma_\omega)$ be a reflection-split representation associated to ω . Let s_ω be a projection in $B(\mathcal{H}_\omega)$ such that the support projection of Ω_ω in $\mathbb{I}_{\mathcal{H}_\omega} \otimes B(\mathcal{H}_\omega)$ is $\mathbb{I}_{\mathcal{H}_\omega} \otimes s_\omega$. Set $\mathcal{M} := s_\omega \otimes B(s_\omega \mathcal{H}_\omega)$, and $p_\omega = s_\omega \otimes s_\omega$. Let J_ω be the modular conjugation on $p_\omega \mathcal{H}_\omega$ associated to $(\mathcal{M}, \Omega_\omega)$. Then we have the following:*

1. *There exists an anti-unitary $\theta : s_\omega \mathcal{H}_\omega \rightarrow s_\omega \mathcal{H}_\omega$ such that*

$$J_\omega (s_\omega \otimes x) J_\omega^* = \theta x \theta^* \otimes s_\omega, \quad (75)$$

for all $x \in B(s_\omega \mathcal{H}_\omega)$.

2. *For any anti-unitary $\theta : s_\omega \mathcal{H}_\omega \rightarrow s_\omega \mathcal{H}_\omega$ satisfying (75), we have*

$$J_\omega (x \otimes s_\omega) J_\omega^* = s_\omega \otimes \theta x \theta^*, \quad (76)$$

for all $x \in B(s_\omega \mathcal{H}_\omega)$.

3. *There exists a $\kappa_\omega \in \{\pm 1\}$ satisfying $\theta^2 = \kappa_\omega s_\omega$ for any anti-unitary $\theta : s_\omega \mathcal{H}_\omega \rightarrow s_\omega \mathcal{H}_\omega$ satisfying (75).*

Proof. Let Δ_ω be the modular operator of $(\mathcal{M}, \Omega_\omega)$. From 5. of Lemma 4.1, $\tilde{\Gamma} := \Gamma_\omega p_\omega$ defines a unitary operator on $p_\omega \mathcal{H}_\omega$. Note that $\tilde{\Gamma} \Omega_\omega = \Omega_\omega$, because of Lemma 4.1 2. and Definition 2.2 4. We claim that

$$J_\omega \tilde{\Gamma} = \tilde{\Gamma} J_\omega. \quad (77)$$

To see this, we recall (40) and (41). For any $x \in \mathcal{M}$, we have

$$\tilde{\Gamma} J_\omega \Delta_\omega^{\frac{1}{2}} x \Omega_\omega = \tilde{\Gamma} x^* \Omega_\omega = \tilde{\Gamma} x^* \tilde{\Gamma}^* \tilde{\Gamma} \Omega_\omega = \tilde{\Gamma} x^* \tilde{\Gamma}^* \Omega_\omega. \quad (78)$$

Note that from (19), the element $\tilde{\Gamma} x^* \tilde{\Gamma}^*$ belongs to \mathcal{M}' . Therefore, from (41), we have

$$(78) = J_\omega \Delta_\omega^{-\frac{1}{2}} \tilde{\Gamma} x \tilde{\Gamma}^* \Omega_\omega = J_\omega \Delta_\omega^{-\frac{1}{2}} \tilde{\Gamma} x \Omega_\omega. \quad (79)$$

Since $\mathcal{M}\Omega_\omega$ is a core of $J_\omega\Delta_\omega^{\frac{1}{2}}$ and $\mathcal{M}'\Omega_\omega$ is a core of $J_\omega\Delta_\omega^{-\frac{1}{2}}$, this means

$$\tilde{\Gamma}J_\omega\Delta_\omega^{\frac{1}{2}} = J_\omega\Delta_\omega^{-\frac{1}{2}}\tilde{\Gamma} = J_\omega\tilde{\Gamma}\tilde{\Gamma}^*\Delta_\omega^{-\frac{1}{2}}\tilde{\Gamma}. \quad (80)$$

By the uniqueness of the polar decomposition, we obtain $\tilde{\Gamma}J_\omega = J_\omega\tilde{\Gamma}$, proving the claim.

Next we note that there are anti*-automorphisms $\Theta_{L \rightarrow R}, \Theta_{R \rightarrow L}$ on $B(s_\omega\mathcal{H}_\omega)$ such that

$$J_\omega(s_\omega \otimes x)J_\omega^* = \Theta_{R \rightarrow L}(x) \otimes s_\omega, \quad (81)$$

$$J_\omega(x \otimes s_\omega)J_\omega^* = s_\omega \otimes \Theta_{L \rightarrow R}(x), \quad (82)$$

for any $x \in B(s_\omega\mathcal{H}_\omega)$. This is because of the Tomita-Takesaki theory, i.e., $J_\omega\mathcal{M}J_\omega^* = \mathcal{M}'$ and $J_\omega\mathcal{M}'J_\omega^* = \mathcal{M}$. Note that $\Theta_{L \rightarrow R} \circ \Theta_{R \rightarrow L} = \text{id} = \Theta_{R \rightarrow L} \circ \Theta_{L \rightarrow R}$, because of $J_\omega^2 = s_\omega$. By (19) and (77), this $\Theta_{L \rightarrow R}$ and $\Theta_{R \rightarrow L}$ coincides. For any $x \in B(s_\omega\mathcal{H}_\omega)$, we have

$$\begin{aligned} \Theta_{R \rightarrow L}(x) \otimes s_\omega &= (\text{Ad } J_\omega)(s_\omega \otimes x) = (\text{Ad } J_\omega\tilde{\Gamma})(x \otimes s_\omega) = (\text{Ad } \tilde{\Gamma}J_\omega)(x \otimes s_\omega) \\ &= \text{Ad } \tilde{\Gamma}(s_\omega \otimes \Theta_{L \rightarrow R}(x)) = \Theta_{L \rightarrow R}(x) \otimes s_\omega. \end{aligned} \quad (83)$$

Hence we have $\Theta_{R \rightarrow L}(x) = \Theta_{L \rightarrow R}(x)$ for any $x \in B(s_\omega\mathcal{H}_\omega)$.

Now let us prove 1.-3. of the Lemma. 1. is shown in Lemma 4.2 4, as $\theta = uc^*$. To prove the second and the third statement, let $\theta : s_\omega\mathcal{H}_\omega \rightarrow s_\omega\mathcal{H}_\omega$ be any anti-unitary such that

$$J_\omega(s_\omega \otimes x)J_\omega^* = \theta x \theta^* \otimes s_\omega, \quad x \in B(s_\omega\mathcal{H}_\omega). \quad (84)$$

From this and (81), we obtain

$$\Theta_{R \rightarrow L}(x) = \Theta_{L \rightarrow R}(x) = \theta x \theta^*, \quad x \in B(s_\omega\mathcal{H}_\omega). \quad (85)$$

Therefore, from (82),

$$J_\omega(x \otimes s_\omega)J_\omega^* = s_\omega \otimes \theta x \theta^*, \quad x \in B(s_\omega\mathcal{H}_\omega). \quad (86)$$

This proves the second statement.

Furthermore, we have

$$s_\omega \otimes x = J_\omega^2(s_\omega \otimes x)(J_\omega^*)^2 = J_\omega(\theta x \theta^* \otimes s_\omega)J_\omega^* = s_\omega \otimes \theta^2 x (\theta^*)^2, \quad (87)$$

for any $x \in B(s_\omega\mathcal{H}_\omega)$. This means $\theta^2 = \tilde{\kappa}_\theta s_\omega$ with some $\tilde{\kappa}_\theta \in \mathbb{T}$. Then by the anti-linearity of θ , we have

$$\tilde{\kappa}_\theta \theta = \theta^2 \theta = \theta \theta^2 = \theta \tilde{\kappa}_\theta = \bar{\tilde{\kappa}}_\theta \theta. \quad (88)$$

This means $\tilde{\kappa}_\theta$ is real, namely $\tilde{\kappa}_\theta = \pm 1$.

This $\tilde{\kappa}_\theta$ is independent of the choice of θ satisfying (75) for if θ_1 is another such anti-unitary, we have

$$\theta_1 x \theta_1^* \otimes s_\omega = J_\omega(s_\omega \otimes x)J_\omega^* = \theta x \theta^* \otimes s_\omega, \quad (89)$$

for all $x \in B(s_\omega\mathcal{H}_\omega)$. Hence $\theta^* \theta_1$ is a unitary operator on $s_\omega\mathcal{H}_\omega$ which commutes with any $x \in B(s_\omega\mathcal{H}_\omega)$, i.e., $\theta = c \theta_1$ for some $c \in \mathbb{T}$. We then have

$$\tilde{\kappa}_\theta s_\omega = \theta^2 = c \theta_1 c \theta_1 = c \bar{c} \theta_1 \theta_1 = \tilde{\kappa}_{\theta_1} s_\omega, \quad (90)$$

and get $\tilde{\kappa}_\theta = \tilde{\kappa}_{\theta_1} =: \kappa_\omega$. This proves the third statement. \square

The sign κ_ω coincides with our \mathbb{Z}_2 -index σ_ω .

Theorem 4.4. *Let ω be a reflection invariant pure state on \mathcal{A} which satisfies the split property with respect to \mathcal{A}_R and \mathcal{A}_L . Let $(\mathcal{H}_\omega, \pi_\omega, \Omega_\omega, \Gamma_\omega)$ be a reflection-split representation associated to ω . Let s_ω be a projection in $B(\mathcal{H}_\omega)$ such that the support projection of Ω_ω in $\mathbb{I}_{\mathcal{H}_\omega} \otimes B(\mathcal{H}_\omega)$ is $\mathbb{I}_{\mathcal{H}_\omega} \otimes s_\omega$. Set $\mathcal{M} := s_\omega \otimes B(s_\omega \mathcal{H}_\omega)$, and $p_\omega = s_\omega \otimes s_\omega$. Let J_ω be the modular conjugation on $p_\omega \mathcal{H}_\omega$ associated to $(\mathcal{M}, \Omega_\omega)$. Let σ_ω be the \mathbb{Z}_2 -index associated to ω in Definition 2.7 and κ_ω the \mathbb{Z}_2 -index associated to ω in Proposition 4.3. Then we have $\kappa_\omega = \sigma_\omega$.*

Proof. We use the notation used in Lemma 4.2. From $\Gamma_\omega \Omega_\omega = \Omega_\omega$, we obtain $u = \sigma_\omega c^* u^* c$: first we have

$$\begin{aligned} \sum_{k \in \Lambda} u \rho_\omega^{\frac{1}{2}} \zeta_k \otimes \zeta_k &= \sum_{k \in \Lambda} \sqrt{\lambda_k} u \zeta_k \otimes \zeta_k = \Omega_\omega = \Gamma_\omega \Omega_\omega = \sum_{k \in \Lambda} \sqrt{\lambda_k} \Gamma_\omega (u \zeta_k \otimes \zeta_k) = \sum_{k \in \Lambda} \sqrt{\lambda_k} \sigma_\omega (\zeta_k \otimes u \zeta_k) \\ &= \sigma_\omega (u^* \otimes u) \sum_{k \in \Lambda} \sqrt{\lambda_k} (u \zeta_k \otimes \zeta_k) = \sigma_\omega (u^* \otimes s_\omega) (s_\omega \otimes u) \Omega_\omega = \sigma_\omega (u^* \otimes s_\omega) \sum_{l \in \Lambda} u \rho_\omega^{\frac{1}{2}} c^* u^* c \zeta_l \otimes \zeta_l \\ &= \sigma_\omega \sum_{l \in \Lambda} \rho_\omega^{\frac{1}{2}} c^* u^* c \zeta_l \otimes \zeta_l. \end{aligned} \quad (91)$$

Here we used Theorem 2.6 and Lemma 4.2 3. From this we obtain

$$u \rho_\omega^{\frac{1}{2}} \zeta_k = \sigma_\omega \rho_\omega^{\frac{1}{2}} c^* u^* c \zeta_k, \quad (92)$$

for all $k \in \Lambda$. Hence we have

$$u \rho_\omega^{\frac{1}{2}} = \sigma_\omega c^* u^* c c^* u \rho_\omega^{\frac{1}{2}} c^* u^* c. \quad (93)$$

By the uniqueness of the polar decomposition, we obtain the claim

$$u = \sigma_\omega c^* u^* c. \quad (94)$$

Now we are ready to complete the proof of the Theorem. By Proposition 4.3 and Lemma 4.2 4., we have $(uc)^2 = (uc^*)^2 = \kappa_\omega s_\omega$. From this and (94), we have

$$\kappa_\omega = \langle (uc)^2 \zeta_k, \zeta_k \rangle = \langle ucuc \zeta_k, \zeta_k \rangle = \langle (uc)^* \zeta_k, uc \zeta_k \rangle = \langle c^* u^* c \zeta_k, u \zeta_k \rangle = \langle \sigma_\omega u \zeta_k, u \zeta_k \rangle = \sigma_\omega, \quad (95)$$

for any $k \in \Lambda$. This completes the proof. \square

Remark 4.5. From the proof, we see that one way to derive the index σ_ω for concrete state ω is to consider the Schmidt decomposition and calculate u . Using (94), we can obtain σ_ω .

5 \mathbb{Z}_2 -index in Matrix Product States

In this section, we prove that the \mathbb{Z}_2 -index σ_ω for a matrix product state ω is the same as the \mathbb{Z}_2 -index found in [PTBO1]. First let us recall known facts on matrix product states. Let $k \in \mathbb{N}$ be a number and $\mathbf{v} = (v_\mu)_{\mu=1, \dots, d} \in M_k^{\times d}$ a d -tuple of $k \times k$ matrices. For each $l \in \mathbb{N}$, we set

$$\mathcal{K}_l(\mathbf{v}) := \text{span} \{ v_{\mu_0} v_{\mu_1} \dots v_{\mu_{l-1}} \mid (\mu_0, \mu_1, \dots, \mu_{l-1}) \subset \{1, \dots, d\}^{\times l} \}. \quad (96)$$

We say \mathbf{v} is primitive if $\mathcal{K}_l(\mathbf{v}) = M_k$ for l large enough. We denote by $\text{Prim}_u(d, k)$ the set of all primitive d -tuples \mathbf{v} of $k \times k$ matrices which are normalized, i.e.,

$$\sum_{\mu=1, \dots, d} v_\mu v_\mu^* = 1.$$

For $\mathbf{v} \in \text{Prim}_u(d, k)$, there exists a unique $T_{\mathbf{v}}$ -invariant state $\hat{\rho}_{\mathbf{v}}$, and it is faithful. (See [W] for example.) We denote the density matrix corresponding to $\hat{\rho}_{\mathbf{v}}$ by $\rho_{\mathbf{v}}$. Each $\mathbf{v} \in \text{Prim}_u(d, k)$ generates a translationally invariant state $\omega_{\mathbf{v}}$ by

$$\omega_{\mathbf{v}} \left(\bigotimes_{i=0}^{l-1} |\psi_{\mu_i}\rangle \langle \psi_{\nu_i}| \right) = \hat{\rho}_{\mathbf{v}} \left(v_{\mu_0} \cdots v_{\mu_{l-1}} v_{\nu_{l-1}}^* \cdots v_{\nu_0}^* \right), \quad \mu_i, \nu_i = 1, \dots, d, \quad i = 0, \dots, l-1, \quad l \in \mathbb{N}. \quad (97)$$

A translationally invariant state which has this representation is called a matrix product state. This representation is unique up to unitary and phase [FNW2]: If both of $\mathbf{v}^{(1)} \in \text{Prim}_u(d, k_1)$ and $\mathbf{v}^{(2)} \in \text{Prim}_u(d, k_2)$ generate the same matrix product state, then $k_1 = k_2$ and there exist a unitary $U : \mathbb{C}^{k_1} \rightarrow \mathbb{C}^{k_2}$ and $e^{i\theta} \in \mathbb{T}$ such that

$$U v_{\mu}^{(1)} = e^{i\theta} v_{\mu}^{(2)} U, \quad \mu = 1, \dots, d. \quad (98)$$

Let ω be a reflection invariant matrix product state generated by $\mathbf{v} \in \text{Prim}_u(d, k)$. It is a unique ground state of some translation invariant finite range interaction. i.e., there is an interaction $\Phi_{\mathbf{v}}$ given by some fixed local positive element $h_{\mathbf{v}} \in \mathcal{A}_{[0, m-1]}$ with some $m \in \mathbb{N}$ as

$$\Phi_{\mathbf{v}}(X) := \begin{cases} \beta_x(h_{\mathbf{v}}), & \text{if } X = [x, x+m-1] \cap \mathbb{Z} \text{ for some } x \in \mathbb{Z} \\ 0, & \text{otherwise} \end{cases} \quad (99)$$

for each $X \in \mathfrak{S}_{\mathbb{Z}}$ and ω is a unique $\tau^{\Phi_{\mathbf{v}}}$ -ground state. (See [FNW] and [O3] Corollary 5.6 [O1] Theorem 1.18, Lemma 3.25.) For this interaction $h_{\mathbf{v}}$, $1 - h_{\mathbf{v}}$ is equal to the support projection of $\omega|_{\mathcal{A}_{[0, m-1]}}$. (Note that $\omega|_{\mathcal{A}_L} = \Xi_L(\hat{\rho}_{\mathbf{v}})$ with the Ξ_L in Lemma 3.14 of [O1] and the $T_{\mathbf{v}}$ -invariant state $\hat{\rho}_{\mathbf{v}}$. From the proof of Lemma 3.19 of [O1] equation (48), we see that $1 - h_{\mathbf{v}}$ is equal to the support projection of $\omega|_{\mathcal{A}_{[0, m-1]}}$. Note that primitive \mathbf{v} belongs to ClassA, Remark 1.16 of [O1]). Therefore, from the reflection invariance of ω , $\Phi_{\mathbf{v}}$ is reflection invariant. The Hamiltonian given by this interaction is frustration-free, i.e., for each finite interval I with $|I| \geq m$, the local Hamiltonian $(H_{\Phi_{\mathbf{v}}})_I$ has a nontrivial kernel, which is the ground state space of $(H_{\Phi_{\mathbf{v}}})_I$. We denote by $G_{I, \mathbf{v}}$, the orthogonal projection onto this kernel. By Lemma 3.19 of [O1], and its proof (equation (48)), the support of the restriction $\omega|_{\mathcal{A}_I}$ is equal to $G_{I, \mathbf{v}}$ and there exists some constant $d_{\mathbf{v}} > 0$ such that

$$\psi \leq d_{\mathbf{v}} \cdot \omega, \quad (100)$$

for any frustration free state ψ on \mathcal{A}_R , i.e., a state ψ satisfying $\psi(\beta_x(h_{\mathbf{v}})) = 0$ for any $0 \leq x \in \mathbb{Z}$. (See proof of Lemma 2.3 of [O2].)

Now let us come back to our problem. With the analogous argument as in [PTBO1], we obtain the following. See [Tas2] for a nice description.

Lemma 5.1. *Let ω be a reflection invariant matrix product state generated by $\mathbf{v} \in \text{Prim}_u(d, k)$. Let $\hat{\rho}_{\mathbf{v}}$ be the $T_{\mathbf{v}}$ -invariant state given by a density matrix $\rho_{\mathbf{v}}$. Then there exist $e^{it} \in \mathbb{T}$ and $\theta : \mathbb{C}^k \rightarrow \mathbb{C}^k$ an anti-unitary such that*

$$v_{\mu} = e^{it} \rho_{\mathbf{v}}^{-\frac{1}{2}} \theta v_{\mu}^* \theta^* \rho_{\mathbf{v}}^{\frac{1}{2}}, \quad \mu = 1, \dots, d. \quad (101)$$

For any $e^{it} \in \mathbb{T}$ and $\theta : \mathbb{C}^k \rightarrow \mathbb{C}^k$ an anti-unitary satisfying (101), we have

$$\theta^2 = \zeta_{\omega} \mathbb{I}, \quad (102)$$

with some $\zeta_{\omega} \in \{\pm 1\}$. The value ζ_{ω} does not depend on the choice of \mathbf{v} , $e^{it} \in \mathbb{T}$ and $\theta : \mathbb{C}^k \rightarrow \mathbb{C}^k$.

Definition 5.2. By this Lemma, we define a \mathbb{Z}_2 -index ζ_{ω} .

Proof. Let $\rho_{\mathbf{v}} = \sum_{j=1}^k \lambda_j |\xi_j\rangle \langle \xi_j|$ be the spectral decomposition of $\rho_{\mathbf{v}}$, where $\lambda_j > 0$ and $\{\xi_j\}_{j=1}^d$ is an orthonormal basis of \mathbb{C}^k . Let $c : \mathbb{C}^k \rightarrow \mathbb{C}^k$ be the complex conjugation such that $c\xi_j = \xi_j$ for all $j = 1, \dots, k$. Note that $c^* = c$.

Set

$$\bar{v}_\mu := cv_\mu c, \quad \tilde{v}_\mu := \rho_{\mathbf{v}}^{-\frac{1}{2}} (\bar{v}_\mu)^* \rho_{\mathbf{v}}^{\frac{1}{2}}, \quad \mu = 1, \dots, d. \quad (103)$$

We claim $\tilde{\mathbf{v}} \in \text{Prim}_u(d, k)$ and it generates ω . Since $\mathcal{K}_l(\mathbf{v}) = \mathbb{M}_k$ for l large enough, $\mathcal{K}_l(\tilde{\mathbf{v}}) = \mathbb{M}_k$ for l large enough. Hence $\tilde{\mathbf{v}}$ is primitive. Using $c\rho_{\mathbf{v}}c = \rho_{\mathbf{v}}$ and $\sum_{\mu} v_\mu^* \rho_{\mathbf{v}} v_\mu = \rho_{\mathbf{v}}$, we have

$$\sum_{\mu} \tilde{v}_\mu \tilde{v}_\mu^* = \sum_{\mu} \rho_{\mathbf{v}}^{-\frac{1}{2}} cv_\mu^* c \rho_{\mathbf{v}} cv_\mu c \rho_{\mathbf{v}}^{-\frac{1}{2}} = \sum_{\mu} \rho_{\mathbf{v}}^{-\frac{1}{2}} cv_\mu^* \rho_{\mathbf{v}} v_\mu c \rho_{\mathbf{v}}^{-\frac{1}{2}} = \rho_{\mathbf{v}}^{-\frac{1}{2}} c \rho_{\mathbf{v}} c \rho_{\mathbf{v}}^{-\frac{1}{2}} = \mathbb{I}_k \quad (104)$$

Hence we have $\tilde{\mathbf{v}} \in \text{Prim}_u(d, k)$. The state $\hat{\rho}_{\tilde{\mathbf{v}}}$ is $T_{\tilde{\mathbf{v}}}$ -invariant because

$$\sum_{\mu} \tilde{v}_\mu^* \rho_{\tilde{\mathbf{v}}} \tilde{v}_\mu = \sum_{\mu} \rho_{\tilde{\mathbf{v}}}^{\frac{1}{2}} cv_\mu c \rho_{\mathbf{v}}^{-\frac{1}{2}} \rho_{\mathbf{v}} \rho_{\mathbf{v}}^{-\frac{1}{2}} cv_\mu^* c \rho_{\mathbf{v}}^{\frac{1}{2}} = \rho_{\mathbf{v}}, \quad (105)$$

from $\sum_{\mu} v_\mu v_\mu^* = 1$. Now we show that $\tilde{\mathbf{v}}$ generates ω . For any $l \in \mathbb{N}$ and $\mu_i, \nu_i = 1, \dots, d$, $i = 0, \dots, l-1$, from the reflection invariance and translation invariance, we have

$$\begin{aligned} \omega \left(\bigotimes_{i=0}^{l-1} |\psi_{\mu_i}\rangle \langle \psi_{\nu_i}| \right) &= \omega \circ \gamma \left(\bigotimes_{i=0}^{l-1} |\psi_{\mu_i}\rangle \langle \psi_{\nu_i}| \right) = \omega \left(\bigotimes_{i=-l}^{-1} |\psi_{\mu_{l-i-1}}\rangle \langle \psi_{\nu_{l-i-1}}| \right) = \omega \left(\bigotimes_{i=0}^{l-1} |\psi_{\mu_{l-i-1}}\rangle \langle \psi_{\nu_{l-i-1}}| \right) \\ &= \hat{\rho}_{\mathbf{v}} \left(v_{\mu_{l-1}} v_{\mu_{l-2}} \cdots v_{\mu_1} v_{\mu_0} v_{\nu_0}^* v_{\nu_1}^* \cdots v_{\nu_{l-1}}^* \right) = \sum_{j=1}^k \lambda_j \left\langle \xi_j, v_{\mu_{l-1}} v_{\mu_{l-2}} \cdots v_{\mu_1} v_{\mu_0} v_{\nu_0}^* v_{\nu_1}^* \cdots v_{\nu_{l-1}}^* \xi_j \right\rangle \end{aligned} \quad (106)$$

Note that

$$\begin{aligned} \left\langle \xi_j, v_{\mu_{l-1}} v_{\mu_{l-2}} \cdots v_{\mu_1} v_{\mu_0} v_{\nu_0}^* v_{\nu_1}^* \cdots v_{\nu_{l-1}}^* \xi_j \right\rangle &= \left\langle c\xi_j, v_{\mu_{l-1}} v_{\mu_{l-2}} \cdots v_{\mu_1} v_{\mu_0} v_{\nu_0}^* v_{\nu_1}^* \cdots v_{\nu_{l-1}}^* c\xi_j \right\rangle \\ &= \left\langle \xi_j, cv_{\nu_{l-1}} v_{\nu_{l-2}} \cdots v_{\nu_1} v_{\nu_0} v_{\mu_0}^* v_{\mu_1}^* \cdots v_{\mu_{l-1}}^* c\xi_j \right\rangle = \left\langle \xi_j, \bar{v}_{\nu_{l-1}} \bar{v}_{\nu_{l-2}} \cdots \bar{v}_{\nu_1} \bar{v}_{\nu_0} \bar{v}_{\mu_0}^* \bar{v}_{\mu_1}^* \cdots \bar{v}_{\mu_{l-1}}^* \xi_j \right\rangle. \end{aligned} \quad (107)$$

Substituting this to (106), we have

$$\begin{aligned} (106) &= \hat{\rho}_{\mathbf{v}} \left(\bar{v}_{\nu_{l-1}} \bar{v}_{\nu_{l-2}} \cdots \bar{v}_{\nu_1} \bar{v}_{\nu_0} \bar{v}_{\mu_0}^* \bar{v}_{\mu_1}^* \cdots \bar{v}_{\mu_{l-1}}^* \right) = \text{Tr} \left(\rho_{\tilde{\mathbf{v}}}^{\frac{1}{2}} \left(\bar{v}_{\nu_{l-1}} \bar{v}_{\nu_{l-2}} \cdots \bar{v}_{\nu_1} \bar{v}_{\nu_0} \rho_{\mathbf{v}}^{-\frac{1}{2}} \rho_{\mathbf{v}} \rho_{\mathbf{v}}^{-\frac{1}{2}} \bar{v}_{\mu_0}^* \bar{v}_{\mu_1}^* \cdots \bar{v}_{\mu_{l-1}}^* \right) \rho_{\tilde{\mathbf{v}}}^{\frac{1}{2}} \right) \\ &= \hat{\rho}_{\tilde{\mathbf{v}}} \left(\tilde{v}_{\mu_0} \tilde{v}_{\mu_1} \cdots \tilde{v}_{\mu_{l-1}} \tilde{v}_{\nu_{l-1}}^* \tilde{v}_{\nu_{l-2}}^* \cdots \tilde{v}_{\nu_1}^* \tilde{v}_{\nu_0}^* \right). \end{aligned} \quad (108)$$

Hence $\tilde{\mathbf{v}}$ generates ω , proving the claim.

Now, as both of \mathbf{v} and $\tilde{\mathbf{v}}$ generates same state ω , by the uniqueness (98), there exist a unitary U on \mathbb{C}^k and $e^{it} \in \mathbb{T}$ such that

$$Uv_\mu = e^{it} \tilde{v}_\mu U, \quad \mu = 1, \dots, d. \quad (109)$$

From the fact that $\hat{\rho}_{\mathbf{v}}$ is $T_{\tilde{\mathbf{v}}}$ -invariant and (109), we see that the state $\hat{\rho}_{\mathbf{v}} \circ \text{Ad } U$ is $T_{\mathbf{v}}$ -invariant. By the uniqueness of $T_{\mathbf{v}}$ -invariant state (by $\mathbf{v} \in \text{Prim}_u(d, k)$), we get

$$U\rho_{\mathbf{v}}U^* = \rho_{\mathbf{v}}. \quad (110)$$

Set $\theta := U^*c : \mathbb{C}^k \rightarrow \mathbb{C}^k$ an anti-unitary operator. From (109) and the definition of $\tilde{\mathbf{v}}$, and (110), we obtain (101):

$$v_\mu = e^{it} U^* \tilde{v}_\mu U = e^{it} U^* \rho_{\mathbf{v}}^{-\frac{1}{2}} (\bar{v}_\mu)^* \rho_{\mathbf{v}}^{\frac{1}{2}} U = e^{it} \rho_{\mathbf{v}}^{-\frac{1}{2}} (U^*c) v_\mu^* (U^*c)^* \rho_{\mathbf{v}}^{\frac{1}{2}} = e^{it} \rho_{\mathbf{v}}^{-\frac{1}{2}} \theta v_\mu^* \theta^* \rho_{\mathbf{v}}^{\frac{1}{2}}. \quad (111)$$

Now for any $e^{it_0} \in \mathbb{T}$ and $\theta_0 : \mathbb{C}^k \rightarrow \mathbb{C}^k$ an anti-unitary satisfying

$$v_\mu = e^{it_0} \rho_v^{-\frac{1}{2}} \theta_0 v_\mu^* \theta_0^* \rho_v^{\frac{1}{2}}, \quad \mu = 1, \dots, d, \quad (112)$$

we show $\theta_0^2 = \mathbb{I}$ or $\theta_0^2 = -\mathbb{I}$. Taking adjoint of (101), we have

$$v_\mu^* = e^{-it_0} \rho_v^{\frac{1}{2}} \theta_0 v_\mu \theta_0^* \rho_v^{-\frac{1}{2}}. \quad (113)$$

Substituting this to (101), we obtain

$$v_\mu = e^{it_0} \rho_v^{-\frac{1}{2}} \theta_0 e^{-it_0} \rho_v^{\frac{1}{2}} \theta_0 v_\mu \theta_0^* \rho_v^{-\frac{1}{2}} \theta_0^* \rho_v^{\frac{1}{2}} = e^{2it_0} \rho_v^{-\frac{1}{2}} \theta_0 \rho_v^{\frac{1}{2}} \theta_0 v_\mu \theta_0^* \rho_v^{-\frac{1}{2}} \theta_0^* \rho_v^{\frac{1}{2}} \quad (114)$$

Since \mathbf{v} is primitive, this means $e^{2it_0} = 1$ and $\rho_v^{-\frac{1}{2}} \theta_0 \rho_v^{\frac{1}{2}} \theta_0 = b\mathbb{I}$ for some $b \in \mathbb{C}$. Decomposing $b = e^{is}|b|$ with $e^{is} \in \mathbb{T}$, we have

$$\theta_0^2 \left(\theta_0^* \rho_v^{\frac{1}{2}} \theta_0 \right) = e^{is} |b| \rho_v^{\frac{1}{2}}. \quad (115)$$

By the uniqueness of the polar decomposition and the faithfulness of ρ_v , we get $\theta_0^2 = e^{is}\mathbb{I}$. But then

$$e^{is} \theta_0 = \theta_0^2 \theta_0 = \theta_0^3 = \theta_0 \theta_0^2 = \theta_0 e^{is} = e^{-is} \theta_0. \quad (116)$$

Therefore, e^{is} is real and we get that $\theta_0^2 = \mathbb{I}$ or $\theta_0^2 = -\mathbb{I}$.

To prove the independence of this sign of \mathbf{v} , e^{it} and θ , let $\omega \in \text{Prim}_u(d, k')$ be a generator of ω , with a T_ω -invariant state $\hat{\rho}_\omega$ given by a density matrix ρ_ω . Let $e^{iu} \in \mathbb{T}$ and $\xi : \mathbb{C}^{k'} \rightarrow \mathbb{C}^{k'}$ an anti-unitary such that

$$\omega_\mu = e^{iu} \rho_\omega^{-\frac{1}{2}} \xi \omega_\mu^* \xi^* \rho_\omega^{\frac{1}{2}}. \quad (117)$$

Since both of \mathbf{v} and ω generates same ω , from the uniqueness (98), $k = k'$ and there exist a unitary V on \mathbb{C}^k and $e^{i\lambda} \in \mathbb{T}$ such that $v_\mu = e^{i\lambda} V^* \omega_\mu V$. From

$$\hat{\rho}_v = \hat{\rho}_v \circ T_v = \hat{\rho}_v \circ \text{Ad}(V^*) \circ T_\omega \circ \text{Ad}(V), \quad (118)$$

$\hat{\rho}_v \circ \text{Ad}(V^*)$ is a T_ω -invariant state. By the uniqueness of a T_ω -invariant state, we get $\rho_\omega = V \rho_v V^*$. Now we have

$$e^{i\lambda} V^* \omega_\mu V = v_\mu = e^{it} \rho_v^{-\frac{1}{2}} \theta v_\mu^* \theta^* \rho_v^{\frac{1}{2}} = e^{it} \rho_v^{-\frac{1}{2}} \theta (e^{i\lambda} V^* \omega_\mu V)^* \theta^* \rho_v^{\frac{1}{2}} = e^{it+i\lambda} \rho_v^{-\frac{1}{2}} \theta V^* \omega_\mu^* V \theta^* \rho_v^{\frac{1}{2}}. \quad (119)$$

From this, (117), and $\rho_\omega = V \rho_v V^*$, we have

$$e^{it} \rho_\omega^{-\frac{1}{2}} V \theta V^* \omega_\mu^* V \theta^* V^* \rho_\omega^{\frac{1}{2}} = e^{it} V \rho_v^{-\frac{1}{2}} \theta V^* \omega_\mu^* V \theta^* \rho_v^{\frac{1}{2}} V^* = \omega_\mu = e^{iu} \rho_\omega^{-\frac{1}{2}} \xi \omega_\mu^* \xi^* \rho_\omega^{\frac{1}{2}}. \quad (120)$$

Therefore, we get

$$e^{it} V \theta V^* \omega_\mu^* V \theta^* V^* = e^{iu} \xi \omega_\mu^* \xi^*. \quad (121)$$

Since ω is primitive, this means $\xi^* V \theta V^* = e^{i\eta} \mathbb{I}$ for some $e^{i\eta} \in \mathbb{T}$. Then we have

$$V \theta^2 V^* = V \theta V^* V \theta V^* = e^{-i\eta} \xi e^{-i\eta} \xi^* = \xi^2. \quad (122)$$

This proves the claim. \square

Since a matrix product state ω generated by a normalized primitive d -tuple is a unique gapped ground state by [FNW],[O3], it is pure and satisfies the split property. Therefore, if furthermore ω is reflection invariant, we can associate ω , our \mathbb{Z}_2 -index σ_ω in Definition 2.7. We then have the following theorem.

Theorem 5.3. *For a reflection invariant matrix product state ω generated by a normalized primitive d -tuple of matrices, we have*

$$\sigma_\omega = \zeta_\omega.$$

Proof. Let ω be a reflection invariant matrix product state generated by $\mathbf{v} \in \text{Prim}_u(d, k)$. Let $(\mathcal{H}_\omega, \pi_\omega, \Omega_\omega, \Gamma_\omega)$ be a reflection-split representation associated to ω . Then

$$(\hat{\mathcal{H}}_\omega := \mathcal{H}_\omega \otimes \mathcal{H}_\omega, \hat{\pi}_\omega := (\pi_\omega \circ \gamma_{L \rightarrow R}) \otimes \pi_\omega, \Omega_\omega)$$

is a GNS triple of ω . Let s_ω be a projection in $B(\mathcal{H}_\omega)$ such that the support projection of Ω_ω in $\mathbb{I}_{\mathcal{H}_\omega} \otimes B(\mathcal{H}_\omega)$ is $\mathbb{I}_{\mathcal{H}_\omega} \otimes s_\omega$. Set $\mathcal{M} := s_\omega \otimes B(s_\omega \mathcal{H}_\omega)$, and $p_\omega = s_\omega \otimes s_\omega$. We define a density matrix ρ_ω on \mathcal{H}_ω by

$$\text{Tr}_{\mathcal{H}_\omega}(\rho_\omega x) = \langle \Omega_\omega, (\mathbb{I}_{\mathcal{H}_\omega} \otimes x) \Omega_\omega \rangle, \quad x \in B(\mathcal{H}_\omega). \quad (123)$$

Since ω is translation invariant, there is a unitary V on $\hat{\mathcal{H}}_\omega$ such that

$$V \hat{\pi}_\omega(A) V^* = \hat{\pi}_\omega \circ \beta_1(A), \quad A \in \mathcal{A}.$$

From this, we obtain a homomorphism from $\hat{\pi}_\omega(\mathcal{A}_R)'' = \mathbb{I}_{\mathcal{H}_\omega} \otimes B(\mathcal{H}_\omega)$ onto $(\hat{\pi}_\omega \circ \beta_1(\mathcal{A}_R))'' \subset \mathbb{I}_{\mathcal{H}_\omega} \otimes B(\mathcal{H}_\omega)$

$$\mathbb{I}_{\hat{\mathcal{H}}_\omega} \otimes B(\mathcal{H}_\omega) \ni \mathbb{I}_{\mathcal{H}_\omega} \otimes x \mapsto V (\mathbb{I}_{\mathcal{H}_\omega} \otimes x) V^* \in \mathbb{I}_{\mathcal{H}_\omega} \otimes B(\mathcal{H}_\omega). \quad (124)$$

Therefore, there exists an endomorphism Θ on $B(\mathcal{H}_\omega)$ such that

$$V (\mathbb{I}_{\mathcal{H}_\omega} \otimes x) V^* = \mathbb{I}_{\mathcal{H}_\omega} \otimes \Theta(x), \quad x \in B(\mathcal{H}_\omega), \quad (125)$$

and $\Theta(B(\mathcal{H}_\omega))' = \pi_\omega(\text{M}_d \otimes \mathbb{I}_{\mathcal{A}_{[1, \infty)}})$. (Recall Lemma 2.6.8 of [BR1].) Note that $\Theta \circ \pi_\omega(A) = \pi_\omega \circ \beta_1(A)$ for any $A \in \mathcal{A}_R$. We recall the following fact:

Lemma 5.4 ([A]). *Let \mathcal{H} be a separable infinite dimensional Hilbert space, and $n \in \mathbb{N}$. Let $\Phi : B(\mathcal{H}) \rightarrow B(\mathcal{H})$ be a unital endomorphism of $B(\mathcal{H})$ such that $(\Phi(B(\mathcal{H})))'$ is isomorphic to M_n . Let $\{E_{ij}\}_{i,j=1,\dots,n}$ be a system of matrix units of $(\Phi(B(\mathcal{H})))'$. Then there exist $S_i \in B(\mathcal{H})$, $i = 1, \dots, n$ such that*

$$S_i^* S_j = \delta_{ij}, \quad E_{ij} = S_i S_j^*, \quad \sum_{j=1}^n S_j x S_j^* = \Phi(x), \quad x \in B(\mathcal{H}). \quad (126)$$

Applying this to our Θ in (125), we obtain operators $S_\mu \in B(\mathcal{H}_\omega)$ with $\mu = 1, \dots, d$ satisfying the following:

$$S_\mu^* S_\nu = \delta_{\mu\nu} \mathbb{I}, \quad (127)$$

$$\sum_{\mu=1,\dots,d} S_\mu \pi_\omega(A) S_\mu^* = \pi_\omega \circ \beta_1(A), \quad A \in \mathcal{A}_R. \quad (128)$$

$$\pi_\omega(e_{\mu\nu} \otimes \mathbb{I}_{[1, \infty)}) = S_\mu S_\nu^* \quad \text{for all } \mu, \nu = 1, \dots, d. \quad (129)$$

(See [A, BJP, BJ], Proof of Proposition 3.5 of [M2] and Lemma 3.5 of [M1].) Here $e_{\mu\nu} \otimes \mathbb{I}_{[1,\infty)}$ indicates an element $e_{\mu\nu}$ in $\mathcal{A}_{\{0\}} = M_d$ embedded into \mathcal{A}_R . From (127), (128) and (129), we have

$$\pi_\omega \left(\bigotimes_{k=0}^{l-1} e_{\mu_k, \nu_k} \right) = S_{\mu_0} \cdots S_{\mu_{l-1}} S_{\nu_{l-1}}^* \cdots S_{\nu_0}^*, \quad (130)$$

for all $l \in \mathbb{N}$, $\mu_k, \nu_k = 1, \dots, d$.

Now we restrict these S_μ to a *frustration-free subspace* \mathcal{K} of \mathcal{H}_ω . Recall that ω is the frustration free ground state of the translation invariant finite range interaction $\Phi_{\mathbf{v}}$ (99). Namely, there is a self-adjoint element $h_{\mathbf{v}} \in \mathcal{A}_{[0, m-1]}$, such that $\omega(\beta_x(h_{\mathbf{v}})) = 0$ for all $x \in \mathbb{Z}$. We consider the following frustration-free subspace of \mathcal{H}_ω :

$$\mathcal{K} := \bigcap_{\mathbb{Z} \ni x \geq 0} \ker \pi_\omega(\beta_x(h_{\mathbf{v}})).$$

Note that the support of ρ_ω defined in (123), is in \mathcal{K} , because ω is frustration-free. Let $P_{\mathcal{K}}$ be the orthogonal projection onto \mathcal{K} . As in [M1] (Lemma 3.2 and the argument in the proof of Lemma 3.6), \mathcal{K} is a finite dimensional space, and S_μ^* preserves \mathcal{K} :

$$S_\mu^* P_{\mathcal{K}} = P_{\mathcal{K}} S_\mu^* P_{\mathcal{K}}, \quad \mu = 1, \dots, d. \quad (131)$$

We denote $(S_\mu^* P_{\mathcal{K}})^* \in B(\mathcal{K})$ by B_μ , $\mu = 1, \dots, d$. Note that ρ_ω is of finite rank because \mathcal{K} is finite dimensional.

We claim that $\mathbb{B} = (B_\mu)_{\mu=1, \dots, d} \in \text{Prim}_u(d, \dim \mathcal{K})$. To prove this, it suffices to show that ρ_ω is faithful on \mathcal{K} and for the completely positive unital map $T_{\mathbb{B}}$ defined by $T_{\mathbb{B}}(x) = \sum_{\mu=1, \dots, d} B_\mu x B_\mu^*$, $x \in B(\mathcal{K})$, we have $T_{\mathbb{B}}^N(x) \rightarrow \text{Tr}_{\mathcal{H}_\omega}(\rho_\omega x) \mathbb{I}$, as $N \rightarrow \infty$, for each $x \in B(\mathcal{K})$. (See Lemma C.5 of [O1].) First we show that ρ_ω is faithful on \mathcal{K} . If ρ_ω is not faithful on \mathcal{K} , then there exists a unit vector $\xi \in \mathcal{K}$ which is orthogonal to the support of ρ_ω . By the definition of \mathcal{K} , this ξ defines a frustration free state $\psi = \langle \xi, \pi_\omega(\cdot) \xi \rangle$ on \mathcal{A}_R . Let p be the orthogonal projection onto the one-dimensional space $\mathbb{C}\xi$. As $\pi_\omega(\mathcal{A}_R)'' = B(\mathcal{H}_\omega)$, by Kaplansky's density Theorem, (Theorem 2.4.16 of [BR1]) there exists a net $\{x_\alpha\}_\alpha$ of positive elements in the unit ball of \mathcal{A}_R such that $\pi_\omega(x_\alpha) \rightarrow p$ in the σw -topology. For this net, we have $\lim_\alpha \omega(x_\alpha) = 0$ and $\lim_\alpha \psi(x_\alpha) = 1$. This contradicts to (100). Hence ρ_ω is faithful on \mathcal{K} . Next we show $T_{\mathbb{B}}^N(x) \rightarrow \text{Tr}_{\mathcal{H}_\omega}(\rho_\omega x) \mathbb{I}$, as $N \rightarrow \infty$ for all $x \in B(\mathcal{K})$. By $\pi_\omega(\mathcal{A}_R)'' = B(\mathcal{H}_\omega)$ and the finite dimensionality of \mathcal{K} , we have $B(\mathcal{K}) = P_{\mathcal{K}} \pi_\omega(\mathcal{A}_R \cap \mathcal{A}_{\text{loc}}) P_{\mathcal{K}}$. Therefore, for each $x \in B(\mathcal{K})$, there is an element $A \in \mathcal{A}_R \cap \mathcal{A}_{\text{loc}}$ such that $x = P_{\mathcal{K}} \pi_\omega(A) P_{\mathcal{K}}$. Since ω is a factor state and translation invariant, we have $\sigma w - \lim_{N \rightarrow \infty} \pi_\omega \circ \beta_N(A) = \omega(A) \mathbb{I}$. Therefore, for any $\eta \in \mathcal{K}$, we have

$$\langle \eta, T_{\mathbb{B}}^N(x) \eta \rangle = \langle \eta, T_{\mathbb{B}}^N(P_{\mathcal{K}} \pi_\omega(A) P_{\mathcal{K}}) \eta \rangle = \langle \eta, \pi_\omega \circ \beta_N(A) \eta \rangle \rightarrow \omega(A) \|\eta\|^2 = \text{Tr}_{\mathcal{H}_\omega}(\rho_\omega x) \|\eta\|^2, \quad N \rightarrow \infty. \quad (132)$$

Hence $\mathbb{B} \in \text{Prim}_u(d, \dim \mathcal{K})$.

The above proof for the primitivity also tells us that ρ_ω is the $T_{\mathbb{B}}$ -invariant state. From (130) and the definition of \mathbb{B} and (131), we see that \mathbb{B} is a d -tuple generating ω . Furthermore, as ρ_ω is faithful on \mathcal{K} , we have $s_\omega = P_{\mathcal{K}}$.

Let J_ω (resp. Δ_ω) be the modular conjugation (resp. modular operator) on $p_\omega \hat{\mathcal{H}}_\omega$ associated to $(\mathcal{M}, \Omega_\omega)$. By Proposition 4.3 there exists an anti-unitary $\theta : s_\omega \mathcal{H}_\omega \rightarrow s_\omega \mathcal{H}_\omega$ such that

$$J_\omega(s_\omega \otimes x) J_\omega^* = \theta x \theta^* \otimes s_\omega, \quad J_\omega(x \otimes s_\omega) J_\omega^* = s_\omega \otimes \theta x \theta^*, \quad (133)$$

for all $x \in B(s_\omega \mathcal{H}_\omega)$. By Proposition 4.3 and Theorem 4.4, we have $\theta^2 = \kappa_\omega s_\omega = \sigma_\omega s_\omega$. Recall also from Lemma 4.2

$$\Delta_\omega^{\frac{1}{2}}(s_\omega \otimes x) \Omega_\omega = \left(s_\omega \otimes \rho_\omega^{\frac{1}{2}} x \rho_\omega^{-\frac{1}{2}} \right) \Omega_\omega, \quad x \in B(s_\omega \mathcal{H}_\omega). \quad (134)$$

Now we prove that for the θ in (133), there exist an $a \in \mathbb{T}$ such that

$$a\rho_\omega^{-\frac{1}{2}}\theta B_\nu^*\theta^*\rho_\omega^{\frac{1}{2}} = B_\nu, \quad \nu = 1, \dots, d. \quad (135)$$

From this, we obtain the claim of the Theorem:

$$\sigma_\omega s_\omega = \theta^2 = \zeta_\omega s_\omega. \quad (136)$$

To show (135), we first show

$$|(S_\nu^* \otimes \mathbb{I}) \Omega_\omega\rangle \langle (S_\mu^* \otimes \mathbb{I}) \Omega_\omega| = |(\mathbb{I} \otimes S_\nu^*) \Omega_\omega\rangle \langle (\mathbb{I} \otimes S_\mu^*) \Omega_\omega|, \quad \mu, \nu = 1, \dots, d. \quad (137)$$

For any $l \in \mathbb{N}$, $\mu, \nu, \mu_{-l}, \dots, \mu_{-1}, \nu_{-l}, \dots, \nu_{-1}, \lambda_0, \dots, \lambda_{l-1}, \eta_0, \dots, \eta_{l-1} = 1, \dots, d$, we have

$$\begin{aligned} & \left\langle (S_\mu^* \otimes \mathbb{I}) \Omega_\omega, \left(\pi_\omega \circ \gamma_{L \rightarrow R} \left(\bigotimes_{j=-l}^{-1} e_{\mu_j, \nu_j} \right) \otimes \pi_\omega \left(\bigotimes_{j=0}^{l-1} e_{\lambda_j, \eta_j} \right) \right) (S_\nu^* \otimes \mathbb{I}) \Omega_\omega \right\rangle \\ &= \left\langle \Omega_\omega, \left(S_\mu S_{\mu_{-1}} \cdots S_{\mu_{-l}} S_{\nu_{-l}}^* \cdots S_{\nu_{-1}}^* S_\nu^* \otimes \pi_\omega \left(\bigotimes_{j=0}^{l-1} e_{\lambda_j, \eta_j} \right) \right) \Omega_\omega \right\rangle \\ &= \left\langle \Omega_\omega, \left(\pi_\omega \circ \gamma_{L \rightarrow R} \left(\bigotimes_{j=-l-1}^{-2} e_{\mu_{j+1}, \nu_{j+1}} \otimes e_{\mu\nu} \right) \otimes \pi_\omega \left(\bigotimes_{j=0}^{l-1} e_{\lambda_j, \eta_j} \right) \right) \Omega_\omega \right\rangle \\ &= \omega \left(\bigotimes_{j=-l-1}^{-2} e_{\mu_{j+1}, \nu_{j+1}} \otimes e_{\mu\nu} \otimes \bigotimes_{j=0}^{l-1} e_{\lambda_j, \eta_j} \right) \end{aligned} \quad (138)$$

In the third and the fourth line $e_{\mu\nu}$ is localized at site $j = -1$. Since ω is translation invariant, we have

$$(138) = \omega \left(\bigotimes_{j=-l}^{-1} e_{\mu_j, \nu_j} \otimes e_{\mu\nu} \otimes \bigotimes_{j=1}^l e_{\lambda_{j-1}, \eta_{j-1}} \right). \quad (139)$$

Here, $e_{\mu\nu}$ is localized at site $j = 0$. Then we have

$$\begin{aligned} (139) &= \left\langle \Omega_\omega, \left(\pi_\omega \circ \gamma_{L \rightarrow R} \left(\bigotimes_{j=-l}^{-1} e_{\mu_j, \nu_j} \right) \otimes \pi_\omega \left(e_{\mu\nu} \otimes \bigotimes_{j=1}^l e_{\lambda_{j-1}, \eta_{j-1}} \right) \right) \Omega_\omega \right\rangle \\ &= \left\langle \Omega_\omega, \left(\pi_\omega \circ \gamma_{L \rightarrow R} \left(\bigotimes_{j=-l}^{-1} e_{\mu_j, \nu_j} \right) \otimes S_\mu S_{\lambda_0} \cdots S_{\lambda_{l-1}} S_{\eta_{l-1}}^* \cdots S_{\eta_0}^* S_\nu^* \right) \Omega_\omega \right\rangle \\ &= \left\langle \Omega_\omega, \left(\pi_\omega \circ \gamma_{L \rightarrow R} \left(\bigotimes_{j=-l}^{-1} e_{\mu_j, \nu_j} \right) \otimes S_\mu \pi_\omega \left(\bigotimes_{j=0}^{l-1} e_{\lambda_j, \eta_j} \right) S_\nu^* \right) \Omega_\omega \right\rangle \\ &= \left\langle (\mathbb{I} \otimes S_\mu^*) \Omega_\omega, \left(\pi_\omega \circ \gamma_{L \rightarrow R} \left(\bigotimes_{j=-l}^{-1} e_{\mu_j, \nu_j} \right) \otimes \pi_\omega \left(\bigotimes_{j=0}^{l-1} e_{\lambda_j, \eta_j} \right) \right) (\mathbb{I} \otimes S_\nu^*) \Omega_\omega \right\rangle \end{aligned} \quad (140)$$

Hence we obtain

$$\begin{aligned} & \left\langle (S_\mu^* \otimes \mathbb{I}) \Omega_\omega, \left(\pi_\omega \circ \gamma_{L \rightarrow R} \left(\bigotimes_{j=-l}^{-1} e_{\mu_j, \nu_j} \right) \otimes \pi_\omega \left(\bigotimes_{j=0}^{l-1} e_{\lambda_j, \eta_j} \right) \right) (S_\nu^* \otimes \mathbb{I}) \Omega_\omega \right\rangle \\ &= \left\langle (\mathbb{I} \otimes S_\mu^*) \Omega_\omega, \left(\pi_\omega \circ \gamma_{L \rightarrow R} \left(\bigotimes_{j=-l}^{-1} e_{\mu_j, \nu_j} \right) \otimes \pi_\omega \left(\bigotimes_{j=0}^{l-1} e_{\lambda_j, \eta_j} \right) \right) (\mathbb{I} \otimes S_\nu^*) \Omega_\omega \right\rangle, \end{aligned} \quad (141)$$

for any $l \in \mathbb{N}$, $\mu, \nu, \mu_{-l}, \dots, \mu_{-1}, \nu_{-l}, \dots, \nu_{-1}, \lambda_0, \dots, \lambda_{l-1}, \eta_0, \dots, \eta_{l-1} = 1, \dots, d$. Since $\hat{\pi}_\omega(\mathcal{A}_{\text{loc}})$ is dense in $B(\mathcal{H}_\omega)$ with respect to the σ -weak topology, this means

$$|(S_\nu^* \otimes \mathbb{I}) \Omega_\omega\rangle \langle (S_\mu^* \otimes \mathbb{I}) \Omega_\omega| = |(\mathbb{I} \otimes S_\nu^*) \Omega_\omega\rangle \langle (\mathbb{I} \otimes S_\mu^*) \Omega_\omega|, \quad (142)$$

proving the claim.

From (137) with $\mu = \nu = 1, \dots, d$, we see that there is $a_\nu \in \mathbb{T}$ such that

$$(S_\nu^* \otimes \mathbb{I}) \Omega_\omega = a_\nu (\mathbb{I} \otimes S_\nu^*) \Omega_\omega. \quad (143)$$

Substituting this to (137), we find $a_\mu = a_\nu =: a$, if $(S_\nu^* \otimes \mathbb{I}) \Omega_\omega, (S_\mu^* \otimes \mathbb{I}) \Omega_\omega$ are not zero. Hence we get a constant $a \in \mathbb{T}$ such that

$$(S_\nu^* \otimes \mathbb{I}) \Omega_\omega = a (\mathbb{I} \otimes S_\nu^*) \Omega_\omega, \quad \nu = 1, \dots, d. \quad (144)$$

By the definition of \mathbb{B} and recalling $s_\omega = P_{\mathcal{K}}$, we obtain

$$(B_\nu^* \otimes \mathbb{I}) \Omega_\omega = a (\mathbb{I} \otimes B_\nu^*) \Omega_\omega, \quad \nu = 1, \dots, d. \quad (145)$$

On the other hand, by (41), (42), (134), (133) we have

$$\begin{aligned} (B_\nu^* \otimes \mathbb{I}) \Omega_\omega &= (B_\nu^* \otimes s_\omega) \Omega_\omega = J_\omega \Delta_\omega^{-\frac{1}{2}} (B_\nu \otimes s_\omega) \Omega_\omega = \Delta_\omega^{\frac{1}{2}} J_\omega (B_\nu \otimes s_\omega) J_\omega^* \Omega_\omega \\ &= \Delta_\omega^{\frac{1}{2}} (s_\omega \otimes \theta B_\nu \theta^*) \Omega_\omega = \left(s_\omega \otimes \rho_\omega^{\frac{1}{2}} \theta B_\nu \theta^* \rho_\omega^{-\frac{1}{2}} \right) \Omega_\omega. \end{aligned} \quad (146)$$

Combining (145) and (146), we obtain

$$\left(s_\omega \otimes \rho_\omega^{\frac{1}{2}} \theta B_\nu \theta^* \rho_\omega^{-\frac{1}{2}} \right) \Omega_\omega = a (s_\omega \otimes B_\nu^*) \Omega_\omega, \quad \nu = 1, \dots, d. \quad (147)$$

Since Ω_ω is separating for \mathcal{M} , we obtain

$$\rho_\omega^{\frac{1}{2}} \theta B_\nu \theta^* \rho_\omega^{-\frac{1}{2}} = a B_\nu^*, \quad \nu = 1, \dots, d. \quad (148)$$

Taking adjoint, we obtain (135). This completes the proof of the Theorem. \square

Acknowledgment.

The author is grateful to Hal Tasaki for fruitful discussion which was essential for the present work, and for the helpful comments on the manuscript. The beginning of the introduction heavily relies on his help. This work was supported by JSPS KAKENHI Grant Number 16K05171.

References

- [AKLT] I. Affleck, T. Kennedy, E.H. Lieb, and H. Tasaki. Valence bond ground states in isotropic quantum antiferromagnets. *Comm. Math. Phys.*, 115, 477–528, 1988.
- [A] W.B. Arveson. Continuous analogues of Fock space I. *Mem. Amer. Math. Soc.*, **409**, 1989.
- [BMNS] S. Bachmann, S. Michalakis, B. Nachtergaele, and R. Sims. Automorphic Equivalence within Gapped Phases of Quantum Lattice Systems. *Communications in Mathematical Physics* **309**, 835–871, 2012.
- [BN] S. Bachmann and B. Nachtergaele. On gapped phases with a continuous symmetry and boundary operators *J. Stat. Phys.* **154**, 91–112, 2014.

- [BJP] O. Bratteli P. Jorgensen, G. Price. Endomorphisms of $B(\mathcal{H})$. Quantization, nonlinear partial differential equations, and operator algebra. 93–138, Proc. Sympos. Pure Math., **59**, 1996.
- [BJ] O. Bratteli, P. E. T. Jorgensen. Endomorphisms of $B(H)$ II. Finitely Correlated States on O_n . Journal of functional analysis. **145**, 323–373 1997.
- [BR1] O. Bratteli, D. W. Robinson. *Operator Algebras and Quantum Statistical Mechanics 1*. Springer-Verlag, 1986.
- [BR2] O. Bratteli, D. W. Robinson. *Operator Algebras and Quantum Statistical Mechanics 2*. Springer-Verlag, 1996.
- [DJP] J. Dereziński, V. Jaksic and C.-A. Pillet: Perturbation theory of W^* -dynamics, Liouvilleans and KMS-states. *Reviews in Mathematical Physics*, 15-05, 447–489, 2003.
- [GW] Z.-C. Gu, and X.-G. Wen, *Tensor-entanglement-filtering renormalization approach and symmetry-protected topological order*, Phys. Rev. B, **80**, 155131 2009.
- [CGW] X. Chen, Z.-C. Gu, and X.-G. Wen, Classification of gapped symmetric phases in one-dimensional spin systems, Phys. Rev. B **83**, 035107 2011.
- [DL] S. Doplicher, R. Longo. Standard and split inclusions of von Neumann algebras. Invent. Math. **75** 493–536. 1984.
- [FNW] M. Fannes, B. Nachtergaele, and R.F. Werner. Finitely correlated states on quantum spin chains. *Comm. Math. Phys.*, **144**, 443–490, 1992.
- [FNW2] M. Fannes, B. Nachtergaele, and R.F. Werner. Finitely correlated pure states. *Journal of functional analysis.*, **120**, 511–534, 1994.
- [Hal1] F.D.M. Haldane, Continuum dynamics of the 1-D Heisenberg antiferromagnet: identification with the $O(3)$ nonlinear sigma model, Phys. Lett. **93A**, 464–468 1983.
- [Hal2] F.D.M. Haldane, Nonlinear field theory of large-spin Heisenberg antiferromagnets: semi-classically quantized solitons of the one-dimensional easy-axis Néel state, Phys. Rev. Lett. **50** 1153–1156 1983.
- [H1] M. Hastings. An area law for one-dimensional quantum systems. Journal of Statistical Mechanics. P08024, 2007.
- [H2] M. Hastings. Quasi-adiabatic Continuation for Disordered Systems: Applications to Correlations, Lieb-Schultz-Mattis, and Hall Conductance. <http://arxiv.org/abs/1001.5280v2> [math-ph], 2010.
- [K] T. Kennedy, Exact diagonalization of open spin 1 chains, J. Phys.: Cond. Matt. **2**, 5737–5745, 1990.
- [KN] T. Koma and B. Nachtergaele The Spectral Gap of the Ferromagnetic XXZ-Chain Letters in Mathematical Physics **40**, 1–16, 1997.
- [KT1] T. Kennedy and H. Tasaki. Hidden $\mathbb{Z}_2 \times \mathbb{Z}_2$ -symmetry breaking in Haldane-gap antiferromagnets. Phys. Rev. B, **45** 304–307, 1992.
- [KT2] T. Kennedy and H. Tasaki. Hidden symmetry breaking and the Haldane phase in $S = 1$ quantum spin chains. Communications in Mathematical Physics, **147** 431–484, 1992.
- [M1] T. Matsui. A characterization of matrix product pure states. Infinite dimensional analysis and quantum probability. **1** 647–661. 1998.

- [M2] T. Matsui. The split property and the symmetry breaking of the quantum spin chain. *Communications in Mathematical Physics*, **218** 393–416, 2001.
- [M3] T. Matsui. Boundedness of entanglement entropy and split property of quantum spin chains. *Reviews in Mathematical Physics*, 1350017, 2013.
- [NOS] B. Nachtergaele, Y. Ogata, and R. Sims. Boundedness of entanglement entropy and split property of quantum spin chains. *J. Stat. Phys.*, **124**, 1–13, 2006.
- [NR] M. den Nijs and K. Rommelse, Preroughening transitions in crystal surfaces and valence-bond phases in quantum spin chains, *Phys. Rev. B* **40**, 4709, 1989.
- [O1] Y. Ogata. A class of asymmetric gapped Hamiltonians on quantum spin chains and its classification I. *Communications in Mathematical Physics*, **348**, 847–895, 2016.
- [O2] Y. Ogata. A class of asymmetric gapped Hamiltonians on quantum spin chains and its classification II. *Communications in Mathematical Physics*, **348**, 897–957, 2016.
- [O3] Y. Ogata. A class of asymmetric gapped Hamiltonians on quantum spin chains and its classification III. *Communications in Mathematical Physics*, **352**, 1205–1263, 2017.
- [O4] Y. Ogata. A \mathbb{Z}_2 -index of symmetry protected topological phases with time reversal symmetry for quantum spin chains [arXiv:1810.01045](#)
- [OT] Y. Ogata and H. Tasaki Lieb-Schultz-Mattis type theorems for quantum spin chains without continuous symmetry. *Communications in Mathematical Physics*, 2019.
- [PTBO1] F. Pollmann, A. Turner, E. Berg, and M. Oshikawa Entanglement spectrum of a topological phase in one dimension. *Phys. Rev. B* **81**, 064439, 2010.
- [PTBO2] F. Pollmann, A. Turner, E. Berg, and M. Oshikawa Symmetry protection of topological phases in one-dimensional quantum spin systems. *Phys. Rev. B* **81**, 075125, 2012.
- [PWSVC] D. Perez-Garcia, M.M. Wolf, M. Sanz, F. Verstraete, and J.I. Cirac, *String order and symmetries in quantum spin lattices*, *Phys. Rev. Lett.* **100**, 167202 2008.
- [T1] M. Takesaki, Theory of operator algebras. I. *Encyclopaedia of Mathematical Sciences*. Springer-Verlag, Berlin, 2002.
- [T2] M. Takesaki, Theory of operator algebras. II. *Encyclopaedia of Mathematical Sciences*. Springer-Verlag, Berlin, 2003.
- [Tas1] H. Tasaki Topological phase transition and \mathbb{Z}_2 index for $S = 1$ quantum spin chains [arXiv:1804.04337](#)
- [Tas2] H. Tasaki, *Physics and mathematics of quantum many-body systems*, (to be published from Springer).
- [W] M.M. Wolf. Quantum channels & operations. Unpublished. 2012.