

A Bayesian Nonparametric Test for Assessing Multivariate Normality

Luai Al-Labadi^{*1}, Forough Fazeli Asl^{†2}, and Zahra Saberi^{‡2}

¹Department of Mathematical and Computational Sciences, University of
Toronto Mississauga, Mississauga, Ontario L5L 1C6, Canada.

²Department of Mathematical Sciences, Isfahan University of
Technology, Isfahan 84156-83111, Iran.

Abstract

A novel Bayesian nonparametric test for assessing multivariate normal models is presented. While there are extensive frequentist and graphical methods for testing multivariate normality, it is challenging to find Bayesian counterparts. The approach is based on the Dirichlet process and Mahalanobis distance. Specifically, the Mahalanobis distance is employed to transform the m -variate problem into a univariate problem. Then the Dirichlet process is used as a prior on the distribution of the distance. The concentration of the distribution of the distance between the posterior process and the chi-square distribution with m degrees of freedom is compared to that between the prior process and the chi-square distribution via a relative belief ratio. The distance between the Dirichlet process and the chi-square distribution is established based on the Anderson-Darling distance. Key

^{*} *Corresponding author:* luai.allabadi@utoronto.ca

[†]forough.fazeli@math.iut.ac.ir

[‡]z_saberi@cc.iut.ac.ir

results of the approach are derived. The procedure is illustrated through several examples, in which it shows excellent performance.

KEYWORDS: Anderson-Darling distance, Dirichlet process, Mahalanobis distance, Multivariate hypothesis testing, Relative belief inferences.

MSC 2010 62F15, 62G10, 62H15

1 Introduction

The assumption of multivariate normality is a key assumption in many statistical applications such as pattern recognition and exploratory multivariate methods [18, 24]. The need to check this assumption is of special importance as if it does not hold, the obtained results based on this assumption may lead to an error. Specifically, for a given m -variate sampled data $\mathbf{y}_{m \times n} = (\mathbf{y}_1, \dots, \mathbf{y}_n)$ with size of n , where $\mathbf{y}_i \in \mathbb{R}^m$ for $i = 1, \dots, n$, the interest is to determine whether $\mathbf{y}_{m \times n}$ comes from a multivariate normal population.

Many tests and graphical methods have been considered to assess the multivariate normality assumption. Healy [27] described an extension of normal plotting techniques to handle multivariate data. Mardia [37] proposed a test based on the asymptotic distribution of measures of multivariate skewness and kurtosis. Tests based on transforming the multivariate problem into the univariate problem were established by Rincón-Gallardo et al.[40], Royston [41], Fattorini [22], and Hasofer and Stein[26]. A class of invariant consistent tests based on a weighted integral of the squared distance between the empirical characteristic function and the characteristic function of the multivariate normal distribution was suggested by Henze and Zirkler [29]. Holgersson [30] presented a simple graphical method based on the scatter plot. Doornik and Hansen [17] developed an omnibus test based on a transformed skewness and kurtosis. They showed that their test is more powerful than the Shapiro-Wilk test proposed by Royston [41]. Alva and Estrada [9] proposed a multivariate normality test based on Shapiro-Wilks

statistic for univariate normality and on an empirical standardization of the observations. Jönsson [33] presented a robust test with powerful performance. Hanusz and Tarasińska [25] proposed two tests based on Small's and Srivastava's graphical methods. A powerful test was offered by Zhou and Shao [46] with good application in biomedical studies. Kim [35] generalized Jarque-Bera univariate normality test to the multivariate case. Kim and Park [34] derived several powerful omnibus tests based on the likelihood ratio and the characterizations of the multivariate normal distribution. Madukaife and Okafor [36] proposed a powerful affine invariant test based on interpoint distances of principal components. Henze and Visagie [28] derived a new test based on a partial differential equation involving the moment generating function.

While there are considerable frequentist and graphical methods for testing multivariate normality, it is difficult to find relevant Bayesian counterparts. Most available Bayesian tests focused on employing Bayesian nonparametric methods for univariate data. See for example Al-Labadi and Zarepour [5, 6, 8]. A remarkable work that covers the multivariate case was developed by Tokdar and Martin [43], where they established a Bayesian test based on characterizing alternative models by Dirichlet process mixture distributions. The authors took the Bayes factor as evidence for or against the null distribution, using an importance sampling for inference. In the current paper, a novel Bayesian nonparametric test for assessing multivariate normality is proposed. The developed test is based on using Mahalanobis distance. Using this distance has the advantage of converting the m -variate problem into a univariate problem. Specifically, whenever $\mathbf{y}_{m \times n}$ comes from a multivariate normal distribution, the distribution of the corresponding Mahalanobis distances, denoted by P , is approximately chi-square with m degrees of freedom [32]. This reduces the problem to test the hypothesis $\mathcal{H}_0 : P = F_{(m)}$, where $F_{(m)}$ denotes the cumulative distribution function of the chi-square distribution with m degrees of freedom. The proposed test deems the Dirichlet process as a prior on P . Then the concentration of the distribution of the distance between the posterior process and $F_{(m)}$ is compared to the concentration of the

distribution of the distance between the prior process and $F_{(m)}$. The distance between the Dirichlet process and $F_{(m)}$ is developed based on the Anderson-Darling distance as an appropriate tool to detect the difference, especially when this difference is due to the tails. This comparison is made via a *relative belief ratio*. A calibration of the relative belief ratio is also presented. The anticipated test is easy to implement with excellent performance, it does not require providing a closed form of the relative belief ratio. In addition, unlike the tests that use p-values, the proposed test permits us to state evidence for the null hypothesis. We point out that the comparison between the concentration of the posterior and the prior distribution of the distance was suggested by Al-Labadi and Evans [4] for model checking of univariate distributions.

The remainder of this paper is organized as follows. A general discussion about the relative belief ratio is given in Section 2. The definition and some fundamental properties of the Dirichlet process are presented in Section 3. An explicit expression for calculating the Anderson-Darling distance between the Dirichlet process and a continuous distribution is derived in Section 4. A Bayesian nonparametric test for assessing multivariate normality and some of its relevant properties are developed in Section 5. A computational algorithm to carry out the proposed test is discussed in Section 6. The performance of the proposed test is discussed in Section 7, where several simulated examples and a real data set are considered. Finally, a summary of the findings is given in Section 8. All technical proofs are included in the Appendix.

2 Relative belief inferences

The relative belief ratio is a common measure of statistical evidence. It leads to a straightforward inference in hypothesis testing problems. For more details, let $\{f_\theta : \theta \in \Theta\}$ denote a collection of densities on a sample space $\mathfrak{X}$ and let π denote a prior on the parameter space Θ . Note that the densities may represent discrete or continuous probability measures but they are all with respect to the same support

measure $d\theta$. After observing the data x , the posterior distribution of θ , denoted by $\pi(\theta | x)$, is a revised prior and is given by the density $\pi(\theta | x) = \pi(\theta)f_\theta(x)/m(x)$, where $m(x) = \int_{\Theta} \pi(\theta)f_\theta(x) d\theta$ is the prior predictive density of x . For a parameter of interest $\psi = \Psi(\theta)$, let Π_Ψ denote the marginal prior probability measure and $\Pi_\Psi(\cdot | x)$ denote the marginal posterior probability measure. It is assumed that Ψ satisfies regularity conditions so that the prior density π_Ψ and the posterior density $\pi_\Psi(\cdot | x)$ of ψ exist with respect to some support measure on the range space for Ψ [20]. The relative belief ratio for a value ψ is then defined by

$$RB_\Psi(\psi | x) = \lim_{\delta \rightarrow 0} \Pi_\Psi(N_\delta(\psi) | x) / \Pi_\Psi(N_\delta(\psi)),$$

where $N_\delta(\psi)$ is a sequence of neighborhoods of ψ converging nicely to ψ as $\delta \rightarrow 0$ [20]. When π_Ψ and $\pi_\Psi(\cdot | x)$ are continuous at ψ , the relative belief ratio is defined by

$$RB_\Psi(\psi | x) = \pi_\Psi(\psi | x) / \pi_\Psi(\psi),$$

the ratio of the posterior density to the prior density at ψ . Therefore, $RB_\Psi(\psi | x)$ measures the change in the belief of ψ being the true value from a *priori* to a *posteriori*. Note that a relative belief ratio is similar to a Bayes factor, as both are measures of evidence, but the latter measures evidence via the change in an odds ratio. In general, when a Bayes factor is defined via a limit in the continuous case, the limiting value equals the corresponding relative belief ratio. For a further discussion about the relationship between relative belief ratios and Bayes factors see, for instance, Evans [20], Chapter 4.

Since $RB_\Psi(\psi | x)$ is a measure of the evidence that ψ is the true value, if $RB_\Psi(\psi | x) > 1$, then the probability of the ψ being the true value from a priori to a posteriori is increased, consequently there is evidence based on the data that ψ is the true value. If $RB_\Psi(\psi | x) < 1$, then the probability of the ψ being the true value from a priori to a

posteriori is decreased. Accordingly, there is evidence against based on the data that ψ being the true value. For the case $RB_{\Psi}(\psi | x) = 1$ there is no evidence either way.

Obviously, $RB_{\Psi}(\psi_0 | x)$ measures the evidence of the hypothesis $\mathcal{H}_0 : \Psi(\theta) = \psi_0$. For a large value of c ($c \gg 1$), $RB_{\Psi}(\psi_0 | x) = c$ provides strong evidence in favor of ψ_0 because belief has increased by a factor of c after seeing the data. However, there may also exist other values of ψ that had even larger increases. Thus, it is also necessary, however, to calibrate whether this is strong or weak evidence for or against $\mathcal{H}_0$. A typical calibration of $RB_{\Psi}(\psi_0 | x)$ is given by the *strength*

$$\Pi_{\Psi} [RB_{\Psi}(\psi | x) \leq RB_{\Psi}(\psi_0 | x) | x]. \quad (1)$$

The value in (1) indicates that the posterior probability that the true value of ψ has a relative belief ratio no greater than that of the hypothesized value ψ_0 . Noticeably, (1) is not a p-value as it has a very different interpretation. When $RB_{\Psi}(\psi_0 | x) < 1$, there is evidence against ψ_0 , then a small value of (1) indicates strong evidence against ψ_0 . On the other hand, a large value for (1) indicates weak evidence against ψ_0 . Similarly, when $RB_{\Psi}(\psi_0 | x) > 1$, there is evidence in favor of ψ_0 , then a small value of (1) indicates weak evidence in favor of ψ_0 , while a large value of (1) indicates strong evidence in favor of ψ_0 . For applications of the use of the relative belief ratio in different univariate hypothesis testing problems, see Evans [19, 20], Al-Labadi and Evans [4] and Al-Labadi et al. [1, 2].

3 Dirichlet process

The Dirichlet process, introduced by Ferguson [23], is the most commonly used prior in the Bayesian nonparametric inferences. In this section, we recall the most relevant properties of this prior. Consider a space $\mathfrak{X}$ with a σ -algebra $\mathcal{A}$ of subsets of $\mathfrak{X}$, let H be a fixed probability measure on $(\mathfrak{X}, \mathcal{A})$, called the *base measure*, and

a be a positive number, called the *concentration parameter*. A random probability measure $P = \{P(A) : A \in \mathcal{A}\}$ is called a Dirichlet process on $(\mathfrak{X}, \mathcal{A})$ with parameters a and H , denoted by $P \sim DP(a, H)$, if for every measurable partition $A_1, \dots, A_k$ of $\mathfrak{X}$ with $k \geq 2$, the joint distribution of the vector $(P(A_1), \dots, P(A_k))$ has the Dirichlet distribution with parameter $aH(A_1), \dots, aH(A_k)$. Also, it is assumed that $H(A_j) = 0$ implies $P(A_j) = 0$ with probability one. Consequently, for any $A \in \mathcal{A}$, $P(A) \sim \text{beta}(aH(A), a(1 - H(A)))$, $E(P(A)) = H(A)$ and $\text{Var}(P(A)) = H(A)(1 - H(A))/(1 + a)$. Accordingly, the base measure H plays the role of the center of P while the concentration parameter a controls variation of the process P around the base measure H . One of the most well-known properties of the Dirichlet process is the conjugacy property. That is, when the sample $x = (x_1, \dots, x_n)$ is drawn from $P \sim DP(a, H)$, the posterior distribution of P given x , denoted by P_x , is also a Dirichlet process with concentration parameter $a + n$ and base measure

$$H_x = a(a + n)^{-1}H + n(a + n)^{-1}F_n, \quad (2)$$

where F_n denotes the empirical cumulative distribution function (cdf) of the sample x . Note that, H_x is a convex combination of the base measure H and the empirical cdf F_n . Therefore, $H_x \rightarrow H$ as $a \rightarrow \infty$ while $H_x \rightarrow F_n$ as $a \rightarrow 0$. A detailed discussion about choosing the hyperparameters a and H will be presented in Section 5.

Following Ferguson [23], $P \sim DP(a, H)$ can be represented as

$$P = \sum_{i=1}^{\infty} L^{-1}(\Gamma_i) \delta_{Y_i} / \sum_{i=1}^{\infty} L^{-1}(\Gamma_i), \quad (3)$$

where $\Gamma_i = E_1 + \dots + E_i$ with $E_i \stackrel{i.i.d.}{\sim} \text{exponential}(1)$, $Y_i \stackrel{i.i.d.}{\sim} H$ independent of the Γ_i , $L^{-1}(y) = \inf\{x > 0 : L(x) \geq y\}$ with $L(x) = a \int_x^{\infty} t^{-1} e^{-t} dt$, $x > 0$, and δ_a the Dirac delta measure. The series representation (3) implies that the Dirichlet process is a discrete probability measure even for the cases with an absolutely continuous base

measure H . Note that, since data is always measured to finite accuracy, the true distribution being sampled from is discrete. This makes the discreteness property of P has no significant limitation. By imposing the weak topology, the support of the Dirichlet process could be quite large. To be more precise, when the support of the base measure is $\mathfrak{X}$, then the space of all probability measures on $\mathfrak{X}$ is the support of the Dirichlet process.

Because no closed form exists for $L^{-1}(\cdot)$, working with representation (3) to sample from the Dirichlet process is not simple. This issue motivates researchers to propose several methods in order to approximately simulate the Dirichlet process. One such efficient method was presented by Zarepour and Al-Labadi [45]. They showed that the Dirichlet process $P \sim DP(a, H)$ can be approximated by

$$P_N = \sum_{i=1}^N J_i \delta_{Y_i}, \quad (4)$$

with the monotonically decreasing weights $J_i = \frac{G_{a/N}^{-1}(\frac{\Gamma_i}{\Gamma_{N+1}})}{\sum_{j=1}^N G_{a/N}^{-1}(\frac{\Gamma_j}{\Gamma_{N+1}})}$, where Γ_i and Y_i are defined as before, N is a positive large integer and $G_{a/N}$ denotes the complement-cdf of the gamma($a/N, 1$) distribution. Note that, $G_{a/N}^{-1}(p)$ is the $(1 - p)$ -th quantile of the gamma($a/N, 1$) distribution. Also, Zarepour and Al-Labadi [45] showed that P_N converges almost surely to (3) as N goes to infinity with a fast rate of convergence. The following algorithm describes how the approximation (4) can be used to generate a sample from $DP(a, H)$.

Algorithm A (*Approximately generating a sample from $DP(a, H)$*):

- i.* Fix a large positive integer N and generate i.i.d. $Y_i \sim H$ for $i = 1, \dots, N$.
- ii.* For $i = 1, \dots, N + 1$, generate i.i.d. E_i from the exponential distribution with rate 1, independent of $(Y_i)_{1 \leq i \leq N}$ and put $\Gamma_i = E_1 + \dots + E_i$.
- iii.* Compute $G_{a/N}^{-1}(\Gamma_i/\Gamma_{N+1})$ for $i = 1, \dots, N$ and return P_N .

For other simulation methods of the Dirichlet process, see Bondesson [13], Sethuraman [42], Wolpert and Ickstadt [44] and Al-Labadi and Zarepour [7].

4 Measuring the distance

Measuring the distance between two distributions is an essential tool in model checking. In this section, two well-known distances, namely *Anderson-Darling* distance and *Mahalanobis* distance, are considered.

4.1 Anderson-Darling distance

The Anderson-Darling distance between two cdf's F and G is given by

$$d_{AD}(F, G) = \int_{-\infty}^{\infty} \frac{(F(t) - G(t))^2}{G(t)(1 - G(t))} dG(t).$$

Anderson-Darling distance can be viewed as a modification of the Cramér-von Mises distance that gives more weight to data points in the tails of the distribution. The next lemma provides an explicit formula to compute the Anderson-Darling distance between a Dirichlet process and a continuous cdf. Throughout this paper, “log” denotes the natural logarithm.

Lemma 1 *Let G be a continuous cdf and $P_N = \sum_{i=1}^N J_i \delta_{Y_i}$ be a discrete distribution, where $(Y_i)_{1 \leq i \leq N}$ are i.i.d. from H . Let $Y_{(1)} \leq \dots \leq Y_{(N)}$ be the order statistics of $(Y_i)_{1 \leq i \leq N}$ and $J'_1, \dots, J'_N$ are the associated jump sizes such that $J_i = J'_j$ when $Y_i = Y_{(j)}$. Then the Anderson-Darling distance between P_N and G is given by*

$$\begin{aligned} d_{AD}(P_N, G) = & 2 \sum_{i=1}^{N-1} \sum_{j=1}^{i-1} \sum_{k=j+1}^i J'_j J'_k C_{i,i+1} + \sum_{i=1}^{N-1} \sum_{j=1}^i J'^2_j C_{i,i+1} + 2 \sum_{i=1}^{N-1} \sum_{j=1}^i J'_j C^*_{i,i+1} \\ & - \sum_{i=1}^{N-1} C^*_{i,i+1} - \log[G(Y_{(N)})(1 - G(Y_{(1)}))] - 1, \end{aligned}$$

where $C_{i,i+1} = \log \frac{G(Y_{(i+1)})(1-G(Y_{(i)}))}{G(Y_{(i)})(1-G(Y_{(i+1)}))}$ and $C_{i,i+1}^* = \log \frac{1-G(Y_{(i+1)})}{1-G(Y_{(i)})}$.

Proof. The proof is given in Appendix A. ■

The next corollary indicates that the distribution of $d_{AD}(P_N, G)$ is independent of G when $G = H$ and $P_N = \sum_{i=1}^N J_i \delta_{Y_i}$.

Corollary 2 Suppose that $(Y_i)_{1 \leq i \leq N}$ are i.i.d. from G , independent of $(J_i)_{1 \leq i \leq N}$ and $P_N = \sum_{i=1}^N J_i \delta_{Y_i}$. Then $d_{AD}(P_N, G) \stackrel{d}{=} 2 \sum_{i=1}^{N-1} \sum_{j=1}^i J'_j U_{i,i+1}^* - \sum_{i=1}^{N-1} U_{i,i+1}^* + \sum_{i=1}^{N-1} \sum_{j=1}^i J_j'^2 U_{i,i+1} + 2 \sum_{i=1}^{N-1} \sum_{j=1}^{i-1} \sum_{k=j+1}^i J'_j J'_k U_{i,i+1} - \log[U_{(N)}(1 - U_{(1)})] - 1$, where $U_{i,i+1} = \log \frac{U_{(i+1)}(1-U_{(i)})}{U_{(i)}(1-U_{(i+1)})}$, $U_{i,i+1}^* = \log \frac{1-U_{(i+1)}}{1-U_{(i)}}$ and $U_{(i)}$ is the i -th order statistic for $(U_i)_{1 \leq i \leq N}$ i.i.d. uniform $[0, 1]$.

Proof. Since for $1 \leq i \leq N$, $U_i = G(Y_i)$ and Y_i is a sequence of i.i.d. random variable with continuous distribution G , the probability integral transformation theorem implies that $(U_i)_{1 \leq i \leq N}$ is a sequence of i.i.d. random variable uniformly distributed on the interval $[0, 1]$. The proof is completed by considering the order statistic of $(U_i)_{1 \leq i \leq N}$ in Lemma 1. ■

4.2 Mahalanobis distance

Mahalanobis distance measures the distance of m -variate point $\mathbf{Y}$ generated from a known distribution F_θ to the mean $\boldsymbol{\mu}_m = E_\theta(\mathbf{Y})$ of the distribution. Let $\Sigma_{m \times m}$ be the covariance matrix of the m -variate distribution F_θ , the Mahalanobis distance is defined as

$$D_M(\mathbf{Y}) = \sqrt{(\mathbf{Y} - \boldsymbol{\mu}_m)^T \Sigma_{m \times m}^{-1} (\mathbf{Y} - \boldsymbol{\mu}_m)}. \quad (5)$$

Note that, (5) is limited to the cases when both $\boldsymbol{\mu}_m$ and $\Sigma_{m \times m}$ are known. However, in most cases, the mean and covariance of F_θ exist but unknown. For such cases, the Mahalanobis distance is defined based on the measuring of distance between a subject's data and the mean of all observations in an observed sample. To be more precise, given a sample of n independent m -variate $\mathbf{y}_1, \dots, \mathbf{y}_n$, the sample Mahalanobis distance of

$\mathbf{y}_i$ to the sample mean $\bar{\mathbf{y}}$, denoted by $d_M(\mathbf{y}_i)$, is defined by

$$d_M(\mathbf{y}_i) = \sqrt{(\mathbf{y}_i - \bar{\mathbf{y}})^T S_y^{-1} (\mathbf{y}_i - \bar{\mathbf{y}})}, \quad (6)$$

where S_y denotes the sample covariance matrix. Some interesting properties of the Mahalanobis distance are presented in [32]. For example, when the parent population is m -variate normal, for large enough values of n and $n - m$ ($n, n - m > 30$), each of the squared distance $d_M^2(\mathbf{y}_1), \dots, d_M^2(\mathbf{y}_n)$ behave like a chi-square random variable with m degrees of freedom.

5 Bayesian nonparametric approach for assessing multivariate normality

In this section, a new test for assessing multivariate normality is presented. For this purpose, consider the family of m -variate normal distribution $\mathcal{F} = \{N_m(\boldsymbol{\mu}_m, \Sigma_{m \times m}) : \boldsymbol{\mu}_m \in \mathbb{R}^m, \det(\Sigma_{m \times m}) > 0\}$. Let $\mathbf{Y}_1, \dots, \mathbf{Y}_n$ be a random sample from m -variate distribution F . The problem under consideration is to test the hypothesis

$$\mathcal{H}_0 : F \in \mathcal{F}, \quad (7)$$

using the Bayesian nonparametric framework. The first step is to reduce the multivariate problem to a univariate problem. One way to accomplish that is through the Mahalanobis distance. For this, assume that $\hat{\boldsymbol{\mu}}_m = \bar{\mathbf{Y}}$ and $\hat{\Sigma}_{m \times m} = S_Y$ are the sample mean and sample covariance matrix, respectively. Then, $N_m(\hat{\boldsymbol{\mu}}_m, \hat{\Sigma}_{m \times m})$ is the best representative of the family $\mathcal{F}$ to compare with distribution F . Define

$$D_M(\mathbf{Y}_i) = \sqrt{(\mathbf{Y}_i - \hat{\boldsymbol{\mu}}_m)^T \hat{\Sigma}_{m \times m}^{-1} (\mathbf{Y}_i - \hat{\boldsymbol{\mu}}_m)} \quad \text{for } 1 \leq i \leq n.$$

Assume that $F_{(m)}$ is the cdf of the chi-square distribution with m degrees of freedom and $(D_M^2(\mathbf{Y}_i))_{1 \leq i \leq n}$ is a sequence of random variables with continuous distribution function P . From Johnson and Wichern [32], if $\mathcal{H}_0$ is true, then we expect that P is (approximately) the same as $F_{(m)}$. Thus, testing (7) is equivalent to testing

$$\mathcal{H}_0 : P = F_{(m)}. \quad (8)$$

For testing (8), let $\mathbf{y}_{m \times n} = (\mathbf{y}_1, \dots, \mathbf{y}_n)$ be an observed sample from F and $d = (d_M^2(\mathbf{y}_1), \dots, d_M^2(\mathbf{y}_n))$ be the corresponding observed squared Mahalanobis distance. If $P \sim DP(a, F_{(m)})$, for a given choice of a , by (2), then $P_d = P|d \sim DP(a + n, H_d)$. From (6), if $\mathcal{H}_0$ is true, then the sampled observations $d_M^2(\mathbf{y}_1), \dots, d_M^2(\mathbf{y}_n)$ would be approximately independent chi-square with m degrees of freedom. Thus, the posterior distribution of the distance between P_d and $F_{(m)}$ should be more concentrated around zero than the prior distribution of the distance between P and $F_{(m)}$. A good reason for choosing $H = F_{(m)}$ is to avoid prior-data conflict. Prior-data conflict means that there is a tiny overlap between the effective support regions of P and H . It can lead to errors in the computation of the prior distribution of the distance between P and $F_{(m)}$ and then the value of the relative belief ratio and its relevant strength when $\mathcal{H}_0$ is not true, see for example Section 7, Table 2. For more discussion about prior-data conflict see [21, 3, 4, 38].

In the proposed test, we use Lemma 1 to compute the Anderson-Darling distances $d_{AD}(P_N, F_{(m)})$ and $d_{AD}(P_{d_N}, F_{(m)})$, where P_N and P_{d_N} are approximations of Dirichlet processes $P \sim DP(a, H)$ and $P_d \sim DP(a + n, H_d)$, respectively. Then, the relative belief ratio is used to compare the concentration of the posterior and the prior distribution of $d_{AD}(P_{d_N}, F_{(m)})$ and $d_{AD}(P_N, F_{(m)})$, respectively, at zero.

Another significant concern to validate the test is determining suitable values of a . The following Lemma presents the expectation and variance of the prior distribution of the Anderson-Darling distance. For the Cramér-von Mises distance d_{CvM} , Al-Labadi

and Evans [4] showed that $E_P(d_{CvM}(P, H)) = 1/6(a+1)$.

Lemma 3 *Let $P \sim DP(a, H)$ and $d_{AD}(P, H)$ be the Anderson-Darling distance between P and H , then*

$$\begin{aligned} i. \quad E_P(d_{AD}(P, H)) &= \frac{1}{a+1}, \\ ii. \quad Var_P(d_{AD}(P, H)) &= \frac{2((\pi^2 - 9)a^2 + (30 - 2\pi^2)a - 3\pi^2 + 36)}{3(a+1)^2(a+2)(a+3)}, \end{aligned}$$

where $E_P(d_{AD}(P, H))$ and $Var_P(d_{AD}(P, H))$ are the expectation and variance of the prior distribution of $d_{AD}(P, H)$ with respect to P , respectively.

Proof. The proof is given in Appendix B. ■

The next corollary highlights the effect of the value a on the prior distribution of the Anderson-Darling distance.

Corollary 4 *Let $P \sim DP(a, H)$ and $H = F_{(m)}$ be the cdf of the chi-square distribution with m degrees of freedom. Suppose that $a \rightarrow \infty$, then*

$$\begin{aligned} i. \quad E_P(d_{AD}(P, H)) &\rightarrow 0, \text{ and, } Var_P(d_{AD}(P, H)) \rightarrow 0. \\ ii. \quad E_P((a+1)d_{AD}(P, H)) &\rightarrow 1, \text{ and, } Var_P((a+1)d_{AD}(P, H)) \rightarrow \frac{2}{3}(\pi^2 - 9). \\ iii. \quad d_{AD} &\xrightarrow{qm} 0, \text{ and, } d_{AD} \xrightarrow{a.s.} 0. \end{aligned}$$

Where “ $\xrightarrow{qm}$ ” denotes convergence in quadratic mean.

Proof. The proof of (i) and (ii) are followed immediately by letting $a \rightarrow \infty$ in part (i) and (ii) of Lemma 3. For (iii), the convergence in quadratic mean is followed by letting $a \rightarrow \infty$ in $E_P(d_{AD}(P, H))^2 = \frac{a(2\pi^2-15)-6(\pi^2-15)}{3(a+3)(a+2)(a+1)}$; see the proof of Lemma 3 in Appendix B. To prove the almost surely convergence, assume that $a = kc$, for $k \in \{1, 2, \dots\}$ and a fixed positive number c . Then, for any $\epsilon > 0$, $Pr\{d_{AD}(P, H) > \epsilon\} \leq \frac{E_P(d_{AD}(P, H))^2}{\epsilon^2}$. Since, $\sum_{k=1}^{\infty} Pr\{d_{AD}(P, H) > \epsilon\} < \infty$, then, by the first Borel-Catelli lemma, $d_{AD} \xrightarrow{a.s.} 0$, as $k \rightarrow \infty$ ($a \rightarrow \infty$). ■

Interestingly, the limit of the expectation and variance in part (ii) of Corollary 4 coincide with the limit of $E_P(n d_{AD}(F_n, H))$ and $Var_P(n d_{AD}(F_n, H))$, given by Anderson and Darling [10], as $n \rightarrow \infty$, where F_n is the empirical cdf of the sample d .

According to part (i) of Corollary 4, with increasing the value of a , the prior distribution of the distance becomes more concentrated about 0. As recommended in Al-Labadi and Zarepour [8], the value of a should be at most $0.5n$, where n is the sample size; otherwise, the prior may become too influential. On the other hand, if $\mathcal{H}_0$ is true, with increasing the sample size n , the expectation of the posterior distribution of the distance between P_d and $F_{(m)}$ converges to zero. Also, if $\mathcal{H}_0$ is not true, this expectation converges to a positive value. The following Lemma indicates the effect of increasing the value of a and sample size n on the expectation of the posterior distribution of $d_{AD}(P_d, F_{(m)})$.

Lemma 5 *Let $P_d \sim DP(a + n, H_d)$ and H_d be as given by (2) with $H = F_{(m)}$, the cdf of the chi-square distribution with m degrees of freedom. We have*

- i. $d_{AD}(P_d, F_{(m)}) \xrightarrow{a.s.} 0$ as $a \rightarrow \infty$ (for fixed n).
- ii. If $\mathcal{H}_0$ is true, then, $d_{AD}(P_d, F_{(m)}) \xrightarrow{a.s.} 0$ as $n \rightarrow \infty$ (for fixed a).
- iii. If $\mathcal{H}_0$ is not true, then, $\liminf d_{AD}(P_d, F_{(m)}) \xrightarrow{a.s.} > 0$ as $n \rightarrow \infty$ (for fixed a).
- iv. If $\mathcal{H}_0$ is not true, then, $\liminf E_{P_d}(d_{AD}(P_d, F_{(m)})) \xrightarrow{a.s.} > 0$ as $n \rightarrow \infty$ (for fixed a).

Proof. The proof is given in Appendix C. ■

Part (i) of Lemma 5 confirms that for a too large value of a (relative to n), we may accept $\mathcal{H}_0$ even when $\mathcal{H}_0$ is not true. More discussion about this point is given in Section 7.

6 Computational algorithm

The steps for implementing the proposed test is detailed in this section. Note that, since no closed forms of the densities of $D = d_{AD}(P_N, F_{(m)})$ and $D_d = d_{AD}(P_{d_N}, F_{(m)})$

are available, simulation is used to approximate relative belief ratio. The following gives a computational algorithm to test $\mathcal{H}_0$. This algorithm is a revised version of Algorithm B of Al-Labadi and Evans [4].

Algorithm B (*Relative belief algorithm for testing multivariate normality*):

1. Use Algorithm A to generate (approximately) a P from $DP(a, F_{(m)})$.
2. Compute $d_{AD}(P_N, F_{(m)})$ as in Lemma 1.
3. Repeat steps (1)-(2) to obtain a sample of r_1 values from the prior of D .
4. Use Algorithm A to generate (approximately) a P_d from $DP(a + n, H_d)$.
5. Compute $d_{AD}(P_{d_N}, F_{(m)})$.
6. Repeat steps (4)-(5) to obtain a sample of r_2 values of D_d .
7. Let M be a positive number. Let $\hat{F}_D$ denote the empirical cdf of D based on the prior sample in step (3) and for $i = 0, \dots, M$, let $\hat{d}_{i/M}$ be the estimate of $d_{i/M}$, the (i/M) -th prior quantile of D . Here $\hat{d}_0 = 0$, and $\hat{d}_1$ is the largest value of the prior sample in step (3). Let $\hat{F}_D(\cdot | d)$ denote the empirical cdf of D_d based on the posterior sample in step (6). For $q \in [\hat{d}_{i/M}, \hat{d}_{(i+1)/M})$, estimate $RB_D(q | d) = \pi_D(q|d)/\pi_D(q)$ by

$$\widehat{RB}_D(q | d) = M\{\hat{F}_D(\hat{d}_{(i+1)/M} | d) - \hat{F}_D(\hat{d}_{i/M} | d)\}, \quad (9)$$

the ratio of the estimates of the posterior and prior contents of $[\hat{d}_{i/M}, \hat{d}_{(i+1)/M})$. It follows that, we estimate $RB_D(0 | d) = \pi_D(0|d)/\pi_D(0)$ by $\widehat{RB}_D(0 | d) = M\hat{F}_D(\hat{d}_{p_0} | d)$ where $p_0 = i_0/M$ and i_0 is chosen so that i_0/M is not too small (typically $i_0/M \approx 0.05$).

8. Estimate the strength $DP_D(RB_D(q | d) \leq RB_D(0 | d) | d)$ by the finite sum

$$\sum_{\{i \geq i_0 : \widehat{RB}_D(\hat{d}_{i/M} | d) \leq \widehat{RB}_D(0 | d)\}} (\hat{F}_D(\hat{d}_{(i+1)/M} | d) - \hat{F}_D(\hat{d}_{i/M} | d)). \quad (10)$$

For fixed M , as $r_1 \rightarrow \infty, r_2 \rightarrow \infty$, then $\hat{d}_{i/M}$ converges almost surely to $d_{i/M}$ and (9) and (10) converge almost surely to $RB_D(q | d)$ and $DP_D(RB_D(q | d) \leq RB_D(0 | d) | d)$,

respectively.

As recommended in the next section, one should try different values of a to make sure the right conclusion has been obtained. The consistency of the proposed test is achieved by Proposition 6 in Al-Labadi and Evans [4].

7 Numerical results

7.1 Simulation Study

In this subsection, the performance of the proposed test is illustrated through several examples. Hereafter, let $\mathbf{c}_m$ be the m -dimensional column vectors of c 's, I_m be the $m \times m$ identity matrix, A_m be an $m \times m$ matrix with 1's on the main diagonal and 0.1's elsewhere, and B_m be an $m \times m$ matrix with 0.25's on the main diagonal and 0.2's elsewhere. The following notations have been used: t_r for the t -Student distribution with r degrees of freedom, χ_r^2 for the chi-square distribution with r degrees of freedom, $E(\lambda)$ for the exponential distribution with rate λ , $C(a, b)$ for the Cauchy distribution with location parameter a and scale parameter b , $P_{VII}(1, 1, r)$ for the Pearson type VII (aka t -Student) distribution with location parameter 1, scale parameter 1 and r degrees of freedom, $t_r(\mathbf{0}_m, I_m)$ for the m -variate t -student distribution with location parameter $\mathbf{0}_m$, scale parameter I_m and r degrees of freedom, $LN_m(\mathbf{0}_m, B_m)$ for the m -variate lognormal distribution with mean vector $\mathbf{0}_m$ and covariance matrix B_m , $\mathcal{S}^m(Q)$ for the m -variate spherical distribution with distribution Q for radii, and $NMIX$ for the mixture distribution $0.9N_m(\mathbf{5}_m, A_m) + .1N_m(-\mathbf{5}_m, A_m)$. Also, for a univariate distribution Q , $(Q)^m$ denotes an m -variate distribution with m identical and independent marginals Q , while, $Q_1 \otimes \cdots \otimes Q_m$ denotes an m -variate distribution with m independent marginals $Q_1, \dots, Q_m$. In all cases, we generate a sample of size 50 from an exact distribution and then we set $N = 500$, $r_1 = r_2 = 1000$ and $M = 20$ in Algorithm B to compute the RB and the corresponding strength. In addition, to study the sensitivity of the approach to the choice of the chosen concentration parameter, various values of

a are considered. The results are reported in Table 1. The RB of the proposed test is compared to the Bayes factor BF^{TM} due to Tokdar and Martin [43]. Recall that, similar to RB , $BF^{\text{TM}} > 1$ supports $\mathcal{H}_0$, $BF^{\text{TM}} < 1$ supports $\mathcal{H}_1$ and $BF^{\text{TM}} = 1$ means no evidence either way. To compute BF^{TM} , we used the code provided by the same authors and posted at <http://www2.stat.duke.edu/~st118/Software/>. From Table 1, it is seen that for the cases where $\mathcal{H}_0$ is true, the value of RB is greater than 1 and the strength of this evidence is 1. This shows a strong evidence to accept $\mathcal{H}_0$. For instance, when $N_2(\mathbf{0}_2, A_2)$ and $a = 15$ in Table 1, the values of RB and its relevant strength are 7.36 and 1, respectively. The prior and posterior density plots of the Anderson-Darling distance for this example are provided by the left side of part (a) of Figure 1 to show that the posterior density of the Anderson-Darling distance is more concentrated around zero than the prior density of the Anderson-Darling distance, when $\mathcal{H}_0$ is true. Also, the Q-Q plot (see Appendix D) of square ordered Mahalanobis distances related to the generated sample $\mathbf{y}_{2 \times n}$ against the square root of the percentage points of the chi-square distribution with $m = 2$ degrees of freedom is provided by the right side of part (a) of Figure 1. This figure shows the close relationship between the distribution of observed points and chi-square distribution with 2 degrees of freedom for this example. Notice that, $(P_{VII}(1, 1, r))^m$, for $r \geq 10$ is a symmetric distribution and has very similar behavior with an m -variate normal distribution, and then $\mathcal{H}_0$ is accepted in all cases of this example. For the cases where $\mathcal{H}_0$ is not true, the value of RB is less than 1 and its relevant strength is small for $a > 1$. For example, when $t_3(\mathbf{0}_2, I_2)$ and $a = 15$, the value of RB and its relevant strength are 0.073 and 0.085, respectively. The prior and posterior density plots of the Anderson-Darling distance are also provided by the left side of part (b-d) of Figure 1 for various alternative examples to show that the prior density of the Anderson-Darling distance is more concentrated around zero than the posterior density of the Anderson-Darling distance, when $\mathcal{H}_0$ is not true. Note that, while the proposed test rejects $\mathcal{H}_0$ in all examples with $a = 15$, the test of Tokdar and Martin [43] fails to reject $\mathcal{H}_0$ in some case. See, for example, the results

for $\mathcal{S}^m(LN(0, 0.25))$, $\mathcal{S}^m(\chi_5^2)$ and $t_3(\mathbf{0}_m, I_m)$ when $a = 15$ and $m > 2$.

Table 1: Relative belief ratios and strength (str)-s for testing the m -variate normality assumption with various alternatives and choices of a .

True distribution	a	$m = 2$		$m = 3$		$m = 4$	
		$RB(\text{str})$	BF^{TM}	$RB(\text{str})$	BF^{TM}	$RB(\text{str})$	BF^{TM}
$N_m(\mathbf{0}_m, A_m)$	1	19.38(1)	0.197	17.12(1)	10^{-7}	18.88(1)	2×10^{-41}
	5	14.52(1)	7.039	13.73(1)	8×10^{-6}	12.99(1)	5×10^{-8}
	10	9.38(1)	91.50	7.09(1)	2×10^{11}	8.41(1)	10^{22}
	15	7.36(1)	590.2	7.92(1)	4×10^{13}	7.88(1)	4×10^{22}
$E(\frac{1}{2}) \otimes (C(0, 1))^{m-1}$	1	0.24(0.012)	3×10^{-14}	0(0)	6×10^{-30}	0(0)	10^{-24}
	5	0.04(0.004)	8×10^{-19}	0(0)	10^{-50}	0(0)	5×10^{-35}
	10	0(0)	2×10^{-20}	0(0)	2×10^{-40}	0(0)	8×10^{-28}
	15	0(0)	2×10^{-18}	0(0)	4×10^{-36}	0(0)	2×10^{-6}
$N(0, 1) \otimes (t_1)^{m-1}$	1	0.76(0.054)	2×10^{-33}	0.1(0.003)	10^{-28}	0(0)	7×10^{-34}
	5	0.70(0.067)	3×10^{-50}	0(0)	10^{-35}	0(0)	9×10^{-45}
	10	0.68(0.058)	3×10^{-34}	0(0)	3×10^{-25}	0(0)	10^{-28}
	15	0.60(0.055)	2×10^{-31}	0(0)	6×10^{-15}	0(0)	10^{-23}
$(P_{VII}(1, 1, 1))^{m\dagger}$	1	0(0)	10^{-9}	0(0)	10^{-20}	0(0)	10^{-39}
	5	0(0)	2×10^{-11}	0(0)	6×10^{-31}	0(0)	8×10^{-45}
	10	0(0)	3×10^{-12}	0(0)	7×10^{-26}	0(0)	4×10^{-27}
	15	0(0)	10^{-10}	0(0)	3×10^{-19}	0(0)	8×10^{-13}
$(P_{VII}(1, 1, 10))^m$	1	19.92(1)	7×10^{-2}	19.82(1)	8×10^{-7}	19.78(1)	10^{-5}
	5	17.68(1)	3×10^2	16.02(1)	5×10^5	15.84(1)	10^{16}
	10	12.88(1)	3×10^4	12.76(1)	6×10^{12}	11.06(1)	6×10^{26}
	15	9.62(1)	5×10^4	10.36(1)	10^{14}	9.80(1)	7×10^{28}
$\mathcal{S}^m(LN(0, 0.25))^{\ddagger}$	1	1.76(0.321)	4×10^{-6}	2.12(0.376)	10^{-10}	1.92(0.363)	10^{-12}
	5	0.36(0.002)	5×10^{-5}	0.54(0.028)	4×10^4	0.16(0.008)	10^{14}
	10	0.11(0.001)	10^{-3}	0.16(0.001)	10^{11}	0.04(0.003)	3×10^{23}
	15	0.01(0.001)	9×10^{-3}	0.02(0.001)	10^{14}	0.06(0.004)	6×10^{21}
$\mathcal{S}^m(\chi_5^2)$	1	2.31(0.341)	0.408	2.11(0.393)	10^{-8}	1.22(0.308)	3×10^{-38}
	5	0.86(0.155)	294.02	0.78(0.143)	296.66	0.34(0.002)	7×10^8
	10	0.56(0.028)	4459.4	0.22(0.011)	10^{10}	0.12(0.003)	5×10^{20}
	15	0.32(0.001)	31538	0.1(0.005)	7×10^{11}	0.05(0)	10^{19}
$LN_m(\mathbf{0}_m, B_m)^*$	1	1.82(0.301)	3×10^{-36}	1.69(0.359)	2×10^{-137}	1.92(0.340)	0
	5	0.22(0.022)	3×10^{-64}	0.06(0.003)	10^{-158}	0.26(0.006)	6×10^{-232}
	10	0.08(0.004)	7×10^{-46}	0.02(0.001)	2×10^{-147}	0.02(0.001)	8×10^{-56}
	15	0.06(0.003)	2×10^{-33}	0.02(0.003)	10^{-76}	0.01(0.008)	3×10^{-34}
$t_3(\mathbf{0}_m, I_m)^{\ddagger}$	1	1.42(0.328)	3×10^{-8}	1.63(0.311)	10^{-5}	1.81(0.337)	9×10^{-24}
	5	0.88(0.105)	3×10^{-5}	0.90(0.124)	10^3	0.82(0.111)	10^{14}
	10	0.81(0.097)	10^{-3}	0.70(0.090)	3×10^8	0.66(0.053)	9×10^{23}
	15	0.73(0.085)	5×10^{-3}	0.50(0.021)	10^{12}	0.43(0.006)	2×10^{27}
$NMIX^{\ddagger}$	1	1.68(0.281)	10^{-53}	2.13(0.312)	0	2.18	0
	5	0.12(0.006)	9×10^{-48}	0.54(0.121)	0	0.67(0.152)	0
	10	0.06(0.003)	10^{-50}	0.23(0.026)	0	0.34(0.006)	0
	15	0(0)	3×10^{-41}	0.09(0.010)	0	0.02(0.007)	0

[†]Required R package to generate sampled data: PearsonDS.

[‡]Required R package to generate sampled data: distrEllipse.

*Required R package to generate sampled data: compositions.





(a) $N_2(\mathbf{0}_2, A_2)$ (b) $(P_{VII}(1, 1, 1))^2$ (c) $t_3(\mathbf{0}_2, I_2)$ (d) $N(0, 1) \otimes t_1$

Figure 1: Left: The solid line represents the prior density of the Anderson-Darling distance and the dashed line represents the posterior density of the Anderson-Darling distance for $a = 15$. Right: Chi-square Q-Q plot for the simulated sample with 95% envelope.

We end this subsection by investigating the effect of the choice of H in the proposed test. We consider the results of the proposed test for a sample of size 50 from true distributions $N_2(\mathbf{0}_2, A_2)$, $t_3(\mathbf{0}_2, I_2)$ and $N(0, 1) \otimes t_1$ for different choices of H in Table 2. Clearly, when $\mathcal{H}_0$ is not true, the results are correct for only $H = \chi_2^2$; otherwise, they are incorrect.

Table 2: $RB(\text{Strength})$ of a sample of size 50 when there is prior-data conflict (a tiny overlap between the effective support regions).

True distribution	H	$a = 5$	$a = 10$	$a = 15$
$N_2(\mathbf{0}_2, A_2)$	χ_2^2	14.52(1)	9.38(1)	7.36(1)
	$C(0, 1)$	20(1)	20(1)	20(1)
	$N(0, 1)$	20(1)	20(1)	20(1)
$t_3(\mathbf{0}_2, I_2)$	χ_2^2	0.88(0.105)	0.81(0.097)	0.73(0.087)
	$C(0, 1)$	20(1)	20(1)	19.38(1)
	$N(0, 1)$	20(1)	20(1)	20(1)
$N(0, 1) \otimes t_1$	χ_2^2	0.76(0.054)	0.70(0.067)	0.68(0.058)
	$C(0, 1)$	20(1)	20(1)	20(1)
	$N(0, 1)$	20(1)	20(1)	20(1)

7.2 Application

In this subsection, we look at the performance of the methodology by using a real data set. For comparison purposes, the results of the Doornik-Hanson (DH) test are also presented [17]. The R package **asbio** is used to compute p-values of the DH test.

We consider the data set of the National track records for women which reprints women's athletic records for 55 countries with seven variables (Atkinson et al. [11]). The variables are 100, 200, 400 meters in seconds, 800, 1500, 3000 meters in minutes and the marathon. The problem is to assess the seven-variate normality assumption for this data set. Atkinson et al. [11] provided the Q-Q plot (see the right side of Figure 2) of squared ordered Mahalanobis distances against the square root of the percentage points of the chi-square distribution with seven degrees of freedom and then suggested the non-normality of this data set. The DH's p-value for this data is 4.042382×10^{-68} ,

which shows strong evidence to reject the multivariate normality assumption. The results of the proposed test are given in Table 3, which show a strong evidence to reject the assumption of multivariate normality .

Table 3: Relative belief ratios and strength (str)-s for testing the seven-variate normality assumption of national track records for women with various choices of a .

	a					
Real data	1	5	6	8	10	15
$RB(str)$	7.480(0.626)	1.140(0.405)	0.700(0.133)	0.480(0.063)	0.240(0.012)	0.120(0.006)




Figure 2: Left: The solid line represents the prior density of the Anderson-Darling distance and the dashed line represents the posterior density of the Anderson-Darling distance for $a = 15$. Right: Chi-square Q-Q plot of the real data set with 95% envelope.

8 Conclusion

A novel Bayesian nonparametric test for assessing multivariate normality has been proposed. The suggested test is developed by using Mahalanobis distance as a good technique to convert the m -variate problem into the univariate problem. The Dirichlet process has been considered as a prior on the distribution of the Mahalanobis distance. Then, the concentration of the distribution of the distance between posterior process and chi-square distribution with m degrees of freedom is compared to the concentration of the distribution of the distance between prior process and chi-square distribution

with m degrees of freedom via relative belief ratio. The distance between the Dirichlet process and the chi-square distribution is developed based on the Anderson-Darling distance. Several theoretical results including consistency have been discussed for the proposed test. The test is illustrated through various simulated examples and a real data set.

Appendix A Proof of Lemma 1

Consider $P_N(x)$ as

$$P_N(x) = \begin{cases} 0 & x < Y_{(1)} \\ P_N(Y_{(i)}) & Y_{(i)} \leq x < Y_{(i+1)}, (i = 1, \dots, N-1). \\ 1 & x \geq Y_{(N)} \end{cases}$$

Let $g(x) = \frac{dG(x)}{dx}$, then

$$\begin{aligned} d_{AD}(P_N, G) &= \int_{-\infty}^{\infty} \frac{(P_N(x) - G(x))^2}{G(x)(1 - G(x))} g(x) dx \\ &= \int_{-\infty}^{Y_{(1)}} \frac{G(x)^2}{G(x)(1 - G(x))} g(x) dx + \int_{Y_{(N)}}^{\infty} \frac{(1 - G(x))^2}{G(x)(1 - G(x))} g(x) dx \\ &\quad + \sum_{i=1}^{N-1} \int_{Y_{(i)}}^{Y_{(i+1)}} \frac{(P_N(Y_{(i)}) - G(x))^2}{G(x)(1 - G(x))} g(x) dx. \end{aligned}$$

Substituting $y = G(x)$, $G(Y_{(i)}) = U_{(i)}$, $G(-\infty) = 0$ and $G(\infty) = 1$, gives

$$\begin{aligned}
d_{AD}(P_N, G) &= \sum_{i=1}^{N-1} \int_{U_{(i)}}^{U_{(i+1)}} \frac{(P_N(Y_{(i)}) - y)^2}{y(1-y)} dy + \int_0^{U_{(1)}} \frac{y}{1-y} dy + \int_{U_{(N)}}^1 \frac{1-y}{y} dy \\
&= \sum_{i=1}^{N-1} \left[P_N^2(Y_{(i)}) \log(y) - \left[(P_N(Y_{(i)}) - 1)^2 \log(1-y) \right] - y \right]_{U_{(i)}}^{U_{(i+1)}} \\
&\quad + \left[-y - \log(1-y) \right]_0^{U_{(1)}} + \left[\log(y) - y \right]_{U_{(N)}}^1 \\
&= I_1 + I_2 + I_3.
\end{aligned}$$

Note that,

$$\begin{aligned}
I_1 &= - \sum_{i=1}^{N-1} (U_{(i+1)} - U_{(i)}) - \sum_{i=1}^{N-1} (P_N(Y_{(i)}) - 1)^2 (\log(1 - U_{(i+1)}) - \log(1 - U_{(i)})) \\
&\quad + \sum_{i=1}^{N-1} P_N^2(Y_{(i)}) (\log(U_{(i+1)}) - \log(U_{(i)})) \\
&= \sum_{i=1}^{N-1} P_N^2(Y_{(i)}) \log \frac{U_{(i+1)} (1 - U_{(i)})}{U_{(i)} (1 - U_{(i+1)})} + \sum_{i=1}^{N-1} (2P_N(Y_{(i)}) - 1) \log \frac{1 - U_{(i+1)}}{1 - U_{(i)}} \\
&\quad - (U_{(N)} - U_{(1)}).
\end{aligned}$$

Also, $I_2 = -U_{(1)} - \log(1 - U_{(1)})$ and $I_3 = -1 - \log U_{(N)} + U_{(N)}$. Therefore, adding I_1 , I_2 and I_3 , gives

$$\begin{aligned}
d_{AD}(P_N, G) &= \sum_{i=1}^{N-1} P_N^2(Y_{(i)}) \log \frac{U_{(i+1)} (1 - U_{(i)})}{U_{(i)} (1 - U_{(i+1)})} + \sum_{i=1}^{N-1} (2P_N(Y_{(i)}) - 1) \log \frac{1 - U_{(i+1)}}{1 - U_{(i)}} \\
&\quad - 1 - \log(U_{(N)}(1 - U_{(1)})). \tag{11}
\end{aligned}$$

The proof is completed by substituting $P_N(Y_{(i)}) = \sum_{j=1}^i J'_j$, $P_N^2(Y_{(i)}) = \sum_{j=1}^i J_j'^2 + 2 \sum_{j=1}^{i-1} \sum_{k=j+1}^i J'_k J'_j$ and $U_{(i)} = G(Y_{(i)})$ in terms on the right-hand side of (11).

Appendix B Proof of Lemma 3

To prove (i), note that, from the property of the Dirichlet process, for any $t \in \mathbb{R}$, $E_P(P(t) - H(t))^2 = \frac{H(t)(1-H(t))}{a+1}$. Then

$$E_P(d_{AD}(P, H)) = \int_{-\infty}^{\infty} \frac{E_P(P(t) - H(t))^2}{H(t)(1-H(t))} dH(t) = \frac{1}{a+1}.$$

To prove (ii), it is enough to compute $E_P(d_{AD}(P, H))^2$. According to the Corollary 2, we consider H to be the cdf of the Uniform distribution on $[0, 1]$. Then

$$\begin{aligned} E_P(d_{AD}(P, H))^2 &= E_P \left(\int_{-\infty}^{\infty} \frac{(P(t) - H(t))^2}{H(t)(1-H(t))} dH(t) \right)^2 \\ &= E_P \left(\int_0^1 \frac{(P(t) - t)^2}{t(1-t)} dt \int_0^1 \frac{(P(s) - s)^2}{s(1-s)} ds \right) \\ &= E_P \left(\int_0^1 \int_0^t \frac{(P(t) - t)^2 (P(s) - s)^2}{t(1-t)s(1-s)} ds dt \right. \\ &\quad \left. + \int_0^1 \int_0^s \frac{(P(t) - t)^2 (P(s) - s)^2}{t(1-t)s(1-s)} dt ds \right) \\ &= 2E_P \left(\int_0^1 \int_0^t \frac{(P(t) - t)^2 (P(s) - s)^2}{t(1-t)s(1-s)} ds dt \right) \\ &= 2 \int_0^1 \int_0^t \frac{E_P \{ (P(s) + P((s, t]) - t)^2 (P(s) - s)^2 \}}{t(1-t)s(1-s)} ds dt. \end{aligned}$$

Note that, from the property of the Dirichlet process, for any $s < t$ and $i, j \in \mathbb{N}$, $E_P(P^i(s)P^j((s, t])) = \frac{\Gamma(a)}{\Gamma(a+i+j)} \frac{\Gamma(as+i)}{\Gamma(as)} \frac{\Gamma(a(t-s)+j)}{\Gamma(a(t-s))}$ and $E_P(P^i(s)) = \prod_{k=0}^{i-1} \frac{as+k}{a+k}$. Then

$$\begin{aligned}
E_P(d_{AD}(P, H))^2 &= \int_0^1 \int_0^t \frac{1}{ts(1-t)(1-s)} \left\{ \frac{2as(a(t-s)+1)(as+1)(t-s)}{(a+3)(a+2)(a+1)} \right. \\
&+ \frac{4as(as+2)(as+1)(t-s)}{(a+3)(a+2)(a+1)} + \frac{2s(as+3)(as+2)(as+1)}{(a+3)(a+2)(a+1)} \\
&- \frac{4as(2s+t)(as+1)(t-s)}{(a+2)(a+1)} - \frac{4as^2(a(t-s)+1)(t-s)}{(a+2)(a+1)} \\
&- \frac{4s(as+2)(as+1)(s+t)}{(a+2)(a+1)} + \frac{2s(s^2+4st+t^2)(as+1)}{a+1} \\
&+ \frac{4as^2(s+2t)(t-s)}{a+1} + \frac{2s^2(t-s)(a(t-s)+1)}{a+1} - 4s^2t(t-s) \\
&\left. - 4s^2t(s+t) + 2s^2t^2 \right\} ds dt.
\end{aligned}$$

After simplification, we get

$$\begin{aligned}
E_P(d_{AD}(P, H))^2 &= \int_0^1 \int_0^t \frac{2((a-6)((3t-2)s-t)-6)}{t(s-1)(a+3)(a+2)(a+1)} ds dt \\
&= \int_0^1 \frac{2}{t(a+3)(a+2)(a+1)} \left\{ (a-6)(3t-2)t - 2i\pi((a-6)t-a+3) \right. \\
&\quad \left. + 2(a(t-1)-6t+3)\log(t-1) \right\} dt \\
&= \frac{a(2\pi^2-15)-6(\pi^2-15)}{3(a+3)(a+2)(a+1)},
\end{aligned}$$

for $\text{Re}(t) < 1$ or $t \notin \mathbb{R}$, where $\text{Re}(t)$ denotes the real part of t and i is the imaginary unit. Then, the variance of $d_{AD}(P, H)$ is given by

$$\begin{aligned}
\text{Var}_P(d_{AD}(P, H)) &= E_P(d_{AD}(P, H))^2 - E_P^2(d_{AD}(P, H)) \\
&= \frac{2((\pi^2-9)a^2+(30-2\pi^2)a-3\pi^2+36)}{3(a+1)^2(a+2)(a+3)}.
\end{aligned}$$

Hence, the proof is completed.

Appendix C Proof of Lemma 5

Assume that $d = (d_M^2(\mathbf{y}_1), \dots, d_M^2(\mathbf{y}_n))$ is the observed square Mahalanobis distance from P where $P \sim DP(a, F_{(m)})$. Note that

$$\begin{aligned} d_{AD}(P_d, F_{(m)}) &= \int_{-\infty}^{\infty} \frac{(P_d(t) - F_{(m)}(t))^2}{F_{(m)}(t)(1 - F_{(m)}(t))} dF_{(m)}(t) \\ &\leq \left(\sup_{t \in \mathbb{R}} |P_d(t) - F_{(m)}(t)| \right)^2 \int_{-\infty}^{\infty} \frac{1}{F_{(m)}(t)(1 - F_{(m)}(t))} dF_{(m)}(t) \\ &\leq \sup_{t \in \mathbb{R}} |P_d(t) - F_{(m)}(t)| \int_{-\infty}^{\infty} \frac{1}{F_{(m)}(t)(1 - F_{(m)}(t))} dF_{(m)}(t) \\ &\leq \int_{-\infty}^{\infty} \frac{1}{F_{(m)}(t)(1 - F_{(m)}(t))} dF_{(m)}(t) \left\{ \sup_{t \in \mathbb{R}} |P_d(t) - H_d(t)| + \right. \\ &\quad \left. \sup_{t \in \mathbb{R}} |H_d(t) - F_{(m)}(t)| \right\}, \end{aligned}$$

where the third inequality holds since $0 \leq |P_d(t) - F_{(m)}(t)| \leq 1$ and the fourth inequality holds by triangle inequality. To prove (i), as $a \rightarrow \infty$, from James [31], $\sup_{t \in \mathbb{R}} |P_d(t) - H_d(t)| \xrightarrow{a.s.} 0$ and by the continuous mapping theorem $\sup_{t \in \mathbb{R}} |H_d(t) - F_{(m)}(t)| \xrightarrow{a.s.} 0$. To prove (ii), since $\sup_{t \in \mathbb{R}} |P_d(t) - H_d(t)| \xrightarrow{a.s.} 0$ as $n \rightarrow \infty$ and $\mathcal{H}_0$ is true, the continuous mapping theorem and Polya's theorem [16] imply $\sup_{t \in \mathbb{R}} |H_d(t) - F_{(m)}(t)| \xrightarrow{a.s.} 0$. Note that, the final results of part (i) and (ii) are concluded by practical assumptions in probability and measure theory given in Section 3.1 of Capiński and Kopp [14].

To prove (iii), note that, $(F_{(m)}(t)(1 - F_{(m)}(t)))^{-1} \geq 4$, then

$$\begin{aligned} d_{AD}(P_d, F_{(m)}) &\geq 4 \int_{-\infty}^{\infty} (P_d(t) - F_{(m)}(t))^2 dF_{(m)}(t) \\ &\geq 4 d_{CvM}(P_d, F_{(m)}). \end{aligned}$$

From Choi and Bulgren [15], since $d_{CvM}(P_d, F_{(m)}) \geq \frac{1}{3} \left(\sup_{t \in \mathbb{R}} |P_d(t) - F_{(m)}(t)| \right)^3$,

$$d_{AD}(P_d, F_{(m)}) \geq \frac{4}{3} \left(\sup_{t \in \mathbb{R}} |P_d(t) - F_{(m)}(t)| \right)^3.$$

Using the triangle inequality gives

$$\sup_{t \in \mathbb{R}} |P_d(t) - F_{(m)}(t)| \geq \sup_{t \in \mathbb{R}} |H_d(t) - F_{(m)}(t)| - \sup_{t \in \mathbb{R}} |P_d(t) - H_d(t)|.$$

Similar to the proof of part (ii), as $n \rightarrow \infty$, $\sup_{t \in \mathbb{R}} |P_d(t) - H_d(t)| \xrightarrow{a.s.} 0$ and $\sup_{t \in \mathbb{R}} |H_d(t) - F_{(m)}(t)| \xrightarrow{a.s.} \sup_{t \in \mathbb{R}} |P_{true}(t) - F_{(m)}(t)|$, where P_{true} is the true distribution of the sample d . Since $\mathcal{H}_0$ is not true, $\liminf_{t \in \mathbb{R}} \sup_{t \in \mathbb{R}} |P_d(t) - F_{(m)}(t)| \xrightarrow{a.s.} > 0$, which implies $\liminf d_{AD}(P_d, F_{(m)}) \xrightarrow{a.s.} > 0$.

To prove (iv), since for any $t \in \mathbb{R}$, $E_{P_d}(P_d(t)) = H_d(t)$ and $E_{P_d}(P_d(t) - H_d(t))^2 = \frac{H_d(t)(1-H_d(t))}{a+n+1}$, then, as $n \rightarrow \infty$

$$\begin{aligned} \liminf E_{P_d}(d_{AD}(P_d, F_{(m)})) &\geq \liminf \int_{-\infty}^{\infty} \frac{H_d(t)(1-H_d(t))}{(a+n+1)F_{(m)}(t)(1-F_{(m)}(t))} dF_{(m)}(t) \\ &\quad + \liminf \int_{-\infty}^{\infty} \frac{(H_d(t) - F_{(m)}(t))^2}{F_{(m)}(t)(1-F_{(m)}(t))} dF_{(m)}(t). \end{aligned}$$

Applying Fatou's lemma gives

$$\begin{aligned} \liminf E_{P_d}(d_{AD}(P_d, F_{(m)})) &\geq \int_{-\infty}^{\infty} \liminf \left(\frac{H_d(t)(1-H_d(t))}{(a+n+1)F_{(m)}(t)(1-F_{(m)}(t))} \right) dF_{(m)}(t) \\ &\quad + \int_{-\infty}^{\infty} \liminf \left(\frac{(H_d(t) - F_{(m)}(t))^2}{F_{(m)}(t)(1-F_{(m)}(t))} \right) dF_{(m)}(t) \\ &= 0 + \int_{-\infty}^{\infty} \inf \left(\frac{(P_{true}(t) - F_{(m)}(t))^2}{F_{(m)}(t)(1-F_{(m)}(t))} \right) dF_{(m)}(t). \end{aligned}$$

Since $\mathcal{H}_0$ is not true, $\inf \left(\frac{(P_{true}(t) - F_{(m)}(t))^2}{F_{(m)}(t)(1-F_{(m)}(t))} \right) \xrightarrow{a.s.} > 0$. Hence, by Theorem 15.2 of Billingsley [12], $\liminf E_{P_d}(d_{AD}(P_d, F_{(m)})) \xrightarrow{a.s.} > 0$.

Appendix D Chi-square Q-Q plot

Chi-square Q-Q plot shows the relationship between the distribution of data points and chi-square distribution. It is a common graphical method to assess multivariate normality. Specifically, for a given m -variate sampled data $\mathbf{y}_{m \times n} = (\mathbf{y}_1, \dots, \mathbf{y}_n)$ with

the size of n , where $\mathbf{y}_i \in \mathbb{R}^m$ for $i = 1, \dots, n$, chi-square Q-Q plot provide the plot of square ordered Mahalanobis distances against the quantile points of the chi-square distribution with m degrees of freedom. From Johnson and Wichern [32], if $\mathbf{y}_{m \times n}$ comes from an m -variate normal distribution, we should see the points of square ordered Mahalanobis distances are nearly a straight line with unit slope (reference line).

In the main paper, we showed that the chi-square Q-Q plot confirms the result of the proposed test. For this, we provided the chi-square Q-Q plot with its approximate confidence envelope around the reference line for the data set of examples of Section 7 in the right side of Figure 1 and Figure 2. Note that, the confidence envelope is plotted to show the central intervals for each quantile of the chi-square distribution with m degrees of freedom. In fact, the envelope in Figure 1 and 2 provides a density estimate of the quantiles drawn from the chi-square distribution with m degrees of freedom. For more detailed studies about the general methods for Q-Q plot see Oldford [39].

References

- [1] Al-Labadi L., Baskurt Z., and Evans M. (2017). Goodness of fit for the logistic regression model using relative belief. *Journal of Statistical Distributions and Applications*, 4(1), 1.
- [2] Al-Labadi L., Baskurt Z., and Evans M. (2018). Statistical reasoning: choosing and checking the ingredients, inferences based on a measure of statistical evidence with some applications. *Entropy*, 20(4), 289.
- [3] Al-Labadi L. and Evans M. (2017). Optimal robustness results for relative belief inferences and the relationship to prior-data conflict. *Bayesian Analysis*, 12(3), 705–728.
- [4] Al-Labadi L. and Evans M. (2018). Prior-based model checking. *Canadian Journal of Statistics*, 46(3), 380–398.

- [5] Al-Labadi L. and Zarepour M. (2013). A Bayesian nonparametric goodness of fit test for right censored data based on approximate samples from the beta-Stacy process. *The Canadian Journal of Statistics*, 41(3), 466–487.
- [6] Al-Labadi L., and Zarepour M. (2014a). Goodness of fit tests based on the distance between the Dirichlet process and its base measure. *Journal of Nonparametric Statistics*, 26(2), 341–357.
- [7] Al-Labadi L., and Zarepour M. (2014b). Simulations from the Two-Parameter Poisson-Dirichlet Process and the Normalized Inverse-Gaussian Process. *Sankhyā A*, 76(1), 158–176.
- [8] Al-Labadi L. and Zarepour M. (2017). Two-sample Kolmogorov-Smirnov test using a Bayesian nonparametric approach. *Mathematical Methods of Statistics*, 26(3), 212–225.
- [9] Alva J. A. V. and Estrada E. G. (2009). A generalization of Shapiro-Wilks test for multivariate normality. *Communications in Statistics-Theory and Methods*, 38(11-12), 1870–1883.
- [10] Anderson T. W., and Darling D. A. (1954). A test of goodness of fit. *Journal of the American Statistical Association*, 49, 765–769.
- [11] Atkinson A. C., Riani M., and Cerioli A. (2004). *Exploring multivariate data with the forward search*. Springer, New York.
- [12] Billingsley P. (1995). *Probability and Measure*. Wiley.
- [13] Bondesson L. (1982). On simulation from infinitely divisible distributions. *Advances in Applied Probability*, 14(4), 885–869.
- [14] Capiński M., and Kopp E. (2004). *Measure, Integral and Probability*. Springer Undergraduate Mathematics Series. Springer, Berlin, second edition.

- [15] Choi K. , and Bulgren W. G. (1988). An estimation procedure for mixtures of distributions. *Journal of the Royal Statistical Society, B*, 30, 444–460.
- [16] Dasgupta A. (2008). *Asymptotic Theory of Statistics and Probability*. Springer, New York.
- [17] Doornik J. and Hansen H. (2008). An omnibus test for univariate and multivariate normality. *Oxford bulletin of economics and statistics*, 70(1), 927–939.
- [18] Dubes R. and Jain A. K. (1980). Clustering methodologies in exploratory data analysis. *Advances in Computers*, 19, 113–228.
- [19] Evans M. (1997). Bayesian inference procedures derived via the concept of relative surprise. *Communications in Statistics-Theory and Methods*, 26(5), 1125–1143.
- [20] Evans M. (2015). *Measuring Statistical Evidence Using Relative Belief*. volume 144 of *Monographs on Statistics and Applied Probability*. CRC Press, Boca Raton, FL.
- [21] Evans M. and Moshonov H. (2006). Checking for prior-data conflict. *Bayesian Analysis*, 1(4), 893–914.
- [22] Fattorini L. (1986). Remarks on the use of Shapiro-Wilk statistic for testing multivariate normality. *Statistica*, 46(2), 209–217.
- [23] Ferguson T. S. (1973). A Bayesian analysis of some nonparametric problems. *The Annals of Statistics*, 1, 209–230.
- [24] Fernandez G. (2010). *Data mining using SAS applications*. USA: CRC Press, Boca Raton, FL, second edition.
- [25] Hanusz Z. and Tarasińska J. (2012). New test for multivariate normality based on Small’s and Srivastava’s graphical methods. *Journal of Statistical Computation and Simulation*, 82(12), 1743–1752.

- [26] Hasofer A. M. and Stein G. Z. (1990). Testing for multivariate normality after coordinate transformation. *Communications in Statistics-Theory and Methods*, 19(4), 1403–1418.
- [27] Healy M. J. R. (1968). Multivariate normal plotting, *Applied Statistics*, 17(2), 157–161.
- [28] Henze N. and Visagie J. (2019). Testing for normality in any dimension based on a partial differential equation involving the moment generating function. *Annals of the Institute of Statistical Mathematics*. <https://doi.org/10.1007/s10463-019-00720-8>
- [29] Henze N. and Zirkler B. (1990). A class of invariant consistent tests for multivariate normality. *Communications in Statistics-Theory and Methods*, 19(10), 3595–3617.
- [30] Holgersson H. (2006). A graphical method for assessing multivariate normality. *Computational Statistics*. 21(1), 141–149.
- [31] James L. F. (2008). Large sample asymptotics for the two-parameter Poisson-Dirichlet process. In *Pushing the Limits of Contemporary Statistics: Contributions in Honor of Jayanta K. Ghosh*, ed. B. Clarke and S. Ghosal, Ohio: Institute of Mathematical Statistics, 187–199.
- [32] Johnson R. A. and Wichern D. W. (2007). *Applied multivariate statistical analysis*. Pearson Prentice Hall, Upper Saddle River, New Jersey: CRC Press, 6th edition.
- [33] Jönsson K. (2011). A robust test for multivariate normality. *Economics Letters*. 113(2), 199–201.
- [34] Kim I. and Park S. (2018). Likelihood ratio tests for multivariate normality. *Communications in Statistics-Theory and Methods*, 47(8), 1923,1934.
- [35] Kim N. (2016). A robustified Jarque-Bera test for multivariate normality. *Economic Letter*, 140, 48–52.

- [36] Madukaife M. S. and Okafor F. C. (2018). A powerful affine invariant test for multivariate normality based on interpoint distances of principal components. *Communications in Statistics-Simulation and Computation*, 47(5), 1264–1275.
- [37] Mardia K.V. (1970). Measures of multivariate skewness and kurtosis with applications. *Biometrika*, 57(3), 519–530.
- [38] Nott D. J., Seah M., AL-Labadi L., Evans M. Ng, H. K., and Englert B. (2020). Using prior expansions for prior-data conflict checking. *Bayesian Analysis*. <https://projecteuclid.org/euclid.ba/1585360930>.
- [39] Oldford R.W. (2016). Self-calibrating quantile-quantile plots. *The American Statistician*, 70, 74–90.
- [40] Rincón-Gallardo S., Quesenberry C. P. , and O'Reilly F. J. (1979). Conditional probability integral transformations and goodness-of-fit tests for multivariate normal distributions. *The Annals of Statistics*, 7(5), 1052–1057.
- [41] Royston J. P. (1983). Some techniques for assessing multivariate normality based on the Shapiro-Wilk W. *Applied Statistics*, 32(2), 121–133.
- [42] Sethuraman J. (1994). A constructive definition of Dirichlet priors. *Statistica Sinica*, 4(2), 639–650.
- [43] Tokdar S. T. and Martin R. (2011). Bayesian test of normality versus a Dirichlet process mixture alternative, arXiv:1108.2883.
- [44] Wolpert R. L., and Ickstadt K., (1998). Simulation of Lévy random fields. In *Practical Nonparametric and Semiparametric Bayesian Statistics*, volume 133 of *Lect. Notes Stat.*, 227–242. Springer, New York.
- [45] Zarepour M., and Al-Labadi L. (2012). On a rapid simulation of the Dirichlet process. *Statistics & Probability Letters*, 82(5), 916–924.

- [46] Zhou M. and Shao Y. (2014). A powerful test for multivariate normality, *Applied Statistics*, 41(2), 351–363.