

Infinite families of 2-designs from a class of cyclic codes with two non-zeros

Xiaoni Du*, Rong Wang* Cuiling Fan*[†]

*College of Mathematics and Statistics, Northwest Normal University, Lanzhou, Gansu 730070, P.R. China

Email: {ymldxn, rongw113}@126.com

[†] School of Mathematics, Southwest Jiaotong University, Chengdu, China
Email: cuilingfan@163.com

Abstract

Combinatorial t -designs have wide applications in coding theory, cryptography, communications and statistics. It is well known that the supports of all codewords with a fixed weight in a code may give a t -design. In this paper, we first determine the weight distribution of a class of linear codes derived from the dual of extended cyclic code with two non-zeros. We then obtain infinite families of 2-designs and explicitly compute their parameters from the supports of all the codewords with a fixed weight in the codes. By simple counting argument, we obtain exponentially many 2-designs.

Key words: Affine-invariant code, cyclic code, exponential sum, linear code, weight distribution, 2-design

1 Introduction

Let k, t and v are positive integers with $1 \leq t \leq k \leq v$. Let \mathcal{P} be a set of v elements and \mathcal{B} be a set of k -subsets of \mathcal{P} . If every t -subset of \mathcal{P} is contained in exactly λ elements of \mathcal{B} , then we call the pair $\mathbb{D} = (\mathcal{P}, \mathcal{B})$ a t -(v, k, λ) design, or simply t -design. The elements of \mathcal{P} are called *points*, and those of \mathcal{B} are referred to as *blocks*. We often define the number of blocks by b and a t -design is simple when there is no repeated blocks in \mathcal{B} . A t -design is called *symmetric* if $v = b$ and trivial if $k = t$ or $k = v$. Hereafter we study only simple t -design with $t < k < v$. We call t -(v, k, λ) design a *Steiner system* when $t \geq 2$ and $\lambda = 1$. It is well known that a t -(v, k, λ) design is also an i -(v, k, λ_i) design for any $0 \leq i < t$, where

$$\lambda_i = \frac{\lambda \binom{v-i}{t-i}}{\binom{k-i}{t-i}}. \quad (1)$$

Combinatorial t -designs have very important applications in coding theory, cryptography, communications and statistics. There has been an interplay between codes and t -designs for both directions. On one hand, a linear code over any finite field can be derived from the incidence matrix of a t -design and much progress has been made and documented in [1, 6, 19, 20]. On the other hand, linear and nonlinear codes might both hold t -designs. Till now, 4-designs and 5-designs with fixed parameters were only derived from binary and ternary Golay codes. The largest t for which the infinite families of t -designs could be obtained from linear codes is $t = 3$. In 2017, Ding and Li [8] obtained infinite families of 2-designs from p -ary Hamming codes, ternary projective cyclic codes, binary codes with two zeros and their duals as well as the infinite families of 3-designs from the extended codes of these codes and RM codes. Afterwards, infinite families of 2-designs and 3-designs were constructed from a class of binary linear codes with five weights [9]. For other constructions of t -designs, for example, we refer to [2, 5, 17, 18].

The rest of this paper is organized as follows. In Section 2, we introduce some notation, together with some preliminary results on affine-invariant codes and 2-designs, which will be used in subsequent sections. In Section 3, we present the weight distributions of a class of linear codes derived from the dual of the extended cyclic codes with two non-zeros. By proving that the derived codes are affine-invariant, we then obtain infinite families of 2-designs and explicitly determine their parameters. The proof of the main results are given in Section 4. Section 5 concludes the paper.

2 Preliminaries

In this section, we present some standard notation and basic facts on affine-invariant codes and 2-designs.

2.1 Some notation

For convenience, we fix the following notation unless otherwise stated in this paper.

- p is an odd prime and $p^* = (-1)^{\frac{p-1}{2}}p$.
- $m \geq 2$, $1 \leq l \leq m-1$ are positive integers, $q = p^m$, $n = p^m - 1$ and $\gcd(m, l) = d > 1$.
- $\zeta_p = e^{2\pi\nu/p}$ is a primitive p -th root of unity, where $\nu = \sqrt{-1}$.
- \mathbb{F}_q denotes the finite field with q elements and $\mathbb{F}_q^* = \mathbb{F}_q \setminus \{0\}$.
- η and η' are the quadratic characters of \mathbb{F}_q^* and \mathbb{F}_p^* , respectively. We extend these quadratic characters by setting $\eta(0) = 0$ and $\eta'(0) = 0$.
- Tr denotes the trace function from \mathbb{F}_q onto \mathbb{F}_p .

2.2 The classical construction of t -designs from affine-invariant codes

Let p be an odd prime, m, l be positive integers and $(m, l) = d > 1$. Let \mathbb{F}_q denote the finite field with $q = p^m$ elements and $\mathbb{F}_q^* = \mathbb{F}_q \setminus \{0\}$. An $[n, k, d]$ linear code \mathcal{C} over \mathbb{F}_p is a k -dimensional subspace of \mathbb{F}_p^n with minimum Hamming distance d , and is *cyclic* provided that any cyclic shift of a codeword is another codeword of \mathcal{C} . Any cyclic code \mathcal{C} can be expressed as $\mathcal{C} = \langle g(x) \rangle$, where $g(x)$ is monic and has the least degree. The polynomial $g(x)$ is called the *generator polynomial* and $h(x) = (x^n - 1)/g(x)$ is referred to as the *parity check polynomial* of \mathcal{C} . If the generator polynomial $g(x)$ (resp. the parity check polynomial $h(x)$) can be factored into a product of s irreducible polynomials over \mathbb{F}_q , then \mathcal{C} is called a cyclic code with s zeros (resp. s non-zeros). The code with the generator polynomial $x^k h(x^{-1})/h(0)$ is called the *dual* of \mathcal{C} and denoted by \mathcal{C}^\perp , which is an $[n, n - k]$ code. The *extended code* of \mathcal{C} is defined by

$$\overline{\mathcal{C}} = \{(c_0, c_1, \dots, c_n) \in \mathbb{F}_p^{n+1} : (c_0, c_1, \dots, c_{n-1}) \in \mathcal{C} \text{ with } \sum_{i=0}^n c_i = 0\}.$$

For a codeword $\mathbf{c} \in \mathcal{C}$, the *support* of \mathbf{c} is defined by

$$\text{Suppt}(\mathbf{c}) = \{0 \leq i \leq n - 1 : c_i \neq 0\}.$$

Let A_i be the number of codewords with Hamming weight i in a code \mathcal{C} . The *weight enumerator* of \mathcal{C} is defined by

$$1 + A_1 z + A_2 z^2 + \dots + A_n z^n,$$

and the sequence $(1, A_1, \dots, A_n)$ is called the *weight distribution* of \mathcal{C} . If $|\{1 \leq i \leq n : A_i \neq 0\}| = w$, then we say \mathcal{C} a w -weight code. Indexed the coordinates of a codeword in \mathcal{C} by $(0, 1, 2, \dots, n - 1)$, then for each i with $A_i \neq 0$, we denote \mathcal{B}_i by the set of the supports of all codewords with weight i . Let $\mathcal{P} = \{0, 1, \dots, n - 1\}$. The pair $(\mathcal{P}, \mathcal{B}_i)$ could give a t -(n, i, λ) design for some positive λ [8] and the parameter λ can be determined explicitly by Eq.(1). Such codes were used to construct 2-designs and 3-designs in [1, 19, 20].

We conclude this section by introducing some known results on affine-invariant codes related to 2-designs.

The set of coordinate permutations that map a code \mathcal{C} to itself forms a group, which is called the *permutation automorphism group* of \mathcal{C} and denoted by $\text{PAut}(\mathcal{C})$. We define the *affine group* $\text{GA}_1(\mathbb{F}_q)$ by the set of all permutations

$$\sigma_{a,b} : x \mapsto ax + b$$

of \mathbb{F}_q , where $a \in \mathbb{F}_q^*$ and $b \in \mathbb{F}_q$. An affine-invariant code is an extended cyclic code $\overline{\mathcal{C}}$ over \mathbb{F}_p satisfying $\text{GA}_1(\mathbb{F}_q) \subseteq \text{PAut}(\overline{\mathcal{C}})$ [13].

The p -adic expansion of each $s \in \mathcal{P}$ is given by

$$s = \sum_{i=0}^{m-1} s_i p^i, \quad 0 \leq s_i \leq p - 1, \quad 0 \leq i \leq m - 1.$$

For any $r = \sum_{i=0}^{m-1} r_i p^i \in \mathcal{P}$, we call that $r \preceq s$ if $r_i \leq s_i$ for all $0 \leq i \leq m-1$.

For any integer $0 \leq j \leq n-1$, the p -cyclotomic coset of j modulo n is defined by

$$C_j = \{jp^i \pmod{n} : 0 \leq i \leq \ell_j - 1\}$$

where ℓ_j is the smallest positive integer such that $j \equiv jp^{\ell_j} \pmod{n}$. The smallest integer in C_j is called the *coset leader* of C_j . Let $g(x) = \prod_j \prod_{i \in C_j} (x - \alpha^i)$, where j runs through some coset leaders of the p -cyclotomic cosets C_j modulo n . The union of these p -cyclotomic cosets $T = \bigcup_j C_j$ is referred to as the *defining set* of \mathcal{C} .

For certain applications, it is important to know whether a given extended primitive cyclic code $\overline{\mathcal{C}}$ is affine-invariant or not. The following Lemma 1 presented by Kasami *et al.* [15] answered the questions by examining the defining set of the code. Lemma 2 proposed by Ding [7] in 2019 point out that one can get the new affine-invariant codes from the known ones.

Lemma 1 *Let $\overline{\mathcal{C}}$ be an extended cyclic code of length p^m over \mathbb{F}_p with defining set \overline{T} . The code $\overline{\mathcal{C}}$ is affine-invariant if and only if whenever $s \in \overline{T}$ then $r \in \overline{T}$ for all $r \in \mathcal{P}$ with $r \preceq s$.*

Lemma 2 *The dual of an affine-invariant code $\overline{\mathcal{C}}$ over \mathbb{F}_p of length $n+1$ is also affine-invariant.*

Lemma 2 and the following Theorem 1 given by Ding in [7] are very powerful tools in constructing 2-designs from linear codes. We will employ them to present our main results in this paper.

Theorem 1 *For each i with $A_i \neq 0$ in an affine-invariant code $\overline{\mathcal{C}}$, the supports of the codewords of weight i form a 2-design.*

The following theorem presented by Ding in [7] reveals the relation of the codewords with the same support in a linear code \mathcal{C} , which will be used together with Eq.(1) to calculate the parameters of 2-designs later.

Theorem 2 *Let \mathcal{C} be a linear code over \mathbb{F}_p with minimum weight d . Let w be the largest integer with $w \leq n$ satisfying*

$$w - \lfloor \frac{w+p-2}{p-1} \rfloor < d.$$

Let \mathbf{c}_1 and \mathbf{c}_2 be two codewords of weight i with $d \leq i \leq w$ and $\text{Suppt}(\mathbf{c}_1) = \text{Suppt}(\mathbf{c}_2)$. Then $\mathbf{c}_1 = a\mathbf{c}_2$ for some $a \in \mathbb{F}_p^$.*

Table 1: The weight distribution of $\overline{\mathcal{C}}^{\perp}$ when both m and m/d are odd

Weight	Multiplicity
0	1
$p^{m-1}(p-1)$	$p(p^{m-1}+1)(p^m-1)$
$p^{m-1}(p-1) + p^{\frac{m-1}{2}}$	$\frac{1}{2}p^m(p-1)(p^m-1)$
$p^{m-1}(p-1) - p^{\frac{m-1}{2}}$	$\frac{1}{2}p^m(p-1)(p^m-1)$
p^m	$p-1$

3 The weight distribution of the codes and their 2-designs

In this section, we present the main results, including the weight distribution and the underlying 2-designs. The proofs of the main results will be provided in the following section. We begin this section by the definition of the code.

Define

$$\overline{\mathcal{C}}^{\perp} := \{(Tr(ax^{p^l+1} + bx)_{x \in \mathbb{F}_q} + h) : a, b \in \mathbb{F}_q, h \in \mathbb{F}_p\}, \quad (2)$$

where \mathcal{C} is the cyclic code of length n and parity check polynomial $h_1(x)h_2(x)$ where $h_1(x), h_2(x)$ are the minimal polynomials of α, α^{p^l+1} over \mathbb{F}_p , respectively, and α is a primitive element of \mathbb{F}_q . In particular, if $l = \frac{m}{2}$ we have

$$\overline{\mathcal{C}}^{\perp} := \{(Tr(ax^{p^{\frac{m}{2}}+1} + bx)_{x \in \mathbb{F}_q} + h) : a \in \mathbb{F}_{p^{\frac{m}{2}}}, b \in \mathbb{F}_q, h \in \mathbb{F}_p\}.$$

Below, we consider only the case $\gcd(m, l) > 1$ since the case for $\gcd(m, l) = 1$ was discussed in [10].

The following Theorems 3-5 are the main results of this paper.

Theorem 3 *Let $m \geq 2$ and $(m, l) = d > 1$. The weight distribution of the code $\overline{\mathcal{C}}^{\perp}$ over \mathbb{F}_p with length $n+1$ are given in Table 1 when both m and $\frac{m}{d}$ are odd, in Table 2 when m is even and $\frac{m}{d}$ is odd, in Table 3 when $\frac{m}{d} \geq 4$ is even and in Table 4 when $m = 2d$, respectively. And the dimension is $\dim(\overline{\mathcal{C}}^{\perp}) = \frac{3m}{2} + 1$ when $m = 2d$ and $\dim(\overline{\mathcal{C}}^{\perp}) = 2m + 1$ otherwise.*

One can see that the code is four-weight if both m and $\frac{m}{d}$ are odd or $m = 2d$ and six-weight if m is even and $\frac{m}{d}$ is odd or $\frac{m}{d} \geq 4$ is even.

Theorem 4 *Let $m \geq 2, l$ be positive integers with $\gcd(m, l) > 1$. Then the supports of the codewords of weight i with $A_i \neq 0$ in $\overline{\mathcal{C}}^{\perp}$ form a 2-design.*

Table 2: The weight distribution of $\overline{\mathcal{C}}^{\perp\perp}$ when m is even and m/d is odd

Weight	Multiplicity
0	1
$p^{m-1}(p-1)$	$p(p^m-1)$
$p^{m-1}(p-1) + p^{\frac{m}{2}-1}$	$\frac{1}{2}p^m(p-1)(p^m-1)$
$p^{m-1}(p-1) - p^{\frac{m}{2}-1}$	$\frac{1}{2}p^m(p-1)(p^m-1)$
$p^{m-1}(p-1) + p^{\frac{m}{2}-1}(p-1)$	$\frac{1}{2}p^m(p^m-1)$
$p^{m-1}(p-1) - p^{\frac{m}{2}-1}(p-1)$	$\frac{1}{2}p^m(p^m-1)$
p^m	$p-1$

Table 3: The weight distribution of $\overline{\mathcal{C}}^{\perp\perp}$ when $m/d \geq 4$ is even

Weight	Multiplicity
0	1
$p^{m-1}(p-1)$	$p(p^{m-d} - p^{m-2d} + 1)(p^m-1)$
$p^{m-1}(p-1) - (-1)^{\frac{m}{2d}}p^{\frac{m}{2}-1}(p-1)$	$p^{m+d}(p^m-1)/(p^d+1)$
$p^{m-1}(p-1) + (-1)^{\frac{m}{2d}}p^{\frac{m}{2}-1}$	$p^{m+d}(p-1)(p^m-1)/(p^d+1)$
$p^{m-1}(p-1) + (-1)^{\frac{m}{2d}}p^{\frac{m}{2}+d-1}(p-1)$	$p^{m-2d}(p^m-1)/(p^d+1)$
$p^{m-1}(p-1) - (-1)^{\frac{m}{2d}}p^{\frac{m}{2}+d-1}$	$p^{m-2d}(p-1)(p^m-1)/(p^d+1)$
p^m	$p-1$

Table 4: The weight distribution of $\overline{\mathcal{C}}^{\perp\perp}$ when $m/d = 2$

Weight	Multiplicity
0	1
$p^{m-1}(p-1)$	$p(p^m-1)$
$p^{m-1}(p-1) - p^{\frac{m}{2}-1}$	$p^m(p^{\frac{m}{2}}-1)(p-1)$
$p^{m-1}(p-1) + p^{\frac{m}{2}-1}(p-1)$	$p^m(p^{\frac{m}{2}}-1)$
p^m	$p-1$

Theorem 5 Let $m \geq 2, l$ be integers, $\mathcal{P} = \{0, 1, \dots, p^m - 1\}$ and \mathcal{B} be the set of the supports of the codewords of $\overline{\mathcal{C}}^{\perp}$ with weight i , where $A_i \neq 0$. Then $\overline{\mathcal{C}}^{\perp}$ gives $2-(p^m, i, \lambda)$ designs for the following pairs:

(1) if both m and m/d are odd,

- $(i, \lambda) = (p^m - p^{m-1}, (p^m - p^{m-1} - 1)(p^{m-1} + 1)).$
- $(i, \lambda) = (p^m - p^{m-1} + p^{\frac{m-1}{2}}, \frac{1}{2}p^{\frac{m-1}{2}}(p^{\frac{m+1}{2}} - p^{\frac{m-1}{2}} + 1)(p^m - p^{m-1} + p^{\frac{m-1}{2}} - 1)).$
- $(i, \lambda) = (p^m - p^{m-1} - p^{\frac{m-1}{2}}, \frac{1}{2}p^{\frac{m-1}{2}}(p^{\frac{m+1}{2}} - p^{\frac{m-1}{2}} - 1)(p^m - p^{m-1} - p^{\frac{m-1}{2}} - 1)).$

(2) if m is even and m/d is odd,

- $(i, \lambda) = (p^m - p^{m-1}, p^m - p^{m-1} - 1).$
- $(i, \lambda) = (p^{m-1}(p-1) + p^{\frac{m}{2}-1}, \frac{1}{2}p^{\frac{m}{2}-1}(p^{\frac{m}{2}+1} - p^{\frac{m}{2}} + 1)(p^m - p^{m-1} + p^{\frac{m}{2}-1} - 1)).$
- $(i, \lambda) = (p^{m-1}(p-1) - p^{\frac{m}{2}-1}, \frac{1}{2}p^{\frac{m}{2}-1}(p^{\frac{m}{2}+1} - p^{\frac{m}{2}} - 1)(p^m - p^{m-1} - p^{\frac{m}{2}-1} - 1)).$
- $(i, \lambda) = (p^{m-1}(p-1) + p^{\frac{m}{2}-1}(p-1), \frac{1}{2}p^{\frac{m}{2}-1}(p^{\frac{m}{2}} + 1)(p^m - p^{m-1} + p^{\frac{m}{2}} - p^{\frac{m}{2}-1} - 1)).$
- $(i, \lambda) = (p^{m-1}(p-1) - p^{\frac{m}{2}-1}(p-1), \frac{1}{2}p^{\frac{m}{2}-1}(p^{\frac{m}{2}} - 1)(p^m - p^{m-1} - p^{\frac{m}{2}} + p^{\frac{m}{2}-1} - 1)).$

(3) if $m/d \geq 4$ is even,

- $(i, \lambda) = (p^m - p^{m-1}, (p^m - p^{m-1} - 1)(p^{m-d} - p^{m-2d} + 1)).$
- $(i, \lambda) = (p^{m-1}(p-1) + (-1)^{\frac{m}{2d}}p^{\frac{m}{2}-1}, p^{\frac{m}{2}+d-1}(p^{\frac{m}{2}+1} - p^{\frac{m}{2}} + (-1)^{\frac{m}{2d}})(p^m - p^{m-1} + (-1)^{\frac{m}{2d}}p^{\frac{m}{2}-1} - 1)/(p^d + 1)).$
- $(i, \lambda) = (p^{m-1}(p-1) - (-1)^{\frac{m}{2d}}p^{\frac{m}{2}-1}(p-1), p^{\frac{m}{2}+d-1}(p^{\frac{m}{2}} - (-1)^{\frac{m}{2d}})(p^m - p^{m-1} - (-1)^{\frac{m}{2d}}p^{\frac{m}{2}} + (-1)^{\frac{m}{2d}}p^{\frac{m}{2}-1} - 1)/(p^d + 1)).$
- $(i, \lambda) = (p^{m-1}(p-1) + (-1)^{\frac{m}{2d}}p^{\frac{m}{2}+d-1}(p-1), p^{\frac{m}{2}-d-1}(p^{\frac{m}{2}-d} + (-1)^{\frac{m}{2d}})(p^m - p^{m-1} + (-1)^{\frac{m}{2d}}p^{\frac{m}{2}+d} - (-1)^{\frac{m}{2d}}p^{\frac{m}{2}+d-1} - 1)/(p^d + 1)).$
- $(i, \lambda) = (p^{m-1}(p-1) - (-1)^{\frac{m}{2d}}p^{\frac{m}{2}+d-1}, p^{\frac{m}{2}-d-1}(p^{\frac{m}{2}-d+1} - p^{\frac{m}{2}-d} - (-1)^{\frac{m}{2d}})(p^m - p^{m-1} - (-1)^{\frac{m}{2d}}p^{\frac{m}{2}+d-1} - 1)/(p^d + 1)).$

(4) if $m/d = 2$,

- $(i, \lambda) = (p^m - p^{m-1}, p^m - p^{m-1} - 1).$
- $(i, \lambda) = (p^m - p^{m-1} - p^{\frac{m}{2}-1}, p^{\frac{m}{2}-1}(p^{\frac{m}{2}+1} - p^{\frac{m}{2}} - 1)(p^m - p^{m-1} - p^{\frac{m}{2}-1} - 1)/(p^{\frac{m}{2}} + 1)).$
- $(i, \lambda) = (p^m - p^{m-1} + p^{\frac{m}{2}} - p^{\frac{m}{2}-1}, p^{\frac{m}{2}-1}(p^m - p^{m-1} + p^{\frac{m}{2}} - p^{\frac{m}{2}-1} - 1)).$

The following examples from Magma program confirm the main results in Theorems 3 and 5.

Example 1 If $(p, l, m) = (3, 2, 4)$, then the code $\overline{\mathcal{C}}^{\perp}$ has parameters $[81, 7, 51]$ and weight enumerator $1 + 1296z^{51} + 240z^{54} + 648z^{60} + 2z^{81}$, It gives 2- $(81, i, \lambda)$ designs with the following pairs (i, λ) :

$$(51, 255), (54, 53), (60, 177),$$

which confirmed the results in Theorems 3 and 5.

Example 2 If $(p, l, m) = (3, 3, 6)$, then the code $\overline{\mathcal{C}}^{\perp}$ has parameters $[729, 10, 477]$ and weight enumerator $1 + 37908z^{477} + 2184z^{486} + 18954z^{504} + 2z^{729}$, which confirmed the results in Theorem 3.

Example 3 If $(p, l, m) = (3, 2, 6)$, then the code $\overline{\mathcal{C}}^{\perp}$ has parameters $[729, 10, 468]$ and weight enumerator $1 + 265356z^{468} + 530712z^{477} + 2184z^{486} + 530712z^{495} + 265356z^{504} + 2z^{729}$, which confirmed the results in Theorem 3.

4 Proofs of the main results

We begin this section by stating some notation and basic facts on exponential sums.

4.1 Exponential sums

An additive character of \mathbb{F}_q is a non-zero function χ from \mathbb{F}_q to the set of complex numbers of absolute value 1 such that $\chi(x + y) = \chi(x)\chi(y)$ for any pair $(x, y) \in \mathbb{F}_q^2$. For each $u \in \mathbb{F}_q$, the function

$$\chi_u(v) = \zeta^{\text{Tr}(uv)}, \quad v \in \mathbb{F}_q$$

denotes an additive character of \mathbb{F}_q . χ_0 is the trivial additive character of \mathbb{F}_q since $\chi_0(v) = 1$ for all $v \in \mathbb{F}_q$. We call χ_1 the canonical additive character of \mathbb{F}_q and we have $\chi_u(x) = \chi_1(ux)$ for all $u \in \mathbb{F}_q$ [16].

The Gauss sum $G(\eta', \chi'_1)$ over \mathbb{F}_p is defined by $G(\eta', \chi'_1) = \sum_{c \in \mathbb{F}_p^*} \eta'(v) \chi'_1(v) = \sum_{v \in \mathbb{F}_p} \eta'(v) \chi'_1(v)$, where χ'_1 is the canonical additive characters of \mathbb{F}_p .

To determine the parameters of the code $\overline{\mathcal{C}}^{\perp}$ defined in Eq.(2), we introduce the following function.

$$S(a, b) = \sum_{x \in \mathbb{F}_q} \chi(ax^{p^l+1} + bx), \quad a, b \in \mathbb{F}_q, \quad (3)$$

if $m \neq 2l$, and otherwise

$$S(a, b) = \sum_{x \in \mathbb{F}_q} \chi(ax^{p^{\frac{m}{2}}+1} + bx), \quad a \in \mathbb{F}_{p^{\frac{m}{2}}}, \quad b \in \mathbb{F}_q.$$

The following Lemmas 3-6 are essential to determine the values of Eq.(3).

Lemma 3 [16] *With the notation above, we have*

$$G(\eta', \chi'_1) = \sqrt{(-1)^{\left(\frac{p-1}{2}\right)^2}} \sqrt{p} = \sqrt{p^*}.$$

Lemma 4 [6] *For each $y \in \mathbb{F}_p^*$, $\eta(y) = 1$ if $m \geq 2$ is even, and $\eta(y) = \eta'(y)$ if $m \geq 1$ is odd.*

Lemma 5 [3] *Let $f(x) = a^{p^l} x^{p^{2l}} + ax \in \mathbb{F}_q[x]$, $\gcd(m, l) = d$ and $b \in \mathbb{F}_q$. There are three cases.*

(1) *If m/d is odd, then $f(x)$ is a permutation polynomial over \mathbb{F}_q and*

$$S(a, b) = \sqrt{p^*}^m \eta(a) \chi_1(-ax_{a,b}^{p^l+1}).$$

(2) *If m/d is even and $a^{\frac{q-1}{p^d+1}} \neq (-1)^{\frac{m}{2d}}$, then $f(x)$ is a permutation polynomial over \mathbb{F}_q and*

$$S(a, b) = (-1)^{\frac{m}{2d}} p^{m/2} \chi_1(-ax_{a,b}^{p^l+1}).$$

(3) *If m/d is even and $a^{\frac{q-1}{p^d+1}} = (-1)^{\frac{m}{2d}}$, then $f(x)$ is not a permutation polynomial over \mathbb{F}_q . We have $S(a, b) = 0$ when the equation $f(x) = -b^{p^l}$ is unsolvable, and*

$$S(a, b) = -(-1)^{\frac{m}{2d}} p^{m/2+d} \chi_1(-ax_{a,b}^{p^l+1})$$

otherwise. Particularly,

$$S(a, 0) = \begin{cases} \sqrt{p^*}^m \eta(a) & \text{if } m/d \text{ is odd,} \\ (-1)^{\frac{m}{2d}} p^{\frac{m}{2}} & \text{if } m/d \text{ is even and } a^{\frac{q-1}{p^d+1}} \neq (-1)^{\frac{m}{2d}}, \\ (-1)^{\frac{m}{2d}+1} p^{\frac{m}{2}+d} & \text{if } m/d \text{ is even and } a^{\frac{q-1}{p^d+1}} = (-1)^{\frac{m}{2d}}. \end{cases}$$

Notice that $x_{a,b}^{p^l+1}$ is a solution to the equation $f(x) = -b^{p^l}$. Moreover, $x_{a,b}^{p^l+1}$ is the unique solution when $f(x)$ is a permutation polynomial over \mathbb{F}_q .

Lemma 6 [4] *For m even and $\gcd(m, l) = d$, the equation $a^{p^l} x^{p^{2l}} + ax = 0$ is solvable for $x \in \mathbb{F}_q^*$ if and only if*

$$a^{\frac{q-1}{p^d+1}} = (-1)^{\frac{m}{2d}}.$$

In such cases there are $p^{2d} - 1$ non-zero solutions.

4.2 Proofs of the main results

Now we are ready to give the proofs of our main results. First we present the proof of the weight distribution of $\overline{\mathcal{C}}^{\perp}$ which is given in Eq.(2).

Proof of Theorem 3. For each non-zero codeword $\mathbf{c}(a, b, h) = (c_0, \dots, c_n)$ in $\overline{\mathcal{C}}^{\perp}$, the Hamming weight of $\mathbf{c}(a, b, h)$ is

$$w_H(\mathbf{c}(a, b, h)) = p^m - T(a, b, h), \quad (4)$$

where

$$T(a, b, h) = |\{x : \text{Tr}(ax^{p^l+1} + bx) + h = 0, x, a, b \in \mathbb{F}_q, h \in \mathbb{F}_p\}|$$

if $l \neq \frac{m}{2}$, and otherwise

$$T(a, b, h) = |\{x : \text{Tr}(ax^{p^{\frac{m}{2}+1}} + bx) + h = 0, a \in \mathbb{F}_{p^{\frac{m}{2}}}, x, b \in \mathbb{F}_q, h \in \mathbb{F}_p\}|.$$

Then

$$\begin{aligned} T(a, b, h) &= \frac{1}{p} \sum_{y \in \mathbb{F}_p} \sum_{x \in \mathbb{F}_q} \zeta_p^{y \text{Tr}(ax^{p^l+1} + bx) + hy} \\ &= p^{m-1} + \frac{1}{p} \sum_{y \in \mathbb{F}_p^*} \zeta_p^{hy} \sum_{x \in \mathbb{F}_q} \zeta_p^{y \text{Tr}(ax^{p^l+1} + bx)}. \end{aligned}$$

If $a = b = h = 0$, then $\mathbf{c}(a, b, h)$ is the zero codeword.

If $a = b = 0, h \neq 0$, then $T(a, b, h) = p^{m-1} + p^{m-1} \sum_{y \in \mathbb{F}_p^*} \zeta_p^{hy} = 0$.

If $a = 0, b \neq 0$, then $T(a, b, h) = p^{m-1} + \frac{1}{p} \sum_{y \in \mathbb{F}_p^*} \zeta_p^{hy} \sum_{x \in \mathbb{F}_q} \zeta_p^{y \text{Tr}(bx)} = p^{m-1}$.

If $a \neq 0$, then from the definition of $S(a, b)$, we have

$$T(a, b, h) = p^{m-1} + \frac{1}{p} \sum_{y \in \mathbb{F}_p^*} \zeta_p^{hy} S(ay, by).$$

According to Lemma 5, the calculation of $T(a, b, h)$ can be divide into the following cases.

(1) When both m and m/d are odd, by Lemma 5, we have

$$T(a, b, h) = \begin{cases} p^{m-1} & \text{if } h = \text{Tr}(ax_{ay,by}^{p^l+1}), \\ p^{m-1} + p^{\frac{m-1}{2}}(-1)^{\frac{(p-1)(m+1)}{4}}\eta(a)\eta'(h - \text{Tr}(ax_{a,b}^{p^l+1})) & \text{if } h \neq \text{Tr}(ax_{ay,by}^{p^l+1}). \end{cases}$$

Moreover, we find that $T(a, b, h) = p^{m-1}$ appears $p^m(p^m - 1)$ times, and both

$$T(a, b, h) = p^{m-1} + p^{\frac{m-1}{2}} \text{ and } T(a, b, h) = p^{m-1} - p^{\frac{m-1}{2}}$$

appear

$$\frac{p^m(p-1)(p^m-1)}{2}$$

times, respectively, from the properties of the quadratic characters η of \mathbb{F}_q^* .

(2) When m is even and m/d is odd, by Lemma 5, we get

$$T(a, b, h) = \begin{cases} p^{m-1} - p^{\frac{m}{2}-1}(-1)^{\frac{(p-1)m}{4}}\eta(a) & \text{if } h \neq \text{Tr}(ax_{ay,by}^{p^l+1}), \\ p^{m-1} + p^{\frac{m}{2}-1}(-1)^{\frac{(p-1)m}{4}}\eta(a)(p-1) & \text{if } h = \text{Tr}(ax_{ay,by}^{p^l+1}). \end{cases}$$

More precisely, by the properties of the quadratic characters η , we have that both

$$T(a, b, h) = p^{m-1} + p^{\frac{m}{2}-1} \text{ and } T(a, b, h) = p^{m-1} - p^{\frac{m}{2}-1}$$

appear

$$\frac{p^m(p-1)(p^m-1)}{2}$$

times, and both

$$T(a, b, h) = p^{m-1} + p^{\frac{m}{2}-1}(p-1) \text{ and } T(a, b, h) = p^{m-1} - p^{\frac{m}{2}-1}(p-1)$$

appear

$$\frac{p^m(p^m-1)}{2}$$

times, respectively.

(3) When m/d is even and $a^{\frac{q-1}{p^d+1}} = (-1)^{\frac{m}{2d}}$, from Lemma 5, we get

$$T(a, b, h) = \begin{cases} p^{m-1} - (-1)^{\frac{m}{2d}} p^{\frac{m}{2}+d-1}(p-1) & \text{if } f(x) = -b^{p^l} \text{ is solvable and} \\ & h = \text{Tr}(ax_{ay,by}^{p^l+1}), \\ p^{m-1} + (-1)^{\frac{m}{2d}} p^{\frac{m}{2}+d-1} & \text{if } f(x) = -b^{p^l} \text{ is solvable, } b \neq 0 \text{ and} \\ & h \neq \text{Tr}(ax_{ay,by}^{p^l+1}) \text{ or } b = 0 \text{ and } h \neq 0, \\ p^{m-1} & \text{if } f(x) = -b^{p^l} \text{ is no solvable.} \end{cases}$$

It follows from Lemma 6 that there are $\frac{q-1}{p^d+1}$ elements $a \in \mathbb{F}_q^*$ satisfying $a^{\frac{q-1}{p^d+1}} = (-1)^{\frac{m}{2d}}$, and p^{m-2d} elements $b \in \mathbb{F}_q$ such that $f(x) = -b^{p^l}$ is solvable. Therefore,

$$T(a, b, h) = p^{m-1} + (-1)^{\frac{m}{2d}+1} p^{\frac{m}{2}+d-1}(p-1)$$

appears $\frac{p^{m-2d}(p^m-1)}{p^d+1}$ times,

$$T(a, b, h) = p^{m-1} + (-1)^{\frac{m}{2d}} p^{\frac{m}{2}+d-1}$$

appears $\frac{p^{m-2d}(p^m-1)(p-1)}{p^d+1}$ times, and

$$T(a, b, h) = p^{m-1}$$

appears $(p^m-1)p^{m-2d+1}(p^d-1)$ times.

(4) When $m/d \geq 4$ is even and $a^{\frac{q-1}{p^d+1}} \neq (-1)^{\frac{m}{2d}}$, by Lemma 5, we have

$$T(a, b, h) = \begin{cases} p^{m-1} + (-1)^{\frac{m}{2d}} p^{\frac{m}{2}-1}(p-1) & \text{if } h = \text{Tr}(ax_{ay,by}^{p^l+1}), \\ p^{m-1} - (-1)^{\frac{m}{2d}} p^{\frac{m}{2}-1} & \text{if } h \neq \text{Tr}(ax_{ay,by}^{p^l+1}). \end{cases}$$

Particularly, there exist $p^m-1 - \frac{p^m-1}{p^d+1} = \frac{p^d(p^m-1)}{p^d+1}$ elements $a \in \mathbb{F}_q^*$ such that

$$a^{\frac{q-1}{p^d+1}} \neq (-1)^{\frac{m}{2d}}.$$

Then

$$T(a, b, h) = p^{m-1} + (-1)^{\frac{m}{2d}} p^{\frac{m}{2}-1}(p-1)$$

appears $\frac{p^{m+d}(p^m-1)}{p^{d+1}}$ times, and

$$T(a, b, h) = p^{m-1} - (-1)^{\frac{m}{2d}} p^{\frac{m}{2}-1}$$

appears $\frac{p^{m+d}(p-1)(p^m-1)}{p^{d+1}}$ times.

(5) When $m = 2d$, it is clear that $a^{p^{\frac{m}{2}-1}} \neq -1$ for all $a \in \mathbb{F}_{p^{\frac{m}{2}}}^*$. Similarly, by Lemma 5, we have

$$\begin{aligned} T(a, b, h) &= |\{x : \text{Tr}(ax^{p^{\frac{m}{2}+1}} + bx) + h = 0, a \in \mathbb{F}_{p^{\frac{m}{2}}}, x, b \in \mathbb{F}_q, h \in \mathbb{F}_p\}| \\ &= \begin{cases} p^{m-1} - p^{\frac{m}{2}-1}(p-1) & \text{if } h = \text{Tr}(ax_{ay,by}^{p^l+1}), \\ p^{m-1} + p^{\frac{m}{2}-1} & \text{if } h \neq \text{Tr}(ax_{ay,by}^{p^l+1}). \end{cases} \end{aligned}$$

More precisely, we get

$$T(a, b, h) = p^{m-1} - p^{\frac{m}{2}-1}(p-1)$$

appears $(p^{\frac{m}{2}} - 1)p^m$ times, and

$$T(a, b, h) = p^{m-1} + p^{\frac{m}{2}-1}$$

appears $(p^{\frac{m}{2}} - 1)(p-1)p^m$ times.

From all the discussions above, we complete the proof of the theorem. \square

Now we prove that $\overline{\mathcal{C}^\perp}$ is affine invariant.

Lemma 7 *The extended code $\overline{\mathcal{C}^\perp}$ is affine-invariant.*

Proof. We will prove the conclusion with Lemma 1. The defining set T of the cyclic code \mathcal{C}^\perp is $T = C_1 \cup C_{p^l+1}$. Since $0 \notin T$, the defining set \overline{T} of $\overline{\mathcal{C}^\perp}$ is given by $\overline{T} = C_1 \cup C_{p^l+1} \cup \{0\}$. Let $s \in \overline{T}$ and $r \in \mathcal{P}$. Assume that $r \preceq s$. We need to prove that $r \in \overline{T}$ by Lemma 1.

If $r = 0$, then obviously $r \in \overline{T}$. Consider now the case $r > 0$. If $s \in C_1$, then the Hamming weight $wt(s) = 1$. As $r \preceq s$, $wt(r) = 1$. Consequently, $r \in C_1 \subset \overline{T}$. If $s \in C_{p^l+1}$, then the Hamming weight $wt(s) = 2$. As $r \preceq s$, either $wt(r) = 1$ or $r = s$. In both cases, $r \in \overline{T}$. The desired conclusion then follows from Lemma 1. Thus we complete the proof of the lemma. \square

Now we are ready to prove Theorems 4 and 5.

Proof of Theorem 4.

By Lemmas 7 and 2, we find that $\overline{\mathcal{C}^\perp}^\perp$ is affine-invariant. Thus the supports of the codewords of weight i with $A_i \neq 0$ in $\overline{\mathcal{C}^\perp}^\perp$ give 2-designs from Theorem 1. \square

Proof of Theorem 5.

By Theorem 2, one can prove that the number of supports of all codewords with weight $i \neq 0$ in the code $\overline{\mathcal{C}^\perp}^\perp$ is equal to $A_i/(p-1)$ for each i , where A_i is given in Tables 1-4. Then the desired conclusions follow from Theorem 4 and Eq.(1). Thus, we complete the proof. \square

5 Concluding remarks

In this paper, first we determined the weight distribution of a class of linear codes derived from the dual of extended cyclic codes with two non-zeros. Using the properties of affine-invariant codes, we then found that the code $\overline{\mathcal{C}}^{\perp}$ holds 2-designs. By computing certain exponential sums, we explicitly determined their parameters. The total number of the 2-designs we obtained is exponential in m .

References

- [1] Assmus Jr. E.F., Key J.D.: Designs and their codes. Cambridge University Press, Cambridge (1992).
- [2] Beth T., Jungnickel D., Lenz H.: Design theory. Cambridge University Press, Cambridge (1999).
- [3] Coulter R.S.: Further evaluations of weil sums. Acta Arith 86, pp.217-226 (1998).
- [4] Coulter R.S.: Explicit evaluations of some weil sums. Mathematics subject classification. 11T24, (1991).
- [5] Colbourn C.J., Maithon R.: Stener systems. In: Colbourn C.J., Dinitz J. (eds.) Handbook of Combinatorial Designs, pp.102-110. CRC Press, Boca Raton (2007).
- [6] Ding K., Ding C., Wang Q.: A class of two-weight and three-weight codes and their applications in secret sharing. IEEE Trans. Inf. Theory 61, pp.5835-5842 (2015).
- [7] Ding C.: Designs from linear codes. World Scientific, Singapore (2019).
- [8] Ding C., Li C.: Infinite families of 2-designs and 3-designs from linear codes. Discret Math. 340, pp.2415-2431 (2017).
- [9] Ding C.: Infinite families of 3-designs from a type of five weight codes. Codes Cryptogr. 86, pp.703-719 (2018).
- [10] Du X., Wang R., Tang C.: Infinite families of 2-designs from two classes of linear codes, preprint.
- [11] Fitzgerald R.W., Yucas J.L.: Sums of Gauss and weight of irreducible codes. Finite Field Appl. 11, pp.89-110 (2005).
- [12] Feng K., Luo J.: Weight distribution of some reducible cyclic codes. Finite Field Appl. 14, pp.390-409 (2008).
- [13] Huffman W.C., Pless V.: Fundamentals of error correcting codes. Cambridge University Press, Cambridge (2003).

- [14] Ireland K., Rosen M.: A classical introduction to modern number theory. In: Graduate Texts in Mathematics, vol. 84, 2nd edn. Springer, New York (1990).
- [15] Kasami T., Lin S., Peterson W.: Some results on cyclic codes which are invariant under the affine group and their applications. Inform. and Control 11, pp.475-496 (1968).
- [16] Lidl R., Niederreiter H.: Finite fields. Cambridge University Press, Cambridge (1997).
- [17] MacWilliams F.J., Sloane N.J.A.: The theory of error-correcting codes. North-Holland, Amsterdam (1977).
- [18] Reid C., Rosa A.: Steiner systems $S(2,4)$ -a survey. Electron. J. Comb. #DS18 (2010).
- [19] Tonchev V.D.: Codes and design. In: Pless V.S., Huffman W.C.(eds.) Handbook of Coding Theory, vol. II, pp.1229-1268. Elsevier, Amsterdam (1998).
- [20] Tonchev V.D.: Code. In: Collbourn C.J., Dinitz J.H.(eds.) Handbook of Combinatorial Designs, 2nd edn, 677-701. CRC Press, New York (2007).
- [21] Van Der Vlugt M., Hasse-Davenport curve.: Gauss sums and weight distribution of irreducible cyclic codes. J. Number Theory 55, pp.145-159 (1995).