

Gradient-Free Multi-Agent Nonconvex Nonsmooth Optimization

Davood Hajinezhad ^{*}

Michael M. Zavlanos [†]

April 1, 2024

Abstract

In this paper we consider the problem of minimizing the sum of nonconvex and possibly nonsmooth functions over a connected multi-agent network, where the agents have partial knowledge about the global cost function and can only access the zeroth-order information (i.e., the functional values) of their local cost functions. We propose and analyze a distributed primal-dual gradient-free algorithm for this challenging problem. We show that by appropriately choosing the parameters, the proposed algorithm converges to the set of first order stationary solutions with a provable global sublinear convergence rate. Numerical experiments demonstrate the effectiveness of our proposed method for optimizing nonconvex and nonsmooth problems over a network.

1 Introduction

Consider a network with N distributed agents that collectively solve the following optimization problem

$$\min_{x \in \mathbb{R}^M} f(x) := \sum_{i=1}^N f_i(x). \quad (1)$$

Here, $f_i : \mathbb{R}^M \rightarrow \mathbb{R}$ is possibly a nonconvex nonsmooth function, that is only available to agent i . Such distributed optimization problems arise in many applications such as machine learning [1, 2], resource allocation [3], robotic networks [4], and signal processing [5]. See [6] for more applications.

Many distributed optimization methods have been proposed to solve problem (1). Many of them rely on consensus between the agents and assume that the cost functions are convex. One of the first such methods is the distributed subgradient (DSG) algorithm [7]. Subsequently, a number of similar consensus-based algorithms were also proposed to solve distributed convex optimization problems in the form of (1); see, e.g., [8–11]. These methods only converge to a neighborhood of the solution set unless they use diminishing

^{*}Department of Mechanical Engineering and Materials Science, Duke University, USA, Email: dhajinezhad@gmail.com

[†]Department of Mechanical Engineering and Materials Science, Duke University, USA, Email: michael.zavlanos@duke.edu

stepsizes, which often makes them slow. Faster algorithms using constant stepsizes include the incremental aggregated gradient (IAG) method [12], the exact first-order algorithm (EXTRA) [13], and Accelerated Distributed Augmented Lagrangian (ADAL) algorithm. See also [14–16] for optimization of convex problems.

Optimization of nonconvex functions is a much more challenging problem. Only recently, have there been developed a few nonconvex distributed optimization algorithms motivated by applications in resource allocation in ad-hoc network [17], sparse PCA [18], and flow control in communication networks [19]; see also [19–27] for additional algorithms developed for nonconvex optimization problems.

Regardless of convexity and/or smoothness, all the aforementioned methods require that either the first order (gradient/subgradient) information or the explicit form of the objective function is available to the agents. However, in many important practical situations such information can be expensive to obtain, or even impossible. Examples include, simulation-based optimization where the objective function can only be evaluated using repeated simulation [28], training deep neural networks where the relationship between the variables and the cost function is too complicated to derive an explicit form of the gradient [29], and bandit optimization where a player optimizes a sequence of cost functions having only knowledge of a single function value each time [30]. In these cases, the zeroth-order information about the objective function values is often readily available. Such zeroth-order information can be obtained through a stochastic zeroth-order oracle ($\mathcal{SZO}$). In particular, suppose that $\hat{x} \in \text{dom}(f_i)$, then the i th $\mathcal{SZO}$ at agent i returns a noisy version of $f_i(\hat{x})$ denoted by $\mathcal{H}_i(\hat{x}, \xi)$ that satisfies

$$\mathbb{E}_{\xi}[\mathcal{H}_i(\hat{x}, \xi)] = f_i(\hat{x}), \quad (2)$$

where ξ is a random variable representing the noise.

Recently, centralized zeroth-order optimization has received significant attention. In [31], Nesterov proposed a general framework for analyzing zeroth-order algorithms and provided the global convergence rates for both convex and nonconvex problems. In [32], the authors established a stochastic zeroth-order method, which again can deal with both convex and nonconvex (but smooth) optimization problems. In [33] a Mirror Descent based zeroth-order algorithm was proposed for solving convex optimization problems. Both these zeroth-order algorithms are centralized and cannot be implemented over a multi-agent network. A few recent works [34–36] considered zeroth-order distributed convex (possibly nonsmooth) problems, but none of these works can address nonconvex problems. For distributed nonconvex but smooth optimization problem using zeroth-order information a primal-dual algorithm has been proposed in [37, 38].

In this paper, we propose a new algorithm for distributed nonconvex and nonsmooth optimization with zeroth-order information. Specifically, we first show that this problem can be reformulated as a linearly constrained optimization problem over a connected multi-agent network. Then, we propose a nonconvex primal-dual based algorithm, which requires only local communication among the agents, and utilizes local zeroth-order information. Theoretically, We show that the current solution converges approximately to a stationary solution of the problem. We also provide numerical results that corroborate the theoretical findings.

Notation. Given a vector a and a matrix A , we use $\|a\|$ and $\|A\|$ to denote the Euclidean norm of vector a , and spectral norm of matrix A , respectively. $A^\top$ represents the transpose of matrix A . We define $\|a\|_A^2 := a^\top A a$. The notation $\langle a, b \rangle$ is used to denote the inner product of two vectors a, b . For matrices A and B , $A \otimes B$ is the Kronecker product of A and B . To denote an $M \times M$ identity matrix we use I_M . $\mathbb{E}[\cdot]$ denotes the expectation with respect to all random variables, and $\mathbb{E}_v[\cdot]$ denotes the expectation with respect to the random variable v .

2 Problem Definition and Proposed Algorithm

Let us define a graph $\mathcal{G} = \{\mathcal{V}, \mathcal{E}\}$, where $\mathcal{V}$ is the node set with $|\mathcal{V}| = N$, and $\mathcal{E} \subseteq \mathcal{V} \times \mathcal{V}$ is the edge set with $|\mathcal{E}| = E$. We assume that $\mathcal{G}$ is undirected, meaning that if $(i, j) \in \mathcal{E}$ then $(j, i) \in \mathcal{E}$. Moreover, every agent i can only communicate with its direct neighbors in the set $\mathcal{N}_i = \{j \in \mathcal{V} \mid (i, j) \in \mathcal{E}\}$, and let $d_i = |\mathcal{N}_i|$ denote the degree of node i . We assume that the graph $\mathcal{G}$ is connected, meaning that there is a path, i.e., a sequence of nodes where consecutive nodes are neighbors, between any two nodes in $\mathcal{G}$.

In order to decompose problem (1) let us introduce N new variables $x_i \in \mathbb{R}^M$ that are local to every agent i . Then, problem (1) can be reformulated as follows:

$$\begin{aligned} \min_{\{x_i\}} \quad & \sum_{i=1}^N f_i(x_i), \\ \text{s.t.} \quad & x_i = x_j \quad \forall (i, j) \in \mathcal{E}. \end{aligned} \quad (3)$$

The set of constraints $x_i = x_j$ enforce consensus on the local variables x_i and x_j for all neighbors $j \in \mathcal{N}_i$. We stack all the local variables x_i in a vector $x := \{x_i\} \in \mathbb{R}^{Q \times 1}$, where $Q = NM$. Moreover, we define the Degree matrix $\tilde{D} \in \mathbb{R}^{N \times N}$ to be a diagonal matrix where $\tilde{D}(i, i) = d_i$; let $D = \tilde{D} \otimes I_M \in \mathbb{R}^{Q \times Q}$. For a given graph $\mathcal{G}$, the incidence matrix $\tilde{A} \in \mathbb{R}^{E \times N}$ is a matrix where $\tilde{A}(k, i) = 1$ and $\tilde{A}(k, j) = -1$, where (i, j) is the k th edge of $\mathcal{G}$; the rest of the entries of $\tilde{A}$ are all zero. Let $A = \tilde{A} \otimes I_M \in \mathbb{R}^{EM \times Q}$. Finally, we define the Signed and Signless Laplacian matrix, denoted by L^- and L^+ , respectively as

$$L^- := A^\top A \in \mathbb{R}^{Q \times Q}, \quad (4)$$

$$L^+ := 2D - A^\top A \in \mathbb{R}^{Q \times Q}. \quad (5)$$

Using the above notations, problem (3) can be written in the following compact form:

$$\min_{x \in \mathbb{R}^Q} f(x), \quad \text{s.t.} \quad Ax = 0. \quad (6)$$

2.1 Preliminaries

In this section, we first introduce some standard techniques presented in [31] for approximating and smoothing the gradient of a given function. Suppose that $\phi \in \mathbb{R}^Q$ is a Gaussian random vector and let $\mu > 0$ be some smoothing parameter. The smoothed version of function f is defined as

$$f_\mu(x) = \mathbb{E}_\phi(f(x + \mu\phi)) = \frac{1}{(2\pi)^{\frac{Q}{2}}} \int f(x + \mu\phi) e^{-\frac{1}{2}\|\phi\|^2} d\phi. \quad (7)$$

Then it can be shown that the function f_μ is differentiable and its gradient is given by Eq. (22) in [31]

$$\nabla f_\mu(x) = \frac{1}{(2\pi)^{\frac{Q}{2}}} \int \frac{f(x + \mu\phi) - f(x)}{\mu} \phi e^{-\frac{1}{2}\|\phi\|^2} d\phi. \quad (8)$$

Further, assuming that the original function f is Lipschitz continuous, i.e., there exists L_0 such that $|f(x) - f(y)| \leq L_0|x - y|$ for all $x, y \in \text{dom}(f)$, it can be shown that (see [31, Lemma 2]) ∇f_μ is also Lipschitz continuous with constant $L_1 = \frac{2L_0\sqrt{Q}}{\mu}$. In other words, for all $x, y \in \text{dom}(f_\mu)$ we have

$$\|\nabla f_\mu(x) - \nabla f_\mu(y)\| \leq L_1\|x - y\|. \quad (9)$$

Let $\mathcal{H}(x, \xi)$ denote the noisy functional value of the function f obtained from an associated $\mathcal{SZO}$ as in equation (2). In view of (8), the gradient of $f_\mu(x)$ can be approximated as

$$G_\mu(x, \phi, \xi) = \frac{\mathcal{H}(x + \mu\phi, \xi) - \mathcal{H}(x, \xi)}{\mu} \phi, \quad (10)$$

where the constant $\mu > 0$ is the smoothing parameter. It can be easily checked that $G_\mu(x, \phi, \xi)$ is an unbiased estimator of $\nabla f_\mu(x)$, i.e.,

$$\mathbb{E}_{\xi, \phi}[G_\mu(x, \phi, \xi)] = \mathbb{E}_\phi[\mathbb{E}_\xi[G_\mu(x, \phi, \xi) \mid \phi]] = \nabla f_\mu(x). \quad (11)$$

For simplicity we define $\zeta := (\xi, \phi)$. For a given number J of independent samples of $\{\zeta_j\}_{j=1}^J$, we define the sample average $\bar{G}_\mu(x, \zeta) := \frac{1}{J} \sum_{j=1}^J G_\mu(x, \zeta_j)$, where $\zeta := \{\zeta_j\}_{j=1}^J$. It is easy to see that for any $J \geq 1$, $\bar{G}_\mu(x, \zeta)$ is also an unbiased estimator of $\nabla f_\mu(x)$.

2.2 The Proposed Algorithm

In this part we propose a primal-dual algorithm for the distributed optimization problem (6). Let $\lambda_{ij} \in \mathbb{R}^M$ be the multiplier associated with the consensus constraint $x_i - x_j = 0$ for each $(i, j) \in \mathcal{E}$. Moreover, stack all λ_{ij} 's in a vector $\lambda = \{\lambda_{ij}\}_{(i,j) \in \mathcal{E}} \in \mathbb{R}^{EM}$. Then, the augmented Lagrangian (AL) function for problem (6) is given by

$$U_\rho(x, \lambda) := f(x) + \langle \lambda, Ax \rangle + \frac{\rho}{2} \|Ax\|_2^2. \quad (12)$$

where $\rho > 0$ is a constant. Moreover, as in (7) define the smoothed version $f_{i,\mu}$ of the local function f_i . At iteration r of the algorithm we obtain an unbiased estimation of the gradient of local function $f_{i,\mu}(x_i^r)$ as follows. For every sample $j \in \{1, 2, \dots, J\}$ we generate a random vector $\phi_{i,j}^r \in \mathbb{R}^M$ from an i.i.d standard Gaussian distribution and calculate $\bar{G}_{\mu,i}(x_i^r, \zeta_i^r) \in \mathbb{R}^M$ similar to (10) by

$$\bar{G}_{\mu,i}(x_i^r, \zeta_i^r) = \frac{1}{J} \sum_{j=1}^J \frac{\mathcal{H}_i(x_i^r + \mu\phi_{i,j}^r, \xi_{i,j}^r) - \mathcal{H}_i(x_i^r, \xi_{i,j}^r)}{\mu} \phi_{i,j}^r. \quad (13)$$

Define $G_\mu^{J,r} := \{\bar{G}_{\mu,i}(x_i^r, \zeta_i^r)\}_{i=1}^N \in \mathbb{R}^Q$. The following theorem bounds the norm of $G_\mu^{J,r}$.

Algorithm 1 The Proposed Algorithm for problem (1)

- 1: **Input:** Degree matrix $D \in \mathbb{R}^{Q \times Q}$, total number of iterations $T \geq 1$, number of samples $J \geq 1$, smoothing parameter $\mu > 0$
 - 2: **Initialize:** Primal variable $x^0 \in \mathbb{R}^Q$, dual variable $\lambda^0 = 0 \in \mathbb{R}^{EM}$
 - 3: **for** $r = 0$ **to** $T - 1$ **do**
Update primal variable x and dual variable λ by
$$x^{r+1} = \underset{x}{\operatorname{argmin}} \langle G_\mu^{J,r} + A^\top \lambda^r + \rho A^\top A x^r, x - x^r \rangle + \rho \|x - x^r\|_D^2, \text{ (Primal Step)} \quad (15)$$

$$\lambda^{r+1} = \lambda^r + \rho A x^{r+1}. \quad \text{ (Dual Step)} \quad (16)$$
 - 4: **end for**
 - 5: Choose uniformly randomly $u \in \{0, 1, \dots, T - 1\}$
 - 6: **Output:** (z^u, λ^u) .
-

Theorem 1 (Theorem 4 [31]) *If f is a Lipschitz continuous function with constant L_0 , then*

$$\mathbb{E}_\zeta[\|G_\mu^{J,r}\|^2] \leq \frac{L_0^2(Q+4)^2}{J^2}. \quad (14)$$

Our proposed algorithm is summarized in Algorithm 1. In the primal step (15), an *approximate gradient descent* step is taken towards minimizing the augmented Lagrangian function with respect to x . In particular, in the first-order approximation of the AL function the true gradient of the function $f(x^r)$ is approximated by the noisy zeroth-order estimate $G_\mu^{J,r}$ and then, a matrix-weighted quadratic penalty $\rho \|x - x^r\|_D$ is used. This term is critical for the algorithm itself, as well as for the analysis. The dual step (16) is then performed, which is a *gradient ascent* step over the dual variable λ .

To see how Algorithm 1 can be implemented in a distributed way, consider the optimality condition for (15) as

$$G_\mu^{J,r} + A^\top (\lambda^r + \rho A x^r) + 2\rho D(x^{r+1} - x^r) = 0. \quad (17)$$

Utilizing (4) and (5), we have

$$x^{r+1} = \frac{1}{2\rho} D^{-1} \left[\rho L^+ x^r - G_\mu^{J,r} + A^\top \lambda^r \right].$$

To implement this primal iteration, each agent i only requires local information as well as information from its neighbors $\mathcal{N}_i$. This is because D is a diagonal matrix and the structure of the matrix L^+ ensures that the i th block vector of $L^+ x^r$ is only related to x_j^r , $j \in \mathcal{N}_i$. For the dual step w.l.o.g we assign the dual variable λ_{ij} to node i and therefore, from (16) we have

$$\lambda_{ij}^{r+1} = \lambda_{ij}^r + \rho(x_i^{r+1} - x_j^{r+1}), \quad (18)$$

which only requires the local information as well as information from the neighbors in $\mathcal{N}_i$.

3 The Convergence Analysis

In this section we study the convergence of Algorithm 1. We make the following assumptions.

Assumptions A. We assume that

- A1. The function f is Lipschitz continuous.
- A2. The function f is lower bounded.

The above assumptions on the objective f are quite standard in the analysis of first order optimization Algorithms (1). To simplify notation let $\mathcal{F}^r := \sigma(\zeta_1, \zeta_2, \dots, \zeta_r)$ be the σ -field generated by the entire history of algorithm up to iteration r , $\sigma_{\min}$ be the smallest nonzero eigenvalue of $A^\top A$, and $w^r := (x^{r+1} - x^r) - (x^r - x^{r-1})$ be the successive difference of the differences of the primal iterates. In the analysis that follows we will make use of the following relations:

- For any given vectors a and b we have

$$\langle b - a, b \rangle = \frac{1}{2}(\|b\|^2 + \|a - b\|^2 - \|a\|^2), \quad (19)$$

$$\langle a, b \rangle \leq \frac{1}{2\epsilon}\|a\|^2 + \frac{\epsilon}{2}\|b\|^2; \quad \forall \epsilon > 0. \quad (20)$$

- For n given vectors a_i we have that

$$\left\| \sum_{i=1}^n a_i \right\|^2 \leq n \sum_{i=1}^n \|a_i\|^2. \quad (21)$$

Our proof consists of a series of lemmas leading to the main convergence rate result. In our presentation we try to provide insights about the analysis steps, while the proofs of the results are relegated to the appendix.

First we bound the difference between $\nabla f_\mu(x^r)$ and its unbiased estimation $G_\mu^{J,r}$ as follows:

$$\begin{aligned} \mathbb{E}_\zeta \|G_\mu^{J,r} - \nabla f_\mu(x^r)\|^2 &= \mathbb{E}_\zeta \|G_\mu^{J,r} - \mathbb{E}_\zeta[G_\mu^{J,r}]\|^2 \\ &\leq \mathbb{E}_\zeta \|G_\mu^{J,r}\|^2 \\ &\leq \frac{L_0^2(Q+4)^2}{J^2}, \end{aligned} \quad (22)$$

where the last inequality follows from (14). The next lemma bounds the change of the dual variables by that of the primal variables. The proofs of the results that follow can be found in the appendix.

Lemma 1 *Suppose Assumptions A hold true. Let L_1 denote the gradient Lipschitz constant for function f_μ . Then we have the following inequity:*

$$\frac{1}{\rho} \mathbb{E} \|\lambda^{r+1} - \lambda^r\|^2 \leq \frac{9L_0^2(Q+4)^2}{\rho\sigma_{\min}J^2} + \frac{6L_1^2}{\rho\sigma_{\min}} \mathbb{E} \|x^r - x^{r-1}\|^2 + \frac{3\rho\|L^+\|}{\sigma_{\min}} \mathbb{E} \|w^r\|_{L^+}^2. \quad (23)$$

The next step is the key in our analysis. We define the smoothed version of the AL function in a similar way as (7) and denote it by $U_{\rho,\mu}(x, \lambda)$. For notational simplicity let us define $U_{\rho,\mu}^{r+1} := U_{\rho,\mu}(x^{r+1}, \lambda^{r+1})$. From equation (9) we know that function f_μ is Lipschitz continuous with constant L_1 . Now let $c > 0$ be some positive constant and set $k := 2(\frac{6L_1^2}{\rho\sigma_{\min}} + \frac{3cL_1}{2})$. Moreover, we define

$$V^{r+1} := \frac{\rho}{2} (\|Ax^{r+1}\|^2 + \|x^{r+1} - x^r\|_B^2),$$

where $B := L^+ + \frac{k}{c\rho}I_Q$. Finally we define the following potential function:

$$P^{r+1} := U_{\rho,\mu}^{r+1} + cV^{r+1}. \quad (24)$$

We study the behavior of the proposed potential function as the algorithm proceeds.

Lemma 2 *Suppose Assumptions A hold true. We have that*

$$\mathbb{E}[P^{r+1} - P^r] \leq -\alpha_1 \mathbb{E}\|x^{r+1} - x^r\|^2 - \alpha_2 \mathbb{E}\|w^r\|_{L^+}^2 + \alpha_3 \frac{L_0^2(Q+4)^2}{J^2}, \quad (25)$$

where

$$\begin{aligned} \alpha_1 &:= \rho^2 - (2cL_1 + L_1/2 + L_1^2/2 + 1/2)\rho - \frac{6L_1^2}{\sigma_{\min}}, \\ \alpha_2 &:= \frac{3\rho\|L^+\|}{\sigma_{\min}} - \frac{c\rho}{2}, \quad \alpha_3 = \frac{9}{\rho\sigma_{\min}} + \frac{6c+1}{L_1}. \end{aligned} \quad (26)$$

Note that the constants α_1 and α_2 in (25) can be made positive as long as we choose constants c and ρ large enough. In particular, the following conditions are sufficient to have positive α_1, α_2

$$c > \frac{6\|L^+\|}{\sigma_{\min}}, \quad \rho > b + \sqrt{b^2 + 6L_1^2/\sigma_{\min}}, \quad (27)$$

where $b = (cL_1 + L_1/4 + L_1^2/4 + 1/4)$.

The key insight obtained from Lemma 2 is that, a proper combination of a primal objective (i.e., the AL function), and the dual gap (i.e., the violation of the feasibility) can be served as the potential function that guides the progress of the algorithm.

In the next lemma we show that P^{r+1} is lower bounded.

Lemma 3 *Suppose Assumptions A hold true, and constant c is selected as $c \geq \frac{2\|L^+\|}{\sigma_{\min}}$. Then there exists a constant $\underline{P}$ that is independent of the total number of iterations T so that*

$$\mathbb{E}[P^{r+1}] \geq \underline{P} > -\infty, \quad \forall r \geq 1. \quad (28)$$

To characterize the convergence rate of Algorithm 1, let us define the *stationarity gap* of the smoothed version of problem (6) as

$$\Phi_\mu(x^r, \lambda^{r-1}) := \mathbb{E} [\|\nabla_x U_{\rho,\mu}(x^r, \lambda^{r-1})\|^2 + \|Ax^r\|^2]. \quad (29)$$

It can be easily checked that $\Phi_\mu(x^*, \lambda^*) = 0$ if and only if (x^*, λ^*) is a KKT point of the smoothed version of problem (6). For simplicity let us denote $\Phi_\mu^r := \Phi_\mu(x^r, \lambda^{r-1})$.

At this point we are ready to combine the previous results to obtain our main theorem.

Theorem 2 *Suppose Assumptions A hold true, the penalty parameter ρ satisfies the condition given in Lemma 2, and the constant c satisfies $c \geq \frac{6\|L^+\|}{\sigma_{\min}}$. Then, there exist constants $\gamma_1, \gamma_2 > 0$ such that*

$$\mathbb{E}_u[\Phi_\mu^u] \leq \frac{\gamma_1}{T} + \gamma_2 \frac{L_0(Q+4)^2}{J^2}. \quad (30)$$

From Theorem 2 we can observe that there exists always a constant term in the right-hand-side of the stationarity gap. Therefore, no matter how many iterations we run the algorithm, we always converge to a neighborhood of a stationary point. However, if we choose the number of samples $J \in \mathcal{O}(\sqrt{T})$, we have the following bound:

$$\mathbb{E}_u[\Phi_\mu^u] \leq \frac{\gamma_1}{T} + \frac{\gamma_2 L_0(Q+4)^2}{T}, \quad (31)$$

which verifies the sublinear convergence rate for the algorithm.

4 Numerical Results

In this section we illustrate the proposed algorithm through numerical simulations. For our experiments we study two nonconvex distributed optimization problems.

First, we consider a simple nonconvex nonsmooth distributed optimization problem defined below:

$$\min_{x \in \mathbb{R}^Q} \sum_{i=1}^N f_i(x_i), \quad \text{s.t. } x_i = x_j, \quad \forall (i, j) \in \mathcal{E}, \quad (32)$$


where for each agent we have

$$f_i(x_i) = |\cos(x_i) + \|x_i\| + \exp(x_i)|.$$


In (32) the problem dimension $M = 1$ and the number of nodes in the network is $N = 10$. Therefore, $Q = MN = 10$. The details of the underlying graph are discussed in [39]. We compare Algorithm 1 using a constant stepsize that satisfies (27) and the Randomize Gradient Free (RGF) algorithm proposed in [34] using diminishing stepsize $\frac{1}{\sqrt{r}}$. Note that in theory RGF only works for the convex problems. However, we include it here for the purpose of comparison only. We compare the two algorithms in terms of the stationarity gap defined in (29) and the constraint violation $\|Ax\|$. The stopping criterion is set to $T = 1000$ iterations and the results are the average over 30 independent trials. Figs 1a and 1b show our comparative results. We observe that the stationarity gap and the consensus error vanish faster for our proposed algorithm than for the RGF algorithm. Next, we study a mini-batch binary classification problem using nonconvex nonsmooth regularizers, where each node stores b (batch size) data points. For this problem the local function is given by

$$f_i(x_i) = \frac{1}{Nb} \left[\sum_{j=1}^b \log(1 + \exp(-y_{ij} x_i^T v_{ij})) + \alpha \log(\epsilon + \|x_i\|_1) \right],$$


where $v_{ij} \in \mathbb{R}^M$ and $y_{ij} \in \{1, -1\}$ are the feature vector and the label for the j th data point of i th agent [40]. The nonconvex nonsmooth regularization term $\log(\epsilon + \|x_i\|_1)$ imposes




(a) Comparison of proposed Algorithm 1, and RGF algorithm [34] in terms of the stationarity gap for the nonconvex nonsmooth distributed optimization problem (32).



(b) Comparison of proposed Algorithm 1, and RGF algorithm [34] in terms of the constraint violation (i.e. $\|Ax\|$) for the nonconvex nonsmooth distributed optimization problem (32).



(a) Comparison of proposed Algorithm 1, and RGF algorithm [34] in terms of the stationarity gap for binary classification problem.



(b) Comparison of proposed Algorithm 1, and RGF algorithm [34] in terms of the constraint violation (i.e. $\|Ax\|$) for binary classification problem.

sparsity to vector x_i , the constant α controls the sparsity level, and $\epsilon > 0$ is a small number. The network has $N = 15$ nodes and each node contains randomly generated $b = 100$ data point. Algorithm 1 and RGF run for $T = 1000$ iterations and Figures 2a and 2b illustrate the stationarity gap and the constraint violation versus the iteration counter. From these plots we can observe that, as in the previous problem, Algorithm 1 is faster than the RGF. Note again that, in theory, RGF is designed for convex problems only. To the best of our knowledge, our algorithm is the first provable distributed zeroth-order method for nonconvex and nonsmooth problems.

5 Conclusion

In this work, we proposed a distributed gradient-free optimization algorithm to solve nonconvex and nonsmooth problems utilizing local zeroth-order information. We rigorously analyzed the convergence rate of the proposed algorithm and demonstrated its performance

via simulation. To the best of our knowledge, this is the first distributed framework for the solution of nonconvex and nonsmooth distributed optimization problems that also has a provable sublinear convergence rate. The proposed framework can be used to solve a variety of problems where access to first or second order information is very expensive or even impossible.

APPENDIX

5.1 Proof of Lemma 1

From equation (16) we have

$$\lambda^{r+1} - \lambda^r = \rho A x^{r+1}. \quad (33)$$

Equation (33) implies that $\lambda^{r+1} - \lambda^r$ lies in the column space of A , therefore we have

$$\sqrt{\sigma_{\min}} \|\lambda^{r+1} - \lambda^r\| \leq \|A^\top (\lambda^{r+1} - \lambda^r)\|, \quad (34)$$

where $\sigma_{\min}$ denotes the smallest non-zero eigenvalue of $A^\top A$. Utilizing equation (33) and equation (17), we obtain

$$G_\mu^{J,r} + A^\top \lambda^{r+1} + \rho L^+ (x^{r+1} - x^r) = 0. \quad (35)$$

Replacing r with $r-1$ in equation (35) and then using the definition of $w^r := (x^{r+1} - x^r) - (x^r - x^{r-1})$ we obtain

$$\begin{aligned} \frac{1}{\rho} \|\lambda^{r+1} - \lambda^r\|^2 &\leq \frac{1}{\rho \sigma_{\min}} \|G_\mu^{J,r} - G_\mu^{J,r-1} + \rho L^+ w^r\|^2 \\ &= \frac{1}{\rho \sigma_{\min}} \|G_\mu^{J,r} - G_\mu^{J,r-1} + \nabla f_\mu(x^r) - \nabla f_\mu(x^{r-1}) + \rho L^+ w^r\|^2 \\ &\leq \frac{3}{\rho \sigma_{\min}} \|G_\mu^{J,r} - \nabla f_\mu(x^r)\|^2 + \frac{3}{\rho \sigma_{\min}} \|\nabla f_\mu(x^r) - G_\mu^{J,r-1}\|^2 + \frac{3\rho^2}{\rho \sigma_{\min}} \|L^+ w^r\|^2, \end{aligned} \quad (36)$$

where the last inequality follows from (21). Adding and subtracting $\nabla f_\mu(x^{r-1})$ to the second term on the r.h.s of (36) and taking the expectation on both sides gives

$$\begin{aligned} \frac{1}{\rho} \mathbb{E} \|\lambda^{r+1} - \lambda^r\|^2 &\leq \frac{3}{\rho \sigma_{\min}} \mathbb{E} \|G_\mu^{J,r} - \nabla f_\mu(x^r)\|^2 \\ &\quad + \frac{6}{\rho \sigma_{\min}} \mathbb{E} \|\nabla f_\mu(x^r) - \nabla f_\mu(x^{r-1})\|^2 + \frac{3\rho^2}{\rho \sigma_{\min}} \mathbb{E} \|L^+ w^r\|^2 \\ &\quad + \frac{6}{\rho \sigma_{\min}} \mathbb{E} \|\nabla f_\mu(x^{r-1}) - G_\mu^{J,r-1}\|^2 \\ &\leq \frac{9L_0^2(Q+4)^2}{\rho \sigma_{\min} J^2} + \frac{6L_1^2}{\rho \sigma_{\min}} \mathbb{E} \|x^r - x^{r-1}\|^2 + \frac{3\rho \|L^+\|}{\sigma_{\min}} \mathbb{E} \|w^r\|_{L^+}^2, \end{aligned} \quad (37)$$

where the last inequity is true because of (22), the fact that $\nabla f_\mu(z)$ is gradient Lipschitz with constant L_1 , and the inequality $\|L^+ w^r\|^2 \leq \|L^+\| \|w^r\|_{L^+}^2$. The proof is complete.

5.2 Proof of Lemma 2

First we prove that the function $g(x) := U_{\rho,\mu}(x, \lambda) + \frac{\rho}{2}\|x - x^r\|_{L^+}^2$ is strongly convex with respect to variable x when $2\rho \geq L_1$. From Assumption A.1 and the fact that $D \succeq I$ we have

$$\begin{aligned} \langle \nabla g(x_1) - \nabla g(x_2), x_1 - x_2 \rangle &= \langle \nabla f_\mu(x_1) - \nabla f_\mu(x_2) + \rho(A^T A + L^+)(x_1 - x_2), x_1 - x_2 \rangle \\ &= \langle \nabla f_\mu(x_1) - \nabla f_\mu(x_2) + 2\rho D(x_1 - x_2), x_1 - x_2 \rangle \\ &= \langle \nabla f_\mu(x_1) - \nabla f_\mu(x_2), x_1 - x_2 \rangle + 2\rho\|x_1 - x_2\|_D^2 \\ &\geq \langle \nabla f_\mu(x_1) - \nabla f_\mu(x_2), x_1 - x_2 \rangle + 2\rho\|x_1 - x_2\|^2 \\ &\geq -L_1\|x_1 - x_2\|^2 + 2\rho\|x_1 - x_2\|^2 = (2\rho - L_1)\|x_1 - x_2\|^2. \end{aligned}$$

This proves that $U_{\rho,\mu}(x, \lambda) + \frac{\rho}{2}\|x - x^r\|_{L^+}^2$ is strongly convex with modulus $2\rho - L_1$. Using this fact, we can bound $U_{\rho,\mu}^{r+1} - U_{\rho,\mu}^r$ as follows:

$$\begin{aligned} U_{\rho,\mu}^{r+1} - U_{\rho,\mu}^r &= U_{\rho,\mu}(x^{r+1}, \lambda^{r+1}) - U_{\rho,\mu}(x^{r+1}, \lambda^r) + U_{\rho,\mu}(x^{r+1}, \lambda^r) - U_{\rho,\mu}(x^r, \lambda^r) \\ &\leq \frac{1}{\rho}\|\lambda^{r+1} - \lambda^r\|^2 + \langle \nabla_x U_{\rho,\mu}(x^{r+1}, \lambda^r) + \rho L^+(x^{r+1} - x^r), x^{r+1} - x^r \rangle \\ &\quad - \frac{2\rho - L_1}{2}\|x^{r+1} - x^r\|^2, \end{aligned} \tag{38}$$

where the last inequity holds true due to the strong convexity of $U_{\rho,\mu}(x, \lambda) + \frac{\rho}{2}\|x - x^r\|_{L^+}^2$ and (33). Now using (35) we have

$$\begin{aligned} U_{\rho,\mu}^{r+1} - U_{\rho,\mu}^r &\leq \langle \nabla f_\mu(x^{r+1}) - G_\mu^{J,r}, x^{r+1} - x^r \rangle + \frac{1}{\rho}\|\lambda^{r+1} - \lambda^r\|^2 - \frac{2\rho - L_1}{2}\|x^{r+1} - x^r\|^2 \\ &\leq \frac{1}{\rho}\|\lambda^{r+1} - \lambda^r\|^2 + \frac{L_1^2 - 2\rho + L_1}{2}\|x^{r+1} - x^r\|^2 + \frac{1}{2L_1^2}\|\nabla f_\mu(x^{r+1}) - G_\mu^{J,r}\|^2, \end{aligned}$$

here the last inequality follows from (20) with $\epsilon = L_1^2$. Taking expectation on both sides we have

$$\begin{aligned} \mathbb{E}[U_{\rho,\mu}^{r+1} - U_{\rho,\mu}^r] &\leq \frac{9L_0^2(Q+4)^2}{\rho\sigma_{\min}J^2} + \frac{6L_1^2}{\rho\sigma_{\min}}\mathbb{E}\|x^r - x^{r-1}\|^2 \\ &\quad + \frac{3\rho\|L^+\|}{\sigma_{\min}}\mathbb{E}\|w^r\|_{L^+}^2 + \frac{L_1^2 - 2\rho + L_1}{2}\mathbb{E}\|x^{r+1} - x^r\|^2 \\ &\quad + \frac{1}{2L_1^2}\mathbb{E}\|\nabla f_\mu(x^{r+1}) - \nabla f_\mu(x^r) + \nabla f_\mu(x^r) - G_\mu^{J,r}\|^2 \\ &\leq \left(\frac{9}{\rho\sigma_{\min}} + \frac{1}{L_1^2}\right)\frac{L_0^2(Q+4)^2}{J^2} + \frac{6L_1^2}{\rho\sigma_{\min}}\mathbb{E}\|x^r - x^{r-1}\|^2 \\ &\quad + \frac{3\rho\|L^+\|}{\sigma_{\min}}\mathbb{E}\|w^r\|_{L^+}^2 + \frac{L_1^2 - 2\rho + L_1 + 1}{2}\mathbb{E}\|x^{r+1} - x^r\|^2, \end{aligned} \tag{39}$$

where the first inequality follows from (37) and the second inequality follows from (22). Now we bound $V^{r+1} - V^r$. From the optimality condition for problem (15) and the dual update (16) we have

$$\langle G_\mu^{J,r} + A^\top \lambda^{r+1} + \rho L^+(x^{r+1} - x^r), x^{r+1} - x \rangle \leq 0, \quad \forall x \in \mathbb{R}^Q.$$

Similarly, for the $(r-1)$ th iteration, we have

$$\langle G_\mu^{J,r-1} + A^\top \lambda^r + \rho L^+(x^r - x^{r-1}), x^r - x \rangle \leq 0, \forall x \in \mathbb{R}^Q.$$

Setting $x = x^r$ in first equation, $x = x^{r+1}$ in second equation, and adding them, we obtain

$$\langle A^\top (\lambda^{r+1} - \lambda^r), x^{r+1} - x^r \rangle \leq -\langle G_\mu^{J,r} - G_\mu^{J,r-1} + \rho L^+ w^r, x^{r+1} - x^r \rangle. \quad (40)$$

The l.h.s can be expressed as follows:

$$\begin{aligned} \langle A^\top (\lambda^{r+1} - \lambda^r), x^{r+1} - x^r \rangle &= \rho \langle Ax^{r+1}, Ax^{r+1} - Ax^r \rangle \\ &= \frac{\rho}{2} \left(\|Ax^{r+1}\|^2 - \|Ax^r\|^2 + \|A(x^{r+1} - x^r)\|^2 \right), \end{aligned} \quad (41)$$

where the first equality follows from (16) and the second equality follows from (19). For the r.h.s of (40) we have

$$\begin{aligned} & -\langle G_\mu^{J,r} - G_\mu^{J,r-1} + \rho L^+ w^r, x^{r+1} - x^r \rangle \\ &= -\langle G_\mu^{J,r} - G_\mu^{J,r-1}, x^{r+1} - x^r \rangle - \langle \rho L^+ w^r, x^{r+1} - x^r \rangle \\ &\leq \frac{1}{2L_1} \|G_\mu^{J,r} - G_\mu^{J,r-1}\|^2 + \frac{L_1}{2} \|x^{r+1} - x^r\|^2 - \rho \langle L^+ w^r, x^{r+1} - x^r \rangle \\ &\leq \frac{3}{2L_1} \left(\|G_\mu^{J,r} - \nabla f_\mu(x^r)\|^2 + \|\nabla f_\mu(x^{r-1}) - G_\mu^{J,r-1}\|^2 + \|\nabla g_\mu(x^r) - \nabla g_\mu(x^{r-1})\|^2 \right) \\ &+ \frac{L_1}{2} \|x^{r+1} - x^r\|^2 - \rho \langle L^+ w^r, x^{r+1} - x^r \rangle, \end{aligned}$$

where the first inequality follows from (20). To get the second inequality we add and subtract $\nabla f_\mu(x^r) + \nabla f_\mu(x^{r-1})$ to $G_\mu^{J,r} - G_\mu^{J,r-1}$ and use (21). Taking expectation on both sides, we have

$$\begin{aligned} & -\mathbb{E}[\langle G_\mu^{J,r} - G_\mu^{J,r-1} + \rho L^+ w^r, x^{r+1} - x^r \rangle] \\ &\leq \frac{6L_0^2(Q+4)^2}{L_1 J^2} + \frac{3L_1}{2} \mathbb{E}\|x^r - x^{r-1}\|^2 + \frac{L_1}{2} \mathbb{E}\|x^{r+1} - x^r\|^2 - \rho \mathbb{E}[\langle L^+ w^r, x^{r+1} - x^r \rangle] \\ &= \frac{6L_0^2(Q+4)^2}{L_1 J^2} + \frac{3L_1}{2} \mathbb{E}\|x^r - x^{r-1}\|^2 + \frac{L_1}{2} \mathbb{E}\|x^{r+1} - x^r\|^2 \\ &+ \frac{\rho}{2} \mathbb{E} \left[\|x^r - x^{r-1}\|_{L^+}^2 - \|x^{r+1} - x^r\|_{L^+}^2 - \|w^r\|_{L^+}^2 \right], \end{aligned} \quad (42)$$

where the inequality follows from (22) and the last equality follows from (19). Combining (41) and (42), we obtain

$$\begin{aligned} & \frac{\rho}{2} \mathbb{E} \left(\|Ax^{r+1}\|^2 - \|Ax^r\|^2 + \|A(x^{r+1} - x^r)\|^2 \right) \\ &\leq \frac{6L_0^2(Q+4)^2}{L_1 J^2} + \frac{3L_1}{2} \mathbb{E}\|x^r - x^{r-1}\|^2 + \frac{L_1}{2} \mathbb{E}\|x^{r+1} - x^r\|^2 \\ &+ \frac{\rho}{2} \mathbb{E} \left(\|x^r - x^{r-1}\|_{L^+}^2 - \|x^{r+1} - x^r\|_{L^+}^2 - \|w^r\|_{L^+}^2 \right). \end{aligned} \quad (43)$$

Rearranging terms in (43), and using the definition of V^r and B , we have

$$\begin{aligned}
\mathbb{E}[V^{r+1} - V^r] &\leq \left(\frac{L_1}{2} + \frac{k}{2c}\right) \mathbb{E}\|x^{r+1} - x^r\|^2 + \frac{6L_0^2(Q+4)^2}{L_1J^2} \\
&\quad + \left(\frac{3L_1}{2} - \frac{k}{2c}\right) \mathbb{E}\|x^r - x^{r-1}\|^2 - \frac{\rho}{2} \mathbb{E}\left(\|w^r\|_{L^+}^2 + \|A(x^{r+1} - x^r)\|^2\right) \\
&\leq \left(\frac{L_1}{2} + \frac{k}{2c}\right) \mathbb{E}\|x^{r+1} - x^r\|^2 + \left(\frac{3L_1}{2} - \frac{k}{2c}\right) \mathbb{E}\|x^r - x^{r-1}\|^2 \\
&\quad - \frac{\rho}{2} \mathbb{E}\|w^r\|_{L^+}^2 + \frac{6L_0^2(Q+4)^2}{L_1J^2}.
\end{aligned} \tag{44}$$

Recall the definition of $P^{r+1} := U_{\rho,\mu}^{r+1} + cV^{r+1}$. Utilizing (39) and (44) eventually we obtain

$$\mathbb{E}[P^{r+1} - P^r] \leq -\alpha \mathbb{E}\|x^{r+1} - x^r\|^2 - \alpha_2 \mathbb{E}\|w^r\|_{L^+}^2 + \alpha_3 \frac{L_0^2(Q+4)^2}{J^2},$$

where we have

$$\begin{aligned}
\alpha_1 &:= \rho^2 - (2cL_1 + L_1/2 + L^2/2 + 1/2)\rho - \frac{6L_1^2}{\sigma_{\min}}, \\
\alpha_2 &:= \frac{3\rho\|L^+\|}{\sigma_{\min}} - \frac{c\rho}{2}, \quad \alpha_3 = \frac{9}{\rho\sigma_{\min}} + \frac{6c+1}{L_1}.
\end{aligned}$$

The lemma is proved.

5.3 Proof of Lemma 3

From (35) we have

$$\|A^\top \lambda^{r+1}\|^2 \leq 2\|G_\mu^{J,r}\|^2 + 2\rho^2\|L^+(x^{r+1} - x^r)\|^2. \tag{45}$$

Recall that $\sigma_{\min}$ is the smallest nonzero eigenvalue of $A^T A$. Also from the fact that $\lambda^0 = 0$ we have that the dual variable lies in the column space of A , thus we get that

$$\|\lambda^{r+1}\|^2 \leq \frac{2}{\sigma_{\min}}\|G_\mu^{J,r}\|^2 + \frac{2\rho^2}{\sigma_{\min}}\|L^+(x^{r+1} - x^r)\|^2. \tag{46}$$

From the definition of the potential function we have

$$\begin{aligned}
P^{r+1} &= f_\mu(x^{r+1}) + \frac{\rho}{2}\|Ax^{r+1}\|^2 + \frac{1}{\rho}\|\lambda^{r+1}\|^2 - \frac{1}{2\rho}\|\lambda^{r+1}\|^2 \\
&\quad + \frac{c\rho}{2}\|Ax^{r+1}\|^2 + \frac{c\rho}{2}\|x^{r+1} - x^r\|_B^2,
\end{aligned} \tag{47}$$

where $B := L^+ + \frac{k}{c\rho}I$. From Assumption A.2 we have that f is lower bounded, therefore, f_μ is lower bounded too, i.e. there exists $\underline{f}$ such that $\underline{f} \leq f_\mu(x)$ for all $x \in \text{dom}(f_\mu)$. Plugging (46) in (47), and utilizing the fact that $\|Ax^{r+1} + \frac{1}{\rho}\lambda^{r+1}\|^2 \geq 0$ we obtain

$$\begin{aligned}
P^{r+1} &\geq \frac{-1}{\rho\sigma_{\min}}\|G_\mu^{J,r}\|^2 - \frac{\rho}{\sigma_{\min}}\|L^+(x^{r+1} - x^r)\|^2 + \frac{c\rho}{2}\|x^{r+1} - x^r\|_B^2 + \underline{f} \\
&\geq \frac{-1}{\rho\sigma_{\min}}\|G_\mu^{J,r}\|^2 + \frac{\rho}{\sigma_{\min}}\|x^{r+1} - x^r\|_{\frac{c\sigma_{\min}}{2}L^+ - (L^+)^2}^2 + \underline{f},
\end{aligned}$$

where the last inequality is due to the fact that $\frac{k}{2}\|x^{r+1} - x^r\|^2 \geq 0$. Notice that L^+ is a symmetric PSD matrix. Therefore, picking constant c large enough such that $c \geq \frac{2\|L^+\|}{\sigma_{\min}}$, we have $\frac{c\sigma_{\min}}{2}L^+ - (L^+)^2 \succeq 0$. Hence, with this choice of c we get the following bound for the potential function

$$P^{r+1} \geq -\frac{1}{\rho\sigma_{\min}}\|G_{\mu}^{J,r}\|^2 + \underline{f}. \quad (48)$$

Taking expectation on both sides we have

$$\begin{aligned} \mathbb{E}[P^{r+1}] &\geq -\frac{1}{\rho\sigma_{\min}}\mathbb{E}\|G_{\mu}^{J,r}\|^2 + \underline{f} \\ &\geq -\frac{L_0(Q+4)^2}{\rho\sigma_{\min}J^2} + \underline{f}, \end{aligned} \quad (49)$$

where the last inequality follows from (14). To complete the proof we only need to set $\underline{P} = -\frac{L_0(Q+4)^2}{\sigma_{\min}J^2} + \underline{f}$.

5.4 Proof of Theorem 2

First we bound the stationarity gap given in (29). We have

$$\begin{aligned} \|\nabla_x U_{\rho,\mu}(x^{r+1}, \lambda^r)\|^2 &= \|\nabla f_{\mu}(x^{r+1}) + A^{\top}\lambda^{r+1}\|^2 \\ &= \|\nabla f_{\mu}(x^{r+1}) - G_{\mu}^{J,r} - \rho L^+(x^{r+1} - x^r)\|^2 \\ &\leq 2\|\nabla f_{\mu}(x^{r+1}) - G_{\mu}^{J,r}\|^2 + 2\rho^2\|L^+(x^{r+1} - x^r)\|^2 \\ &\leq 2\|\nabla f_{\mu}(x^{r+1}) - \nabla f_{\mu}(x^r) + \nabla f_{\mu}(x^r) - G_{\mu}^{J,r}\|^2 + 2\rho^2\|L^+(x^{r+1} - x^r)\|^2 \\ &\leq 4\|\nabla f_{\mu}(x^{r+1}) - \nabla f_{\mu}(x^r)\|^2 + 4\|\nabla f_{\mu}(x^r) - G_{\mu}^{J,r}\|^2 \\ &\quad + 2\rho^2\|L^+(x^{r+1} - x^r)\|^2, \end{aligned}$$

where the first equality follows from (16), the second equality follows from (35), and the first inequality follows from (21). Taking expectation on both sides gives

$$\mathbb{E}\|\nabla_x U_{\rho,\mu}(x^{r+1}, \lambda^r)\|^2 \leq 4L_1^2\mathbb{E}\|x^{r+1} - x^r\|^2 + \frac{4L_0^2(Q+4)^2}{J^2} + 2\rho^2\mathbb{E}\|L^+(x^{r+1} - x^r)\|^2, \quad (50)$$

where the inequality follows from (22). Next, we bound the expected value of the constraint violation. Utilizing the equation (16) we have

$$\|Ax^{r+1}\|^2 = \frac{1}{\rho^2}\|\lambda^{r+1} - \lambda^r\|^2.$$

Taking expectation and utilizing (23) we reach

$$\begin{aligned} \mathbb{E}\|Ax^{r+1}\|^2 &= \frac{1}{\rho^2}\mathbb{E}\|\lambda^{r+1} - \lambda^r\|^2 \leq \frac{9L_0^2(Q+4)^2}{\rho^2\sigma_{\min}J^2} \\ &\quad + \frac{6L_1^2}{\rho^2\sigma_{\min}}\mathbb{E}\|x^r - x^{r-1}\|^2 + \frac{3\|L^+\|}{\sigma_{\min}}\mathbb{E}\|w^r\|_{L^+}^2. \end{aligned} \quad (51)$$

Summing (50) and (51), we have the following bound for the stationarity gap

$$\Phi_\mu^{r+1} \leq \beta_1 \mathbb{E} \|x^{r+1} - x^r\|^2 + \beta_2 \mathbb{E} \|x^r - x^{r-1}\|^2 + \beta_3 \mathbb{E} \|w^r\|_{L^+}^2 + \beta_4 \frac{L_0^2(Q+4)^2}{J^2}, \quad (52)$$

where $\beta_1, \beta_2, \beta_3, \beta_4$ are positive constants given by

$$\begin{aligned} \beta_1 &= 4L_1^2 + 2\rho^2 \|L^+\|^2, \quad \beta_2 = \frac{6L_1^2}{\rho^2 \sigma_{\min}}, \\ \beta_3 &= \frac{3\|L^+\|}{\sigma_{\min}}, \quad \beta_4 = \frac{9 + 4\rho^2 \sigma_{\min}}{\rho \sigma_{\min}}. \end{aligned}$$

Summing both sides of (52) over T iterations, we get

$$\begin{aligned} \sum_{r=1}^T \Phi_\mu^{r+1} &\leq \sum_{r=1}^{T-1} (\beta_1 + \beta_2) \mathbb{E} \|x^{r+1} - x^r\|^2 + \sum_{r=1}^T \beta_3 \mathbb{E} \|w^r\|_{L^+}^2 \\ &\quad + \beta_2 \mathbb{E} \|x^1 - x^0\|^2 + \beta_1 \mathbb{E} \|x^{T+1} - x^T\|^2 + T\beta_4 \frac{L_0^2(Q+4)^2}{J^2}. \end{aligned} \quad (53)$$

Applying Lemma 2 and summing both sides of (25) over T iterations, we obtain

$$\begin{aligned} \mathbb{E}[P^1 - P^{T+1}] &\geq \sum_{r=1}^{T-1} \alpha_1 \mathbb{E} \|x^{r+1} - x^r\|^2 + \sum_{r=1}^T \alpha_2 \mathbb{E} \|w^r\|_{L^+}^2 \\ &\quad + \alpha_1 \mathbb{E} \|x^{T+1} - x^T\|^2 - T\alpha_3 \frac{L_0^2(Q+4)^2}{J^2}. \end{aligned} \quad (54)$$

Let us set $\tau = \frac{\max(\beta_1 + \beta_2, \beta_3)}{\min(\alpha_1, \alpha_2)}$. Combining the two inequalities (53) and (54) and utilizing the fact that $\mathbb{E}[P^{T+1}]$ is lower bounded by $\underline{P}$, we arrive at the following inequality

$$\sum_{r=1}^T \Phi_\mu^{r+1} \leq \tau \mathbb{E}[P^1 - \underline{P}] + \beta_2 \mathbb{E} \|x^1 - x^0\|^2 + T(\tau\alpha_3 + \beta_4) \frac{L_0^2(Q+4)^2}{J^2}. \quad (55)$$

Because u is a uniform random number from $\{0, 1, \dots, T-1\}$ we have

$$\mathbb{E}_u[\Phi_\mu^u] = \frac{1}{T} \sum_{r=1}^T \Phi_\mu^{r+1}. \quad (56)$$

Combining (55) and (56) implies the following

$$\mathbb{E}_u[\Phi_\mu^u] \leq \frac{\tau \mathbb{E}[P^1 - \underline{P}] + \beta_2 \mathbb{E} \|x^1 - x^0\|^2}{T} + (\tau\alpha_3 + \beta_4) \frac{L_0^2(Q+4)^2}{J^2}.$$

Setting

$$\gamma_1 = \tau \mathbb{E}[P^1 - \underline{P}] + \beta_2 \mathbb{E} \|x^1 - x^0\|^2, \quad \gamma_2 = \tau\alpha_3 + \beta_4, \quad (57)$$

we conclude the proof.

References

- [1] M. Hong, M. Razaviyayn, Z.-Q. Luo, and J.-S. Pang, “A unified algorithmic framework for block-structured optimization involving big data,” *IEEE Signal Processing Magazine*, vol. 33, no. 1, pp. 57–77, 2016.
- [2] D. Hajinezhad, M. Hong, T. Zhao, and Z. Wang, “NESTT: A nonconvex primal-dual splitting method for distributed and stochastic optimization,” in *Advances in Neural Information Processing Systems 29*. NIPS.
- [3] E. Bjornson and E. Jorswieck, “Optimal resource allocation in coordinated multi-cell systems,” *Foundations and Trends in Communications and Information Theory*, vol. 9, 2013.
- [4] M. Zavlanos, A. Ribeiro, and George J Pappas, “Network integrity in mobile robotic networks,” *IEEE Transactions on Automatic Control*, vol. 58, no. 1, pp. 3–18, 2013.
- [5] V. Cevher, S. Becker, and M. Schmidt, “Convex optimization for big data: Scalable, randomized, and parallel algorithms for big data analytics,” *IEEE Signal Processing Magazine*, vol. 31, no. 5, pp. 32–43, 2014.
- [6] S. Boyd, N. Parikh, E. Chu, B. Peleato, and J. Eckstein, “Distributed optimization and statistical learning via the alternating direction method of multipliers,” *Foundations and Trends in Machine Learning*, vol. 3, no. 1, pp. 1–122, 2011.
- [7] A. Nedic and A. Ozdaglar, “Distributed subgradient methods for multi-agent optimization,” *IEEE Transactions on Automatic Control*, vol. 54, no. 1, pp. 48–61, 2009.
- [8] A. Nedic, A. Olshevsky, A. Ozdaglar, and J. Tsitsiklis, “On distributed averaging algorithms and quantization effects,” *Automatic Control, IEEE Transactions on*, vol. 54, no. 11, pp. 2506–2517, 2009.
- [9] I. Lobel and A. Ozdaglar, “Distributed subgradient methods for convex optimization over random networks,” *IEEE Transactions on Automatic Control*, vol. 56, no. 6, pp. 1291–1306, 2011.
- [10] A. Nedic and A. Olshevsky, “Distributed optimization over time-varying directed graphs,” *IEEE Transactions on Automatic Control*, vol. 60, no. 3, pp. 601–615, 2015.
- [11] S. Lee and M. M. Zavlanos, “Approximate projections for decentralized optimization with sdp constraints,” in *2016 IEEE 55th Conference on Decision and Control (CDC)*, Dec 2016, pp. 1030–1035.
- [12] M. Gurbuzbalaban, A. Ozdaglar, and P. Parrilo, “On the convergence rate of incremental aggregated gradient algorithms,” *arXiv preprint arXiv:1506.02081*, 2015.
- [13] W. Shi, Q. Ling, G. Wu, and W. Yin, “Extra: An exact first-order algorithm for decentralized consensus optimization,” *SIAM Journal on Optimization*, vol. 25, no. 2, pp. 944–966, 2014.

- [14] N. Chatzipanagiotis, D. Dentcheva, and M.M. Zavlanos, “An augmented lagrangian method for distributed optimization,” *Mathematical Programming*, vol. 152, no. 1-2, pp. 405–434, 2015.
- [15] S. Lee, N. Chatzipanagiotis, and M. M. Zavlanos, “Complexity certification of a distributed augmented lagrangian method,” *IEEE Transactions on Automatic Control*, vol. 63, no. 3, pp. 827–834, March 2018.
- [16] N. Chatzipanagiotis and M. M. Zavlanos, “A distributed algorithm for convex constrained optimization under noise,” *IEEE Transactions on Automatic Control*, vol. 61, no. 9, pp. 2496–2511, 2016.
- [17] G. Tychogiorgos, A. Gkelias, and K. Leung, “A non-convex distributed optimization framework and its application to wireless ad-hoc networks,” *IEEE Transactions on Wireless Communications*, vol. 12, no. 9, pp. 4286–4296, 2013.
- [18] D. Hajinezhad and M. Hong, “Nonconvex alternating direction method of multipliers for distributed sparse principal component analysis,” in *IEEE Global Conference on Signal and Information Processing (GlobalSIP)*. IEEE, 2015.
- [19] Y. Sun, G. Scutari, and D. Palomar, “Distributed nonconvex multiagent optimization over time-varying networks,” in *50th Asilomar Conference on Signals, Systems and Computers*, 2016, pp. 788–794.
- [20] P. Bianchi and J. Jakubowicz, “Convergence of a multi-agent projected stochastic gradient algorithm for non-convex optimization,” *IEEE Transactions on Automatic Control*, vol. 58, no. 2, pp. 391–405, 2013.
- [21] T. Tatarenko and B. Touri, “Non-convex distributed optimization,” *IEEE Transactions on Automatic Control*, vol. 62, no. 8, pp. 3744–3757, Aug 2017.
- [22] P. D. Lorenzo and G. Scutari, “Next: In-network nonconvex optimization,” *IEEE Transactions on Signal and Information Processing over Networks*, vol. 2, no. 2, pp. 120–136, 2016.
- [23] M. Hong, D. Hajinezhad, and M-M. Zhao, “Prox-PDA: The proximal primal-dual algorithm for fast distributed nonconvex optimization and learning over networks,” in *Proceedings of the 34th International Conference on Machine Learning (ICML)*, 2017, vol. 70, pp. 1529–1538.
- [24] N. Chatzipanagiotis and M. M. Zavlanos, “On the convergence of a distributed augmented lagrangian method for nonconvex optimization,” *IEEE Transactions on Automatic Control*, vol. 62, no. 9, pp. 4405–4420, Sept 2017.
- [25] D. Hajinezhad and Q. Shi, “Alternating direction method of multipliers for a class of nonconvex bilinear optimization: convergence analysis and applications,” *Journal of Global Optimization*, pp. 1–28, 2018.

- [26] D. Ataee. Tarzanagh, P. Nazari, and M. R. Peyghami, “A nonmonotone PRP conjugate gradient method for solving square and under-determined systems of equations,” *Computers & Mathematics with Applications*, vol. 73, no. 2, pp. 339–354, 2017.
- [27] M. R. Peyghami and D. Ataee. Tarzanagh, “A relaxed nonmonotone adaptive trust region method for solving unconstrained optimization problems,” *Computational Optimization and Applications*, vol. 61, no. 2, pp. 321–341, 2015.
- [28] J. C. Spall, *Introduction to Stochastic Search and Optimization: Estimation, Simulation, and Control*, Wiley, 2003.
- [29] X. Lian, H. Zhang, C-J. Hsieh, Y. Huang, and J. Liu, “A comprehensive linear speedup analysis for asynchronous stochastic parallel optimization from zeroth-order to first-order,” *arXiv preprint arXiv:1606.00498*, 2016.
- [30] A. Agarwal, O. Dekel, and L. Xiao, “Optimal algorithms for online convex optimization with multi-point bandit feedback,” in *COLT*, 2010, pp. 28–40.
- [31] Y. Nesterov and V. Spokoiny, “Random gradient-free minimization of convex functions,” *Foundations of Computational Mathematics*, pp. 1–40, 2011.
- [32] S. Ghadimi and G. Lan, “Stochastic first-and zeroth-order methods for nonconvex stochastic programming,” *SIAM Journal on Optimization*, vol. 23, no. 4, pp. 2341–2368, 2013.
- [33] J. Duchi, M. Jordan, M. Wainwright, and A. Wibisono, “Optimal rates for zero-order convex optimization: the power of two function evaluations,” *IEEE Transactions on Information Theory*, vol. 61, no. 5, pp. 2788–2806, 2015.
- [34] D. Yuan and D.W.C Ho, “Randomized gradient-free method for multiagent optimization over time-varying networks,” *IEEE Transactions on Neural Networks and Learning Systems*, vol. 26, no. 6, pp. 1342–1347, 2015.
- [35] D. Yuan, S. Xu, and J. Lu, “Gradient-free method for distributed multi-agent optimization via push-sum algorithms,” *International Journal of Robust and Nonlinear Control*, vol. 25, no. 10, pp. 1569–1580, 2015.
- [36] D. Yuan, D. W. Ho, and S. Xu, “Zeroth-order method for distributed optimization with approximate projections,” *IEEE transactions on neural networks and learning systems*, vol. 27, no. 2, pp. 284–294, 2016.
- [37] D. Hajinezhad, M. Hong, and A. Garcia, “Zeroth order nonconvex multi-agent optimization over networks,” *arXiv preprint arXiv:1710.09997*, 2017.
- [38] D. Hajinezhad, M. Hong, and A. Garcia, “Zone: Zeroth order nonconvex multi-agent optimization over networks,” *IEEE Transactions on Automatic Control*, 2019.
- [39] M. E. Yildiz and A. Scaglione, “Coding with side information for rate-constrained consensus,” *IEEE Transactions on Signal Processing*, vol. 56, no. 8, pp. 3753–3764, 2008.

- [40] A. Antoniadis, I. Gijbels, and M. Nikolova, “Penalized likelihood regression for generalized linear models with non-quadratic penalties,” *Annals of the Institute of Statistical Mathematics*, vol. 63, no. 3, pp. 585–615, 2009.