

The cosmological constant and the size of the Causal Universe

Enrique Gaztañaga

*Institute of Space Sciences (ICE, CSIC), 08193 Barcelona, Spain
Institut d'Estudis Espacials de Catalunya (IEEC), 08034 Barcelona, Spain*

(Dated: March 2, 2022)

The cosmological constant Λ is a free parameter in Einstein's equations of gravity. We propose to fix its value with a boundary condition: test particles should be free when outside causal contact, e.g. at infinity. Under this condition, we show that constant vacuum energy does not change cosmic expansion and there can not be cosmic acceleration for an infinitely large and uniform Universe. The observed acceleration requires either a large Universe with evolving Dark Energy (DE) and equation of state $\omega > -1$ or a finite causal boundary (that we call Causal Universe) without DE. The former can't explain why $\Omega_\Lambda \simeq 2.3\Omega_m$ today, something that comes naturally with a finite Causal Universe. This boundary condition, combined with the anomalous lack of correlations observed above 60 degrees in the CMB predicts $\Omega_\Lambda \simeq 0.70$ for a flat universe, with independence of any other measurements. This solution provides new clues and evidence for inflation and removes the need for Dark Energy or Modified Gravity.

I. INTRODUCTION

Measurements of cosmic expansion (see e.g. [1–5]) point to a model with Λ , that we refer to as Λ CDM. For a flat homogeneous metric $ds^2 = a^2(d\eta^2 - d\chi^2)$, the standard expanding equations are:

$$H(a)^2 \equiv \left(\frac{\dot{a}}{a}\right)^2 = \frac{8\pi G}{3}\rho(a) + \Lambda \quad (1)$$

$$\rho(a) = \rho_m a^{-3} + \rho_r a^{-4} + \rho_{vac}$$

and its derivative. Where ρ_m is the pressureless matter density today ($a = 1$), ρ_r corresponds to radiation (with pressure $p_r = \rho_r/3$) and ρ_{vac} represents vacuum energy ($p_{vac} = -\rho_{vac}$). How this equation emerge (or can be valid) in a Universe of finite age and therefore a finite causal size? How can $\rho(a)$ be the same everywhere at a fix a ? We will argue here that this problem relates to cosmic acceleration. Both ρ_{vac} and Λ produce cosmic acceleration but we can only measure the combination:

$$\rho_\Lambda \equiv \rho_{vac} + \frac{\Lambda}{8\pi G}. \quad (2)$$

The measured ρ_Λ is extremely small compared to what we expect for ρ_{vac} . Moreover, $\rho_\Lambda \simeq 2.3\rho_m$, which is a remarkable and puzzling coincidence. Possible solutions are: I) $\Lambda = 0$, II) $\rho_{vac} = 0$ or III) a cancellation between them (for a review see [6–13]). In option I) the observed ρ_Λ originates only from ρ_{vac} or dark energy (DE). But in quantum field theory (QFT) $\rho_{vac} = \infty$ and observables depend only on energy differences. If this is also true for gravity, only Λ contributes to ρ_Λ . This is option II), which includes modified gravity models.

In this paper we take Λ to be a fundamental part of the gravitational interactions. To understand the role of Λ consider first the implications in classical physics. This corresponds to adding a Hooke term, i.e. proportional to distance, to the gravitational acceleration:

$$\vec{g} = -\left(\frac{Gm}{r^3} - \frac{\Lambda}{3}\right)\vec{r} \quad (3)$$

or equivalently a Poisson equation $\nabla^2\phi = 4\pi G\rho - \Lambda$ (see Eq.8). This generalization retains a key property required for gravity, that a spherical mass shell of arbitrary density produces a gravitational field which is identical to a point source of equal mass in its center [14, 15]. Here, in addition, we require that test particles should be free when outside causal contact. If we request $\vec{g} \Rightarrow 0$ for $r \Rightarrow \infty$ in the equation above, we obtain $\Lambda = 0$. The observational fact that $\Lambda > 0$ is therefore indicating that $r \Rightarrow r_\S < \infty$ (in agreement with the finite age of the Universe) and gives $\Lambda = 4\pi G\rho(r < r_\S)$, which is related to the coincidence and horizon problems.

Particles separated by distances larger than the comoving Hubble radius $d_H(t) = c/[a(t)H(t)]$ can't communicate at time t . Distances larger than the horizon

$$\eta(a) = c \int_0^t \frac{dt}{a(t)} = c \int_0^a d\ln(a) d_H(a), \quad (4)$$

have never communicated. This either means that the initial conditions were acausally smooth to start with or that there is a mechanism like inflation [16–19] which inflates causally connected regions outside the Hubble radius. This allows the full observable Universe to originate from a very small causally connected homogeneous patch, which here we call the Causal Universe, χ_\S . During inflation, d_H decreases which freezes out communication on comoving scales larger than the horizon $\chi_\S \simeq \eta(a_i) = d_H(t_i)$ when inflation begins, at $a_i = a(t_i)$. When inflation ends, radiation from reheating makes d_H grow again. When χ_\S re-enters causal contact, we will see that the Universe starts another inflationary epoch so that χ_\S keeps frozen. Thus, causality can only play a role for comoving scales $\chi < \chi_\S$. The Causal Universe χ_\S is therefore fixed in comoving coordinates and is the same for all times, while the horizon η and d_H change with time. Fig.1 illustrates this situation. We conclude that Eq.1 only makes sense for comoving scales $\chi < \chi_\S$.

II. FIXING THE VALUE OF Λ

The symmetries of Einstein's gravitational field equations allow a cosmological constant Λ ([20]):

$$R_{\mu\nu} + \Lambda g_{\mu\nu} = -8\pi G (T_{\mu\nu} - \frac{1}{2}g_{\mu\nu}T^\sigma_\sigma), \quad (5)$$

For a homogeneous and isotropic perfect relativistic fluid with density $\bar{\rho}$ and pressure $\bar{p} \equiv \omega\bar{\rho}$:

$$T_{\mu\nu} = \bar{p}g_{\mu\nu} + (\bar{p} + \bar{\rho})u_\mu u_\nu \quad (6)$$

As explained in the introduction, causality can only be efficient for $\chi < \chi_\S$. How can we implement this condition in $T_{\mu\nu}$? Larger scales can have no effect on the metric, which is equivalent to say that $T_{\mu\nu}$ becomes zero for $\chi > \chi_\S$. This corresponds to:

$$\bar{\rho} = \rho \mathcal{H}(\chi_\S - \chi) \quad (7)$$

where \mathcal{H} is the Heaviside step function and ρ is the mean density inside χ_\S . This does not mean that space is empty for $\chi > \chi_\S$, but just that it can have no effect on the metric as seen by our observer. An observer situated close to the causal boundary of our first observer will find a similar solution, but could measure different values for ρ , p and Λ depending on the initial conditions. The solutions from different acausal regions could be matched [22] which creates a smooth but inhomogeneous background across disconnected regions with an infrared cutoff in the spectrum of homogeneities for $\chi > \chi_\S$.

On scales $\chi < \chi_\S$ we have a homogeneous expanding universe. On larger scales we look for a solution as a perturbation around Minkowski space. In the weak field limit, $g_{00} \simeq -(1+2\phi)$, where ϕ is the Newtonian potential (see eg [20]). The field equations become a covariant generalization of Poisson equation:

$$R_{00} \simeq -\partial^\mu \partial_\mu \phi = -4\pi G (\bar{\rho} + 3\bar{p}) + \Lambda, \quad (8)$$

This is the same as Eq.6.13 in [21], keeping the time derivative term to have a covariant 4D d'Alambert operator. We can use Stokes theorem to estimate the invariant Gauss flux of the 4D acceleration $g_\mu \equiv \partial_\mu \phi$ in a 3D hyper-surface ∂M of a 4D volume M :

$$\Phi = \oint_{\partial M} dx_\mu g^\mu = - \int_M d^4x [4\pi G (\bar{\rho} + 3\bar{p}) - \Lambda], \quad (9)$$

where d^4x and dx_μ are the invariant 4D volume element and normal surface element of ∂M .

A. Causal Boundary condition

We require next that a test particle should be free (i.e. the metric is Minkowski) outside causal contact. Thus




FIG. 1. Comoving Hubble radius $d_H = c/(aH)$ (blue line) as a function of the scale factor a . The Causal Universe χ_\S is identified with the region inside the largest causally connected scale at the beginning of inflation (red dashed line). During inflation d_H decreases so scales $\chi > d_H$ can no longer communicate. After inflation d_H grows again (during HOT=radiation and COLD=matter domination) and χ_\S re-enters d_H at some time a_\S , which creates another inflation.

$g_\mu = 0$ and $\Phi = 0$ for $\chi > \chi_\S$. This fixes Λ :

$$\Phi = 0 \Rightarrow \frac{\Lambda}{8\pi G} = \frac{1}{2M_\S} \int_{M_\S} d^4x (\rho + 3p), \quad (10)$$

where M_\S is the volume inside the lightcone to the surface ∂M_\S and we have used Eq.7. As we approach the boundary ∂M_\S there is no gravitational field and there is no energy associated with it. Because of energy conservation, the Λ term has to be constant and its value is the same at any cosmic time for our arbitrary observer.

B. Vacuum Energy does not gravitate

Inside $\chi < \chi_\S$, we can use Eq.1 with $\rho = \rho_m + \rho_r + \rho_{vac}$ and $p = \rho_r/3 - \rho_{vac}$, so that we can write Eq.10 as:

$$\frac{\Lambda}{8\pi G} = \frac{\rho_m(\S)}{2} + \rho_r(\S) - \rho_{vac} \equiv \rho_\S - \rho_{vac}, \quad (11)$$

where ρ_\S is the matter and radiation contribution in the integral of Eq.10. The values of ρ_m and ρ_r evolve with space-time, so that ρ_\S is the average contribution inside the volume M_\S , while the vacuum density contribution is constant. As pointed out in Eq.2, ρ_{vac} has the same effect in Eq.5 as Λ , and the only observable is:

$$\rho_\Lambda = \frac{\Lambda}{8\pi G} + \rho_{vac} = \rho_\S, \quad (12)$$

where in the last equality we have used our causality condition in Eq.11. So we see how vacuum energy cancels out and can not change the observed value of ρ_Λ , even for $\rho_{vac} = \infty$, as predicted by QFT. If vacuum energy suffers a phase transition or changes in some other way, as is believed to have happened during inflation, then this cancellation will not necessarily happen and ρ_{vac} could contribute to the effective value of ρ_Λ .




FIG. 2. Space-time diagram $ds^2 = a^2(d\eta^2 - d\chi^2)$ for an expanding universe after inflation with two options (labeled §1 and §2). The causal boundary (blue dot) re-enters the Horizon some time that can be earlier (§1) or later (§2) than our present time (red dot). Note how in case §1 we can observe regions outside the Causal Universe at the time when CMB was produce (see section III B).

C. Effective Dark Energy (DE)

The general case considered here is:

$$\begin{aligned} \rho_{DE}(a) &= \rho_{vac} + \rho_{DE} a^{-3(1+\omega)} \\ p_{DE}(a) &= -\rho_{vac} + \omega \rho_{DE} a^{-3(1+\omega)}, \end{aligned} \quad (13)$$

where only one component of DE is evolving. We then have from Eq.10 and Eq.2:

$$\rho_\Lambda = \rho_\S + \rho_{DE} \left[1 + \frac{1+3\omega}{2} \hat{a}_\S^{-3(1+\omega)} \right]. \quad (14)$$

where \hat{a}_\S is some mean value of a in the past light-cone of a_\S in Eq.10. This reduces to $\rho_\Lambda = \rho_\S$ for $\omega = -1$, which indicates that we need a finite χ_\S to explain cosmic acceleration. For $a_\S \Rightarrow \infty$ we have $\rho_\S \Rightarrow 0$ because $\rho_m(a)$ and $\rho_r(a)$ tend to zero as we increase a_\S . The same happens with $\hat{a}_\S^{-3(1+\omega)}$ for $\omega > -1$, so that:

$$\rho_\Lambda = \rho_{DE} \text{ for } a_\S \Rightarrow \infty \text{ \& } \omega > -1. \quad (15)$$

So evolving DE could produce the observed cosmic acceleration in an infinitely large Universe. This solution does not explain why $\rho_\Lambda = \rho_{DE} \simeq 2.3\rho_m$. The original motivation to introduce DE was to understand why the vacuum energy ρ_{vac} can be as small as the measured ρ_Λ [6, 8, 10]. The causal boundary condition shows that ρ_{vac} does not contribute to ρ_Λ , which removes the motivation to have DE. So we will explore here a different way to get $\Omega_\Lambda \simeq 0.70$ without DE, i.e. for $\omega = -1$.

III. THE SIZE OF THE CAUSAL UNIVERSE

We assume in this section that vacuum energy is constant after inflation ($\omega = -1$). In this case Eq.10 gives:

$$\rho_\Lambda = \rho_\S = \frac{1}{2M_\S} \int_{M_\S} d^4x (\rho_m + 2\rho_r). \quad (16)$$

From Eq.4, the horizon after inflation is:

$$\chi(a) = \eta(a) - \eta(a_e) \quad (17)$$

where a_e represents the end of inflation. We then have $\chi_\S = \chi(a_\S) = \eta(a_i)$ where a_\S is the time when the causal boundary enters the horizon after inflation and a_i the beginning of inflation. Fig.1-2 illustrate this. We calculate ρ_\S in Eq.16 by integrating within the light-cone of χ_\S :

$$\rho_\S = \frac{\int_0^{\chi_\S} d\chi \chi^2 a^3 (\rho_m a^{-3} + 2\rho_r a^{-4})}{2 \int_0^{\chi_\S} d\chi \chi^2 a^3}, \quad (18)$$

where $a = a(\chi)$ in Eq.17. For $H(a)$ we use Eq.1 with $\Omega \equiv \rho/\rho_c$, $\rho_c = 3H_0^2/8\pi G$, $\Omega_r = 4.2 \times 10^{-5}$ [1] and flat Universe $\Omega_m = 1 - \Omega_\Lambda - \Omega_r$. We use Eq.18 to solve $\Omega_\Lambda = \Omega_\S$ numerically for $\Omega_\Lambda = 0.69 \pm 0.01$ [1]:

$$\chi_\S = (3.149 \pm 0.004) \frac{c}{H_0} \quad (19)$$

$$a_\S = 0.933 \pm 0.004. \quad (20)$$

This scale factor corresponds to an age:

$$t_\S = (0.887 \pm 0.004) \frac{1}{H_0} \simeq 12.5 \text{ Gyr}, \quad (21)$$

compared to $t_{age} \simeq 0.955/H_0$ today, i.e. about $\Delta t \simeq 0.84$ Gyr into our past. We can't observe this boundary χ_\S today (see Fig.2) but we will be able to observe it in the future and in our past (see section III B).

A. Inflation and the coincidence problem

Eq.16 indicates that when the causal boundary re-enters the Horizon the expansion becomes dominated by ρ_Λ . This is because $\rho_m(a_\S) < \rho_\S = \rho_\Lambda$, as density decreases with the expansion. This results in another inflationary epoch at $a = a_\S$ which keeps the Causal Universe frozen (see Fig.1). We can now recast the coincidence problem (why $\rho_\Lambda \simeq 2.3\rho_m$?) into a new question: why we live at a time which is close to a_\S ? Looking at Fig.1, we can see that the best time to host observers is a time close to a_\S as the Universe is dominated by ρ_m (so there are galaxies) and the Hubble radius is the largest. There is nothing too special about this coincidence.




FIG. 3. Points with error-bars show the two-point correlation function of measured CMB temperature fluctuations in WMAP (points with errorbars) as a function of angular separation (adopted from Fig.12 in [25]). There is no correlations above $\theta > 60$ deg, which corresponds to the causal boundary (vertical dashed line) for $\Omega_\Lambda \simeq 0.69$. The black continuous line shows the Λ CDM prediction for an infinite Universe. The shaded (red) region shows Λ CDM for a finite universe.

The reason why $\chi_\S \sim 3c/H_0$ and not some other value could reside in the details of inflation: when inflation begins a_i and ends a_e (see Fig.1). This recasts the coincidence problem into an opportunity to better understand inflation and the origin of homogeneity. We propose to identify $\chi_\S = \eta(a_i)$ with the comoving horizon before inflation begins at time t_i , $H_i = H(t_i)$ or $a_i = a(t_i)$:

$$a_i H_i = c \chi_\S^{-1} \simeq (0.321 \pm 0.004) H_0 \quad (22)$$

The Hubble rate during inflation H_I is proportional to the energy of inflation. During reheating this energy is converted into radiation: $H_I^2 \simeq \Omega_r H_0^2 a_e^{-4}$, with $a_e \equiv e^N a_i$. We can combine with Eq.22 to find:

$$a_i \chi_\S = \frac{H_i}{H_I} e^{-2N} \Omega_r^{1/2} (\chi_\S^2 H_0 / c) \simeq 4 \times 10^8 l_{\text{Planck}} \quad (23)$$

where for the second equality we have used the canonical value of $N \simeq 60$ and $H_i \simeq H_I$, which also yields $a_i \simeq 1.56 \times 10^{-53}$ and $H_I \sim 10^{10}$ GeV. The condition $a_i \chi_\S > l_{\text{Planck}}$ requires $N < 70$, close to the value found in [28].

B. Implications for CMB

The (look-back) comoving distance to the surface of last scattering $a_* \simeq 9.2 \times 10^{-4}$ [1] is $\chi_{\text{CMB}} = \eta(1) - \eta(a_*) \simeq 3.145 \frac{c}{H_0}$. This is shown as the horizontal dashed line in the bottom of Fig.2. This is just slightly smaller than our estimate of the scale when the causal boundary re-enters the Horizon. Thus, we would expect to see no correlations in the CMB on angular scales $\theta > \theta_\S$, where:

$$\theta_\S \equiv 2 \arcsin \frac{\chi_\S/2}{\chi_{\text{CMB}}} \simeq 60.1 \text{ deg.} \quad (24)$$

The lack of structure seen in the CMB on these large scales is one of the well known anomalies in the CMB data (e.g. see [23, 24] and references therein). Fig.3 (from [25]) shows a comparison of the measured CMB temperature correlations (points with error-bars) with the Λ CDM prediction for an infinite Universe (continuous line). There is a very clear discrepancy, which [23] estimates to happen in only 0.025 per cent of the realizations of the infinite Λ CDM model. If we suppress the large scale modes above $\simeq 60$ deg in the Λ CDM simulations, the agreement is much better (shaded red area in Fig.3).

We can also predict Ω_Λ from the lack of CMB correlations. From Fig.3 we estimate $\theta_\S = 62.2 \pm 4.0$ deg. to find (using Eq.24 and Eq.18):

$$\theta_\S = 62.2 \pm 4.0 \text{ deg.} \Rightarrow \Omega_\Lambda = 0.63 \pm 0.11 \quad (25)$$

Note that the above estimate does not take into account the foreground ISW and lensing effects [26, 27], which will typically reduce θ_\S slightly. The value most used in the literature, $\theta_\S \simeq 60$ deg., corresponds to $\Omega_\Lambda \simeq 0.70$.

Note that there are temperature differences on scales larger θ_\S , but they are not correlated, as expected in causality disconnected regions. Nearby regions are connected which creates a smooth background across disconnected regions.

IV. DISCUSSION AND CONCLUSIONS

Λ CDM in Eq.1 assumes that ρ is constant everywhere at a fixed comoving time. This requires acausal initial conditions [17] unless there is inflation, where a tiny homogeneous and causally connected patch, the Causal Universe χ_\S , was inflated to be very large today. Regions larger than χ_\S are out of causal contact. Here we require that test particles become free as we approach χ_\S . This leads to Eq.10, which is the main result in this paper.

If we ignore the vacuum for now, this condition requires: $\Lambda = 8\pi G \rho_\S$, where ρ_\S is the matter and radiation inside χ_\S (Eq.16). For an infinite Universe ($a_\S \rightarrow \infty$) we have $\rho_\S \Rightarrow 0$ which requires $\Lambda \Rightarrow 0$. This is also what we find in classical gravity with a Λ term, because Hooke's term diverges at infinity (see Eq.3). So

the fact that $\rho_\Lambda \neq 0$ could indicate that χ_\S is not infinite. Adding vacuum ρ_{vac} does not change this as we have that $\Lambda = 8\pi G\rho_\S - \rho_\S$ so that $\rho_\Lambda = \Lambda/8\pi G + \rho_{\text{vac}} = \rho_\S$ turns out to be independent of ρ_{vac} . Thus, whether the Causal Universe is finite or not, ρ_{vac} can not gravitate.

For constant vacuum ($\omega = -1$), we find $\chi_\S \simeq 3.15c/H_0$ for $\Omega_\S = \Omega_\Lambda \simeq 0.69$. We can also estimate χ_\S as $c/(a_i H_i)$ when inflation begins, see Eq.22. After inflation χ_\S freezes out until it re-enters causality at $a_\S \simeq 0.93$, close to now ($a = 1$). This starts a new inflation (as $\rho_\Lambda = \rho_\S > \rho_m$) which keeps the causal boundary frozen. Thus a finite χ_\S explains why $\rho_\Lambda \simeq 2.3\rho_m$ and looking at Fig.1, we argued that a_\S is the best time for observers like us to exist.

For $\omega = -1$, the measured value of $\Omega_\Lambda \simeq 0.70$ predicts that CMB temperature should not be correlated above $\theta > \theta_\S \simeq 60$ deg, a prediction that matches observations (see Fig.3). One can also reverse the argument and use the lack of CMB correlations above $\theta_\S \simeq 60$ deg, to predict the size of $\chi_\S \simeq 2\chi_{\text{CMB}} \sin(\theta_\S/2)$. Together with condition $\rho_\S = \rho_\Lambda$, this provides a prediction of $\Omega_\Lambda \simeq 0.70$, which is independent of other measurements for Ω_Λ .

If DE exists, we have shown that only the evolving

component of DE is observable. A universe with $\omega < -1$ violates our causality condition. In the limit of an infinite Universe with $\omega > -1$, we find that $\rho_\Lambda = \rho_{\text{DE}}$ (see Eq.15). But DE gives no clue as to why $\rho_{\text{DE}} \simeq 2.3\rho_m$ and can not explain the lack of CMB correlations for $\theta > 60$ deg. We apply Occam's razor to argue that there is no need for DE: measurements of cosmic acceleration and CMB can be explained by the finite size of our Causal Universe, as predicted by inflation.

ACKNOWLEDGMENTS

I want to thank A.Alarcon, J.Barrow, C.Baugh, R.Brandenberger, G.Bernstein, M.Bruni, S.Dodelson, E.Elizalde, J.Frieman, M.Gatti, L.Hui, D.Huterer, A.Liddle, P.J.E. Peebles, R.Scoccimarro and S.Weinberg for their feedback. This work has been supported by MINECO grants CSD2007-00060 and AYA2015-71825, LACEGAL Marie Skłodowska-Curie grant No 734374 with ERDF funds from the European Union Horizon 2020 Programme. IEEC is partially funded by the CERCA program of the Generalitat de Catalunya.

-
- [1] Planck Collaboration, 2018, arXiv:1807.06209
 - [2] DES Collaboration, 2018, arXiv:1811.02375
 - [3] Tutusaus, I., et al 2017, A&A, 602, A73
 - [4] Gaztañaga, E., Miquel, R., & Sánchez, E. 2009, Physical Review Letters, 103, 091302
 - [5] Gaztañaga, E., Manera, M., & Multamäki, T. 2006, MNRAS, 365, 171
 - [6] Weinberg, S. 1989, Reviews of Modern Physics, 61, 1
 - [7] Elizalde, E., & Gaztañaga, E. 1990, Phys.Let.B, 234, 265
 - [8] Huterer D., Turner M. S., 1999, PhRvD, 60, 081301
 - [9] Carroll, S. M., Press, W. H., & Turner, E. L. 1992, ARAA, 30, 499
 - [10] Martin, J. 2012, arXiv:1205.3365
 - [11] Gaztañaga, E., & Lobo, J. A. 2001, ApJ, 548, 47
 - [12] Gaztañaga, E., et al 2002, Phys. Rev. D, 65, 023506
 - [13] Lue, A., Scoccimarro, R., & Starkman, G. D. 2004, Phys.Rev.D, 69, 124015
 - [14] Wilkins, D. 1986, American Journal of Physics, 54, 726
 - [15] Calder, L., & Lahav, O. 2008, Astron. & Geoph., 49, 1.13
 - [16] Liddle, A. R. 1999, astro-ph/9901124
 - [17] Brandenberger, R. 2017, Int.J.Mod.Phys.D, 26, 1740002-126
 - [18] Martin, J. 2019, arXiv:1902.05286
 - [19] Dodelson, S. 2003, Modern cosmology, Academic Press. ISBN 0-12-219141-2, 2003, XIII + 440 p.
 - [20] Weinberg S., 1972, Gravitation and Cosmology, ISBN 0-471-92567-5. Wiley-VCH, 688
 - [21] Peebles, P. J. E. 1980, The Large-Scale Structure of the Universe, Princeton University Press.
 - [22] Sanghai, V. A. A., & Clifton, T. 2015, Phys. Rev. D, 91, 103532
 - [23] Copi, C. J., Huterer, D., Schwarz, D. J., & Starkman, G. D. 2009, MNRAS, 399, 295
 - [24] Schwarz, D. J., Copi, C. J., Huterer, D., & Starkman, G. D. 2016, Classical and Quantum Gravity, 33, 184001
 - [25] Gaztañaga, E., Wagg, J., Multamäki, T., Montaña, A., & Hughes, D. H. 2003, MNRAS, 346, 47
 - [26] Fosalba, P., Gaztañaga, E., & Castander, F. J. 2003, ApJL, 597, L89
 - [27] Das, S., & Souradeep, T. 2014, JCAP, 2, 002
 - [28] Dodelson, S., & Hui, L. 2003, Phys Rev Let, 91, 131301