

A SCHWARZ LEMMA FOR TWO FAMILIES OF DOMAINS AND COMPLEX GEOMETRY

SOURAV PAL AND SAMRIDDHO ROY

ABSTRACT. We make sharp estimates to obtain a Schwarz type lemma for the symmetrized polydisc \mathbb{G}_n and for the extended symmetrized polydisc $\tilde{\mathbb{G}}_n$. We explicitly construct an interpolating function under certain condition. To do so, we followed the methods described in [35]. Also we find a few geometric interplay between the members of the family $\tilde{\mathbb{G}}_n$ and its closure $\tilde{\Gamma}_n$.

1. INTRODUCTION

This article is a sequel of [28]. Being motivated by the inspiring works due to Bharali, Costara, Edigarian, Kosinski, Nikolov, Zwonek [8, 14, 18, 19, 20, 21, 22, 24, 33] and few others (see references there in), we descend one more step into the depth of studying complex geometry and function theory of the symmetrized n -disk \mathbb{G}_n for $n \geq 3$. The *symmetrized n -disk* \mathbb{G}_n or simply the *symmetrized polydisc*, which consists of symmetric polynomials, is defined by

$$\mathbb{G}_n = \left\{ \left(\sum_{1 \leq i \leq n} z_i, \sum_{1 \leq i < j \leq n} z_i z_j, \dots, \prod_{i=1}^n z_i \right) : |z_i| < 1, i = 1, \dots, n \right\}.$$

This domain arises in the famous μ -synthesis problem, which is a part of the theory of robust control of systems comprising of interconnected electronic devices whose outputs are linearly dependent on the inputs. Given a *structure* E , which is a linear subspace of $\mathcal{M}_{m \times n}(\mathbb{C})$, the space of all $m \times n$ matrices, the functional

$$\begin{aligned} &\mu_E(B) \\ &:= (\inf \{ \|X\| : X \in E \text{ and } (I - BX) \text{ is singular} \})^{-1}, \quad B \in \mathcal{M}_{m \times n}(\mathbb{C}), \end{aligned}$$

2010 *Mathematics Subject Classification.* 30C80, 32A10, 32A60, 32E20, 32E30.

Key words and phrases. Symmetrized polydisc, Extended symmetrized polydisc, Complex geometry, Schwarz lemma, Interpolating function.

The first author is supported by the Seed Grant of IIT Bombay, the CPDA and the INSPIRE Faculty Award (Award No. DST/INSPIRE/04/2014/001462) of DST, India. The second author is supported by a Ph.D fellowship from the University Grand Commission of India.

is called a *structured singular value*. If $m = n$ and if E is the space of all scalar multiples of the identity matrix I , then $\mu_E(B)$ is equal to the spectral radius $r(B)$. Also if $E = \mathcal{M}_{m \times n}(\mathbb{C})$, then $\mu_E(B)$ is precisely the operator norm $\|B\|$. Naturally if E is any linear subspace of $\mathcal{M}_n(\mathbb{C})$ that contains the identity matrix, then $r(B) \leq \mu_E(A) \leq \|B\|$. For the control-theory motivations behind μ_E , we refer to the pioneering work of Doyle [15]. The μ -synthesis problem aims to find an analytic function f from the open unit disk \mathbb{D} of the complex plane \mathbb{C} to $\mathcal{M}_{m \times n}(\mathbb{C})$ subject to a finite number of interpolation conditions such that $\mu_E(f(\lambda)) < 1$, for all $\lambda \in \mathbb{D}$. If $E = \{\lambda I : \lambda \in \mathbb{C}\} \subseteq \mathcal{M}_n(\mathbb{C})$, then $\mu_E(B) = r(B) < 1$ if and only if $\pi_n(\nu_1, \dots, \nu_n) \in \mathbb{G}_n$ (see [14]); here ν_1, \dots, ν_n are eigenvalues of B and π_n is the symmetrization map defined on \mathbb{C}^n by

$$\pi_n(z) = (s_1(z), \dots, s_{n-1}(z), p(z)), \quad z = (z_1, \dots, z_n),$$

where

$$s_i(z) = \sum_{1 \leq k_1 \leq k_2 \leq \dots \leq k_i \leq n} z_{k_1} \cdots z_{k_i} \quad \text{and} \quad p(z) = \prod_{i=1}^n z_i.$$

It is merely mentioned that π_n is a proper holomorphic map and $\pi_n(\mathbb{D}^n) = \mathbb{G}_n$, where \mathbb{D}^n is the open polydisc defined by

$$\mathbb{D}^n = \{(z_1, \dots, z_n) : |z_i| < 1, i = 1, \dots, n\}.$$

The *closed symmetrized polydisc* Γ_n , which is the closure of \mathbb{G}_n , is given by

$$\begin{aligned} \Gamma_n &:= \left\{ \left(\sum_{1 \leq i \leq n} z_i, \sum_{1 \leq i < j \leq n} z_i z_j, \dots, \prod_{i=1}^n z_i \right) : |z_i| \leq 1, i = 1, \dots, n \right\} \\ &= \pi_n(\overline{\mathbb{D}^n}). \end{aligned}$$

The set Γ_n is polynomially convex but not convex (see [18]). It is evident from the definition that $\mathbb{G}_1 = \mathbb{D}$ and below we provide an explicit form of \mathbb{G}_2 and \mathbb{G}_3 for the convenience of the readers.

$$\mathbb{G}_2 = \{(z_1 + z_2, z_1 z_2) : z_1, z_2 \in \mathbb{D}\},$$

$$\mathbb{G}_3 = \{(z_1 + z_2 + z_3, z_1 z_2 + z_2 z_3 + z_3 z_1, z_1 z_2 z_3) : z_1, z_2, z_3 \in \mathbb{D}\}.$$

The symmetrized polydisc has attracted considerable attentions in past two decades because of its rich function theory [2, 3, 7, 14, 19, 23, 32], complex geometry [13, 18, 20, 21, 22], associated operator theory [4, 9, 10, 12, 25, 26, 29]. An interested reader can also see the articles referred there.

The classical Schwarz lemma in one variable is stated in the following way.

Theorem 1.1. *Let f be an analytic function on \mathbb{D} such that $|f(z)| \leq 1$, for all $z \in \mathbb{D}$ and $f(0) = 0$. Then*

- (a) $|f(z)| \leq |z|$, for all $z \in \mathbb{D}$,
- (b) $|f'(0)| \leq 1$.

Moreover, if $|f'(0)| = 1$ or if $|f(z_0)| = |z_0|$ for some $z_0 \neq 0$, then there is a constant c such that $|c| = 1$ and $f(w) = cw$ for all $w \in \mathbb{D}$.

In [28], we obtained an analogue of the first part of Theorem 1.1 for the symmetrized polydisc. The main aim of this article is to continue the same program to find an analogous part-(b) for \mathbb{G}_n of the classical Schwarz lemma.

To study the complex geometry of \mathbb{G}_n (and Γ_n) more deeply and for proving a Schwarz lemma for \mathbb{G}_n , we introduced a new family of domains in [28], which we named *extended symmetrized polydisc* and defined as

$$\tilde{\mathbb{G}}_n := \left\{ (y_1, \dots, y_{n-1}, q) \in \mathbb{C}^n : q \in \mathbb{D}, y_j = \beta_j + \bar{\beta}_{n-j}q, \beta_j \in \mathbb{C} \text{ and } |\beta_j| + |\beta_{n-j}| < \binom{n}{j}, j = 1, \dots, n-1 \right\}.$$

We called the closure of $\tilde{\mathbb{G}}_n$, the *closed extended symmetrized polydisc* and denoted it by $\tilde{\Gamma}_n$. We proved in [28] that

$$\tilde{\Gamma}_n := \left\{ (y_1, \dots, y_{n-1}, q) \in \mathbb{C}^n : q \in \mathbb{D}, y_j = \beta_j + \bar{\beta}_{n-j}q, \beta_j \in \mathbb{C} \text{ and } |\beta_j| + |\beta_{n-j}| \leq \binom{n}{j}, j = 1, \dots, n-1 \right\}.$$

The purpose of introducing the family $\tilde{\mathbb{G}}_n$ was to make a few sharp estimates which provides a Schwarz lemma for \mathbb{G}_n , [28]. Also we obtained a variety of characterizations for the points in \mathbb{G}_n and Γ_n via a similar set of characterizations for $\tilde{\mathbb{G}}_n$ and $\tilde{\Gamma}_n$ respectively, [28].

In [14], Costara showed that

$$\mathbb{G}_n = \left\{ (s_1, \dots, s_{n-1}, p) : p \in \mathbb{D}, s_j = \beta_j + \bar{\beta}_{n-j}p \text{ \& } (\beta_1, \dots, \beta_{n-1}) \in \mathbb{G}_{n-1} \right\}$$

and

$$\Gamma_n = \{(s_1, \dots, s_{n-1}, p) : p \in \overline{\mathbb{D}}, s_j = \beta_j + \bar{\beta}_{n-j}p \text{ \& } (\beta_1, \dots, \beta_{n-1}) \in \Gamma_{n-1}\}.$$

It is obvious that if $(\beta_1, \dots, \beta_{n-1}) \in \mathbb{G}_{n-1}$, then $|\beta_j| + |\beta_{n-j}| < \binom{n}{j}$. Therefore, it follows that $\mathbb{G}_n \subseteq \widetilde{\mathbb{G}}_n$. In fact, $\mathbb{G}_2 = \widetilde{\mathbb{G}}_2$ but $\mathbb{G}_n \subsetneq \widetilde{\mathbb{G}}_n$ for $n \geq 3$ (see [28], Lemma 3.0.2).

We introduced $n - 1$ fractional linear transformations $\Phi_1, \dots, \Phi_{n-1}$ and with their help we made some sharp estimates to find necessary conditions for the existence of an interpolating function from \mathbb{D} to \mathbb{G}_n and since $\mathbb{G}_n \subseteq \widetilde{\mathbb{G}}_n$, the estimates became necessary for a Schwarz lemma for \mathbb{G}_n (see [28]). Since the maximum modulus of each co-ordinate of a point in $\widetilde{\mathbb{G}}_n$ does not exceed that of a point in \mathbb{G}_n , these estimates are sharp for \mathbb{G}_n too. Moreover, the functions $\Phi_1, \dots, \Phi_{n-1}$ are specially designed for $\widetilde{\mathbb{G}}_n$ and they characterize the points in $\widetilde{\mathbb{G}}_n$ and $\widetilde{\Gamma}_n$.

In this article, we first prove an analogue of part-(b) of Theorem 1.1 for $\widetilde{\mathbb{G}}_n$, which is Theorem 3.1 and it is one of the main results of this paper. As a consequence the desired Schwarz lemma for \mathbb{G}_n (Theorem 3.2) follows. We also show in Theorem 3.1 that under certain condition, the achieved estimates are sufficient for the existence of an interpolating function from \mathbb{D} to $\widetilde{\mathbb{G}}_n$. In Section 4, we explicitly construct such an interpolating function. Section 5 deals with some geometric interplay between the members of $\widetilde{\mathbb{G}}_n$ and $\widetilde{\Gamma}_n$. In Section 2, we accumulate few results from the literature which are used in the subsequent sections.

Note. The main idea and applied techniques to the results of Sections 3 and 4 of this article are borrowed from the paper [35], where analogous results for the tetrablock \mathbb{E} are achieved. The primary reason for which the techniques of [35] are applicable here is that $\widetilde{\mathbb{G}}_3$ is linearly isomorphic to \mathbb{E} .

2. BACKGROUND MATERIALS AND PREPARATORY RESULTS

We begin with a set of $(n-1)$ fractional linear transformations $\Phi_1, \dots, \Phi_{n-1}$ which we introduced in [28] to characterize the points in the extended symmetrized polydisc $\widetilde{\mathbb{G}}_n$.

Definition 2.1. For $z \in \mathbb{C}$, $y = (y_1, \dots, y_{n-1}, q) \in \mathbb{C}^n$ and for any $j \in \{1, \dots, n-1\}$, let us define

$$\Phi_j(z, y) = \begin{cases} \frac{\binom{n}{j}qz - y_j}{y_{n-j}z - \binom{n}{j}} & \text{if } y_{n-j}z \neq \binom{n}{j} \text{ and } y_j y_{n-j} \neq \binom{n}{j}^2 q \\ \frac{y_j}{\binom{n}{j}} & \text{if } y_j y_{n-j} = \binom{n}{j}^2 q. \end{cases} \quad (1)$$

It was shown in [28] that if $|y_{n-j}| < \binom{n}{j}$, then

$$\|\Phi_j(\cdot, y)\|_{H^\infty} = \frac{\binom{n}{j} |y_j - \bar{y}_{n-j}q| + |y_j y_{n-j} - \binom{n}{j}^2 q|}{\binom{n}{j}^2 - |y_{n-j}|^2}. \quad (2)$$

Clearly $|y_{n-j}| < \binom{n}{j}$, for any point $y \in \tilde{\mathbb{G}}_n$. Thus equation (2) holds for any $y \in \tilde{\mathbb{G}}_n$.

For $n \geq 3$, we introduced in [28] the following subset \mathcal{J}_n of $\tilde{\mathbb{G}}_n$ as follows:

$$\mathcal{J}_n = \begin{cases} \mathcal{J}_n^{\text{odd}} & \text{if } n \text{ is odd} \\ \mathcal{J}_n^{\text{even}} & \text{if } n \text{ is even,} \end{cases} \quad (3)$$

where

$$\mathcal{J}_n^{\text{odd}} = \left\{ (y_1, \dots, y_{n-1}, y_n) \in \tilde{\mathbb{G}}_n : y_j = \frac{\binom{n}{j}}{n} y_1, y_{n-j} = \frac{\binom{n}{j}}{n} y_{n-1}, \right. \\ \left. \text{for } j = 2, \dots, \left\lfloor \frac{n}{2} \right\rfloor \right\}$$

and

$$\mathcal{J}_n^{\text{even}} = \left\{ (y_1, \dots, y_{n-1}, y_n) \in \tilde{\mathbb{G}}_n : y_{\lfloor \frac{n}{2} \rfloor} = \binom{n}{\lfloor \frac{n}{2} \rfloor} \frac{y_1 + y_{n-1}}{2n}, y_j = \frac{\binom{n}{j} y_1}{n}, \right. \\ \left. y_{n-j} = \frac{\binom{n}{j} y_{n-1}}{n}, \text{ for } j = 2, \dots, \left\lfloor \frac{n}{2} \right\rfloor - 1 \right\}.$$

The following theorem provides a few characterizations for the points in $\tilde{\mathbb{G}}_n$.

Theorem 2.2 ([28], Theorem 3.1.4). *For a point $y = (y_1, \dots, y_{n-1}, q) \in \mathbb{C}^n$, the following are equivalent:*

- (1) $y \in \tilde{\mathbb{G}}_n$;

(2) $\binom{n}{j} - y_j z - y_{n-j} w + \binom{n}{j} q z w \neq 0$, for all $z, w \in \overline{\mathbb{D}}$ and for all $j = 1, \dots, \lfloor \frac{n}{2} \rfloor$;

(3) for all $j = 1, \dots, \lfloor \frac{n}{2} \rfloor$ either

$$\binom{n}{j} |y_j - \bar{y}_{n-j} q| + \left| y_j y_{n-j} - \binom{n}{j}^2 q \right| < \binom{n}{j}^2 - |y_{n-j}|^2$$

or

$$\binom{n}{j} |y_{n-j} - \bar{y}_j q| + \left| y_j y_{n-j} - \binom{n}{j}^2 q \right| < \binom{n}{j}^2 - |y_j|^2;$$

(4) $|y_{n-j} - \bar{y}_j q| + |y_j - \bar{y}_{n-j} q| < \binom{n}{j} (1 - |q|^2)$ for all $j = 1, \dots, \lfloor \frac{n}{2} \rfloor$;

(5) there exist $\lfloor \frac{n}{2} \rfloor$ number of 2×2 matrices $B_1, \dots, B_{\lfloor \frac{n}{2} \rfloor}$ such that $\|B_j\| < 1$, $y_j = \binom{n}{j} [B_j]_{11}$, $y_{n-j} = \binom{n}{j} [B_j]_{22}$ for all $j = 1, \dots, \lfloor \frac{n}{2} \rfloor$ and

$$\det B_1 = \det B_2 = \dots = \det B_{\lfloor \frac{n}{2} \rfloor} = q.$$

In [28], we obtained several equivalent necessary conditions which established a Schwarz type lemma for $\tilde{\mathbb{G}}_n$. Here we mention a few of them.

Theorem 2.3 ([28], Theorem 5.2.1). *Let $\lambda_0 \in \mathbb{D} \setminus \{0\}$ and let $y^0 = (y_1^0, \dots, y_{n-1}^0, q^0) \in \tilde{\mathbb{G}}_n$. Then in the set of following conditions, (1) implies (2) and (3).*

(1) *There exists an analytic function $\psi : \mathbb{D} \rightarrow \tilde{\mathbb{G}}_n$ such that $\psi(0) = (0, \dots, 0)$ and $\psi(\lambda_0) = y^0$.*

(2)

$$\max_{1 \leq j \leq n-1} \{ \|\Phi_j(\cdot, y^0)\|_{H^\infty} \} \leq |\lambda_0|.$$

(3) *There exist $\lfloor \frac{n}{2} \rfloor$ number of functions $F_1, F_2, \dots, F_{\lfloor \frac{n}{2} \rfloor}$ in the Schur class such that $F_j(0) = \begin{bmatrix} 0 & * \\ 0 & 0 \end{bmatrix}$, and $F_j(\lambda_0) = B_j$, for $j = 1, \dots, \lfloor \frac{n}{2} \rfloor$, where $\det B_1 = \dots = \det B_{\lfloor \frac{n}{2} \rfloor} = q^0$, $y_j^0 = \binom{n}{j} [B_j]_{11}$ and $y_{n-j}^0 = \binom{n}{j} [B_j]_{22}$.*

Furthermore, if $y^0 \in \mathcal{J}_n$ then all the conditions (1) – (3) are equivalent.

The following result is known as Parrott's Theorem and it will be used in sequel. One can see Theorem 12.22 in [34] for a proof to this result.

Theorem 2.4 ([30], Theorem 1). *Let H_i, K_i are Hilbert spaces for $i = 1, 2$, and let*

$$\begin{bmatrix} Q \\ S \end{bmatrix} : H_2 \longrightarrow K_1 \oplus K_2, \quad \begin{bmatrix} R & S \end{bmatrix} : H_1 \oplus H_2 \longrightarrow K_2$$

be contractions. Then there exists $P \in \mathcal{L}(H_1, K_1)$ such that

$$\begin{bmatrix} P & Q \\ R & S \end{bmatrix} : H_1 \oplus H_2 \longrightarrow K_1 \oplus K_2$$

is a contraction.

3. A SCHWARZ LEMMA FOR $\tilde{\mathbb{G}}_n$ AND \mathbb{G}_n

Let B_1, \dots, B_k be 2×2 contractive matrices such that $\det B_1 = \det B_2 = \dots = \det B_k$. We define two functions π_{2k+1} and π_{2k} in the following way:

$$\begin{aligned} & \pi_{2k+1}(B_1, \dots, B_k) \\ &= \left(\binom{n}{1} [B_1]_{11}, \dots, \binom{n}{k} [B_k]_{11}, \binom{n}{k} [B_k]_{22}, \dots, \binom{n}{1} [B_1]_{22}, \det B_1 \right) \end{aligned}$$

and

$$\begin{aligned} & \pi_{2k}(B_1, \dots, B_k) \\ &= \left(\binom{n}{1} [B_1]_{11}, \dots, \binom{n}{k-1} [B_{k-1}]_{11}, \binom{n}{k} \frac{([B_k]_{11} + [B_k]_{22})}{2}, \right. \\ & \quad \left. \binom{n}{k-1} [B_{k-1}]_{22}, \dots, \binom{n}{1} [B_1]_{22}, \det B_1 \right). \end{aligned}$$

Then by Theorem 2.2, we have

$$\pi_{2k}(B_1, \dots, B_k) \in \tilde{\mathbb{G}}_{2k} \quad \text{and} \quad \pi_{2k+1}(B_1, \dots, B_k) \in \tilde{\mathbb{G}}_{2k+1}.$$

For $n \geq 3$, let \mathcal{K}_n be the following subset of \mathbb{C}^n :

$$\mathcal{K}_n = \begin{cases} \mathcal{K}_n^{odd} & \text{if } n \text{ is odd} \\ \mathcal{K}_n^{even} & \text{if } n \text{ is even,} \end{cases}$$

where

$$\begin{aligned} \mathcal{K}_n^{odd} = \left\{ (y_1, \dots, y_{n-1}, y_n) \in \mathbb{C}^n : y_j = \frac{\binom{n}{j}}{n} y_1, y_{n-j} = \frac{\binom{n}{j}}{n} y_{n-1}, \right. \\ \left. \text{for } j = 2, \dots, \left\lfloor \frac{n}{2} \right\rfloor \text{ \& } \max \left\{ \frac{|y_1|}{n}, \frac{|y_{n-1}|}{n} \right\} + |y_n| \leq 1 \right\} \end{aligned}$$

and

$$\mathcal{K}_n^{\text{even}} = \left\{ (y_1, \dots, y_{n-1}, y_n) \in \mathbb{C}^n : y_{[\frac{n}{2}]} = \binom{n}{[\frac{n}{2}]} \frac{y_1 + y_{n-1}}{2n}, y_j = \frac{\binom{n}{j} y_1}{n}, \right. \\ \left. y_{n-j} = \frac{\binom{n}{j} y_{n-1}}{n}, \text{ for } j = 2, \dots, \left[\frac{n}{2}\right] - 1 \text{ \& } \max\left\{\frac{|y_1|}{n}, \frac{|y_{n-1}|}{n}\right\} + |y_n| \leq 1 \right\}.$$

For any $Z \in \mathbb{C}^{2 \times 2}$ with $\|Z\| < 1$, let $D_Z = (1 - Z^*Z)^{\frac{1}{2}}$. We denote the unit ball of $\mathbb{C}^{2 \times 2}$ by $R_I(2, 2)$. Consider the following function:

$$\mathcal{M}_Z(X) = -Z + D_Z^* X (1 - Z^* X)^{-1} D_Z \quad \text{for } X \in R_I(2, 2).$$

The function \mathcal{M}_Z is a matrix Möbius transformation that maps Z to 0. The transformation \mathcal{M}_Z is an automorphism of $R_I(2, 2)$, and $(\mathcal{M}_Z)^{-1} = \mathcal{M}_{-Z}$.

We now present a Schwarz type lemma for $\widetilde{\mathbb{G}}_n$.

Theorem 3.1. *Let $x = (x_1, \dots, x_n) \in \mathbb{C}^n$ and there exists an analytic map $\psi : \mathbb{D} \longrightarrow \widetilde{\mathbb{G}}_n$ such that $\psi(0) = (0, \dots, 0)$ and $\psi'(0) = x$. Then*

$$\max_{1 \leq j \leq n-1} \left\{ \frac{|x_j|}{\binom{n}{j}} \right\} + |x_n| \leq 1. \quad (4)$$

The converse holds if $x \in \mathcal{K}_n$.

Proof. We already know that $\widetilde{\mathbb{G}}_2 = \mathbb{G}_2$ and this theorem was proved by Agler and Young for \mathbb{G}_2 (see Theorem 1.1 in [5]). For this reason we shall consider $n \geq 3$ when proving the converse part of this theorem.

Let $\psi : \mathbb{D} \longrightarrow \widetilde{\mathbb{G}}_n$ is an analytic map such that $\psi(0) = (0, \dots, 0)$ and $\psi'(0) = x$. Write $\psi = (\psi_1, \dots, \psi_n)$. Then, by Theorem 2.3, for each $\lambda \in \mathbb{D} \setminus \{0\}$ we have

$$\max_{1 \leq j \leq n-1} \{ \|\Phi_j(\cdot, \psi(\lambda))\|_{H^\infty} \} \leq |\lambda|,$$

which is same as saying

$$\max_{1 \leq j \leq n-1} \left\{ \frac{\binom{n}{j} |\psi_j(\lambda) - \bar{\psi}_{n-j}(\lambda) \psi_n(\lambda)| + |\psi_j(\lambda) \psi_{n-j}(\lambda) - \binom{n}{j}^2 \psi_n(\lambda)|}{\binom{n}{j}^2 - |\psi_{n-j}(\lambda)|^2} \right\} \\ \leq |\lambda|, \quad (5)$$

for each $\lambda \in \mathbb{D} \setminus \{0\}$. Note that, using L-Hospital's rule and the fact that $\psi(0) = (\psi_1(0), \dots, \psi_n(0)) = (0, \dots, 0)$, we have

$$\begin{aligned} & \lim_{\lambda \rightarrow 0} \frac{\binom{n}{j} |\psi_j(\lambda) - \bar{\psi}_{n-j}(\lambda) \psi_n(\lambda)|}{|\lambda| \left(\binom{n}{j}^2 - |\psi_{n-j}(\lambda)|^2 \right)} \\ &= \binom{n}{j} \left| \lim_{\lambda \rightarrow 0} \frac{\psi_j(\lambda) - \bar{\psi}_{n-j}(\lambda) \psi_n(\lambda)}{\lambda} \right| \lim_{\lambda \rightarrow 0} \frac{1}{\left(\binom{n}{j}^2 - |\psi_{n-j}(\lambda)|^2 \right)} = \frac{|\psi'_j(0)|}{\binom{n}{j}} \end{aligned}$$

and

$$\lim_{\lambda \rightarrow 0} \frac{|\psi_j(\lambda) \psi_{n-j}(\lambda) - \binom{n}{j}^2 \psi_n(\lambda)|}{|\lambda| \left(\binom{n}{j}^2 - |\psi_{n-j}(\lambda)|^2 \right)} = \frac{1}{\binom{n}{j}^2} \left| -\binom{n}{j}^2 \psi'_n(0) \right| = |\psi'_n(0)|.$$

So

$$\lim_{\lambda \rightarrow 0} \frac{\binom{n}{j} |(\psi_j - \bar{\psi}_{n-j} \psi_n)(\lambda)| + |(\psi_j \psi_{n-j} - \binom{n}{j}^2 \psi_n)(\lambda)|}{|\lambda| \left(\binom{n}{j}^2 - |\psi_{n-j}(\lambda)|^2 \right)} = \frac{|\psi'_j(0)|}{\binom{n}{j}} + |\psi'_n(0)|.$$

Since inequality (5) is true for all $\lambda \in \mathbb{D} \setminus \{0\}$, by dividing both side of (5) by $|\lambda|$ and letting $\lambda \rightarrow 0$, we have

$$\max_{1 \leq j \leq n-1} \left\{ \frac{|\psi'_j(0)|}{\binom{n}{j}} + |\psi'_n(0)| \right\} \leq 1.$$

Since $\psi'(0) = x$, that is, $(\psi'_1(0), \dots, \psi'_n(0)) = (x_1, \dots, x_n)$, we have

$$\max_{1 \leq j \leq n-1} \left\{ \frac{|x_j|}{\binom{n}{j}} \right\} + |x_n| \leq 1.$$

We divide the converse part into two cases, $n = 3$ and $n > 3$.

Case-I. Suppose $n = 3$ and condition (4) holds for $x = (x_1, x_2, x_3) \in \mathcal{K}_3$, that is,

$$\max \left\{ \frac{|x_1|}{3}, \frac{|x_2|}{3} \right\} + |x_3| \leq 1. \quad (6)$$

We show that there exists an analytic map $\psi : \mathbb{D} \rightarrow \tilde{\mathbb{G}}_3$ such that $\psi(0) = (0, 0, 0)$ and $\psi'(0) = (x_1, x_2, x_3)$. We shall follow similar technique as in Theorem 2.1 of [35] to construct such a function. We first assume that $|x_2| \leq |x_1|$. So, if $x_1 = 0$, then we have $x_2 = 0$ and hence from the inequality (6), we obtain $|x_3| \leq 1$. Now consider the function $\psi(\lambda) = (0, 0, \lambda x_3)$. Clearly ψ is analytic and satisfies

$$\psi(0) = (0, 0, 0) \quad \text{and} \quad \psi'(0) = \psi'(\lambda)|_{\lambda=0} = (0, 0, x_3) = x.$$

Next, assume $x_1 \neq 0$. According to Theorem 2.2, for any $W \in \mathcal{S}_{2 \times 2}$ the function $\psi = \pi_3 \circ W = (3W_{11}, 3W_{22}, W_{11}W_{22} - W_{12}W_{21})$ is an analytic map from \mathbb{D} to $\widetilde{\mathbb{G}}_3$. Hence it is enough to show the existence of a function $W = [W_{ij}] \in \mathcal{S}_{2 \times 2}$ such that $(\pi_3 \circ W)(0) = (0, 0, 0)$ and $(\pi_3 \circ W)'(0) = x$. Suppose W is a 2×2 matrix valued function such that

$$W(0) = \begin{bmatrix} 0 & \sigma \\ 0 & 0 \end{bmatrix}, \quad (7)$$

where $\sigma \in \mathbb{D}$ will be chosen later. Then, $\psi(0) = (\pi_3 \circ W)(0) = (0, 0, 0)$ and

$$\psi'(0) = (3W_{11}, 3W_{22}, W_{11}W_{22} - W_{12}W_{21})'(0) = (3W'_{11}, 3W'_{22}, -\sigma W'_{21})(0).$$

Accordingly, $\psi'(0) = x$ if and only if

$$W'(0) = \begin{bmatrix} x_1/3 & * \\ -x_3/\sigma & x_2/3 \end{bmatrix}. \quad (8)$$

We shall find a function $W \in \mathcal{S}_{2 \times 2}$ that satisfies equations (7) and (8).

For any $W \in \mathcal{S}_{2 \times 2}$, we have

$$(\mathcal{M}_Z \circ W)(\lambda) = -Z + D_{Z^*}W(\lambda)(1 - Z^*W(\lambda))^{-1}D_Z \quad \text{for } \lambda \in \mathbb{D}$$

and

$$\begin{aligned} (\mathcal{M}_Z \circ W)' &= D_{Z^*}[1 + W(1 - Z^*W)^{-1}Z^*]W'(1 - Z^*W)^{-1}D_Z \\ &= D_{Z^*}(1 - WZ^*)^{-1}W'(1 - Z^*W)^{-1}D_Z. \end{aligned} \quad (9)$$

For a fixed $\sigma \in \mathbb{D}$, let

$$Z = \begin{bmatrix} 0 & \sigma \\ 0 & 0 \end{bmatrix}. \quad (10)$$

Then $\|Z\| < 1$,

$$D_Z = (1 - Z^*Z)^{\frac{1}{2}} = \begin{bmatrix} 1 & 0 \\ 0 & (1 - |\sigma|^2)^{\frac{1}{2}} \end{bmatrix} \quad \text{and} \quad D_{Z^*} = \begin{bmatrix} (1 - |\sigma|^2)^{\frac{1}{2}} & 0 \\ 0 & 1 \end{bmatrix}.$$

Now if $W \in \mathcal{S}_{2 \times 2}$ satisfies (7) and (8), then $W(0) = Z$ and hence we have

$$\begin{aligned} (\mathcal{M}_Z \circ W)'(0) &= D_{Z^*}(1 - ZZ^*)^{-1}W'(0)(1 - Z^*Z)^{-1}D_Z \\ &= \begin{bmatrix} \frac{x_1}{3(1 - |\sigma|^2)^{\frac{1}{2}}} & \frac{W'_{12}(0)}{1 - |\sigma|^2} \\ \frac{-x_3}{\sigma} & \frac{x_2}{3(1 - |\sigma|^2)^{\frac{1}{2}}} \end{bmatrix}. \end{aligned} \quad (11)$$

Note that if $W \in \mathcal{S}_{2 \times 2}$ then by (7) and (10), the map $\mathcal{M}_Z \circ W : \mathbb{D} \longrightarrow R_I(2, 2)$ satisfies the following condition

$$(\mathcal{M}_Z \circ W)(0) = \mathcal{M}_Z(W(0)) = \mathcal{M}_Z(Z) = 0.$$

Therefore, by Schwarz lemma for $R_I(2, 2)$, we have $\|(\mathcal{M}_Z \circ W)'(0)\| < 1$. Thus, if there exists a function W in $\mathcal{S}_{2 \times 2}$ which satisfies (7) and (8), then the matrix in the right hand side of the equation (11) must be a strict contraction.

Now choose $\sigma = \sqrt{1 - \frac{|x_1|}{3}}$. By (6), $|x_3| < 1$ hence $\sigma \in \mathbb{D}$. For a $\rho \in \mathbb{C}$, define a matrix B_ρ by

$$B_\rho = \begin{bmatrix} \frac{x_1}{\sqrt{3|x_1|}} & \rho \\ -x_3 & \frac{x_2}{\sqrt{3|x_1|}} \end{bmatrix}. \quad (12)$$

For a fixed ρ (which is to be determined), define a function

$$V_\rho(\lambda) = \lambda B_\rho, \quad \lambda \in \mathbb{D}. \quad (13)$$

Then, $V_\rho(0) = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$ and $V'_\rho(0) = B_\rho$. We define B_ρ in such a fashion because, for a suitable choice of ρ , the matrix $V'_\rho(0)$ is analogous to the matrix in equation (11) with above choice of σ . Since we have assumed $|x_2| \leq |x_1|$ and since condition (6) holds, we have that $|x_1|/3 + |x_3| \leq 1$. Thus, the norm of first column of B_ρ is equal to

$$\left(\frac{|x_1|}{3} + \frac{|x_3|^2}{(1 - |x_1|/3)} \right)^{\frac{1}{2}} \leq \left(\frac{|x_1|}{3} + \frac{|x_3|(1 - |x_1|/3)}{(1 - |x_1|/3)} \right)^{\frac{1}{2}} = \left(\frac{|x_1|}{3} + |x_3| \right)^{\frac{1}{2}} \leq 1$$

and also the norm of the second row of B_ρ is

$$\left(\frac{|x_3|^2}{(1 - |x_1|/3)} + \frac{|x_2|^2}{3|x_1|} \right)^{\frac{1}{2}} \leq \frac{|x_3|^2}{(1 - |x_1|/3)} + \frac{|x_1|^2}{3|x_1|} \leq 1 \text{ (since } |x_2| \leq |x_1| \text{)}.$$

Then, by Theorem 2.4, there exists a $\rho \in \mathbb{C}$ such that $\|B_\rho\| \leq 1$. Consequently, there exists a $\rho \in \mathbb{C}$ such that $V_\rho \in \mathcal{S}_{2 \times 2}$. A choice of such ρ is

$$\rho = \frac{x_1 x_2 \bar{x}_3 \sqrt{3 - |x_1|}}{\sqrt{3}|x_1|(3 - |x_1| - 3|x_3|^2)}. \quad (14)$$

Now we define a function $W = \mathcal{M}_{-Z} \circ V_\rho$, where Z is as in (10). Then $W \in \mathcal{S}_{2 \times 2}$ and $W(0) = \mathcal{M}_{-Z}(0) = Z$. So W satisfies equation (7). Since $V_\rho(0)$ is the zero matrix and $V'_\rho(0) = B_\rho$, by equation (9) we have

$$\begin{aligned} W'(0) &= D_{Z^*}(1 + V_\rho(0)Z^*)^{-1}V'_\rho(0)(1 + Z^*V_\rho(0))^{-1}D_Z \\ &= D_{Z^*}V'_\rho(0)D_Z = \begin{bmatrix} x_1/3 & \rho|x_1|/3 \\ -x_3/\sigma & x_2/3 \end{bmatrix}. \end{aligned}$$

Hence the function W also satisfies equation (8). Thus, there exists a function $W \in \mathcal{S}_{2 \times 2}$ that satisfies (7) and (8). The case $|x_1| \leq |x_2|$ can be dealt in similar way. Hence condition (6) is sufficient for the existence of an analytic map $\psi : \mathbb{D} \longrightarrow \widetilde{\mathbb{G}}_3$ such that $\psi(0) = (0, \dots, 0)$ and $\psi'(0) = (x_1, x_2, x_3)$.

Case-II. Let $n > 3$. First assume that n is odd. Suppose $x = (x_1, \dots, x_n) \in \mathcal{K}_n$. Then for all $j = 2, \dots, \lfloor \frac{n}{2} \rfloor$, we have $x_j = \frac{\binom{n}{j}x_1}{n}$ and $x_{n-j} = \frac{\binom{n}{j}x_{n-1}}{n}$. Therefore, condition (4) reduces to

$$\max \left\{ \frac{|x_1|}{n}, \frac{|x_{n-1}|}{n} \right\} + |x_n| \leq 1 \quad \Leftrightarrow \quad \max \left\{ \frac{\left| \frac{3x_1}{n} \right|}{3}, \frac{\left| \frac{3x_{n-1}}{n} \right|}{3} \right\} + |x_n| \leq 1.$$

Hence by Case-I, there exists a function $\widehat{W} \in \mathcal{S}_{2 \times 2}$ such that

$$\widehat{W}(0) = \begin{bmatrix} 0 & \sigma \\ 0 & 0 \end{bmatrix} \quad \text{and} \quad \widehat{W}'(0) = \begin{bmatrix} x_1/n & * \\ -x_n/\sigma & x_{n-1}/n \end{bmatrix}. \quad (15)$$

Now consider $\lfloor \frac{n}{2} \rfloor$ number of 2×2 matrix valued functions $W_1, \dots, W_{\lfloor \frac{n}{2} \rfloor}$, where $W_j(\lambda) = \widehat{W}(\lambda)$ for each $j = 1, \dots, \lfloor \frac{n}{2} \rfloor$. Since n is odd, $2\lfloor \frac{n}{2} \rfloor + 1 = n$. So, we have

$$\begin{aligned} &(\pi_{2\lfloor \frac{n}{2} \rfloor + 1} \circ (W_1, \dots, W_{\lfloor \frac{n}{2} \rfloor}))'(0) \\ &= \left(x_1, \dots, \frac{\binom{n}{\lfloor \frac{n}{2} \rfloor}}{n} x_1, \frac{\binom{n}{\lfloor \frac{n}{2} \rfloor}}{n} x_{n-1}, \dots, x_{n-1}, x_n \right) = x \end{aligned}$$

and

$$\left(\pi_n \circ (W_1, \dots, W_{\lfloor \frac{n}{2} \rfloor}) \right)(0) = \pi_{2\lfloor \frac{n}{2} \rfloor + 1} \left(\widehat{W}(0), \dots, \widehat{W}(0) \right) = (0, \dots, 0).$$

Note that, each $W_j \in \mathcal{S}_{2 \times 2}$ and $\det W_i = \det W_j$ for each i, j . Clearly, the function $\psi = \pi_n \circ (W_1, \dots, W_{\lfloor \frac{n}{2} \rfloor})$ is analytic which maps \mathbb{D} into $\widetilde{\mathbb{G}}_n$.

Now suppose n is even. In this case, $x = (x_1, \dots, x_n) \in \mathcal{K}_n^{even}$. Then, $x_j = \frac{\binom{n}{j}x_1}{n}$, $x_{n-j} = \frac{\binom{n}{j}x_{n-1}}{n}$ for all $j = 2, \dots, [\frac{n}{2}] - 1$ and $x_{[\frac{n}{2}]} = \frac{\binom{n}{[\frac{n}{2}]}(x_1 + x_{n-1})}{2n}$. Hence condition (4) is reduced to

$$\max \left\{ \frac{|x_1|}{n}, \frac{|x_{n-1}|}{n}, \frac{|x_1| + |x_{n-1}|}{2n} \right\} + |x_n| \leq 1,$$

which is same as

$$\max \left\{ \frac{|x_1|}{n}, \frac{|x_{n-1}|}{n} \right\} + |x_n| \leq 1.$$

So in a similar fashion as if n is odd, there exists a function $\widehat{W} \in \mathcal{S}_{2 \times 2}$ which satisfies condition (15). Again consider $\frac{n}{2}$ number of 2×2 matrix valued functions $W_1, \dots, W_{\frac{n}{2}}$, where $W_j(\lambda) = \widehat{W}(\lambda)$ for each $j = 1, \dots, \frac{n}{2}$. So, we have

$$\begin{aligned} & \left(\pi_{2[\frac{n}{2}]} \circ (W_1, \dots, W_{\frac{n}{2}}) \right)'(0) \\ &= \left(x_1, \dots, \frac{\binom{n}{\frac{n}{2}-1}}{n}x_1, \left(\frac{n}{\frac{n}{2}} \right) \frac{x_1 + x_{n-1}}{2}, \frac{\binom{n}{\frac{n}{2}-1}}{n}x_{n-1}, \dots, x_{n-1}, x_n \right) = x \end{aligned}$$

and

$$(\pi_n \circ (W_1, \dots, W_{\frac{n}{2}}))(0) = (0, \dots, 0).$$

The function $\psi = \pi_n \circ (W_1, \dots, W_{\frac{n}{2}})$ is analytic and it maps \mathbb{D} into $\mathcal{K}_n^{even} \cap \widetilde{\mathbb{G}}_n$. Thus, for any $n \in \mathbb{N}$ if condition (4) holds for $x \in \mathcal{K}_n$ then there exists an analytic map $\psi : \mathbb{D} \rightarrow \widetilde{\mathbb{G}}_n$ such that $\psi(0) = (0, \dots, 0)$ and $\psi'(0) = x$. The proof is now complete. ■

Remark. In particular if $n = 3$, then the condition (4) is necessary and sufficient for the existence of such an interpolating function ψ for any $x \in \mathbb{C}^3$.

The following is a Schwarz type lemma for the symmetrized polydisc.

Theorem 3.2. *Let $x = (x_1, \dots, x_n) \in \mathbb{C}^n$. If there exists an analytic map $\psi : \mathbb{D} \rightarrow \mathbb{G}_n$ such that $\psi(0) = (0, \dots, 0)$ and $\psi'(0) = x$, then*

$$\max_{1 \leq j \leq n-1} \left\{ \frac{|x_j|}{\binom{n}{j}} \right\} + |x_n| \leq 1.$$

Proof. Follows from Theorem 3.1 as $\mathbb{G}_n \subset \tilde{\mathbb{G}}_n$. ■

4. AN EXPLICIT INTERPOLATING FUNCTION

In Theorem 3.1, we proved the existence of an analytic function ψ mapping origin to origin and satisfying $\psi'(0) = (x_1, \dots, x_n) \in \mathcal{K}_n$. In this section, we show an explicit construction of such a function ψ .

Theorem 4.1. *Let $x \in \mathcal{K}_n$. If a function $\psi = (\psi_1, \dots, \psi_n)$ is given by*

$$\psi(\lambda) = \frac{\lambda}{1 + \lambda \bar{x}_n r_x} \left(x_1, \dots, \binom{n}{[\frac{n}{2}]} \frac{x_1}{n}, \binom{n}{[\frac{n}{2}]} \frac{x_{n-1}}{n}, \dots, x_{n-1}, x_n + \lambda r_x \right)$$

when n is odd, and

$$\psi(\lambda) = \frac{\lambda}{1 + \lambda \bar{x}_n r_x} \left(x_1, \dots, \binom{n}{\frac{n-2}{2}} \frac{x_1}{n}, \binom{n}{\frac{n}{2}} \frac{x_1 + x_{n-1}}{2n}, \right. \\ \left. \binom{n}{\frac{n-2}{2}} \frac{x_{n-1}}{n}, \dots, x_{n-1}, x_n + \lambda r_x \right).$$

when n is even, where

$$r_x = \begin{cases} 0 & \text{if } x_1 = 0 = x_{n-1} \\ \frac{x_1 x_{n-1} (n - |x_1|)}{n |x_1| (n - |x_1| - n |x_n|^2)} & \text{if } |x_{n-1}| \leq |x_1| \neq 0 \\ \frac{x_1 x_{n-1} (n - |x_{n-1}|)}{n |x_{n-1}| (n - |x_{n-1}| - n |x_n|^2)} & \text{if } |x_1| \leq |x_{n-1}| \neq 0. \end{cases}$$

Then ψ is an analytic map from \mathbb{D} into $\tilde{\mathbb{G}}_n$ with $\psi(0) = (0, \dots, 0)$ and $\psi'(0) = x$.

Proof. We divide the proof into two cases, $n = 3$ and $n > 3$ as in Theorem 3.1. The idea and technique that are used in constructing an interpolating function ψ when $n = 3$ are borrowed from Theorem 2.2 in [35].

Case-I: Suppose $n = 3$. Then $\mathcal{K}_3 = \mathbb{C}^3$. Let $x \in \mathbb{C}^3$ be such that $\max \left\{ \frac{|x_1|}{3}, \frac{|x_2|}{3} \right\} + |x_3| \leq 1$. For this particular case, we denote r_x by l_x . Then

$$\psi(\lambda) = \frac{\lambda}{1 + \lambda \bar{x}_3 l_x} (x_1, x_2, l_x \lambda + x_3), \quad (16)$$

where

$$l_x = \begin{cases} 0 & \text{if } x_1 = 0 = x_2 \\ \frac{x_1 x_2 (3 - |x_1|)}{3|x_1|(3 - |x_1| - 3|x_3|^2)} & \text{if } |x_2| \leq |x_1| \neq 0 \\ \frac{x_1 x_2 (3 - |x_2|)}{3|x_2|(3 - |x_2| - 3|x_3|^2)} & \text{if } |x_1| \leq |x_2| \neq 0. \end{cases} \quad (17)$$

We shall show that the function ψ given by (16) and (17) is analytic, $\psi(\mathbb{D}) \subset \tilde{\mathbb{G}}_3$, $\psi(0) = (0, 0, 0)$ and $\psi'(0) = x$. Suppose $x_1 = 0 = x_2$, then $|x_3| \leq 1$ and $l_x = 0$. Consider the function $\psi(\lambda) = (0, 0, \lambda x_3)$, $\lambda \in \mathbb{D}$. Clearly ψ is analytic, $\psi(0) = (0, 0, 0)$ and $\psi'(0) = (0, 0, x_3) = x$. Now suppose $|x_2| \leq |x_1| \neq 0$. The function ψ , given by (16) and (17), clearly satisfies $\psi(0) = 0$. Note that

$$\psi'(\lambda) = \left(\frac{x_1}{(1 + \lambda \bar{x}_3 l_x)^2}, \frac{x_2}{(1 + \lambda \bar{x}_3 l_x)^2}, \frac{x_3 + \lambda l_x (2 + \lambda \bar{x}_3 l_x)}{(1 + \lambda \bar{x}_3 l_x)^2} \right).$$

Hence $\psi'(0) = x$. We shall show that ψ is analytic and $\psi(\mathbb{D}) \subset \tilde{\mathbb{G}}_3$.

Consider Z as in equation (10) with $\sigma = \sqrt{1 - \frac{|x_1|}{3}}$. Then

$$D_Z = \begin{bmatrix} 1 & 0 \\ 0 & \sqrt{\frac{|x_1|}{3}} \end{bmatrix} \text{ and } D_{Z^*} = \begin{bmatrix} \sqrt{\frac{|x_1|}{3}} & 0 \\ 0 & 1 \end{bmatrix}.$$

Also consider B_ρ as in equation (12), where ρ is given by (14). Then $\|B_\rho\| \leq 1$ (as we observe in the proof of Theorem 3.1). Consider the function $W(\lambda) = \mathcal{M}_{-Z}(\lambda B_\rho)$, $\lambda \in \mathbb{D}$. Then $W \in \mathcal{S}_{2 \times 2}$. For a contraction Z , we know that $Z^* D_{Z^*} = D_Z Z^*$ and hence $D_Z^{-1} Z^* = Z^* D_{Z^*}^{-1}$ when $\|Z\| < 1$. Here $D_Z = (1 - Z^* Z)^{\frac{1}{2}}$. Thus

$$\begin{aligned} W(\lambda) &= Z + D_{Z^*} \lambda B_\rho (1 + Z^* \lambda B_\rho)^{-1} D_Z \\ &= Z + \lambda (D_{Z^*} B_\rho D_Z) (1 + \lambda Z^* D_{Z^*}^{-1} B_\rho D_Z)^{-1}. \end{aligned}$$

Note that,

$$D_{Z^*} B_\rho D_Z = \begin{bmatrix} \frac{x_1}{3} & \frac{\rho |x_1|}{3} \\ \frac{-x_3}{\sigma} & \frac{x_2}{3} \end{bmatrix} \text{ and } D_{Z^*}^{-1} B_\rho D_Z = \begin{bmatrix} \frac{x_1}{|x_1|} & \rho \\ \frac{-x_3}{\sigma} & \frac{x_2}{3} \end{bmatrix}.$$

So,

$$(1 + \lambda Z^* D_{Z^*}^{-1} B_\rho D_Z)^{-1} = \frac{1}{1 + \lambda \rho \sigma} \begin{bmatrix} 1 + \lambda \rho \sigma & 0 \\ \frac{-x_1 \lambda \sigma}{|x_1|} & 1 \end{bmatrix}.$$

Thus

$$W(\lambda) = \begin{bmatrix} 0 & \sigma \\ 0 & 0 \end{bmatrix} + \frac{\lambda}{1 + \lambda \rho \sigma} \begin{bmatrix} \frac{x_1}{3} & \frac{\rho |x_1|}{3} \\ v & \frac{x_2}{3} \end{bmatrix}, \quad (18)$$

where

$$v = v(\lambda) = -\frac{x_3}{\sigma}(1 + \lambda \rho \sigma) - \frac{\lambda \sigma x_1 x_2}{3|x_1|} = -\frac{x_3}{\sigma} - \lambda \sigma l_x. \quad (19)$$

From (18) we have

$$\det W(\lambda) = \frac{\lambda^2(x_1 x_2 - 3v \rho |x_1|)}{9(1 + \lambda \rho \sigma)^2} - \frac{\lambda \sigma v}{(1 + \lambda \rho \sigma)}.$$

Again

$$\rho \sigma = \frac{x_1 x_2 \bar{x}_3 (3 - |x_1|)}{3|x_1|(3 - |x_1| - 3|x_3|^2)} = \bar{x}_3 l_x, \quad \frac{\rho}{\sigma} = \frac{x_1 x_2 \bar{x}_3}{|x_1|(3 - |x_1| - 3|x_3|^2)}$$

and thus

$$\begin{aligned} x_1 x_2 - 3v \rho |x_1| &= x_1 x_2 + 3x_3 \frac{x_1 x_2 \bar{x}_3 |x_1|}{|x_1|(3 - |x_1| - 3|x_3|^2)} + 3\lambda \bar{x}_3 (l_x)^2 |x_1| \\ &= 3|x_1| l_x (1 + \lambda \bar{x}_3 l_x). \end{aligned}$$

Then

$$\det W(\lambda) = \frac{3\lambda^2 |x_1| l_x (1 + \lambda \bar{x}_3 l_x)}{9(1 + \lambda \bar{x}_3 l_x)^2} - \frac{\lambda \sigma v}{(1 + \lambda \bar{x}_3 l_x)} = \frac{\lambda}{1 + \lambda \bar{x}_3 l_x} (x_3 + \lambda l_x).$$

From the equation (18), we have

$$[W(\lambda)]_{11} = \frac{\lambda x_1}{3(1 + \lambda \bar{x}_3 l_x)} \quad \text{and} \quad [W(\lambda)]_{22} = \frac{\lambda x_2}{3(1 + \lambda \bar{x}_3 l_x)}.$$

Consider the function

$$\psi(\lambda) = \pi \circ W(\lambda) = ([3W(\lambda)]_{11}, [3W(\lambda)]_{22}, \det W(\lambda)), \quad \lambda \in \mathbb{D}.$$

Then

$$\psi(\lambda) = \left(\frac{\lambda x_1}{1 + \lambda \bar{x}_3 l_x}, \frac{\lambda x_2}{1 + \lambda \bar{x}_3 l_x}, \frac{\lambda}{1 + \lambda \bar{x}_3 l_x} (x_3 + \lambda l_x) \right),$$

which is of the form given in (16). Since $W \in \mathcal{S}_{2 \times 2}$, by Theorem 2.2 the map ψ is an analytic and $\psi(\mathbb{D}) \subset \widetilde{\mathbb{G}}_3$. The case when $|x_1| \leq |x_2| \neq 0$, can be dealt in a similar way. Hence we are done for $n = 3$.

Case-II: Suppose $n > 3$. Let $x = (x_1, \dots, x_n) \in \mathcal{K}_n$. First suppose n is odd. Then, $x_j = \frac{\binom{n}{j}}{n} x_1$ and $x_{n-j} = \frac{\binom{n}{j}}{n} x_{n-1}$ for all $j = 2, \dots, \left[\frac{n}{2}\right]$. Hence,

$$\max_{1 \leq j \leq n-1} \left\{ \frac{|x_j|}{\binom{n}{j}} \right\} + |x_n| \leq 1 \iff \max \left\{ \frac{|x_1|}{n}, \frac{|x_{n-1}|}{n} \right\} + |x_n| \leq 1.$$

Consider $y = (y_1, y_2, y_3) = \left(\frac{3x_1}{n}, \frac{3x_{n-1}}{n}, x_n \right)$. Therefore, by hypothesis

$$\max \left\{ \frac{|y_1|}{3}, \frac{|y_2|}{3} \right\} + |y_3| \leq 1.$$

So, by Case-I, there exists $\widehat{W} \in \mathcal{S}_{2 \times 2}$ such that

$$[\widehat{W}(\lambda)]_{11} = \frac{\lambda y_1}{3(1 + \lambda \bar{y}_3 l_y)}, \quad [\widehat{W}(\lambda)]_{22} = \frac{\lambda y_2}{3(1 + \lambda \bar{y}_3 l_y)}$$

and $\det \widehat{W}(\lambda) = \frac{\lambda}{1 + \lambda \bar{y}_3 l_y} (y_3 + \lambda l_y)$, where

$$l_y = \begin{cases} 0 & \text{if } y_1 = 0 = y_2 \\ \frac{y_1 y_2 (3 - |y_1|)}{3|y_1|(3 - |y_1| - 3|y_3|^2)} & \text{if } |y_2| \leq |y_1| \neq 0 \\ \frac{y_1 y_2 (3 - |y_2|)}{3|y_2|(3 - |y_2| - 3|y_3|^2)} & \text{if } |y_1| \leq |y_2| \neq 0. \end{cases}$$

Substituting the values of y , we get the following:

$$(i) \quad l_y = r_x$$

$$(ii) \quad \det \widehat{W}(\lambda) = \frac{\lambda}{1 + \lambda \bar{x}_n r_x} (x_n + \lambda r_x) \quad (20)$$

$$(iii) \quad [\widehat{W}(\lambda)]_{11} = \frac{\lambda x_1}{n(1 + \lambda \bar{x}_n r_x)}, \quad [\widehat{W}(\lambda)]_{22} = \frac{\lambda x_{n-1}}{n(1 + \lambda \bar{x}_n r_x)}. \quad (21)$$

Consider the 2×2 matrix valued functions $W_1, \dots, W_{\left[\frac{n}{2}\right]}$, where $W_j(\lambda) = \widehat{W}(\lambda)$ for each $j = 1, \dots, \left[\frac{n}{2}\right]$. Then each $W_j \in \mathcal{S}_{2 \times 2}$. Since n is odd,

$2 \left\lfloor \frac{n}{2} \right\rfloor + 1 = n$. Therefore,

$$\begin{aligned} & \left(\pi_{2 \left\lfloor \frac{n}{2} \right\rfloor + 1} \circ (W_1, \dots, W_{\left\lfloor \frac{n}{2} \right\rfloor}) \right) (\lambda) \\ &= \frac{\lambda}{(1 + \lambda \bar{x}_n r_x)} \left(x_1, \dots, \binom{n}{\left\lfloor \frac{n}{2} \right\rfloor} \frac{x_1}{n}, \binom{n}{\left\lfloor \frac{n}{2} \right\rfloor} \frac{x_{n-1}}{n}, \dots, x_{n-1}, (x_n + \lambda r_x) \right). \end{aligned}$$

Now suppose n is even. So, $\left\lfloor \frac{n}{2} \right\rfloor = \frac{n}{2}$. In this case, $x = (x_1, \dots, x_n) \in \mathcal{K}_n^{\text{even}}$. Thus $x_j = \frac{\binom{n}{j}}{n} x_1$, $x_{n-j} = \frac{\binom{n}{j}}{n} x_{n-1}$ for all $j = 2, \dots, \frac{n}{2} - 1$ and $x_{\frac{n}{2}} = \frac{\binom{n}{\frac{n}{2}}}{2n} \frac{x_1 + x_{n-1}}{2n}$. Again,

$$\max_{1 \leq j \leq n-1} \left\{ \frac{|x_j|}{\binom{n}{j}} \right\} + |x_n| \leq 1 \Leftrightarrow \max \left\{ \frac{|x_1|}{n}, \frac{|x_{n-1}|}{n} \right\} + |x_n| \leq 1.$$

Therefore, as in the case when n is odd, there exists $\widehat{W} \in \mathcal{S}_{2 \times 2}$ such that (20) and (21) hold. Now consider the 2×2 matrix valued functions $W_1, \dots, W_{\frac{n}{2}}$, where $W_j(\lambda) = \widehat{W}(\lambda)$ for each $j = 1, \dots, \frac{n}{2}$. Then

$$\begin{aligned} & \left(\pi_{2 \left\lfloor \frac{n}{2} \right\rfloor} \circ (W_1, \dots, W_{\frac{n}{2}}) \right) (\lambda) \\ &= \frac{\lambda}{(1 + \lambda \bar{x}_n r_x)} \left(x_1, \dots, \binom{n}{\frac{n-2}{2}} \frac{x_1}{n}, \binom{n}{\frac{n}{2}} \frac{x_1 + x_{n-1}}{2n}, \right. \\ & \quad \left. \binom{n}{\frac{n-2}{2}} \frac{x_{n-1}}{n}, \dots, x_{n-1}, x_n + \lambda r_x \right). \end{aligned}$$

In both cases $\psi = \pi_n \circ (W_1, \dots, W_{\left\lfloor \frac{n}{2} \right\rfloor})$. Clearly, the function ψ is an analytic map from \mathbb{D} into $\widetilde{\mathbb{G}}_n$ and $\psi(0) = (0, \dots, 0)$. If n is odd, then

$$\begin{aligned} \psi'(\lambda) &= \frac{1}{(1 + \lambda \bar{x}_n r_x)^2} \left(x_1, \dots, \binom{n}{\left\lfloor \frac{n}{2} \right\rfloor} \frac{x_1}{n}, \binom{n}{\left\lfloor \frac{n}{2} \right\rfloor} \frac{x_{n-1}}{n}, \right. \\ & \quad \left. \dots, x_{n-1}, x_n + \lambda r_x (2 + \lambda \bar{x}_n r_x) \right). \end{aligned}$$

If n is even, we have

$$\begin{aligned} \psi'(\lambda) &= \frac{1}{(1 + \lambda \bar{x}_n r_x)^2} \left(x_1, \dots, \binom{n}{\frac{n-2}{2}} \frac{x_1}{n}, \binom{n}{\frac{n}{2}} \frac{(x_1 + x_{n-1})}{2n}, \right. \\ & \quad \left. \binom{n}{\frac{n-2}{2}} \frac{x_{n-1}}{n}, \dots, x_{n-1}, x_n + \lambda r_x (2 + \lambda \bar{x}_n r_x) \right). \end{aligned}$$

It is evident that in either cases $\psi'(0) = x$ and the proof is complete. \blacksquare

5. GEOMETRIC INTERPLAY BETWEEN THE MEMBERS OF $\widetilde{\mathbb{G}}_n$ AND $\widetilde{\Gamma}_n$

In [28], we have witnessed several important geometric properties of $\widetilde{\mathbb{G}}_n$ and $\widetilde{\Gamma}_n$, e.g., $\widetilde{\Gamma}_n$ is polynomially convex but not convex, $\widetilde{\mathbb{G}}_n$ is starlike but not circled etc. In this section, we shall see some interplay between $\widetilde{\mathbb{G}}_n$ (or $\widetilde{\Gamma}_n$) and $\widetilde{\mathbb{G}}_{n+1}$ (or, $\widetilde{\Gamma}_{n+1}$).

Theorem 5.1. *Let $n \in \mathbb{N}$. Suppose $y = (y_1, \dots, y_{n-1}, q) \in \mathbb{C}^n$*

- (1) *If n is an even number, then the point $y \in \widetilde{\mathbb{G}}_n$ (or, $\in \widetilde{\Gamma}_n$) if and only if $\hat{y} \in \widetilde{\mathbb{G}}_{n+1}$ (or, $\in \widetilde{\Gamma}_{n+1}$), where*

$$\hat{y} = \left(\frac{n+1}{n}y_1, \dots, \frac{n+1}{n+1-j}y_j, \dots, \frac{n+1}{\frac{n}{2}+1}y_{\frac{n}{2}}, \frac{n+1}{\frac{n}{2}+1}y_{\frac{n}{2}}, \right. \\ \left. \underbrace{\frac{n+1}{\frac{n}{2}+2}y_{\frac{n}{2}+1}}_{(\frac{n}{2}+2)\text{-th position}}, \dots, \underbrace{\frac{n+1}{n+1-j}y_{n-j}}_{n+1-j\text{-th position}}, \dots, \frac{n+1}{n}y_{n-1}, q \right).$$

- (2) *If n is an odd number, then the point $y \in \widetilde{\mathbb{G}}_n$ (or, $\in \widetilde{\Gamma}_n$) if and only if $y^* \in \widetilde{\mathbb{G}}_{n+1}$ (or, $\in \widetilde{\Gamma}_{n+1}$), where*

$$y^* = \left(\frac{n+1}{n}y_1, \dots, \frac{n+1}{n+1-j}y_j, \dots, \frac{2(n+1)}{n+3}y_{[\frac{n}{2}]}, (y_{[\frac{n}{2}]} + y_{[\frac{n}{2}]+1}), \right. \\ \left. \underbrace{\frac{2(n+1)}{n+3}y_{[\frac{n}{2}]+1}}_{([\frac{n}{2}]+2)\text{-th position}}, \dots, \underbrace{\frac{n+1}{n+1-j}y_{n-j}}_{n+1-j\text{-th position}}, \dots, \frac{n+1}{n}y_{n-1}, q \right).$$

Proof. (1). First note that $\binom{n+1}{j} = \frac{n+1}{n+1-j}\binom{n}{j}$. As n is even, $[\frac{n}{2}] = \frac{n}{2} = \lceil \frac{n+1}{2} \rceil$. Suppose $y \in \widetilde{\mathbb{G}}_n$. Then, by Theorem 2.2, we have

$$\binom{n}{j} |y_j - \bar{y}_{n-j}q| + \left| y_j y_{n-j} - \binom{n}{j}^2 q \right| < \binom{n}{j}^2 - |y_{n-j}|^2 \quad \text{for } j = 1, \dots, \frac{n}{2}.$$

Consider the point $\hat{y} = (\hat{y}_1, \dots, \hat{y}_n, \hat{q})$, where $\hat{q} = q$ and

$$\hat{y}_j = \frac{n+1}{n+1-j}y_j, \quad \hat{y}_{n+1-j} = \frac{n+1}{n+1-j}y_{n-j} \quad \text{for } j = 1, \dots, \frac{n}{2}.$$

Then, the following holds for each $j = 1, \dots, \frac{n}{2}$.

$$\begin{aligned}
& \binom{n+1}{j} |\hat{y}_j - \bar{\hat{y}}_{n+1-j} \hat{q}| + \left| \hat{y}_j \hat{y}_{n+1-j} - \binom{n+1}{j}^2 \hat{q} \right| \\
&= \binom{n+1}{j} \frac{n+1}{n+1-j} |y_j - \bar{y}_{n-j} q| + \left(\frac{n+1}{n+1-j} \right)^2 \left| y_j y_{n-j} - \binom{n}{j}^2 q \right| \\
&= \left(\frac{n+1}{n+1-j} \right)^2 \left[\binom{n}{j} |y_j - \bar{y}_{n-j} q| + \left| y_j y_{n-j} - \binom{n}{j}^2 q \right| \right] \\
&< \left(\frac{n+1}{n+1-j} \right)^2 \left[\binom{n}{j}^2 - |y_{n-j}|^2 \right] \\
&= \binom{n+1}{j}^2 - |\hat{y}_{n+1-j}|^2.
\end{aligned}$$

Therefore, by Theorem 2.2, $\hat{y} \in \tilde{\mathbb{G}}_{n+1}$.

Conversely, suppose $\hat{y} \in \tilde{\mathbb{G}}_{n+1}$. Then for each $j = 1, \dots, \left\lfloor \frac{n+1}{2} \right\rfloor$, we have

$$\binom{n+1}{j} |\hat{y}_j - \bar{\hat{y}}_{n+1-j} \hat{q}| + \left| \hat{y}_j \hat{y}_{n+1-j} - \binom{n+1}{j}^2 \hat{q} \right| < \binom{n+1}{j}^2 - |\hat{y}_{n+1-j}|.$$

Similarly, we have

$$\binom{n}{j} |y_j - \bar{y}_{n-j} q| + \left| y_j y_{n-j} - \binom{n}{j}^2 q \right| < \binom{n}{j}^2 - |y_{n-j}|^2$$

for any $j = 1, \dots, \frac{n}{2}$. Consequently, by Theorem 2.2, $y \in \tilde{\mathbb{G}}_n$. In a similar fashion one can prove that $y \in \tilde{\Gamma}_n$ if and only if $\hat{y} \in \tilde{\Gamma}_{n+1}$.

(2). Suppose n is odd and suppose $y = (y_1, \dots, y_{n-1}, q) \in \tilde{\mathbb{G}}_n$. Then $|q| < 1$ and there exists $(\beta_1, \dots, \beta_{n-1}) \in \mathbb{C}^{n-1}$ such that

$$y_j = \beta_j + \bar{\beta}_{n-j} q, \quad y_{n-j} = \beta_{n-j} + \bar{\beta}_j q \quad \text{and} \quad |\beta_j| + |\beta_{n-j}| < \binom{n}{j},$$

for each $j = 1, \dots, \left\lfloor \frac{n}{2} \right\rfloor$. Since n is odd, $\left\lfloor \frac{n}{2} \right\rfloor = \frac{n-1}{2}$ and $\left\lfloor \frac{n+1}{2} \right\rfloor = \frac{n+1}{2}$. Consider $(\gamma_1, \dots, \gamma_n) \in \mathbb{C}^n$, where

$$\gamma_{\frac{n+1}{2}} = \beta_{\frac{n-1}{2}} + \beta_{\frac{n+1}{2}}, \quad \gamma_j = \frac{n+1}{n+1-j} \beta_j \quad \text{and} \quad \gamma_{n+1-j} = \frac{n+1}{n+1-j} \beta_{n-j},$$

for $j = 1, \dots, \left\lfloor \frac{n}{2} \right\rfloor$. Then, we have

$$2|\gamma_{\frac{n+1}{2}}| \leq 2\left(|\beta_{\left\lfloor \frac{n}{2} \right\rfloor}| + |\beta_{\left\lfloor \frac{n}{2} \right\rfloor+1}|\right) < \frac{n+1}{n+1-\left\lfloor \frac{n+1}{2} \right\rfloor} \binom{n}{\left\lfloor \frac{n}{2} \right\rfloor} = \binom{n+1}{\left\lfloor \frac{n+1}{2} \right\rfloor}$$

and

$$|\gamma_j| + |\gamma_{n+1-j}| = \frac{n+1}{n+1-j} (|\beta_j| + |\beta_{n-j}|) < \frac{n+1}{n+1-j} \binom{n}{j} = \binom{n+1}{j},$$

for all $j = 1, \dots, \left\lfloor \frac{n}{2} \right\rfloor$. Therefore,

$$(\gamma_1 + \overline{\gamma}_n q, \dots, \underbrace{\gamma_j + \overline{\gamma}_{n+1-j} q}_{j\text{-th position}}, \dots, \gamma_n + \overline{\gamma}_1 q, q) \in \widetilde{\mathbb{G}}_{n+1}.$$

Also

$$\gamma_{\frac{n+1}{2}} + \overline{\gamma}_{\frac{n+1}{2}} q = \beta_{\left\lfloor \frac{n}{2} \right\rfloor} + \overline{\beta}_{\left\lfloor \frac{n}{2} \right\rfloor+1} q + \beta_{\left\lfloor \frac{n}{2} \right\rfloor+1} + \overline{\beta}_{\left\lfloor \frac{n}{2} \right\rfloor} q = y_{\left\lfloor \frac{n}{2} \right\rfloor} + y_{\left\lfloor \frac{n}{2} \right\rfloor+1}$$

and

$$\begin{aligned} \gamma_j + \overline{\gamma}_{n+1-j} q &= \frac{n+1}{n+1-j} (\beta_j + \overline{\beta}_{n-j} q) = \frac{n+1}{n+1-j} y_j, \\ \gamma_{n+1-j} + \overline{\gamma}_j q &= \frac{n+1}{n+1-j} (\beta_{n-j} + \overline{\beta}_j q) = \frac{n+1}{n+1-j} y_{n-j}, \end{aligned}$$

for all $j = 1, \dots, \left\lfloor \frac{n}{2} \right\rfloor$. Thus,

$$\begin{aligned} &\left(\frac{n+1}{n} y_1, \dots, \frac{n+1}{n+1-j} y_j, \dots, \frac{2(n+1)}{n+3} y_{\left\lfloor \frac{n}{2} \right\rfloor}, (y_{\left\lfloor \frac{n}{2} \right\rfloor} + y_{\left\lfloor \frac{n}{2} \right\rfloor+1}), \right. \\ &\quad \left. \underbrace{\frac{2(n+1)}{n+3} y_{\left\lfloor \frac{n}{2} \right\rfloor+1}, \dots, \frac{n+1}{n+1-j} y_{n-j}, \dots, \frac{n+1}{n} y_{n-1}}_{(\left\lfloor \frac{n}{2} \right\rfloor+2)\text{-th position}}, q \right) \in \widetilde{\mathbb{G}}_{n+1}. \end{aligned}$$

Conversely, suppose $y^* = (y_1^*, \dots, y_n^*, q) \in \widetilde{\mathbb{G}}_{n+1}$. Then

$$y_{\frac{n+1}{2}}^* = y_{\left\lfloor \frac{n}{2} \right\rfloor} + y_{\left\lfloor \frac{n}{2} \right\rfloor+1}, \quad y_j^* = \frac{n+1}{n+1-j} y_j, \quad \text{and} \quad y_{n+1-j}^* = \frac{n+1}{n+1-j} y_{n-j},$$

for all $j = 1, \dots, \frac{n-1}{2}$. By definition, there exists $(\gamma_1, \dots, \gamma_n) \in \mathbb{C}^n$ such that

$$y_j^* = \gamma_j + \bar{\gamma}_{n+1-j}q, \quad y_{n+1-j}^* = \gamma_{n+1-j} + \bar{\gamma}_j q \quad \text{and} \quad |\gamma_j| + |\gamma_{n+1-j}| < \binom{n+1}{j},$$

for each $j = 1, \dots, \left\lfloor \frac{n+1}{2} \right\rfloor$. Consider $(\beta_1, \dots, \beta_{n-1}) \in \mathbb{C}^{n-1}$, where

$$\beta_j = \frac{n+1-j}{n+1} \gamma_j, \quad \beta_{n-j} = \frac{n+1}{n+1-j} \gamma_{n+1-j} \quad \text{for } j = 1, \dots, \frac{n-1}{2}.$$

Then

$$|\beta_j| + |\beta_{n-j}| = \frac{n+1-j}{n+1} (|\gamma_j| + |\gamma_{n+1-j}|) < \frac{n+1-j}{n+1} \binom{n+1}{j} = \binom{n}{j},$$

for each $j = 1, \dots, \left\lfloor \frac{n}{2} \right\rfloor$. Therefore,

$$(\beta_1 + \bar{\beta}_{n-1}q, \dots, \beta_j + \bar{\beta}_{n-j}q, \dots, \beta_{n-1} + \bar{\beta}_1q, q) \in \tilde{\mathbb{G}}_n$$

Note that,

$$\begin{aligned} \beta_j + \bar{\beta}_{n-j}q &= \frac{n+1-j}{n+1} (\gamma_j + \bar{\gamma}_{n+1-j}q) = \frac{n+1-j}{n+1} y_j^* = y_j, \\ \beta_{n-j} + \bar{\beta}_j q &= \frac{n+1-j}{n+1} (\gamma_{n+1-j} + \bar{\gamma}_j q) = \frac{n+1-j}{n+1} y_{n+1-j}^* = y_{n-j} \end{aligned}$$

for each $j = 1, \dots, \left\lfloor \frac{n}{2} \right\rfloor$. Therefore, $y = (y_1, \dots, y_{n-1}, q) \in \tilde{\mathbb{G}}_n$. The proof of $y \in \tilde{\Gamma}_n$ if and only if $\hat{y} \in \tilde{\Gamma}_{n+1}$ is similar. ■

Theorem 5.2. *Let $y = (y_1, \dots, y_{n-1}, q) \in \mathbb{C}^n$.*

- (1) *If n is even and $y \in \tilde{\mathbb{G}}_n$ (or, $\in \tilde{\Gamma}_n$), then $\check{y} \in \tilde{\mathbb{G}}_{n-1}$ (or, $\in \tilde{\Gamma}_{n-1}$), where*

$$\begin{aligned} \check{y} = \left(\frac{n-1}{n} y_1, \dots, \frac{n-j}{n} y_j, \dots, \frac{\frac{n}{2}+1}{n} y_{\frac{n}{2}-1}, \underbrace{\frac{\frac{n}{2}+1}{n} y_{\frac{n}{2}+1}}_{\frac{n}{2}\text{-th position}}, \right. \\ \left. \frac{\frac{n}{2}+2}{n} y_{\frac{n}{2}+2}, \dots, \underbrace{\frac{n-j}{n} y_{n-j}}_{(n-1-j)\text{-th position}}, \dots, \frac{n-1}{n} y_{n-1}, q \right). \end{aligned}$$

(2) If n is odd and $y \in \tilde{\mathbb{G}}_n$ (or, $\in \tilde{\Gamma}_n$), then $\tilde{y} \in \tilde{\mathbb{G}}_{n-1}$ (or, $\in \tilde{\Gamma}_{n-1}$), where

$$\tilde{y} = \left(\frac{n-1}{n}y_1, \dots, \frac{n-j}{n}y_j, \dots, \frac{n+3}{2n}y_{\frac{n-3}{2}}, \underbrace{\frac{n+1}{2n} \left(\frac{y_{\frac{n-1}{2}} + y_{\frac{n+1}{2}}}{2} \right)}_{\frac{n-1}{2}\text{-th position}}, \right. \\ \left. \frac{n+3}{2n}y_{\frac{n+3}{2}}, \dots, \underbrace{\frac{n-j}{n}y_{n-j}}_{(n-j-1)\text{-th position}}, \dots, \frac{n-1}{n}y_{n-1}, q \right).$$

Proof. (1). Let $n \in \mathbb{N}$ be even. Then, $\lfloor \frac{n-1}{2} \rfloor = \frac{n}{2} - 1$. Also it is merely mentioned that $\binom{n}{j} = \frac{n}{n-j} \binom{n-1}{j}$. Now suppose $y \in \tilde{\mathbb{G}}_n$. Then by Theorem 2.2, we have for each $j \in \left\{1, \dots, \lfloor \frac{n}{2} \rfloor\right\}$

$$\binom{n}{j} |y_j - \bar{y}_{n-j}q| + \left| y_j y_{n-j} - \binom{n}{j}^2 q \right| < \binom{n}{j}^2 - |y_{n-j}|^2.$$

Consider the point

$$\tilde{y} = (\tilde{y}_1, \dots, \tilde{y}_{n-2}, \tilde{q}) \\ = \left(\frac{n-1}{n}y_1, \dots, \frac{\frac{n}{2}+1}{n}y_{\frac{n}{2}-1}, \underbrace{\frac{\frac{n}{2}+1}{n}y_{\frac{n}{2}+1}}_{\frac{n}{2}\text{-th position}}, \frac{\frac{n}{2}+2}{n}y_{\frac{n}{2}+2}, \dots, \frac{n-1}{n}y_{n-1}, q \right).$$

Then $\tilde{q} = q$ and

$$\tilde{y}_j = \frac{n-j}{n}y_j, \quad \hat{y}_{n-1-j} = \frac{n-j}{n}y_{n-j} \quad \text{for } j = 1, \dots, \frac{n}{2} - 1.$$

Note that for each $j = 1, \dots, \frac{n}{2} - 1$, we have

$$|\tilde{y}_j - \bar{\tilde{y}}_{n-1-j}\tilde{q}| = \frac{n-j}{n} |y_j - \bar{y}_{n-j}q| \\ \text{and} \quad \left| \tilde{y}_j \tilde{y}_{n-1-j} - \binom{n-1}{j}^2 \tilde{q} \right| = \left(\frac{n-j}{n} \right)^2 \left| y_j y_{n-j} - \binom{n}{j}^2 q \right|.$$

Hence, for any $j = 1, \dots, \frac{n}{2} - 1$ ($= \lfloor \frac{n-1}{2} \rfloor$), we have

$$\binom{n-1}{j} |\tilde{y}_j - \bar{\tilde{y}}_{n-1-j}\tilde{q}| + \left| \tilde{y}_j \tilde{y}_{n-1-j} - \binom{n-1}{j}^2 \tilde{q} \right| \\ = \left(\frac{n-j}{n} \right)^2 \left[\binom{n}{j} |y_j - \bar{y}_{n-j}q| + \left| y_j y_{n-j} - \binom{n}{j}^2 q \right| \right]$$

$$\begin{aligned}
&< \left(\frac{n-j}{n}\right)^2 \left[\binom{n}{j}^2 - |y_{n-j}|^2 \right] \\
&= \binom{n-1}{j}^2 - |\check{y}_{n-1-j}|.
\end{aligned}$$

Therefore, by Theorem 2.2, we conclude that $\check{y} \in \tilde{\mathbb{G}}_{n-1}$. Similarly if $y \in \tilde{\Gamma}_n$, then $\check{y} \in \tilde{\Gamma}_{n-1}$.

(2). Suppose n is odd and let $y \in \tilde{\mathbb{G}}_n$. Then $|q| < 1$ and there exists $(\beta_1, \dots, \beta_{n-1}) \in \mathbb{C}^{n-1}$ such that

$$y_j = \beta_j + \bar{\beta}_{n-j}q, \quad y_{n-j} = \beta_{n-j} + \bar{\beta}_j q \quad \text{and} \quad |\beta_j| + |\beta_{n-j}| < \binom{n}{j},$$

for each $j \in \left\{1, \dots, \left\lfloor \frac{n}{2} \right\rfloor\right\}$. Consider the given point $\tilde{y} = (\tilde{y}_1, \dots, \tilde{y}_{n-2}, \tilde{q}) \in \mathbb{C}^{n-1}$. Then $\tilde{q} = q$,

$$\tilde{y}_{\frac{n-1}{2}} = \frac{n+1}{2n} \left(\frac{y_{\frac{n-1}{2}} + y_{\frac{n+1}{2}}}{2} \right),$$

$$\tilde{y}_j = \frac{n-j}{n} y_j \quad \text{and} \quad \tilde{y}_{n-j-1} = \frac{n-j}{n} y_{n-j}, \quad \text{for } j = 1, \dots, \frac{n-3}{2}.$$

Consider $(\gamma_1, \dots, \gamma_{n-2}) \in \mathbb{C}^{n-2}$, where $\gamma_{\frac{n-1}{2}} = \frac{n+1}{2n} \left(\frac{\beta_{\frac{n-1}{2}} + \beta_{\frac{n+1}{2}}}{2} \right)$,

$$\gamma_j = \frac{n-j}{n} \beta_j \quad \text{and} \quad \gamma_{n-1-j} = \frac{n-j}{n} \beta_{n-j}, \quad \text{for } j = 1, \dots, \frac{n-3}{2}.$$

Then, we have

$$2 \left| \gamma_{\frac{n-1}{2}} \right| = \frac{n+1}{2n} \left| \beta_{\frac{n-1}{2}} + \beta_{\frac{n+1}{2}} \right| < \frac{n - \frac{n-1}{2}}{n} \binom{n}{\frac{n-1}{2}} = \binom{n-1}{\frac{n-1}{2}}$$

$$\text{and} \quad |\gamma_j| + |\gamma_{n-1-j}| = \frac{n-j}{n} (|\beta_j| + |\beta_{n-j}|) < \frac{n-j}{n} \binom{n}{j} = \binom{n-1}{j},$$

for $j = 1, \dots, \frac{n-3}{2}$. Therefore,

$$(\gamma_1 + \bar{\gamma}_{n-2}q, \dots, \underbrace{\gamma_j + \bar{\gamma}_{n+1-j}q}_{j\text{-th position}}, \dots, \gamma_{n-2} + \bar{\gamma}_1 q, q) \in \tilde{\mathbb{G}}_{n-1}.$$

Also we have,

$$\begin{aligned}\gamma_{\frac{n-1}{2}} + \bar{\gamma}_{\frac{n-1}{2}}q &= \frac{n+1}{2n} \left(\frac{\beta_{\frac{n-1}{2}} + \bar{\beta}_{\frac{n+1}{2}}q + \beta_{\frac{n+1}{2}} + \bar{\beta}_{\frac{n-1}{2}}q}{2} \right) \\ &= \frac{n+1}{2n} \frac{(y_{\frac{n-1}{2}} + y_{\frac{n+1}{2}})}{2} = \tilde{y}_{\frac{n-1}{2}},\end{aligned}$$

$$\begin{aligned}\gamma_j + \bar{\gamma}_{n-1-j}q &= \frac{n-j}{n} (\beta_j + \bar{\beta}_{n-j}q) = \frac{n-j}{n} y_j = \tilde{y}_j \\ \text{and } \gamma_{n-1-j} + \bar{\gamma}_j q &= \frac{n-j}{n} (\beta_{n-j} + \bar{\beta}_j q) = \frac{n-j}{n} y_{n-j} = \tilde{y}_{n-j-1},\end{aligned}$$

for $j = 1, \dots, \frac{n-3}{2}$. Hence, $|\tilde{q}| < 1$ and there exists $(\gamma_1, \dots, \gamma_{n-2}) \in \mathbb{C}^{n-2}$ such that

$$\tilde{y}_j = \gamma_j + \bar{\gamma}_{n-1-j}\tilde{q}, \quad \tilde{y}_{n-j-1} = \gamma_{n-1-j} + \bar{\gamma}_j\tilde{q} \quad \text{and} \quad |\gamma_j| + |\gamma_{n-1-j}| < \binom{n-1}{j},$$

for each $j = 1, \dots, \frac{n-1}{2}$. Consequently $\tilde{y} = (\tilde{y}_1, \dots, \tilde{y}_{n-2}, \tilde{q}) \in \tilde{\mathbb{G}}_{n-1}$.

The proof of $y \in \tilde{\Gamma}_n$ implies $\tilde{y} \in \tilde{\Gamma}_{n-1}$ is similar. \blacksquare

Theorem 5.3. *Let $n \in \mathbb{N}$ be even.*

(1) *Let the point $y = (y_1, \dots, y_{n-1}, q) \in \tilde{\mathbb{G}}_n$ (or, $\in \tilde{\Gamma}_n$). Then the point*

$$\underline{y} = (y_1, \dots, y_{\frac{n}{2}}, y_{\frac{n}{2}}, \underbrace{y_{\frac{n}{2}+1}, \dots, y_{n-1}}_{(\frac{n}{2}+2)\text{-th}}, q) \in \tilde{\mathbb{G}}_{n+1} \text{ (or, } \in \tilde{\Gamma}_{n+1})$$

and the map $f : \tilde{\mathbb{G}}_n \rightarrow \tilde{\mathbb{G}}_{n+1}$ that maps y to \underline{y} is an analytic embedding.

(2) *Let $y = (y_1, \dots, y_n, q) \in \tilde{\mathbb{G}}_{n+1}$ (or, $\in \tilde{\Gamma}_{n+1}$). Then the point $\tilde{y} \in \tilde{\mathbb{G}}_n$ (or, $\in \tilde{\Gamma}_n$), where*

$$\begin{aligned}\tilde{y} = \left(\frac{n}{n+1}y_1, \dots, \frac{n+1-j}{n+1}y_j, \dots, \frac{\frac{n}{2}+2}{n+1}y_{\frac{n}{2}-1}, \frac{\frac{n}{2}+1}{2(n+1)}(y_{\frac{n}{2}} + y_{\frac{n}{2}+1}), \right. \\ \left. \underbrace{\frac{\frac{n}{2}+2}{n+1}y_{\frac{n}{2}+2}, \dots, \frac{n+1-j}{n+1}y_{n+1-j}, \dots, \frac{n}{n+1}y_n}_{(\frac{n}{2}+1)\text{-th}}, q \right).\end{aligned}$$

The map $g : \tilde{\mathbb{G}}_{n+1} \rightarrow \tilde{\mathbb{G}}_n$ that maps y to \tilde{y} is analytic.

(3) Let $y = (y_1, \dots, y_n, q) \in \tilde{\mathbb{G}}_{n+1}$ (or, $\in \tilde{\Gamma}_{n+1}$). Then the point

$$\hat{y} = \left(\frac{y_1}{\binom{n+1}{1}}, \dots, \frac{y_j}{\binom{n+1}{j}}, \dots, \frac{y_n}{\binom{n+1}{1}}, q \right) \in \tilde{\mathbb{G}}_{n+1} \text{ (or, } \in \tilde{\Gamma}_{n+1})$$

and the point

$$\hat{y}_{\sharp} = \left(\hat{y}_1, \dots, \hat{y}_{\frac{n}{2}-1}, \underbrace{\frac{\hat{y}_{\frac{n}{2}} + \hat{y}_{\frac{n}{2}+1}}{2}}_{\frac{n}{2}-th}, \hat{y}_{\frac{n}{2}+2}, \dots, \hat{y}_n, q \right) \in \tilde{\mathbb{G}}_n \text{ (or, } \in \tilde{\Gamma}_n),$$

where $\hat{y}_j = \frac{y_j}{\binom{n+1}{j}}$ for each j . Also the function $h : \tilde{\mathbb{G}}_{n+1} \rightarrow \tilde{\mathbb{G}}_n$ that maps y to \hat{y}_{\sharp} is analytic.

Proof.

(1) . Since $y = (y_1, \dots, y_{n-1}, q) \in \tilde{\mathbb{G}}_n$, there exists a unique $(\beta_1, \dots, \beta_{n-1}) \in \mathbb{C}^{n-1}$ such that

$$y_j = \beta_j + \bar{\beta}_{n-j}q, \quad y_{n-j} = \beta_{n-j} + \bar{\beta}_j q \quad \text{and} \quad |\beta_j| + |\beta_{n-j}| < \binom{n}{j},$$

for each $j = 1, \dots, \frac{n}{2}$. Note that $y_{\frac{n}{2}} = \beta_{\frac{n}{2}} + \bar{\beta}_{\frac{n}{2}}q$. Consider the given point $\underline{y} = (\underline{y}_1, \dots, \underline{y}_n, \underline{q}) \in \mathbb{C}^{n+1}$. Then $\underline{q} = q$,

$$\underline{y}_j = y_j \quad \text{and} \quad \underline{y}_{n+1-j} = y_{n-j} \quad \text{for } j = 1, \dots, \frac{n}{2}.$$

Define $\gamma_j = \beta_j$ and $\gamma_{n+1-j} = \beta_{n-j}$ for $j = 1, \dots, \frac{n}{2}$. Then,

$$\gamma_1 = \beta_1, \dots, \gamma_{\frac{n}{2}} = \beta_{\frac{n}{2}}, \quad \gamma_{\frac{n}{2}+1} = \beta_{\frac{n}{2}}, \quad \gamma_{\frac{n}{2}+2} = \beta_{\frac{n}{2}+1}, \dots, \gamma_n = \beta_{n-1}.$$

Evidently,

$$|\gamma_j| + |\gamma_{n+1-j}| = |\beta_j| + |\beta_{n-j}| < \binom{n}{j} \leq \binom{n+1}{j}, \quad \text{for } j = 1, \dots, \frac{n}{2}.$$

For $j = 1, \dots, \frac{n}{2}$, we have

$$\gamma_j + \bar{\gamma}_{n+1-j}\underline{q} = y_j = \underline{y}_j \quad \text{and} \quad \gamma_{n+1-j} + \bar{\gamma}_j \underline{q} = y_{n-j} = \underline{y}_{n+1-j}.$$

Hence, there exists $(\gamma_1, \dots, \gamma_n) \in \mathbb{C}^n$ such that

$$\underline{y}_j = \gamma_j + \bar{\gamma}_{n+1-j}\underline{q}, \quad \underline{y}_{n+1-j} = \gamma_{n+1-j} + \bar{\gamma}_j \underline{q} \quad \text{and} \quad |\gamma_j| + |\gamma_{n+1-j}| < \binom{n+1}{j},$$

for each $j = 1, \dots, \left[\frac{n+1}{2}\right]$. Hence $y \in \tilde{\mathbb{G}}_{n+1}$. The map f is clearly an analytic embedding of $\tilde{\mathbb{G}}_n$ into $\tilde{\mathbb{G}}_{n+1}$.

(2) Let $y = (y_1, \dots, y_n, q) \in \widetilde{\mathbb{G}}_{n+1}$. Then, there exists $(\beta_1, \dots, \beta_n) \in \mathbb{C}^n$ such that

$$y_j = \beta_j + \bar{\beta}_{n+1-j}q, \quad y_{n+1-j} = \beta_{n+1-j} + \bar{\beta}_j q \quad \text{and} \quad |\beta_j| + |\beta_{n+1-j}| < \binom{n+1}{j},$$

for $j = 1, \dots, n$. Consider the point $\tilde{y} = (\tilde{y}_1, \dots, \tilde{y}_{n-1}, \tilde{q}) \in \mathbb{C}^n$, where $\tilde{q} = q$, $\tilde{y}_{\frac{n}{2}} = \frac{\frac{n}{2} + 1}{2(n+1)} (y_{\frac{n}{2}} + y_{\frac{n}{2}+1})$ and

$$\tilde{y}_j = \frac{n+1-j}{n+1} y_j, \quad \tilde{y}_{n-j} = \frac{n+1-j}{n+1} y_{n+1-j} \quad \text{for } j = 1, \dots, \frac{n}{2} - 1.$$

Define $(\gamma_1, \dots, \gamma_{n-1}) \in \mathbb{C}^{n-1}$ as follows

$$\gamma_j = \frac{n+1-j}{n+1} \beta_j, \quad \gamma_{n-j} = \frac{n+1-j}{n+1} \beta_{n+1-j} \quad \text{for } j = 1, \dots, \frac{n}{2} - 1$$

$$\text{and } \gamma_{\frac{n}{2}} = \frac{n+1-\frac{n}{2}}{2(n+1)} (\beta_{\frac{n}{2}} + \beta_{\frac{n}{2}+1}) = \frac{n+2}{4(n+1)} (\beta_{\frac{n}{2}} + \beta_{\frac{n}{2}+1}).$$

Then for $j = 1, \dots, \frac{n}{2} - 1$, we have

$$|\gamma_j| + |\gamma_{n-j}| = \frac{n+1-j}{n+1} (|\beta_j| + |\beta_{n+1-j}|) < \frac{n+1-j}{n+1} \binom{n+1}{j} = \binom{n}{j}$$

$$\text{and } 2|\gamma_{\frac{n}{2}}| \leq \frac{n+2}{2(n+1)} (|\beta_{\frac{n}{2}}| + |\beta_{\frac{n}{2}+1}|) < \frac{n+1-\frac{n}{2}}{(n+1)} \binom{n+1}{\frac{n}{2}} = \binom{n}{\frac{n}{2}}.$$

Therefore, $(\gamma_1, \dots, \gamma_{n-1}) \in \mathbb{C}^{n-1}$ where $|\gamma_j| + |\gamma_{n-j}| < \binom{n}{j}$, for all $j = 1, \dots, \frac{n}{2}$. Also

$$\tilde{y}_j = \frac{n+1-j}{n+1} y_j = \frac{n+1-j}{n+1} (\beta_j + \bar{\beta}_{n+1-j}q) = \gamma_j + \bar{\gamma}_{n-j}\tilde{q},$$

$$\tilde{y}_{n-j} = \frac{n+1-j}{n+1} y_{n+1-j} = \frac{n+1-j}{n+1} (\beta_{n+1-j} + \bar{\beta}_j q) = \gamma_{n-j} + \bar{\gamma}_j \tilde{q},$$

for $j = 1, \dots, \frac{n}{2} - 1$. Since $y_{\frac{n}{2}} = \beta_{\frac{n}{2}} + \bar{\beta}_{\frac{n}{2}+1}q$ and $y_{\frac{n}{2}+1} = \beta_{\frac{n}{2}+1} + \bar{\beta}_{\frac{n}{2}}q$, we have that

$$\begin{aligned} \tilde{y}_{\frac{n}{2}} &= \frac{n+2}{4(n+1)} (y_{\frac{n}{2}} + y_{\frac{n}{2}+1}) \\ &= \frac{n+2}{4(n+1)} [(\beta_{\frac{n}{2}} + \beta_{\frac{n}{2}+1}) + (\bar{\beta}_{\frac{n}{2}} + \bar{\beta}_{\frac{n}{2}+1})q] \\ &= \gamma_{\frac{n}{2}} + \bar{\gamma}_{\frac{n}{2}}\tilde{q}. \end{aligned}$$

Thus, there exists $(\gamma_1, \dots, \gamma_{n-1}) \in \mathbb{C}^{n-1}$ such that

$$\tilde{y}_j = \gamma_j + \bar{\gamma}_{n-j}\tilde{q}, \quad \tilde{y}_{n-j} = \gamma_{n-j} + \bar{\gamma}_j\tilde{q} \quad \text{and} \quad |\gamma_j| + |\gamma_{n-j}| < \binom{n}{j},$$

for all $j = 1, \dots, \frac{n}{2}$. Hence $\tilde{y} \in \tilde{\mathbb{G}}_n$. The map g is obviously analytic.

(3) Let $y = (y_1, \dots, y_n, q) \in \tilde{\mathbb{G}}_{n+1}$. Then there exists $(\beta_1, \dots, \beta_n) \in \mathbb{C}^n$ such that

$$y_j = \beta_j + \bar{\beta}_{n+1-j}q, \quad y_{n+1-j} = \beta_{n+1-j} + \bar{\beta}_j q \quad \text{and} \quad |\beta_j| + |\beta_{n+1-j}| < \binom{n+1}{j},$$

for $j = 1, \dots, n$. Consider the point $\hat{y} = (\hat{y}_1, \dots, \hat{y}_n, \hat{q})$, where $\hat{q} = q$, and

$$\hat{y}_j = \frac{y_j}{\binom{n+1}{j}} \quad \text{for all } j = 1, \dots, n. \quad \text{Let } \alpha_j = \frac{\beta_j}{\binom{n+1}{j}} \quad \text{for } j = 1, \dots, n. \quad \text{Then}$$

$$|\alpha_j| + |\alpha_{n+1-j}| = \frac{|\beta_j| + |\beta_{n+1-j}|}{\binom{n+1}{j}} < 1 \leq \binom{n+1}{j}.$$

Also we have

$$\hat{y}_j = \frac{y_j}{\binom{n+1}{j}} = \alpha_j + \bar{\alpha}_{n+1-j}q, \quad \hat{y}_{n+1-j} = \frac{y_{n+1-j}}{\binom{n+1}{j}} = \alpha_{n+1-j} + \bar{\alpha}_j q,$$

for $j = 1, \dots, n$. Thus, $\hat{y} \in \tilde{\mathbb{G}}_{n+1}$. Next define $(\gamma_1, \dots, \gamma_{n-1}) \in \mathbb{C}^{n-1}$ in the following way:

$$\gamma_{\frac{n}{2}} = \frac{(\alpha_{\frac{n}{2}} + \alpha_{\frac{n}{2}+1})}{2}, \quad \gamma_j = \alpha_j \quad \text{and} \quad \gamma_{n-j} = \alpha_{n+1-j}$$

for $j = 1, \dots, \frac{n}{2} - 1$. Then, we have

$$2|\gamma_{\frac{n}{2}}| \leq |\beta_{\frac{n}{2}}| + |\beta_{\frac{n}{2}+1}| < 1 \leq \binom{n}{\frac{n}{2}}$$

and $|\gamma_j| + |\gamma_{n-j}| = |\alpha_j| + |\alpha_{n+1-j}| < 1 \leq \binom{n}{j} \quad \text{for } j = 1, \dots, \frac{n}{2} - 1.$

Therefore, $(\gamma_1, \dots, \gamma_{n-1}) \in \mathbb{C}^{n-1}$ where $|\gamma_j| + |\gamma_{n-j}| < \binom{n}{j}$, for all $j = 1, \dots, \frac{n}{2}$. Note that,

$$\begin{aligned} \hat{y}_j &= \gamma_j + \bar{\gamma}_{n-j}q \\ \hat{y}_{n-j} &= \gamma_{n-j} + \bar{\gamma}_j q \quad \text{for } j = 1, \dots, \frac{n}{2} - 1. \end{aligned}$$

Since $\hat{y}_{\frac{n}{2}} = \alpha_{\frac{n}{2}} + \bar{\alpha}_{\frac{n}{2}+1}q$ and $\hat{y}_{\frac{n}{2}+1} = \alpha_{\frac{n}{2}+1} + \bar{\alpha}_{\frac{n}{2}}q$, we also have

$$\begin{aligned} \frac{\hat{y}_{\frac{n}{2}} + \hat{y}_{\frac{n}{2}+1}}{2} &= \frac{(\alpha_{\frac{n}{2}} + \alpha_{\frac{n}{2}+1})}{2} + \frac{(\bar{\alpha}_{\frac{n}{2}} + \bar{\alpha}_{\frac{n}{2}+1})}{2}q \\ &= \gamma_{\frac{n}{2}} + \bar{\gamma}_{\frac{n}{2}}q. \end{aligned}$$

Hence, the point

$$\hat{y}_{\sharp} = \left(\hat{y}_1, \dots, \hat{y}_{\frac{n}{2}-1}, \frac{\hat{y}_{\frac{n}{2}} + \hat{y}_{\frac{n}{2}+1}}{2}, \hat{y}_{\frac{n}{2}+2}, \dots, \hat{y}_n, q \right) \in \tilde{\mathbb{G}}_n.$$

Clearly, the map h is analytic. ■

Theorem 5.4. *Let $n \in \mathbb{N}$ be odd.*

(1) *Let the point $y = (y_1, \dots, y_{n-1}, q) \in \tilde{\mathbb{G}}_n$ (or, $\in \tilde{\Gamma}_n$). Then the point*

$$\underline{y} = \left(y_1, \dots, y_{[\frac{n}{2}]}, \frac{y_{[\frac{n}{2}]} + y_{[\frac{n}{2}]+1}}{2}, \underbrace{y_{\frac{n}{2}+1}}_{(\frac{n}{2}+2)\text{-th}}, \dots, y_{n-1}, q \right) \in \tilde{\mathbb{G}}_{n+1} \text{ (or, } \in \tilde{\Gamma}_{n+1})$$

and the map $f : \tilde{\mathbb{G}}_n \rightarrow \tilde{\mathbb{G}}_{n+1}$ that maps y to \underline{y} is analytic.

(2) *Let $y = (y_1, \dots, y_n, q) \in \tilde{\mathbb{G}}_{n+1}$ (or, $\in \tilde{\Gamma}_{n+1}$). Then the point $\tilde{y} \in \tilde{\mathbb{G}}_n$ (or, $\in \tilde{\Gamma}_n$), where*

$$\tilde{y} = \left(\frac{n}{n+1}y_1, \dots, \frac{n+1-j}{n+1}y_j, \dots, \frac{n+3}{2(n+1)}y_{[\frac{n}{2}]}, \right. \\ \left. \underbrace{\frac{n+3}{2(n+1)}y_{[\frac{n}{2}]+2}, \dots, \frac{n+1-j}{n+1}y_{n+1-j}, \dots, \frac{n}{n+1}y_n, q}_{([\frac{n}{2}]+1)\text{-th}} \right).$$

The map $g : \tilde{\mathbb{G}}_{n+1} \rightarrow \tilde{\mathbb{G}}_n$ that maps y to \tilde{y} is an analytic embedding.

(3) *Let $y = (y_1, \dots, y_n, q) \in \tilde{\mathbb{G}}_{n+1}$ (or, $\in \tilde{\Gamma}_{n+1}$). Then the point*

$$\hat{y} = \left(\frac{y_1}{\binom{n+1}{1}}, \dots, \frac{y_j}{\binom{n+1}{j}}, \dots, \frac{y_n}{\binom{n+1}{1}}, q \right) \in \tilde{\mathbb{G}}_{n+1} \text{ (or, } \in \tilde{\Gamma}_{n+1})$$

and the point

$$\hat{y}_{\sharp} = \left(\hat{y}_1, \dots, \hat{y}_{[\frac{n}{2}]-1}, \underbrace{\hat{y}_{[\frac{n}{2}]}}_{\frac{n}{2}\text{-th}}, \hat{y}_{[\frac{n}{2}]+2}, \dots, \hat{y}_n, q \right) \in \tilde{\mathbb{G}}_n \text{ (or, } \in \tilde{\Gamma}_n),$$

where $\hat{y}_j = \frac{y_j}{\binom{n+1}{j}}$. The map $h : \tilde{\mathbb{G}}_{n+1} \rightarrow \tilde{\mathbb{G}}_n$ that maps y to \hat{y}_{\sharp} is an analytic embedding.

Proof. The proof is similar to that of Theorem 5.3. ■

REFERENCES

- [1] A. A. Abouhajar, M. C. White and N. J. Young, A Schwarz lemma for a domain related to μ -synthesis, *J. Geom. Anal.* 17 (2007), 717 – 750.
- [2] J. Agler, Z. A. Lykova and N. J. Young, Extremal holomorphic maps and the symmetrized bidisc, *Proc. Londo Math. Soc.*, 106 (2013), 781 – 818.
- [3] J. Agler, Z. A. Lykova and N. J. Young, Algebraic and geometric aspects of rational -inner functions, *Adv. Math.*, 328 (2018), 133 – 159.
- [4] J. Agler and N. J. Young, A commutant lifting theorem for a domain in \mathbb{C}^2 and spectral interpolation, *J. Funct. Anal.* 161 (1999), 452 – 477.
- [5] J. Agler and N. J. Young, A schwarz lemma for the symmetrized bidisc, *Bull. London Math. Soc.*, 33 (2001), 175 – 186.
- [6] J. Agler and N. J. Young, The hyperbolic geometry of the symmetrized bidisc, *J. Geom. Anal.* 14 (2004), no.3, 375 – 403.
- [7] G. Bharali, Some new observations on interpolation in the spectral unit ball, *Integral Equa. Oper. Theory*, 59 (2007), 329 – 343.
- [8] G. Bharali, A family of domains associated with μ -synthesis, *Integral Equa. Oper. Theory* 82 (2015), 267 – 285.
- [9] T. Bhattacharyya, S. Pal and S. Shyam Roy, Dilations of Γ -contractions by solving operator equations, *Adv. Math.*, 230 (2012), 577 – 606.
- [10] T. Bhattacharyya and S. Pal, A functional model for pure Γ -contractions, *J. Operator Thoery*, 71 (2014), 327 – 339.
- [11] T. Bhattacharyya and Haripada Sau, Holomorphic functions on the symmetrized bidisk – realization, interpolation and extension., *J. Funct. Anal.*, 274 (2018), 504 – 524.
- [12] S. Biswas and S. Shyam Roy, Functional models for Γ_n -contractions and characterization of Γ_n -isometries, *J. Func. Anal.*, 266 (2014), 6224 – 6255.
- [13] C. Costara, The symmetrized bidisc and Lempert’s theorem, *Bull. London Math. Soc.*, 36 (2004) 656 – 662.
- [14] C. Costara, On the spectral Nevanlinna-Pick problem, *Studia Math.*, 170 (2005), 23 – 55.
- [15] J. Doyle, Analysis of feedback systems with structured uncertainties, *IEEE Proceedings D, Control Theory and Applications*, 129 (1982), 242 – 250.
- [16] A. Edigarian, A note on C. Costara’s paper: ”The symmetrized bidisc and Lempert’s theorem” *Ann. Polon. Math*, 83 (2004), no. 2, 189 – 191
- [17] The Lempert theorem and the tetrablock, *J. Geom. Anal.*, 23 (2013), 1818 – 1831.
- [18] A. Edigarian and W. Zwonek, Geometry of the symmetrized polydisc, *Arch. Math. (Basel)*, 84 (2005), 364 – 374.
- [19] L. Kosiński and W. Zwonek, Extremal holomorphic maps in special classes of domains, *Annals of the Scuola Normale Superiore di Pisa - Science Class*, doi: 10.2422/2036-2145.201401 003.
- [20] N. Nikolov, The symmetrized polydisc cannot be exhausted by domains biholomorphic to convex domains, *Ann. Pol. Math.* 88 (2006) , 279 – 283.
- [21] N. Nikolov, P. Pflug, P. J. Thomas and W. Zwonek, Estimates of the Carathodory metric on the symmetrized polydisc, *J. Math. Anal. Appl.* 341 (2008), 140 – 148.

- [22] N. Nikolov, P. Pflug and W. Zwonek, The Lempert function of the symmetrized polydisc in higher dimensions is not a distance, *Proc. Amer. Math. Soc.*, 135 (2007), 2921 – 2928.
- [23] N. Nikolov, P. J. Thomas and T. Duc-Anh, Lifting maps from the symmetrized polydisc in small dimensions, *Complex Anal. Oper. Theory*, 10 (2016), 921 – 941.
- [24] N. Nikolov and W. Zwonek, The Bergman kernel of the symmetrized polydisc in higher dimensions has zeros, *Arch. Math. (Basel)*, 87 (2006), 412 – 416.
- [25] S. Pal, From Stinespring dilation to Sz.-Nagy dilation on the symmetrized bidisc and operator models, *New York Jour. Math.*, 20 (2014), 545 – 564.
- [26] S. Pal, Canonical decomposition of operators associated with the symmetrized polydisc, *Complex Anal. Oper. Theory*, 12 (2018), 931 – 943.
- [27] S. Pal, A Nagy-Foias program for C_0 operator tuples associated with the symmetrized polydisc, *Preprint*.
- [28] S. Pal and S. Roy, The complex geometry and a two-point interpolation problem for two domains in \mathbb{C}^n , preprint, available at <https://arxiv.org/abs/1904.03745>.
- [29] S. Pal and O. M. Shalit, Spectral sets and distinguished varieties in the symmetrized bidisc, *J. Funct. Anal.*, 266 (2014), 5779 - 5800.
- [30] S. Parrott, On a quotient norm and the Sz.-Nagy-Foias lifting theorem, *J. Funct. Anal.*, 30 (1978), 311 – 328.
- [31] P. Pflug and W. Zwonek, Exhausting domains of the symmetrized bidisc, *Ark. Mat.*, 50 (2012), 397 – 402.
- [32] P. Pflug and W. Zwonek, Description of all complex geodesics in the symmetrized bidisc, *Bull. London Math. Soc.*, 37 (2005), 575 – 584.
- [33] P. Thomas, N. V. Trao, W. Zwonek, Green functions of the spectral ball and symmetrized polydisk, *J. Math. Anal. Appl.*, 377 (2011), 624 – 630.
- [34] N. J. Young, An introduction to Hilbert space (Cambridge University Press, Cambridge, 1988).
- [35] N. J. Young, The automorphism group of the tetrablock, *J. Lond. Math. Soc.(2)*, 77 (2008), 757 – 770.

(Sourav Pal) MATHEMATICS DEPARTMENT, INDIAN INSTITUTE OF TECHNOLOGY BOMBAY, POWAI, MUMBAI - 400076, INDIA.

E-mail address: `sourav@math.iitb.ac.in`

(Samriddho Roy) MATHEMATICS DEPARTMENT, INDIAN INSTITUTE OF TECHNOLOGY BOMBAY, POWAI, MUMBAI - 400076, INDIA.

E-mail address: `samriddhoroy@gmail.com`