Tautological Rings of Fibrations

Nils Prigge

Abstract. We study the analogue of tautological rings of fibre bundles in the context of fibrations with Poincaré fibre, i.e. the ring obtained by fibre integrating powers of the fibrewise Euler class. We approach this ring of characteristic classes using rational homotopy theory. More specifically, we discuss a cdga model for the universal fibration and chain level representatives of fibre integration and the fibrewise Euler class. This is particularly useful for rationally elliptic spaces and as an example we compute the tautological ring for complex projective spaces and products of odd spheres.

Contents

1			2
	1.1	Algebraic models of fibrations	3
	1.2	Models of fibre integration and fibrewise Euler class	3
	1.3	Computational results	
2	The	dg coalgebra model of the universal fibration	5
	2.1	Preliminaries and conventions	6
	2.2	Classification of coalgebra bundles	7
	2.3	Model and comparison with Tanré	
	2.4	The cdga model of the universal 1-connected fibration	15
3	Fibre integration in rational homotopy theory		17
	3.1	ϕ -integration	17
	3.2	Chain level ϕ -integration	
	3.3	Relation to parametrized stable homotopy theory	19
4			20
	4.1	Leray–Hirsch fibrations	22
5	Computations and observations		23
		The Euler ring of complex projective space	24
		The Euler ring of products of odd spheres	

1 Introduction

Consider a fibration $X \hookrightarrow E \xrightarrow{\pi} B$ where all spaces are Poincaré duality spaces and assume it is oriented, i.e. that $H^d(X)$ is a trivial $\pi_1(B)$ -module where d is the formal dimension of X. Then the diagonal $\Delta : E \to E \times_B E$ has an Umkehr map $\Delta_! = D_{E \times_B E}^{-1} \Delta_* D_E : H^*(E) \to H^{*+d}(E \times_B E)$ where D_E and $D_{E \times_B E}$ denote the corresponding Poincaré duality isomorphisms. We can define the **fibrewise Euler class** as

$$e^{fW}(\pi) := \Delta^* \Delta_!(1) \in H^d(E). \tag{1.1}$$

This definition can be extended to to all oriented (Hurewicz) fibrations with Poincaré fibre *X* [HLLR17]. In particular, one can associate a fibrewise Euler class to the universal oriented *X*-fibration

$$X \hookrightarrow BhAut_{*}^{+}(X) \xrightarrow{\pi} BhAut^{+}(X),$$
 (1.2)

which classifies oriented fibrations with fibre X [Sta63, May75]. Here, hAut $^+(X)$ denotes the topological monoid of orientation preserving homotopy automorphism of X and hAut $^+(X)$ is the submonoid of homotopy automorphisms fixing a base point.

If we assume that X is an oriented manifold and $\pi: E \rightarrow B$ is an oriented fibre bundle, then the fibrewise Euler class agrees with the Euler class of the vertical tangent bundle $T_{\pi}E \rightarrow E$ because this vector bundle is equivalent to a regular neighbourhood of $E \hookrightarrow E \times_B E$. Hence, the characteristic classes obtained by fibre integrating powers of the fibrewise Euler class agree with the generalized Miller-Morita-Mumford classes $\kappa_{e^{i+1}} := \pi_!(e^{i+1}(T_{\pi}E)) \in H^{i\cdot d}(B)$, where $\pi_!: H^*(E) \rightarrow H^{*-d}(B)$ denotes fibre integration. These classes can be defined for fibre bundles for all $c \in H^*(B\operatorname{SO}(d))$ analogously as $\kappa_c := \pi_!(c(T_{\pi}E))$ and the subring of $H^*(B\operatorname{Diff}^+(X))$ generated by the generalized MMM-classes of the universal X-bundle is called the tautological ring. It has been originally introduced for surfaces [Mum83] and more recently for higher dimensional manifolds [GGRW17, Gri17]. In both cases, it has been an area of fruitful research and motivated by progress on tautological rings of high-dimensional manifolds [RW18, GGRW17, Gri17] we want to study the tautological ring of oriented fibrations with Poincaré fibre, obtained by fibre integrating powers of the fibrewise Euler class.

A rational oriented Poincaré duality space of formal dimension d is a pair (X, ε_X) where X is a connected space and $\varepsilon_X : H^d(X; \mathbb{Q}) \to \mathbb{Q}$ is a linear map such that $(H^*(X; \mathbb{Q}), \varepsilon_X)$ is an oriented Poincaré duality algebra, i.e. the pairing

$$H^q(X;\mathbb{Q}) \otimes H^{d-q}(X;\mathbb{Q}) \xrightarrow{\smile} H^d(X;\mathbb{Q}) \xrightarrow{\varepsilon_X} \mathbb{Q}$$

is non-degenerate for all *q*.

Definition 1.1. Let X be a rational oriented Poincaré duality space of formal dimension d. The **Euler ring** $E^*(X)$ is the subring of $H^*(B hAut^+(X); \mathbb{Q})$ generated by the generalized MMM-classes

$$\kappa_i := \pi_!(e^{\mathrm{fw}}(\pi)^{i+1}) \in H^{id}(\mathrm{B}\,\mathrm{hAut}^+(X);\mathbb{Q}),$$

where $e^{\text{fw}}(\pi) \in H^d(B \text{ hAut}^+_*(X); \mathbb{Q})$ is the fibrewise Euler class of the universal oriented *X*-fibration.

One can also define the Euler ring $E^*_{CAT}(M)$ for smooth or topological fibre bundles (or the corresponding block versions) for $M \in CAT$ as the subring of $H^*(BCAT^+(M))$ which is the image of the Euler ring under the pull back along $BCAT^+(M) \rightarrow BhAut^+(M)$.

1.1 Algebraic models of fibrations. There are essentially two different algebraic approaches to rational homotopy theory, using either the Quillen equivalence of the category of 1-connected rational spaces of finite type with the (opposite) category of 1-connected commutative differential graded algebras, or with the category of connected differential graded Lie algebras of finite type. These two algebraic categories are Quillen equivalent with right adjoint given by the the cdga of Chevalley-Eilenberg cochains $C^*_{CE}(-;\mathbb{Q})$ of a dg Lie algebra. Since B hAut $^+_{(*)}(X)$ is not 1-connected in general, we start by studying the universal 1-connected X-fibration

$$X \hookrightarrow B hAut_{0,*}(X) \xrightarrow{\pi} B hAut_0(X),$$
 (1.3)

for a simply connected space X, where $hAut_0(X)$ and $hAut_{0,*}(X)$ denote the respective connected components containing the identity. Sullivan proved in [Sul77] that for a 1-connected space of finite type with minimal Sullivan model $(\Lambda V, d)$ the dg Lie algebra $Der^+(\Lambda V)$ of derivations is a Lie model of $BhAut_0(X)$. However, in order to compute the Euler ring we need to know an algebraic model of the map $BhAut_{0,*}(X) \to BhAut_0(X)$. Tanré gave such a description in [Tan83] in the category of dg Lie algebras. More precisely, let $L = (\mathbb{L}V, d)$ be a minimal dg Lie algebra model of X. Then the adjoint ad : $L \to Der^+(L)$ is a dg Lie algebra homomorphism and the mapping cone $Der^+(L)$ // ad L can be given the structure of a dg Lie algebra. Tanré showed that isomorphism classes of semidirect products of dg Lie algebras $L \rtimes L'$ are in a 1-1 correspondence with maps $\varphi: L' \to Der^+(L)$ // ad L, and that the semidirect product corresponding to the identity

$$L \longrightarrow L \rtimes_{\mathrm{id}} (\mathrm{Der}^+(L) \ /\!\!/ \ \mathrm{ad} \ L) \longrightarrow \mathrm{Der}^+(L) \ /\!\!/ \ \mathrm{ad} \ L$$
 (1.4)

is a dg Lie algebra model of (1.3). We can apply $C_{CE}^*(-;\mathbb{Q})$ to this sequence to get a cdga model of (1.3). Our first result is an analogous description of (1.3) in the category of cdga's in terms of a Sullivan model of X.

Theorem A. Let X be a 1-connected space of finite type with minimal Sullivan model $(\Lambda V, d)$ and unit $\eta : \mathbb{Q} \rightarrow \Lambda V$. Then

$$C_{CE}^*(\operatorname{Der}^+(\Lambda V); \mathbb{Q}) \xrightarrow{\eta_*} C_{CE}^*(\operatorname{Der}^+(\Lambda V); \Lambda V)$$
 (1.5)

is a cdga model of the universal oriented 1-connected fibration (1.3).

This result is known to experts, but we do not know a reference for it. It is an analogue of Tanré's result, in that it is based on a classification of the algebraic model of a fibration in the (opposite) category of cdga's with given fibre–or rather in the equivalent category of cocommutative dg coalgebras.

We remark that Theorem A turns out to be particularly useful for rationally elliptic spaces, because the algebraic model of the universal 1-connected fibration in Theorem A is much smaller than Tanré's model.

1.2 Models of fibre integration and fibrewise Euler class. Let $\pi: E \rightarrow B$ be a 1-connected fibration with Poincaré fibre X of dimension d. Then fibre integration $\pi_!: H^*(E) \rightarrow H^{*-d}(B)$ is an $H^*(B)$ -module map. By abuse of notation, we denote by $\pi^*: B \rightarrow E$ a cdga model of the fibration. We want to find a B-module homomorphism $\Pi: E \rightarrow B$ that induces $\pi_!$ on

cohomology. We show that there is a unique "derived" chain homotopy class $[\Pi]$ inducing fibre integration, and that it is determined by the orientation ε_X . More precisely, let $B \to E' = (B \otimes \Lambda V, D)$ be a relative Sullivan model of π^* and assume that B and E' are simply connected. Then $E' \otimes_B \mathbb{Q} \cong (\Lambda V, d)$ is a cdga model of X and the orientation ε_X has a representative in $H^{-d}(\operatorname{Hom}(\Lambda V, \mathbb{Q})) \cong \operatorname{Hom}(H^d(X; \mathbb{Q}), \mathbb{Q})$ which we denote the same. We show that there is a unique chain homotopy class of B-module chain maps $\Pi : E' \to B$ satisfying $\Pi(1 \otimes \chi) = \varepsilon_X(\chi)$ for all $\chi \in (\Lambda V)^d$.

Knowing $\Pi: E' \to B$ enables us to find a representative $\Delta_!: E' \to E' \otimes_B E'[-d]$ of the Umkehr map in the homotopy category of B-modules, which we need to determine a representative of the fibrewise Euler class in E'. Denote the shift of a B-module M by M[n]. We prove that Π induces a quasi-isomorphism of B-modules $\bar{\Pi}: E'[-d] \to \operatorname{Hom}_B(E', B)$ which sends $e \in E'$ to the morphism $e' \mapsto (-1)^{d+d\cdot|e|}\Pi(e\cdot e')$, and that the Umkehr map is determined in the homotopy category of B-modules by the diagram

$$E'[-d] \xrightarrow{\Delta_!} E' \otimes_B E'[-2d]$$

$$\bar{\Pi} \downarrow \simeq \qquad \qquad \overline{\Pi \otimes \Pi} \downarrow \simeq$$

$$Hom_B(E', B) \xrightarrow{\Delta^*} Hom_B(E' \otimes_B E', B)$$

where $\Delta : E' \otimes E' \to E'$ denotes the multiplication. Finally, we show that $\Delta^* \Delta_![1] \in H^d(E')$ is a representative of the fibrewise Euler class defined in [HLLR17], so that we can compute the Euler ring in terms of the algebraic models from rational homotopy theory.

1.3 Computational results. We focus on two classes of elliptic spaces: complex projective spaces and products of odd dimensional simply connected spheres. It was pointed out to us by Alexander Berglund after completion of this work that the following model of the universal 1-connected fibration for complex projective space was already determined in [Kur10, Prop.1.3] with different methods.

Proposition B. A cdga model of the universal 1-connected $\mathbb{C}P^n$ -fibration is given by

$$B_n := \left(\mathbb{Q}[x_2, \dots, x_{n+1}], |x_i| = 2i, d = 0 \right) \xrightarrow{\pi} E_n := \left(B_n[x] / (x^{n+1} - \sum_{i=2}^{n+1} x_i \cdot x^{n+1-i}), |x| = 2, d = 0 \right).$$

In particular, the universal fibration is formal.

We observe that the universal 1-connected $\mathbb{C}P^n$ -fibration is Leray–Hirsch. This simplifies the description of fibre integration and the Euler class as described in Section 4.1. But actually, we can exploit that E_n is a complete intersection over B_n to determine a representative of the fibrewise Euler class.

We show that a complete intersection $E = B[x_1, ..., x_n]/(f_1, ..., f_n)$ over a polynomial ring B in positive even degrees corresponds to a fibration whose fibre X is a rational Poincaré space and that the fibrewise Euler class, with a suitable choice of orientation $\varepsilon_X : H^d(X; \mathbb{Q}) \to \mathbb{Q}$, is given by

$$e^{\text{fw}}(\pi) = \det\left(\frac{\partial f_i}{\partial x_j}\right) \in E.$$
 (1.6)

We can apply this to the universal 1-connected $\mathbb{C}P^n$ -fibration to determine a representative of the fibrewise Euler class $e^{\mathrm{fw}}(\pi) \in E_n$, and we can compute the polynomials $\kappa_i = \kappa_i(x_2, \ldots, x_{n+1}) \in B_n$ by fibre integrating powers of it. We prove that $\kappa_1, \ldots, \kappa_{n-1}, \kappa_{n+1}$ generate the Euler ring by applying the Cayley–Hamilton theorem as explained in [RW18] and proved there in much greater generality. It remains to identify relations between these polynomials in B_n . By analysing terms of maximal power in x_{n+1} we can show that they are algebraically independent.

Theorem C. The Euler ring of complex projective space is $E^*(\mathbb{C}P^n) \cong \mathbb{Q}[\kappa_1, \ldots, \kappa_{n-1}, \kappa_{n+1}]$.

For n = 2 this has been proved in [RW18] and extended by Dexter Chua for $n \le 4$ using the natural torus action $T^n \curvearrowright \mathbb{C}P^n$ and localisation in equivariant cohomology.

The second class of elliptic spaces we study are rationally equivalent to products of odd dimensional simply connected spheres $X \simeq_{\mathbb{Q}} \prod_i S^{2n_i+1}$. In this case the universal 1-connected fibration is not Leray–Hirsch so that we have to use the techniques from section 3 and 4 to find representatives of fibre integration and the fibrewise Euler class. However, the minimal model of X is finite dimensional as a vector space which simplifies the problem significantly.

Proposition D. Let X be rationally equivalent to a product of odd dimensional simply connected spheres. Then the fibrewise Euler class $e^{\text{fw}}(\pi) \in H^{\dim X}(B \text{ hAut}_{0,*}(X); \mathbb{Q})$ vanishes.

We denote by $E_0^*(X) \subset H^*(B \operatorname{hAut}_0(X); \mathbb{Q})$ the image of the Euler ring under the pullback of $\operatorname{B}\operatorname{hAut}_0(X) \to \operatorname{B}\operatorname{hAut}^+(X)$. The above result implies $E_0^*(X) = \mathbb{Q}$ by naturality of the Euler class and fibre integration. Moreover, if $X \simeq_{\mathbb{Q}} \prod_i S^{2n_i+1}$ is a finite CW complex then $\pi_0(\operatorname{hAut}^+(X))$ is arithmetic [Sul77, Thm.10.3], and one can argue as in [GGRW17, Prop.5.2] that $E^*(X)/\sqrt{0} = \mathbb{Q}$. In some cases we can extend this result.

Theorem E. Let X be either rationally equivalent to $S^{2k+1} \times ... \times S^{2k+1}$ for $k \ge 1$ or a finite CW complex rationally equivalent to product of two odd dimensional simply connected spheres of different dimension. Then $E^*(X) = \mathbb{Q}$.

It remains an open problem to extend this result to all spaces rationally equivalent to a product of odd dimensional spheres.

Acknowledgements. This paper is part of my PhD project and I would like to thank my supervisor Oscar Randal-Williams for many enlightening discussions and his help and patience over the past years. I would also like to thank Alexander Berglund for many useful discussions.

2 The dg coalgebra model of the universal fibration

Rational models of fibrations $F \rightarrow E \rightarrow B$ of 1-connected spaces are well studied and are often given in terms of relative Sullivan algebras or Koszul–Sullivan models [Sul77, Hal83, Tho81, FHT01]. The main goal of this section is to find a cdga model of

$$X \hookrightarrow B \, hAut_{0,*}(X) \longrightarrow B \, hAut_0(X)$$

for a 1-connected space X using its minimal Sullivan model (ΛV , d). Sullivan proved that for a space X of finite type the dg Lie algebra $\operatorname{Der}^+(\Lambda V)$ is a model of $\operatorname{BhAut}_0(X)$, so that the natural starting point is to build a relative Sullivan algebra from it. However, note that this dg Lie algebra is not of finite type in general and thus it is technically involved to build a cdga model from it. One does not have these difficulties when considering the differential graded coalgebra $C_*^{CE}(\operatorname{Der}^+(\Lambda V);\mathbb{Q})$, and consequently, we will build a coalgebra model of the universal fibration. In the process, we will give in section 2.2 an algebraic classification of coalgebra bundles similar in spirit to Tanré's classification of semidirect products of dg Lie algebras. We should note that this classification it not needed in the proof of Theorem 2.19 but we included it as we find it conceptually quite clarifying.

Remark 2.1. Most of the results in this section as well as the main Theorem 2.19 are known to experts. But they are not easily deduced or found in the literature. In fact, we do not know sources for Theorem 2.19 nor for an algebraic classification of coalgebra bundles analogous to Tanré's work. We hope that this fact and later applications justify the rather extensive elaboration of known results.

2.1 Preliminaries and conventions. Throughout this paper, we denote by **dgcc** the category of differential graded cocommutative coalgebras over \mathbb{Q} . More precisely, an object $C \in \mathsf{dgcc}$ is $(C, \Delta_C, d_C, \varepsilon_C)$ where (C, d_C) is a chain complex over \mathbb{Q} and $(C, \Delta_C, \varepsilon_C)$ is a cocommutative counital coalgebra so that the comultiplication $\Delta_C : C \to C \otimes C$ and counit $\varepsilon_C : C \to \mathbb{Q}$ are chain maps. We will omit the subscripts when it is clear from the context which coalgebras we are referring to. The morphisms in **dgcc** are counital coalgebra maps of degree zero that commute with the differential.

Then dgcc has a symmetric monoidal structure given by the tensor product of the underlying chain complexes: For $C, C' \in \text{dgcc}$ the chain complex $C \otimes_{\mathbb{Q}} C'$ has a comultiplication $\Delta_{C \otimes C'} := (\operatorname{Id}_C \otimes T \otimes \operatorname{Id}_{C'})(\Delta_C \otimes \Delta_{C'})$ and counit $\epsilon_C \otimes \epsilon_{C'}$, where $T: C \otimes C' \to C' \otimes C$ is the chain map $T(c \otimes c') = (-1)^{|c||c'|}c' \otimes c$. The monoidal unit **1** is the coalgebra on the 1-dimensional chain complex concentrated in degree zero $(\mathbb{Q}, d = 0)$ with comultiplication determined by $\Delta(1) = 1 \otimes 1$ and counit $\epsilon_1 = \operatorname{Id}_{\mathbb{Q}}$. Note that the monoidal unit is also the terminal object in dgcc.

Lemma 2.2. Let $C \in \mathsf{dgcc}$, then the comultiplication $\Delta : C \to C \otimes C$ and counit $\epsilon : C \to \mathbb{Q}$ are maps in dgcc . Hence dgcc is cartesian monoidal, i.e. the monoidal structure is the product in dgcc .

We will denote by $\operatorname{dgcc}^{\operatorname{conil}}$ the full subcategory of coaugmented conilpotent coalgebras. A coaugmentation η is a section of the counit in dgcc and we denote $\eta(1)$ by $1 \in C$. A coaugmented coalgebra C is conilpotent if $\overline{C} := \ker(\epsilon)$ is a coalgebra with nilpotent comultiplication $\overline{\Delta}(c) := \Delta(c) - c \otimes 1 - 1 \otimes c$ (see [LV12, Ch.1.2] for details). Let dgLie be the category of differential graded Lie algebras. Then there is an adjunction [Qui69]

$$dgLie \xrightarrow{\tau}_{\mathcal{L}} dgcc^{conil}. \tag{2.1}$$

Let us recall the definition of $C: \mathsf{dgLie} \to \mathsf{dgcc}^\mathsf{conil}$ from [Qui69, App.B]: For $L \in \mathsf{dgLie}$ there is an acyclic dg Lie algebra $sL \oplus L$ containing L as a dg Lie subalgebra with brackets and differential D determined by

$$[sl, sl'] = 0$$
 $[sl, l'] = s[l, l']$ $D(sl) = l - sd_1 l.$ (2.2)

Then the enveloping algebra $U(sL \oplus L)$ is an acyclic dg Hopf algebra that is a free U(L)-module via the inclusion $L \hookrightarrow sL \oplus L$. Quillen's defintion of the functor C is

$$C(L) := U(sL \oplus L) \otimes_{U(L)} \mathbb{Q}. \tag{2.3}$$

Note that $U(sL \oplus L)$ is a free U(L)-module on the envelopping algebra of the abelian Lie algebra sL so that ignoring differentials $C(L) \cong U(sL) = \Lambda sL$ as a graded coalgebra. We choose C(L) as our defintion of the Chevalley-Eilenberg complex $C_*^{CE}(L;\mathbb{Q})$. More generally, if M is a left U(L)-module then the Chevalley-Eilenberg complex of L with coefficients in M is $C_*^{CE}(L;M) := U(sL \oplus L) \otimes_{U(L)} M$.

Remark 2.3. This definition of C is different than the one used for example in [FHT01]. The differential differs by a sign on the quadratic corestriction. The different choices are isomorphic as dgc coalgebras via a coalgebra isomorphism which only has a non-trivial linear corestriction given by $-\operatorname{Id}_{sL}: sL \to sL$.

Denote by dgLie_c the full subcategory of connected dg Lie algebras, i.e. Lie algebras concentrated in positive degrees; and by dgcc_1 the full subcategory of simply connected coalgebras, i.e. coalgebras whose counit is an isomorphism in degree 0 and \overline{C} is concentrated in degrees >1. Then the adjunction (2.1) induces a Quillen equivalence $\mathcal{L} \dashv C$ between dgLie_c and $\operatorname{dgcc}_1^{\operatorname{conil}}$ according to [Qui69].

Remark 2.4. The category dgcc^{conil} has a model category structure [Qui69] so in particular it has all small limits and colimits. The analogous statement of Lemma 2.2 holds in dgcc^{conil}₍₁₎.

2.2 Classification of coalgebra bundles. We prove a classification of coalgebra bundles which are the algebraic models for fibrations. This classification is inspired by the geometric classification of fibre bundles via associated principal bundles. The counter part of principal bundles in rational homotopy theory are principal *L*-bundles for a dg Lie algebra *L*. This has been introduced by Quillen in [Qui69, App.B] where he also describes the classification of such principal *L*-bundles.

Definition 2.5. A **coalgebra bundle** with fibre $C \in \mathsf{dgcc}^{\mathsf{conil}}$ is a map $\pi : E \to B$ in $\mathsf{dgcc}^{\mathsf{conil}}$ such that there exists an isomorphism $\phi : B \otimes C \to E$ of graded coaugmented coalgebras satisfying $\pi \phi|_{B \otimes 1} = \mathrm{Id}_B$ and $\phi|_{1 \otimes C} \in \mathrm{Hom}_{\mathsf{dgcc}^{\mathsf{conil}}}(C, E)$. We call such an isomorphism a **local trivialization**, and a pair $(\pi : E \to B, \phi)$ a trivialized coalgebra bundle with fibre C. Note that a local trivialization does not need to be a chain map. If it is then the bundle is trivial.

We want to formulate the classification of such bundles via the pullback of a universal bundle, which is also how the classification of semidirect products of dg Lie algebras in [Tan83] can be phrased. The universal coalgebra bundle with fibre $C \in \mathsf{dgcc}^{\mathsf{conil}}$ is obtained from the universal principal bundles that Quillen described in [Qui69]. In the following, we set

$$L = \text{CoDer}(C) \in \text{dgLie}.$$

Then C and \mathbb{Q} are U(L)-modules and the counit $\epsilon: C \to \mathbb{Q}$ is a U(L)-module map. We will show that the induced map

$$\operatorname{Id} \otimes \epsilon : U(sL \oplus L) \otimes_{U(L)} C \longrightarrow U(sL \oplus L) \otimes_{U(L)} \mathbb{Q} = C(L)$$
 (2.4)

is the universal coalgebra bundle with fibre C and local trivialization

$$\phi_L: C(L) \otimes C \stackrel{\cong}{\longrightarrow} U(sL) \otimes C \stackrel{j \otimes \operatorname{Id}_C}{\longrightarrow} U(sL \oplus L) \otimes_{U(L)} C,$$

where j is the map on enveloping algebras induced by inclusion of the abelian Lie algebra $j: sL \rightarrow sL \oplus L$. Note that ϕ_L is only a map of graded coalgebras and does not commute with the differential in general because neither the isomorphism $C(L) \cong U(sL)$ nor the inclusion of Hopf algebras $U(sL) \rightarrow U(sL \oplus L)$ are chain maps.

Given a coalgebra bundle $\pi: E \to B$ with fibre C and a map $f: X \to B$ the pullback $f^*E := \lim(X \to B \leftarrow E) \in \mathsf{dgcc}^{\mathsf{conil}}$ exists. We will show that the canonical map $\pi: f^*E \to X$ is a coalgebra bundle with the same fibre. The limit of $X \to B \leftarrow E$ in the category of graded cocommutative coalgebras is the underlying coalgebra of f^*E . Thus for a local trivialization $\phi: B \otimes C \to E$ we get $f\pi_X = \pi \circ \phi \circ (f \otimes \mathrm{Id}_C): X \otimes C \to B$ and hence there is an induced map of graded coalgebras $f^*\phi: X \otimes C \to f^*E$ which defines a local trivialization of the pullback.

Theorem 2.6. Let $C \in \mathsf{dgcc}^{conil}$ and $L = \mathsf{CoDer}(C) \in \mathsf{dgLie}$. For $B \in \mathsf{dgcc}^{conil}$ denote the set of isomorphism classes of trivialized coalgebra bundles with fibre C over B by $\mathsf{Bun}_B(C)$, then

$$\operatorname{Hom}_{\operatorname{dgcc}}(B,C(L)) \longrightarrow \operatorname{Bun}_B(C)$$
$$f \longmapsto \left[\pi: f^*(U(sL \oplus L) \otimes_{U(L)} C) {\to} B, f^* \phi_L\right]$$

is a one-to-one correspondence, where $f^*\phi_L$ is the induced local trivialization as described above. If $B \in \mathsf{dgcc}_1^{conil}$ then the above statement is also true if one replaces L by $L^+ = \mathsf{CoDer}^+(C)$.

The transition from principal L-bundles for L = CoDer(C) to coalgebra bundles with fibre C as in (2.4) corresponds to the construction of associated bundles in geometric topology. Following this analogy, the proof of Theorem 2.6 is based on the algebraic analogue of the reverse process, that associates to a smooth fibre bundle $M \rightarrow E \rightarrow B$ the principal Diff(M)-bundle $\coprod_{b \in B} \text{Diff}(M, E_b) \rightarrow B$. We implement this correspondence and construct a principal L-bundle associated to a coalgebra bundle with fibre C by using the classification of principal L-bundles due to Quillen.

Let us briefly recall some definitions and important properties from Appendix B in [Qui69].

Definition 2.7 ([Qui69]). Given $L \in \mathsf{dgLie}$ and $B \in \mathsf{dgcc}^{\mathsf{conil}}$ then a **principal** L-**bundle** with base B is a triple (E, m, π) , where $E \in \mathsf{dgcc}^{\mathsf{conil}}$ is a right U(L)-module with module map $m : E \otimes U(L) \longrightarrow E$ that is a map of dgc coalgebras, $\pi : E {\to} B$ is a map in $\mathsf{dgcc}^{\mathsf{conil}}$ satisfying $\pi(e {\cdot} u) = \pi(e) {\cdot} \epsilon(u)$ and there exists a map of graded coalgebras $\rho : B {\to} E$ so that $\varphi(b {\otimes} u) = \rho(b) {\cdot} u$ defines an isomorphism $\varphi : B \otimes U(L) {\to} E$ of graded cocommutative coalgebras and right U(L)-modules. Note however that φ is not a chain map in general. Such a map ρ is called a **local trivialization**.

Quillen proved that a principal *L*-bundle over *B* with a choice of local trivialization is determined by a **twisting function**, a linear map $\tau : B \rightarrow L$ of degree -1 such that

$$d_L \tau + \tau d_B + \frac{1}{2} [,] \circ (\tau \otimes \tau) \circ \Delta = 0 \quad \text{and} \quad \tau(1) = 0.$$
 (2.5)

In particular, the principal L-bundle over B corresponding to a twisting function $\tau: B \to L$ has as total space $E = B \otimes U(L)$ with the obvious module structure and projection and (co)differential on E defined in terms of τ as

$$d_{E}(b \otimes u) = d_{E}(b \otimes 1) \cdot u + (-1)^{|b|} b \otimes d_{U(L)}(u)$$

$$d_{E}(b \otimes 1) = d_{C}(b) + \sum_{i} (-1)^{|b_{(1)}|} b_{(1)} \otimes \tau(b_{(2)}).$$
(2.6)

Denote by $\mathcal{T}(B, L)$ the set of twisting functions. The functor $\mathcal{T}(-, -)$: $(\mathsf{dgcc}^{\mathrm{conil}})^{\mathrm{op}} \times \mathsf{dgLie} \to \mathsf{Set}$ is representable for a fixed B or L via $\mathsf{Hom}_{\mathsf{dgLie}}(\mathcal{L}(B), -)$ respectively $\mathsf{Hom}_{\mathsf{dgcc}}(-, C(L))$, i.e. there is a commutative diagram

$$C(L)$$

$$\exists ! \varphi_{B} \nearrow \tau$$

$$T$$

$$\exists ! \varphi_{L}$$

$$\mathcal{L}(B)$$

$$(2.7)$$

where φ_B and φ_L are maps in the respective categories. If we consider the principal bundles $E(B, \mathcal{L}(B), \tau_B)$ and $E(C(L), L, \tau_L) := U(sL \oplus L)$ corresponding to the universal twisting functions (see [Qui69, App. B] for definitions), this can be rephrased as a classification of principal bundles similar to geometric topology: For fixed L, every principal L-bundle over B is obtained as a pullback from the universal bundle $E(C(L), L, \tau_L)$ along a map $B \rightarrow C(L)$ in $\mathsf{dgcc}^{\mathsf{conil}}$.

We can now implement the construction of the principal L = CoDer(C)-bundle associated to a coalgebra bundle with fibre C.

Proposition 2.8. Let $\pi: E \rightarrow B$ be a coalgebra bundle with fibre $C \in \mathsf{dgcc}^{conil}$ and let ϕ be a local trivialization. Then

$$\tau_{E|_{\overline{B}}} : \overline{B} \longrightarrow \text{CoDer}(C),$$

$$b \longmapsto \left(c \mapsto \pi_{C} \phi^{-1} d_{E} \phi(b \otimes c)\right)$$
(2.8)

extends to a twisting function τ_E by setting $\tau_E(1) = 0$

Proof. Since we fix a local trivialization, we can assume that $E = B \otimes C$ as graded coalgebra with differential D such that the projection $\pi = \pi_B : (B \otimes C, D) \rightarrow (B, d_B)$ and inclusion $i : (C, d_C) \rightarrow (B \otimes C, D)$ are maps in dgcc. Then $\tau(b)(c) = \pi_C \circ D(b \otimes c)$ and we need to check that this defines in fact a coderivation and that τ satisfies the twisting condition.

Once can check that $\tau(b) := \pi_C \circ D(b \otimes -)$ defines a coderivation of C by using that π_C is a map of graded coalgebras. Moreover, it is clear from the definition that τ has degree -1 so it remains to check $d_L\tau + \tau d_B + \frac{1}{2}[\ ,\] \circ (\tau \otimes \tau) \circ \Delta = 0$. Let us first compute $D(b \otimes c)$. Note that $(\pi_B \otimes \pi_C)\Delta_{B \otimes C} = \mathrm{Id}_{B \otimes C}$, which, combined with D being a coderivation, gives

$$D(b \otimes c) = \sum_{A \in D} \pi_B D(b \otimes c_{(1)}) \otimes c_{(2)} + \sum_{A \in D} (-1)^{|b_{(1)}|} b_{(1)} \otimes \pi_C D(b_{(2)} \otimes c)$$
$$= d_B(b) \otimes c + \sum_{A \in D} (-1)^{|b_{(1)}|} b_{(1)} \otimes \pi_C D(b_{(2)} \otimes c)$$

where the second equality uses that π_B is a chain map. When we apply D again, the first summand gives

$$D(d_B b \otimes c) = \sum d_B(b_{(1)}) \otimes \pi_C D(b_{(2)} \otimes c) + (-1)^{|b_{(1)}|} b_{(1)} \otimes \pi_C D(d_B b_{(2)} \otimes c)$$

as $\Delta_B d_B b = d_B b_{(1)} \otimes b_{(2)} + (-1)^{|b_{(1)}|} b_{(1)} \otimes d_B b_{(2)}$, and the second summand becomes

$$\sum (-1)^{|b_{(1)}|} d_B(b_{(1)}) \otimes \pi_C D(b_{(2)} \otimes c) + \sum (-1)^{|b_{(1,2)}|} b_{(1,1)} \otimes \pi_C D(b_{(1,2)} \otimes \pi_C D(b_{(2)} \otimes c))$$

where we have set $\Delta b_{(1)} = \sum b_{(1,1)} \otimes b_{(1,2)}$. Combining these contributions and applying π_C gives

$$0 = \pi_C D^2(b \otimes c) = \pi_C D(d_B b \otimes c) + \sum_{(-1)^{|b_{(1)}|}} \pi_C D(b_{(1)} \otimes \pi_C D(b_{(2)} \otimes c)). \tag{2.9}$$

We compare with the twisting condition. Its first two contributions are

$$\begin{split} [d_C,\tau(b)](c) &= d_C \pi_C D(b \otimes c) - (-1)^{|b|-1} \pi_C D(b \otimes d_C c) \\ &= \pi_C D(1 \otimes \pi_C D(b \otimes c)) + (-1)^{|b|} \pi_C D(b \otimes \pi_C D(1 \otimes c)), \\ \tau(d_B b)(c) &= \pi_C D(d_B b \otimes c), \end{split}$$

where we have rewritten d_C in terms of D using that the inclusion $C \rightarrow C \otimes B$ is a chain map by assumption; the third and last contribution of the twisting condition is

$$\begin{split} \left(\frac{1}{2}\sum_{|b_{(1)}|<|b|}[\tau(b_{(1)}),\tau(b_{(2)})]\right)(c) &= \\ &\frac{1}{2}\sum_{|b_{(i)}|<|b|}\pi_{C}D(b_{(1)}\otimes\pi_{C}D(b_{(2)}\otimes c)) - (-1)^{(|b_{(1)}|-1)(|b_{(2)}|-1)}\pi_{C}D(b_{(2)}\otimes\pi_{C}D(b_{(1)}\otimes c)) \end{split}$$

Note that $\tau(1) = 0$ so the sum does not include the contribution from $b \otimes 1 + 1 \otimes b$. One can check using the cocommutativity of B that the sum of these three terms gives the right side of (2.9) and thus vanishes. Hence, τ is a twisting function.

Corollary 2.9. Let C and L be as above and denote by L-Bun $_B$ the set of isomorphism classes of trivialized principal L-bundles over B, then

$$L\text{-}Bun_B \longrightarrow Bun_B(C)$$

$$[(E, m, \pi), \rho] \longmapsto [\operatorname{Id} \otimes \epsilon : E \otimes_{U(L)} C {\rightarrow} E \otimes_{U(L)} \mathbb{Q}, \rho \otimes \operatorname{Id}_C]$$

is a one-to-one correspondence. We call the coalgebra bundle $\mathrm{Id} \otimes \varepsilon : E \otimes_{U(L)} \mathbb{C} \to E \otimes_{U(L)} \mathbb{Q}$ the **associated coalgebra bundle**. Hence, we can think of both principal L-bundles and coalgebra bundles just as a choice of local trivialization and a twisting function $\mathcal{T}(B,L)$.

Proof. We first show that the map $E \otimes_{U(L)} C \to E \otimes_{U(L)} \mathbb{Q}$ defines a coalgebra bundle. Note that the forgetful functor $\mathcal{U}: \mathsf{dgcc}^{\mathsf{conil}} \to \mathsf{dgVect}$ has as right adjoint $\Lambda^c: \mathsf{dgVect} \to \mathsf{dgcc}^{\mathsf{conil}}$, where $\Lambda^c V$ is the cofree conilpotent cocommutative coalgebra on V with differential determined by its linear part. Hence, \mathcal{U} preserves colimits. If H is a cocommutative dg Hopf algebra and M and N are right respectively left H-modules for which the module maps $M \otimes H \to M$ and $H \otimes N \to N$ are maps in $\mathsf{dgcc}^{\mathsf{conil}}$, then $M \otimes H \otimes N \rightrightarrows M \otimes N$ is a diagram in $\mathsf{dgcc}^{\mathsf{conil}}$, and

 $\mathcal{U}(\text{coeq}(M \otimes H \otimes N \Rightarrow M \otimes N)) = M \otimes_H N \in \mathsf{dgVect}$. This should be interpreted as defining a coalgebra structure on the tensor product as a coequalizer in $\mathsf{dgcc}^{\text{conil}}$. If $N_1 \rightarrow N_2$ is a map in $\mathsf{dgcc}^{\text{conil}}$ and a H-module map, the induced map $M \otimes_H N_1 \rightarrow M \otimes_H N_2$ is a map in $\mathsf{dgcc}^{\text{conil}}$.

The module map $E \otimes U(L) \rightarrow E$ is a map in $\mathsf{dgcc}^{\mathsf{conil}}$ by definition, and the same is true for \mathbb{Q} as a (left) U(L)-module for every $L \in \mathsf{dgLie}$. One can check that the U(L)-module structure on C given by

$$m: U(L) \otimes C \longrightarrow C$$

$$(\theta_1 \cdot \ldots \cdot \theta_n) \otimes c \longmapsto \theta_1 \circ \ldots \circ \theta_n(c)$$
(2.10)

is a map in $\operatorname{dgcc}^{\operatorname{conil}}$. Since the counit $\varepsilon: C \to \mathbb{Q}$ is a U(L)-module map, $\operatorname{Id} \otimes \varepsilon: E \otimes_{U(L)} C \to E \otimes_{U(L)} \mathbb{Q}$ is a map in $\operatorname{dgcc}^{\operatorname{conil}}$. If $\rho: B \to E$ is a local trivialization, then $\phi: B \otimes C \to E \otimes_{U(L)} C$ given by $\phi(b \otimes c) := \rho(b) \otimes c$ defines a local trivialization of the coalgebra bundle. Hence, the above map is well-defined.

By [Qui69, App.B Prop.5.3], there is a one-to-one correspondence L-Bun $_B \to \mathcal{T}(B,L)$ that associates to a trivialized principal L-bundle its twisting function. This twisting function is the same as the one obtained from applying Proposition 2.8 to the associated bundle. Moreover, given $[\pi: E \to B, \phi] \in \text{Bun}_B(C)$ then there is a trivialized principal L-bundle associated to τ_E and its associated coalgebra bundle is isomorphic to $(\pi: E \to B, \phi)$.

Corollary 2.10. *The map defined in* (2.4) *is a coalgebra bundle with fibre C.*

There is one last technical statement we need for the proof of Theorem 2.6 in order to show that the pullback of a coalgebra bundle is a coalgebra bundle.

Lemma 2.11. Let $(C, \otimes, 1)$ be a cartesian monoidal category and suppose $f : X \rightarrow B$ and $\pi : E \rightarrow B$ are two maps in C. Then $\lim(X \rightarrow B \leftarrow E)$ if it exists is isomorphic to the equalizer of

$$X \otimes E \xrightarrow[X \otimes \pi]{f \otimes E} X \otimes B \otimes E$$

Proof. In a cartesian monoidal category every object is a coalgebra object. So if there are $h: Z \rightarrow X$ and $h': Z \rightarrow E$ satisfying $fh = \pi h'$ then $(h \otimes h')\Delta_Z: Z \longrightarrow X \otimes E$ equalizes the above diagram. Hence, there is a unique map $H: Z \longrightarrow \operatorname{eq}(X \otimes E \rightrightarrows X \otimes B \otimes E)$ so that the composition of H with $\operatorname{eq}(X \otimes E \rightrightarrows X \otimes B \otimes E) \rightarrow X \otimes E$ and the respective projections to X respectively E commute with E respectively E. Hence, the equalizer has the universal property of the limit.

Proof of Theorem 2.6. By Corollary 2.9, the set $\operatorname{Bun}_B(C)$ is in one-to-one correspondence with the set of twisting functions which is in one-to-one correspondence with $\operatorname{Hom}_{\operatorname{dgcc}}(B, C(L))$ by [Qui69]. Thus is remains to show that the pullback of coalgebra bundles in Theorem 2.6 induces this correspondence. Let $f \in \operatorname{Hom}_{\operatorname{dgcc}}(B, C(L))$. The pullback in the Theorem is the limit

$$\lim(B \xrightarrow{f} C(L) \longleftarrow U(sL \oplus L) \otimes_{U(L)} C) \in \mathsf{dgcc}^{\mathrm{conil}}.$$

Since $dgcc^{conil}$ is a cartesian monoidal category with all small limits, the above Lemma implies that the pullback in $dgcc^{conil}$ of a diagram $X \rightarrow B \leftarrow E$ is the cotensor product

 $X \square_B E = \operatorname{eq}(X \otimes E \rightrightarrows X \otimes B \otimes E)$. If $E \rightarrow B$ is a coalgebra bundle with fibre C, then $E \cong B \otimes C$ as graded coalgebras. The B-comodule structure induced by π_B makes E into an extended B-comodule so that $X \square_B E \cong X \otimes C$ as graded coalgebras [EM66, Prop2.1]. This isomorphism is a local trivialization and it coincides with the local trivialization $f^*\phi$ obtained using the universal property of the pullback. In conclusion, $f^*E \rightarrow X$ is a coalgebra bundle with fibre C and local trivialization $f^*\phi$. We can apply this to the pullback of the universal coalgebra bundle along $f: B \rightarrow C(L)$. One can check that the twisting function from Proposition 2.8 is $\tau_L \circ f$. Hence, the pullback induces a one-to-one correspondence as claimed.

The second part of the statement for simply connected base follows from the following observation about twisting functions $\tau: B \to L = \text{CoDer}(C)$. If $b \in B$ is primitive and $d_B(b) = 0$, then it follows from the twisting equation that $[d_C, \tau(b)] = 0$. Consequently, if $B \in \mathsf{dgcc}_1^{\mathrm{conil}}$, then $\tau: \overline{B} \to \mathrm{CoDer}^+(C)$ as every element in B_2 is primitive and a cycle. In particular, $\mathcal{T}(B, \mathrm{CoDer}(C)) = \mathcal{T}(B, \mathrm{CoDer}^+(C))$ and so we can consider the 1-truncation if the base is simply connected.

2.3 Model and comparison with Tanré. If we think of the dg Lie algebra $L = \operatorname{CoDer}^+(C)$ for $C \in \operatorname{dgcc}_1^{\operatorname{conil}}$ as a model for BhAut₀(X) for a space $X = \langle C \rangle$, the universal principal L-bundle plays the role of EhAut₀(X) with an action of Ω BhAut₀(X) = hAut₀(X). This resembles the homotopy theoretic classification of fibrations, where the universal fibration can be thought of as the associated bundle EhAut₀(X) $\times_{\operatorname{hAut}_0(X)} X$. Of course, we can only expect this to be true for coalgebras that behave well with respect to homotopy theory in $\operatorname{dgcc}_1^{\operatorname{conil}}$. Hence, if $(\Lambda V, d)$ is a minimal Sullivan model of a space X of finite type with dual coalgebra model $(\Lambda^c V^{\vee}, d^{\vee})$, then based on the above analogy we expect the associated coalgebra bundle for the universal principal L-bundle for $L = (\operatorname{CoDer}^+(\Lambda^c V^{\vee}), [d^{\vee}, -]) \in \operatorname{dgLie}_c$ to be a model of the universal 1-connected fibration.

Theorem 2.12. Let X be a 1-connected space of finite type with minimal Sullivan model $(\Lambda V, d)$ and dual coalgebra $(\Lambda^c V^{\vee}, d^{\vee}) \in \mathsf{dgcc}_1^{\mathrm{conil}}$. Then the associated coalgebra bundle

$$\pi := C_*^{CE}(\operatorname{CoDer}^+(\Lambda^c V^{\vee}); \Lambda^c V^{\vee}) \xrightarrow{\epsilon_*} C_*^{CE}(\operatorname{CoDer}^+(\Lambda^c V^{\vee}); \mathbb{Q})$$
 (2.11)

is a coalgebra model for the universal 1-connected fibration $X \to B \, hAut_{0,*}(X) \to B \, hAut_0(X)$.

A possible strategy of proof is probably to do a classification of coalgebra bundles up to fibre homotopy equivalence similar as in [Tan83]. Two coalgebra bundles $\pi_i : E_i \rightarrow B_i$ for = 1,2 are **fibre homotopy equivalent** if there are quasi-isomorphisms $F : E_1 \rightarrow E_2$ and $f : B_1 \rightarrow B_2$ in dgcc so that $\pi_2 F = f \pi_1$. But we prefer to simply compare the map in (2.11) with Tanré's model by applying C(-) to the fibre sequence (1.4), which is a coalgebra bundle over $C(\text{Der}^+(L) / \text{ad } L)$ with fibre C(L) and therefore there is a classifying map

$$\Phi : C(\operatorname{Der}^+(L) /\!\!/ \operatorname{ad} L) \rightarrow C(\operatorname{CoDer}^+(C(L)))$$

by Theorem 2.6. We show that if L is a free dg Lie algebra ($\mathbb{L}V,d$) then Φ is a quasi-isomorphism and induces a fibre homotopy equivalence of the corresponding coalgebra bundles with fibre $C(\mathbb{L}V,d)$. This almost gives a full proof of Theorem 2.12. The last step will be to compare coalgebra bundles with fibre $C(\mathbb{L}V,d)$ for a Quillen model of X with coalgebra bundles with fibre (Λ^cV^\vee,d^\vee) for the dual of a Sullivan model of X.

Proposition 2.13. Let Φ be the map above. Then the only non-trivial corestriction is linear, i.e. it is induced by a map

$$\phi : \operatorname{Der}^+(L) /\!\!/ \operatorname{ad} L \longrightarrow \operatorname{CoDer}^+(C(L))$$

of dg Lie algebras. Moreover, ϕ is a quasi-isomorphism if L is a free dg Lie algebra.

Remark 2.14. This proposition has been proved in [SS12, Thm.3.17]. We have included it in this discussion because it is derived from the classification of coalgebra bundles, which has the (small) advantage that we do not need to check that it induces a map of dg Lie algebras. Also note that there is a dual statement involving the left adjoint \mathcal{L} in [Gat97].

Proof. The sequence

$$C(L) \rightarrow C(L \rtimes_{id} (\operatorname{Der}^+(L) /\!\!/ \operatorname{ad} L)) \rightarrow C(\operatorname{Der}^+(L) /\!\!/ \operatorname{ad} L)$$

is a coalgebra bundle with fibre C(L) and Proposition 2.8 describes the corresponding twisting function $\tau: C(\operatorname{Der}^+(L) /\!\!/ \operatorname{ad} L) \to \operatorname{CoDer}^+(C(L))$. Note that the total space can be identified with $C^{CE}_*(\operatorname{Der}^+(L) /\!\!/ \operatorname{ad} L) \to \operatorname{CoDer}^+(C(L))$. Since the differential on the total space is the differential of the Chevalley-Eilenberg complex of a dg Lie algebra, it only has nontrivial linear and quadratic corestriction. Hence, one can check that $\tau(\chi) = \pi_{C(L)}D(\chi \otimes -)$ is only non-trivial on elements $\chi \in \Lambda^1 s(\operatorname{Der}^+(L) /\!\!/ \operatorname{ad} L)$. Denote by \overline{D} the differential on the Chevalley-Eilenberg complex of the semidirect product from [Tan83, VII.2.(11)], then the corestrictions of the coderivation $\tau(\chi)$ of C(L) for $\chi \in \Lambda^1 s(\operatorname{Der}^+(L) /\!\!/ \operatorname{ad} L)$ are

$$\tau(s^2l)_n(sl_1 \wedge \ldots \wedge sl_n) = \pi_{\Lambda^1 sL} \overline{D}(s^2l \wedge sl_1 \wedge \ldots \wedge sl_n)$$

$$= \begin{cases} \pi_{\Lambda^1 sL} \overline{D}(s^2l) = sl & n = 0\\ 0 & n > 0 \end{cases}$$

and

$$\tau(s\theta)_n(sl_1 \wedge \ldots \wedge sl_n) = \pi_{\Lambda^1 sL} \overline{D}(s\theta \wedge sl_1 \wedge \ldots \wedge sl_n)$$

$$= \begin{cases} \pi_{\Lambda^1 sL} \overline{D}(s\theta \wedge sl_1) = (-1)^{|\theta|} s[\theta, l_1] = (-1)^{|\theta|} s\theta(l_1) & n = 1\\ 0 & n \neq 1 \end{cases}$$

By (2.7) τ factors through a dgc coalgebra map $\Phi: C(\operatorname{Der}^+(L) /\!\!/ \operatorname{ad} L) \longrightarrow C(\operatorname{CoDer}^+(C(L)),$ which in turn has also only a linear part $\Phi_1: s(\operatorname{Der}^+(L) /\!\!/ \operatorname{ad} L) \longrightarrow s(\operatorname{CoDer}^+(C(L)))$. The desuspension of Φ_1 is described by

$$(s^{-1}\Phi_1)(sl) \in \text{CoDer}^+(C(L))$$
 has correstriction : $\Lambda sL \to \Lambda^0 sL = \mathbb{Q} \xrightarrow{sl} sL$
 $(s^{-1}\Phi_1)(\theta) \in \text{CoDer}^+(C(L))$ has correstriction : $\Lambda sL \to \Lambda^1 sL = sL \xrightarrow{(-1)^{|\theta|} s\theta} sL$

Since we already know that Φ is a dgc coalgebra map on the Chevalley-Eilenberg complexes of the two dg Lie algebras with only linear part, the desuspendsion $s^{-1}\Phi_1 =: \phi$ has to be a map of dg Lie algebras, which is also readily checked.

Before we prove the second part of the statement, observe that there are isomorphisms of graded vector spaces

$$CoDer(C(L)) \cong Hom_{arVect}(\Lambda sL, sL) \cong Hom_{U(L)}(U(sL \oplus L), sL)$$

where the right side is the graded vector space of U(L)-module homomorphisms and sL is the suspension of L considered as a module over itself. We will discuss differential homological algebra and semifree resolutions in more detail in section 3.2 and we refer the reader to that section for definitions. To finish this proof, we note that $\operatorname{Hom}_{U(L)}(U(sL \oplus L), sL)$ is a differential U(L)-module and one can check that the above isomorphism of graded vector spaces is in fact an isomorphism of chain complexes. As $U(sL \oplus L) \to \mathbb{Q}$ is a semifree resolution of \mathbb{Q} [FHT01, Prop.22.4], the homology of $\operatorname{CoDer}(C(L))$ can be identified with (a shift of) $H^*_{CF}(L;L) = \operatorname{Ext}_{U(L)}(\mathbb{Q},L)$.

If $L = (\mathbb{L}V, d)$ is a free dg Lie algebra, there is a small free resolution of \mathbb{Q} as described in [Wei94, Prop.7.2.4] given by $I \to U(L) \stackrel{\varepsilon}{\to} \mathbb{Q}$, where I is the kernel of the augmentation. Then I is a free U(L)-module on a basis of V because L is free. We can build a semifree resolution of \mathbb{Q} from it by setting $P = sI \oplus U(L)$ with standard differential on U(L) and $d(sI) = l - sd_{U(L)}l$ for $l \in I$. Hence, $Hom_{U(L)}(U(sL \oplus L), sL)$ is quasi-isomorphic to $Hom_{U(L)}(P, sL)$ which is given as a bicomplex by

$$\operatorname{Hom}_{U(\mathbb{L}V)}(s\mathcal{I}, s\mathbb{L}V) \cong \operatorname{Hom}_{\operatorname{grVect}}(V, \mathbb{L}V) \longleftarrow \operatorname{Hom}_{U(\mathbb{L}V)}(U(\mathbb{L}V), s\mathbb{L}V) \cong s\mathbb{L}V.$$

This can be identified with $\operatorname{Der}(\mathbb{L}V)$ // ad $\mathbb{L}V$. Moreover, there is an obvious inclusion $i: P \to U(s\mathbb{L}V \oplus \mathbb{L}V)$ which is a map between semifree resolutions and therefore induces a quasi-isomorphism $i^*: \operatorname{Hom}_{U(\mathbb{L}V)}(P, s\mathbb{L}V) \to \operatorname{Hom}_{U(\mathbb{L}V)}(U(s\mathbb{L}V \oplus \mathbb{L}V), s\mathbb{L}V)$ [FHT01, Prop.6.7] which agrees with ϕ under the identification of $\operatorname{Der}(\mathbb{L}V)$ // ad $\mathbb{L}V \cong \operatorname{Hom}_{U(\mathbb{L}V)}(P, s\mathbb{L}V)$ and $\operatorname{CoDer}(C(\mathbb{L}V, d)) \cong \operatorname{Hom}_{U(\mathbb{L}V)}(U(s\mathbb{L}V \oplus \mathbb{L}V), s\mathbb{L}V)$. Hence, the map of 1-truncations in the statement is a quasi-isomorphism.

Corollary 2.15. The map $\phi_* : C_*^{CE}(\operatorname{Der}^+(\mathbb{L}V) / \operatorname{ad} \mathbb{L}V; C(\mathbb{L}V)) \longrightarrow C_*^{CE}(\operatorname{CoDer}^+(C(\mathbb{L}V)); C(\mathbb{L}V))$ is a quasi-isomorphism in $\operatorname{dgcc}_1^{conil}$ and is compatible with the map induced by the change of coefficients $\epsilon : C(\mathbb{L}V) \to \mathbb{Q}$. In particular, ϕ_* is a fibre homotopy equivalence of coalgebra bundles with fibre $C(\mathbb{L}V)$.

We need the following lemma to finish the proof of Theorem 2.12. It is proved similarly as [BM14, Lem.3.5].

Lemma 2.16. Let $\phi: C \to C'$ be a map in $\operatorname{dgcc}_1^{conil}$ and define $\operatorname{CoDer}_{\phi}(C, C')$ as the linear maps $\eta: C \to C'$ satisfying $\Delta_{C'} \eta = (\eta \otimes \phi + T(\eta \otimes \phi)T)\Delta_C$ and differential $D(\eta) = d_{C'} \eta - (-1)^{|\eta|} \eta d_C$. Then

$$\phi_* : \operatorname{CoDer}(C) \longrightarrow \operatorname{CoDer}_{\phi}(C, C') \qquad \qquad \phi^* : \operatorname{CoDer}(C') \longrightarrow \operatorname{CoDer}_{\phi}(C, C')$$

$$\eta \longmapsto \phi \circ \eta \qquad \qquad \eta' \longmapsto \eta' \circ \phi$$

are chain maps and quasi-isomorphisms if C and C' are cofree.

Proof of 2.12. Let ($\mathbb{L}X$, d) be a minimal Lie model of X and ($\Lambda^c V^{\vee}$, d^{\vee}) \in dgcc $_1^{\mathrm{conil}}$ the dual of a minimal Sullivan model of X. Fix an injective quasi-isomorphism $\psi: \Lambda^c V^{\vee} \hookrightarrow C(\mathbb{L}X)$. Then by the above lemma there is a zig-zag of quasi-isomorphisms

$$\operatorname{CoDer}(\Lambda^{c}V^{\vee}) \xrightarrow{\simeq} \operatorname{CoDer}_{\psi}(\Lambda^{c}V^{\vee}, \mathcal{C}(\mathbb{L}X)) \xleftarrow{\simeq}_{\psi^{*}} \operatorname{CoDer}(\mathcal{C}(\mathbb{L}X)).$$

Note that ψ^* is surjective, and hence by the same argument as in the proof of Theorem 3.10 in [BM14] the pullback $P = \{(\eta, \eta') \in \text{CoDer}(\Lambda^c V^{\vee}) \times \text{CoDer}(C(\mathbb{L}X)) \mid \eta \psi = \psi \eta' \}$ in chain complexes

of the above diagram is a differential graded Lie algebra satisfying that the projections of 1-truncations $\pi_1: P^+ \to \operatorname{CoDer}^+(\Lambda^c V^\vee)$ and $\pi_2: P^+ \to \operatorname{CoDer}^+(C(\mathbb{L}X))$ are quasi-isomorphisms in dgLie. Note that both $\Lambda^c V^\vee$ and $C(\mathbb{L}X)$ are U(P)-modules via these projections and importantly, ψ is a U(P)-module homomorphism. Hence, we have a sequence of induced maps

$$C_{*}^{CE}(\mathsf{CoDer}^{+}(C(\mathbb{L}X));C(\mathbb{L}X)) \xleftarrow{(\pi_{2})_{*}} C_{*}^{CE}(P;C(\mathbb{L}X))$$

$$\downarrow^{\psi_{*}}\uparrow^{\simeq}$$

$$C_{*}^{CE}(P;\Lambda^{c}V^{\vee}) \xrightarrow{(\pi_{1})_{*}} C_{*}^{CE}(\mathsf{CoDer}^{+}(\Lambda^{c}V^{\vee});\Lambda^{c}V^{\vee})$$

which are quasi-isomorphisms as they are induced by quasi-isomorphisms, and which are all compatible with the map to $C^{CE}(-;\mathbb{Q})$ induced by the counits. Hence, the above describes a sequence of fibre homotopy equivalences of coalgebra bundles. Combining with Corollary 2.15 gives the desired comparison with Tanré's model of the universal fibration.

2.4 The cdga model of the universal 1-connected fibration. The dual of the universal coalgebra fibration in (2.11) determines a cdga model of the universal 1-connected fibration. We will describe a simpler description of this dual where we do not need consider the dual coalgebra of a cdga. This model will also be more closely related to the the dg Lie algebra model of B hAut₀(X) of Sullivan in terms of derivations of a minimal Sullivan model of X.

Let $L \in \mathsf{dgLie}$ and $C \in \mathsf{dgcc}$ be a U(L)-module. Recall that if the module map $U(L) \otimes C \to C$ is a map in dgcc or equivalently that C acts by coderivations, then $C^{CE}_*(L;C) := U(sL \oplus L) \otimes_{U(L)} C$ is in dgcc. Thus, the dual $C^{CE}_*(L;C)^{\vee}$ is a cdga and isomorphic to

$$\begin{aligned} \operatorname{Hom}_{\operatorname{\mathsf{grVect}}} \left(U(sL \oplus L) \otimes_{U(L)} C, \mathbb{Q} \right) & \cong \operatorname{Hom}_{R-U(L)} \left(U(sL \oplus L), \operatorname{Hom}_{\operatorname{\mathsf{grVect}}} (C, \mathbb{Q}) \right) \\ & \cong \operatorname{Hom}_{L-U(L^{\operatorname{op}})} \left(U(sL \oplus L), \operatorname{Hom}_{\operatorname{\mathsf{grVect}}} (C, \mathbb{Q}) \right), \end{aligned}$$

where the first isomorphism follows from adjunction (C^{\vee} := $\operatorname{Hom}_{\operatorname{grVect}}(C, \mathbb{Q})$ is a right U(L)-module so that we have to consider right U(L)-morphisms) and the second isomorphism follows from the fact that every right U(L)-module determines a left $U(L^{\operatorname{op}})$ -module. Here, L^{op} denotes the opposite dg Lie algebra (L^{op} , $[,]_{L^{\operatorname{op}}}$, $d_{L^{\operatorname{op}}}$) on the same vector space as L with $[x,y]_{L^{\operatorname{op}}}=[y,x]_L$ and $d_{L^{\operatorname{op}}}(x)=-(-1)^{|x|}d_L(x)$. It is a basic fact that L is isomorphic as dg Lie algebra to its opposite.

Lemma 2.17. Let $(L, [,], d) \in \mathsf{dgLie}$, then $(L, [,]^{op}, d^{op})$ is a differential graded Lie algebra and

$$\phi: (L, [\,,\,], d) \longrightarrow (L, [\,,\,]^{op}, d_{L^{op}})$$

$$l \longmapsto -(-1)^{\lfloor |l|/2\rfloor} l$$

is an isomorphism of differential graded Lie algebras.

Consequently, $U(sL \oplus L)$ is isomorphic to $U(sL^{op} \oplus L^{op})$ as a left $U(L^{op})$ -module and we get an isomorphism of cdga's

$$\operatorname{Hom}_{\operatorname{\mathsf{grVect}}} \left(U(sL \oplus L) \otimes_{U(L)} C, \mathbb{Q} \right) \cong \operatorname{\mathsf{Hom}}_{L-U(L^{\operatorname{op}})} \left(U(sL^{\operatorname{op}} \oplus L^{\operatorname{op}}), C^{\vee} \right).$$

This is the definition of $C^*_{CE}(L^{op}, C^{\vee})$ using left-modules. By the above Lemma, this is isomorphic to Lie algebra cohomology via right modules. But it is not necessary to explicitly work out the isomorphism.

Lemma 2.18. Let $C \in \operatorname{dgcc}$ and $L = \operatorname{CoDer}(C) \in \operatorname{dgLie}$. Then $L^{op} \cong \operatorname{Der}(C^{\vee})$ and under this isomorphism the left $U(L^{op})$ -module structure is given by evaluating the derivation.

Thus, we can avoid dualizing the cdga models of spaces and their derivations completely when we want to compute Lie algebra cohomology, and we arrive at the following reformulation of Theorem 2.12.

Theorem 2.19. Let X be a 1-connected space of finite type with minimal Sullivan model $(\Lambda V, d)$ and unit $\eta : \mathbb{Q} \rightarrow \Lambda V$. Then

$$C_{CE}^*(\operatorname{Der}^+(\Lambda V); \mathbb{Q}) \xrightarrow{\eta_*} C_{CE}^*(\operatorname{Der}^+(\Lambda V); \Lambda V)$$
 (2.12)

is a cdga model of the universal oriented 1-connected fibration (1.3).

We can give a more explicit description of (2.12). Let L be a connected dg Lie algebra that acts on a connected cgda A of finite type through positive derivations. Note that we have to define positive degree derivations of a cdga as derivations that *lower* the degree if we don't fix our convention for chain complexes to be homologically or cohomologically graded. In order to be consistent (for the adjunction as well) we consider a cdga A in this section as negatively graded, so that both grading issues are resolved. In particular, the cgda $\operatorname{Hom}_{U(L)}(U(sL \oplus L), A)$ as the dual of a connected dgc coalgebra is concentrated in negative degrees. If A is of finite type, then

$$\operatorname{Hom}_{L-U(L)}(U(sL \oplus L), A) \cong \bigoplus_{n < 0} \prod_{k > 0} \operatorname{Hom}_{\mathsf{Vect}}((\Lambda sL)^k, A^{k+n}) \cong \operatorname{Hom}_{L-U(L)}(U(sL \oplus L), \mathbb{Q}) \otimes A$$

is an isomorphism of commutative algebras concentrated in negative degrees. We can work out the differential on the right side $C^*_{CE}(L;\mathbb{Q})\otimes A$ under this isomorphism. Denote by $\{l_i\}$ a basis of L and by $\{a_j\}$ a basis of A, and denote by $\{a_j^*\}$ and $\{(sl_i)^*\}$ the corresponding dual bases of A^\vee and $\Lambda^1 sL \subset C^*_{CE}(L;\mathbb{Q})$. Furthermore, we denote the differential on $C^*_{CE}(L;\mathbb{Q})\otimes A$ by D and by d on $C^*_{CE}(L;\mathbb{Q})$. Then under the above isomorphism D is given by

$$D(x \otimes 1) = d(x) \otimes 1$$

$$D(1 \otimes a) = 1 \otimes d_A(a) - \sum_j \prod_i \left(a_j^* (l_i \cdot a) \cdot (sl_i)^* \right) \otimes a_j.$$
(2.13)

for $x \in C^*_{CE}(L; \mathbb{Q})$ and $a \in A$. Finally, note that if A is a simply connected Sullivan algebra, then $\eta_* : C^*_{CE}(L; \mathbb{Q}) \to C^*_{CE}(L; A)$ is a relative Sullivan algebra. In the following, we will always interpret the above cdga's as positively graded.

Example 2.20. Let $X = S^{2n}$ with Sullivan model $A_n = (\Lambda(x, y), |x| = 2n, |y| = 4n-1, d = x^2 \cdot \partial/\partial y)$. Then $Der^+(A_n)$ is 3-dimensional with basis $\eta_{2n-1} := x \cdot \partial/\partial y$, $\eta_{2n} := \partial/\partial x$ and $\eta_{4n-1} := \partial/\partial y$ and differential $[d, \eta_{2n}] = -2\eta_{2n-1}$. Hence, the inclusion of the abelian Lie algebra with trivial differential $g := \mathbb{Q}\{\eta_{4n-1}\} \hookrightarrow Der^+(A_n)$ is a quasi-isomorphism of dg Lie algebras and we get a cdga quasi-isomorphism $C^*_{CE}(Der^+(A_n); \mathbb{Q}) \to C^*_{CE}(g; \mathbb{Q}) = (\Lambda z_{4n}, d = 0)$ as well as with coefficients in A_n . Thus, the cgda model of the universal 1-connected S^{2n} -fibration in Theorem 2.19 is equivalent to

$$(\Lambda(z_{4n}),d=0)\longrightarrow (\Lambda(z_{4n})\otimes\Lambda(x,y),D(x)=0,D(y)=x^2-z_{4n}),$$

where we use (2.13) with basis $\{x^k, yx^l\}_{k,l\geq 0}$ of A_n and $\{\eta_{4n-1}\}$ of g.

3 Fibre integration in rational homotopy theory

3.1 ϕ -integration. Let $\pi: E \to B$ be a fibration with fibre F. We want to introduce fibre integration as a special case of a more general construction. Let k be a commutative ring. Suppose $H^*(F;k)$ vanishes for *>d and is non-trivial in degree d. Note that $H^*(F;k)$ is a $k[\pi_1(B)]$ -module and let $\phi: H^d(F;k) \to k$ be a $k[\pi_1(B)]$ -module homomorphism, where k is the trivial $k[\pi_1(B)]$ -module.

Since $H^*(F;k)$ vanishes in degrees higher than d one can project $H^*(E;k)$ onto the d-th row of the E_{∞} -page of the Serre spectral sequence, and as there are no differentials into this row we have $E_{\infty}^{*-d,d} \subset E_2^{*-d,d}$. Thus we can define ϕ -integration as the composition

$$\Phi_*: H^*(E;k) \to E_{\infty}^{*-d,d} \subset E_2^{*-d,d} = H^{*-d}(B;H^d(F;k)) \xrightarrow{H(\phi)} H^{*-d}(B;k). \tag{3.1}$$

Since the cohomological Serre spectral sequence is compatible with cup product, there is a push-pull identity

$$\phi_*(\pi^*(x) \smile y) = x \smile \phi_*(y) \tag{3.2}$$

for $x \in H^*(B;k)$ and $y \in H^*(E;k)$, and thus ϕ -integration is a $H^*(B;k)$ -module map.

Definition 3.1. Let $\pi: E \to B$ be a fibration with oriented Poincaré fibre X of formal dimension d and suppose π is oriented, i.e. the orientation $\varepsilon_X: H^d(X;k) \to k$ is an isomorphism of $k[\pi_1(B)]$ -modules. Then

$$\pi_1 := (\mathcal{E}_X)_* : H^*(E;k) \to H^{*-d}(B;k)$$

is called **fibre integration** of $\pi : E \rightarrow B$.

3.2 Chain level ϕ **-integration.** The goal of this section is to find formulas for fibre integration and more generally ϕ -integration in terms of the algebraic models that we have introduced in the previous section. We will show that a homomorphism $\phi: H^d(F; \mathbb{Q}) \to \mathbb{Q}$ as above defines a unique "derived" chain homotopy class $\Phi: C^*(E; \mathbb{Q}) \to C^{*-d}(B; \mathbb{Q})$ of $C^*(B; \mathbb{Q})$ -modules that induces ϕ -integration.

We don't work directly with the associative differential algebra of singular cochains but with a cdga model of $\pi: E \to B$ as above which we denote by $\pi^*: B \to E$. This makes E a differential B-module. For differential (left) B-modules M and N we denote by $\operatorname{Hom}_B(M,N)$ the subcomplex of $\operatorname{Hom}_{\operatorname{grVect}}(M,N)=\oplus_i \prod_n \operatorname{Hom}_{\operatorname{Vect}}(M^{n+i},N^n)$ that are B-module homomorphisms, i.e. linear maps that satisfy $f(b\cdot m)=(-1)^{|f|\cdot|b|}b\cdot f(m)$. Then $\operatorname{Hom}_B(M,N)$ is a differential B-module with differential $D(f):=d_Nf-(-1)^{|f|}fd_M$ and module structure $(b\cdot f)(m):=b\cdot f(m)$.

We cannot expect to find a representative of ϕ -integration in $\mathsf{Hom}_B(\mathsf{E},\mathsf{B})$ for every cgda model of π . Instead one should consider *derived* mapping spaces, which will be sufficient for us to define as the (left) derived functor of $\mathsf{Hom}_B(\mathsf{-},N):\mathsf{Mod}_B\to\mathsf{Mod}_B^{op}$. The relevant resolutions that will appear in proofs and computations are defined as follows.

Definition 3.2. A B-module P is called **semifree** if it is the union of an increasing sequence of differential B-submodules $P(0) \subset P(1) \subset ...$ such that P(0) and each P(k)/P(k-1) are free B-modules on a basis of cycles. A **semifree resolution** of $M \in \mathsf{Mod}_B$ is a quasi-isomorphism $\varphi : P {\rightarrow} M$ of differential B-modules where P is semifree. The differential Ext groups are defined as $\mathsf{Ext}_B^*(M,N) := H^*(\mathsf{Hom}_B(P,N))$ for a semifree resolution of M.

The most important example of semifree modules in this context are relative Sullivan models of fibrations. A **relative Sullivan algebra** is a cdga of the form $(B \otimes \Lambda V, D)$ so that $id_B \otimes 1 : B \to B \otimes \Lambda V$ is a map of cdga's, and $V = \bigoplus_{p \geq 1} V^p$ is a graded vector space with an exhaustive filtration $V(0) \subset V(1) \subset \ldots$ of graded subspaces so that $D|_{1 \otimes V(0)} : V(0) \to B$ and $D|_{1 \otimes V(k)} : V(k) \to B \otimes \Lambda V(k-1)$. A relative Sullivan model of $\pi^* : B \to E$ is a quasi-isomorphism $E' \to E$ of B-algebras where E' is a relative Sullivan algebra.

The following properties of relative Sullivan models can be found in [FHT01] (with some additional assumptions about finiteness): A relative Sullivan algebra is a semifree B-module, so a Sullivan model $E' \to E$ is a semifree resolution of E. If B is a simply connected cdga and either the fibre or the base are of finite type then $E' \otimes_B \mathbb{Q} \cong (\Lambda V, d)$ is a Sullivan model of the fibre. Moreover, every cdga map $\pi^* : B \to E$ admits a relative Sullivan model $\varphi : E' \xrightarrow{\simeq} E$, which is a cofibrant replacement in the standard model structure on cdga. We denote the relative Sullivan model by $E' \to E$ to indicate this fact.

Proposition 3.3. Let $\pi: E \rightarrow B$ be a fibration with fibre F of 1-connected spaces and assume that $H^*(F; \mathbb{Q})$ is of finite type and nontrivial in degree d and vanishes above. Let $\pi^*: B \rightarrow E$ be a cdga model of π where B is simply connected, and denote by $E' \xrightarrow{\simeq} E$ a relative Sullivan model. Then the augmentation of B induces an isomorphism

$$\operatorname{Ext}^{-d}(\operatorname{E},\operatorname{B}) = H^{-d}(\operatorname{Hom}_{\operatorname{B}}(\operatorname{E}',\operatorname{B})) \xrightarrow{\cong} H^{-d}(\operatorname{Hom}_{\mathbb{Q}}(\operatorname{E}' \otimes_{\operatorname{B}} \mathbb{Q},\mathbb{Q})) \cong \operatorname{Hom}(H^{d}(F;\mathbb{Q}),\mathbb{Q})$$
$$[\Phi : \operatorname{E}' \to B] \longmapsto [\Phi \otimes_{\operatorname{B}} \mathbb{Q} : \operatorname{E}' \otimes_{\operatorname{B}} \mathbb{Q} \to \operatorname{B} \otimes_{\operatorname{B}} \mathbb{Q} \cong \mathbb{Q}]$$

In particular, for every $\phi \in \text{Hom}(H^d(F), \mathbb{Q})$ there is a unique element $[\Phi] \in H^{-d}(\text{Hom}_B(E', B))$ and the composition $H^*(E) \cong H^*(E') \xrightarrow{H(\Phi)} H^{*-d}(B) \cong H^{*-d}(B)$ is the same as ϕ -integration.

Proof. We assume that B is simply connected so that $B^{\geq p} \subset B$ is a B-submodule. Consider the exhaustive filtration of $Hom_B(E', B)$ given by $F^p = Hom_B(E', B^{\geq p})$. According to [Boa99, Thm.9.3] the corresponding spectral sequence is conditionally convergent to the completion

$$\begin{split} \lim_{\longleftarrow p} \operatorname{Hom}_{B}(E',B) / \operatorname{Hom}_{B}(E',B^{\geq p}) &\cong \lim_{\longleftarrow p} \operatorname{Hom}_{B}(E',B / B^{\geq p}) \\ &\cong \operatorname{Hom}_{B}(E',\lim_{\longleftarrow p} B / B^{\geq p}) \\ &\cong \operatorname{Hom}_{B}(E',B) \end{split}$$

where we have used that E' is a projective B-module for the first isomorphism. The E_1 -page is $H(\operatorname{Hom}_B(\mathrm{E}',\mathbb{Q}))\otimes B^p$. By assumption E' is a relative Sullivan algebra so that E' = $(\mathrm{B}\otimes\Lambda V,D)$ and E' $\otimes_B\mathbb{Q}=(\Lambda V,d)$ is a cdga model of the fibre. Hence, the E_1 -page can be simplified as $E_1^{p,q}=H^q((\Lambda V)^\vee)\otimes B^p$. Since B is simply connected, the differential on the E_1 -page is given by id $\otimes d_\mathrm{B}:E_1^{p,q}\to E_1^{p+1,q}$ and thus the spectral sequence has E_2 -page $E_2^{p,q}=H^p(\mathrm{B})\otimes H^q((\Lambda V)^\vee)$. In particular, the spectral sequence vanishes for q<-d. Since the gradings are such that the differentials are $d_r:E_r^{p,q}\to E_r^{p+r,q-r+1}$, there are only finitely many nontrivial differentials. This implies that the derived E_∞ -page is zero and so by [Boa99, Thm.7.1] the spectral sequence converges strongly

$$E_2^{p,q} = H^p(\mathsf{B}) \otimes H^q((\Lambda V)^\vee) \cong H^p(B) \otimes H_{-q}(F) \Rightarrow H(\mathsf{Hom}_\mathsf{B}(\mathsf{E}',\mathsf{B}))$$

and we can recover $H(\operatorname{Hom}_B(E',B))$ from the entries of the E_∞ -page. The only contribution with total degree -d comes from $E_\infty^{0,-d} \cong E_2^{0,-d} \cong \operatorname{Hom}(H^d(F;\mathbb{Q}),\mathbb{Q})$ which proves the first part of the statement.

It remains to show that for $\phi = H^d(\Phi \otimes_B \mathbb{Q}) : H^d(\Lambda V, d) \to \mathbb{Q}$ the induced ϕ -integration map coincides with $H(\Phi) : H^*(E') \to H^*(B)$. First, we note that $E' = (B \otimes \Lambda V, D)$ has a filtration $F^p = B^{\geq p} \otimes \Lambda V$ and that the corresponding spectral sequence converges as $E_2^{p,q} = H^p(B) \otimes H^q(\Lambda V) \Rightarrow H^{p+q}(E')$. In fact, B also has an analogous filtration $G^p = B^{\geq p}$ with only nontrivial differential on the E_1 -page. Then Φ induces a map between these two filtrations and the map on E_2 -pages is precisely ϕ -integration defined using this spectral sequence. As we have defined ϕ -integration using the Serre spectral sequence, it remains to show that this spectral sequence is isomorphic to the Serre spectral sequence. Grivel has shown in [Gri79] that $A_{PL}(B) \to A_{PL}(E)$ has a filtration which gives rise to the Serre spectral squence, and the construction is based on the construction of the Serre spectral sequence by Dress [Dre67]. Moreover, in the proof of [Gri79, Thm.6.4] he shows that the comparison map of $B \to E'$ with $A_{PL}(B) \to A_{PL}(E)$ is compatible with the above filtration on E' and the filtration on $A_{PL}(E)$ and induces an isomorphism on the E_2 -pages. This concludes the proof.

We can use Proposition 3.3 to build a representative of fibre integration for an oriented fibration $\pi: E \to B$ and oriented Poincaré fibre (X, ε_X) as follows: Consider a relative Sullivan model $\pi^*: B \to (B \otimes \Lambda V, D)$. Pick a chain level representative $\varepsilon_X \in \operatorname{Hom}^{-d}(\Lambda V, \mathbb{Q})$. The above Proposition implies that there is a cycle $\Pi \in \operatorname{Hom}^{-d}(E', B)$ unique up to chain homotopy that satisfies

$$\Pi(1 \otimes \chi) = \varepsilon_X(\chi) \in \mathbb{Q} = \mathbb{B}^0 \tag{3.3}$$

for all $\chi \in (\Lambda V)^d$, and that this chain map induces fibre integration on cohomology

$$\pi_{!}: H^{*}(E; \mathbb{O}) \cong H^{*}(E') \stackrel{H(\Pi)}{\longrightarrow} H^{*-d}(B) \cong H^{*-d}(B; \mathbb{O}).$$

We demonstrate this technique in the following example.

Example 3.4. Recall the relative Sullivan model of the universal 1-connected fibration for an even dimensional sphere $X = S^{2n}$ as discussed in Example 2.20. We choose as orientation $\varepsilon_X : H^d(X; \mathbb{Q}) \to \mathbb{Q}$ the homomorphism determined by $\varepsilon_X(x) = 1$. For degree reasons $\Pi(yx^k) = 0$ and since Π has to be a chain map we have $0 = \Pi(D(yx^k)) = \Pi(x^{k+2} - z_{4n}x^k)$. This determines a $\Lambda(z_{4n})$ -module map $\Pi : (\Lambda(z_{4n}, x, y), D) \to (\Lambda(z_{4n}), d = 0)$ by

$$\Pi(yx^k) = 0$$
 and $\Pi(x^n) = \begin{cases} 0 & n = 2k \\ z_{4n}^k & n = 2k + 1 \end{cases}$

which is a chain map by construction and induces fibre integration on cohomology as it satisfies (3.3).

3.3 Relation to parametrized stable homotopy theory. Fibre integration can also be viewed as a construction in fibrewise stable homotopy theory. More precisely, if $\pi : E \rightarrow B$ is a fibration then $\pi_+ := \pi \coprod Id_B : E_+ = E \coprod B \rightarrow B_+ = B \coprod B$ is a map of ex-spaces over B, both of which are exfibrations over B. The fibrewise suspension spectra $\Sigma_B^{\infty} E_+$ is strongly dualizable in the category

 $\operatorname{Sp}_{/B}$ of parametrized spectra over B if the fibre of π is equivalent to a finite CW complex [MS06]. The fibrewise suspension spectra $\Sigma_B^{\infty}B_+ = \mathbf{S}_B$ is self dual so that, if the fibre of π is equivalent to a finite CW complex, the dual of $\Sigma_B^{\infty}\pi_+$ is $D(\Sigma_B^{\infty}\pi_+): \Sigma_B^{\infty}B_+ \to D(\Sigma_B^{\infty}E_+) = F_B(\Sigma_B^{\infty}E_+, \mathbf{S}_B)$. If the fibre of $\pi: E \to B$ is Poincaré, we can combine this with

$$F_B(\Sigma_B^{\infty}E_+, \mathbf{S}_B) \rightarrow F_B(\Sigma_B^{\infty}E_+, H_B\mathbb{Z}) \rightarrow \Sigma_B^{\infty-d}E_+ \wedge_B H_B\mathbb{Z},$$

where the second map is an equivalence [HLLR17, Section 3.1], to obtain the version of fibre integration in parametrized stable homotopy theory

$$\pi_!: \Sigma_B^{\infty} B_+ {\longrightarrow} \Sigma_B^{\infty-d} E_+ \wedge_B H_B \mathbb{Z}.$$

Rationally, $H_B\mathbb{Z}$ is equivalent to S_B so that we get a map of rational parametrized suspension spectra. The set of homotopy classes of such maps has been computed in [FMT10, Thm.1.1] as a differential Ext group. This is consistent with our result. However, we would like to note that in order to apply Theorem 1.1 from [FMT10] to describe $\pi_!$ as an element in an differential Ext group, it is essential to use the equivalence $F_B(\Sigma_B^\infty E_+, H_B\mathbb{Z}) \to \Sigma_B^{\infty-d} E_+ \wedge_B H_B\mathbb{Z}$ so as to identify $\pi_!$ as a map of fibrewise suspension spectra. The equivalence of parametrizes spectra needs Poincaré duality of the fibre. In contrast, we don't need to assume this in order to find a unique representative of $\pi_!$ in the differential Ext groups because the construction works just as well for finding representatives of ϕ -integration in general.

Finally, we want to remark that fibre integration has also been described as elements in differential Ext groups in [FT09, Thm.A] for fibrations over Poincaré duality spaces and for pullbacks from such fibrations.

4 Algebraic definition of the fibrewise Euler class

We have defined the Euler class of fibrations with Poincaré fibre over Poincaré spaces in the introduction in (1.1). In this section, we will give a definition for arbitrary base spaces and show that it agrees with the definition of the fibrewise Euler class in [HLLR17]. In the following, we denote by $\mathbb{Q}[n]$ the graded vector space with \mathbb{Q} in degree n and for a B-module M define $M[n] := M \otimes \mathbb{Q}[n]$.

Proposition 4.1. Let $B \to E'$ be a relative Sullivan model of a fibration of 1-connected spaces $\pi : E \to B$ with Poincaré fibre X of formal dimension d. Let ε_X be the orientation and $\Pi \in \operatorname{Hom}_B^{-d}(E', B)$ be a corresponding cycle representing fibre integration. Then the map

$$\bar{\Pi}: \mathrm{E}'[-d] \longrightarrow \mathrm{Hom}_{\mathrm{B}}(\mathrm{E}',\mathrm{B})$$

$$e \longmapsto (e' \mapsto (-1)^{d+d\cdot|e|} \Pi(e\cdot e'))$$

is a quasi-isomorphism of B-modules.

Proof. It is a simple check that $\bar{\Pi}$ defines a B-module homomorphism. By assumption, $E' = (B \otimes \Lambda, D)$ is a relative Sullivan algebra and thus has a filtration which induces the Serre spectral sequence as discussed in the proof of Proposition 3.3. In the same proof we have described a filtration of $Hom_B(E', B)$ which strongly converges because there is a horizontal

vanishing line. The map $\bar{\Pi}$ is compatible with the two filtrations and induces a map of the associated spectral sequences. The induced map on the E_2 -page is given by

$$E_2^{p,q} = H^p(\mathsf{B}) \otimes H^q(\Lambda V, d) \overset{\mathrm{Id} \otimes \bar{\varepsilon}_X}{\longrightarrow} H^p(\mathsf{B}) \otimes \mathrm{Hom}(H^{q-d}(\Lambda V), \mathbb{Q})$$

where $\bar{\varepsilon}_X: H^q(\Lambda V) \to \operatorname{Hom}(H^{d-q}(\Lambda V), \mathbb{Q})$ is the adjoint of $H^q(\Lambda V) \otimes H^{d-q}(\Lambda V) \overset{\cup}{\to} H^d(\Lambda V) \overset{\varepsilon_X}{\to} \mathbb{Q}$. Since $(H^*(X; \mathbb{Q}), \varepsilon_X)$ is an oriented Poincaré duality algebra, $\bar{\Pi}$ induces an isomorphism of E_2 -pages.

With this we get an algebraic model of the Umkehr map of $\Delta : E \to E \times_B E$. Let $B \to E'$ be a relative Sullivan model of $\pi : E \to B$ as above. Then $E' \otimes_B E'$ is a Sullivan model of $E \times_B E$ and $\Pi \otimes \Pi : E' \otimes_B E' \to B \otimes_B B = B$ is a chain level representative of fibre integration for the parametrized product as it is a cycle and restricts to the induced orientation of the fibre $X \times X$. If we denote by $\Delta : E' \otimes E' \to E'$ the multiplication, we can provide the Umkehr map as the (dashed) lift in

$$E'[-d] \xrightarrow{---} E' \otimes_{B} E'[-2d]$$

$$\prod_{\square} \cong \qquad \qquad \overline{\square \otimes \square} \downarrow \simeq \qquad (4.1)$$

$$Hom_{B}(E', B) \xrightarrow{\Delta^{*}} Hom_{B'}(E' \otimes_{B} E', B)$$

uniquely up to homotopy, since E' is semifree. This determines the Umkehr map in the homotopy category of B-modules for any cdga model of the fibration, and enables us to identify a representative of the fibrewise Euler class in terms of the algebraic models.

Definition 4.2. Let $\pi: E \to B$ be a 1-connected oriented fibration with Poincaré fibre (X, ε_X) of formal dimension d. Let $B \to E'$ be a relative Sullivan model, then $e^{\mathrm{fw}}(\pi) := \Delta^* \Delta_!(1) \in H^d(E')$.

It remains to show that this coincides with the naive definition from the introduction which was only given when the base is a Poincaré complex. This has been carried out in greater generality in [RW17] and [HLLR17], but in our case there is a simple algebraic proof.

Lemma 4.3. Let $\pi: E \rightarrow B$ be an oriented fibration with fibre and base simply connected Poincaré complexes of formal dimension d respectively b. Let $B \rightarrow E'$ be a relative Sullivan model. Then the Umkehr map $\Delta_!: E' \rightarrow E' \otimes_B E'$ agrees on cohomology with $D_{E \times_B E}^{-1} \Delta_* D_E: H^*(E) \rightarrow H^{*+d}(E \times_B E)$ where $\Delta: E \rightarrow E \times_B E$ and $D_E = [E] \cap -$ and $D_{E \times_B E} = [E \times_B E] \cap -$ denote the Poincaré duality isomorphisms.

Proof. Since B is a Poincaré space, $B \rightarrow *$ is a fibration with Poincaré fibre and we can apply the characterisation of the previous section to get a chain level representative of fibre integration map $\Pi_B \in \operatorname{Hom}_{\mathbb{Q}}(B, \mathbb{Q})$ corresponding to evaluating a fundamental class. Suppose $B \rightarrow E'$ is a Sullivan model of the fibration and let $\Pi \in \operatorname{Hom}_{\mathbb{B}}(E', B)$ be a model of fibre integration of π . It is well known that $([B] \cap -) \circ \pi_! : H^{b+d}(E) \rightarrow H^b(B) \rightarrow \mathbb{Q}$ is an orientation of the Poincaré algebra $H^*(E; \mathbb{Q})$ itself. In particular, $\Pi_B \circ \Pi \in \operatorname{Hom}_{\mathbb{Q}}(E', \mathbb{Q})$ is a cycle that induces this orientation thus has to be fibre integration of $E \rightarrow *$. Denote $\Pi_E := \Pi_B \circ \Pi$ so that

$$\operatorname{Hom}_{B}(E',B) \stackrel{\simeq}{\longleftarrow} E'[-d] \stackrel{\simeq}{\longrightarrow} \operatorname{Hom}_{\mathbb{Q}}(E'),\mathbb{Q})$$

is a commuting diagram, where $(\Pi_B)_*\varphi = \Pi_B \circ \varphi$ for $\varphi \in \operatorname{Hom}_B(E',B)$). We can choose chain level representatives of fibre integration of $E \times_B E \to B$ and $E \times_B E \to *$ as $\Pi \otimes \Pi : E' \otimes_B E' \to B \otimes_B B \cong B$ and $\Pi_{E \times_B E} := \Pi_B \circ (\Pi \otimes \Pi)$ by the same arguments as above. We therefore have a commutative diagram

where the dashed maps denote the maps in the Umkehr maps in the homotopy category from (4.1) and using the Poincaré duality of E. The diagram shows that they induce the same map up to homotopy and therefore the same on cohomology.

Corollary 4.4. Let $\pi: E \to B$ be an oriented fibration of 1-connected spaces with Poincaré fibre X and relative Sullivan model $B \to E'$. Then $e^{fw}(\pi) = \Delta^* \Delta_![1] \in H^d(E')$ agrees with the fibrewise Euler class constructed in [HLLR17] in $H^d(E') \cong H^d(E; \mathbb{Q})$.

Proof. Since rational homology of a space B agrees with rational stably framed bordism, it suffices to evaluate the Euler class on bordism classes $[f:M\rightarrow B,\xi]\in\Omega^{fr}(B)\otimes\mathbb{Q}$, where we can assume by performing framed surgery that M is simply connected. The construction in [HLLR17] is natural under pullback $f^*: \mathrm{Sp}_{/B}\rightarrow\mathrm{Sp}_{/M}$, and they show in section 3.2 that the Euler class they define coincides for base spaces which are Poincaré with the class defined using the cohomological Umkehr maps from Poincaré duality.

The class defined Definition 4.2 is natural with respect to pullbacks as well: Let $f: B' \to B$ be a map of simply connected spaces and denote the pullback fibration by $\pi': f^*E \to B'$. Let $f^*: B \to B'$ be a cdga model of f. Then $f^*E':= B' \otimes_B E'$ is a relative Sullivan model of π' . If Π denotes a chain level representative of fibre integration then $f^*\Pi:= B' \otimes \Pi: B' \otimes_B E' \to B' \otimes_B B \cong B'$ is a representative of fibre integration of the pullback and if $\Delta_!$ denotes a choice of lift for the Umkehr map then

$$f^*\Delta_! := B' \otimes \Delta_! : f^*E' = B' \otimes_B E' \longrightarrow B' \otimes_B E \otimes_B E[-d] \cong (f^*E' \otimes_{B'} f^*E')[-d]$$

is a lift for the pullback which is natural with respect to f^* . If B' = M is a simply connected manifold, the Euler class agrees with class defined using the cohomological Umkehr maps from Poincaré duality by Lemma 4.3. Hence, the definitions of the fibrewise Euler class in Definition 4.2 and in [HLLR17] are both natural and agree on rational framed bordism classes. Therefore, our definition of the fibrewise Euler class agrees with that in [HLLR17].

4.1 Leray–Hirsch fibrations. In the case of fibrations $\pi: E \rightarrow B$ with oriented Poincaré fibre (X, ε_X) which are Leray–Hirsch, i.e. where the restriction map $H^*(E) \rightarrow H^*(X)$ is surjective, the definition of fibre integration and the fibrewise Euler class can be significantly simplified. Surjectivity of the restriction map implies that $H^*(E)$ is a free $H^*(B)$ -module, and we may denote by $1, e_1, \ldots, e_k \in H^*(E)$ a $H^*(B)$ -basis of the cohomology of E that restricts to a basis

 $1, x_1 = i^*(e_1), \dots, x_k = i^*(e_k) \in H^*(X)$ of the fibre. If X is a Poincaré complex of formal dimension d, we can order the basis such that $|e_k| = d$ and all other $|e_i|$ have lower degree. Since fibre integration is a $H^*(B)$ -module map. It suffices to determine $\pi_!$ on a basis and for degree reasons $\pi_!(e_i) = 0$ for i < k. If we set $\pi_!(e_k) = \varepsilon_X(x_k)$ this restricts to the orientation on the fibre hence determines fibre integration as

$$\pi_! \left(\sum_{i=0}^k b_i \cdot e_i \right) = \varepsilon_X(x_k) \cdot b_k \tag{4.2}$$

for $b_i \in H^*(B)$. Since the fibre is Poincaré, the (fibrewise) intersection pairing

$$\langle -, - \rangle : H^*(E) \otimes_{H^*(B)} H^*(E) \xrightarrow{\pi_!(-\cup -)} H^*(B)$$

$$\tag{4.3}$$

is non-degenerate. This enables us to mimic the construction of the Euler class as the dual of the fibrewise diagonal.

Proposition 4.5 ([RW18]). Let $\pi : E \to B$ be an oriented fibration with Poincaré fibre which is Leray–Hirsch. Let $e_0, \ldots, e_k \in H^*(E)$ be an $H^*(B)$ -module basis and denote by $e_0^{\sharp}, \ldots, e_k^{\sharp} \in H^*(E)$ the dual basis under the non-degenerate pairing (4.3). Then the fibrewise Euler class is

$$e^{\text{fw}}(\pi) = \sum_{i=0}^{k} (-1)^{|e_i|} e_i e_i^{\#} \in H^d(E; \mathbb{Q}).$$
(4.4)

Remark 4.6. We have already pointed out that Theorem 2.19 is particularly useful for elliptic spaces as the models are smaller than the model of Tanré. An elliptic space X with positive Euler characteristic is called **positively elliptic** and for such spaces, Halperin conjectured that the universal 1-connected fibration $B \mapsto Aut_{0,*}(X) \to B \mapsto Aut_0(X)$ is Leray–Hirsch. This is known to be true for a large number of examples [Mei82, ST87, Tho81]. Note that a positively elliptic space always satisfies (rational) Poincaré duality so that one can ask about Euler rings for these spaces. Moreover, if a positively elliptic satisfies the Halperin conjecture, fibre integration and the fibrewise Euler class are determined by (4.2) and (4.4) which is significantly easier than the general case we have discussed before.

Example 4.7. Let $X = S^{2n}$ be an even dimensional sphere and recall the cdga model and fibre integration from the Examples 2.20 and 3.4. Then 1 and x are a $H^*(B \, hAut_0(S^{2n}); \mathbb{Q})$ -basis of the cohomology of the total space that restricts to a basis of $H^*(S^{2n})$ on the fibre, i.e. the fibration is Leray-Hirsch. Note that the formula for fibre integration in (4.2) gives the same result as our construction of Π in Example 3.4. We can apply the above Proposition to find a representative of the fibrewise Euler class. The dual basis with respect to the pairing induced by $\pi_! = H(\Pi)$ is $x^\# = 1$ and $1^\# = x$ since $\pi_!(x \cdot 1^\#) = \pi_!(x^2) = 0$, and we find that the fibrewise Euler class is represented by $e^{fw}(\pi) = 2x$.

5 Computations and observations

First, we combine our previous computations on the universal 1-connected fibration of an even dimensional sphere to compute the Euler ring in this simple case.

Proposition 5.1. The Euler ring of an even dimensional sphere is $E^*(S^{2n}) \cong \mathbb{Q}[\kappa_2]$ and the relations are determined by $\kappa_2^k = 2^{k-1}\kappa_{2k}$. In particular, all odd κ_{2i+1} vanish.

Proof. We have seen in Example 4.7 that $e^{\mathrm{fw}}(\pi) = 2x \in \Lambda(x, y, z_{4n})$ and in Example 3.4 that fibre integration is given by $\Pi(x^{2k}) = 0$ and $\Pi(x^{2k+1}) = z_{4n}^k$. Thus $\kappa_{2k} = 2^{1-k}(2^3z_{4n})^k = 2^{1-k}\kappa_2^k$. Since $\pi_0(\mathrm{hAut}^+(S^{2n}))$ is trivial, the 1-connected universal fibration is the universal fibration and the result follows.

Of course, this is well known and there are easier ways to prove this. For example, it follows from Example 2.20 that BhAut⁺(S^{2n}) is rationally equivalent to $K(\mathbb{Q},4n)$, which reduces to proving this statement on linear bundles.

5.1 The Euler ring of complex projective space. We start by applying the results from section 2 to get a model of the universal 1-connected fibration π : B hAut_{0,*}($\mathbb{C}P^n$) \rightarrow B hAut₀($\mathbb{C}P^n$) with fibre $\mathbb{C}P^n$. In the following, we denote the minimal model of $\mathbb{C}P^n$ by $P_n := (\Lambda(x, y), |x| = 2, |y| = 2n + 1, d = x^{n+1} \partial/\partial y)$.

Proposition 5.2. A cdga model of the universal 1-connected $\mathbb{C}P^n$ -fibration is given by

$$B_n := \left(\mathbb{Q}[x_2, \dots, x_{n+1}], |x_i| = 2i, d = 0 \right) \xrightarrow{\pi} E_n := \left(B_n[x] / (x^{n+1} - \sum_{i=2}^{n+1} x_i \cdot x^{n+1-i}), |x| = 2, d = 0 \right). \tag{5.1}$$

In particular, the universal fibration is formal.

Proof. Note that $\operatorname{Der}^+(P_n)$ has a (vector space) basis given by $\theta_i := x^{n+1-i}\partial/\partial y$ for $i=1,\ldots,n+1$ of degree 2i-1 and $\eta:=\partial/\partial x$ of degree 2. The only non-trivial differential on the derivation Lie algebra is given by $[d,\eta]=-(n+1)\theta_1$. Denote by $\mathfrak{a}_n\subset\operatorname{Der}^+(P_n)$ the vector space spanned by the cycles $\theta_2,\ldots,\theta_{n+1}$ which generate the homology. This is an abelian dg Lie subalgebra with trivial differential and the inclusion $\mathfrak{a}_n\hookrightarrow\operatorname{Der}^+(P_n)$ induces a quasi-isomorphism of dg Lie algebras. Hence, the inclusion induces a quasi-isomorphism of cdga's

$$C_{CE}^*(\operatorname{Der}^+(P_n); \mathbb{Q}) \xrightarrow{\simeq} C_{CE}^*(\mathfrak{a}_n; \mathbb{Q}) \cong (\mathbb{Q}[x_2, \dots, x_{n+1}], d = 0),$$

where $x_i = (s\theta_i)^*$ and has degree 2i. Combining Theorem 2.19 with the above quasi-isomorphism shows that

$$C_{CE}^*(\mathfrak{a}_n; \mathbb{Q}) \xrightarrow{\eta_*} C_{CE}^*(\mathfrak{a}_n; P_n) \cong (C_{CE}^*(\mathfrak{a}_n; \mathbb{Q}) \otimes P_n, D)$$

is a relative Sullivan model of the universal 1-conneted fibration, where $\eta: \mathbb{Q} \to P_n$ is the unit. Here P_n is a left \mathfrak{a}_n -module by restricting the $\mathrm{Der}^+(P_n)$ -action. With respect to the chosen basis of \mathfrak{a}_n and the basis $\{x^k, yx^l\}_{k,l \geq 0}$, the formulas in (2.13) are

$$D(x) = d(x) - \sum_{i,k} (x^k)^* (\theta_i(x)) \cdot x_i x^k + (yx^k)^* (\theta_i(x)) \cdot x_i y x^k = 0,$$

$$D(y) = d(y) - \sum_{i,k} (x^k)^* ((\theta_i(y))) \cdot x_i x^k + (yx^k)^* (\theta_i(y)) \cdot x_i y x^k$$

$$= x^{n+1} - \sum_{i=2}^{n+1} x_i \cdot x^{n+1-i}.$$

Consider the map of B_n -algebras $C: C^*_{CE}(\mathfrak{a}_n; P_n) \to E_n$ determined by C(y) = 0 and C(x) = x. The cohomology of $C^*_{CE}(\mathfrak{a}_n; P_n)$ can be computed via the spectral sequence of a relative Sullivan algebra corresponding to the Serre spectral sequence. The spectral sequence degenerates on the E_2 -page and thus the cohomology of $C^*_{CE}(\mathfrak{a}_n; P_n)$ is a free module over $H^*_{CE}(\mathfrak{a}_n; \mathbb{Q}) \cong B_n$. Moreover, C sends the B_n -basis $\{1, [x], \ldots, [x^n]\}$ of $H^*_{CE}(\mathfrak{a}_n; P_n)$ to a B_n -basis of E_n . Hence, C is a quasi-isomorphism of B_n -algebras and in particular of cgda's.

Remark 5.3. The Halperin conjecture implies that one can always find an abelian Lie subalgebra $\mathfrak a$ of the derivations of the minimal model of a positively elliptic space as above that is quasi-isomorphic to it. This has been explained to the author by Alexander Berglund and it gives a shorter proof of the above statement than our original one. In particular, it implies that the spaces in the universal 1-connected fibration are rationally equivalent to $\prod_{i=1}^{N} K(\mathbb Q, 2n_i)$ so that the universal fibration is determined by a map of polynomial rings. This was already observed in [Kur10, Thm.1.1] with different methods. The paper [Kur10, Prop.1.3] also states the algebraic model for complex projective spaces, which we learned after the completion of this work.

Since the universal 1-connected fibration for $\mathbb{C}P^n$ is Leray–Hirsch we could compute the fibrewise Euler class in E_n using Proposition 4.5. But actually E_n is a complete intersection over B_n and we can use this to determine the fibrewise Euler class in a simpler way, using properties of complete intersections. More generally, the Halperin conjecture implies that $H^*(B \text{ hAut}_{0,*}(X))$ is a complete intersection over $H^*(B \text{ hAut}_0(X))$ for any positively elliptic space [Kur10, Thm.1.1] so that we get a new description of the fibrewise Euler class in this case. First, we will recall some properties from [dSL97] with notation adapted to our situation.

Definition 5.4. Let B be a commutative ring. A B-algebra E is called finite if it is finitely generated as a B-module. A finite B-algebra E is a **complete intersection** if there exists an $n \in \mathbb{N}$ and $f_1, \ldots, f_n \in B[x_1, \ldots, x_n]$ such that E is isomorphic to $B[x_1, \ldots, x_n]/(f_1, \ldots, f_n)$.

For any E-module M there is an E-module structure on the B-linear dual $\operatorname{Hom}_B(M,B)$ given by $(e \cdot \varphi)(m) := \varphi(e \cdot m)$ for $e \in E$, $m \in M$ and $\varphi \in \operatorname{Hom}_B(M,B)$. For a finite B-algebra E there is an isomorphism $\operatorname{Hom}_B(E,E) \cong E \otimes_B \operatorname{Hom}_B(E,B)$ so that we can define the trace of any B-endomorphism of E via the evaluation map $E \otimes_B \operatorname{Hom}_B(E,B) \to B$. In particular, any element in E defines an endomorphism by multiplication of which one can take the trace, giving an element $\operatorname{Tr}_{E/B} \in \operatorname{Hom}_B(E,B)$.

Proposition 5.5 ([dSL97]). Let B be a commutative ring and $f_1, \ldots, f_n \in B[x_1, \ldots, x_n]$ for a non-negative integer n. Assume that $E = B[x_1, \ldots, x_n]/(f_1, \ldots, f_n)$ is a finite B-algebra. Then

- (i) E is a projective B-module;
- (ii) $\operatorname{Hom}_{B}(E, B)$ is a free of rank 1 as E-module;
- (iii) there is a generator λ of $\operatorname{Hom}_B(E,B)$ as an E-module such that $\operatorname{Tr}_{E/B} = \det(\partial f_i/\partial x_i) \cdot \lambda$.

The equation in (iii) is the analogue for a complete intersection of the relation $\operatorname{trf}_{\pi}^*(-) = \pi_!(e^{\mathrm{fw}}(\pi) \cdot -)$ between the (Becker-Gottlieb) transfer $\operatorname{trf}_{\pi}^*: H^*(E) \to H^*(B)$, fibre integration and the fibrewise Euler class for a fibration $\pi: E \to B$ with Poincaré fibre X.

We make this precise as follows: let B be a polynomial ring over \mathbb{Q} on finitely many generators in positive even degrees and augmentation ε . Consider a complete intersection $E = B[x_1, \ldots, x_n]/(f_1, \ldots, f_n)$ where the x_i are all in positive even degree. Then E is a projective B-module by (i) and since B is a connected graded algebra, E is a free B-module. Denote by $\pi : \langle E \rangle \rightarrow \langle B \rangle$ the geometric realization [BG76, Ch.5] of a relative Sullivan model of $B \rightarrow E$.

Theorem 5.6. Let B and E be as above. Then the homotopy fibre of $\pi: \langle E \rangle \to \langle B \rangle$ has the rational homotopy type of the complete intersection $X = \mathbb{Q}[x_1, \ldots, x_n]/(\bar{f_1}, \ldots, \bar{f_n})$ over \mathbb{Q} with $\bar{f_i} = \epsilon \circ f_i$. In particular, X is a rational Poincaré space with an orientation defined by $\varepsilon_X(\det(\partial \bar{f_i}/\partial x_j)) := \chi(\langle X \rangle)$ and the fibrewise Euler class is represented by

$$e^{\text{fw}}(\pi) = \det\left(\frac{\partial f_i}{\partial x_i}\right) \in E.$$
 (5.2)

Proof. Consider the differential *B*-algebra $E' := (B \otimes \Lambda(x_i, y_i), D)$ with $|y_i| = |f_i| - 1$ and $D(y_i) := f_i$. This is relative version of a pure Sullivan model and the *B*-algebra map $C : E' \rightarrow E$ determined by $C(y_i) = 0$ and $C(x_i) = x_i$ is a quasi-isomorphism of differential *B*-algebras by the same argument as in the proof of Proposition 5.2. Moreover, E' and B are 1-connected Sullivan algebras so that $B \rightarrow A_{PL}(\langle B \rangle)$ and $E' \rightarrow A_{PL}(\langle E' \rangle)$ are quasi-isomorphisms and hence $B \rightarrow E'$ is a relative Sullivan model of the fibration $\pi : \langle E' \rangle \rightarrow \langle B \rangle$. Because B is of finite type, the map $\langle E' \rangle \rightarrow \langle B \rangle$ is algebraic and thus a cdga model of the fibre is given by $\mathbb{Q} \otimes_B E'$ [Tho81] which is a pure Sullivan model. Hence, the homotopy fibre has the rational homotopy type of a positively elliptic space [FHT01] with top degree generator given by $\det(\partial \bar{f_i}/\partial x_j) \in X$ [Mur93, ST87], so that ε_X as defined above is a valid choice of orientation.

Note that $\langle E' \rangle \rightarrow \langle B \rangle$ is formal so that fibre integration $\pi_! : H(\langle E' \rangle; \mathbb{Q}) \cong E \rightarrow H(\langle B \rangle; \mathbb{Q}) \cong B$ determines a map of B-modules with respect to our choice of orientation. According to Proposition 5.5, there is a B-module map $\lambda : E \rightarrow B$ satisfying (iii). It follows for degree reasons that λ and $\pi_!$ agree up to a scalar in \mathbb{Q}^\times . One can show that the trace and the transfer agree using the general theory in [DP80], or by directly checking that $\mathrm{Tr}_{E/B}(x) = \pi_!(e^{\mathrm{fw}}(\pi) \cdot x)$, as has been done in [RW18, Lemma 2.3] using that E is a finitely generated free B-algebra. Hence, if we evaluate for x = 1 we find that $\lambda(\det(\partial f_i/\partial x_j)) = \mathrm{Tr}_{E/B}(1) = \pi_!(e^{\mathrm{fw}}(\pi)) = \chi(\langle X \rangle)$, which agrees with $\pi_!(\det(\partial f_i/\partial x_j))$ by our choice of orientation ε_X . Hence, it follows that $\pi_! = \lambda$ and consequently

$$\det(\partial f_i/\partial x_j)\cdot \pi_! = \det(\partial f_i/\partial x_j)\cdot \lambda = \operatorname{Tr}_{E/B} = \operatorname{tr} f_\pi^* = e^{\operatorname{fw}}(\pi)\cdot \pi_!$$

under the identification of $H^*(\langle E' \rangle; \mathbb{Q})$ with E and $H^*(\langle B \rangle; \mathbb{Q})$ with B. The claim follows because B-linear dual of E is a free E-module by (ii).

We can apply the above theorem to the complete intersection in Proposition 5.2 where we use the integral Poincaré duality to fix an orientation on $\mathbb{C}P^n$ by $\varepsilon_{\mathbb{C}P^n}(x^n) = 1$.

Corollary 5.7. The fibrewise Euler class in E_n is represented by

$$e^{\text{fw}}(\pi) = (n+1) \cdot x^n - \sum_{i=2}^n (n+1-i) \cdot x_i \cdot x^{n-i} \in E_n.$$
 (5.3)

Remark 5.8. In all examples of positively elliptic spaces X with cohomology ring $H^*(X; \mathbb{Q}) = \mathbb{Q}[x_1, \ldots, x_n]/(f_1, \ldots, f_n)$ we have encountered, one can always scale the functions such that they all have integral coefficients and such that $\mathbb{Z}[x_1, \ldots, x_n]/(f_1, \ldots, f_n)$ is a complete intersection over \mathbb{Z} , i.e. there seems to be a distinguished choice for f_1, \ldots, f_n . Moreover, this seems to describe the image $H^*(X; \mathbb{Z}) \to H^*(X; \mathbb{Q})$. We don't know if this is true in general.

Finally, we are able to determine the Euler ring of complex projective spaces.

Theorem 5.9. The Euler ring is generated by $\kappa_1, \ldots, \kappa_{n-1}, \kappa_{n+1}$ and they are algebraically independent. Hence, $E^*(\mathbb{C}P^n) \cong \mathbb{Q}[\kappa_1, \ldots, \kappa_{n-1}, \kappa_{n+1}]$.

Proof. We can compute the Euler ring of the universal 1-connected fibration $E_0^*(\mathbb{C}P^n)$ using the algebraic models and the representative of the fibrewise Euler class and fibre integration. In order to extend this computation to the full Euler ring in B hAut⁺($\mathbb{C}P^n$), we need to study the spectral sequence of the fibre sequence B hAut₀(X) \to B hAut⁺(X) \to B π_0 (hAut⁺(X)) which describes the universal covering of B hAut⁺(X). The inclusion $\mathbb{C}P^n \to \mathbb{C}P^\infty$ induces a bijection $[\mathbb{C}P^n, \mathbb{C}P^n] \to [\mathbb{C}P^n, \mathbb{C}P^\infty] \cong H^2(\mathbb{C}P^n; \mathbb{Z})$ and the homotopy equivalences correspond to generators of $H^2(\mathbb{C}P^n; \mathbb{Z})$. Thus π_0 (hAut⁺($\mathbb{C}P^n$)) is finite and it follows by transfer that $H^*(\mathbb{B} \text{ hAut}^+(\mathbb{C}P^n); \mathbb{Q})$ injects into $H^*(\mathbb{B} \text{ hAut}_0(X); \mathbb{Q})$. In particular, the pullback induces an isomorphism $E^*(\mathbb{C}P^n) \xrightarrow{\cong} E_0^*(\mathbb{C}P^n)$.

The Euler ring is defined as a subring generated by κ_i for $i \geq 1$ and we will first identify a finite generating set following [RW18]. Consider the B_n -endomorphism given by $-\cdot e^{\mathrm{fw}}(\pi)$: $E_n \rightarrow E_n$. As both algebras are concentrated in even degrees and E_n is a free B_n -module, we can apply the Cayley–Hamilton theorem to get that $e^{\mathrm{fw}}(\pi)$ is a root of its characteristic polynomial p(e), which is monic and of degree n+1. The coefficients are certain polynomials in $\mathrm{Tr}_{E_n/B_n}(e^{\mathrm{fw}}(\pi)^j)$. Since $\mathrm{Tr}_{E_n/B_n}(c) = \pi_!((e^{\mathrm{fw}}(\pi) \cdot c))$ for all $c \in E_n$ as discussed in the the proof of Theorem 5.6, it follows by fibre integrating the characteristic polynomial $p(e^{\mathrm{fw}}(\pi))$ that κ_n lies in the ring generated by $\kappa_1, \ldots, \kappa_{n-1}$. The same does not hold for κ_{n+1} because the constant term of p(e) contains a κ_{n+1} and fibre integrating $p(e^{\mathrm{fw}}(\pi)) \cdot e^{\mathrm{fw}}(\pi)$ will not give a new relation. Fibre integrating $p(e^{\mathrm{fw}}(\pi)) \cdot e^{\mathrm{fw}}(\pi)$ for $k \geq 2$ expresses κ_{n+k} in terms of lower degree κ_i and thus $\kappa_1, \ldots, \kappa_{n-1}, \kappa_{n+1}$ generate the Euler ring.

It remains to identify relations between the polynomials $\kappa_i = \kappa_i(x_2, \dots, x_{n+1}) \in B_n$ which we get by fibre integrating the representative of the fibrewise Euler class in (5.3). We will show that they are in fact algebraically independent by showing that $\det(\partial \kappa_i/\partial x_j)$ is non-zero. It turns out that the polynomials representing the κ_i are quite complicated so that it is difficult to give a closed formula for the determinant of the Jacobian. We will resolve this issue by focussing on the terms containing x_{n+1} because it is the variable of the highest degree and it is not contained in $e^{fw}(\pi)$ so that it only arises through fibre integrating x^k for k > n. It will be sufficient to consider elements modulo decomposables, i.e. for $x, y \in B_n$ then $x \sim y$ if $x - y \in (B_n^+)^2$. We will start with the following observation about fibre integration.

1. If
$$k = 2, ..., n + 1$$
 then $\pi_!(x^{n+k}) \sim x_k \in B_n$.

Proof. Rewriting x^{n+2} in terms of the module basis $\{1, x, ..., x^n\}$, one can see that $\pi_!(x^{n+2}) = x_2$. Then $\pi_!(x^{n+k}) = \pi_!(x^{n+1} \cdot x^{k-1}) = \sum_{i=2}^{n+1} x_i \cdot \pi_!(x^{n+k-i})$ and by induction over k, the only indecomposable contribution is for i = k.

2. For i = 1, ..., n-1 the highest power of x_{n+1} in $\kappa_i(x_2, ..., x_{n+1})$ is i-1 and the coefficient $c_i \in \mathbb{Q}[x_2, ..., x_n]$ of x_{n+1}^{i-1} satisfies $c_i \sim i(n+1)^i(n-i) \cdot x_{n+1-i}$.

Proof. It follows from degree considerations that the highest power of x_{n+1} is i-1 and $c_i = A \cdot x_{n+1-i}$ + decomposables. It remains to determine the coefficient A. When expanding $e^{\text{fw}}(\pi)$ using (5.3), the only relevant contributions are

$$(n+1)^{i+1}x^{n(i+1)} - (i+1)(n+1-(n+1-i))x_{n+1-i}x^{n-(n+1-i)} \cdot (n+1)^{i}x^{ni}$$

= $(n+1)^{i+1}(x^{n+1})^{i-1} \cdot x^{2n-i+1} - i(i+1)(n+1)^{i}x_{n+1-i} \cdot (x^{n+1})^{i-1} \cdot x^{n}$

Now rewrite $x^{n+1} = \sum_{i=2}^{n+1} x_i x^{n+1-i}$ and collect all terms containing x_{n+1}^{i-1} and $x_{n+1}^{i-2} x_{n+1-i}$ (we can ignore the rest because it cannot contribute to A) to get

$$(n+1)^{i+1}(x_{n+1}^{i-1}+(i-1)x_{n+1}^{i-2}x_{n+1-i}\cdot x^i)\cdot x^{2n-i+1}-i(i+1)(n+1)^ix_{n+1-i}\cdot x_{n+1}^{i-1}\cdot x^n$$

The statement follows by fibre integrating and discarding decomposables as in 1 above.

3. The highest contribution of x_{n+1} in κ_{n+1} is $(n+1)^{n+2} \cdot x_{n+1}^n$.

Proof. The expression for $e^{\text{fw}}(\pi)^{n+2}$ contains the summand $(n+1)^{n+2} \cdot x^{n(n+2)} = (n+1)^{n+2} \cdot (x^{n+1})^n \cdot x^n$. This is the only summand that fibre integrates to a multiple of x_{n+1}^n , i.e. $\kappa_{n+1} = (n+1)^{n+2} \cdot x_{n+1}^n + \dots$ where we can ignore all other terms.

We can now analyze $\det(\partial \kappa_i/\partial x_i)$ which contains the summand

$$\frac{\partial \kappa_1}{\partial x_n} \cdot \frac{\partial \kappa_2}{\partial x_{n-1}} \cdot \dots \cdot \frac{\partial \kappa_{n-1}}{\partial x_2} \cdot \frac{\partial \kappa_{n+1}}{\partial x_{n+1}}.$$

It follows from **2** and **3** that the above expression contains $C \cdot x_{n+1}^N$, where C is a non-zero constant and $N = \frac{1}{2}n(n-1)$. This is the only possible way to get a monomial in $\det\left(\frac{\partial \kappa_i}{\partial x_j}\right)$ that contains only x_{n+1} . Hence, the determinant does not vanish and the generating set $\kappa_1, \ldots, \kappa_{n-1}, \kappa_{n+1}$ is algebraically independent.

Remark 5.10. Theorem 5.9 has been studied in the smooth case for n = 2 in [RW18] by studying the natural smooth 2-torus action on $\mathbb{C}P^2$ and that implies our result in this case as well. This has been extended by Dexter Chua to $n \le 4$, but for large n the algebra becomes intractable.

Remark 5.11. Consider $H^*(B \text{ hAut}_0(\mathbb{C}P^n); \mathbb{Q})$ as a module over $E^*(\mathbb{C}P^n)$. By [Smi95, Cor.6.7.11] it is finitely generated as a $E^*(\mathbb{C}P^n)$ -module if and only if $\kappa_1, \ldots, \kappa_{n-1}, \kappa_{n+1} \in \mathbb{Q}[x_2, \ldots, x_{n+1}] \cong H^*(B \text{ hAut}_0(\mathbb{C}P^n); \mathbb{Q})$ is a regular sequence. This can be checked by computing the radical of $(\kappa_1, \ldots, \kappa_{n-1}, \kappa_{n+1})$, as for a parameter ideal the radical is the unique graded maximal ideal $H^+(B \text{ hAut}_0(\mathbb{C}P^n); \mathbb{Q})$. With this criterion one can check that $H^*(B \text{ hAut}_0(\mathbb{C}P^n); \mathbb{Q})$ is finite over the Euler ring for n = 1, 2 (for n = 1 this is obvious) but not for n = 3, 4. We don't know how to check this condition for $n \geq 5$ in general but we expect that the cohomology of B hAut $_0(\mathbb{C}P^n)$ is not finite over the Euler ring in this case. This shows that one cannot hope to find a version of Theorem 3.1 in [RW18] with Euler rings instead of tautological rings in order to find a lower bound on the Krull dimensions via Torus actions.

We end this section with the following observation: the natural map $PU(n+1) \rightarrow hAut_0(\mathbb{C}P^n)$ induces a map between classifying spaces, and it is a classical result that $H^*(BPU(n+1);\mathbb{Q}) \cong \mathbb{Q}[q_2,\ldots,q_{n+1}]$ with generators of degree $|q_i|=2i$ which are related to Chern classes. Thus BPU(n+1) has the same rational cohomology ring as $BhAut_0(\mathbb{C}P^n)$, and in particular both are rationally equivalent to $\prod_{i=2}^{n+1} K(\mathbb{Q},2i)$ by its intrinsic rational formality. Hence, $BPU(n+1) \simeq_{\mathbb{Q}} BhAut_0(\mathbb{C}P^n)$ and it has been shown that the natural map induces a rational equivalence.

Theorem ([Sas74, Kur11]). *The natural map* B PU(n + 1) \longrightarrow B hAut₀($\mathbb{C}P^n$) *is a rational homotopy equivalence.*

Sasao proves this statement using classical homotopy theory to study the fibration sequence $\operatorname{Map}_{l_m}(\mathbb{C}P^m,\mathbb{C}P^n) \to \operatorname{Map}_{l_{m-1}}(\mathbb{C}P^{m-1},\mathbb{C}P^n)$ for m < n, where the subscript denote the connected component of the inclusion $\iota_m:\mathbb{C}P^m \hookrightarrow \mathbb{C}P^n$. In particular, he doesn't just obtain results about rational homotopy groups but also statements about the connectivity of the map. For example, he proves that the inclusion $\operatorname{PU}(n+1) \to \operatorname{Map}_{\operatorname{id}}(\mathbb{C}P^n,\mathbb{C}P^n)$ induces an isomorphism on fundamental groups and $\pi_1(\operatorname{PU}(n+1)) \cong \mathbb{Z}/(n+1)\mathbb{Z}$. In contrast, Kuribayahshi uses tools from rational homotopy theory as well. His result is based on a rational model of the evaluation map $\operatorname{hAut}_0(X) \times X \to X$.

We want sketch a simple proof of this result that uses the algebraic models of the $\mathbb{C}P^n$ -fibrations over $B \text{ hAut}_0(\mathbb{C}P^n)$ and B PU(n+1). It will be easier to prove that the natural map $B \text{ SU}(n+1) \to B \text{ hAut}_0(\mathbb{C}P^n)$ is a rational equivalence. This implies the theorem as the natural map $B \text{ SU}(n+1) \to B \text{ PU}(n+1)$ is a rational equivalence because $SU(n+1) \to PU(n+1)$ is an (n+1)-fold normal covering.

Proof. The statement is a direct consequence of the projective bundle formula which describes the algebraic structure of the projectivization of a complex vector bundle in terms of the Chern classes. Let $i: BSU(n+1) \rightarrow BU(n+1)$ be the natural map induced by inclusion and $\gamma_{n+1} \rightarrow BU(n+1)$ the tautological vector bundle. Then there is an algebra isomorphism

$$H^*(\mathbb{P}(i^*\gamma_{n+1});\mathbb{Q}) \cong H^*(\mathbb{B}\operatorname{SU}(n+1);\mathbb{Q})[t]/(t^{n+1} + \sum_{k=2}^{n+1} (-1)^k i^* c_k \cdot t^{n+1-k})$$

where t is the Euler class of the canonical line bundle $L \to \mathbb{P}(i^*\gamma_{n+1})$ and $i^*c_k \in H^{2k}(B\,SU(n+1);\mathbb{Q})$ is the restriction of the k-th Chern class. The projectivization $\mathbb{P}(i^*\gamma_{n+1})$ is the pullback of the universal $\mathbb{C}P^n$ -fibration along the natural map $c: B\,SU(n+1) \to B\,hAut_0(\mathbb{C}P^n)$. As both E_n and $H^2(\mathbb{P}(i^*\gamma_{n+1});\mathbb{Q})$ are 1-dimensional in degree 2, the pullback of $x \in E_n^2$ agrees with t up to a scalar, which is 1 for our choice of orientation. It follows that $c^*(x_k) = -(-1)^k i^* c_k$ and hence c^* induces an isomorphism of complete intersections.

The above theorem has the following non-obvious consequence.

Corollary 5.12. The natural map B Diff⁺($\mathbb{C}P^n$) \longrightarrow B hAut⁺($\mathbb{C}P^n$) induces and injection on rational cohomology.

Proof. Consider the composition $BPU(n+1) \rightarrow BDiff^+(\mathbb{C}P^n) \rightarrow BhAut^+(\mathbb{C}P^n)$ which factors through the respective connected components of the identity. As $\pi_0(hAut^+(\mathbb{C}P^n))$ is finite, the cohomology of $BhAut^+(\mathbb{C}P^n)$ injects into the cohomology of $BhAut_0(\mathbb{C}P^n)$ as the ring of $\pi_0(hAut^+(\mathbb{C}P^n))$ -invariants. The statement then follows from the previous theorem.

5.2 The Euler ring of products of odd spheres. In the following, let X be simply connected and rationally equivalent to a product of (simply connected) odd dimensional spheres $\prod_{i=1}^{2N} S^{2n_i+1}$. Denote by F the set of all factors, then X has a minimal Sullivan model $A_F = (\Lambda(x_f)_{f \in F}, |x_f| = 2n_f + 1, d = 0)$. We choose a total ordering < of F so that f < f' implies $|f| \le |f'|$ where $|f| = 2n_f + 1$ is the dimension of the corresponding sphere. This determines a basis $\{x_S := \prod_{f \in S} x_f\}_{S \subset F}$ where the product $\prod_{f \in S} x_f$ is with respect to the induced total order on S. We define $x_\emptyset := 1$. The dg Lie algebra of derivations of A_F is graded and has the following basis

$$\operatorname{Der}^+(A_F) = \bigoplus_{n \ge -1} \mathbb{Q} \{ \eta_S^f \mid S \subset (F, <), \#S = n + 1 \text{ and } |S| < |f| \}$$

where $|S| = \sum_{f \in S} |f|$ and $\eta_S^f = x_S \cdot \frac{\partial}{\partial x_f}$. The bracket is given by

$$[\eta_S^f, \eta_{S'}^{f'}] = \begin{cases} \operatorname{sgn}(S, f, S', f') \eta_{S \sqcup (S' \setminus f)}^{f'} & \text{if } f \in S' \\ -(-1)^{|\eta_S^f| \cdot |\eta_{S'}^{f'}|} \cdot \operatorname{sgn}(S', f', S, f) \eta_{S' \sqcup (S \setminus f')}^f & \text{if } f' \in S \\ 0 & \text{otherwise} \end{cases}$$

where the exact form of the signs can easily be worked out but will not matter for the Euler ring, and hence we omit a discussion here. We denote the corresponding dual basis of $(s \operatorname{Der}^+(A_F))^\vee$ by y_S^f . In the following, denote by $B_F := C_{CE}^*(\operatorname{Der}^+(A_F); \mathbb{Q})$ the cdga model of the base with differential d and by $E_F := (B_F \otimes A_F, D)$ the cdga model of the total space. Using (2.13), the we find that the relative Sullivan model of the universal fibration is determined by

$$d(y_S^f) = (-1)^{|f|-|S|} \sum_{S_1 \sqcup S_2 = S, g \in S_{< f} \setminus S} \operatorname{sgn}(S_1, g, S_2 \cup g, f) \cdot y_{S_2 \cup g}^f y_{S_1}^g$$

$$D(x_f) = -\sum_{S \subset S_{< f} \text{ with } |S| < |f|} y_S^f \wedge x_S.$$

Recall that a chain level representative of fibre integration $\Pi \in \operatorname{Hom}_{B_F}(E_F, B_F)$ is uniquely determined up to homotopy by (3.3). We choose as orientation of X the homomorphism determined by $\varepsilon_X(x_F) = 1$. Since Π lowers degree by |F|, it can only be nontrivial on x_F so that setting

$$\Pi(1 \otimes x_S) := \begin{cases} 1 & \text{if } S = F \\ 0 & \text{otherwise} \end{cases}$$
 (5.4)

determines the unique element in the 1-dimensional vector space $\operatorname{Hom}_{B_F}^{-d}(E_F, B_F)$ that satisfies (3.3). Hence, by Proposition 3.3 is has to be a representative of fibre integration and in particular is a cycle, which can also be easily checked. We construct a representative of $\Delta_!: E_F[-|F|] \to E_F \otimes_{B_F} E_F[-2|F|]$ from section 4 to compute the fibrewise Euler class.

But first, note that the integral fibrewise Euler class of a Poincaré complex of odd formal dimension is a 2-torsion element. From the point of view of rational homotopy theory, there is no structural difference in the algebraic model of the universal fibration of X if #F is odd or even, so that one can expect the following.

Proposition 5.13. Let X be rationally equivalent to a product of odd dimensional spheres. Then the fibrewise Euler class $e^{fw}(\pi) \in H^{|X|}(B \, hAut_{0,*}(X); \mathbb{Q})$ vanishes. In particular, $E_0^*(X) = \mathbb{Q}$.

Proof. Recall that the description of $\Delta_!$: $E_F[-|F|] \to E_F \otimes_{B_F} E_F[-2|F|]$ uses the quasi-isomorphism $\bar{\Pi}: E_F \to \operatorname{Hom}_{B_F}(E_F, B_F)$. Because A_F is finite dimensional, $\operatorname{Hom}_{B_F}(E_F, B_F)$ is a semifree module itself and it is minimal because A_F is a minimal Sullivan model. Hence, $\bar{\Pi}$ and $\overline{\Pi \otimes \Pi}$ are isomorphisms of B_F -modules by uniqueness of minimal free resolutions [FHT01, Ex.8,Ch.6] so that

$$\Delta_{!}: E_{F}[-|F|] \xrightarrow{\bar{\Pi}} \operatorname{Hom}_{B_{F}}(E_{F}, B_{F}) \xrightarrow{\Delta^{*}} \operatorname{Hom}_{B_{F}}(E_{F} \otimes_{B_{F}} E_{F}, B_{F}) \xrightarrow{(\overline{\Pi \otimes \Pi})^{-1}} E_{F} \otimes_{B_{F}} E_{F}[-2|F|].$$

The composition of $\bar{\Pi}$ with the vector space isomorphism $\operatorname{Hom}_{B_F}(E_F, B_F) \cong (A_F)^{\vee} \otimes B_F$ is given by $\bar{\varepsilon}_X \otimes \operatorname{Id}_{B_F}$ where $\bar{\varepsilon}_X : A_F \to (A_F)^{\vee}$ is the adjoint of $\varepsilon_X : A_F \otimes A_F \to \mathbb{Q}$. The same statement holds for $\overline{\Pi \otimes \Pi}$ with the appropriate choice of orientation on $X \times X$ given by $\varepsilon_{X \times X} := \varepsilon_X \otimes \varepsilon_X : (A_F \otimes A_F)^{\otimes 2} \to \mathbb{Q}$. This is very special to the products of odd dimensional spheres and holds because the Sullivan model is finite dimensional, which is only ever true for this situation.

Note that $\Delta^*\bar{\Pi}(1)$ is contained in $(A_F \otimes A_F)^{\vee} \otimes 1$ so that $(\overline{\Pi \otimes \Pi})^{-1}\Delta^*\bar{\Pi}(1)$ is in $A_F \otimes A_F \otimes 1 \subset E_F \otimes_{B_F} E_F$. A direct computation shows that

$$\Delta_!(1) = \sum_{S_1 \sqcup S_2 = F} \pm x_{S_1} \otimes x_{S_2} \in E_F \otimes_{B_F} E_F$$

for some signs that can be worked out. Hence, the fibrewise Euler class is $e^{\mathrm{fw}}(\pi) = \Delta^* \circ \Delta_!(1) = \sum_{S_1 \sqcup S_2 = F} \pm x_F$ and since $\Pi(e^{fw}(\pi)) = \chi(X) = 0$, the summands must cancel.

In order to understand the fibrewise Euler class in $H^d(B hAut_*^+(X))$; $\mathbb{Q})$ we need to study the universal covering spectral sequence of $B hAut_{0,*}^+(X) \to B hAut_*^+(X) \to B \pi_0(hAut_*^+(X))$. We do not yet know how to do this in general, but in some cases we can deduce that $E^*(X) = \mathbb{Q}$.

Theorem 5.14. Let X be either rationally equivalent to $(S^{2k+1})^{\times n}$ or a finite CW complex rationally equivalent to $S^{2k+1} \times S^{2l+1}$ for 1 < k < l and n even. Then $E^*(X) = \mathbb{Q}$.

Proof. In the second case, we will show that $\pi_0(hAut^+(X))$ is finite so that the universal covering spectral sequence collapses and the cohomology of $BhAut^+_*(X)$ injects as the invariants of $H^*(BhAut_{*,0}(X);\mathbb{Q})$ with respect to the action of $\pi_0(hAut^+_0(X))$. In particular, the fibrewise Euler class is trivial in $H^{2(k+l+1)}(BhAut^+_*(X);\mathbb{Q})$ and therefore the Euler ring is \mathbb{Q} .

By [Sul77, Thm.10.3] the group $\pi_0(\text{hAut}^+(X))$ is commensurable with an arithmetic subgroup of the homotopy classes automorphisms of A_F . If $X \simeq_{\mathbb{Q}} S^{2k+1} \times S^{2l+1}$ then the group of automorphisms of A_F modulo homotopy is $\mathbb{Q}^\times \times \mathbb{Q}^\times$ and the arithmetic subgroups of this linear algebraic group are finite and hence by commensurability so is $\pi_0(\text{hAut}^+(X))$.

In the first case, the cdga model E_F of BhAut_{0,*}(X) is a free algebra on $x_1, \ldots, x_n, y^1, \ldots, y^n$ with differential $D(x_i) = -y^i$. Hence $H(E_F) = \mathbb{Q}$ and thus BhAut_{0,*}(X) is rationally contractible. Therefore, BhAut⁺(X) \to B π_0 (hAut⁺(X)) is a rational equivalence and thus every class in $H^*(BhAut^+_*(X);\mathbb{Q})$ is pulled back from B π_0 (hAut⁺(X)). In particular, $e^{fw}(\pi) = H^*e$ for some

 $e \in H^{n \cdot (2k+1)}(B \pi_0(hAut^+(X)); \mathbb{Q})$. By commutativity of the following diagram

$$B hAut_{*}^{+}(X) \xrightarrow{\mathcal{H}} B \pi_{0}(hAut_{*}^{+}(X))$$

$$\downarrow^{\pi} \qquad \qquad \parallel$$

$$B hAut^{+}(X) \xrightarrow{h} B \pi_{0}(hAut^{+}(X)),$$

the fibrewise Euler class $e^{\text{fw}}(\pi) = H^*e = \pi^*h^*e$ is pulled back from the base. Hence, fibre integrating powers of the fibrewise Euler class gives $\pi_!(\pi^*(h^*e)^k) = (h^*e)^k \cdot \pi_!(1) = 0$.

We expect that one can extend Proposition 5.13 to show that $e^{\mathrm{fw}}(\pi) \in H^d(\mathrm{B}\,\mathrm{hAut}^+_*(X);\mathbb{Q})$ always vanishes by studying the universal fibration of $X_\mathbb{Q}$ instead. As rationalization is functorial there is a (continuous) map $\mathrm{B}\,\mathrm{hAut}^+_*(X) \to \mathrm{B}\,\mathrm{hAut}^+_*(X_\mathbb{Q})$ and the Euler class in the cohomology of $\mathrm{B}\,\mathrm{hAut}^+_*(X_\mathbb{Q})$ pulls back to the Euler class in $\mathrm{B}\,\mathrm{hAut}^+_*(X)$. Since the homotopy automorphism of $X_\mathbb{Q}$ are easier to study, we expect that one can find a differential graded algebra model of $C^*(\mathrm{B}\,\mathrm{hAut}^+_*(X_\mathbb{Q});\mathbb{Q})$, and further that one can find models of fibre integration similarly as we have discussed to show that the fibrewise Euler class vanishes.

References

- [BG76] A. K. Bousfield and V. K. A. M. Gugenheim, *On PL De Rham theory and rational homotopy type*, Mem. Amer. Math. Soc. **8** (1976), no. 179, ix+94.
- [BM14] A. Berglund and I. Madsen, Rational homotopy theory of automorphisms of manifolds, ArXiv e-prints (2014).
- [Boa99] J. M. Boardman, Conditionally convergent spectral sequences, Homotopy invariant algebraic structures (Baltimore, MD, 1998), Contemp. Math., vol. 239, Amer. Math. Soc., Providence, RI, 1999, pp. 49–84.
- [DP80] A. Dold and D. Puppe, *Duality, trace, and transfer*, Proceedings of the International Conference on Geometric Topology (Warsaw, 1978), PWN, Warsaw, 1980, pp. 81–102.
- [Dre67] A. Dress, Zur Spectralsequenz von Faserungen, Invent. Math. 3 (1967), 172–178.
- [dSL97] B. de Smit and H. W. Lenstra, Jr., Finite complete intersection algebras and the completeness radical, J. Algebra 196 (1997), no. 2, 520–531.
- [EM66] S. Eilenberg and J. C. Moore, Homology and fibrations. I. Coalgebras, cotensor product and its derived functors, Comment. Math. Helv. 40 (1966), 199–236.
- [FHT01] Y. Félix, S. Halperin, and J.-C. Thomas, *Rational homotopy theory*, Graduate Texts in Mathematics, vol. 205, Springer-Verlag, New York, 2001.
- [FMT10] Y. Félix, A. Murillo, and D. Tanré, Fibrewise stable rational homotopy, J. Topol. 3 (2010), no. 4, 743–758.
- [FT09] Y. Félix and J.-C. Thomas, String topology on Gorenstein spaces, Math. Ann. 345 (2009), no. 2, 417–452.
- [Gat97] J.-B. Gatsinzi, *The homotopy Lie algebra of classifying spaces*, J. Pure Appl. Algebra **120** (1997), no. 3, 281–289.
- [GGRW17] S. Galatius, I. Grigoriev, and O. Randal-Williams, *Tautological rings for high-dimensional manifolds*, Compos. Math. 153 (2017), no. 4, 851–866.
- [Gri79] P.-P. Grivel, Formes différentielles et suites spectrales, Ann. Inst. Fourier (Grenoble) **29** (1979), no. 3, ix, 17–37.
- [Gri17] I. Grigoriev, Relations among characteristic classes of manifold bundles, Geom. Topol. 21 (2017), no. 4, 2015–2048.
- [Hal83] S. Halperin, Lectures on minimal models, Mém. Soc. Math. France (N.S.) (1983), no. 9-10, 261.
- [HLLR17] F. Hebestreit, M. Land, W. Lück, and O. Randal-Williams, *A vanishing theorem for tautological classes of aspherical manifolds*, ArXiv e-prints (2017).
- [Kur10] K. Kuribayashi, *On the rational cohomology of the total space of the universal fibration with an elliptic fibre*, Homotopy theory of function spaces and related topics, Contemp. Math., vol. 519, Amer. Math. Soc., Providence, RI, 2010, pp. 165–179.

- [Kur11] _____, Rational visibility of a Lie group in the monoid of self-homotopy equivalences of a homogeneous space, Homology Homotopy Appl. 13 (2011), no. 1, 349–379.
- [LV12] J.-L. Loday and B. Vallette, *Algebraic operads*, Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], vol. 346, Springer, Heidelberg, 2012.
- [May75] J. P. May, Classifying spaces and fibrations, Mem. Amer. Math. Soc. 1 (1975), no. 1, 155, xiii+98.
- [Mei82] W. Meier, Rational universal fibrations and flag manifolds, Math. Ann. 258 (1981/82), no. 3, 329–340.
- [MS06] J. P. May and J. Sigurdsson, *Parametrized homotopy theory*, Mathematical Surveys and Monographs, vol. 132, American Mathematical Society, Providence, RI, 2006.
- [Mum83] D. Mumford, *Towards an enumerative geometry of the moduli space of curves*, Arithmetic and geometry, Vol. II, Progr. Math., vol. 36, Birkhäuser Boston, Boston, MA, 1983, pp. 271–328.
- [Mur93] A. Murillo, The top cohomology class of certain spaces, J. Pure Appl. Algebra 84 (1993), no. 2, 209–214.
- [Qui69] D. Quillen, Rational homotopy theory, Ann. of Math. (2) 90 (1969), 205–295.
- [RW17] O. Randal-Williams, An upper bound for the pseudoisotopy stable range, Math. Ann. **368** (2017), no. 3-4, 1081–1094.
- [RW18] _____, Some phenomena in tautological rings of manifolds, Selecta Math. (N.S.) 24 (2018), no. 4, 3835–3873.
- [Sas74] S. Sasao, *The homotopy of Map* (*CP*^m, *CP*ⁿ), J. London Math. Soc. (2) 8 (1974), 193–197.
- [Smi95] L. Smith, *Polynomial invariants of finite groups*, Research Notes in Mathematics, vol. 6, A K Peters, Ltd., Wellesley, MA, 1995.
- [SS12] M. Schlessinger and J. Stasheff, Deformation theory and rational homotopy type, ArXiv e-prints (2012).
- [ST87] H. Shiga and M. Tezuka, *Rational fibrations, homogeneous spaces with positive Euler characteristics and Jacobians*, Ann. Inst. Fourier (Grenoble) **37** (1987), no. 1, 81–106.
- [Sta63] J. Stasheff, A classification theorem for fibre spaces, Topology 2 (1963), 239–246.
- [Sul77] D. Sullivan, *Infinitesimal computations in topology*, Inst. Hautes Études Sci. Publ. Math. (1977), no. 47, 269–331 (1978).
- [Tan83] D. Tanré, *Homotopie rationnelle: modèles de Chen, Quillen, Sullivan,* Lecture Notes in Mathematics, vol. 1025, Springer-Verlag, Berlin, 1983.
- [Tho81] J.-C. Thomas, Rational homotopy of Serre fibrations, Ann. Inst. Fourier (Grenoble) 31 (1981), no. 3, v, 71–90.
- [Wei94] C. A. Weibel, *An introduction to homological algebra*, Cambridge Studies in Advanced Mathematics, vol. 38, Cambridge University Press, Cambridge, 1994.