

Geometric Estimates in Interpolation by Linear Functions on a Euclidean Ball

Mikhail Nevskii*

May 9, 2019

Abstract

Let B_n be the Euclidean unit ball in \mathbb{R}^n given by the inequality $\|x\| \leq 1$, $\|x\| := \left(\sum_{i=1}^n x_i^2 \right)^{\frac{1}{2}}$. By $C(B_n)$ we mean the space of continuous functions $f : B_n \rightarrow \mathbb{R}$ with the norm $\|f\|_{C(B_n)} := \max_{x \in B_n} |f(x)|$. The symbol $\Pi_1(\mathbb{R}^n)$ denotes the set of polynomials in n variables of degree ≤ 1 , i. e., the set of linear functions upon \mathbb{R}^n . Assume $x^{(1)}, \dots, x^{(n+1)}$ are the vertices of an n -dimensional nondegenerate simplex $S \subset B_n$. The interpolation projector $P : C(B_n) \rightarrow \Pi_1(\mathbb{R}^n)$ corresponding to S is defined by the equalities $Pf(x^{(j)}) = f(x^{(j)})$. Denote by $\|P\|_{B_n}$ the norm of P as an operator from $C(B_n)$ onto $C(B_n)$. We describe the approach in which $\|P\|_{B_n}$ can be estimated from below via the volume of S .

Keywords: simplex, ball, linear interpolation, projector, norm, estimate

1 Main Definitions

We always suppose $n \in \mathbb{N}$. An element $x \in \mathbb{R}^n$ will be written in the form $x = (x_1, \dots, x_n)$. By definition,

$$\|x\| := \left(\sum_{i=1}^n x_i^2 \right)^{\frac{1}{2}}, \quad B_n := \{x \in \mathbb{R}^n : \|x\| \leq 1\}, \quad Q_n := [0, 1]^n.$$

*Department of Mathematics, P.G. Demidov Yaroslavl State University, Sovetskaya str., 14, Yaroslavl, 150003, Russia orcid.org/0000-0002-6392-7618 mnevsk55@yandex.ru

The notation $L(n) \asymp M(n)$ means that there exist $c_1, c_2 > 0$ not depending on n such that $c_1 M(n) \leq L(n) \leq c_2 M(n)$. By $\Pi_1(\mathbb{R}^n)$ we mean the set of polynomials in n variables ≤ 1 , i.e., the set of linear functions upon \mathbb{R}^n .

Let S be a nondegenerate simplex in \mathbb{R}^n with vertices $x^{(j)} = (x_1^{(j)}, \dots, x_n^{(j)})$, $1 \leq j \leq n+1$. Consider *the vertex matrix (or the node matrix)*

$$\mathbf{A} := \begin{pmatrix} x_1^{(1)} & \dots & x_n^{(1)} & 1 \\ x_1^{(2)} & \dots & x_n^{(2)} & 1 \\ \vdots & \vdots & \vdots & \vdots \\ x_1^{(n+1)} & \dots & x_n^{(n+1)} & 1 \end{pmatrix}.$$

We have the equality $\text{vol}(S) = \frac{|\det(\mathbf{A})|}{n!}$. Put $\mathbf{A}^{-1} = (l_{ij})$. Let us define λ_j as polynomials from $\Pi_1(\mathbb{R}^n)$ whose coefficients form the columns of \mathbf{A}^{-1} , i.e., $\lambda_j(x) := l_{1j}x_1 + \dots + l_{nj}x_n + l_{n+1,j}$. We call λ_j *the basic Lagrange polynomials* corresponding to this simplex. The numbers $\lambda_j(x)$ are the barycentric coordinates of a point $x \in \mathbb{R}^n$ with respect to S .

For a convex body $\Omega \subset \mathbb{R}^n$, denote by $C(\Omega)$ the space of continuous functions $f : \Omega \rightarrow \mathbb{R}$ with the uniform norm

$$\|f\|_{C(\Omega)} := \max_{x \in \Omega} |f(x)|.$$

We say that an interpolation projector $P : C(\Omega) \rightarrow \Pi_1(\mathbb{R}^n)$ corresponds to a simplex $S \subset \Omega$ if interpolation nodes of P coincide with vertices $x^{(j)}$ of this simplex. Projector P is given by the equalities $Pf(x^{(j)}) = f(x^{(j)})$. We have the following analogue of the Lagrange interpolation formula:

$$Pf(x) = \sum_{j=1}^{n+1} f(x^{(j)}) \lambda_j(x). \quad (1)$$

Denote by $\|P\|_\Omega$ the norm of P as an operator from $C(\Omega)$ to $C(\Omega)$. It follows from (1) that

$$\|P\|_\Omega = \max_{x \in \Omega} \sum_{j=1}^{n+1} |\lambda_j(x)|.$$

If Ω is a convex polytope in \mathbb{R}^n (e.g., $\Omega = Q_n$), this equality is equivalent to the formula

$$\|P\|_\Omega = \max_{x \in \text{ver}(\Omega)} \sum_{j=1}^{n+1} |\lambda_j(x)|,$$

where $\text{ver}(\Omega)$ is the set of vertices of Ω .

Let us define $\theta_n(\Omega)$ as the minimal value of $\|P\|_\Omega$ under the conditions $x^{(j)} \in \Omega$. In the case $\Omega = Q_n$ various relations for the numbers $\theta_n(\Omega)$, including the equivalence $\theta_n(\Omega) \asymp \sqrt{n}$, were obtained by the author earlier. This results are systematized in [1]. Further on some estimates were improved (see [2], [3], [4] and references in these papers).

This paper deals with the case $\Omega = B_n$. We describe the approach when the norm of an interpolation projector $P : C(B_n) \rightarrow \Pi_1(\mathbb{R}^n)$ can be estimated from below through the volume of the corresponding simplex. The essential feature of this approach is the application of the classical Legendre polynomials. We prove that $\theta_n(B_n) \asymp \sqrt{n}$. In other words, the interpolation projector corresponding to a regular simplex inscribed into the boundary sphere has the norm equivalent to the minimal possible.

2 Estimation of $\|P\|_{B_n}$ via the Volume of S

The *standardized Legendre polynomial of degree n* is the function

$$\chi_n(t) := \frac{1}{2^n n!} [(t^2 - 1)^n]^{(n)}$$

(the Rodrigues formula). For properties of χ_n , see [6], [7]. The Legendre polynomials are orthogonal on $[-1, 1]$ with respect to the weight $w(t) = 1$. It is known that $\chi_n(1) = 1$; if $n \geq 1$, then $\chi_n(t)$ strictly increases for $t \geq 1$. Denote by χ_n^{-1} the function inverse to χ_n on the halfline $[1, +\infty)$.

The appearance of Legendre polynomials in our questions is connected with such their property. For $\gamma \geq 1$, consider the set

$$E_{n,\gamma} := \left\{ x \in \mathbb{R}^n : \sum_{j=1}^n |x_j| + \left| 1 - \sum_{j=1}^n x_j \right| \leq \gamma \right\}.$$

In 2003 the author established that

$$\text{mes}_n(E_{n,\gamma}) = \frac{\chi_n(\gamma)}{n!}. \quad (2)$$

(the proof is also given in [1]). Utilizing this equality he managed to obtain the lower estimates for projector's norms related to linear interpolation on the unit cube Q_n . The following relations take place:

$$\theta_n(Q_n) \geq \chi_n^{-1} \left(\frac{1}{\nu_n} \right) = \chi_n^{-1} \left(\frac{n!}{h_n} \right). \quad (3)$$

Here ν_n is the maximum volume of a simplex contained in Q_n and h_n is the maximum value of 0/1-determinant of order n . Applying the properties of

χ_n the author got from (3) some more visible inequalities, e. g., $\theta_n(Q_n) > \frac{1}{e}\sqrt{n-1}$. This estimate occurred to be sharp concerning to n which led to the equivalence $\theta_n(Q_n) \asymp \sqrt{n}$.

Further on we will extend this approach to linear interpolation of functions given on the unit ball B_n . Denote $\varkappa_n := \text{vol}(B_n)$. By σ_n we mean the volume of a regular simplex inscribed into B_n .

Theorem 1. *Assume $P : C(B_n) \rightarrow \Pi_1(\mathbb{R}^n)$ is an arbitrary interpolation projector. Then for the corresponding simplex $S \subset B_n$ and the node matrix \mathbf{A} we have*

$$\|P\|_{B_n} \geq \chi_n^{-1} \left(\frac{\varkappa_n}{\text{vol}(S)} \right) = \chi_n^{-1} \left(\frac{n! \varkappa_n}{|\det(\mathbf{A})|} \right). \quad (4)$$

Proof. As known, a regular simplex which is inscribed into a ball has the maximum possible volume among all simplices being contained in this ball. Therefore, $\text{vol}(S) = \frac{|\det(\mathbf{A})|}{n!} \leq \sigma_n$. For each $i = 1, \dots, n$, let us subtract from the i th row of \mathbf{A} its $(n+1)$ th row. Denote by \mathbf{B} the submatrix of order n which stands in the first n rows and columns of the result matrix. Then

$$|\det(\mathbf{B})| = |\det(\mathbf{A})| = n! \text{vol}(S) \leq n! \sigma_n.$$

In other words,

$$\frac{n! \sigma_n}{|\det(\mathbf{B})|} \geq 1. \quad (5)$$

Let $x^{(j)}$ be the vertices and λ_j be the basic Lagrange polynomials of the simplex S . Since $\lambda_1(x), \dots, \lambda_{n+1}(x)$ are the barycentric coordinates of a point x , we have

$$\|P\|_{B_n} = \max_{x \in B_n} \sum_{j=1}^{n+1} |\lambda_j(x)| = \max \left\{ \sum_{j=1}^{n+1} |\beta_j| : \sum_{j=1}^{n+1} \beta_j = 1, \sum_{j=1}^{n+1} \beta_j x^{(j)} \in B_n \right\}.$$

Let us replace β_{n+1} with the equal value $1 - \sum_{j=1}^n \beta_j$. The condition $\sum_{j=1}^{n+1} \beta_j x^{(j)} \in B_n$ is equivalent to $\sum_{j=1}^n \beta_j (x^{(j)} - x^{(n+1)}) \in B' := B_n - x^{(n+1)}$. This means

$$\|P\|_{B_n} = \max \left\{ \sum_{j=1}^n |\beta_j| + \left| 1 - \sum_{j=1}^n \beta_j \right| \right\}. \quad (6)$$

Maximum in (6) is taken upon β_j such that $\sum_{j=1}^n \beta_j (x^{(j)} - x^{(n+1)}) \in B'$.

Consider the nondegenerate linear operator $F : \mathbb{R}^n \rightarrow \mathbb{R}^n$ which maps a point $\beta = (\beta_1, \dots, \beta_n)$ into a point $x = (x_1, \dots, x_n)$ according to the rule

$$x = F(\beta) := \sum_{j=1}^n \beta_j (x^{(j)} - x^{(n+1)}).$$

We have the matrix equality $F(\beta) = (\beta_1, \dots, \beta_n)\mathbf{B}$, where \mathbf{B} is the above introduced $(n \times n)$ -matrix with the elements $b_{ij} = x_j^{(i)} - x_j^{(n+1)}$. Put

$$\gamma^* := \chi_n^{-1} \left(\frac{n! \varkappa_n}{|\det(\mathbf{B})|} \right).$$

Since $\varkappa_n \geq \sigma_n$, it follows from (5) that γ^* is defined correctly. Note the equality

$$\chi_n(\gamma^*) = \frac{n! \varkappa_n}{|\det(\mathbf{B})|}. \quad (7)$$

Now suppose $1 \leq \gamma < \gamma^*$,

$$E_{n,\gamma} = \left\{ \beta = (\beta_1, \dots, \beta_n) \in \mathbb{R}^n : \sum_{j=1}^n |\beta_j| + \left| 1 - \sum_{j=1}^n \beta_j \right| \leq \gamma \right\}.$$

Let us show that $B' \not\subset F(E_{n,\gamma})$. It is sufficient to get the inequality $\text{mes}_n(F(E_{n,\gamma})) < \text{mes}_n(B') = \varkappa_n$. This is really so:

$$\begin{aligned} \text{mes}_n(F(E_{n,\gamma})) &< \text{mes}_n(F(E_{n,\gamma^*})) = |\det \mathbf{B}| \cdot \text{mes}_n(E_{n,\gamma^*}) = \\ &= |\det \mathbf{B}| \cdot \frac{\chi_n(\gamma^*)}{n!} = \varkappa_n. \end{aligned}$$

We have applied (2) and (7). Thus, for every $\varepsilon > 0$, there exists a point $x^{(\varepsilon)}$ with the properties

$$x^{(\varepsilon)} = \sum_{j=1}^n \beta_j^{(\varepsilon)} (x^{(j)} - x^{(n+1)}) \in B', \quad \sum_{j=1}^n |\beta_j^{(\varepsilon)}| + \left| 1 - \sum_{j=1}^n \beta_j^{(\varepsilon)} \right| \geq \gamma^* - \varepsilon.$$

In view of (6) this gives $\|P\|_{B_n} \geq \gamma^* - \varepsilon$. Since $\varepsilon > 0$ is an arbitrary, we obtain

$$\|P\| \geq \gamma^* = \chi_n^{-1} \left(\frac{n! \varkappa_n}{|\det(\mathbf{B})|} \right) = \chi_n^{-1} \left(\frac{n! \varkappa_n}{|\det(\mathbf{A})|} \right) = \chi_n^{-1} \left(\frac{\varkappa_n}{\text{vol}(S)} \right).$$

The theorem is proved. □

Corollary 1. *For each n ,*

$$\theta_n(B_n) \geq \chi_n^{-1} \left(\frac{\varkappa_n}{\sigma_n} \right). \quad (8)$$

Proof. Let $P : C(B_n) \rightarrow \Pi_1(\mathbb{R}^n)$ be an arbitrary interpolation projector. Since the volume of the corresponding simplex is not greater than σ_n , inequality (4) yields

$$\|P\|_{B_n} \geq \chi_n^{-1} \left(\frac{\varkappa_n}{\text{vol}(S)} \right) \geq \chi_n^{-1} \left(\frac{\varkappa_n}{\sigma_n} \right).$$

This gives (8). □

It is known that

$$\varkappa_n = \frac{\pi^{\frac{n}{2}}}{\Gamma\left(\frac{n}{2} + 1\right)}, \quad \sigma_n = \frac{1}{n!} \sqrt{n+1} \left(\frac{n+1}{n} \right)^{\frac{n}{2}}, \quad (9)$$

$$\varkappa_{2k} = \frac{\pi^k}{k!}, \quad \varkappa_{2k+1} = \frac{2^{k+1}\pi^k}{(2k+1)!!} = \frac{2(k!)(4\pi)^k}{(2k+1)!}. \quad (10)$$

Therefore, the estimate (8) can be made more concrete.

Corollary 2. *For every n ,*

$$\theta_n(B_n) \geq \chi_n^{-1} \left(\frac{\pi^{\frac{n}{2}} n!}{\Gamma\left(\frac{n}{2} + 1\right) \sqrt{n+1} \left(\frac{n+1}{n}\right)^{\frac{n}{2}}} \right). \quad (11)$$

If $n = 2k$, then (11) is equivalent to the inequality

$$\theta_{2k}(B_{2k}) \geq \chi_{2k}^{-1} \left(\frac{\pi^k (2k)!}{k! \sqrt{2k+1} \left(\frac{2k+1}{2k}\right)^k} \right). \quad (12)$$

For $n = 2k + 1$ we have

$$\theta_{2k+1}(B_{2k+1}) \geq \chi_{2k+1}^{-1} \left(\frac{2(k!)(4\pi)^k}{\sqrt{2k+2} \left(\frac{2k+2}{2k+1}\right)^{\frac{2k+1}{2}}} \right). \quad (13)$$

Proof. It is sufficient to apply (8), (9), and (10). □

Corollary 3. *Suppose P is a minimal interpolation projector, i. e., $\|P\|_{B_n} = \theta_n(B_n)$. Then*

$$\text{vol}(S) \geq \frac{\varkappa_n}{\chi_n(\sqrt{n+1})} = \frac{\pi^{\frac{n}{2}}}{\Gamma\left(\frac{n}{2} + 1\right) \chi_n(\sqrt{n+1})}. \quad (14)$$

Proof. It was proved in [5] that $\theta_n(B_n) \leq \sqrt{n+1}$. Hence, if P is a minimal projector, then

$$\sqrt{n+1} \geq \|P\|_{B_n} \geq \chi_n^{-1} \left(\frac{\varkappa_n}{\text{vol}(S)} \right).$$

We have made use of (4). It remains to compare the boundary values and take into account (9). \square

Since $|\det(\mathbf{A})| = n! \text{vol}(S)$, relation (14) also implies an estimate for the determinant of the node matrix corresponding to a minimal projector. Namely, $|\det(\mathbf{A})| \geq \varepsilon_n$, where ε_n is $n!$ times more than the right-hand part of (14). Without giving the value of ε_n this restriction was used in [5].

3 The Relation $\theta_n(B_n) \asymp \sqrt{n}$

The Stirling formula $n! = \sqrt{2\pi n} \left(\frac{n}{e}\right)^n e^{\frac{\zeta_n}{12n}}$, $0 < \zeta_n < 1$, yields

$$\sqrt{2\pi n} \left(\frac{n}{e}\right)^n < n! < \sqrt{2\pi n} \left(\frac{n}{e}\right)^n e^{\frac{1}{12n}}. \quad (15)$$

Also we will need the following estimates which were proved in [1, Section 3.4.2]:

$$\chi_{2k}^{-1}(s) > \left(\frac{(k!)^2 s}{(2k)!} \right)^{\frac{1}{2k}}, \quad \chi_{2k+1}^{-1}(s) > \left(\frac{(k+1)! k! s}{(2k+1)!} \right)^{\frac{1}{2k+1}}. \quad (16)$$

Theorem 2. *There exist a constant $c > 0$ not depending on n such that*

$$\theta_n(B_n) > c\sqrt{n}. \quad (17)$$

The inequality (17) takes place, e. g., with the constant

$$c = \frac{\sqrt[3]{\pi}}{\sqrt{12e} \cdot \sqrt[6]{3}} = 0.2135\dots$$

Proof. First let $n = 2k$ be even. Making use of (12) and (16), we get

$$\theta_{2k}(B_{2k}) \geq \chi_{2k}^{-1} \left(\frac{\pi^k (2k)!}{k! \sqrt{2k+1} \left(\frac{2k+1}{2k}\right)^k} \right) > \left(\frac{\pi^k (2k)! (k!)^2}{k! \sqrt{2k+1} \left(\frac{2k+1}{2k}\right)^k (2k)!} \right)^{\frac{1}{2k}} =$$

$$= \left(\frac{\pi^k k!}{\sqrt{2k+1} \left(\frac{2k+1}{2k} \right)^k} \right)^{\frac{1}{2k}}.$$

Let us estimate $k!$ from below applying (15):

$$\theta_{2k}(B_{2k}) > \left(\frac{\pi^k \sqrt{2\pi k} \left(\frac{k}{e} \right)^k}{\sqrt{2k+1} \left(\frac{2k+1}{2k} \right)^k} \right)^{\frac{1}{2k}} = \sqrt{\frac{\pi}{e}} \left(\frac{2\pi k}{2k+1} \right)^{\frac{1}{4k}} \sqrt{\frac{2k}{2k+1}} \cdot \sqrt{k}.$$

Since $n = 2k$, for each even n ,

$$\theta_n(B_n) > \sqrt{\frac{\pi}{2e}} \left(\frac{\pi n}{n+1} \right)^{\frac{1}{2n}} \sqrt{\frac{n}{n+1}} \cdot \sqrt{n} > \sqrt{\frac{\pi}{3e}} \cdot \sqrt{n}. \quad (18)$$

We have utilized the inequalities $\frac{\pi n}{n+1} > 1$, $\frac{n}{n+1} > \frac{2}{3}$. Remark that $\sqrt{\frac{\pi}{3e}} = 0.6206\dots$

Now suppose n is odd, i. e., $n = 2k + 1$. In this case (13) and (16) give

$$\begin{aligned} \theta_{2k+1}(B_{2k+1}) &\geq \chi_{2k+1}^{-1} \left(\frac{2(k!)(4\pi)^k}{\sqrt{2k+2} \left(\frac{2k+2}{2k+1} \right)^{\frac{2k+1}{2}}} \right) > \\ &> \left(\frac{2(k!)(4\pi)^k (k+1)!k!}{\sqrt{2k+2} \left(\frac{2k+2}{2k+1} \right)^{\frac{2k+1}{2}} (2k+1)!} \right)^{\frac{1}{2k+1}} = A^{\frac{1}{2k+1}} \cdot B^{\frac{1}{2k+1}}; \\ A &:= \frac{(k!)^2 (k+1)!}{(2k+1)!}, \quad B := \frac{\sqrt{2}(4\pi)^k}{\sqrt{k+1} \left(\frac{2k+2}{2k+1} \right)^{\frac{2k+1}{2}}}. \end{aligned}$$

From (15), it follows that

$$\begin{aligned} A &> \frac{2\pi k \left(\frac{k}{e} \right)^{2k} \cdot \sqrt{2\pi(k+1)} \left(\frac{k+1}{e} \right)^{k+1}}{\sqrt{2\pi(2k+1)} \left(\frac{2k+1}{e} \right)^{2k+1} e^{\frac{1}{12(2k+1)}}} = \\ &= 2\pi e^{-k - \frac{1}{12(2k+1)}} k^{2k+1} (k+1)^{k+1+\frac{1}{2}} (2k+1)^{-\frac{1}{2}-(2k+1)}, \\ A^{\frac{1}{2k+1}} &> (2\pi)^{\frac{1}{2k+1}} e^{-\frac{k}{2k+1} - \frac{1}{12(2k+1)^2}} k(k+1)^{\frac{k+1+\frac{1}{2}}{2k+1}} (2k+1)^{-\frac{1}{2(2k+1)}-1}. \end{aligned}$$

Also

$$\begin{aligned} (2\pi)^{\frac{1}{2k+1}} &> 1, \quad \frac{k}{2k+1} + \frac{1}{12(2k+1)^2} < \frac{1}{2}, \\ k(2k+1)^{-\frac{1}{2(2k+1)}-1} &= \frac{k}{2k+1} (2k+1)^{-\frac{1}{2(2k+1)}} > \frac{1}{3} (2k+1)^{-\frac{1}{2(2k+1)}} > \frac{1}{3\sqrt[6]{3}}. \end{aligned}$$

Consequently,

$$A^{\frac{1}{2k+1}} > \frac{1}{3\sqrt[6]{3}\sqrt{e}}(k+1)^{\frac{k+1+\frac{1}{2}}{2k+1}} > \frac{1}{3\sqrt[6]{3}\sqrt{e}}\sqrt{k+1}.$$

Now let us estimate $B^{\frac{1}{2k+1}}$:

$$\begin{aligned} B^{\frac{1}{2k+1}} &= \left(\frac{\sqrt{2}(4\pi)^k}{\sqrt{k+1} \left(\frac{2k+2}{2k+1}\right)^{\frac{2k+1}{2}}} \right)^{\frac{1}{2k+1}} = \\ &= (\sqrt{2})^{\frac{1}{2k+1}} (4\pi)^{\frac{k}{2k+1}} \sqrt{\frac{2k+1}{2k+2}} (k+1)^{-\frac{1}{2(2k+1)}} > \\ &> \sqrt[3]{4\pi} \cdot \sqrt{\frac{3}{4}} \cdot 2^{-\frac{1}{6}} = \sqrt{\frac{3}{2}} \cdot \sqrt[3]{\pi}. \end{aligned}$$

Therefore,

$$\theta_{2k+1}(B_{2k+1}) > A^{\frac{1}{2k+1}} \cdot B^{\frac{1}{2k+1}} > \sqrt{\frac{3}{2}} \cdot \frac{\sqrt[3]{\pi}}{3\sqrt[6]{3}\sqrt{e}} \cdot \sqrt{k+1} = \frac{\sqrt[3]{\pi}}{\sqrt{6e}\sqrt[6]{3}} \cdot \sqrt{k+1}.$$

Since $n = 2k + 1$, we have $\sqrt{k+1} = \frac{1}{\sqrt{2}}\sqrt{2k+2} > \frac{1}{\sqrt{2}}\sqrt{n}$. For every odd n ,

$$\theta_n(B_n) > \frac{\sqrt[3]{\pi}}{\sqrt{12e} \cdot \sqrt[6]{3}} \cdot \sqrt{n}. \quad (19)$$

The constant 0.2135... from the right-hand part of (19) is less than the constant from the above inequality (18) for even n . Hence, (19) is true for all $n \in \mathbb{N}$. This completes the proof. \square

Corollary 4. $\theta_n(B_n) \asymp \sqrt{n}$.

Proof. In [5] it was proved that $\theta_n(B_n) \leq \sqrt{n+1}$. Consequently, the lower estimate $\theta_n(B_n) > c\sqrt{n}$ is precise with respect to dimension n . \square

Corollary 5. Assume $P : C(B_n) \rightarrow \Pi_1(\mathbb{R}^n)$ is the interpolation projector whose nodes coincide with vertices of a regular simplex being inscribed into the boundary sphere $\|x\| = 1$. When $\|P\|_{B_n} \asymp \theta_n(B_n)$.

Proof. As it was shown in [5], $\sqrt{n} \leq \|P\|_{B_n} \leq \sqrt{n+1}$. It remains to utilize the previous corollary. \square

Our results mean that any interpolation projector P corresponding to an inscribed regular simplex has the norm equivalent to the minimal possible. The equality $\|P\|_{B_n} = \theta_n(B_n)$ remains proved only for $1 \leq n \leq 4$.

The author is grateful to A. Yu. Ukhlov for some useful computer calculations.

References

1. Nevskii, M. V., *Geometricheskie ocenki v polinomialnoi interpolyacii* (Geometric Estimates in Polynomial Interpolation), Yaroslavl': Yarosl. Gos. Univ., 2012 (in Russian).
2. Nevskii, M. V., and Ukhalov, A. Yu., On numerical characteristics of a simplex and their estimates, *Model. Anal. Inform. Sist.*, 2016, vol. 23, no. 5, pp. 603–619 (in Russian). English transl.: *Aut. Control Comp. Sci.*, 2017, vol. 51, no. 7, pp. 757–769.
3. Nevskii, M. V., and Ukhalov, A. Yu., New estimates of numerical values related to a simplex, *Model. Anal. Inform. Sist.*, 2017, vol. 24, no. 1, pp. 94–110 (in Russian). English transl.: *Aut. Control Comp. Sci.*, 2017, vol. 51, no. 7, pp. 770–782.
4. Nevskii, M. V., and Ukhalov, A. Yu., On optimal interpolation by linear functions on an n -dimensional cube, *Model. Anal. Inform. Sist.*, 2018, vol. 25, no. 3, pp. 291–311 (in Russian). English transl.: *Aut. Control Comp. Sci.*, 2018, vol. 52, no. 7, pp. 828–842.
5. Nevskii, M. V., and Ukhalov, A. Yu., Linear interpolation on a Euclidean ball in \mathbb{R}^n (to appear).
6. Szego, G., *Orthogonal Polynomials*, New York: American Mathematical Society, 1959.
7. Suetin, P. K. *Klassicheskie ortogonal'nye mnogochleny*, Moscow: Nauka, 1979 (in Russian).