

# When a spherical body of constant diameter is of constant width?

Marek Lassak

**Abstract.** Let  $D$  be a convex body of diameter  $\delta$ , where  $0 < \delta < \frac{\pi}{2}$ , on the  $d$ -dimensional sphere. We prove that  $D$  is of constant diameter  $\delta$  if and only if it is of constant width  $\delta$  in the following two cases. The first case is when  $D$  is smooth. The second case is when  $d = 2$ .

**Keywords:** spherical geometry, convex body, lune, width, constant width, constant diameter

**MSC:** 52A55, 82D25

## 1 Introduction

The subject of this paper is spherical geometry (for a larger contexts see the monographs [2], [8] and [9]).

In the next section we recall the notion of a spherical convex body of constant width. Shortly speaking, for a convex body  $C$  on the  $d$ -dimensional sphere  $S^d$  and any hemisphere  $K$  supporting  $C$  we define the *width of  $C$  determined by  $K$*  as the thickness of any narrowest lune  $K \cap K^*$  containing  $C$ . By a *body of constant width* we mean a spherical convex body whose all widths are equal.

Let  $C \subset S^d$  be a convex body of diameter  $\delta$ . If the spherical distance  $|pq|$  of points  $p, q \in C$  is  $\delta$ , we call  $pq$  a *diametral chord* of  $C$  and we say that  $p, q$  are *diametrically opposed points* of  $C$ . Clearly,  $p, q \in \text{bd}(C)$ . After Part 4 of [7] we say that a convex body  $D \subset S^d$  of diameter  $\delta$  is *of constant diameter  $\delta$*  provided for every point  $p \in \text{bd}(D)$  there exists at least one point  $p' \in \text{bd}(D)$  such that  $|pp'| = \delta$  (in other words, that  $pp'$  is a diametral chord of  $D$ ). For the known analogous notion in  $E^d$  see [1].

Recall that in [7] it is proved that a convex body on  $S^d$  is of diameter  $\delta \geq \frac{\pi}{2}$  is of constant diameter if and only if it is a body of constant width  $\delta$ . Moreover, there is observed that the “if” part holds also for  $\delta < \frac{\pi}{2}$ , and the problem is put *if every spherical body of constant diameter  $\delta < \frac{\pi}{2}$  on  $S^d$  is a body of constant width  $\delta$ ?*

Our aim is to prove that every smooth spherical convex body of constant diameter  $\delta$  is of

constant width  $\delta$ , and that a body on the two-dimensional sphere is of constant diameter  $\delta$  if and only if it is of constant width  $\delta$ . As a consequence of these facts, the above problem remains now open only for non-smooth bodies of constant diameter below  $\frac{\pi}{2}$ .

By the way, in [3], [4] and [6] spherical bodies of constant width and constant diameter  $\frac{\pi}{2}$  are applied for recognizing if a Wulff shape is self-dual.

## 2 On spherical geometry

By  $S^d$  denote the unit sphere in the  $(d + 1)$ -dimensional Euclidean space  $E^{d+1}$ , where  $d \geq 2$ . The intersection of  $S^d$  with any  $(k + 1)$ -dimensional Euclidean space, where  $0 \leq k \leq d - 1$ , is called a *k-dimensional subsphere of  $S^d$* . For  $k = 1$  we call it a *great circle*, and for  $k = 0$  a *pair of antipodes*. If different points  $a, b \in S^d$  are not antipodes, by the *arc  $ab$*  connecting them we mean this part of the great circle containing  $a$  and  $b$ , which does not contain any pair of antipodes. By the *spherical distance  $|ab|$* , or shortly *distance*, of these points we understand the length of the arc connecting them.

By a *d-dimensional spherical ball of radius  $\rho \in (0, \frac{\pi}{2}]$* , or shorter a *ball*, we mean the set of points of  $S^d$  which are at the distance at most  $\rho$  from a fixed point, called the *center* of this ball. For  $d = 2$  it is called a *disk*, and its boundary is called a *circle of radius  $\delta$* . Spherical balls of radius  $\frac{\pi}{2}$  are called *hemispheres*. In other words, by a *hemisphere of  $S^d$*  we mean the common part of  $S^d$  with any closed half-space of  $E^{d+1}$ . We denote by  $H(c)$  the hemisphere whose center is  $c$ . Two hemispheres whose centers are antipodes are called *opposite hemispheres*.

By a *spherical  $(d - 1)$ -dimensional ball of radius  $\rho \in (0, \frac{\pi}{2}]$*  we mean the set of points of a  $(d - 1)$ -dimensional great sphere of  $S^d$  at the distance at most  $\rho$  from a point, called the *center* of this ball. The  $(d - 1)$ -dimensional balls of radius  $\frac{\pi}{2}$  are called  *$(d - 1)$ -dimensional hemispheres*.

Let a set  $C \subset S^d$  does not contain any pair of antipodes. We say that  $C$  is *convex* if together with every two its points,  $C$  contains the whole arc connecting them. If the interior of a closed convex set  $C$  is non-empty, we call  $C$  a *convex body*. Its boundary is denoted by  $\text{bd}(C)$ .

If a hemisphere  $H$  contains a convex body  $C$  and if  $p \in \text{bd}(H) \cap C$ , we say that  $H$  *supports  $C$  at  $p$*  or that  $H$  is a *supporting hemisphere of  $C$  at  $p$* . If exactly one hemisphere supports a convex body  $C$  at its boundary point  $p$ , we say that  $p$  is a *smooth point of  $\text{bd}(C)$* , and in the opposite case we say that  $p$  is an *acute point of  $\text{bd}(C)$* . If every boundary point of  $C \subset S^d$  is smooth, then  $C$  is called *smooth*. We call  $C$  *strictly convex* if  $\text{bd}(C)$  does not contain any arc.

If hemispheres  $G$  and  $H$  of  $S^d$  are different and not opposite, then  $L = G \cap H$  is called a *lune of  $S^d$* . This notion is considered in many books and papers. The parts of  $\text{bd}(G)$  and  $\text{bd}(H)$  contained in  $G \cap H$  are denoted by  $G/H$  and  $H/G$ , respectively. By the *thickness  $\Delta(L)$  of the lune  $L = G \cap H \subset S^d$*  we mean the spherical distance of the centers of the  $(d - 1)$ -dimensional

hemispheres  $G/H$  and  $H/G$ .

For any convex body  $C \subset S^d$  and any hemisphere  $K$  supporting  $C$  we define the *width*  $\text{width}_K(C)$  of  $C$  determined by  $K$  as the thickness of any narrowest lune  $K \cap K^*$  containing  $C$  (so that no lune of the form  $K \cap K'$  with a smaller thickness contains  $C$ ). By the *thickness*  $\Delta(C)$  of  $C$  we mean the minimum of  $\text{width}_K(C)$  over all hemispheres  $K$  supporting  $C$ . Clearly,  $\Delta(C)$  is nothing else but the thickness of a “narrowest” lune containing  $C$ . We say that  $C$  is of *constant width*  $w$  if all its widths are  $w$ .

The above notions are given and a few properties of lunes and convex bodies in  $S^d$  are presented in [5] and [7].

**Lemma.** *Let  $K$  be a hemisphere of  $S^d$  and let  $p \in \text{bd}(K)$ . Moreover, let  $pq \subset K$  be an arc orthogonal to  $\text{bd}(K)$  with  $q$  in the interior of  $K$  and  $|pq| < \frac{\pi}{2}$ . Then from amongst all the lunes of the form  $K \cap M$ , with  $q$  in the boundary of the hemisphere  $M$ , the lune  $K \cap K_{\perp}$  such that  $pq$  is orthogonal to  $\text{bd}(K_{\perp})$  at  $q$  has the smallest thickness.*

An easy proof is left to the reader.

### 3 Spherical bodies of constant diameter

The notion of a spherical body of constant diameter is recalled in the Introduction. In this section we present a few propositions on bodies of constant diameter.

**Proposition 1.** *Every convex body  $D \subset S^d$  of constant diameter  $\delta < \frac{\pi}{2}$  is strictly convex.*

*Proof.* Assume the opposite that  $D$  is not strictly convex. Then  $\text{bd}(D)$  contains an arc  $xz$ . Denote by  $y$  its midpoint. Clearly,  $y \in \text{bd}(D)$ . Since  $D$  is of constant diameter  $\delta$ , there is a point  $y' \in \text{bd}(D)$  such that  $|yy'| = \delta$ . This,  $y \in xz$  and  $\delta < \frac{\pi}{2}$  imply that  $|y'x| > \delta$  or  $|y'z| > \delta$ . Thus  $\text{diam}(D) > \delta$  in contradiction to the fact that  $D$  is of constant diameter  $\delta$ . Consequently,  $D$  is strictly convex.  $\square$

**Proposition 2.** *Let  $D \subset S^2$  be a body of constant diameter. Then every two diametral chords of  $D$  intersect.*

*Proof.* Denote the diameter of  $D$  by  $\delta$ . Suppose that some diametral chords  $ab$  and  $cd$  of  $D$  do not intersect (let for instance  $a, b, d, c$  be in this order on  $\text{bd}(D)$ ). Then  $abdc$  is a convex spherical non-degenerate quadrangle. Hence  $ad$  and  $bc$  intersect at exactly one point. Denote it by  $x$ . Since  $|ab| = \delta$  and  $|cd| = \delta$ , by the triangle inequality we get  $|ax| + |xb| \geq \delta$  and  $|cx| + |xd| \geq \delta$ . What is more, since  $x \notin ab$  and  $x \notin cd$ , we get  $|ax| + |xb| > \delta$  and  $|cx| + |xd| > \delta$ . This leads to  $|ax| + |xb| + |cx| + |xd| > 2\delta$ . So  $|ax| + |xd| > \delta$  or  $|cx| + |xb| > \delta$ . In other words,

$|ad| > \delta$  or  $|bc| > \delta$  in contradiction to  $\text{diam}(D) = \delta$ . Consequently, every two diametral chords of  $D$  intersect.  $\square$

**Proposition 3.** *If a hemisphere  $K$  supports a convex body  $D \subset S^d$  of constant diameter  $\delta < \frac{\pi}{2}$  at a smooth point of its boundary, then  $\text{width}_K(D) = \delta$ .*

*Proof.* Let  $p$  be a smooth point of  $\text{bd}(D)$ . Since  $D$  is of constant diameter, there exists a diametrically opposed point  $p' \in \text{bd}(D)$ . Hence the ball  $B$  of radius  $\delta$  and center  $p'$  contains  $D$ . Clearly,  $p \in \text{bd}(B)$ .

Since  $p$  is a smooth point of  $\text{bd}(D)$ , we conclude that  $K$  supports  $B$  at  $p$ .

From  $\text{diam}(D) = \delta$  and  $p' \in \text{bd}(D)$  we see that the spherical ball  $B'$  of radius  $\delta$  centered at  $p$  contains  $D$  and supports it at  $p'$ .

Denote by  $K'$  the hemisphere supporting  $B'$  at  $p'$ . Clearly, we have  $D \subset B \cap B' \subset K \cap K'$ .




FIGURE 1. Illustration to the proof of Proposition 3

Since our chord  $pp'$  of length  $\delta$  is orthogonal to  $\text{bd}(K)$  at  $p$  and to  $\text{bd}(K')$  at  $p'$ , by the definition of the thickness of a lune we see that the lune  $K \cap K'$  has thickness  $|pp'| = \delta$ . Consequently, by  $D \subset K \cap K'$  we conclude that  $\text{width}_K(D) \leq \delta$ .

By Lemma the lune  $K \cap K'$  is the narrowest lune from the family  $\mathcal{L}_1$  of lunes of the form  $K \cap M$  with  $p' \in \text{bd}(M)$  containing  $pp'$ . From  $pp' \subset D \subset K \cap K'$  we conclude that  $K \cap K'$  is also the narrowest lune from the family  $\mathcal{L}_2$  of all lunes the form  $K \cap M$  containing  $D$ . Since every lune from  $\mathcal{L}_2$  contains a lune from  $\mathcal{L}_1$ , every lune from  $\mathcal{L}_2$  is of thickness at least  $\delta$ . Consequently,  $\text{width}_K(D) \geq \delta$ .

From the above two paragraphs we conclude that  $\text{width}_K(D) = \delta$ .  $\square$

## 4 Two cases in which a spherical convex body of constant diameter is of constant width

Since the question is answered for  $\delta \geq \frac{\pi}{2}$  and since, as mentioned in the Introduction, every spherical body of constant width is of constant diameter, now we concentrate on checking when a spherical body of constant diameter  $\delta < \frac{\pi}{2}$  is of constant width. The following theorem gives a partial answer. It results immediately from Proposition 3 and from the fact that every body of constant width  $\delta$  is of constant diameter  $\delta$ .

**Theorem 1.** *Let  $0 < \delta < \frac{\pi}{2}$ . A spherical smooth convex body on  $S^d$  is of constant diameter  $\delta$  if and only if it is of constant width  $\delta$ .*

Below is our main theorem. Since in its proof we apply polar sets, let us recall this notion. For a convex body  $C \subset S^d$  by its *polar* we mean the set  $C^\circ = \{r : C \subset H(r)\}$ . It is easy to show that  $C^\circ$  is a convex body. Recall that  $\text{bd}(C^\circ)$  is the set of points  $r$  such that  $H(r)$  is a supporting hemisphere of  $C$ .

**Theorem 2.** *Let  $0 < \delta < \frac{\pi}{2}$ . A convex body on the two-dimensional sphere is of constant diameter  $\delta$  if and only if it is of constant width  $\delta$ .*

*Proof.* In [7] there is observed that every body of constant width on  $S^d$  is a body of constant diameter.

It remains to show that every body  $D \subset S^2$  of constant diameter  $\delta < \frac{\pi}{2}$  is of constant width  $\delta$ , i.e., that  $\text{width}_K(D) = \delta$  for every supporting hemisphere  $K$  of  $D$ .

By Proposition 1 there is exactly one point  $p$  of support of  $D$  by  $K$ .

When  $p$  is a smooth point of  $\text{bd}(D)$ , then we apply Proposition 3.

Consider the case when  $p$  is an acute point of  $\text{bd}(D)$ . Denote by  $H_1 = H(r_1)$  the first supporting hemisphere and by  $H_2 = H(r_2)$  the last supporting hemisphere of  $D$  at  $p$ , as we go counterclockwise with the center  $r$  of  $H(r)$  on  $\text{bd}(D^\circ)$ , see Fig 2.

For  $i = 1, 2$  provide the arc  $pp_i \subset H_i$  of length  $\delta$  orthogonal to  $\text{bd}(H_i)$  at  $p$ . So  $p'_i \in pr_i$  for  $i = 1, 2$ . By  $B_i$  denote the ball of radius  $\delta$  centered at  $p_i$ , where  $i = 1, 2$ .

Of course,  $D \subset B_1 \cap B_2 \subset H_1 \cap H_2$ .




FIGURE 2. Illustration to the proof of Theorem 2

Consider the piece  $P$  of the circle with center  $p$  and radius  $\delta$  from  $p_1$  to  $p_2$ . Denote by  $s_i$  the point of  $\text{bd}(D)$  in the arc  $pp_i$ , where  $i = 1, 2$ , and by  $S$  the piece of  $\text{bd}(D)$ , when going from  $s_1$  counterclockwise to  $s_2$ . Take any  $s \in S$ . There exists a diametrically opposed point  $s'$  of  $D$  in  $\text{bd}(D)$ . By Proposition 2 the chord  $ss'$  intersects the chords  $pp_1$  and  $pp_2$ . Hence  $s' = p$ . This and  $|ss'| = \delta$  imply that  $s$  belongs to the circle  $F$  with center  $p$  and radius  $\delta$ .

What is more,  $s \in P$ . The reason is that if  $s \in F \setminus P$ , then the disk of center  $s$  and radius  $\delta$  contains  $D$ . Consequently, the hemisphere supporting the ball bounded by  $F$  at  $p$  is a supporting hemisphere of  $D$  at  $p$ . This contradicts the fact that  $H_1$  and  $H_2$  are the first and the last supporting hemispheres of  $D$  at  $p$ .

Consequently, for every hemisphere  $H$  supporting  $D$  at  $p$  the chord of  $D$  orthogonal to  $H$  at  $p$  is a diametral chord of  $D$ . Thus by Lemma we get  $\text{width}_H(D) = \delta$ .

In particular,  $\text{width}_K(D) = \delta$ , as required.  $\square$

By the proof of Theorem 2 and also Proposition 3 any supporting hemisphere  $H = H(r)$  of  $D$  determines a unique diametral chord  $pp'$  of  $D$ ; it is orthogonal to the great circle bounding  $H$ . Moreover, the center  $r$  of  $H$  belongs to  $\text{bd}(D^\circ)$ . On the other hand, take any  $r \in \text{bd}(D^\circ)$ . Then  $H(r)$  supports  $D$  at exactly one point  $p$ . Clearly,  $D \subset H(r)$ . Consequently, for any body  $D \subset S^2$  we have the one-to-one correspondence between the following objects:

- the supporting hemispheres of  $D$ ,
- the points of  $\text{bd}(D^\circ)$ ,
- the diametral chords of  $D$ .

Every position of diametral chord is determined by a supporting hemisphere of  $D$ . It is also determined by a point of  $\text{bd}(D^\circ)$ ; still such a point  $r$  determines exactly one point  $p$ , and the considered diametral chord is in the arc  $pp'$ .

A forthcoming paper is devoted to show the analogous facts as in Propositions 1–3 and Theorems 1–2 in  $E^d$  (they are true also in the hyperbolic space). The proofs of these theorems apply the parallelism, which cannot be used here for spherical bodies.

## References

- [1] G. D. Chakerian, H. Groemer, Convex bodies of constant width, In *Convexity and its Applications*, pp. 49–96, Birkhauser, Basel (1983).
- [2] J. W. Harris, H. Stocker, Spherical Geometry, 4.9 in *Handbook of Mathematics and Computational Science*. New York: Springer-Verlag, pp. 108–113, 1998.
- [3] H. Han, T. Nishimura. Self-dual Wulff shapes and spherical convex bodies of constant width  $\pi/2$ , *J. Math. Soc. Japan* **69** (2017), 1475–1484.
- [4] H. Han, T. Nishimura Spherical method for studying Wulff shapes and related topics. Singularities in Generic geometry, pp. 1–53 in *Adv. Stud. Pure Math.* **78** (2018) Math. Soc. Japan, Tokyo.
- [5] M. Lassak, Width of spherical convex bodies, *Aequationes Math.* **89** (2015), 555–567.
- [6] M. Lassak, Application of spherical geometry for Wulff shape (to appear), temporarily see at the repository arXiv:1903.04148v2.
- [7] M. Lassak, M. Musielak, Spherical bodies of constant width, *Aequationes Math.* **92** (2018), 627–640
- [8] A. Papadopoulos, On the works of Euler and his followers on spherical geometry, *Ganita Bharati* **36** (2014), no.1, 53–108.
- [9] G. Van Brummelen, Heavenly mathematics. The forgotten art of spherical trigonometry. Princeton University Press (Princeton, 2013).

Marek Lassak  
 University of Science and Technology  
 al. Kaliskiego 7, Bydgoszcz 85-796  
 e-mail: lassak@utp.edu.pl