

ON THE CONVEX CONES ARISING FROM CLASSIFICATIONS OF PARTIAL ENTANGLEMENT IN THE THREE QUBIT SYSTEM

KYUNG HOON HAN AND SEUNG-HYEOK KYE

ABSTRACT. In order to classify partial entanglement of multi-partite states, it is natural to consider the convex hulls, intersections and differences of basic convex cones obtained from partially separable states with respect to partitions of systems. In this paper, we consider convex cones consisting of X -shaped three qubit states arising in this way. The class of X -shaped states includes important classes like Greenberger-Horne-Zeilinger diagonal states. We find all the extreme rays of those convex cones to exhibit corresponding partially separable states. We also give characterizations for those cones which give rise to necessary criteria in terms of diagonal and anti-diagonal entries for general three qubit states.

1. INTRODUCTION

The notion of entanglement is now considered as an indispensable resource in the current quantum information theory. In the multi-partite systems, there are various notions of separability according to partitions of systems, which give rise to different kinds of partial entanglement. In the tri-partite system, we may consider three kinds of bi-partitions $A-BC$, $B-CA$ and $C-AB$ of systems. In this way, a tri-partite state may be considered as a bi-partite state with respect to one of the above bi-partitions. It was shown in [1] that a three qubit state may be entangled even though it is separable as a bi-partite state with respect to any bi-partitions. Therefore, it is natural to classify partial separability in multi-partite systems, as they were suggested in the literature [2, 3, 4, 5, 6, 7, 8, 9, 10]. We recall that a multi-partite state is (fully) separable if it is a convex sum of pure product states, and entangled if it is not separable.

We will work in the real vector space of all three qubit self-adjoint matrices, and consider the convex cones \mathcal{A} , \mathcal{B} and \mathcal{C} consisting of all unnormalized separable states with respect to the bi-partitions $A-BC$, $B-CA$ and $C-AB$, respectively. Recall that a subset C of a real vector space is called a *convex cone* when $C + C \subset C$ and $aC \subset C$ for $a \geq 0$. We note that the sum $C_1 + C_2$ of two convex cones C_1 and C_2 is again a convex cone which coincides with the convex hull of C_1 and C_2 , that is, the smallest convex set containing C_1 and C_2 .

The above mentioned result [1] tells us that the convex cone $\mathcal{A} \cap \mathcal{B} \cap \mathcal{C}$ is strictly bigger than the convex cone of all fully separable states as tri-partite states. The differences $\mathcal{A} \setminus (\mathcal{B} \cup \mathcal{C})$ and $(\mathcal{A} \cap \mathcal{B}) \setminus \mathcal{C}$ have been considered in [4, 5], together with similar sets obtained by permuting \mathcal{A} , \mathcal{B} and \mathcal{C} . On the other hand, the convex hull

1991 *Mathematics Subject Classification.* 81P40, 52A20, 15A69.

Both KHH and SHK were partially supported by NRF-2017R1A2B4006655, Korea.

$\mathcal{A} + \mathcal{B} + \mathcal{C}$ and the difference $(\mathcal{A} + \mathcal{B} + \mathcal{C}) \setminus (\mathcal{A} \cup \mathcal{B} \cup \mathcal{C})$ have been also considered in [2] and [7], respectively. More recently, all the possible classes

$$(1) \quad [C_1 \cap \dots \cap C_k] \setminus [C_{k+1} \cup \dots \cup C_\ell]$$


have been considered in [10], where C_i is one of the following convex cones

$$(2) \quad \mathcal{A}, \mathcal{B}, \mathcal{C}, \mathcal{A} + \mathcal{B}, \mathcal{B} + \mathcal{C}, \mathcal{C} + \mathcal{A}, \mathcal{A} + \mathcal{B} + \mathcal{C}.$$

Nontrivial classes of three qubit states obtained by (1) are known to be nonempty only recently [11].

The main purposes of this note are twofold: Exhibiting three qubit states in the above classes in (2) and giving criteria for states to be members of the cones. We will do these for so called *X-shaped* three qubit states, whose entries are zero, by definition, except for diagonal and anti-diagonal entries. Many important states like GHZ diagonal states are of this form. An X-shaped state is also called as an *X-state* for brevity. In order to exhibit all the three qubit X-states in a given cone, we find all the extreme rays of the convex cone. Recall that an element x of a convex cone C generates an extreme ray whenever $x = x_1 + x_2$ with $x_i \in C$ implies that x_i is a nonnegative multiple of x for $i = 1, 2$. By the abuse of the terminology, we say that x itself is an extreme ray in this case. Then all the elements of convex cones in (2) are the nonnegative sums of extreme rays. Those extreme rays also play essential roles to find criteria for the dual cone. Those criteria will be expressed in terms of algebraic inequalities with the entries of X-states, which give rise to necessary criteria for general three qubit states to be a member of a given cone, in terms of diagonal and anti-diagonal entries. As for the corresponding results for full separability, we refer to the papers [12, 13, 14, 15].

Our main tool is the duality between closed convex cones in real vector spaces, and so, we will also consider the dual cones, whose members play roles of witnesses, of the cones in (2). Since the intersection and the convex hull are dual operations, we also naturally consider the intersections as well as convex hulls through the discussion. Therefore, we will consider the convex cones appearing in the following diagram



which shows us partial order relations by inclusion among convex cones we are considering. The dual cones will be also discussed.

After we explain briefly the duality in the next section, we will consider the convex cones \mathcal{A} , \mathcal{B} and \mathcal{C} in Section 3 together with their dual cones. We will also consider the convex cones $\mathcal{A} + \mathcal{B} + \mathcal{C}$ and $\mathcal{A} \cap \mathcal{B} \cap \mathcal{C}$ in Section 4. Conditions for those convex cones with X-shaped matrices are already scattered in the literature [16, 17, 18, 19, 20]. Here, we give an alternative proof in the context of duality, together with exhibition of all extreme rays in the convex cone of X-shaped matrices in each of them. In Section 5, we deal with convex hulls and intersections of two convex cones like $\mathcal{A} + \mathcal{B}$, $\mathcal{A} \cap \mathcal{B}$ and their duals. We will summarize our results in the final section.

The authors are grateful to the referee for careful reading and valuable suggestions to improve presentation.

2. DUALITY

Let C be a subset of a finite dimensional real vector space V with a non-degenerating bilinear pairing $\langle \cdot, \cdot \rangle$, that is, $\langle x, y \rangle = 0$ for every $y \in V$ implies $x = 0$. We define the *dual cone* C° by

$$C^\circ = \{x \in V : \langle x, y \rangle \geq 0 \text{ for every } y \in C\}.$$

Then C° is a closed convex cone of V in general, and $C^{\circ\circ}$ is the smallest closed convex cone containing C by the Hahn-Banach type separation theorem. If C itself is a closed convex cone then we have $C = C^{\circ\circ}$, and so we see that the following are equivalent:

- $x \notin C$;
- there exists $y \in C^\circ$ such that $\langle x, y \rangle < 0$.

For example, if \mathcal{S} is the closed convex cone consisting of unnormalized fully separable states in the real vector space V of self-adjoint matrices in $M_2 \otimes M_2 \otimes M_2$, then we see by this principle that ϱ is non-separable, that is, entangled if and only if there exists $W \in \mathcal{S}^\circ$ such that $\langle W, \varrho \rangle < 0$. Such a W must be non-positive, and called an entanglement witness [21]. Here, the bilinear pairing is given by $\langle a, b \rangle = \text{Tr}(ba^t)$ for matrices a and b , as usual. On the other hand, the closed convex cone \mathcal{P} of all positive matrices is self-dual, that is, $\mathcal{P}^\circ = \mathcal{P}$, by the Hadamard theorem.

We note that the two operations, convex hull and intersection, are dual to each other. In other words, the following identities


$$(C_1 + C_2)^\circ = C_1^\circ \cap C_2^\circ, \quad (C_1 \cap C_2)^\circ = C_1^\circ + C_2^\circ$$

hold for closed convex cones C_1 and C_2 . The first identity follows from the definition. See [22]. The second one follows from the first one and the fact that the convex hull of two closed convex cones is closed. This is an easy consequence of Carathéodory theorem which tells us that the convex hull of a compact set is compact. We note that a convex cone C spans the whole space V if and only if $C + (-C) = V$. If we apply

the above duality to the four closed convex cones C , $-C$, $\{0\}$ and V , then we see that the following two properties

- (C₁) C spans the whole space;
- (C₂) $C \cap (-C) = \{0\}$

are dual to each other. In other words, a closed convex cone C satisfies (C₁) if and only if C° satisfies (C₂). Recall that the real vector space $(M_{n_1} \otimes M_{n_2})_{\text{sa}}$ of all self-adjoint matrices in the tensor product $M_{n_1} \otimes M_{n_2}$ coincides with the tensor product $(M_{n_1})_{\text{sa}} \otimes (M_{n_2})_{\text{sa}}$ of the self-adjoint parts. See [15, Section 7]. This is also true for multi-tensor products by induction. Therefore, the convex cone $\mathcal{S} = M_2^+ \otimes M_2^+ \otimes M_2^+$ spans the whole space $V = (M_2 \otimes M_2 \otimes M_2)_{\text{sa}}$. Since $\mathcal{P} = (M_2 \otimes M_2 \otimes M_2)^+$ satisfies $\mathcal{P} \cap (-\mathcal{P}) = \{0\}$, we see that all the convex cones \mathcal{X} in the diagram (3) also satisfy both conditions, by the relation $\mathcal{S} \subset \mathcal{X} \subset \mathcal{P}$. We list up the dual cones of the cones in (3) as follows:



We note that all the convex cones in the diagram (4) also satisfy both conditions (C₁) and (C₂), as dual cones of the convex cones satisfying the conditions. An important consequence of (C₂) is that every element of the convex cone is a nonnegative sum of extreme rays. See [23, Theorem 18.5].

The duality is also very useful to find all the candidates for extreme rays. We say that a subset S of a closed convex cone C is a *generating set* for C if every element of C is the limit of nonnegative sums of finitely many elements in S . This happens if and only if $S^{\circ\circ} = C$ if and only if $S^\circ = C^\circ$. In other words, we have to show that the following two statements

- $y \in C^\circ$, that is, $\langle x, y \rangle \geq 0$ for each $x \in C$;
- $\langle x, y \rangle \geq 0$ for each $x \in S$

are equivalent to each other, in order to show that S is a generating set for C . This equivalence, in turn, gives rise to a criterion for the convex cone C° in terms of algebraic

inequalities arising from members in the generating set S . This principle will be the main tool of our discussion throughout this paper.

We note that generating sets of a convex cone are not determined uniquely. For example, the convex cone C itself is also a generating set for C . Furthermore, a generating set need not contain all the extreme rays. If a generating set S for C is closed, then its convex hull is also closed by Carathéodory theorem, and so every element of C is the sum of finitely many elements in S . Therefore, we conclude that a *closed* generating set for C contains all the extreme rays of C . In this way, we are looking for the set $\text{Ext}(C)$ of all extreme rays of the convex cone C . We summarize as follows:

Proposition 2.1. *For a subset S of a closed convex cone C in a finite dimensional real vector space V , the following are equivalent:*

- (i) S is a generating set for C ;
- (ii) if $y \in V$ and $\langle x, y \rangle \geq 0$ for each $x \in S$, then $y \in C^\circ$.

If S is a closed generating set for C , then we have $\text{Ext}(C) \subset S$.

In this paper, we will concentrate on the three qubit system, and so we will work in the real vector space V of all 8×8 self-adjoint matrices. The space V has an important subspace, denoted by \mathbf{X} , consisting of all \mathbf{X} -shaped matrices whose entries are zero except for diagonal and anti-diagonal entries. In the three qubit case, an \mathbf{X} -shaped self-adjoint matrix is of the form

$$\mathbf{X}(a, b, z) = \begin{pmatrix} a_1 & & & & & & & z_1 \\ & a_2 & & & & & & z_2 \\ & & a_3 & & & & & z_3 \\ & & & a_4 & z_4 & & & \\ & & & \bar{z}_4 & b_4 & & & \\ & & & \bar{z}_3 & & b_3 & & \\ & & & \bar{z}_2 & & & b_2 & \\ \bar{z}_1 & & & & & & & b_1 \end{pmatrix},$$

for $a, b \in \mathbb{R}^4$ and $z \in \mathbb{C}^4$, where $\mathbb{C}^2 \otimes \mathbb{C}^2 \otimes \mathbb{C}^2$ is identified with the space \mathbb{C}^8 using the lexicographic order of indices. Many important multi-qubit states arise in this form. For example, GHZ diagonal states [24] are in this form, and an \mathbf{X} -state $\mathbf{X}(a, b, z)$ is a GHZ diagonal if and only if $a = b$ and $z \in \mathbb{R}^4$.

Note that V and \mathbf{X} are of 64 and 16-dimensional spaces, respectively. For a given matrix $\varrho \in V$, we denote by $\varrho_{\mathbf{X}}$ the \mathbf{X} -part of ϱ . The map $\varrho \mapsto \varrho_{\mathbf{X}}$ from V onto \mathbf{X} has the following important property.

Proposition 2.2. *For every convex cone \mathcal{X} in the diagram (3), we have the following:*

- (i) if $\varrho \in \mathcal{X}$, then $\varrho_{\mathbf{X}} \in \mathcal{X}$;
- (ii) if $W \in \mathcal{X}^\circ$, then $W_{\mathbf{X}} \in \mathcal{X}^\circ$.

Proof. It suffices to prove for the convex cone \mathcal{A} . For the statement (i), it also suffices to show for a vector state ϱ associated with a product vector $|x\rangle \otimes |y\rangle \in \mathbb{C}^2 \otimes \mathbb{C}^4$, where $|x\rangle = (x_1, x_2)^t$ and $|y\rangle = (y_1, y_2, y_3, y_4)^t$. We consider the following product vectors

$$\begin{aligned} & (+x_1, +x_2)^t \otimes (+y_1, +y_2, +y_3, +y_4)^t, \\ & (+x_1, +x_2)^t \otimes (+y_1, -y_2, -y_3, +y_4)^t, \\ & (+x_1, -x_2)^t \otimes (+y_1, -y_2, +y_3, -y_4)^t, \\ & (+x_1, -x_2)^t \otimes (+y_1, +y_2, -y_3, -y_4)^t. \end{aligned}$$

We take the average of four vector states associated with these four product vectors, to recover the \mathbf{X} -part of ϱ . This proves (i). For the statement (ii) with $\mathcal{X} = \mathcal{A}$, take $W \in \mathcal{A}^\circ$. For every $\varrho \in \mathcal{A}$, we see that $\langle W_{\mathbf{X}}, \varrho \rangle = \langle W, \varrho_{\mathbf{X}} \rangle$ is nonnegative, because $W \in \mathcal{A}^\circ$ and $\varrho_{\mathbf{X}} \in \mathcal{A}$ by (i). Therefore, we have $W_{\mathbf{X}} \in \mathcal{A}^\circ$. \square

Corresponding results for full separability are found in Section 3 of [12]. See also Proposition 4.1 of [15] for multi-qubit cases. If ϱ is an \mathbf{X} -state, then $\langle W, \varrho \rangle = \langle W_{\mathbf{X}}, \varrho \rangle$, and so we have the following:

Corollary 2.3. *For a convex cone \mathcal{X} in the diagram (3), we have the following:*

- (i) *for a three qubit \mathbf{X} -state ϱ , we have $\varrho \in \mathcal{X}$ if and only if $\langle W, \varrho \rangle \geq 0$ for every \mathbf{X} -shaped $W \in \mathcal{X}^\circ$;*
- (ii) *for a three qubit \mathbf{X} -shaped W , we have $W \in \mathcal{X}^\circ$ if and only if $\langle W, \varrho \rangle \geq 0$ for every \mathbf{X} -state $\varrho \in \mathcal{X}$.*

Corollary 2.4. *For convex cones \mathcal{X}_1 and \mathcal{X}_2 in diagrams (3) or (4), we have the relation $(\mathcal{X}_1 + \mathcal{X}_2) \cap \mathbf{X} = (\mathcal{X}_1 \cap \mathbf{X}) + (\mathcal{X}_2 \cap \mathbf{X})$.*

Once we characterize \mathbf{X} -shaped matrices in the convex cones in (3) or (4), these conditions will give rise to necessary conditions for general three qubit self-adjoint matrices to belong to those convex cones, by Proposition 2.2. On the other hand, Corollary 2.3 tells us that we may restrict ourselves on the bi-linear pairing in the real vector space \mathbf{X} for this purpose.

3. BASIC PARTIAL SEPARABILITY

In this section, we consider the three basic convex cones \mathcal{A} , \mathcal{B} , \mathcal{C} and their dual cones \mathcal{A}° , \mathcal{B}° , \mathcal{C}° . It was shown in [20, Proposition 5.2] that an \mathbf{X} -shaped multi-qubit state is separable with respect to a bi-partition of systems if and only if it is of positive partial transpose with respect to the same bi-partition. The PPT condition is easily checked for three qubit \mathbf{X} -shaped states by the following inequalities

$$S_1[i, j] : \quad \min\{\sqrt{a_i b_i}, \sqrt{a_j b_j}\} \geq \max\{|z_i|, |z_j|\},$$

for $i, j = 1, 2, 3, 4$. If $\varrho = \mathsf{X}(a, b, (z_1, z_2, z_3, z_4))$, then the partial transposes are given by

$$\begin{aligned}\varrho^{\Gamma A} &= \mathsf{X}(a, b, (\bar{z}_4, \bar{z}_3, \bar{z}_2, \bar{z}_1)), \\ \varrho^{\Gamma B} &= \mathsf{X}(a, b, (z_3, z_4, z_1, z_2)), \\ \varrho^{\Gamma C} &= \mathsf{X}(a, b, (z_2, z_1, z_4, z_3)).\end{aligned}$$

Therefore, we have the following:

Proposition 3.1. [20, Proposition 5.2] *For a three qubit X -state $\varrho = \mathsf{X}(a, b, z)$, we have the following:*

- (i) $\varrho \in \mathcal{A}$ if and only if both $S_1[1, 4]$ and $S_1[2, 3]$ hold;
- (ii) $\varrho \in \mathcal{B}$ if and only if both $S_1[1, 3]$ and $S_1[2, 4]$ hold;
- (iii) $\varrho \in \mathcal{C}$ if and only if both $S_1[1, 2]$ and $S_1[3, 4]$ hold.

We note that inequalities $S_1[i, j]$'s give us necessary criteria for general three qubit states to belong to \mathcal{A} , \mathcal{B} and \mathcal{C} , respectively, by Proposition 2.2. Now, we proceed to provide generating sets for the convex cones $\mathcal{A} \cap \mathsf{X}$, $\mathcal{B} \cap \mathsf{X}$ and $\mathcal{C} \cap \mathsf{X}$. To be motivated, we decompose an X -state $\varrho = \mathsf{X}(a, b, z)$ in \mathcal{A} by

$$\begin{aligned}\varrho &= \mathsf{X}((a_1, 0, 0, a_4), (b_1, 0, 0, b_4), (z_1, 0, 0, z_4)) \\ &\quad + \mathsf{X}((0, a_2, a_3, 0), (0, b_2, b_3, 0), (0, z_2, z_3, 0)),\end{aligned}$$

then two summands satisfy both $S_1[1, 4]$ and $S_1[2, 3]$. Therefore, we may assume that $a_k = b_k = z_k = 0$ for $k = 2, 3$. If $z_1 = 0$, then ϱ is the average of two states

$$\mathsf{X}((a_1, 0, 0, a_4), (b_1, 0, 0, b_4), (z_4, 0, 0, z_4)), \quad \mathsf{X}((a_1, 0, 0, a_4), (b_1, 0, 0, b_4), (-z_4, 0, 0, z_4))$$

in $\mathcal{A} \cap \mathsf{X}$. If $0 < |z_1| < |z_4|$, then ϱ is a convex combination of

$$\mathsf{X}((a_1, 0, 0, a_4), (b_1, 0, 0, b_4), (\frac{|z_4|}{|z_1|}z_1, 0, 0, z_4)), \quad \mathsf{X}((a_1, 0, 0, a_4), (b_1, 0, 0, b_4), (-\frac{|z_4|}{|z_1|}z_1, 0, 0, z_4))$$

in $\mathcal{A} \cap \mathsf{X}$.

By subtracting a suitable diagonal state, it is natural to consider the following conditions

$$S_1^e[i, j] : \quad \sqrt{a_i b_i} = |z_i| = \sqrt{a_j b_j} = |z_j| = 1, \quad \text{the others are zero,}$$

for each $i, j = 1, 2, 3, 4$ with $i \neq j$. We define

$$\mathcal{E}_A = \{\varrho = \mathsf{X}(a, b, z) : S_1^e[1, 4] \text{ or } S_1^e[2, 3] \text{ holds}\},$$

$$\mathcal{E}_B = \{\varrho = \mathsf{X}(a, b, z) : S_1^e[1, 3] \text{ or } S_1^e[2, 4] \text{ holds}\},$$

$$\mathcal{E}_C = \{\varrho = \mathsf{X}(a, b, z) : S_1^e[1, 2] \text{ or } S_1^e[3, 4] \text{ holds}\}.$$

We also denote by Δ the set of all extreme diagonal states, that is,

$$\Delta = \{\mathsf{X}(E_i, 0, 0) : i = 1, 2, 3, 4\} \cup \{\mathsf{X}(0, E_i, 0) : i = 1, 2, 3, 4\},$$

where $\{E_i : i = 1, 2, 3, 4\}$ denotes the canonical basis of \mathbb{R}^4 .

We have $\mathcal{E}_A \subset \mathcal{A}$ by Proposition 3.1, and \mathcal{E}_A is parameterized by four real variables. The same comments also hold for \mathcal{B} and \mathcal{C} . We also consider the following inequalities:

$$W_1[i, j] : \quad \sqrt{s_i t_i} + \sqrt{s_j t_j} \geq |u_i| + |u_j|,$$

for $i, j = 1, 2, 3, 4$ with $i \neq j$, in order to characterize the dual cones $\mathcal{A}^\circ \cap \mathcal{X}$, $\mathcal{B}^\circ \cap \mathcal{X}$ and $\mathcal{C}^\circ \cap \mathcal{X}$. We denote $\mathbf{X}_i(s_i, t_i, u_i) := \mathbf{X}(s_i E_i, t_i E_i, u_i E_i)$ for $i = 1, 2, 3, 4$.

Lemma 3.2. *For a given self-adjoint \mathbf{X} -shaped matrix $W = \mathbf{X}(s, t, u)$, the following are equivalent:*

- (i) $\langle W, \varrho \rangle \geq 0$ for each $\varrho \in \mathcal{E}_{\mathcal{A}} \cup \Delta$;
- (ii) $s_i, t_i \geq 0$ for $i = 1, 2, 3, 4$, and the inequalities $W_1[1, 4]$ and $W_1[2, 3]$ hold;
- (iii) $\langle W, \varrho \rangle \geq 0$ for each $\varrho \in \mathcal{A}$.

Proof. For the direction (i) \implies (ii), we obtain $s_i, t_i \geq 0$ from $\langle W, \varrho \rangle \geq 0$ for $\varrho \in \Delta$. Suppose that both s_i and t_i are nonzero for each $i = 1, 2, 3, 4$. Then, we can consider the following states

$$\varrho_{i,j} := \mathbf{X}_i \left(\sqrt{\frac{t_i}{s_i}}, \sqrt{\frac{s_i}{t_i}}, -e^{-i\theta_i} \right) + \mathbf{X}_j \left(\sqrt{\frac{t_j}{s_j}}, \sqrt{\frac{s_j}{t_j}}, -e^{-i\theta_j} \right)$$

for $(i, j) = (1, 4), (2, 3)$, with $\theta_k = \arg u_k$. Since $\varrho_{i,j} \in \mathcal{E}_{\mathcal{A}}$, we have

$$0 \leq \frac{1}{2} \langle W, \varrho_{i,j} \rangle = \sqrt{s_i t_i} + \sqrt{s_j t_j} - |u_i| - |u_j|,$$

by (i). When one of s_i or t_i is zero, we apply the result to $W + \varepsilon I$ to get the inequality $\sqrt{(s_i + \varepsilon)(t_i + \varepsilon)} + \sqrt{(s_j + \varepsilon)(t_j + \varepsilon)} \geq |u_i| + |u_j|$ for each $\varepsilon > 0$.

For the implication (ii) \implies (iii), it is enough to prove the following by Corollary 2.3 and Proposition 3.1:

$$(5) \quad S_1[1, 4], S_1[2, 3], W_1[1, 4], W_1[2, 3] \implies \langle \mathbf{X}(s, t, u), \mathbf{X}(a, b, z) \rangle \geq 0.$$

Indeed, we have

$$\begin{aligned} \sum_{i=1}^4 \sqrt{s_i t_i} \sqrt{a_i b_i} &= (\sqrt{s_1 t_1} \sqrt{a_1 b_1} + \sqrt{s_4 t_4} \sqrt{a_4 b_4}) + (\sqrt{s_2 t_2} \sqrt{a_2 b_2} + \sqrt{s_3 t_3} \sqrt{a_3 b_3}) \\ &\geq (\sqrt{s_1 t_1} + \sqrt{s_4 t_4}) \max\{|z_1|, |z_4|\} + (\sqrt{s_2 t_2} + \sqrt{s_3 t_3}) \max\{|z_2|, |z_3|\} \\ &\geq (|u_1| + |u_4|) \max\{|z_1|, |z_4|\} + (|u_2| + |u_3|) \max\{|z_2|, |z_3|\} \\ &\geq \sum_{i=1}^4 |u_i| |z_i|, \end{aligned}$$

which implies

$$(6) \quad \begin{aligned} \frac{1}{2} \langle \mathbf{X}(s, t, u), \mathbf{X}(a, b, z) \rangle &= \frac{1}{2} \sum_{i=1}^4 (s_i a_i + t_i b_i + 2 \operatorname{Re}(u_i z_i)) \\ &\geq \sum_{i=1}^4 (\sqrt{s_i t_i} \sqrt{a_i b_i} - |u_i| |z_i|) \geq 0. \end{aligned}$$

as it was required. The direction (iii) \implies (i) is clear since $\mathcal{E}_{\mathcal{A}} \cup \Delta \subset \mathcal{A}$ by Proposition 3.1. \square

The equivalence between (ii) and (iii) of Lemma 3.2 gives rise to a characterization of the convex cone $\mathcal{A}^\circ \cap \mathcal{X}$, whose members are the Choi matrix of $(1, 2, 2)$ -positive bilinear maps between 2×2 matrices in the sense of [19]. Therefore, Lemma 3.2 recovers Theorem 6.2 in [19], as follows:

Proposition 3.3. [19, Theorem 6.2] *For a self-adjoint $W = \mathsf{X}(s, t, u)$ with nonnegative diagonals, we have the following:*

- (i) $W \in \mathcal{A}^\circ$ if and only if both $W_1[1, 4]$ and $W_1[2, 3]$ hold;
- (ii) $W \in \mathcal{B}^\circ$ if and only if both $W_1[1, 3]$ and $W_1[2, 4]$ hold;
- (iii) $W \in \mathcal{C}^\circ$ if and only if both $W_1[1, 2]$ and $W_1[3, 4]$ hold.

The implication (i) \implies (iii) of Lemma 3.2 tells us that the set $\mathcal{E}_{\mathcal{A}} \cup \Delta$ is a generating set for the convex cone $\mathcal{A} \cap \mathcal{X}$ by Proposition 2.1. We also note that the set $\mathcal{E}_{\mathcal{A}} \cup \Delta$ is closed, and so we conclude that every extreme ray of $\mathcal{A} \cap \mathcal{X}$ must be an element of $\mathcal{E}_{\mathcal{A}} \cup \Delta$. We show that the converse actually holds. Because states in Δ generate extreme rays in the cone \mathcal{P} , they also generate extreme rays of the smaller convex cones listed in the diagram (3). In order to prove that every state in the set $\mathcal{E}_{\mathcal{A}}$ generates an extreme ray of the convex cone $\mathcal{A} \cap \mathcal{X}$, we first prove a technical lemma which will play a key role in characterization of extreme rays of the other cones.

Lemma 3.4. *Suppose that a three qubit X -state $\rho = \mathsf{X}(a, b, z)$ in \mathcal{A} (respectively, \mathcal{B} and \mathcal{C}) is decomposed as*

$$\rho = \mathsf{X}(a', b', z') + \mathsf{X}(a'', b'', z'')$$

in $\mathcal{A} \cap \mathcal{X}$ (respectively, $\mathcal{B} \cap \mathcal{X}$ and $\mathcal{C} \cap \mathcal{X}$). If

$$\sqrt{a_i b_i} = \sqrt{a_j b_j} = |z_i|$$

for $\{i, j\} = \{1, 4\}$ or $\{2, 3\}$ (respectively, $\{i, j\} = \{1, 3\}$ or $\{2, 4\}$, and $\{i, j\} = \{1, 2\}$ or $\{3, 4\}$), then we have

$$(a'_i, a'_j, b'_i, b'_j, z'_i) \parallel (a''_i, a''_j, b''_i, b''_j, z''_i).$$

Proof. Let $k = i, j$. We have

$$|z_i| = |z'_i + z''_i| \leq |z'_i| + |z''_i| \leq \sqrt{a'_k b'_k} + \sqrt{a''_k b''_k} \leq \sqrt{a'_k + a''_k} \sqrt{b'_k + b''_k} = \sqrt{a_k b_k}$$

by $S_1[i, j]$ and the Cauchy-Schwartz inequality. Since $|z_i| = \sqrt{a_k b_k}$, we have

$$z'_i \parallel z''_i, \quad \sqrt{a'_k b'_k} = |z'_i|, \quad \sqrt{a''_k b''_k} = |z''_i|, \quad (\sqrt{a'_k}, \sqrt{a''_k}) \parallel (\sqrt{b'_k}, \sqrt{b''_k}).$$

Let $(\sqrt{a'_k}, \sqrt{a''_k}) = \lambda_k (\sqrt{b'_k}, \sqrt{b''_k})$ for $\lambda_k > 0$. Then we have

$$\frac{z''_i}{z'_i} = \frac{|z''_i|}{|z'_i|} = \frac{\sqrt{a''_k b''_k}}{\sqrt{a'_k b'_k}} = \frac{\lambda_k b''_k}{\lambda_k b'_k} = \frac{b''_k}{b'_k} = \frac{a''_k / \lambda_k^2}{a'_k / \lambda_k^2} = \frac{a''_k}{a'_k},$$

as it was required. \square

Theorem 3.5. *We have*

$$\text{Ext}(\mathcal{A} \cap \mathbf{X}) = \mathcal{E}_{\mathcal{A}} \cup \Delta, \quad \text{Ext}(\mathcal{B} \cap \mathbf{X}) = \mathcal{E}_{\mathcal{B}} \cup \Delta \quad \text{and} \quad \text{Ext}(\mathcal{C} \cap \mathbf{X}) = \mathcal{E}_{\mathcal{C}} \cup \Delta.$$

Proof. It suffices to show that every state in the set $\mathcal{E}_{\mathcal{A}}$ generates an extreme ray of the convex cone $\mathcal{A} \cap \mathbf{X}$. Suppose that $\varrho = \mathbf{X}(a, b, z)$ satisfies the condition $S_1^e[1, 4]$ and

$$\varrho = \mathbf{X}(a', b', z') + \mathbf{X}(a'', b'', z'') \quad \text{in } \mathcal{A} \cap \mathbf{X}.$$

For $j = 2, 3$, we see that $a_j = b_j = 0$ implies $a'_j = b'_j = z'_j = a''_j = b''_j = z''_j = 0$. Applying Lemma 3.4 with $(i, j) = (1, 4)$ and $(i, j) = (4, 1)$, we get $(a', b', z') \parallel (a'', b'', z'')$, as it was required. The same argument works for the case of $S_1^e[2, 3]$. \square

In the remainder of this section, we look for extreme rays of $\mathcal{A}^\circ \cap \mathbf{X}$, $\mathcal{B}^\circ \cap \mathbf{X}$ and $\mathcal{C}^\circ \cap \mathbf{X}$. To do this, we consider the condition

$$W_1^e[i, j] : \quad \sqrt{s_i t_i} = |u_j| = 1, \quad \text{the others are zero,}$$

for $i \neq j$, and define

$$\mathcal{E}_{\mathcal{A}^\circ} = \{W = \mathbf{X}(s, t, u) : W_1^e[i, j] \text{ holds for some } (i, j) = (1, 4), (4, 1), (2, 3), (3, 2)\},$$

$$\mathcal{E}_{\mathcal{B}^\circ} = \{W = \mathbf{X}(s, t, u) : W_1^e[i, j] \text{ holds for some } (i, j) = (1, 3), (3, 1), (2, 4), (4, 2)\},$$

$$\mathcal{E}_{\mathcal{C}^\circ} = \{W = \mathbf{X}(s, t, u) : W_1^e[i, j] \text{ holds for some } (i, j) = (1, 2), (2, 1), (3, 4), (4, 3)\}.$$

We also consider the following set

$$\mathcal{W}^\Delta = \{W = \mathbf{X}_i(r, r^{-1}, e^{i\theta}) : i = 1, 2, 3, 4, r > 0, \theta \in \mathbb{R}\}.$$

Lemma 3.6. *For a given self-adjoint \mathbf{X} -shaped matrix $\varrho = \mathbf{X}(a, b, z)$, the following are equivalent:*

- (i) $\langle W, \varrho \rangle \geq 0$ for each $W \in \mathcal{E}_{\mathcal{A}^\circ} \cup \Delta \cup \mathcal{W}^\Delta$;
- (ii) ϱ is a state satisfying the inequalities $S_1[1, 4]$ and $S_1[2, 3]$;
- (iii) $\langle W, \varrho \rangle \geq 0$ for each $W \in \mathcal{A}^\circ$.

Proof. The equivalence between (ii) and (iii) follows from Proposition 3.1. Therefore, it suffices to show the direction (i) \implies (ii). Since $\langle W, \varrho \rangle \geq 0$ for $W \in \Delta$, we have $a_i, b_i \geq 0$. By taking $\varrho + \varepsilon I$ into account as in the proof of Lemma 3.2, we may assume that $a_i, b_i > 0$ without loss of generality. Then, we can consider

$$W_{i,j} := \mathbf{X}_i \left(\sqrt{\frac{b_i}{a_i}}, \sqrt{\frac{a_i}{b_i}}, 0 \right) + \mathbf{X}_j(0, 0, -e^{i\theta_j}) \in \mathcal{E}_{\mathcal{A}^\circ} \cup \mathcal{W}^\Delta$$

for $(i, j) = (1, 4), (4, 1), (2, 3), (3, 2)$ or $i = j$. Note that $W_{i,i} \in \mathcal{W}^\Delta$. We see that ϱ is a state by $\langle W_{i,i}, \varrho \rangle \geq 0$, and the inequalities $S_1[1, 4]$ and $S_1[2, 3]$ follow from $\langle W_{i,j}, \varrho \rangle \geq 0$ for $(i, j) = (1, 4), (4, 1), (2, 3), (3, 2)$. \square

As for extreme rays of the dual cones, we also begin with a technical lemma which is a witness counterpart to Lemma 3.4.

Lemma 3.7. *Suppose that a three qubit self-adjoint X-shaped matrix $W = \mathsf{X}(s, t, u)$ in \mathcal{A}° (respectively, \mathcal{B}° and \mathcal{C}°) is decomposed as*

$$W = \mathsf{X}(s', t', u') + \mathsf{X}(s'', t'', u'')$$

in $\mathcal{A}^\circ \cap \mathsf{X}$ (respectively, $\mathcal{B}^\circ \cap \mathsf{X}$ and $\mathcal{C}^\circ \cap \mathsf{X}$). If

$$\sqrt{s_i t_i} = |u_k| \quad \text{and} \quad s_j = t_j = 0$$

for $\{i, j\} = \{1, 4\}$ or $\{2, 3\}$ (respectively, $\{i, j\} = \{1, 3\}$ or $\{2, 4\}$, and $\{i, j\} = \{1, 2\}$ or $\{3, 4\}$) and $k \in \{i, j\}$, then we have

$$s'_j = t'_j = 0 = s''_j = t''_j, \quad u'_\ell = 0 = u''_\ell \quad \text{and} \quad (s'_i, t'_i, u'_k) \parallel (s''_i, t''_i, u''_k)$$

for ℓ with $\{k, \ell\} = \{i, j\}$.

Proof. The condition $s_j = t_j = 0$ implies $s'_j = t'_j = 0 = s''_j = t''_j$. We have

$$\begin{aligned} |u_k| &= |u'_k + u''_k| \\ &\leq |u'_k| + |u''_k| \\ &\leq \sqrt{s'_i} \sqrt{t'_i} - |u'_\ell| + \sqrt{s''_i} \sqrt{t''_i} - |u''_\ell| \\ &\leq \sqrt{s'_i + s''_i} \sqrt{t'_i + t''_i} - |u'_\ell| - |u''_\ell| \\ &= \sqrt{s_i t_i} - |u'_\ell| - |u''_\ell| \\ &\leq \sqrt{s_i t_i} \end{aligned}$$

by $W_1[i, j] = W_1[k, \ell]$ and the Cauchy-Schwartz inequality. Since $|u_k| = \sqrt{s_i t_i}$, we have

$$u'_k \parallel u''_k, \quad u'_\ell = 0 = u''_\ell, \quad \sqrt{s'_i t'_i} = |u'_k|, \quad \sqrt{s''_i t''_i} = |u''_k|, \quad (\sqrt{s'_i}, \sqrt{s''_i}) \parallel (\sqrt{t'_i}, \sqrt{t''_i}).$$

Let $(\sqrt{s'_i}, \sqrt{s''_i}) = \lambda(\sqrt{t'_i}, \sqrt{t''_i})$ for $\lambda > 0$. Then we have

$$\frac{u''_k}{u'_k} = \frac{|u''_k|}{|u'_k|} = \frac{\sqrt{s''_i t''_i}}{\sqrt{s'_i t'_i}} = \frac{\lambda t''_i}{\lambda t'_i} = \frac{t''_i}{t'_i} = \frac{s''_i / \lambda^2}{s'_i / \lambda^2} = \frac{s''_i}{s'_i}.$$

□

Theorem 3.8. *We have*

$$\text{Ext}(\mathcal{A}^\circ \cap \mathsf{X}) = \mathcal{E}_{\mathcal{A}^\circ} \cup \Delta \cup \mathcal{W}^\Delta, \quad \text{Ext}(\mathcal{B}^\circ \cap \mathsf{X}) = \mathcal{E}_{\mathcal{B}^\circ} \cup \Delta \cup \mathcal{W}^\Delta, \quad \text{Ext}(\mathcal{C}^\circ \cap \mathsf{X}) = \mathcal{E}_{\mathcal{C}^\circ} \cup \Delta \cup \mathcal{W}^\Delta.$$

Proof. It suffices to show that every ray in $\mathcal{E}_{\mathcal{A}^\circ} \cup \Delta \cup \mathcal{W}^\Delta$ is extreme by Proposition 2.1. It is easy to see that diagonal states in Δ generate extreme rays of the convex cone $\mathcal{A}^\circ \cap \mathsf{X}$ by the conditions $W_1[1, 4]$ and $W_1[2, 3]$. For the remaining cases for $\mathcal{E}_{\mathcal{A}^\circ} \cup \mathcal{W}^\Delta$, we take $i = 1, 4$ and may assume that $W = \mathsf{X}(s, t, u)$ satisfies

$$\sqrt{s_1 t_1} = |u_i| > 0 \quad \text{and} \quad s_k = t_k = u_\ell = 0 \text{ for } k \neq 1 \text{ and } \ell \neq i.$$

Suppose that $W = \mathsf{X}(s', t', u') + \mathsf{X}(s'', t'', u'')$ in $\mathcal{A}^\circ \cap \mathsf{X}$. For $k = 2, 3, 4$, the condition $s_k = t_k = 0$ implies $s'_k = t'_k = 0 = s''_k = t''_k$. Combining this with $W_1[2, 3]$, we also have $u'_2 = u'_3 = 0 = u''_2 = u''_3$. Applying Lemma 3.7 with $(i, j, k, \ell) = (1, 4, 1, 4), (1, 4, 4, 1)$, we get $(s', t', u') \parallel (s'', t'', u'')$. □

4. FULL BI-SEPARABILITY AND BI-SEPARABILITY

In this section, we consider convex cones $\mathcal{A} \cap \mathcal{B} \cap \mathcal{C}$ for full bi-separable states and $\mathcal{A} + \mathcal{B} + \mathcal{C}$ for bi-separable states, together with their dual cones $\mathcal{A}^\circ + \mathcal{B}^\circ + \mathcal{C}^\circ$ and $\mathcal{A}^\circ \cap \mathcal{B}^\circ \cap \mathcal{C}^\circ$, respectively. We first note that $\rho = \mathbf{X}(a, b, z) \in \mathcal{A} \cap \mathcal{B} \cap \mathcal{C}$ if and only if $S_1[i, j]$ holds for every $i, j = 1, 2, 3, 4$, which is equivalent to the PPT condition of ρ [20, Theorem 5.3]. In order to find extreme rays of the cone $\mathcal{A} \cap \mathcal{B} \cap \mathcal{C} \cap \mathbf{X}$, we consider the condition

$$S_3^e : \quad \sqrt{a_i b_i} = |z_i| = 1, \quad i = 1, 2, 3, 4,$$

and define

$$\mathcal{E}_{\mathcal{A} \cap \mathcal{B} \cap \mathcal{C}} = \{\rho = \mathbf{X}(a, b, z) : S_3^e \text{ holds}\}.$$

We also recall the inequality

$$W_3 : \quad \sum_{i=1}^4 \sqrt{s_i t_i} \geq \sum_{i=1}^4 |u_i|,$$

which appears in the characterization of decomposability of \mathbf{X} -shaped entanglement witnesses in [20, Theorem 5.5].

Lemma 4.1. *For a given self-adjoint \mathbf{X} -shaped matrix $W = \mathbf{X}(s, t, u)$, the following are equivalent.*

- (i) $\langle W, \rho \rangle \geq 0$ for each $\rho \in \mathcal{E}_{\mathcal{A} \cap \mathcal{B} \cap \mathcal{C}} \cup \Delta$;
- (ii) $s_i, t_i \geq 0$ for $i = 1, 2, 3, 4$, and the inequality W_3 holds;
- (iii) $\langle W, \rho \rangle \geq 0$ for each $\rho \in \mathcal{A} \cap \mathcal{B} \cap \mathcal{C}$.

Proof. For the direction (i) \implies (ii), we first obtain $s_i, t_i \geq 0$ from $\langle W, \rho \rangle \geq 0$ for $\rho \in \Delta$. In order to prove the inequality W_3 , we may assume that $s_i, t_i > 0$ as in the proof of Lemma 3.2. We can consider the state ρ defined by

$$\rho := \mathbf{X}((r_1, r_2, r_3, r_4), (r_1^{-1}, r_2^{-1}, r_3^{-1}, r_4^{-1}), (-e^{-i\theta_1}, -e^{-i\theta_2}, -e^{-i\theta_3}, -e^{-i\theta_4}))$$

with $r_i = \sqrt{\frac{t_i}{s_i}}$ and $\theta_k = \arg u_k$. This state ρ belongs to $\mathcal{E}_{\mathcal{A} \cap \mathcal{B} \cap \mathcal{C}}$, and so $\langle W, \rho \rangle \geq 0$ gives rise to the inequality W_3 . For (ii) \implies (iii), it suffices to show that $\langle W, \rho \rangle \geq 0$ for $\rho = \mathbf{X}(a, b, z)$ satisfying $S_1[i, j]$ for all i, j and $W = \mathbf{X}(s, t, u)$ satisfying W_3 by Corollary 2.3. Indeed, taking M satisfying $\sqrt{a_i b_i} \geq M \geq |z_j|$ for each $i, j = 1, 2, 3, 4$, we have

$$\sum_{i=1}^4 \sqrt{s_i t_i} \sqrt{a_i b_i} \geq M \sum_{i=1}^4 \sqrt{s_i t_i} \geq M \sum_{i=1}^4 |u_i| \geq \sum_{i=1}^4 |z_i| |u_i|,$$

which implies $\langle W, \rho \rangle \geq 0$, as in (6). \square

Since $(\mathcal{A} \cap \mathcal{B} \cap \mathcal{C})^\circ = \mathcal{A}^\circ + \mathcal{B}^\circ + \mathcal{C}^\circ$, the equivalence (ii) \iff (iii) in Lemma 4.1 gives another proof for [20, Theorem 5.5] which uses the duality principle.

Proposition 4.2. [20, Theorem 5.5] *An X-shaped self-adjoint matrix $W = \mathsf{X}(s, t, u)$ with nonnegative diagonals belongs to $\mathcal{A}^\circ + \mathcal{B}^\circ + \mathcal{C}^\circ$ if and only if the inequality W_3 holds.*

For convex cones C_1 and C_2 , it is clear that $\text{Ext}(C_1 + C_2) \subset \text{Ext}(C_1) \cup \text{Ext}(C_2)$ in general. Therefore, we see that $\text{Ext}((\mathcal{A}^\circ + \mathcal{B}^\circ + \mathcal{C}^\circ) \cap \mathsf{X})$ is contained in the union of $\text{Ext}(\mathcal{A}^\circ \cap \mathsf{X})$, $\text{Ext}(\mathcal{B}^\circ \cap \mathsf{X})$ and $\text{Ext}(\mathcal{C}^\circ \cap \mathsf{X})$ by Corollary 2.4. We show that they actually coincide.

Theorem 4.3. *We have the following:*

- (i) $\text{Ext}(\mathcal{A} \cap \mathcal{B} \cap \mathcal{C} \cap \mathsf{X}) = \mathcal{E}_{\mathcal{A} \cap \mathcal{B} \cap \mathcal{C}} \cup \Delta$;
- (ii) $\text{Ext}((\mathcal{A}^\circ + \mathcal{B}^\circ + \mathcal{C}^\circ) \cap \mathsf{X}) = \text{Ext}(\mathcal{A}^\circ \cap \mathsf{X}) \cup \text{Ext}(\mathcal{B}^\circ \cap \mathsf{X}) \cup \text{Ext}(\mathcal{C}^\circ \cap \mathsf{X})$.

Proof. For (i), it remains to show that every PPT state in $\mathcal{E}_{\mathcal{A} \cap \mathcal{B} \cap \mathcal{C}}$ generates an extreme ray of the cone $\mathcal{A} \cap \mathcal{B} \cap \mathcal{C}$. Suppose that $\varrho = \mathsf{X}(a, b, z)$ satisfies the condition S_3^e and

$$\varrho = \mathsf{X}(a', b', z') + \mathsf{X}(a'', b'', z'') \quad \text{in } \mathcal{A} \cap \mathcal{B} \cap \mathcal{C} \cap \mathsf{X}.$$

Applying Lemma 3.4 with all the pairs (i, j) , we conclude $(a', b', z') \parallel (a'', b'', z'')$, and this completes the proof of (i).

In order to prove (ii), it suffices to show $\text{Ext}(\mathcal{A}^\circ \cap \mathsf{X}) \subset \text{Ext}((\mathcal{A}^\circ + \mathcal{B}^\circ + \mathcal{C}^\circ) \cap \mathsf{X})$. It is easy to see that diagonal states in Δ generate extreme rays in the convex cone $(\mathcal{A}^\circ + \mathcal{B}^\circ + \mathcal{C}^\circ) \cap \mathsf{X}$ by the condition W_3 . We will show that $W = \mathsf{X}(rE_1, r^{-1}E_1, e^{i\theta}E_j)$ generates an extreme ray of the cone $(\mathcal{A}^\circ + \mathcal{B}^\circ + \mathcal{C}^\circ) \cap \mathsf{X}$ for $j = 1, 4$. Suppose that

$$(7) \quad W = \mathsf{X}(s', t', u') + \mathsf{X}(s'', t'', u'') \quad \text{in } (\mathcal{A}^\circ + \mathcal{B}^\circ + \mathcal{C}^\circ) \cap \mathsf{X}.$$

For $i = 2, 3, 4$, $s_i = 0 = t_i$ implies that $s'_i = t'_i = 0 = s''_i = t''_i$. By W_3 , we have

$$\begin{aligned} 1 = |e^{i\theta}| &= |u'_j + u''_j| \leq |u'_j| + |u''_j| \leq \sum_{k=1}^4 (|u'_k| + |u''_k|) \\ &\leq \sqrt{s'_1 t'_1} + \sqrt{s''_1 t''_1} \leq \sqrt{s'_1 + s''_1} \sqrt{t'_1 + t''_1} = 1, \end{aligned}$$

and so it follows that $u'_k = u''_k = 0$ for $k \neq j$. Therefore, the summands in (7) belong to the cone $\mathcal{A}^\circ \cap \mathsf{X}$ by W_3 again, and we may apply Theorem 3.8. \square

Now, we turn our attention to the cone $\mathcal{A}^\circ \cap \mathcal{B}^\circ \cap \mathcal{C}^\circ$ and its dual cone. For each $i = 1, 2, 3, 4$, we consider the condition

$$\boxed{W_3^e[i] : \quad |u_i| = \sqrt{s_j t_j} = \sqrt{s_k t_k} = \sqrt{s_\ell t_\ell} = 1, \quad \text{the others are zero,}}$$

where j, k, ℓ are chosen so that i, j, k, ℓ are mutually distinct, and define

$$\mathcal{E}_{\mathcal{A}^\circ \cap \mathcal{B}^\circ \cap \mathcal{C}^\circ} = \{W = \mathsf{X}(s, t, u) : W_3^e[i] \text{ holds for some } i = 1, 2, 3, 4\}.$$

We also consider the following inequality

$$S_3 : \quad \sum_{j \neq i} \sqrt{a_j b_j} \geq |z_i|, \quad i = 1, 2, 3, 4.$$

These are exactly the inequalities which appear in the necessary criteria [16] for bi-separability. We also refer to [17] for necessary criteria of multi-qubit bi-separable states. If ρ itself is X-shaped, then the converse is also true [18]. The authors have shown in [20, Corollary 3.4] that even a PPT mixture satisfies the multi-qubit analogue of S_3 , to recover the above characterization of bi-separability of multi-qubit X-states. We give here another alternative proof using the duality.

Lemma 4.4. *For a given self-adjoint X-shaped matrix $\rho = \mathbf{X}(a, b, z)$, the following are equivalent.*

- (i) $\langle W, \rho \rangle \geq 0$ for each $W \in \mathcal{E}_{\mathcal{A}^\circ \cap \mathcal{B}^\circ \cap \mathcal{C}^\circ} \cup \mathcal{W}^\Delta \cup \Delta$;
- (ii) ρ is a state satisfying the inequality S_3 ;
- (iii) $\langle W, \rho \rangle \geq 0$ for each $W \in \mathcal{A}^\circ \cap \mathcal{B}^\circ \cap \mathcal{C}^\circ$.

Proof. For the direction (i) \implies (ii), we first note that ρ is a state as in the proof of Lemma 3.6. Now, we consider

$$W := \mathbf{X} \left(\left(0, \sqrt{\frac{a_2}{b_2}}, \sqrt{\frac{a_3}{b_3}}, \sqrt{\frac{a_4}{b_4}} \right), \left(0, \sqrt{\frac{b_2}{a_2}}, \sqrt{\frac{b_3}{a_3}}, \sqrt{\frac{b_4}{a_4}} \right), (-e^{-i\theta_1}, 0, 0, 0) \right),$$

which belongs to $\mathcal{E}_{\mathcal{A}^\circ \cap \mathcal{B}^\circ \cap \mathcal{C}^\circ}$, where $\theta_1 = \arg z_1$. Then, we have

$$0 \leq \frac{1}{2} \langle \rho, W \rangle = \sum_{j \neq 1} \sqrt{a_j b_j} - |z_1|.$$

The other inequalities come out by the same way.

For the direction (ii) \implies (iii), it suffices to show the following:

$$S_3, W_1[i, j] \text{ for } i, j = 1, 2, 3, 4 \text{ with } i \neq j \implies \langle \mathbf{X}(s, t, u), \mathbf{X}(a, b, z) \rangle \geq 0$$

by Corollary 2.3 and Proposition 3.3. The inequality $\langle W, \rho \rangle \geq 0$ is trivial when W is positive, that is, $\sqrt{s_i t_i} \geq |u_i|$ for all $i = 1, 2, 3, 4$. Suppose that W is not positive, and so there exists $i_0 \in \{1, 2, 3, 4\}$ such that $\sqrt{s_{i_0} t_{i_0}} < |u_{i_0}|$, say $\sqrt{s_1 t_1} < |u_1|$ without loss of generality. We have

$$(\sqrt{s_1 t_1} + \sqrt{s_i t_i}) \sqrt{a_i b_i} \geq (|u_1| + |u_i|) \sqrt{a_i b_i}, \quad i = 2, 3, 4,$$

by $W_1[1, i]$. Summing up, we also have

$$\begin{aligned} \sqrt{s_1 t_1} \sum_{i=2}^4 \sqrt{a_i b_i} + \sum_{i=2}^4 \sqrt{s_i t_i} \sqrt{a_i b_i} &\geq |u_1| \sum_{i=2}^4 \sqrt{a_i b_i} + \sum_{i=2}^4 |u_i| \sqrt{a_i b_i} \\ &\geq |u_1| \sum_{i=2}^4 \sqrt{a_i b_i} + \sum_{i=2}^4 |u_i| |z_i|, \end{aligned}$$

which implies

$$\begin{aligned} \sum_{i=2}^4 \left(\sqrt{s_i t_i} \sqrt{a_i b_i} - |u_i| |z_i| \right) &\geq (|u_1| - \sqrt{s_1 t_1}) \sum_{i=2}^4 \sqrt{a_i b_i} \\ &\geq (|u_1| - \sqrt{s_1 t_1}) |z_1| \geq |u_1| |z_1| - \sqrt{s_1 t_1} \sqrt{a_1 b_1}, \end{aligned}$$

by S_3 and $\sqrt{s_1 t_1} < |u_1|$. Therefore, we have $\sum_{i=1}^4 \sqrt{s_i t_i} \sqrt{a_i b_i} \geq \sum_{i=1}^4 |u_i| |z_i|$, which completes the proof by (6). \square

Since the dual cone of $\mathcal{A}^\circ \cap \mathcal{B}^\circ \cap \mathcal{C}^\circ$ is just $\mathcal{A} + \mathcal{B} + \mathcal{C}$, we recover the following characterization of biseparability of three qubit states. Especially, every three qubit biseparable state with the X-part $\mathsf{X}(a, b, z)$ must satisfy the inequalities S_3 , as it was observed in [16].

Proposition 4.5. [16, 20, 18] *For a three qubit X-state $\rho = \mathsf{X}(a, b, z)$, the following are equivalent:*

- (i) ρ belongs to $\mathcal{A} + \mathcal{B} + \mathcal{C}$;
- (ii) the inequality S_3 holds.

As for extreme rays, we also have the following:

Theorem 4.6. *We have the following:*

- (i) $\text{Ext}((\mathcal{A} + \mathcal{B} + \mathcal{C}) \cap \mathsf{X}) = \text{Ext}(\mathcal{A} \cap \mathsf{X}) \cup \text{Ext}(\mathcal{B} \cap \mathsf{X}) \cup \text{Ext}(\mathcal{C} \cap \mathsf{X})$;
- (ii) $\text{Ext}(\mathcal{A}^\circ \cap \mathcal{B}^\circ \cap \mathcal{C}^\circ \cap \mathsf{X}) = \mathcal{E}_{\mathcal{A}^\circ \cap \mathcal{B}^\circ \cap \mathcal{C}^\circ} \cup \mathcal{W}^\Delta \cup \Delta$.

Proof. For (i), it suffices to show $\text{Ext}(\mathcal{A} \cap \mathsf{X}) \subset \text{Ext}((\mathcal{A} + \mathcal{B} + \mathcal{C}) \cap \mathsf{X})$. Suppose that $\rho = \mathsf{X}(a, b, z)$ satisfies the condition $S_1^e[1, 4]$, and

$$(8) \quad \rho = \mathsf{X}(a', b', z') + \mathsf{X}(a'', b'', z'') \quad \text{in } (\mathcal{A} + \mathcal{B} + \mathcal{C}) \cap \mathsf{X}.$$

Then we have $a'_i = a''_i = b'_i = b''_i = 0$ for $i = 2, 3$, which also implies that $z'_i = z''_i = 0$ for $i = 2, 3$. By the inequality S_3 , the summands in (8) must belong to the cone \mathcal{A} . Therefore, we can apply Theorem 3.5.

As for (ii), we note that matrices in Δ and \mathcal{W}^Δ generate extreme rays in $\mathcal{A}^\circ \cap \mathsf{X}$, and so they also generate extreme rays in the smaller cone $\mathcal{A}^\circ \cap \mathcal{B}^\circ \cap \mathcal{C}^\circ \cap \mathsf{X}$. Suppose that $W = \mathsf{X}(s, t, u)$ satisfies $W_3^e[1]$ and

$$W = \mathsf{X}(s', t', u') + \mathsf{X}(s'', t'', u'') \quad \text{in } \mathcal{A}^\circ \cap \mathcal{B}^\circ \cap \mathcal{C}^\circ \cap \mathsf{X}.$$

Applying Lemma 3.7 with $(i, j, k, \ell) = (2, 1, 1, 2), (3, 1, 1, 3), (4, 1, 1, 4)$, we get $(s', t', u') // (s'', t'', u'')$. \square

It was shown in [20, Theorem 4.1] that $W \in \mathcal{E}_{\mathcal{A}^\circ \cap \mathcal{B}^\circ \cap \mathcal{C}^\circ}$ is an optimal genuine entanglement witness. This means that the set $\{\rho \in \mathcal{P} : \langle W, \rho \rangle < 0\}$ of genuine entanglement detected by W is maximal with respect to the inclusion. It is easy to see that extremeness implies optimality. We have shown in Theorem 4.6 that $W \in \mathcal{E}_{\mathcal{A}^\circ \cap \mathcal{B}^\circ \cap \mathcal{C}^\circ}$ is extreme in the cone $\mathcal{A}^\circ \cap \mathcal{B}^\circ \cap \mathcal{C}^\circ \cap \mathsf{X}$. It would be interesting to ask if they are extreme in the much bigger convex cone $\mathcal{A}^\circ \cap \mathcal{B}^\circ \cap \mathcal{C}^\circ$.

5. INTERSECTIONS AND CONVEX HULLS OF TWO BASIC CONES

In this section, we consider the following convex cones

$$\mathcal{A} \cap \mathcal{B}, \quad \mathcal{B} \cap \mathcal{C}, \quad \mathcal{C} \cap \mathcal{A}, \quad \mathcal{A} + \mathcal{B}, \quad \mathcal{B} + \mathcal{C}, \quad \mathcal{C} + \mathcal{A},$$

together with their dual cones:

$$\mathcal{A}^\circ + \mathcal{B}^\circ, \quad \mathcal{B}^\circ + \mathcal{C}^\circ, \quad \mathcal{C}^\circ + \mathcal{A}^\circ, \quad \mathcal{A}^\circ \cap \mathcal{B}^\circ, \quad \mathcal{B}^\circ \cap \mathcal{C}^\circ, \quad \mathcal{C}^\circ \cap \mathcal{A}^\circ.$$

We look for inequalities characterizing the above convex cones, together with extreme rays of the cones. As for intersections of two cones, we just put together inequalities for both cones. For a three qubit X-state $\varrho = \mathbf{X}(a, b, z)$, we have the following:

- $\varrho \in \mathcal{A} \cap \mathcal{B}$ if and only if $S_1[1, 4], S_1[2, 3], S_1[1, 3], S_1[2, 4]$ hold;
- $\varrho \in \mathcal{B} \cap \mathcal{C}$ if and only if $S_1[1, 3], S_1[2, 4], S_1[1, 2], S_1[3, 4]$ hold;
- $\varrho \in \mathcal{C} \cap \mathcal{A}$ if and only if $S_1[1, 2], S_1[3, 4], S_1[1, 4], S_1[2, 3]$ hold.

For an X-shaped $W = \mathbf{X}(s, t, u)$, we also have

- $W \in \mathcal{A}^\circ \cap \mathcal{B}^\circ$ if and only if $W_1[1, 4], W_1[2, 3], W_1[1, 3], W_1[2, 4]$ hold;
- $W \in \mathcal{B}^\circ \cap \mathcal{C}^\circ$ if and only if $W_1[1, 3], W_1[2, 4], W_1[1, 2], W_1[3, 4]$ hold;
- $W \in \mathcal{C}^\circ \cap \mathcal{A}^\circ$ if and only if $W_1[1, 2], W_1[3, 4], W_1[1, 4], W_1[2, 3]$ hold.

In order to find extreme rays of the cones $\mathcal{B} \cap \mathcal{C} \cap \mathbf{X}$, $\mathcal{C} \cap \mathcal{A} \cap \mathbf{X}$ and $\mathcal{A} \cap \mathcal{B} \cap \mathbf{X}$, we consider the condition

$$S_2^e[i, j] : \quad |z_i| = \sqrt{a_i b_i} = \sqrt{a_k b_k} = \sqrt{a_\ell b_\ell} = 1, \quad \text{the others are zero,}$$

for $i, j = 1, 2, 3, 4$ with $i \neq j$, where k, ℓ are chosen so that i, j, k, ℓ are mutually distinct.

Here, we point out that $a_j = 0 = b_j$. We define

$$\mathcal{E}_{\mathcal{B} \cap \mathcal{C}} = \{\varrho = \mathbf{X}(a, b, z) : S_2^e[i, j] \text{ holds for some } (i, j) = (1, 4), (4, 1), (2, 3), (3, 2)\},$$

$$\mathcal{E}_{\mathcal{C} \cap \mathcal{A}} = \{\varrho = \mathbf{X}(a, b, z) : S_2^e[i, j] \text{ holds for some } (i, j) = (1, 3), (3, 1), (2, 4), (4, 2)\},$$

$$\mathcal{E}_{\mathcal{A} \cap \mathcal{B}} = \{\varrho = \mathbf{X}(a, b, z) : S_2^e[i, j] \text{ holds for some } (i, j) = (1, 2), (2, 1), (3, 4), (4, 3)\},$$

and consider the following inequalities

$$W_2[i, j] : \quad \sum_{k \neq j} \sqrt{s_k t_k} \geq |u_i|, \quad \sum_{k \neq i} \sqrt{s_k t_k} \geq |u_j|,$$

for $i, j = 1, 2, 3, 4$ with $i \neq j$.

Lemma 5.1. *For a given self-adjoint X-shaped matrix $W = \mathbf{X}(s, t, u)$, the following are equivalent.*

- (i) $\langle W, \varrho \rangle \geq 0$ for each $\varrho \in \mathcal{E}_{\mathcal{B} \cap \mathcal{C}} \cup \mathcal{E}_{\mathcal{A} \cap \mathcal{B} \cap \mathcal{C}} \cup \Delta$ (respectively, $\mathcal{E}_{\mathcal{C} \cap \mathcal{A}} \cup \mathcal{E}_{\mathcal{A} \cap \mathcal{B} \cap \mathcal{C}} \cup \Delta$ and $\mathcal{E}_{\mathcal{A} \cap \mathcal{B}} \cup \mathcal{E}_{\mathcal{A} \cap \mathcal{B} \cap \mathcal{C}} \cup \Delta$);
- (ii) $s_i, t_i \geq 0$ for $i = 1, 2, 3, 4$ and the inequalities W_3 and $W_2[1, 4], W_2[2, 3]$ (respectively, $W_2[1, 3], W_2[2, 4]$ and $W_2[1, 2], W_2[3, 4]$) hold;
- (iii) $\langle W, \varrho \rangle \geq 0$ for each $\varrho \in \mathcal{B} \cap \mathcal{C}$ (respectively, $\mathcal{C} \cap \mathcal{A}$ and $\mathcal{A} \cap \mathcal{B}$).

Proof. The inequalities $s_i, t_i \geq 0$ and W_3 follow from Lemma 4.1. We will prove for $\mathcal{B} \cap \mathcal{C}$. The others follow by applying the operator $x_A \otimes x_B \otimes x_C \mapsto x_{\sigma(A)} \otimes x_{\sigma(B)} \otimes x_{\sigma(C)}$ for permutations σ on $\{A, B, C\}$. To prove (i) \implies (ii), we may assume that all the diagonal elements s_i and t_i are nonzero, and consider four X -states

$$\begin{aligned}\varrho_{1,4} &:= \mathsf{X} \left(\left(\sqrt{\frac{t_1}{s_1}}, \sqrt{\frac{t_2}{s_2}}, \sqrt{\frac{t_3}{s_3}}, 0 \right), \left(\sqrt{\frac{s_1}{t_1}}, \sqrt{\frac{s_2}{t_2}}, \sqrt{\frac{s_3}{t_3}}, 0 \right), (-e^{-i\theta_1}, 0, 0, 0) \right), \\ \varrho_{2,3} &:= \mathsf{X} \left(\left(\sqrt{\frac{t_1}{s_1}}, \sqrt{\frac{t_2}{s_2}}, 0, \sqrt{\frac{t_4}{s_4}} \right), \left(\sqrt{\frac{s_1}{t_1}}, \sqrt{\frac{s_2}{t_2}}, 0, \sqrt{\frac{s_4}{t_4}} \right), (0, -e^{-i\theta_2}, 0, 0) \right), \\ \varrho_{3,2} &:= \mathsf{X} \left(\left(\sqrt{\frac{t_1}{s_1}}, 0, \sqrt{\frac{t_3}{s_3}}, \sqrt{\frac{t_4}{s_4}} \right), \left(\sqrt{\frac{s_1}{t_1}}, 0, \sqrt{\frac{s_3}{t_3}}, \sqrt{\frac{s_4}{t_4}} \right), (0, 0, -e^{-i\theta_3}, 0) \right), \\ \varrho_{4,1} &:= \mathsf{X} \left(\left(0, \sqrt{\frac{t_2}{s_2}}, \sqrt{\frac{t_3}{s_3}}, \sqrt{\frac{t_4}{s_4}} \right), \left(0, \sqrt{\frac{s_2}{t_2}}, \sqrt{\frac{s_3}{t_3}}, \sqrt{\frac{s_4}{t_4}} \right), (0, 0, 0, -e^{-i\theta_4}) \right),\end{aligned}$$

with $\theta_k = \arg u_k$. These states belong to $\mathcal{E}_{\mathcal{B} \cap \mathcal{C}}$. We expand $\langle W, \varrho_{i,j} \rangle \geq 0$ to obtain $W_2[1, 4]$ and $W_2[2, 3]$.

For (ii) \implies (iii), it suffices to show $\langle W, \varrho \rangle \geq 0$ when $W = \mathsf{X}(s, t, u)$ satisfies $W_2[1, 4]$, $W_2[2, 3]$, W_3 , and $\varrho = \mathsf{X}(a, b, z)$ satisfies $S_1[1, 3]$, $S_1[2, 4]$, $S_1[1, 2]$, $S_1[3, 4]$ by Corollary 2.3 and Proposition 3.1. If $\varrho \in \mathcal{A} \cap \mathcal{B} \cap \mathcal{C}$, then this is trivial by W_3 and Proposition 4.2. So, we may assume that $\varrho \notin \mathcal{A}$, especially $|z_4| > \sqrt{a_1 b_1}$, without loss of generality. We begin with

$$\begin{aligned}\frac{1}{2} \langle W, \varrho \rangle &\geq \sum_{i=1}^4 \left(\sqrt{s_i t_i} \sqrt{a_i b_i} - |u_i| |z_i| \right) \\ &= \left(\sqrt{s_2 t_2} \sqrt{a_2 b_2} + \sqrt{s_3 t_3} \sqrt{a_3 b_3} + \sqrt{s_4 t_4} \sqrt{a_4 b_4} - |u_4| |z_4| \right) \\ &\quad + \left(\sqrt{s_1 t_1} \sqrt{a_1 b_1} - |u_1| |z_1| - |u_2| |z_2| - |u_3| |z_3| \right),\end{aligned}$$

as in (6). We have $\sqrt{a_i b_i} \geq |z_4|$ for $i = 2, 3, 4$ by $S_1[2, 4]$, $S_1[3, 4]$, and $\sqrt{a_1 b_1} \geq |z_i|$ for $i = 1, 2, 3$ by $S_1[1, 2]$, $S_1[1, 3]$. By the inequality $W_2[1, 4]$ and the assumption $|z_4| > \sqrt{a_1 b_1}$, we have

$$\begin{aligned}\frac{1}{2} \langle W, \varrho \rangle &\geq (\sqrt{s_2 t_2} + \sqrt{s_3 t_3} + \sqrt{s_4 t_4} - |u_4|) |z_4| + (\sqrt{s_1 t_1} - |u_1| - |u_2| - |u_3|) \sqrt{a_1 b_1} \\ &\geq (\sqrt{s_2 t_2} + \sqrt{s_3 t_3} + \sqrt{s_4 t_4} - |u_4|) \sqrt{a_1 b_1} + (\sqrt{s_1 t_1} - |u_1| - |u_2| - |u_3|) \sqrt{a_1 b_1} \\ &= \sqrt{a_1 b_1} \left(\sum_{i=1}^4 \sqrt{s_i t_i} - \sum_{i=1}^4 |u_i| \right).\end{aligned}$$

This is nonnegative by the inequality W_3 , as it was desired. \square

By the equivalence (ii) \iff (iii), we have the following criteria for the convex hull of two basic dual cones:

Theorem 5.2. *For a self-adjoint $W = \mathsf{X}(s, t, u)$ with nonnegative diagonals, we have the following:*

- (i) $W \in \mathcal{B}^\circ + \mathcal{C}^\circ$ if and only if $W_2[1, 4]$, $W_2[2, 3]$ and W_3 hold;

- (ii) $W \in \mathcal{C}^\circ + \mathcal{A}^\circ$ if and only if $W_2[1, 3]$, $W_2[2, 4]$ and W_3 hold;
- (iii) $W \in \mathcal{A}^\circ + \mathcal{B}^\circ$ if and only if $W_2[1, 2]$, $W_2[3, 4]$ and W_3 hold.

If W is a self-adjoint three qubit matrix with the \mathbf{X} -part $\mathbf{X}(s, t, u)$, then the ‘only if’ parts hold.

Theorem 5.3. *We have the following:*

- (i) $\text{Ext}(\mathcal{B} \cap \mathcal{C} \cap \mathbf{X}) = \mathcal{E}_{\mathcal{B} \cap \mathcal{C}} \cup \mathcal{E}_{\mathcal{A} \cap \mathcal{B} \cap \mathcal{C}} \cup \Delta$,
 $\text{Ext}(\mathcal{C} \cap \mathcal{A} \cap \mathbf{X}) = \mathcal{E}_{\mathcal{C} \cap \mathcal{A}} \cup \mathcal{E}_{\mathcal{A} \cap \mathcal{B} \cap \mathcal{C}} \cup \Delta$,
 $\text{Ext}(\mathcal{A} \cap \mathcal{B} \cap \mathbf{X}) = \mathcal{E}_{\mathcal{A} \cap \mathcal{B}} \cup \mathcal{E}_{\mathcal{A} \cap \mathcal{B} \cap \mathcal{C}} \cup \Delta$;
- (ii) $\text{Ext}((\mathcal{B}^\circ + \mathcal{C}^\circ) \cap \mathbf{X}) = \text{Ext}(\mathcal{B}^\circ \cap \mathbf{X}) \cup \text{Ext}(\mathcal{C}^\circ \cap \mathbf{X})$,
 $\text{Ext}((\mathcal{C}^\circ + \mathcal{A}^\circ) \cap \mathbf{X}) = \text{Ext}(\mathcal{C}^\circ \cap \mathbf{X}) \cup \text{Ext}(\mathcal{A}^\circ \cap \mathbf{X})$,
 $\text{Ext}((\mathcal{A}^\circ + \mathcal{B}^\circ) \cap \mathbf{X}) = \text{Ext}(\mathcal{A}^\circ \cap \mathbf{X}) \cup \text{Ext}(\mathcal{B}^\circ \cap \mathbf{X})$.

Proof. (i). We will prove the first identity. Suppose that $\varrho = \mathbf{X}(a, b, z)$ satisfies $S_2^e[1, 4]$ and

$$\varrho = \mathbf{X}(a', b', z') + \mathbf{X}(a'', b'', z'') \quad \text{in } \mathcal{B} \cap \mathcal{C} \cap \mathbf{X}.$$

The condition $a_4 = b_4 = 0$ implies $a'_4 = b'_4 = z'_j = 0 = a''_4 = b''_4 = z''_j$ for $j = 2, 3, 4$ by $S_1[2, 4]$, $S_1[3, 4]$. Applying Lemma 3.4 with $(i, j) = (1, 2)$ and $(1, 3)$, we get $(a', b', z') \parallel (a'', b'', z'')$.

Next, suppose that $\varrho = \mathbf{X}(a, b, z)$ satisfies S_3^e and

$$\varrho = \mathbf{X}(a', b', z') + \mathbf{X}(a'', b'', z'') \quad \text{in } \mathcal{B} \cap \mathcal{C} \cap \mathbf{X}.$$

Applying Lemma 3.4 with $(i, j) = (1, 2), (2, 1), (3, 1), (4, 2)$, we get $(a', b', z') \parallel (a'', b'', z'')$.

(ii). States in $\text{Ext}(\mathcal{B}^\circ \cap \mathbf{X}) \cup \text{Ext}(\mathcal{C}^\circ \cap \mathbf{X})$ generate extreme rays of the convex cone $(\mathcal{A}^\circ + \mathcal{B}^\circ + \mathcal{C}^\circ) \cap \mathbf{X}$ by Theorem 4.3. Therefore, they also generate extreme rays in the smaller cone $(\mathcal{B}^\circ + \mathcal{C}^\circ) \cap \mathbf{X}$. \square

Now, we look for extreme rays of $\mathcal{B}^\circ \cap \mathcal{C}^\circ \cap \mathbf{X}$ (respectively, $\mathcal{C}^\circ \cap \mathcal{A}^\circ \cap \mathbf{X}$ and $\mathcal{A}^\circ \cap \mathcal{B}^\circ \cap \mathbf{X}$) to get conditions for the cone $\mathcal{B} + \mathcal{C}$ (respectively, $\mathcal{C} + \mathcal{A}$ and $\mathcal{A} + \mathcal{B}$). To do this, we consider the condition

$$W_2^e[i, j] : \quad \sqrt{s_i t_i} = \sqrt{s_j t_j} = |u_k| = |u_\ell| = 1, \quad \text{the others are zero,}$$

for $i \neq j$, where k and ℓ are chosen so that i, j, k, ℓ are mutually distinct, and define

$$\mathcal{E}_{\mathcal{B}^\circ \cap \mathcal{C}^\circ} = \{W = \mathbf{X}(s, t, u) : W_2^e[i, j] \text{ holds for } (i, j) = (1, 4) \text{ or } (2, 3)\},$$

$$\mathcal{E}_{\mathcal{C}^\circ \cap \mathcal{A}^\circ} = \{W = \mathbf{X}(s, t, u) : W_2^e[i, j] \text{ holds for } (i, j) = (1, 3) \text{ or } (2, 4)\},$$

$$\mathcal{E}_{\mathcal{A}^\circ \cap \mathcal{B}^\circ} = \{W = \mathbf{X}(s, t, u) : W_2^e[i, j] \text{ holds for } (i, j) = (1, 2) \text{ or } (3, 4)\}.$$

We also consider the following inequalities

$$S_2[i, j] : \quad \min \{ \sqrt{a_i b_i} + \sqrt{a_j b_j}, \sqrt{a_k b_k} + \sqrt{a_\ell b_\ell} \} \geq \max \{ |z_i| + |z_j|, |z_k| + |z_\ell| \},$$

for $i \neq j$, where k, ℓ are chosen so that i, j, k, ℓ are mutually distinct.

These inequalities have been used in [11] to get necessary conditions for a three state ρ with the \mathbf{X} -part $\mathbf{X}(a, b, z)$ to belong to $\mathcal{B} + \mathcal{C}$, $\mathcal{C} + \mathcal{A}$ and $\mathcal{A} + \mathcal{B}$ respectively. We show in Theorem 5.5 that they provide actually sufficient conditions when ρ itself \mathbf{X} -shaped. Note that

- $S_2[1, 4]$ holds if and only if $S_2[2, 3]$ holds;
- $S_2[1, 3]$ holds if and only if $S_2[2, 4]$ holds;
- $S_2[1, 2]$ holds if and only if $S_2[3, 4]$ holds.

Lemma 5.4. *For a given self-adjoint \mathbf{X} -shaped matrix $\rho = \mathbf{X}(a, b, z)$, the following are equivalent.*

- (i) $\langle W, \rho \rangle \geq 0$ for each $W \in \mathcal{E}_{\mathcal{B}^\circ \cap \mathcal{C}^\circ} \cup \mathcal{W}^\Delta \cup \Delta$ (respectively, $\mathcal{E}_{\mathcal{C}^\circ \cap \mathcal{A}^\circ} \cup \mathcal{W}^\Delta \cup \Delta$ and $\mathcal{E}_{\mathcal{A}^\circ \cap \mathcal{B}^\circ} \cup \mathcal{W}^\Delta \cup \Delta$);
- (ii) ρ is a state satisfying the inequalities $S_2[1, 4]$ (respectively, $S_2[1, 3]$ and $S_2[1, 2]$) hold;
- (iii) $\langle W, \rho \rangle \geq 0$ for each $W \in \mathcal{B}^\circ \cap \mathcal{C}^\circ$ (respectively, $\mathcal{C}^\circ \cap \mathcal{A}^\circ$ and $\mathcal{A}^\circ \cap \mathcal{B}^\circ$).

Proof. Although the proof of the direction (i) \implies (ii) already appears in [11], we include it here for the completeness. We consider \mathbf{X} -shaped three qubit self-adjoint matrices

$$W := \mathbf{X} \left(\left(\sqrt{\frac{b_1}{a_1}}, 0, 0, \sqrt{\frac{b_4}{a_4}} \right), \left(\sqrt{\frac{a_1}{b_1}}, 0, 0, \sqrt{\frac{a_4}{b_4}} \right), (0, -e^{-i\theta_2}, -e^{-i\theta_3}, 0) \right)$$

$$W' := \mathbf{X} \left(\left(0, \sqrt{\frac{b_2}{a_2}}, \sqrt{\frac{b_3}{a_3}}, 0 \right), \left(0, \sqrt{\frac{a_2}{b_2}}, \sqrt{\frac{a_3}{b_3}}, 0 \right), (-e^{-i\theta_1}, 0, 0, -e^{-i\theta_4}) \right),$$

for $\theta_i = \arg z_i$. Then, both W and W' belong to $\mathcal{E}_{\mathcal{B}^\circ \cap \mathcal{C}^\circ}$. We have

$$0 \leq \frac{1}{2} \langle \rho, W \rangle = \sqrt{a_1 b_1} + \sqrt{a_4 b_4} - |z_2| - |z_3|,$$

$$0 \leq \frac{1}{2} \langle \rho, W' \rangle = \sqrt{a_2 b_2} + \sqrt{a_3 b_3} - |z_1| - |z_4|.$$

For the implication (ii) \implies (iii), suppose that $\rho = \mathbf{X}(a, b, z)$ satisfies $S_2[1, 4]$. By Corollary 2.3, it suffices to show $\langle W, \rho \rangle \geq 0$ for every $W = \mathbf{X}(s, t, u) \in \mathcal{B}^\circ \cap \mathcal{C}^\circ$. This is trivial when $\mathbf{X}(s, t, u)$ is positive, that is, $\sqrt{s_i t_i} \geq |u_i|$ for all $i = 1, 2, 3, 4$. We may assume without loss of generality that

$$(9) \quad 0 > \sqrt{s_1 t_1} - |u_1| = \min\{\sqrt{s_i t_i} - |u_i| : i = 1, 2, 3, 4\}.$$

Then we have

$$(\sqrt{s_1 t_1} + \sqrt{s_2 t_2}) \sqrt{a_2 b_2} \geq (|u_1| + |u_2|) \sqrt{a_2 b_2},$$

$$(\sqrt{s_1 t_1} + \sqrt{s_3 t_3}) \sqrt{a_3 b_3} \geq (|u_1| + |u_3|) \sqrt{a_3 b_3},$$

$$(|u_1| + \sqrt{s_4 t_4}) |z_4| \geq (\sqrt{s_1 t_1} + |u_4|) |z_4|,$$

where the first and second inequalities follow from $W_1[1, 2]$ and $W_1[1, 3]$, respectively, and the last one comes out from the equality in (9). Put $M = \sqrt{a_2 b_2} + \sqrt{a_3 b_3} - |z_4|$. Summing up the above three inequalities, we have

$$\sqrt{s_1 t_1} M + \sum_{i=2}^3 \sqrt{s_i t_i} \sqrt{a_i b_i} + \sqrt{s_4 t_4} |z_4| \geq |u_1| M + \sum_{i=2}^3 |u_i| \sqrt{a_i b_i} + |u_4| |z_4|,$$

which implies

$$\sqrt{s_1 t_1} M + \sum_{i=2}^4 \sqrt{s_i t_i} \sqrt{a_i b_i} \geq |u_1| M + \sum_{i=2}^4 |u_i| |z_i|.$$

Because $M \geq |z_1|$ by $S_2[1, 4]$ and $|u_1| - \sqrt{s_1 t_1} > 0$ by (9), we have

$$\begin{aligned} \sum_{i=2}^4 (\sqrt{s_i t_i} \sqrt{a_i b_i} - |u_i| |z_i|) &\geq (|u_1| - \sqrt{s_1 t_1}) M \\ &\geq (|u_1| - \sqrt{s_1 t_1}) |z_1| \geq |u_1| |z_1| - \sqrt{s_1 t_1} \sqrt{a_1 b_1}. \end{aligned}$$

This gives $\sum_{i=1}^4 \sqrt{s_i t_i} \sqrt{a_i b_i} \geq \sum_{i=1}^4 |u_i| |z_i|$, and $\langle W, \rho \rangle \geq 0$ by (6). \square

Because $\mathcal{B} + \mathcal{C}$ (respectively, $\mathcal{C} + \mathcal{A}$ and $\mathcal{A} + \mathcal{B}$) is the dual of $\mathcal{B}^\circ \cap \mathcal{C}^\circ$ (respectively, $\mathcal{C}^\circ \cap \mathcal{A}^\circ$ and $\mathcal{A}^\circ \cap \mathcal{B}^\circ$), the equivalence between (ii) and (iii) of Lemma 5.4 gives rise to the following characterization of the cone $\mathcal{B} + \mathcal{C}$ (respectively, $\mathcal{C} + \mathcal{A}$ and $\mathcal{A} + \mathcal{B}$) for X-states.

Theorem 5.5. *For a three qubit X-state $\rho = \mathsf{X}(a, b, z)$, we have the following:*

- (i) $\rho \in \mathcal{B} + \mathcal{C}$ if and only if $S_2[1, 4]$ holds;
- (ii) $\rho \in \mathcal{C} + \mathcal{A}$ if and only if $S_2[1, 3]$ holds;
- (iii) $\rho \in \mathcal{A} + \mathcal{B}$ if and only if $S_2[1, 2]$ holds.

For a general three qubit state ρ with the X-part $\mathsf{X}(a, b, z)$, the ‘only if’ parts hold.

Theorem 5.6. *We have the following:*

- (i) $\text{Ext}(\mathcal{B}^\circ \cap \mathcal{C}^\circ \cap \mathsf{X}) = \mathcal{E}_{\mathcal{B}^\circ \cap \mathcal{C}^\circ} \cup \mathcal{W}^\Delta \cup \Delta$,
 $\text{Ext}(\mathcal{C}^\circ \cap \mathcal{A}^\circ \cap \mathsf{X}) = \mathcal{E}_{\mathcal{C}^\circ \cap \mathcal{A}^\circ} \cup \mathcal{W}^\Delta \cup \Delta$,
 $\text{Ext}(\mathcal{A}^\circ \cap \mathcal{B}^\circ \cap \mathsf{X}) = \mathcal{E}_{\mathcal{A}^\circ \cap \mathcal{B}^\circ} \cup \mathcal{W}^\Delta \cup \Delta$;
- (ii) $\text{Ext}((\mathcal{B} + \mathcal{C}) \cap \mathsf{X}) = \text{Ext}(\mathcal{B} \cap \mathsf{X}) \cup \text{Ext}(\mathcal{C} \cap \mathsf{X})$,
 $\text{Ext}((\mathcal{C} + \mathcal{A}) \cap \mathsf{X}) = \text{Ext}(\mathcal{C} \cap \mathsf{X}) \cup \text{Ext}(\mathcal{A} \cap \mathsf{X})$,
 $\text{Ext}((\mathcal{A} + \mathcal{B}) \cap \mathsf{X}) = \text{Ext}(\mathcal{A} \cap \mathsf{X}) \cup \text{Ext}(\mathcal{B} \cap \mathsf{X})$.

Proof. (i). We will prove the first identity. Since elements in Δ and \mathcal{W}^Δ are extremal in $\mathcal{B}^\circ \cap \mathsf{X}$, they are also extremal in the smaller cone $\mathcal{B}^\circ \cap \mathcal{C}^\circ \cap \mathsf{X}$. Suppose that $W = \mathsf{X}(s, t, u)$ satisfies $W_2^e[1, 4]$ and

$$W = \mathsf{X}(s', t', u') + \mathsf{X}(s'', t'', u'') \quad \text{in } \mathcal{B}^\circ \cap \mathcal{C}^\circ \cap \mathsf{X}.$$

Applying Lemma 3.7 with $(i, j, k, \ell) = (1, 2, 2, 1), (4, 2, 2, 4), (1, 3, 3, 1), (4, 3, 3, 4)$, we get $(s', t', u') // (s'', t'', u'')$.

criteria	extreme	states	witnesses	extreme	criteria
$S_1[1, 4]$, $S_1[2, 3]$ Prop. 3.1	$S_1^e[1, 4]$ $S_1^e[2, 3]$ Th. 3.5	$\mathcal{A} \cap \mathcal{X}$	$\mathcal{A}^\circ \cap \mathcal{X}$	$W_1^e[1, 4]$ $W_1^e[4, 1]$ $W_1^e[2, 3]$ $W_1^e[3, 2]$ Th. 3.8	$W_1[1, 4]$ $W_1[2, 3]$ Prop. 3.3
$S_1[i, j]$ Prop. 3.1	S_3^e Th. 4.3(i)	$\mathcal{A} \cap \mathcal{B} \cap \mathcal{C} \cap \mathcal{X}$	$(\mathcal{A}^\circ + \mathcal{B}^\circ + \mathcal{C}^\circ) \cap \mathcal{X}$	$W_1^e[i, j]$ Th. 4.3(ii)	W_3 Prop. 4.2
S_3 Prop. 4.5	$S_1^e[i, j]$ Th. 4.6(i)	$(\mathcal{A} + \mathcal{B} + \mathcal{C}) \cap \mathcal{X}$	$\mathcal{A}^\circ \cap \mathcal{B}^\circ \cap \mathcal{C}^\circ \cap \mathcal{X}$	$W_3^e[j]$ Th. 4.6(ii)	$W_1[i, j]$ Prop. 3.3
$S_1[1, 4]$ $S_1[2, 3]$ $S_1[1, 3]$ $S_1[2, 4]$ Prop. 3.1	S_3^e $S_2^e[1, 2]$ $S_2^e[2, 1]$ $S_2^e[3, 4]$ $S_2^e[4, 3]$ Th. 5.3(i)	$\mathcal{A} \cap \mathcal{B} \cap \mathcal{X}$	$(\mathcal{A}^\circ + \mathcal{B}^\circ) \cap \mathcal{X}$	$W_1^e[1, 4]$ $W_1^e[4, 1]$ $W_1^e[2, 3]$ $W_1^e[3, 2]$ $W_1^e[1, 3]$ $W_1^e[3, 1]$ $W_1^e[2, 4]$ $W_1^e[4, 2]$ Th. 5.3(ii)	W_3 $W_2[1, 2]$ $W_2[3, 4]$ Th. 5.2
$S_2[1, 2]$ Th. 5.5	$S_1^e[1, 4]$ $S_1^e[2, 3]$ $S_1^e[1, 3]$ $S_1^e[2, 4]$ Th. 5.6(ii)	$(\mathcal{A} + \mathcal{B}) \cap \mathcal{X}$	$\mathcal{A}^\circ \cap \mathcal{B}^\circ \cap \mathcal{X}$	$W_2^e[1, 2]$ $W_2^e[3, 4]$ Th. 5.6(i)	$W_1[1, 4]$ $W_1[2, 3]$ $W_1[1, 3]$ $W_1[2, 4]$ Prop. 3.3

TABLE 1. Criteria and extreme rays of convex cones: Conditions for ‘criteria’ and ‘extreme’ are connected by ‘and’ and ‘or’, respectively.

(ii). Since states in $\text{Ext}(\mathcal{B} \cap \mathcal{X}) \cup \text{Ext}(\mathcal{C} \cap \mathcal{X})$ are extremal in the convex cone $(\mathcal{A} + \mathcal{B} + \mathcal{C}) \cap \mathcal{X}$ by Theorem 4.6, they are also extremal in the smaller cone $(\mathcal{B} + \mathcal{C}) \cap \mathcal{X}$.
□

6. SUMMARY

In this paper, we have considered the convex cones in the diagrams (3) and (4) arising from classification of partial separability/entanglement of three qubit states and their witnesses. For those convex cones, we got the following results:

- characterization for \mathcal{X} -shaped matrices by algebraic inequalities, which give rise to necessary criteria for general three qubit states/witnesses in terms of diagonal and anti-diagonal entries;
- finding all the extreme rays of the cones consisting of \mathcal{X} -shaped matrices, with which we may exhibit all \mathcal{X} -shaped matrices in the cones.

We summarize the results in Table 1. We note our characterization is one of very few cases when we may check separability by inequalities, without decomposing into the sum of pure product states. For example, we may check separability for $2 \otimes 2$ and $2 \otimes 3$

cases by the PPT condition. We may also check full separability of multi-qubit X-states by inequalities [14, 15, 12]. Checking separability with inequalities in this paper was possible through the duality and characterizing extreme rays of the dual cones.

This work has been partly motivated by the questions [10] on the existence of states in the seven classes arising in the classification of partial entanglement, including the following classes:

$$\begin{aligned}\mathcal{C}^{2,6,1} &:= \mathcal{A} \cap (\mathcal{B} + \mathcal{C}) \cap \mathcal{B}^c \cap \mathcal{C}^c, \\ \mathcal{C}^{2,4} &:= (\mathcal{A} + \mathcal{B}) \cap (\mathcal{B} + \mathcal{C}) \cap (\mathcal{C} + \mathcal{A}) \cap \mathcal{A}^c \cap \mathcal{B}^c \cap \mathcal{C}^c, \\ \mathcal{C}^{2,3,1} &:= (\mathcal{A} + \mathcal{B}) \cap (\mathcal{C} + \mathcal{A}) \cap \mathcal{A}^c \cap (\mathcal{B} + \mathcal{C})^c,\end{aligned}$$

together with the convex cones obtained by permuting systems. Here, $\mathcal{C}^{2,6,1}$, $\mathcal{C}^{2,4}$ and $\mathcal{C}^{2,3,1}$ are notations in [10]. The authors [11] gave examples of X-shaped states belonging to those classes. In this paper, we gave complete necessary and sufficient conditions for X-states to be members of the classes. For example, an X-state $\rho = X(a, b, z)$ belongs to the class $\mathcal{C}^{2,6,1}$ if and only if the following hold:

- ρ satisfies the inequalities $S_1[1, 4]$, $S_1[2, 3]$ and $S_2[1, 4]$;
- ρ violates $S_1[1, 3]$ or $S_1[2, 4]$;
- ρ violates $S_1[1, 2]$ or $S_1[3, 4]$.

The example $\rho = X((0, 1, 1, 2), (0, 1, 1, 2), (0, 1, 1, 0))$ given in [11] satisfies $S_1[1, 4]$, $S_1[2, 3]$ and $S_2[1, 4]$, but violates $S_1[1, 3]$ and $S_1[1, 2]$.

It is natural to ask what happens in the four qubit system, or arbitrary qubit systems. We began with the result [20] that an X-shaped multi-qubit state is separable with respect to a *bi-partition* of systems if and only if it is of positive partial transpose with respect to the same *bi-partition*. This was crucial to give characterizations in terms of diagonal entries and the *modulus* of anti-diagonal entries. But this is not the case for tri-partitions. In the three qubit system, considering tri-partition is amount to full separability. We need the phase parts, that is, the angular parts of anti-diagonal entries, as well as the modulus parts to characterize full separability of three qubit X-states. See [14, 15, 13]. We note that all kinds of partial separability come out from bi-partitions in the three qubit case. But, it is necessary to consider tri-partitions as well as bi-partitions in the four qubit case. See [9, 25]. Therefore, exploring partial separability/entanglement in general qubit system must be a very challenging project even for X-shaped states.

REFERENCES

- [1] C. H. Bennett, D. P. DiVincenzo, T. Mor, P. W. Shor, J. A. Smolin and B. M. Terhal, *Unextendible product bases and bound entanglement*, Phys. Rev. Lett. **82** (1999), 5385–5388.
- [2] A. Acin, D. Bruß, M. Lewenstein and A. Sanpera, *Classification of mixed three-qubit states*, Phys. Rev. Lett. **87** (2001), 040401.
- [3] V. Coffman, J. Kundu and W. K. Wootters, *Distributed entanglement*, Phys. Rev. A **61** (2000), 052306.

- [4] W. Dür and J. I. Cirac, *Classification of multi-qubit mixed states: separability and distillability properties*, Phys. Rev. A **61** (2000), 042314.
- [5] W. Dür, J. I. Cirac and R. Tarrach, *Separability and Distillability of Multiparticle Quantum Systems*, Phys. Rev. Lett. **83** (1999), 3562–3565.
- [6] W. Dür, G. Vidal and J. I. Cirac, *Three qubits can be entangled in two inequivalent ways*, Phys. Rev. A **62** (2000), 062314.
- [7] M. Seevinck and J. Uffink, *Partial separability and entanglement criteria for multiqubit quantum states*, Phys. Rev. A **78** (2008), 032101.
- [8] S. Szalay, *Separability criteria for mixed three-qubit states*, Phys. Rev. A **83** (2011), 062337.
- [9] S. Szalay, *Multipartite entanglement measures*, Phys. Rev. A **92** (2015), 042329.
- [10] S. Szalay and Z. Kökényesi, *Partial separability revisited: Necessary and sufficient criteria*, Phys. Rev. A **86** (2012), 032341.
- [11] K. H. Han and S.-H. Kye, *Construction of three-qubit biseparable states distinguishing kinds of entanglement in a partial separability classification*, Phys. Rev. A, **99** (2019), 032304.
- [12] K. H. Han and S.-H. Kye, *Separability of three qubit Greenberger-Horne-Zeilinger diagonal states*, J. Phys. A: Math. Theor. **50** (2017), 145303.
- [13] K. H. Han and S.-H. Kye, *The role of phases in detecting three qubit entanglement*, J. Math. Phys. **58** (2017), 102201.
- [14] L. Chen, K. H. Han and S.-H. Kye, *Separability criterion for three-qubit states with a four dimensional norm*, J. Phys. A: Math. Theor. **50** (2017) 345303.
- [15] K.-C. Ha, K. H. Han and S.-H. Kye, *Separability of multi-qubit states in terms of diagonal and anti-diagonal entries*, Quantum Inf. Process. **18** (2019), 34.
- [16] O. Gühne and M. Seevinck, *Separability criteria for genuine multiparticle entanglement*, New J. Phys. **12** (2010), 053002.
- [17] T. Gao and Y. Hong, *Separability criteria for several classes of n -partite quantum states*, Eur. Phys. J. D **61** (2011), 765–771.
- [18] S. M. H. Rafsanjani, M. Huber, C. J. Broadbent and J. H. Eberly, *Genuinely multipartite concurrence of N -qubit X matrices*, Phys. Rev. A **86** (2012), 062303.
- [19] K. H. Han and S.-H. Kye, *Various notions of positivity for bi-linear maps and applications to tri-partite entanglement*, J. Math. Phys. **57** (2016), 015205.
- [20] K. H. Han and S.-H. Kye, *Construction of multi-qubit optimal genuine entanglement witnesses*, J. Phys. A: Math. Theor. **49** (2016), 175303.
- [21] B. M. Terhal, *Bell Inequalities and the Separability Criterion*, Phys. Lett. A **271** (2000), 319–326.
- [22] M.-H. Eom and S.-H. Kye, *Duality for positive linear maps in matrix algebras*, Math. Scand. **86** (2000), 130–142.
- [23] R. T. Rockafellar, *Convex Analysis*, Princeton Univ. Press, 1970.
- [24] D. M. Greenberger, M. A. Horne and A. Zeilinger, *Going beyond Bell's theorem*, in Kafatos M. (eds) Bell's Theorem, Quantum Theory and Conceptions of the Universe. Fundamental Theories of Physics, Vol. 37, Springer, Dordrecht, 1989.
- [25] S. Szalay, *The classification of multipartite quantum correlation*, J. Phys. A: Math. Theor. **51** (2018), 485302 .

KYUNG HOON HAN, DEPARTMENT OF DATA SCIENCE, THE UNIVERSITY OF SUWON, GYEONGGI-DO 445-743, KOREA

E-mail address: kyunghoon.han at gmail.com

SEUNG-HYEOK KYE, DEPARTMENT OF MATHEMATICS AND INSTITUTE OF MATHEMATICS, SEOUL NATIONAL UNIVERSITY, SEOUL 151-742, KOREA

E-mail address: kye at snu.ac.kr