

A GENERALIZATION OF THE GRAM DETERMINANT OF TYPE A

RHEA PALAK BAKSHI, DIONNE IBARRA, SUJOY MUKHERJEE, AND JÓZEF H. PRZYTICKI

ABSTRACT. The Gram determinant of type A was introduced by Lickorish in his work on invariants of 3 - manifolds. We generalize the theory of the Gram determinant of type A by evaluating, in the annulus, a bilinear form of non-intersecting connections in the disc. The main result provides a closed formula for this Gram determinant. We conclude the paper by discussing Chen's conjecture about the Gram determinant of type Mb evaluated in the Klein bottle.

CONTENTS

1. Introduction	1
1.1. Relative Skein Modules	2
1.2. The Temperley - Lieb algebra and the Jones - Wenzl idempotent	2
1.3. The Gram Determinant of Type A	4
1.4. The Gram Determinant of Type B	5
2. The Gram Determinant of Generalized Type A	6
2.1. Theta Nets	7
2.2. A Change of Basis for the Temperley - Lieb Algebra	9
2.3. Proof of the Main Theorem	11
3. Future Directions: The Gram Determinant of Type Mb	12
Acknowledgments	16
References	16

1. INTRODUCTION

Modern work on Gram determinants in knot theory began with the construction of the Witten-Reshetikhin-Turaev invariants of 3-manifolds by Lickorish. [Wit, RT, Lic1]. Lickorish's Gram determinant (also called the Gram determinant of type A) was computed in the series of papers [KS], [Wes], and [DiF]. For our work, Cai's proof of the closed formula of the determinant using Jones - Wenzl idempotents is important [Cai].

In fact, in knot theory, matrices whose determinants are related to the Gram determinants, are prevalent. The earliest bilinear forms and their related Gram determinants (or their modifications) in knot theory are as follows:

- (1) the Seifert matrix of a link, L , denoted by S_L , and its modification which gives the Alexander matrix $tS_L - t^{-1}S_L^T$. Recall that the determinant of the Alexander matrix is the Alexander polynomial of the link L ,

Date: January 19, 2022.

2010 Mathematics Subject Classification. Primary: 57M27. Secondary: 57M25, 05A19.

Key words and phrases. Catalan numbers, Gram determinants, Klein bottle, knot theory, Möbius band, Temperley - Lieb algebra, Witten - Reshetikhin - Turaev invariants.

- (2) the determinant of the Goeritz matrix of a link which is called the determinant of the link. If the determinant is not zero, then it is the rank of the first homology group of the double branched cover of \mathbb{S}^3 branched along the link, and
- (3) the linking matrix of a framed link. If the determinant is not zero, then it is the rank of the first homology group of the 3 - manifold obtained from surgery along the link.

Rodica Simion introduced the Gram determinant of type B and its closed form was computed in [CP] and [MS].

1.1. Relative Skein Modules. Relative skein modules allow the theory of the Gram determinants to be developed for 3 - manifolds using any skein relation (for example, the HOMFLYPT skein relation). See [Prz1, Prz2]. In our work, we use the relative Kauffman bracket skein module to study Gram determinants.

Definition 1.1.

Let M be an oriented 3 - manifold and $\{x_1, x_2, \dots, x_{2n}\}$ be a set of $2n$ framed points on ∂M . Let $\mathcal{L}_{fr}(2n)$ be the set of all relative framed links in $(M, \partial M)$ considered up to ambient isotopy keeping ∂M fixed, such that $L \cap \partial M = \partial L = \{x_i\}$. Let R be a commutative ring with unity, A an invertible element in R , and $S_{2,\infty}(2n)$, the submodule of $R\mathcal{L}_{fr}(2n)$, generated by all the Kauffman bracket skein relations. Then, the relative Kauffman bracket skein module of M is the quotient:

$$S_{2,\infty}(M, \{x_i\}_1^{2n}; R, A) = \frac{R\mathcal{L}_{fr}(2n)}{S_{2,\infty}(2n)}.$$

Theorem 1.2. [Prz2]

Let F be a surface and $M = F \times I$ (or $F \hat{\times} I$, if F is unoriented), such that $\partial F \neq \emptyset$. Then, $S_{2,\infty}(M, \{x_i\}_1^{2n}; R, A)$ is a free R -module whose basis consists of relative links in F , without trivial components.

The following corollary discusses well known combinatorial notions using relative skein modules.

Corollary 1.3.


- (1) $S_{2,\infty}(D^2 \times I, \{x_i\}_1^{2n}; R, A)$ is a free R -module with $c_n = \frac{1}{n+1} \binom{2n}{n}$ basic elements. Here c_n denotes the n^{th} Catalan number.
- (2) $S_{2,\infty}(A^2 \times I, \{x_i\}_1^{2n}; R, A)$ is a free $R[z]$ - module with $\binom{2n}{n}$ basic elements, where z denotes the homotopically non - trivial curve on the annulus.

1.2. The Temperley - Lieb algebra and the Jones - Wenzl idempotent. The relative Kauffman bracket skein module of $D^2 \times I$ can be equipped with an algebra structure which leads to the classical Temperley - Lieb algebra (see [TL]).

Definition 1.4.

Let R be a commutative ring with unity and an invertible element A . Further, let $d = -A^{-2} - A^2 \in R$. The Temperley - Lieb Algebra, denoted by $TL_n(d)$, is an R - algebra generated by n elements, $\{1, e_1, e_2, \dots, e_{n-1}\}$ with relations:

- (1) $e_i^2 = de_i$ for $1 \leq i \leq n - 1$,
- (2) $e_i e_j e_i = e_i \ \forall |i - j| = 1$, and
- (3) $e_i e_j = e_j e_i \ \forall |i - j| > 1$.


 FIGURE 1. The generators of $TL_n(d)$.

Diagrammatically, the elements of the basis of $TL_n(d)$, as an R -module, can be represented as crossingless connections between the $2n$ marked points on the boundary of a rectangle with n points each on its top edge and bottom edge (see Figure 1). For further reading, see [Kau].

There is a natural bilinear form on the elements of $TL_n(d)$ which Lickorish introduced to construct the Witten - Reshetikhin - Turaev invariants of 3 - manifolds. The determinant of this bilinear form is called the Gram Determinant of type A which we will discuss in the following subsection.

Definition 1.5. [Jon1, Wen]

Let $p : \mathbb{B}_n \longrightarrow S_n$ be a map, where \mathbb{B}_n denotes the Artin braid group and S_n , the permutation group. This function sends a braid word to the induced permutation. Define $F_n = \sum_{\pi \in S_n} (A^3)^{|\pi|} b_\pi \in \mathbb{Z}[A^{\pm 1}]\mathbb{B}_n$, where $|\pi|$ is the length of the permutation π and b_π is the unique, minimal, positive braid such that $p(b_\pi) = \pi$. The n^{th} Jones - Wenzl idempotent, denoted by f_n , is defined to be $\frac{F_n}{(\{n\}_q)!} \in \mathbb{Q}(A)\mathbb{B}_n$, where $\{n\}_q = 1 + q + q^2 + \cdots + q^{n-1}$, and $(\{n\}_q)! = \{1\}_q \cdot \{2\}_q \cdots \{n\}_q$.¹

The Jones - Wenzl idempotent is evaluated in $TL_n(d)$ by taking the quotient of $\mathbb{Q}(A)\mathbb{B}_n$ by the Kauffman bracket skein relations.


Theorem 1.6. [Jon1, KL, Lic2]

The n^{th} Jones - Wenzl idempotent, $f_n \in TL_n(d)$, satisfies the following properties:

- (1) $f_n e_i = 0 = e_i f_n$, for $1 \leq i \leq n-1$,
- (2) $(f_n - \mathbb{1})$ belongs to the algebra generated by $\{e_1, e_2, \dots, e_{n-1}\}$, and
- (3) $f_n f_n = f_n$.


The n^{th} Jones-Wenzl idempotent is represented by a rectangular box (see Figure 2) with n curves entering and n curves exiting it. The integer n beside a curve represents n parallel copies of it.

¹The extension of $\mathbb{Z}[A^{\pm 1}]$ to the field of rational functions, $\mathbb{Q}(A)$, does not change the Gram determinant.


 FIGURE 2. The Jones - Wenzl idempotent f_n .

Theorem 1.7. [Wen]

The Jones - Wenzl idempotent f_{n+1} , for $n > 1$, satisfies recursive relation shown in Figure 3.


 FIGURE 3. The Wenzl formula for f_{n+1} .

1.3. The Gram Determinant of Type A.


Definition 1.8.

Consider the disc D^2 with $2n$ framed points on its boundary. Let $\mathcal{A}_n = \{a_1, a_2, \dots, a_{c_n}\}$ be the set of all diagrams with crossingless connections, up to ambient isotopy, between the $2n$ framed points in D^2 . Define a bilinear form $\langle \cdot, \cdot \rangle$ in the following way:

$$\langle \cdot, \cdot \rangle : \mathcal{S}_{2,\infty}(D^2 \times I, \{x_i\}_1^{2n}) \times \mathcal{S}_{2,\infty}(D^2 \times I, \{x_i\}_1^{2n}) \longrightarrow R.$$

Let $a_i, a_j \in \mathcal{A}_n$. Glue a_i with the inversion of a_j along the marked circle, respecting the labels of the framed points. The resulting picture is that of a disc with disjoint null homotopic circles. Thus, we define, $\langle a_i, a_j \rangle = d^m$ where m denotes the number of these circles. Example 1.9 illustrates an example of the bilinear form when $n = 4$.

The Gram matrix of type A is defined as $G_n^A = (\langle a_i, a_j \rangle)_{1 \leq i, j \leq c_n}$. Its determinant D_n^A is called the Gram determinant of type A.

Example 1.9.

Theorem 1.10. [Wes, DiF, Cai]

Let $R = \mathbb{Z}[A^{\pm 1}]$. Then,

$$D_n^A(d) = \Delta_1^{c_n} \prod_{i=1}^n \left(\frac{\Delta_i}{\Delta_{i-1}} \right)^{\alpha_i}, \text{ where } \Delta_i = (-1)^i \frac{A^{2i+2} - A^{-2i-2}}{A^2 - A^{-2}} \text{ and } \alpha_i = \binom{2n}{n-i} - \binom{2n}{n-i-1}.$$

Note that $\Delta_1 = -A^2 - A^{-2} = d$. Furthermore, $\Delta_i(d)$ is the Chebyshev polynomial of the second kind (see Theorem 2.3). In [Cai], the author used Jones - Wenzl idempotents to construct a new basis of $TL_n(d)$ and provided a new proof of the formula for the Gram determinant of type A using it.

1.4. The Gram Determinant of Type B. The Gram determinant of type B was first considered by Rodica Simion while working with matrices of chromatic joins [Sim, Sch]. Jones - Wenzl idempotents and Chebyshev polynomials of the first kind were used to find the closed formula of the Gram determinant of type B in [CP].

Definition 1.11.

Let A^2 be an annulus with $2n$ marked points on its outer boundary. Let $\mathcal{B}_n = \{b_1, b_2, \dots, b_{\binom{2n}{n}}\}$ be the set of all diagrams of crossingless connections between these $2n$ points. Define a bilinear form \langle , \rangle in the following way:

$$\langle , \rangle : \mathcal{S}_{2,\infty}(A^2 \times I, \{x_i\}_1^{2n}; R, A) \times \mathcal{S}_{2,\infty}(A^2 \times I, \{x_i\}_1^{2n}; R, A) \longrightarrow R[z].$$

Given $b_i, b_j \in \mathcal{B}_n$, glue b_i with the inversion of b_j along the marked circle, respecting the labels of the marked points. The resulting picture has disjoint circles which are either homotopically non-trivial or null homotopic. Then, $\langle b_i, b_j \rangle = z^k d^m$, where k and m denote the number of these circles, respectively.

The Gram matrix of type B is defined as $G_n^B = (\langle b_i, b_j \rangle)_{1 \leq i, j \leq \binom{2n}{n}}$. Its determinant D_n^B is called the Gram determinant of type B.

The following theorem, evaluating this determinant, answers Rodica Simion's question.

Theorem 1.12. [CP, MS]

Let $R = \mathbb{Z}[A^{\pm 1}]$. Then, $D_n^B(d, z) = \prod_{i=1}^n (T_i(d)^2 - z^2)^{\binom{2n}{n-i}}$, where T_i denotes the Chebyshev polynomial of the first kind defined recursively by the equation $T_{n+1}(d) = d \cdot T_n(d) - T_{n-1}(d)$, with the initial conditions $T_0(d) = 2$ and $T_1(d) = d$.

The idea of the proof is as follows. For a positive integer n , there are $\binom{2n}{n}$ annular Catalan states. The evaluation of the Gram matrix of type B at $z = (-1)^{k-1} T_k(d)$ is closely related to the evaluation of the Gram matrix of type A. A Hopf link is formed with one of its components, the

annulus used in the Gram determinant of type B , and the other decorated with the k^{th} Jones - Wenzl idempotent. For details, see [CP]. With this decoration, the number of annular Catalan states is equal to $\binom{2n}{n} - \binom{2n}{n-k}$ and thus, the nullity of the matrix is at least $\binom{2n}{n-k}$. Therefore, $D_n^B(d, z)$ is divisible by $(T_k(d) + (-1)^k z)^{\binom{2n}{n-k}}$. By a simple analysis of involution, which changes z to $-z$, it follows that $D_n^B(d, z)$ is also divisible by $(T_k(d) - (-1)^k z)^{\binom{2n}{n-k}}$. Thus, the theorem holds up to a multiplicative factor. By analysing the term having the highest degree which is obtained from the diagonal of the matrix, it follows that this factor is equal to 1.




FIGURE 4. Depiction of the bilinear form $\langle a_i, a_j \rangle = \text{Tr}(a_i \bar{a}_j)$ of generalized type A, where $a_i \in \mathcal{A}_n$ and \bar{a}_j is the reflection of $a_j \in \mathcal{A}_n$ about the horizontal axis.

2. THE GRAM DETERMINANT OF GENERALIZED TYPE A

We now introduce a generalized version of the Gram determinant of type A.

Definition 2.1.

Let the disc D^2 , with $2n$ marked points on its boundary, be considered as a rectangle with n points on the top edge and n points on the bottom edge. Define a bilinear form $\langle \cdot, \cdot \rangle$ in the following way:

$$\langle \cdot, \cdot \rangle : \mathcal{S}_{2,\infty}(D^2 \times I, \{x_i\}_1^{2n}) \times \mathcal{S}_{2,\infty}(D^2 \times I, \{x_i\}_1^{2n}) \longrightarrow \mathbb{Z}[d, z].$$

For $a_i, a_j \in \mathcal{A}_n$, glue a_i with the reflection, about the horizontal axis, of a_j which is denoted by \bar{a}_j , such that the bottom edge of a_i is identified with the top edge of \bar{a}_j . Connect the marked points on the top edge of a_i with those on the bottom edge of \bar{a}_j , in the annulus, respecting the ordering of the

marked points (see Figure 4). Recall that the described operation is the same as taking the trace of the image of the bilinear form.

The result is an annulus with two types of disjoint circles, homotopically trivial and non-trivial. Thus, we define, $\langle a_i, a_j \rangle = d^k z^m$ where k and m denote the number of these circles respectively. We define the Gram matrix of generalized type A as $G_n^{A^{gen}} = (\langle a_i, a_j \rangle)_{1 \leq i, j \leq c_n}$, and denote its determinant by $D_n^{A^{gen}}$.

Remark 2.2.

The Temperley - Lieb algebra leads to the Frobenius algebra by defining the Frobenius form to be the trace of the elements of the Temperley - Lieb algebra (see [Koc]).

Theorem 2.3.

Let $R = \mathbb{Z}[A^{\pm 1}]$. Then, the Gram determinant of generalized type A is given by the following formula:

$$D_n^{A^{gen}}(d, z) = D_n^A(d) \prod_{i=0}^{\lfloor \frac{n}{2} \rfloor} \left(\frac{S_{n-2i}(z)}{\Delta_{n-2i}(d)} \right) \left(\binom{n}{i} - \binom{n}{i-1} \right)^2,$$


where $S_k(z)$ denotes the Chebyshev polynomial of the second kind, defined recursively by the equation $S_{k+1}(z) = z \cdot S_k(z) - S_{k-1}(z)$, with the initial conditions $S_0(z) = 1$ and $S_1(z) = z$.

Before giving the proof of this theorem in Section 2.3, we give an outline here. To compute the Gram determinant $D_n^{A^{gen}}$, we change the basis of the Temperley - Lieb algebra, $TL_n(d)$, so that in the new basis, the Gram matrix is a diagonal matrix. It follows from [Cai] that the change of basis is given by an upper triangular matrix with 1's on the diagonal. Therefore, the Gram determinant is unchanged by the change of basis. To find the diagonal entries of the new Gram matrix, we use the standard properties of theta nets. The difference with the Gram determinant of generalized type A is that we finally evaluate the theta nets in the annulus and not in the disc. Hence, we use the fact that the trace of the Jones - Wenzl idempotents, in the annulus, is the Chebyshev polynomial of the second kind in the variable z . Finally, we use the combinatorial properties of mountain paths to find a closed formula for the Gram determinant of generalized type A.

2.1. Theta Nets. This subsection serves as a preparation for the proof of the main theorem.

Definition 2.4.

Let $a = y + z$, $b = x + z$, and $c = x + y$, that is, $x = \frac{(b+c-a)}{2}$, $y = \frac{(a+c-b)}{2}$, and $z = \frac{(a+b-c)}{2}$, so that



Notice that $(a + b - c)$, $(b + c - a)$, and $(a + c - b)$ are all even integers. The Temperley - Lieb category TL consists of objects $n \in \mathbb{N}$, morphism spaces $TL_{m,n}$, and multiplication $TL_{m,n} \times TL_{l,m} \longrightarrow TL_{l,n}$ (see, for example, [QW]). Then, the following figure depicts the Markov trace of the multiplication of $TL_{c,a+b} \times TL_{a+b,c} \longrightarrow TL_{a+b,a+b}$.²

²Recall that trace in the disk is called the Markov trace.

The diagram shows the trace of two trivalent vertices. On the left, two vertices are shown, each with three legs labeled a , b , and c . They are connected by their c legs. This is equated to a bubble diagram on the right, which consists of two vertices connected by three arcs labeled c , a , and b .

Therefore, we obtain the following picture and can compute its unreduced Kauffman bracket using Theorem 2.5.

The diagram shows the equation $\Theta(a, b, c) = \Gamma_{\mathbb{R}^2}(x, y, z)$. On the right, there is a diagram with three loops labeled x , y , and z . Each loop has a small square box on it, labeled c , b , and a respectively.

Theorem 2.5 ([MV, KL, Lic2]).

Let $\Delta_n!$ denote the product $\Delta_n \Delta_{n-1} \cdots \Delta_1$. Then

$$\Gamma_{\mathbb{R}^2}(x, y, z) = \frac{\Delta_{x+y+z}! \Delta_{x-1}! \Delta_{y-1}! \Delta_{z-1}!}{\Delta_{x+y-1}! \Delta_{x+z-1}! \Delta_{y+z-1}!}.$$

In particular, for $b = x + z = 1$, that is, $|a - c| = 1$, we get $\Gamma_{\mathbb{R}^2}(x, y, z) = \Delta_{y+1}$.

Corollary 2.6.


The following formula reduces digons in the net (called bubble reduction),

The diagram shows the bubble reduction formula. On the left, a digon (a loop with two vertices) is shown with legs labeled a' , b , and a . This is equated to a fraction $\frac{\delta_{a,a'} \Theta(a, b, c)}{\Delta_a}$ multiplied by a square box labeled a .

where $\delta_{a,a'}$ is the Kronecker delta. Compare with [KL, Lic2].

Proof.

If $a \neq a'$ then without loss of generality we can assume that $a' > a$. Therefore, the corresponding diagram has a return on the part with a' in it. So the value of a associated to the diagram on the left is equal to 0. Otherwise $a = a'$, and by the property of Jones - Wenzl idempotents, we get



After closing the diagram in \mathbb{R}^2 , the left side is a decorated theta curve with evaluation $\Theta(a, b, c)$, while the right side gives us $\lambda\Delta_a$. Thus, $\lambda = \frac{\Theta(a,b,c)}{\Delta_a}$. □

Corollary 2.7.


The formula for the decorated theta curve in \mathbb{R}^2 can be naturally generalized to the decorated theta curve in the annulus. That is,

$$\Gamma_{Ann}(x, y, z) = \frac{S_{x+y}(x)}{\Delta_{x+y}} \Gamma_{\mathbb{R}^2}(x, y, z),$$

where we start from a pair of pants with holes ∂_x, ∂_y and ∂_z . Then, we cap off ∂_z , so that, ∂_x is homotopic to ∂_y .

Proof.

Using Corollary 2.6, we get



which in the Kauffman bracket skein module, $S_{2,\infty}(A^2 \times I)$, is equal to the Chebyshev polynomial $S_{x+y}(\alpha) \frac{\Theta(a,b,c)}{\Delta_{x+y}}$, where $a = y + z$, $b = x + z$, and $c = x + y$. □

2.2. A Change of Basis for the Temperley - Lieb Algebra. Following [Cai], we make a change of basis so that our bilinear form is orthogonal. Hence, in the new basis, the Gram matrix is diagonal.

Definition 2.8.

Consider the finite sequence $\{a_1, a_2, \dots, a_{2n-1}\}$ of natural numbers which satisfies the following two conditions:

- (1) $a_1 = a_{2n-1} = 1$ and
- (2) $|a_i - a_{i-1}| = 1 \ \forall i$.

The elements of $TL_n(d)$, denoted by $D_{a_1, a_2, \dots, a_{2n-1}}$, are depicted in Figure 5. The triple points in the diagram are admissible.





FIGURE 5. The depiction of $D_{a_1, a_2, \dots, a_{2n-1}} \in TL_n(d)$ using a_1, \dots, a_{2n-1} (Definition 2.8).

Example 2.9.

Pictured below is the basis for $TL_3(d)$, decorated with Jones - Wenzl idempotents.



In order to compute the Gram determinant of generalized type A, we will need the following results.

Theorem 2.10. [Cai]

Let \mathcal{D}_n be the collection of all elements $D_{a_1, a_2, \dots, a_{2n-1}}$. Then, \mathcal{D}_n is a basis for $TL_n(d)$. The change of basis matrix between \mathcal{A}_n and \mathcal{D}_n is an upper triangular matrix with 1's on the diagonal. In particular, the matrix has determinant 1.

Theorem 2.11.

The elements $D_{a_1, a_2, \dots, a_{2n-1}}$ are orthogonal with respect to the bilinear form of generalized type A.

Proof.

We adapt the proof for the Gram determinant of type A [Cai] to the Gram determinant of generalized type A. In particular, we use Corollary 2.6 for $a \neq a'$ to obtain orthogonality. \square

Corollary 2.12.

The Gram determinant of generalized type A is unchanged under the change of basis from \mathcal{A}_n to \mathcal{D}_n .

Proof.

The proof follows from Theorem 2.10. \square

2.3. Proof of the Main Theorem. By Proposition 2.11 and Corollary 2.12, it suffices to find the diagonal entries of the Gram matrix of generalized type A in the basis \mathcal{D}_n . We follow the proof in [Cai] in the evaluation using Corollaries 2.6 and 2.7.

Since we work in the annulus, we need to modify the part of Cai's proof which takes into account the terms appearing after bubble reduction (using Corollary 2.6). The distinction in our case is that after bubble reduction, we obtain $S_{a_n}(z)$, while in Cai's proof the bubble reduction gives $S_{a_n}(d)$.




FIGURE 6. Bubble reduction.


Recall that a_n is the middle term of the sequence in Definition 2.8. This term will give the factor $S_{a_n}(z)$ in the Gram determinant of generalized type A. Therefore, to see the difference between the Gram determinant of type A and the Gram determinant of generalized type A, for fixed i , we have to count the number of states with the term which has the value $a_n = n - 2i$, for $0 \leq i \leq \frac{n}{2}$. We find the exponents of $S_{n-2i}(z)$ using the Desiré - André reflection principle. We show that this number is equal to $\left(\binom{n}{i} - \binom{n}{i-1}\right)^2$. This follows from the next lemma.

Lemma 2.13.

Consider all mountain paths from $(0, 0)$ to $(n, n - 2i)$ that lie completely in the first quadrant of the xy - plane. The number of such paths is $\binom{n}{i} - \binom{n}{i-1}$.

Proof.

Consider all mountain paths from $(0, 0)$ to $(n, n - 2i)$. These are paths whose piecewise smooth components have slope ± 1 . See Figure 7 for an example. There are $\binom{n}{i}$ such paths. Note that


 FIGURE 7. Mountain paths and their reflection about the horizontal axis $y = -1$.

these paths can have portions that lie below the x - axis. We now find the number of mountain paths that have portions below this axis ('bad' paths). Travel along the path from $(0, 0)$ till you reach the point that has the y coordinate equal to -1 for the first time. Reflect the path till this point about the horizontal line $y = -1$, keeping the remaining portion of the path untouched. We obtain the path from $(0, -2)$ till $(n, n - 2i)$. See Figure 7. This construction gives a bijection from all such bad paths to all paths from $(0, -2)$ to $(n, n - 2i)$. The number of these paths is $\binom{n}{i-1}$. Therefore, the number of paths lying completely above the x - axis is $\binom{n}{i} - \binom{n}{i-1}$. \square




FIGURE 8. The five different simple closed curves on the Klein bottle.

3. FUTURE DIRECTIONS: THE GRAM DETERMINANT OF TYPE MB

The fourth author proposed the notion of the Gram determinant of type Mb evaluated in the Klein bottle [Prz3], that is, the study of the Gram determinant based on crossingless connections on a Möbius band. In this case, the bilinear form is defined by identifying two Möbius bands along their boundaries (see Definition 3.2). The identification produces a Klein bottle which has five homotopically distinct curves and thus the Gram matrix given by this bilinear form on the

elements of $S_{2,\infty}(Mb \hat{\times} I, \{x_i\}_1^{2n})$ has entries in the five variables that denote these curves. After some preliminary calculations, Chen proposed the following conjecture [Che] for the Gram determinant of the Möbius band.

Conjecture 3.1 (Chen).

Let $R = \mathbb{Z}[A^{\pm 1}]$. Then, the Gram determinant of type Mb for $n \geq 1$, denoted $D_n^{(Mb)}$, is

$$\begin{aligned} D_n^{(Mb)}(d, w, x, y, z) = & \prod_{k=1}^n (T_k(d) + (-1)^k z)^{\binom{2n}{n-k}} \\ & \prod_{\substack{k=1 \\ k \text{ odd}}}^n (T_k(d) - (-1)^k z) T_k(w) - 2xy)^{\binom{2n}{n-k}} \\ & \prod_{\substack{k=1 \\ k \text{ even}}}^n (T_k(d) - (-1)^k z) T_k(w) - 2(2-z))^{\binom{2n}{n-k}} \\ & \prod_{i=1}^n D_{n,i}, \end{aligned}$$

where $D_{n,i} = \prod_{k=1+i}^n (T_{2k}(d) - 2)^{\binom{2n}{n-k}}$, and i represents the number of curves passing through the crosscap.

In the remaining part of this paper, we discuss some of our work on this project and state a few results supporting Conjecture 3.1 (Proposition 3.4, Theorem 3.5, and Theorem 3.6). Compare Theorem 3.5 and Theorem 3.6 with analogous cases in [CP] and [PZ], respectively.

Definition 3.2.

Consider a Möbius band with $2n$ marked points on its boundary, and n crossingless curves connecting these $2n$ marked points. Denote the set of all diagrams of crossingless connections on the Möbius band with $2n$ points by Mb_n . Then the number of distinct crossingless connections is,

$$|Mb_n| = \sum_{k=0}^n \binom{2n}{k} = 2^{2n-1} + \frac{1}{2} \binom{2n}{n}.$$

Notice that in the formula $\sum_{k=0}^n \binom{2n}{k}$, k tells us how many curves are disjoint from the crosscap (see Figure 9).

Define a bilinear form $\langle \cdot, \cdot \rangle$ in the following way:

$$\langle \cdot, \cdot \rangle : S_{2,\infty}(Mb \hat{\times} I, \{x_i\}_1^{2n}) \times S_{2,\infty}(Mb \hat{\times} I, \{x_i\}_1^{2n}) \longrightarrow \mathbb{Z}[d, w, x, y, z],$$

where d, w, x, y , and z denote the five homotopically distinct simple closed curves on the Klein bottle which are illustrated in Figure 8.³

Given $m_i, m_j \in Mb_n$, identify the boundary component of m_i with that of the inversion of m_j , respecting the labels of the marked points. The result is a collection of disjoint simple closed curves on the Klein bottle which are of five different types, as shown in Figure 8. In particular, given two

³The variable d denotes the homotopically trivial curve, while x intersects the inner crosscap, y intersects the outer crosscap, z separates the two crosscaps, and w intersects both crosscaps.

elements in Mb_n , the image of its bilinear form is a monomial in $\mathbb{Z}[d, x, y, z, w]$, that is, $\langle m_i, m_j \rangle = d^m x^n y^k z^l w^h$ where m, n, k, l and h denote the number of these simple closed curves, respectively.

The Gram matrix of type Mb is defined as $G_n^{Mb} = (\langle m_i, m_j \rangle)_{1 \leq i, j \leq |Mb_n|}$ and its determinant is denoted by D_n^{Mb} . Since the Gram matrix is given by the bilinear form, the size of the matrix, $|G_n^{Mb}|$, is equal to the number of different crossingless connections.

Example 3.3.

For $n = 2$, there are eleven different crossingless connections in the Möbius band.




FIGURE 9. In the eleven different crossingless connections Möbius band, the first six, $\binom{4}{2}$, do not intersect the crosscap, the next four, $\binom{4}{1}$, pass through the crosscap once, and finally, the last one, $\binom{4}{0}$, passes through the crosscap twice.

The Gram matrix given by the bilinear form is as follows:

$\langle \ , \ \rangle$												
	d^2	dz	z^2	z	d	z	dy	y	yz	y	z	
	dz	d^2	dz	d	z	d	dy	y	dy	y	d	
	z^2	dz	d^2	z	d	z	yz	y	dy	y	z	
	z	d	z	d^2	dz	z^2	y	yz	y	dy	z	
	d	z	d	dz	d^2	dz	y	dy	y	dy	d	
	z	d	z	z^2	dz	d^2	y	dy	y	yz	z	
	dx	dx	xz	x	x	x	dw	w	xy	w	x	
	x	x	x	xz	dx	dx	w	dw	w	xy	x	
	xz	dx	dx	x	x	x	xy	w	dw	w	x	
	x	x	x	dx	dx	xz	w	xy	w	dw	x	
	z	d	z	z	d	z	y	y	y	y	w^2	

The determinant of the Gram matrix $\text{Det}(G_2^{Mb})$ is equal to :

$$(d - z)^4 [(d + z)w - 2xy]^4 (d^2(d^2 - 4))(d^2 - 2 + z)[(d^2 - 2 - z)(w^2 - 2) - 2(2 - z)].$$

In terms of Chebyshev polynomials of the first kind, the formula coincides with Chen's conjecture and $\text{Det}(G_2^{Mb})$ is equal to:

$$(T_1(d) - z)^4 [(T_1(d) + z)T_1(w) - 2xy]^4 (T_4(d) - 2)(T_2(d) + z)[(T_2(d) - z)T_2(w) - 2(2 - z)]$$




FIGURE 10. The crossingless connection $b_3 \in Mb_n$ with the connection α_3 in b_3 passing through the crosscap and the crossingless connections $b_1, b_2 \in Mb_n$ obtained from b_3 by modifying α_3 so that it does not pass through the crosscap.

Proposition 3.4.

$\text{Det}(G_n^{Kb})$ is divisible by $(w(d + z) - 2xy)^{\binom{2n}{n-1}}$.

Proof.

Consider a crossingless connection, that is, an element $b_3 \in Mb_n$, which cuts the crosscap exactly once (denoted by α_3). Let b_1 and b_2 be obtained from b_3 by modifying α_3 so that it does not pass through the crosscap (see Figure 10). Consider the block associated with these three elements. For some monomial u , we get:

$$\begin{bmatrix} ud & uz & uy \\ uz & ud & uy \\ ux & ux & uw \end{bmatrix}$$

If we replace b_3 by $(d + z)b_3 - y(b_1 + b_2)$, denoted by b'_3 , we obtain the matrix:

$$\begin{bmatrix} ud & uz & 0 \\ uz & ud & 0 \\ ux & ux & u((d + z)w - 2xy) \end{bmatrix}$$

with determinant $(d + z)$ times the determinant of the previous matrix. Now, we observe that for any element $c \in Mb_n$ the row $\langle c, b_1 \rangle, \langle c, b_2 \rangle, \langle c, b_3 \rangle$ is proportional (by a monomial) to (d, z, y) , (z, d, y) , or (x, x, y) . Thus, the column operation we did earlier produces either a zero or a monomial multiple of $((d + z)w - 2xy)$. Therefore, the column under b'_3 is divisible by $((d + z)w - 2xy)$. Now, we can perform the same operation, as on b_3 , on any element (crossingless connection) which cuts the crosscap exactly once. There are $\binom{2n}{n-1}$ such elements and thus we obtain a matrix with $\binom{2n}{n-1}$ columns divisible by $((d + z)w - 2xy)$. Additionally, note that $d + z$ and $((d + z)w - 2xy)$ are co-prime. Therefore, $\text{Det}(G_n^{Kb})$ is divisible by $(w(d + z) - 2xy)^{\binom{2n}{n-1}}$. \square

Theorem 3.5.

$$\prod_{k=1}^n (T_k(d) + (-1)^k z)^{\binom{2n}{n-k}} \text{ divides } D_n^{(Mb)}.$$

Theorem 3.6.

$(D_{n-1}^{(Mb)})^2$ divides $D_n^{(Mb)}$, if $(d^2 - 1)$ is invertible.

ACKNOWLEDGMENTS

The third author was partially supported by the CCAS Dean's Dissertation Completion Fellowship. The fourth author was partially supported by the Simons Foundation Collaboration Grant for Mathematicians - 316446 and the CCAS Dean's Research Chair award.

REFERENCES

- [Cai] X. Cai, A Gram determinant of Lickorish's bilinear form. *Math. Proc. Cambridge Philos. Soc.*, 151(1), 2011, 83-94. [arXiv:1006.1297](#) [math.GT].
- [Che] Q. Chen, Personal communication (email) with J. H. Przytycki, April 3, 2009.
- [CP] Q. Chen, J. H. Przytycki, The Gram determinant of the type B Temperley-Lieb algebra, *Adv. in Appl. Math.*, 43(2), 2009, 156-161. [arXiv:0802.1083v2](#) [math.GT].
- [DiF] P. Di Francesco, Meander determinants. *Comm. Math. Phys.*, 191, 1998, 543-583. [arXiv:hep-th/9612026v1](#).
- [Jon1] V. F. R. Jones, Index for subfactors. *Invent. Math.* 72, 1983, 1-25.
- [Jon2] V. F. R. Jones, An affine Hecke algebra quotient in the Brauer Algebra. *l'Enseignement Mathématique* 40, 1994, 313-344.
- [Kau] L. H. Kauffman. Knot diagrammatics. *Handbook of Knot Theory*, W. W. Menasco and M. Thistlethwaite (editors), Elsevier, 2005. [arXiv:math/0410329v5](#) [math.GN].
- [KL] L. H. Kauffman, S. L. Lins, Temperley - Lieb recoupling theory and invariants of 3-manifolds. *Annals of Mathematics Studies*, Number 134, Princeton University Press, 1994.
- [KS] K. H. Ko, L. Smolinsky, A combinatorial matrix in 3-manifold theory. *Pacific J. Math.*, 149(2), 1991, 319-336.
- [Koc] J. Kock, Frobenius algebras and 2D topological quantum field theories. *London Mathematical Society Student Texts*, 59. Cambridge University Press, Cambridge, 2004, xiv+240.
- [Lic1] W. B. R. Lickorish, Invariants for 3-manifolds from the combinatorics of the Jones polynomial. *Pacific Journ. Math.*, 149(2), 1991, 337-347.
- [Lic2] W. B. R. Lickorish, An introduction to Knot Theory. Graduate Texts in Mathematics, 175, Springer 1997.
- [MS] P. Martin, H. Saleur, On an algebraic approach to higher dimensional statistical mechanics. *Commun. Math. Phys.*, 158, 1993, 155-190.
- [MV] G. Masbaum, P. Vogel, 3-valent graphs and the Kauffman bracket. *Pacific J. Math.*, 164(2), 1994, 361-381.
- [Prz1] J. H. Przytycki, Skein modules of 3-manifolds. *Bull. Polish Acad. Science*, 39(1-2), 1991, 91-100. [arXiv:math/0611797v1](#) [math.GT].
- [Prz2] J. H. Przytycki, Fundamentals of Kauffman bracket skein modules. *Kobe Math. J.*, 16(1), 1999, 45-66. [arXiv:math/9809113](#) [math.GT].
- [Prz3] J. H. Przytycki, Personal communication (email) with Q. Chen, November 21, 2008.
- [PZ] J. H. Przytycki, X. Zhu, Gram determinant of planar curves. *Involve*, Vol. 3, No. 2, 2010, 149-169. [arXiv:0810.4649v1](#) [math.GT].
- [QW] H. Queffelec, P. Wedrich, Extremal weight projectors. *Mathematical Research Letters*, Volume 25 (2018), No. 6, 1911-1936. [arXiv:1803.09883v1](#) [math.RT].
- [RT] N. Y. Reshetikhin, V. Turaev, Invariants of three manifolds via link polynomials and quantum groups. *Invent. Math.* 103, 1991, 547-597.
- [Sch] F. Schmidt, Problems related to type-A and type-B matrices of chromatic joins. Special issue on the Tutte polynomial, *Adv. in Appl. Math.* 32 (2004), no. 1-2, 380-390.
- [Sim] R. Simion, A type-B associahedron. Formal power series and algebraic combinatorics (Scottsdale, AZ, 2001). *Adv. in Appl. Math.* 30 (2003), no. 1-2, 2-25.
- [TL] H. N. V. Temperley, E. H. Lieb, Relations between the percolation and colouring problem and other graph-theoretical problems associated with regular planar lattices: some exact results for the percolation problem. *Proc. Roy. Soc. London Ser. A*, 322 (1549) : 251280, 1971.
- [Wen] H. Wenzl, On sequences of projections. *C. R. Math. Rep. Acad. Sci. Canada*, 9(1) : 5-9, 1987.
- [Wes] B. W. Westbury, The representation theory of the Temperley-Lieb algebras. *Math. Z.*, 219(4), 1995, 539-565.

- [Wit] E. Witten, Quantum field theory and the Jones polynomial. Braid group, knot theory and statistical mechanics, II, 361-451, *Adv. Ser. Math. Phys.*, 17, *World Sci. Publ.*, River Edge, NJ, 1994.

DEPARTMENT OF MATHEMATICS, THE GEORGE WASHINGTON UNIVERSITY, WASHINGTON DC, USA
E-mail address: rhea.palak@gwu.edu

DEPARTMENT OF MATHEMATICS, THE GEORGE WASHINGTON UNIVERSITY, WASHINGTON DC, USA
E-mail address: dfkunkel@gwu.edu

DEPARTMENT OF MATHEMATICS, THE GEORGE WASHINGTON UNIVERSITY, WASHINGTON DC, USA
E-mail address: sujoymukherjee@gwu.edu

DEPARTMENT OF MATHEMATICS, THE GEORGE WASHINGTON UNIVERSITY, WASHINGTON DC, USA AND
DEPARTMENT OF MATHEMATICS, UNIVERSITY OF GDAŃSK, POLAND
E-mail address: przytyck@gwu.edu