

INTERPOLATION OF QUASI NONCOMMUTATIVE L_p -SPACES

JUAN GU, ZHI YIN, AND HAONAN ZHANG

ABSTRACT. Let \mathcal{M} be a $(\sigma$ -finite) von Neumann algebra associated with a normal faithful state ϕ . We prove a complex interpolation result for a couple of two (quasi) Haagerup noncommutative L_p -spaces $L_{p_0}(\mathcal{M}, \phi)$ and $L_{p_1}(\mathcal{M}, \phi)$, $0 < p_0 < p_1 \leq \infty$, which has further applications to the sandwiched p -Rényi divergence.

1. INTRODUCTION

Let \mathcal{M} be a semifinite von Neumann algebra equipped with a normal semifinite faithful (n.s.f.) trace τ , and $\Lambda_p(\mathcal{M}, \tau)$ be the tracial noncommutative L_p -space associated with (\mathcal{M}, τ) . It is well-known [31] that the complex interpolation space of a couple of two Banach spaces $(\Lambda_{p_0}(\mathcal{M}, \tau), \Lambda_{p_1}(\mathcal{M}, \tau))$ is isometric to $\Lambda_{p_\theta}(\mathcal{M}, \tau)$, i.e.,

$$(1.1) \quad [\Lambda_{p_0}(\mathcal{M}, \tau), \Lambda_{p_1}(\mathcal{M}, \tau)]_\theta = \Lambda_{p_\theta}(\mathcal{M}, \tau),$$

where $1/p_\theta = (1 - \theta)/p_0 + \theta/p_1$, $1 \leq p_0 < p_1 \leq \infty$. Through this line of research, the complex interpolation for weighted noncommutative L_p -spaces [32], noncommutative Hardy spaces [29, 8] and noncommutative Orlicz spaces [6, 7] have been studied. Because noncommutative L_p -spaces naturally appear in the study of noncommutative analysis, the interpolation of noncommutative L_p -spaces, together with the Riesz-Thorin theorem and Marcinkiewicz theorem, become fundamental and powerful tools in proving noncommutative martingale inequalities [17, 30], maximal ergodic inequalities [18], and the duality properties of noncommutative function spaces [24].

In the early 1990s, Xu [37] showed that the above interpolation result (1.1) holds for the couple of quasi-Banach spaces $(\Lambda_{p_0}(\mathcal{M}, \tau), \Lambda_{p_1}(\mathcal{M}, \tau))$, $0 < p_0 < p_1 \leq \infty$. However, it didn't draw too much attention to the community, contrary to the Banach space case ($1 \leq p_0 < p_1 \leq \infty$). In this paper, we will extend Xu's results to a more general setting. Namely, let \mathcal{M} be a

Date: January 12, 2022.

2010 Mathematics Subject Classification. Primary 46L51; Secondary 46M35, 81P45.

Key words and phrases. Noncommutative L_p -spaces, complex interpolation, quantum relative entropies, data processing inequality.

σ -finite von Neumann algebra equipped with a normal faithful (n.f.) state ϕ , we would like to pursue a complex interpolation result for a couple of (quasi) noncommutative L_p -spaces associated with (\mathcal{M}, ϕ) .

There are two different but equivalent approaches of constructing noncommutative L_p -spaces associated with (\mathcal{M}, ϕ) . One was introduced by Haagerup [15]. We denote by $L_p(\mathcal{M}, \phi)$ the Haagerup noncommutative L_p -space (see Section 2 for definition). However, it is well-known that for two different indices p_0 and p_1 , the natural intersection $L_{p_0}(\mathcal{M}, \phi) \cap L_{p_1}(\mathcal{M}, \phi)$ is trivial [35], thus the couple $(L_{p_0}(\mathcal{M}, \phi), L_{p_1}(\mathcal{M}, \phi))$ is not compatible for interpolation. To overcome this difficulty, Kosaki [22] introduced another kind of noncommutative L_p -space, denoted by $C_p(\mathcal{M}, \phi)$, $1 \leq p \leq \infty$. It was defined as the complex interpolation of the couple $(h_\phi^{1/2} \mathcal{M} h_\phi^{1/2}, L_1(\mathcal{M}, \phi))$ (see Section 2 for the definition of h_ϕ). More precisely, for $1 \leq p \leq \infty$ we have

$$C_p(\mathcal{M}, \phi) := \left[h_\phi^{1/2} \mathcal{M} h_\phi^{1/2}, L_1(\mathcal{M}, \phi) \right]_{\frac{1}{p}},$$

where $C_\infty(\mathcal{M}, \phi) = h_\phi^{1/2} \mathcal{M} h_\phi^{1/2}$. By the reiteration theorem [1, Theorem 4.6.1], one has [22]

$$[C_{p_0}(\mathcal{M}, \phi), C_{p_1}(\mathcal{M}, \phi)]_\theta = C_{p_\theta}(\mathcal{M}, \phi),$$

where $\frac{1}{p_\theta} = \frac{1-\theta}{p_0} + \frac{\theta}{p_1}$ and $1 \leq p_0 < p_1 \leq \infty$. The crucial point of Kosaki's interpolation is that $C_\infty(\mathcal{M}, \phi)$ naturally embeds into $L_1(\mathcal{M}, \phi)$, which makes $C_\infty(\mathcal{M}, \phi)$ and $L_1(\mathcal{M}, \phi)$ a compatible couple of interpolation. Hence the following Kosaki's embedding $i_p(x) = h_\phi^{\frac{1}{2p}} x h_\phi^{\frac{1}{2p}}, x \in \mathcal{M}$ embeds \mathcal{M} into $L_p(\mathcal{M}, \phi)$ (see Proposition 2.6). In this way any pair of Haagerup's noncommutative L_p -spaces is compatible.

In this paper, We show that for $0 < p_0 < p_1 \leq \infty$,

$$(1.2) \quad [L_{p_0}(\mathcal{M}, \phi), L_{p_1}(\mathcal{M}, \phi)]_\theta = L_{p_\theta}(\mathcal{M}, \phi),$$

where $1/p_\theta = (1-\theta)/p_0 + \theta/p_1$ (see Theorem 4.1). The key idea of our proof is to use Xu's reversed Hölder's inequality (see Lemma 3.1 and Corollary 3.2) for tracial noncommutative L_p -spaces [37] and a Hadamard three lines theorem for (quasi) Haagerup noncommutative L_p -spaces (see Proposition 3.6).

Recently, people in the quantum information community introduced a noncommutative L_p -space, denoted by $L_{p,\sigma}(H)$ [3, 26, 23]. Recall that for a fixed positive operator $\sigma \in B(H)$, the space $L_{p,\sigma}(H)$ is defined as the completion of $(B(H), \|\cdot\|_{p,\sigma})$, where the (quasi) norm $\|\cdot\|_{p,\sigma}$ is defined as

$$\|x\|_{p,\sigma} := \left\| \sigma^{\frac{1}{2p}} x \sigma^{\frac{1}{2p}} \right\|_p, \quad x \in B(H)$$

for $0 < p < \infty$. Here $\|\cdot\|_p$ is the Schatten p -norm on $B(H)$. In fact, due to the Tomita-Takesaki theory [36], $L_{p,\sigma}(H)$ is nothing but a special case of Haagerup noncommutative L_p -space. And it was used by Beigi [3] to study the data processing inequality (DPI) of the sandwiched p -Rényi divergence \tilde{D}_p introduced by Müller Lennert et al. [25]. This motivates us to define the following generalization of p -Rényi divergence for $p \in (0, 1) \cup (1, \infty)$:

$$(1.3) \quad \tilde{D}_p(\psi\|\phi) := \frac{1}{p-1} \log \left\| h_\phi^{\frac{1-p}{2p}} h_\psi h_\phi^{\frac{1-p}{2p}} \right\|_{L_p(\mathcal{M}, \phi)}^p,$$

where ϕ, ψ are two normal faithful states on \mathcal{M} . This definition fits well with the one by Jenčová [21] (for $1 < p < \infty$) and the one by Berta et al. [9] (for $1/2 \leq p < 1$), where they firstly generalized the sandwiched p -Rényi divergence \tilde{D}_p to the general von Neumann algebra setting. As an application of our interpolation result, we can easily cover Beigi's results by adapting his argument to our setting. Moreover, we also derive a simple sufficient condition for the equality in DPI for all $p \in (0, 1) \cup (1, \infty)$ (see Theorem 5.8). We end this introduction by the following remark: We have realized that the complex interpolation for noncommutative L_p -spaces has been used in many works about the sandwiched p -Rényi divergence, see for instance [3, 9, 19]. However, they all concern with the Banach noncommutative L_p -spaces ($1 \leq p < \infty$). Our interpolation result for the quasi ones will serve as a great complement to this topic.

Our paper is organized as follows. In Section 2 we summarize necessary preliminaries on tracial noncommutative L_p -spaces, Haagerup noncommutative L_p -spaces, Haagerup's reduction theorem and complex interpolation of tracial noncommutative L_p -spaces. Section 3 is devoted to the Hadamard three lines theorem for Haagerup noncommutative L_p -spaces, which is the key ingredient for proving our main interpolation result. We also present a direct application of our Hadamard three lines theorem to matrix inequalities. In Section 4 we show the main result about the complex interpolation. Section 5 presents some applications to the sandwiched p -Rényi divergence.

2. PRELIMINARIES

2.1. Tracial noncommutative L_p -spaces. In this subsection we concentrate ourselves to noncommutative L_p -spaces associated with semifinite von Neumann algebras, which were firstly laid out in the early 1950's by Segal [33] and Dixmier [11]. Our reference is [31].

Let \mathcal{M} be a semifinite von Neumann algebra equipped with a n.s.f. trace τ . Denote by \mathcal{M}_+ the positive cone of \mathcal{M} and by \mathcal{S}_+ the set of all $x \in \mathcal{M}_+$

such that $\tau[\text{supp}(x)] < \infty$, where $\text{supp}(x)$ denotes the support projection of x . Let \mathcal{S} be the linear span of \mathcal{S}_+ . Then \mathcal{S} is a weak*-dense *-subalgebra of \mathcal{M} . Given $0 < p < \infty$, we define

$$\|x\|_p := [\tau(|x|^p)]^{\frac{1}{p}}, \quad x \in \mathcal{S},$$

where $|x| = (x^*x)^{\frac{1}{2}}$ is the modulus of x . One can show that $\|\cdot\|_p$ is a norm on \mathcal{S} if $1 \leq p < \infty$, and a quasi-norm if $0 < p < 1$. In particular, we have for $p \geq 1$

$$(2.1) \quad \|x + y\|_p \leq \|x\|_p + \|y\|_p, \quad x, y \in \mathcal{S};$$

and for $0 < p < 1$:

$$(2.2) \quad \|x + y\|_p^p \leq \|x\|_p^p + \|y\|_p^p, \quad x, y \in \mathcal{S}.$$

The completion of $(\mathcal{S}, \|\cdot\|_p)$ is denoted by $\Lambda_p(\mathcal{M}, \tau)$, or simply $\Lambda_p(\mathcal{M})$. It is called *tracial noncommutative L_p -space associated with (\mathcal{M}, τ)* . As usual, we set $\Lambda_\infty(\mathcal{M}, \tau) = \mathcal{M}$ equipped with the operator norm.

The elements of $\Lambda_p(\mathcal{M})$ can be viewed as closed densely defined operators on H (H being the Hilbert space on which \mathcal{M} acts). A linear closed operator x is said to be *affiliated with \mathcal{M}* if it commutes with all unitary elements in \mathcal{M}' , i.e., $xu = ux$ for any unitary $u \in \mathcal{M}'$, where \mathcal{M}' denotes the commutant of \mathcal{M} . Note that x can be unbounded on H . An operator x affiliated with \mathcal{M} is said to be *measurable with respect to (\mathcal{M}, τ)* , or simply *measurable* if for any $\varepsilon > 0$, there exists a projection $e \in \mathcal{M}$ such that

$$e(H) \subset \mathcal{D}(x) \text{ and } \tau(e^\perp) \leq \varepsilon,$$

where $e^\perp = 1 - e$ and $\mathcal{D}(x)$ denotes the domain of x . We denote by $L_0(\mathcal{M}, \tau)$, or simply $L_0(\mathcal{M})$ the family of measurable operators. For such an operator x , we define the *distribution function* of x as

$$\lambda_s(x) := \tau[\chi_{(s, \infty)}(|x|)], \quad s \geq 0,$$

where $\chi_{(s, \infty)}(|x|)$ is the spectral projection of $|x|$ corresponding to the interval (s, ∞) , and define the *generalized singular numbers* of x as

$$\mu_t(x) := \inf\{s > 0 : \lambda_s(x) < t\}, \quad t \geq 0.$$

It is easy to check that

$$\|x\| = \lim_{t \rightarrow 0^+} \mu_t(x),$$

and for $0 < p < \infty$

$$(2.3) \quad \|x\|_p^p = \int_0^\infty \mu_t(x)^p dt.$$

2.2. Haagerup noncommutative L_p -spaces. For Haagerup noncommutative L_p -spaces we refer to [35]. The Haagerup noncommutative L_p -space is defined on a general (σ -finite) von Neumann algebra \mathcal{M} associated with a n.f. state ϕ . In fact, the Haagerup noncommutative L_p -space can be defined for any n.s.f. weight, but we will consider only the state case.

Let \mathcal{R} denote the crossed product $\mathcal{M} \rtimes_{\sigma^\phi} \mathbb{R}$, where $\{\sigma_t^\phi\}_{t \in \mathbb{R}}$ is the *modular automorphism group* associated with ϕ on \mathcal{M} (for general modular theory see [36]).

Recall that if \mathcal{M} acts on a Hilbert space H , \mathcal{R} is the von Neumann algebra acting on $L_2(\mathbb{R}, H)$ generated by the operators $\pi(x), x \in \mathcal{M}$, and $\lambda(s), s \in \mathbb{R}$, such that: for every $\xi \in L_2(\mathbb{R}, H)$ and $t \in \mathbb{R}$,

$$\pi(x)(\xi)(t) = \sigma_{-t}(x)\xi(t), \quad \lambda(s)(\xi)(t) = \xi(t - s).$$

π is a normal faithful representation of \mathcal{M} on $L_2(\mathbb{R}; H)$, so we can identify $\pi(\mathcal{M})$ with \mathcal{M} . There is a dual action $\{\hat{\sigma}_t^\phi\}_{t \in \mathbb{R}}$ of \mathcal{R} uniquely given by

$$\hat{\sigma}_t^\phi(x) = x, \quad \hat{\sigma}_t^\phi(\lambda(s)) = e^{-ist} \lambda(s),$$

for $x \in \mathcal{M}$ and $s, t \in \mathbb{R}$.

It is known that \mathcal{R} is semifinite and there is a canonical n.s.f. trace τ on \mathcal{R} satisfying

$$\tau \circ \hat{\sigma}_t^\phi = e^{-t} \tau, \quad t \in \mathbb{R},$$

Any normal positive functional $\psi \in \mathcal{M}_*^+$ induces a dual normal semifinite weight $\hat{\psi}$ on \mathcal{R} defined by

$$(2.4) \quad \hat{\psi}(x) = \psi \left[\int_{-\infty}^{\infty} \hat{\sigma}_t^\phi(x) dt \right], \quad x \in \mathcal{R}^+.$$

The dual weight $\hat{\psi}$ has a *Radon-Nikodym derivative* with respect to τ , denoted by h_ψ :

$$\tau(h_\psi x) = \hat{\psi}(x), \quad x \in \mathcal{R}^+.$$

Here $\tau(h_\psi \cdot)$ is understood as $\tau(h_\psi^{\frac{1}{2}} \cdot h_\psi^{\frac{1}{2}})$. In particular, the dual weight $\hat{\phi}$ of our distinguished n.s.f. state ϕ has a Radon-Nikodym derivative h_ϕ with respect to τ . We will call h_ϕ the *density operator* of ϕ .

Let $L_0(\mathcal{R}, \tau)$ denote the topological $*$ -algebra of all operators on $L_2(\mathbb{R}, H)$ which are measurable with respect to (\mathcal{R}, τ) . Then for $0 < p \leq \infty$, the Haagerup L_p -space is defined as

$$(2.5) \quad L_p(\mathcal{M}, \phi) = \{x \in L_0(\mathcal{R}, \tau) : \hat{\sigma}_t^\phi(x) = e^{-t/p} x, \quad \forall t \in \mathbb{R}\}.$$

Recall that for any $\psi \in \mathcal{M}_*^+$, we have $h_\psi \in L_0(\mathcal{R}, \tau)$ and

$$\hat{\sigma}_t^\phi(h_\psi) = e^{-t} h_\psi, \quad \forall t \in \mathbb{R}.$$

Thus $h_\psi \in L_1(\mathcal{M}, \phi)_+$. This correspondence between \mathcal{M}_*^+ and $L_1(\mathcal{M}, \phi)_+$ extends to a bijection between \mathcal{M}_* and $L_1(\mathcal{M}, \phi)$. More precisely, for any $\psi \in \mathcal{M}_*$, if $\psi = u|\psi|$ is its polar decomposition, then we have

$$h_\psi = uh_{|\psi|}.$$

Thus we may define a norm on $L_1(\mathcal{M}, \phi)$ by

$$\|h_\psi\|_1 := \|\psi\|_{\mathcal{M}_*} = |\psi|(1), \quad \psi \in \mathcal{M}_*.$$

Hence $L_1(\mathcal{M}, \phi) = \mathcal{M}_*$ isometrically. Due to this correspondence, there is a linear functional on $L_1(\mathcal{M}, \phi)$, denoted by tr , given through

$$\text{tr}(x) = \psi_x(1), \quad x \in L_1(\mathcal{M}, \phi),$$

where ψ_x is the unique normal functional associated with x by the identification between \mathcal{M}_* and $L_1(\mathcal{M}, \phi)$ we discussed earlier.

Now for $x \in L_p(\mathcal{M}, \phi)$, $0 < p < \infty$, we can define

$$(2.6) \quad \|x\|_p := \| |x|^p \|_1^{\frac{1}{p}} = [\text{tr}(|x|^p)]^{\frac{1}{p}}.$$

Then $\|\cdot\|_p$ is a norm (quasi-norm if $p < 1$) on $L_p(\mathcal{M}, \phi)$. For Haagerup L_p -spaces we have the following Hölder's inequality which will be used frequently: for every $p, q, r > 0$ with $\frac{1}{r} = \frac{1}{p} + \frac{1}{q}$, we have

$$(2.7) \quad \|xy\|_r \leq \|x\|_p \|y\|_q, \quad x \in L_p(\mathcal{M}, \phi), y \in L_q(\mathcal{M}, \phi).$$

Moreover, from (2.6) it follows directly that

$$(2.8) \quad \|x\|_{2p}^2 = \|x^*x\|_p = \|xx^*\|_p, \quad x \in L_{2p}(\mathcal{M}, \phi).$$

Note that from (2.5), for different p_0 and p_1 , the intersection of $L_{p_0}(\mathcal{M}, \phi)$ and $L_{p_1}(\mathcal{M}, \phi)$ is trivial.

Let \mathcal{M}_a be the family of *analytic elements* in \mathcal{M} . Recall that for any $x \in \mathcal{M}$, it belongs to \mathcal{M}_a iff $t \mapsto \sigma_t^\phi(x)$ extends to an analytic function from \mathbb{C} to \mathcal{M} . We will use the following technical lemma proved by Junge and Xu [17].

Lemma 2.1. [17, Lemma 1.1]. *Let $0 < p < \infty, 0 \leq \eta \leq 1$. Then:*

$$(i) \quad h_\phi^{\frac{1-\eta}{p}} \mathcal{M}_a h_\phi^{\frac{\eta}{p}} = \mathcal{M}_a h_\phi^{\frac{1}{p}};$$

$$(ii) \quad \mathcal{M}_a h_\phi^{\frac{1}{p}} \text{ is dense in } L_p(\mathcal{M}, \phi).$$

We note that here $h_\phi^{\frac{1-\eta}{p}} \mathcal{M}_a h_\phi^{\frac{\eta}{p}} = \mathcal{M}_a h_\phi^{\frac{1}{p}}$ follows from

$$(2.9) \quad x h_\phi^{\frac{1}{p}} = h_\phi^{\frac{1-\eta}{p}} \sigma_{\frac{i(1-\eta)}{p}}^\phi(x) h_\phi^{\frac{\eta}{p}}, \quad x \in \mathcal{M}_a.$$

Now we mention two basic facts about Haagerup L_p -spaces. The first is, when \mathcal{M} is a semifinite von Neumann algebra associated with a n.f. tracial

state τ , Haagerup L_p -space $L_p(\mathcal{M}, \tau)$ and tracial L_p -space $\Lambda_p(\mathcal{M}, \tau)$ can be isometrically identified [35, Chapter II]. In fact, in this case, $\mathcal{M} \rtimes_{\sigma\tau} \mathbb{R} = \mathcal{M} \overline{\otimes} L_\infty(\mathbb{R})$ and the density operator of τ is: $h_\tau = id \otimes \exp(\cdot)$. For each $x \in L_p(\mathcal{M}, \tau)$ there exists unique $x' \in \Lambda_p(\mathcal{M}, \tau)$ such that $x = x' \otimes \exp(\cdot/p)$. Moreover, $\|x\|_{L_p(\mathcal{M}, \tau)} = \|x'\|_{\Lambda_p(\mathcal{M}, \tau)}$. Thus the map $x \mapsto x'$ yields an isometry between $L_p(\mathcal{M}, \tau)$ and $\Lambda_p(\mathcal{M}, \tau)$.

The second well-known fact is that Haagerup L_p -spaces $L_p(\mathcal{M}, \phi)$ are independent of the choice of state ϕ up to isometry [35, Chapter II]. Namely, suppose that φ_0, φ_1 are two n.f. states on a von Neumann algebra \mathcal{M} . Recall that $\mathcal{R}_i := \mathcal{M} \rtimes_{\sigma\varphi_i} \mathbb{R}$ is generated by $\pi_i(x)$, $x \in \mathcal{M}$ and $\lambda(s)$, $s \in \mathbb{R}$, where

$$\pi_i(x)\xi(t) = \sigma_{-t}^{\varphi_i}(x)\xi(t), \quad \lambda(s)\xi(t) = \xi(t-s), \quad \xi \in L_2(\mathbb{R}, H), \quad s, t \in \mathbb{R},$$

for $i = 0, 1$. Then there is an isometry $\kappa : L_p(\mathcal{M}, \varphi_0) \rightarrow L_p(\mathcal{M}, \varphi_1)$ such that [35, Theorem 37]

$$\kappa[\pi_0(x)] = \pi_1(x), \quad \kappa[\lambda(t)] = \pi_0(u_t^*)\lambda(t), \quad t \in \mathbb{R}, x \in \mathcal{M},$$

where $u_t = (h_{\varphi_1} : h_{\varphi_0})_t$ is the *Radon-Nikodym cocycle* of φ_1 relative to φ_0 :

$$u_{s+t} = u_s \sigma_t^{\varphi_0}(u_t), \quad \sigma_t^{\varphi_1}(x) = u_t \sigma_t^{\varphi_0}(x) u_t^*,$$

where $s, t \in \mathbb{R}$, $x \in \mathcal{M}$, and h_{φ_i} is the density operator of φ_i , $i = 0, 1$. In particular, if $\varphi_1 = \varphi_0(h \cdot)$, then $u_t = h^{it}$. Thus we have $\kappa[\pi_0(h)h_{\varphi_0}] = h_{\varphi_1}$.

Then combining the above two facts, we obtain

Lemma 2.2. *Let \mathcal{M} be a von Neumann algebra associated with a n.f. tracial state τ . Let ϕ and ψ be two n.f. states on \mathcal{M} defined by:*

$$\psi(x) := \tau(\rho x), \quad \phi(x) := \tau(\sigma x), \quad x \in \mathcal{M},$$

where $\rho, \sigma \geq 0$ and $\tau(\rho) = \tau(\sigma) = 1$. Then there exist an isometry $\theta : L_p(\mathcal{M}, \tau) \rightarrow L_p(\mathcal{M}, \phi)$ and an isometry $\theta' : \Lambda_p(\mathcal{M}, \tau) \rightarrow L_p(\mathcal{M}, \phi)$ such that

$$(2.10) \quad h_\psi = \theta[h_\tau \pi_{\sigma\tau}(\rho)] = \theta'(\rho),$$

where h_ψ is the Radon-Nikodym derivative of ψ with respect to the canonical trace τ^ϕ on $\mathcal{M} \rtimes_{\sigma\phi} \mathbb{R}$ and h_τ is the density operator of τ . In particular, we have

$$(2.11) \quad h_\phi = \theta[h_\tau \pi_{\sigma\tau}(\sigma)] = \theta'(\sigma).$$

Proof. The equation (2.11) is a direct consequence of two facts as above. Note that $\psi = \phi(x^* \cdot x)$, with $x = \rho^{\frac{1}{2}} \sigma^{-\frac{1}{2}}$. Then (2.10) follows immediately from (2.11) and the fact [35, Chapter II, Proposition 4] that $h_{\phi(x^* \cdot x)} = x h_\phi x^*$. \square

Using the above lemma, we can show that the noncommutative L_p -space $L_{p,\sigma}(H)$ is a Haagerup L_p -space.

Definition 2.3. Let \mathcal{M} be a σ -finite von Neumann algebra equipped with a n.f. state ϕ . For $0 < p < \infty$, define a (quasi) norm $\|\cdot\|_{p,\phi}$ on \mathcal{M} by

$$\|x\|_{p,\phi} := \left\| h_{\phi}^{\frac{1}{2p}} x h_{\phi}^{\frac{1}{2p}} \right\|_{L_p(\mathcal{M},\phi)}, \quad x \in \mathcal{M}.$$

Recall [3] that for a fixed positive operator $\sigma \in B(H)$, the space $L_{p,\sigma}(H)$ is defined as the completion of $(B(H), \|\cdot\|_{p,\sigma})$, where the (quasi) norm $\|\cdot\|_{p,\sigma}$ is defined by

$$\|x\|_{p,\sigma} = \left\| \sigma^{\frac{1}{2p}} x \sigma^{\frac{1}{2p}} \right\|_p, \quad x \in B(H)$$

for $0 < p < \infty$. Here $\|\cdot\|_p$ is the Schatten p -norm on $B(H)$.

Proposition 2.4. Let \mathcal{M} be a finite von Neumann algebra with a n.f. tracial state τ , and ϕ be a n.f. state on \mathcal{M} defined by $\phi = \tau(\sigma \cdot)$, where $\sigma \geq 0$ and $\tau(\sigma) = 1$. Then

$$(2.12) \quad \|x\|_{p,\phi} = \left[\tau \left(\left| \sigma^{\frac{1}{2p}} x \sigma^{\frac{1}{2p}} \right|^p \right) \right]^{\frac{1}{p}}, \quad x \in \mathcal{M}$$

for $0 < p < \infty$.

Proof. Using Lemma 2.2, we have

$$\begin{aligned} \left\| h_{\phi}^{\frac{1}{2p}} x h_{\phi}^{\frac{1}{2p}} \right\|_{L_p(\mathcal{M},\phi)}^p &= \left\| \theta' \left(\sigma^{\frac{1}{2p}} x \sigma^{\frac{1}{2p}} \right) \right\|_{L_p(\mathcal{M},\phi)}^p \\ &= \left\| \sigma^{\frac{1}{2p}} x \sigma^{\frac{1}{2p}} \right\|_{\Lambda_p(\mathcal{M},\tau)}^p \\ &= \tau \left(\left| \sigma^{\frac{1}{2p}} x \sigma^{\frac{1}{2p}} \right|^p \right), \end{aligned}$$

where θ' is the isometry from $\Lambda_p(\mathcal{M},\tau)$ to $L_p(\mathcal{M},\phi)$. \square

2.3. Haagerup's reduction theorem. The idea of the Haagerup reduction theorem is to approximate a general von Neumann algebra by finite ones. It was firstly proposed by Haagerup and published in [14] after amending more details and applications to noncommutative martingale inequalities and maximal inequalities. In this paper, we will use the Haagerup reduction theorem for noncommutative L_p -spaces, which briefly states that a Haagerup L_p -space associated with a general von Neumann algebra can be approximated by a sequence of tracial L_p -spaces.

Let G denotes the subgroup $\bigcup_{n \geq 1} 2^{-n}\mathbb{Z}$ of \mathbb{R} . Let \mathcal{M} be a σ -finite von Neumann algebra associated with a normal faithful state ϕ . Denote $\mathcal{R} := \mathcal{M} \rtimes_{\sigma\phi} G$. Then we have the following reduction theorem.

Theorem 2.5. [14, Theorem 3.1]. *For $0 < p < \infty$, let $L_p(\mathcal{M}, \phi)$ be the Haagerup noncommutative L_p -space. Then there exist a sequence $\{\mathcal{R}_n\}_{n \geq 1}$ of finite von Neumann algebras, each equipped with a normal faithful finite trace τ_n , and for each $n \geq 1$ an isometric embedding $\theta_n : \Lambda_p(\mathcal{R}_n, \tau_n) \rightarrow L_p(\mathcal{R}, \hat{\phi})$ such that*

- (i) *the sequence $\{\theta_n[\Lambda_p(\mathcal{R}_n, \tau_n)]\}_{n \geq 1}$ is increasing;*
- (ii) *$\bigcup_{n \geq 1} \theta_n[\Lambda_p(\mathcal{R}_n, \tau_n)]$ is dense in $L_p(\mathcal{R}, \hat{\phi})$;*
- (iii) *$L_p(\mathcal{M}, \phi)$ is isometric to a subspace Y_p of $L_p(\mathcal{R}, \hat{\phi})$;*
- (iv) *Y_p and all $\theta_n[\Lambda_p(\mathcal{R}_n, \tau_n)]$ are 1-complemented in $L_p(\mathcal{R}, \hat{\phi})$ for $1 \leq p < \infty$.*

Here $\Lambda_p(\mathcal{R}_n, \tau_n)$ is the tracial noncommutative L_p -space associated with (\mathcal{R}_n, τ_n) .

2.4. Complex interpolation. We refer to [1] for more information in complex interpolation theory. Let S denote the strip $\{z \in \mathbb{C} : 0 < \operatorname{Re} z < 1\}$, and \bar{S} its closure. Let $A(S)$ be the set of all bounded functions which are analytic in S and continuous in \bar{S} . Let (X_0, X_1) be an interpolation couple of (quasi) Banach spaces. Set

$$\mathcal{F}(X_0, X_1) := \left\{ f : f(z) = \sum_{k=1}^m f_k(z) x_k, x_k \in X_0 \cap X_1, f_k \in A(S), m \in \mathbb{N} \right\}.$$

Equipped with the norm

$$\|f\|_{\mathcal{F}(X_0, X_1)} = \sup_{t \in \mathbb{R}} \{ \|f(it)\|_{X_0}, \|f(1+it)\|_{X_1} \},$$

$\mathcal{F}(X_0, X_1)$ becomes a (quasi) Banach space. For $0 < \theta < 1$ we define the complex interpolation (quasi) norm $\|\cdot\|_\theta$ on $X_0 \cap X_1$ as follows

$$(2.13) \quad \|x\|_\theta = \inf \{ \|f\|_{\mathcal{F}(X_0, X_1)} : f(\theta) = x, f \in \mathcal{F}(X_0, X_1) \}, \quad x \in X_0 \cap X_1.$$

Then the complex interpolation space of X_0 and X_1 is defined as the completion of $(X_0 \cap X_1, \|\cdot\|_\theta)$, denoted by $[X_0, X_1]_\theta$.

For the tracial noncommutative L_p -space $\Lambda_p(\mathcal{M}, \tau)$, it is well-known that

$$(2.14) \quad [\Lambda_{p_0}(\mathcal{M}, \tau), \Lambda_{p_1}(\mathcal{M}, \tau)]_\theta = \Lambda_{p_\theta}(\mathcal{M}, \tau),$$

where $1 \leq p_0 < p_1 \leq \infty$, and $\frac{1}{p_\theta} = \frac{1-\theta}{p_0} + \frac{\theta}{p_1}$ (see [31]). By showing a type of reversed Hölder's inequality (see Lemma 3.1), Xu [37] proved the above complex interpolation result for the quasi-Banach space $\Lambda_p(\mathcal{M}, \tau)$:

$$(2.15) \quad [\Lambda_{p_0}(\mathcal{M}, \tau), \Lambda_{p_1}(\mathcal{M}, \tau)]_\theta = \Lambda_{p_\theta}(\mathcal{M}, \tau),$$

where $0 < p_0 < p_1 \leq \infty$ and $\frac{1}{p_\theta} = \frac{1-\theta}{p_0} + \frac{\theta}{p_1}$.

2.5. Compatibility of Haagerup L_p -spaces. As we have mentioned in the introduction, the compatibility of two (quasi) Haagerup noncommutative L_p -spaces is due to Kosaki's construction. More precisely, we have the following proposition:

Proposition 2.6. *Let \mathcal{M} be a σ -finite von Neumann algebra with a n.f. state ϕ . For $0 < p_0 < p_1 \leq \infty$, and $x \in \mathcal{M}$, we have*

$$\|x\|_{p_0, \phi} \leq \|x\|_{p_1, \phi}.$$

Proof. For any $x \in \mathcal{M}$, $h_\phi^{\frac{1}{2p_0}} x h_\phi^{\frac{1}{2p_0}}$ can be decomposed as

$$h_\phi^{\frac{1}{2p_0}} x h_\phi^{\frac{1}{2p_0}} = h_\phi^{\frac{1}{q}} h_\phi^{\frac{1}{2p_1}} x h_\phi^{\frac{1}{2p_1}} h_\phi^{\frac{1}{q}} = h_\phi^{\frac{1}{q}} y h_\phi^{\frac{1}{q}},$$

where $\frac{1}{q} + \frac{1}{2p_1} = \frac{1}{2p_0}$ and $y = h_\phi^{\frac{1}{2p_1}} x h_\phi^{\frac{1}{2p_1}}$. By Hölder's inequality (2.7), we have

$$\begin{aligned} \left\| h_\phi^{\frac{1}{2p_0}} x h_\phi^{\frac{1}{2p_0}} \right\|_{L_{p_0}(\mathcal{M}, \phi)} &\leq \|y\|_{L_{p_1}(\mathcal{M}, \phi)} \left\| h_\phi^{\frac{1}{q}} \right\|_{L_q(\mathcal{M}, \phi)}^2 \\ &= \|y\|_{L_{p_1}(\mathcal{M}, \phi)} \|h_\phi^{\frac{1}{q}}\|_{L_1(\mathcal{M}, \phi)}^{\frac{2}{q}} = \|y\|_{L_{p_1}(\mathcal{M}, \phi)}. \quad \square \end{aligned}$$

Due to the above proposition, for $0 < p_0 < p_1 \leq \infty$, $L_{p_1}(\mathcal{M}, \phi)$ can be embedded into $L_{p_0}(\mathcal{M}, \phi)$ via the map $h_\phi^{\frac{1}{2p_1}} x h_\phi^{\frac{1}{2p_1}} \mapsto h_\phi^{\frac{1}{2p_0}} x h_\phi^{\frac{1}{2p_0}}$. With this embedding, $(L_{p_0}(\mathcal{M}, \phi), L_{p_1}(\mathcal{M}, \phi))$ forms an interpolation couple.

3. HADAMARD THREE LINES THEOREM FOR (QUASI) NONCOMMUTATIVE L_p -SPACES

3.1. The tracial case. In this subsection, we denote by \mathcal{M} a semifinite von Neumann algebra associated with a n.s.f. trace τ . For $\theta \in [0, 1]$, and $0 < p_0 \leq p_1 \leq \infty$, define p_θ as follows:

$$(3.1) \quad \frac{1}{p_\theta} = \frac{1-\theta}{p_0} + \frac{\theta}{p_1}.$$

Recall that $\bar{S} = \{z \in \mathbb{C} : 0 \leq \operatorname{Re} z \leq 1\}$. Denote by $AF(X)$ the set of all bounded functions from \bar{S} to X which are analytic in S and continuous on \bar{S} . The following reversed Hölder's inequality is due to Xu.

Lemma 3.1. [37, Lemma 2.2]. *Let \mathcal{M} be a von Neumann algebra associated with a n.s.f. trace τ such that $\tau(1) < \infty$. Fix $0 < \lambda < 1$. Then for any $f \in AF(\mathcal{M})$ and any $\epsilon > 0$, there exist $g, h \in AF(\mathcal{M})$ such that $f = gh$ and for all $z \in \partial S, t \geq 0$ we have*

$$\begin{cases} \mu_t(g(z)) \leq \mu_t(f(z))^{1-\lambda} + \epsilon, \\ \mu_t(h(z)) \leq \mu_t(f(z))^\lambda + \epsilon. \end{cases}$$

Remark here that our statement is slightly different from Xu's original form. Indeed, in [37] \mathcal{M} -valued analytic functions in $AF(\mathcal{M})$ are defined on the interval $[0, 2\pi]$. But they are equivalent, since one can identify $[0, 2\pi]$ with the unit circle and then apply Riemann mapping theorem. As a direct consequence of (2.3) and Lemma 3.1 we have

Corollary 3.2. *Let (\mathcal{M}, τ) and λ be as above. Then for any $f \in AF(\mathcal{M})$ and any $\epsilon, r > 0$, there exist $g, h \in AF(\mathcal{M})$ such that $f = gh$ and for all $z \in \partial S$ we have*

$$\begin{cases} \|g(z)\|_{\frac{r}{1-\lambda}} \leq \|f(z)\|_r^{1-\lambda} + \epsilon, \\ \|h(z)\|_{\frac{r}{\lambda}} \leq \|f(z)\|_r^\lambda + \epsilon. \end{cases}$$

Remark 3.3. The above corollary is a special case of the Riesz factorization theorem, which was proved in the general case of subdiagonal algebras by Blecher-Labuschagne [4] and Bekjan-Xu [5].

Proposition 3.4. *Let \mathcal{M} be a semifinite von Neumann algebra associated with a n.s.f. trace τ . Let $G \in AF(\mathcal{M})$. Assume that $0 < p_0 < p_1 \leq \infty$, and for $\theta \in [0, 1]$ define p_θ as in the equation (3.1). For $k = 0, 1$, define*

$$M_k := \sup_{t \in \mathbb{R}} \|G(k + it)\|_{p_k}.$$

Then we have

$$\|G(\theta)\|_{p_\theta} \leq M_0^{1-\theta} M_1^\theta.$$

Proof. The proof of Banach spaces case $1 \leq p_0 < p_1 \leq \infty$ is a standard argument, using Hadamard three lines theorem for analytic functions. We refer to [3, Theorem 2] and [21, Theorem 2.10] for the proof of (not necessarily tracial) state case for $1 \leq p_0 < p_1 \leq \infty$. For the proof for quasi-Banach spaces case $0 < p_0 < p_1 \leq \infty$, choose $n \in \mathbb{N}$ such that $np_0 \geq 1$. Then for any $\epsilon > 0$, by Corollary 3.2 there exist $G_1, \dots, G_n \in AF(\mathcal{M})$ such that $G = \prod_{j=1}^n G_j$ and for $1 \leq j \leq n$

$$(3.2) \quad \sup_{t \in \mathbb{R}} \|G_j(k + it)\|_{np_k} \leq \sup_{t \in \mathbb{R}} \|G(k + it)\|_{p_k}^{\frac{1}{n}} + \epsilon, \quad k = 0, 1.$$

Hence by Hölder's inequality and the desired result for Banach spaces case, we have

$$\|G(\theta)\|_{p_\theta} \leq \prod_{j=1}^n \|G_j(\theta)\|_{np_\theta} \leq \prod_{j=1}^n \left(\sup_{t \in \mathbb{R}} \|G_j(it)\|_{np_0} \right)^{1-\theta} \left(\sup_{t \in \mathbb{R}} \|G_j(1 + it)\|_{np_1} \right)^\theta.$$

Applying (3.2), we obtain

$$\begin{aligned} \|G(\theta)\|_{p_\theta} &\leq \prod_{j=1}^n \left(\sup_{t \in \mathbb{R}} \|G(it)\|_{p_0}^{\frac{1}{n}} + \epsilon \right)^{1-\theta} \left(\sup_{t \in \mathbb{R}} \|G(1+it)\|_{p_1}^{\frac{1}{n}} + \epsilon \right)^\theta \\ &\leq \left(\sup_{t \in \mathbb{R}} \|G(it)\|_{p_0} \right)^{1-\theta} \left(\sup_{t \in \mathbb{R}} \|G(1+it)\|_{p_1} \right)^\theta + O(\epsilon). \end{aligned}$$

Letting $\epsilon \rightarrow 0^+$, we deduce our desired result. \square

3.2. The Haagerup noncommutative L_p -spaces case.

Lemma 3.5. *Let \mathcal{M} be a semifinite von Neumann algebra with a n.f. trace τ . Let $G \in AF(\mathcal{M})$. Assume that $0 < p_0 < p_1 \leq \infty$, and for $\theta \in [0, 1]$ define p_θ as in the equation (3.1). For $k = 0, 1$ define*

$$M_k := \sup_{t \in \mathbb{R}} \|G(k+it)\|_{p_k, \tau}.$$

Then we have

$$\|G(\theta)\|_{p_\theta, \tau} \leq M_0^{1-\theta} M_1^\theta.$$

Proof. It is obvious that

$$\|G(z)\|_{p_z, \tau} = \left\| h_\tau^{\frac{1}{2p_z}} G(z) h_\tau^{\frac{1}{2p_z}} \right\|_{L_{p_z}(\mathcal{M}, \tau)} = \|G(z)\|_{\Lambda_{p_z}(\mathcal{M}, \tau)},$$

where $\frac{1}{p_z} = \frac{1-\text{Re}(z)}{p_0} + \frac{\text{Re}(z)}{p_1}$. Then our result can be deduced from Proposition 3.4. \square

The following proposition extends Beigi's result [3, Theorem 2] (see also Jenčová's result [21, Theorem 2.10]) to the quasi-Banach space case.

Proposition 3.6. *Let \mathcal{M} be a (σ -finite) von Neumann algebra with a n.s.f. state ϕ . Let $G \in AF(\mathcal{M})$. Assume that $0 < p_0 < p_1 \leq \infty$, and for $\theta \in [0, 1]$ define p_θ as in the equation (3.1). For $k = 0, 1$ define*

$$M_k := \sup_{t \in \mathbb{R}} \|G(k+it)\|_{p_k, \phi}.$$

Then we have

$$\|G(\theta)\|_{p_\theta, \phi} \leq M_0^{1-\theta} M_1^\theta.$$

Proof. The key idea is to use the Haagerup reduction theorem. Recall that $\mathcal{R} = \mathcal{M} \rtimes_{\sigma_\phi} \cup_{n \geq 1} 2^{-n} \mathbb{Z}$, and $L_p(\mathcal{M}, \phi)$ can be isometrically embedded into $L_p(\mathcal{R}, \hat{\phi})$ for any $0 < p < \infty$. Hence it suffices to show that for $G \in AF(\mathcal{R})$

$$\|G(\theta)\|_{p_\theta, \hat{\phi}} \leq M_0^{1-\theta} M_1^\theta,$$

where $M_k := \sup_{t \in \mathbb{R}} \|G(k+it)\|_{p_k, \hat{\phi}}$, $k = 0, 1$.

According to Theorem 2.5, $\bigcup_n L_p(\mathcal{R}_n, \tau_n)$ is dense in $L_p(\mathcal{R}, \hat{\phi})$. Hence without loss of generality, we can assume that there exists an integer N , such that $x = G(\theta) \in \mathcal{R}$, and $M_k \leq 1$ for $k = 0, 1$. Suppose that

$$G(z) = \sum_m G_m(z) x_m,$$

where $G_m \in A(S)$ and $x_m \in \mathcal{R}_N$. Then for $\epsilon > 0$, there exists a sequence $\{y_m\} \subset \mathcal{R}_{N'}$ such that for each m

$$(3.3) \quad \|x_m - y_m\|_{p_1, \tau_{N'}} < \frac{\epsilon}{2^m K_m},$$

where $N' > N$ and $K_m := \sup\{|G_m(z)| : z \in \overline{S}\} < \infty$. Hence

$$G = \sum_m G_m y_m + \sum_m G_m (x_m - y_m) =: g + h,$$

where $g(z) = \sum_m G_m(z) y_m$ and $h(z) = \sum_m G_m(z) (x_m - y_m)$. Then if $p_0 \geq 1$, using the inequality (2.1) we have

$$\begin{aligned} \|h(it)\|_{p_0, \tau_{N'}} &\leq \sum_m |G_m(it)| \|x_m - y_m\|_{p_0, \tau_{N'}} \\ &\leq \sum_m |G_m(it)| \|x_m - y_m\|_{p_1, \tau_{N'}} \\ &\leq \sum_m \frac{\epsilon}{2^m} = O(\epsilon). \end{aligned}$$

If $0 < p_0 < 1$, then we apply the inequality (2.2) instead of (2.1) and obtain also

$$\|h(it)\|_{p_0, \tau_{N'}}^{p_0} \leq \sum_m \frac{\epsilon^{p_0}}{2^{mp_0}} = O(\epsilon^{p_0}).$$

So in any case, we always have $\|h(it)\|_{p_0, \tau_{N'}} \leq O(\epsilon)$. Similarly we have

$$\|h(1 + it)\|_{p_1, \tau_{N'}} \leq O(\epsilon).$$

Thus using Lemma 3.5, we obtain

$$\|h(\theta)\|_{p_\theta, \tau_{N'}} \leq O(\epsilon),$$

and moreover

$$\begin{cases} \|g(it)\|_{p_0, \tau_{N'}} \leq 1 + O(\epsilon), \\ \|g(1 + it)\|_{p_1, \tau_{N'}} \leq 1 + O(\epsilon). \end{cases}$$

Set $\tilde{G}(z) := g(z) + h(\theta)$. Then it is easy to see that $\tilde{G} \in AF(\mathcal{R}_{N'})$ and $\tilde{G}(\theta) = G(\theta)$. Moreover,

$$\begin{cases} \|\tilde{G}(it)\|_{p_0, \tau_{N'}} \leq 1 + O(\epsilon), \\ \|\tilde{G}(1 + it)\|_{p_1, \tau_{N'}} \leq 1 + O(\epsilon). \end{cases}$$

Hence from Lemma 3.5 we have $\|\tilde{G}(\theta)\|_{p_\theta, \tau_{N'}} \leq 1 + O(\epsilon)$. Thus

$$\|G(\theta)\|_{p_\theta, \hat{\phi}} = \|G(\theta)\|_{p_\theta, \tau_{N'}} = \|\tilde{G}(\theta)\|_{p_\theta, \tau_{N'}} \leq 1 + O(\epsilon).$$

By letting $\epsilon \rightarrow 0^+$ we deduce our result. \square

3.3. Applications to some matrix inequalities. In this subsection, let \mathcal{M} denote the set of all $d \times d$ complex matrices. The following proposition extends Sutter et al.'s result [34, Theorem 3.1] (see also [19, Lemma 3.3]) to the quasi-Banach spaces case, which generalizes Hirschman's strengthening of the Hadamard three lines theorem [13].

Proposition 3.7. *Let $G \in AF(\mathcal{M})$. Assume that $0 < p_0 < p_1 \leq \infty$, and for $\theta \in [0, 1]$ define p_θ as in the equation (3.1). Then we have*

$$(3.4) \quad \log \|G(\theta)\|_{p_\theta} \leq \int_{-\infty}^{\infty} dt \left(\beta_{1-\theta}(t) \log \|G(it)\|_{p_0}^{1-\theta} + \beta_\theta(t) \log \|G(1+it)\|_{p_1}^\theta \right),$$

where $\beta_\theta(t) = \frac{\sin(\pi\theta)}{2\theta(\cosh(\pi t) + \cos(\pi\theta))}$.

Proof. The proof is similar to that of Proposition 3.4. Choose an integer $n \in \mathbb{N}$ such that $np_0 \geq 1, np_1 \geq 1$, then we have

$$\begin{aligned} \log \|G(\theta)\|_{p_\theta} &\leq \sum_{k=1}^n \log \|G_k(\theta)\|_{np_\theta} \\ &\leq \sum_{k=1}^n \int_{-\infty}^{\infty} dt \left(\beta_{1-\theta}(t) \log \|G_k(it)\|_{np_0}^{1-\theta} + \beta_\theta(t) \log \|G_k(1+it)\|_{np_1}^\theta \right) \\ &= \int_{-\infty}^{\infty} dt \left(\beta_{1-\theta}(t) \log \prod_{k=1}^n \|G_k(it)\|_{np_0}^{1-\theta} + \beta_\theta(t) \log \prod_{k=1}^n \|G_k(1+it)\|_{np_1}^\theta \right) \\ &\leq \int_{-\infty}^{\infty} dt \beta_{1-\theta}(t) \log \left(\|G(it)\|_{p_0}^{1/n} + \epsilon \right)^{n(1-\theta)} \\ &\quad + \int_{-\infty}^{\infty} dt \beta_\theta(t) \log \left(\|G(1+it)\|_{p_1}^{1/n} + \epsilon \right)^{n\theta}, \end{aligned}$$

where $G_k \in AF(\mathcal{M}), k = 1, \dots, n$, are functions that are deduced by Corollary 3.2. Note that for the first inequality we have used Hölder's inequality, and for the second one, we have used Theorem 3.1 in [34], which states that the inequality (3.4) is true for the Banach spaces case, i.e., $1 \leq p_0, p_1 \leq \infty$. We complete our proof by letting $\epsilon \rightarrow 0^+$. \square

Corollary 3.8. *Let $0 < p \leq \infty, r \in (0, 1], n \in \mathbb{N}$, then for a family of positive semidefinite matrices $\{A_k\}_{k=1}^n \subset \mathcal{M}$ we have*

$$(3.5) \quad \log \left\| \left\| \prod_{k=1}^n A_k^r \right\|_p^{\frac{1}{r}} \right\|_p \leq \int_{-\infty}^{\infty} dt \beta_r(t) \log \left\| \prod_{k=1}^n A_k^{1+it} \right\|_p.$$

Proof. We use the same idea of [34, Theorem 3.2]. Define the function

$$G(z) := \prod_{k=1}^n A_k^z.$$

It satisfies the conditions of Proposition 3.7. Furthermore we pick $\theta = r, p_0 = \infty$ and $p_1 = p$. Then $p_\theta = p/r$ and we have

$$\begin{cases} \log \|G(it)\|_{p_0}^{1-\theta} = (1-r) \log \left\| \prod_{k=1}^n A_k^{it} \right\|_\infty = 0, \\ \log \|G(1+it)\|_{p_1}^\theta = r \log \left\| \prod_{k=1}^n A_k^{1+it} \right\|_p, \\ \text{and } \log \|G(\theta)\|_{p_\theta} = r \log \left\| \prod_{k=1}^n A_k^r \right\|_{p_\theta}^{\frac{1}{r}}. \end{cases}$$

Plugging the above equations into Proposition 3.7 deduces the corollary. \square

Remark 3.9. In [34], Hiai et al. proved the equation (3.5) for any unitarily invariant norm $\|\cdot\|$ on \mathcal{M} in an alternative way, which includes the conclusion of Corollary 3.8.

4. INTERPOLATION OF NONCOMMUTATIVE L_p -SPACES.

Let \mathcal{M} be a (σ -finite) von Neumann algebra with a n.s.f. state ϕ . In this section, we will use the notation $L_p(\mathcal{M})$ for $L_p(\mathcal{M}, \phi)$ if there is no ambiguity. By embedding \mathcal{M}_a into $L_p(\mathcal{M})$ via the map $x \mapsto h_\phi^{1/2p} x h_\phi^{1/2p}$, we can define our interpolation space as follows. For $x \in \mathcal{M}_a$ Set

$$\mathcal{F}(L_{p_0}(\mathcal{M}), L_{p_1}(\mathcal{M})) := \left\{ f : f(z) = \sum_{k=1}^m f_k(z) x_k, x_k \in \mathcal{M}_a, f_k \in A(S), m \in \mathbb{N} \right\},$$

equipped with the norm

$$\|f\|_{\mathcal{F}(L_{p_0}(\mathcal{M}), L_{p_1}(\mathcal{M}))} = \sup_{t \in \mathbb{R}} \{ \|f(it)\|_{p_0, \phi}, \|f(1+it)\|_{p_1, \phi} \},$$

For $0 < \theta < 1$ we define the complex interpolation (quasi) norm $\|\cdot\|_\theta$ on \mathcal{M}_a as follows: for $x \in \mathcal{M}_a$,

$$\|x\|_\theta := \inf \{ \|f\|_{\mathcal{F}(L_{p_0}(\mathcal{M}), L_{p_1}(\mathcal{M}))} : f(\theta) = x, f \in \mathcal{F}(L_{p_0}(\mathcal{M}), L_{p_1}(\mathcal{M})) \}.$$

Then the complex interpolation space $[L_{p_0}(\mathcal{M}), L_{p_1}(\mathcal{M})]_\theta$ is defined as the completion of $(\mathcal{M}_a, \|\cdot\|_\theta)$. The following theorem extends Xu's result [37] to (non-tracial) state case.

Theorem 4.1. *Let \mathcal{M} be a (σ -finite) von Neumann algebra with a n.f. state ϕ . For $0 < p_0 < p_1 \leq \infty$, we have*

$$(4.1) \quad [L_{p_0}(\mathcal{M}), L_{p_1}(\mathcal{M})]_\theta = L_{p_\theta}(\mathcal{M}),$$

where $\frac{1}{p_\theta} = \frac{1-\theta}{p_0} + \frac{\theta}{p_1}$.

Proof. The case of $1 \leq p_0 < p_1 \leq \infty$ is basically Kosaki's interpolation theorem [22].

The proof for the case $0 < p_0 < 1$ is an adaption of Xu' result [37]. On one hand, we prove that

$$(4.2) \quad L_{p_\theta}(\mathcal{M}) \subseteq [L_{p_0}(\mathcal{M}), L_{p_1}(\mathcal{M})]_\theta.$$

For this it is sufficient to show $\|x\|_\theta \leq \|x\|_{p_\theta, \phi}$ for any $x \in \mathcal{M}_a$.

Firstly we consider the case $\frac{1}{2} \leq p_0 < 1$. To this end, using the polar decomposition, $x \in \mathcal{M}_a$ can be written as

$$x = u|x| = u|x|^{\frac{1}{2}}|x|^{\frac{1}{2}} = x_1 x_2,$$

where $x_1 = u|x|^{\frac{1}{2}}$ and $x_2 = |x|^{\frac{1}{2}}$. Since \mathcal{M}_a is w^* -dense in \mathcal{M} , for any $\epsilon > 0$ and $k = 0, 1$, there exists $y_k \in \mathcal{M}_a$ such that

$$\|x_k - y_k\| \leq \epsilon.$$

Thus

$$\left\| h_\phi^{\frac{1}{2p_\theta}} (x_1 - y_1) \right\|_{L_{2p_\theta}(\mathcal{M})} \leq \|x_1 - y_1\| \left\| h_\phi^{\frac{1}{2p_\theta}} \right\|_{L_{2p_\theta}(\mathcal{M})} = \|x_1 - y_1\| \leq \epsilon,$$

and similarly

$$\left\| (x_2 - y_2) h_\phi^{\frac{1}{2p_\theta}} \right\|_{L_{2p_\theta}(\mathcal{M})} \leq \|x_2 - y_2\| \leq \epsilon.$$

On the other hand, by (2.8), we have

$$\left\| h_\phi^{\frac{1}{2p_\theta}} x_1 \right\|_{L_{2p_\theta}(\mathcal{M})} = \left\| h_\phi^{\frac{1}{2p_\theta}} u|x|u^* h_\phi^{\frac{1}{2p_\theta}} \right\|_{L_{p_\theta}(\mathcal{M})}^{\frac{1}{2}} = \|x\|_{p_\theta, \phi}^{\frac{1}{2}},$$

and

$$\left\| x_2 h_\phi^{\frac{1}{2p_\theta}} \right\|_{L_{2p_\theta}(\mathcal{M})} = \left\| h_\phi^{\frac{1}{2p_\theta}} |x| h_\phi^{\frac{1}{2p_\theta}} \right\|_{L_{p_\theta}(\mathcal{M})}^{\frac{1}{2}} = \|x\|_{p_\theta, \phi}^{\frac{1}{2}}.$$

Therefore

$$\left\| h_\phi^{\frac{1}{2p_\theta}} y_1 \right\|_{L_{2p_\theta}(\mathcal{M})} \leq \|x\|_{p_\theta, \phi}^{\frac{1}{2}} + \epsilon,$$

and

$$\left\| y_2 h_\phi^{\frac{1}{2p_\theta}} \right\|_{L_{2p_\theta}(\mathcal{M})} \leq \|x\|_{p_\theta, \phi}^{\frac{1}{2}} + \epsilon.$$

Let $z_1 = \sigma_{-\frac{i}{4p_\theta}}^\phi(y_1)$ and $z_2 = \sigma_{\frac{i}{4p_\theta}}^\phi(y_2)$, then due to the equality (2.9) we have $h_{\frac{1}{\phi}}^{\frac{1}{4p_\theta}} z_1 h_{\frac{1}{\phi}}^{\frac{1}{4p_\theta}} = h_{\frac{1}{\phi}}^{\frac{1}{2p_\theta}} y_1$ and $h_{\frac{1}{\phi}}^{\frac{1}{4p_\theta}} z_2 h_{\frac{1}{\phi}}^{\frac{1}{4p_\theta}} = y_2 h_{\frac{1}{\phi}}^{\frac{1}{2p_\theta}}$. Moreover, $z_k \in \mathcal{M}_a$ and

$$\|z_k\|_{2p_\theta, \phi} \leq \|x\|_{p_\theta, \phi}^{\frac{1}{2}} + \epsilon, k = 1, 2.$$

By the first part, $[L_{2p_0}(\mathcal{M}), L_{2p_1}(\mathcal{M})]_\theta = L_{2p_\theta}(\mathcal{M})$. Then for any $\epsilon > 0$ and $k = 1, 2$, there exists $f_k \in \mathcal{F}(L_{2p_0}(\mathcal{M}), L_{2p_1}(\mathcal{M}))$, such that $f_k(\theta) = z_k$, and

$$\begin{aligned} \|f_k\|_{\mathcal{F}(L_{2p_0}(\mathcal{M}), L_{2p_1}(\mathcal{M}))} &\leq \|z_k\|_{2p_\theta, \phi} + \epsilon \\ &\leq \|x\|_{p_\theta, \phi}^{\frac{1}{2}} + \epsilon. \end{aligned}$$

Now consider

$$f(z) := \sigma_{\frac{i}{4p_z}}^\phi(f_1(z)) \sigma_{-\frac{i}{4p_z}}^\phi(f_2(z)) - y_1 y_2 + x.$$

It is easy to see that f is analytic in S , continuous on \bar{S} , and $f(\theta) = x$. Moreover, from (2.2) and Hölder's inequality (2.7) it follows that

$$\begin{aligned} \|f(it)\|_{p_0, \phi}^{p_0} &\leq \left\| h_{\frac{1}{\phi}}^{\frac{1}{2p_0}} \sigma_{\frac{i}{4p_0}}^\phi(f_1(it)) \sigma_{-\frac{i}{4p_0}}^\phi(f_2(it)) h_{\frac{1}{\phi}}^{\frac{1}{2p_0}} \right\|_{L_{p_0}(\mathcal{M})}^{p_0} + \|x - y_1 y_2\|_{p_0, \phi}^{p_0} \\ &\leq \left\| h_{\frac{1}{\phi}}^{\frac{1}{4p_0}} f_1(it) h_{\frac{1}{\phi}}^{\frac{1}{2p_0}} f_2(it) h_{\frac{1}{\phi}}^{\frac{1}{4p_0}} \right\|_{L_{p_0}(\mathcal{M})}^{p_0} + \|x - y_1 y_2\|_{p_0, \phi}^{p_0} \\ &\leq \|f_1(it)\|_{2p_0, \phi}^{p_0} \|f_2(it)\|_{2p_0, \phi}^{p_0} + \|x - y_1 y_2\|_{p_0, \phi}^{p_0} \\ &\leq \|x\|_{p_\theta, \phi}^{p_0} + \|x - y_1 y_2\|_{p_0, \phi}^{p_0} + O(\epsilon) \\ &\leq \|x\|_{p_\theta, \phi}^{p_0} + \|x - y_1 y_2\|_{p_0, \phi}^{p_0} + O(\epsilon) \\ &\leq \|x\|_{p_\theta, \phi}^{p_0} + O(\epsilon). \end{aligned}$$

Similarly we have (replace (2.2) with (2.1) if necessary)

$$\|f(1 + it)\|_{p_1, \phi} \leq \|x\|_{p_\theta, \phi} + O(\epsilon).$$

Therefore

$$\|x\|_\theta \leq \|f\|_{\mathcal{F}(L_{p_0}(\mathcal{M}), L_{p_1}(\mathcal{M}))} \leq \|x\|_{p_\theta, \phi} + O(\epsilon).$$

Thus we can conclude the inclusion (4.2) for $\frac{1}{2} \leq p_0 < 1$ by letting $\epsilon \rightarrow 0^+$.

By repeating the above arguments we can obtain the inclusion (4.2) for $\frac{1}{4} \leq p_0 < \frac{1}{2}$. Thus by iterating this procedure, we get the inclusion (4.2) for all $0 < p_0 < 1$.

On the other hand, we show that

$$(4.3) \quad [L_{p_0}(\mathcal{M}), L_{p_1}(\mathcal{M})]_\theta \subseteq L_{p_\theta}(\mathcal{M}).$$

To this end, it suffices to show that $\|x\|_{p_\theta, \phi} \leq \|x\|_\theta$ for any $x \in \mathcal{M}_a$. Thus for any $f \in \mathcal{F}(L_{p_0}(\mathcal{M}), L_{p_1}(\mathcal{M}))$ such that $f(\theta) = x$, we will show

$$\|x\|_{p_\theta, \phi} \leq \|f\|_{\mathcal{F}(L_{p_0}(\mathcal{M}), L_{p_1}(\mathcal{M}))}.$$

Without loss of generality, suppose $\|f\|_{\mathcal{F}(L_{p_0}(\mathcal{M}), L_{p_1}(\mathcal{M}))} \leq 1$. Therefore using Proposition 3.6, we have

$$\|x\|_{p_\theta, \phi} = \|f(\theta)\|_{p_\theta, \phi} \leq \|f(it)\|_{p_0, \phi}^{1-\theta} \|f(1+it)\|_{p_1, \phi}^\theta \leq 1,$$

which completes our proof. \square

Remark 4.2. (i) Instead of the symmetric embedding for \mathcal{M} into $L_p(\mathcal{M}, \phi)$, our interpolation also holds for any embedding $x \mapsto h_\phi^{\frac{1-\eta}{2p}} x h_\phi^{\frac{\eta}{2p}}$ as in Lemma 2.1, $0 \leq \eta \leq 1$.

(ii) The n.f. state ϕ can be replaced by a weight.

We have the following direct corollaries.

Corollary 4.3. *Let $0 < p_0 < p_1 \leq \infty$, and for $\theta \in [0, 1]$ define p_θ as in the equation (3.1). For any $x \in \mathcal{M}$ we have*

$$\left\| h_\phi^{\frac{1}{2p_\theta}} x h_\phi^{\frac{1}{2p_\theta}} \right\|_{L_{p_\theta}(\mathcal{M})} \leq \left\| h_\phi^{\frac{1}{2p_0}} x h_\phi^{\frac{1}{2p_0}} \right\|_{L_{p_0}(\mathcal{M})}^{1-\theta} \left\| h_\phi^{\frac{1}{2p_1}} x h_\phi^{\frac{1}{2p_1}} \right\|_{L_{p_1}(\mathcal{M})}^\theta.$$

Proof. By taking the map $G(z) = x$ in Proposition 3.6 we can deduce our result. \square

Let \mathcal{M}, \mathcal{N} be two (σ -finite) von Neumann algebras, and ϕ (resp. ψ) be a n.f. state on \mathcal{M} (resp. \mathcal{N}). Let $T : \mathcal{M} \rightarrow \mathcal{N}$ be a linear map. Then we define an operator norm $\|T\|_{(p, \phi) \rightarrow (q, \psi)}$ by

$$\|T\|_{(p, \phi) \rightarrow (q, \psi)} = \sup_{X \neq 0} \frac{\|T(X)\|_{q, \psi}}{\|X\|_{p, \phi}}.$$

Corollary 4.4. *Let \mathcal{M}, \mathcal{N} be two (σ -finite) von Neumann algebras, and ϕ (resp. ψ) be a n.s.f. state on \mathcal{M} (resp. \mathcal{N}). Let $T : \mathcal{M} \rightarrow \mathcal{N}$ be a linear map. Assume that $0 < p_0 < p_1 \leq \infty, 0 < q_0 < q_1 \leq \infty$ and p_θ, q_θ satisfy the equation (3.1). Then we have*

$$\|T\|_{(p_\theta, \phi) \rightarrow (q_\theta, \psi)} \leq \|T\|_{(p_0, \phi) \rightarrow (q_0, \psi)}^{1-\theta} \cdot \|T\|_{(p_1, \phi) \rightarrow (q_1, \psi)}^\theta.$$

Proof. It is a direct corollary of Theorem 4.1 using the standard approach in the interpolation theory [1]. \square

5. APPLICATIONS TO THE SANDWICHED RÉNYI RELATIVE ENTROPY

The history of the generalization of p -Rényi divergence can be traced back to Petz's work [27, 28], where he extended the notion of the p -Rényi divergence to the general von Neumann algebra context. Regarding to the sandwiched p -Rényi divergence, there are mainly two approaches.

One is introduced by Jencová [20, 21] as follows

$$(5.1) \quad \tilde{D}_p^J(\psi||\phi) := \frac{1}{p-1} \log \|h_\psi\|_{C_p(\mathcal{M}, \phi)}^p, \quad p \in (1, \infty),$$

where $C_p(\mathcal{M}, \phi)$ is the Kosaki's noncommutative L_p -space associated with (\mathcal{M}, ϕ) . Recall that

$$C_p(\mathcal{M}, \phi) := \left[h_\phi^{\frac{1}{2}} \mathcal{M} h_\phi^{\frac{1}{2}}, L_1(\mathcal{M}, \phi) \right]_{1/p},$$

where $1 \leq p \leq \infty$. Note that $C_\infty(\mathcal{M}, \phi) = h_\phi^{\frac{1}{2}} \mathcal{M} h_\phi^{\frac{1}{2}}$, and for any $x \in \mathcal{M}$, $\left\| h_\phi^{\frac{1}{2}} x h_\phi^{\frac{1}{2}} \right\|_{C_\infty(\mathcal{M}, \phi)} = \|x\|$. Since the definition (5.1) relies on Kosaki's construction of noncommutative L_p -space, the index p can not go beyond the interval $(1, \infty)$.

Another definition is derived by Berta, Scholz and Tomamichel [9] via a so-called *noncommutative vector valued L_p -space*, denoted by $L_p^{BST}(\mathcal{M}, \phi)$. The definition of spaces $L_p^{BST}(\mathcal{M}, \phi)$ relies on the *spatial derivative* $\Delta(\xi/\phi)$, where ξ is a vector in $L_2(\mathcal{M}, \phi)$. The definition of the space $L_p^{BST}(\mathcal{M}, \phi)$ is subtle. We omit the details and refer to [9]. They defined the divergence as follows

$$(5.2) \quad \tilde{D}_p^{BST}(\psi||\phi) := \frac{2p}{p-1} \log \|\xi_\psi\|_{L_{2p}^{BST}(\mathcal{M}, \phi)}, \quad p \in [1/2, 1) \cup (1, \infty),$$

where ξ_ψ is a vector representation of ψ for a $*$ -representation $\pi : \mathcal{M} \rightarrow B(H)$.

Now let \mathcal{M} be a σ -finite von Neumann algebra acting on H , and ϕ, ψ be two normal faithful states on \mathcal{M} . For $p \in (0, 1) \cup (1, \infty)$ we define

$$(5.3) \quad \begin{aligned} \tilde{D}_p(\psi||\phi) &:= \frac{1}{p-1} \log \left\| h_\phi^{\frac{1-p}{2p}} h_\psi h_\phi^{\frac{1-p}{2p}} \right\|_{L_p(\mathcal{M}, \phi)}^p \\ &= \frac{1}{p-1} \log \left\| h_\phi^{-\frac{1}{2}} h_\psi h_\phi^{-\frac{1}{2}} \right\|_{p, \phi}^p, \end{aligned}$$

where h_ψ and h_ϕ are Randon-Nikodym derivatives of dual weights $\hat{\psi}$ and $\hat{\phi}$ with respect to the canonical trace τ on $\mathcal{M} \rtimes_{\sigma\phi} \mathbb{R}$, respectively.

Using Lemma 2.2, we can show that the definition of (5.3) extends the notion of the sandwiched p -Rényi divergence to the σ -finite von Neumann algebra case. We recall that for two positive operators $\sigma, \rho \in B(H)$, the sandwiched p -Rényi divergence is defined as [25]

$$(5.4) \quad \tilde{D}_p(\rho||\sigma) = \frac{1}{p-1} \log \left\| \sigma^{\frac{1-p}{2p}} \rho \sigma^{\frac{1-p}{2p}} \right\|_p^p, \quad p \in (0, 1) \cup (1, \infty),$$

where $\|\cdot\|_p$ is the Schatten p -norm on $B(H)$.

Proposition 5.1. *Let \mathcal{M} be a finite von Neumann algebra with a n.f. tracial state τ , and ψ, ϕ be two n.f. states on \mathcal{M} defined by*

$$\psi(x) = \tau(\rho x), \quad \phi(x) = \tau(\sigma x), \quad x \in \mathcal{M},$$

where ρ and σ are two positive operators with unit trace. Then

$$(5.5) \quad \tilde{D}_p(\psi||\phi) = \frac{1}{p-1} \log \tau \left[\left(\sigma^{\frac{1-p}{2p}} \rho \sigma^{\frac{1-p}{2p}} \right)^p \right].$$

Proof. By Lemma 2.2, there exists an isometry $\theta' : \Lambda_p(\mathcal{M}, \tau) \rightarrow L_p(\mathcal{M}, \phi)$ such that

$$\theta'(\sigma) = h_\phi, \quad \theta'(\rho) = h_\psi.$$

Then we have

$$\begin{aligned} \left\| h_\phi^{\frac{1-p}{2p}} h_\psi h_\phi^{\frac{1-p}{2p}} \right\|_{L_p(\mathcal{M}, \phi)}^p &= \left\| \theta' \left(\sigma^{\frac{1-p}{2p}} \rho \sigma^{\frac{1-p}{2p}} \right) \right\|_{L_p(\mathcal{M}, \phi)}^p \\ &= \left\| \sigma^{\frac{1-p}{2p}} \rho \sigma^{\frac{1-p}{2p}} \right\|_{\Lambda_p(\mathcal{M}, \tau)}^p \\ &= \tau \left[\left(\sigma^{\frac{1-p}{2p}} \rho \sigma^{\frac{1-p}{2p}} \right)^p \right], \end{aligned}$$

which completes our proof. \square

Proposition 5.2. *Our definition $\tilde{D}_p(\psi||\phi)$ fits well with the ones defined by Jenčová and Berta et al.. More precisely, we have*

- (i) $\tilde{D}_p(\psi||\phi) = \tilde{D}_p^J(\psi||\phi)$ for $1 < p < \infty$;
- (ii) $\tilde{D}_p(\psi||\phi) = \tilde{D}_p^{BST}(\psi||\phi)$ for $1/2 \leq p < 1$.

Proof. We note that (ii) has been proved by Jenčová in [20, Theorem 5].

For (i), we will use the following fact proved by Kosaki [22]: For $1 \leq p \leq \infty$ and $1/p + 1/p' = 1$, the map $i_p : x \mapsto h_\phi^{1/2p'} x h_\phi^{1/2p'}$ is an isometric isomorphism from $L_p(\mathcal{M}, \phi)$ onto $C_p(\mathcal{M}, \phi)$. From this it follows that

$$(5.6) \quad \left\| h_\phi^{\frac{1-p}{2p}} h_\psi h_\phi^{\frac{1-p}{2p}} \right\|_{L_p(\mathcal{M}, \phi)} = \left\| i_p \left(h_\phi^{\frac{1-p}{2p}} h_\psi h_\phi^{\frac{1-p}{2p}} \right) \right\|_{C_p(\mathcal{M}, \phi)} = \|h_\psi\|_{C_p(\mathcal{M}, \phi)}.$$

Hence for $1 < p < \infty$, the equation (5.3) coincides with Jenčová's definition, i.e.,

$$(5.7) \quad \tilde{D}_p(\psi||\phi) = \frac{1}{p-1} \log \|h_\psi\|_{C_p(\mathcal{M}, \phi)}^p = \tilde{D}_p^J(\psi||\phi).$$

\square

Proposition 5.3. *(Monotonicity in p). For all n.f. states ψ, ϕ on \mathcal{M} , the function $p \mapsto \tilde{D}_p(\psi||\phi)$ is increasing for $p \in (0, 1) \cup (1, \infty)$.*

Proof. For $p \in (1, \infty)$, by a direct modification of the proof in [3, Theorem 7.1] we can deduce our result. It was also proved by Jenčová [21].

Since the logarithm function is non-decreasing, it remains to show that when $0 < p_0 < p_1 < 1$ or $0 < p_0 < 1 < p_1$, we have

$$(5.8) \quad \left\| h_\phi^{\frac{1-p_0}{2p_0}} h_\psi h_\phi^{\frac{1-p_0}{2p_0}} \right\|_{L_{p_0}(\mathcal{M}, \phi)}^{p'_0} \leq \left\| h_\phi^{\frac{1-p_1}{2p_1}} h_\psi h_\phi^{\frac{1-p_1}{2p_1}} \right\|_{L_{p_1}(\mathcal{M}, \phi)}^{p'_1}.$$

Here we use s' to denote the conjugate number of s . Note that for $s \in (0, 1)$, $s' = \frac{s}{s-1} < 0$.

If $0 < p_0 < p_1 < 1$, then there exists $0 < \theta < 1$ such that

$$(5.9) \quad \frac{1}{p_1} = \frac{1-\theta}{p_0} + \frac{\theta}{1}.$$

Using Corollary 4.3 (by letting $x = h_\phi^{-\frac{1}{2}} h_\psi h_\phi^{-\frac{1}{2}}$) we have

$$\left\| h_\phi^{\frac{1-p_1}{2p_1}} h_\psi h_\phi^{\frac{1-p_1}{2p_1}} \right\|_{L_{p_1}(\mathcal{M}, \phi)} \leq \left\| h_\phi^{\frac{1-p_0}{2p_0}} h_\psi h_\phi^{\frac{1-p_0}{2p_0}} \right\|_{L_{p_0}(\mathcal{M}, \phi)}^{1-\theta} \|h_\psi\|_{L_1(\mathcal{M}, \phi)}^\theta.$$

Since $\|h_\psi\|_{L_1(\mathcal{M}, \phi)} = 1$ and $p'_1 < 0$, we have

$$\left\| h_\phi^{\frac{1-p_1}{2p_1}} h_\psi h_\phi^{\frac{1-p_1}{2p_1}} \right\|_{L_{p_1}(\mathcal{M}, \phi)}^{p'_1} \geq \left\| h_\phi^{\frac{1-p_0}{2p_0}} h_\psi h_\phi^{\frac{1-p_0}{2p_0}} \right\|_{L_{p_0}(\mathcal{M}, \phi)}^{(1-\theta)p'_1}.$$

By equation (5.9), we have $(1-\theta)p'_1 = p'_0$, which gives (5.8).

The proof for $0 < p_0 < 1 < p_1$ is similar. In this case there exists $0 < \theta < 1$ such that

$$(5.10) \quad \frac{1}{1} = \frac{1-\theta}{p_0} + \frac{\theta}{p_1}.$$

Using Corollary 4.3 we have

$$1 = \|h_\psi\|_{L_1(\mathcal{M}, \phi)} \leq \left\| h_\phi^{\frac{1-p_0}{2p_0}} h_\psi h_\phi^{\frac{1-p_0}{2p_0}} \right\|_{L_{p_0}(\mathcal{M}, \phi)}^{1-\theta} \left\| h_\phi^{\frac{1-p_1}{2p_1}} h_\psi h_\phi^{\frac{1-p_1}{2p_1}} \right\|_{L_{p_1}(\mathcal{M}, \phi)}^\theta.$$

Recall that $p'_0 < 0$, then

$$\left\| h_\phi^{\frac{1-p_0}{2p_0}} h_\psi h_\phi^{\frac{1-p_0}{2p_0}} \right\|_{L_{p_0}(\mathcal{M}, \phi)}^{p'_0} \leq \left\| h_\phi^{\frac{1-p_1}{2p_1}} h_\psi h_\phi^{\frac{1-p_1}{2p_1}} \right\|_{L_{p_1}(\mathcal{M}, \phi)}^{\frac{\theta}{\theta-1} p'_0}.$$

From (5.10) we have $(\theta-1)p'_1 = \theta p'_0$ and then obtain (5.8). \square

We fix some notations that will be used in the remaining part of the paper. Let \mathcal{M} be a σ -finite von Neumann algebra with a normal faithful state ϕ . Let \mathcal{N} be a von Neumann subalgebra of \mathcal{M} and set $\phi_{\mathcal{N}} := \phi|_{\mathcal{N}}$. Let τ (resp. $\tau_{\mathcal{N}}$) be the canonical trace on $\mathcal{R} := \mathcal{M} \rtimes_{\sigma\phi} \mathbb{R}$ (resp. $\mathcal{R}_{\mathcal{N}} := \mathcal{N} \rtimes_{\sigma\phi_{\mathcal{N}}} \mathbb{R}$). Let h_ψ (resp. $h_{\psi_{\mathcal{N}}}$) be the Randon-Nikodym derivative of dual weight $\hat{\phi}$ (resp. $\hat{\phi}_{\mathcal{N}}$) with respect to τ (resp. $\tau_{\mathcal{N}}$).

Theorem 5.4. *Let \mathcal{N} be a von Neumann subalgebra of \mathcal{M} and ψ, ϕ be n.f. states on \mathcal{M} . Then for $p \in (1, \infty)$, we have the following data processing inequality (DPI):*

$$\tilde{D}_p(\psi_{\mathcal{N}} \parallel \phi_{\mathcal{N}}) \leq \tilde{D}_p(\psi \parallel \phi).$$

Proof. Basically it is an adaption of Beigi's proof [3]. It is sufficient to prove that

$$(5.11) \quad \left\| h_{\phi_{\mathcal{N}}}^{\frac{1-p}{2p}} h_{\psi_{\mathcal{N}}} h_{\phi_{\mathcal{N}}}^{\frac{1-p}{2p}} \right\|_{L_p(\mathcal{N}, \phi_{\mathcal{N}})} \leq \left\| h_{\phi}^{\frac{1-p}{2p}} h_{\psi} h_{\phi}^{\frac{1-p}{2p}} \right\|_{L_p(\mathcal{M}, \phi)},$$

for $1 < p < \infty$.

Recall that $\mathcal{M}_* = L_1(\mathcal{M}, \phi)$. Then for any $x = h_{\omega} \in L_1(\mathcal{M}, \phi)$ with $\omega \in \mathcal{M}_*$, define a map $T : L_1(\mathcal{M}, \phi) \rightarrow L_1(\mathcal{N}, \phi_{\mathcal{N}})$ by

$$T(x) := h_{\omega_{\mathcal{N}}}.$$

It is easy to see that T is positive and tr-preserving on $L_1(\mathcal{M}, \phi)$. Consider the map $S : L_1(\mathcal{M}, \phi) \rightarrow L_1(\mathcal{N}, \phi_{\mathcal{N}})$ given by

$$S(x) = h_{\phi_{\mathcal{N}}}^{-\frac{1}{2}} T \left(h_{\phi}^{\frac{1}{2}} x h_{\phi}^{\frac{1}{2}} \right) h_{\phi_{\mathcal{N}}}^{-\frac{1}{2}}.$$

Clearly $S \left(h_{\phi}^{-\frac{1}{2}} h_{\psi} h_{\phi}^{-\frac{1}{2}} \right) = h_{\phi_{\mathcal{N}}}^{-\frac{1}{2}} h_{\psi_{\mathcal{N}}} h_{\phi_{\mathcal{N}}}^{-\frac{1}{2}}$. Using Corollary 4.4, to show (5.11) it is sufficient to prove that

$$\|S\|_{(p, \phi) \rightarrow (p, \phi_{\mathcal{N}})} \leq 1,$$

for $p = 1$ and $p = \infty$.

For $p = 1$, since T is tr-preserving, we have

$$(5.12) \quad \|S(x)\|_{1, \phi_{\mathcal{N}}} = \left\| T \left(h_{\phi}^{\frac{1}{2}} x h_{\phi}^{\frac{1}{2}} \right) \right\|_{1, \phi_{\mathcal{N}}} = \left\| h_{\phi}^{\frac{1}{2}} x h_{\phi}^{\frac{1}{2}} \right\|_{L_1(\mathcal{M}, \phi)} = \|x\|_{1, \phi}.$$

For $p = \infty$, using the positivity of T , we have

$$\|S\|_{(\infty, \phi) \rightarrow (\infty, \phi_{\mathcal{N}})} = \|S(1)\|_{\mathcal{N}} = \|h_{\phi_{\mathcal{N}}}^{-\frac{1}{2}} h_{\phi_{\mathcal{N}}} h_{\phi_{\mathcal{N}}}^{-\frac{1}{2}}\|_{\mathcal{N}} = 1. \quad \square$$

Remark 5.5. It is known that [12] the sandwiched p -Rényi divergence satisfies DPI for $p \in (\frac{1}{2}, 1) \cup (1, \infty)$. DPI for a more general family of p -Rényi divergences, knowns as α - z Rényi relative entropies, was determined by the last author in [38]. See also a survey paper [10] for more information of DPI.

Suppose \mathcal{N} is invariant under the automorphism group σ_t^{ϕ} , i.e., $\sigma_t^{\phi}(\mathcal{N}) = \mathcal{N}$ for all $t \in \mathbb{R}$. Then due to the modular theory [36], there exists a conditional expectation $\mathcal{E} : \mathcal{M} \rightarrow \mathcal{N}$ such that $\phi \circ \mathcal{E} = \phi$. Moreover [14, Theorem 4.1], \mathcal{E} can be naturally extended to a conditional expectation $\hat{\mathcal{E}} : \mathcal{R} \rightarrow \mathcal{R}_{\mathcal{N}}$ satisfying

$$(5.13) \quad \tau_{\mathcal{N}}[\hat{\mathcal{E}}(y)] = \tau(y),$$

for any $y \in \mathcal{R}$.

Denote by $\iota : \mathcal{R}_{\mathcal{N}} \rightarrow \mathcal{R}$ the natural embedding. By (5.13), we have

$$(5.14) \quad \tau[\iota(x)y] = \tau_{\mathcal{N}}[\hat{\mathcal{E}}(\iota(x)y)] = \tau_{\mathcal{N}}[\hat{\mathcal{E}}(xy)] = \tau_{\mathcal{N}}[x\hat{\mathcal{E}}(y)],$$

for any $x \in \mathcal{R}_{\mathcal{N}}, y \in \mathcal{R}$.

Proposition 5.6. *Let \mathcal{N} be a von Neumann subalgebra of \mathcal{M} and ψ, ϕ be n.f. states on \mathcal{M} . Suppose that $\sigma_t^\phi(\mathcal{N}) = \mathcal{N}$ for all $t \in \mathbb{R}$, and let $\mathcal{E} : \mathcal{M} \rightarrow \mathcal{N}$ be the ϕ -preserving conditional expectation. Then we have*

$$\hat{\mathcal{E}}(h_\psi) = h_{\psi_{\mathcal{N}}}.$$

Proof. Since $\phi \circ \mathcal{E} = \phi$, we have $\mathcal{E} \circ \sigma_t^\phi = \sigma_t^{\phi_{\mathcal{N}}} \circ \mathcal{E}$. Thus by a result of Haagerup, Junge and Xu [14, Theorem 4.1], we have $\hat{\mathcal{E}} \circ \hat{\sigma}_t^\phi = \hat{\sigma}_t^{\phi_{\mathcal{N}}} \circ \hat{\mathcal{E}}$. It follows immediately that

$$\hat{\sigma}_t^\phi|_{\mathcal{N}} = \hat{\sigma}_t^{\phi_{\mathcal{N}}}.$$

Therefore for any $x \in \mathcal{R}_{\mathcal{N}}, x \geq 0$,

$$\begin{aligned} \tau_{\mathcal{N}}[\hat{\mathcal{E}}(h_\psi)x] &= \tau(h_\psi x) \\ &= \psi \left[\int_{-\infty}^{\infty} \hat{\sigma}_t^\phi(x) dt \right] = \psi \left[\int_{-\infty}^{\infty} \hat{\mathcal{E}} \circ \hat{\sigma}_t^{\phi_{\mathcal{N}}}(x) dt \right] \\ &= \psi_{\mathcal{N}} \left[\int_{-\infty}^{\infty} \hat{\sigma}_t^{\phi_{\mathcal{N}}}(x) dt \right] = \tau_{\mathcal{N}}[h_{\psi_{\mathcal{N}}}x]. \end{aligned}$$

This completes our proof, since $\tau_{\mathcal{N}}$ is faithful. \square

Proposition 5.7. *Let \mathcal{N} be a von Neumann subalgebra of \mathcal{M} and ψ, ϕ be n.f. states on \mathcal{M} . Suppose that $\sigma_t^\phi(\mathcal{N}) = \mathcal{N}$ for all $t \in \mathbb{R}$, and let $\mathcal{E} : \mathcal{M} \rightarrow \mathcal{N}$ be the ϕ -preserving conditional expectation. Then we have*

$$\iota(h_{\psi_{\mathcal{N}}}) = h_{\psi_{\mathcal{N}} \circ \mathcal{E}}.$$

Proof. Since $\mathcal{E} \circ \sigma_t^\phi = \sigma_t^{\phi_{\mathcal{N}}} \circ \mathcal{E}$, we have by [14, Theorem 4.1] that

$$\hat{\mathcal{E}} \circ \hat{\sigma}_t^\phi = \hat{\sigma}_t^{\phi_{\mathcal{N}}} \circ \hat{\mathcal{E}}.$$

Therefore for any $y \in \mathcal{R}, y \geq 0$,

$$\begin{aligned} \tau[h_{\psi_{\mathcal{N}} \circ \mathcal{E}}y] &= \psi_{\mathcal{N}} \circ \mathcal{E} \left[\int_{-\infty}^{\infty} \hat{\sigma}_t^\phi(y) dt \right] \\ &= \psi_{\mathcal{N}} \left[\int_{-\infty}^{\infty} \hat{\mathcal{E}} \circ \hat{\sigma}_t^\phi(y) dt \right] \\ &= \psi_{\mathcal{N}} \left[\int_{-\infty}^{\infty} \hat{\sigma}_t^{\phi_{\mathcal{N}}} \circ \hat{\mathcal{E}}(y) dt \right] \\ &= \tau_{\mathcal{N}}[h_{\psi_{\mathcal{N}}} \hat{\mathcal{E}}(y)] = \tau[\iota(h_{\psi_{\mathcal{N}}})(y)], \end{aligned}$$

where the last equality uses (5.14). This completes our proof, since τ is faithful. \square

Theorem 5.8. *Let \mathcal{N} be a von Neumann subalgebra of \mathcal{M} and ψ, ϕ be n.f. states on \mathcal{M} . Suppose that $\sigma_t^\phi(\mathcal{N}) = \mathcal{N}$, and $\mathcal{E} : \mathcal{M} \rightarrow \mathcal{N}$ is the induced conditional expectation. If $\psi_{\mathcal{N}} \circ \mathcal{E} = \psi$, then we have*

$$\tilde{D}_p(\psi_{\mathcal{N}} \|\phi_{\mathcal{N}}) = \tilde{D}_p(\psi \|\phi), \quad p \in (0, 1) \cup (1, \infty).$$

Proof. Obviously we have $\phi_{\mathcal{N}} \circ \mathcal{E} = \phi \circ \mathcal{E} = \phi$. Thus using Proposition 5.7 and the fact that ι naturally embeds $L_p(\mathcal{N}, \phi_{\mathcal{N}})$ into $L_p(\mathcal{M}, \phi)$ for all $0 < p < \infty$ [14], we have

$$\begin{aligned} \left\| h_{\phi_{\mathcal{N}}}^{\frac{1-p}{2p}} h_{\psi_{\mathcal{N}}} h_{\phi_{\mathcal{N}}}^{\frac{1-p}{2p}} \right\|_{L_p(\mathcal{N}, \phi_{\mathcal{N}})} &= \left\| \iota \left(h_{\phi_{\mathcal{N}}}^{\frac{1-p}{2p}} h_{\psi_{\mathcal{N}}} h_{\phi_{\mathcal{N}}}^{\frac{1-p}{2p}} \right) \right\|_{L_p(\mathcal{M}, \phi)} \\ &= \left\| h_{\phi_{\mathcal{N}} \circ \mathcal{E}}^{\frac{1-p}{2p}} h_{\psi_{\mathcal{N}} \circ \mathcal{E}} h_{\phi_{\mathcal{N}} \circ \mathcal{E}}^{\frac{1-p}{2p}} \right\|_{L_p(\mathcal{M}, \phi)} \\ &= \left\| h_{\phi}^{\frac{1-p}{2p}} h_{\psi} h_{\phi}^{\frac{1-p}{2p}} \right\|_{L_p(\mathcal{M}, \phi)}. \quad \square \end{aligned}$$

Acknowledgements. We would like to thank Quanhua Xu and Simeng Wang for helpful comments and discussions. JG and ZY are partially supported by NSFC No. 11771106, No. 11431011 and No. 11826012. HZ is partially supported by the French project ISITE-BFC (contract ANR-15-IDEX-03) and the NCN (National Centre of Science) grant 2014/14/E/ST1/00525.

REFERENCES

- [1] J. Bergh, J. Löfström, *Interpolation spaces: an introduction*, Springer (1976).
- [2] R. Bhatia, *Matrix analysis*, Springer (2010).
- [3] S. Beigi, *Sandwiched Rényi divergence satisfies data processing inequality*, J. Math. Phys. **54** (2013) 122202.
- [4] D. P. Blecher, L. E. Labuschagne, *Applications of the Fuglede-Kadison Determinant: Szegő's Theorem and Outers for Noncommutative H^p* , Trans. Amer. Math. Soc. **360** (2008) 6131-6147.
- [5] T. N. Bekjan, Q. Xu, *Riesz and Szegő type factorizations for noncommutative Hardy spaces*, J. Operat. Theor. **62** (2009) 215-231.
- [6] T. N. Bekjan, Z. Chen, *Interpolation and Φ -moment inequalities of noncommutative martingales*, Probab Theory Relat Fields. **152** (2012) 179-206.
- [7] T. N. Bekjan, Z. Chen, A. Osekowski, *Noncommutative maximal inequalities associated with convex functions*, Trans. Amer. Math. Soc. **369** (2017) 409-427.
- [8] T. N. Bekjan, K. N. Ospanov, *Complex interpolation of noncommutative Hardy spaces associated semifinite von Neumann algebra*, arXiv:1711.01532 (2017).
- [9] M. Berta, V. B. Scholz, M. Tomamichel, *Rényi divergences as weighted non-commutative vector valued L_p spaces*, Ann. Henri Poincaré **19** (2018) 1843-1867.

- [10] E. A. Carlen, R. L. Frank, E. H. Lieb. *Inequalities for quantum divergences and the Audenaert-Datta conjecture*, J. Phys. A. **51** (2018) 483001.
- [11] J. Dixmier. *Formes linéaires sur un anneau d'opérateurs*, Bull. Soc. Math. France, **81** (1953) 9–39.
- [12] R. L. Frank, E. H. Lieb. *Monotonicity of a relative Rényi entropy*, J. Math. Phys. **54** (2013) 122201.
- [13] I. I. Hirschman, *A convexity theorem for certain groups of transformations*, J. Anal. Math. **2** (1952) 209-218.
- [14] U. Haagerup, M. Junge, Q. Xu, *A reduction method for noncommutative L_p spaces and applications*, Trans. Amer. Math. Soc. **362** (2010) 2125-2165.
- [15] U. Haagerup, “ *L_p spaces associated with an arbitrary von Neumann algebra*” in *Algèbres d'opérateurs et leurs applications en physique mathématique (Marseille, 1977)*, Proc. Colloq. CNRS 274, CNRS, Paris, (1979) 175-184.
- [16] F. Hiai, R. König, M. Tomamichel, *Generalized log-majorization and multivariate trace inequalities*. Ann. Henri Poincaré. **18** (2017), 2499-2521.
- [17] M. Junge, Q. Xu, *Noncommutative Burkholder/Rosenthal inequalities*, Ann. Probab. **31** (2003) 948-995.
- [18] M. Junge, Q. Xu, *Noncommutative maximal ergodic theorems*, J. Amer. Math. Soc. **20** (2007) 385-439.
- [19] M. Junge, R. Renner, D. Sutter, M. M. Wilde, A. Winter, *Universal recovery from a decrease of quantum relative entropy*, Ann. Henri Poincaré. **19** (2018) 2955-2978.
- [20] A. Jenčová, *Rényi relative entropies and noncommutative L_p spaces II*, arXiv:1707.00047 (2017).
- [21] A. Jenčová, *Rényi relative entropies and noncommutative L_p spaces*, Ann. Henri Poincaré. **19** (2018) 2513-2542.
- [22] H. Kosaki, *Applications of the complex interpolation method to a von Neumann algebra: non-commutative L_p spaces*, J. Funct. Anal. **56** (1984) 26-78.
- [23] M. J. Kastoryano, K. Temme, *Quantum logarithmic Sobolev inequalities and rapid mixing*, J. Math. Phys. **54** (2013) 052202.
- [24] T. Mei, *Operator valued Hardy spaces*, Mem. Amer. Math. Soc. **188** (2007).
- [25] M. Müller Lennert, F. Dupuis, O. Szehr, S. Fehr, M. Tomamichel, *On quantum Rényi entropies: a new generalization and some properties*, J. Math. Phys. **54** (2013) 122203.
- [26] R. Olkiewicz, B. Zegarlinski, *Hypercontractivity in noncommutative L_p spaces*, J. Funct. Anal. **161** (1999) 246-285.
- [27] D. Petz, *Quasi-entropies for finite quantum systems*, Rep. Math. Phys. **23** (1986) 57-65.
- [28] M. Ohya, D. Petz, *Quantum entropy and its use*, Springer (2004).

- [29] G. Pisier, *Interpolation between H_p spaces and non-commutative generalizations I*, Pacific J. Math. **155** (1992) 341-368.
- [30] G. Pisier, Q. Xu, *Non-commutative martingale inequalities*, Comm. Math. Phys. **189** (1997) 667-698.
- [31] G. Pisier, Q. Xu, *Noncommutative L^p -spaces. Handbook of the geometry of Banach spaces Vol. 2*, ed. W.B.Johnson and J.Lindenstrauss, 2003, 1459-1517, North-Holland, Amsterdam.
- [32] E. Ricard, Q. Xu, *Complex interpolation of weighted noncommutative L_p spaces*, Houston J. Math. **37** (2011) 1165-1179.
- [33] I. E. Segal. *A non-commutative extension of abstract integration*, Ann. Math. **57** (1953) 401-457.
- [34] D. Sutter, M. Berta, M. Tomamichel, *Multivariate trace inequalities*, Commun. Math. Phys. **352** (2017) 37-58.
- [35] M. Terp, *L_p spaces associated with von Neumann algebras*, Notes, Math. Institute, Copenhagen Univ. (1981).
- [36] M. Takesaki, *Theory of operator algebras. II*, volume **125** of Encyclopaedia of Mathematical Sciences. Springer-Verlag, Berlin, (2003).
- [37] Q. Xu, *Applications du théorème de factorisation pour des fonctions à valeurs opérateurs*, Studia Math. **95** (1990) 273-292.
- [38] H. Zhang, *Carlen-Frank-Lieb conjecture and monotonicity of α -z Rényi relative entropy*, arXiv:1811.01205. (2018)

DEPARTMENT OF MATHEMATICS, HARBIN INSTITUTE OF TECHNOLOGY, HARBIN 150006, CHINA, AND DEPARTMENT OF MATHEMATICS, HEILONGJIANG UNIVERSITY OF SCIENCE AND TECHNOLOGY, HARBIN, 150022, CHINA
E-mail address: gujuan-1977@163.com

INSTITUTE OF ADVANCED STUDY IN MATHEMATICS, HARBIN INSTITUTE OF TECHNOLOGY, HARBIN 150006, CHINA
E-mail address: hustyinzhzhi@163.com

LABORATOIRE DE MATHÉMATIQUES, UNIVERSITÉ DE FRANCHE-COMTÉ, 25030 BESANÇON, FRANCE AND INSTITUTE OF MATHEMATICS, POLISH ACADEMY OF SCIENCES, UL. ŚNIADECKICH 8, 00-656 WARSZAWA, POLAND
E-mail address: haonan.zhang@edu.univ-fcomte.fr