

# One-Shot Quantum State Exchange

Yonghae Lee,<sup>1,\*</sup> Hayata Yamasaki,<sup>2,†</sup> Gerardo Adesso,<sup>3,‡</sup> and Soojoon Lee<sup>1,3,§</sup>

<sup>1</sup> Department of Mathematics and Research Institute for Basic Sciences, Kyung Hee University, Seoul 02447, Korea

<sup>2</sup> Photon Science Center, Graduate School of Engineering,  
The University of Tokyo, Bunkyo-ku, Tokyo 113-8656, Japan

<sup>3</sup> School of Mathematical Sciences and Centre for the Mathematics and Theoretical Physics of Quantum Non-Equilibrium Systems,  
University of Nottingham, University Park, Nottingham NG7 2RD, United Kingdom

(Dated: March 18, 2022)

The quantum state exchange is a quantum communication task in which two users exchange their respective quantum information in the asymptotic setting. In this work, we consider a one-shot version of the quantum state exchange task, in which the users hold a single copy of the initial state, and they exchange their parts of the initial state by means of entanglement-assisted local operations and classical communication. We first derive lower bounds on the least amount of entanglement required for carrying out this task, and provide conditions on the initial state such that the protocol succeeds with zero entanglement cost. Based on these results, we reveal two counter-intuitive phenomena in this task, which make it different from a conventional SWAP operation. One tells how the users deal with their symmetric information in order to reduce the entanglement cost. The other shows that it is possible for the users to gain extra shared entanglement after this task.

PACS numbers: 03.67.Hk, 89.70.Cf, 03.67.Mn

**Introduction.**— In quantum information theory, the quantum state exchange [1, 2] is a quantum communication task, in which two users, Alice and Bob, exchange their quantum information by means of local operations and classical communication (LOCC) assisted by shared entanglement. A main research aim in the study of the quantum state exchange is to evaluate the least amount of entanglement needed for the task, as in other quantum communication tasks, such as quantum state merging [3, 4] and quantum state redistribution [5, 6].

Most quantum communication tasks [3–8] including the quantum state exchange usually assume the *asymptotic scenario*, in which users can have an unbounded number of independent and identically distributed copies of an initial state, and they carry out their task with the copies. On the other hand, it is not easy in a realistic situation to prepare a sufficiently large number of state copies, and the amount of non-local resources available for the users is limited. To reflect these practical difficulties, quantum information research has focused more recently on the *one-shot scenario* [9–17].

In this work, we introduce and study the *one-shot quantum state exchange* (OSQSE) task. This is not only a useful quantum communication task, but can also have a potential application in quantum computation. Let us consider a specific situation as follows. Alice and Bob want to carry out the SWAP gate [18], which plays an important role in universal quantum computation [19]. The problem is that they cannot directly apply the SWAP gate, because they are far apart. If Alice and Bob are sharing prior entanglement, then the OSQSE can be a method to non-locally perform the SWAP gate, as both operationally provide the same result. However, the OSQSE has unique properties which we reveal in this work.

We formally define the OSQSE and its optimal entanglement cost, and derive computable lower bounds on the latter, which in turn yield bounds for the asymptotic quantum state exchange [1, 2]. In addition, we provide two useful conditions to decide whether a given initial state enables OSQSE with zero entanglement cost. We then show that there exist counter-intuitive phenomena in the OSQSE task that set it apart from the conventional SWAP operation.

**One-shot quantum state exchange.**— Consider two users, Alice and Bob, holding parts  $A$  and  $B$  of the initial state  $|\psi\rangle \equiv |\psi\rangle_{A_1 B_1 A_2 B_2 R}$  with systems  $A = A_1 A_2$  and  $B = B_1 B_2$ , respectively. Alice's and Bob's goal is either to exchange their parts  $A_1$  and  $B_1$  or to exchange their whole parts  $A$  and  $B$ .

Specifically, let  $\psi_{f_1}$  and  $\psi_{f_{12}}$  be the final states of the task,

$$\begin{aligned}\psi_{f_1} &= (\mathbb{1}_{A_1 \rightarrow A'_1} \otimes \mathbb{1}_{B_1 \rightarrow B'_1} \otimes \mathbb{1}_{A_2 B_2 R})(\psi), \\ \psi_{f_{12}} &= (\mathbb{1}_{A \rightarrow A'} \otimes \mathbb{1}_{B \rightarrow B'} \otimes \mathbb{1}_R)(\psi),\end{aligned}$$

where  $\psi = |\psi\rangle\langle\psi|$ , and the dimension of system  $X'$  is identical to that of system  $X$ . Note that  $B'_1$ ,  $B'$  and  $A'_1$ ,  $A'$  are Alice's and Bob's systems, respectively. Then three joint operations

$$\begin{aligned}\mathcal{E}_{\psi,K,L}^1 &: A_1 E_A^{\text{in}} \otimes B_1 E_B^{\text{in}} \longrightarrow B'_1 E_A^{\text{out}} \otimes A'_1 E_B^{\text{out}}, \\ \mathcal{E}_{\psi,K,L}^{1|2} &: A E_A^{\text{in}} \otimes B E_B^{\text{in}} \longrightarrow B'_1 A_2 E_A^{\text{out}} \otimes A'_1 B_2 E_B^{\text{out}}, \\ \mathcal{E}_{\psi,K,L}^{12} &: A E_A^{\text{in}} \otimes B E_B^{\text{in}} \longrightarrow B' E_A^{\text{out}} \otimes A' E_B^{\text{out}},\end{aligned}\quad (1)$$

are called the OSQSE protocols of  $|\psi\rangle$ , if they are performed by LOCC between Alice and Bob, and satisfy

$$\begin{aligned}\psi_{f_1} \otimes \Phi &= (\mathcal{E}_{\psi,K,L}^1 \otimes \mathbb{1}_{A_2 B_2 R})(\psi \otimes \Psi) \\ &= (\mathcal{E}_{\psi,K,L}^{1|2} \otimes \mathbb{1}_R)(\psi \otimes \Psi), \\ \psi_{f_{12}} \otimes \Phi &= (\mathcal{E}_{\psi,K,L}^{12} \otimes \mathbb{1}_R)(\psi \otimes \Psi),\end{aligned}\quad (2)$$

where  $\Psi$  and  $\Phi$  are pure maximally entangled states with Schmidt rank  $K$  and  $L$  on systems  $E_A^{\text{in}} E_B^{\text{in}}$  and  $E_A^{\text{out}} E_B^{\text{out}}$ , respectively. It is possible to generalize the above definitions by

\*Electronic address: yonghaelee@khu.ac.kr

†Electronic address: yamasaki@qi.t.u-tokyo.ac.jp

‡Electronic address: gerardo.adesso@nottingham.ac.uk

§Electronic address: level@khu.ac.kr

adding errors for approximation to Eq. (2), but it suffices to only consider error-free protocols to obtain our main results.

At this point, it is instructive to inform differences among the three protocols in Eq. (1) as follows: The first two protocols  $\mathcal{E}_{\psi,K,L}^1$  and  $\mathcal{E}_{\psi,K,L}^{1|2}$  indicate that only the parts  $A_1$  and  $B_1$  are exchanged, while the whole parts  $A_1A_2$  and  $B_1B_2$  are exchanged in the third protocol  $\mathcal{E}_{\psi,K,L}^{12}$ . In addition, the parts  $A_2$  and  $B_2$  can be used for exchanging  $A_1$  and  $B_1$  in the protocol  $\mathcal{E}_{\psi,K,L}^{1|2}$ , while  $A_2$  and  $B_2$  are untouched in the protocol  $\mathcal{E}_{\psi,K,L}^1$ .

Depending on the types of OSQSE protocols, we define three *optimal entanglement costs*

$$\begin{aligned} \mathbf{e}_{A_1 \leftrightarrow B_1}(\psi) &= \inf_{\mathcal{E}_{\psi,K,L}^1} (\log K - \log L), \\ \mathbf{e}_{A_1 \leftrightarrow B_1}^{A_2B_2}(\psi) &= \inf_{\mathcal{E}_{\psi,K,L}^{1|2}} (\log K - \log L), \\ \mathbf{e}_{A \leftrightarrow B}(\psi) &= \inf_{\mathcal{E}_{\psi,K,L}^{12}} (\log K - \log L), \end{aligned} \quad (3)$$

where the quantity  $\log K - \log L$  is called the *entanglement cost* of the OSQSE protocol, and the infimums are taken over all joint protocols  $\mathcal{E}_{\psi,K,L}^1$ ,  $\mathcal{E}_{\psi,K,L}^{1|2}$ , and  $\mathcal{E}_{\psi,K,L}^{12}$ , respectively.

By the definitions of the optimal entanglement costs, we obtain the following proposition.

**Proposition 1.** *For any input state  $\psi$ ,  $\mathbf{e}_{A_1 \leftrightarrow B_1}(\psi) \geq \mathbf{e}_{A_1 \leftrightarrow B_1}^{A_2B_2}(\psi)$ .*

*Converse bounds.*— A real number  $r$  is called a *converse bound* of the optimal entanglement cost if it is upper bounded by the entanglement cost of any OSQSE protocol. We derive converse bounds of the optimal entanglement costs as follows.

As in the asymptotic scenario [1, 2], we consider a one-shot version of the  $R$ -assisted quantum state exchange task, in which the reference system  $R$  is divided into two systems  $R_A$  and  $R_B$ , and then Alice and Bob receive the divided parts  $R_A$  and  $R_B$ , respectively, so that the initial state  $|\tilde{\psi}\rangle_{A_1B_1A_2B_2R_AR_B}$  is divided into Alice's parts  $AR_A$  and Bob's parts  $BR_B$ . This can be realized by using a quantum channel  $\mathcal{N} : R \rightarrow R_A$  and its complementary channel  $\mathcal{N}^c : R \rightarrow R_B$  [20]. Let  $\mathcal{E}_{\tilde{\psi},K,L}$  be an  $R$ -assisted OSQSE protocol,

$$\mathcal{E}_{\tilde{\psi},K,L} : AR_A E_{A_1}^{\text{in}} \otimes BR_B E_{B_1}^{\text{in}} \rightarrow B'_1 A_2 R_A E_{A_2}^{\text{out}} \otimes A'_1 B_2 R_B E_{B_2}^{\text{out}},$$

with the entanglement cost  $\log K - \log L$  such that  $\tilde{\psi}_f \otimes \Phi = \mathcal{E}_{\tilde{\psi},K,L}(\tilde{\psi} \otimes \Psi)$ , where  $\tilde{\psi}_f = (\mathbb{1}_{A_1 \rightarrow A'_1} \otimes \mathbb{1}_{B_1 \rightarrow B'_1} \otimes \mathbb{1}_{A_2B_2R_AR_B})(\tilde{\psi})$ . Note that  $\mathcal{E}_{\tilde{\psi},K,L}$  is an LOCC protocol by Alice and Bob. Let  $\sigma^M$  be the maximally mixed state with rank  $M$ . From the majorization condition for LOCC convertibility [21, 22], the state  $\rho_{B'_1A_2R_A} \otimes \sigma_{E_{B_2}^{\text{out}}}^L$  majorizes the state  $\rho_{AR_A} \otimes \sigma_{E_{B_1}^{\text{in}}}^K$ . Let  $F$  be an additive and Schur concave function [23] such that  $F(\sigma^M) = \log M$  for any  $M$ . From the Schur concavity of the function  $F$ , the inequality  $F(\rho_{B'_1A_2R_A} \otimes \sigma_{E_{B_2}^{\text{out}}}^L) \leq F(\rho_{AR_A} \otimes \sigma_{E_{B_1}^{\text{in}}}^K)$  holds. Since  $F$  is additive and  $\rho_{B'_1A_2R_A} = \rho_{B_1A_2R_A}$ , it follows that  $\log K - \log L \geq F(\rho_{B_1A_2R_A}) - F(\rho_{AR_A})$ . Since any protocol  $\mathcal{E}_{\psi,K,L}^{1|2}$  is also an  $R$ -assisted OSQSE protocol for the initial state  $|\psi\rangle$ , we obtain the following theorem.

**Theorem 2.** *For any input state  $\psi$ , the optimal entanglement cost  $\mathbf{e}_{A_1 \leftrightarrow B_1}^{A_2B_2}(\psi)$  is lower bounded by*

$$l_{1|2}(\psi) = \sup_{F, \mathcal{N}} [F(\mathcal{N}(\psi)_{B_1A_2R_A}) - F(\mathcal{N}(\psi)_{AR_A})],$$

where  $F$  is an additive and Schur concave function such that  $F(\sigma^M) = \log M$  for any  $M$  and  $\mathcal{N}(\rho)$  is a quantum channel from  $R$  to  $R_A$ .

In Theorem 2, if  $R$  is directly sent to either Alice or Bob without splitting, and we restrict the function  $F$  to the quantum Rényi entropy  $S_\alpha(\rho)$  of order  $\alpha$  [23] for a quantum state  $\rho$ , then we obtain the following computable converse bounds.

**Corollary 3.** *For any input state  $\psi$ ,  $\mathbf{e}_{A_1 \leftrightarrow B_1}^{A_2B_2}(\psi) \geq l_{1|2}^c(\psi) = \max_{\alpha \in [0, \infty]} f_\psi(\alpha)$ , where  $f_\psi(\alpha)$  is a function of  $|\psi\rangle$  and  $\alpha$  defined by  $f_\psi(\alpha) = \max\{S_\alpha(\rho_{A_1B_2}) - S_\alpha(\rho_B), S_\alpha(\rho_{B_1A_2}) - S_\alpha(\rho_A)\}$ .*

We refer the reader to Appendix A for the proof of Corollary 3. Remark that the converse bound  $l_{1|2}^c$  can be easily computed by means of analytical or numerical methods, since the function  $f_\psi(\alpha)$  is one-variable and differentiable on  $(0, \infty)$ . For the different types of the OSQSE protocols, we can also obtain a similar computable converse bound as follows:

$$\mathbf{e}_{X \leftrightarrow Y}(\psi) \geq \max_{\alpha \in [0, \infty]} |S_\alpha(\rho_X) - S_\alpha(\rho_Y)|, \quad (4)$$

where the pair  $(X, Y)$  can refer either to  $(A_1, B_1)$  or to  $(A, B)$ .

We also remark that in Theorem 2, if  $F$  is chosen as the von Neumann entropy [20], then the converse bound  $l_{1|2}$  recovers a theoretical converse bound in Refs. [1, 2]. In addition, a computable converse bound therein is just  $f_\psi(1)$  in Corollary 3. By virtue of the additivity of  $F$ , it is clear that  $l_{1|2}$  and  $l_{1|2}^c$  are also converse bounds of the optimal entanglement cost for the asymptotic quantum state exchange task. Hence, our converse bounds improve the existing bounds in Refs. [1, 2]. For example, if the initial state  $|\psi_1\rangle \equiv |\psi_1\rangle_{A_1B_1A_2B_2R}$  is

$$|\psi_1\rangle = \frac{1}{5} |00000\rangle + \sqrt{\frac{3}{50}} |00010\rangle + \frac{3}{5} |01001\rangle + \sqrt{\frac{27}{50}} |11100\rangle, \quad (5)$$

then we can find a value  $\alpha_0 \in [0, \infty]$  such that  $l_{1|2}^c(\psi_1) = f_{\psi_1}(\alpha_0) > f_{\psi_1}(1)$  as depicted in Fig. 1. This example shows that our bound  $l_{1|2}^c(\psi)$  is tighter than the existing bound  $f_\psi(1)$ .

*Conditions for zero entanglement cost.*— We now present conditions for OSQSE with zero entanglement cost.

By the converse bound in Eq. (4), it is obvious that if there exist Alice's and Bob's local isometries performing the OSQSE task, then the optimal entanglement cost is zero. We first characterize this type of strategy. Let  $(X, Y)$  be a pair of two systems, which can be either  $(A_1, B_1)$  or  $(A, B)$ , and consider a spectral decomposition of the reduced state  $\rho_{XY}$  for  $|\psi\rangle$ ,  $\rho_{XY} = \sum_{i=1}^N \lambda_i |\xi_i\rangle \langle \xi_i|_{XY}$ , where  $\lambda_i > 0$  with  $\sum_{i=1}^N \lambda_i = 1$ . For each  $i$ , we define the matrix  $\Omega_{XY}^i(\psi)$  by

$$\Omega_{XY}^i(\psi) = \sum_{j,k} (\langle j|_X \otimes \langle k|_Y) |\xi_i\rangle_{XY} |j\rangle \langle k|,$$

where  $\{|j\rangle\}$  and  $\{|k\rangle\}$  indicate the computational bases on Alice's and Bob's systems, respectively. Then we obtain the following sufficient condition.




FIG. 1: The graph of the function  $f_{\psi_1}(\alpha)$  for a specific initial state  $|\psi_1\rangle$  in Eq. (5). The maximum of  $f_{\psi_1}(\alpha)$  is attained at the point  $\alpha_0$  ( $\approx 3.362$ ). In the graph,  $\alpha_0$  is represented as the yellow dashed line, and 1 is represented as the red dashed dotted line.

**Theorem 4.** Let  $(X, Y)$  be either  $(A_1, B_1)$  or  $(A, B)$ . If there exist isometries  $U$  and  $V$  such that, for each  $i$ ,  $(\Omega_{XY}^i(\psi))^i = U\Omega_{XY}^i(\psi)V$ , then  $\mathbf{e}_{X \leftrightarrow Y}(\psi) = 0$ .

Here, the isometries  $U$  and  $V$  indicate Alice's and Bob's local operations exchanging the parts  $X$  and  $Y$  without shared entanglement. The proof of Theorem 4 is in Appendix B.

From the converse bound in Eq. (4), observe that if the spectrum of Alice's state is different from that of Bob's state, then the optimal entanglement cost cannot be zero. Based on this observation, we obtain the following theorem, whose proof can be found in Appendix C.

**Theorem 5.** Let  $(X, Y)$  be either  $(A_1, B_1)$  or  $(A, B)$ . If  $\mathbf{e}_{X \leftrightarrow Y}(\psi) = 0$ , then there exists an isometry  $U_{X \rightarrow Y}$  such that  $\rho_Y = U_{X \rightarrow Y} \rho_X (U_{X \rightarrow Y})^\dagger$ .

We remark that the converse of Theorem 5 is not true in general. Let us consider the following simple initial state

$$|\psi_2\rangle_{A_1 B_1 A_2 B_2} = \frac{1}{2}(|0000\rangle + |0101\rangle + |1010\rangle + |1111\rangle),$$

then  $\mathbf{e}_{A_1 \leftrightarrow B_1}(\psi_2) \geq \mathbf{e}_{A_1 \leftrightarrow B_1}^{A_2 B_2}(\psi_2) \geq 2$ , from Proposition 1 and Corollary 3. However, the state  $|\psi_2\rangle$  satisfies the necessary condition in Theorem 5, since its reduced states  $\rho_{A_1}$  and  $\rho_{B_1}$  are identical.

*Counter-intuitive phenomena.*— We are now in the position to present two phenomena which show the important differences between the OSQSE task and the SWAP operation.

(1) *Symmetric information.*— For the initial state  $|\psi\rangle$ , let us consider a scenario in which Alice and Bob exchange their whole information  $A$  and  $B$ . Assume that their parts  $A_2$  and  $B_2$  are symmetric, while the remaining parts  $A_1$  and  $B_1$  are not symmetric, i.e., the initial state  $|\psi\rangle$  satisfies  $(\text{SWAP}_{A_1 \leftrightarrow B_1})(\psi) \neq \psi$  and  $(\text{SWAP}_{A_2 \leftrightarrow B_2})(\psi) = \psi$  where  $\text{SWAP}_{X \leftrightarrow Y}$  is the operation swapping quantum states in systems  $X$  and  $Y$ .

From a viewpoint of the SWAP operation, if Alice and Bob want to exchange  $A$  and  $B$ , then it suffices for them to exchange  $A_1$  and  $B_1$ , since  $A_2$  is identical to  $B_2$ . This situation can be more easily understood by using a cargo exchange as a




FIG. 2: Illustration of the cargo exchange task. The cargoes  $A_1 A_2$  and  $B_1 B_2$  belong to Alice and Bob, respectively. Assume that the cargoes  $A_2$  and  $B_2$  are symmetric, but  $A_1$  and  $B_1$  are not symmetric. When Alice and Bob exchange their whole cargoes  $A_1 A_2$  and  $B_1 B_2$ , it suffices for them to exchange  $A_1$  and  $B_1$ , since  $A_2$  is identical to  $B_2$ . The truck indicates the cost needed for exchanging  $A_1$  and  $B_1$ .

metaphor for the SWAP operation as depicted in Fig. 2. In the cargo exchange, assume that Alice and Bob want to exchange their whole cargoes, and some of the cargoes are symmetric. In terms of efficiency, it is reasonable for them to exchange only  $A_1$  and  $B_1$  in order to reduce the cargo exchange cost, because the cargoes  $A_2$  and  $B_2$  are the same.

On the other hand, in the OSQSE, the proper use of the symmetric parts  $A_2$  and  $B_2$  can more efficiently reduce the entanglement cost compared to exchanging only  $A_1$  and  $B_1$  without using  $A_2$  and  $B_2$ . To be specific, there exists an initial state  $|\psi\rangle$  such that the parts  $A_2$  and  $B_2$  are symmetric and  $\mathbf{e}_{A \leftrightarrow B}(\psi) = 0$  while the rest parts  $A_1$  and  $B_1$  are not symmetric. Consider the specific initial state

$$|\phi_1\rangle_{A_1 B_1 A_2 B_2 R} = \frac{1}{\sqrt{2}}(|00000\rangle + |01111\rangle),$$

where  $A_2$  and  $B_2$  are symmetric but  $A_1$  and  $B_1$  are not. Since  $\Omega_{AB}^1(\phi_1) = |00\rangle\langle 00|$  and  $\Omega_{AB}^2(\phi_1) = |01\rangle\langle 11|$ , we can show that  $\Omega_{AB}^1(\phi_1)$  and  $\Omega_{AB}^2(\phi_1)$  satisfy the condition in Theorem 4, by setting

$$U = V = |00\rangle\langle 00| + |01\rangle\langle 11| + |10\rangle\langle 10| + |11\rangle\langle 01|. \quad (6)$$

Thus we obtain that  $\mathbf{e}_{A \leftrightarrow B}(\phi_1) = 0$ , which means that  $A$  and  $B$  can be exchanged by means of LOCC without consuming any non-local resource. As mentioned above, this phenomenon cannot occur when using the SWAP operation.

The above example also shows that the use of the symmetric parts  $A_2$  and  $B_2$  can reduce the entanglement cost for exchanging  $A_1$  and  $B_1$ . Since the initial state  $|\phi_1\rangle$  does not satisfy the necessary condition in Theorem 5, we obtain  $\mathbf{e}_{A_1 \leftrightarrow B_1}(\phi_1) > 0$ . Observe that the isometry  $U$  ( $V$ ) in Eq. (6) represents Alice's (Bob's) local operation  $\text{CNOT}_A$  ( $\text{CNOT}_B$ ) whose target and controlled systems are  $A_1$  ( $B_1$ ) and  $A_2$  ( $B_2$ ), respectively. This implies that Alice and Bob can exchange  $A_1$  and  $B_1$  by using local operations. It follows that  $0 \geq \mathbf{e}_{A_1 \leftrightarrow B_1}^{A_2 B_2}(\phi_1)$ . In fact,  $\mathbf{e}_{A_1 \leftrightarrow B_1}^{A_2 B_2}(\phi_1) = 0$  from Corollary 3. Therefore, we obtain  $\mathbf{e}_{A_1 \leftrightarrow B_1}(\phi_1) > \mathbf{e}_{A_1 \leftrightarrow B_1}^{A_2 B_2}(\phi_1)$ .

When  $A_2$  and  $B_2$  are symmetric, we can show the following relation between the optimal entanglement costs by definition.

**Proposition 6.**  $\mathbf{e}_{A \leftrightarrow B}(\psi) = \mathbf{e}_{A_1 \leftrightarrow B_1}^{A_2 B_2}(\psi)$ , if the parts  $A_2$  and  $B_2$  of  $|\psi\rangle$  are symmetric.




FIG. 3: Illustration of the one-shot quantum state exchange protocol of  $|\phi_2\rangle$  in Eq. (7). (a) In order to exchange  $A_1$  and  $B_1$ , Alice and Bob locally prepare an ebit each, and they apply Bell measurements to the shaded areas. (b) By performing local operations corresponding to the measurement outcomes, the parts  $A_1$  and  $B_1$  can be exchanged. At the same time, Alice and Bob can share two ebits.

From Proposition 6, we can see that, when Alice and Bob exchange systems  $A$  and  $B$  of  $|\psi\rangle$  with symmetric parts  $A_2$  and  $B_2$ , they can achieve the optimal entanglement cost by exchanging only  $A_1$  and  $B_1$ , making the most of this symmetry.

(2) *Negative entanglement cost.*— As in the asymptotic quantum state exchange task [1, 2], there exist initial states to show that the entanglement cost of the OSQSE task can be negative. Assume that Alice and Bob exchange the parts  $A_1$  and  $B_1$  of the initial state

$$|\phi_2\rangle_{A_1 B_1 A_2 B_2} = \frac{1}{2} \sum_{i,j=0}^1 |i\rangle_{A_1} |j\rangle_{B_1} |j\rangle_{A_2} |i\rangle_{B_2}, \quad (7)$$

where  $|\phi_2\rangle$  consists of two ebits  $|e\rangle_{A_1 B_2}$  and  $|e\rangle_{B_1 A_2}$ . To exchange  $A_1$  and  $B_1$ , both Alice and Bob prepare an ebit, respectively, and they locally implement entanglement swapping [24] by performing two Bell measurements on  $A_2$ ,  $B_2$ , and the parts of the ebits, as described in Fig. 3. Then they can exchange  $A_1$  and  $B_1$ , and can share two ebits at the same time. This means that the entanglement cost can be negative. In fact, we have  $\mathbf{e}_{A_1 \leftrightarrow B_1}^{A_2 B_2}(\phi_2) = -2$  from Corollary 3. This is in stark contrast with the SWAP operation, which cannot lead to creation of shared entanglement between Alice and Bob.

From Proposition 6, we can know that if  $A_2$  and  $B_2$  are symmetric, then  $\mathbf{e}_{A_1 \leftrightarrow B_1}^{A_2 B_2}(\psi)$  cannot be negative. One may ask the question: Is there any condition that implies the non-negativity of the optimal entanglement cost  $\mathbf{e}_{A_1 \leftrightarrow B_1}^{A_2 B_2}$ ? To answer this question, we present the following inequalities.

**Proposition 7.**

$$\begin{aligned} \mathbf{e}_{A_1 \leftrightarrow B_1}^{A_2 B_2}(\psi) + \mathbf{e}_{B'_1 \leftrightarrow A'_1}^{A_2 B_2}(\psi_{f_1}) &\geq 0, \\ \mathbf{e}_{A_1 \leftrightarrow B_1}^{A_2 B_2}(\psi) + \mathbf{e}_{A_2 \leftrightarrow B_2}^{B'_1 A'_1}(\psi_{f_1}) &\geq \mathbf{e}_{A \leftrightarrow B}(\psi), \end{aligned}$$

where  $\mathbf{e}_{B'_1 \leftrightarrow A'_1}^{A_2 B_2}(\psi_{f_1})$  is the optimal entanglement cost for exchanging  $B'_1$  and  $A'_1$  when using  $A_2$  and  $B_2$ , and  $\mathbf{e}_{A_2 \leftrightarrow B_2}^{B'_1 A'_1}(\psi_{f_1})$  is the optimal entanglement cost for exchanging  $A_2$  and  $B_2$  when using  $B'_1$  and  $A'_1$ .

In Proposition 7, the first inequality comes from the fact that Alice and Bob cannot increase the amount of entanglement between them by means of LOCC [25], while the second one is straightforward from the definitions of the optimal entanglement costs. From Proposition 7, we can see that if  $\mathbf{e}_{B'_1 \leftrightarrow A'_1}^{A_2 B_2}(\psi_{f_1})$  or  $\mathbf{e}_{A_2 \leftrightarrow B_2}^{B'_1 A'_1}(\psi_{f_1})$  is non-positive, then  $\mathbf{e}_{A_1 \leftrightarrow B_1}^{A_2 B_2}(\psi)$  cannot be negative. Moreover, if the condition  $\mathbf{e}_{A_2 \leftrightarrow B_2}^{B'_1 A'_1}(\psi_{f_1}) \leq \mathbf{e}_{A \leftrightarrow B}(\psi)$  holds, then Proposition 7 implies  $\mathbf{e}_{A_1 \leftrightarrow B_1}^{A_2 B_2}(\psi) \geq 0$ .

In particular, let us assume that  $A_1$  and  $B_1$  are symmetric. Then it is obvious that  $0 \geq \mathbf{e}_{A_1 \leftrightarrow B_1}^{A_2 B_2}(\psi)$ , from Proposition 1. If  $0 > \mathbf{e}_{A_1 \leftrightarrow B_1}^{A_2 B_2}(\psi)$  then it follows from Proposition 7 that  $\mathbf{e}_{B'_1 \leftrightarrow A'_1}^{A_2 B_2}(\psi_{f_1}) > 0$ . However, since  $B'_1$  and  $A'_1$  are also symmetric, Proposition 1 implies  $\mathbf{e}_{B'_1 \leftrightarrow A'_1}^{A_2 B_2}(\psi_{f_1}) \leq 0$ , which leads to a contradiction. Therefore, we obtain the following corollary.

**Corollary 8.**  $\mathbf{e}_{A_1 \leftrightarrow B_1}^{A_2 B_2}(\psi) = 0$ , if  $A_1$  and  $B_1$  are symmetric.

This tells us that if  $A_1$  and  $B_1$  are symmetric, Alice and Bob cannot increase the amount of shared entanglement after the OSQSE task, even if they make use of the parts  $A_2$  and  $B_2$ .

*Conclusion.*— In this work, we have considered a one-shot version of the original quantum state exchange task, and have formally defined the OSQSE task and its optimal entanglement costs. We have derived converse bounds on the optimal entanglement costs, and have presented conditions on the initial state to achieve zero entanglement cost. As a related open problem, we can ask the following question: If  $\mathbf{e}_{A \leftrightarrow B}(\psi) = 0$ , then is it possible to exchange the parts  $A$  and  $B$ , without classical communication and entanglement, that is, are there local operations  $L_A$  and  $L_B$  such that  $\psi_{f_{12}} = (L_A \otimes L_B)(\psi)$ ?

The two counter-intuitive phenomena are the most interesting contribution of this work, showing the major difference between the SWAP operation and the OSQSE. One phenomenon tells us that it is worth using the symmetric parts in order to optimally perform the OSQSE. The other shows that the entanglement cost of the OSQSE can be negative. By observing the aforementioned examples involving the phenomena, we can provide another interesting open problem: If  $\mathbf{e}_{A_1 \leftrightarrow B_1}^{A_2 B_2}(\psi) \leq 0$ , do there exist Alice's and Bob's local operations  $L'_A$  and  $L'_B$  such that  $\psi_{f_1} \otimes \Phi = (L'_A \otimes L'_B)(\psi)$ ?

A further open problem is whether the catalytic use of entanglement [26–28] can reduce the optimal entanglement cost for the OSQSE. To be more specific, for the initial state  $|\psi\rangle$ , do there exist a bipartite entangled state  $|\psi_c\rangle_{A_3 B_3}$  shared by Alice and Bob and a OSQSE protocol  $C_{K,L} : AA_3 E_A^{in} \otimes BB_3 E_B^{in} \rightarrow B'_A A_3 E_A^{out} \otimes A'_B B_3 E_B^{out}$  such that  $\psi_{f_{12}} \otimes \psi_c \otimes \Phi = (C_{K,L} \otimes \mathbb{1}_R)(\psi \otimes \psi_c \otimes \Psi)$  and  $\log K - \log L < \mathbf{e}_{A \leftrightarrow B}(\psi)$ ?

Theoretically, the OSQSE is a powerful two-user quantum communication task, which includes quantum teleportation [29] and quantum state merging [3, 4] as special cases. Practically, this task can be a fundamental building block for applications involving multiple users, such as distributed quantum computation [30, 31] and quantum network [32–34].

## Acknowledgments

We would like to thank Ryuji Takagi and Bartosz Regula for fruitful discussion. This research was supported by Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Science and ICT (NRF-2019R1A2C1006337) and the MSIT (Ministry of Science and ICT), Korea, under the ITRC (Information Technology Research Center) support program (IITP-2019-2018-

0-01402) supervised by the IITP (Institute for Information & communications Technology Promotion). H. Y. acknowledges Grant-in-Aid for JSPS Research Fellow, JSPS KAKENHI Grant No. 18J10192, Cross-ministerial Strategic Innovation Promotion Program (SIP) (Council for Science, Technology and Innovation (CSTI)), and CREST (Japan Science and Technology Agency) JPMJCR1671. G. A. acknowledges support from the ERC Starting Grant GQCOP (Grant Agreement No. 637352).

- 
- [1] J. Oppenheim and A. Winter, arXiv:quant-ph/0511082.
  - [2] Y. Lee, R. Takagi, H. Yamasaki, G. Adesso, and S. Lee, Phys. Rev. Lett. **122**, 010502 (2019).
  - [3] M. Horodecki, J. Oppenheim, and A. Winter, Nature (London) **436**, 673 (2005).
  - [4] M. Horodecki, J. Oppenheim, and A. Winter, Commun. Math. Phys. **269**, 107 (2006).
  - [5] I. Devetak and J. Yard, Phys. Rev. Lett. **100**, 230501 (2008).
  - [6] J. T. Yard and I. Devetak, IEEE Trans. Inf. Theory **55**, 5339 (2009).
  - [7] B. Schumacher, Phys. Rev. A **51**, 2738 (1995).
  - [8] A. Abeyesinghe, I. Devetak, P. Hayden, and A. Winter, Proc. R. Soc. A **465**, 2537 (2009).
  - [9] M. Berta, M. Christandl, and R. Renner, Commun. Math. Phys. **306**, 579 (2011).
  - [10] J. M. Renes and R. Renner, IEEE Trans. Inf. Theory **58**, 1985 (2012).
  - [11] L. Wang and R. Renner, Phys. Rev. Lett. **108**, 200501 (2012).
  - [12] N. Datta and M.-H. Hsieh, IEEE Trans. Inf. Theory **59**, 1929 (2013).
  - [13] M. Berta, M. Christandl, and D. Touchette, IEEE Trans. Inf. Theory **62**, 1425 (2016).
  - [14] Q. Zhao, Y. Liu, X. Yuan, E. Chitambar, and X. Ma, Phys. Rev. Lett. **120**, 070403 (2018).
  - [15] A. Anshu, R. Jain, and N. A. Warsi, IEEE Trans. Inf. Theory **64**, 1425 (2018).
  - [16] B. Regula, K. Fang, X. Wang, and G. Adesso, Phys. Rev. Lett. **121**, 010401 (2018).
  - [17] H. Yamasaki and M. Murao, IEEE Trans. Inf. Theory [arXiv:1806.07875 (to be published)].
  - [18] M. A. Nielsen and I. L. Chuang, *Quantum Computation and Quantum Information* (Cambridge University Press, 2010).
  - [19] R. Jozsa and A. Miyake, Proc. R. Soc. A **464**, 3089 (2008).
  - [20] M. M. Wilde, *Quantum Information Theory* (Cambridge University Press, Cambridge, England, 2013).
  - [21] M. A. Nielsen, Phys. Rev. Lett. **83**, 436 (1999).
  - [22] A. Mari, V. Giovannetti, and A. S. Holevo, Nat. Commun. **5** (2014).
  - [23] W. van Dam and P. Hayden, arXiv:quant-ph/0204093.
  - [24] M. Żukowski, A. Zeilinger, M. A. Horne, and A. K. Ekert, Phys. Rev. Lett. **71**, 4287 (1993).
  - [25] V. Vedral, M. B. Plenio, M. A. Rippin, and P. L. Knight, Phys. Rev. Lett. **78**, 2275 (1997).
  - [26] D. Jonathan and M. B. Plenio, Phys. Rev. Lett. **83**, 3566 (1999).
  - [27] J. Eisert and M. Wilkens, Phys. Rev. Lett. **85**, 437 (2000).
  - [28] C. Majenz, M. Berta, F. Dupuis, R. Renner, and M. Christandl, Phys. Rev. Lett. **118**, 080503 (2017).
  - [29] C. H. Bennett, G. Brassard, C. Crépeau, R. Jozsa, A. Peres, and W. K. Wootters, Phys. Rev. Lett. **70**, 1895 (1993).
  - [30] J. I. Cirac, A. K. Ekert, S. F. Huelga, , and C. Macchiavello, Phys. Rev. A **59**, 4249 (1999).
  - [31] D. Bruß, G. M. D'Ariano, M. Lewenstein, C. Macchiavello, A. Sen(De), and U. Sen, Phys. Rev. Lett. **93**, 210501 (2004).
  - [32] J. I. Cirac, P. Zoller, H. J. Kimble, and H. Mabuchi, Phys. Rev. Lett. **78**, 3221 (1997).
  - [33] K. Azuma, A. Mizutani, and H.-K. Lo, Nat. Commun. **7** (2016).
  - [34] A. Pirker, J. Wallnöfer, and W. Dür, New J. Phys. **20**, 053054 (2018).

### Appendix A: Proof of Corollary 3

We show that there exists a number  $\alpha_0 \in [0, \infty]$  such that  $I_{1|2}^c(\psi) = f_\psi(\alpha_0)$ . Note that the function  $f_\psi(\alpha)$  is continuous on the compact set  $[0, 1]$ . So the extreme value theorem implies that there exists a number  $\alpha_1 \in [0, 1]$  such that  $f_\psi(\alpha_1) \geq f_\psi(\alpha)$  for all  $\alpha \in [0, 1]$ . Let us consider the function  $g(x)$  on the interval  $[0, 1]$  defined as

$$g(x) = \begin{cases} f_\psi(\infty) & \text{if } x = 0 \\ f_\psi(\frac{1}{x}) & \text{otherwise,} \end{cases}$$

then  $g(x)$  is continuous on  $[0, 1]$ . By using the extreme value theorem again, there exists a number  $x_0 \in [0, 1]$  such that  $g(x_0) \geq g(x)$  for all  $x \in [0, 1]$ . It follows that there exists a number  $\alpha_2 \in [1, \infty]$  such that  $f_\psi(\alpha_2) \geq f_\psi(\alpha)$  for all  $\alpha \in [1, \infty]$ . By setting  $\alpha_0 = \max\{\alpha_1, \alpha_2\}$ , we obtain that  $I_{1|2}^c(\psi) = f_\psi(\alpha_0) \geq f_\psi(\alpha)$  for all  $\alpha \in [0, \infty]$ .

### Appendix B: Proof of Theorem 4

When  $X = A$  and  $Y = B$ , consider the Schmidt decompositions of  $|\psi\rangle$ ,

$$|\psi\rangle_{ABR} = \sum_{i=1}^N \sqrt{\lambda_i} |\xi_i\rangle_{AB} \otimes |i\rangle_R,$$

where  $\lambda_i > 0$  with  $\sum_{i=1}^N \lambda_i = 1$ . For the computational bases  $\{|j\rangle\}$  and  $\{|k\rangle\}$  on the systems  $A$  and  $B$ , respectively, we have

$$\begin{aligned} |\psi\rangle_{ABR} &= \sum_{i=1}^N \sqrt{\lambda_i} \sum_{j,k} [\Omega_{AB}^i(\psi)]_{jk} |j\rangle_A \otimes |k\rangle_B \otimes |i\rangle_R, \end{aligned}$$

where  $[\Omega_{AB}^i(\psi)]_{jk} = (\langle j|_A \otimes \langle k|_B) |\xi_i\rangle_{AB}$ . If the parts  $A$  and  $B$  are perfectly exchanged, then Alice and Bob hold the final state

$$\begin{aligned} |\psi\rangle_{BAR} &= \sum_{i=1}^N \sqrt{\lambda_i} \sum_{j,k} [\Omega_{AB}^i(\psi)]_{kj} |j\rangle_B \otimes |k\rangle_A \otimes |i\rangle_R, \end{aligned}$$

By the hypothesis, there exist isometries  $U$  and  $V$  such that for each  $i$ ,

$$(\Omega_{AB}^i(\psi))^t = U \Omega_{AB}^i(\psi) V.$$

So we have, for each  $i$ ,

$$[\Omega_{AB}^i(\psi)]_{kj} = \sum_{l,m} [\Omega_{AB}^i(\psi)]_{lm} \langle j| U |l\rangle \langle k| V^t |m\rangle,$$

which implies that

$$\begin{aligned} |\psi\rangle_{BAR} &= \sum_{i=1}^N \sqrt{\lambda_i} \sum_{l,m} [\Omega_{AB}^i(\psi)]_{lm} \sum_j |j\rangle \langle j| U |l\rangle \\ &\quad \otimes \sum_k |k\rangle \langle k| V^t |m\rangle \otimes |i\rangle_R, \\ &= \sum_{i=1}^N \sqrt{\lambda_i} \sum_{l,m} [\Omega_{AB}^i(\psi)]_{lm} U |l\rangle \otimes V^t |m\rangle \otimes |i\rangle_R, \\ &= (U \otimes V^t \otimes I_R) |\psi\rangle_{ABR}. \end{aligned}$$

Hence,  $\mathbf{e}_{A \leftrightarrow B}(\psi) = 0$ .

Similarly, we can show that  $\mathbf{e}_{A_1 \leftrightarrow B_1}(\psi) = 0$  by using isometries  $U'$  and  $V'$  such that for each  $i$ ,  $(\Omega_{A_1 B_1}^i(\psi))^t = U' \Omega_{A_1 B_1}^i(\psi) V'$ .

### Appendix C: Proof of Theorem 5

We use the following lemma in order to prove Theorem 5.

**Lemma 9.** *Let  $Z$  and  $W$  be any discrete random variables on alphabets  $\mathcal{Z}$  and  $\mathcal{W}$  with  $|\mathcal{Z}| = N$  and  $|\mathcal{W}| = M$ . Let  $\{p_i\}_{i=1}^N$  and  $\{q_i\}_{i=1}^M$  be probability distributions for  $X$  and  $Y$ , respectively. If the following equality holds for all  $\alpha \in [0, \infty]$ ,*

$$H_\alpha(Z) = H_\alpha(W),$$

*where  $H_\alpha(\cdot)$  is the Rényi entropy of classical random variables, then  $|\mathcal{Z}| = |\mathcal{W}|$  and there exists a permutation  $\sigma \in S_N$  such that  $p_i = q_{\sigma(i)}$  for all  $i \in [N]$ , where  $S_N$  is the set of all permutations on  $[N] = \{1, \dots, N\}$ .*

Note that  $H_\alpha(Z) = \lim_{x \rightarrow \alpha} H_x(Z)$  and  $S_\alpha(\rho_A) = \lim_{x \rightarrow \alpha} S_x(\rho_A)$  for each  $\alpha \in [0, \infty]$ .

*Proof.* Suppose that  $H_\alpha(Z) = H_\alpha(W)$  for all  $\alpha \in [0, \infty]$ . Since  $H_0(Z) = H_0(W)$ , it holds that  $|\mathcal{Z}| = |\mathcal{W}|$ . For convenience, we assume that any probability distribution  $\{r_i\}_{i=1}^N$  satisfies  $r_1 \geq r_i$  for all  $i \in [N]$ .

We now prove the statement by using mathematical induction on  $N$ .

(i) If  $N = 2$ , then  $H_\infty(Z) = H_\infty(W)$  implies  $p_1 = q_1$  and so  $p_2 = 1 - p_1 = 1 - q_1 = q_2$ . Thus the statement is true.

(ii) Suppose that the statement is true for  $N = k - 1$ . Let  $Z_1$  and  $Z_2$  be discrete random variables on alphabets  $\mathcal{Z}_1$  and  $\mathcal{Z}_2$  with  $|\mathcal{Z}_1| = |\mathcal{Z}_2| = k$ . Let  $\{p_i\}_{i=1}^k$  and  $\{q_i\}_{i=1}^k$  be probability distributions for  $Z_1$  and  $Z_2$ , respectively. Since  $H_\infty(Z_1) = H_\infty(Z_2)$ ,  $p_1 = q_1$ . By setting  $p'_i = \frac{p_{i+1}}{1-p_1}$  and  $q'_i = \frac{q_{i+1}}{1-q_1}$  for each  $i \in [k-1]$ , we can construct random variables  $Z'_1$  and  $Z'_2$  on alphabets  $\mathcal{Z}'_1$  and  $\mathcal{Z}'_2$  whose probability distributions are  $\{p'_i\}_{i=1}^{k-1}$  and  $\{q'_i\}_{i=1}^{k-1}$ , respectively. Obviously,  $|\mathcal{Z}'_1| = |\mathcal{Z}'_2| = k - 1$ , and so  $H_0(Z'_1) = H_0(Z'_2)$ . Observe that

for  $\alpha \in (0, 1) \cup (1, \infty)$

$$\begin{aligned}
H_\alpha(Z_1) &= H_\alpha(Z_2) \\
\Rightarrow \frac{1}{1-\alpha} \log \left( \sum_{i=1}^k p_i^\alpha \right) &= \frac{1}{1-\alpha} \log \left( \sum_{i=1}^k q_i^\alpha \right) \\
\Rightarrow \sum_{i=2}^k p_i^\alpha &= \sum_{i=2}^k q_i^\alpha \\
\Rightarrow \sum_{i=1}^{k-1} \left( \frac{p_{i+1}}{1-p_1} \right)^\alpha &= \sum_{i=1}^{k-1} \left( \frac{q_{i+1}}{1-p_1} \right)^\alpha \\
\Rightarrow \frac{1}{1-\alpha} \log \left( \sum_{i=1}^{k-1} (p'_i)^\alpha \right) &= \frac{1}{1-\alpha} \log \left( \sum_{i=1}^{k-1} (q'_i)^\alpha \right) \\
\Rightarrow H_\alpha(Z'_1) &= H_\alpha(Z'_2).
\end{aligned}$$

In addition, if  $\alpha = 1$ , then

$$\begin{aligned}
H_1(Z_1) &= H_1(Z_2) \\
\Rightarrow \sum_{i=1}^k p_i \log \frac{1}{p_i} &= \sum_{i=1}^k q_i \log \frac{1}{q_i} \\
\Rightarrow \sum_{i=2}^k p_i \log \frac{1}{p_i} &= \sum_{i=2}^k q_i \log \frac{1}{q_i} \\
\Rightarrow (1-p_1) \log (1-p_1) + \sum_{i=2}^k p_i \log \frac{1}{p_i} \\
&= (1-p_1) \log (1-p_1) + \sum_{i=2}^k q_i \log \frac{1}{q_i} \\
\Rightarrow \sum_{i=2}^k \frac{p_i}{1-p_1} \log \frac{1-p_1}{p_i} &= \sum_{i=2}^k \frac{q_i}{1-p_1} \log \frac{1-p_1}{q_i} \\
\Rightarrow \sum_{i=1}^{k-1} p'_i \log \frac{1}{p'_i} &= \sum_{i=1}^{k-1} q'_i \log \frac{1}{q'_i} \\
\Rightarrow H_1(Z'_1) &= H_1(Z'_2).
\end{aligned}$$

Finally, we have

$$\begin{aligned}
H_\infty(Z'_1) - H_\infty(Z'_2) &= \lim_{\alpha \rightarrow \infty} H_\alpha(Z'_1) - \lim_{\alpha \rightarrow \infty} H_\alpha(Z'_2) \\
&= \lim_{\alpha \rightarrow \infty} (H_\alpha(Z'_1) - H_\alpha(Z'_2)) = 0.
\end{aligned}$$

It follows that  $H_\alpha(Z'_1) = H_\alpha(Z'_2)$  for all  $\alpha \in [0, \infty]$ . By the induction hypothesis, there exists a permutation  $\sigma' \in S_{k-1}$  such that  $p'_i = q'_{\sigma'(i)}$  for all  $i \in [k-1]$ . Define  $\sigma(1) = 1$  and  $\sigma(i) = \sigma'(i-1)$  with  $i \neq 1$ . Then  $\sigma \in S_k$  and  $p_i = q_{\sigma(i)}$  for all  $i \in [k]$ . Therefore, the statement is true for  $N = k$ .  $\square$

In fact, we can prove Lemma 9 by assuming a weaker condition as follows. Let  $S$  be a subset of  $[0, \infty]$  including 0, the extended real number  $\infty$ , and a sequence  $\{s_n\}_{n \in \mathbb{N}}$  such that  $\lim_{n \rightarrow \infty} s_n = \infty$ . Then we can show that if  $H_\alpha(Z) = H_\alpha(W)$  holds for all  $\alpha \in S$ , then  $Z$  and  $W$  have the same probability distribution.

The contrapositive of the following lemma proves Theorem 5.

**Lemma 10** (Sufficient conditions on the initial state  $|\psi\rangle$  with  $\mathbf{e}_{X \leftrightarrow Y}(\psi) > 0$ ). Let  $(X, Y)$  be the pair of two systems, which can be either  $(A_1, B_1)$  or  $(A, B)$ . Let  $\{\lambda_i\}_{i=1}^N$  and  $\{\tau_i\}_{i=1}^M$  be non-zero eigenvalues for the reduced states  $\rho_X$  and  $\rho_Y$  of  $|\psi\rangle$ , respectively, which satisfy  $\lambda_1 \geq \dots \geq \lambda_N$ ,  $\tau_1 \geq \dots \geq \tau_M$ , and  $\sum_{i=1}^N \lambda_i = \sum_{i=1}^M \tau_i = 1$ . Then  $\mathbf{e}_{X \leftrightarrow Y} > 0$ , if one of the following conditions holds:

- (i)  $N \neq M$ .
- (ii)  $N = M$  and  $\lambda_{i'} \neq \tau_{i'}$  for some  $i' \in [N] = \{1, \dots, N\}$ .

*Proof.* (i) If  $N \neq M$ , then  $\text{rank}(\rho_X) \neq \text{rank}(\rho_Y)$ , which means

$$\mathbf{e}_{X \leftrightarrow Y}(\psi) \geq |S_0(\rho_X) - S_0(\rho_Y)| > 0,$$

by the converse bound in Eq. (4).

(ii) Suppose that  $|\psi\rangle$  satisfies  $N = M$  and  $\lambda_{i'} \neq \tau_{i'}$  for some  $i' \in [N]$ . Let  $Z$  and  $W$  be discrete random variables on alphabets  $\mathcal{Z}$  and  $\mathcal{W}$  with  $|\mathcal{Z}| = |\mathcal{W}| = N$ , whose probability distributions are  $\{\lambda_i\}_{i=1}^N$  and  $\{\tau_i\}_{i=1}^N$ , respectively. Let us consider the set

$$A = \{i \in [N] | \lambda_i \neq \tau_i\},$$

then  $A$  is a non-empty subset of  $[N]$ , since  $i' \in A$ . So we can choose the largest element in  $A$ , say  $j$ . Then  $\lambda_j \neq \tau_j$  and  $\lambda_i = \tau_i$  for all  $i > j$  by the definition of the set  $A$ . If  $\lambda_j > \tau_j$  (or  $\lambda_j < \tau_j$ ) then  $\lambda_i > \tau_j$  (or  $\lambda_j < \tau_i$ ) for all  $i \in [j]$ . Thus  $\lambda_i \neq \tau_j$  (or  $\lambda_j \neq \tau_i$ ) for all  $i \in [j]$ , which shows that for each  $\sigma \in S_j$ , there exists  $i \in [j]$  such that  $\lambda_i \neq \tau_{\sigma(i)}$ . It follows that for each  $\sigma \in S_N$ , there exists  $i \in [N]$  such that  $\lambda_i \neq \tau_{\sigma(i)}$ . From the contrapositive of Lemma 9, there exists  $\alpha' \in [0, \infty]$  such that  $H_{\alpha'}(X) \neq H_{\alpha'}(Y)$ . Therefore, from the converse bound in Eq. (4), we obtain

$$\mathbf{e}_{X \leftrightarrow Y}(\psi) \geq |S_{\alpha'}(\rho_X) - S_{\alpha'}(\rho_Y)| > 0.$$

$\square$