WHITNEY NUMBERS FOR POSET CONES

GALEN DORPALEN-BARRY, JANG SOO KIM AND VICTOR REINER

ABSTRACT. Hyperplane arrangements dissect \mathbb{R}^n into connected components called chambers, and a wellknown theorem of Zaslavsky counts chambers as a sum of nonnegative integers called Whitney numbers of the first kind. His theorem generalizes to count chambers within any cone defined as the intersection of a collection of halfspaces from the arrangement, leading to a notion of Whitney numbers for each cone. This paper focuses on cones within the braid arrangement, consisting of the reflecting hyperplanes $x_i = x_j$ inside \mathbb{R}^n for the symmetric group, thought of as the type A_{n-1} reflection group. Here,

• cones correspond to posets,

• chambers within the cone correspond to linear extensions of the poset,

• the Whitney numbers of the cone interestingly refine the number of linear extensions of the poset.

We interpret this refinement for all posets as counting linear extensions according to a statistic that generalizes the number of left-to-right maxima of a permutation. When the poset is a disjoint union of chains, we interpret this refinement differently, using Foata's theory of cycle decomposition for multiset permutations, leading to a simple generating function compiling these Whitney numbers.

1. INTRODUCTION

This paper concerns arrangements $\mathcal{A} = \{H_1, \ldots, H_m\}$ of hyperplanes H_i , which are affine-linear codimension one subspaces of $V = \mathbb{R}^n$. Each such arrangement dissects V into the connected components of its complement $V \setminus \bigcup_{i=1}^m H_i$, called *chambers*. Let $\mathcal{C}(\mathcal{A})$ denote the collection of all such chambers.

The theory of hyperplane arrangements is rich and well-explored, with connections to reflection groups, braid groups, random walks, card-shuffling, and discrete geometry of polytopes and oriented matroids; see [19, 27]. In particular, the number of chambers $\#\mathcal{C}(\mathcal{A})$ has a famous formula due to Zaslavsky [31], expressed in terms of the *intersection poset* $\mathcal{L}(\mathcal{A})$, consisting of all *intersection subspaces* $X = H_{i_1} \cap H_{i_2} \cap \cdots \cap H_{i_k}$, ordered via reverse inclusion. This poset is known to have the property that every *lower interval*

 $[V, X] := \{ Y \in \mathcal{L}(\mathcal{A}) : V \le Y \le X \}$

from its unique bottom element V to any intersection space X forms a geometric lattice. In particular, each such [V, X] is a ranked poset, where the rank of X is $\operatorname{codim}(X) := n - \dim(X)$. Zaslavsky's result says that

(1)
$$\#\mathcal{C}(\mathcal{A}) = \sum_{X \in \mathcal{L}(\mathcal{A})} |\mu(V, X)| = \sum_{k=0}^{n} c_k(\mathcal{A}) = [\mathsf{Poin}(\mathcal{A}, t)]_{t=1}$$

where $\mu(-,-)$ denotes the *Möbius function* of $\mathcal{L}(\mathcal{A})$, while the nonnegative integers

$$c_k(\mathcal{A}) := \sum_{\substack{X \in \mathcal{L}(\mathcal{A}):\\ \operatorname{codim}(X)=k}} |\mu(V, X)|,$$

are often called the (signless) Whitney numbers of the first kind for \mathcal{A} , and their generating function

$$\mathsf{Poin}(\mathcal{A},t) := \sum_{k \ge 0} c_k(\mathcal{A}) t^k$$

is called the *Poincaré polynomial*¹.

Date: June 18, 2021.

¹⁹⁹¹ Mathematics Subject Classification. 05Axx, 06A07.

Key words and phrases. hyperplane, arrangement, cone, poset, Whitney, Catalan, Stirling, Foata, multiset, permutation. The first and third authors were partially supported by NSF grant DMS-1601961. The second author was supported by

NRF grants #2019R1F1A1059081 and #2016R1A5A1008055.

¹This is because it is the generating function for the Betti numbers of the complexified complement $\mathbb{C}^n \setminus \mathcal{A}$; see [19, Chap. 5]

Our starting point is a less widely-known generalization of equation (1), also due to Zaslavsky [32]. It applies more generally to count the chambers of \mathcal{A} that lie within a *cone* \mathcal{K} , defined to be the intersection of any collection of open halfspaces for hyperplanes of \mathcal{A} ; said differently, a cone \mathcal{K} of \mathcal{A} is a chamber in $\mathcal{C}(\mathcal{A}')$ for some subarrangement $\mathcal{A}' \subseteq \mathcal{A}$. Results on the set $\mathcal{C}(\mathcal{K})$ of all chambers of \mathcal{A} inside a cone \mathcal{K} have appeared more recently in work of Brown on random walks [8], and in work of Gente on Varchenko determinants [13, Section 2.4], and independently Aguiar and Mahajan [1, Theorem 8.22]. Define the *poset* of interior intersections for \mathcal{K} to be the following order ideal in $\mathcal{L}(\mathcal{A})$:

$$\mathcal{L}^{\mathrm{int}}(\mathcal{K}) = \{ X \in \mathcal{L}(\mathcal{A}) \mid X \cap \mathcal{K} \neq \emptyset \}.$$

Zaslavsky observed in [32, Example A, p. 275] that (1) generalizes to cones \mathcal{K} , asserting

(2)
$$#\mathcal{C}(\mathcal{K}) = \sum_{X \in \mathcal{L}^{\text{int}}(\mathcal{K})} |\mu(V, X)| = \sum_{k=0}^{n} c_k(\mathcal{K}) = [\mathsf{Poin}(\mathcal{K}, t)]_{t=1}$$

Here we again define nonnegative integers, the (signless) Whitney numbers of the first kind for the cone \mathcal{K}

$$c_k(\mathcal{K}) := \sum_{\substack{X \in \mathcal{L}^{\text{int}}(\mathcal{K}):\\ \text{codim}(X) = k}} |\mu(V, X)|,$$

with generating function $\mathsf{Poin}(\mathcal{K}, t) := \sum_k c_k(\mathcal{K}) t^k$, which we call the *Poincaré polynomial* of \mathcal{K} .

For example, inside $\mathcal{A} = \{H_1, H_2, H_3, H_4, H_5\}$ in \mathbb{R}^2 shown below on the left, we have shaded one of four possible cones \mathcal{K} defined by the subarrangement $\mathcal{A}' = \{H_4, H_5\}$, containing $\#\mathcal{C}(\mathcal{K}) = 5$ chambers of \mathcal{A} :



Zaslavsky's formula (2) computes this as follows. The poset of interior intersections $\mathcal{L}^{int}(\mathcal{K})$ has Hasse diagram:



Here $\mu(V, X) = (-1)^{\text{codim}(X)}$ for all X, so that $(c_0(\mathcal{K}), c_1(\mathcal{K}), c_2(\mathcal{K})) = (1, 3, 1)$, and

$$\begin{array}{rcl} \operatorname{\mathsf{Poin}}(\mathcal{K},t) &=& c_0(\mathcal{K}) + c_1(\mathcal{K})t + c_2(\mathcal{K})t^2 &=& 1 + 3t + t^2, \\ \#\mathcal{C}(\mathcal{K}) &=& \left[\operatorname{\mathsf{Poin}}(\mathcal{K},t)\right]_{t=1} &=& c_0(\mathcal{K}) + c_1(\mathcal{K}) + c_2(\mathcal{K}) &=& 5. \end{array}$$

The object of this paper is to understand the distribution of the signless Whitney numbers as a refinement of $\#C(\mathcal{K})$ as in equation (2), for cones \mathcal{K} in the *braid arrangement*. The braid arrangement $A_{n-1} = \{H_{ij}\}_{1 \le i < j \le n}$ is the set of $\binom{n}{2}$ reflecting hyperplanes

$$H_{ij} = \{ (x_1, \dots, x_n) \in V = \mathbb{R}^n \mid x_i - x_j = 0 \}$$

for the symmetric group \mathfrak{S}_n on *n* letters, thought of as the reflection group of type A_{n-1} . There is a wellknown and easy bijection between the chambers $\mathcal{C}(A_{n-1})$ and the permutations $\sigma = [\sigma_1, \sigma_2, \ldots, \sigma_n]$ in \mathfrak{S}_n , sending σ to the chamber:

(3)
$$\mathcal{K}_{\sigma} := \{ x \in V = \mathbb{R}^n : x_{\sigma_1} < x_{\sigma_2} < \dots < x_{\sigma_n} \}.$$

More generally, one has an easy bijection, reviewed in Section 2, between posets on the set $[n] := \{1, 2, ..., n\}$ and cones in the braid arrangement A_{n-1} , sending a poset P to the cone

$$\mathcal{K}_P := \{ x \in V = \mathbb{R}^n : x_i < x_j \text{ for } i <_P j \}.$$

It is readily checked that the chamber \mathcal{K}_{σ} lies in the cone $\mathcal{C}(\mathcal{K}_P)$ if and only if σ is a *linear extension* of P, meaning that $i <_P j$ implies $i <_{\sigma} j$, regarding σ as a total order $\sigma_1 < \sigma_2 < \cdots < \sigma_n$ on [n]. Letting LinExt(P) denote the set of all linear extensions of P, this shows that $\#\mathcal{C}(\mathcal{K}_P) = \#\operatorname{LinExt}(P)$, and hence equation (2) becomes

(4)
$$\# \operatorname{LinExt}(P) = \sum_{k \ge 0} c_k(P) = [\operatorname{\mathsf{Poin}}(P, t)]_{t=1}$$

abbreviating $c_k(P) := c_k(\mathcal{K}_P)$ and $\mathsf{Poin}(P,t) := \mathsf{Poin}(\mathcal{K}_P,t)$ from now on.

A motivating special case occurs when $P = \text{Antichain}_n$ is the *antichain poset* on [n] that has no order relations. In this case $\text{LinExt}(\text{Antichain}_n) = \mathfrak{S}_n$ itself, and the signless Whitney number $c_{n-k}(\text{Antichain}_n)$ of the first kind is well-known [28, Prop. 1.3.7] to be the signless Stirling number of the first kind c(n, k) that counts the permutations in \mathfrak{S}_n having k cycles. Consequently, (4) becomes the easy summation formula

(5)
$$n! = \#\mathfrak{S}_n = \sum_{k \ge 0} c(n,k).$$

Our main results will generalize three well-known expressions for $\mathsf{Poin}(\mathsf{Antichain}_n, t)$, explained below:

(6)
$$\mathsf{Poin}(\mathsf{Antichain}_n, t) = \sum_{k \ge 0} c(n, k) t^{n-k} = \sum_{\sigma \in \mathfrak{S}_n} t^{n-\mathsf{cyc}(\sigma)}$$

(7)
$$= \sum_{\sigma \in \mathfrak{S}} t^{n-\mathsf{LRmax}(\sigma)}$$

(8)
$$= 1(1+t)(1+2t)\cdots(1+(n-1)t).$$

Equation (6) comes from the definition of c(n, k) mentioned above, where $cyc(\sigma)$ is the number of cycles of σ . This interpretation of the Poincaré polynomial will be generalized to all posets in Theorem 1.1, and generalized in a different way to posets which are *disjoint unions of chains* in Theorem 1.3.

Equation (7) arises from a well-known bijection $\tau \mapsto \sigma$ from \mathfrak{S}_n to itself, called the fundamental bijection², such that $\mathsf{cyc}(\tau) = \mathsf{LRmax}(\sigma)$, see [28, Proposition 1.3.1]; here $\mathsf{LRmax}(\sigma)$ is the number of *left-to-right maxima* of $\sigma = \sigma_1 \sigma_2 \dots \sigma_n$, i.e., the number of integers $1 \leq j \leq n$ such that $\sigma_j = \max\{\sigma_1, \sigma_2, \dots, \sigma_j\}$. This bijection and interpretation of the Poincaré polynomial will be generalized to all posets in Theorem 1.2.

Equation (8) is well-known [28, Proposition 1.3.7], and will be generalized to a generating function compiling the Poincaré polynomials of all posets which are disjoint unions of chains in Theorem 1.4 below.

The remainder of this paper is organized as follows.

Section 2 gives preliminaries on the intersection lattice and cones in braid arrangements. In particular, for a poset P on [n], it gives an explicit combinatorial description of the interior intersections $\mathcal{L}^{int}(\mathcal{K}_P)$ for a poset cone \mathcal{K}_P , as the subset $\Pi^{\uparrow}(P)$ of P-transverse partitions inside the lattice Π_n of set partitions of [n]; roughly speaking, these are the set partitions π for which the quotient preposet P/π does not collapse any strict order relations $i <_P j$ into equalities. With this in hand, it defines the subset $\mathfrak{S}^{\uparrow}(P)$ of P-transverse permutations inside \mathfrak{S}_n as those permutations σ whose cycles form a P-transverse partition, in order to prove the following generalization of equation (6).

Theorem 1.1. For any poset P on [n], one has

$$\mathsf{Poin}(P,t) = \sum_{\sigma \in \mathfrak{S}^{\pitchfork}(P)} t^{n - \mathsf{cyc}(\sigma)}$$

In particular, $\# \operatorname{LinExt}(P) = \#\mathfrak{S}^{\uparrow}(P).$

This result even generalizes to a statement (Theorem 2.30) about cones in any real reflection arrangement.

In light of Theorem 1.1, one expects bijections between $\mathfrak{S}^{\uparrow}(P)$ and $\operatorname{LinExt}(P)$. One such bijection is the goal of Section 3, which defines a notion of *P*-left-to-right maximum for a linear extension of *P*, and then generalizes the fundamental bijection $\tau \mapsto \sigma$ and equation (7) as follows.

²This bijection, often called *Foata's first fundamental transformation*, seems to have appeared also in work of Rényi [24, §4].

Theorem 1.2. For any poset P on [n], one has a bijection

$$\begin{array}{ccc} \mathfrak{S}^{\pitchfork}(P) & \stackrel{\Phi}{\longrightarrow} & \operatorname{LinExt}(P) \\ \tau & \longmapsto & \sigma \end{array}$$

such that $cyc(\tau) = LRmax_P(\sigma)$, where $LRmax_P(\sigma)$ is the number of P-left-to-right maxima of σ . Therefore,

$$\mathsf{Poin}(P,t) = \sum_{\sigma \in \mathrm{LinExt}(P)} t^{n-\mathsf{LRmax}_P(\sigma)}$$

Section 4 examines posets P which are disjoint unions of chains, and produces a second interesting bijection $\mathfrak{S}^{\uparrow}(P) \to \operatorname{LinExt}(P)$. Given any *composition* $\overline{a} = (a_1, \ldots, a_\ell)$ of n, meaning that $\overline{a} \in \{0, 1, 2, \ldots\}^\ell$ and $|\overline{a}| := \sum_{i=1}^{\ell} a_i = n$, let a_i denote a chain (totally ordered) poset on a_i elements, and then

$$P_{\overline{a}} := \mathsf{a}_1 \sqcup \mathsf{a}_2 \sqcup \cdots \sqcup \mathsf{a}_\ell$$

is a disjoint union of incomparable chains of sizes a_1, a_2, \ldots, a_ℓ . Here one can identify elements in $\text{LinExt}(P_{\overline{a}})$ with *multiset permutations* of $\{1^{a_1}, 2^{a_2}, \ldots, \ell^{a_\ell}\}$. Section 4 reviews the beautiful theory of *prime cycle decompositions* for such multiset permutations due to Foata [9], and then reinterprets it as giving a bijection $\text{LinExt}(P_{\overline{a}}) \to \mathfrak{S}^{\uparrow}(P_{\overline{a}})$, and the following (second) generalization of equation (6).

Theorem 1.3. For any composition \overline{a} of n, the disjoint union $P_{\overline{a}}$ of chains has a bijection

$$\begin{array}{ccc} \operatorname{LinExt}(P_{\overline{a}}) & \longrightarrow & \mathfrak{S}^{\pitchfork}(P_{\overline{a}}) \\ \sigma & \longmapsto & \tau \end{array}$$

with $cyc(\tau) = pcyc(\sigma)$, the number of prime cycles in Foata's unique decomposition for σ . Thus

$$\mathsf{Poin}(P_{\overline{a}}, t) = \sum_{\sigma \in \mathrm{LinExt}(P_{\overline{a}})} t^{n - \mathsf{pcyc}(\sigma)}.$$

Foata's theory is then used to prove the following generating function, generalizing equation (8).

Theorem 1.4. For $\ell = 1, 2, \ldots$, one has

$$\sum_{e \in \{1,2,\dots\}^{\ell}} \mathsf{Poin}(P_{\overline{a}}, t) \cdot \mathbf{x}^{\overline{a}} = \frac{1}{1 - \sum_{j=1}^{\ell} e_j(\mathbf{x}) \cdot (t-1)(2t-1) \cdots ((j-1)t-1)}$$

where $\mathbf{x}^{\overline{a}} := x_1^{a_1} \cdots x_\ell^{a_\ell}$ and $e_j(\mathbf{x}) := \sum_{1 \le i_1 < \cdots < i_j \le \ell} x_{i_1} \cdots x_{i_j}$ is the j^{th} elementary symmetric function.

Section 5 then examines posets of width two, that is, posets P decomposable as $P = P_1 \cup P_2$ where the subposets P_1, P_2 are chains (i.e. totally ordered subsets) inside P. Here the Whitney numbers $c_k(P)$ are interpreted by a descent-like statistic on σ in LinExt(P):

$$\mathsf{des}_{P_1,P_2}(\sigma) := \#\{i \in [n-1] : \sigma_i \in P_2, \ \sigma_{i+1} \in P_1, \text{ with } \sigma_i, \sigma_{i+1} \text{ incomparable in } P\}.$$

Theorem 1.5. For a width two poset decomposed into two chains as $P = P_1 \cup P_2$, one has

$$\mathsf{Poin}(P,t) = \sum_{\sigma \in \mathrm{LinExt}(P)} t^{\mathsf{des}_{P_1,P_2}(\sigma)}.$$

Theorem 1.5 will be used to show (Corollary 5.3) that, for certain very special width two posets P, the Poincaré polynomial $\mathsf{Poin}(P, t)$ coincides with the *P*-Eulerian polynomial,

(9)
$$\sum_{\sigma \in \text{LinExt}(P)} t^{\text{des}(\sigma)}$$

which counts linear extensions σ of P according to their number of (usual) descents

$$\mathsf{les}(\sigma) := \#\{i \in [n-1] : \sigma_i > \sigma_{i+1}\},\$$

assuming that P has been naturally labeled in the sense that the identity permutation $\sigma = [1, 2, ..., n]$ lies in LinExt(P). In particular, Example 5.4 uses this to deduce that for $P = 2 \times n$, the Cartesian product of chains having sizes 2 and n, the Whitney numbers $c_k(2 \times n)$ are Narayana numbers, counting $2 \times n$ standard tableaux according to their number of descents. This P-Eulerian polynomial has many interpretations, e.g., as the h-polynomial of the order complex for the distributive lattice J(P) of order ideals in P, or of the P-partition triangulation of the order polytope for P; see [23, Proposition 2.1, Proposition 2.2] and [28, Sections 3.4, 3.8, 3.13] for more on this. Unfortunately, in general the *P*-Eulerian polynomial differs from the Poincaré polynomial Poin(P, t) considered here. For example, when *P* is an antichain with three elements, the *P*-Eulerian polynomial is $1 + 4t + t^2$, while $Poin(P, t) = 1 + 3t + 2t^2$.

Lastly, we note that the cone \mathcal{K}_P associated to a poset P has appeared in many places in the literature, for example, implicitly in Stanley's theory of P-partitions [28, §3.15] and related work of Gessel on quasisymmetric functions [12], more explicitly in work of Björner and Wachs [4, 5, 6], and as the ranking COMs of Bandelt, Chepoi and Knauer [2]. A pointed version of this cone, called an order cone is discussed in Beck and Sanyal [3, Chap. 6]. When the underlying Hasse diagram of P is a tree, results on the number of chambers $\#\mathcal{C}(\mathcal{K}_P) = \#\operatorname{LinExt}(P)$ in these cones appear in work of K. Saito [25].

2. The braid arrangement, its intersection lattice, and its cones

2.1. Preliminaries on arrangements. We begin with some preliminaries on hyperplane arrangements, focusing on braid arrangements. Good references include [19], [27], [28, §3.11], [20, §3.3].

Definition 2.1. A hyperplane in $V = \mathbb{R}^n$ is an affine linear subspace of codimension one. An arrangement of hyperplanes in \mathbb{R}^n is a finite collection $\mathcal{A} = \{H_1, \ldots, H_m\}$ of distinct hyperplanes. A chamber of \mathcal{A} is an open, connected component of $\mathbb{R}^n \setminus \bigcup_{H \in \mathcal{A}} H$. The set of all chambers of \mathcal{A} is denoted by $\mathcal{C}(\mathcal{A})$.

Example 2.2. The type A reflection arrangement, A_{n-1} , also called the braid arrangement, consists of the $\binom{n}{2}$ hyperplanes of the form

$$H_{ij} = \{(x_1, \dots, x_n) \in \mathbb{R}^n \mid x_i - x_j = 0\}$$

for integers $1 \leq i < j \leq n$. There are n! chambers $\mathcal{K}_{\sigma} = \{x \in \mathbb{R}^n : x_{\sigma_1} < \cdots < x_{\sigma_n}\}$ of A_{n-1} , naturally indexed by the permutations $\sigma = [\sigma_1, \sigma_2, \ldots, \sigma_n]$ of [n], that give the strict inequalities ordering the coordinates within the chamber, as in (3). For example, when n = 4,

$$\mathcal{K}_{1243} = \{ x \in \mathbb{R}^4 : x_1 < x_2 < x_4 < x_3 \}, \\ \mathcal{K}_{4213} = \{ x \in \mathbb{R}^4 : x_4 < x_2 < x_1 < x_3 \}$$

are two out of the 4! = 24 chambers of $\mathcal{C}(A_{4-1})$.

Definition 2.3. Let \mathcal{A} be a hyperplane arrangement in \mathbb{R}^n . An *intersection* of \mathcal{A} is a nonempty subspace of the form $X = H_{i_1} \cap H_{i_2} \cap \cdots \cap H_{i_k}$ where $\{H_{i_1}, H_{i_2}, \ldots, H_{i_k}\} \subseteq \mathcal{A}$. Here the ambient vector space $V = \mathbb{R}^n$ is considered to be the intersection $\bigcap_{H \in \mathcal{A}} H$ of the empty set of hyperplanes. We denote the set of intersections of \mathcal{A} by $\mathcal{L}(\mathcal{A})$.

Example 2.4. The intersections of A_{n-1} are described by equalities between the variables.

- For all $n \ge 1$, the line $x_1 = x_2 = \cdots = x_n$ is the intersection of all the hyperplanes of A_{n-1} .
- When n = 4 the intersection of H_{12} and H_{34} in the subspace of \mathbb{R}^4 in which $x_1 = x_2$ and $x_3 = x_4$. On the other hand, the intersection of H_{12} and H_{13} is the subspace of \mathbb{R}^4 in which $x_1 = x_2 = x_3$.

More generally, there is a bijection $\pi \mapsto X_{\pi}$ between the collection Π_n of all set partitions $\pi = \{B_1, \ldots, B_k\}$ of $[n] = \{1, 2, \ldots, n\}$ and the set of all intersections of A_{n-1} . The bijection sends the set partition π to the subspace X_{π} where one has equal coordinates $x_i = x_j$ whenever i, j lie in a common block B_k of π . We sometimes denote the set partition $\pi = \{B_1, \ldots, B_k\}$ with the notation $\pi = B_1|B_2|\cdots|B_k$, and may or may not include commas and set braces around the elements of each block B_i . E.g., $1 \mid 23 \mid 456$ and $\{\{1\}, \{2,3\}, \{4,5,6\}\}$ represent the same set partition of [6].

- For example, the set partition $1|2|\cdots|n$ in which all elements appear as singletons corresponds to $X_{1|2|\cdots|n} = V = \mathbb{R}^n$, the empty intersection, which is the ambient space.
- For all $n \ge 1$, the set partition $123 \cdots n$ having all the elements in the same block corresponds to the line $X_{123\cdots n}$ defined by $x_1 = x_2 = \cdots = x_n$.
- When n = 4, one has $X_{12|34} = H_{12} \cap H_{34}$ and $X_{123|4} = H_{12} \cap H_{13}$.

The collection $\mathcal{L}(\mathcal{A})$ of all intersections of an arrangement \mathcal{A} will be partially ordered by reverse inclusion, and called the *intersection poset* of \mathcal{A} . It has a unique minimal element, namely the intersection

$$\bigcap_{H\in\mathscr{O}}H=V=\mathbb{R}^n$$

Let Π_n denote the lattice of set partitions on [n], ordered via refinement: $\pi_1 \leq \pi_2$ if π_1 refines π_2 . For the braid arrangement A_{n-1} , the intersection poset $\mathcal{L}(A_{n-1})$ is isomorphic to Π_n .

Proposition 2.5 ([27, pp. 26-27]). The map $\pi \mapsto X_{\pi}$ in Example 2.4 is a poset isomorphism $\Pi_n \cong \mathcal{L}(A_{n-1})$.

For any hyperplane arrangement \mathcal{A} , each of the lower intervals $[V, X] := \{Y \in \mathcal{L}(\mathcal{A}) : V \leq Y \leq X\}$ forms a geometric lattice [28, Prop. 3.11.2]. In particular, this implies that each such lower interval is a ranked poset, with rank function given by the codimension $\operatorname{codim}(X) = \dim(V) - \dim(X)$. Furthermore, this implies that its *Möbius function* values $\mu(V, X)$, defined recursively by $\mu(V, V) := 1$ and $\mu(V, X) :=$ $-\sum_{Y:V \leq Y < X} \mu(V, Y)$, will alternate in sign in the sense that $(-1)^{\operatorname{codim}(X)} \mu(V, X) \geq 0$. For the braid arrangement, these Möbius function values have a simple expression.

Proposition 2.6 ([28, Example 3.10.4]). For any set partition $\pi = \{B_1, \ldots, B_k\}$ in Π_n , one has

$$\mu(V, X_{\pi}) = (-1)^{n-k} \prod_{i=1}^{k} (\#B_i - 1)! \quad (= \mu(1|2|\cdots|n, \pi))$$

with the convention 0! := 1. Here $\mu(V, X_{\pi}), \mu(1|2| \cdots |n, \pi)$ are $\mu(-, -)$ values in $\mathcal{L}(A_{n-1}), \Pi_n$, respectively.

For the sake of our later discussion, we point out a re-interpretation of this formula involving permutations. Given a permutation σ in \mathfrak{S}_n , when considered as acting on $V = \mathbb{R}^n$, its fixed space $V^{\sigma} := \{ \mathbf{x} \in \mathbb{R}^n : \sigma(\mathbf{x}) = \{ \mathbf{x} \in \mathbb{R}^n : \sigma(\mathbf{x}) = \{ \mathbf{x} \in \mathbb{R}^n : \sigma(\mathbf{x}) \in \mathbb{R}^n : \sigma(\mathbf{x}) \in \mathbb{R}^n \} \}$ **x**} will be the intersection subspace $V^{\sigma} = X_{\pi}$, where $\pi = \{B_1, B_2, \dots, B_k\}$ is the set partition in Π_n given by the cycles of σ . Consequently, one can re-interpret

(10)
$$\begin{aligned} |\mu(V, X_{\pi})| &= \prod_{i=1}^{k} (\#B_{i} - 1)! \\ &= \#\{\sigma \in \mathfrak{S}_{n} \text{ with cycle partition } \pi\} \\ &= \#\{\sigma \in \mathfrak{S}_{n} : V^{\sigma} = X_{\pi}\}, \end{aligned}$$

since each block B_i of π has $(\#B_i - 1)!$ choices of a cyclic orientation.

Definition 2.7. Let \mathcal{A} be an arrangement of hyperplanes in \mathbb{R}^n . For $0 \leq k \leq n$, the kth signless Whitney number of $\mathcal{L}(\mathcal{A})$ of the first kind is

$$c_k(\mathcal{A}) = \sum_{\substack{X \in \mathcal{L}(\mathcal{A}):\\ \operatorname{codim}(X) = k}} |\mu(V, X)| = (-1)^k \sum_{\substack{X \in \mathcal{L}(\mathcal{A}):\\ \operatorname{codim}(X) = k}} \mu(V, X).$$

Henceforth, we call $c_k(\mathcal{A}), 0 \leq k \leq n$, the Whitney numbers of \mathcal{A} . One of the standard ways to compile them into a generating function is their *Poincaré polynomial* $\mathsf{Poin}(\mathcal{A}, t) := \sum_{k=0}^{n} c_k(\mathcal{A}) t^k$; see [19, §2.3].

As mentioned in the Introduction, we aim to understand the chambers, intersections, and Whitney numbers for cones in \mathcal{A} ; the chambers, intersections, and Whitney numbers for \mathcal{A} are a special case.

Definition 2.8. Let \mathcal{A} be an arrangement of hyperplanes in $V = \mathbb{R}^n$. A cone³ \mathcal{K} of \mathcal{A} is any nonempty intersection $\emptyset \neq \mathcal{K} \subseteq V = \mathbb{R}^n$ of (open) halfspaces defined by a subset \mathcal{A}' of the hyperplanes from \mathcal{A} . That is, a cone \mathcal{K} is any one of the (open) chambers from the set of all chambers $\mathcal{C}(\mathcal{A}')$ for some subarrangement $\mathcal{A}' \subseteq \mathcal{A}$. For example, in the following arrangement in \mathbb{R}^2 there are four cones defined by the dashed hyperplanes. One such cone \mathcal{K} is shaded below.



³Aguiar and Mahajan [1] call these objects top-cones.

Each cone \mathcal{K} of \mathcal{A} has its collection of *chambers*, namely those chambers in $\mathcal{C}(\mathcal{A})$ that lie inside \mathcal{K} :

$$\mathcal{C}(\mathcal{K}) = \{ C \in \mathcal{C}(\mathcal{A}) : C \subset \mathcal{K} \}.$$

The poset of *interior intersections* of the cone \mathcal{K} is the following order ideal within the poset $\mathcal{L}(\mathcal{A})$:

$$\mathcal{L}^{\rm int}(\mathcal{K}) = \mathcal{L}^{\rm int}_{\mathcal{A}}(\mathcal{K}) = \{ X \in \mathcal{L}(\mathcal{A}) \mid X \cap \mathcal{K} \neq \emptyset \}.$$

For each X in $\mathcal{L}^{int}_{\mathcal{A}}(\mathcal{K})$, its lower interval [V, X] is still a geometric lattice, with same rank function $\operatorname{codim}(X)$, so that one can define the k^{th} (signless) Whitney number of \mathcal{K} by

$$c_k(\mathcal{K}) = \sum_{\substack{X \in \mathcal{L}^{\mathrm{int}}(\mathcal{K}):\\ \mathrm{codim}(X) = k}} |\mu(V, X)| = (-1)^k \sum_{\substack{X \in \mathcal{L}^{\mathrm{int}}(\mathcal{K}):\\ \mathrm{codim}(X) = k}} \mu(V, X),$$

along with their generating function $\mathsf{Poin}(\mathcal{K},t) := \sum_{k=0}^{n} c_k(\mathcal{K}) t^k$, the *Poincaré polynomial* for \mathcal{K} .

The starting point for our study is the following result of Zaslavsky [32] counting the number $\#C(\mathcal{K})$ of chambers of an arrangement \mathcal{A} lying inside one of its cones \mathcal{K} .

Theorem 2.9 ([32, Example A, p. 275]). Let \mathcal{K} be a cone of an arrangement \mathcal{A} in $V = \mathbb{R}^n$. Then

$$#\mathcal{C}(\mathcal{K}) = \sum_{X \in \mathcal{L}^{\text{int}}(\mathcal{K})} |\mu(V, K)| = c_0(\mathcal{K}) + c_1(\mathcal{K}) + \dots + c_n(\mathcal{K}) = [\mathsf{Poin}(\mathcal{K}, t)]_{t=1}.$$

Zaslavsky proved in his doctoral thesis [31] the better-known special case of Theorem 2.9 for the full arrangement, that is, where $\mathcal{K} = V = \mathbb{R}^n$.

The following two examples illustrate Theorem 2.9 for two cones in A_3 .

Example 2.10. Consider the braid arrangement $\mathcal{A} = A_4 = \{H_{12}, H_{13}, H_{14}, H_{23}, H_{24}, H_{34}\}$ inside $V = \mathbb{R}^4$. On the left below we have drawn a linearly equivalent picture of its intersection with the hyperplane where $x_1 + x_2 + x_3 + x_4 = 0$, isomorphic to \mathbb{R}^3 , and depicted the intersection of the hyperplanes with the unit 2-sphere in this 3-dimensional space. Here we pick the cone \mathcal{K} to be the one defined by the halfspace $x_3 < x_4$ for the hyperplane H_{34} , and draw the intersection of H_{34} with the unit sphere as the equatorial circle, with the other five hyperplanes H_{ij} depicted as great circles intersecting the hemisphere where $x_3 < x_4$. On the right below the non-hyperplane interior intersection subspaces X_{π} are labeled.



Therefore the intersection poset $\mathcal{L}^{int}(\mathcal{K})$ of this cone is



We have $(c_0(\mathcal{K}), c_1(\mathcal{K}), c_2(\mathcal{K})) = (1, 5, 6)$. Summing these gives 1 + 5 + 6 = 12, and a quick visual verification assures that there are 12 chambers in this cone.

Example 2.11. Consider the cone \mathcal{K} of A_3 in which $x_3 < x_4$ and $x_1 < x_2$. On the left below we have drawn the same picture as Example 2.10 with the cone corresponding to \mathcal{K} shaded. We depict $\mathcal{L}^{\text{int}}(\mathcal{K})$ on the right.



We have $c_0(\mathcal{K}) = 1$, $c_1(\mathcal{K}) = 4$, and $c_2(\mathcal{K}) = 1$. Summing these gives $1 + 4 + 1 = 6 = \#\mathcal{C}(\mathcal{K})$.

For the remainder of this paper, we focus on cones \mathcal{K} inside braid arrangements A_{n-1} . It is well-known (see [20, §3.3], for example) and easy to see that such cones correspond bijectively with posets P on [n]via this rule: one has $x_i < x_j$ for all points in the cone \mathcal{K} if and only if $i <_P j$. We will denote the cone associated to P by \mathcal{K}_P , and abbreviate $c_k(P) := c_k(\mathcal{K}_P)$ and $\mathsf{Poin}(P,t) := \mathsf{Poin}(\mathcal{K}_P, t)$.

Example 2.12. The cone inside A_3 in Example 2.10 given by the inequality $x_3 < x_4$ on $V = \mathbb{R}^4$ has defining poset P_1 with order relation $3 <_{P_1} 4$ on $[4] = \{1, 2, 3, 4\}$, while the cone in Example 2.11 given by the inequalities $x_1 < x_2$ and $x_3 < x_4$ has defining poset P_2 with order relations $1 <_{P_2} 2$ and $3 <_{P_2} 4$. These posets P_1, P_2 are shown here:

$$P_1 = \begin{array}{c} (4) \\ (1) (2) (3) \end{array} \qquad P_2 = \begin{array}{c} (2) (4) \\ (1) (3) \end{array}$$

2.2. **Preposets, posets, cones and a characterization of** $\mathcal{L}^{\text{int}}(\mathcal{K}_P)$ **.** By Theorem 2.5, the intersection poset $\mathcal{L}(A_{n-1})$ is isomorphic to the set partition lattice Π_n , and hence for each cone \mathcal{K}_P in \mathcal{A}_{n-1} , one should be able to identify the interior intersection poset $\mathcal{L}^{\text{int}}(\mathcal{K}_P)$ as some order ideal inside Π_n . This is our next goal, which will be aided by recalling some facts about preposets, posets, binary relations, and cones.

Definition 2.13. Recall that a preposet Q on [n] is a binary relation $Q \subseteq [n] \times [n]$ which is both

- reflexive, meaning $(i, i) \in Q$ for all i, and
- transitive, meaning $(i, j), (j, k) \in Q$ implies $(i, k) \in Q$.

If in addition Q is antisymmetric, meaning $(i, j), (j, i) \in Q$ implies i = j, then Q is called a poset on [n]; in this case, we sometimes write $i \leq_Q j$ when $(i, j) \in Q$.

A set partition $\pi \in \Pi_n$ is identified with an *equivalence relation* $\pi \subseteq [n] \times [n]$ having $(i, j) \in \pi$ when i, j appear in the same block of π . That is, π is reflexive, transitive, and *symmetric*, meaning $(i, j) \in \pi$ implies $(j, i) \in \pi$. We will sometimes write this binary relation as $i \equiv_{\pi} j$ when $(i, j) \in \pi$.

The union $Q_1 \cup Q_2 \subseteq [n] \times [n]$ of two preposets will be reflexive, but possibly not transitive, so not always a preposet. However, the transitive closure operation $Q \mapsto \overline{Q}$ lets one complete it to a preposet $\overline{Q_1 \cup Q_2}$.

We will use a slight rephrasing of the folklore *cone-preposet dictionary*, as discussed by Postnikov, Reiner, and Williams in [20, Section 3.3]. This dictionary is a bijection between preposets Q on [n] and *closed* cones of any dimension that are intersections in $V = \mathbb{R}^n$ of closed halfspaces of the form $\{x_i \leq x_j\}$. Under this bijection, any such closed cone C corresponds to a preposet Q_C via

$$C \mapsto Q_C := \{(i, j) \mid x_i \le x_j \text{ for all } \mathbf{x} \in C\}.$$

Conversely, any preposet Q on [n] corresponds to a closed cone C_Q via

$$Q \mapsto C_Q := \bigcap_{(i,j) \in Q} \{ x_i \le x_j \} = \{ \mathbf{x} \in \mathbb{R}^n \mid x_i \le x_j \text{ for all } (i,j) \in Q \}.$$

For a subset $A \subseteq \mathbb{R}^n$, denote its *interior* and *relative interior* by int(A), relint(A). Then for a preposet Q,

(11)
$$\operatorname{relint}(C_Q) = \left\{ \mathbf{x} \in \mathbb{R}^n : \frac{x_i < x_j \text{ if } (i,j) \in Q \text{ but } (j,i) \notin Q,}{x_i = x_j \text{ if both } (i,j), (j,i) \in Q} \right\}$$

Also, one has the following assertions, using the notation of this dictionary:

• for π in Π_n , the subspace denoted X_{π} is the (non-pointed) cone C_{π} , regarding π as a preposet, and

• for any poset P on [n], the open n-dimensional cone denoted \mathcal{K}_P earlier is $\operatorname{relint}(C_P)$ (= $\operatorname{int}(C_P)$). We will need one further dictionary fact.

Proposition 2.14 ([20, Proposition 3.5]). For preposets Q, Q', one has $C_Q \cap C_{Q'} = C_{\overline{Q \cup Q'}}$.

The following definition will help to characterize the set partitions π having X_{π} in $\mathcal{L}^{\text{int}}(\mathcal{K}_P)$.

Definition 2.15. Given a poset P on [n] and a set partition $\pi = \{B_1, \ldots, B_k\}$ in Π_n , define a preposet P/π on the set $\{B_1, \ldots, B_k\}$ as the transitive closure of the (reflexive) binary relation having $(B_i, B_j) \in P/\pi$ whenever there exist $p \in B_i$ and $q \in B_j$ with $p \leq_P q$.

Proposition 2.16. For P a poset on [n] and $\pi = \{B_1, \ldots, B_k\}$ a set partition in Π_n , the following are equivalent.

- (i) $X_{\pi} \in \mathcal{L}^{int}(\mathcal{K}_P)$, that is, one has a nonempty intersection $X_{\pi} \cap \mathcal{K}_P \neq \emptyset$.
- (ii) If $i <_P j$, meaning that $i \leq_P j$ and $i \neq j$, then $(j,i) \notin \overline{P \cup \pi}$.
- (iii) Every block $B_i \in \pi$ is an antichain of P, and the preposet P/π is actually a poset.

We give a proof of Proposition 2.16 toward the end of this subsection, after some discussion and examples.

Definition 2.17. Let P be a poset on [n]. A set partition π of P is called a P-transverse partition if it satisfies one of the equivalent conditions in Proposition 2.16. We denote by $\Pi^{\uparrow}(P)$ the induced subset of Π_n consisting of P-transverse partitions.

Remark 2.18. Aguiar and Mahajan [1, p.230] have a similar concept, which they call a prelinear extension of P. A prelinear extension of P is equivalent to a P-transverse partition π together with a linear ordering on the blocks of π that extends the partial order P/π from Proposition 2.16(iii).

Proposition 2.16 and Corollary 2.6 immediately imply the following corollary.

Corollary 2.19. Let P be a poset on [n]. Then $\Pi^{\uparrow}(P)$ and $\mathcal{L}^{int}(\mathcal{K}_P)$ are isomorphic as posets. Consequently,

$$\mathsf{Poin}(P,t) = \sum_{\pi \in \Pi^{\pitchfork}(P)} |\mu(V,X_{\pi})| \cdot t^{n-\#\mathsf{blocks}(\pi)} = \sum_{\substack{\pi = \{B_1,\dots,B_k\}\\in \ \Pi^{\pitchfork}(P)}} \prod_{i=1}^{\kappa} (\#B_i - 1)! \cdot t^{n-k}.$$

Example 2.20. Let $P := P_2$ be the second poset from Example 2.12, with $x_1 <_P x_2$ and $x_3 <_P x_4$. Then

- $\pi = 13|24$ is *P*-transverse.
- $\pi = 12|3|4$ is not P-transverse as it fails condition (ii): $1 <_P 2$, but $(2,1) \in \pi \subset \overline{P \cup \pi}$.
- $\pi = 14|23$ is not P-transverse, failing condition (ii): $1 <_P 2$, but $(2,1) \in \overline{P \cup \pi}$, though $(2,1) \notin P \cup \pi$.

The six *P*-transverse partitions give this subposet $\Pi^{\uparrow}(P)$ of Π_4 isomorphic to $\mathcal{L}^{\text{int}}(\mathcal{K}_P)$, as in Example 2.11:



It happens that here $|\mu(V, X_{\pi})| = 1$ for all π in $\Pi^{\uparrow}(P)$, so that $\mathsf{Poin}(P, t) = 1 + 4t + t^2$.

Example 2.21. Let P be the following poset:

Then $\pi = \{\{1, 4, 7\}, \{2, 5, 10\}, \{3, 6, 8\}, \{9\}\}$ in Π_{10} is *P*-transverse, represented here by shading the blocks:



Viewed in this way, Proposition 2.16(iii), roughly speaking, states that π is *P*-transverse if and only if its blocks are antichains that can be "stacked without crossings" with respect to the Hasse diagram for *P*.

Proof of Proposition 2.16. We will show a cycle of implications: (i) \Rightarrow (ii) \Rightarrow (iii) \Rightarrow (iii) \Rightarrow (i).

(i) implies (ii):

Assume (i), so that there exists some \mathbf{x} in \mathbb{R}^n lying in the nonempty set

$$\begin{split} X_{\pi} \cap \mathcal{K}_{P} &= X_{\pi} \cap \operatorname{int}(C_{P}) = \operatorname{relint}(X_{\pi} \cap C_{P}) = \operatorname{relint}(C_{\overline{P \cup \pi}}) \\ &= \left\{ \mathbf{x} \in \mathbb{R}^{n} : \begin{array}{c} x_{i} < x_{j} \text{ if } (i,j) \in \overline{P \cup \pi} \text{ but } (j,i) \notin \overline{P \cup \pi}, \\ x_{i} = x_{j} \text{ if both } (i,j), (j,i) \in \overline{P \cup \pi}, \end{array} \right\}, \end{split}$$

where the first equality comes from the definition of \mathcal{K}_P and C_P , the second from the fact that \mathcal{K}_P, C_P are full *n*-dimensional, the third from Proposition 2.14, and the fourth from equation (11) above. Now to see that (ii) holds, given any pair i, j with $i <_P j$, then $x_i < x_j$ since $\mathbf{x} \in \mathcal{K}_P$, but then since $(i, j) \in P \subseteq \overline{P \cup \pi}$, the conditions above imply $(j, i) \notin \overline{P \cup \pi}$, as desired for (ii).

(ii) implies (iii):

Assume (ii) holds. Then every block B of π must be an antichain in P, else there exists $i \neq j$ in B with $i <_P j$, and then $(j, i) \in \pi \subseteq \overline{P \cup \pi}$, contradicting (ii).

Now suppose for the sake of contradiction that P/π is not a poset. Since P/π is a preposet, it can only fail to be antisymmetric, that is, there are blocks $B \neq B'$ of π having both (B, B'), (B', B) in P/π . Since both P, π are transitive binary relations, this means there must exist a (periodic) sequence of elements of the form

$$\cdots \equiv_{\pi} p_1 <_P p_2 \equiv_{\pi} p_3 <_P p_4 \equiv_{\pi} \cdots <_P p_{m-2} \equiv_{\pi} p_{m-1} <_P p_m \equiv_{\pi} p_1 <_P p_2 \equiv \cdots$$

alternating relations (p_i, p_{i+1}) lying in P and in π . Then $p_1 <_P p_2$ and $(p_2, p_1) \in \overline{P \cup \pi}$, contradicting (ii).

(iii) implies (i):

Assume (iii), that is, the blocks of π are antichains of P, and P/π is a poset. One can then reindex the blocks of π such that (B_1, B_2, \ldots, B_k) is a linear extension of P/π . Use this indexing to define a point $\mathbf{x} \in \mathbb{R}^n$ whose p^{th} coordinate $x_p = i$ if p lies in block B_i of π .

We claim \mathbf{x} lies in $X_{\pi} \cap \mathcal{K}_P$, verifying (i). By construction \mathbf{x} lies in X_{π} , since its coordinates are constant within the blocks of π . To verify $\mathbf{x} \in \mathcal{K}_P$, given $p <_P q$, one must check that $x_p < x_q$. Assume that p, q lie in blocks B_i, B_j of π , so that $x_p = i$ and $x_q = j$. Since the blocks of π are antichains in P and $p <_P q$, one has $i \neq j$, and since (B_1, B_2, \ldots, B_k) is a linear extension of P/π , one must have i < j, that is, $x_p < x_q$. \Box

It will help later in identifying P-transverse partitions to also have the following recursive characterization.

Proposition 2.22. Let P be a poset on [n], and π a set partition of [n]. Then π lies in $\Pi^{\uparrow}(P)$ if and only if it contains a block B with these two properties:

- (a) B is a subset of the minimal elements of P, and
- (b) if $\hat{\pi} := \pi \setminus \{B\}$ and \hat{P} is the poset on $[n] \setminus B$ obtained from P by removing the elements in B, then $\hat{\pi} \in \Pi^{\uparrow}(\hat{P})$.

Proof. For the forward implication, assume $\pi \in \Pi^{\uparrow}(P)$ and $B \in \pi$ is a minimal block. We use Proposition 2.16(iii) to show that (a) and (b) hold. For (a), assume there exists some $x \in B$ which is *not* minimal in P, i.e., there is some $y \in P$ with $x >_P y$. The block B is an antichain, so necessarily $y \notin B$ and so y lies in a block $B' \neq B$ of π . But then $B >_{P/\pi} B'$ contradicts the minimality of B in P/π .

For (b), note that the blocks of $\hat{\pi}$ are a subset of the blocks of π , so they are antichains in P and necessarily also antichains in \hat{P} . Furthermore, the preposet $\hat{P}/\hat{\pi}$ on the blocks of $\hat{\pi}$ must be a poset, else if B', B'' were two blocks of $\hat{\pi}$ having $B' \geq_{\hat{P}/\hat{\pi}} B''$ and $B'' \geq_{\hat{P}/\hat{\pi}} B'$, then these same two blocks B', B'' in π and would have have $B' \geq_{P/\pi} B''$ and $B'' \geq_{P/\pi} B'$, a contradiction. For the backward implication, assume π in Π_n has a block B satisfying properties (a), (b). We use Proposition 2.16(i) to show that $\pi \in \Pi^{\uparrow}(P)$. Since $\hat{\pi} \in \Pi^{\uparrow}(\hat{P})$, there is a point $\hat{\mathbf{x}} \in \mathbb{R}^{[n]\setminus B}$ in the (nonempty) set $X_{\hat{\pi}} \cap \mathcal{K}_{\hat{P}}$, i.e., the coordinates of \mathbf{x} are constant within each block of $\hat{\pi}$, and $\hat{\mathbf{x}}_p < \hat{\mathbf{x}}_q$ whenever $p <_{\hat{P}} q$. Let $\mathfrak{m} := \min_{\mathbb{R}} \{ \hat{\mathbf{x}}_p : p \in [n] \setminus B \}$ be the smallest coordinate of $\hat{\mathbf{x}}$, and then extend $\hat{\mathbf{x}} \in \mathbb{R}^{[n]\setminus B}$ to a point $\mathbf{x} \in \mathbb{R}^n$ by assigning all of the new coordinates x_p for $p \in B$ to have the same value, but strictly smaller than \mathfrak{m} , e.g., $x_p := \mathfrak{m} - 1$ for all $p \in B$. One then checks that this \mathbf{x} lies in $X_{\pi} \cap \mathcal{K}_P$: it lies in \mathcal{K}_p due to the fact that B is a subset of the minimal elements of P, and it lies in X_{π} because it is constant on the new block B of π not already in $\hat{\pi}$, as well as constant on the blocks of $\hat{\pi}$.

2.3. More examples of $\Pi^{\uparrow}(P)$.

Example 2.23. Given a poset P on [n], its *dual* or *opposite* poset P^{opp} has the same underlying set [n], but with opposite order relation: $i \leq_P j$ if and only if $j \leq_{P^{opp}} i$. One can readily check that conditions (ii) and (iii) in Proposition 2.16 are self-dual in the sense that π in Π_n is P-transverse if and only if it is P^{opp} -transverse. Consequently, $\Pi^{\uparrow}(P^{opp}) = \Pi^{\uparrow}(P)$, and hence $\mathsf{Poin}(P^{opp}, t) = \mathsf{Poin}(P, t)$.

Example 2.24. Given posets P_1, P_2 , respectively, their ordinal sum $P_1 \oplus P_2$ is the poset whose underlying set is the disjoint union $P_1 \sqcup P_2$, and having order relations $x \leq_{P_1 \oplus P_2} y$ if either

- $x, y \in P_i$ and $x_{\leq P_i} y$ for some i = 1, 2, or
- $x \in P_1$ and $y \in P_2^i$.

If the underlying sets for P_1, P_2 are $[n_1], [n_2]$, one can readily check from either of Proposition 2.16 (ii) or (iii) that a partition π of $[n_1] \sqcup [n_2]$ is $P_1 \oplus P_2$ -transverse if and only if it is of form $\pi = \{A_1, \ldots, A_k, B_1, \ldots, B_\ell\}$ where $\pi_1 = \{A_i\}_{i=1}^k$ and $\pi_2 = \{B_j\}_{j=1}^\ell$ are P_1 -transverse and P_2 -transverse partitions of $[n_1]$ and $[n_2]$, respectively. Bearing in mind that $V = \mathbb{R}^{n_1+n_2} = V_1 \oplus V_2$ where $V_i = \mathbb{R}^{n_i}$ for i = 1, 2, this gives isomorphisms

$$\Pi^{\uparrow}(P \oplus P_2) \cong \Pi^{\uparrow}(P_1) \times \Pi^{\uparrow}(P_2)$$
$$[V, X_{\pi}] \cong [V_1, X_{\pi_1}] \times [V_2, X_{\pi_2}]$$

and therefore also

$$\mathsf{Poin}(P_1 \oplus P_2, t) = \mathsf{Poin}(P_1, t) \cdot \mathsf{Poin}(P_2, t)$$

2.4. Linear extensions, *P*-transverse permutations, and proof of Theorem 1.1. We recall here the bijection between the chambers of braid arrangement A_{n-1} inside a cone \mathcal{K}_P and the linear extensions of *P*. We then define *P*-transverse permutations and use them to combinatorially re-interpret Poin(P, t).

Definition 2.25. Given two posets P, Q on [n], say that Q extends P if $i \leq_P j$ implies $i \leq_Q j$, that is, $P \subseteq Q$ as binary relations on [n], or equivalently, their cones satisfy $\mathcal{K}_Q \subseteq \mathcal{K}_P$. When Q is a total or linear order $\sigma_1 < \cdots < \sigma_n$ on [n], we identify it with a permutation $\sigma = \sigma_1 \ldots \sigma_n$, and call σ a *linear extension* of P. Let LinExt(P) denote the set of all linear extensions of P.

Example 2.2 noted that chambers of the braid arrangement A_{n-1} are of the form \mathcal{K}_{σ} for permutations σ . Then \mathcal{K}_{σ} is a chamber lying in the cone $\mathcal{C}(\mathcal{K}_P)$ if and only if σ lies in LinExt(P), giving a bijection

$$\begin{array}{rcl} \operatorname{LinExt}(P) & \longrightarrow & \mathcal{C}(\mathcal{K}_P) \\ \sigma & \longmapsto & \mathcal{K}_{\sigma}. \end{array}$$

See also [27, Example 1.3]. Consequently, as noted in (4) one has

$$\#\operatorname{LinExt}(P) = \sum_{k \ge 0} c_k(P) = [\operatorname{\mathsf{Poin}}(P, t)]_{t=1}.$$

Example 2.26. The poset P defined by $1 <_P 2$ and $3 <_P 4$ from Example 2.11 has six linear extensions, shown here labeling the chambers in $C(\mathcal{K}_P)$:



Recall our goal of finding combinatorial interpretations for $\mathsf{Poin}(P, t)$. Comparing the expression for $\mathsf{Poin}(P, t)$ given by Corollary 2.19 in terms of *P*-transverse partitions π , and the interpretation of $\mu(V, X_{\pi})$ in terms of permutations given in (10) motivates the following definition.

Definition 2.27. Given a poset P on [n], a P-transverse permutation is a permutation σ in \mathfrak{S}_n for which the set partition π given by the cycles of σ is a P-transverse partition. Denote by $\mathfrak{S}^{\pitchfork}(P)$ the set of all P-transverse permutations.

Corollary 2.19 and equation (10) then immediately imply this interpretation for Poin(P, t).

Thereom 1.1. For any poset P on [n], one has

$$\mathsf{Poin}(P,t) = \sum_{\sigma \in \mathfrak{S}^{\pitchfork}(P)} t^{n - \mathsf{cyc}(\sigma)}.$$

In particular, setting t = 1, one has $\# \operatorname{LinExt}(P) = \#\mathfrak{S}^{\pitchfork}(P)$.

2.5. An aside on cones in reflection arrangements. We digress here to generalize Theorem 1.1 from posets P parametrizing cones in the type A_{n-1} reflection arrangement, to any real reflection arrangement.

We start first at the level of generality of a *complex reflection group* W acting on $V = \mathbb{C}^n$. This means that W is a finite subgroup of $GL(V) \cong GL_n(\mathbb{C})$ generated by (complex, unitary, pseudo-) reflections, which are elements w in W whose fixed space $V^w = H$ is a hyperplane, that is, a codimension one (\mathbb{C}) -linear subspace. Let \mathcal{A}_W denote the arrangement of all such reflecting hyperplanes, and $\mathcal{L}(\mathcal{A}_W)$ its poset of intersections, as before. This generalizes the type A_{n-1} setting, where $W = \mathfrak{S}_n$ and $\mathcal{L}(\mathcal{A}_W) \cong \Pi_n$ is the poset of set partitions of [n]. There is also a well-known generalization of the map $\mathfrak{S}_n \to \Pi_n$ that sends a permutation σ to the set partition $\pi = \{B_1, B_2, \ldots, B_k\}$ whose blocks B_i are the cycles of σ , given by

(12)
$$\begin{array}{ccc} W & \longrightarrow & \mathcal{L}(\mathcal{A}_W) \\ w & \longmapsto & V^w. \end{array}$$

Orlik and Solomon proved [18, §4] the following facts about this map.

Proposition 2.28. For any finite complex reflection group W, the map defined in (12) has these properties:

- (a) The map is well-defined: V^w is an intersection of reflecting hyperplanes, so it lies in $\mathcal{L}(\mathcal{A}_W)$.
- (b) The map surjects $W \twoheadrightarrow \mathcal{L}(\mathcal{A}_W)$.
- (c) The Möbius function values for lower intervals [V, X] in $\mathcal{L}(\mathcal{A}_W)$ can be expressed via this map as

$$\mu(V,X) = \sum_{w \in W: V^w = X} \det(w)$$

Proof. Parts (a) and (b) are rephrasings of [18, Lemma (4.4)] and part (c) is [18, Lemma (4.7)]. However, we include here a shorter proof, due to C. Athanasiadis⁴. Recall that the values $\mu(V, X)$ are uniquely determined by the identity $\delta_{X,V} = \sum_{Y:V \supseteq Y \supseteq X} \mu(V, Y)$ where $\delta_{X,V} = 1$ if X = V and 0 otherwise. It therefore suffices

 $^{^{4}}$ A version of this proof for *real* reflection groups appears (in Greek) within the proof of Theorem 5.1 on pages 33-34 in the Masters Thesis of Athanasiadis' student C. Savvidou.

to check these equalities:

$$\sum_{Y:V\supseteq Y\supseteq X} \left(\sum_{w\in W:V^w=Y} \det(w)\right) = \sum_{w\in W:V^w\supseteq X} \det(w) = \sum_{w\in W_X} \det(w) = \delta_{X,V}$$

where here W_X denotes the subgroup of W that fixes X pointwise. The last equality follows from Steinberg's Theorem [29, Thm. 1.5]: he showed W_X is generated by the reflections whose hyperplane contains X, so that $W_X = \{1\}$ when X = V (implying $\sum_{w \in W_X} \det(w) = \det(1) = 1$), and otherwise if $X \neq V$, summing the (nontrivial) character $\det(-)$ over W_X yields $\sum_{w \in W_X} \det(w) = 0$.

For *real* reflection groups, part (c) above has the following reformulation, generalizing equation (10) above; see [1, Lemma 5.17], [17, §2, pp. 413-414], and [22, Prop. 7.2].

Corollary 2.29. Let W be a finite real reflection group acting on $V = \mathbb{R}^n$. For any X in $\mathcal{L}(\mathcal{A}_W)$, one has

$$\mu(V, X) = (-1)^{n - \dim(X)} \# \{ w \in W : V^w = X \}.$$

Proof. Note W acts orthogonally. We claim that any w acting orthogonally on \mathbb{R}^n has det $(w) = (-1)^{n-\dim(V^w)}$; given this, Lemma 2.28(c) would finish the proof. To see this claim, note that the eigenvalues λ of w lie on the unit circle in \mathbb{C} , so $\lambda \overline{\lambda} = 1$. If the eigenvalue λ has multiplicity m_{λ} , then $m_{\overline{\lambda}} = m_{\lambda}$. Thus

$$\det(w) = \prod_{\lambda \in \mathbb{C}} \lambda^{m_{\lambda}} = (+1)^{m_1} (-1)^{m_{-1}} \prod_{\text{pairs}\{\lambda \neq \overline{\lambda}\}} (\lambda \overline{\lambda})^{m_{\lambda}} = (-1)^{m_{-1}}.$$

Modulo two, however, we have

$$m_{-1} = n - \sum_{\lambda \neq -1} m_{\lambda} = n - m_{+1} - \sum_{\lambda \neq \overline{\lambda}} 2m_{\lambda} = n - m_{+1} = n - \dim(V^w),$$

and so $\det(w) = (-1)^{n - \dim(V^w)}$.

We specialize now to real reflection groups W. Here it is known (see, e.g., [14, Chap. 1]) that W permutes the chambers $\mathcal{C}(\mathcal{A}_W)$ simply transitively. Thus by fixing a choice of base chamber C_0 , every other chamber wC_0 has a unique label by some w in W, giving a bijection $\mathcal{C}(\mathcal{A}_W) \leftrightarrow W$.

Cones \mathcal{K} inside the reflection arrangement \mathcal{A}_W correspond to what were called *parsets* by the third author [21, Chap. 3], or *Coxeter cones* by Stembridge [30], where they were studied as well-behaved generalizations of posets P on [n]. In particular, the set of chambers $\mathcal{C}(\mathcal{K})$ inside \mathcal{K} generalizes the set LinExt(P) of linear extensions of P. For a cone \mathcal{K} in \mathcal{A}_W , we consider as before the subposet $\mathcal{L}^{\text{int}}(\mathcal{K})$ of intersection subspaces interior to \mathcal{K} , playing the role of the P-transverse set partitions $\Pi^{\uparrow}(P)$ in type A_{n-1} . Generalizing the P-transverse permutations $\mathfrak{S}^{\uparrow}(P)$, define the subset

$$W^{\uparrow}(\mathcal{K}) := \{ w \in W : V^w \in \mathcal{L}^{\text{int}}(\mathcal{K}) \}.$$

The real reflection group generalization of Theorem 1.1 is the following.

Theorem 2.30. Any cone \mathcal{K} in the reflection arrangement \mathcal{A}_W for a finite real reflection group W has

$$\mathsf{Poin}(\mathcal{K},t) = \sum_{w \in W^{\pitchfork}(\mathcal{K})} t^{n - \dim(V^w)}$$

In particular, setting t = 1, one has $\#C(\mathcal{K}) = \#W^{\uparrow}(\mathcal{K})$.

Proof.

$$\begin{split} \mathsf{Poin}(\mathcal{K},t) &= \sum_{X \in \mathcal{L}^{\mathrm{int}}(\mathcal{K})} |\mu(V,X)| \cdot t^{n-\dim(X)} \\ &= \sum_{X \in \mathcal{L}^{\mathrm{int}}(\mathcal{K})} \#\{w \in W : V^w = X\} \cdot t^{n-\dim(X)} = \sum_{w \in W^{\oplus}(\mathcal{K})} t^{n-\dim(V^w)} \end{split}$$

where the second equality used Corollary 2.29, and the last equality used the definition of $W^{\uparrow}(\mathcal{K})$.

3. Bijecting P-transverse permutations and linear extensions: proof of Theorem 1.2

The goal in the next few subsections is to define mutually inverse bijections

$$\mathfrak{S}^{\pitchfork}(P) \xrightarrow{\Phi} \operatorname{LinExt}(P)$$

LinExt(P) $\xrightarrow{\Psi} \mathfrak{S}^{\pitchfork}(P)$

along with the notion of *P*-left-to-right maxima for σ in LinExt(*P*), to prove this result from the Introduction.

Theorem 1.2. For any poset P on [n], one has a bijection $\Phi : \mathfrak{S}^{\uparrow}(P) \to \operatorname{LinExt}(P)$ sending the number of cycles to the number of P-left-to-right-maxima. Therefore,

$$\mathsf{Poin}(P,t) = \sum_{\sigma \in \mathrm{LinExt}(P)} t^{n - \mathsf{LRmax}_P(\sigma)}$$

where $\mathsf{LRmax}_P(\sigma)$ denotes the number of P-left-to-right maxima of σ .

To this end, we first recall the special case the antichain poset $P = \text{Antichain}_n$ on [n], where Φ is known as the fundamental bijection [28, Proposition 1.3.1]. A permutation τ in \mathfrak{S}_n sending $i \mapsto \tau(i)$ may be written

- in a one-line notation as $\tau = [\tau(1), \tau(2), \dots, \tau(n)]$, or in a two-line notation $\tau = \begin{pmatrix} 1 & 2 & \cdots & n \\ \tau(1) & \tau(2) & \cdots & \tau(n) \end{pmatrix}$, or
- in various cycle notations that list the τ -orbits on [n], called its cycles, in some arbitrarily chosen order, with each cycle listed as $(j, \tau(j), \tau^2(j), \ldots)$ for some arbitrary choice of the first element j.

One way to make the choices non-arbitrary and put the cycle notation in standard form insists that the first element *i* listed within each cycle $\tau^{(i)}$ is the maximum element of the cycle, and then insists that the cycles $\tau^{(1)}, \tau^{(2)}, \ldots$ are listed with their maximum elements in increasing order as integers, that is, $j_1 < \mathbb{Z}$ $j_2 < \mathbb{Z} \cdots$. The fundamental bijection $\mathfrak{S}_n \to \mathfrak{S}_n$ sends τ to $\Phi(\tau) := \sigma = [\sigma_1, \sigma_2, \ldots, \sigma_n]$ by erasing the parentheses around the standard form cycle notation for τ .

Example 3.1. The permutation $\tau = [7, 5, 9, 4, 2, 8, 3, 6, 1]$ in \mathfrak{S}_9 in one-line notation can also be written in two-line notation and factored according to its τ -orbits or cycles

$$\tau = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 & 9 \\ 7 & 5 & 9 & 4 & 2 & 8 & 3 & 6 & 1 \end{pmatrix} = \begin{pmatrix} 1 & 3 & 7 & 9 \\ 7 & 9 & 3 & 1 \end{pmatrix} \begin{pmatrix} 2 & 5 \\ 5 & 2 \end{pmatrix} \begin{pmatrix} 4 \\ 4 \end{pmatrix} \begin{pmatrix} 6 & 8 \\ 8 & 6 \end{pmatrix}$$

Its cycle notation in standard form and image $\sigma = \Phi(\tau)$ are then

$$\begin{aligned} \tau &= (4)(5,2)(8,6)(9,1,7,3) \\ \Phi(\tau) &= \sigma &= [4,5,2,8,6,9,1,7,3]. \end{aligned}$$

The inverse map Ψ starts with $\sigma = [\sigma_1, \ldots, \sigma_n]$ in one-line notation, and determines where to re-insert the parenthesis pairs in the sequence to obtain the standard form for the cycles of τ . One only needs to know the locations of the left parentheses, since then the right parenthesis locations are determined. There will be one left parenthesis just to the left of each σ_i which is a *left-to-right maximum* (or *LR-maximum* for short) in σ , meaning $\sigma_i = \max_{\mathbb{Z}} \{\sigma_1, \sigma_2, \dots, \sigma_i\}$. It is not hard to check that Φ, Ψ are mutual inverses, and if $\sigma = \Phi(\tau)$, one has $cyc(\tau) = LRmax(\sigma)$, the number of LR-maxima of σ .

3.1. The map $\Phi : \mathfrak{S}^{\uparrow}(P) \to \operatorname{LinExt}(P)$. To define the map $\Phi : \mathfrak{S}^{\uparrow}(P) \to \operatorname{LinExt}(P)$ on a *P*-transverse permutation τ , we will first use the P-transverse partition π whose blocks are the cycles of τ to separate the blocks of π and the elements of P into levels, and then define a notion of essential elements. Recall that because σ lies in $\mathfrak{S}^{\uparrow}(P)$, meaning π lies in $\Pi^{\uparrow}(P)$, the quotient preposet P/π on the blocks of π is actually a poset. This leads to the following definition:

Definition 3.2. Say that a block of π which is minimal in P/π is of Level 1. For $k \geq 2$, the blocks of π of Level k are the minimal ones in the poset obtained from P/π by removing all blocks of Level less than k.

In other words, a block B of π is of Level k if and only if

 $k = \max\{\ell : \text{ there exists a chain } B =: B_1 >_{P/\pi} B_2 >_{P/\pi} \cdots >_{P/\pi} B_\ell\},\$

or even more concretely, k is the maximum among integers ℓ with the property that there exist blocks $B =: B_1, B_2, \dots, B_\ell$ of π and elements $x_i >_P y_{i+1}$ with $x_i \in B_i, y_{i+1} \in B_{i+1}$ for each $i = 1, 2, \dots, \ell - 1$.

For an element x in [n], define the Level of x to be the Level of the unique block of π containing x.

Definition 3.3. An element x is essential if x has Level k and there exists some y of Level k - 1 with $x >_P y$; by convention, all Level 1 elements x are essential.

In order to define the map $\Phi : \mathfrak{S}^{\uparrow}(P) \to \operatorname{LinExt}(P)$, we first introduce a standard form for a P-transverse permutation τ . Let τ have cycle partition $\pi = \{B_1, \ldots, B_m\}$, lying in $\Pi^{\uparrow}(P)$. List the cycles of τ in order $\tau^{(1)}, \tau^{(2)}, \ldots, \tau^{(m)}$, with the block B_i of π corresponding to the cycle $\tau^{(i)}$, and write the cycle $\tau^{(i)}$ as $\tau^{(i)} = (x_i, \tau(x_i), \tau^2(x_i), \ldots)$ for some x_i in B_i . Then this listing is the standard form of τ if and only if it has the following properties:

- If blocks B_i, B_j have Levels k, k+1 in π , respectively, then their indices satisfy $i \leq \mathbb{Z} j$.
- For each *i*, the first element x_i listed in the cycle $\tau^{(i)} = (x_i, \tau(x_i), \tau^2(x_i), \dots)$ is the maximal essential element of B_i ; by Lemma 3.5(b), below, every block B_i contains an essential element.
- If B_i, B_j are blocks of Level k with $i <_{\mathbb{Z}} j$ then $x_i <_{\mathbb{Z}} x_j$.

Following the fundamental bijection, the map $\Phi : \mathfrak{S}^{\uparrow}(P) \to \operatorname{LinExt}(P)$ is defined by putting $\tau \in \mathfrak{S}^{\uparrow}(P)$ into standard form and erasing parenthesis. The following example illustrates this process.

Example 3.4. Let P be the following poset on [13]:



Let

$$\begin{aligned} \tau &= (4)(6,3)(9)(10)(11,7)(12,5,8,2)(13,1) &\in \mathfrak{S}^{\pitchfork}(P), \text{ so that} \\ \pi &= \{\{4\},\{3,6\},\{9\},\{7,10\},\{2,5,8,12\},\{1,13\}\} &\in \Pi^{\pitchfork}(P), \end{aligned}$$

and the poset P/π , drawn as a poset on the cycles of τ , looks as follows:



One can check that

- the Level 1 cycles are (13, 1), (9), (4), with essential elements 1, 4, 9, 13,
- there is one Level 2 cycle (11,7), with one essential element 7,
- the Level 3 cycles are (6,3), (12,5,8,2), with essential elements 2,3,5,
- there is one Level 4 cycle (10), with one essential element 10.

Here is τ in standard form, with essential elements overlined, and Levels separated by bars:

$$\tau = (\overline{4})(\overline{9})(\overline{13},\overline{1}) \mid (\overline{7},11) \mid (\overline{3},6)(\overline{5},8,\overline{2},12) \mid (\overline{10}),$$

Removing the parentheses (and bars), one obtains its image under Φ :

 $\Phi(\tau) = \sigma = [4, 9, 13, 1, 7, 11, 3, 6, 5, 8, 2, 12, 10] \in \text{LinExt}(P).$

The next lemma is used to prove the image of Φ lies in $\operatorname{LinExt}(P)$, and $\mathfrak{S}^{\pitchfork}(P) \xrightarrow{\Phi} \operatorname{LinExt}(P)$ is bijective.

Lemma 3.5. Let P be a poset on [n] and $\tau \in \mathfrak{S}^{\uparrow}(P)$. Then the following properties hold:

- (a) For each $k \ge 1$, the Level k elements of [n] form an antichain in P.
- (b) Every cycle contains at least one essential element.
- (c) The image $\Phi(\tau)$ of τ is a linear extension of P.

Proof. For (a), assume that there were two comparable elements $x <_P y$ with x, y both of Level k. Either x, y lie in the same block of π , contradicting P-transversality, or they lie in *different* blocks of π , which would be comparable in P/π , contradicting both blocks being of Level k.

For (b), note that our most concrete description of a block *B* having Level *k* shows that there exist blocks $B =: B_1, B_2, \ldots, B_k$ of π and elements $x_i >_P y_{i+1}$ with $x_i \in B_i, y_{i+1} \in B_{i+1}$ for each $i = 1, 2, \ldots, k-1$. But then this shows that the block B_2 must be of Level k-1, and $x_1 >_P y_2$ shows x_1 is essential in $B_1(=B)$.

For (c), one can show via induction on k that the restriction of $\Phi(\tau)$ to the order ideal of elements of P having Level at most k is a linear extension. In both the base case k = 1, and in the inductive step, one notes that one can add in the elements of Level k in any order, because they form an antichain by part (a).

3.2. The inverse map Ψ . To define the inverse map $\Psi = \Phi^{-1}$ on a linear extension σ , we proceed similarly to the previous subsection. We first cut σ into consecutive strings, suggestively called *Levels*, define a notion of *essential* elements of σ , and a notion of *P-left-to-right-maximum*.

Definition 3.6. Given $\sigma = [\sigma_1, \ldots, \sigma_n]$ in LinExt(P), we recursively break σ into disjoint contiguous sequences $[\sigma_i, \sigma_{i+1}, \ldots, \sigma_{i+j}]$, each forming an antichain of P, and each maximal in the sense that the slightly longer sequence $[\sigma_i, \sigma_{i+1}, \ldots, \sigma_{i+j}, \sigma_{i+j+1}]$ is not an antichain of P:

- Let $[\sigma_1, \sigma_2, \ldots, \sigma_r]$ be the longest initial segment of σ whose underlying set $\{\sigma_1, \sigma_2, \ldots, \sigma_r\}$ is an antichain of P; call $\{\sigma_1, \sigma_2, \ldots, \sigma_r\}$ the *Level* 1 elements of σ .
- The Level 2 elements of σ are $\{\sigma_{r+1}, \sigma_{r+2}, \ldots, \sigma_s\}$, where $[\sigma_{r+1}, \sigma_{r+2}, \ldots, \sigma_s]$ is the longest initial segment of $[\sigma_{r+1}, \sigma_{r+2}, \ldots, \sigma_n]$ that forms an antichain in P.
- Similarly, for $k \ge 3$, the Level k elements of σ are defined as follows: if the union of all elements of Levels $1, 2, \ldots, k 1$ are $\{\sigma_1, \sigma_2, \ldots, \sigma_t\}$, then the set of Level k elements is the underlying set of the longest initial segment in $[\sigma_{t+1}, \sigma_{t+2}, \ldots, \sigma_n]$ that forms an antichain in P.

Definition 3.7. As in the previous section, say x in σ is *essential* if x has Level k in σ and there exists an element y of Level k - 1 in σ with $x >_P y$; again by convention, Level 1 elements of σ are all essential.

Definition 3.8. Say that an element x is a P-left-to-right-maximum of σ , or P-LR-maximum for short, if x is essential in σ , and x appears as a LR-maximum in the usual sense among the subsequence of essential elements of σ having the same Level as x. In other words, if x has Level k, it is a P-LR-maximum if the subsequence of essential Level k elements is $[\sigma_{i_1}, \sigma_{i_2}, \ldots, \sigma_{i_r}]$ for some indices $i_1 < i_2 < \cdots < i_r$, and there is some j with $1 \le j \le r$ for which $\sigma_{i_j} = x = \max\{\sigma_{i_1}, \sigma_{i_2}, \ldots, \sigma_{i_j}\}$. We denote the number of P-LR maxima of σ by LRmax_P(σ).

A map Ψ : LinExt $(P) \to \mathfrak{S}_n$ can now be defined in much the same way as for the fundamental bijection: starting with $\sigma = [\sigma_1, \ldots, \sigma_n]$ in LinExt(P), one must determine where to re-insert the parenthesis pairs in the sequence to recover the cycles of τ . In fact, one only needs to know the locations of the left parentheses, since then the right parenthesis locations are determined, and there will be one left parenthesis just to the left of each $x = \sigma_i$ which is a *P*-LR maximum.

Example 3.9. Let P be the poset in Example 3.4 and let

 $\sigma = [4, 9, 13, 1, 7, 11, 3, 6, 5, 8, 2, 12, 10] \in \text{LinExt}(P),$

which is $\Phi(\tau)$ from Example 3.4. The Level decomposition of σ , with essential elements overlined, looks like

$$\overline{4}, \overline{9}, \overline{13}, \overline{1} \mid \overline{7}, 11 \mid \overline{3}, 6, \overline{5}, 8, \overline{2}, 12 \mid \overline{10}.$$

Now create cycles by placing left parentheses just before each *P*-LR maximum:

 $(\overline{4})(\overline{9})(\overline{13},\overline{1}) \mid (\overline{7},11) \mid (\overline{3},6)(\overline{5},8,\overline{2},12) \mid (\overline{10}).$

The resulting cycle structure gives the *P*-transverse permutation $\Psi(\sigma)$:

 $\Psi(\sigma) = (4)(9)(13,1)(7,11)(3,6)(5,8,2,12)(10),$

which is the *P*-transverse permutation τ in Example 3.4.

Note that it is not yet clear that the image of Ψ lies in $\mathfrak{S}^{\uparrow}(P)$, but it will follow from the proof of Theorem 1.2. First we need a technical lemma.

Lemma 3.10. Let $\sigma \in \text{LinExt}(P)$ and $\tau \in \mathfrak{S}^{\uparrow}(P)$ such that $\sigma = \Phi(\tau)$. Then the set of Level k elements of τ is precisely the set of Level k elements of σ .

Proof. We prove this by induction on k. For the base case (k = 1), note that the Level 1 elements of τ will form an initial segment $\{\sigma_1, \sigma_2, \ldots, \sigma_\ell\}$ of $\sigma = [\sigma_1, \ldots, \sigma_n]$, by the definition of Φ . These Level 1 elements of τ will also form an antichain of P by Lemma 3.5(a). On the other hand, we claim that the longer initial segment $\{\sigma_1, \sigma_2, \ldots, \sigma_\ell, \sigma_{\ell+1}\}$ cannot form an antichain in P, because $\sigma_{\ell+1}$ is of Level 2 in τ by definition of Φ , and it is also essential in τ because it is leftmost in its cycle in the standard form for τ , and all such elements are essential. Thus $\sigma_{\ell+1} >_P \sigma_i$ for some $i = 1, 2, \ldots, \ell$, showing that the longer segment is not an antichain of P. By definition of Ψ , this means that the Level 1 elements of σ will be those in the shorter segment $\{\sigma_1, \sigma_2, \ldots, \sigma_\ell\}$.

For the inductive step $(k \ge 2)$, we perform the same argument as in the base case, but replace P, τ, σ with their counterparts $\hat{P}, \hat{\tau}, \hat{\sigma}$ in which the elements of Level $1, 2, \ldots, k-1$ have been removed. One must check that $\hat{\tau}$ lies in $\mathfrak{S}^{\uparrow}(\hat{P})$, but this is straightforward, because if $\sigma, \hat{\sigma}$ have cycle partitions $\pi, \hat{\pi}$, then $\hat{P}/\hat{\pi}$ is obtained from P/π by removing its minimal blocks.

Proof of Theorem 1.2. Note $\#\mathfrak{S}^{\pitchfork}(P) = \#\operatorname{LinExt}(P)$ by Theorem 1.1, and Φ maps $\mathfrak{S}^{\pitchfork}(P) \to \operatorname{LinExt}(P)$ by Lemma 3.5(c). We therefore claim that it suffices to check $\Psi(\Phi(\tau)) = \tau$ for all τ in $\mathfrak{S}^{\pitchfork}(P)$. This will imply that Φ is injective, hence bijective, with Ψ its inverse bijection, and thus the image of Ψ is $\mathfrak{S}^{\pitchfork}(P)$. Note that by construction, if $\Psi(\sigma) = \tau$, then $\operatorname{LRmax}_P(\sigma) = \operatorname{cyc}(\tau)$, so Theorem 1.2 would follow.

By Lemma 3.10, we have if $\sigma = \Phi(\tau)$, then the sets of Level k elements of σ and τ coincide. It follows immediately from Lemma 3.10 that the *essential* elements in σ and τ coincide, since in each case, their definition uses only the order P and the partition by Levels. Therefore, we can focus our attention on each Level k separately, where the definition of Φ and Ψ coincides almost exactly with their definition in the fundamental bijection, ignoring the non-essential elements carried along in each Level. It then follows that $\Psi(\Phi(\tau)) = \tau$ via the same argument as for the fundamental bijection.

4. Disjoint Unions of Chains and proofs of Theorems 1.3 and 1.4

The goal of this section is to understand Poin(P, t) for a poset P that is a disjoint union of chains. Although Theorem 1.2 applies to *any* poset, when P is a disjoint union of chains, there turns out to be another elegant expression for Poin(P, t) stemming from Foata's theory of multiset permutations, generalizing equation (8) for the antichain poset P.

In Subsection 4.1 we review Foata's theory of multiset permutations, in particular his work with the *inter*calation product and prime cycle decompositions. Subsection 4.2 reviews its relation to partial commutation monoids. Subsection 4.3 shows how the results in Section 3 can be rephrased in terms of multiset permutations when P is a disjoint union of chains. Theorem 1.3 is also proved in this subsection. Finally Subsection 4.4 employs Foata's theory to give a generalization of MacMahon's Master Theorem which specializes to Theorem 1.4, a generating function compiling the Poincaré polynomials for disjoint unions of chains.

4.1. Multiset Permutations. This subsection gives background on the theory of multiset permutations as introduced by Foata in his PhD thesis [9, Section 3.2], and extended in later publications [10, Chapters 3-5]. It also appears in Knuth [15, Section 5.1.2].

Definition 4.1. Recall that a *(weak) composition* $\overline{a} = (a_1, \ldots, a_\ell)$ of n is a sequence of nonnegative integers having sum $|a| := \sum_i a_i = n$. We will regard \overline{a} as specifying the multiplicities in a *multiset* $M(\overline{a}) := \{1^{a_1}, 2^{a_2}, \ldots, \ell^{a_\ell}\}$, that is, a set with repetitions

$$M(\overline{a}) = \{\underbrace{1, 1, \dots, 1}_{a_1 \text{ times}}, \underbrace{2, 2, \dots, 2}_{a_2 \text{ times}}, \dots, \underbrace{\ell, \ell, \dots, \ell}_{a_\ell \text{ times}}\}.$$

A multiset permutation $\sigma = [\sigma_1, \ldots, \sigma_n]$ is a rearrangement of the elements of $M(\overline{a})$, which we will often write in a two-line notation that generalizes that of permutations:

$$\sigma = \begin{pmatrix} 1 & \cdots & 1 & 2 & \cdots & 2 & \cdots & \ell & \cdots & \ell \\ \sigma_1 & \cdots & \sigma_{a_1} & \sigma_{a_1+1} & \cdots & \sigma_{a_1+a_2} & \cdots & \sigma_{a_1+\dots+a_{\ell-1}+1} & \cdots & \sigma_n \end{pmatrix}.$$

We denote the set of all multiset permutations of $M(\overline{a})$ by $\mathfrak{S}_{M(\overline{a})}$. For any $\sigma \in \mathfrak{S}_{M(\overline{a})}$, we call $M(\overline{a})$ the support of σ , and write $M(\overline{a}) = \operatorname{supp}(\sigma)$.

Example 4.2. The composition $\overline{a} = (2, 3, 2, 3)$ gives the multiplicities of the multiset

$$M(\overline{a}) = \{1^2, 2^3, 3^2, 4^3\} = \{1, 1, 2, 2, 2, 3, 3, 4, 4, 4\}.$$

Then the following multiset permutation σ is an element of $\mathfrak{S}_{M(\overline{\alpha})}$:

$$\sigma = \begin{pmatrix} 1 & 1 & 2 & 2 & 2 & 3 & 3 & 4 & 4 \\ 2 & 4 & 4 & 3 & 1 & 2 & 1 & 3 & 4 & 2 \end{pmatrix}$$

Foata [9, §3.2] defined an associative *intercalation product* operation on multiset permutations $(\sigma, \rho) \mapsto \sigma \intercal \rho$. Knuth [15, §5.1.2] describes it algorithmically: think of σ, ρ in two-line notation as sequences of columns $\binom{i}{j}$, and juxtapose these sequences of columns. Then perform swaps to *sort* the columns according to their top entries, never swapping two with the *same* top entry. For example,

$$\begin{pmatrix} 2 & 3 & 4 \\ 4 & 2 & 3 \end{pmatrix} \mathsf{T} \begin{pmatrix} 1 & 1 & 2 & 2 & 3 & 4 & 4 \\ 2 & 4 & 3 & 1 & 1 & 4 & 2 \end{pmatrix} = \begin{pmatrix} 2 & 3 & 4 \\ 4 & 2 & 3 \\ 2 & 4 & 3 & 1 & 1 & 4 & 2 \end{pmatrix}$$
$$= \begin{pmatrix} 1 & 1 & 2 & 2 & 3 & 3 & 4 & 4 \\ 2 & 4 & 4 & 3 & 1 & 2 & 1 & 3 & 4 & 2 \end{pmatrix}.$$

Definition 4.3. For each ℓ , the *intercalation monoid* Int_{ℓ} is the submonoid of all multiset permutations σ whose support $M = \{1^{a_1}, 2^{a_2}, \ldots, \ell^{a_\ell}\}$ involves only the letters in $\{1, 2, \ldots, \ell\}$. The empty permutation () is the identity element for T , since () $\mathsf{T} \sigma = \sigma = \sigma \mathsf{T}$ ().

Note that, just as permutations in the symmetric group \mathfrak{S}_n do not commute in general, the monoid $\operatorname{Int}_{\ell}$ is not commutative. For example

$$\begin{pmatrix} 1 & 2 \\ 2 & 1 \end{pmatrix} \mathsf{T} \begin{pmatrix} 1 & 3 \\ 3 & 1 \end{pmatrix} = \begin{pmatrix} 1 & 1 & 2 & 3 \\ 2 & 3 & 1 & 1 \end{pmatrix} \neq \begin{pmatrix} 1 & 1 & 2 & 3 \\ 3 & 2 & 1 & 1 \end{pmatrix} = \begin{pmatrix} 1 & 3 \\ 3 & 1 \end{pmatrix} \mathsf{T} \begin{pmatrix} 1 & 2 \\ 2 & 1 \end{pmatrix}.$$

However, one can check that $\sigma \intercal \rho = \rho \intercal \sigma$ when σ, ρ are *disjoint*, that is, $supp(\sigma) \cap supp(\rho) = \emptyset$.

Definition 4.4. Say σ in Int_{ℓ} is *prime* if the only factorizations $\sigma = \rho \tau \tau$ have either $\rho = ()$ or $\tau = ()$.

Example 4.5. The permutation $\begin{pmatrix} 2 & 4 & 5 & 7 \\ 5 & 7 & 4 & 2 \end{pmatrix}$ is prime. However, $\begin{pmatrix} 1 & 1 & 2 & 3 \\ 2 & 3 & 1 & 1 \end{pmatrix}$ is not prime, since $\begin{pmatrix} 1 & 1 & 2 & 3 \\ 2 & 3 & 1 & 1 \end{pmatrix} = \begin{pmatrix} 1 & 2 \\ 2 & 1 \end{pmatrix} \intercal \begin{pmatrix} 1 & 3 \\ 3 & 1 \end{pmatrix}$.

On the other hand $\begin{pmatrix} 2 & 4 & 5 & 7 \\ 5 & 7 & 2 & 4 \end{pmatrix}$ is not prime, even though its support is multiplicity free, since

$$\begin{pmatrix} 2 & 4 & 5 & 7 \\ 5 & 7 & 2 & 4 \end{pmatrix} = \begin{pmatrix} 2 & 5 \\ 5 & 2 \end{pmatrix} \mathsf{T} \begin{pmatrix} 4 & 7 \\ 7 & 4 \end{pmatrix} = \begin{pmatrix} 4 & 7 \\ 7 & 4 \end{pmatrix} \mathsf{T} \begin{pmatrix} 2 & 5 \\ 5 & 2 \end{pmatrix}.$$

It is not obvious, but turns out to be true that σ is prime if and only if both

- $supp(\sigma) = M$ is multiplicity free, that is, M is a set not a multiset, and
- σ has only one cycle when considered as an ordinary permutation of the set M.

We therefore call prime elements prime cycles. More generally, one has the following.

Theorem 4.6 (Foata, 1969 [10, 15]). Let σ be a multiset permutation. Then σ has a decomposition into a product of prime cycles. That is, there exist $k \ge 0$ prime cycles $\sigma^{(1)}, \ldots, \sigma^{(k)}$ such that

(13)
$$\sigma = \sigma^{(1)} \mathsf{T} \sigma^{(2)} \mathsf{T} \cdots \mathsf{T} \sigma^{(k)}.$$

Further, this cycle decomposition of σ is unique up to successively interchanging pairs of adjacent prime cycles with disjoint support. In particular k is unique.

Definition 4.7. Call $pcyc(\sigma) := k$ the number of prime cycles in the decomposition of σ from Theorem 4.6. **Example 4.8.** The element σ from Example 4.2 has $pcyc(\sigma) = 4$ and two prime cycle decompositions

$$\begin{pmatrix} 1 & 1 & 2 & 2 & 2 & 3 & 3 & 4 & 4 & 4 \\ 2 & 4 & 4 & 3 & 1 & 2 & 1 & 3 & 4 & 2 \end{pmatrix} = \begin{pmatrix} 2 & 3 & 4 \\ 4 & 2 & 3 \end{pmatrix} \mathsf{T} \begin{pmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \end{pmatrix} \mathsf{T} \begin{pmatrix} 4 \\ 4 \end{pmatrix} \mathsf{T} \begin{pmatrix} 1 & 2 & 4 \\ 4 & 1 & 2 \end{pmatrix}$$
$$= \begin{pmatrix} 2 & 3 & 4 \\ 4 & 2 & 3 \end{pmatrix} \mathsf{T} \begin{pmatrix} 4 \\ 4 \end{pmatrix} \mathsf{T} \begin{pmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \end{pmatrix} \mathsf{T} \begin{pmatrix} 1 & 2 & 4 \\ 4 & 1 & 2 \end{pmatrix} .$$

We describe here an algorithm to find a prime cycle decomposition of a multiset permutation, which can be deduced from [15, §5.1.2], and illustrate how it produces the first of the two decompositions in Example 4.8. Encode a multiset permutation σ in $\mathfrak{S}_{M(\overline{\alpha})}$ as two pieces of data:

- a directed graph D_{σ} on vertex set $\{1, 2, \dots, \ell\}$ having one copy of the directed arc $i \to j$ for each occurrence of the column $\binom{i}{j}$ in its two-line notation, along with
- specification for each vertex x in $\{1, 2, ..., \ell\}$ the linear ordering of the arcs $x \to y$ emanating from x, indicating the left-to-right ordering of the corresponding columns in the two-line notation.

The resulting digraphs are those with the outdegree equal to the indegree equal to a_i for each *i*. E.g., the σ from Example 4.8 has this directed graph D_{σ} , with linear orderings indicated on the arcs out of each vertex:



With this identification, one factors σ recursively. First produce a prime cycle $\sigma^{(1)}$ for which

(14)
$$\sigma = \sigma^{(1)} \mathsf{T} \hat{\sigma}$$

via the following algorithm that takes a directed walk in D_{σ} .

- Start at the smallest vertex i_0 in $\{1, 2, ..., \ell\}$ with outdegree $a_{i_0} \ge 1$, and follow its first outward arc $i_0 \rightarrow i_1$. Then follow i_1 's first outward arc $i_1 \rightarrow i_2$, follow i_2 's first outward arc $i_2 \rightarrow i_3$, etc.
- Repeat until first arriving at a previously-visited⁵ vertex i_s , say $i_s = i_r$ with r < s; possibly r = s 1.
- The directed *circuit* C of arcs $i_r \to i_{r+1} \to i_{r+2} \to \cdots \to i_{s-1} \to i_s (=i_r)$ corresponds to a prime cycle $\sigma^{(1)}$ that one can factor out to the left as in (14): by construction, each of its corresponding columns $\binom{i_t}{i_t}$ occurs as the *leftmost* column of σ having i_t as its top element.
- Complete the factorization recursively, replacing σ by $\hat{\sigma}$, removing the arcs C from D_{σ} to give $D_{\hat{\sigma}}$.

Example 4.9. Here is the algorithm for σ above, with dotted arrows showing the directed walks in D_{σ} :



⁵During the process, when one enters a new vertex i_j along an arc $i_{j_1} \rightarrow i_j$, there will always be at least one outward arc $i_j \rightarrow i_{j+1}$ leaving i_j , because each vertex started with its indegree matching its outdegree.

4.2. Partial Commutation Monoids. It will be helpful to view the intercalation monoid Int_{ℓ} as a partial commutation monoid. We briefly review some relevant facts about partial commutation monoids.

Definition 4.10. Given a set \mathbb{A} , which we call an *alphabet* and a subset of its pairs $C \subseteq {\binom{\mathbb{A}}{2}}$, the associated *partial commutation monoid* \mathcal{M} is defined to be the set of equivalence classes on words $\alpha_1 \alpha_2 \dots \alpha_k$ in the alphabet \mathbb{A} under the equivalence relation

(15)
$$\alpha_1 \alpha_2 \dots \alpha_i \alpha_{i+1} \dots \alpha_k \equiv \alpha_1 \alpha_2 \dots \alpha_{i+1} \alpha_i \dots \alpha_k$$

if $\{\alpha_i, \alpha_{i+1}\} \in C$.

From this perspective, Foata's Theorem 4.6 asserts that Int_{ℓ} is a partial commutation monoid, whose associated alphabet A is the set of all prime cycles, and C the pairs of prime cycles with disjoint supports.

For later use, we point out the following (nontrivial) proposition, see [15, §5.1.2, Exercise 11] and [28, Exercise 3.123]. Given a factorization of an element $\alpha = \alpha_1 \alpha_2 \dots \alpha_k$ in \mathcal{M} a partial commutation monoid, define a poset \mathcal{P}_{α} on [k] as the transitive closure of the binary relation containing $(i, j) \in \mathcal{P}_{\alpha}$ when $i <_{\mathbb{Z}} j$ and either $\alpha_i = \alpha_j$ or $\alpha_i \alpha_j \neq \alpha_j \alpha_i$.

Proposition 4.11. Given a factorization of $\alpha = \alpha_1 \alpha_2 \dots \alpha_k$ in \mathcal{M} a partial commutation monoid,

- (1) \mathcal{P}_{α} does not depend on the choice of factorization of α , and
- (2) there is a bijection between $\operatorname{LinExt}(\mathcal{P}_{\alpha})$ and the factorizations of α given by

$$(i_1,\ldots,i_k)\mapsto \alpha_{i_1}\ldots\alpha_{i_k}$$

Example 4.12. The multiset permutation σ from Example 4.8 had two prime cycle factorizations

$$\sigma = \sigma^{(1)} \mathsf{T} \sigma^{(2)} \mathsf{T} \sigma^{(3)} \mathsf{T} \sigma^{(4)}$$
$$= \sigma^{(1)} \mathsf{T} \sigma^{(3)} \mathsf{T} \sigma^{(2)} \mathsf{T} \sigma^{(4)}$$

corresponding to the two linear extensions of the poset \mathcal{P}_{σ} on [4] with this Hasse diagram:



4.3. Connection with linear extensions and *P*-transverse permutations. We wish to use Foata's prime cycle decomposition to define a bijection $\text{LinExt}(P_{\overline{a}}) \to \mathfrak{S}^{\uparrow}(P_{\overline{a}})$, and use this to prove Theorem 1.3.

We begin with an easy identification of $\operatorname{LinExt}(P_{\overline{a}})$ with $\mathfrak{S}_{M(\overline{a})}$. For this purpose, given any weak composition $\overline{a} = (a_1, \ldots, a_\ell)$ of n, consider two labelings of $P_{\overline{a}}$, one by the elements $\{1, 2, \ldots, n\}$ which we will call the *standardized labeling*, and the second by the elements of the multiset $M(\overline{a})$, which we will call the *multiset labeling*. The standardized labeling labels the first chain \mathbf{a}_1 by $1, 2, \ldots, a_1$ from bottom-to-top, then the second chain \mathbf{a}_2 by $a_1 + 1, \ldots, a_1 + a_2$ from bottom-to-top, and so on. The multiset labeling labels the elements in the first chain \mathbf{a}_1 all by 1, the second chain \mathbf{a}_2 all by 2, etc.

Example 4.13. For n = 10 and $\overline{a} = (2, 3, 2, 3)$, the standardized and multiset labelings of $P_{\overline{a}}$ are

(5) (10)	$(2) \qquad (4)$
2479	1 2 3 4
(1) (3) (6) (8)	(1) (2) (3) (4)

With this in hand, the following proposition is a straightforward observation.

Proposition 4.14. For any weak composition \overline{a} of n, one has a bijection

$$\begin{array}{rcl} \operatorname{LinExt}(P_{\overline{a}}) & \longrightarrow & \mathfrak{S}_{M(\overline{a})} \\ \lambda = [\lambda_1, \dots, \lambda_n] & \longmapsto & \sigma = [\sigma_1, \dots, \sigma_n] \end{array}$$

replacing λ_i by its corresponding multiset label σ_i , that is, if λ_i lies on the j^{th} chain a_j in $P_{\overline{a}}$, then $\sigma_i := j$.

Proof. The inverse map recovers λ from σ by labeling the a_j occurrences of the value j within σ from left-to-right with the integers in the interval $[a_1 + a_2 + \cdots + a_{j-1} + 1, a_1 + a_2 + \cdots + a_{j-1} + a_j]$.

Example 4.15. For $\overline{a} = (2, 3, 2, 3)$, this bijection maps $\lambda = [3, 8, 9, 6, 1, 4, 2, 7, 10, 5]$ in LinExt($P_{\overline{a}}$) to

$$\sigma = [2, 4, 4, 3, 1, 2, 1, 3, 4, 2] = \begin{pmatrix} 1 & 1 & 2 & 2 & 2 & 3 & 3 & 4 & 4 & 4 \\ 2 & 4 & 4 & 3 & 1 & 2 & 1 & 3 & 4 & 2 \end{pmatrix}.$$

We can now define a map φ : LinExt $(P_{\overline{a}}) \to \mathfrak{S}_n$, which will turn out to be a bijection onto $\mathfrak{S}^{\uparrow}(P_{\overline{a}})$.

Definition 4.16. Fix a weak composition \overline{a} of n. Given λ in LinExt $(P_{\overline{a}})$,

- let $\sigma \in \mathfrak{S}_{M(\overline{\alpha})}$ be its corresponding multiset permutation from Proposition 4.14,
- label the entries in the top line of σ 's two-line notation with subscripts $1, 2, \ldots, n$ from left-to-right,
- use Foata's Theorem 4.6 to decompose $\sigma = \sigma^{(1)} \intercal \cdots \intercal \sigma^{(\ell)}$ into prime cycles $\sigma^{(i)}$, carrying along the subscripts in the top line, and finally
- replace each prime cycle $\sigma^{(i)}$ with the cyclic permutation $\tau^{(i)}$ of the subscripts of its top line.

Then $\varphi(\lambda) := \tau = \tau^{(1)} \cdots \tau^{(\ell)}$ in \mathfrak{S}_n .

Example 4.17. We continue Example 4.15. Let $\overline{a} = (2,3,2,3)$ and $\lambda = [3,8,9,6,1,4,2,7,10,5]$ in LinExt($P_{\overline{a}}$). Subscript the top line of its corresponding σ in $\mathfrak{S}_{M(\overline{a})}$, and factor as in Theorem 4.6, carrying along subscripts:

$$\sigma = \begin{pmatrix} 1_1 & 1_2 & 2_3 & 2_4 & 2_5 & 3_6 & 3_7 & 4_8 & 4_9 & 4_{10} \\ 2 & 4 & 4 & 3 & 1 & 2 & 1 & 3 & 4 & 2 \end{pmatrix}$$
$$= \begin{pmatrix} 2_3 & 3_6 & 4_8 \\ 4 & 2 & 3 \end{pmatrix} \mathsf{T} \begin{pmatrix} 1_1 & 2_4 & 3_7 \\ 2 & 3 & 1 \end{pmatrix} \mathsf{T} \begin{pmatrix} 4_9 \\ 4 \end{pmatrix} \mathsf{T} \begin{pmatrix} 1_2 & 2_5 & 4_{10} \\ 4 & 1 & 2 \end{pmatrix}$$
$$= (2_3, 4_8, 3_6) \mathsf{T} (1_1, 2_4, 3_7) \mathsf{T} (4_9) \mathsf{T} (1_2, 4_{10}, 2_5)$$
$$= \sigma^{(1)} \mathsf{T} \sigma^{(2)} \mathsf{T} \sigma^{(3)} \mathsf{T} \sigma^{(4)}$$

Replacing each prime cycle $\sigma^{(i)}$ with the cycle $\tau^{(i)}$ on its subscripts gives $\varphi(\lambda) = \tau \in \mathfrak{S}_{10}$:

$$\varphi(\lambda) = \tau = \tau^{(1)} \tau^{(2)} \tau^{(3)} \tau^{(4)} = (3, 8, 6)(1, 4, 7)(9)(2, 10, 5).$$

We can now prove Theorem 1.3, whose statement we recall here.

Theorem 1.3. For any composition \overline{a} of n, the disjoint union $P_{\overline{a}}$ of chains has a bijection

$$\begin{array}{cccc} \operatorname{LinExt}(P_{\overline{a}}) & \longrightarrow & \mathfrak{S}^{\pitchfork}(P_{\overline{a}}) \\ \sigma & \longmapsto & \tau \end{array}$$

with $cyc(\tau) = pcyc(\sigma)$, the number of prime cycles in Foata's unique decomposition for σ . Thus

$$\mathsf{Poin}(P_{\overline{a}}, t) = \sum_{\sigma \in \mathrm{LinExt}(P_{\overline{a}})} t^{n - \mathsf{pcyc}(\sigma)}.$$

Proof. We claim that the above map φ : LinExt $(P_{\overline{a}}) \to \mathfrak{S}_n$ is the desired bijection. Since Theorem 1.1 showed $\# \operatorname{LinExt}(P_{\overline{a}}) = \#\mathfrak{S}^{\oplus}(P_{\overline{a}})$, it suffices to show that the image of φ lies in $\mathfrak{S}^{\oplus}(P_{\overline{a}})$, and that φ is injective.

To see that every λ in LinExt $(P_{\overline{a}})$ has $\varphi(\lambda) = \tau$ lying in $\mathfrak{S}^{\pitchfork}(P_{\overline{a}})$, we will induct on the number ℓ of cycles in $\tau = \tau^{(1)} \cdots \tau^{(\ell)}$, which is also the number of prime cycles in the decomposition $\sigma = \sigma^{(1)} \mathsf{T} \cdots \mathsf{T} \sigma^{(\ell)}$. By definition of $\mathfrak{S}^{\pitchfork}(P_{\overline{a}})$, we must check that the cycle partition $\pi = \{B_1, \ldots, B_\ell\}$ of τ lies in $\Pi^{\pitchfork}(P_{\overline{a}})$, where B_i is the set underlying the cycle $\tau^{(i)}$. To this end, assume that we produce the factorization $\sigma = \sigma^{(1)} \mathsf{T} \cdots \mathsf{T} \sigma^{(\ell)}$ according to the algorithm presented in Subsection 4.1, and let us check that the block $B = B_1$ underlying $\tau^{(1)}$ satisfies the two properties (a), (b) in the recursive characterization of $\Pi^{\pitchfork}(P_{\overline{a}})$ from Proposition 2.22:

- For (a), the elements of *B* are all minimal in $P_{\overline{a}}$ because, in the initial factorization step $\sigma = \sigma^{(1)} \mathsf{T} \hat{\sigma}$, each column $\binom{i}{j}$ in the two-line notation of $\sigma^{(1)}$ is the *leftmost* column of σ having *i* as its top element, so it corresponds to the *bottom* element in the *i*th chain a_i of $P_{\overline{a}}$.
- For (b), note that after that initial factorization step, the poset $\hat{P}_{\overline{a}}$ and partition $\hat{\pi} = \{B_2, \ldots, B_\ell\}$ will correspond to $\hat{\sigma}$ in the above factorization, coming from a $\hat{\lambda}$ in $\text{LinExt}(\hat{P}_{\overline{a}})$ with $\varphi(\hat{\lambda}) = \hat{\sigma}$ for which the result holds by induction on ℓ .

To show that φ is injective, we must give an algorithm to recover λ from $\tau = \varphi(\lambda)$. It would be equivalent to recover τ 's multiset labeled image σ in $\mathfrak{S}_{M(\overline{\alpha})}$ from the bijection in Proposition 4.14. Factoring τ lets us recover its *unordered* set of cycles $\{\tau^{(1)}, \ldots, \tau^{(\ell)}\}$, and hence also the *unordered* set of prime cycles $\{\sigma^{(1)}, \ldots, \sigma^{(\ell)}\}$ that will appear in an intercalation factorization of σ . We would like to know how to properly index $\{\sigma^{(i)}\}_{i=1,\ldots,\ell}$, up to interchanging commuting elements, so that we could recover σ as their intercalation product $\sigma = \sigma^{(1)} \mathsf{T} \cdots \mathsf{T} \sigma^{(\ell)}$. We claim that this (partial) ordering is already contained in the information of the unordered set of cycles $\{\tau^{(i)}\}_{i=1,\ldots,\ell}$ as follows. When two prime cycles $\sigma^{(r)}, \sigma^{(s)}$ do not commute, it is because they share a common element *i*, so there must exist two elements x, y in [*n*] that come from the i^{th} chain a_i in the standardized labeling of $P_{\overline{a}}$, with $x \in \tau^{(r)}, y \in \tau^{(s)}$. If $x <_{\mathbb{Z}} y$, then $\sigma^{(r)}$ must occur to the left of $\sigma^{(s)}$ in the intercalation product.

4.4. **Proof of Theorem 1.4.** Our goal here is to find a generating function compiling the Poincaré polynomials $Poin(P_{\overline{a}}, t)$ for all compositions \overline{a} of length ℓ . This uses more of Foata's theory for the intercalation monoid Int_{ℓ} , similar to his deduction of MacMahon's *Master Theorem*.

Since each multiset permutation σ has only finitely many intercalation factorizations $\sigma = \rho \tau \tau$, one can define a convolution algebra on the set of functions $\phi : \text{Int}_{\ell} \to \mathbb{Z}$ with pointwise addition:

$$(\phi_1 * \phi_2)(\sigma) := \sum_{\rho \intercal \tau = \sigma} \phi_1(\rho) \cdot \phi_2(\tau).$$

Let $\zeta : \operatorname{Int}_{\ell} \to \mathbb{Z}$ denote the *zeta function* defined by $\zeta(\sigma) = 1$ for all σ in $\operatorname{Int}_{\ell}$. The zeta function has a unique convolutional inverse μ , called the *Möbius function*. Foata proved that the Möbius function can be expressed by the following explicit formula

$$\mu(\sigma) = \begin{cases} (-1)^{\mathsf{pcyc}(\sigma)} & \text{if } \sigma \text{ is simple,} \\ 0 & \text{else,} \end{cases}$$

where $\sigma \in \operatorname{Int}_{\ell}$ is simple if all the letters of σ are distinct, that is, $\operatorname{supp}(\sigma)$ is a set, not a multiset. This may be formulated as an identity in a completion $\mathbb{Z}[[\operatorname{Int}_{\ell}]] := \{\sum_{\sigma \in \operatorname{Int}_{\ell}} z_{\sigma} \sigma : z_{\sigma} \in \mathbb{Z}\}$ of the monoid algebra $\mathbb{Z}[\operatorname{Int}_{\ell}]$, allowing infinite \mathbb{Z} -linear combinations of elements of $\operatorname{Int}_{\ell}$ (see [10, Théorème 2.4]):

(16)
$$1 = \left(\sum_{\sigma \in \operatorname{Int}_{\ell}} \sigma\right) \left(\sum_{\sigma \in \operatorname{Int}_{\ell}} \mu(\sigma) \sigma\right) = \left(\sum_{\sigma \in \operatorname{Int}_{\ell}} \sigma\right) \left(\sum_{\text{simple } \sigma \in \operatorname{Int}_{\ell}} (-1)^{\operatorname{\mathsf{pcyc}}(\sigma)} \sigma\right).$$

Now introduce an $\ell \times \ell$ matrix $B := (b_{ij})_{i,j=1,2,...,\ell}$ of indeterminates, and let $\mathbb{Z}[[b_{ij},t]]$ be the (usual, commutative) power series ring in $\{b_{ij}\}_{i,j=1}^{\ell}$ along with one further indeterminate t. One can then define a ring homomorphism

$$\begin{aligned} \mathbb{Z}[[\operatorname{Int}_{\ell}]] & \xrightarrow{u_t} & \mathbb{Z}[[b_{ij}, t]] \\ \sigma & \longmapsto & t^{\mathsf{pcyc}(\sigma)} \cdot \underline{b}_{\sigma} \end{aligned}$$

where if $\sigma = \begin{pmatrix} i_1 & i_2 & \cdots & i_n \\ \sigma_1 & \sigma_2 & \cdots & \sigma_n \end{pmatrix}$ then $\underline{b}_{\sigma} := \prod_{k=1}^n b_{i_k \sigma_k}$.

Applying the homomorphism u_t to both sides of (16) gives a t-version of MacMahon's Master Theorem.

Theorem 4.18. In $\mathbb{Z}[[b_{ij}, t]]$ one has the identity

$$\sum_{\sigma \in \operatorname{Int}_{\ell}} t^{\mathsf{pcyc}(\sigma)} \underline{b}_{\sigma} = \left(\sum_{simple \ \sigma \in \operatorname{Int}_{\ell}} (-t)^{\mathsf{pcyc}(\sigma)} \underline{b}_{\sigma} \right)^{-1} = \left(\sum_{H \subseteq [\ell]} \sum_{\sigma \in \mathfrak{S}_{H}} (-t)^{\mathsf{pcyc}(\sigma)} \underline{b}_{\sigma} \right)^{-1}$$

Remark 4.19. Setting t = 1 in Theorem 4.18 gives an identity in $\mathbb{Z}[[b_{ij}]]$:

(17)
$$\sum_{\sigma \in \text{Int}_{\ell}} \underline{b}_{\sigma} = \left(\sum_{H \subseteq [\ell]} \sum_{\sigma \in \mathfrak{S}_{H}} (-1)^{\mathsf{pcyc}(\sigma)} \underline{b}_{\sigma} \right)^{-1}$$

which is equivalent to an identity in Foata's proof of the (commutative) MacMahon Master Theorem, as we recall here. Introduce two sets of ℓ variables $\mathbf{x} = (x_1, \ldots, x_\ell), \mathbf{y} = (y_1, \ldots, y_\ell)$ related by the matrix B of indeterminates as follows: $\mathbf{y} = B\mathbf{x}$, that is, $y_i = \sum_j b_{ij} x_j$. Then MacMahon's Master Theorem is this identity in $\mathbb{Z}[[b_{ij}]]$:

(18)
$$\sum_{\overline{a} \in \{0,1,2,\dots\}^{\ell}} \left(\text{coefficient of } \mathbf{x}^{\overline{a}} \text{ in } \mathbf{y}^{\overline{a}} \right) = \det(I_{\ell} - B)^{-1},$$

where $\mathbf{x}^{\overline{a}} := x_1^{a_1} \cdots x_{\ell}^{a_{\ell}}$. It is not hard to check that the left sides and right sides of (18) and (17) are the same: the left side of (17) needs to be grouped according to the multiplicity vector \overline{a} giving the support $\mathsf{supp}(\sigma)$, and the right side must be reinterpreted in terms of the permutation expansion of a determinant.

Remark 4.20. Theorem 4.18 is similar in spirit to Garoufalidis-Lê-Zeilberger's quantum MacMahon Master Theorem [11, Theorem 1] (see also Konvalinka-Pak [16, Theorem 1.2]). Their quantum version inserts a $(-q)^{-inv(\sigma)}$ in order to produce a q-determinant, but $inv(\sigma) \neq pcyc(\sigma)$.

We now specialize $b_{ij} = x_j$ in Theorem 4.18 to deduce Theorem 1.4, whose statement we recall here.

Theorem 1.4. For $\ell = 1, 2, ...,$ one has

70

$$\sum_{\substack{\in\{1,2,\ldots\}^{\ell}}} \operatorname{Poin}(P_{\overline{a}},t) \cdot \mathbf{x}^{\overline{a}} = \frac{1}{1 - \sum_{j=1}^{\ell} e_j(\mathbf{x}) \cdot (t-1)(2t-1) \cdots ((j-1)t-1)}$$

where $e_j(\mathbf{x}) := \sum_{1 \le i_1 < \dots < i_j \le \ell} x_{i_1} \cdots x_{i_j}$ is the jth elementary symmetric function.

Proof. Setting $b_{ij} = x_j$ in Theorem 4.18 gives

(19)
$$\sum_{\sigma \in \operatorname{Int}_{\ell}} t^{\mathsf{pcyc}(\sigma)} \prod_{k} x_{\sigma_{k}} = \left(\sum_{H \subseteq [\ell]} \sum_{\sigma \in \mathfrak{S}_{H}} (-t)^{\mathsf{pcyc}(\sigma)} \prod_{k \in H} x_{k} \right)^{-1}$$

Let us manipulate both sides of equation (19). On the left side, grouping terms according to $supp(\sigma)$ gives

$$\sum_{\{0,1,2,\ldots\}^{\ell}} \mathbf{x}^{\overline{a}} \sum_{\sigma \in \mathfrak{S}_{M(\overline{a})}} t^{\mathsf{pcyc}(\sigma)}$$

On the right side of (19), any subset $H \subseteq [\ell]$ of cardinality $j \ge 1$ satisfies

 \overline{a}

$$\sum_{\sigma \in \mathfrak{S}_H} (-t)^{\mathsf{pcyc}(\sigma)} = \sum_{\sigma \in \mathfrak{S}_j} (-t)^{\mathsf{cyc}(\sigma)} = (-t)(1-t)(2-t)\cdots(j-1-t)$$

by (8). Therefore grouping according to j = #H, and noting $\sum_{\substack{H \subseteq [\ell]: \\ \#H=j}} \prod_{k \in H} x_k = e_j(\mathbf{x})$ lets one rewrite the sum inside the parentheses on the right side of (19) as this:

$$1 + \sum_{j=1}^{\ell} (-t)(1-t)(2-t) \cdots (j-1-t) \cdot e_j(\mathbf{x}).$$

So far this gives

$$\sum_{\overline{a}\in\{0,1,2,\dots\}^{\ell}} \mathbf{x}^{\overline{a}} \sum_{\sigma\in\mathfrak{S}_{M(\overline{a})}} t^{\mathsf{pcyc}(\sigma)} = \left(1 + \sum_{j=1}^{\ell} (-t)(1-t)(2-t)\cdots(j-1-t)\cdot e_{j}(\mathbf{x})\right)^{-1}.$$

Now perform two more substitutions: first replace t by t^{-1} , giving this

$$\sum_{\overline{a} \in \{0,1,2,\ldots\}^{\ell}} \mathbf{x}^{\overline{a}} \sum_{\sigma \in \mathfrak{S}_{M(\overline{a})}} t^{-\mathsf{pcyc}(\sigma)} = \left(1 + \sum_{j=1}^{\ell} (-t^{-1})(1-t^{-1})(2-t^{-1})\cdots(j-1-t^{-1}) \cdot e_j(\mathbf{x}) \right)^{-1},$$

and then replace x_i by tx_i for $i = 1, 2, ..., \ell$, so that $\mathbf{x}^{\overline{a}} \mapsto t^{|\overline{a}|} \mathbf{x}^{\overline{a}}$ and $e_j(\mathbf{x}) \mapsto t^j e_j(\mathbf{x})$, giving this

$$\sum_{\overline{a}\in\{0,1,2,\ldots\}^{\ell}} \mathbf{x}^{\overline{a}} \sum_{\sigma\in\mathfrak{S}_{M(\overline{a})}} t^{|\overline{a}|-\mathsf{pcyc}(\sigma)} = \left(1 - \sum_{j=1}^{\ell} (t-1)(2t-1)\cdots((j-1)t-1)\cdot e_j(\mathbf{x})\right)^{-1}$$

Comparison of the left side with Theorem 1.3 shows that this last equation is Theorem 1.4.

Remark 4.21. We justify here the claim from the Introduction that Theorem 1.4 generalizes the formula (8):

$$\mathsf{Poin}(\mathsf{Antichain}_{\ell}, t) = 1(1+t)(1+2t)\cdots(1+(\ell-1)t).$$

Since $\operatorname{Antichain}_{\ell} = P_{\overline{a}}$ where $\overline{a} = (1, 1, \ldots, 1)$, we seek to explain why the coefficient of $x_1 \ldots x_{\ell}$ in the power series on the right side in Theorem 1.4 should be $1(1+t)(1+2t)\cdots(1+(\ell-1)t)$. Introducing the abbreviation $\langle t \rangle_j := (t-1)(2t-1)\cdots((j-1)t-1)$, the right side in Theorem 1.4 can be rewritten and expanded as

(20)
$$\frac{1}{1 - \sum_{j=1}^{\ell} e_j(\mathbf{x}) \langle t \rangle_j} = \sum_{n \ge 0} \left(\sum_{j=1}^{\ell} e_j(\mathbf{x}) \langle t \rangle_j \right)^n$$

If we let A_{ℓ} denote the coefficient of $x_1 \cdots x_{\ell}$ in this series, then it suffices to explain why

$$A_{\ell+1} = (1+\ell t) \cdot A_{\ell}$$

For this coefficient extraction, it is safe to replace each $e_j(\mathbf{x}) = e_j(x_1, x_2, \dots, x_\ell)$ in (20) with an infinite variable version $e_j(\mathbf{x}) = e_j(x_1, x_2, \dots)$. Extracting the coefficient of $x_1 \cdots x_\ell$ on the right side in (20) shows

$$A_{\ell} = \sum_{\substack{\text{ordered set partitions}\\ \pi = (B_1, \dots, B_n) \text{ of } [\ell]}} w(\pi), \qquad \text{where } w(\pi) := \prod_{B_i \in \pi} \langle t \rangle_{|B_i|}.$$

To explain (21), note that each ordered set partition $\hat{\pi}$ of $[\ell + 1]$ can be obtained from a unique ordered set partition $\pi = (B_1, \ldots, B_n)$ of $[\ell]$ as follows: either $\hat{\pi}$ has added $\ell + 1$ into one of the preexisting blocks B_i of π , or $\hat{\pi}$ has a singleton block $\{\ell + 1\}$, inserted into one of the n + 1 locations in the sequence (B_1, \ldots, B_n) . Thus having fixed an ordered set partition π of $[\ell]$, the sum of $w(\hat{\pi})$ over $\hat{\pi}$ which correspond to π is this sum:

$$w(\pi) \cdot (|B_1|t-1) + \dots + w(\pi) \cdot (|B_n|t-1) + \underbrace{w(\pi) + w(\pi) + \dots + w(\pi)}_{n+1 \text{ times}}$$
$$= w(\pi) \left(\sum_{i=1}^n (|B_i|t-1) + n + 1 \right) = w(\pi) \left(\ell t - n + n + 1\right) = w(\pi) \cdot (1 + \ell t).$$

Summing this over all possible π gives (21).

5. Posets of width two and proof of Theorem 1.5

The width of a poset P is the maximum size of an antichain in P. A famous result of Dilworth from 1950 (see [28, Ch 3, Exer 77(d)]), asserts that the width d of P is the minimum number of chains required in a chain decomposition $P = P_1 \cup P_2 \cup \cdots \cup P_d$, that is, where each P_i is a totally ordered subset $P_i \subseteq P$.

Consequently, a poset P of width two can be decomposed into two chains $P = P_1 \cup P_2$, possibly with some order relations between elements of P_1 and P_2 .

Recall that in the Introduction we defined a descent-like statistic on $\sigma = [\sigma_1, \ldots, \sigma_n]$ in LinExt(P), as the cardinality $\operatorname{des}_{P_1, P_2}(\sigma) := \#\operatorname{Des}_{P_1, P_2}(\sigma)$ of this set,

(22)
$$\mathsf{Des}_{P_1,P_2}(\sigma) := \{ i \in [n-1] : \sigma_i \in P_2, \ \sigma_{i+1} \in P_1, \text{ with } \sigma_i, \sigma_{i+1} \text{ incomparable in } P \},\$$

in order to state the following result.

Theorem 1.5. For a width two poset decomposed into two chains as $P = P_1 \cup P_2$, one has

$$\mathsf{Poin}(P,t) = \sum_{\sigma \in \mathrm{LinExt}(P)} t^{\mathsf{des}_{P_1,P_2}(\sigma)}$$

To prove this, we start with the following observation.

Corollary 5.1. For posets P of width two, one has

$$\mathsf{Poin}(P,t) = \sum_{\pi \in \Pi^{\pitchfork}(P)} t^{\mathsf{pairs}(\pi)}$$

where $pairs(\pi)$ is the number of two-element blocks B_i in π . In particular, setting t = 1,

$$\#\operatorname{LinExt}(P) = \#\Pi^{\oplus}(P)(=\#\mathfrak{S}^{\oplus}(P))$$

Proof. Antichains in P have at most two elements, so Proposition 2.16(iii) implies that P-transverse permutations have only 1-cycles and 2-cycles. But then this implies that the map $\mathfrak{S}^{\pitchfork}(P) \to \Pi^{\pitchfork}(P)$ sending a P-transverse permutation τ to the set partition π given by its cycles is a bijection, with $\mathsf{pairs}(\pi) = n - \mathsf{cyc}(\tau)$. The result then follows from Theorem 1.1.

Example 5.2. Let $P = a \sqcup b$ be a poset which is a disjoint union of two chains a, b having a, b elements respectively. One can check that a *P*-transverse partition having $pairs(\pi) = k$ is completely determined by the choice of a k element subset $x_1 <_P \cdots <_P x_k$ from a and a k element subset $y_1 <_P \cdots <_P y_k$ from b to constitute the two-element blocks, as follows: $\{x_1, y_1\}, \ldots, \{x_k, y_k\}$. This implies

$$\mathsf{Poin}(\mathsf{a} \sqcup \mathsf{b}, t) = \sum_{k=0}^{\min(a,b)} \binom{a}{k} \binom{b}{k} t^k.$$

This is consistent with $\# \operatorname{LinExt}(\mathsf{a} \sqcup \mathsf{b}) = \binom{a+b}{a}$, since setting t = 1 in the equation above gives

$$\binom{a+b}{a} = \sum_{k=0}^{\min(a,b)} \binom{a}{k} \binom{b}{k}$$

which is an instance of the Chu-Vandermonde summation.

In light of Corollary 5.1, to prove Theorem 1.5, one would need a bijection from LinExt(P) to $\Pi^{\uparrow}(P)$ (or $\mathfrak{S}^{\uparrow}(P)$) that sends the statistic $\mathsf{des}_{(P_1,P_2)}(-)$ to the number of pairs or number of 2-cycles. Unfortunately, there does not seem to be a consistent labeling of a width two poset $P = P_1 \cup P_2$ to make the bijection Ψ from Section 3 play this role. Nevertheless, having fixed the chain decomposition⁶ $P = P_1 \sqcup P_2$, we provide in the proof below such a bijection $\Omega : \text{LinExt}(P) \to \Pi^{\uparrow}(P)$.

Proof of Theorem 1.5. We describe Ω and Ω^{-1} recursively, via induction on n := #P. There are two cases, based on whether P has one or two minimal elements.

Case 1. There is a unique minimum element $p_0 \in P$.

In this case, given $\sigma = [\sigma_1, \ldots, \sigma_n]$ in LinExt(P), we must have $\sigma_1 = p_0$, so that $\{p_0\}$ should be a singleton block of $\pi = \Omega(\sigma)$, and one produces the remaining blocks of π by applying Ω recursively to $[\sigma_2, \ldots, \sigma_n]$. This is depicted schematically here:



For the inverse map Ω^{-1} , given a *P*-transverse partition π , since the blocks of π are antichains in *P*, the unique minimum element p_0 of *P* must lie in a singleton block $\{p_0\}$ in π . So make $\Omega^{-1}(\pi) = \sigma$ have $\sigma_1 = p_0$, and construct $[\sigma_2, \ldots, \sigma_n]$ by applying Ω^{-1} recursively to the $(P - \{p_0\})$ -transverse partition obtained from π by removing the block $\{p_0\}$.

Case 2. There are two minimal elements of P.

Label these two minimal elements p_1, p_2 of P so that $p_i \in P_i$ for i = 1, 2. Note that this implies that every $\sigma = [\sigma_1, \sigma_2, \ldots, \sigma_n]$ in LinExt(P) has either $\sigma_1 = p_1$ or $\sigma_1 = p_2$. Note also that any P-transverse partition π only has blocks of cardinality 1 or 2, which yields two subcases for defining Ω and Ω^{-1} :

- The Subcase 2a for
 - defining Ω occurs when $\sigma_1 = p_1$,
 - defining Ω^{-1} occurs when $\{p_1\}$ appears as a singleton block within π .
- The Subcase 2b for
 - defining Ω occurs when $\sigma_1 = p_2$,

⁶So we assume here that $P_1 \cap P_2 = \emptyset$, but there may be order relations between elements of P_1 and P_2 .

- defining Ω^{-1} occurs when p_1 appears in a two-element block within π .

Subcase 2a.

When defining Ω , if $\sigma_1 = p_1$, then make $\{p_1\}$ a singleton block of $\pi = \Omega(\sigma)$, and produce the remaining blocks of π by applying Ω recursively to $[\sigma_2, \ldots, \sigma_n]$.



To define Ω^{-1} , if $\{p_1\}$ is a singleton block of π , make $\Omega^{-1}(\pi) = \sigma$ have $\sigma_1 = p_1$, and construct $[\sigma_2, \ldots, \sigma_n]$ by applying Ω^{-1} recursively to the $(P - \{p_1\})$ -transverse partition obtained from π by removing the block $\{p_1\}$.

Subcase 2b.

When defining Ω , if $\sigma_1 = p_2$, then p_1 appears elsewhere in σ , say $p_1 = \sigma_{i+1}$ where $i \geq 1$. Because σ lies in LinExt(P) and $\sigma_{i+1} = p_1$ is the minimum element of P_1 , this forces $\sigma_1, \sigma_2, \ldots, \sigma_i$ to all be elements of P_2 . In this case, add to π the singleton blocks $\{\sigma_1\}, \{\sigma_2\}, \ldots, \{\sigma_{i-1}\}$ along with the two-element block $\{\sigma_i, \sigma_{i+1}\} = \{\sigma_i, p_1\}$, and compute the rest of $\Omega(\sigma) = \pi$ recursively by replacing (P, σ) with $(P - \{\sigma_1, \sigma_2, \ldots, \sigma_{i+1}\}, [\sigma_{i+2}, \sigma_{i+3}, \ldots, \sigma_n])$. Here is the schematic picture:



When defining $\Omega^{-1}(\pi)$, if p_1 appears in some two-element block of π , then it appears in some block $\{p_1, p'_2\}$ for some p'_2 in P_2 . We claim that π being *P*-transverse then forces any elements $p <_P p'_2$ in P_2 to lie in singleton blocks $\{p\}$ of π . To see this claim, assume not, so that some such p lies in a two-element block of π , necessarily of the form $\{p'_1, p\}$ for some p'_1 in P_1 with $p_1 <_P p'_1$. This leads to a contradiction of Proposition 2.16(ii), since (p'_1, p_1) would then be a relation in $\overline{P \cup \pi}$ via this transitive chain of relations: $p'_1 \equiv_{\pi} p <_P p'_2 \equiv_{\pi} p_1$.

In this subcase, list the totally ordered (and possibly empty) collection of all elements p in P_2 with $p <_P p'_2$ at the beginning of σ as $\sigma_1, \sigma_2, \ldots, \sigma_{i-1}$, followed by $\sigma_i \sigma_{i+1} = p'_2 p_1$. Then compute the rest of $\Omega^{-1}(\pi) = \sigma$ recursively, by applying Ω^{-1} to the $(P - \{\sigma_1, \sigma_2, \ldots, \sigma_{i+1}\})$ -transverse partition obtained from π by removing the singleton blocks $\{\sigma_1\}, \{\sigma_2\}, \ldots, \{\sigma_{i-1}\}$ and the two-element block $\{\sigma_i, \sigma_{i+1}\} = \{p'_2, p_1\}$.

It is not hard to check that the two maps Ω, Ω^{-1} defined recursively in this way are actually mutually inverse bijections. By construction, Ω has the property that the two-element blocks of $\pi = \Omega(\sigma)$ are exactly those containing *P*-incomparable pairs $\{\sigma_i, \sigma_{i+1}\}$ for which $\sigma_i \in P_1$ and $\sigma_{i+1} \in P_2$, as claimed. \Box

Recall from the Introduction that the number of usual descents of a permuation σ is defined as $des(\sigma) = #Des(\sigma)$ where

$$\mathsf{Des}(\sigma) := \{ i \in [n-1] \mid \sigma_i > \sigma_{i+1} \}.$$

This was used to define the *P*-Eulerian polynomial in equation (9) as $\sum_{\sigma \in \text{LinExt}(P)} t^{\text{des}(\sigma)}$, assuming that *P* is naturally labeled, that is, LinExt(P) contains the identity permutation $\sigma = [1, 2, ..., n]$.

Corollary 5.3. When P is a width two poset having a chain decomposition $P_1 \cup P_2$ with P_1 an order ideal of P, then the Poincaré polynomial for P coincides with the P-Eulerian polynomial:

$$\mathsf{Poin}(P,t) = \sum_{\sigma \in \mathrm{LinExt}(P)} t^{\mathsf{des}(\sigma)}$$

Proof. Let $\#P_i = n_i$ for i = 1, 2, so that $n = \#P = n_1 + n_2$. One can then choose a natural labeling for P by [n] such that the elements of the order ideal P_1 are labeled by the initial segment $[n_1] = \{1, 2, \ldots, n_1\}$, and P_2 is labeled by $\{n_1+1, n_1+2, \ldots, n\}$. We claim that with this natural labeling, one has $\mathsf{Des}_{(P_1, P_2)}(w) = \mathsf{Des}(w)$. This is because the labeling renders one of the conditions in the definition (22) of $\mathsf{Des}_{(P_1, P_2)}(\sigma)$ superfluous: assuming that $\sigma_i \in P_2$ and $\sigma_{i+1} \in P_1$, then σ_i, σ_{i+1} must already be incomparable in P, because otherwise $\sigma_i <_P \sigma_{i+1}$ (since σ was a linear extension of P) and then P_1 being an order ideal would force $\sigma_i \in P_1$, a contradiction. Now since P_1, P_2 are totally ordered in P, and σ lies in $\mathsf{LinExt}(P)$, one has $\sigma_i \in P_2$ and σ_{i+1} in P_1 if and only if $\sigma_i >_{\mathbb{Z}} \sigma_{i+1}$. That is $\mathsf{Des}_{(P_1, P_2)}(w) = \mathsf{Des}(w)$.

Example 5.4. An interesting family of posets to which Corollary 5.3 applies are the posets $P(\lambda/\mu)$ associated with two-row skew Ferrers diagrams λ/μ . A *Ferrers diagram* associated to a partition (of a number) $\lambda = (\lambda_1, \ldots, \lambda_\ell)$ has λ_i square cells drawn left-justified in row *i*. A *skew Ferrers diagram* λ/μ for two partitions λ, μ having $\lambda_i \geq \mu_i$ is the diagram for λ with the cells occupied by the diagram for μ removed. There is a poset structure $P(\lambda/\mu)$ on the cells of λ/μ in which a cell (i, j) in row *i* and column *j* has $(i, j) \leq_{P(\lambda/\mu)} (i', j')$ if both $i \leq i'$ and $j \leq j'$.

When λ/μ has only two parts, we will call it a *two-row skew Ferrers diagram*. Three examples of such λ/μ and their associated $P(\lambda/\mu)$ are shown below.



The decomposition $P(\lambda/\mu) = P_1 \cup P_2$ where P_i correspond to the cells in row *i* of λ/μ shows that $P(\lambda/\mu)$ has width two, and furthermore P_1 forms an order ideal. Therefore Corollary 5.3 implies that for any two-row skew Ferrers diagram λ/μ one has

(23)
$$\mathsf{Poin}(P(\lambda/\mu), t) = \sum_{\sigma \in \mathrm{LinExt}(P(\lambda/\mu))} t^{\mathsf{des}(\sigma)}.$$

On the other hand, there is a well-known bijection between linear extensions σ of $P(\lambda/\mu)$ and the standard Young tableaux Q of shape λ/μ , which are (bijective) labelings of the cells of the diagram by [n] where $n = \sum_i \lambda_i - \sum_i \mu_i$, with the numbers increasing left-to-right in rows and top-to-bottom in columns; see [26, §7.10]. There is also a notion of descent set Des(Q) for such tableaux, having $i \in \text{Des}(Q)$ whenever i + 1 labels a cell in a lower row of Q than i. However, in general when σ corresponds to Q, one does not have $\text{des}(\sigma) = \text{des}(Q)$, so that $\text{Poin}(P(\lambda/\mu), t)$ differs from the generating function $\sum_{Q} t^{\text{des}(Q)}$ of standard tableaux Q shape λ/μ by des(Q). For example, there are two standard tableaux of shape $\lambda/\mu = (2,1)/(0,0)$

$$Q_1 = \boxed{\begin{array}{c}1 & 2\\3\end{array}} \qquad Q_2 = \boxed{\begin{array}{c}1 & 3\\2\end{array}}$$

both having $des(Q_i) = 1$, however $Poin(P(\lambda/\mu), t) = 1 + t$. In two special cases, however, they (essentially) coincide.

When P_{λ/μ} = a ⊔ b is a disjoint union of two chains, as in Example 5.2, one can check that, if one (naturally) labels a ⊔ b so that the elements of the order ideal b are labeled 1, 2, ..., b while a is labeled b + 1, b + 2, ..., b + a, then one does have des(σ) = des(Q), and hence

$$\sum_{Q} t^{\mathsf{des}(Q)} = \mathsf{Poin}(P_{\mathsf{a}\sqcup\mathsf{b}}, t) = \sum_{k} \binom{a}{k} \binom{b}{k} t^{k}.$$

- When λ/μ is a $2 \times n$ rectangle, so that $P_{\lambda/\mu} = 2 \times n$ is a Cartesian product poset, then σ in LinExt(P) and standard Young tableaux Q of shape $2 \times n$ can both be identified with *Dyck paths of semilength* n, that is, lattice paths from (0,0) to (2n,0) in \mathbb{Z}^2 taking steps northeast or southeast and staying weakly above the x-axis. One can check that
 - $\mathsf{Des}(\sigma)$ corresponds to valleys (i.e. southwest steps followed by a northeast step), while
 - Des(Q) correspond to *peaks* (i.e. northeast steps followed by a southwest step).

In general, such a Dyck path has one more peak than valley [26, Exercises 6.19(i, ww, aaa)]. Hence one has

$$\mathsf{Poin}(2 \times \mathsf{n}, t) = \frac{1}{t} \sum_{Q} t^{\mathsf{des}(Q)} = \sum_{k=0}^{n-1} \frac{1}{n} \binom{n}{k} \binom{n}{k+1} t^{k},$$

which is the generating function for the Narayana numbers $N(n,k) := \frac{1}{n} \binom{n}{k-1} \binom{n}{k}$ (see [7, p.2] and [26, Exer. 6.36(a)]). Upon setting t = 1, the Naryana numbers sum to the Catalan number

$$\#\operatorname{LinExt}(2\times \mathsf{n}) = \frac{1}{n+1} \binom{2n}{n}.$$

Note that for any (non-skew) partition λ , the celebrated hook-length formula of Frame, Robinson and Thrall [26, Corollary 7.21.6] gives a simple product formula for $\# \operatorname{LinExt}(P(\lambda)) = [\operatorname{Poin}(P(\lambda), t)]_{t=1}$.

Open Problem 5.5. Combinatorially interpret $Poin(P(\lambda), t)$ for other partitions λ , and in particular, for $m \times n$ rectangular partitions, where $P(\lambda) = m \times n$ is a Cartesian product of chains.

Below we give $\mathsf{Poin}(3 \times \mathsf{n}, t)$ for $2 \le n \le 8$.

n	Poin(3 imes n, t)
2	$1 + 3t + t^2$
3	$1 + 9t + 19t^2 + 11t^3 + 2t^4$
4	$1 + 18t + 92t^2 + 174t^3 + 133t^4 + 40t^5 + 4t^6$
5	$1 + 30t + 280t^2 + 1091t^3 + 1987t^4 + 1746t^5 + 731t^6 + 132t^7 + 8t^8$
6	$1 + 45t + 665t^2 + 4383t^3 + 14603t^4 + 25957t^5 + 25064t^6 + 12965t^7 + 3413t^8 + 404t^9$
7	$1 + 63t + 1351t^2 + 13475t^3 + 71305t^4 + 213539t^5 + 373651t^6 + 385578t^7 + 232310t^8$
	$+79023t^9 + 14174t^{10} + 1168t^{11} + 32t^{12}$
8	$1 + 84t + 2464t^2 + 34608t^3 + 266470t^4 + 1206826t^5 + 3343958t^6 + 5782699t^7$
	$+ 6275503t^8 + 4240489t^9 + 1743730t^{10} + 417622t^{11} + 53884t^{12} + 3232t^{13} + 64t^{14} + 54t^{14} + $

Remark 5.6. Since equation (8) shows that the Poincaré polynomial Poin(P, t) for the antichain poset $P = P_{(1^n)} = Antichain_n$ has only real roots, one might wonder whether this holds for some more general class of posets. It does not hold for all posets, and not even for all disjoint unions of chains $P_{\overline{a}}$, since

$$\mathsf{Poin}(P_{(2,2,2)},t) = 1 + 12t + 43t^2 + 30t^3 + 4t^4$$

has a pair of non-real complex roots. It can fail even for rectangular Ferrers posets, e.g., $\lambda = (3, 3, 3)$ has $Poin(P(\lambda), t) = Poin(3 \times 3, t) = 1 + 9t + 19t^2 + 11t^3 + 2t^4$ in the above table, with two non-real complex roots.

On the other hand, computations show that Poin(P, t) is real-rooted for all posets P of width two having at most 9 elements. This leads to the following question.

Question 5.7. Is Poin(P,t) real-rooted when the poset P has width two?

Acknowledgements

The authors gratefully acknowledge Dennis Stanton for conversations about MacMahon's Master Theorem, as well as Anders Björner, Jesus DeLoera, Theo Douvropoulos, Michael Falk, Ira Gessel, Benjamin Steinberg, Volkmar Welker, Chi-Ho Yuen for enlightening discussions and references. They thank Philip Zhang for asking them questions about real-rootedness at the 2019 Mid-Atlantic Algebra, Geometry, and Combinatorics Workshop. Finally, the authors thank an anonymous referee for helpful comments.

References

- [1] Marcelo Aguiar and Swapneel Mahajan. Topics in Hyperplane Arrangements. American Mathematica Society, 2017.
- [2] Hans-Jürgen Bandelt, Victor Chepoi, and Kolja Knauer. COMs: complexes of oriented matroids. J. Combin. Theory Ser. A, 156:195–237, 2018.
- [3] Matthias Beck and Raman Sanyal. Combinatorial reciprocity theorems: An invitation to enumerative geometric combinatorics Graduate Studies in Mathematics 195, American Mathematical Society, 2018.
- [4] A. Björner and M.L. Wachs. Generalized quotients in Coxeter groups. Trans. Amer. Math. Soc. 308 (1988), 1–37.
- [5] A. Björner and M.L. Wachs. q-hook length formulas for forests. J. Combin. Theory Ser. A 52 (1989) 165–187.
- [6] A. Björner and M.L. Wachs. Permutation statistics and linear extensions of posets. J. Combin. Theory Ser. A 58 (1991), 85–114.
- [7] Petter Brändén. q-Narayana numbers and the flag h-vector of $J(2 \times n)$. Discrete Math., 281(1-3):67–81, 2004.
- [8] Kenneth Brown. Semigroups, Rings, and Markov Chains. Journal of Theoretical Probability, 13(3):342–351, 2000.
- [9] Dominique Foata. Etude algébrique de certains problèmes d'analyse combinatoire et du calcul des probabilités. PhD thesis, Publ. Inst. Statist. Univ. Paris, 1965.
- [10] Dominique Foata and Pierre Cartier. Problèmes combinatoires de commutation et réarrangements. Springer Verlag, Lecture Notes in Mathematics, 1969.
- [11] Stavros Garoufalidis, Thang T.Q. Lê, and Doron Zeilberger. The Quantum MacMahon Master Theorem. J. Algebra, March 2003.
- [12] I.M. Gessel, Multipartite P-partitions and inner products of skew Schur functions. Combinatorics and algebra (Boulder, Colo., 1983), 289–317, Contemp. Math. 34, Amer. Math. Soc., Providence, RI, 1984.
- [13] Regina Gente. The Varchenko Matrix for Cones. PhD thesis, Universität Marburg, 2013.
- [14] James Humphreys. Reflection groups and Coxeter groups. Cambridge Studies in Advanced Mathematics 29. Cambridge University Press, Cambridge, 1990.
- [15] Donald Ervin Knuth. The Art of Computer Programming. Addison-Wesley, 2015.
- [16] Matjaz Konvalinka and Igor Pak. Non-commutative Extensions of MacMahon's Master Theorem. Advances in Mathematics, 216(1), 29–61. 2007.
- [17] Gustav Lehrer and Louis Solomon, On the action of the symmetric group on the cohomology of the complement of its reflecting hyperplanes. J. Algebra 104(2):410–424, 1986).
- [18] Peter Orlik and Louis Solomon, Unitary reflection groups and cohomology. Invent. Math. 59(1):77-94. 1980.
- [19] Peter Orlik and Hiroaki Terao. Arrangements of hyperplanes, volume 300 of Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]. Springer-Verlag, Berlin, 1992.
- [20] Alexander Postnikov, Victor Reiner, and Lauren Williams. Faces of Generalized Permutohedra. Documenta Mathematica, 13:207–273, 2008.
- [21] Victor Reiner, Quotients of Coxeter complexes and P-partitions. Mem. Amer. Math. Soc. 95, no. 460, 1992.
- [22] Victor Reiner, Saliola, Franco, and Volkmar Welker, Spectra of symmetrized shuffling operators. (English summary) Mem. Amer. Math. Soc. 228, no. 1072, 2014.
- [23] Victor Reiner and Volkmar Welker. On the Charney-Davis and Neggers-Stanley conjectures. Journal of Combinatorial Theory, Series A, 109(2):247–280, February 2005.
- [24] A. Rényi. Théorie des éléments saillants d?une suite d?observations. Ann. Fac. Sci. Univ. Clermont-Ferrand 8 (1962), 7–13.
- [25] Kyoji Saito. Principal Γ-cone for a tree. Adv. Math., 212(2):645-668, 2007.
- [26] Richard Stanley. Enumerative Combinatorics, volume 2. Cambridge University Press, Cambridge, UK, 1999.
- [27] Richard Stanley. An Introduction to Hyperplane Arrangements. Geometric Combinatorics IAS/Park City Mathematics Series, pages 389–496, 2007.
- [28] Richard Stanley. Enumerative Combinatorics, volume 1. Cambridge University Press, New York, NY, USA, 2 edition, 2012.
- [29] Robert Steinberg, Differential equations invariant under finite reflection groups. Trans. Amer. Math. Soc. 112:392–400, 1964.
- [30] John Stembridge, Coxeter cones and their h-vectors. Adv. Math. 217(5): 1935–1961, 2008.
- [31] Thomas Zaslavsky, Facing up to arrangements: face-count formulas for partitions of space by hyperplanes. Mem. Amer. Math. Soc. 1(1), no. 154, 1975.

[32] Thomas Zaslavsky. A Combinatorial Analysis of Topological Dissections. Advances in Mathematics, 25(3):267–285, 1977.

School of Mathematics, University of Minnesota, Minneapolis, MN 55455, USA $E\text{-}mail\ address:\ dorpa003@umn.edu$

School of Mathematics, University of Minnesota, Minneapolis, MN 55455, USA $E\text{-}mail\ address:\ \texttt{reiner@math.um.edu}$