

Homotopy of the space of initial values satisfying the dominant energy condition strictly

Jonathan Glöckle

June 4, 2019

The dominant energy condition imposes a restriction on initial value pairs found on a spacelike hypersurface of a Lorentzian manifold. In this article, we study the space of initial values that satisfy this condition strictly. To this aim, we introduce a Lorentzian α -invariant for initial value pairs and compare it to its classical counterpart. Recent non-triviality results for the latter will then imply that this space has non-trivial homotopy groups.

1 Introduction

1.1 Dominant energy condition for initial values

According to general relativity, the universe can be modeled by a time-oriented Lorentzian manifold (N, \bar{g}) whose large-scale behavior is governed by the Einstein equation

$$T = \text{ric}^{\bar{g}} - \frac{1}{2} \text{scal}^{\bar{g}} \bar{g},$$

where T denotes the energy-momentum tensor. This does not only apply to the dynamics, the field equations also constraint the physical quantities experienced on a time-slice. More precisely, suppose that (N, \bar{g}) contains M as a spacelike hypersurface. On M , the induced Riemannian metric g and the second fundamental form K , defined with respect to the future-pointing unit normal e_0 , form a so-called initial value pair (g, K) . The Gauß-Codazzi equations imply that it is subject to the Einstein constraints (cf. [BI04])

$$\begin{aligned} 2\rho &= \text{scal}^g + (\text{tr } K)^2 - \|K\|^2 \\ j &= \text{div } K - \text{d tr } K, \end{aligned} \tag{1}$$

where energy density $\rho = T(e_0, e_0)$ and momentum density $j = T(e_0, -)|_{TM}$ are components of the energy-momentum tensor.

For physical reasons, the energy-momentum tensor is assumed to always satisfy the dominant energy condition, which implies that $\rho \geq \|j\|$. We will say that an initial value pair (g, K) satisfies the dominant energy condition if $\rho \geq \|j\|$, when ρ and j are defined by (1). This condition plays a vital role in the positive mass theorem [SY81; Wit81] stating that for an asymptotically Euclidean manifold (M, g) with K tending to zero at infinity, the ADM-mass is non-negative if (g, K) satisfies the dominant energy condition.

In this article, we consider the case that M is a compact spin manifold of dimension $n \geq 2$. Our aim is to study the space $\mathcal{I}^+(M)$ of initial value pairs (g, K) for which the dominant energy condition holds strictly, i.e. $\rho > \|j\|$. This is a subspace of the space $\mathcal{I}(M)$ of all initial value pairs, with C^∞ -topology. The reason for restricting to the strict version of the dominant energy condition is that it nicely connects to positive scalar curvature, which in turn is rather well-studied.

1.2 Connection to positive scalar curvature and main result

It is a simple observation that if $K \equiv 0$, then the strict dominant energy condition for (g, K) reduces to the condition that g has positive scalar curvature. However, whereas existence of positive scalar curvature metrics imposes a condition on the manifold, this is not true for the strict dominant energy condition. Taking any metric g , the pair $(g, \frac{1}{n}\tau g)$ satisfies the dominant energy condition strictly as long as the absolute value of the constant $\tau \in \mathbb{R}$ is large enough. Moreover, such a τ can be chosen in a way that it continuously depends on the metric g (in C^2 -topology). This allows to define a comparison map $\Phi: \Sigma \mathcal{R}^+(M) \simeq \mathcal{R}^+(M) \times [-1, 1] \cup \mathcal{R}(M) \times \{-1, 1\} \rightarrow \mathcal{I}^+(M)$ by $(g, t) \mapsto (g, \frac{1}{n}\tau(g)tg)$, where $\mathcal{R}(M)$ is the C^∞ -space of metrics and $\mathcal{R}^+(M)$ its subspace of positive scalar curvature metrics.

One of the main approaches to positive scalar curvature is by index theoretic methods. Assume that (M, g) is compact and spin. Then, there is a spinor bundle $\Sigma_{Cl}M$ with a right Cl_n -action, called Cl_n -linear spinor bundle of M . Its Dirac operator D commutes with the Cl_n -action and thus gives rise to a Cl_n -Fredholm operator, which has a KO -valued index called α -index $\alpha(M)$. The Schrödinger-Lichnerowicz formula

$$D^2 = \nabla^* \nabla + \frac{1}{4} \text{scal}$$

implies that it is invertible if g has positive scalar curvature and so its index vanishes. By homotopy invariance of the index, it is independent of g , and so the α -index provides an obstruction to existence of positive scalar curvature metrics on M if it is non-zero for some spin structure on M .

In the case when there is a positive scalar curvature metric on M , this invariant can be refined to a secondary invariant called index difference or α -invariant that allows to detect non-trivial homotopy groups in the space of positive scalar curvature metrics. As before, the Cl_n -linear Dirac operator defines a map assigning to each metric a Cl_n -Fredholm operator, which is invertible if the metric is of positive scalar curvature. Then applying the KO -valued index, we obtain the map

$$\alpha: \pi_k(\mathcal{R}^+(M)) \cong \pi_{k+1}(\mathcal{R}(M), \mathcal{R}^+(M)) \rightarrow KO^{-n}(D^{k+1}, S^k) \cong KO^{-n-k-1}(\{*\}).$$

A similar invariant exists in the case of initial values. For this, the Cl_n -linear spinor bundle has to be replaced by the $Cl_{n,1}$ -linear hypersurface spinor bundle $\overline{\Sigma}_{Cl}M$. To define it, we embed M as spacelike hypersurface into a time-oriented Lorentzian manifold (N, \overline{g}) such that the pair (g, K) arises as induced metric and second fundamental form. Then $\overline{\Sigma}_{Cl}M$ is the restriction of the $Cl_{n,1}$ -linear spinor bundle of N to M . It turns out that this bundle can be defined intrinsically – without reference to N – by $\overline{\Sigma}_{Cl}M = \Sigma_{Cl}M \otimes_{Cl_n} Cl_{n,1}$, i.e. it is given by two copies of $\Sigma_{Cl}M$. The role of the Dirac operator is now played by the Dirac-Witten operator \overline{D} , which is $Cl_{n,1}$ -linear in our case. There is a Schrödinger-Lichnerowicz type formula for \overline{D}

$$\overline{D}^2 = \overline{\nabla}^* \overline{\nabla} + \frac{1}{2}(\rho - e_0 \cdot j^\# \cdot),$$

which ensures that \overline{D} is invertible if (g, K) satisfies the dominant energy condition strictly. With these changes, the same construction as before yields an α -invariant for initial values

$$\overline{\alpha}: \pi_k(\mathcal{I}^+(M)) \cong \pi_{k+1}(\mathcal{I}(M), \mathcal{I}^+(M)) \rightarrow KO^{-n+1}(D^{k+1}, S^k) \cong KO^{-n-k}(\{*\}).$$

The main theorem of this paper compares these two α -invariants and relates the π_0 -part of the Lorentzian α -invariant to the α -index.

Main Theorem. 1. For $g_0 \in \mathcal{R}^+(M)$ and all $k \geq 0$, the diagram

$$\begin{array}{ccccc} \pi_k(\mathcal{R}^+(M), g_0) & \xrightarrow{\Sigma} & \pi_{k+1}(\Sigma \mathcal{R}^+(M), [g_0, 0]) & \xrightarrow{\Phi_*} & \pi_{k+1}(\mathcal{I}^+(M), (g_0, 0)) \\ & \searrow \alpha & & & \swarrow \overline{\alpha} \\ & & KO^{-n-k-1}(\{*\}) & & \end{array}$$

commutes, where $n = \dim M \geq 2$.

2. For $g_0 \in \mathcal{R}(M)$,

$$\overline{\alpha}: \pi_0 \left(\mathcal{I}^+(M), \left(g_0, \frac{1}{n} \tau(g_0) g_0 \right) \right) \longrightarrow KO^{-n}(\{*\})$$

sends the class defined by $S^0 = \{\pm 1\} \ni t \mapsto \left(g_0, \frac{1}{n} \tau(g_0) t g_0 \right)$ to $\alpha(M)$.

The idea of the proof is the following: For a pair of the form $(g, \frac{1}{n}\tau(g)tg)$, the Cl_{n+1} -linear Dirac-Witten operator is given by $\overline{D} = D \otimes_{Cl_n} Cl_{n,1} - \tau(g)tL(e_0)$, where D is the Cl_n -linear Dirac operator from before and $L(e_0)$ is left multiplication with the future-pointing unit normal on M when M is considered as spacelike hypersurface of N as above. Now, we observe that the $Cl_{n,1}$ -structure of $\overline{\Sigma}_{Cl}M$ given by right multiplication can be extended to a $Cl_{n+1,1}$ -structure by setting the right multiplication by the additional basis vector as $\tilde{R}(e_{n+1}) := L(e_0)\alpha$, where α is the even-odd grading operator. With this $Cl_{n+1,1}$ -structure, $\overline{\Sigma}_{Cl}M$ corresponds to $\Sigma_{Cl}M$ and $D \otimes_{Cl_n} Cl_{n,1}$ to D under the first Morita equivalence, and, by definition, the index map is invariant under this kind of equivalence. The second summand can be understood as coming from the Bott map, which assigns to a $Cl_{n+1,1}$ -Fredholm operator F the family of $Cl_{n,1}$ -Fredholm operators $[-1, 1] \ni t \mapsto F + t\tilde{R}(e_{n+1})\alpha = F + tL(e_0)$. Again, invariance of the index map under this assignment is a consequence of its definition, but an extra sign has to be taken into account resulting from the fact that in the definition of the index map Morita equivalence and Bott map are applied in the reverse order.

As a consequence of the main theorem, every element in $\pi_k(\mathcal{R}^+(M), g_0)$ with non-trivial α -invariant gives rise to a non-zero element in $\pi_{k+1}(\mathcal{I}^+(M), (g_0, 0))$. Such elements have been constructed for example by Hitchin [Hit74], Hanke, Schick and Steimle [HSS14], Botvinnik, Ebert and Randal-Williams [BER14] as well as Crowley, Schick and Steimle [CSS18] using different techniques. In particular, we obtain the following corollary.

- Corollary 1.1.** *1. If M is a compact spin manifold of dimension $n \geq 6$ that admits a metric of positive scalar curvature, then $\mathcal{I}^+(M)$ is not contractible.*
- 2. If M is a compact spin manifold of dimension $n \geq 2$ with $\alpha(M) \neq 0$ (in particular, M does not carry a positive scalar curvature metric), then $\mathcal{I}^+(M)$ is not connected.*

The structure of the article is as follows. In the first chapter, we review the KO -valued index map and the construction of the classical α -invariant. Much of this material is owed to Ebert [Ebe13]. The second chapter is devoted to the construction of the Lorentzian α -invariant. To this end, the $Cl_{n,1}$ -linear hypersurface spinor bundle and its Dirac-Witten operator are introduced. We discuss the $Cl_{n,1}$ -linear version of the Dirac-Witten operator in some detail, as it seems not to have been studied before. In the last chapter, we construct the comparison map and prove the main theorem.

Acknowledgements

I would like to thank Bernd Ammann for the idea for this project as well as his support during its execution.

2 The classical α -invariant

2.1 KO -theory via Fredholm operators

This section is devoted to the KO -valued index map, a map that associates to a family of Clifford-linear Fredholm operators an element in KO -theory. In its description, we will stick closely to the framework presented in Ebert [Ebe13] that we briefly recall. All Hilbert spaces are understood as being real and separable. A $Cl_{n,k}$ -Hilbert space H is always $\mathbb{Z}/2\mathbb{Z}$ -graded. Typically, the $\mathbb{Z}/2\mathbb{Z}$ -grading is given in terms of a grading operator $\iota: H \rightarrow H$, and the Clifford action is determined by a Clifford multiplication $c: \mathbb{R}^{n,k} \rightarrow \text{End}(H)$, where $\mathbb{R}^{n,k}$ is the pseudo-Euclidean vector space $\mathbb{R}^n \oplus \mathbb{R}^k$ with the standard inner product that is positive definite on the first factor and negative definite on the second one.

If (H, ι, c) is a $Cl_{n,k}$ -Hilbert space, then c gives rise to a representation $Cl_{n,k} \rightarrow \text{End}(H)$, which can be decomposed into irreducible ones. (H, ι, c) is called *ample*, if it contains each irreducible representation infinitely often. By the structure theory for real Clifford representations, this just means that H is infinite dimensional if $n - k \not\equiv 0 \pmod{4}$, and amounts to the condition that both the $+1$ - and the -1 -eigenspace of the volume element $\omega_{n,k} := \iota c(e_1) \cdots c(e_{n+k})$ are infinite dimensional if $n - k \equiv 0 \pmod{4}$.

Definition 2.1. Let (H, ι, c) be an ample $Cl_{n,k}$ -Hilbert space. Then a $Cl_{n,k}$ -Fredholm operator is a (bounded) Fredholm operator on H that is self-adjoint, odd with respect to ι , $Cl_{n,k}$ -linear and, in the case $n - k \equiv -1 \pmod{4}$, satisfies the additional condition that $\omega_{n,k} F \iota$ is neither essentially positive nor essentially negative. We denote by $\text{Fred}^{n,k}(H)$ the space of $Cl_{n,k}$ -Fredholm operators with operator norm topology. Furthermore, we write $G^{n,k}(H) \subseteq \text{Fred}^{n,k}(H)$ for the subspace of invertible elements.

The additional condition in the case where $n - k \equiv -1 \pmod{4}$ is needed to ensure bijectivity of the index map. Note that both this condition and the ampleness condition for $n - k \equiv 0 \pmod{4}$ become immediate if H is infinite dimensional and the Clifford multiplication extends, so that H becomes a $Cl_{n+1,k}$ - or $Cl_{n,k+1}$ -Hilbert space and the Fredholm operator F is Clifford-linear with respect to the extended Clifford multiplication; for then the additional generator anti-commutes with $\omega_{n,k}$ if $n - k \equiv 0 \pmod{4}$ and with $\omega_{n,k} F \iota$ if $n - k \equiv -1 \pmod{4}$. In consequence, $\text{Fred}^{n+1,k}(H) \subseteq \text{Fred}^{n,k}(H)$ and $\text{Fred}^{n,k+1}(H) \subseteq \text{Fred}^{n,k}(H)$.

Example 2.2. The archetypical example of a $Cl_{n,0}$ -Fredholm operator is (the bounded transform of) the Cl_n -linear Dirac operator on a closed Riemannian spin manifold (M, g) of dimension $n > 0$: Let $P_{\text{Spin}(n)}M \rightarrow P_{\text{SO}(n)}M$ be a spin structure of M . The Cl_n -linear spinor bundle is $\Sigma_{Cl}M := P_{\text{Spin}(n)}M \times_{\ell} Cl_n$, where $\ell: \text{Spin}(n) \rightarrow \text{End}(Cl_n)$ is given by left multiplication. Its name derives from the fact that right multiplication in Cl_n induces a *right Clifford multiplication* $R: \mathbb{R}^n \rightarrow \text{End}(\Sigma_{Cl}M)$, which commutes with the

left Clifford multiplication by tangent vectors. Furthermore, it carries a $\mathbb{Z}/2\mathbb{Z}$ -grading α induced by $Cl_n \rightarrow Cl_n$, $\mathbb{R}^n \ni v \mapsto -v$, the *even-odd-grading*. The bundle metric induced by the metric on Cl_n that makes the standard basis $(e_{i_1} \dots e_{i_l})_{0 \leq l \leq n+k, 1 \leq i_1 < \dots < i_l \leq n+k}$ orthonormal allows to define an L^2 -scalar product and the space of L^2 -sections $H := L^2(M, \Sigma_{Cl}M)$. Both α and R descend to H , turning (H, α, R) into an ample Cl_n -Hilbert space. The Cl_n -linear Dirac operator D , i.e. the Dirac operator of $\Sigma_{Cl}M$ w.r.t. the connection induced by the Levi-Civita connection, can be viewed as unbounded operator on H . By standard results on the analysis of Dirac operators, its bounded transform $F := \frac{D}{\sqrt{1+D^2}}$ is a Fredholm operator on H , and as D is Cl_n -linear (w.r.t. R) and odd (w.r.t. α), so is F . Thus, $F \in \text{Fred}^{n,0}(H)$, whereby the additional condition for $n \equiv -1 \pmod{4}$ follows from the Weyl asymptotic formula for Dirac operators. It is worth noting, that the Schrödinger-Lichnerowicz formula implies that F is invertible, so $F \in G^{n,0}(H)$, if g is a metric of positive scalar curvature.

The following consequence of Kuiper's theorem is proven in [Ebe13]. It is one of the main ingredients for translating the classical results from [AS69] into the present framework.

Proposition 2.3. *The space $G^{n,k}(H)$ is contractible for all $n, k \geq 0$.*

Theorem 2.4 (Index map). *If H is an ample $Cl_{n,k}$ -Hilbert space, then $\text{Fred}^{n,k}(H)$ represents KO -theory: For compact relative CW-complexes (X, Y) , there is a natural (in (X, Y)) bijection*

$$\text{ind}: [(X, Y), (\text{Fred}^{n,k}(H), G^{n,k}(H))] \longrightarrow KO^{k-n}(X, Y)$$

called index map. Moreover, ind is invariant under $Cl_{n,k}$ -Hilbert space isomorphisms, i.e. if $U: H \rightarrow H'$ is an isomorphism of $Cl_{n,k}$ -Hilbert spaces, then

$$\begin{array}{ccc} [(X, Y), (\text{Fred}^{n,k}(H), G^{n,k}(H))] & \xrightarrow{\cong} & [(X, Y), (\text{Fred}^{n,k}(H'), G^{n,k}(H'))] \\ \searrow \text{ind} & & \swarrow \text{ind} \\ & KO^{k-n}(X, Y) & \end{array}$$

commutes, where the upper map is induced by $\text{Fred}^{n,k}(H) \ni F \mapsto UFU^{-1}$.

The index map is constructed inductively, the starting point being the index of a family of $Cl_{0,0}$ -Fredholm operators, i.e. odd Fredholm operators on a $\mathbb{Z}/2\mathbb{Z}$ -graded Hilbert space. Here, the corresponding statement is known as Atiyah-Jänich theorem (cf. [Glö19, Thm. 2.17] for a detailed derivation from the version in [AS69]).

The generalization to arbitrary n (but still with $k = 0$) is provided by the Bott map.

Theorem 2.5 (Bott map, [AS69, Thm. A(k)]). *For compact CW-pairs (X, Y) , the map*

$$\begin{aligned} [(X, Y), (\text{Fred}^{n+1,k}(H), G^{n+1,k}(H))] &\longrightarrow [(X, Y) \times (I, \partial I), (\text{Fred}^{n,k}(H), G^{n,k}(H))] \\ [x \mapsto F_x] &\longmapsto [(x, t) \mapsto F_x + tc(e)\iota] \end{aligned}$$

is a natural bijection.¹ Thereby, e is the additional basis vector of $\mathbb{R}^{n+1,k}$ compared to $\mathbb{R}^{n,k}$ and $I = [-1, 1]$.

As $(X \times I)/(Y \times I \cup X \times \partial I) \cong \Sigma_{red} X/Y$ the right hand isomorphism in the following diagram exists, and the definition of the index map can be extended inductively by requiring that it commutes:

$$\begin{array}{ccc} [(X, Y), (\text{Fred}^{n,0}(H), G^{n,0}(H))] & \xrightarrow{\text{ind}} & KO^{-n}(X, Y) \\ \downarrow \cong & & \downarrow \cong \\ [(X, Y) \times (I, \partial I), (\text{Fred}^{n-1,0}(H), G^{n-1,0}(H))] & \xrightarrow{\text{ind}} & KO^{-n+1}(X \times I, X \times \partial I \cup Y \times I). \end{array} \quad (2)$$

The extension to arbitrary k uses periodicity statements in the theory of $Cl_{n,k}$ -Hilbert spaces known as Morita equivalences. The first Morita equivalence states that the categories of $Cl_{n,k}$ -Hilbert spaces and $Cl_{n+1,k+1}$ -Hilbert spaces are equivalent. Its construction is the following: A $Cl_{n,k}$ -Hilbert space (H, ι, c) defines a $Cl_{n+1,k+1}$ -Hilbert space structure on $H \oplus H$ by

$$\begin{aligned} \tilde{\iota} &= \begin{pmatrix} \iota & 0 \\ 0 & -\iota \end{pmatrix} \\ \tilde{c}(v) &= \begin{pmatrix} c(v) & 0 \\ 0 & -c(v) \end{pmatrix} \quad \text{for all } v \in R^{n+k} \oplus 0 \\ \tilde{c}(e) &= \begin{pmatrix} 0 & -\mathbb{1} \\ \mathbb{1} & 0 \end{pmatrix} \\ \tilde{c}(\varepsilon) &= \begin{pmatrix} 0 & \mathbb{1} \\ \mathbb{1} & 0 \end{pmatrix}, \end{aligned}$$

where we view $\mathbb{R}^{n+1,k+1}$ as $R^{n,k} \oplus \mathbb{R}e \oplus \mathbb{R}\varepsilon$. And a morphism $F: H \rightarrow H'$ of $Cl_{n,k}$ -Hilbert spaces gives rise to a morphism

$$\tilde{F} = \begin{pmatrix} F & 0 \\ 0 & F \end{pmatrix} : H \oplus H \rightarrow H' \oplus H'$$

of the corresponding $Cl_{n+1,k+1}$ -Hilbert spaces. Conversely, for a $Cl_{n+1,k+1}$ -Hilbert space (H, ι, c) , the restrictions of the structure maps to $H_0 := \ker(c(\varepsilon)c(e) - \mathbb{1})$ yield a $Cl_{n,k}$ -Hilbert space, and morphisms of $Cl_{n+1,k+1}$ -Hilbert spaces restrict to morphisms of these $Cl_{n,k}$ -Hilbert spaces. These constructions are seen to be mutually inverse up to natural isomorphism.

The second Morita equivalence is an equivalence between $Cl_{n+4,k}$ -Hilbert spaces and $Cl_{n,k+4}$ -Hilbert spaces. For this, we regard both $\mathbb{R}^{n+4,k}$ and $\mathbb{R}^{n,k+4}$ as $\mathbb{R}^n \oplus \mathbb{R}^k \oplus$

¹For two pairs (X, A) and (Y, B) , we write $(X, A) \times (Y, B) := (X \times Y, X \times B \cup A \times Y)$.

$\text{span}\{e_1, e_2, e_3, e_4\}$, where e_1, \dots, e_4 are the last four basis vectors of \mathbb{R}^{n+4} or the last four basis vectors of \mathbb{R}^{k+4} , respectively. Given a $Cl_{n+4, k}$ -Hilbert space (H, ι, c) , we can define a $Cl_{n, k+4}$ -Hilbert space (H, ι, \tilde{c}) by $\tilde{c}|_{\mathbb{R}^{n, k}} = c|_{\mathbb{R}^{n, k}}$ and $\tilde{c}(e_i) = \eta c(e_i)$ for $\eta = c(e_1) \cdots c(e_4)$. Morphisms are mapped to the morphisms defined by the same underlying bounded linear maps. The inverse procedure is given similarly, by assigning to a $Cl_{n, k+4}$ -Hilbert space (H, ι, \tilde{c}) the $Cl_{n+4, k}$ -Hilbert space (H, ι, c) with $c|_{\mathbb{R}^{n, k}} = \tilde{c}|_{\mathbb{R}^{n, k}}$ and $c(e_i) = \tilde{\eta} \tilde{c}(e_i)$, where $\tilde{\eta} = \tilde{c}(e_1) \cdots \tilde{c}(e_4)$.

These equivalences are accompanied by homoemorphisms between the spaces of Clifford-linear Fredholm operators.

Proposition 2.6. *The Morita equivalences induce homeomorphisms of pairs*

$$\begin{aligned} (\text{Fred}^{n, k}(H), G^{n, k}(H)) &\longrightarrow (\text{Fred}^{n+1, k+1}(H \oplus H), G^{n+1, k+1}(H \oplus H)) \\ F &\longmapsto \begin{pmatrix} F & 0 \\ 0 & F \end{pmatrix} \end{aligned}$$

and

$$\begin{aligned} (\text{Fred}^{n+4, k}(H), G^{n+4, k}(H)) &\longrightarrow (\text{Fred}^{n, k+4}(H), G^{n, k+4}(H)) \\ F &\longmapsto F. \end{aligned}$$

In particular, there is a homoemorphism

$$\begin{aligned} (\text{Fred}^{n, k}(H), G^{n, k}(H)) &\longrightarrow (\text{Fred}^{n+8, k}(H \otimes \mathbb{R}^{16}), G^{n+8, k}(H \otimes \mathbb{R}^{16})) \\ F &\longmapsto F \otimes \mathbb{1}_{\mathbb{R}^{16}}. \end{aligned}$$

The index map is then defined inductively for all (n, k) with $0 \leq k \leq n$ by the requirement that

$$\begin{array}{ccc} [(X, Y), (\text{Fred}^{n, k}(H), G^{n, k}(H))] & \xrightarrow{\text{ind}} & KO^{k-n}(X, Y) \\ \cong \uparrow & & \parallel \\ [(X, Y), (\text{Fred}^{n-1, k-1}(H_0), G^{n-1, k-1}(H_0))] & \xrightarrow{\text{ind}} & KO^{k-n}(X, Y) \end{array} \quad (3)$$

commutes. Lastly, it is extended to the missing (n, k) with $0 \leq n, k$ by commutativity of

$$\begin{array}{ccc} [(X, Y), (\text{Fred}^{n, k}(H), G^{n, k}(H))] & \xrightarrow{\text{ind}} & KO^{k-n}(X, Y) \\ \downarrow \cong & & \downarrow \cdot x \\ [(X, Y), (\text{Fred}^{n+8, k}(H \otimes \mathbb{R}^{16}), G^{n+8, k}(H \otimes \mathbb{R}^{16}))] & \xrightarrow{\text{ind}} & KO^{k-n-8}(X, Y), \end{array} \quad (4)$$

where x denotes a generator of $KO^{-8}(\{*\})$.

Remark 2.7. The commutativity of (4) does not only hold for $n < k$ (where it is true by definition), but is also true for $k \leq n$ provided that the right generator $x \in KO^{-8}(\{*\})$ is chosen. This follows from the last remark in [AS69].

Example 2.8. In the setting of Example 2.2, we can define the α -index of M by $\alpha(M) = \text{ind}(F) \in KO^{-n}(\{*\})$. This invariant was first defined by Hitchin [Hit74] and is a well-known obstruction to positive scalar curvature: From the continuity of the assignment $g \mapsto F_g$ discussed in the next section, it follows that $\alpha(M)$ is independent of the metric on M (in fact, it is even spin-bordism invariant) and so has to vanish for every spin structure if M carries a positive scalar curvature metric.

2.2 Construction of the α -invariant

Let M be a compact spin manifold of dimension $n > 0$ that has a positive scalar curvature metric g_0 . The α -invariant, also introduced by Hitchin [Hit74], is a family version of the α -index. More precisely, $\alpha: \pi_k(\mathcal{R}^+(M), g_0) \rightarrow KO^{-n-k-1}(\{*\})$ arises in the following way: As $\mathcal{R}(M)$ is contractible, the long exact sequence for homotopy groups implies $\pi_k(\mathcal{R}^+(M), g_0) \cong \pi_{k+1}(\mathcal{R}(M), \mathcal{R}^+(M), g_0)$. For each metric g , the Cl_n -linear Dirac operator D_g defines a Cl_n -linear Fredholm operator

$$F_g = \frac{D_g}{\sqrt{1 + D_g^2}},$$

which is invertible if $g \in \mathcal{R}^+(M)$. The assignment $g \mapsto F_g$ gives rise to a map $(\mathcal{R}(M), \mathcal{R}^+(M)) \rightarrow (\text{Fred}^{n,0}, G^{n,0})$, which induces a map to $\pi_{k+1}(\text{Fred}^{n,0}, G^{n,0}, F_{g_0})$. Applying the index map from the last section, we obtain an element in $KO^{-n}(D^{k+1}, S^k) \cong KO^{-n-k-1}(\{*\})$.

In this outline, however, we glossed over the detail that the Cl_n -linear spinor bundles and hence the L^2 -spaces, on which the Fredholm operators F_g act, depend on the metric g . These L^2 -spaces form a Hilbert bundle over $\mathcal{R}(M)$, which, by Kuiper's theorem, can be trivialized. Such a trivialization allows to define the map $(\mathcal{R}(M), \mathcal{R}^+(M)) \rightarrow (\text{Fred}^{n,0}, G^{n,0})$. We will make this more explicit: The Cl_n -linear spinor bundles for different metrics can be identified using the method of generalized cylinders due to Bär, Gauduchon and Moroianu [BGM05]. This gives rise to a specific trivialization of the Hilbert bundle of L^2 -spaces.

Let us start with this construction by fixing a topological spin structure on M , i.e. a double covering

$$P_{\widetilde{GL}^+(n)} M \rightarrow P_{GL^+(n)} M$$

over the principal bundle of positively oriented frames of TM . This defines, for any $g \in \mathcal{R}(M)$, a spin structure for (M, g) by pullback

$$\begin{array}{ccc} P_{\text{Spin}(n)}(M, g) & \longrightarrow & P_{\widetilde{GL}^+(n)} M \\ \downarrow & & \downarrow \\ P_{\text{SO}(n)}(M, g) & \longrightarrow & P_{GL^+(n)} M, \end{array}$$

where $P_{\text{SO}(n)}(M, g)$ is the principal bundle of positively oriented orthonormal frames with respect to g . Moreover, pulling back over the canonical projection $M \times [0, 1] \rightarrow M$, we obtain

$$\begin{array}{ccc} P_{\widetilde{GL}^+(n)} M \times [0, 1] & \longrightarrow & P_{\widetilde{GL}^+(n)} M \\ \downarrow & & \downarrow \\ P_{GL^+(n)} M \times [0, 1] & \longrightarrow & P_{GL^+(n)} M \\ \downarrow & & \downarrow \\ M \times [0, 1] & \longrightarrow & M. \end{array}$$

This gives rise a topological spin structure $P_{\widetilde{GL}^+(n+1)} M \times [0, 1] \rightarrow P_{GL^+(n+1)} M \times [0, 1]$ on $M \times [0, 1]$ by extension along the standard embedding

$$\begin{aligned} GL^+(n) &\longrightarrow GL^+(n+1) \\ A &\longmapsto \begin{pmatrix} A & 0 \\ 0 & 1 \end{pmatrix} \end{aligned}$$

and its double covering.

Now, given a metric $g \in \mathcal{R}(M)$, we can define a family of metrics by $g_t = (1-t)g_0 + tg$. Such a family in turn defines the generalized cylinder $(M \times [0, 1], g_t + dt^2)$, t being the variable in $[0, 1]$ -direction. As above, the topological spin structure induces a spin structure $P_{\text{Spin}(n+1)}(M \times [0, 1], g_t + dt^2) \rightarrow P_{\text{SO}(n+1)}(M \times [0, 1], g_t + dt^2)$ on the generalized cylinder. This has the property that for all $t_0 \in [0, 1]$ it restricts to the spin structure of (M, g_{t_0}) in the sense that

$$\begin{array}{ccc} P_{\text{Spin}(n)}(M, g_{t_0}) & \longrightarrow & P_{\text{Spin}(n+1)}(M \times [0, 1], g_t + dt^2) \\ \downarrow & & \downarrow \\ P_{\text{SO}(n)}(M, g_{t_0}) & \longrightarrow & P_{\text{SO}(n+1)}(M \times [0, 1], g_t + dt^2) \end{array}$$

is a pullback, where the lower map is the inclusion $(e_1, \dots, e_n) \mapsto (e_1, \dots, e_n, \frac{\partial}{\partial t})$.

The reason, why we do this is that on $P_{\text{Spin}(n+1)}(M \times [0, 1], g_t + dt^2)$ the Levi-Civita connection induces a canonical connection ∇ , which provides parallel transports

$$P_{\gamma_x}^\nabla : P_{\text{Spin}(n+1)}(M \times [0, 1], g_t + dt^2)|_{(x,0)} \longrightarrow P_{\text{Spin}(n+1)}(M \times [0, 1], g_t + dt^2)|_{(x,1)}$$

along the curves $\gamma_x: [0, 1] \rightarrow M \times [0, 1]$, $t \mapsto (x, t)$ for all $x \in M$. These assemble into an isomorphism of principle bundles

$$P^\nabla: P_{\text{Spin}(n+1)}(M \times [0, 1], g_t + dt^2)|_{M \times \{0\}} \xrightarrow{\cong} P_{\text{Spin}(n+1)}(M \times [0, 1], g_t + dt^2)|_{M \times \{1\}}.$$

The fact that $\frac{\partial}{\partial t}$ is parallel along the curves γ_x implies that P^∇ restricts to

$$P^\nabla: P_{\text{Spin}(n)}(M, g_0) \xrightarrow{\cong} P_{\text{Spin}(n)}(M, g),$$

and this induces an isomorphism on the associated Cl_n -linear spinor bundles

$$\begin{aligned} P^\nabla: \Sigma_{Cl}(M, g_0) &\xrightarrow{\cong} \Sigma_{Cl}(M, g). \\ [\tilde{\varepsilon}, \tilde{\phi}] &\mapsto [P^\nabla \tilde{\varepsilon}, \tilde{\phi}] \end{aligned}$$

Furthermore, it is immediate that P^∇ is a point-wise isometry with respect to the standard scalar products $\langle -, - \rangle$ defined on the Cl_n -linear spinor bundles.

We want to promote this to a unitary transformation between the associated L^2 -spaces. As the L^2 -norm also depends on the volume element, we first compare those: There exists a positive function $\beta \in C^\infty(M)$ such that $d\text{vol}^g = \beta d\text{vol}^{g_0}$. Then $\sqrt{\beta}P^\nabla: \Sigma_{Cl}(M, g_0) \rightarrow \Sigma_{Cl}(M, g)$ induces a unitary transformation

$$\Phi_g: H := L^2(M, \Sigma_{Cl}(M, g_0)) \xrightarrow{\cong} L^2(M, \Sigma_{Cl}(M, g))$$

as

$$(\Phi_g(\phi), \Phi_g(\psi))_{L^2} = \int_M \langle \sqrt{\beta}P^\nabla(\phi), \sqrt{\beta}P^\nabla(\psi) \rangle d\text{vol}^g = \int_M \langle \phi, \psi \rangle d\text{vol}^{g_0} = (\phi, \psi)_{L^2}.$$

Moreover, it is clear that Φ_g preserves $\mathbb{Z}/2\mathbb{Z}$ -grading and the right Clifford multiplication. The left Clifford multiplication by a vector field $X \in \mathfrak{X}(M)$ satisfies $\Phi_g(X \cdot \phi) = P^\nabla(X) \cdot \Phi_g(\phi)$ for any $\phi \in H$, where $P^\nabla(X)$ is the vector field obtained from X by parallel transport along the curves $(\gamma_x)_{x \in M}$.

It is not surprising that using this identification of the L^2 -spaces (the bounded transforms of) the Dirac operators depend continuously on the metric. For a detailed proof of the following statement see [Glö19, Thm. 2.22].

Theorem 2.9. *The map*

$$\begin{aligned} (\mathcal{R}(M), \mathcal{R}^+(M)) &\longrightarrow (\text{Fred}^{n,0}(H), G^{n,0}(H)) \\ g &\longmapsto \Phi_g^{-1} \circ \frac{D_g}{\sqrt{1 + D_g^2}} \circ \Phi_g \end{aligned}$$

is well-defined and continuous with respect to the C^1 -topology on the space of smooth metrics $\mathcal{R}(M)$. In particular, it is continuous if $\mathcal{R}(M)$ carries the C^∞ -topology.

Definition 2.10. The map from Theorem 2.9 gives rise to the composition

$$\begin{aligned} \alpha: \pi_k(\mathcal{R}^+(M), g_0) &\cong \pi_{k+1}(\mathcal{R}(M), \mathcal{R}^+(M), g_0) \rightarrow [(D^{k+1}, S^k), (\mathcal{R}(M), \mathcal{R}^+(M))] \\ &\rightarrow [(D^{k+1}, S^k), (\text{Fred}^{n,0}, G^{n,0})] \cong KO^{-n-k-1}(\{*\}) \end{aligned}$$

called α -invariant.

The α -invariant detects non-trivial homotopy groups in the space of metrics of positive scalar curvature. The following two results of this kind were independently obtained by different methods:

Theorem 2.11 (Crowley, Schick, Steimle [CSS18]). *Let (M, g_0) be a compact Riemannian spin manifold of positive scalar curvature and $n = \dim(M) \geq 6$. For all $k \geq 0$ with $k + n + 1 \equiv 1, 2 \pmod{8}$, the α -invariant*

$$\alpha: \pi_k(\mathcal{R}^+(M), g_0) \longrightarrow KO^{-n-k-1}(\{*\}) \cong \mathbb{Z}/2\mathbb{Z}$$

is split surjective.

Theorem 2.12 (Botvinnik, Ebert, Randal-Williams [BER14]). *Let (M, g_0) be a compact Riemannian spin manifold of positive scalar curvature and $n = \dim(M) \geq 6$. For all $k \geq 0$ with $k + n + 1 \equiv 1, 2 \pmod{8}$, the α -invariant*

$$\alpha: \pi_k(\mathcal{R}^+(M), g_0) \longrightarrow KO^{-n-k-1}(\{*\}) \cong \mathbb{Z}/2\mathbb{Z}$$

is surjective, and for all $k \geq 0$ with $k + n + 1 \equiv 0, 4 \pmod{8}$, the localized α -invariant

$$\alpha \otimes \mathbb{1}_{\mathbb{Q}}: \pi_k(\mathcal{R}^+(M), g_0) \otimes \mathbb{Q} \longrightarrow KO^{-n-k-1}(\{*\}) \otimes \mathbb{Q} \cong \mathbb{Q}$$

is surjective.

We will use these results to construct non-trivial homotopy groups in the space of initial value pairs satisfying the dominant energy condition. The detection of these groups then uses an α -invariant for initial values that will be defined in the next chapter.

3 An α -invariant for initial values

3.1 The $Cl_{n,1}$ -linear hypersurface spinor bundle

In this section, we want to study the bundle obtained by restricting the $Cl_{n,1}$ -linear spinor bundle of a space- and time-oriented Lorentzian spin manifold (N, \bar{g}) to a spacelike hypersurface $M \subseteq N$. Especially, we want to describe it intrinsically, only in terms of

quantities of M , the induced metric g and the second fundamental form K . This will be of use later, when defining the α -invariant for initial values and comparing it to the classical α -invariant.

The first step is to construct compatible spin structures on M and N . Fixing a spin structure on (N, \bar{g}) , we obtain a spin structure on (M, g) by pulling back the one from N :

$$\begin{array}{ccc} P_{\text{Spin}(n)}(M) & \longrightarrow & P_{\text{Spin}_0(n,1)}(N)|_M \\ \downarrow & & \downarrow \\ P_{\text{SO}(n)}(M) & \longrightarrow & P_{\text{SO}_0(n,1)}(N)|_M. \end{array} \quad (5)$$

Thereby, the lower map is given by $(e_1, \dots, e_n) \mapsto (e_0, e_1, \dots, e_n)$, where e_0 is the future-pointing unit normal on M . As the right hand map is a double covering, so is the left hand one, and it suffices to construct a compatible $\text{Spin}(n)$ -action. This, we obtain by pulling back the action maps. More explicitly, there is a commutative diagram

$$\begin{array}{ccc} P_{\text{Spin}(n)}(M) \times \text{Spin}(n) & \longrightarrow & P_{\text{Spin}_0(n,1)}(N)|_M \times \text{Spin}_0(n,1) \\ \downarrow & & \downarrow \\ P_{\text{SO}(n)}(M) \times \text{SO}(n) & \longrightarrow & P_{\text{SO}_0(n,1)}(N)|_M \times \text{SO}_0(n,1). \end{array} \quad (6)$$

and the desired map is the unique map from its upper-left corner to the upper-left corner of (5) building, together with the other action maps, a commutative cube out of (5) and (6). Note, that this commutative cube shows that $P_{\text{Spin}(n)}(M)$ is not only a $\text{Spin}(n)$ -reduction of $P_{\text{SO}(n)}(M)$ but also a reduction of $P_{\text{Spin}_0(n,1)}(N)|_M$ with respect to the inclusion $i: \text{Spin}(n) \hookrightarrow \text{Spin}_0(n,1)$.

Next, we study associated bundles. The $Cl_{n,1}$ -linear spinor bundle

$$\Sigma_{Cl}N = P_{\text{Spin}_0(n,1)}(N) \times_{\ell} Cl_{n,1}$$

is defined via the representation induced by left multiplication on $Cl_{n,1}$:

$$\ell: \text{Spin}_0(n,1) \hookrightarrow Cl_{n,1} \longrightarrow \text{End}(Cl_{n,1}).$$

As noted above, $P_{\text{Spin}(n)}(M) \rightarrow P_{\text{Spin}_0(n,1)}(N)|_M$ is a $\text{Spin}(n)$ -reduction. Hence, from the theory of principal bundles (e.g. [Bau14, Satz 2.18]), it follows that

$$\Sigma_{Cl}N|_M = P_{\text{Spin}_0(n,1)}(N)|_M \times_{\ell} Cl_{n,1} \cong P_{\text{Spin}(n)}(M) \times_{\ell i} Cl_{n,1}, \quad (7)$$

so the bundle $\Sigma_{Cl}N|_M \rightarrow M$ only depends on the Riemannian manifold (M, g) and its chosen spin structure.

Definition 3.1. The bundle $\Sigma_{Cl}N|_M$ from above is called $Cl_{n,1}$ -linear hypersurface spinor bundle and denoted by $\bar{\Sigma}_{Cl}M$.

Remark 3.2. It is also possible to express $\overline{\Sigma}_{Cl}M$ in terms of the Cl_n -linear spinor bundle on M defined in Example 2.2: In view of (7), the canonical isomorphism $Cl_{n,1} \cong Cl_n \otimes_{Cl_n} Cl_{n,1}$, where Cl_n acts by right multiplication on Cl_n and by left multiplication on $Cl_{n,1}$, implies that $\overline{\Sigma}_{Cl}M \cong P_{\text{Spin}(n)}(M) \times_{\ell_i} Cl_{n,1} \cong \Sigma_{Cl}M \otimes_{Cl_n} Cl_{n,1}$. Informally spoken, this means the $Cl_{n,1}$ -linear hypersurface spinor bundle consists of two copies of the Cl_n -linear spinor bundle. In fact, we will see that $\overline{\Sigma}_{Cl}M$ canonically carries a $Cl_{n+1,1}$ -linear structure, and corresponds to $\Sigma_{Cl}M$ under the first Morita equivalence.

Similarly to the case of the Cl_n -linear spinor bundle, the $Cl_{n,1}$ -linear hypersurface spinor bundle carries a right Clifford multiplication $R: \mathbb{R}^{n,1} \rightarrow \text{End}(\overline{\Sigma}_{Cl}M)$ and an even-odd grading $\alpha: \overline{\Sigma}_{Cl}M \rightarrow \overline{\Sigma}_{Cl}M$ as the corresponding notions for $Cl_{n,1}$ are $\text{Spin}_0(n,1)$ -invariant. Despite not being $\text{Spin}_0(n,1)$ -invariant, the scalar product $\langle -, - \rangle$ on $Cl_{n,1}$ for which the basis² $(e_{i_1}e_{i_2} \cdots e_{i_k})_{0 \leq k \leq n, 0 \leq i_1 < \cdots < i_k \leq n}$ is orthonormal can be extended to $\overline{\Sigma}_{Cl}M$: Due to (7), $\text{Spin}(n)$ -invariance of $\langle -, - \rangle$ is sufficient. This scalar product gives rise to a space of L^2 -sections $\overline{H} := L^2(M, \overline{\Sigma}_{Cl}M)$, on which R and α define a $Cl_{n,1}$ -Hilbert space structure.

Yet, the trivialization of $TN|_M$ by e_0 allows us to do better. We immediately obtain the following result:

Proposition 3.3. *Setting*

$$\Psi \cdot e_{n+1} := e_0 \cdot \alpha(\Psi)$$

for all $\Psi \in \overline{\Sigma}_{Cl}M$, R extends to a $Cl_{n+1,1}$ -multiplication

$$\tilde{R}: \mathbb{R}^{n+2} \rightarrow \text{End}(\overline{\Sigma}_{Cl}M).$$

that commutes with left multiplication by any $X \in TM$. Moreover, $(\overline{H}, \alpha, \tilde{R})$ is an ample $Cl_{n+1,1}$ -Hilbert space.

This $Cl_{n+1,1}$ -Hilbert space structure establishes the connection to the space H of L^2 -sections of the Cl_n -linear spinor bundle $\Sigma_{Cl}M$.

Proposition 3.4. *The $Cl_{n+1,1}$ -Hilbert space $(\overline{H}, \alpha, \tilde{R})$ corresponds to the Cl_n -Hilbert space (H, α, R) under the first Morita equivalence.*

Proof. Via the first Morita equivalence, the $Cl_{n+1,1}$ -Hilbert space \overline{H} corresponds to the $Cl_{n,0}$ -Hilbert space $\overline{H}_0 = \ker(\tilde{R}(e_0)\tilde{R}(e_{n+1}) - \mathbb{1})$ with the structure obtained by

²For consistency with Lorentzian geometry, the basis vector of the negative definite part of $\mathbb{R}^{n,1}$ is called e_0 rather than e_{n+1} .

restriction. $\tilde{R}(e_0)\tilde{R}(e_{n+1})$ is induced by a map $Cl_{n+1} \rightarrow Cl_{n+1}$, which in turn is induced by the endomorphism

$$\begin{aligned}\mathbb{R}^{n,1} &\longrightarrow \mathbb{R}^{n,1} \\ v &\longmapsto -e_0 v e_0\end{aligned}$$

reflecting at the hyperplane orthogonal to the line $\mathbb{R}e_0$. So the 1-eigenspace of the map on Cl_{n+1} is given by $Cl_n \subseteq Cl_{n+1}$ and the -1 -eigenspace is $R(e_0)Cl_n \subseteq Cl_{n+1}$, where Cl_n is viewed as the subalgebra generated by e_1, \dots, e_n . This implies that

$$\overline{H}_0 = L^2(M, \ker(\tilde{R}(e_0)\tilde{R}(e_{n+1}) - \mathbb{1})) = L^2(M, P_{\text{Spin}(n)}M \times_{\ell} Cl_n) = H.$$

So \overline{H} and H correspond to each other under the first Morita equivalence. \square

As a consequence of (7), the $Cl_{n,1}$ -linear hypersurface spinor bundle possesses two natural connections: On the one hand, the Levi-Civita connection (N, \overline{g}) induces a connection $\overline{\nabla}$ on $P_{\text{Spin}_0(n,1)}N|_M$ and $\overline{\Sigma}_{Cl}M$. On the other hand, as bundle associated to $P_{\text{Spin}(n)}M$, the bundle $\overline{\Sigma}_{Cl}M$ carries a connection ∇ induced by the Levi-Civita connection of (M, g) . They are related by the Weingarten map (also known as shape operator):

Lemma 3.5. *For all $X \in TM$ and $\psi \in \Gamma(\overline{\Sigma}_{Cl}M)$*

$$\overline{\nabla}_X \psi = \nabla_X \psi - \frac{1}{2}e_0 \cdot W(X) \cdot \psi$$

holds, where $W(X) = \overline{\nabla}_X e_0$ is the Weingarten map³.

Proof. Let $\tilde{\varepsilon}$ be a local section of $P_{\text{Spin}(n)}M$, and (e_1, \dots, e_n) its projection to $P_{SO(n)}M$. Writing a spinor locally as $\psi = [\tilde{\varepsilon}, \tilde{\psi}]$ and using the local formula for the spinorial connection, we perform the following local calculation:

$$\begin{aligned}\overline{\nabla}_X \psi - \nabla_X \psi &= \left[\tilde{\varepsilon}, \partial_X \tilde{\psi} + \frac{1}{2} \sum_{0 \leq i < j} \varepsilon_i \overline{g}(\overline{\nabla}_X e_i, e_j) e_i \cdot e_j \cdot \tilde{\psi} \right] \\ &\quad - \left[\tilde{\varepsilon}, \partial_X \tilde{\psi} + \frac{1}{2} \sum_{1 \leq i < j} \overline{g}(\nabla_X e_i, e_j) e_i \cdot e_j \cdot \tilde{\psi} \right] \\ &= \left[\tilde{\varepsilon}, \frac{1}{2} \sum_{0 < j} (-1) g(\overline{\nabla}_X e_0, e_j) e_0 \cdot e_j \cdot \tilde{\psi} \right] \\ &= -\frac{1}{2} e_0 \cdot W(X) \cdot \psi.\end{aligned}$$

\square

³The sign of W is different than in Riemannian geometry. It is chosen such that for all $X, Y \in TM$, $K(X, Y) = -\overline{g}(\mathbb{I}(X, Y), e_0) = -\overline{g}(\overline{\nabla}_X Y, e_0) = \overline{g}(Y, \overline{\nabla}_X e_0) = g(Y, W(X))$ holds, so $W = K^\sharp$.

By the way α , R and $\langle -, - \rangle$ are defined, it is clear that they are ∇ -parallel. The left Clifford multiplication $L: TN|_M \otimes \overline{\Sigma}_{Cl}M \rightarrow \overline{\Sigma}_{Cl}M$ is ∇ -parallel as well, where ∇ is defined on $TN|_M$ by viewing it as bundle associated to $P_{SO(n)}M$ via the lower map of (5). This can be reexpressed by saying that both the restricted left Clifford multiplication $TM \otimes \overline{\Sigma}_{Cl}M \rightarrow \overline{\Sigma}_{Cl}M$ and the endomorphism $\overline{\Sigma}_{Cl}M \rightarrow \overline{\Sigma}_{Cl}M$ given by left multiplication with e_0 are ∇ -parallel. As a consequence, the extended right Clifford multiplication \tilde{R} is ∇ -parallel as well.

With respect to the other connection, the following can be said. α , R and L are $\overline{\nabla}$ -parallel. The scalar product $\langle -, - \rangle$, however, in general is not, as it does not originate from a $\text{Spin}_0(n, 1)$ -invariant scalar product on $Cl_{n,1}$. Instead, it satisfies the following formula that follows from ∇ -parallelism together with Lemma 3.5:

$$\partial_X \langle \phi, \psi \rangle = \langle \overline{\nabla}_X \phi, \psi \rangle + \langle \phi, \overline{\nabla}_X \psi \rangle + \langle e_0 \cdot W(X) \cdot \phi, \psi \rangle.$$

3.2 $Cl_{n,1}$ -linear Dirac-Witten operator and α -invariant for initial values

As before, let M be a spacelike hypersurface of a space- and time-oriented Lorentzian spin manifold (N, \overline{g}) . The Dirac-Witten operator is a kind of Dirac operator on the hypersurface spinor bundle. In the case of classical spinor bundles, it was first defined by Witten [Wit81] in order to give his spinorial proof of the positive mass theorem (cf. [PT82] for a rigorous formulation of the proof) and later studied in more detail by Hijazi and Zhang [HZ03]. We are interested in its $Cl_{n,1}$ -linear version and use it to define a kind of α -invariant for initial values. Furthermore, we compare it to the $Cl_{n,1}$ -linear Dirac operator, which will be of later use.

Definition 3.6. The composition

$$\overline{D}: \Gamma(\overline{\Sigma}_{Cl}M) \xrightarrow{\overline{\nabla}} \Gamma(T^*M \otimes \overline{\Sigma}_{Cl}M) \xrightarrow{\# \otimes 1} \Gamma(TM \otimes \overline{\Sigma}_{Cl}M) \xrightarrow{L} \Gamma(\overline{\Sigma}_{Cl}M)$$

defines the $Cl_{n,1}$ -linear Dirac-Witten operator. The composition (with $\overline{\nabla}$ replaced by ∇)

$$D: \Gamma(\overline{\Sigma}_{Cl}M) \xrightarrow{\nabla} \Gamma(T^*M \otimes \overline{\Sigma}_{Cl}M) \xrightarrow{\# \otimes 1} \Gamma(TM \otimes \overline{\Sigma}_{Cl}M) \xrightarrow{L} \Gamma(\overline{\Sigma}_{Cl}M)$$

is the $Cl_{n,1}$ -linear Dirac operator.

The following lemma justifies the names of these operators. It is a direct consequence of the parallelism discussion at the end of the last section.

Lemma 3.7. \overline{D} and D are both $Cl_{n,1}$ -linear with respect to the right Clifford multiplication R and odd with respect to α . Furthermore, D is $Cl_{n+1,1}$ -linear with respect to the extended right Clifford multiplication \tilde{R} .

Lemma 3.8. $\overline{D} = D - \frac{1}{2}\tau L(e_0)$ holds, where $\tau = \text{tr } W = \text{tr } K$ is the mean curvature of M in N . Both D and \overline{D} are formally self-adjoint.

Proof. For $\psi \in \Gamma(\overline{\Sigma}_{Cl}M)$ and a local orthonormal frame e_1, \dots, e_n we perform the following local calculation:

$$\begin{aligned} \overline{D}\psi - D\psi &= \sum_{i=1}^n e_i \cdot (\overline{\nabla}_{e_i} - \nabla_{e_i})\psi \\ &= -\frac{1}{2} \sum_{i=1}^n e_i \cdot e_0 \cdot W(e_i) \cdot \psi \\ &= \frac{1}{2} \sum_{i,j=1}^n g(W(e_i), e_j) e_i \cdot e_j \cdot e_0 \cdot \psi \\ &= -\frac{1}{2} \sum_{i=1}^n g(W(e_i), e_i) e_0 \cdot \psi. \end{aligned}$$

Here, we used that $g(W(e_i), e_j) = K(e_i, e_j)$ is symmetric in i and j . Being a Dirac operator, D is formally self-adjoint. As left multiplication with e_0 is self-adjoint as well, the same holds true for \overline{D} . \square

The utility of the Dirac-Witten operator to general relativity results from following observation due to Witten [Wit81]:

Proposition 3.9. *The Dirac-Witten operator satisfies the Schrödinger-Lichnerowicz type formula*

$$\overline{D}^2 = \overline{\nabla}^* \overline{\nabla} + \frac{1}{2}(\rho - e_0 \cdot j^\sharp),$$

with

$$\begin{aligned} 2\rho &= \text{scal} + \tau - \|K\|^2 \\ j &= -d\tau + \text{div } K. \end{aligned}$$

From now on, we assume that M is compact.

Corollary 3.10. *If the pair (g, K) satisfies the strict dominant energy condition, i.e. if $\rho > \|j\|$, then \overline{D} has empty kernel.*

Proof. For any $\psi \in \Gamma(\overline{\Sigma}_{Cl}M)$ with $\psi \neq 0$

$$\begin{aligned} \|\overline{D}\psi\|_{L^2}^2 &= (\psi, \overline{D}\overline{D}\psi) = \|\overline{\nabla}\psi\|_{L^2}^2 + \frac{1}{2}(\psi, \rho\psi) - \frac{1}{2}(\psi, e_0 \cdot j^\sharp \cdot \psi) \\ &\geq \frac{1}{2}(\psi, \rho\psi) - \frac{1}{2}(\psi, \|j\|\psi) = \frac{1}{2}(\psi, (\rho - \|j\|)\psi) > 0 \end{aligned}$$

holds as $|\langle \psi, e_0 \cdot j^\sharp \cdot \psi \rangle| \leq \|j\| \|\psi\|^2$. Here, $\| - \|$ (without subscript L^2) denotes the pointwise norm. \square

Proposition 3.11. \overline{D} and D extend to densely defined operators

$$D, \overline{D}: L^2(M, \overline{\Sigma}_{Cl}M) \supseteq H^1(M, \overline{\Sigma}_{Cl}M) \rightarrow L^2(M, \overline{\Sigma}_{Cl}M)$$

admitting a spectral decomposition with discrete spectrum and finite dimensional eigenspaces.

Proof. Recall that a generalized Dirac operator on a vector bundle $\Sigma \rightarrow M$ in the sense of Roe [Roe99] is a formally self-adjoint operator \tilde{D} with

$$\tilde{D}^2 = \nabla^* \nabla + A$$

for a metric connection ∇ on Σ and some bounded operator $A: L^2(M, \Sigma) \rightarrow L^2(M, \Sigma)$. D is such an operator by the Schrödinger-Lichnerowicz formula and due to

$$\begin{aligned} \overline{D}^2 \psi &= \left(D - \frac{1}{2} \tau e_0 \cdot \right) \left(D - \frac{1}{2} \tau e_0 \cdot \right) \psi \\ &= D^2 \psi - \frac{1}{2} D(\tau e_0 \cdot \psi) - \frac{1}{2} \tau e_0 \cdot D\psi + \frac{1}{4} \tau^2 \psi \\ &= D^2 \psi + \frac{1}{2} e_0 \cdot \text{grad } \tau \cdot \psi + \frac{1}{4} \tau^2 \psi \end{aligned}$$

the same holds for \overline{D} . Now the proposition is just a special case of the corresponding statement for generalized Dirac operators [Roe99, Thm 5.27]. \square

Corollary 3.12. If $n = \dim(M) > 0$ and $\overline{H} := L^2(M, \overline{\Sigma}_{Cl}M)$, then there are well-defined elements

$$\overline{F} := \frac{\overline{D}}{\sqrt{1 + \overline{D}^2}} \in \text{Fred}^{n,1}(\overline{H})$$

and

$$F := \frac{D}{\sqrt{1 + D^2}} \in \text{Fred}^{n+1,1}(\overline{H}) \subseteq \text{Fred}^{n,1}(\overline{H}).$$

Furthermore, \overline{F} is invertible if (g, K) satisfies the strict dominant energy condition and F is invertible if g has positive scalar curvature.

Proof. \overline{H} is ample as $Cl_{n+1,1}$ -Hilbert space, so it is ample as $Cl_{n,1}$ -Hilbert space with the restricted Clifford action as well. As \overline{D} is odd and $Cl_{n,1}$ -linear, so is \overline{F} . From the proposition above, we conclude that \overline{F} is a Fredholm operator. The additional condition in the case $n - 1 \equiv -1 \pmod{4}$ is again a consequence of the Weyl asymptotics. Invertibility for (g, K) satisfying the strict dominant energy condition follows from Corollary 3.10 and coker $\overline{F} = \ker \overline{F}$. The argumentation for F is completely analogous. Invertibility here uses the classical Schrödinger-Lichnerowicz formula. \square

If the mean curvature τ is constant, we can relate the spectral decompositions of \overline{D} and D and refine the invertibility result.

Proposition 3.13. *The spectral decomposition of D can be written as*

$$D = \sum_{k=0}^{\infty} \lambda_k \pi_{E_k} + \sum_{k=0}^{\infty} (-\lambda_k) \pi_{\alpha(E_k)}$$

where all $\lambda_k > 0$ are pairwise disjoint and π_{E_k} and $\pi_{\alpha(E_k)}$ are the orthogonal projections on the finite dimensional subspaces E_k and $\alpha(E_k)$, respectively. If the mean curvature τ is constant, then there are decompositions $F_k \oplus \alpha(F_k) = E_k \oplus \alpha(E_k)$ for all $k \geq 0$ and $K \oplus \alpha(K) = \ker D$ such that the spectral decomposition of \overline{D} is given by

$$\overline{D} = \sum_{k=0}^{\infty} \sqrt{\lambda_k^2 + \frac{1}{4}\tau^2} \pi_{F_k} + \sum_{k=0}^{\infty} \left(-\sqrt{\lambda_k^2 + \frac{1}{4}\tau^2} \right) \pi_{\alpha(F_k)} + \frac{1}{2}\tau \pi_K - \frac{1}{2}\tau \pi_{\alpha(K)}$$

In particular, \overline{D} is invertible for all constants $\tau \neq 0$.

Proof. As α anti-commutes with D , for any eigenvector ϕ to the eigenvalue λ

$$D\alpha(\phi) = -\alpha(D\phi) = -\alpha(\lambda\phi) = -\lambda\alpha(\phi).$$

So $\alpha(\phi)$ is an eigenvector to the eigenvalue $-\lambda$. This implies that the spectral decomposition can be written in the stated form. With the same argument, we also expect the spectral decomposition of \overline{D} to be of that form.

\tilde{R} anti-commutes with D , so the eigenspaces are invariant under $\tilde{R}(v)$ for all $v \in \mathbb{R}^{n+2}$. In particular,

$$\alpha(E_k) = \tilde{R}(e_{n+1})\alpha(E_k) = L(e_0)(E_k)$$

for all $k \geq 0$. Thus we can identify E_k with $\alpha(E_k)$ via the map $E_k \rightarrow \alpha(E_k)$, $\phi \mapsto L(e_0)(\phi)$ and get $E_k \oplus \alpha(E_k) \cong E_k \oplus E_k \cong E_k \otimes \mathbb{R}^2$. Under this identification, by Lemma 3.8, the restriction of the Dirac-Witten operator corresponds to

$$\mathbb{1}_{E_k} \otimes \begin{pmatrix} \lambda_k & -\frac{1}{2}\tau \\ -\frac{1}{2}\tau & -\lambda_k \end{pmatrix}.$$

The characteristic polynomial of the 2×2 -matrix is $x^2 - \lambda_k^2 - \frac{1}{4}\tau^2$, so it is diagonalizable with eigenvalues $\pm\sqrt{\lambda_k^2 + \frac{1}{4}\tau^2}$. This gives rise to a diagonalization of $\overline{D}|_{E_k \oplus \alpha E_k}$ with the same eigenvalues, and we call the positive eigenspace F_k .

Now, we turn our attention to $\ker D$. As $L(e_0) = \tilde{R}(e_{n+1})\alpha$ anti-commutes with D , $L(e_0)$ operates on $\ker D$. This operation is self-adjoint and squares to $\mathbb{1}_{\ker D}$, so by the

spectral theorem $L(e_0)|_{\ker D}$ is diagonalizable and its eigenvalues must be contained in $\{1, -1\}$. Let K be the -1 -eigenspace. Then $\alpha(K)$ is the 1 -eigenspace. Due to

$$\overline{D}|_{\ker D} = -\frac{1}{2}\tau L(e_0)|_{\ker D},$$

K and $\alpha(K)$ become the $\frac{1}{2}\tau$ - and $-\frac{1}{2}\tau$ -eigenspaces of \overline{D} , respectively. \square

Remark 3.14. That \overline{D} is invertible for constant mean curvature $\tau \neq 0$, can also be seen directly from the fact that D anti-commutes with $L(e_0)$: As $L(e_0)^2 = \mathbb{1}$,

$$\overline{D}^2 = \left(D - \frac{1}{2}\tau L(e_0)\right)^2 = D^2 + \frac{1}{4}\tau^2 \mathbb{1}$$

and so $\text{coker } \overline{D} = \ker \overline{D} = 0$.

In the remainder of this section, we want to use the $Cl_{n,1}$ -linear Dirac-Witten operator to define an α -invariant for initial values. For this, let M be compact, spin and of dimension $n > 0$. We need no longer assume that it is embedded into a manifold N , as we succeeded in expressing all the relevant structures in terms of M and the pair (g, K) . In fact, the $Cl_{n,1}$ -linear hypersurface spinor bundle $\overline{\Sigma}_{Cl}(M, g) \cong \Sigma_{Cl}(M, g) \otimes_{Cl_n} Cl_{n,1}$ (cf. Remark 3.2) depends on the metric g alone, whereas its connection $\overline{\nabla}$ and thus its $Cl_{n,1}$ -linear Dirac-Witten operator \overline{D} is effected by K as well.

In analogy to the case of the classical α -invariant, we need to compare the spaces of L^2 -sections of the hypersurface spinor bundles for different initial value pairs (g, K) . Adopting the notation from Section 2.2, there is a bundle map

$$\sqrt{\beta}P^\nabla \otimes \mathbb{1}_{Cl_{n,1}} : \Sigma_{Cl}(M, g_0) \otimes_{Cl_n} Cl_{n,1} \rightarrow \Sigma_{Cl}(M, g) \otimes_{Cl_n} Cl_{n,1},$$

which induces

$$\overline{\Phi}_g : \overline{H} := L^2(M, \overline{\Sigma}_{Cl}(M, g_0)) \xrightarrow{\cong} L^2(M, \overline{\Sigma}_{Cl}(M, g)).$$

This allows to produce a continuous map from initial values to the space of Fredholm operators.

Theorem 3.15 (cf. [Glö19, Thm. 3.19]). *The map*

$$\begin{aligned} (\mathcal{I}(M), \mathcal{I}^+(M)) &\longrightarrow (\text{Fred}^{n,1}(\overline{H}), G^{n,1}(\overline{H})) \\ (g, K) &\longmapsto \overline{\Phi}_g^{-1} \circ \frac{\overline{D}_{(g,K)}}{\sqrt{1 + \overline{D}_{(g,K)}^2}} \circ \overline{\Phi}_g \end{aligned}$$

is well-defined and continuous with respect to the C^1 -topology on the space of smooth initial value pairs $\mathcal{I}(M)$. In particular, it is continuous if $\mathcal{I}(M)$ carries the C^∞ -topology.

Definition 3.16. The α -invariant for initial values is defined by the composition

$$\begin{aligned}\bar{\alpha}: \pi_k(\mathcal{I}^+(M), (g_0, K_0)) &\cong \pi_{k+1}(\mathcal{I}(M), \mathcal{I}^+(M), (g_0, K_0)) \\ &\rightarrow [(D^{k+1}, S^k), (\mathcal{I}(M), \mathcal{I}^+(M))] \\ &\rightarrow [(D^{k+1}, S^k), (\text{Fred}^{n,1}(\bar{H}), G^{n,1}(\bar{H}))] \cong KO^{n-k}(\{*\}).\end{aligned}$$

In the next chapter, $\bar{\alpha}$ will be compared to the classical α -invariant. The first step will be to establish a comparison map between the space of metrics of positive scalar curvature and the space of initial value pairs satisfying the dominant energy condition strictly.

4 Comparing the α -invariants

4.1 Positive scalar curvature and initial values

In the following, M is a compact smooth manifold of dimension $n \geq 2$. The aim of this section is to construct a continuous map $\Phi: \Sigma \mathcal{R}^+(M) \rightarrow \mathcal{I}^+(M)$ such that $\Phi|_{\mathcal{R}^+(M) \times \{0\}}$ is the inclusion $g \mapsto (g, 0)$. This comparison map will be used later to relate the α -invariants.

Lemma 4.1. *For every $C > 0$, the function*

$$\begin{aligned}\tau: \mathcal{R}(M) &\longrightarrow \mathbb{R} \\ g &\longmapsto \sqrt{\frac{n}{n-1} \max\{0, \sup_{x \in M} -\text{scal}^g(x)\}} + C\end{aligned}$$

is continuous.

Proof. It suffices to show that the assignment $g \mapsto \sup_{x \in M} \text{scal}^g(x)$ is continuous. This breaks into two pieces: Firstly, the function $C^0(M) \rightarrow \mathbb{R}$, $f \mapsto \sup_{x \in M} f(x)$ is (Lipschitz-)continuous, because for all $f, g \in C^0(M)$

$$\begin{aligned}\sup_{x \in M} f(x) - \sup_{x \in M} g(x) &= \sup_{x \in M} \left(f(x) - g(x) + g(x) - \sup_{y \in M} g(y) \right) \\ &\leq \sup_{x \in M} (f(x) - g(x)) \leq \|f - g\|_{C^0}.\end{aligned}$$

and likewise $\sup_{x \in N} g(x) - \sup_{x \in N} f(x) \leq \|f - g\|_{C^0}$. Secondly, continuity of $\mathcal{R}(M) \rightarrow C^0(M)$, $g \mapsto -\text{scal}^g$ follows from the fact that the scalar curvature can be expressed locally as a function of the coefficients of the metric and their first and second derivatives. \square

Proposition 4.2. *For any $C > 0$, the following is a well-defined continuous map of pairs:*

$$\begin{aligned} \phi: (\mathcal{R}(M), \mathcal{R}^+(M)) \times (I, \partial I) &\longrightarrow (\mathcal{I}(M), \mathcal{I}^+(M)) \\ (g, t) &\longmapsto \left(g, \frac{\tau(g)}{n}tg \right). \end{aligned}$$

Moreover, its homotopy class $[\phi] \in [(\mathcal{R}(M), \mathcal{R}^+(M)) \times (I, \partial I), (\mathcal{I}(M), \mathcal{I}^+(M))]$ is independent of $C > 0$.

Proof. Continuity directly follows from the lemma above. Moreover, varying the parameter $C > 0$ defines a continuous homotopy between different such maps. Thus, it only remains to prove that $\mathcal{R}(M) \times \partial I \cup \mathcal{R}^+(M) \times I$ is mapped into $\mathcal{I}^+(M)$. To this aim, we first observe that for a pair of the form $(g, \frac{\tau}{n}g)$ with $\tau \in \mathbb{R}$

$$\begin{aligned} 2\rho &= \text{scal} + \frac{n-1}{n}\tau^2 \\ j &= \frac{1-n}{n} \text{grad } \tau = 0 \end{aligned}$$

holds. Hence, such a pair fulfills the strict dominant energy condition if and only if

$$\tau^2 > -\frac{n}{n-1} \text{scal}.$$

But by definition of the function τ , this is the case for $(g, \pm \frac{\tau(g)}{n}g)$, which shows that $\mathcal{R}(M) \times \partial I$ maps into $\mathcal{I}^+(M)$. Moreover, the condition is automatically satisfied if g has positive scalar curvature, so $\mathcal{R}^+(M) \times I$ is sent to $\mathcal{I}^+(M)$ as well. \square

Proposition 4.3. *Let $C > 0$ and $h \in \mathcal{R}(M)$ a Riemannian metric. Then the composition*

$$\Phi: \Sigma \mathcal{R}^+(M) \longrightarrow \mathcal{R}(M) \times \partial I \cup \mathcal{R}^+(M) \times I \xrightarrow{\phi} \mathcal{I}^+(M),$$

where the first map is given by

$$[g, t] \longmapsto \begin{cases} ((-2t-1)h + 2(1+t)g, -1) & t \in [-1, -\frac{1}{2}] \\ (g, 2t) & t \in [-\frac{1}{2}, \frac{1}{2}] \\ ((2t-1)h + 2(1-t)g, 1) & t \in [\frac{1}{2}, 1], \end{cases}$$

is a well-defined, continuous map. Its homotopy class is independent of $C > 0$ and $h \in \mathcal{R}(M)$.

Proof. By the previous proposition, we just need to study the first map: Plugging in $t = \pm \frac{1}{2}$, we see that the different definitions agree on the intersections, and for the special

values $t = \pm 1$ we observe that the result is independent of g , i.e. the map descends to the suspension. This shows well-definedness. Continuity can now be checked on each domain of definition, where it is obvious. Moreover, this map continuously depends on $h \in \mathcal{R}(M)$, so by connectedness of $\mathcal{R}(M)$, its homotopy class is independent of h . \square

Corollary 4.4. *The inclusion $\mathcal{R}^+(M) \rightarrow \mathcal{I}^+(M)$, $g \mapsto (g, 0)$ is null-homotopic. In particular, if there exists a metric $g_0 \in \mathcal{R}^+(M)$, the induced map on homotopy groups $\pi_k(\mathcal{R}^+(M), g_0) \rightarrow \pi_k(\mathcal{I}^+(M), (g_0, 0))$ is the zero-map for all k .*

Proof. Using the map defined above, we get a factorization of the inclusion map as follows

$$\mathcal{R}^+(M) \hookrightarrow C\mathcal{R}^+(M) \hookrightarrow \Sigma\mathcal{R}^+(M) \xrightarrow{\Phi} \mathcal{I}^+(M),$$

where the first two maps are the canonical inclusions of a space into its cone and of the cone into the suspension as upper half. As cones are contractible, the composition is null-homotopic. \square

This shows that we cannot find non-trivial elements of homotopy groups in the space initial data with strict dominant energy condition by simply considering the space of positive scalar curvature metrics as subspace. However, the map Φ defined above allows for a better construction: In the remaining section, we will show that under certain conditions the composition

$$\pi_k(\mathcal{R}^+(M), g_0) \xrightarrow{\Sigma} \pi_{k+1}(\Sigma\mathcal{R}^+(M), [g_0, 0]) \xrightarrow{\Phi_*} \pi_{k+1}(\mathcal{I}^+(M), (g_0, 0))$$

has non-trivial image.

4.2 Main theorem

Let M be a compact spin manifold of dimension $n \geq 2$. The aim of this section is to relate the α -invariant for initial values $\bar{\alpha}: \pi_k(\mathcal{I}^+(M), (g_0, 0)) \rightarrow KO^{-n-k}(\{*\})$, where g_0 is a metric of positive scalar curvature, to the classical α -invariant using the map from Proposition 4.3. This will lead to a non-triviality result for $\pi_k(\mathcal{I}^+(M), (g_0, 0))$. Moreover, the same argument shows that the α -invariant for initial values detects that $\mathcal{I}^+(M)$ has least two connected components if $\alpha(M) \neq 0$.

Theorem 4.5 (Main Theorem). *1. If M carries a metric g_0 of positive scalar curvature, then for all $k \geq 0$, the diagram*

$$\begin{array}{ccccc} \pi_k(\mathcal{R}^+(M), g_0) & \xrightarrow{\Sigma} & \pi_{k+1}(\Sigma\mathcal{R}^+(M), [g_0, 0]) & \xrightarrow{\Phi_*} & \pi_{k+1}(\mathcal{I}^+(M), (g_0, 0)) \\ & \searrow \alpha & & \swarrow \bar{\alpha} & \\ & & KO^{-n-k-1}(\{*\}) & & \end{array}$$

commutes. Thereby, Σ is the suspension homomorphism and Φ is the map from Proposition 4.3.

2. For any metric g_0 ,

$$\bar{\alpha}: \pi_0 \left(\mathcal{I}^+(M), \left(g_0, \frac{\tau(g_0)}{n} g_0 \right) \right) \longrightarrow KO^{-n}(\{*\})$$

sends the class defined by $S^0 = \{\pm 1\} \ni t \mapsto \left(g_0, \frac{\tau(g_0)}{n} t g_0 \right)$ to $\alpha(M)$, where τ is defined as in Lemma 4.1.

Proof. For the first part, we start by exploring the effect of the upper composition. The claim is that

$$\begin{array}{ccc} \pi_k(\mathcal{R}^+(M), g_0) & \xrightarrow{\Sigma} & \pi_{k+1}(\Sigma \mathcal{R}^+(M), [g_0, 0]) & \xrightarrow{\Phi_*} & \pi_{k+1}(\mathcal{I}^+(M), (g_0, 0)) \\ \cong \uparrow & & & & \cong \uparrow \\ \pi_{k+1}(\mathcal{R}(M), \mathcal{R}^+(M), g_0) & \xrightarrow{\phi_*} & \pi_{k+2}(\mathcal{I}(M), \mathcal{I}^+(M), (g_0, 0)) & & \\ \downarrow & & \downarrow & & \\ [(D^{k+1}, S^k), (\mathcal{R}(M), \mathcal{R}^+(M))] & \xrightarrow{\phi_*} & [(D^{k+1}, S^k) \times (I, \partial I), (\mathcal{I}(M), \mathcal{I}^+(M))] & & \end{array} \quad (8)$$

commutes, where the middle and the lower map are both induced by

$$\begin{aligned} \phi: (\mathcal{R}(M), \mathcal{R}^+(M)) \times (I, \partial I) &\longrightarrow (\mathcal{I}(M), \mathcal{I}^+(M)) \\ (g, t) &\longmapsto \left(g, \frac{\tau(g)}{n} t g \right). \end{aligned}$$

Note that ϕ preserves the base point, if the base point of $(D^{k+1}, S^k) \times (I, \partial I)$ is chosen to be $(*, 0)$ when $*$ is the base point of S^k , so the middle map is well-defined. The lower square obviously commutes. For the upper square, we start with a class $[g] \in \pi_k(\mathcal{R}^+(M), g_0)$. Then the preimage under the boundary isomorphism is represented by

$$\begin{aligned} \tilde{g}: (D^{k+1}, S^k, *) &\longrightarrow (\mathcal{R}(M), \mathcal{R}^+(M), g_0) \\ rx &\longmapsto (1-r)g_0 + rg(x) \end{aligned}$$

for $r \in [0, 1]$ and $x \in S^k$. Applying the horizontal map and restricting to the boundary yields the class of

$$\begin{aligned} (\partial(D^{k+1} \times I), (*, 0)) &\longrightarrow (\mathcal{I}^+(M), (g_0, 0)) \\ (x, t) &\longmapsto \left(\tilde{g}(x), -\frac{\tau(\tilde{g}(x))}{n} t \tilde{g}(x) \right). \end{aligned}$$

Using the homeomorphism

$$(\Sigma S^k, [*], 0) \cong (\partial(D^{k+1} \times I), (*, 0))$$

$$[x, t] \mapsto \begin{cases} (2(1+t)x, -1) & t \in [-1, -\frac{1}{2}] \\ (x, 2t) & t \in [-\frac{1}{2}, \frac{1}{2}] \\ (2(1-t)x, 1) & t \in [\frac{1}{2}, 1], \end{cases}$$

this precisely gives the formula for $\Phi \circ \Sigma g$ (cf. Proposition 4.3).

The core of the proof is showing that the following diagram commutes:

$$\begin{array}{ccc} [(D^{k+1}, S^k), (\mathcal{R}(M), \mathcal{R}^+(M))] & \xrightarrow{\phi_*} & [(D^{k+1}, S^k) \times (I, \partial I), (\mathcal{I}(M), \mathcal{I}^+(M))] \\ \downarrow & & \downarrow \\ [(D^{k+1}, S^k), (\text{Fred}^{n,0}(H), G^{n,0}(H))] & & [(D^{k+1}, S^k) \times (I, \partial I), (\text{Fred}^{n,1}(\overline{H}), G^{n,1}(\overline{H}))] \\ & \searrow \cong & \nearrow \cong \\ & [(D^{k+1}, S^k), (\text{Fred}^{n+1,1}(\overline{H}), G^{n+1,1}(\overline{H}))]. & \end{array} \tag{9}$$

The first lower map is the one from Proposition 2.6, recalling that H and \overline{H} correspond to each other under the first Morita equivalence according to Proposition 3.4. The second lower map is Theorem 2.5, with $e = -e_{n+1}$.

Before doing so, let us show that

$$\begin{array}{ccccc}
& & & & [(D^{k+1}, S^k) \times (I, \partial I), (\text{Fred}^{n,1}(\overline{H}), G^{n,1}(\overline{H}))] \\
& & & \nearrow \cong & \downarrow \cong \\
& & [(D^{k+1}, S^k), (\text{Fred}^{n+1,1}(\overline{H}), G^{n+1,1}(\overline{H}))] & & \\
& \nearrow \cong & & \searrow \cong & \\
[(D^{k+1}, S^k), (\text{Fred}^{n,0}(H), G^{n,0}(H))] & & & & [(D^{k+1}, S^k) \times (I, \partial I), (\text{Fred}^{n,1}(\overline{H}), G^{n,1}(\overline{H}))] \\
\downarrow \cong & & & & \downarrow \cong \\
\text{ind} & & & & \text{ind} \\
& & [(D^{k+1}, S^k) \times (I, \partial I), (\text{Fred}^{n-1,0}(H), G^{n-1,0}(H))] & & \\
& \nearrow \cong & & \searrow \text{ind} & \\
& & & \cong & \\
KO^{-n}(D^{k+1}, S^k) & \xrightarrow{\cong} & KO^{-n+1}((D^{k+1}, S^k) \times (I, \partial I)) & & \\
& \searrow \cong & & \nearrow \cong & \\
& & KO^{-n-k-1}(\{*\}) & &
\end{array} \tag{10}$$

commutes, where the maps forming the central diamond are the Bott maps associated to $e = e_n$ along with the maps induced by the Morita equivalences, and the topmost right hand map is induced by a $Cl_{n,1}$ -Hilbert space isomorphism to be defined later. Notice that the right hand vertical composition is the index map, which follows from the invariance of the index map under $Cl_{n,1}$ -Hilbert space isomorphisms. So stitching the diagrams (8)-(10) together, we obtain the diagram from the first claim.

Moreover, setting $k = -1$, the commutative diagram composed of (9) and (10) implies the second assertion. Then $(D^{k+1}, S^k) = (\{*\}, \emptyset)$ and the upper left corner of the diagram is the one-point set $\{*\}, \mathcal{R}(M)$. Now the left hand vertical composition maps this point to the α -index of M , whereas the composition through the upper right corner is seen to map it to the Lorentzian α -invariant of the π_0 -class from the claim.

The lower half of (10) commutes by the definition of the index map, cf. (2) and (3). The middle diamond commutes as well, this is obvious from the way its constituting maps are defined. We are left with the upper triangle. Note first that we are dealing with two different $Cl_{n,1}$ -Hilbert space structures on \overline{H} : Since the map from the center upwards is the Bott map for $e = -e_{n+1}$, the $Cl_{n,1}$ -structure is the one obtained by forgetting the $\tilde{R}(e_{n+1})$ -action, whereas in the lower Hilbert space, we forget the multiplication by e_n .

These are connected by the $Cl_{n,1}$ -Hilbert space isomorphism

$$U: \overline{H} \longrightarrow \overline{H}$$

$$\phi \mapsto \frac{1}{\sqrt{2}} \tilde{R}(e_{n+1}) \tilde{R}(e_n + e_{n+1}).$$

Indeed, $\alpha \in B(\overline{H})$ corresponds via U to $\alpha = U\alpha U^{-1}$, $\tilde{R}(e_i)$ to $\tilde{R}(e_i)$ for $i < n$ and $\tilde{R}(e_n)$ to $\tilde{R}(e_{n+1})$. The right hand map in the triangle is defined to be the map induced by $\text{Fred}^{n,1}(\overline{H}) \ni F \mapsto U F U^{-1}$. As the analogous map on $\text{Fred}^{n+1,1}(\overline{H})$ is the identity, the diagram relating the Bott maps gets the shape of a triangle rather than a square. Its commutativity follows from

$$U \tilde{R}(-e_{n+1}) U^{-1} = \frac{1}{2} \tilde{R}(e_{n+1}) \tilde{R}(e_n + e_{n+1}) \tilde{R}(-e_{n+1}) \tilde{R}(e_n + e_{n+1}) \tilde{R}(e_{n+1})$$

$$= \frac{1}{2} (\tilde{R}(e_{n+1}) + \tilde{R}(e_n) + \tilde{R}(e_n) - \tilde{R}(e_{n+1})) = \tilde{R}(e_n).$$

It only remains prove that (9) commutes. The first two maps of the lower composition map $[g] \in [(D^{k+1}, S^k), (\mathcal{R}(M), \mathcal{R}^+(M))]$ to the class of

$$(D^{k+1}, S^k) \longrightarrow (\text{Fred}^{n+1,1}(\overline{H}), G^{n+1,1}(\overline{H}))$$

$$x \longmapsto \Phi_{g(x)}^{-1} \frac{D_{g(x)}}{\sqrt{1 + D_{g(x)}^2}} \Phi_{g(x)}.$$

This is because it restricts to the correct map on $H = \ker(\tilde{R}(e_0) \tilde{R}(e_{n+1}) - \mathbb{1}) \subseteq \overline{H}$, i.e. the Cl_n -Hilbert space associated to \overline{H} via the first Morita equivalence. The remaining map sends it to the class of

$$(D^{k+1}, S^k) \times (I, \partial I) \longrightarrow (\text{Fred}^{n,1}(\overline{H}), G^{n,1}(\overline{H}))$$

$$(x, t) \longmapsto \Phi_{g(x)}^{-1} \frac{D_{g(x)}}{\sqrt{1 + D_{g(x)}^2}} \Phi_{g(x)} - t \tilde{R}(e_{n+1}) \alpha$$

$$= \Phi_{g(x)}^{-1} \left(\frac{D_{g(x)}}{\sqrt{1 + D_{g(x)}^2}} - t L(e_0) \right) \Phi_{g(x)}.$$

In contrast, the result of the upper composition is represented by

$$(D^{k+1}, S^k) \times (I, \partial I) \longrightarrow (\text{Fred}^{n,1}(\overline{H}), G^{n,1}(\overline{H}))$$

$$(x, t) \longmapsto \Phi_{g(x)}^{-1} \frac{\overline{D}_{(g(x), K(x,t))}}{\sqrt{1 + \overline{D}_{(g(x), K(x,t))}^2}} \Phi_{g(x)}$$

$$\text{with } K(x, t) = \frac{\tau(g(x))}{n} t g(x).$$

Remembering that $\overline{D}_{(g,K)} = D_g - \frac{1}{2} \tau L(e_0)$, these do not look too much different, and we show that the following is a well-defined homotopy between them:

$$(D^{k+1}, S^k) \times (I, \partial I) \times [0, 1] \rightarrow (\text{Fred}^{n,1}(\overline{H}), G^{n,1}(\overline{H}))$$

$$(x, t, s) \mapsto \Phi_{g(x)}^{-1} \left(a_{(x,t,s)}(D_{g(x)}) D_{g(x)} - b_{(x,t,s)}(D_{g(x)}) t L(e_0) \right) \Phi_{g(x)}$$

for

$$a_{(x,t,s)}(\lambda) = \frac{s}{\sqrt{1+\lambda^2}} + \frac{1-s}{\sqrt{1+\lambda^2 + \frac{1}{4}t^2\tau(g(x))}}$$

$$b_{(x,t,s)}(\lambda) = s + \frac{(1-s)\frac{1}{2}\tau(g(x))}{\sqrt{1+\lambda^2 + \frac{1}{4}t^2\tau(g(x))}}.$$

As this operator family is obtained by linearly interpolating between two continuous operator families, it is again continuous. So it remains to see that its target is indeed $(\text{Fred}^{n,1}(\overline{H}), G^{n,1}(\overline{H}))$. It is clear, that all the operators are bounded, self-adjoint, odd and $Cl_{n,1}$ -linear. To show that the operator $F_{(x,t,s)}$ associated to (x, t, s) is Fredholm, we use the spectral decomposition of $D_{g(x)}$ from Proposition 3.13: The restriction of $F_{(x,t,s)}$ to $E_k \oplus \alpha(E_k) \cong E_k \otimes \mathbb{R}^2$ is given by

$$\mathbb{1}_{E_k} \otimes \begin{pmatrix} a_{(x,t,s)}(\lambda_k)\lambda_k & -b_{(x,t,s)}(\lambda_k)t \\ -b_{(x,t,s)}(\lambda_k)t & -a_{(x,t,s)}(\lambda_k)\lambda_k \end{pmatrix}.$$

This is diagonalizable with eigenvalues $\pm\sqrt{a_{(x,t,s)}(\lambda_k)^2\lambda_k^2 + b_{(x,t,s)}(\lambda_k)^2t^2}$. Note that due to $\sqrt{a_{(x,t,s)}(\lambda_k)^2\lambda_k^2 + b_{(x,t,s)}(\lambda_k)^2t^2} \geq a_{(x,t,s)}(\lambda_k)|\lambda_k|$, their absolute values, for any $t \in I$ and $s \in [0, 1]$, are bounded away from zero by

$$\frac{\lambda_0}{\sqrt{1+\lambda_0^2 + \frac{1}{4}\tau(g(x))}} > 0,$$

where $\lambda_0 > 0$ denotes the smallest positive eigenvalue of $D_{g(x)}$. A similar consideration as in Proposition 3.13 shows that $F_{(x,t,s)}$ restricted to $\ker(D_{g(x)})$ is diagonalizable as well, with eigenvalues $\pm b_{(x,t,s)}(0)t$. Putting this together, we find that $F_{(x,t,s)}$ has finite dimensional kernel, co-kernel and closed image (for this, the boundedness away from zero is needed). Furthermore, $F_{(x,t,s)}$ is invertible if $D_{g(x)}$ is invertible or $t > 0$, one of which is the case on $\partial(D^{k+1} \times I)$.

In the case $n-1 \equiv -1 \pmod{4}$ one more tiny bit of thought is necessary. The space self-adjoint $Cl_{n,1}$ -linear Fredholm operators has three components (cf. [AS69]): Those F for which $\omega_{n,1}F\iota$ is essentially positive, those for which it is essentially negative and the rest. As for $s=0$ (or $s=1$) all operators $F_{(x,t,s)}$ fall into the last category, the same has to be true for all $s \in [0, 1]$ by continuity. \square

Together with the non-triviality results for the classical α -invariant from Theorems 2.11 and 2.12, we obtain the following conclusions:

Corollary 4.6. *If M is a closed spin manifold of dimension $n \geq 6$ that carries a metric g_0 of positive scalar curvature, then for all $k \geq 1$ with $k+n \equiv 1, 2 \pmod{8}$ the α -invariant for initial values $\bar{\alpha}: \pi_k(\mathcal{I}^+(M), (g_0, 0)) \rightarrow KO^{-n-k}(\{*\}) \cong \mathbb{Z}/2\mathbb{Z}$ is split surjective.*

Corollary 4.7. *If M is a closed spin manifold of dimension $n \geq 6$ that carries a metric g_0 of positive scalar curvature, then for all $k \geq 1$ with $k + n \equiv 1, 2 \pmod{8}$ the α -invariant for initial values $\bar{\alpha}: \pi_k(\mathcal{I}^+(M), (g_0, 0)) \rightarrow KO^{-n-k}(\{*\}) \cong \mathbb{Z}/2\mathbb{Z}$ is surjective and for all $k \geq 1$ with $k + n \equiv 0, 4 \pmod{8}$ the localized α -invariant for initial values $\bar{\alpha} \otimes \mathbb{1}_{\mathbb{Q}}: \pi_k(\mathcal{I}^+(M), (g_0, 0)) \otimes \mathbb{Q} \rightarrow KO^{-n-k}(\{*\}) \otimes \mathbb{Q} \cong \mathbb{Q}$ is surjective.*

In particular, under the assumptions of the corollaries above, $\pi_k(\mathcal{I}^+(M), (g_0, 0)) \neq 0$, which shows Corollary 1.1. Note that the main theorem provides an explicit construction of the non-trivial elements, provided that in $\pi_{k-1}(\mathcal{R}^+(M), g_0)$ the non-trivial elements detected by the α -invariant are known.

References

- [AS69] Michael F. Atiyah and Isadore M. Singer. “Index theory for skew-adjoint Fredholm operators”. In: *Publications Mathématiques de l’IHÉS* 37 (1969), pp. 5–26.
- [Bau14] Helga Baum. *Eichfeldtheorie*. Jan. 2014. DOI: 10.1007/978-3-642-38539-1.
- [BER14] Boris Botvinnik, Johannes Ebert, and Oscar Randal-Williams. “Infinite loop spaces and positive scalar curvature”. In: *Inventiones mathematicae* (Nov. 2014). DOI: 10.1007/s00222-017-0719-3.
- [BGM05] Christian Bär, Paul Gauduchon, and Andrei Moroianu. “Generalized cylinders in semi-Riemannian and spin geometry”. In: *Mathematische Zeitschrift* 249.3 (Mar. 2005), pp. 545–580. DOI: 10.1007/s00209-004-0718-0.
- [BI04] Robert Bartnik and Jim Isenberg. “The constraint equations”. In: *The Einstein equations and the large scale behavior of gravitational fields*. Birkhäuser, Basel, 2004, pp. 1–38. URL: <https://arxiv.org/abs/gr-qc/0405092>.
- [CSS18] Diarmuid Crowley, Thomas Schick, and Wolfgang Steimle. “Harmonic spinors and metrics of positive curvature via the Gromoll filtration and Toda brackets”. In: *Journal of Topology* 11.4 (2018), pp. 1077–1099. DOI: 10.1112/topo.12081.
- [Ebe13] Johannes Ebert. “The two definitions of the index difference”. In: *Transactions of the American Mathematical Society* 369 (Aug. 2013). DOI: 10.1090/tran/7133.
- [Glö19] Jonathan Glöckle. “Initial Value Spaces in General Relativity”. Master Thesis. Universität Regensburg, Jan. 2019.
- [Hit74] Nigel Hitchin. “Harmonic Spinors”. In: *Advances in Mathematics* 14.1 (1974), pp. 1–55. ISSN: 0001-8708. DOI: 10.1016/0001-8708(74)90021-8.
- [HSS14] Bernhard Hanke, Thomas Schick, and Wolfgang Steimle. “The space of metrics of positive scalar curvature”. In: *Publications mathématiques de l’IHÉS* 120.1 (Nov. 2014), pp. 335–367. ISSN: 1618-1913. DOI: 10.1007/s10240-014-0062-9.

- [HZ03] Oussama Hijazi and Xiao Zhang. “The Dirac-Witten Operator on Spacelike Hypersurfaces”. In: *Communications in Analysis and Geometry* 11 (Sept. 2003). DOI: 10.4310/CAG.2003.v11.n4.a5.
- [PT82] Thomas Parker and Clifford H. Taubes. “On Witten’s proof of the positive energy theorem”. In: *Communications in Mathematical Physics* 84.2 (June 1982), pp. 223–238. ISSN: 1432-0916. DOI: 10.1007/BF01208569.
- [Roe99] John Roe. *Elliptic Operators, Topology, and Asymptotic Methods*. 2nd ed. Chapman and Hall/CRC, New York, 1999. ISBN: 0582325021.
- [SY81] Richard Schoen and Shing-Tung Yau. “Proof of the positive mass theorem. II”. In: *Communications in Mathematical Physics* 79.2 (Mar. 1981), pp. 231–260. ISSN: 1432-0916. DOI: 10.1007/BF01942062.
- [Wit81] Edward Witten. “A new proof of the positive energy theorem”. In: *Communications in Mathematical Physics* 80.3 (1981), pp. 381–402. URL: <https://projecteuclid.org:4>