

Torsion braneworlds in a tensor-vector gravity

J. E. G. Silva,^{a,d} L. J. S. Sousa,^b W. T. Cruz,^c C. A. S. Almeida^{d,1}

^a*Universidade Federal do Cariri (UFCA) - Centro de Ciências e Tecnologia.
Av. Tenente Raimundo Rocha S/N, Cidade universitária
63048-080 Juazeiro do Norte - Ceará - Brazil*

^b*Instituto Federal de Educação Ciência e Tecnologia do Ceará
(IFCE) - Campus de Maracanaú Av. Contorno Norte, 10 - Distrito Industrial, Maracanaú,
61936-000 - Ceará - Brazil*

^c*Instituto Federal de Educação, Ciência e Tecnologia do Ceará (IFCE), Campus Juazeiro do Norte
- 63040-540 Juazeiro do Norte, CE, Brazil*

^d*Departamento de Física - Universidade Federal do Ceará - UFC
C.P. 6030, 60455-760 Fortaleza - Ceará - Brazil*

*E-mail: euclides.silva@ufca.edu.br, luisjose@fisica.ufc.br,
wilami@ifce.edu.br, carlos@fisica.ufc.br*

ABSTRACT: We study the properties of gravity and bulk fields living in a torsion warped braneworld. The torsion is driven by a background vector whose norm provides a source for the bulk cosmological constant. For a vector as the derivative of a scalar field, we find new isotropic and anisotropic thick brane geometries. We analyse the features of bosonic and fermionic fields in this isotropic and in a standing wave scenarios. The background vector provides nonminimal coupling between the field and the geometry leading to modifications in the Kaluza-Klein states. The spinor connection is modified by the torsion and a derivative Yukawa-like coupling is proposed. The effects of these new couplings are investigated.

¹Corresponding author.

Contents

1	Introduction	1
2	Torsion in Lyra geometry	3
3	Warped braneworld solutions	5
3.1	Isotropic brane	5
3.2	Anisotropic standing wave branes	8
4	Bulk fields	10
4.1	Scalar field	10
4.2	Gauge field	13
4.3	Fermion field	14
5	Final remarks and perspectives	17

1 Introduction

In braneworld paradigm our $4D$ world is view as a brane embedded in a higher dimensional bulk spacetime. These models provide geometrical solutions for some of greatest problems in physics, such as the the hierarchy problem [1], the origin of dark matter [2], the cosmological constant problem [3] and the Big Bang singularity [4].

The main idea here is to define the gravity and field dynamics in the bulk and obtain the effective dynamics on the brane by dimensional reduction. Several mechanisms were proposed for bosonic, fermionic and gravity fields [5–11]. The dimensional reduction yields to the standard model (SM) action on the brane and a tower of massive Kaluza-Klein (KK) states. The KK spectrum signals for a physics beyond the standard model [12].

The extra dimensions also allow us to propose non-standard bulk dynamics and seek for their phenomenological effects on the brane. Notably, models involving supersymmetry [13], Lorentz violation [14], noncommutative geometry [15], orbifold internal space [16], folded branes [17] and minetic gravity [18] were investigated.

An important geometric extension in Braneworld models is achieved considering a Weyl geometry [19–21]. In Weyl geometry, the covariant derivative of the metric is proportional to a vector field, altering the metric compatibility. This background vector breaks the spacetime integrability which can be restored provided the Weyl vector be the gradient of a scalar function. For a particular choice of the Weyl scalar potential, thick brane solutions were found in this vacuum Weyl spacetime [19–22]. This integrable Weyl braneworld share many properties with the Einstein-dilaton brane models [23]. Moreover, Weyl geometries are vacuum sources for bouncing cosmologies [24].

Another interesting braneworld extension is due to the inclusion of a torsional bulk. Torsion breaks the symmetry of the connection while can preserves the metric compatibility [25, 26]. In string theory, notably the heterotic theory, the skew-symmetric Kalb-Ramond gauge field can be regarded as a torsion source [28–31]. Thus, the open string spectrum provides bulk extensions of Riemannian geometry. Torsion models can also be found in M-theory context [32]. Among the torsion effects on the brane we outline the modified black hole horizon [33], a correction to the brane cosmological constant [34], a topological mass for the gauge field [35], *CPT* violation [36] and the split of the brane in $F(T)$ teleparallel gravity [37]. Furthermore, the coupling of torsion with standard model fields can yields to Lorentz symmetry breaking [38?].

In this article we propose a torsional braneworld whose torsion is driven by a background vector, in the so called Lyra geometry [40, 41]. The asymmetric Lyra connection can be recast into a symmetric (compactible) connection by including the torsion terms in the connection coefficients. Likewise the Weyl geometry, the resulting gravitational action can be regarded as a tensor-vector theory [42]. The torsion in Lyra geometry provides interesting isotropic and anisotropic cosmological solutions [43, 44], as well as modified Kaluza-Klein models [45, 46].

We construct isotropic and anisotropic branes in $5D$ as vacuum solutions of tensor-vector dynamics. We show that the Lyra background field can be a source for the bulk cosmological constant. By assuming the background vector as the derivative of a scalar field and an independent bulk cosmological constant, pure geometric thick brane solutions are found. We also obtain an anisotropic standing wave braneworld [52–54] solution in this pure geometric torsion geometry.

The torsion effects on the bulk bosonic and fermionic fields are also investigated. The presence of torsion enables non-minimal couplings providing dissipative and massive terms. Unlike in usual Einstein-Cartan theories, where the torsion breaks the $U(1)$ gauge symmetry, by using the associated symmetric connection, the gauge symmetry is preserved and a scalar coupling similar to a derivative dilaton-Maxwell term is allowed. For massless fermion, besides the modification in the spinor connection, torsion allows a geometric Yukawa-like interaction relating torsion, metric and the fermion. These derivative terms strong modifies the Kaluza-Klein (KK) modes nearby the brane.

This work is organized as follows: in section (2) we review the main properties of torsion in Lyra geometry, discussing the gravitational action and the deriving the modified Einstein equation. In section (4) thin and thick braneworld solutions are obtained with a background torsion and their geometric features studied. In section (4.3), the bulk fields dynamics is analysed in both isotropic and anisotropic torsion solutions. Finally, section (5) is devoted to additional comments and possible future investigations.

Throughout this text we adopt the mostly minus metric signature $(+, -, \dots, -)$. Further, we use capital latin for the bulk space-time indexes $A = 1, \dots, D$, and greek indexes for the brane $\mu = 0, \dots, 3$.

2 Torsion in Lyra geometry

In this section we review the definitions and the main properties of the torsional Lyra geometry.

Firstly, let us introduce the notion of torsion in spacetime. In a parallel transport from the point x^A to $x^A + dx^A$, a vector v^M changes as $dv^M = -\Gamma_{AB}^M v^A dx^B$. The connection coefficients induce a covariant derivative in the form $\nabla_A v^M = \partial_A v^M + \Gamma_{AB}^M v^B$. The antisymmetric part of the connection coefficients yields the torsion tensor $T_{AB}^M := \Gamma_{AB}^M - \Gamma_{BA}^M$.

Now, we pointed out that, we can introduce the Levi-Civita connection given by the Christoffel symbols of the second kind $\{_{AB}^M\} := \frac{g^{MN}}{2}(\partial_A g_{NB} + \partial_B g_{NA} - \partial_N g_{AB})$ [25], which is a torsion free connection $T_{AB}^M = 0$ satisfying the metric compatibility condition $\nabla_A g_{BC} = 0$.

In the bosonic spectrum of string theories the antisymmetric Kalb-Ramond gauge field $B_2 = \frac{1}{2}B_{MN}dx^M \wedge dx^N$ has the field strength $H_3 = dB_2$ [28]. The Kalb-Ramond field strength provides a source for the spacetime torsion in the form $T_{AB}^M = H_{AB}^M$ [29].

The possible torsion geometries can be classified by means of the components of the torsion tensor. A general torsion tensor can be decomposed as [26]

$$T_{MNP} = \frac{1}{(D-1)}(g_{MP}T_N - g_{MN}T_P) - \frac{1}{(D-3)2!}\epsilon_{MNPQ_1\dots Q_{D-3}}S^{Q_1\dots Q_{D-3}} + q_{MNP}, \quad (2.1)$$

where $T_A := T_{AB}^B$ is the trace vector, $S^{Q_1\dots Q_{D-3}} := \epsilon^{MNPQ_1\dots Q_{D-3}}T_{MNP}$ is the torsion axial tensor, q_{MNP} is the null trace tensor, and D is the space-time dimension [26, 27].

A special example of torsion space-time is given in the so-called Lyra geometry, where the torsion tensor T_{AB}^M is generated by a vector field $b_1 = b_M dx^M$ in the form [42]

$$T_{AB}^M = \tilde{\Gamma}_{AB}^M - \tilde{\Gamma}_{BA}^M = -\frac{1}{2}(\delta_A^M b_B - \delta_B^M b_A), \quad (2.2)$$

whereas a vector V^M is modified by a parallel transport as

$$\delta V^M = -\Omega(x)\tilde{\Gamma}_{AB}^M V^A dx^B. \quad (2.3)$$

Therefore, the Lyra torsion is a pure trace vector torsion geometry with $T_M = \frac{(D-1)}{2}b_M$, where the torsion vector b_M is called the displacement vector [42].

Besides torsion, Lyra geometry has a rich notion of scale invariance gauge. In fact, in addition to a coordinate system x^M , a reference frame is endowed with a scale function φ , i.e., (φ, x^M) , such that a vector V^M transforms as $V'^M = \Omega \frac{\partial x'^M}{\partial x^N} V^N$ and $\Omega := \frac{\varphi(x')}{\varphi(x)}$ [41]. The scale function Ω is an extra geometric gauge freedom which induces a conformal transformation in the form $ds^2 = \Omega^2(x)g_{MN}dx^M dx^N$ [42]. The tensor vector b_M changes under a scale transformation as $b'_M = \Omega^{-1}(b_M + \varphi^{-1}\partial_M(\log \Omega^2))$ [41].

Assuming the Lyra connection is compactible with the metric $\tilde{\nabla}_A g_{MN} = 0$, we can obtain an associated torsion-free Lyra connection ∇_A^L whose components are given by [42],

$$\Gamma_{AM}^M = \Omega^{-1}\{_{AM}^M\} + \frac{1}{2}(\delta_A^M b_B + \delta_B^M b_A - g_{AB}b^M). \quad (2.4)$$

Note that for the gauge choice $\Omega = 1$ the torsion-free Lyra connection is similar to the Weyl connection [41].

The Lyra connection modifies the Ricci tensor in the form $R_{MN}^L = R_{(MN)}^L + R_{[MN]}^L$, where the symmetric part is given by [41]

$$R_{(MN)}^L = \Omega^{-2} R_{MN} + \frac{\Omega^{-1}}{2} (2\nabla_{(M} b_{N)} + \nabla_C b^C g_{MN}) - \frac{1}{2} (b_M b_N - b_C b^C g_{MN}), \quad (2.5)$$

where R_{MN} is the Ricci tensor obtained from the Levi-Civita connection and the skew-symmetric part has the form [41]

$$R_{[MN]}^L = \Omega^{-1} f_{MN}, \quad (2.6)$$

where $f_{MN} := \partial_M b_N - \partial_N b_M$ [41]. The appearance of a skew-symmetric term in the Ricci tensor is an ubiquitous feature of torsion theories and in Einstein-Cartan models is related to the spin interaction [25]. In Lyra torsion model the $R_{[MN]}^L$ term seems linked to the dynamics of the torsion vector b_M .

The Ricci scalar curvature derived from the Lyra connection $R^L := g^{MN} R_{MN}^L$ for $\Omega = 1$ can be written as [42],

$$R^L = R + \frac{1}{4} (D-2)(D-1) b^A b_A + (D-1) \nabla^A b_A \quad (2.7)$$

where D is space-time dimension, R is the Ricci scalar build from the torsion-free Levi-Civita connection and the capital letters represent the indexes in bulk. Likewise Weyl geometry, the torsion tensor can be regarded as an external field modifying the space-time curvature. Notwithstanding, note that the skew-symmetric component of the Lyra-Ricci tensor vanishes when contracted with the metric.

The presence of space-time torsion enlarges the possible gravitational dynamics, by allowing new interactions and possibly considering both the metric and connection as independent variables [25]. Nonetheless, in the present work we consider the torsion vector as a background field and assume a metric theory described by an extension of the Einstein-Hilbert gravitation action in the form

$$S_g = \frac{1}{\kappa_g} \int \Omega^D R^L \sqrt{|g|} dx^1 \dots dx^D, \quad (2.8)$$

which yields to the modified Einstein equation

$$R_{MN} - \frac{1}{2} g_{MN} R = -\frac{\Omega^2}{4} (D-2)(D-1) (b_M b_N - \frac{1}{2} g_{MN} b_C b^C). \quad (2.9)$$

The Einstein equation (2.9) may be interpreted as a tensor-vector-scalar theory of gravity or only a tensor-scalar theory by assuming $b_M = \partial_M h$. This theory is similar to the Brans-Dicke gravity in some special regimes [42].

It is worthwhile to mention that a kinetic term for the torsion vector can be obtained as

$$S_b = \alpha \int \Omega^D R_{[MN]} R^{[MN]} \sqrt{|g|} dx^1 \dots dx^D, \quad (2.10)$$

which yields to a Lagrangian similar to the dilaton-Maxwell term $\alpha\Omega^{D-2}f_{MN}f^{MN}$. Accordingly, considering only linear terms for the curvatures, i.e., without high-order terms in the gravitational Lagrangian, yields to background torsion vector. Further, for $b_M = \partial_M h$, i.e., in the tensor-scalar regime, the term $R_{[MN]}R^{[MN]}$ vanishes identically.

The effects of the torsion in Lyra geometry were investigated in black holes modified solutions [42], cosmology [43, 44] and Kaluza-Klein (KK) models [45, 46]. In next section we explore the torsion effects in warped braneworld models assuming the torsion as a source for the thick brane configurations. Furthermore, we point out a possible relation between the Lyra torsion and the Kalb-Ramond field in 5D.

3 Warped braneworld solutions

In this section we study some brane geometries which are solutions of the modified Einstein equation (2.9).

For a covariant constant displacement vector $\nabla_M b^M = 0$ the torsion provides the bulk cosmological constant. In fact, the norm $b_M b^M$ is a global constant throughout the bulk and hence we can define the bulk cosmological constant Λ as

$$\Lambda = \frac{(D-1)(D-2)}{8} b_M b^M. \quad (3.1)$$

Thus, for a AdS_D bulk, b_M ought to be a spacelike vector.

Despite being considered here as a background field, the dynamics for the torsion vector b_M might be provided from the Kalb-Ramond field. A noteworthy fact is that the Kalb-Ramond is equivalent to a vector field in $5D$. In fact, $\star H_3$ is a 2-form which can be written as $\star H_3 = \frac{f_{MN}}{2} dx^M \wedge dx^N$, where $f_2 = d\tilde{A}_1$ is a pseudo electromagnetic field. Thus, the Kalb-Ramond kinetic term $S_B = \int \star H_3 \wedge H_3$ is equivalent to a dual pseudo electromagnetic term $S_{\tilde{A}} = \int \star f_2 \wedge f_2$. The Kalb-Ramond free field equation of motion $d(\star H_3) = 0$ and $dH_3 = 0$ are ready satisfied, and the field strength can be written as

$$H_{MNP} = \epsilon_{MNP}^{ST} \partial_S \tilde{A}_T. \quad (3.2)$$

From the Kalb-Ramond field strength we can obtain the pseudo electric and magnetic fields $\tilde{E}_y = \epsilon_{0y}^{ijk} H_{ijk}$ and $\tilde{B}_i = H_{0iy}$ and write the Lyra vector as $b^M = (0, \vec{\tilde{B}}, \tilde{E}^y)$. For a pure-gauge configuration $\tilde{A}_1 = d\phi$ the kinetic term vanishes resulting in a torsion-free gravitational theory. By including the dilaton interactions, this torsion free geometry exhibits conformal symmetry and provides a source for a Weyl geometry [27]. Other possible configurations can arise in vector and tensor spontaneous Lorentz violating theories [38]. Therefore, we speculate that five dimensional space-times are interesting scenarios to seek for torsion Lyra geometry with a Kalb-Ramond source. Henceforward we concern ourselves to $D = 5$ models.

3.1 Isotropic brane

Consider a warped brane metric in the form

$$ds_5^2 = e^{-2A(y)} \eta_{\mu\nu} dx^\mu dx^\nu - dy^2. \quad (3.3)$$

For $\Lambda = 3b^2/2 < 0$ let us consider $b_M = (0, 0, 0, 0, b_4)$. The vacuum Einstein equation in the extra dimension yields to $A(y) = \pm \left(\frac{b_4}{4}y\right) + A_0$, providing a thin 3-brane immersed in the AdS_5 RS model [1].

For a spacelike vector $b_A = (0, b_1, b_2, b_3, b_4)$, the contravariant components are $b^i = e^{2A}b_i$. Therefore, the background vector b^i is exponentially suppressed on the 3-brane at the origin. Since the background vector can couple with the standard model fields, as we will discuss in next section, the suppression of this vector might explain the small departure of Lorentz symmetry on the brane [38, 39].

An interesting interpretation of the torsion in Lyra spacetimes is achieved for $b_M = \partial_M h(x)$ from which the Einstein equation with a bulk cosmological constant reads

$$G_{MN} = -3\tilde{T}_{MN} + \Lambda g_{MN}, \quad (3.4)$$

where $\tilde{T}_{MN} = (\nabla_M h \nabla_M h - \frac{1}{2}g_{MN} \nabla^C h \nabla_C h)$ is analogous to the stress energy tensor of a massless real scalar field $h(x)$. Assuming that the Lyra scalar field $h(x)$ does not propagate on the brane, the free field equation of motion $g^{MN} \nabla_M \nabla_N h = 0$ yields to $-h''(y) + 4A'(y)h'(y) = 0$ whose solution is $h = \int e^{(4A+c_1)} dy + c_2$.

The Einstein equation in the extra dimension yields to

$$6A'^2 = -\frac{3}{2}h'^2 - \Lambda, \quad (3.5)$$

thus, the torsion provides a negative term in Eq.(3.5), which possesses solution provided $h' \leq (2\Lambda/3)^{1/2}$.

For the free Lyra scalar $A = \ln \left[\frac{-2\Lambda}{h_0^2} \operatorname{sech}^2 \left(2\sqrt{\frac{-2\Lambda}{3}}y \right) \right]^{\frac{1}{8}}$, and the Lyra scalar field has the form $h = tg^{-1} \left(\tanh \left[2\sqrt{\frac{-2\Lambda}{3}}y + c_2 \right] \right)$. Therefore, a free Lyra scalar field exhibits a kink-like profile with a localized energy density around the brane at $y = 0$ and a nonvanishing value asymptotically. In a similar form, accelerated cosmological solutions in 4D were found in bounce models generated by a Weyl scalar [24]. This free field configuration provides a divergent warp factor and thus, an additional mechanism is required in order to trap gravity. Note, however, that a constant derivative $h' = h_0$ satisfies the free field equation with a constant energy density and negative pressure. Thus, the Lyra scalar provides a positive cosmological constant term which opposes to the bulk cosmological constant.

The profile of the free Lyra scalar motivated us to seek for a background Lyra scalar localized near the origin in the form

$$h(y) = \frac{1}{\lambda} \sqrt{\frac{-2\Lambda}{3}} tg^{-1} (\tanh \lambda y). \quad (3.6)$$

Using this *ansatz*, the Einstein equations (3.5) provides the warp factor

$$A(y) = \frac{1}{\lambda} \sqrt{\frac{-\Lambda}{6}} \log \cosh \lambda y, \quad (3.7)$$

where λ measures the width of the Lyra scalar non-constant region. In figure (1) we compare the RS Z_2 -orbifolded solution $A(y) = k|y|$ (thin line) to the smooth warp factor in Eq.(3.7).




Figure 1. Warp function for $\lambda = 1$ (dashed line) and for the RS model (thin line).




Figure 2. Warp factor for $\lambda = 1$ (dashed line) and for the RS model (thin line).

Thus, remarkably, this pure-geometric braneworld scenarios is equivalent to the thick brane models.

The resulting geometry is a family of asymptotically AdS_5 spacetimes with warped metrics parametrized by the ratio between Λ and λ in the form

$$ds^2 = \text{sech}^2 \sqrt{-\Lambda/6\lambda^2} (\lambda y) \eta_{\mu\nu} dx^\mu dx^\nu - dy^2. \quad (3.8)$$

This braneworld solution, including the scalar field pattern, is rather similar to one found in teleparallel $f(T)$ gravity [37]. The profile of the energy density and the scalar curvature are shown, respectively in Figures 3 and 4. The torsion term in Einstein equation provides a negative pressure that smooths the brane.




Figure 3. Energy density for $\lambda = 1$ (dashed line) and $\lambda = 2$ (thin line) using $\Lambda = -\lambda^2$.




Figure 4. Ricci scalar for $\lambda = 1$ (dashed line) and $\lambda = 2$ (thin line) using $\Lambda = -\lambda^2$.

Considering $b_M = (0, 0, 0, 0, E_y)$, the Lyra scalar h behaves as a static potential function $E_y = \partial_y h$ whose profile is shown in figure (6). Thus, the torsion interpolates between these two asymptotic values and it is concentrated around the brane. On the other hand, although being considered here as a background field, a noteworthy property is that the Lyra scalar

field in Eq.(3.6) could also be regarded as a solution for the EOM $g^{MN}\nabla_M h \nabla_N h = V'(h)$, where $V(h)$ is a exotic modified sine-Gordon potential of form

$$V(h) = \frac{-2\Lambda}{\kappa_5 \lambda} (\lambda + 4\sqrt{-\Lambda}) \left\{ \sqrt{\cos(2ch)} - \sqrt{2} \cos(ch) \log[\sqrt{2} \cos(ch) + \sqrt{\cos(2ch)}] \right\} \sec(2ch) \sqrt{1 - \tan(ch)^2}, \quad (3.9)$$

and $c = \sqrt{\frac{-\lambda^2 \kappa_5}{2\Lambda}}$. In figure (5) we plotted this dual potential comparing to the sine-Gordon potential (dashed line). The modified potential has a compact support only near the critical points of the sine-Gordon potential.




Figure 5. Lyra scalar potential for $c = 1$ (thin), $c = 2$ (thick) and the sine-gordon potential (dashed).




Figure 6. Lyra scalar for $\lambda = 1$ (dashed line) and $\lambda = 2$ (thin line).

3.2 Anisotropic standing wave branes

We consider a warped anisotropic metric ansatz in the form

$$ds^2 = e^{A(y)} \left(dt^2 - e^{u(y,t)} dx^2 - e^{u(y,t)} d\tilde{y}^2 - e^{-2u(y,t)} dz^2 \right) - dy^2, \quad (3.10)$$

where $A(y)$ and $u = u(y, t)$. The Einstein equations yields to the system of differential equations

$$-\frac{1}{2}e^{A+u} \left(2A'^2 + A'' - e^{-A}\ddot{u} + 2A'u' + u'' \right) = -\frac{2}{3}e^{A+u}\Lambda, \quad (3.11)$$

$$-\frac{1}{2}e^{A-2u} \left(2A'^2 + A'' + 2e^{-A}\ddot{u} - 4A'u' - 2u'' \right) = -\frac{2}{3}e^{A-2u}\Lambda, \quad (3.12)$$

$$\frac{1}{2}e^A \left(2A'^2 + A'' - 3e^{-A}\dot{u}^2 \right) = -\kappa_5^2 \dot{h}^2 + \frac{2}{3}e^A \Lambda, \quad (3.13)$$

$$-\frac{3}{2}\dot{u}u' = -\kappa_5^2 \dot{h}h', \quad (3.14)$$

$$\frac{1}{2} \left(-2A'^2 - 4A'' - 3u'^2 \right) = -\kappa_5^2 h'^2 - \frac{2}{3}\Lambda. \quad (3.15)$$




Figure 7. Profile of the Lyra scalar for a flat Lorentzian brane (thin line) and for an anisotropic brane (thick line).

Note that Eq. (3.14) leads to $\dot{h} = \sqrt{\frac{3}{2\kappa_5^2}}\dot{u}$ and $h' = \sqrt{\frac{3}{2\kappa_5^2}}u'$. For $A''(r) = 0$ and $A'^2(y) = \frac{2}{3}\Lambda$, where $\Lambda > 0$, the $u(y, t)$ satisfies

$$-e^{-A}\ddot{u} + 2A'u' + u'' = 0. \quad (3.16)$$

Choosing a standing wave *ansatz* $u(y, t) = Q(y) \sin(\omega t)$ gives

$$Q''(y) + 2A'(y)Q'(y) + \omega^2 e^{-A(y)}Q(y) = 0. \quad (3.17)$$

For $A(y) = \sqrt{\frac{2}{3}}\Lambda|y|$ the general solution to this equation is

$$Q(y) = C_1 e^{-\sqrt{\frac{2}{3}}\Lambda|y|} J_2 \left(\sqrt{\frac{6}{\Lambda}} \omega e^{-\sqrt{\frac{\Lambda}{6}}|y|} \right) + C_2 e^{-\sqrt{\frac{2}{3}}\Lambda|y|} Y_2 \left(\sqrt{\frac{6}{\Lambda}} \omega e^{-\sqrt{\frac{\Lambda}{6}}|y|} \right), \quad (3.18)$$

where J_2, Y_2 are first and second kind Bessel function of order 2, respectively, and C_1, C_2 are integration constants [52–54].

Even though the exponential function allows a finite Y_2 , at the origin, let us consider the Bessel function of the first kind Y_1 only, resulting in

$$Q(y) = C_1 e^{-\sqrt{\frac{2}{3}}\Lambda|y|} J_2 \left(\sqrt{\frac{6}{\Lambda}} \omega e^{-\sqrt{\frac{\Lambda}{6}}|y|} \right). \quad (3.19)$$

The boundary condition $Q(0) = 0$ allows only the frequencies satisfying

$$\omega \sqrt{\frac{6}{\Lambda}} = X_{2,n}, \quad (3.20)$$

where $X_{2,n}$ is the n -st root of the Bessel function J_2 , [52]. Insofar as $A(y) > 0$, the number of roots of Eq. (3.20) is finite and it can be restricted to one zero at the origin by assuming $\omega \sqrt{\frac{6}{\Lambda}} = X_1 = 5.14$ [52].

In order to analyse the behaviour of the standing wave geometry, We plotted the function $Q(r)$ in Fig. (1) for $\sqrt{\frac{6}{\Lambda}}\omega = 5.14$ (thin line) and for $\sqrt{\frac{6}{\Lambda}}\omega = 1$ (thick line). Once

the torsion scalar field is given by $h(y, t) = \sqrt{\frac{3}{2\kappa_5}} Q(y) \sin(\omega t)$, the solutions where $Q(0) = 0$ represents a torsion-free standing wave branes.

In the next section the properties of bulk fields in the isotropic and standing wave solutions are investigated. Once space-time time-dependence enhances the complexity of the field equations, a suitable regime to seek for the geometry effects upon the fields is achieved assuming that the field frequency is much smaller than the standing wave frequency ω . Further, considering the average of the configurations [52]

$$\langle u \rangle = \langle \dot{u} \rangle = \langle u' \rangle = \langle (e^{bu}) \rangle = 0, \quad (3.21)$$

the only non-vanishing terms are

$$\langle e^{bu} \rangle = I_0, \quad (3.22)$$

$$\langle (e^{bu})' \rangle = b^2 Z Z' \left[I_0 + \frac{\pi}{2} (I_0 L_1 - I_1 L_0) \right], \quad (3.23)$$

where I_0, I_1 are the modified Bessel functions and Struve L_0, L_1 functions for the arguments bZ .

4 Bulk fields

In this section we investigate the properties of Bulk fields in this torsion geometry. The background Lyra vector provides non-minimal couplings whose effects are analysed.

4.1 Scalar field

In order to study the behaviour of scalar field consider the action for a non-minimally coupled scalar

$$S_\Phi = \frac{1}{2} \int d^5x \sqrt{-g} \left(g^{MN} \nabla_M^L \Phi \nabla_N^L \Phi - \zeta R \frac{\Phi^2}{2} \right). \quad (4.1)$$

The equation of motion (EOM) is $g^{MN} \nabla_M^L \nabla_N^L \Phi - \zeta R \Phi = 0$. Using the Lyra connection expression Eq.(2.4), the EOM reads

$$\partial_M (\sqrt{-g} g^{MN} \partial_N \Phi) + \frac{3}{4} T^P \partial_P \Phi + \zeta R \Phi = 0. \quad (4.2)$$

The torsion coupling $\frac{3}{4} T^P \partial_P \Phi$ provides a dissipative-like term depending on the background vector b_M .

Consider the isotropic brane determined by the warp factor in Eq.(3.7). The Kaluza-Klein decomposition $\Phi(x, y) = \tilde{\Phi}(x) \chi(y)$, where the scalar field on the brane satisfies $\partial_\mu \partial^\mu \tilde{\Phi} = m^2 \tilde{\Phi}$, yields the EOM Eq.4.2 to

$$\chi'' - \left(4\sqrt{-\Lambda} \tanh \lambda y - \frac{3}{2} b_y \right) \chi' + \left\{ m^2 \cosh^p \lambda y - 2\zeta \Lambda \left(\frac{5}{3} + \text{Sech}^2 \lambda y \right) \right\} \chi = 0, \quad (4.3)$$

where $p := \sqrt{-\Lambda}/\lambda$ is a dimensionless parameter. Near the brane, the massive modes satisfy $\chi''(y) + \frac{3}{2} \lambda \sqrt{\frac{2}{\kappa_5}} \chi'(y) + m'^2 \chi(y) = 0$, where $m'^2 = m^2 - \frac{16\zeta\Lambda}{3}$. Thus, the torsion coupling adds a damping factor $\exp(-\frac{3}{4} \lambda \sqrt{\frac{2}{\kappa_5}} y)$ into the massive modes around the brane.

For $\Lambda = -6\lambda^2$, the change of coordinate $z = (1/\lambda) \sinh \lambda y$ turns the metric into the conformal form $ds^2 = \frac{1}{1+\lambda^2 z^2} (\eta_{\mu\nu} dx^\mu dx^\nu - dz^2)$. In this coordinates, the KK equation takes a Schrödinger-like form $-\ddot{\Psi} + U_a(z)\Psi = m^2\Psi$, where

$$U_\Phi = \frac{3\lambda^2}{4(1+\lambda^2 z^2)^2} (5\lambda^2 z^2 - 2) + \frac{2\Lambda\zeta}{(1+\lambda^2 z^2)} \left(\frac{5}{3} + \frac{1}{1+\lambda^2 z^2} \right), \quad (4.4)$$

and the dot stands for d/dz . For $\zeta = 0$, the analogue Schrödinger potential U_Φ has a SUSY-like structure $U_\Phi = W^2 - \dot{W}$, where the superpotential is $W = \frac{3}{2} \left(\dot{A} - \frac{1}{2}\dot{h} \right)$. In this regime, this potential enables a normalized massless mode

$$\Psi_0 = \frac{N}{(1+\lambda^2 z^2)^{3/4}}, \quad (4.5)$$

and it shows a gapless KK massive spectrum. For $\zeta \neq 0$ the analogue SUSY symmetry is




Figure 8. Potential for the scalar field without torsion coupling.




Figure 9. Potential for the scalar field with torsion coupling.

broken, albeit the massive tower remains not localized. Further, for a scalar field without the torsion coupling, the potential exhibits a symmetric profile where the higher ζ is the deeper the potential well around the origin becomes, as shown in Fig.(8). Thus, for a strong non-minimal coupling the potential approaches a reflectionless potential, as the Pöschl-Teller found in [21]. The torsion coupling breaks the Z_2 symmetry of the potential producing a barrier for $z < 0$ and a well for $z > 0$, as shown in Fig.(9). Moreover, the massless mode is modified by the torsion coupling by

$$\Psi_0 = N e^{3h/4} e^{-3A/2}, \quad (4.6)$$

thereby amplifying the wave function near the brane and breaking the Z_2 symmetry.

Now let us consider the standing wave geometry. For the sake of simplicity, we assume a minimal coupling to the Bulk geometry. By setting $\sqrt{\frac{2}{3}}\Lambda y = 2ay$ and $y > 0$, results in the KK equation

$$[\partial_t^2 - e^{-u}(\partial_x^2 + \partial_y^2) - e^{2u}\partial_z^2] \Phi = e^{2ay} \cdot (e^{4ay}\Phi)'. \quad (4.7)$$




Figure 10. Scalar field massless mode for $\lambda = 1$ with the torsion coupling (thin line) and without the torsion coupling (dashed line).

Assuming the KK decomposition $\Phi(x^\nu, y) = e^{ip_\nu x^\nu} \xi(y)$, equation (4.7) takes the form

$$(e^{4ay} \xi')' = e^{2ay} [(p_x^2 + p_y^2)e^{-u} + p_z^2 e^{2u} - p_t^2] \xi. \quad (4.8)$$

Once the *eikonal* functions p_ν are unknown, we employ the condition whereupon the standing wave frequencies are much bigger than the field frequencies, thereby leading to

$$(e^{4ar} \xi')' - e^{2ar} P^2(r) \xi = 0, \quad (4.9)$$

where $P^2(r) = (\langle e^{-u} \rangle - 1)(p_x^2 + p_y^2) + (\langle e^{2u} \rangle - 1)p_z^2$. Due to the complexity of the KK Eq. (4.9) we investigate this equation near the brane $y \rightarrow 0$ and in the asymptotic regime $y \rightarrow \infty$. Nearby the brane $P^2(y)|_{y \rightarrow 0} \sim y^2$, whereas asymptotically $P^2(y)|_{y \rightarrow \infty} \text{const.}$ Thus, the scalar field exhibits the respective behaviour

$$\xi(y)|_{y \rightarrow 0} \sim \text{constant}, \quad (4.10)$$

$$\xi(y)|_{y \rightarrow \infty} \sim e^{-4ay}. \quad (4.11)$$

Therefore, the standing wave scenario enables a normalized massless mode.

Considering the the non-minimal coupling the equation for the extra dimension turns into

$$(e^{4ay} \xi')' - e^{2ay} \bar{P}^2(y) \xi = 0, \quad (4.12)$$

where

$$\bar{P}^2(y) = (\langle e^{-u} \rangle - 1)(p_x^2 + p_y^2) + (\langle e^{2u} \rangle - 1)p_z^2 - \frac{5}{3} \Lambda \epsilon \kappa_5^2 e^{-2ay}. \quad (4.13)$$

The inclusion of the non-minimal term $\frac{5}{3} \Lambda \epsilon \kappa_5^2 e^{-2ay}$ provides only a constant for $y \rightarrow 0$ and for $y \rightarrow \infty$ there is no difference from the results obtained for Eq. (4.9).

4.2 Gauge field

We begin considering a general gauge invariant Lagrangian for a vector field in the form

$$\mathcal{L}_A = \mathcal{L}_{A,T} + \mathcal{L}_{A,S} \quad (4.14)$$

where the a trace-torsion vector coupling is

$$\mathcal{L}_{A,T} := \sqrt{g}(T_C T^C) g^{MN} g^{PR} F_{MP} F_{NR} \quad (4.15)$$

and the axial coupling is

$$\mathcal{L}_{A,S} := \sqrt{g} S^{MP} S^{NR} F_{MP} F_{NR}. \quad (4.16)$$

The field strength is defined as $F_{MP} = \nabla_M^L A_P - \nabla_P^L A_M$ and ∇_P^L is the torsion-free connection. The presence of torsion usually breaks the gauge symmetry [47]. Nevertheless, the existence of an associated torsion-free connection in Lyra geometry preserves the $U(1)$ gauge symmetry. Further, in Lyra geometry, the tensor S_{MN} vanishes and then we consider the $\mathcal{L}_{A,T}$ term only. The coupling between the gauge field and the background vector b_C provides a birefringence interaction for the vector field. The norm of the trace torsion vector T_C can be interpreted as a kind of dilaton interacting with the spin one field. For $b_C = \partial_C h$, this term can be regarded as a derivative dilaton-Maxwell theory. The resulting equation of motion is

$$\partial_M ((b_C b^C \sqrt{g} g^{MN} g^{PR} F_{NR}) = 0. \quad (4.17)$$

In the isotropic scenario with warp factor Eq.(3.7), by performing the KK dimensional reduction $A_\mu(x, y) = \tilde{A}_\mu(x) \Phi(y)$ and choosing $\partial_\mu A^\mu = A_y = 0$, we obtain the KK equation

$$m^2 b^2 \Phi - (b^2 e^{-2A} \Phi)' = 0, \quad (4.18)$$

where the KK mass arises from $\partial_\mu \partial^\mu A^\nu = m^2 A^\nu$ and $b^2 = b_C b^C$. The massless mode is given by

$$\Phi_0 = \int^y \frac{e^{2A}}{b^2} dy' + c. \quad (4.19)$$

Although the first term diverges, a constant massless mode $\Phi_0 = c$ is localized provided that $\int b^2 dy$ converges, as for $b^2 = h'^2$ in Eq.(3.6).

Performing the change of coordinate $z' = \int \frac{e^A}{b} dz''$, the Eq.(4.18) turns into $-\ddot{\Psi}_A + U_A(z') \Psi_A = m^2 \Psi_A$, where

$$U_A = \frac{1}{4} \left(\dot{A} - \frac{\dot{b}}{b} \right)^2 + \frac{1}{2} \frac{d}{dz'} \left(\dot{A} - \frac{\dot{b}}{b} \right). \quad (4.20)$$

Unlike for the scalar field, the Schrödinger equation is not written in the conformal coordinate due to the presence of the non-minimal coupling. In spite of the analytical form of the function $z(y)$ can not be found in general, the regularity and positivity of $z'(y)$ guarantees that some asymptotic and qualitative features can be addressed. Notably, since $\dot{A} = b e^{-A} A'$ the potential vanishes asymptotically provided $b e^{-A} \rightarrow 0$ as $z \rightarrow \infty$. This condition is fulfilled for a covariant constant background Lyra vector $b_M = e^{-A}(0, 0, 0, 0, b_4)$ and for the scalar Lyra $b_M = \partial_M h$ as well.

The corresponding massless bound state has the form

$$\Psi_{0A} = N_A \sqrt{b} e^{-\frac{1}{2}A}, \quad (4.21)$$

which for a covariant constant background vector provides a constant that can be absorbed into the normalization constant. For $b_M = \partial_M h$ the massless is $\Psi_{0A} = N_A e^{\frac{1}{2}A} \dot{h}$ which vanishes asymptotically, for $h' = e^A \dot{h} \rightarrow 0$ as $z \rightarrow \infty$, thereby showing a damping effect of the non-minimal torsion interaction to the massless gauge field.

Now let us study the behaviour of the gauge field in the standing wave model. We assume a minimal coupling and the anisotropic ansatz

$$A_M = v(r)(a_t(x^\mu), e^u a_x(x^\mu), e^u a_{\bar{y}}(x^\mu), e^{-2u} a_z(x^\mu), 0). \quad (4.22)$$

Taking the time average of the y component of Eq. (4.17) yields to $\partial_\alpha (g^{\alpha\beta} A'_\beta) = 0$ and $g^{\alpha\beta} \partial_\alpha A_\beta = \eta^{\alpha\beta} \partial_\alpha a_\beta = 0$, which represent the gauge conditions. The EOM (4.17) on the brane are

$$\partial_\mu (g^{\mu\nu} g^{\alpha\beta} F_{\nu\beta}) - \frac{1}{\sqrt{g}} (\sqrt{g} g^{\alpha\beta} A'_\alpha)' = 0. \quad (4.23)$$

Considering the time average of Eq.(4.23) with the gauge conditions provides

$$v g^{\alpha\beta} \partial_\alpha \partial_\beta a_\mu + e^{-2a|y|} (e^{-2a|y|} v')' a_\mu = 0. \quad (4.24)$$

In addition, assuming the *eikonal* form $a_\mu(x^\nu) \sim \epsilon_\mu e^{ip_\nu x^\nu}$, the Eq. (4.24) results in

$$(e^{2a|y|} v')' - P^2(y)v = 0. \quad (4.25)$$

The asymptotic behaviour of the massless mode is

$$v(y)|_{y \rightarrow 0} \sim \text{constant}, \quad (4.26)$$

$$v(y)|_{y \rightarrow \infty} \sim e^{-2a|y|}. \quad (4.27)$$

Therefore, the anisotropy driven by the standing wave brane is enough to trap gauge massless mode.

4.3 Fermion field

We propose the following non-minimal coupling of the massless bulk spinor and the torsional geometry

$$S_\Psi = \int d^5x \sqrt{-g} \left[\bar{\Psi} i \Gamma^M D_M \Psi - (\epsilon_1 h^p \sqrt{R} - \epsilon_2 \square h) \bar{\Psi} \Psi \right], \quad (4.28)$$

where $p \in \mathbb{N}$, $\Gamma^M = e^M_{\bar{M}} \gamma^{\bar{M}}$ are the Dirac matrices in a curved spacetime, $\gamma^{\bar{M}}$ are the flat Dirac matrices and the *vielbeins* $e^M_{\bar{M}}$ satisfy $g_{MN} = \eta_{\bar{M}\bar{N}} e^{\bar{M}}_M e^{\bar{N}}_N$. The spinorial covariant derivative in torsional Lyra geometry has the form [49–51]

$$D_M = \partial_M + \Omega_M^L, \quad (4.29)$$

where $\Omega_M^L = \Omega_M + \Omega'_M$ is the torsion spinor connection, with $\Omega_M = \frac{1}{4}\Gamma_M^{\bar{M}\bar{N}}\gamma_{\bar{M}}\gamma_{\bar{N}}$ being the torsion-free spinor connection and $\Gamma_M^{\bar{M}\bar{N}}$ are 1-form Levi-Civita connection coefficients, i.e., $\omega_{\bar{N}}^{\bar{M}} = \Gamma_{\bar{M}\bar{N}}^{\bar{M}}dx^M$. The torsion provides a new spin connection term $\Omega'_M = -\frac{1}{2}T_M$ which for $b_M = \partial_M h$ yields to $\Omega'_M = -(3/2)\partial_M h$. Further, the background vector provides a non-minimal mass term of form

$$F(h, g) = (\epsilon_1 h^p \sqrt{R} - \epsilon_2 \square h), \quad (4.30)$$

which can be regarded as a kind of geometric Yukawa-like derivative interaction between the fermion, the Lyra scalar (torsion) and the metric. The coupling constants $\epsilon_{1,2}$ are dimensionless real numbers.

For the isotropic scenario in the conformal coordinate the metric can be written as $ds^2 = \eta_{\bar{M}\bar{N}}\hat{\theta}^{\bar{M}} \otimes \hat{\theta}^{\bar{N}}$ where $\hat{\theta}^{\bar{M}} = e^{-A}\delta_M^{\bar{M}}dx^M$. The Cartan equation $d\hat{\theta}^{\bar{M}} + \omega_{\bar{N}}^{\bar{M}}\hat{\theta}^{\bar{N}} = 0$ yields to the nonvanishing 1-form connections coefficients $\Gamma_{5\nu}^{\mu} = -\dot{A}\delta_{\nu}^{\mu}$ and $\Gamma_{\bar{\mu}\bar{\nu}}^5 = -\dot{A}\eta_{\bar{\mu}\bar{\nu}}$. Thus, the Dirac equation reads

$$\left[\gamma^{\mu}(\partial_{\mu} + \frac{3}{2}b_{\mu}) + \gamma^z(\partial_z - 2\dot{A} - \frac{3}{2}b_z) - e^{-A}(\epsilon_1 h^p \sqrt{R} - \epsilon_2 \square h) \right] \Psi = 0. \quad (4.31)$$

For $\gamma^{\mu}(\partial_{\mu} + \frac{3}{2}b_{\mu})\Psi = m\Psi$ and $\Psi(x, z) = f(z) \sum \psi_{4R}(x)\alpha_R(z) + \psi_{4L}\alpha_L(z)$, where $f(z) = e^{2A + \frac{3}{2}\int b_z dz'}$ the Dirac equation leads to

$$\begin{aligned} (\partial_z + e^{-A}(\epsilon_1 h^p \sqrt{R} - \epsilon_2 \square h))\alpha_L &= m\alpha_R \\ (\partial_z - e^{-A}(\epsilon_1 h^p \sqrt{R} - \epsilon_2 \square h))\alpha_R &= -m\alpha_L. \end{aligned} \quad (4.32)$$

By decoupling the system (4.32), we obtain $-\ddot{\alpha}_{R,L} + U_{R,L}(z)\alpha_{R,L} = m^2\alpha_{R,L}$, where

$$U_{R,L} = e^{-2A}(\epsilon_1 h^p \sqrt{R} - \epsilon_2 \square h)^2 \pm \frac{d}{dz}[e^{-A}(\epsilon_1 h^p \sqrt{R} - \epsilon_2 \square h)] \quad (4.33)$$

Due to the complexity of the fermion potential, we study the properties for each interaction separately. Let us first consider the $\epsilon_2 = 0$ case for $p = 1$. We plotted the analogue quantum potential for the left chirality using $\lambda = 2$ in figure (11) and the massless mode in figure (12). Unlike the scalar field, the non-minimal coupling intensify the interaction between the fermion and the brane by enhancing the potential barrier and the massless mode decay.

By varying the parameter p the width of the potential well increases, as well as the high of the barriers. Besides, figure (13) shows two asymmetric minima inside the volcano well. These minima deforms the massless bound state by shifting their peaks from the origin, as shown in figure (14). Moreover, the increase of the potential barrier due to p augmented the massless mode localization.

Finally, let us consider the derivative coupling for $\epsilon_1 = 0$. This interaction strongly modifies the volcano potential by reducing the width and the depth of the potential well, though keeping only one minimum inside the well, as shown in figure (15). The corresponding bound state is localized inside the well and its amplitude increase as the brane width decreases, as shown in figure (16).




Figure 11. Potential for the left chirality.




Figure 12. Fermion massless mode.




Figure 13. Left-handed potential with $p = 3$ (dotted line) and $p = 5$ (thick line).




Figure 14. Left-handed massless mode with $p = 3$ (dotted line) and $p = 5$ (thick line).




Figure 15. Left-handed potential with $\epsilon_1 = 0$ for $\lambda = 1$ (dotted line), $\lambda = 2$ (thin line) and $\lambda = 3$ (thick line).




Figure 16. Left-handed massless mode with $\epsilon_1 = 0$ for $\lambda = 1$ (dotted line), $\lambda = 2$ (thin line) and $\lambda = 3$ (thick line).

In the standing wave background, the spin connection has the following nonvanishing

components

$$\begin{aligned}\Omega_t^{\bar{t}\bar{y}} &= -\left(e^{a|y|}\right)'; & \Omega_x^{\bar{x}\bar{y}} &= \Omega_y^{\bar{y}\bar{y}} = -\left(e^{a|y|+u/2}\right)'; \\ \Omega_z^{\bar{z}\bar{y}} &= -\left(e^{a|y|-u}\right)'; & \Omega_x^{\bar{x}\bar{t}} &= \Omega_y^{\bar{y}\bar{t}} = \left(e^{u/2}\right)'; & \Omega_z^{\bar{z}\bar{t}} &= \left(e^{-u}\right)'. \end{aligned} \quad (4.34)$$

Assuming the chiral KK decomposition $\Psi(x^\nu, y) = \psi_L(x^\nu)\lambda(y) + \psi_R(x^\nu)\rho(y)$ and considering the standing wave average time regime, the equations for the extra dimension variable are, respectively [52],

$$\lambda'' - \left[5a \operatorname{sgn}(y) - \frac{P'}{P}\right] \lambda' + \left[4a\delta(y) + 6a^2 - 2a \operatorname{sgn}(y)\frac{P'}{P} - P^2 e^{-2a|y|}\right] \lambda = 0, \quad (4.35)$$

and

$$\rho(y)D = e^{a|y|} \frac{\sigma^i P_i(y)}{P^2(y)} [2a \operatorname{sgn}(y) + \partial_y] \lambda(y)L. \quad (4.36)$$

where σ^i ($i = x, y, z$) represent the standard 2×2 Pauli matrices. The average momentum components are

$$\begin{aligned}P_x(y) &= [I_0(|C_1|Q(y)/2) - 1] p_x, \\ P_y(y) &= [I_0(|C_1|Q(y)/2) - 1] p_{\bar{y}}, \\ P_z(y) &= [I_0(|C_1|Q(y)) - 1] p_z, \end{aligned} \quad (4.37)$$

where the function $Q(y)$ is the standing wave solution given in Eq. (3.19), L and D are constant 2-spinors. Likewise the bosonic fields considered above, the Eq.(4.35) is too complicated to be addressed algebraically. Yet, we only need to consider its asymptotic form close and far from the brane. Nearby the brane the left and right fermions are $\lambda(y)|_{y \rightarrow \pm 0} = B e^{-2a|y|}$ and $\rho(y)|_{y \rightarrow \pm 0} = 0$, whilst asymptotically they behave as $\lambda(y)|_{y \rightarrow \pm \infty} \sim e^{-3a|y|}$ and $\rho(y)|_{y \rightarrow \pm \infty} \sim e^{-2a|y|}$, respectively. The relations above show the standing wave scenario allows a bound massless left fermion on the brane while the right chirality is kept a free bulk state.

5 Final remarks and perspectives

In this work we proposed torsional 5D isotropic and anisotropic braneworld scenarios whose dynamics is equivalent to a tensor-vector gravity theory. The torsion steams from a background vector in the so-called Lyra geometry.

Inasmuch as the tensor vector is covariantly constant, the squared norm of this Lyra vector can be regarded as a source for the bulk cosmological constant. Assuming the Lyra vector as the derivative of a scalar field we obtained an isotropic scenario similar to the coupled two field Bloch brane [10]. This configuration can be obtained from a self-interacting modified sine-Gordon potential. The profile of such potential exhibits a compact support around the vacua of the usual sine-Gordon potential. The existence of BPS configurations of this potential and the relation with compactons solutions are interesting for future developments [55, 56]. Moreover, the Lyra scalar also acts as a source for a smooth

and anisotropic standing wave model. In the isotropic model the Lyra scalar vanishes at the origin whereas the standing wave solution supports a non-zero Lyra scalar on the brane whose cosmological can be further investigated.

The background vector allow us to define non-minimal couplings to the bulk fields. For a scalar field in the isotropic model a curvature mass-like term modifies the interaction between the KK modes and the brane. In fact, for high values of the non-minimal coupling parameter the analogue KK quantum potential tends to a reflectionless potential, as one found in [21]. Thus, this non-minimal coupling reduces the chances to detect massive modes on the brane. For the Abelian gauge vector we proposed a modified kinetic term with a birefringence-like factor proportional to the norm of the Lyra vector. If $b_M = \partial_M h$, this coupling resembles a derivative dilaton-gauge vector interaction. Though more involved, the anisotropy in standing wave geometry enables the existence of a normalized massless mode even without the non-minimal coupling for both the scalar and vector fields.

The spinor connection is modified by the torsion that yields to a derivative $\bar{\Psi}i\Gamma^M\partial_M h\Psi$ coupling. In addition, we proposed a non-minimal coupling of form $hR\bar{\Psi}\Psi$, which can be considered as a pure-geometric Yukawa-like coupling. It turns out that only the non-minimal coupling leads to relevant modifications to the fermionic KK modes. Indeed, the volcano potential is everywhere positive, whilst a gapless massive tower and a localized massless left-chiral mode are still allowed. It is worthwhile to mention that the KK reduction provides a CPT-odd SME Lorentz violating term $i\bar{\psi}\gamma^\mu b_\mu\psi$ [38, 39]. We argue that by considering more non-minimal couplings other SME terms could be generated and their experimental test could be used to obtain upper bound to the bulk-brane torsion. Furthermore, in the standing wave model the Lyra scalar is non zero on the brane what could lead to interesting cosmological effects.

The close relationship between Lyra and Weyl geometries in braneworld models is another topic which was addressed. In a integrable Weyl geometry and using a general conformal invariant action, some pure-geometric thick branes were found [20, 21]. Therefore, the investigation of a pure-geometric thick branes in a conformal Lyra geometry, and the effects upon the fields thereof, it is a worth extension of the present work.

Acknowledgments

The authors thank the Fundação Cearense de Apoio ao Desenvolvimento Científico e Tecnológico (FUNCAP)(PNE-0112-00061.01.00/16), and the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) for grants Nos. 307688/2016-0 (WTC), 312356/2017-0 (JEGS) and 308638/2015-8 (CASA).

References

- [1] L. Randall and R. Sundrum, “A Large mass hierarchy from a small extra dimension,” *Phys. Rev. Lett.* **83**, 3370 (1999) [hep-ph/9905221].
- [2] N. Arkani-Hamed, S. Dimopoulos and G. R. Dvali, “Phenomenology, astrophysics and cosmology of theories with submillimeter dimensions and TeV scale quantum gravity,” *Phys. Rev. D* **59**, 086004 (1999) [hep-ph/9807344].

- [3] J. W. Chen, M. A. Luty and E. Ponton, “A Critical cosmological constant from millimeter extra dimensions,” *JHEP* **0009**, 012 (2000) [hep-th/0003067].
- [4] J. Khoury, B. A. Ovrut, P. J. Steinhardt and N. Turok, “The Ekpyrotic universe: Colliding branes and the origin of the hot Big Bang,” *Phys. Rev. D* **64**, 123522 (2001) [hep-th/0103239].
- [5] I. Oda, “Bosonic fields in the string - like defect model,” *Phys. Rev. D* **62**, 126009 (2000) [hep-th/0008012]. [hep-th/0008012].
- [6] T. Gherghetta and M. E. Shaposhnikov, “Localizing gravity on a string - like defect in six-dimensions,” *Phys. Rev. Lett.* **85**, 240 (2000) [hep-th/0004014].
- [7] A. Kehagias and K. Tamvakis, “Localized gravitons, gauge bosons and chiral fermions in smooth spaces generated by a bounce,” *Phys. Lett. B* **504**, 38 (2001) [hep-th/0010112].
- [8] C. A. S. Almeida, M. M. Ferreira, Jr., A. R. Gomes and R. Casana, “Fermion localization and resonances on two-field thick branes,” *Phys. Rev. D* **79**, 125022 (2009) [arXiv:0901.3543 [hep-th]].
- [9] W. T. Cruz, A. R. P. Lima and C. A. S. Almeida, “Gauge field localization on the Bloch Brane,” *Phys. Rev. D* **87**, no. 4, 045018 (2013) [arXiv:1211.7355 [hep-th]].
- [10] D. Bazeia and A. R. Gomes, “Bloch brane,” *JHEP* **0405**, 012 (2004) [hep-th/0403141].
- [11] W. T. Cruz, L. J. S. Sousa, R. V. Maluf and C. A. S. Almeida, “Graviton resonances on two-field thick branes,” *Phys. Lett. B* **730**, 314 (2014) [arXiv:1310.4085 [hep-th]].
- [12] T. Flacke, D. W. Kang, K. Kong, G. Mohlabeng and S. C. Park, “Electroweak Kaluza-Klein Dark Matter,” *JHEP* **1704**, 041 (2017) [arXiv:1702.02949 [hep-ph]].
- [13] M. Zucker, “Supersymmetric brane world scenarios from off-shell supergravity,” *Phys. Rev. D* **64**, 024024 (2001) [hep-th/0009083].”
- [14] T. G. Rizzo, T. G. Rizzo, “Lorentz Violation in Warped Extra Dimensions,” *JHEP* **1011**, 156 (2010) [arXiv:1008.0380 [hep-ph]].
- [15] L. Pilo and A. Riotto, “The noncommutative brane world,” *JHEP* **0103**, 015 (2001) [hep-ph/0012174].
- [16] J. E. G. Silva and C. A. S. Almeida, “Scalar field localization on 3-branes placed at a warped resolved conifold,” *Phys. Rev. D* **84**, 085027 (2011) [arXiv:1110.1597 [hep-th]].
- [17] N. Arkani-Hamed, S. Dimopoulos, G. R. Dvali and N. Kaloper, “Many fold universe,” *JHEP* **0012**, 010 (2000) [hep-ph/9911386].
- [18] Y. Zhong, Y. P. Zhang, W. D. Guo and Y. X. Liu, “Gravitational resonances in mimetic thick branes,” *JHEP* **1904**, 154 (2019) [arXiv:1812.06453 [gr-qc]].
- [19] O. Arias, R. Cardenas and I. Quiros, “Thick brane worlds arising from pure geometry,” *Nucl. Phys. B* **643**, 187 (2002) [hep-th/0202130].
- [20] Y. X. Liu, X. H. Zhang, Y. X. Liu, X. H. Zhang, L. D. Zhang and Y. S. Duan, “Localization of Matters on Pure Geometrical Thick Branes,” *JHEP* **0802**, 067 (2008) [arXiv:0708.0065 [hep-th]].
- [21] Y. X. Liu, K. Yang and Y. Zhong, “de Sitter Thick Brane Solution in Weyl Geometry,” *JHEP* **1010**, 069 (2010). [arXiv:0911.0269 [hep-th]].

- [22] H. Guo, A. Herrera-Aguilar, Y. X. Liu, D. Malagon-Morejon and R. R. Mora-Luna, “Localization of bulk matter fields, the hierarchy problem and corrections to Coulomb’s law on a pure de Sitter thick braneworld,” *Phys. Rev. D* **87**, no. 9, 095011 (2013).
- [23] Tao-Tao Sui, Li Zhao, Yu-Peng Zhang, Qun-Ying Xie, Localization and mass spectra of various matter fields on Weyl thin brane, *Eur. Phys. J. C* **77** (2017) no.6, 411.
- [24] M. Novello and S. E. P. Bergliaffa, “Bouncing Cosmologies,” *Phys. Rept.* **463**, 127 (2008), [arXiv:0802.1634 [astro-ph]].
- [25] F. W. Hehl, P. Von Der Heyde, G. D. Kerlick and J. M. Nester, “General Relativity with Spin and Torsion: Foundations and Prospects,” *Rev. Mod. Phys.* **48**, 393 (1976).
- [26] I. L. Shapiro, “Physical aspects of the space-time torsion,” *Phys. Rept.* **357**, 113 (2002) [hep-th/0103093].
- [27] J. A. Helayel-Neto, A. Penna-Firme and I. L. Shapiro, “Conformal symmetry, anomaly and effective action for metric-scalar gravity with torsion,” *Phys. Lett. B* **479**, 411 (2000).
- [28] B. Mukhopadhyaya, S. Sen and S. SenGupta, “Does a Randall-Sundrum scenario create the illusion of a torsion free universe?,” *Phys. Rev. Lett.* **89**, 121101 (2002) Erratum: [*Phys. Rev. Lett.* **89**, 259902 (2002)]
- [29] B. Mukhopadhyaya, S. Sen and S. SenGupta, “A Randall-Sundrum scenario with bulk dilaton and torsion,” *Phys. Rev. D* **79**, 124029 (2009).
- [30] M. Kalb and P. Ramond, Classical direct interstring action, *Phys.Rev.D* **9**, 2273 (1974).
- [31] K. Becker and K. Dasgupta, “Heterotic strings with torsion,” *JHEP* **0211**, 006 (2002). [hep-th/0209077].
- [32] J. Gaillard and J. Schmude, “The Lift of type IIA supergravity with D6 sources: M-theory with torsion,” *JHEP* **1002**, 032 (2010). [arXiv:0908.0305 [hep-th]].
- [33] J. M. Hoff da Silva and R. da Rocha, “Torsion Effects in Braneworld Scenarios,” *Phys. Rev. D* **81**, 024021 (2010). [arXiv:0912.5186 [hep-th]].
- [34] J. M. Hoff da Silva and R. da Rocha, “Braneworld remarks in Riemann-Cartan manifolds,” *Class. Quant. Grav.* **26**, 055007 (2009). Erratum: [*Class. Quant. Grav.* **26**, 179801 (2009)] [arXiv:0804.4261 [gr-qc]].
- [35] N. R. F. Braga and C. N. Ferreira, “Topological mass term in effective brane-world scenario with torsion,” *JHEP* **0503**, 039 (2005). [hep-th/0410186].
- [36] J. Ellis, N. E. Mavromatos and S. Sarkar, “Environmental CPT Violation in an Expanding Universe in String Theory,” *Phys. Lett. B* **725**, 407 (2013) [arXiv:1304.5433 [gr-qc]].
- [37] J. Yang, Y. L. Li, Y. Zhong and Y. Li, “Thick Brane Split Caused by Spacetime Torsion,” *Phys. Rev. D* **85**, 084033 (2012). [arXiv:1202.0129 [hep-th]].
- [38] V. A. Kostelecky, “Gravity, Lorentz violation, and the standard model,” *Phys. Rev. D* **69**, 105009 (2004).
- [39] V. A. Kostelecky, N. Russell and J. Tasson, “New Constraints on Torsion from Lorentz Violation,” *Phys. Rev. Lett.* **100**, 111102 (2008), [arXiv:0712.4393 [gr-qc]].
- [40] Lyra, G., Über eine Modifikation der Riemannschen Geometrie, *Math. Z.* **54**, 52 (1951).
- [41] D. K. Sen and J. R. Vanstone, On Weyl and Lyra Manifolds, *Jour. Math. Phys.* **13**, 990 (1972).

- [42] D. K. Sen and K. A. Dunn, A Scalar Tensor Theory of Gravitation in a Modified Riemannian Manifold, *Jour. Math. Phys.* **12**, 578 (1971).
- [43] W. D. Halford, Cosmological theory based on Lyra's geometry, *Aust. Jour. Phys.*, **23**, 863 (1970).
- [44] V. K. Shchigolev, E. A. Semenova, Scalar field cosmology in Lyra's geometry. *Int. Jour. Adv. Astr.*, **3** (2015) 117-122.
- [45] F. Rahaman, J. K Bera, Higher dimensional cosmological model in Lyra geometry. *Int. Jour. Mod. Phys. D*, **10**, 729-733 (2001).
- [46] S. D. Katore, S. P. Hatkar, Kaluza Klein universe with magnetized anisotropic dark energy in general relativity and Lyra manifold. *New Astr.*, **34**, 172-177 (2015).
- [47] R. Casana, J. T. Lunardi, B. M. Pimentel and R. G. Teixeira, "Spin 1 fields in Riemann-Cartan space-times via Duffin-Kemmer-Petiau theory," *Gen. Rel. Grav.* **34**, 1941 (2002), [gr-qc/0203068].
- [48] R. Casana, C. A. M. de Melo, B. M. Pimentel, Electromagnetic field in Lyra manifold: a first order approach. *Braz. Jour. Phys.*, **35**, 1151-1154 (2005).
- [49] H. T. Nieh and M. L. Yan, "Quantized Dirac Field in Curved Riemann-Cartan Background. 1. Symmetry Properties, Green's Function," *Annals Phys.* **138**, 237 (1982).
- [50] R. Casana, C. A. M. de Melo, B. M. Pimentel, Spinorial field and Lyra geometry *Astrophys. Space Sci.* **305**, 125 (2006).
- [51] R. Casana, C. A. M. de Melo and B. M. Pimentel, "Massless DKP field in Lyra manifold," *Class. Quant. Grav.* **24**, 723 (2007).
- [52] M. Gogberashvili, Localization of Matter Fields in the 5D Standing Wave Braneworld. *JHEP* **1209** (2012) 056. arXiv:1204.2448 [hep-th]
- [53] M. Gogberashvili, P. Midodashvili and L. Midodashvili, Localization of gauge bosons in the 5D standing wave braneworld, *Phys. Lett. B* **707**, 169 (2012). [arXiv:1105.1866 [hep-th]].
- [54] L. J. S. Sousa, W. T. Cruz, C. A. S. Almeida, Six-dimensional standing-wave braneworld with normal matter as source. *Phys. Rev. D*, **89**, 064006 (2014).
- [55] P. Rosenau, J. Hyman, Compactons: Solitons with finite wavelength, *Phys. Rev. Lett.* **70**, 564 (1993)
- [56] Philip Rosenau and Alon Zilburg, Compactons, *Journal of Physics A* **51**, 343001 (2018)