

# On the energy of a non-singular black hole solution satisfying the weak energy condition

I. Radinschi<sup>\*1</sup>, Th. Grammenos<sup>†2</sup>, F. Rahaman<sup>‡3</sup>, M. M. Cazacu<sup>§1</sup>, A. Spanou<sup>¶4</sup>, and J. Chakraborty<sup>||3</sup>

<sup>1</sup>Department of Physics , “Gheorghe Asachi” Technical University, Iasi, 700050, Romania

<sup>2</sup>Department of Civil Engineering, University of Thessaly, 383 34 Volos, Greece

<sup>3</sup>Department of Mathematics, Jadavpur University, Kolkata 700 032, West Bengal, India.

<sup>4</sup>School of Applied Mathematics and Physical Sciences, National Technical University of Athens, 157 80 Athens, Greece

June 6, 2019

## Abstract

The energy-momentum localization for a new four-dimensional and spherically symmetric, charged black hole solution that through a coupling of general relativity with non-linear electrodynamics is everywhere non-singular while it satisfies the weak energy condition is investigated. The Einstein and Møller energy-momentum complexes have been employed in order to calculate the energy distribution and the momenta for the aforesaid solution. It is found that the energy distribution depends explicitly on the mass and the charge of the black hole, on two parameters arising from the space-time geometry considered, and on the radial coordinate. Further, in both prescriptions all the momenta vanish. In addition, a comparison of the results obtained by the two energy-momentum complexes is made, whereby some limiting and particular cases are pointed out.

---

\*radinschi@yahoo.com

†thgramme@civ.uth.gr

‡rahaman@associates.iucaa.in

§marius.cazacu@tuiasi.ro

¶aspanou@central.ntua.gr

||joydeep\_sweety2010@rediffmail.com

# 1 Introduction

Even though the problem of energy-momentum localization has triggered a lot of interesting research work, it still remains not fully answered for more than a century. The only step one can hope to make forward is to find a more powerful tool for accessing this issue of general relativity.

The use of different tools for energy-momentum localization, like super-energy tensors [1]-[4], quasi-local definitions [5]-[9] and the energy-momentum complexes [10]-[16] has led to the development of several interesting works. In particular, the energy-momentum complexes of Einstein [10], Landau-Lifshitz [12], Papapetrou [13], Bergmann-Thomson [14] and Weinberg [16] are pseudotensorial quantities and coordinate dependent. They can be used in Cartesian and quasi-Cartesian coordinates, more precisely in Schwarzschild Cartesian coordinates and in Kerr-Schild Cartesian coordinates, and have yielded so far many physically meaningful results [17]-[23]. The Møller energy-momentum complex [15] allows the calculation of energy and momenta in any coordinate system, including Schwarzschild Cartesian coordinates and Kerr-Schild Cartesian coordinates, and has also provided physically interesting results for many space-time geometries, in particular for  $(3 + 1)$ ,  $(2 + 1)$  and  $(1 + 1)$  space-times [24]-[44]. It is worth noting that different pseudotensors yield the same energy value for any metric of the Kerr-Schild class and also for solutions more general than those of the Kerr-Schild class (for reviews and references, see the works [48], [49] and [50]). Moreover, the localization of energy-momentum has been studied in the context of teleparallel theory of gravitation whereby many similar results have been obtained [51]-[62].

The Einstein, Landau-Lifshitz, Papapetrou, Bergmann-Thomson, Weinberg and Møller prescriptions are in agreement with the definition of the quasi-local mass given by Penrose [45] and developed by Tod [46] for some gravitating systems. Recently, the new concept in the localization of energy is that of quasi-local energy-momentum associated with a closed 2-surface. In fact, it has been demonstrated that “the quasi-local quantities could provide a more detailed characterization of the states of the gravitational field” [47]. The energy localization is also connected with the quasi-local energy given by Wang and Yau [63], [64]. The rehabilitation of energy-momentum complexes concerns the searching for a common quasi-local energy value. Recently, an important discovery has been made, namely, by considering pseudotensors and quasi-local approaches in the context of the Hamiltonian formulation and with the choice of a 4D isometric Minkowski reference geometry on the boundary, it is found that for any closed 2-surface there is a common value for the quasi-local energy for all expressions that are in agreement (to linear order) with the Freud superpotential or, put simply, all the quasi-local expressions in a large class yield the same energy-momentum [65], [66].

This paper is organized as follows. In Section 2 we briefly present the non-singular black hole solution that satisfies the weak energy condition. In Section 3 the two prescriptions of Einstein and Møller used for the calculation of the energy distribution and momenta are introduced. In Section 4 we show the results obtained for the aforementioned non-singular black hole solution by applying the pseudotensorial definitions given in Section 3. Finally, in the Discussion we summarize the results and we focus on commenting about some limiting and particular cases. Throughout we have used geometrized units ( $c = G = 1$ ) and for the signature the choice has been  $(+, -, -, -)$ . In the case of the Einstein prescription we have used the Schwarzschild Cartesian coordinates  $\{t, x, y, z\}$  and for the Møller prescription the Schwarzschild coordinates  $\{t, r, \theta, \phi\}$ , respectively. Further, Greek indices take values from 0 to 3, while Latin indices run from 1 to 3.

## 2 The Non-Singular Black Hole Solution Satisfying the Weak Energy Condition

In this section the new spherically symmetric and charged non-singular black hole solution that satisfies the weak energy condition developed by L. Balart and E. C. Vagenas [67] is introduced. This black hole solution has been found in the context of general relativity coupled to non linear electrodynamics via a term  $L(F)$  in the action, with the Lorentz-invariant scalar  $F = \frac{1}{4}F^{\mu\nu}F_{\mu\nu}$ , and  $F_{\mu\nu}$  is the electromagnetic tensor that reduces to the Maxwellian field tensor in the weak field case. Here,  $F_{\mu\nu}$  is restricted only to the electric field. The black hole metric has been constructed by employing the Dagum distribution [68], a continuous probability distribution, given by

$$\sigma(x) = \frac{ap x^{ap-1}}{b^{ap} [1 + (\frac{x}{b})^a]^{p+1}}, \quad (1)$$

where  $x > 0$ , while the parameters  $a, b, p \in \mathbb{R}^+$ . For  $p = \frac{1}{a}$  and  $b = 1$ ,  $\sigma(x)$  can be written as

$$\sigma(x) = \frac{1}{(1 + x^a)^{\frac{a+1}{a}}}. \quad (2)$$

Here, with the transformation  $x \rightarrow \frac{q^2}{Mr}$  the black hole metric function reads

$$f(r) = 1 - \frac{2M}{r} \left\{ \frac{1}{\left[1 + \gamma \left(\frac{q^2}{Mr}\right)^a\right]^{\frac{a+1}{a}}} \right\}^\beta, \quad (3)$$

with  $\gamma \in \mathbb{R}^+$ , while  $M$  and  $q$  are the mass and the charge of the black hole, respectively. For  $\beta \geq \frac{3}{(a+1)}$  this black hole metric is non-singular everywhere,

while for  $\beta \leq \frac{3}{(a+1)}$  it satisfies the weak energy condition  $T_{\mu\nu}u^\mu u^\nu \geq 0$  for every timelike vector  $u^\mu$ , according to which the local energy density must be positive-definite. This general inequality is shown in [69] to be equivalent to two inequalities on the first and second derivative of the mass function w.r.t.  $r$ . In order to have a non-singular black hole solution that also satisfies the weak energy condition, one imposes the condition  $\beta = \frac{3}{(a+1)}$ , so that the obtained new spherically symmetric, static and charged non-singular black hole metric has a line element of the form

$$ds^2 = B(r)dt^2 - A(r)dr^2 - r^2(d\theta^2 + \sin^2\theta d\phi^2), \quad (4)$$

with  $B(r) = f(r)$ ,  $A(r) = \frac{1}{f(r)}$ , and the metric function now reads

$$f(r) = 1 - \frac{2M}{r} \left[ \frac{1}{1 + \gamma \left( \frac{q^2}{Mr} \right)^a} \right]^{3/a}. \quad (5)$$

The corresponding electric field is

$$E(r) = \frac{3q}{2r^2} \frac{\gamma(3+a) \left( \frac{q^2}{Mr} \right)^{a-1}}{\left[ 1 + \gamma \left( \frac{q^2}{Mr} \right)^a \right]^{2+3/a}}. \quad (6)$$

For small values of the radial coordinate  $r$  the family of metrics derived from (5) behaves as a de Sitter black hole

$$f(r) \approx 1 - \frac{2M^4}{\gamma^{3/a} q^6} r^2, \quad (7)$$

while asymptotically it behaves as the Schwarzschild black hole.

It is important to notice that a particular case is obtained for  $a = 1$  (hence  $\beta = 3/2$ ) and  $\gamma = \frac{1}{6}$ , so that the metric function (5) reads

$$f(r) = 1 - \frac{2M}{r} \left( \frac{1}{1 + \frac{q^2}{6Mr}} \right)^3 \quad (8)$$

and the black hole solution asymptotically behaves as the Reissner-Nordström solution. For all the other values of the parameters  $a$  and  $\gamma$  the non-singular black hole space-time geometries with the metric function (5) asymptotically do not have the behaviour of the Reissner-Nordström solution. It must also be pointed out that only for the values  $a = 1$  and  $\gamma = \frac{1}{6}$  one gets a nonlinear electrodynamics model leading to Maxwell's electrodynamics in the weak field approximation.

### 3 Einstein and Møller Prescriptions

The Einstein energy-momentum complex [10] defined for a (3 + 1) dimensional space-time is given by

$$\theta_\nu^\mu = \frac{1}{16\pi} h_{\nu,\lambda}^{\mu\lambda}. \quad (9)$$

The superpotentials  $h_\nu^{\mu\lambda}$  are expressed as

$$h_\nu^{\mu\lambda} = \frac{1}{\sqrt{-g}} g_{\nu\sigma} \left[ -g(g^{\mu\sigma} g^{\lambda\kappa} - g^{\lambda\sigma} g^{\mu\kappa}) \right]_{,\kappa} \quad (10)$$

and satisfy the necessary antisymmetric property

$$h_\nu^{\mu\lambda} = -h_\nu^{\lambda\mu}. \quad (11)$$

In the Einstein prescription the local conservation law holds:

$$\theta_{\nu,\mu}^\mu = 0. \quad (12)$$

The energy and momentum can be calculated with

$$P_\mu = \iiint \theta_\mu^0 dx^1 dx^2 dx^3, \quad (13)$$

where  $\theta_0^0$  and  $\theta_i^0$  represent the energy and momentum density components, respectively.

Applying Gauss' theorem, the energy-momentum becomes

$$P_\mu = \frac{1}{16\pi} \iint h_\mu^{0i} n_i dS, \quad (14)$$

with  $n_i$  the outward unit normal vector over the surface  $dS$ . In eq. (14)  $P_0$  represents the energy.

The expression for the Møller energy-momentum complex [15] is given by

$$\mathcal{J}_\nu^\mu = \frac{1}{8\pi} M_{\nu,\lambda}^{\mu\lambda}, \quad (15)$$

with the Møller superpotentials  $M_\nu^{\mu\lambda}$

$$M_\nu^{\mu\lambda} = \sqrt{-g} \left( \frac{\partial g_{\nu\sigma}}{\partial x^\kappa} - \frac{\partial g_{\nu\kappa}}{\partial x^\sigma} \right) g^{\mu\kappa} g^{\lambda\sigma}. \quad (16)$$

The Møller superpotentials  $M_\nu^{\mu\lambda}$  are also antisymmetric

$$M_\nu^{\mu\lambda} = -M_\nu^{\lambda\mu}. \quad (17)$$

Møller's energy-momentum complex also satisfies the local conservation law

$$\frac{\partial \mathcal{J}_\nu^\mu}{\partial x^\mu} = 0. \quad (18)$$

In (15)  $\mathcal{J}_0^0$  is the energy density and  $\mathcal{J}_i^0$  represents the momentum density components.

For the Møller prescription, the energy and momentum distributions are given by

$$P_\mu = \iiint \mathcal{J}_\mu^0 dx^1 dx^2 dx^3 \quad (19)$$

and the energy distribution can be evaluated with

$$E = \iiint \mathcal{J}_0^0 dx^1 dx^2 dx^3. \quad (20)$$

Using Gauss' theorem one obtains

$$P_\mu = \frac{1}{8\pi} \iint M_\mu^{0i} n_i dS. \quad (21)$$

## 4 Energy-Momentum Distribution of the Non-Singular Black Hole Solution Satisfying the Weak Energy Condition

In order to perform the calculations using the Einstein prescription, the metric given by the line element (4) is transformed into Schwarzschild Cartesian coordinates applying the coordinate transformation  $x = r \sin \theta \cos \varphi$ ,  $y = r \sin \theta \sin \varphi$ ,  $z = r \cos \theta$ . Then, the line element takes the following form:

$$ds^2 = B(r)dt^2 - (dx^2 + dy^2 + dz^2) - \frac{A(r) - 1}{r^2} (xdx + ydy + zdz)^2. \quad (22)$$

The calculations for  $\mu = 0, 1, 2, 3$  and  $i = 1, 2, 3$  for the components of the superpotential  $h_\mu^{0i}$  (needed in (14)) in quasi-Cartesian coordinates are

$$\begin{aligned} h_1^{01} &= h_1^{02} = h_1^{03} = 0, \\ h_2^{01} &= h_2^{02} = h_2^{03} = 0, \\ h_3^{01} &= h_3^{02} = h_3^{03} = 0 \end{aligned} \quad (23)$$

while the non-vanishing components of the superpotential are

$$h_0^{01} = \frac{4Mx}{r^3} \left[ \frac{1}{1 + \gamma \left( \frac{q^2}{Mr} \right)^a} \right]^{3/a}, \quad (24)$$

$$h_0^{02} = \frac{4My}{r^3} \left[ \frac{1}{1 + \gamma \left( \frac{q^2}{Mr} \right)^a} \right]^{3/a}, \quad (25)$$

$$h_0^{03} = \frac{4Mz}{r^3} \left[ \frac{1}{1 + \gamma \left( \frac{q^2}{Mr} \right)^a} \right]^{3/a}. \quad (26)$$

With the aid of the line element (22), the expression for the energy-momentum distribution (14) and the expressions (24)-(26) for the superpotentials, the energy distribution in the Einstein prescription for the new charged non-singular black hole solution satisfying the weak energy condition is obtained as

$$E_E = M \left[ \frac{1}{1 + \gamma \left( \frac{q^2}{Mr} \right)^a} \right]^{3/a}, \quad (27)$$

while, due to (23), we find that all the momentum components vanish:

$$P_x = P_y = P_z = 0. \quad (28)$$




Figure 1: Einstein energy vs.  $r$  for various values of the parameter  $a$  and  $\gamma = 1$ .

In Fig. 1, the energy distribution (27) obtained in the Einstein prescription is plotted as a function of  $r$  for four different values of the parameter  $a$  and  $M = 1$ ,  $q = 0.1$ , and  $\gamma = 1$ .




Figure 2: Einstein energy vs.  $r$  near the origin for various values of the parameter  $a$  and  $\gamma = 1$ .

Fig. 2 exhibits the behaviour of the Einstein energy distribution as a function of  $r$  near the origin for four different values of  $a$  and  $M = 1$ ,  $q = 0.1$ , and  $\gamma = 1$ .

Next, we apply the Møller prescription using Schwarzschild coordinates  $\{t, r, \theta, \phi\}$ , for the line element (4) and the metric function (5). The only non-vanishing component of the Møller superpotential (16) is found to be

$$M_0^{01} = M \left[ \frac{1}{1 + \gamma \left( \frac{q^2}{Mr} \right)^a} \right]^{3/a} \left[ 2 - \frac{6\gamma \left( \frac{q^2}{Mr} \right)^a}{1 + \gamma \left( \frac{q^2}{Mr} \right)^a} \right] \sin \theta \quad (29)$$

while all the other components vanish.

Inserting the above expression (29) of the Møller superpotential into (21), we obtain the energy distribution in the Møller prescription:

$$\begin{aligned} E_M &= M \left[ \frac{1}{1 + \gamma \left( \frac{q^2}{Mr} \right)^a} \right]^{3/a} \left[ 1 - \frac{3\gamma \left( \frac{q^2}{Mr} \right)^a}{1 + \gamma \left( \frac{q^2}{Mr} \right)^a} \right] \\ &= E_E \left[ 1 - \frac{3\gamma \left( \frac{q^2}{Mr} \right)^a}{1 + \gamma \left( \frac{q^2}{Mr} \right)^a} \right], \end{aligned} \quad (30)$$

from which we infer that the energy in the Møller prescription can be given in terms of the energy in the Einstein prescription.

Furthermore, as expected from the vanishing of the spatial components of the Møller superpotential, all the momentum components are found to




Figure 3: Møller energy vs.  $r$  for various values of the parameter  $a$  and  $\gamma = 1$ .




Figure 4: Møller energy vs.  $r$  near the origin for various values of the parameter  $a$  and  $\gamma = 1$ .

be zero:

$$P_r = P_\theta = P_\phi = 0. \quad (31)$$

In Fig. 3 the energy distribution in the Møller prescription for four different values of the parameter  $a$  and  $M = 1$ ,  $q = 0.1$ , and  $\gamma = 1$  is plotted, while in Fig. 4 the behaviour of the Møller energy distribution near the origin is presented for four different values of the parameter  $a$  and  $M = 1$ ,




Figure 5: Energy distributions in the Einstein and Møller prescriptions for  $\gamma = 1$  and  $a = 0.5$ .




Figure 6: Energy distributions in the Einstein and Møller prescriptions for  $\gamma = 1$  and  $a = 0.5$  near origin.

$q = 0.1$ , and  $\gamma = 1$ .

A comparison of the energy distributions in the Einstein and Møller prescriptions is given in Fig. 5 for  $a = 0.5$  and  $M = 1$ ,  $q = 0.1$ , and  $\gamma = 1$ . Additionally, in Fig. 6 an analogous comparison of the energy distributions in the Einstein and Møller prescriptions near the origin is presented for the same values of the parameters  $a$ ,  $M$ ,  $q$ , and  $\gamma$ .

Case	$r \rightarrow 0$	$r \rightarrow \infty$	$q = 0$	$a = 1, \gamma = \frac{1}{6}$
$E_E$	0	$M$	$M$	$M \left( \frac{1}{1 + \frac{q^2}{6Mr}} \right)^3$
$E_M$	0	$M$	$M$	$E_E \left[ 1 - \frac{\frac{q^2}{Mr}}{2 \left( 1 + \frac{q^2}{6Mr} \right)} \right]$

Table 1: Limiting and particular values for the Einstein energy  $E_E$  and the Møller energy  $E_M$ .

## 5 Discussion

The aim of this work is the study of the energy-momentum distribution for a new spherically symmetric and charged, non-singular black hole solution satisfying the weak energy condition.

To this purpose, the Einstein and Møller energy-momentum complexes have been applied and it has been found that all the momenta vanish in both pseudotensorial prescriptions. Additionally, the energy distributions obtained have well-defined expressions showing a dependence on the mass  $M$ , the charge  $q$ , the two parameters  $\gamma$  and  $a$ , introduced in Section 2, and on the radial coordinate  $r$ . Furthermore, the Møller energy is expressed in terms of the Einstein energy. Both energies acquire the same value  $M$  (ADM mass) for  $r \rightarrow \infty$  or for  $q = 0$ .

We have also studied the limiting behavior of the energy distribution for  $r \rightarrow 0$  and  $r \rightarrow \infty$ , and for the particular cases  $q = 0$  and  $a = 1$  and  $\gamma = \frac{1}{6}$ , respectively. In Table 1 we summarize the physically meaningful results for these limiting and particular cases.

At this point some remarks are needed to clarify the results. From Table 1 we conclude that the energy distribution in both prescriptions vanishes near the origin  $r \rightarrow 0$ , while for either  $r \rightarrow \infty$  or  $q = 0$  both energies are equal to the ADM mass  $M$ , in agreement to the result obtained by Virbhadra for the energy distribution of the Schwarzschild black hole solution [50]. For the particular case  $a = 1$  and  $\gamma = \frac{1}{6}$  we get an expression for the energy distribution that depends on the mass  $M$  and the charge  $q$  of the black hole, as well as on the radial coordinate  $r$  for a model of general relativity coupled to nonlinear electrodynamics, the latter corresponding to Maxwell's theory in the weak field approximation. Furthermore, the non-singular black hole solution with  $\beta = \frac{3}{2}$ ,  $a = 1$  and  $\gamma = \frac{1}{6}$  satisfies the weak energy condition and is described by the metric function  $f(r) = 1 - \frac{2M}{r} \left( \frac{1}{1 + \frac{q^2}{6Mr}} \right)^3$ , while asymptotically it behaves as the Reissner-Nordström solution. In fact, for

$r \rightarrow 0$  this non-singular and charged black hole solution exhibits a de Sitter behaviour near zero.




Figure 7: Energy distributions in the Einstein and Møller prescriptions for  $a = 1$  and  $\gamma = \frac{1}{6}$ .




Figure 8: Energy distributions in the Einstein and Møller prescriptions near the origin for  $a = 1$  and  $\gamma = \frac{1}{6}$ .

In Fig. 7 the energy distributions in the Einstein prescription  $E_E$  and in the Møller prescription  $E_M$  are presented, as a function of  $r$ , for the particular case  $a = 1$  and  $\gamma = \frac{1}{6}$ , while in Fig. 8 the same functions for the




Figure 9: Energy distributions in the Einstein and Møller prescriptions near the origin for  $a = 0.7$  and  $\gamma = 1$ .

same values of the parameters are presented near the origin.

Looking closer at the behaviour of the energy distribution near zero, it is seen from Fig. 2 that the Einstein energy tends to zero from positive values, as expected from expression (27), and it is an increasing function of  $r$ . The Møller energy, on the other hand, tends also to zero but, according to Fig. 4, close enough to zero, i.e. for  $r \lesssim 0.04$ , it acquires negative values before reaching zero, a result which is supported by (30). For values of  $r$  greater than 0.04, the Møller energy is positive and increasing.

In Fig. 5 and Fig. 6, we consider the particular value  $a = 0.5$  as an example in order to compare the two energies obtained and we conclude that the Einstein energy  $E_E$  is everywhere greater than the Møller energy. In fact, one can advocate that the positive energy region can be used for the effect of a convergent gravitational lens [70]-[72].

The negativity of the energy distribution in the case of the Møller prescription, for a range of values of  $r$ ,  $a$ ,  $\gamma$ ,  $M$  and  $q$  pinpoints the difficulty of a physically meaningful interpretation of the energy in certain regions. The values of  $r$  where the energy distribution becomes negative depend on the

roots of the equation  $1 - \frac{3\gamma\left(\frac{q^2}{Mr}\right)^a}{1+\gamma\left(\frac{q^2}{Mr}\right)^a} = 0$  (see Eq. (30)). In fact, the energy

becomes negative for  $r^a < 2\gamma\left(\frac{q^2}{M}\right)^a$ . As an example, in Fig. 9 we see that the energy distribution becomes negative for  $0 < r < 0.0275$  if we choose  $a = 0.7$  and  $\gamma = 1$ . For larger values of  $r$ , the energy distribution becomes positive and then it increases.

As pointed out in the Introduction, a modern viewpoint regarding the

energy-momentum localization is represented by a quasi-local energy-momentum associated with a closed 2-surface. The pseudotensors have this property, mostly since Penrose developed his definition of quasi-local mass [45]. Further, according to [47], since a generally accepted expression for the energy-momentum has not been found until today, some important criteria for the quasi-local energy-momentum expressions must be satisfied, basically (a) the expressions should yield the standard values at spatial infinity, and (b) they should not give negative values for the energy. In the case of the black hole solution that satisfies the weak energy condition and is under study in the present work, both the Einstein and Møller energy-momentum complexes do yield the standard values at infinity. In fact, in the Einstein prescription the energy takes only positive values, while in the case of the Møller prescription the energy takes negative values when  $r$  satisfies the inequality  $r^a < 2\gamma(\frac{q^2}{M})^a$ . We believe that this apparent weakness of the Møller prescription can be justified by the properties of the particular metric considered. Indeed, a similar behaviour of the Møller energy-momentum complex was found in [73] and [74]. This strange behaviour is attributed to the particularities of these black hole solutions originating in the coupling of the gravitational field to non-linear electrodynamics.

Although the energy distribution in the Møller prescription does not exhibit the desired behaviour for every value of  $r$ , we can conclude that the two prescriptions used in the present work still constitute instructive tools for the energy-momentum localization. Given the problem that arises from the Møller prescription, it is challenging to employ other pseudo-tensorial prescriptions available, as well as the teleparallel equivalent theory of general relativity, for further investigation of the energy-momentum localization in the context of the four-dimensional spherically symmetric and charged, non-singular black hole solution satisfying the weak energy condition. We consider facing this challenge in a future work.

## References

- [1] L. Bel, “Définition d’une densité d’énergie et d’un état de radiation totale généralisée”, *Comptes Rendus de l’Académie des Sciences*, vol. 246, pp. 3015-3018, 1958.
- [2] I. Robinson, “On the Bel-Robinson tensor”, *Classical and Quantum Gravity*, vol. 14, no. 1, pp. A331-A333, 1997.
- [3] M. A. G. Bonilla and J. M. M. Senovilla, “Some properties of the Bel and Bel-Robinson tensors”, *General Relativity and Gravitation*, vol. 29, no. 1, pp. 91-116, 1997.
- [4] J. M. Senovilla, “Super-energy tensors”, *Classical and Quantum Gravity*, vol. 17, no. 14, pp. 2799-2841, 2000.

- [5] J. D. Brown and J. W. York, “Quasilocal energy and conserved charges derived from the gravitational action”, *Physical Review D*, vol. 47, no. 4, p. 1407, 1993.
- [6] S. A. Hayward, “Quasilocal gravitational energy”, *Physical Review D*, vol. 49, no. 2, pp. 831-839, 1994.
- [7] C.-M. Chen and J.M. Nester, “Quasilocal quantities for general relativity and other gravity theories”, *Classical and Quantum Gravity*, vol. 16, no. 4, pp. 1279-1304, 1999.
- [8] C.-C. M. Liu and S.-T. Yau, “Positivity of quasilocal mass”, *Physical Review Letters*, vol. 90, no. 23, Article ID 231102, 4 pages, 2003.
- [9] L. Balart, “Quasilocal energy, Komar charge and horizon for regular black holes”, *Physics Letters B*, vol. 687, no. 4-5, pp. 280- 285, 2010.
- [10] A. Einstein, “On the general theory of relativity”, *Sitzungsberichte der Königlich Preussischen Akademie der Wissenschaften*, vol. 47, pp. 778-786, 1915, Addendum: *Sitzungsberichte der Königlich Preussischen Akademie der Wissenschaften.*, vol. 47, pp. 799-801, 1915. (Online at <https://einsteinpapers.press.princeton.edu/vol6-doc/>)
- [11] A. Trautman, “Conservation laws in general relativity”, in *Gravitation: an Introduction to Current Research*, L. Witten (ed.), p. 169, John Wiley & Sons, New York, USA, 1962.
- [12] L. D. Landau and E. M. Lifshitz, *The Classical Theory of Fields*, p. 280, Pergamon Press, New York, USA, 1987.
- [13] A. Papapetrou, “Einstein’s theory of gravitation and flat space”, *Proceedings of the Royal Irish Academy. Section A: Mathematical and Physical Sciences*, vol. 52, pp. 11-23, 1948.
- [14] P. G. Bergmann and R. Thomson, “Spin and angular momentum in general relativity”, *Physical Review Letters*, vol. 89, pp. 400-407, 1953.
- [15] C. Møller, “On the localization of the energy of a physical system in the general theory of relativity”, *Annals of Physics*, vol. 4, no. 4, pp. 347-371, 1958.
- [16] S. Weinberg, *Gravitation and Cosmology: Principles and Applications of General Theory of Relativity*, p. 165, John Wiley & Sons, New York, USA, 1972.
- [17] K. S. Virbhadra, “Energy associated with a Kerr-Newman black hole”, *Physical Review D*, vol. 41, no. 4, pp. 1086-1090, 1990.

- [18] K. S. Virbhadra, “Energy distribution in Kerr-Newman spacetime in Einstein’s as well as Møller’s prescriptions”, *Physical Review D*, vol. 42, no. 8, pp. 2919-2921, 1990.
- [19] N. Rosen and K. S. Virbhadra, “Energy and momentum of cylindrical gravitational waves”, *General Relativity and Gravitation*, vol. 25, no. 4, pp. 429-433, 1993.
- [20] K. S. Virbhadra and J. C. Parikh, “A conformal scalar dyon black hole solution”, *Physics Letters B*, vol. 331, 3-4, pp. 302-304, 1994.
- [21] I. Radinschi, “Energy associated with the Bianchi type VI<sub>0</sub> universe”, *Chinese Journal of Physics*, vol. 39, no. 5, pp. 393-396, 2001.
- [22] P. K. Sahoo, K. L. Mahanta, D. Goit, A.K. Sinha, S.S. Xulu, U.R. Das, A. Prasad and R. Prasad, “Einstein energy-momentum complex for a phantom black hole metric”, *Chinese Physics Letters*, vol. 32, no. 2, Article ID 020402, 2015.
- [23] S. K. Tripathy, B. Mishra, G. K. Pandey, A. K. Singh, T. Kumar, and S. S. Xulu, “Energy and momentum of Bianchi type VI<sub>h</sub> universes”, *Advances in High Energy Physics*, vol. 2015, Article ID 705262, 8 pages, 2015.
- [24] I.-C. Yang and I. Radinschi, “On the difference of energy between the Einstein and Møller prescription”, *Chinese Journal of Physics*, vol. 42, no. 1, p. 40, 2004.
- [25] I. Radinschi and T. Grammenos, “Møller’s energy-momentum complex for a spacetime geometry on a noncommutative curved D3-brane”, *International Journal of Theoretical Physics*, vol. 47, no. 5, pp. 1363-1372, 2008.
- [26] I.-C. Yang, C.-L. Lin, and I. Radinschi, “The energy of a regular black hole in general relativity coupled to nonlinear electrodynamics”, *International Journal of Theoretical Physics*, vol. 48, no. 1, pp. 248-255, 2009.
- [27] I. Radinschi, F. Rahaman, and A. Ghosh, “On the energy of charged black holes in generalized dilaton-axion gravity”, *International Journal of Theoretical Physics*, vol. 49, no. 5, pp. 943-956, 2010.
- [28] I. Radinschi, F. Rahaman, and A. Banerjee, “On the energy of Hořava-Lifshitz black holes”, *International Journal of Theoretical Physics*, vol. 50, no. 9, pp. 2906-2916, 2011.
- [29] I. Radinschi, F. Rahaman, and A. Banerjee, “The energy distribution of Hořava-Lifshitz black hole solutions”, *International Journal of Theoretical Physics*, vol. 51, no. 5, pp. 1425-1434, 2012.

- [30] M. Abdel-Megied and R. M. Gad, “Møller’s energy in the Kantowski-Sachs space-time”, *Advances in High Energy Physics*, vol. 2010, Article ID 379473, 6 pages, 2010.
- [31] T. Bringley, “Energy and momentum of a stationary beam of light”, *Modern Physics Letters A*, vol. 17, no. 3, pp. 157-161, 2002.
- [32] I. Radinschi, F. Rahaman, and U. F. Mondal, “Energy distribution for non-commutative radiating Schwarzschild black holes”, *International Journal of Theoretical Physics*, vol. 52, no. 1, pp. 96-104, 2013.
- [33] M. Sharif and Tasnim Fatima, “Energy distribution associated with static axisymmetric solutions”, *Astrophysics and Space Science*, vol. 302, no. 1-4, pp. 217224, 2006.
- [34] M. Sharif and M. Azam, “Energy-momentum distribution: some examples”, *International Journal of Modern Physics A*, vol. 22, no. 10, pp. 1935-1951, 2007.
- [35] P. Halpern, “Energy of the Taub cosmological solution”, *Astrophysics and Space Science*, vol. 306, no. 4, pp. 279-283, 2006.
- [36] E. C. Vagenas, “Energy distribution in 2d stringy black hole backgrounds”, *International Journal of Modern Physics A*, vol. 18, no. 31, pp. 5781-5794, 2003.
- [37] E. C. Vagenas, “Effective mass of a radiating charged particle in Einstein’s universe”, *Modern Physics Letters A*, vol. 19, no. 3, pp. 213-222, 2004.
- [38] E. C. Vagenas, “Energy distribution in a BTZ black hole spacetime”, *International Journal of Modern Physics D*, vol. 14, no. 03n04, pp. 573-585, 2005.
- [39] E. C. Vagenas, “Energy distribution in the dyadosphere of a Reissner-Nordström black hole in Møller’s prescription”, *Modern Physics Letters A*, vol. 21, no. 25, pp. 1947-1956, 2006.
- [40] T. Multamaki, A. Putaja, E. C. Vagenas, and I. Vilja, “Energy-momentum complexes in  $f(R)$  theories of gravity”, *Classical and Quantum Gravity*, vol. 25, no. 7, Article ID 075017, 2008.
- [41] L. Balart, “Energy distribution of (2+1)-dimensional black holes with nonlinear electrodynamics”, *Modern Physics Letters A*, vol. 24, no. 34, pp. 2777-2785, 2009.
- [42] A. M. Abbassi, S. Mirshekari, and A. H. Abbassi, “Energy-momentum distribution in static and nonstatic cosmic string space-times”, *Physical Review D*, vol. 78, no. 6, Article ID 064053, 2008.

- [43] Gamal G. L. Nashed, “Møller’s Energy of Kerr-NUT Metric”, *Chinese Physics Letters*, vol. 25, no. 4, pp. 1202-1204, 2008.
- [44] J. Matyjasek, “Some remarks on the Einstein and Møller pseudotensors for static and spherically-symmetric configurations”, *Modern Physics Letters A*, vol. 23, no. 8, pp. 591-601, 2008.
- [45] R. Penrose, “Quasi-local mass and angular momentum in general relativity”, *Proceedings of the Royal Society of London A: Mathematical, Physical and Engineering Sciences*, vol. 381, no. 1780, pp. 53-63, 1982.
- [46] K. P. Tod, “Some examples of Penrose’s quasilocal mass construction”, *Proceedings of the Royal Society of London A: Mathematical, Physical and Engineering Sciences*, vol. 388, no. 1795, pp. 457-477, 1983.
- [47] L. B. Szabados, “Quasi-local energy-momentum and angular momentum in general relativity”, *Living Reviews in Relativity*, vol. 12, p. 4, 2009.
- [48] J. M. Aguirregabiria, A. Chamorro, and K. S. Virbhadra, “Energy and angular momentum of charged rotating black holes”, *General Relativity and Gravitation*, vol. 28, no. 11, pp. 1393-1400, 1996.
- [49] S. S. Xulu, “Bergmann-Thomson energy-momentum complex for solutions more general than the Kerr-Schild class”, *International Journal of Theoretical Physics*, vol. 46, no. 11, pp. 2915- 2922, 2007.
- [50] K. S. Virbhadra, “Naked singularities and Seifert’s conjecture”, *Physical Review D*, vol. 60, no. 10, Article ID 104041, 1999.
- [51] C. Møller, “The four-momentum of an insular system in general relativity”, *Nuclear Physics*, vol. 57, pp. 330-338, 1964.
- [52] K. Hayashi and T. Shirafuji, “New general relativity”, *Physical Review D* vol. 19, p. 3524, 1979.
- [53] J. M. Nester, Lau Loi So, and T. Vargas, “Energy of homogeneous cosmologies”, *Physical Review D* vol. 78, p. 044035, 2008.
- [54] Gamal G. L. Nashed, “Kerr-Newman solution and energy in teleparallel equivalent of Einstein theory”, *Modern Physics Letters A*, vol. 22, no. 14, pp. 1047-1056, 2007.
- [55] Gamal G. L. Nashed, “Energy and angular momentum of general four-dimensional stationary axi-symmetric space-time in teleparallel geometry”, *International Journal of Modern Physics A*, vol. 23, no. 12, p.19031918, 2008.

- [56] Gamal G. L. Nashed, “Energy of spherically symmetric space-times on regularizing teleparallelism”, *International Journal of Modern Physics A*, vol. 25, no. 14, pp. 28-83, 2010.
- [57] M. Sharif, and Abdul Jawad, “Energy contents of some well-known solutions in teleparallel gravity”, *Astrophysics and Space Science*, vol. 331, no. 1, pp. 257-263, 2011.
- [58] J. W. Maluf, M. V. O. Veiga, and J. F. da Rocha-Neto, “Regularized expression for the gravitational energy-momentum in teleparallel gravity and the principle of equivalence”, *Classical and Quantum Gravity*, vol. 39, no. 3, pp. 227-240, 2007.
- [59] M. Sharif and S. Taj, “Energy contents of some non-vacuum spacetimes in teleparallel gravity”, *Astrophysics and Space Science*, vol. 325, no. 1, pp. 325-375, 2010.
- [60] A. Sousa, R. B. Pereira and A. C. Silva, “Energy and angular momentum densities in a Gödel-type universe in teleparallel geometry”, *Gravitation and Cosmology*, vol. 16, no. 1, pp. 25-33, 2010.
- [61] S. Aygün, H. Baysal, C. Aktaş, I. Yılmaz, P. K. Sahoo and I. Tarhan, “Teleparallel energy-momentum distribution of various black hole and wormhole metrics”, *International Journal of Modern Physics A*, vol. 33, no. 30, p. 1850184, 2018.
- [62] M. G. Ganiou, M. J. S. Houndjo and J. Tossa,  $f(T)$  gravity and energy distribution in Landau-Lifshitz prescription, *International Journal of Modern Physics D*, vol. 27, no. 04, p. 1850039, 2018.
- [63] M.-T. Wang and S.-T. Yau, “Quasilocal Mass in General Relativity”, *Physical Review Letters*, vol. 102, p. 021101, 2009.
- [64] M.-T. Wang and S.-T. Yau, “Isometric Embeddings into the Minkowski Space and New Quasi-Local Mass”, *Communications in Mathematical Physics*, vol. 288, no. 3, pp. 919942, 2009.
- [65] C.-M. Chen, J.-L. Liu and J. M. Nester, “Gravitational energy is well defined”, *International Journal of Modern Physics D*, vol. 27 p.1847017, 2018.
- [66] C.-M. Chen, J.-L. Liu and J. M. Nester, “Quasi-local energy from a Minkowski reference”, *General Relativity and Gravitation*, vol. 50, no.12, 158, 2018.
- [67] L. Balart and E. C. Vagenas, “Regular black holes with a non linear electrodynamics source” *Physical Review D* vol. 90, p. 124045, 2014.

- [68] C. Dagum, “A new model of personal income distribution: Specification and estimation”, *Economie Appliquée*, vol. 30, p. 413437, 1977.
- [69] L. Balart and E. C. Vagenas, “Regular black hole metrics and the weak energy condition” *Physic Letters B* vol. 730, pp. 14-17, 2014.
- [70] C.-M. Claudel, K. S. Virbhadra, and G. F. R. Ellis, “The geometry of photon surfaces” *Journal of Mathematical Physics*, vol. 42, no. 2, p. 818-838, 2001.
- [71] K. S. Virbhadra and G. F. R. Ellis, “Schwarzschild black hole lensing”, *Physical Review D*, vol. 62, no. 8, p. 084003. 2000.
- [72] K. S. Virbhadra, “Relativistic images of Schwarzschild black hole lensing”, *Physical Review D*, vol. 79, no. 8, p. 083004, 2009.
- [73] I. Radinschi, Th. Grammenos, F. Rahaman, A. Spanou, S. Islam, S. Chattopadhyay and A. Pasqua, “Energy-momentum for a charged non-singular black hole solution with a nonlinear mass function”, *Advances in High Energy Physics*, vol.2017, Article ID 7656389, pp.1-11, 2017.
- [74] I. Radinschi, F. Rahaman, Th. Grammenos and S. Islam, “Einstein and Møller energy-momentum complexes for a regular black hole solution with a nonlinear electrodynamics source”, *Advances in High Energy Physics*, vol. 2016, Article ID 9049308, pp.1-9, 2016.