

CPT-violating Effects on Neutron Gravitational Bound States

Zhi Xiao

*Department of Mathematics and Physics, North China Electric Power University,
Beijing 102206, China*

Analytical solutions with effective CPT-violating spin-gravity corrections to the neutron's gravitational bound states are obtained. The helicity-dependent phase evolution due to $\vec{\sigma} \cdot \vec{b}$ and $\vec{\sigma} \cdot \hat{p}$ couplings not only leads to spin precession, but also to transition-frequency shifts between different gravitational bound states. Utilizing transition frequencies measured in the qBounce experiment,¹ we obtain the rough bound $|\vec{b}| < 6.9 \times 10^{-21} \text{ GeV}$. Incorporating known systematic errors may lead to more robust and tighter constraints.

1. Introduction

Lorentz-violating matter-gravity couplings² open a broad and interesting avenue for testing Lorentz symmetry. Recently, spin-independent Lorentz-violating neutron-gravity couplings have been thoroughly studied³ in an attempt to analyze the GRANIT experiment.⁴ However, to our best knowledge, an extensive study of spin-dependent fermion-gravity couplings is still under development.⁵ Here, we provide a first glimpse at CPT-violating spin-dependent neutron-gravity couplings. A more detailed and complete analysis can be found in Ref. 6.

2. LV corrections due to spin-dependent interactions

The main vertical hamiltonian after averaging over the horizontal degrees of freedom is³

$$\hat{H} = \frac{\hat{p}_z^2}{2m_I} + m_G g z - \vec{\sigma} \cdot \vec{b} (1 + \Phi_0), \quad (1)$$

where we have started with the hamiltonian in Ref. 7 and performed a series of redefinitions and approximations. The stationary solution of (1) is

$$\Psi_{\perp}(t, z) = \frac{1}{\sqrt{2}} \begin{pmatrix} [\cos(\frac{\Omega}{2}t) + ir_+ \sin(\frac{\Omega}{2}t)] e^{-\frac{i\omega}{2}t} \\ [\cos(\frac{\Omega}{2}t) - ir_- \sin(\frac{\Omega}{2}t)] e^{+\frac{i\omega}{2}t} \end{pmatrix} \phi_n(z) e^{-iE_n t}, \quad (2)$$

where $\phi_n(z) \equiv \frac{\text{Ai}[\frac{z}{L_c} - x_{n+1}]}{L_c^{1/2} |\text{Ai}'[-x_{n+1}]|}$, $\Omega \equiv \sqrt{\omega^2 + 4\omega B_0 \cos \theta + 4B_0^2}$, and $r_{\pm} \equiv [\omega + 2B_0(\cos \theta \pm \sin \theta e^{-i\phi})] / \Omega$; the initial state is assumed to be an eigenstate of σ_x , $|X \uparrow\rangle$.




Fig. 1. (a) Probability to remain in the initial state $|X \uparrow\rangle$. We have chosen an unrealistically large $B_0 = \pi/120$ to make the probability variation apparent. The other parameters are $\theta = \pi/3$ and $\omega = 0$; the ϕ dependence is clearly discernible from the three curves. (b) Evolution of the spin vector for the initial state $|X \uparrow\rangle$. Again, an unrealistically large $B_0 = \pi/13$ has been chosen. The other parameters are $\theta = \pi/6$ and $\phi = \omega = 0$.

The probability profile for unrealistically large $B_0 \equiv |\vec{b}|$ as well as the spin-precession on the Bloch sphere are shown in Fig. 1. The eigensolution of $\hat{H} = \hat{p}_z^2/(2m_I) + m_G g z - \bar{b}_{\text{eff}} (1 + \Phi_0) \sigma_z \hat{p}_z / m_I$ is

$$\Psi(t, z) = \begin{pmatrix} c_1 e^{i[\bar{b}_{\text{eff}}(1+\Phi_0)]z} \\ c_2 e^{-i[\bar{b}_{\text{eff}}(1+\Phi_0)]z} \end{pmatrix} \text{Ai}\left[\frac{z}{L_c} - x_{n+1}\right] e^{-i\{E_n - \frac{[\bar{b}_{\text{eff}}(1+\Phi_0)]^2}{2m_I}\}t}, \quad (3)$$

where the parity-odd nature of $\vec{\sigma} \cdot \hat{p}$ again dictates opposite phase evolutions for the two helicity components, and c_1 and c_2 are constants to be determined.

For $\hat{H}_{LV} = -\vec{\sigma} \cdot \vec{b} (1 + \Phi_0 + g z) - \bar{b}_{\text{eff}} (1 + \Phi_0) \sigma_z \hat{p}_z / m_I$, we can use the matrix elements

$$\begin{aligned} & \begin{pmatrix} \langle n+ | \hat{H}_{LV} | n, + \rangle & \langle n+ | \hat{H}_{LV} | n, - \rangle \\ \langle n- | \hat{H}_{LV} | n, + \rangle & \langle n- | \hat{H}_{LV} | n, - \rangle \end{pmatrix} \\ &= -B_0 \left[(1 + \Phi_0) + \frac{2}{3} g L_c x_{n+1} \right] \begin{pmatrix} \cos \theta & \sin \theta e^{-i\phi} \\ \sin \theta e^{i\phi} & -\cos \theta \end{pmatrix} \end{aligned} \quad (4)$$

to calculate the shift in the eigenenergies

$$\delta E_n = \mp B_0 \left[(1 + \Phi_0) + \frac{2}{3} g L_c x_{n+1} \right], \quad (5)$$

where the upper or lower sign depends on whether the spin state is $|\hat{n}+\rangle = (e^{-i\phi} \cos \frac{1}{2}\theta, \sin \frac{1}{2}\theta)$ or $|\hat{n}-\rangle = (\sin \frac{1}{2}\theta, -e^{i\phi} \cos \frac{1}{2}\theta)$, respectively. From (5), the transition-frequency shift is given by $\delta \nu_{mn}^\pi = \mp \frac{2g}{3\hbar} |\vec{b}| L_c (x_{m+1} - x_{n+1})$. Comparing $\delta \nu_{mn}^\pi$ with the precisely measured frequencies in the qBounce experiment,¹ we obtain the rough upper bound $|\vec{b}| < 6.946 \times 10^{-21}$ GeV.

3. Summary

In this work, we have discussed CPT-violating spin-gravity corrections on the neutron's gravitational bound states. With several analytical solutions, we have demonstrated that the phase evolution depends on the helicity of the wave-function components. The resulting phenomena are spin precession and θ - and ϕ -dependent probability variations. Using degenerate perturbation theory, we have also calculated the transition-frequency shift. Comparison with the measurements in Ref. 1 has yielded the rough bound $|\vec{b}| < 6.9 \times 10^{-21}$ GeV, which can be improved further if systematic errors from known physics are taken into account, or if polarized neutrons are used in the future.

Acknowledgments

The author appreciates valuable encouragement and helpful discussions with M. Snow and A. Kostelecký as well as help from many others. This work is partially supported by the National Science Foundation of China under grant no. 11605056, no. 11875127, and no. 11575060.

References

1. G. Cronenberg *et al.*, Nature Phys. **14**, 1022 (2018).
2. V.A. Kostelecký and J.D. Tasson, Phys. Rev. Lett. **102**, 010402 (2009); Phys. Rev. D **83**, 016013 (2011).
3. A. Martín-Ruiz and C.A. Escobar, Phys. Rev. D **97**, 095039 (2018).
4. V.V. Nesvizhevsky *et al.*, Nature **415**, 297 (2002).
5. V.A. Kostelecký and Z. Li, to appear. See also Z. Li, these proceedings.
6. Z. Xiao, arXiv:1906.00146 [hep-ph].
7. Y. Bonder, Phys. Rev. D **88**, 105011 (2013).