

BPS solutions for generalised Wess-Zumino models and their applications

Steven Abel^{a,b}, Quentin Bonnefoy^c and Debtosh Chowdhury^{c,d}

^a*Institute for Particle Physics Phenomenology, Durham University, South Road, Durham, U.K.*

^b*Department of Mathematical Sciences, Durham University, South Road, Durham, U.K.*

^c*Centre de Physique Théorique, CNRS, École Polytechnique, IP Paris, 91128 Palaiseau, France*

^d*Laboratoire de Physique Théorique (UMR8627), CNRS, Univ. Paris-Sud, Université Paris-Saclay, 91405 Orsay, France*

E-mail: s.a.abel@durham.ac.uk, quentin.bonnefoy@polytechnique.edu,
debtosh.chowdhury@polytechnique.edu

ABSTRACT: We present BPS solutions to a general class of Wess-Zumino models which extend previous results in the literature. We discuss their relation to amplitudes on threshold, and their application to scalar domain walls in Supersymmetric QCD. We also find partial expressions for Wess-Zumino models with softly broken supersymmetry.

Contents

1	Introduction and summary	1
2	Multiparticle amplitudes in generalised Wess-Zumino models	3
2.1	Recursion relations and classical solutions	3
2.2	Soft terms in the Wess-Zumino model	6
2.3	Towards a solution for symmetric soft masses	7
3	SQCD with quartic couplings	8
4	Conclusion	10
A	The BPS condition	11
B	Link with softly broken $O(2)$ models	12
C	Derivation of the solution	12

1 Introduction and summary

The purpose of this paper is to present and discuss a rather general solution to the Bogomol’nyi-Prasad-Sommerfield (BPS) equations, for a rather general class of Wess-Zumino (WZ) models. As we shall see the solution has applications in several areas, including multiparticle amplitudes on threshold, and scalar domain walls in Supersymmetric QCD (SQCD) duality.

Consider the following superpotential for a chiral superfield Φ :

$$W = \frac{1}{2}\Phi^2 + \frac{1}{p}\Phi^p, \quad (1.1)$$

where we do not place a restriction on the allowed value of the index p (except $p > 2$), and where couplings can be trivially reinstated by scaling. The associated scalar potential (where ϕ is the scalar component) is

$$V(\phi) = |\phi + \phi^{p-1}|^2, \quad (1.2)$$

and if p is positive one might seek domain wall solutions between the supersymmetric minimum at $\phi = 0$ and the $p - 2$ supersymmetric minima at $\phi = e^{i\frac{n\pi}{p-2}}$, $n \in \mathbb{Z}$. Because the potential is a complete square, the equations of motion can be integrated once and factorised, yielding the familiar BPS equation (see Appendix A for a brief discussion of the latter):

$$\frac{d\phi}{dt} = e^{2i\theta}(\bar{\phi} + \bar{\phi}^{p-1}), \quad (1.3)$$

where t is the coordinate across the wall and θ is an arbitrary constant angle. If we restrict ϕ to be real then solving eq. (1.3) is trivial, however the conjugation on the right hand side makes it difficult to find the general complex solution for arbitrary p . Our central result is the following solution to eq. (1.3):

$$\phi(z, \bar{z}) = \frac{z \left(1 + \frac{\bar{z}^{p-2} - z^{p-2}}{2p} \right)}{\left(\left(1 + \frac{\bar{z}^{p-2} - z^{p-2}}{2p} \right)^p + \frac{\bar{z}^{p-2} \left(\left(1 - \frac{\bar{z}^{p-2} - z^{p-2}}{2p} \right)^p - \left(1 + \frac{\bar{z}^{p-2} - z^{p-2}}{2p} \right)^p \right)}{\bar{z}^{p-2} - z^{p-2}} \right)^{\frac{1}{p-2}}}, \quad (1.4)$$

where $z = e^{t+i\theta}$ (see Appendix C for a few words on the derivation).

This is a generalisation of the BPS domain wall solution of Ref.[1] (with appropriate shifts in ϕ) which considered $p = 3$ and real ϕ . Indeed taking $\theta = \frac{\pi}{p-2}$ we find

$$\phi(t) = \left(\frac{-e^{(p-2)t}}{1 + e^{(p-2)t}} \right)^{\frac{1}{p-2}}, \quad (1.5)$$

which reduces, for $p = 3$, to the non-singular domain wall solution,

$$\phi(t) = -\frac{e^t}{1 + e^t}, \quad (1.6)$$

connecting the two minima ($\phi(-\infty) = 0$ and $\phi(\infty) = -1$) of the WZ model. As an illustration, in Figure 1 we plot the generalised BPS solution as given in eq. (1.4) by setting $p = 3$. There, we see that, even though $\phi(z, \bar{z})$ is singular for most values of θ , there exist




Figure 1. Plot of the solution in eq. (1.4) when $p = 3$. In the left panel, the colour bar denotes the argument of the function $\phi(z, \bar{z})$. In the right panel, $\theta = \pi$ as in eq. (1.6).

smooth configurations (along the dashed lines) that correspond to domain walls connecting two minima of the function $V(\phi)$, consistent with eqs. (1.5) and (1.6). For $p = 3$, when $\theta = \pm\pi$, the domain wall connects the two minima at $\phi(-\infty) = 0$ and $\phi(\infty) = -1$ passing through $t = 0$, as shown in the right panel of Figure 1.

The expression in eq. (1.4) is also related to the softly broken $O(2)$ models of Ref.[2] that were examined in the context of multiparticle amplitudes on threshold (which took $p = 3$, see Appendix B), and some other work in this area (which typically considered real ϕ). However, the solution above has a richer structure and is more general than those that have been previously considered in the literature. Indeed to derive it we imposed only that ϕ scales as e^t as t goes to infinity, which is enough/required for amplitudes.

In the following section we discuss the application of our solution in the amplitude context, with particular emphasis on the recursion relations of multiparticle amplitudes on threshold, and their relationship with classical solutions of the equations of motion. After spending some time reviewing and discussing the classical ways of obtaining these amplitudes in the WZ model, we demonstrate that the general complex solution presented above translates into the ability to distinguish chiral fields and their conjugates in the possible final multiparticle states.

The arbitrariness of the exponent p also makes eq. (1.4) applicable to situations in which the second term in the superpotential of eq. (1.1) is generated non-perturbatively. In Section 3 we show that this allows one to find exact (classical) domain wall solutions for the scalar mesons in the magnetic duals of Supersymmetric QCD theories with a quartic coupling. In $SU(N_c)$ theories with N_f flavours of quark/antiquark, this is relevant in the free-magnetic window, where $N_c + 1 < N_f < \frac{3}{2}N_c$. The exponent is given by $p = N_f/(N_f - N_c)$, so that p is generally a rational number between 3 and $N_f/2$. These non-perturbatively generated domain walls interpolate between two supersymmetric minima, going from the unbroken magnetic dual at the origin, to one of $2N_c - N_f$ pure Yang-Mills minima with meson vacuum expectation values (VEVs). This configuration is of general interest, and would appear for example in the duality cascade.

2 Multiparticle amplitudes in generalised Wess-Zumino models

Multiparticle amplitudes have been investigated for a long time [2–10], and have been the subject of renewed scrutiny recently within discussions of the so-called Higgspllosion mechanism [11–14]. The quantities of interest include the tree-level threshold amplitudes, which describe the decay of an off-shell particle to many on-shell ones, all taken to be at rest. Our solution in eq. (1.4) can be understood in this respect as the generating function of such tree-level multiparticle amplitudes at kinematic threshold for the generalised Wess-Zumino models of eq. (1.1). One can indeed show that such a generating function must satisfy a BPS condition (see Appendix A for more details), consistent with the fact that a specific limit of eq. (1.4) has been previously identified as a BPS domain wall solution [1, 15]. As we will also see, eq. (1.4) can be extended to softly broken SUSY scenarios, yielding either a complete or a partial solution depending on the choice of soft terms.

2.1 Recursion relations and classical solutions

In order to review standard techniques while simultaneously applying them to our specific problem, we will begin this section by following a diagrammatic approach to tree-level multiparticle amplitudes at kinematic threshold before linking it to classical solutions of

the equations of motion. We will then show that eq. (1.4) indeed generates the amplitudes for the model of eq. (1.1), for specified numbers of emitted particles/anti-particles. In the next section we will extend the discussion to WZ models with specific sets of soft terms.

We are interested in evaluating tree-level amplitudes connecting an ingoing off-shell particle to outgoing on-shell ones, all taken to be at rest¹, for generalised Wess-Zumino models of a chiral superfield Φ . Let us take a canonical Kähler potential and for this discussion reinstate the couplings in the superpotential,

$$W = \frac{M}{2}\Phi^2 + \frac{\lambda}{p}\Phi^p, \quad (2.1)$$

where $p-3 \in \mathbb{N}$, giving rise to the following scalar potential for the complex scalar excitation ϕ :

$$V(\phi) = |M\phi + \lambda\phi^{p-1}|^2. \quad (2.2)$$

The kinematic situation is summed up in Figure 2. Since there are two possible kinds of scalar excitation, the outgoing state is labelled by two integers m and n , denoting the number of particles and antiparticles respectively.




Figure 2. Kinematic setup (particles/anti-particles are represented using direct/reversed arrows).

The WZ model also includes scalar-fermion interactions. However, since we will be interested in tree-level amplitudes with initial and final states only made out of scalars, those interactions (which preserve fermion number) will not play any role.

Following earlier works on multiparticle amplitudes [3, 4], one can recursively calculate such amplitudes following the scheme of Figure 3. From this, after working out the correct combinatorics, one finds the following recursion relation:

$$\begin{aligned} \frac{a_{nm}}{m!n!} = & - \frac{(p-1)i^{2p-2}|\lambda|^2}{|M|^{2(2p-3)}} \sum_{\substack{\sum n_i = n \\ \sum m_i = m}} \frac{b_{n_1 m_1} b_{n_2 m_2} \dots b_{n_{p-2} m_{p-2}} a_{n_{p-1} m_{p-1}} \dots a_{n_{2p-3} m_{2p-3}}}{\prod_{i=1..2p-3} n_i! m_i! [(n_i + m_i)^2 - 1]} \\ & - \frac{i^p}{|M|^{2(p-1)}} \sum_{\substack{\sum n_i = n \\ \sum m_i = m}} \frac{\lambda \bar{M} a_{n_1 m_1} a_{n_2 m_2} \dots a_{n_{p-1} m_{p-1}} + (p-1) \bar{\lambda} M a_{n_1 m_1} b_{n_2 m_2} \dots b_{n_{p-1} m_{p-1}}}{\prod_{i=1..p-1} n_i! m_i! [(n_i + m_i)^2 - 1]}, \end{aligned} \quad (2.3)$$

where a_{nm} symbolises the amplitude $\phi \longrightarrow n \times \phi + m \times \bar{\phi}$ and b_{nm} the amplitude $\bar{\phi} \longrightarrow n \times \phi + m \times \bar{\phi}$. Viewing a_{nm} as a function of λ and M , we can immediately deduce that $b_{nm}(\lambda, M) = a_{mn}(\bar{\lambda}, \bar{M})$ since $V(\phi)$ is hermitian.

¹Exact results are much harder to obtain at loop-level or in the out of threshold regime [7–10, 16–18].




Figure 3. Recursion scheme for the amplitudes, drawn here for $p = 3$.

Inspection of the lowest amplitudes shows that the recursion is correctly initialised by imposing the following conditions:

$$\left. \frac{a_{nm}}{[(n+m)^2 - 1]} \right|_{n=1, m=0} = -i|M|^2, \quad \left. \frac{a_{nm}}{[(n+m)^2 - 1]} \right|_{n=0, m=1 \text{ or } n=0, m=0} = 0, \quad (2.4)$$

which, combined with eq. (2.3), imply that $b_{nm} = -\overline{a_{mn}}$.

A convenient factorisation can be performed:

$$a_{nm} = -i|M|^2 A_{nm} n!m! [(n+m)^2 - 1] \left(\frac{\lambda}{M} \right)^{\frac{n-1}{p-2}} \left(\frac{\bar{\lambda}}{\bar{M}} \right)^{\frac{m}{p-2}} \quad (2.5)$$

with coefficients A_{nm} satisfying

$$\begin{cases} ((n+m)^2 - 1)A_{nm} = & (p-1) \sum A_{m_1 n_1} \dots A_{m_{p-2} n_{p-2}} A_{n_{p-1} m_{p-1}} \dots A_{n_{2p-3} m_{2p-3}} \\ & + \sum (A_{n_1 m_1} \dots A_{n_{p-1} m_{p-1}} + (p-1)A_{n_1 m_1} A_{m_2 n_2} \dots A_{m_{p-1} n_{p-1}}) , \\ A_{10} = 1, \quad A_{01} = A_{00} = 0, \end{cases} \quad (2.6)$$

where the summations over indices match those in eq. (2.3). In particular, it implies that all A_{nm} are real and positive. The fact that all coupling constants disappeared from the above relation is a consequence of the (R-)symmetries of eq. (2.1) and of holomorphicity². Defining a generating function

$$A(z, \bar{z}) = \sum_{n,m} A_{nm} z^n \bar{z}^m, \quad (2.7)$$

²Indeed, the effective superpotential generating tree-level diagrams can only take the form

$$W = \frac{M}{2} \Phi^2 \sum_n \left(\frac{\lambda \Phi^{p-2}}{M} \right)^n,$$

and each amplitude has a dependence on λ, M fixed by this expression.

the recursion yields the differential equation³

$$\begin{cases} [(z\partial_z + \bar{z}\partial_{\bar{z}})^2 - 1]A = (p-1)A^{p-1}\bar{A}^{p-2} + A^{p-1} + (p-1)A\bar{A}^{p-2} \\ A(z=0, \bar{z}) = 0, \quad \partial_z A(0,0) = 1, \quad \partial_{\bar{z}} A(0,0) = 0 \end{cases}. \quad (2.8)$$

Finally, defining $z = e^{t+i\theta}$, this system becomes

$$\begin{cases} \partial_t^2 A = (p-1)A^{p-1}\bar{A}^{p-2} + A^{p-1} + (p-1)A\bar{A}^{p-2} + A = \frac{\partial}{\partial A}V(A) \\ A(t=-\infty, \theta) = 0, \quad \partial_t A(t=-\infty, \theta) = e^{i\theta} \end{cases}, \quad (2.9)$$

where the potential is as in eq. (1.2). The last equality in the first line illustrates the method of classical solutions of Ref.[6], which states that tree-level multiparticle amplitudes can be derived from the expansion of classical solutions with specific initial conditions. Besides, integrating it once and taking the square root yields the condition in eq. (1.3).

One can verify that our solution in eq. (1.4) indeed satisfies all of the conditions listed in eq. (2.9). Consequently $\phi \equiv A$ is the generating function of the diagrams of Figure 2. That is, Taylor expanding it with respect to z and \bar{z} yields the amplitudes A_{nm} .

2.2 Soft terms in the Wess-Zumino model

Given the generality of the solution in eq. (1.4), it is natural to ask if one might be able to extend the analysis to non-supersymmetric cases, by deforming the theory with supersymmetry breaking operators. The renormalizable $p = 3$ WZ model allows for the following soft terms [19] in addition to the supersymmetric potential:

$$V = |\lambda\phi^2 + m\phi|^2 + \delta m^2|\phi|^2 + (\mu_3\phi^3 + \mu_2\phi^2 + h.c.), \quad (2.10)$$

which can be expressed in terms of real parameters by defining $\mu_3 = c_3 + id_3$, $\mu_2 = c_2 + id_2$ and $\phi = \varphi + i\chi$ as

$$\begin{aligned} V = & \lambda^2(\varphi^2 + \chi^2)^2 + (2\lambda m + 2c_3)\varphi^3 - 6d_3\varphi^2\chi + (2\lambda m - 6c_3)\varphi\chi^2 + 2d_3\chi^3 \\ & + (m^2 + \delta m^2 + 2c_2)\varphi^2 - 4d_2\varphi\chi + (m^2 + \delta m^2 - 2c_2)\chi^2, \end{aligned} \quad (2.11)$$

where we take λ and m to be real by making a suitable U(1) rotation on ϕ .

Specific choices for the soft terms can be related to the softly broken O(2) model described in Appendix B. Starting with eq. (B.2) and performing rotations and shifts on φ and χ , one can only generate

$$V = |\lambda\phi^2 + m\phi|^2 + \frac{\delta m^2}{2} \left(\frac{\phi - \bar{\phi}}{2i} \right)^2. \quad (2.12)$$

³The first condition on the second row is a slight generalisation of $A_{01} = A_{00} = 0$ since, due to the ϕ dependence of $V(\phi)$, the number of ϕ or $\bar{\phi}$ can only increase in a tree-level diagram.

Then implementing the same series of rotations and shifts on the solution obtained for the softly broken $O(2)$ model yields a classical solution of the model in eq. (2.12):

$$\phi(z, \bar{z}) = \frac{z + \frac{\lambda}{m}i(z - \bar{z}) \frac{i(z - \bar{z}) + i\left(\sqrt{2\frac{m^2 \text{Im}(\phi)}{|m|}} - 1\right)^2 (z + \bar{z})}{4\left(2\frac{m^2 \text{Im}(\phi)}{m^2} - 1\right)}}{1 - \frac{\lambda}{m} \frac{z + \bar{z}}{2} + \left(\frac{\lambda}{m}\right)^2 \frac{(z - \bar{z})^2}{4\left(2\frac{m^2 \text{Im}(\phi)}{m^2} - 1\right)} - \left(\frac{\lambda}{m}\right)^3 \frac{\left(\sqrt{2\frac{m^2 \text{Im}(\phi)}{|m|}} - 1\right)^4 (z - \bar{z})^2 (z + \bar{z})}{8\left(2\frac{m^2 \text{Im}(\phi)}{m^2} - 1\right)^3}}, \quad (2.13)$$

where $m_{\text{Im}(\phi)}^2 = 2m^2 + \delta m^2$. It reduces to eqs. (1.4) and (B.1), if $\delta m^2 = 0$ (and $\lambda = m = 1$). Its limit when $m \rightarrow 0$, which both cancels the cubic vertices and makes $\text{Re}(\phi)$ massless, is the usual “ φ^4 ” real scalar solution, where “ φ ” is here the imaginary part of ϕ :

$$\lim_{m \rightarrow 0} \phi(z, \bar{z}) = i \frac{\text{Im}(z)}{1 - \frac{\lambda}{2m_{\text{Im}(\phi)}^2} \text{Im}(z)^2}. \quad (2.14)$$

2.3 Towards a solution for symmetric soft masses

As stated in Section 2.2, there are more general soft terms than those of eq. (2.12). In particular, it is tempting to consider soft masses for the full complex scalar ϕ ,

$$V = |\lambda\phi^2 + m\phi|^2 + \frac{\delta m^2}{2} |\phi|^2, \quad (2.15)$$

if we, for instance, want to leave some state light and decouple its superpartner. Thus far, we have not found a closed form solution, but we have been able to identify various limits of it. This could be used to either check or guess a more complete expression.

For simplicity, up to redefinitions in the recursion relation like the one we performed in eq. (2.5), we can restrict ourselves to the study of

$$V = |A^2 + A|^2 + \frac{1 - \alpha}{\alpha} |A|^2, \quad (2.16)$$

and of the associated recursion relation/differential equation:

$$\begin{cases} ((n + m)^2 - 1)A_{nm} = 2\alpha^3 \sum A_{m_1 n_1} A_{n_2 m_2} A_{n_3 m_3} + \alpha^2 \sum (A_{n_1 m_1} A_{n_2 m_2} + 2A_{n_1 m_1} A_{m_2 n_2}) \\ [(z\partial_z + \bar{z}\partial_{\bar{z}})^2 - 1]A = 2\alpha^3 A^2 \bar{A} + \alpha^2 (A^2 + 2A\bar{A}) \end{cases}. \quad (2.17)$$

Then, one can solve for real A , or use only the vertices $A\bar{A}^2$ or $A^2\bar{A}$ (see Appendix C for details), to determine the properties of the solution in various limits:

$$\begin{aligned} A(-\rho, -\rho) &= -\frac{\rho}{1 + \alpha\rho - \alpha\frac{1-\alpha}{4}\rho^2}, \\ A(z, 0) &= \frac{z}{\alpha(1 - \frac{\alpha z}{6})^2}, \\ (A/z)(z = 0, \bar{z}) &= \frac{(1 + \frac{\alpha \bar{z}}{6})}{\alpha(1 - \frac{\alpha \bar{z}}{6})^3}. \end{aligned} \quad (2.18)$$

Note that, the first of these is no longer a domain wall solution: depending on the value of α , it either diverges or it describes a regular solution oscillating once in a potential well. Indeed, if $0 < \alpha < 1$ (the positive soft mass case), the denominator vanishes for $\rho = \frac{2(\alpha \pm \sqrt{\alpha})}{\alpha(\alpha-1)}$. On the other hand, when $1 \leq \alpha$, the potential has three extrema: $A = 0, \frac{-3 + \sqrt{\frac{9\alpha-8}{\alpha}}}{4}, \frac{-3 - \sqrt{\frac{9\alpha-8}{\alpha}}}{4}$. The last one is the true minimum, whereas the other two being a local minimum and a local maximum, respectively. In this case the solution corresponds to the field rolling on the inverse potential, from $\phi = 0$ in the direction of the global minimum until it gets blocked by the potential barrier, then settling back at $\phi = 0$. Like a domain wall solution, it has a finite action $\int dt \left(\left| \frac{dA}{dt} \right|^2 + V \right)$ (for $\alpha = 2$ it is ≈ 0.06).

3 SQCD with quartic couplings

The solution of eq. (1.4) is also of interest in SQCD, whose dynamics for N_c colours and N_f flavours has been studied in great detail over the years (for reviews see [20–26]). We are particularly interested in the free magnetic regime, $N_c + 1 < N_f < \frac{3}{2}N_c$, in which there exist WZ domain walls described by eq. (1.4), as we shall now see (see [1, 15, 27, 28] for other studies of domain walls in SQCD theories).

Consider SQCD in such a phase, with a quartic superpotential

$$W^{(\text{el})} = \frac{1}{\mu_X} \text{Tr}[(Q \cdot \tilde{Q})^2] , \quad (3.1)$$

where the dot indicates colour contractions, and the trace is over flavours of quarks and antiquarks Q_i^a, \tilde{Q}_a^j , which are respectively in the fundamental and anti-fundamental representations of $\text{SU}(N_c)$. This operator could be generated by the integrating out of heavier fields of mass $\mathcal{O}(\mu_X)$, as happens generically in the duality cascade [26]. For physical consistency we will therefore require that $\mu_X > \Lambda$, with Λ being the dynamical scale of the electric theory. Below the scale Λ , the electric SQCD theory described above becomes strongly coupled, and physics is best described by its magnetic dual. This theory also has N_f flavours, but $\text{SU}(N)$ gauge group, where $N = N_f - N_c$, and a classical superpotential

$$W_{\text{cl}}^{(\text{mag})} = h q \Phi \tilde{q} + \frac{\mu_\Phi}{2} \text{Tr}(\Phi^2) . \quad (3.2)$$

Here Φ_j^i are the flavour mesons of the infrared (IR) free theory, h is a Yukawa coupling of order unity, and q_i^a, \tilde{q}_a^j are fundamental and anti-fundamental quarks of $\text{SU}(N)$. The Φ mass term is $\mu_\Phi \approx \Lambda^2/\mu_X \ll \Lambda$.

This theory has supersymmetric minima at the origin. In order to be able to count them and compare with the original $\text{SU}(N_c)$ theory, it is useful to also allow the addition of a mass term for the quarks in the electric theory, $W^{(\text{el})} \supset m_Q \text{Tr}(Q \cdot \tilde{Q})$ which must have $m_Q < \Lambda$ (to avoid the quarks being integrated out of the electric theory before we ever reach the scale Λ). In the magnetic theory this becomes a linear meson term, $W_{\text{cl}}^{(\text{mag})} \supset m_Q \Lambda \text{Tr} \Phi$. The conditions for supersymmetric minima then become

$$F_{\Phi_j^i} = h q_i \cdot \tilde{q}^j + \mu_\Phi \phi_i^j + m_Q \Lambda \delta_i^j = 0 , \quad (3.3)$$

along with the $F_q = F_{\bar{q}} = 0$ condition, which has solutions at $\langle q \rangle = \langle \bar{q} \rangle = 0$ and $\langle \phi_i^j \rangle = -\delta_i^j m_Q \Lambda / \mu_\Phi$, parametrically close to the origin (whereas earlier we use ϕ_i^j to denote the scalar component of the superfield). This VEV gives a mass $|hm_Q \Lambda / \mu_\Phi|$ to all the magnetic quarks, and therefore by the usual Witten index theorem, we expect N vacua corresponding to the low energy pure $SU(N)$ Yang-Mills theory.

The remaining supersymmetric minima are separated from the origin by domain walls, beyond which ϕ develops a much larger VEV. Along this direction one is still in a pure $SU(N)$ Yang-Mills theory, but non-perturbative contributions to the superpotential become important. Including these (and neglecting the quark mass term), the complete superpotential for the mesons is as in eq. (1.1)

$$W^{(\text{mag})} = \frac{\mu_\Phi}{2} \text{Tr}(\Phi^2) + N \left(\frac{h^{N_f} \det_{N_f} \Phi}{\Lambda^{N_f - 3N}} \right)^{\frac{1}{N}}, \quad (3.4)$$

where the effective exponent, $p \equiv \frac{N_f}{N}$, is generically a rational number. In the regime of interest, $N_c + 2 \leq N_f < \frac{3}{2}N_c$, we have

$$3 < p \leq \frac{N_f}{2}. \quad (3.5)$$

In principle using eq. (1.4) one can get the exact domain wall solutions for this magnetic theory, for any p .

To find them let us first locate the minima which are along $\phi_i^j = \delta_i^j \phi$ (where we use ϕ to also stand for the trace component of the scalar). Setting $F_\Phi = 0$, we find non-perturbatively generated SUSY preserving minima at

$$\langle \phi_i^j \rangle = \delta_i^j \phi_0 = \delta_i^j \Lambda \left(-h^{\frac{N_f}{N_f - N_c}} \frac{\Lambda}{\mu_\Phi} \right)^{\frac{N_f - N_c}{N_f - 2N_c}}. \quad (3.6)$$

The exponent here is negative so that $\langle \phi \rangle < \Lambda$ as required for the minima to be found in the IR theory. Also note that, as there are no massless quarks, there are generically $2N_c - N_f$ solutions corresponding to the roots of -1 . Together with the $N = N_f - N_c$ minima near the origin this gives the full complement of N_c vacua predicted by the Witten index theorem.

For the domain walls we define

$$\hat{\Phi} = \frac{\Phi}{|\phi_0|}; \quad \hat{W} = \frac{W^{(\text{mag})}}{\mu_\Phi |\phi_0|^2}, \quad (3.7)$$

giving $\hat{W} = \frac{\hat{\Phi}^2}{2} + \frac{\hat{\Phi}^p}{p}$ with $p = N_f/N$. We will henceforth drop the hats.

In order to determine the possible phases of the solution to the BPS condition in eq. (1.3), letting $\phi(t) = |\phi|e^{i(\theta+\eta)}$ we find two equations:

$$\begin{aligned} \partial_t \eta &= -\sin((p-2)\theta + p\eta)|\phi|^{p-2} - \sin(2\eta), \\ \partial_t \log |\phi| &= \cos((p-2)\theta + p\eta)|\phi|^{p-2} + \cos(2\eta), \end{aligned} \quad (3.8)$$

where we recall that now both W and Φ are dimensionless. It is clear that domain wall solutions with constant phase require $\eta = 0$ and $\theta = n\pi/(p-2)$ for integer n . Eq. (1.4) then has $\bar{z}^{p-2} - z^{p-2} \rightarrow 0$ along this direction, and we find

$$\phi(t) = \frac{e^{i\theta} e^t}{(1 - (-1)^n e^{(p-2)t})^{\frac{1}{p-2}}}, \quad (3.9)$$

which is non-singular if n is odd. Hence, there is a domain wall with constant phase between each minimum and the origin. To illustrate this, we show a solution in Figure 4 with $p = 15$. In the large p limit these solutions tend to a universal form, $\phi(t) \stackrel{p \rightarrow \infty}{\approx} 1 + \vartheta(-t)(e^t - 1)$, where ϑ is the Heaviside theta function.




Figure 4. Domain wall solution for SQCD with $p = 15$. In the right panel we plot the potential as a function of t .

4 Conclusion

In this paper, we have presented an exact classical BPS solution of generalised Wess-Zumino models. It extends expressions previously known in the literature by being a solution for a complex scalar field (or equivalently for two real scalar fields), which goes beyond the usual single real scalar field limit via its two-parameters dependence, and by corresponding to a general class of dynamics which encompasses the few two-fields solutions which were obtained earlier. Moreover, the derivation of the complete result required the combined use of several non-trivial techniques.

We have discussed the applications of our solution as a generating function for multi-particle tree-level amplitudes on threshold, where its complex nature is essential to describe both particles and antiparticles, and as a generalisation of known expressions for domain walls in Wess-Zumino models, which are for instance relevant for the vacuum structure of Supersymmetric QCD or for tunneling in the metastable ISS scenario.

We have also pointed out natural extensions of our work, mostly in the context of models with spontaneously or softly broken supersymmetry. There, our methods yield partial expressions which would be interesting to complete, since they would be for instance of relevance for supersymmetric versions of the standard model.

Acknowledgments

We would like to thank Emilian Dudas for his collaboration at the early stages of this work, and Valya Khoze for useful discussions. This research is supported by Royal-Society/CNRS International Cost Share Award IE160590 and by the (Indo-French) CEFIPRA/IFCPAR Project No. 5404-2. Q.B. and D.C. would like to thank the Institute for Particle Physics Phenomenology at Durham University, where part of this work has been conducted, for its hospitality.

A The BPS condition

In this appendix we recap some facts about the BPS condition that underlie the discussion in the main text. A field configuration is BPS [29, 30] if it preserves some amount of supersymmetry. For scalar field configurations (transformed into fermions by supersymmetry), it amounts to requiring that fermions remain equal to zero when the preserved supersymmetry generators act. For a chiral superfield Φ such as the one in the WZ model, the fermion variation is⁴:

$$\delta_\xi \psi = i\sqrt{2}\sigma^m \bar{\xi} \partial_m \phi + \sqrt{2}\xi F, \quad (\text{A.1})$$

for $\Phi = \phi + \sqrt{2}\theta\psi + \theta^2 F$. When calculating multiparticle amplitudes or domain wall profiles, we are interested in one-dimensional problems, so without loss of generality we choose $\phi(x^\mu) = \phi(x)$, x being the spatial coordinate along which the wall extends. Then, demanding that $\delta_\xi \psi = 0$ translates into

$$\bar{\xi}^2 \frac{d\phi}{dx} = i\xi_1 F \text{ and } \bar{\xi}^1 \frac{d\phi}{dx} = i\xi_2 F. \quad (\text{A.2})$$

Whenever the scalars verify $\frac{d\phi}{dx} = -e^{i2\theta} F$ for some real number θ , eq. (A.2) can be satisfied. Using the on-shell value for F , for a trivial Kähler potential and a superpotential W , we find

$$\frac{d\phi}{dx} = e^{2i\theta} \frac{d\bar{W}}{d\bar{\phi}}. \quad (\text{A.3})$$

For the WZ model in eq. (1.1), this reduces to eq. (1.3).

Equation (A.3) can also be understood as a factorisation of the equations of motion. Indeed, imposing the former is enough to satisfy the latter:

$$\frac{d^2\phi}{dx^2} = e^{2i\theta} \frac{d^2\bar{W}}{d\bar{\phi}^2} \frac{d\bar{\phi}}{dx} = \frac{d^2\bar{W}}{d\bar{\phi}^2} \frac{dW}{d\phi} = \frac{dV}{d\bar{\phi}}, \quad (\text{A.4})$$

since $V = \left| \frac{dW}{d\phi}(\phi) \right|^2$ for a chiral superfield.

Equation (A.3) can finally be understood as the condition that minimises the energy per unit surface of a time-independent wall [32–34]:

$$\mathcal{E} = \int dx \left(\left| \frac{d\phi}{dx} \right|^2 + \left| \frac{dW}{d\phi} \right|^2 \right) = \int dx \left| \frac{d\phi}{dx} - e^{2i\theta} \frac{d\bar{W}}{d\bar{\phi}} \right|^2 + 2 \operatorname{Re}(e^{-2i\theta} \Delta W), \quad (\text{A.5})$$

⁴We use the conventions of Ref.[31].

where $\Delta W = W(x = +\infty) - W(x = -\infty)$. The fact that this condition is valid regardless of θ implies the so-called BPS bound:

$$\mathcal{E} \geq 2|\Delta W|. \quad (\text{A.6})$$

In order to saturate this bound, one must again enforce eq. (A.3).

The fact that the generating function of multiparticle amplitudes verifies a BPS condition can be understood from [35]: smooth field configurations which solve the equations of motion and originate from a supersymmetric vacuum state must verify the BPS condition. Eq. (2.9), which defines the generating function in eq. (1.4), thus implies eq. (A.3).

B Link with softly broken O(2) models

Here we draw links with the special case in [2]. When $p = 3$, eq. (1.4) reduces to:

$$\begin{aligned} A(z, \bar{z}) &= \frac{z(1 + \frac{\bar{z}-z}{6})}{1 - \frac{z+\bar{z}}{2} + \frac{(z-\bar{z})^2}{12} - \frac{(z+\bar{z})(z-\bar{z})^2}{216}} \\ &= \left|_{z=e^{t+i\theta}} \frac{e^{t+i\theta}(1 - \frac{2ie^t \sin(\theta)}{6})}{1 - e^t \cos(\theta) - \frac{e^{2t} \sin^2(\theta)}{3} + \frac{e^{3t} \cos(\theta) \sin^2(\theta)}{27}} \right|. \end{aligned} \quad (\text{B.1})$$

Equation (B.1) can be identified with generating functions in softly broken O(2) models [2] of two real scalar fields φ and χ , with potential

$$V(\varphi, \chi) = \mu(\varphi^2 + \chi^2)^2 + \frac{m_1^2}{2}\varphi^2 + \frac{m_2^2}{2}\chi^2. \quad (\text{B.2})$$

Indeed, defining $B = A + \frac{1}{2}$, $V = |A^2 + A|^2 = |B^2 - \frac{1}{4}|^2$ matches (up to the constant term) with $V(\varphi = \text{Re}(B), \chi = \text{Im}(B))$ if we take $\mu = m_2^2 = -m_1^2 = 1$. Then, the “broken reflection symmetry” solution given in [2] matches eq. (B.1) once we identify $A = \varphi - \frac{1}{2} + i\chi$.

C Derivation of the solution

Here, we outline the way eq. (1.4) was found. Although one can check from the solution itself that it solves the BPS condition for the model of eq. (1.2), different methods have been used in its derivation, so we quickly list them here, following our chronological progression.

First, for the $p = 3$ case one can start by solving eq. (2.8) with $\theta = 0$ or π (i.e. z real), which makes $A(z, \bar{z} = z)$ real, giving

$$A(z, \bar{z} = z) = \frac{z}{1 - z}. \quad (\text{C.1})$$

In order to derive this expression, we impose that A scales as z as z goes to 0, which is enough/required for walls or amplitudes. One then seeks the multiparticle amplitudes where an incoming ϕ goes into n ϕ 's (and no $\bar{\phi}$'s) in the final states. This corresponds to graphs where only ϕ propagates since, at each vertex, the number of ϕ 's, or $\bar{\phi}$'s, in the

out-states is always larger (or equal) than the one in the in-states. It amounts to solving the equation $\partial_t^2 A = A + A^2$, which determines $A(z, 0)$. The solution is

$$A(z, \bar{z} = 0) = \frac{z}{\left(1 - \frac{z}{6}\right)^2} . \quad (\text{C.2})$$

Then, one can make an educated guess of the form

$$A(z, \bar{z} = 0) = \frac{z}{\left(1 - \frac{z}{6}\right)^2 + \bar{z}f(z, \bar{z})} , \quad (\text{C.3})$$

and numerically solve the first steps of the recursion relation in eq. (2.6) to get the (z, \bar{z}) expansion of $f(z, \bar{z})$, from which one can surmise the following fully resummed expression:

$$\left(A/z\right)(z = 0, \bar{z}) = \frac{\left(1 + \frac{\bar{z}}{6}\right)}{\left(1 - \frac{\bar{z}}{6}\right)^3} . \quad (\text{C.4})$$

After some more recursive steps one can deduce the full $p = 3$ solution:

$$A(z, \bar{z}) = \frac{z\left(1 + \frac{\bar{z}-z}{6}\right)}{1 - \frac{z+\bar{z}}{2} + \frac{(z-\bar{z})^2}{12} - \frac{(z+\bar{z})(z-\bar{z})^2}{216}} . \quad (\text{C.5})$$

This solution turns out to be a reshuffling of the one found in [2].

Higher p solutions are derived in the following way: the Hamilton-Jacobi equation for the WZ model with $p = 4$ can be solved with a variable separation, as in [2], by defining:

$$A = \sqrt{2(\xi^2 - 1)(1 - \eta^2)} + i\sqrt{2}\xi\eta . \quad (\text{C.6})$$

Ultimately this gives

$$A(z, \bar{z}) = \frac{z\left(1 + \frac{\bar{z}^2 - z^2}{8}\right)}{\sqrt{\left[1 - \frac{(z-\bar{z})^2}{4} + \frac{(z^2 - \bar{z}^2)^2}{64}\right]\left[1 - \frac{(z+\bar{z})^2}{4} + \frac{(z^2 - \bar{z}^2)^2}{64}\right]}} . \quad (\text{C.7})$$

From this example one can guess that, for general p ,

$$A(z, \bar{z}) = \frac{z\left(1 + \frac{\bar{z}^{p-2} - z^{p-2}}{2p}\right)}{P(z, \bar{z})} , \quad (\text{C.8})$$

with $P(z, \bar{z})$ being a real function. This parametrisation makes it possible to solve the BPS condition of eq. (1.3), which gives a simple first order equation for $P(z, \bar{z})$ whose solution, with our boundary conditions, is eq. (1.4). The latter yields expressions in particular limits that match the results of derivations similar to the discussions for eqs. (C.1) and (C.2):

$$A(z, \bar{z} = z) = \frac{z}{(1 - z^{p-2})^{\frac{1}{p-2}}} , \quad A(z, \bar{z} = 0) = \frac{z}{\left(1 - \frac{z^{p-2}}{2p}\right)^{\frac{2}{p-2}}} . \quad (\text{C.9})$$

References

- [1] B. Chibisov and M. A. Shifman, “BPS saturated walls in supersymmetric theories,” *Phys. Rev. D* **56** (1997) 7990–8013, [arXiv:hep-th/9706141 \[hep-th\]](#). [Erratum: *Phys. Rev. D* 58, 109901 (1998)].
- [2] M. V. Libanov, V. A. Rubakov, and S. V. Troitsky, “Tree amplitudes at multiparticle threshold in a model with softly broken $O(2)$ symmetry,” *Nucl. Phys. B* **412** (1994) 607–620, [arXiv:hep-ph/9307213 \[hep-ph\]](#).
- [3] M. B. Voloshin, “Multiparticle amplitudes at zero energy and momentum in scalar theory,” *Nucl. Phys. B* **383** (1992) 233–248.
- [4] E. N. Argyres, R. H. P. Kleiss, and C. G. Papadopoulos, “Amplitude estimates for multi - Higgs production at high-energies,” *Nucl. Phys. B* **391** (1993) 42–56.
- [5] M. B. Voloshin, “Estimate of the onset of nonperturbative particle production at high-energy in a scalar theory,” *Phys. Lett. B* **293** (1992) 389–394.
- [6] L. S. Brown, “Summing tree graphs at threshold,” *Phys. Rev. D* **46** (1992) R4125–R4127, [arXiv:hep-ph/9209203 \[hep-ph\]](#).
- [7] M. V. Libanov, V. A. Rubakov, D. T. Son, and S. V. Troitsky, “Exponentiation of multiparticle amplitudes in scalar theories,” *Phys. Rev. D* **50** (1994) 7553–7569, [arXiv:hep-ph/9407381 \[hep-ph\]](#).
- [8] D. T. Son, “Semiclassical approach for multiparticle production in scalar theories,” *Nucl. Phys. B* **477** (1996) 378–406, [arXiv:hep-ph/9505338 \[hep-ph\]](#).
- [9] M. V. Libanov, V. A. Rubakov, and S. V. Troitsky, “Multiparticle processes and semiclassical analysis in bosonic field theories,” *Phys. Part. Nucl.* **28** (1997) 217–240.
- [10] V. V. Khoze and J. Reiness, “Review of the semiclassical formalism for multiparticle production at high energies,” [arXiv:1810.01722 \[hep-ph\]](#).
- [11] V. V. Khoze, “Multiparticle Higgs and Vector Boson Amplitudes at Threshold,” *JHEP* **07** (2014) 008, [arXiv:1404.4876 \[hep-ph\]](#).
- [12] V. V. Khoze and M. Spannowsky, “Higgspllosion: Solving the Hierarchy Problem via rapid decays of heavy states into multiple Higgs bosons,” *Nucl. Phys. B* **926** (2018) 95–111, [arXiv:1704.03447 \[hep-ph\]](#).
- [13] V. V. Khoze, “Multiparticle production in the large n limit: realising Higgspllosion in a scalar QFT,” *JHEP* **06** (2017) 148, [arXiv:1705.04365 \[hep-ph\]](#).
- [14] V. V. Khoze, “Semiclassical computation of quantum effects in multiparticle production at large λn ,” [arXiv:1806.05648 \[hep-ph\]](#).
- [15] G. R. Dvali and M. A. Shifman, “Dynamical compactification as a mechanism of spontaneous supersymmetry breaking,” *Nucl. Phys. B* **504** (1997) 127–146, [arXiv:hep-th/9611213 \[hep-th\]](#).
- [16] M. B. Voloshin, “Summing one loop graphs at multiparticle threshold,” *Phys. Rev. D* **47** (1993) R357–R361, [arXiv:hep-ph/9209240 \[hep-ph\]](#).
- [17] A. S. Gorsky and M. B. Voloshin, “Nonperturbative production of multiboson states and quantum bubbles,” *Phys. Rev. D* **48** (1993) 3843–3851, [arXiv:hep-ph/9305219 \[hep-ph\]](#).

- [18] M. V. Libanov, D. T. Son, and S. V. Troitsky, “Exponentiation of multiparticle amplitudes in scalar theories. 2. Universality of the exponent,” *Phys. Rev.* **D52** (1995) 3679–3687, [arXiv:hep-ph/9503412](#) [hep-ph].
- [19] L. Girardello and M. T. Grisaru, “Soft Breaking of Supersymmetry,” *Nucl. Phys.* **B194** (1982) 65.
- [20] J. Terning, *Modern supersymmetry: Dynamics and duality*. 2006.
- [21] M. Dine, *Supersymmetry and String Theory*. Cambridge University Press, 2016.
<http://www.cambridge.org/academic/subjects/physics/particle-physics-and-nuclear-physics/supersymmetry-and-string-theory-beyond-standard-model-2nd-edition?format=AR&isbn=9781316405307>.
- [22] K. A. Intriligator and N. Seiberg, “Lectures on supersymmetric gauge theories and electric-magnetic duality,” *Nucl. Phys. Proc. Suppl.* **45BC** (1996) 1–28, [arXiv:hep-th/9509066](#) [hep-th]. [,157(1995)].
- [23] M. E. Peskin, “Duality in supersymmetric Yang-Mills theory,” in *Fields, strings and duality. Proceedings, Summer School, Theoretical Advanced Study Institute in Elementary Particle Physics, TASI’96, Boulder, USA, June 2-28, 1996*, pp. 729–809. 1997.
[arXiv:hep-th/9702094](#) [hep-th]. <http://www-public.slac.stanford.edu/sciDoc/docMeta.aspx?slacPubNumber=SLAC-PUB-7393>.
- [24] M. A. Shifman, “Nonperturbative dynamics in supersymmetric gauge theories,” *Prog. Part. Nucl. Phys.* **39** (1997) 1–116, [arXiv:hep-th/9704114](#) [hep-th]. [,345(1995)].
- [25] K. A. Intriligator, N. Seiberg, and D. Shih, “Supersymmetry breaking, R-symmetry breaking and metastable vacua,” *JHEP* **07** (2007) 017, [arXiv:hep-th/0703281](#) [hep-th].
- [26] M. J. Strassler, “The Duality cascade,” in *Progress in string theory. Proceedings, Summer School, TASI 2003, Boulder, USA, June 2-27, 2003*, pp. 419–510. 2005.
[arXiv:hep-th/0505153](#) [hep-th].
- [27] I. Bandos, F. Farakos, S. Lanza, L. Martucci, and D. Sorokin, “Three-forms, dualities and membranes in four-dimensional supergravity,” *JHEP* **07** (2018) 028, [arXiv:1803.01405](#) [hep-th].
- [28] I. Bandos, S. Lanza, and D. Sorokin, “Supermembranes and domain walls in $\mathcal{N} = 1$, $D = 4$ SYM,” [arXiv:1905.02743](#) [hep-th].
- [29] E. B. Bogomolny, “Stability of Classical Solutions,” *Sov. J. Nucl. Phys.* **24** (1976) 449. [*Yad. Fiz.*24,861(1976)].
- [30] M. K. Prasad and C. M. Sommerfield, “An Exact Classical Solution for the ’t Hooft Monopole and the Julia-Zee Dyon,” *Phys. Rev. Lett.* **35** (1975) 760–762.
- [31] J. Wess and J. Bagger, *Supersymmetry and supergravity*. Princeton University Press, Princeton, NJ, USA, 1992.
- [32] J. A. de Azcarraga, J. P. Gauntlett, J. M. Izquierdo, and P. K. Townsend, “Topological Extensions of the Supersymmetry Algebra for Extended Objects,” *Phys. Rev. Lett.* **63** (1989) 2443.
- [33] P. Fendley, S. D. Mathur, C. Vafa, and N. Warner, “Integrable Deformations and Scattering Matrices for the $N = 2$ Supersymmetric Discrete Series,” *Phys. Lett.* **B243** (1990) 257–264.

- [34] S. Cecotti and C. Vafa, “On classification of N=2 supersymmetric theories,” *Commun. Math. Phys.* **158** (1993) 569–644, [arXiv:hep-th/9211097 \[hep-th\]](#).
- [35] D. Bazeia, J. Menezes, and M. M. Santos, “Complete factorization of equations of motion in supersymmetric field theories,” *Nucl. Phys.* **B636** (2002) 132–140, [arXiv:hep-th/0103041 \[hep-th\]](#).