

Magnetic field effects on neutrino oscillations

Andrea Erdas* and Zachary Metzler†

Department of Physics, Loyola University Maryland 4501 N Charles Street, Baltimore, MD 21210

(Dated: July 4, 2019)

Using the exact propagators in a constant magnetic field, the neutrino self-energy has been calculated to all orders in the field strength B within the minimal extension of the Weinberg-Salam model with massive Dirac neutrinos. A neutrino dispersion relation, effective potential and effective mass have been obtained that depend on B . The consequences of this effective potential on neutrino oscillations have been explored, and resonance conditions have been obtained for magnetic fields $0 < eB \leq 4m_e^2$, where e is the elementary charge and m_e is the electron mass. The oscillation length has also been obtained, showing that ν_e - ν_μ resonant oscillations in magnetic field are likely to occur in the proximity of most stars generating high magnetic fields.

PACS numbers: 13.15.+g, 13.40.-f, 14.60.Lm, 14.60.Pq, 95.30.Cq

I. INTRODUCTION

The study of neutrinos in the presence of a magnetic field is important in astrophysics and cosmology [1, 2]. The neutrino self-energy and its dispersion relation are modified in a magnetic field, and such modifications have been studied extensively in the literature [3–12]. There is a natural magnetic field scale that is required to significantly impact quantum processes, dependent on the electron mass m_e and the elementary charge e , $B_e = m_e^2/e = 4.41 \times 10^{13}$ G. Large magnetic fields are present in many astrophysical sites such as supernovae, neutron stars, and white dwarfs. Fields larger than B_e can appear during the explosion of a supernova or the coalescing of a neutron star. Magnetars are young neutron stars that generate even stronger magnetic fields, $10^{14} - 10^{16}$ G [13–15]. Magnetic fields as high as $10^{22} - 10^{24}$ G have been hypothesized to exist during the electroweak phase transition of the early universe [16–18]. Although it is rare for neutrinos to encounter magnetic fields larger than 10^{16} G, in many situations neutrinos come upon astrophysical sites where magnetic field strengths are at or around B_e .

While the electromagnetic properties of massless neutrinos have been studied extensively and for a long time [3–6], the nontrivial electromagnetic properties of massive neutrinos have only been addressed more recently [19–21]. A recent investigation on the role of a neutrino magnetic moment in flavor, spin and spin-flavor oscillations has produced interesting results [22, 23], but does not consider magnetic effects beyond those linear in B . In this paper we will focus on massive Dirac neutrinos within the minimally extended Standard Model and take a rigorous field theoretical approach to calculating their self-energy in a homogeneous magnetic field. We will accomplish this task by using Schwinger’s proper time method [24] and the exact fermion, scalar, and W -propagator in a constant magnetic field [6, 7].

In Section II the notation for the fermion, gauge boson and scalar propagators in magnetic field is reviewed and the one-loop neutrino self-energy is set up. In Section III we calculate the self-energy operator and obtain a simple expression for the self-energy in the limit of $eB \ll m_W^2$, where m_W is the mass of the W -boson, exact up to order B^2 . In Section IV we use the self-energy operator we obtained to calculate the dispersion relation, effective potential and effective mass for massive Dirac neutrinos in a magnetic field and show that terms in the self-energy of order B^2 are often dominant when compared to linear terms such as the magnetic dipole moment term. In Section V we use the effective potential to investigate neutrino oscillation in a magnetic field and obtain resonance condition and oscillation length.

II. PROPAGATORS AND NEUTRINO SELF-ENERGY IN A CONSTANT MAGNETIC FIELD

The metric used in this paper is $g^{\mu\nu} = \text{diag}(-1, +1, +1, +1)$ and the z -axis points in the direction of the constant magnetic field $\mathbf{B}$. Therefore the electromagnetic field strength tensor $F^{\mu\nu}$ has only two non-vanishing components $F^{12} = -F^{21} = B$.

*Electronic address: aerdas@loyola.edu

†Electronic address: zbmetzler@loyola.edu

For the purpose of this work, it would seem convenient to work in the unitary gauge where the unphysical scalars disappear. However, the W -propagator is quite cumbersome in this gauge, and we prefer to work in the Feynman gauge, where the W -propagator has a much simpler expression. The following expressions for the charged lepton $S(x', x'')$ [24], W -boson $G^{\mu\nu}(x', x'')$ and scalar propagators $D(x', x'')$ [6] in a constant magnetic field have been written using Schwinger's proper time method:

$$S(x', x'') = \Omega^*(x', x'') \int \frac{d^4 k}{(2\pi)^4} e^{ik \cdot (x' - x'')} S(k) \quad , \quad (1)$$

$$G^{\mu\nu}(x', x'') = \Omega(x', x'') \int \frac{d^4 k}{(2\pi)^4} e^{ik \cdot (x' - x'')} G^{\mu\nu}(k) \quad , \quad (2)$$

$$D(x', x'') = \Omega(x', x'') \int \frac{d^4 k}{(2\pi)^4} e^{ik \cdot (x' - x'')} D(k) \quad . \quad (3)$$

In the Feynman gauge, the translationally invariant parts of the propagators are

$$S(k) = \int_0^\infty \frac{ds}{\cos eBs} \exp \left[-is \left(m_\ell^2 + k_\parallel^2 + k_\perp^2 \frac{\tan eBs}{eBs} \right) \right] \left[(m_\ell - \not{k}_\parallel) e^{-ieBs\sigma_3} - \frac{\not{k}_\perp}{\cos eBs} \right], \quad (4)$$

$$G^{\mu\nu}(k) = \int_0^\infty \frac{ds}{\cos eBs} \exp \left[-is \left(m_W^2 + k_\parallel^2 + k_\perp^2 \frac{\tan eBs}{eBs} \right) \right] [g_\parallel^{\mu\nu} + (e^{2eFs})^{\mu\nu}], \quad (5)$$

$$D(k) = \int_0^\infty \frac{ds}{\cos eBs} \exp \left[-is \left(m_W^2 + k_\parallel^2 + k_\perp^2 \frac{\tan eBs}{eBs} \right) \right], \quad (6)$$

where $-e$ and m_ℓ are the charge and mass of the charged lepton ℓ , and m_W is the W -mass. It is convenient to use the notation

$$a_\parallel^\mu = (a^0, 0, 0, a^3), \quad a_\perp^\mu = (0, a^1, a^2, 0) \quad (7)$$

and

$$(ab)_\parallel = -a^0 b^0 + a^3 b^3, \quad (ab)_\perp = a^1 b^1 + a^2 b^2 \quad (8)$$

for arbitrary four-vectors a and b . Using this notation we write the metric tensor as

$$g^{\mu\nu} = g_\parallel^{\mu\nu} + g_\perp^{\mu\nu}. \quad (9)$$

The 4×4 matrix σ_3 that appears in the charged lepton propagator (4), is

$$\sigma_3 = \frac{i}{2} [\gamma^1, \gamma^2]. \quad (10)$$

When writing the W -propagator (5), we use the notation

$$(e^{2eFs})^{\mu\nu} = g_\perp^{\mu\nu} \cos(2eBs) + \frac{F^{\mu\nu}}{B} \sin(2eBs) \quad . \quad (11)$$

We choose the electromagnetic vector potential to be $A_\mu = -\frac{1}{2} F_{\mu\nu} x^\nu$ and therefore the phase factor which appears in Eqs. (2), (3) is given by

$$\Omega(x', x'') = \exp \left(i \frac{e}{2} x'_\mu F^{\mu\nu} x''_\nu \right) \quad , \quad (12)$$

with $\Omega^*(x', x'')$ in Eq. (1) being its complex conjugate.

Perturbatively, the self-energy operator in the Feynman gauge corresponds to the sum of two diagrams, a bubble diagram with the gauge boson and a bubble diagram with the scalar

$$\Sigma(p) = \Sigma_W(p) + \Sigma_\Phi(p). \quad (13)$$

The translationally non-invariant phase factors $\Omega(x', x'')$ are identical for all propagators and the product of phase factors in the two-vertex loop is

$$\Omega^*(x', x'')\Omega(x', x'') = 1, \quad (14)$$

therefore, within the minimally extended version of the standard model of electroweak interactions with an $SU(2)$ -singlet right-handed neutrino, the two bubble diagrams can be written as [6, 25]

$$\Sigma_W(p) = -i\frac{g^2}{2}R\gamma_\alpha \int \frac{d^4k}{(2\pi)^4} S(p-k)G^{\beta\alpha}(k)\gamma_\beta L, \quad (15)$$

$$\Sigma_\Phi(p) = -i\frac{g^2}{2m_W^2}[m_\ell R - m_\nu L] \int \frac{d^4k}{(2\pi)^4} S(p-k)D(k)[m_\ell L - m_\nu R], \quad (16)$$

where g is the $SU(2)$ coupling constant, $L = \frac{1}{2}(1 - \gamma_5)$ and $R = \frac{1}{2}(1 + \gamma_5)$ are the left-handed and right-handed projectors and neutrino mixing is allowed by taking a nondiagonal neutrino mass matrix m_ν in Eq. (16).

III. THE SELF-ENERGY OPERATOR

Inserting the expression for the propagators from Eqs. (4) and (5) into the self-energy, we write $\Sigma_W(p)$ as

$$\begin{aligned} \Sigma_W(p) = & -\frac{ig^2}{2} \int \frac{d^4k}{(2\pi)^4} \int_0^\infty \frac{ds_1}{\cos z_1} \int_0^\infty \frac{ds_2}{\cos z_2} e^{-is_1(m_\ell^2 + q_\parallel^2 + q_\perp^2 \frac{\tan z_1}{z_1})} e^{-is_2(m_W^2 + k_\parallel^2 + k_\perp^2 \frac{\tan z_2}{z_2})} \times \\ & R\gamma_\alpha \left[(m_\ell - \not{q}_\parallel) e^{-iz_1\sigma_3} - \frac{\not{q}_\perp}{\cos z_1} \right] [g_\parallel^{\beta\alpha} + (e^{2eFs_2})_\perp^{\beta\alpha}] \gamma_\beta L \end{aligned} \quad (17)$$

where

$$q = p - k \quad , \quad z_1 = eBs_1 \quad , \quad z_2 = eBs_2 \quad . \quad (18)$$

We do the straightforward γ -algebra, change variables from s_i to z_i , translate the k variables of integration and do the four gaussian integrals over the shifted variables k , to obtain

$$\begin{aligned} \Sigma_W(p) = & \frac{g^2}{(4\pi)^2} \int_0^\infty \int_0^\infty \frac{dz_1 dz_2}{(z_1 + z_2) \sin(z_1 + z_2)} e^{-i[z_1 m_\ell^2 + z_2 m_W^2 + \mathcal{P}]/eB} \times \\ & \left[\frac{z_2}{z_1 + z_2} \not{p}_\parallel e^{i\sigma_3(z_1 + 2z_2)} + \frac{\sin z_2}{\sin(z_1 + z_2)} \not{p}_\perp \right] L + (\text{c.t.})_W \end{aligned} \quad (19)$$

where

$$\mathcal{P} = \frac{z_1 z_2}{(z_1 + z_2)} p_\parallel^2 + \frac{\sin z_1 \sin z_2}{\sin(z_1 + z_2)} p_\perp^2 \quad , \quad (20)$$

and the appropriate counter-terms (c.t.) are added. Next it is convenient to change integration variables from (z_1, z_2) to (s, u) defined by

$$z_1 = eBs u = zu \quad \text{and} \quad z_2 = eBs(1 - u) = z(1 - u) \quad . \quad (21)$$

The result is

$$\Sigma_W(p) = \frac{g^2}{(4\pi)^2} \int_0^\infty \frac{dz}{\sin z} \int_0^1 du e^{-is\Lambda^2} R \left[(1 - u) e^{iz(2-u)\sigma_3} \not{p}_\parallel + \frac{\sin[z(1 - u)]}{\sin z} \not{p}_\perp \right] L + (\text{c.t.})_W \quad (22)$$

where

$$\Lambda^2 = um_\ell^2 + (1 - u)m_W^2 + u(1 - u)p_\parallel^2 + \frac{\sin zu \sin(z - zu)}{z \sin z} p_\perp^2. \quad (23)$$

The scalar-loop contribution to the self-energy is found similarly

$$\Sigma_\Phi(p) = \frac{g^2}{(4\pi)^2} \frac{\delta_l}{2} \int_0^\infty \frac{dz}{\sin z} \int_0^1 du e^{-is\Lambda^2} R \left[(1-u)e^{-izu\sigma_3} \not{p}_\parallel + \frac{\sin[z(1-u)]}{\sin z} \not{p}_\perp \right] L + (\text{c.t.})_\Phi \quad (24)$$

where we used

$$\delta_l = \frac{m_l^2}{m_W^2}, \quad (25)$$

and neglected the matrix m_ν from Eq. (16) whose elements are all of the order of 2eV or less.

At this point we focus on magnetic fields $eB \ll m_W^2$, since magnetic fields $eB \sim m_W^2$ are not found in the universe. The expression of $\Sigma_W(p)$ in the limit of $eB \ll m_W^2$ is readily found by taking $z \ll 1$, and we obtain

$$\begin{aligned} \Sigma_W(p) = & \frac{g^2}{(4\pi)^2} \int_0^\infty \frac{ds}{s} \int_0^1 du e^{-is\Lambda_0^2} (1-u) R \left[i(2-u) \frac{z}{m_W^2} \sigma_3 \not{p}_\parallel + \frac{z^2}{6m_W^4} \not{p} - \frac{z^2}{2m_W^4} (2-u)^2 \not{p}_\parallel \right. \\ & \left. + \frac{z^2}{6m_W^4} u(2-u) \not{p}_\perp - is \frac{z^2}{3m_W^4} u^2 (1-u)^2 \left(\frac{p_\perp^2}{m_W^2} \right) \not{p} \right] L, \end{aligned} \quad (26)$$

where we neglected terms that do not depend on the magnetic field since they will be absorbed by the neutrino wavefunction and mass renormalization, and we used the notation

$$\Lambda_0^2 = (1-u) + u \frac{m_l^2}{m_W^2} + (1-u)u \frac{p^2}{m_W^2}. \quad (27)$$

Since $p^2 \sim m_\nu^2$, where m_ν is the mass of the neutrino, we can take $\Lambda_0^2 \simeq (1-u) + u\delta_l$ and, after doing the s and u integration, we obtain

$$\begin{aligned} \Sigma_W(p) = & \frac{g^2}{(4\pi)^2} R \left[\frac{3eB}{2m_W^2} \sigma_3 \not{p}_\parallel + \frac{(eB)^2}{6m_W^4} \left(\frac{5}{2} + \frac{4p_\perp^2}{3m_W^2} + \ln \delta_l \right) \not{p} \right. \\ & \left. + \frac{(eB)^2}{2m_W^4} (1 - \ln \delta_l) \not{p}_\parallel + \frac{(eB)^2}{6m_W^4} (\ln \delta_l) \not{p}_\perp \right] L. \end{aligned} \quad (28)$$

Undertaking similar steps, we find

$$\begin{aligned} \Sigma_\Phi(p) = & \frac{g^2}{(4\pi)^2} \delta_l R \left[-\frac{eB}{4m_W^2} \sigma_3 \not{p}_\parallel + \frac{(eB)^2}{3m_W^4} \left(\frac{1}{4} + \frac{p_\perp^2}{3m_W^2} + \frac{1}{4} \ln \delta_l \right) \not{p} \right. \\ & \left. - \frac{(eB)^2}{4m_W^4} \left(\frac{5}{2} + \ln \delta_l \right) \not{p}_\parallel + \frac{(eB)^2}{12m_W^4} \left(\frac{3}{2} + \ln \delta_l \right) \not{p}_\perp \right] L, \end{aligned} \quad (29)$$

where each term is suppressed by a factor of δ_l when compared to the similar term in $\Sigma_W(p)$, thus making $\Sigma_\Phi(p)$ negligible when compared to $\Sigma_W(p)$. The self-energy operator is therefore $\Sigma(p) \simeq \Sigma_W(p)$.

IV. DISPERSION RELATION, EFFECTIVE POTENTIAL AND EFFECTIVE MASS

We intend to interpret the effect of the magnetic field on the self-energy operator as an effective neutrino mass [6, 26]. To do so we must first evaluate the average value of $\Sigma(p)$ over the neutrino spinor in the mass basis, $\langle \Sigma \rangle = \bar{u}_i \Sigma(p) u_i$, since neutrino mass is defined in the mass basis and not in the flavor basis. We find,

$$\bar{u}_i R \sigma_3 \not{p}_\parallel L u_i = -\frac{m_i s_3}{2} \quad (30)$$

where $s_3 = \pm 1$, depending on the neutrino spin orientation relative to the magnetic field and, similarly,

$$\bar{u}_i R \not{p} L u_i = -\frac{m_i}{2}, \quad (31)$$

and

$$\bar{u}_i R \not{p}_\perp L u_i = \chi \bar{u}_i \not{p}_\perp u_i, \quad (32)$$

where $\chi = \frac{1}{2} \langle 1 - \frac{\vec{\sigma} \cdot \mathbf{p}}{E + m_i} \rangle$ in the Dirac representation and E and $\mathbf{p}$ are the neutrino energy and momentum. Indicating with U the neutrino mixing matrix

$$\nu_\ell = \sum_{i=1}^3 U_{\ell i} \nu_i, \quad (33)$$

and using Eqs. (30), (31), and (32), we write

$$\langle \Sigma \rangle = -\frac{g^2}{64\pi^2} \frac{eB}{m_W^2} \sum_\ell U_{\ell i}^* U_{\ell i} \left[3s_3 m_i + \frac{eB}{3m_W^2} \left(\frac{11}{2} + \frac{4p_\perp^2}{3m_W^2} - 2 \ln \delta_\ell \right) m_i + \frac{2eB}{m_W^2} \left(1 - \frac{4}{3} \ln \delta_\ell \right) \chi \bar{u}_i \not{p}_\perp u_i \right], \quad (34)$$

where $\ell = e, \mu, \tau$. Eq (34) agrees with [6] in the absence of mixing and neutrino masses. Notice that transitions of the type $\nu_i \rightarrow \nu_j$ are suppressed [27] by a GIM type mechanism, and therefore quantities such as $\bar{u}_j \Sigma(p) u_i$ can be neglected. We obtain the dispersion relation by setting

$$\not{p} + m_i - \langle \Sigma \rangle = 0, \quad (35)$$

and find the following energy-momentum relation for the massive neutrino in a magnetic field

$$E^2 = \mathbf{p}^2 + 2\Lambda p_\perp^2 + (1 + 2\mu)m_i^2, \quad (36)$$

with

$$\Lambda = \frac{g^2}{32\pi^2} \frac{(eB)^2}{m_W^4} \sum_\ell U_{\ell i}^* U_{\ell i} \left(1 - \frac{4}{3} \ln \delta_\ell \right) \chi, \quad (37)$$

and

$$\mu = \frac{g^2}{64\pi^2} \frac{eB}{m_W^2} \sum_\ell U_{\ell i}^* U_{\ell i} \left[3s_3 + \frac{eB}{3m_W^2} \left(\frac{11}{2} + \frac{4p_\perp^2}{3m_W^2} - 2 \ln \delta_\ell \right) \right]. \quad (38)$$

Eq. (38) is valid in the case of weak ($eB \ll m_e^2$) and moderate ($m_e^2 \ll eB \ll m_W^2$) magnetic field, while Eq. (37) is valid only in the case of weak magnetic field. In the case of moderate magnetic field Λ becomes [7]

$$\Lambda = \frac{g^2}{32\pi^2} \frac{(eB)^2}{m_W^4} \sum_\ell U_{\ell i}^* U_{\ell i} \left[1 - \frac{4}{3} \ln \left(\frac{eB}{6m_W^2} + \delta_\ell \right) \right] \chi, \quad (39)$$

which is more general than Eq. (37) and renders obsolete the distinction between weak and moderate magnetic field.

The dispersion relation of Eq (36) yields the following effective neutrino mass

$$m_{eff} = \sqrt{(1 + 2\mu)m_i^2 + 2\Lambda p_\perp^2}. \quad (40)$$

Notice that, since $m_i \leq 2\text{eV}$, for $p_\perp \sim 1\text{MeV}$ or higher the Λ term in Eq. (40) dominates over the μ term, which can be safely neglected in many situations.

The effective potential is most useful in the flavor basis and, neglecting terms similar to μ of Eq. (38) for the reason stated above, we find

$$V_\ell = \lambda_\ell p_\perp^2 / E, \quad (41)$$

with

$$\lambda_\ell = \frac{g^2}{32\pi^2} \frac{(eB)^2}{m_W^4} \left[1 - \frac{4}{3} \ln \left(\frac{eB}{6m_W^2} + \delta_\ell \right) \right] \chi, \quad (42)$$

where $\ell = e, \mu, \tau$.

V. RESONANCE CONDITION AND OSCILLATION LENGTH

Our main results, Eqs. (40) - (42), have significant consequences on neutrino oscillations of the type $\nu_e \leftrightarrow \nu_{\mu,\tau}$. While the original MSW effect [28–30] showed that the presence of a medium alone can produce neutrino oscillations, in this work we will explore the role of a lone magnetic field in neutrino oscillations. The mixing angle θ_B in a magnetic field is determined by the following

$$\sin^2 2\theta_B = \frac{\sin^2 2\theta_{1i}}{\left[\cos 2\theta_{1i} - \frac{2E(V_e - V_\ell)}{\Delta m_{i1}^2} \right]^2 + \sin^2 2\theta_{1i}} \quad (43)$$

where Δm_{i1}^2 and θ_{1i} are the squared-mass splitting and vacuum mixing angle in the ν_e, ν_ℓ system, with $\ell = \mu, \tau$ and $i = 2, 3$ for $\ell = \mu, \tau$, and V_e and V_ℓ are the ν_e and ν_ℓ effective potentials in a magnetic field obtained in Eq. (41). Resonant oscillations will occur if

$$\Delta m_{i1}^2 \cos 2\theta_{i1} = 2E(V_e - V_\ell) \quad (44)$$

and, using Eq. (41), we find that the resonance conditions is

$$\Delta m_{i1}^2 \cos 2\theta_{i1} \simeq 2(\lambda_e - \lambda_\ell)p_\perp^2, \quad (45)$$

where we neglected the smaller term proportional to μ in (41) and

$$\lambda_e - \lambda_\ell = \frac{g^2}{24\pi^2} \frac{(eB)^2}{m_W^4} \chi \ln \left(\frac{eB + 6m_\ell^2}{eB + 6m_e^2} \right). \quad (46)$$

We take $\chi \sim 1$, the newest and most accurate values of Δm_{i1}^2 and $\cos 2\theta_{i1}$ [31], and in Figures 1 and 2 plot the value of $p_\perp$ for which resonance occurs as a function of B , for $0 < B \leq 4B_e$, where $B_e = m_e^2/e = 4.4 \times 10^{13}$ G. Figure 1 shows our result for ν_e - ν_μ oscillations, Figure 2 shows it for ν_e - ν_τ oscillations.

Neutrino oscillatory behavior is prominent when $L \sim L^{osc}$, where L is the distance travelled inside the magnetic field and L^{osc} is the oscillation length. The oscillation length for neutrinos in a magnetic field is

$$L^{osc} = \frac{L_0^{osc}}{\sqrt{\left[\cos 2\theta_{1i} - \frac{2(\lambda_e - \lambda_\ell)p_\perp^2}{\Delta m_{i1}^2} \right]^2 + \sin^2 2\theta_{1i}}} \quad (47)$$

where L_0^{osc} is the vacuum oscillation length, given by

$$L_0^{osc} = \frac{4\pi E}{\Delta m_{i1}^2} = 2.5 \frac{E}{\Delta m_{i1}^2} \text{ meter} \quad (48)$$




FIG. 1: Plot of $p_\perp$ for which resonance occurs as a function of B , in the case of ν_e - ν_μ oscillations. $p_\perp$ is in GeV and $0 < B \leq 4B_e$.

where the neutrino energy E is in MeV and Δm_{i1}^2 is in eV^2 . The neutrino energy is easily related to $p_\perp$ by assuming that the three components of the neutrino momentum are of similar magnitude, $p_1 \simeq p_2 \simeq p_3$, which leads to $E \sim \sqrt{\frac{3}{2}} p_\perp$. The oscillation length at resonance is

$$L_R^{osc} = \frac{L_0^{osc}}{\sin 2\theta_{1i}}, \quad (49)$$

and, in the case of $\nu_e\text{-}\nu_\mu$ oscillations when $B = B_e$ and $E \sim 270$ MeV, we find $L_R^{osc} \sim 10^4$ Km. For $\nu_e\text{-}\nu_\mu$ oscillations, $B = 4B_e$, and $E \sim 70$ MeV, the oscillation length is $L_R^{osc} \sim 2.5 \times 10^3$ Km. Many astrophysical sites such as supernovae, neutron stars, white dwarfs and magnetars have magnetic fields as large as B_e or larger, even larger by more than one order of magnitude, and therefore $\nu_e\text{-}\nu_\mu$ resonant oscillations in magnetic field are likely to occur in the proximity of such objects.

-
- [1] G. G. Raffelt, *Stars as Laboratories for Fundamental Physics* (University of Chicago Press, Chicago, 1996).
 - [2] C. Giunti and C. W. Kim, *Fundamentals of Neutrino Physics and Astrophysics* (Oxford, UK: University Press, 2007).
 - [3] A. Erdas, C. W. Kim, and T. H. Lee, Phys. Rev. D **58**, 085016 (1998).
 - [4] J. C. D'Olivo, J. F. Nieves and P. B. Pal, Phys. Rev. D **40**, 3679 (1989).
 - [5] P. Elmfors, D. Grasso and G. Raffelt, Nucl. Phys. B **479**, 3 (1996).
 - [6] A. Erdas and G. Feldman, Nucl. Phys. B **343**, 597 (1990).
 - [7] A. Erdas, Phys. Rev. D **80**, 113004 (2009).
 - [8] A. V. Kuznetsov, N. V. Mikheev, G. G. Raffelt and L. A. Vassilevskaya, Phys. Rev. D **73**, 023001 (2006).
 - [9] G. McKeon, Phys. Rev. D **24**, 2744 (1981).
 - [10] E. Elizalde, E. J. Ferrer and V. de la Incera, Annals Phys. **295**, 33 (2002).
 - [11] E. Elizalde, E. J. Ferrer, and V. de la Incera Phys. Rev. D **70**, 043012 (2004).
 - [12] J. F. Nieves and S. Sahu, Eur. Phys. J. C **78**, no. 7, 547 (2018).
 - [13] R. C. Duncan and C. Thompson, Astrophys. J. **392**, L9 (1992).
 - [14] C. Thompson and R. C. Duncan, Mon. Not. Roy. Astron. Soc. **275**, 255 (1995).
 - [15] C. Thompson and R. C. Duncan, Astrophys. J. **473**, 322 (1996).
 - [16] A. Brandenburg, K. Enqvist and P. Olesen, Phys. Rev. D **54**, 1291 (1996).
 - [17] M. Joyce and M. E. Shaposhnikov, Phys. Rev. Lett. **79**, 1193 (1997).
 - [18] D. Grasso and H. R. Rubinstein, Phys. Rept. **348**, 163 (2001).
 - [19] C. Giunti and A. Studenikin, Rev. Mod. Phys. **87**, 531 (2015).
 - [20] A. Studenikin, PoS EPS **-HEP2017**, 137 (2017).
 - [21] C. G. Tarazona, A. Castillo, R. A. Diaz and J. Morales, arXiv:1706.08614 [hep-ph].
 - [22] A. Popov and A. Studenikin, Eur. Phys. J. C **79**, no. 2, 144 (2019).
 - [23] A. Popov and A. Studenikin, arXiv:1803.05755 [hep-ph].
 - [24] J. Schwinger, Phys. Rev. **82**, 664 (1951).




FIG. 2: Plot of $p_\perp$ for which resonance occurs as a function of B , in the case of $\nu_e\text{-}\nu_\tau$ oscillations. $p_\perp$ is in GeV and $0 < B \leq 4B_e$.

- [25] A. V. Kuznetsov and N. V. Mikheev, arXiv:0605114 [hep-ph].
- [26] H. A. Bethe, Phys. Rev. Lett. **56**, 1305 (1986).
- [27] A. V. Borisov, I. M. Ternov and L. A. Vasilevskaya, Phys. Lett. B **273**, 163 (1991).
- [28] L. Wolfenstein, Phys. Rev. D **17**, 2369 (1978).
- [29] L. Wolfenstein, Phys. Rev. D **20**, 2634 (1979).
- [30] S. P. Mikheyev and A. Y. Smirnov, Sov. J. Nucl. Phys. **42**, 913 (1985) [Yad. Fiz. **42**, 1441 (1985)].
- [31] M. Tanabashi *et al.* [Particle Data Group], Phys. Rev. D **98**, no. 3, 030001 (2018).