

Higher Derivative Theory For Curvature Term Coupling With Scalar Field

Pawan Joshi^{1,*} and Sukanta Panda^{1,†}

¹*Department of Physics, Indian Institute of Science Education and Research, Bhopal, India*

Higher order derivative theories, generally suffer from instabilities, known as Ostrogradsky instabilities. This issue can be resolved by removing any existing degeneracy present in such theories. We consider a model involving at most second order derivatives of scalar field non-minimally coupled to curvature terms. Here we perform (3+1) decomposition of Lagrangian to separate second order time derivative from the rest. This is useful to check the degeneracy hidden in the Lagrangian will help us to find conditions under which Ostrogradsky instability do not appear. In our case, we find no such non trivial conditions which can stop the appearance of the Ostrogradsky ghost.

I. INTRODUCTION

Observations suggest that our current universe is in an accelerating phase[1] and explanation of this acceleration can be provided by dark energy. The dark energy is generally modeled by modifying gravity part of Einstein Hilbert action[2]. One of the brands of this modified gravity model is scalar-tensor theories[3, 4]. When we consider scalar-tensor theories with higher order derivative terms. If higher derivative Lagrangian is non-degenerate there exist a ghost-like instability known as Ostrogradsky instability[5] in which Hamiltonian contains such terms which are linear in momentum. In degenerate theory, the higher derivative is present in the Lagrangian but they cancel such a way that does not appear in the equation of motion[6]. In 1974 Horndeski proposed a general action for scalar field that contain the higher derivative term in the Lagrangian but the equation of motion is of second order[7]. Horndeski Lagrangian takes a form,

$$L = L_2^H + L_3^H + L_4^H + L_5^H, \quad (1)$$

where,

$$L_2^H = G_2(\phi, X), \quad L_3^H = G_3(\phi, X)\square\phi, \quad L_4^H = G_4(\phi, X)R - 2G_{4,X}(\phi, X)(\square\phi^2 - \phi_{\mu\nu}\phi^{\mu\nu}), \quad (2)$$

$$L_5^H = G_5(\phi, X)G_{\mu\nu}\phi^{\mu\nu} + \frac{1}{3}G_5(\phi, X)R - (\square\phi^3 - 3\square\phi\phi_{\mu\nu}\phi^{\mu\nu} + 2\phi_{\mu\nu}\phi_\rho^\nu\phi^{\mu\rho}), \quad (3)$$

and $\phi_\mu = \nabla_\mu\phi$, $\phi_{\mu\nu} = \nabla_\mu\nabla_\nu\phi$, $X = \nabla_\mu\phi\nabla^\mu\phi$, R , $R_{\mu\nu}$, $G_{\mu\nu}$ is Ricci scalar, Ricci Tensor and Einstein tensor respectively. Here we notice that In L_2^H is some combination scalar field and its first derivative and L_3^H additionally contain the second derivative of the scalar field, L_4^H , L_5^H contain curvature term and first and second derivative of ϕ . In the case, we have constructed a Lagrangian containing higher derivatives of a scalar field with non-minimal coupling to curvature. Our motivation is to find a degeneracy condition to get rid of Ostrogradsky instability in our higher derivative Lagrangian.

II. POSSIBLE TERMS FOR $\nabla_\mu\nabla_\nu\phi\nabla_\rho\nabla_\sigma\phi$

Consider an action of the form,

$$S = \int d^4x\sqrt{-g}\tilde{C}^{\mu\nu,\rho\sigma}\nabla_\mu\nabla_\nu\phi\nabla_\rho\nabla_\sigma\phi, \quad (4)$$

where $\tilde{C}^{\mu\nu,\rho\sigma}$ contain metric tensor and curvature term and it is only possible term for non-minimal coupling of second derivative of scalar field with the curvature term. Its simple form is,

$$\tilde{C}^{\mu\nu,\rho\sigma} = (D_1g^{\mu\rho}g^{\nu\sigma} + D_2g^{\mu\sigma}g^{\nu\rho})R + (D_3g^{\eta\rho}g^{\mu\nu}g^{\beta\sigma} + D_4g^{\mu\eta}g^{\beta\rho}g^{\nu\sigma})R_{\eta\beta} + (D_5g^{\mu\eta}g^{\nu\beta}g^{\gamma\rho}g^{\delta\sigma} + D_6g^{\mu\eta}g^{\sigma\beta}g^{\gamma\rho}g^{\delta\nu})R_{\eta\beta\gamma\delta}.$$

Now we rewrite the action replace first derivative of scalar field by a new field A_μ i.e. $\nabla_\mu\phi = A_\mu$. So the new form of action is given as,

$$S = \int d^4x\sqrt{-g}[\tilde{C}^{\mu\nu,\rho\sigma}\nabla_\mu A_\nu\nabla_\rho A_\sigma + \lambda^\mu(\nabla_\mu\phi - A_\mu)], \quad (5)$$

* pawanjoshi697@iiserb.ac.in

† sukanta@iiserb.ac.in

this action can be written as,

$$S = \int d^4x \sqrt{-g} [\mathcal{L}_1 + \mathcal{L}_2 + \mathcal{L}_3 + \mathcal{L}_4 + \mathcal{L}_5 + \mathcal{L}_6] + \lambda^\mu (\nabla_\mu \phi - A_\mu), \quad (6)$$

where,

$$\mathcal{L}_1 = D_1 R g^{\mu\rho} g^{\nu\sigma} \nabla_\mu A_\nu \nabla_\rho A_\sigma, \quad (7)$$

$$\mathcal{L}_2 = D_2 R g^{\mu\nu} g^{\rho\sigma} \nabla_\mu A_\nu \nabla_\rho A_\sigma, \quad (8)$$

$$\mathcal{L}_3 = D_3 g^{\eta\mu} g^{\beta\rho} g^{\nu\sigma} R_{\eta\beta} \nabla_\mu A_\nu \nabla_\rho A_\sigma, \quad (9)$$

$$\mathcal{L}_4 = D_4 g^{\eta\rho} g^{\mu\nu} g^{\beta\sigma} R_{\eta\beta} \nabla_\mu A_\nu \nabla_\rho A_\sigma, \quad (10)$$

$$\mathcal{L}_5 = D_5 g^{\mu\eta} g^{\nu\beta} g^{\gamma\rho} g^{\delta\sigma} R_{\eta\beta\gamma\delta} \nabla_\mu A_\nu \nabla_\rho A_\sigma, \quad (11)$$

$$\mathcal{L}_6 = D_6 g^{\mu\eta} g^{\nu\beta} g^{\gamma\rho} g^{\delta\sigma} R_{\eta\beta\gamma\delta} \nabla_\mu A_\nu \nabla_\rho A_\sigma, \quad (12)$$

and λ^μ are Lagrange multiplier. Here it is noted that due to the symmetry property of Riemann tensor L_5 term vanishes.

III. REPRESENTATION OF ACTION IN (3+1) DECOMPOSITION

In this section, we want to find degeneracy condition on Lagrangian, using a (3+1) decomposition. Our (3+1) convention and notation similar to ref.[8, 9]. in (3+1) decomposition $\nabla_a A_b$ is,

$$\nabla_a A_b = \mathcal{D}_a A_b - A_* K_{ab} + n_a (K_{bc} A^c - \mathcal{D}_b A_*) + n_b (K_{ac} A^c - \mathcal{D}_a A_*) + \frac{1}{N} n_a n_b (\dot{A}_* - N^c D_c A_* - N \mathcal{A}_c a^c). \quad (13)$$

Now we introduced $X_{ab} = (D_a A_b - A_* K_{ab})$, $Y_b = (K_{bc} A^c - D_b A_*)$ and $Z = (N^c D_c A_* + N \mathcal{A}_c a^c)$ for simplifying the calculation so eq.(13) becomes,

$$\nabla_a A_b = \frac{1}{N} n_a n_b (\dot{A}_* - Z) - X_{ab} - n_a Y_b - n_b Y_a, \quad (14)$$

where \mathcal{D}_a spatial derivative associated with spatial metric h_{ab} . \mathcal{A}_a and A_* spatial and normal projection of 4-vector A_a given as $\mathcal{A}_a = h_a^b A_b$ and $A_* = n^a A_a$, N^a is shift vector, N is lapse function, $a_b = n^c \nabla_c n_b$ is the acceleration vector and K_{ab} is Extrinsic curvature tensor related to first derivative of metric. In (3+1) formalism $R, R_{\mu\nu}, R_{\mu\nu\rho\sigma}$ are,

$$R = \mathcal{R} + K^2 - 3K_{ab} K^{ab} + 2h^{ab} L_n K_{ab} - 2\mathcal{D}_b a^b - 2a_b a^b, \quad (15)$$

$$\perp R_{ab} = \mathcal{R}_{ab} + K_{ab} K - K_{as} K_b^s + L_n K_{ab} - \mathcal{D}_a a_b - a_a a_b, \quad (16)$$

$$\perp R_{bn} = \mathcal{D}_s K_b^s - \mathcal{D}_b K, \quad (17)$$

$$\perp R_{nn} = K_{st} K^{st} - h^{st} L_n K_{st} + \mathcal{D}_s a^s + a_s a^s, \quad (18)$$

$$\perp R_{abcd} = \mathcal{R}_{abcd} + K_{ac} K_{ad} - K_{ad} K_{bc}, \quad (19)$$

$$\perp R_{abcn} = \mathcal{D}_a K_{bc} - \mathcal{D}_b K_{ac}, \quad (20)$$

$$\perp R_{abnn} = K_{au} K_b^u - L_n K_{ab} + \mathcal{D}_a a_b - a_a a_b, \quad (21)$$

where $L_n K_{ab}$ is Lie derivative of Extrinsic curvature tensor and related to second order derivative of metric. $\perp R_{ab}, \perp R_{bn}, \perp R_{nn}$ is spatial, one normal and two normal projection of Ricci tensor and $\perp R_{abcd}, \perp R_{abcn}, \perp R_{abnn}$ is spatial, one normal and two normal projection of Riemann tensor known as Gauss, Codazzi and Ricci relations respectively.

By using these relation we decompose the action (6) in (3+1) formalism and separate out second order derivative of metric. Next we derive conditions that no second order derivative of metric appear in the action. Here we are not analysing Ostrogradsky instability arising from the higher derivative of scalar field.

IV. CONDITION FOR NO SECOND ORDER DERIVATIVE OF METRIC

In this section, after substituting (13-20) in eq.(6) and keeping the terms that are second derivative of metric.

$$\begin{aligned} \mathcal{L} = h^{ab}L_nK_{ab} & \left(\frac{\dot{A}_*^2}{N^2} - \frac{2\dot{A}_*Z}{N^2} + \frac{Z^2}{N^2} \right) (2D_1 + 2D_2 + D_3 + D_4) + h^{ab}L_nK_{ab}Y_cY^c(-4D_1 - D_3) + h^{ab}L_nK_{ab} \\ & \frac{(\dot{A}_* - Z)X}{N} (4D_2 + D_4) + 2h^{ab}L_nK_{ab}(D_1X^{cd}X_{cd} + D_2X^2) + L_nK_{ab}Y^aY^b(-D_3 - 2D_6) \\ & + L_nK_{ab} \frac{(\dot{A}_* - Z)X^{ab}}{N} (D_4 - 2D_6) + L_nK_{ab}(D_3X_d^aX^{bd} + D_4XX^{ab}) + \text{other terms.} \end{aligned} \quad (22)$$

To be free from terms containing second derivative of metric in the Lagrangian, we require the coefficient of $h^{ab}L_nK_{ab}$ and L_nK_{ab} to vanish. This amounts to a trivial condition $D_1 = D_2 = D_3 = D_4 = D_5 = D_6 = 0$. There is no non trivial condition exists in this case.

V. UNITARY GAUGE

Here we check the possibility to get rid of second derivative of metric in unitary gauge, this gauge gives the condition $\phi(x, t) = \phi_o(t)$. In this case constant time hypersurfaces coincide with uniform scalar field hypersurfaces. When we apply unitary gauge ($\mathcal{A}^a = 0$), demand that the values of X_{ab} , Y_b and Z become, $X_{ab} = -A_*K_{ab}$, $Y_b = -D_bA_*$ and $Z = 0$ respectively, after substituting this result into eq.(22), we get

$$\begin{aligned} \mathcal{L} = h^{ab}L_nK_{ab} \frac{\dot{A}_*^2}{N^2} & (2D_1 + 2D_2 + D_3 + D_4) + \frac{A_*}{N} \dot{A}_*K_c^c(4D_2 + D_4) + 2h^{ab}L_nK_{ab}A_*^2(D_1K_c^dK_d^c + D_2A_*^2K_c^cK_d^d) \\ & + D_4L_nK_{ab} \frac{A_*}{N} \dot{A}_*K^{ab}(D_4 + 2D_6) + L_nK_{ab}(D_3A_*^2K_c^aK^{bc} + D_4A_*^2K_c^cK^{ab}) + \text{other terms.} \end{aligned} \quad (23)$$

In this case also, we have find condition on the condition $D_1 = D_2 = D_3 = D_4 = D_5 = D_6 = 0$. There is no non trivial condition exists in this case.

VI. CONCLUSION

In this paper, we work with higher derivative model where both second derivative of metric and scalar field arise in the Lagrangian. Then using (3+1) decomposition, we have shown that no non trivial conditions can be found under which all the terms containing second order derivative of metric disappear from the Lagrangian.

VII. ACKNOWLEDGEMENT

This work was partially funded by DST (Govt. of India), Grant No. SERB/PHY/2017041. Calculations were performed using xAct packages of Mathematica.

-
- [1] A. G. Riess *et al.* (Supernova Search Team), *Astron. J.* **116**, 1009 (1998), arXiv:astro-ph/9805201 [astro-ph].
 - [2] S. Nojiri and S. D. Odintsov, *Theoretical physics: Current mathematical topics in gravitation and cosmology. Proceedings, 42nd Karpacz Winter School, Ladek, Poland, February 6-11, 2006*, eConf **C0602061**, 06 (2006), [Int. J. Geom. Meth. Mod. Phys.4,115(2007)], arXiv:hep-th/0601213 [hep-th].
 - [3] J. Gleyzes, D. Langlois, and F. Vernizzi, *Int. J. Mod. Phys.* **D23**, 1443010 (2015), arXiv:1411.3712 [hep-th].
 - [4] B. Ratra and P. J. E. Peebles, *Phys. Rev. D* **37**, 3406 (1988).
 - [5] M. Ostrogradsky, *Mem. Acad. St. Petersburg* **6**, 385 (1850).
 - [6] A. Nicolis, R. Rattazzi, and E. Trincherini, *Phys. Rev.* **D79**, 064036 (2009), arXiv:0811.2197 [hep-th].
 - [7] G. W. Horndeski, *International Journal of Theoretical Physics* **10**, 363 (1974).
 - [8] T. W. Baumgarte and S. L. Shapiro, *Numerical relativity: solving Einstein's equations on the computer* (Cambridge University Press, 2010).
 - [9] D. Langlois and K. Noui, *Journal of Cosmology and Astroparticle Physics* **2016**, 034 (2016).