

Comment (2) on “Quantum Transfiguration of Kruskal Black Holes”

Martin Bojowald*

*Institute for Gravitation and the Cosmos, The Pennsylvania State University,
104 Davey Lab, University Park, PA 16802, USA*

An explicit calculation shows that the model constructed in [1] violates general covariance. It is therefore unphysical and does not determine properties of space-time or black holes.

The authors of [1] attempt to modify the dynamics of gravitational collapse by viewing the static, spherically symmetric exterior space-time region of a Kruskal black hole as a homogeneous model with timelike hypersurfaces “evolving” in the direction of the radial coordinate. However, the authors never consider the spherically symmetric, static description after they modify the dynamics. This description, in fact, is no longer equivalent, owing to a breakdown of general covariance in the model.

An \hbar -dependent effective theory is covariant if and only if it is covariant order by order in \hbar . It is therefore sufficient to consider only leading-order modifications, parameterized in [1] by δ_b and δ_c . This leads to the cosmological Hamiltonian

$$H_{\text{hom}}[N] = -\frac{N}{2G} \left(2bc\sqrt{p_c} + (1-b^2)\frac{p_b}{\sqrt{p_c}} \right. \quad (1)$$

$$\left. + \frac{1}{3}b^3(c\sqrt{p_c} - bp_b/\sqrt{p_c})\delta_b^2 - \frac{1}{3}bc^3\sqrt{p_c}\delta_c^2 \right)$$

for an arbitrary lapse function N , Newton’s constant G , and canonical pairs (p_b, b) and (c, p_c) . (In the specific choice of [1], δ_c is of higher order in \hbar than δ_b and could therefore be eliminated here without changing our result. We assume $p_c > 0$ and ignore parameters such as γ and L_0 that are irrelevant for our argument.)

The meaning of p_b and p_c is shown by how they appear in the space-time line element

$$ds^2 = -\frac{p_b^2}{p_c}dx^2 + N^2dt^2 + p_c d\Omega^2, \quad (2)$$

while b and c are related to time derivatives of p_b and p_c according to equations of motion generated by (1). Inverting these equations, we obtain

$$b = \frac{\dot{p}_c}{2N\sqrt{p_c}} - \frac{1}{6}b^3\delta_b^2 + \frac{1}{2}bc^2\delta_c^2 \quad (3)$$

$$c = -\frac{\dot{p}_b}{N\sqrt{p_c}} + b\frac{p_b}{p_c} - \frac{1}{6}b^2(3c - 4bp_b/p_c)\delta_b^2 + \frac{1}{6}c^3\delta_c^2 \quad (4)$$

They are valid if δ_b and δ_c depend on p_b or p_c . If δ_b and δ_c also depend on b or c , (3) and (4) acquire extra terms that do not change our result of non-covariance. We work with (3), (4) because their solutions (to all orders in δ_b and δ_c) have been used in [1]; see also [2].

We now evaluate the same dynamics from the perspective of a spherically symmetric model with line element

$$ds^2 = -M^2dT^2 + \frac{(E^\varphi)^2}{E^x}dX^2 + E^x d\Omega^2 \quad (5)$$

where E^x and E^φ are the components of a densitized triad analogous to p_c and p_b . Comparing (2) and (5) such that $T = x$ and $X = t$, we obtain the correspondence $p_c = E^x$, $p_b = M\sqrt{E^x}$, $N = E^\varphi/\sqrt{E^x}$, while b and c follow from (3) and (4) observing that the dot amounts to an X -derivative (a prime) in spherical symmetry.

If there is a covariant space-time underlying this model, the Hamiltonian constraint (1), using our correspondence, must vanish when expressed in spherically symmetric models. After several cancellations, we obtain

$$-2GH_{\text{hom}}[N] = -\frac{M'(E^x)'\sqrt{E^x}}{E^\varphi} + \frac{ME^\varphi}{\sqrt{E^x}} - \frac{M((E^x)')^2}{4\sqrt{E^x}E^\varphi} \quad (6)$$

$$+ \frac{1}{8}\frac{(M\sqrt{E^x})'((E^x)')^3}{(E^\varphi)^3}\delta_b^2 - \frac{1}{2}\frac{(M')^3(E^x)'(E^x)^{3/2}}{(E^\varphi)^3}\delta_c^2.$$

The lapse function M of the spherically symmetric model is not independent but rather determined (non-locally) by E^x and E^φ through staticity (zero momenta). This condition amounts to a differential equation

$$M' + \frac{1}{4}\frac{(E^x)'}{E^x}M + \frac{1}{2}\frac{(E^\varphi)^2}{\sqrt{E^x}(E^x)'}V(E^x)M = 0, \quad (7)$$

which, for full generality, we use here for a general dilaton model [3] with potential $V(E^x)$; see for instance [4, 5]. (These models are broad and include, for instance, spherically symmetric Palatini- $f(R)$ theories through their equivalence with scalar-tensor metric theories with a non-dynamical scalar [6].) Classically, (6) implies

$$-2GH_{\text{hom}}[N] = \frac{ME^\varphi}{\sqrt{E^x}} \left(1 + \frac{1}{2}\sqrt{E^x}V(E^x) \right) = 0 \quad (8)$$

which selects the correct dilaton potential $V(E^x) = -2/\sqrt{E^x}$ of classical spherically symmetric gravity. Non-zero δ_b and δ_c imply additional terms in (6) which depend on M and E^φ . They cannot be absorbed in a modified dilaton potential $V(E^x)$. Since dilaton gravity actions present the most general class of generally covariant 1+1-dimensional theories, the model of [1] violates covariance.

Note added: A transformation of the cosmological-type solutions given in [1] to spherically symmetric slices has already been performed in [7], focusing on implications for the asymptotic structure. In the present comment, an extension of such a transformation to the model’s constraints and equations of motion makes it possible to draw conclusions about covariance.

The author thanks Norbert Bodendorfer and Sudhasattwa Brahma for comments. This work was supported in part by NSF grant PHY-1607414.

* Electronic address: bojowald@gravity.psu.edu

[1] A. Ashtekar, J. Olmedo, and P. Singh, Phys. Rev. Lett. **121**, 241301 (2018).

[2] N. Bodendorfer and J. Mele, F. M. abd Münch, arXiv:1902.04032.

[3] T. Strobl, hep-th/0011240.

[4] M. Bojowald and J. D. Reyes, Class. Quantum Grav. **26**, 035018 (2009).

[5] M. Bojowald and S. Brahma, Phys. Rev. D **95**, 124014 (2017).

[6] G. J. Olmo, Phys. Rev. Lett. **95**, 261102 (2005).

[7] M. Bouhmadi-López *et al.*, arXiv:1902.07874.