

How the modified Bertrand theorem explains regularities of the periodic table I. From conformal invariance to Hopf mapping

Arkady Kholodenko^{a1} and Louis Kauffman^b

^a 375 H.L. Hunter Laboratories, Clemson University, Clemson, SC 29634-0973, USA.

^b Department of Mathematics, Statistics and Computer Science, University of Illinois at Chicago, 851 S.Morgan Street, Chicago, IL, 60607-7045, USA

Abstract

Bertrand theorem permits closed orbits in 3d Euclidean space only for 2 types of central potentials. These are of Kepler- Coulomb and harmonic oscillator type. Volker Perlick recently extended Bertrand theorem. He designed new static spherically symmetric (Bertrand) spacetimes obeying Einstein's equations and supporting closed orbits. In this work we demonstrate that the topology and geometry of these spacetimes permits to solve quantum many-body problem for any atom of periodic system exactly. The computations of spectrum for any atom of periodic system becomes analogous to that for hydrogen atom. Initially the exact solution of the Schrödinger equation for any multielectron atom was obtained by Tietz in 1956. However, neither himself nor others fully comprehended what actually was obtained. We recalculated Tietz results by applying the methodology consistent with new (different from that developed by Fock in 1936) way of solving Schrödinger's equation for hydrogen atom. In the light of this new result it had become possible to demonstrate rigorously that the Tietz- type Schrödinger's equation is in fact describing the quantum motion in Bertrand spacetime. As a bonus, we obtained the analytical proof of the Madelung rule defined in the text.

Keywords:

Maxwell's fish-eye potential,
conformally invariant classical and quantum equations,
superintegrability, quantum Bertrand spacetimes

MSC 2010: 81Q05, 81Q99, 83C15, 83C20

1. Background and summary

Although quantum mechanical description of multielectron atoms and molecules is considered to be a well established domain of research, recently published book [1] indicates that

¹Corresponding author.

E-mail address: string@clemson.edu

there are still many topics left for development. As is well known, the quantum mechanical description of multielectron atom (with atomic number Z and infinitely heavy nucleus) begins with writing down the stationary Schrödinger equation

$$\hat{H}\Psi(\mathbf{r}_1, \mathbf{r}_2, \dots, \mathbf{r}_Z) = E\Psi(\mathbf{r}_1, \mathbf{r}_2, \dots, \mathbf{r}_Z) \quad (1.1a)$$

with the Hamiltonian

$$\hat{H} = -\sum_{i=1}^Z \frac{\hbar^2}{2m} \nabla_i^2 - \sum_{i=1}^Z \frac{Ze^2}{r_i} + \frac{1}{2} \sum_{\substack{i,j=1 \\ i \neq j}}^Z \frac{e^2}{r_{ij}}. \quad (1.1b)$$

Following Bohr's *Aufbauprinzip* the atom with atomic number Z is made up of electrons added in succession to the bare atomic nucleus. At the initial stages of this process electrons are assumed to occupy the one-electron levels of lowest energy. This process is described in terms of the one electron eigenvalue problem

$$\hat{H}_i \psi_{\square_i}(r_i) = \left[-\frac{\hbar^2}{2m} \nabla_i^2 + V_{eff}(\mathbf{r}_i) \right] \psi_{\square_i}(\mathbf{r}_i) = \varepsilon_{nl}(i) \psi_{\square_i}(\mathbf{r}_i), i = 1 \div Z, \quad (1.2)$$

where $V_{eff}(\mathbf{r}_i)$ is made of the combined nuclear potential $-\frac{Ze^2}{r_i}$ for the i -th electron and the centrally symmetric Hartree-Fock type potential $\mathcal{F}(\mathbf{r}_i)$ coming from the presence of the rest of atomic electrons. The fact that $\mathcal{F}(\mathbf{r}_i)$ is centrally symmetric was discussed, for example, in the book by Bethe and Jackiw [2]. Later on in the text and, especially, in the Appendix G, we provide much more details on this topic. The symbol \square_i indicates the i -th entry into the set made out of hydrogen-like quantum numbers characterizing individual electrons. Recall that the concept of an *orbital* is determined by the major quantum number n having its origin in studies of hydrogen atom. The number of electrons allowed to sit on a given orbital is determined by the *Pauli exclusion principle*. Thus, with increasing Z the electrons are occupying successive orbitals according to Bohr's Aufbau scheme until the *final ground state electron configuration* is reached. Since electrons are indistinguishable, the hydrogen-like quantum numbers n, l, m and m_s cannot be associated with a particular electron. Therefore, the symbol \square_i should be understood as representing a specific set of quantum numbers otherwise used for description of individual (that is not collectivized) electrons.

The problem with just described *Aufbauprinzip* lies in the assumption that the guiding principle in designing the final ground state electron configuration is made out of two components: a) knowledge of hydrogen atom-like wave functions supplying the quantum boxes/numbers \square_i and, b) the Pauli principle which is mathematically restated in the form of the fully antisymmetric wavefunction $\Psi(\mathbf{r}_1, \mathbf{r}_2, \dots, \mathbf{r}_Z)$. Should these requirements be sufficient then, we would be able to replace $V_{eff}(\mathbf{r}_i)$ by $-\frac{Ze^2}{r_i}$ so that the filling of electronic levels would occur according to the Fock n -rule

Fock n -rule: *With increasing Z the nl orbitals are filled in order of increasing n .*

This rule leads to the problems already for the lithium as explained in [1], page 330. As result, the (n, l) rule was proposed instead.

The hydrogenic (n,l) rule: *With increasing Z , the orbitals are filled in order of increasing n while for a fixed n the orbitals are filled in order of increasing l .*

After $Z = 18$ the (n,l) rule breaks down though. Therefore, it was subsequently replaced by the $(n+l,n)$ rule suggested by Madelung- the person who reformulated Schrödinger's equation in hydrodynamic form [3].

The Madelung $(n+l,n)$ rule: *With increasing Z , the orbitals are filled in order of increasing $n+l = N$. For fixed N , the orbitals are filled in order of increasing n .*

All the above rules are empirical. As such, they require theoretical explanation. This fact brings us to the

Löwdin's challenge problem: Find a way to derive the Madelung rule ab initio. ².

The essence of Mendeleev's periodic system of elements lies exactly in the discovered periodicity of properties of chemical elements. Although there are 100's of ways this periodicity is exhibited³, the commonly accepted periodic table of elements consists of seven periods: 2-8-8-18-18-32-32. Notice that all period lengths occur in pairs (period doubling), except for the very first period of size 2. To determine whether this exception is intrinsic or not, analysis of work by Charles Janet on periodic table done in 1930 (6 years before work by Madelung!) is the most helpful, [1], pages 336-340. Although initially Janet developed his version of periodic table without guidance of quantum mechanics, eventually he did make a connection with Bohr's results. Janet's periodic table has 8 periods. The periods in Janet's table are characterized (without exception) by the constant value of $N = n + l$. It is in excellent agreement with the Madelung rule suggesting elevation of $N = n + l$ to the rank of new quantum number. By organizing the elements in periods of constant $n + l$ and groups of constant l, m_l and m_s , the period doubling emerges naturally and leads to the sequence of periods: 2-2-8-8-18-18-32-32.

A concise and convincing explanation of the period doubling and its connection with the Madelung rule is given in [6]. The Madelung rule had attracted the attention of Demkov and Ostrovsky in 1971 resulting in their publication [7] of major importance. Nevertheless, subsequently Kitagawara and Barut [8,9] found apparent flaws in the logic of Demkov-Ostrovsky calculations. The authors of the book [1] also expressed their objections to the Demkov-Ostrovsky cycle of works. On page 381 of [1] we found the following statement: "Demkov and Ostrovsky developed an atomic physics model that incorporates the Madelung rule, but by replacing the quantization of level energies with quantization of coupling constants at zero energy." Furthermore, Demkov and Ostrovsky, while obtained the correct results, had been unable to provide their rigorous justification since their effective potential $V_{eff}(\mathbf{r}_i)$ was correctly guessed. The authors of [1] conclude that, even though the Demkov -Ostrovsky results do reproduce the Madelung rule correctly, the way these results were obtained cannot be considered as solution of the Löwdin challenge.

²This problem was posed by Per-Olov Löwdin [4]. Additional details and references are given in [5].

³Results of www searches indicate that this process is still ongoing.

In this work we demonstrate that the objections raised in [1, 8, 9] are coming from the lack of knowledge of needed mathematical apparatus. Thus, one of the tasks of this work is to describe this apparatus. By doing so, a number of problems of major practical importance is solved and all of the objections raised in [1, 8, 9] are removed. Some additional objections to be described shortly below are also removed. In their seminal work [7] Demkov-Ostrovsky (D-O) realized that the key to success of solving problems of periodic table lies is Eq.(1.2) where $V_{eff}(\mathbf{r}_i)$ should be correctly chosen. The Bertrand theorem of classical mechanics [10] imposes seemingly insurmountable restrictions on selection of $V_{eff}(\mathbf{r}_i)$ since for spherically symmetric potentials only the Coulombic $-\frac{Ze^2}{r_i}$ and the harmonic oscillator kr^2 potentials allow dynamically closed orbits. D-O believed that, in spite of the indistinguishability of electrons, the discrete spectrum of multielectron atoms should be associated with closed orbits. From the point of view of theory of chaos⁴ at the semiclassical level of description the role of closed orbits recently was discussed, e.g. in [11] and, much earlier, in the seminal book by Gutzwiller [12]⁵.

Hoping to bypass limitations of the Bertrand theorem, D-O employed the optical-mechanical analogy in their calculations. It permitted them to use the Maxwell fish-eye potential (and, later on, its conformally deformed modifications) instead of the Coulombic potential. They demonstrated in [7] the equivalence at the classical level between the Hamilton-Jacobi equations employing these two, seemingly different, potentials. D-O hoped that use of the fish-eye potential (or its conformal modification) might allow to bypass limitations of the Bertrand theorem. This fact attracted attention of John Wheeler who in [14, 16] and, with his student, in [15], studied classically and semiclassically the motion in the fish-eye and conformally deformed fish-eye potentials used by D-O in [7]. The dynamics of electrons in such conformally deformed potentials happens to involve orbits which are closed, planar and have self-intersections. In his paper [17] Ostrovski argued that the self-intersections of orbits do not contradict the Bertrand theorem. This statement by Ostrovski happens to be wrong as demonstrated in section 5. In it, we mention works by other authors who obtained the same results prior to the discovery of curved Bertrand spacetimes. None of these authors was concerned with the limitations of Bertrand theorem. Very recently, in 2017, the same self-intersecting patterns were obtained for dynamical trajectories in Bertrand spacetimes, e.g. see page 3362 of [18]. The results were obtained without any reference to atomic physics.

According to work by Little [19] there are only 3 distinct regular homotopy classes of oriented closed curves on S^2 . These are: a) those for curves without self-intersections, b) those for curves with just 1 self-intersection and, c) those with 2 self-intersections. The b)-type homotopy class of curves were obtained by Wheeler and by other authors as discussed in section 5. Thus, from the work by Little it follows that the effects of curvature and presence of self-intersection in dynamical trajectories are connected with each other. The Kepler-Coulomb dynamics in flat 3d Euclidean space does not allow self-intersections. The self-intersections

⁴Beginning from the motion of electrons in helium, all atomic multielectron systems are believed to be chaotic,

⁵See also the book by Cvitanovic [13].

are allowed if the Bertrand theorem is extended to motions on curved (Bertrand) manifolds [18]. The sphere S^2 is conformally flat⁶. The results of the modified Bertrand theorem are valid exactly for the conformally flat manifolds. Details are given in section 5. These are corollaries of results obtained in [20]. Thus, flatness of self-intersecting patterns obtained in [18] is in fact conformal flatness.

To connect Bertrand spacetimes with atomic physics we begin with D-O statement made in [7] : "The Maxwell's fish-eye problem is **closely related** to the Coulomb problem". Being aware of the book by Luneburg [21]⁷ D-O nevertheless underestimated the nature of connection between the Coulombic and optical (fish-eye) problems described in [21]. This fact is discussed in detail in section 2. The assumption of only "close relationship" caused D-O to replace Eq.(1.2) by

$$[-\frac{\hbar^2}{2m}\nabla_i^2 + V_{eff}(\mathbf{r}_i)]\psi(\mathbf{r}_i) = 0. \quad (1.3)$$

Eq.(1.3) is looking differently from Eq.(1.2). Eq.(1.2) is the eigenvalue problem while Eq.(1.3) is the Sturmian problem. For such type of problem to make sense the parameters entering into $V_{eff}(\mathbf{r}_i)$ must be quantized. Such quantization of parameters is making Sturmian and eigenvalue problems equivalent. This fact is nontrivial. It is considered in detail in sections 2 and 4 and, in Appendices A,B, E and F. Without further explanations this equivalence represents the major reason for using the fish-eye-type potentials in treatments of atomic systems. In fact, in many ways (discussed in section 3 and Appendices C, D and F) to use Eq.(1.3) is more advantageous than Eq.(1.2).⁸

In [1], page 377, we read that Eq.(1.3) "does not describe the bound states of the atom". That this is not the case could be concluded already by D-O themselves should they read the corresponding places in books by Luneburg [21] and Caratheodory [22]. They do quote these works though in [7]. This misunderstanding of importance of Eq.(1.3) and its relation with Eq.(1.2) resulted in critique and neglect of D-O works, e.g. read [1], page 377. In this paper we are going to demonstrate that Maxwell's fish-eye and related to the fish-eye classical and quantum Coulombic (hydrogen atom) problems are not **closely related**. Instead, **they are isomorphic**. This is illustrated in section 4. By overlooking the Coulombic-fish-eye isomorphism at the quantum level D-O argued, nevertheless, that Schrödinger's Eq.(1.3) with Coulombic and fish-eye-type potentials **both** possess the $O(4,2)$ (or $SO(4,2)$) dynamical symmetry known for the hydrogen atom and later established for the rest of atoms of the periodic table [1], [3]. Selection of the fish-eye-like potentials by D-O was guided in part by their desire to describe the *atoms other than hydrogen*. In addition, they believed that: a) $V_{eff}(\mathbf{r}_i)$ in Eq.(1.3) can be represented by the conformally deformed fish-eye potential so that b) use of such potential removes restrictions posed by the Bertrand theorem in flat space. Ostrovsky concludes his paper [17] with the following remark:

⁶This concept was defined in our work [3] and is going to be also explained in the bulk of this work.

⁷Ref.4 in [7].

⁸Mathematically, there is alternative proof of this fact which is interesting on its own. It will be discussed in a separate publication.

"It ⁹ leaves a very interesting question unresolved, the question of why the interaction of a number of electrons with each other and with an atomic nucleus leads to an effective one-electron potential having some approximate hidden symmetry. The method of solution of this question can hardly be envisaged at the present time." Thus, with complete formal success of quantum description of atoms of the whole periodic system [7] culminating in formal proof of the Madelung rule, Ostrovsky admits that with all his results published to date, the Löwdin's challenge problem still remains out of reach. This is so because the deformed fish-eye potential used in D-O calculations had no visible connection with $V_{eff}(\mathbf{r}_i)$ coming from the Hartree-Fock calculations. In addition to their inability of solving the Löwdin challenge, D-O also failed to solve the Bertrand challenge problem¹⁰: What makes the deformed fish-eye potential used by D-O as substitute of $V_{eff}(\mathbf{r}_i)$ so good that it removes the restrictions of the classical Bertrand theorem?

In their cycle of works on proving the Madelung rule D-O used the fish-eye ($\gamma = 1$) and conformally deformed ($\gamma \neq 1$) fish-eye potentials

$$V(x, y, z) = - \left(\frac{a}{r} \right)^2 \left[\frac{n_0}{(r/a)^{-\gamma} + (r/a)^\gamma} \right]^2, \quad (1.4)$$

$r^2 = x^2 + y^2 + z^2, a = const, \gamma$ is a rational number, as an alternative to the $V_{eff}(\mathbf{r}_i)$ Hartree-Fock-type potentials routinely used in atomic physics literature. Such a replacement required them to switch from Eq.(1.2) to Eq.(1.3) for reasons explained in detail below, in sections 2 and 3 and Appendices A and B. Since Eq.(1.3) seemingly allows only to look for eigenfunctions with zero eigenvalue, both D-O and the rest of researchers in the field considered this limitation as a serious deficiency. In this work we demonstrate that this restriction is harmless and, in fact, very helpful. We remind to our readers that use of the potential, Eq.(1.4), was made by D-O for the purpose of taking care of limitations of the classical Bertrand theorem. No other authors were concerned with these limitations. Apparently, this fact allowed D-O to neglect works by other authors on the same or related subjects and to draw attention of others [1] only to their own works. This happens to be a fundamental drawback. Details are given in section 5 and Appendices F and G.

Below, we provide a summary of the results just sketched. D-O realized that when used in Eq.(1.3) the constant n_0 acquires discrete values as it happens in all Sturmian type problems (Appendix F). Furthermore, for $\gamma = 1/2$ the solution of Eq.(1.3) provides results compatible with the Madelung rule. The apparent limitation, $E = 0$, along with (in their expositions) no apparent relationship between the potential $V(x, y, z)$ and $V_{eff}(\mathbf{r})$ caused Ostrovski to acknowledge (in [17]) that all D-O results to date do not solve the Löwdin challenge problem. Thus, we are left with the following (unproven yet) facts:

- a) use of the potential, Eq.(1.4), removes the limitations of the classical Bertrand theorem;
- b) the choice $\gamma = 1/2$ in Eq.(1.4) apparently explains the Madelung rule;

⁹...that is the group-theoretical (our insert from previous discussion) consideration...

¹⁰We are posing the Bertrand challenge problem at this point in addition to the Löwdin challenge problem known for some time

c) finding of the spectral results beyond $E = 0$ requires (according to D-O) uses of sophisticated perturbational methods;

d) the choice $\gamma = 1/2$ in Eq.(1.4) is detached from known Hartree-Fock results;

e) the case $\gamma = 1$ corresponds to the standard Maxwell's fish-eye potential. Classical dynamics in such

potential is isomorphic to that in Kepler-Coulomb potential. However, because of the apparent $E = 0$ limitation,

neither D-O nor other researchers reproduced known eigenvalue spectrum for the hydrogen atom.

Subsequently, other authors studied Eq.(1.3) with D-O potential, Eq.(1.4), in 2 dimensions where use of conformal transformations leaves Eq.(1.3) form-invariant. This fact provides many advantages. Still, no attempts to reproduce known 2 dimensional results for hydrogen atom were made.

In our work we use the observation, in section 4, that results on \mathbf{R}^2 can be lifted to S^2 and then lifted further to S^3 via inverse Hopf mapping. These lifting procedures will be discussed in detail in our next paper, Part II. Using stereographic projection: from S^3 to \mathbf{R}^3 , it is possible then to reobtain D-O results done in \mathbf{R}^3 . Even though the connection between the Hartree-Fock and the D-O potential, Eq.(1.4), will be discussed in detail from geometrical and topological perspective in Part II, already available results allow us to discuss rigorously such a connection in Part I, that is this paper. Details are given in Appendix G and section 5. The noticed connection is inseparably linked with the validity of the Madelung rule as explained in Appendix G and section 5.

Going back to a) we direct our readers attention to work [23] by Volker Perlick. In it the results of classical Bertrand theorem [10] valid in Euclidean 3 space had been generalized to static spherically symmetric spacetimes of general relativity. By design, the motion in such spacetimes takes place on closed orbits. In section 5 we demonstrate that the potential, Eq.(1.4), indeed removes limitations of the classical Bertrand theorem. However, it is used not in flat Euclidean \mathbf{R}^3 but in curved Bertrand spacetime.

The choice $\gamma = 1/2$ listed in b) and d) is indeed connected directly with the results of Hartree-Fock calculations and with the Madelung rule. In atomic physics literature the potential, Eq.(1.4), $\gamma = 1/2$, is known as the Tietz potential¹¹. Its origin and many properties are discussed in the book by Flugge [24]. Its remarkable numerical coincidence with the corresponding Hartree-Fock potentials were discussed in many places, e.g. read [25], p.664, fig.10. Tietz, the author who invented the Tietz potential, was initially driven by the desire to simplify the Thomas -Fermi (T-F) calculations. Much more analytically cumbersome T-F type potentials were used by Latter [26] in his numerical study of the Schrödinger equation spectra of low lying excitations for all atoms of periodic system. Numerical results of Latter had been subsequently analyzed by March. On page 76 of [27] without explicit mention of the Madelung rule he described results by Latter in terms of the Madelung rule. After discovery of his potential Tietz used it in the stationary Schrödinger equation, Eq.(1.2), in which $V_{eff}(\mathbf{r}_i)$ was replaced by the Tietz potential, that is by Eq.(1.4) with $\gamma = 1/2$ [28].

¹¹It is bearing the name of its creator.

Tietz used Eq.(1.2) in which $E \neq 0$ instead of D-O version of this equation, Eq.(1.3). In the light of results of Appendix F, this happens to be permissible. Tietz also recognized that Eq.(1.2) with his potential can be solved exactly. His first attempt to do so was initially made in 1956 [28]. The rest of his attempts is summarized in [29]. The exact solution methods in these works differ in style from that used by D-O. Besides, Tietz used these exact solutions only to check them against the Hartree-Fock results for the Mercury ($Z = 80$). In doing so he got a good agreement with the Hartree-Fock results. In this work we prove that the Tietz potential used in [28], [29] is describing the classical and quantum motion in Bertand spacetimes. Unlike Tietz and, in accord with D-O, we used Eq.(1.3) for solving the corresponding eigenvalue problem. Our method of solving this equation differs somewhat from that used by D-O though since their method is unable to reproduce the hydrogen atom spectrum (the problem e)). Our way of solving Eq.(1.3) with $\gamma = 1/2$ is consistent with our new way (different from that developed by Fock in 1936 (discussed in [3])) of obtaining the spectrum of hydrogen atom. We were able to reproduce this spectrum correctly also in 2 dimensions. These facts allowed us to bypass entirely the item c) present in D-O works. They also allow us to obtain the low lying spectrum of any atom of periodic system consistent with the Madelung rule. In another paper of ours [30], we discuss some exceptions to the Madelung rule¹² and theoretical means of explaining these exceptions. More details are still forthcoming.

2. Known and unknown properties of Maxwell's fish-eye lenses and potentials associated with them. Their role in optics and quantum mechanics

Maxwell's fish eye lens was invented by James Clerk Maxwell. It is nicely described in [6],[21],[22]. Its invention was motivated by the desire to make perfect (absolute) optical instrument. In such an instrument it is expected that *all* rays leaving point \mathbf{x} meet again at the (*conjugated or focal*) point \mathbf{x}' . Mathematically this can be expressed with help of the Fermat principle. It is described as follows. Introduce the *optical path* length $S(\mathbf{x}, \mathbf{x}')$ via the length functional

$$S(\mathbf{x}, \mathbf{x}') = \int_0^t dl, \quad (2.1)$$

where the conformally flat metric dl^2 is given by $dl^2 = n^2(\mathbf{x}(t)) \left(\frac{d\mathbf{x}}{dt} \right)^2 dt^2$ with $n(\mathbf{x}(t))$ being position-dependent refractive index of the medium.

Fermat principle: Light rays through spatial points $\mathbf{x}(0)$ and $\mathbf{x}'(t)$ move on optically shortest paths .

Based on this definition, it follows that minimization of the functional, Eq.(2.1), results in the Euler-Lagrange equations for geodesics in the optical medium in analogy with equations

¹²https://en.wikipedia.org/wiki/Aufbau_principle

for the point-like particle moving in the gravitational field [31],[32]. We shall demonstrate below that such noticed known optical-mechanical-gravitational correspondence is capable of producing new results. Being armed with the above information, now we are in the position to describe the origins of Maxwell's fish-eye lens (additional information is given in Appendix A). For this purpose, let us consider the n -dimensional sphere S^n in the flat Euclidean space \mathbf{R}^{n+1} described by

$$S^n = \{(z_1, \dots, z_n, y) \mid \sum_{i=1}^n z_i^2 + y^2 = 1\}. \quad (2.2)$$

The stereographic projection: from \mathbf{R}^n to S^n , is given by

$$z_i = \frac{x_i}{1 + \frac{1}{4}\mathbf{x}^2}, y = \frac{1 - \frac{1}{4}\mathbf{x}^2}{1 + \frac{1}{4}\mathbf{x}^2}, i = 1 \div n, x_i \in \mathbf{R}^n, \mathbf{x}^2 = x_i x^i. \quad (2.3)$$

Here and below the summation over repeated indices is assumed. From the mathematical literature [33], [34] it is known that the stereographic projection is a conformal mapping since it is angle-preserving. This means that using Eq.(2.3) it is possible to rewrite the metric ds^2 on the sphere S^n in terms of the metric on \mathbf{R}^n , that is

$$ds_{S^n}^2 = \rho(\{x_i\}) (d\mathbf{x})^2, \quad (2.4a)$$

where

$$\rho(\{x_i\}) = \left(\frac{1}{1 + \frac{1}{4}\mathbf{x}^2} \right)^2. \quad (2.4b)$$

An example of such relation, e.g. between the metric of the 2-sphere S^2 and the metric of the 2-plane \mathbf{R}^2 is described in detail in [34], page 88. Since $\mathbf{R}^2 + \{\infty\} = S^2$. The same is true for the spheres of any dimension. The geodesics on S^n are the great circles. All great circles passing through some point P on S^n intersect again in the conjugate point P' , that is in the diametrically opposite point $-P$ (the antipodal map [33]). This circumstance can be used for design of perfect optical instruments. Specifically, let us adopt Eq.(2.4a) to 4 and 3 dimensions respectively. Then, in view of Eq.(2.1), the conformal factor $\rho(\{x_i\})$ can be interpreted as the square of refractive index $n^2(\mathbf{x}(t))$. This explains the origin of Maxwell's fish-eye refractive index. Written with account of dimensionality arguments, the refractive index is given by

$$n(\mathbf{x}) = \frac{n_0}{1 + \mathbf{x}^2/a^2}. \quad (2.5)$$

The light rays in 3 dimensional Maxwell's fish-eye medium are images of the great circles, that is of *the geodesics*, in S^3 . The antipodal map $P \rightarrow -P, P \in S^3$, corresponds to the negative inversion map $x'_i = -\frac{x_i}{|\mathbf{x}|^2}, x_i \in \mathbf{R}^3$ from \mathbf{R}^3 to \mathbf{R}^3 . The antipodal points on S^3 correspond to points \mathbf{x}_1 and \mathbf{x}_2 in \mathbf{R}^3 which lie on a common line through the origin, and satisfy $\mathbf{x}_1 \cdot \mathbf{x}_2 = -1$. The great circles on S^3 are just those circles (or lines through the origin in \mathbf{R}^3) which map into themselves under the negative inversion operation [33].

These facts, we hope, remind our readers about the classical (by Moser [35, 36]) and quantum mechanical (Fock [37]) treatments of hydrogen atom respectively. They were discussed to some extent in our previous work [3]. The connection between the description of optical (fish-eye refractive index) instruments and the classical/quantum mechanical description of hydrogen atom was discovered by Luneburg [21]. Some important additional details (not of quantum mechanical nature though) were independently developed by Caratheodory [22]. They are discussed in Appendix A. Contrary to the D-O claim [7] that the Maxwell's fish-eye and the Coulomb (that is hydrogen atom) problems are "closely related", Luneburg found the exact mapping between these problems. To avoid further literature interpretation ambiguities, we present Luneburg's arguments below.

To shorten our exposition to the absolute minimum, we extensively use the results of our previous work [3]. It also discusses results of Luneburg's book [21], albeit with different purpose. Thus, we begin with Eq.(4.11) of [3] for the wavefront. We write it in Luneburg's notations (e.g. see Eq.(28.13) of [21]),

$$\psi_x^2 + \psi_y^2 + \psi_z^2 = n^2(x, y, z) \quad (2.6)$$

and select for the refractive index

$$n^2(x, y, z) = C + \frac{1}{\sqrt{x^2 + y^2 + z^2}}. \quad (2.7)$$

If this equation is to be used quantum mechanically, Eq.(2.6) must be rewritten in the form of Eq.(4.21) of [21], that is in the form

$$\psi_x^2 + \psi_y^2 + \psi_z^2 = 2m(E - V). \quad (2.8a)$$

Then, following Schrödinger, the fully reversible substitution: $\psi \rightleftharpoons \hbar \ln \varphi$ done in Eq.(2.8a) is converting it into Eq.(4.22) of [3]. That is into

$$\varphi_x^2 + \varphi_y^2 + \varphi_z^2 = \frac{2m}{\hbar^2}(E - V)\varphi^2. \quad (2.8b)$$

Eq.s (2.8a) and (2.8b) are *equivalent classical* Hamilton Jacobi (H-J) equations *even though* Eq.(2.8b) *has* \hbar *in it!* The rationale of rewriting Eq.(2.8a) into Eq.(2.8b) lies in the fact that the stationary Schrödinger equation is obtainable from Eq.(2.8b) variationally. That is by minimization of the functional

$$J[\varphi] = \frac{1}{2} \int d^3x [(\nabla \varphi)^2 - \frac{2m}{\hbar^2}(E - V)\varphi^2] \quad (2.9)$$

under subsidiary condition

$$\int d^3x \varphi^2 = 1 \quad (2.10)$$

the Schrödinger equation

$$\left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} + \frac{\partial^2}{\partial z^2}\right)\varphi + \frac{2m}{\hbar^2}(E - V)\varphi = 0 \quad (2.11)$$

is obtained. As discussed in [3] the quantum solutions of Eq.(2.11) are also *exact solutions* of the classical Eq.s(2.8a), (2.8b). Because of this, it is sufficient to look for solutions of quantum Schrodinger's equation by using classical H-J Eq.s (2.8). This **fundamentally nontrivial** fact was exploited by Luneburg and, apparently, was overlooked by Demkov and Ostrovsky [7] who wrote that "Maxwell's fish-eye problem is *closely related to the Coulomb* (that is hydrogen atom) problem". By comparing Eq.s(2.6),(2.7) with Eq.(2.8a) we realize that Luneburg selected the system of units in which $C = E, 2m = 1$ and $e^2 = 1$. We hope that his choice should not cause any difficulty in reading. Therefore, we shall proceed along with Luneburg's choices for the time being. Furthermore, in addition he performs the following nontrivial Legendre transformation (which is canonical in nature as it will be explained below)

$$\begin{aligned} \xi &= \psi_x, \eta = \psi_y, \zeta = \psi_z, \\ x &= \omega_\xi, y = \omega_y, z = \omega_z, \\ \psi + \omega &= x\xi + y\eta + z\zeta \end{aligned} \quad (2.12)$$

resulting in

$$\xi^2 + \eta^2 + \zeta^2 = C + \frac{1}{\sqrt{\omega_\xi^2 + \omega_y^2 + \omega_z^2}} \quad (2.13a)$$

or, equivalently,

$$\omega_\xi^2 + \omega_y^2 + \omega_z^2 = \left(\frac{1}{-C + \xi^2 + \eta^2 + \zeta^2}\right)^2. \quad (2.13b)$$

By comparing Eq.s (2.5)-(2.7) and (2.13) we realize that for negative energies, $C = -|E|$. That is presence of the constant C accounts for the spectrum of bound states of the Coulomb (hydrogen atom) problem. The r.h.s. of Eq.(2.13) is describing the fish-eye potential. Luneburg's choice $C = -1$ apparently caused Demkov and Ostrovsky to conclude that the Schrödinger equation originating from the quantum version of the H-J Eq.(2.13b) employing the fish-eye potential should be written with $E = 0$ (not to be confused with the choices for C , though). This conclusion is correct and is of fundamental importance to all discussions which follow. However, it is still fundamentally misleading since the condition $E = 0$ for the Schrödinger equation with the fish-eye potential is **not** related to the energy of the Coulombic problem as we just have demonstrated! Much more on this topic is presented in section 4 and Appendix F. Alternative arguments are presented below.

Specifically, we write the Schrödinger equation with the fish-eye potential as

$$\left(-\left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} + \frac{\partial^2}{\partial z^2}\right) - \left(\frac{n_0}{|E| + x^2 + y^2 + z^2}\right)^2\right)\psi = 0 \quad (2.14a)$$

or, accounting for the discreteness of $|E| \equiv A$, as

$$\left(-\left(\frac{\partial^2}{\partial \frac{x^2}{A}} + \frac{\partial^2}{\partial \frac{y^2}{A}} + \frac{\partial^2}{\partial \frac{z^2}{A}}\right) \frac{1}{A} - \left(\frac{n_0}{A}\right)^2 \left(\frac{1}{1 + \frac{x^2}{A} + \frac{y^2}{A} + \frac{z^2}{A}}\right)\right)\psi = 0,$$

or, equivalently,

$$\left(-\left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} + \frac{\partial^2}{\partial z^2}\right) - \beta_n \left(\frac{1}{1 + x^2 + y^2 + z^2}\right)\right)\psi = 0 \quad (2.14b)$$

resulting in the manifestly discrete constant $\beta_n \simeq \frac{n_0^2}{A}$ ¹³. To make a comparison with the Schrödinger equation for hydrogen atom, in view of Eq.s (2.11),(2.13), we have: a) to choose system of units in which $\hbar = 1$, b) to replace n_0 by Ze^2 , c) to multiply the Laplacian by factor of 1/2 (kinetic energy standard form). The combination $\beta_n \simeq \frac{n_0^2}{A}$ is then being replaced by $\beta_n = \frac{(Ze^2)^2}{|E_n|}$. In the case of 2 dimensions the discrete values of β_n were calculated by Makowski and Gorska [38]. The reason why this was done in two dimensions is worthy of discussion. We shall do so in sections 3 and 4 and Appendix F.

The extension of idea of designing perfect optical instrument is based on the assumption that typically such instruments are axially symmetric. Also, the Euclidean space \mathbf{R}^3 in which the fish-eye potential lives can be foliated by the two-spheres S^2 . The stereographic projection, Eq.s(2.2),(2.3) adopted to S^2 in notations of Luneburg [21] reads

$$X = \frac{2x}{1+r^2}, Y = \frac{2y}{1+r^2}, Z = \frac{1-r^2}{1+r^2}, r^2 = x^2 + y^2, \quad (2.15)$$

so that the metric is converted into

$$ds^2 = dX^2 + dY^2 + dZ^2 = 4 \left(d\frac{x}{1+r^2}\right)^2 + 4 \left(d\frac{y}{1+r^2}\right)^2 + \left(d\frac{r^2-1}{r^2+1}\right)^2 = \frac{4}{(1+r^2)^2}(dx^2 + dy^2). \quad (2.16)$$

In view of axial symmetry it is possible to replace Eq.s(2.4a),(2.4b) by their 2-dimensional analog

$$ds^2 = n^2(x, y)(dx^2 + dy^2), n^2(x, y) = \frac{4}{(1+r^2)^2}. \quad (2.17)$$

Let now

$$x = u(\xi, \eta), y = v(\xi, \eta). \quad (2.18)$$

and introduce a complex z-plane variable via

$$z = f(\xi + i\eta) = u(\xi, \eta) + iv(\xi, \eta). \quad (2.19)$$

¹³The problem of solving the Schrödinger equation resulting in discreteness of β_n is known as Sturmian problem. It is discussed in detail in section 4 and Appendix F.

The Cauchy-Riemann equations imply

$$|dz|^2 = (dx^2 + dy^2) = |f_z(z)|^2 (d\xi^2 + d\eta^2) \text{ with } |f_z(z)|^2 = u_\xi^2 + v_\xi^2 = u_\eta^2 + v_\eta^2 \quad (2.20)$$

By comparing Eq.s(2.17),(2.20) we obtain:

$$n(\xi, \eta) = \frac{2}{1+r^2} |f_z(z)| = \frac{2}{1+|f(z)|^2} |f_z(z)|. \quad (2.21)$$

Consider now $f(z) = z^\gamma$, $\gamma \geq 1$. Using Eq.(2.21), we obtain,

$$n(r) = \frac{2\gamma r^{\gamma-1}}{1+r^{2\gamma}}. \quad (2.22)$$

For $\gamma = 1$ we recover the fish-eye potential reduced to 2-dimensional space defined in Eq.(2.17). Since $r = \sqrt{x^2 + y^2}$, it is apparently permissible¹⁴ by continuity and complementarity to extend this result to 3 dimensions as it was done by D-O and the rest of authors who followed D-O to obtain the extended fish-eye like potential (index of refraction) in 3 dimensions

$$n(r) = \frac{\beta_n (r/a)^{\gamma-1}}{1 + (r/a)^{2\gamma}} = \frac{a}{r} \frac{\beta_n}{(r/a)^{-\gamma} + (r/a)^\gamma}. \quad (2.23)$$

Here $r = \sqrt{x^2 + y^2 + z^2}$. The constant a making the ratio dimensionless is specified in section 4. The $n(r)$ defined by Eq.(2.23) should not be confused with the analogous result used in 2 dimensional calculations. For $\gamma = 1$, by design, such a potential coincides with that presented in Eq.(2.5) as required. The D-O potential, used for the whole periodic system, is obtained by selecting $\gamma = 1/2$ in Eq.(2.23). This choice is motivated by some classical and semiclassical arguments apparently allowing to by-pass restrictions of the classical Bertrand theorem. This hypothesis by D-O is incorrect in the form discussed by D-O. Instead, it is studied in detail in section 5 from the point of view of Bertrand spaces discovered by Volker Perlick. There are yet other, less sophisticated, arguments causing selection of the exponent $\gamma = 1/2$. First of them lies in the fact that *only for such exponent it is possible to repeat word-for-word scaling calculations done in* Eq.s(2.14 a) and (2.14b) and to get the quantization of the coupling constant β_n in a manner used for the fish-eye potential. Such a quantization **was postulated** by D-O. They entirely overlooked the consequences of the scaling analysis. Second of them lies in the observation that, when used in the Schrödinger Eq.(2.14), such a potential coincides exactly (up to scaling) with that proposed by Tietz [28],[29]¹⁵, as discussed in section where it was also mentioned that the Tietz potential wonderfully coincides with that obtained with help of the Hartee-Fock calculations [25].Now,

we would like to connect these results with those by Konstantin Caratheodory presented in Appendix A.

¹⁴Actually, it is not permissible!. The rest of our 2 parts work is devoted to correction of this unintentional error by D-O.

¹⁵See also a paper by Belokolos [39] who used the WKB method applied to Schrodinger's equation with Tietz potential for proving the validity of the Madelung rule.

Eq.(A.5) of this appendix A coincides exactly with Eq.(3.4) of lecture notes by Dirac [40] on quantum mechanics of the dynamical systems with constraints. It is also given in Part II, section 4, of the classical book by Arnol'd [41]. More is presented in the book by Oliver Johns[42] and in [43]. We follow, in addition, Ref.s[44] and [45] in which modern treatment of constrained systems is given in the context of gauge fields and time-dependent processes. Thus, in Eq.(A.5) we shall assume that $t = q_0(\tau)$ so that $\frac{dt}{d\tau} = \dot{q}_0$. Using these changes of notations, we obtain as well: $\dot{q}_i = \frac{dq_i}{d\tau} = \dot{q}_i \frac{dt}{d\tau}$, $i = 1 \div n$, so that $L'(t', q'_i, \frac{dq'_i}{dt}) = L^*(\tau, q_i, \dot{q}_i)$. If initially $L(t, q_i, \dot{q}_i) = \frac{1}{2}g_{ij}\dot{q}_i\dot{q}_j - V(\{q_i\})$ then, finally we obtain: $L^*(q_0, q_i, \dot{q}_0, \dot{q}_i) = (\frac{1}{2}g_{ij}\dot{q}_i\dot{q}_j / (\dot{q}_0)^2 - V(\{q_i\}))\dot{q}_0$. The canonical momenta now are given as follows:

$$p_i = \frac{\partial L^*}{\partial \dot{q}_i} = g_{ij} \frac{\dot{q}_j}{\dot{q}_0} \text{ and } p_0 = \frac{\partial L^*}{\partial \dot{q}_0} = -\frac{1}{2}g_{ij}\dot{q}_i\dot{q}_j / (\dot{q}_0)^2 - V. \quad (2.24)$$

Since the first of these equations yields

$$\frac{\dot{q}_j}{\dot{q}_0} = g^{ji} p_i, \quad (2.25)$$

its use in the second of Eq.s(2.24) produces

$$p_0 + \frac{1}{2}g^{ij}p_i p_j + V = 0 \text{ or } p_0 + H(q_i, p_i) = 0. \quad (2.26)$$

According to known rules of mechanics, the "true" Hamiltonian $\mathcal{H}(q_\alpha, p_\beta)$, $\alpha, \beta = 0 \div n$, is obtainable now via prescription

$$\mathcal{H}(q_\alpha, p_\beta) = p_i \dot{q}_i + p_0 \dot{q}_0 - L^*(q_0, q_i, \dot{q}_0, \dot{q}_i) = \dot{q}_0(p_0 + \frac{1}{2}g^{ij}p_i p_j + V) = 0. \quad (2.27)$$

Evidently, the quantized version of just obtained result can formally be written as

$$\mathcal{H}(\hat{q}_\alpha, \hat{p}_\beta)\psi(\{\hat{q}_\alpha\}) = 0, \quad (2.28)$$

where the hats denote the Hilbert space operators. Eq.(2.14a) belongs to the class of "timeless" equations of the type defined by Eq.(2.28), e.g. read [45]-[47]. This circumstance provides us with the opportunity to illustrate general results described by Eq.s (2.27),(2.28) in sufficient detail. These are to be used in the rest of this paper. Our readers should be warned

that the description of timeless equations in quantum mechanics and gravity in full generality requires many volumes. For the latest effort, e.g. read [45]. In gravitation the story also begins with Eq.(2.26), e.g. read [46], Eq. (1.11). A condensed summary is given in [47] by the same author. To keep focus on tasks to be completed in this paper, we squeeze general description of time problem to the absolute minimum. We shall study time-related issues using Eq.(2.14b) first because at least in 2 dimensions (rigorously) the fish-eye potential,

Eq.(2.5), can be replaced by the Lenz potential, Eq.(2.23), without D-O restrictions to the case of $\gamma = 1/2$. This can be achieved because the two dimensional Laplacian is conformally invariant. The conformal transformations leading to extension of the fish-eye potential discussed in Eq.s(2.15)-(2.22) are fully compatible with those for the two-dimensional Laplacian. This fact allowed both classical and quantum mechanical problems to be studied in detail in [38] and [48]. In 3 dimensions the situation seems intractable because the Laplacian is no longer conformally invariant. Therefore, **by completely ignoring the issue of conformal invariance** the fish-eye potential in Eq.(2.14a) was formally replaced by its 3-dimensional Lenz extension by Ostrovsky [17] and by Barut and Kitagawara [8,9] for the special case $\gamma = 1/2$. For arbitrary $\gamma \geq 1$ it was studied already in [7] and, later on, in [49], again, by totally neglecting the issue of conformal invariance. Thus, in this work we are confronted with the task of relating conformally the Schrödinger Eq.(2.14a), where the fish-eye potential is used legitimately, with the analogous 3-dimensional Schrodinger's equation with the Lenz fish-eye-like potential whose use in 3-dimensions is lacking justification thus far. Our study begins with rewriting Eq.s(2.13),(2.14) *classically* first. We obtain,

$$(A + |\mathbf{r}|^2)^2 |\mathbf{p}|^2 = n_0^2. \quad (2.29)$$

Here $|\mathbf{r}|^2 = x^2 + y^2 + z^2$ and $|\mathbf{p}|^2 = \omega_\xi^2 + \omega_y^2 + \omega_z^2$. Up to rescaling the factor $(A + |\mathbf{r}|^2)^2 |\mathbf{p}|^2$ can be identified with the classical Hamiltonian for the Kepler problem as discussed in [50], Eq.(7). Upon quantization Eq.(2.29) becomes a Sturmian-type problem. Our task, however, is to bring this result into the form coinciding with Eq.(2.27) and to study the consequences.

For this purpose following [50] we have to perform canonical transformations analogous to those displayed in (2.12). This time, following [50] and [51], we would like to arrive at our final result more systematically. Thus, we begin with Kepler's Hamiltonian

$$H = \frac{1}{2} |\mathbf{p}|^2 - \frac{\alpha}{|\mathbf{r}|} \quad (2.30)$$

and, using Eq.s (2.26),(2.27), we write

$$0 = (H - E) \frac{|\mathbf{r}|}{\alpha} = (|\mathbf{p}|^2 - 2E) \frac{|\mathbf{r}|}{2\alpha} - 1 \equiv H_0. \quad (2.31)$$

In the above equation, following Moser [35],[36], we used the Levi-Civita time change

$$\frac{dt}{d\tau} = \dot{q}_0 = \frac{|\mathbf{r}|}{\alpha} \quad (2.32)$$

whose physical meaning is discussed in detail in Appendix B. Again, following Moser [35], [36], and [51], we introduce the new Hamiltonian function \mathcal{F} via

$$\mathcal{F} = \frac{1}{2} (1 + H_0)^2 = \frac{1}{8\alpha^2} (|\mathbf{p}|^2 - 2E)^2 |\mathbf{r}|^2. \quad (2.33)$$

But $H_0 = 0$! Therefore, on isoenergetic surface $H = E$ or, which is the same, on $\mathcal{F} = \frac{1}{2}$, the trajectories of the Hamiltonian flow of H traversed in time t coincide with trajectories of the Hamiltonian flow on \mathcal{F} traversed in time τ . Details are provided in Appendix B.

In this subsection, however, we still need to demonstrate that the Hamiltonian $(A + |\mathbf{r}|^2)^2 |\mathbf{p}|^2$ introduced in Eq.(2.29) and the Hamiltonian $\frac{1}{8\alpha^2}(|\mathbf{p}|^2 - 2E)^2 |\mathbf{r}|^2$ introduced in Eq.(2.33) are canonically related. Already in subsection 2.1. we mentioned that the stereographic projection is a conformal mapping transformation. It can be proven that for dimensionality greater than 2 the most general conformal transformation in \mathbf{R}^n is made out of sphere inversions, euclidean motions and scale transformations [52]. Following [50] we perform the inversion of coordinates

$$(x')^i = \frac{x^i}{|\mathbf{x}|^2}, \quad (2.34)$$

so that the corresponding canonical momenta $p'_i = \left(\partial x^j / \partial (x')^i \right) p_j$ become

$$p'_i = |\mathbf{x}|^2 p_i - 2x^i (\mathbf{x} \cdot \mathbf{p}). \quad (2.35)$$

It can be checked explicitly that just described transformations are canonical. Next, one should perform the additional (scale) canonical transformation

$$\bar{x}^i = \frac{p'_i}{b}, \quad \bar{p}_i = -b (x')^i \quad (2.36)$$

where $b > 0$ is an adjustable parameter. With these conformal transformations and by adjusting the value of the parameter b , the Hamiltonian $\frac{1}{8\alpha^2}(|\mathbf{p}|^2 - 2E)^2 |\mathbf{r}|^2$, $E < 0$, $|E| = A$, is converted into Hamiltonian, Eq.(2.29), for the fish-eye potential. Quantization of the Schrödinger equation with such potential is discussed in detail in section 4. Additional independent information is provided in Appendix F.

3. Known and unknown roles of conformal invariance in quantum mechanical problems involving fish-eye potentials

In two dimensions by using conformal transformations we obtained Eq.(2.21) for the refractive index. Adapted to two dimensions, Schrödinger's Eq.(2.14a) can be rewritten as

$$-\left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2}\right)\psi - V(x, y)\psi = 0. \quad (3.1a)$$

In such a form this equation was studied by Makowski [53] for the fish-eye and related potentials. By performing conformal transformation on Eq.(3.1a) and using Eq.(2.21), Eq.(3.1a) can be rewritten in equivalent form

$$-|f_z(z)|^2 \left(\frac{\partial^2}{\partial \xi^2} + \frac{\partial^2}{\partial \eta^2}\right)\psi - |f_z(z)|^2 U(\xi, \eta)\psi = 0 \quad (3.1b)$$

allowing the conformal factor $|f_z(z)|^2$ to drop out. Such an omission of the conformal factor is possible because the 2 dimensional Laplacian is conformally invariant. In dimensions higher than two this is no longer true [54]. Thus, beginning with the two-dimensional version of the fish-eye potential and by applying the conformal transformations to this potential, e.g. to Eq.(2.22), and to the Laplacian, Makowski [53] studied the Schrödinger- type¹⁶ equations for a large class of fish-eye-type potentials. Other authors, e.g. see [7-9],[17],[49] used the 3-dimensional version of the generalized fish-eye potential, Eq.(2.23),¹⁷ in the corresponding 3-dimensional Schrödinger-type equations without any concern about the mathematical correctness of these equations. By ignoring the issue of conformal invariance, these authors had lost some fundamentally important physics essential not only for solution of the Löwdin problem in atomic physics but, more broadly, for solution of dynamical problems in non flat Bertrand spaces. The role of Lie sphere and conformal geometry in finding the solution of the Schrödinger, the Klein-Gordon and other wave equations was discussed in detail in our previous work [3]. Here we are going to add some new details to the topic of conformal invariance by making the unexpected connections between results of section 2 and our previous work [3].

We begin this section by connecting the results of Ref.[55]¹⁸ with those discussed in section 2 and Appendices A and B. Starting with the Lagrangian

$$L(t, q_i, \dot{q}_i) = \frac{1}{2} g_{ij} \dot{q}^i \dot{q}^j - V(\{q_i\}) \quad (3.2)$$

we are defining the conformal transformation

$$\bar{g}_{ij}(q_i) = \omega^2(q_i) g_{ij}(q_i), \quad (3.3)$$

where $\omega^2(q_i)$ is some positive function of coordinates $\{q_i\}$. Instead of previously used $\frac{dt}{d\tau} = \dot{q}_0$ we write now

$$d\tau = \omega^2(q_i) dt. \quad (3.4)$$

In view of this result, the action

$$S = \int dt L(t, q_i, \dot{q}_i) \quad (3.5a)$$

should be rewritten now as

$$S = \int \frac{d\tau}{\omega^2(x)} \left[\frac{1}{2} \omega^4(q_i) g_{ij} q'^i q'^j - V(\{q_i\}) \right], \quad q'_i = \frac{dq_i}{d\tau}, \quad q'^i = g^{ij} q'_j, \quad (3.5b)$$

or, equivalently,

$$S = \int d\tau \left[\frac{1}{2} \bar{g}_{ij} q'^i q'^j - \bar{V}(\{q_i\}) \right] \equiv \int d\tau \bar{L}(t, q_i, q'_i), \quad (3.5c)$$

¹⁶Actually, the Sturmian-type.

¹⁷For $\gamma = 1/2$ in [7-9],[17] and for arbitrary $\gamma > 1$ in [49]. In view of scaling analysis of Eq.(2.14), arbitrary γ 's are not physically realizable though.

¹⁸This paper is containing many typos causing us to rewrite needed equations

where $\bar{g}_{ij} = \bar{g}_{ij}(q)$ is defined by Eq.(3.3) and $\bar{V}(\{q_i\}) = \frac{V(\{q_i\})}{\omega^2(q_i)}$. In Appendix B we demonstrate that the Euler-Lagrange equation of motion for $L(t, q_i, \dot{q}_i)$ and $\bar{L}(t, q_i, q'_i)$ produce the same trajectories. It is of interest to reobtain this result now using different method for a reason which will become clear upon reading.

For $L(t, q_i, \dot{q}_i)$ the Euler-Lagrange equations are

$$\ddot{q}^i + \Gamma_{jk}^i \dot{q}^j \dot{q}^k + V^{,i} = 0 \quad (3.6)$$

while for $\bar{L}(t, q_i, q'_i)$ these equation acquire the following form

$$q''^i + \hat{\Gamma}_{jk}^i q'^j q'^k + \frac{V^{,i}}{\omega^4(q_i)} - \frac{2V}{\omega^5(q_i)} \omega(q_i)^{,i} = 0. \quad (3.7)$$

Here, according to [52], we have for the Christoffel symbols Γ_{jk}^i the following transformation law:

$$\hat{\Gamma}_{jk}^i = \Gamma_{jk}^i + \delta_j^i (\ln \omega)_{,k} + \delta_k^i (\ln \omega)_{,j} - g_{jk} (\ln \omega)^{,i}. \quad (3.8)$$

Since

$$q'^i = \frac{dq^i}{dt} \frac{dt}{d\tau} \equiv \dot{q}^i \frac{1}{\omega^2(q_i)} \quad (3.9a)$$

and

$$q''^i = \frac{1}{\omega^4(q_i)} [\ddot{q}^i - 2\dot{q}^i \dot{q}^j (\ln \omega)_{,j}], \quad (3.9b)$$

by replacing in Eq.(3.7) q''^i and q'^i by Eq.s(3.9a) and (3.9b) and using Eq.(3.8) we finally obtain after some algebra

$$\ddot{q}^i + \Gamma_{jk}^i \dot{q}^j \dot{q}^k + V^{,i} - 2(\ln \omega)^{,i} \left(\frac{1}{2} g_{ij} \dot{q}^i \dot{q}^j + V(\{q_i\}) \right) = 0. \quad (3.10)$$

Here

$$\frac{1}{2} g_{ij} \dot{q}^i \dot{q}^j + V(\{q_i\}) = E \quad (3.11)$$

is an energy. By reversing arguments we also obtain the equations of motion for primed variables as well as the analog of E rewritten in primed variables. Thus, we just obtained the

Theorem 3.1. *Two dynamical systems are conformally invariant at the level of their respective equations of motion if and only if they both have vanishing total energies.*

Corollary 3.2. From results of Appendix B it follows that the dynamics of Reeb vector fields is conformally invariant. It is describing the motion on geodesics.

Corollary 3.3. Conformal invariance is result of time changing transformations. They make time a canonical variable with energy/Hamiltonian being its conjugate. Use of such transformations brings the Newtonian mechanics into covariant form used for description of

dynamics of general relativity. More on this is presented in section 5 in connection with description of the Eisenhart lift and Bertrand spaces.

Extension of these results to nonquantized field theories proceeds analogously. Some representative examples are discussed in [55]. The question arises: How just obtained results can be transferred to quantum mechanics/quantum field theories? In 2 dimensions Eq.(3.1b) demonstrates that this is indeed possible in the case of nonrelativistic quantum mechanics. Therefore, we would like now to demonstrate that since the Theorem 3.1. holds in higher dimensional spaces as well, there should be an analog of this theorem in quantum mechanics/quantum field theories in higher dimensions. If we would follow standard textbooks on

quantum mechanics and replace the classical kinetic energy term by the term involving the Laplacian (perhaps defined on the curved space) we would not get the correct answer for conformally related quantum analogs of conformally related classical systems. This is so because above two dimensions the Laplacian is not going to remain form-invariant under the conformal transformations. Instead, the *conformal* (Yamabe) Laplacian \square_g must be used¹⁹. It is defined as follows

$$\square_g = -\Delta_g + \alpha(d)R(g), \quad (3.12a)$$

where $\alpha(d) = \frac{d-2}{4(d-1)}$ and $R(g)$ is the scalar curvature of d -dimensional Riemannian manifold \mathcal{M}_d whose metric tensor is g_{ij} . In the case when the Laplacian Δ_g is acting on scalar function ψ , its action is defined as [60]:

$$\Delta_g \psi = g^{ij} \left[\frac{\partial^2 \psi}{\partial x^i \partial x^j} - \Gamma_{ij}^k \frac{\partial \psi}{\partial x^k} \right]. \quad (3.12b)$$

Under the conformal change of metric $\tilde{g} = g \exp(2f)$ the conformal Laplacian is transforming as follows

$$\square_{\tilde{g}} = e^{-(\frac{d}{2}+1)f} \square_g e^{(\frac{d}{2}-1)f}. \quad (3.13)$$

Ovsyannikov [59] and Ibragimov [60] considered in detail the solutions of partial differential equations of the following type

$$F[u] = g^{ij} u_{ij} + b^i u_i + cu = 0. \quad (3.14)$$

Here $u_i = \frac{\partial u}{\partial x_i}$, $u_{ij} = \frac{\partial^2 u}{\partial x_i \partial x_{ij}}$, g^{ij} , b^i and c are the known functions of x_i , and $g^{ij} = g^{ji}$.

Following ideas of Hadamard discussed in our previous work [3], Ibragimov [60], using work by Ovsyannikov [59], demonstrated that for spaces of Riemannian signature use of Hadamard-type transformations of the type discussed in [3] converts Eq.(3.14) into conformally invariant equation

$$\square_g \Psi = 0. \quad (3.15)$$

¹⁹For a quick introduction to this topic we refer our readers to our works [58,59].

In spaces of Lorentzian signature it is known as well, e.g. read Appendix B of [3] and [60], section 12.2, Lemma 1., that the wave equation²⁰

$$\square u \equiv \left(\frac{\partial^2}{\partial t^2} - \Delta \right) u = 0 \quad (3.16)$$

adapted for spaces of Lorentzian signature having the pp-type metric (the Penrose limit metric) is conformally invariant and obeys the Huygens principle. Thus, in spaces with pp metrics the conformal invariance is inseparably linked with the Huygens principle [60] for such an equation. In [60] it is demonstrated that equations of the type

$$\square u + b^i u_i + cu = 0 \quad (3.17)$$

with help of the Hadamard-like transformations are reducible to the standard wave equation

$$\square \Phi = 0. \quad (3.18)$$

This property causes Eq.(3.17) to belong to the class of equations obeying Huygens' principle. Summarizing, we obtain the following

Corollary 3.4. *In spaces of Riemannian and pseudo Riemannian signature Eq.s(3.14) and, respectively, (3.17) are conformally invariant. For spaces of Lorentzian signature the conformal invariance is inseparably linked with the validity of Huygens' principle. These equations are quantum analogs of the conformally invariant dynamical equations of classical mechanics.*

The 3 dimensional analog of the 2-dimensional Schrödinger Eq.(3.1a) reads as follows

$$-\left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} + \frac{\partial^2}{\partial z^2} \right) \psi + V(x, y, z) \psi = 0. \quad (3.19a)$$

Here, in view of Eq.(2.23), the potential $V(x, y, z)$ is given by

$$V(x, y, z) = - \left(\frac{a}{r} \right)^2 \left[\frac{n_0}{(r/a)^{-\gamma} + (r/a)^{\gamma}} \right]^2, r^2 = x^2 + y^2 + z^2, a = \text{const}. \quad (3.19b)$$

The quantum mechanical eigenvalue (Sturmian) problem for Eq.(3.19a) with such a potential was formally solved in the foundational work Ref.[7]. Details were further elaborated in [49] for any γ . The case $\gamma = 1$ corresponds to the Maxwell's fish-eye potential (and, hence, to the hydrogen atom) while the $\gamma = 1/2$ case corresponds to the D-O potential used by these authors for description of electronic structure of the whole periodic system of elements. Corollary 3.4. guarantees that for *any* γ these equations are conformally equivalent to each other. But in section 2 we found that Eq.s(2.14a) and (2.14b) make sense physically only for $\gamma = 1$ or $1/2$.

²⁰Here Δ is the flat Laplacian

Given these facts, two questions arise: a) If these Eq.s(3.19) are conformally equivalent, which one of these should be considered as the seed from which others are being obtained? b) What makes Maxwell's fish-eye ($\gamma = 1$) and D-O ($\gamma = 1/2$) potentials so special if, in fact, they are conformally equivalent?

To answer a) we have to remind to our readers some general facts about the conformally related spaces. We begin with

Definition 3.5. The (pseudo) Riemannian space \mathcal{M}_d of dimension d is *flat* if in some region around any point $x \in \mathcal{M}_d$ there is a metric-preferred coordinate system such that

$$ds^2 = g_{ij}(x)dx^i dx^j \quad (3.20)$$

in which the metric tensor $g_{ij}(x) = e_i \delta_{ij}$, $e_i = \pm 1$.

Definition 3.6. A (pseudo) Riemannian space \mathcal{M}_d is called *conformally flat* if it can be obtained from the flat space via conformal change of metric tensor according to Eq.(3.3), that is with help of transformation $\bar{g}_{ij}(x) = \omega^2(x)g_{ij}(x)$ with $g_{ij}(x)$ defined in Eq.(3.20).

Definition 3.7. When $\omega^2(x) = \text{const}$, the above conformal transformation is called *homothety* (or *similarity*), when $\omega^2(x) = 1$ the homothety is *isometry*. The isometric-type of motion on \mathcal{M}_d is described in terms of Killing's equations. The Killing vectors \mathbf{X} are solutions of Killing's equations

$$\mathcal{L}_{\mathbf{X}}g_{ij} = 0, \quad (3.21)$$

where the Lie derivative $\mathcal{L}_{\mathbf{X}}$ is defined in Eq.(B.5).

Definition 3.8. The conformal type of motion on \mathcal{M}_d is described in terms of the *conformal Killing equations*

$$\mathcal{L}_{\mathbf{X}}g_{ij} = 2\Phi(x)g_{ij} \quad (3.22a)$$

in which \mathbf{X} is the *conformal* Killing vector [61]²¹.

Consider now two conformally related spaces, e.g. let $\bar{g}_{ij}(x) = e^{2\sigma(x)}g_{ij}(x)$. It is of interest to study the relationship between Eq.(3.22a) and

$$\mathcal{L}_{\mathbf{X}}\bar{g}_{ij} = 2\bar{\Phi}(x)\bar{g}_{ij}. \quad (3.22b)$$

It can be demonstrated [60], [62] that

$$\bar{\Phi}(x) = \Phi(x) + 2\xi^i \frac{\partial \sigma}{\partial x^i}. \quad (3.23)$$

Here the (Killing) vector ξ^i is defined by Eq.s (C.2),(C.3) in Appendix C. Eq.(3.23) reminds us about the analogous cases in gauge theories associated with the choices of gauges. In

²¹For reader's convenience some basic information on Killing's equations is presented in the Appendix C

particular, the choice $\bar{\Phi}(x) = 0$ implies that the underlying space with the metric \bar{g}_{ij} is completely rigid. It admits only isometric (non bendable/ non stretchable) motions. Although physically non realistic, such spaces do exist mathematically [60] and can be constructed based on solutions to the the equation

$$\Phi(x) + 2\xi^i \frac{\partial \sigma}{\partial x^i} = 0. \quad (3.24)$$

These spaces are serving as boundaries of conformally flexibe/bendable spaces or can occur as subspaces of these spaces. In Appendix C we established that for the Riemannian spaces \mathcal{M}_d of dimensionality d the isometric motion is described in terms of the (maximum) $\frac{d(d+1)}{2}$ parameters. The maximum is achieved for spaces of constant curvature.

The question arises: What is the maximal number of parameters required for description of conformally related spaces?

To answer this question, we have to extend the analysis presented in Appendix C. In particular, we have to count d Killing vectors ξ^i as well as the factor Φ . Altogether there are $d+1$ such terms. In addition, however, we should account for d^2 terms like $\nabla_k \xi^i$ and d terms like $\partial_k \Phi$. Their role is nicely explained in [62], section 1. Thus, altogether we obtain: $d^2 + d + d + 1 = (d+1)^2$ parameters. The truly nontrivial conformal motion is described in terms of $(d+1)^2 - \frac{d(d+1)}{2} = \frac{1}{2}(d+1)(d+2)$ parameters. In [60] there is a proof of the following remarkable

Theorem 3.9. *(Pseudo) Riemannian space \mathcal{M}_d has nontrivial conformal group of motions only if \mathcal{M}_d is conformally equivalent to the conformally flat space²²*

Definition 3.10. The conformal group is *trivial* if use of the relation $\bar{g}_{ij}(x) = \omega^2(x)g_{ij}(x)$ leads to Eq.(3.24) possessing a nonzero solution.

Corollary 3.11. According to Theorem C.1. The maximum number of parameters $\frac{d(d+1)}{2}$ is possible only for spaces \mathcal{M}_d of full isometry group which are of constant scalar curvature. But, according to Corollary C.3., all spaces of constant scalar curvature are conformally flat. This means that such manifolds have exactly $\frac{1}{2}(d+1)(d+2)$ conformal parameters involved in nontrivial conformal group according to Theorem 3.9.

This fact can be illustrated with help of Eq.s(3.12a) and (3.13). Suppose that in Eq.(3.12a) the metric \tilde{g} is describing the space of constant scalar curvature. Because such a space is conformally flat, we can write

$$e^{-(\frac{d}{2}+1)f} (-\Delta_{g=1}) e^{(\frac{d}{2}-1)f} = \square_{\tilde{g}}, \quad (3.25)$$

²²This theorem is compatible with the existence of conformally flat Bertrand spaces [18] described in section 5 and Appendix E.

where $g = 1$ is actually $g = \delta_{ij}$ is the Euclideanized version of the metric defined in Eq.(3.20). Thus, we are having two reference manifolds : a) The manifolds of constant scalar (nonzero) curvature and b) The manifolds whose scalar curvature is exactly zero. All other conformally flat manifolds whose scalar curvatures are nonconstant can be related (deformed) either to a) or to b)-type of manifolds (these are conformally equivalent, of course). This was the point of departure of Yamabe's work [63]. An excellent modern exposition of Yamabe ideas is given in [64]. For a rapid introduction to this topic, please, consult our previous works [56] and [57].

Based on just described results, we argue below that the Schrödinger equation for the Maxwell fish-eye potential ($\gamma = 1$) should be taken as a seed. This will provide us an answer to the question a). Foregoing development makes perfect sense physically since such an equation corresponds to the stationary Schrödinger equation for the hydrogen atom. This is demonstrated in detail in section 4 and also below. It means as well that **the equation proposed by D-O ($\gamma = 1/2$) for the description of spectra of other multielectron atoms is obtainable as conformal deformation of the Schrödinger equation for the hydrogen atom.**

Such a deformation is to be described below. It is also discussed in the Appendix F from the alternative standpoint. The results are fully compatible with the modified Betrand theorem discussed in section 5. Following Ovsyannikov [59] we apply the Hadamard-type

transformations described in some detail in our previous work [3] culminating there in Theorem 4.1. of that work. Below these transformations are described in much greater detail with the purpose of transforming Eq.(3.14) into the D-O form, Eq.(3.19) (firstly with $\gamma = 1$).

In compliance with Hadamard method [3], we consider now the 3 types of Hadamard transformations in full generality.

(α) The coordinate transformation

$$x'^i = p^i(x), i = 1, \dots, d. \quad (3.26a)$$

It leads to $u = u' \circ p$ implying

$$F'[u'] = g'^{ij} u'_{ij} + b'^i u'_i + c' u' = 0. \quad (3.26b)$$

In this equation $u'_i = \frac{\partial}{\partial x'_i} u'$, etc., $g'^{kl} \circ p = g^{ij} p_i^k p_j^l$, $b'^k \circ p = g^{ij} p_{ij}^k + b^i p_i^k$, $c' \circ p = c$, $p_i^k = \frac{\partial}{\partial x_i} p^k$, etc.

(β) The scale transformation

$$F'[u] = e^{-\theta} F[e^\theta u]. \quad (3.27a)$$

It is converting Eq.(3.14) into

$$F'[u] = g^{ij} u_{ij} + b^i u_i + c' u = 0. \quad (3.27b)$$

Here $b'^i = b^i + 2g^{ij}\theta_j$ and $c' = c + b^i\theta_i + g^{ij}\theta_i\theta_j + g^{ij}\theta_i\theta_j$.

(γ) The conformal transformation

$$F'[u] = e^{-2\theta} F[u]. \quad (3.28a)$$

It results in

$$F'[u] = \tilde{g}^{ij}u_{ij} + \tilde{b}^i u_i + \tilde{c}u = 0. \quad (3.28b)$$

Here $g^{ij} = e^{2\theta}\tilde{g}^{ij}$, $b^i = e^{2\theta}\tilde{b}^i$ and $c = e^{2\theta}\tilde{c}$.

Transformations (α) and (β) leave Eq.(3.14) in the same (pseudo) Riemannian space \mathcal{M}_d . The transformation (γ) conformally relates \mathcal{M}_d to $\tilde{\mathcal{M}}_d$.

These statements require proofs. We shall not reproduce these proofs referring our readers to Ovsyannikov's book [59]. Nevertheless, results of these proofs cause us to introduce the scalar H and the tensor K_{ij} to be defined momentarily. For this purpose, following Ovsyannikov [59], we have to define the contravariant vector a^i via

$$a^i = b^i \pm g^{kl}\Gamma_{kl}^i, i = 1, \dots, d. \quad (3.29a)$$

Ovsyannikov uses only sign "+" in his calculations but below we shall argue that, in fact, both signs should be used. Since under (α)-type of transformations Christoffel's symbols transform as

$$\Gamma_{ij}^l p_l^k = p_i^k + \Gamma_{ml}^k p_i^m p_j^l, \quad (3.30)$$

it can be demonstrated that the contravariant vector a^i obeys the standard rule of transformation for vectors:

$$a'^k = b'^k \pm g'^{ml}\Gamma_{ml}^k = (b^i \pm g^{kl}\Gamma_{kl}^i)p_i^k = a^i p_i^k. \quad (3.29b)$$

With help of such defined vector a^i it becomes possible to introduce a skew-symmetric tensor K_{ij} via

$$K_{ij} = a_{i,j} - a_{j,i}, \quad (3.31)$$

where $a_{i,j} = \frac{\partial a_i}{\partial x^j} - a_l \Gamma_{ij}^l$ and the raising and lowering of indices is made with help of the metric tensor g_{ij} , as usual. Accordingly, the scalar H can now be defined as

$$H = -2c + a^i_{,i} + \frac{1}{2}a^i a_i + 2\alpha(d)R(g). \quad (3.32)$$

Thus defined H and K_{ij} allow us to prove the following

Theorem 3.12. *H and K_{ij} remain unchanged under transformations of (β)-type converting Eq.(3.14) into Eq.(3.27b). They are the invariants of (β)-type transformations.*

It can be also proven that with respect to transformations (γ) the tensor K_{ij} will remain unchanged. So it is also an invariant of (γ)-type transformations. At the same time, the

scalar H is transformed into $\tilde{H} = e^{-2\theta}H$. Therefore, it loses its invariance property. The conformal Killing Eq.s(3.22a) can be rewritten in a more familiar form now as

$$\nabla_i \xi_j + \nabla_j \xi_i = \mu g_{ij}, \quad \mu = 2\Phi. \quad (3.33)$$

With help of these equations, it also can be demonstrated that

$$\xi^k H_{,k} + \mu H = 0. \quad (3.34a)$$

If $H \neq 0$, it also can be demonstrated that it is always possible to select θ in Eq.(3.28a) so that $H = e^{2\theta}$. Such a selection leads to $\tilde{H} = 1$. The transformed equation

$$\xi^k \tilde{H}_{,k} + \tilde{\mu} \tilde{H} = 0 \quad (3.34b)$$

is implying that $\tilde{\mu} = 0$. Therefore, whenever $H = e^{2\theta}$, we are back to the Killing Eq.s (C.4) implying that we are dealing with the isometry subgroup containing $\frac{(d+1)d}{2}$ generators *inside* larger *nontrivial* conformal group. This fact is in accord with Yamabe's ideas [63] that "more flexible" spaces \mathcal{M}_d are conformally deformable into spaces of constant scalar curvature.

To make further progress, from now on we shall deal only with the conformally flat metric, that is with the metric of the type $g^{ij}(x) = \rho(x)\delta^{ij}$ (or with $g_{ij}(x) = \frac{1}{\rho(x)}\delta_{ij}$). To illustrate the value of just described results, we begin with the simplest but instructive example.

Instead of considering Eq.(3.14), we shall focus now on Schrödinger's Eq.(3.19a) written in the form

$$\rho \Delta u + cu = 0 \quad (3.35)$$

in which we initially do not require $c(x)$ to be a constant and use the notation $\Delta = \Delta_{g=1}$ as before. This equation plays a significant role also in section 5. By looking at Eq.(3.27b) and comparing it with Eq.(3.14), we realize that in Eq.(3.35) $b'^i = 0$ and c in this equation should be actually relabelled as c' . By looking at Eq.(3.29b) and by noticing that for the (β) -type transformations $g^{ij} = g'^{ij}$ causing $\Gamma_{ml}^k = \Gamma_{kl}^i$, the combination $g^{kl}\Gamma_{kl}^i = \frac{d-2}{2}\frac{\partial}{\partial x^i} \ln \rho$. This result is obtained in the Appendix D. From Eq.(3.29b) we obtain,

$$a'_i = \pm \frac{d-2}{2} \frac{\rho_i}{\rho} \text{ or } a^i = \pm \frac{d-2}{2} \delta^{ij} \rho_j. \quad (3.36)$$

In view of Eq.(3.32) we have to calculate

$$a^i_{,i} = \frac{\partial a^i}{\partial x^i} + a^k \Gamma_{ki}^i = \pm \left(\frac{d-2}{2} \delta^{ij} \rho_{ij} - \frac{d(d-2)}{4} \delta^{ij} \frac{\rho_i \rho_j}{\rho} \right), \quad (3.37)$$

where we used Eq.(D.8) of the Appendix D for calculation of Γ_{ki}^i . Besides,

$$\frac{1}{2} a^i a'_i = \frac{(d-2)^2}{8} \delta^{ij} \frac{\rho_i \rho_j}{\rho}. \quad (3.38)$$

In view of Eq.(3.32), by combining Eq.s(3.37) and (3.38) we obtain:

$$a'_{,i} + \frac{1}{2}a'^i a'_i = \frac{(d-2)^2}{8}\delta^{ij}\frac{\rho_i\rho_j}{\rho} - \frac{d^2-4}{4}\delta^{ij}\frac{\rho_i\rho_j}{\rho}. \quad (3.39)$$

In calculating this result we selected the sign "+" in Eq.(3.36). This is caused by the following considerations. For spaces of constant curvature the conformal factor ρ was obtained in Appendix C. By adopting the Euclidean version of the metric given in Eq.(C.7), small calculation using Eq.s(3.36),(3.37) results in

$$a'_{,i} + \frac{1}{2}a'^i a'_i = \frac{d(d-2)}{2}K. \quad (3.40)$$

Going back to Eq.(3.32) and taking into account that $\frac{d-2}{2(d-1)} = 2\alpha(d)$ and that, according to [65], for spaces of constant scalar curvature $R(g) = -d(d-1)K$, we obtain: $2\alpha(d)R(g) = -\frac{d(d-2)}{2}K$. By combining this result with Eq.(3.40) and using it in Eq.(3.32) we obtain:

$$H' = -2c' + a'_{,i} + \frac{1}{2}a'^i a'_i + 2\alpha(d)R(g) = -2c'. \quad (3.41)$$

But since H is invariant of (β) -transformations, by construction, we must have $H' = H$. In view of Eq.(3.41), this is only possible when c' is a constant. In such a case Eq.(3.35) is the stationary Schrödinger's equation for the hydrogen atom according to Eq.s(3.19a) and (3.19b) where we have to select $\gamma = 1$.

Using Eq.s(3.12a) and (3.12b) we consider now the following equation

$$F[u] = \square_g u = -\rho u'_{,i} + \frac{d-2}{2}\rho^i u_i + \alpha(d)R(\rho)u = 0. \quad (3.42)$$

Clearly, it is of exactly the same type as Eq.(3.14). Therefore, in Eq.(3.42) we must have $b^i = -\frac{d-2}{2}\rho^i$, $c = -\alpha(d)R(\rho)$. At first, we consider this equation for the space of constant scalar curvature. We already know that in such a case $2\alpha(d)R(g) = -\frac{d(d-2)}{2}K$. Using this result we can replace the term $c = -\alpha(d)R(\rho)$ by $c = \frac{d(d-2)}{4}K$. Now we want to compare Eq.(3.42) with Eq.(3.35) since they both have the same ρ factor. Recall that the (β) -type transformations do not change the metric of the underlying space. Since this is so, the equality of invariants $H = H'$ leads to the result:

$$a'_{,i} + \frac{1}{2}a'^i a'_i = -2c' + d(d-2)K = \text{const}. \quad (3.43)$$

To check this result we notice that the contravariant vector a^i is defined by Eq.(3.29a). The term $g^{kl}\Gamma_{kl}^i$ in this equation is calculated in the Appendix D, Eq.(D.2). Therefore, by selecting the "+" sign we obtain,

$$a^i = b^i + g^{kl}\Gamma_{kl}^i = -\frac{d-2}{2}\rho^i + \frac{d-2}{2}\rho^i = 0, \quad (3.44)$$

implying $c' = \frac{d(d-2)}{2}K$. Thus, $c' = 2c$ which makes physical sense. Notice, in arriving at this result Eq.(3.44) was used in which we have not made any specification of the conformal factor. The result $a^i = 0$ was obtained by Ibragimov [60], page 120, by different set of arguments. Just obtained result gives us an opportunity to take also a look at the Yamabe Laplacian for the metric *other* than that of constant scalar curvature. To do so we put the factor ψ in Eq.(D.7) temporarily to zero in view of Eq.(3.44). Then, in view of Eq.s (3.13),(3.27a), we write $e^\theta = (e^{2f})^{\frac{d-2}{4}}$. To utilize fully Eq.(3.13) we have to use Eq.(3.28a), that is we have to use the (γ) transformation which is connecting equations with conformally related metrics. To facilitate matters, we notice that the combination $e^{2f}e^{-\theta}F[e^\theta u]$ in which $e^\theta = (e^{2f})^{\frac{d-2}{4}}$ produces exactly the same result $e^{-(\frac{d}{2}+1)f}F[e^{(\frac{d}{2}-1)f}u]$ as was recorded already in Eq.(3.13) for the Yamabe Laplacian. This provides us with an independent check of correctness of our computations since the conformal factor e^{2f} is present in the equation $\bar{g}_{ij}(x) = e^{2f(x)}g_{ij}(x)$ defining conformally related spaces.

We still have to comment on several topics. First, we need to comment on the role of

(α) -transformations. Such type of transformations are described in detail in the monumental work, [66], Chapter 3. These are needed to bring the PDE's, Eq.(3.14), into the canonical/standard form. Also these transformations had been used for defining the contravariant vector a^i . Since we are using only the metric of the type $g^{ij}(x) = \rho(x)\delta^{ij}$, this spares us from the necessity to consider (α) transformations further. Second, in using the (β) -type transformations we are replacing the coefficients b^i and c by b'^i and c' using the same factor θ . This means that by properly selecting θ we can always replace the factor c by the pre assigned factor c' . However, in such a case the factor b'^i cannot be forced to become zero. For this to happen, in addition to parameter θ , the parameter ψ defined in Eq.(D.5) should be used. If we would select the sign "-" initially in Eq.(3.29a), the result $a^i = 0$ in Eq.(3.44) will change into

$$a^i = -(d-2)\rho^i. \quad (3.45)$$

This is helpful if we want to use Eq.s(D.3)-(D.7) from Appendix D.

These equations should be used as follows.

- a) Using the equation $c' = c + b^i\theta_i + g^{ij}\theta_{ij} + g^{ij}\theta_i\theta_j$ with pre assigned c' factor we can determine the factor θ . For this purpose we need to solve the PDE equation $g^{ij}\theta_{ij} + g^{ij}\theta_i\theta_j + b^i\theta_i = c' - c$. Its solution is greatly facilitated by our use of spherical symmetry.
- b) Use of this result in Eq.(D.7) determines the factor ψ .
- c) After these two factors are determined, equations $b'^i = b^i + 2g^{ij}\theta_j$ and $c' = c + b^i\theta_i + g^{ij}\theta_{ij} + g^{ij}\theta_i\theta_j$ yield the one-to-one relations between c' and c and b'^i and b^i in such a way that we can put $b'^i = 0$.

In treating the hydrogen atom case to keep $b'^i = 0$ is essential. The hydrogen atom case was treated based on Eq.(3.19) where we had selected $\gamma = 1$. But for $\gamma \neq 1$ the same equation

can also be brought into canonical form

$$\rho(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} + \frac{\partial^2}{\partial z^2})u + \tilde{c}'u = 0 \quad (3.46a)$$

in which

$$\rho = [\tilde{V}(x, y, z)]^{-1}, \quad \tilde{c}' = \text{const.} \quad (3.46b)$$

In such a form it is used in section 5. Eq.(3.35) is our physical analog of Eq.(3.27b). In Eq.(3.27b), we have: $\rho = [V(x, y, z)]^{-1}$, $b^i = 0$ and $\tilde{c}' = \text{const.}$ Next, we use Eq.(3.42) which is an analog of Eq.(3.14). In it, we use the ρ - factor just defined. Further, in this equation we have $b^i = -\frac{d-2}{2}\rho^i$ and $c = -\alpha(d)R(\rho)$, as required. Now, we need to relate Eq.(3.42) with Eq.(3.46a) using (β) -type transformations. This is achievable as we just described above. Next, by inverting these operations and using transformation described in Eq.(3.13) **we conformally relate the D-O Eq.(3.46a) describing the multielectron atoms with Eq.(3.35) describing the hydrogen atom.** To answer the earlier posed question b), we follow analysis made by Ovsyannikov [59] in Chr.8, section 10. In it he considers the Lie algebras of two versions of Eq.(3.46a). One- with $\rho \neq 1$ and, another-with $\rho \equiv 1$. In the last case the Lie algebra is made of two types of generators: a) translations: $\frac{\partial}{\partial x_i}$ and, b) rotations $x^i \partial_j - x^j \partial_i$. While in the first case it is **made of the same translations** as in the second, but the rotation generators change now to $(2x_i x^k + (1 - \rho^2)\delta_i^k)\partial_k - (d - 2)x_i u \partial_u$, $i = 1, \dots, d$. These two sets of Lie algebras cannot be smoothly converted into each other. Only the Lie algebra for Eq.(3.46a) with $\rho \neq 1$ makes physical sense. It conformally relates the metric of constant scalar curvature (the hydrogen atom case) with the modified metric induced by the presence of the rest of electrons.

4. From analysis to synthesis

Already in their first seminal work [7], Demkov and Ostrovsky discussed method of solving Eq.s(3.19a),(3.19b) in full generality. Subsequently, their results were further developed in [8, 17, 49, 67], *etc.* These papers differ from each other by the extent of generality of employing the potential, Eq.(3.19b), in Eq.(3.19a)²³. In this section we shall entirely reconsider all these results since none of them are without mistakes. One of the purposes of this section is to correct these mistakes. As a by product new quantum mechanical formalism of dealing with discrete spectrum of hydrogen and many-electron atoms is developed. It is tested on known examples and, subsequently, it is extended along the directions outlined in [7]. While doing so, we shall connect results obtained in section 3 with those to be presented in this section. We begin with the following observation.

The 3-dimensional flat space Laplace equation

$$\Delta \Psi(\mathbf{x}) = 0 \quad (4.1)$$

²³E.g. some of the papers discussed only the physically relevant case: $\gamma = 1/2$, *etc.*

according to [68],[69] admits 11 separable coordinate systems. At the same time, there are 17 coordinate systems for the same Laplace equation permitting R-separation of variables.

Definition 4.1. An R-separable coordinate system $\{u, v, w\}$ for Eq.(4.1) is a coordinate system which permits a family of solutions

$$\Psi_{\lambda, \mu}(\mathbf{x}) = R(u, v, w) U_{\lambda, \mu}(u) V_{\lambda, \mu}(v) W_{\lambda, \mu}(w), \quad (4.2)$$

where λ and μ are separation constants and R is a fixed factor such that it is either $R \equiv 1$ (*pure separation*) or $R \neq 1$ and R cannot then be written in the form $R = R_1(u)R_2(v)R_3(w)$.

This property of R-separation for the Laplace equation happens to be a general trend. The PDE equations admit more non equivalent R-separable coordinate systems than separable coordinate systems. Here we shall be mainly concerned with the separation options for the conformal Laplacian, Eq.s(3.12a),(3.15), and its classical Hamilton-Jacobi analog- the equation for the *null geodesics*

$$g^{ij} \partial_i W \partial_j W = 0. \quad (4.3)$$

Our interest in conformal (Yamabe) Laplacian and null geodesics equations is coming from the results of Appendix B. In it we demonstrated that the Reeb dynamics is always taking place on null geodesics. Quantum mechanically this leads to study of solutions of the conformal (Yamabe) Laplacian equation. The Eq.(4.3) is *additively separable*. This means that there is a system of coordinates $\{q_i\}$ in which it can be split into the system of ordinary differential equations implying the solution in the form

$$W = \sum_{i=1}^d W(q_i; c_\alpha), c_\alpha \in \mathbf{R}, \alpha = 1, \dots, d. \quad (4.4)$$

Here c_α are some known (pre assigned) constants. For the conformal Laplacian and in Eq.(4.3) it is assumed that: a) $g^{ii} \neq 0$ and b) $g^{ij} = 0$ for $i \neq j$. This leads to separability in terms of orthogonal coordinates. Both Eq.s(3.15) and (4.3) are manifestly conformally invariant. Suppose now that Ψ is R-separable solution of Eq.(3.15), i.e. of the equation $\square_g \Psi = 0$. Then, $e^{-(\frac{d}{2}-1)f} \Psi = \tilde{\Psi}$ is also a solution of $\square_{\tilde{g}} \tilde{\Psi} = 0$.

Corollary 4.2. If $\square_g \Psi = 0$ possesses an R-separable solution, then $\square_{\tilde{g}} \tilde{\Psi} = 0$ also possesses an R-separable solution. Thus, R-separability of the conformal Laplacian is conformally invariant property.

Corollary 4.3. The equation $\square_g \Psi = 0$ is R-separable only if the metric g is conformally flat. A necessary condition for the R-separability of Eq.(3.15) is for the Hamilton-Jacobi equation for the null geodesic, Eq.(4.3), to be additively separable in the same separable coordinates [70].

Remark 4.4. Pure separation is a special case of the R-separation. This follows from Eq.(3.25).

Remark 4.5. R-separability is tightly linked with the superintegrability. The superintegrability in turn is tightly linked with the fact that for superintegrable systems all dynamical trajectories are closed [71]²⁴. Superintegrable $2d$ dimensional dynamical system has more than d integrals of motion. Those systems which have d integrals of motion are just integrable. Since for superintegrable systems all dynamical trajectories are closed brings us instantly to the problematics of Bertrand spacetimes [20],[74]. Dynamics in such spaces is superintegrable by definition. Accordingly, such dynamics admits quantization. Further details on superintegrability and Bertrand spaces are presented in section 5.

Based on the results of previous section these properties single out Schrödinger's Eq.(3.19a) (in which we have to put $\gamma = 1$) for the hydrogen atom. Other physically meaningful solutions are obtainable from that for hydrogen atom with help of conformal transformations by using Eq.(3.13).

In [7, 8, 17, 49, 67] the spherical symmetry of the potential, Eq.(3.19b) (for any γ) was taken into account at the outset leading to the standard protocol of separation of variables just like that for the hydrogen atom. Unfortunately, all these treatments are mathematically incorrect. This is explained in detail below, in this section. Nevertheless, the account of spherical symmetry was done correctly leading to separation of variables. Use of Corollaries 4.2. and 4.3. allows then to obtain different integrable cases from the one (seed) solution, e.g. that for the hydrogen atom.

Relevant fundamentals of contact geometry are described in Appendix B. The essence of the Eisenhart lift is nicely summarized in [75]. At the level of classical mechanics the Eisenhart lift is a method of enlargement of the phase space of dynamical system resulting in the associated system whose dynamics is taking place on geodesics. Because of this, such a lift serves as an alternative to the Jacobi- Maupertius formulation of classical dynamics as described in Eq.(B.16) of Appendix B. A slight elaboration on Jacobi- Maupertius formulation of mechanics leads to the equation describing the Reeb dynamics on null geodesics, that is to Eq.(4.3). From here it follows that use of the Eisenhart lift is facilitating separability and superintegrability [76]. In the formalism of Eisenhart lift Eq.(4.3) is interpreted as the null constraint $\mathcal{H} = 0$ imposed on new Hamiltonian \mathcal{H} . This constraint is essential for recovery of the original dynamics from the lifted one [77]. The equation $\mathcal{H} = 0$ coincides with that for the Reeb vector field as explained in Appendix B (after Eq.(B.17)). The additive solution for the null geodesics, Eq.(4.3), depends on c_α constants defined in Eq.(4.4). As mentioned above, the additivity of the solution of Eq.(4.3) is essential for the separability (or R-separability) of the solution of equation $\square_g \Psi = 0$. Different sets of admissible constants

²⁴The latest applications of superintegrability in optics is given, for example, in [72]. Closed geodesics sometimes are called periodic geodesics in mathematical literature [73]

for Eq.(4.3) cause different type of R-separability in Eq.(3.15). It happens that existence of separable coordinates has a coordinate-free characterization [78]. This property leads to the following

Definition 4.6. *Stäckel transform* maps the constants of motion for one orthogonally separable system to the constants of motion for another orthogonally separable system. Two systems related by a sequence of Stäckel transforms are *Stäckel-equivalent*.

Remark 4.7. Stäckel equivalence and superintegrability are inseparably linked to each other [79]. Recently Tsiganov demonstrated [80] how to determine Stäckel matrices preserving superintegrability. Since the number of R-separable coordinates is always finite (for a given dynamical system), the number of Stäckel transforms is also finite. Evidently, none of them changes physics. But all of them are linked with the topology of the underlying manifolds through the issue of closed/periodic geodesics [81]. Although this topic is still open for study, references [76],[82] might serve as good starting point for studying this issue.

The superintegrability for Kepler-Coulomb mechanical problem was studied recently in connection with Bertrand spaces in [20]. This fact is helpful. It will be further considered in section 5. It cannot be used immediately though for reasons which will become evident upon reading. It appears, that the topic of obtaining solutions for the Schrödinger equation for hydrogen atom is so well known that there is nothing which can be added to this subject. This prevailing opinion happens to be incorrect. Indeed, what is taught in any class on quantum mechanics is that the hydrogenic wave functions Ψ are given by the product [83]

$$\Psi_{nlm}(r, \theta, \varphi) = \text{const} \rho^l e^{-\frac{r}{2}} L_{n+l}^{2l+1}(\rho) Y_{lm}(\theta, \varphi). \quad (4.5)$$

Here $\rho = \frac{2r}{n}$, $n = \sqrt{\frac{1}{-2E}}$, r is the radial coordinate, E is the energy, $E < 0$, $Y_{lm}(\theta, \varphi)$ are the spherical harmonics, $L_{n+l}^{2l+1}(\rho)$ are the generalized Laguerre polynomials. The system of units was chosen in which Bohr's radius $a_0 = 1$. The constant in Eq.(4.5) is determined by the normalization of wave function. Since these results are standard, no more details are provided. Much lesser known are four dimensional representations of the hydrogen wave functions taking into account explicitly the hidden $O(4)$ symmetry²⁵. According to [84],[87] in 4 dimensions the wave function is looking as follows

$$Y_{nlm}(\alpha, \theta, \varphi) = 2^{l+1} l! \left[\frac{n(n-l-1)!}{2\pi(n+l)!} \right]^{\frac{1}{2}} \left[\sin^l \alpha \right] C_{n-l-1}^{l+1}(\cos \alpha) Y_{lm}(\theta, \varphi). \quad (4.6a)$$

²⁵Surely, the total dynamical symmetry of the hydrogen atom is $SO(4,2)$. This fact is discussed at length in [1] and [3]. Nevertheless, the account of $O(4)$ symmetry is sufficient for now [84]. Once the results for wave functions with $O(4)$ symmetry are known, the $SO(4,2)$ results are easily obtainable with help of the wavefunctions with $O(4)$ symmetry [1], [85], [86].

Here the spherical harmonics $Y_{lm}(\theta, \varphi)$ are the same as in Eq.(4.5) while $C_{n-l-1}^{l+1}(\cos \alpha)$ are the Gegenbauer polynomials to be discussed at length below. Unlike 3 dimensional spherical system of coordinates which is well known, the 4 dimensional system of coordinates

$$\begin{aligned} x_1 &= r \sin \alpha \sin \theta \cos \varphi, \\ x_2 &= r \sin \alpha \sin \theta \sin \varphi, \\ x_3 &= r \sin \alpha \cos \theta, \\ x_4 &= r \cos \alpha. \end{aligned} \quad (4.7)$$

is known in standard textbook literature much less. It should be noticed, that already in 3 dimensions there is a number of separable coordinate systems. In addition to spherical, there are cylindrical, parabolic, etc. coordinate systems. These are Stäckel-equivalent and, hence, make the hydrogen atom a superintegrable system [20]. The choice of coordinate system is dictated by the physics of the problem. In 4 dimensions the number of separable coordinate systems increases remarkably. Many of them are listed in [84]. However, in [87] yet another coordinate system, introduced by Bateman in 1905, is discussed²⁶. It will be described in section 5.

In the meantime, the normalization conditions for $Y_{nlm}(\alpha, \theta, \varphi)$ are

$$\int_0^\pi \sin^2 \alpha d\alpha \int_0^\pi \sin \theta d\theta \int_0^{2\pi} d\varphi \bar{Y}_{n'l'm'} Y_{nlm} = \delta_{n,n'} \delta_{m,m'} \delta_{l,l'}. \quad (4.8)$$

Interestingly enough, Eq.s (4.5), (4.6) can be compared with each other thanks to the result published as Appendix 5 of the book [89]. By introducing the Fourier transform of $\Psi_{nlm}(r, \theta, \varphi)$ via

$$\Phi_{nlm}(\mathbf{p}, \theta, \varphi) = (2\pi)^{-\frac{3}{2}} \int d^3 r e^{-i\mathbf{p}\cdot\mathbf{r}} \Psi_{nlm}(r, \theta, \varphi), \quad (4.9)$$

where we had put $\hbar = 1$, after some very detailed calculation described in [92], the final result is looking as follows: $\Phi_{nlm}(\mathbf{p}, \theta, \varphi) = F_{nl}(\mathbf{p}) Y_{lm}(\theta, \varphi)$,

$$F_{nl}(\mathbf{p}) = \left[\frac{2}{\pi} \frac{(n-l-1)!}{(n+l)!} \right]^{\frac{1}{2}} n^2 2^{2l+2} l! \frac{(np)^l}{[(np)^2 + 1]^{l+2}} C_{n-l-1}^{l+1} \left(\frac{1 - (np)^2}{1 + (np)^2} \right). \quad (4.10a)$$

To compare this result with Eq.(4.6a) we need to consult [84], page 649. According to this reference we should get an equality²⁷ if we multiply the 3-dimensional result, Eq.(4.9), by the p -factor defined by

$$Y_{nlm}(\alpha, \theta, \varphi) = \frac{p_0^{-1}(p^2 + p_0^2)^2}{4} \Phi_{nlm}(\mathbf{p}, \theta, \varphi). \quad (4.10b)$$

²⁶E.g. read [88].

²⁷It should be noted that in [84] the combination $p_0^{-1}(p^2 + p_0^2)^2$ was written as $p_0^{-\frac{5}{2}}(p^2 + p_0^2)^2$. This is caused by the fact that, instead of the constant factor $[\frac{n(n-l-1)!}{2\pi(n+l)!}]^{\frac{1}{2}}$ taken from [87] in Eq.(4.6a) the factor $[\frac{(l+1)(n-l-1)!}{2\pi(n+l)!}]^{\frac{1}{2}}$ was used erroneously.

To check the correctness of this proposition, we adopt the spherical coordinates defined by Eq.(4.7) to the case of 3-sphere of unit radius and use the Fock parametrization of this 3-sphere [37],[93]²⁸. Specifically, in this parametrization

$$x_4 = \frac{p_0^2 - p^2}{p_0^2 + p^2} = \frac{1 - (np)^2}{1 + (np)^2} = \cos \alpha. \quad (4.11)$$

We also took into account that $p_0 = \sqrt{-E}$, $E < 0$, $-E = 1/n$, $m = 1/2$, m is the electron mass in selected system of units. Based on this information, we obtain:

$$(np)^2 = \frac{1 - \cos \alpha}{1 + \cos \alpha}. \quad (4.12)$$

Next, using this result, we obtain as well:

$$(np)^2 + 1 = \frac{2}{1 + \cos \alpha}, \text{ and } \left(\frac{1 - \cos \alpha}{1 + \cos \alpha} \right)^l \left(\frac{1 + \cos \alpha}{2} \right)^{l+2}$$

and, therefore,

$$p_0^{-3} \left[(np)^2 + 1 \right]^2 \frac{(np)^l}{\left[(np)^2 + 1 \right]^{l+2}} = p_0^{-3} \left(\frac{1 - \cos \alpha}{1 + \cos \alpha} \right)^{\frac{l}{2}} \left(\frac{1 + \cos \alpha}{2} \right)^l = p_0^{-3} \sin^l \alpha. \quad (4.13)$$

Using this result in Eq.(4.10a), accounting for Eq.(4.10b) and the comparing the result with Eq.(4.6) we obtain:

$$Y_{nlm}(\alpha, \theta, \varphi) = 2^{l+1} l! \left[\frac{n(n-l-1)!}{2\pi(n+l)!} \right]^{\frac{1}{2}} \left[\sin^l \alpha \right] C_{n-l-1}^{l+1}(\cos \alpha) Y_{lm}(\theta, \varphi). \quad (4.6b)$$

The obtained result is in perfect agreement with Eq.(4.6a), as required. These results allow us now to move further. In particular, we would like to consider a solution of the Schrödinger Eq.(3.19a) with the potential Eq.(3.19b). D-O formally obtained its solution already in their first work, [7], for any γ . We found, however, more convenient to discuss the same equation and its solution following [49]. In Eq.s(3.19a),(3.19b) and in [7],[49] this equation was treated as 3-dimensional while in [17] Ostrovsky did consider the 4 dimensional version of the same equation, but the obtained solution is not consistent with mathematically rigorous results of [87]. Therefore his results are to be reconsidered below.

In section 1 we noticed that initially Demkov and Ostrovsky [7] believed that "Maxwell's fish-eye problem is **closely related to** the Coulomb problem". In section 3 we discussed the geometrical meaning of the stationary Schrödinger's equation for the hydrogen atom.

Now it is time to reproduce Eq.(4.6a) with help of Eq.s(3.19a),(3.19b) ($\gamma = 1$). By doing so

²⁸See also Eq.s(2.2),(2.3) in addition.

we are going to replace the statement of D-O about "close relationship" by the statement "isomorphic". Following [49], we reproduce Eq.s(3a),(3b) and (6) from this reference. Thus, we have

$$\Psi(\rho, \theta, \varphi) = \text{const} \frac{\rho^{l+1}}{(1 + \rho^2)^{\frac{2l+1}{2}}} C_{n-l-1}^{l+1}(\xi(\rho)) Y_{lm}(\theta, \varphi), \quad (4.14)$$

where $\xi(\rho) = \frac{1 - \rho^2}{1 + \rho^2}$. A comparison between Eq.s(4.6a) and (4.14) suggests that we have to rewrite the factors containing the variable ρ accordingly. Fortunately, this can be done almost immediately by analogy with results displayed in Eq.s(4.11)-(4.14). Now we have to write $(np)^2 = \rho^2$, so that we obtain:

$$\cos \alpha = \frac{1 - \rho^2}{1 + \rho^2}; \rho^2 = \frac{1 - \cos \alpha}{1 + \cos \alpha}; \rho^2 + 1 = \frac{2}{1 + \cos \alpha}. \quad (4.15a)$$

With help of these results we establish that $\xi(\rho) = \cos \alpha$,

$$\left(\frac{1 + \cos \alpha}{2} \right)^{\frac{2l+1}{2}} \left(\frac{1 - \cos \alpha}{1 + \cos \alpha} \right)^{\frac{l+1}{2}} = 2^{\frac{2l+1}{2}} \sin^l \alpha \quad (4.15b)$$

Using these results in Eq.(4.14) brings us back to Eq.(4.6a) so that needed isomorphism is established, provided that we resolve the apparent paradox. Eq.s(3.19a),(3.19b) as well as Eq.s(3a),(3b) and (6) of [49] are manifestly 3 dimensional while the result, Eq.(4.6a), is 4 dimensional. It is also rather weird to have the 3-dimensional spherical harmonics $Y_{lm}(\theta, \varphi)$ in the 4-dimensional result, Eq.(4.6). These circumstances prompt us to dig deeper into the issues just mentioned.

In the entire section 3 we dealt exclusively with the 3 dimensional versions of Eq.s (3.19a),(3.19b). In the already cited references, e.g.[7],[49], the 3 dimensional versions of Eq.s (3.19a),(3.19b) were treated and solved. Just above we noticed that these solutions carry information about the 4th dimension. It was left unnoticed in [7],[49] while in [17] Ostrovsky made an attempt to solve the 3-dimensional Eq.s(3.19a),(3.19b) by using the 3-sphere living in 4 dimensional Euclidean space (without justification for this step whatsoever). His results are not mathematically rigorous for reasons just described. Below, we provide a detailed explanation. Notice that our demonstration has almost nothing in common with the celebrated 4-dimensional solution of the Schrödinger equation for the hydrogen atom developed by Fock [37]. Nevertheless, the end result of our calculation will reproduce the 4-dimensional Eq.(4.6a) taken from [87]. Thus, we are about to uncover the "invisible" influence of the 4th dimension on solutions of 3-dimensional equations (3.19a), (3.19b). Our arguments are complementary to those by Fock [37].

We begin with description of pentaspherical coordinates introduced in section 7 of our work [3]. This time, however, we shall provide additional details following [69] and [90]. The pentaspherical coordinates had been discussed in accessible form in the monograph by Felix Klein cited in [3]. Mathematical usefulness of these coordinates was recognized already

by Böcher at the end of 19th century and was rediscovered at the end of the 20th century and further developed by Miller and collaborators [69]. Physical usefulness of pentaspherical coordinates was explained in [3]. For uninterrupted reading, we provide here some excerpts from [3] on this topic. We begin with consideration of a sphere S^2 in \mathbf{R}^3 . Analytically, the S^2 is described as

$$x^2 + y^2 + z^2 - 2ax - 2by - 2cz + D = 0. \quad (4.16a)$$

The radius r of the sphere is defined via

$$r^2 = a^2 + b^2 + c^2 - D. \quad (4.16b)$$

The introduced parameters are considered as coordinates in the space of spheres. In addition, it is convenient to introduce the homogenous (projective) coordinates

$$a = \frac{\xi}{\nu}, b = \frac{\eta}{\nu}, c = \frac{\zeta}{\nu}, r = \frac{\lambda}{\nu}, D = \frac{\mu}{\nu}. \quad (4.17)$$

These projective coordinates allow us to embed S^2 into projective space \mathbf{RP}^5 resulting in the equation

$$\xi^2 + \pi^2 + \zeta^2 - \lambda^2 - \nu\mu = 0 \quad (4.18a)$$

or, with $\nu\mu = \alpha^2 - \beta^2$, its analog

$$\xi^2 + \pi^2 + \zeta^2 + \beta^2 - \lambda^2 - \alpha^2 = 0. \quad (4.18b)$$

Although in pentaspherical coordinates (x_1, \dots, x_5) one uses a collection of 5 orthogonally intersecting spheres, just described mathematics remains very much the same. Specifically, the quadric, Eq. (4.18), now is written as the cone equation

$$\sum_{i=1}^5 x_i^2 = 0. \quad (4.19)$$

Let $(x, y, z) \in \mathbf{R}^3$. Via stereographic projection (e.g. see Eq.s (2.2), (2.3)) these Euclidean coordinates are related to the spherical coordinates $(s_1, \dots, s_4) \in S^3$, $\sum_{i=1}^4 s_i^2 = 1$. As in Eq.s (2.2) and (2.3), we obtain analogously:

$$x = \frac{s_1}{1 + s_4}, y = \frac{s_2}{1 + s_4}, z = \frac{s_3}{1 + s_4} \quad (4.20a)$$

so that

$$s_1 = \frac{2x}{r^2 + 1}, s_2 = \frac{2y}{r^2 + 1}, s_3 = \frac{2z}{r^2 + 1}, s_4 = \frac{1 - r^2}{1 + r^2}, r^2 = x^2 + y^2 + z^2. \quad (4.20b)$$

Next, it is convenient to introduce the analog of Eq.s (4.17) by taking into account Eq. (4.19). Thus, we write:

$$x_1 = 2XT, x_2 = 2YT, x_3 = 2ZT, x_4 = X^2 + Y^2 + Z^2 - T^2, x_5 = i(X^2 + Y^2 + Z^2 - T^2). \quad (4.21)$$

Accordingly,

$$\begin{aligned} x &= \frac{X}{T} = \frac{-x_1}{x_4 + ix_5}, \quad y = \frac{Y}{T} = \frac{-x_2}{x_4 + ix_5}, \quad z = \frac{Z}{T} = \frac{-x_3}{x_4 + ix_5}, \\ x^2 + y^2 + z^2 - 1 &= \frac{-2x_4}{x_4 + ix_5}, \quad x^2 + y^2 + z^2 + 1 = \frac{2ix_5}{x_4 + ix_5}. \end{aligned} \quad (4.22)$$

Eq.s(4.21) imply (via duality) the following chain of equalities (taken from [90],page 403):

$$\partial_X = 2T\partial_{x_1} + 2X\partial_{x_4} + 2iX\partial_{x_5}, \quad \partial_Y = 2T\partial_{x_1} + 2Y\partial_{x_4} + 2iY\partial_{x_5}, \quad \partial_Z = 2T\partial_{x_1} + 2Z\partial_{x_4} + 2iZ\partial_{x_5}. \quad (4.23)$$

Also,

$$\partial_x = T\partial_X, \quad \partial_y = T\partial_Y, \quad \partial_z = T\partial_Z \quad (4.24)$$

and, following [69], [90], we obtain as well:

$$\begin{aligned} \partial_x &= -(x_4 + ix_5)\partial_{x_1} + x_1(\partial_{x_4} + i\partial_{x_5}), \quad \partial_y = -(x_4 + ix_5)\partial_{x_2} + x_2(\partial_{x_4} + i\partial_{x_5}), \\ \partial_z &= -(x_4 + ix_5)\partial_{x_3} + x_3(\partial_{x_4} + i\partial_{x_5}). \end{aligned} \quad (4.25a)$$

By replacing the quantum $-i\partial_x$ by the classical p_x , *etc.* we obtain the classical analog of Eq.(4.25a)

$$\begin{aligned} p_x &= -(x_4 + ix_5)p_{x_1} + x_1(p_{x_4} + ip_{x_5}), \quad p_y = -(x_4 + ix_5)p_{x_2} + x_2(p_{x_4} + ip_{x_5}), \\ p_z &= -(x_4 + ix_5)p_{x_3} + x_3(p_{x_4} + ip_{x_5}) \end{aligned} \quad (4.25b)$$

The conformally flat space classical Hamiltonian \mathcal{H} now reads

$$\mathcal{H} = p_x^2 + p_y^2 + p_z^2 + V(\mathbf{x}) = 0 = (x_4 + ix_5)^2 \{p_{x_1}^2 + p_{x_2}^2 + p_{x_3}^2 + p_{x_4}^2 + p_{x_5}^2 + \tilde{V}(x)\} = 0 \quad (4.26a)$$

so that $(x_4 + ix_5)^2 \tilde{V}(x) = V(\mathbf{x})$. Now we adopt the quadric, Eq.(4.19), to S^3 . This is done by requiring $x_5 = -i$. Accordingly, using Eq.(4.19), now we obtain:

$$\sum_{i=1}^4 \frac{x_i^2}{-x_5^2} = 1,$$

implying $s_1 = x_1$, *etc.* In view of this, Eq.(4.26a) can be rewritten as

$$(x_4 + 1)^2 \{p_{x_1}^2 + p_{x_2}^2 + p_{x_3}^2 + p_{x_4}^2 + \tilde{V}(x)\} = 0. \quad (4.26b)$$

By comparing Eq.s(4.26a),(4.26b) with Eq.s(3.19a),(3.19b) adapted to the case $\gamma = 1$, we have to rewrite the potential, Eq.(3.19b) taking also into account Eq.(2.14b) and discussion which follow after this equation. Thus, we obtain:

$$V(\mathbf{x}) = \frac{(Ze^2)^2}{|E_n|} \left(\frac{1}{1 + r^2} \right)^2 \Big|_{S^3} = (x_4 + 1)^2 \frac{(Ze^2)^2}{4|E_n|} \quad (4.27)$$

In Eq.(4.26b) the combination $p_{x_1}^2 + p_{x_2}^2 + p_{x_3}^2 + p_{x_4}^2$ must be multiplied by the factor 1/2 in accord with the discussion following Eq.(2.14b). Furthermore, this combination represents not the flat Laplacian. This fact is fundamentally nontrivial. Its origin is based on the following chain of arguments.

Using Eq.(4.26a) we have to require that p_{x_i} and x_i to be the canonically conjugate pair of variables. If, indeed, this is so then, say, at the classical level of description we have to calculate the Poisson bracket (P.B). Thus, we obtain:

$$\left\{ \sum_{i=1}^5 x_i^2, \mathcal{H} \right\}_{P.B.} = 2(x_4 + ix_5)^2 \sum_{i=1}^5 x_i p_i = 0. \quad (4.28)$$

Adopted to S^3 , Eq.(4.28) reads:

$$\sum_{i=1}^4 x_i p_i = 0. \quad (4.29)$$

Thus, we have to solve Eq.(4.26b) subject to the constraint, Eq.(4.29). The solution of this problem is nontrivial. It is presented in the Appendix E. With help of obtained results, below we shall recover the 4-dimensional hydrogen wave function, Eq.(4.6a).

The 4-dimensional wave function, Eq.(4.6a), is presented in [84],[87]. In both references it was given without derivation. Therefore, we would like to restore needed details being guided by the results of our Appendix E, Ref.[91], and known quantum mechanical results for the 3-dimensional rigid rotator. At this point we need to extend these 3 dimensional results to the 4th dimension. In 3 dimensions the well known Casimir operator $\hat{\mathbf{L}}^2 = \hat{J}_x^2 + \hat{J}_y^2 + \hat{J}_z^2$ is used together with \hat{J}_z to label the spherical wavefunctions $Y_{lm}(\theta, \varphi)$. The first task now lies in finding the 4-d Casimir operator (or operators)-the analog(s) of $\hat{\mathbf{L}}^2$. Fortunately, the results can be found in [87], pages 112-113. These are taken from [91],page 248. In Appendix E we argue that the 4-dimensional analogs of \hat{J}_x, \hat{J}_y and \hat{J}_z are the quantized Plücker coordinates, e.g. $\hat{l}_{12}, \hat{l}_{13}, \hat{l}_{14}, \hat{l}_{23}, \hat{l}_{24}$ and \hat{l}_{34} so that $\hat{\mathbf{L}}^2$ is replaced now by $\hat{\mathcal{L}}^2 = \hat{l}_{12}^2 + \hat{l}_{13}^2 + \hat{l}_{14}^2 + \hat{l}_{23}^2 + \hat{l}_{24}^2 + \hat{l}_{34}^2$. Explicitly, in terms of spherical coordinates defined in Eq.(4.7) we obtain:

$$\hat{\mathcal{L}}^2 = -\frac{\partial^2}{\partial \alpha^2} - 2 \cot \alpha \frac{\partial}{\partial \alpha} - \frac{1}{\sin^2 \alpha} \left[\frac{\partial^2}{\partial \theta^2} + \cot \theta \frac{\partial}{\partial \theta} + \frac{1}{\sin^2 \theta} \frac{\partial^2}{\partial \varphi^2} \right]. \quad (4.30a)$$

Notice, however, that $\hat{\mathbf{L}}^2 Y_{lm}(\theta, \varphi) = \hbar^2 l(l+1) Y_{lm}(\theta, \varphi)$. Because of the known relation

$$\frac{\hat{\mathbf{L}}^2}{\hbar^2} = -\left[\frac{1}{\sin \theta} \frac{\partial}{\partial \theta} (\sin \theta \frac{\partial}{\partial \theta}) + \frac{1}{\sin^2 \theta} \frac{\partial^2}{\partial \varphi^2} \right] = -\left[\frac{\partial^2}{\partial \theta^2} + \cot \theta \frac{\partial}{\partial \theta} + \frac{1}{\sin^2 \theta} \frac{\partial^2}{\partial \varphi^2} \right],$$

Eq.(4.30a) should be rewritten accordingly as

$$\hat{\mathcal{L}}^2 = -\frac{\partial^2}{\partial \alpha^2} - 2 \cot \alpha \frac{\partial}{\partial \alpha} + \frac{1}{\sin^2 \alpha} \frac{\hat{\mathbf{L}}^2}{\hbar^2}. \quad (4.30b)$$

By analogy with 3 dimensions, in 4 dimensions we write instead:

$$\hat{\mathcal{L}}^2 Y_{nlm}(\alpha, \theta, \varphi) = I_{nl} Y_{nlm}(\alpha, \theta, \varphi). \quad (4.31a)$$

The eigenvalue I_{nl} is to be determined from the equation

$$\left[\frac{l(l+1)}{\sin^2 \alpha} - \frac{\partial^2}{\partial \alpha^2} - 2 \cot \alpha \frac{\partial}{\partial \alpha} \right] \Psi_{nl}(\alpha) = I_{nl} \Psi_{nl}(\alpha). \quad (4.32)$$

in which, in view of Eq.s(4.30b),(4.31a), we have to present $Y_{nlm}(\alpha, \theta, \varphi)$ as $\Psi_{nl}(\alpha)Y_{lm}(\theta, \varphi)$. Since the obtained equation is equivalent to Eq.(4.26b), we should make an identification:

$I_{nl} = \frac{-(Ze^2)^2}{2|E_n|}$. In arriving at this result we took into account that the combination $p_{x_1}^2 + p_{x_2}^2 + p_{x_3}^2 + p_{x_4}^2$ is coming along with the factor of $1/2$. This was explained after Eq.(2.14b).

The combined use of Eq.s (4.26b),(4.30b), (4.31),(4.32) then leads to $I_{nl} = \frac{-(Ze^2)^2}{2|E_n|}$.

Next, we let $x = \cos \alpha$ in Eq.(4.32). Then, we write $\Psi_{nl}(\alpha) = (1-x^2)^{\frac{1}{2}} F_{nl}(\alpha)$. With help of these substitutions Eq.(4.32) acquires the following look

$$(1-x^2) \frac{d^2}{dx^2} F_{nl} - (2l+3)x \frac{d}{dx} F_{nl} + [I_{nl} - l(l+2)] F_{nl} = 0. \quad (4.33)$$

This equation should be compared against the canonical equation for the Gegenbauer polynomials [92] :

$$(1-x^2) \frac{d^2}{dx^2} C_n^\lambda(x) - (2\lambda+1)x \frac{d}{dx} C_n^\lambda(x) + n(n+2\lambda) C_n^\lambda(x) = 0. \quad (4.34)$$

Upon comparison between these two equations, we obtain: $(2l+3) \rightleftharpoons (2\lambda+1)$ implying $\lambda = l+1$. Also, $n(n+2\lambda) \rightleftharpoons I_{nl} - l(l+2)$ implying $I_{nl} = (n+l+1)^2 - 1$. Let now $n+l+1 = 2F+1$ then, $(n+l+1)^2 - 1 = 4F(F+1)$. This result should be compared with those presented in [84],[87]. For this, it is sufficient to relabel $n+l+1$ as \hat{n} . By comparing with standard physics literature on hydrogen atom, we have to relabel n as n_r , where n_r is the radial quantum number. Such an identification between n as n_r is plausible but misleading even though it is present in all works by D-O and those who used the D-O works. Explanation is given at the end of section 5. With such a background, we obtain: $\hat{n}^2 - 1 = 4F(F+1)$. On page 5 of [87] we find that the 4 dimensional wave functions $Y_{\hat{n}lm}$ (e.g. see Eq.(4.31)) are belonging to the eigenvalue $\hat{n}^2 - 1$ as required. Next, we obtain : $n_r = 2F - l$ implying $F_{\hat{n}l}(\alpha) = C_{2F-l}^{l+1}(\cos \alpha)$. Since $\Psi_{\hat{n}l}(\alpha) = (1-x^2)^{\frac{1}{2}} F_{\hat{n}l}(\alpha)$, we obtain as well $\Psi_{\hat{n}l}(\alpha) = \text{const}(\sin \alpha)^l C_{2F-l}^{l+1}(\cos \alpha)$. Finally, since $Y_{nlm}(\alpha, \theta, \varphi) = \Psi_{\hat{n}l}(\alpha)Y_{lm}(\theta, \varphi)$, we end up (accounting for the normalization, Eq.(4.8)) with the wave function, Eq.(4.6a), as required.

Calculation of the eigenvalue spectrum for the hydrogen atom can be found in any text-book on quantum mechanics. Here, we approach this problem nontraditionally. In doing so, we borrow some ideas from the obscure paper by Schrödinger [93] (circa 1940) superimposed

with results of our Appendix F. Schrödinger was interested in finding the extent to which the effects of curvature of Universe may affect the spectrum of hydrogen atom. Schrödinger choose to consider the Keplerian motion on the hypersphere S^3 of radius R . Fundamentally interesting (for us) is to read about the rationale of his calculation. In his paper we find the following passages:

”It may appear foolish to pay attention to the extremely feeble curvature of the Universe in dealing with the hydrogen atom, because even the influence of those much stronger fields of gravitation in which all our observations are actually made (if the frame is properly chosen) entirely negligible. But this problem, by obliterating the sharp cut between ” elliptic and hyperbolic orbits”(the classical orbits are all closed) and by resolving the continuum spectrum into intensely crowded line spectrum, has extremely interesting features, well worth investigating...”

By looking for spaces *in which all classical orbits are closed*, Schrödinger actually was looking at

- a) Bläshke manifolds discussed in Appendices A and B,
- b) Superintegrable dynamical systems discussed in this and next sections,
- c) Bertrand spaces to be discussed in the next section.

He begins his calculations with the equation on S^3 (not on \mathbf{R}^3 !)

$$\frac{\hbar^2}{2mR^2} \left[-\frac{1}{\sin^2 \alpha} \frac{d}{d\alpha} (\sin^2 \alpha \frac{dS}{d\alpha}) + \frac{l(l+1)}{\sin^2 \alpha} S \right] - \frac{e^2}{R} S \cot \alpha - ES = 0. \quad (4.35a)$$

A quick look at our Eq.(4.30b) allows us to rewrite Eq.(4.35a) into more familiar form as follows

$$\frac{\hbar^2}{2mR^2} \hat{\mathcal{L}}^2 S = ES + \left[\frac{e^2}{R} \cot \frac{\alpha}{2} \right] S. \quad (4.35b)$$

In this equation we corrected Scrodinger’s result: in going from \mathbf{R}^3 to S^3 of radius R he replaced the Coulombic 3-dimensional term $\frac{e^2}{R}$ by $\frac{e^2}{R} \cot \alpha$. The correct result is: $\frac{e^2}{R} \cot \frac{\alpha}{2}$. It is worthy of demonstrating how this result is actually obtained. Using Eq.(4.20b) we write:

$$r^2 = \frac{1 - s_4}{1 + s_4}. \quad (4.36)$$

Hence,

$$\frac{1}{r} = \left(\sqrt{\frac{1 - s_4}{1 + s_4}} \right)^{-1} = \sqrt{\frac{(1 + s_4)^2}{1 - s_4^2}} \quad (4.37a)$$

Taking into account that $s_4 = x_4 = R \cos \alpha$, let $R = 1$, then we obtain:

$$\frac{1}{r} = \frac{1 + \cos \alpha}{\sin \alpha} = \frac{2 \cos^2 \frac{\alpha}{2}}{2 \sin \frac{\alpha}{2} \cos \frac{\alpha}{2}} = \cot \frac{\alpha}{2}. \quad (4.37b)$$

In our calculations we want to take advantage of the Coulomb-fish-eye isomorphism. Therefore, being guided by this isomorphism, instead of the factor $\frac{e^2}{R} \cot \frac{\alpha}{2}$ we systematically

used the factor $-\frac{(Ze^2)^2}{2|E_n|}$. This caused us to obtain the eigenvalue spectrum I_{nl} correctly in agreement with [84], [87]. However, this is not yet the spectrum for hydrogen atom. To obtain this spectrum correctly, we use the results of Appendix F. These are also allowing us to make a connection between our current results and those by Schrödinger. This is achieved with the amended Eq.(4.31a), that is, in view of Eq.(4.35b), with

$$\hat{\mathcal{L}}^2 Y = (I_{nl} - E)Y \quad (4.31b)$$

The choice of E is determined by replacing $I_{nl} = \hat{n}^2 - 1 = 4F(F+1)$ by $I_{nl} - E = 4F(F+1)$. From Appendix F we know that the parameter E can be chosen once and for all based on physical considerations. In our case, it is sufficient to choose $-E = -1$. With this choice we obtain:

$$-\frac{(Ze^2)^2}{2|E_n|} = (2F+1)^2 \text{ or } E_n = -\frac{(Ze^2)^2}{2(2F+1)^2}, F = 0, \frac{1}{2}, 1, \dots \quad (4.38)$$

This result coincides with that given on p.645 of [84] as required.

Remark 4.8. In the limit $R \rightarrow \infty$ the spectrum obtained by Schrödinger correctly reproduces the spectrum of hydrogen atom. However, the calculated wavefunctions differ from the standard, Eq.(4.6a). Since in this section we obtained both the eigenfunctions and eigenvalues for hydrogen atom correctly using methods and arguments different from those used by Fock [37], we leave for our readers to investigate further two topics. a) What logic made Schrödinger to use 4-dimensional calculations instead of 3-dimensional²⁹? Unlike Fock, who clearly stated the reason for using 4 dimensional calculations for hydrogen atom, Schrödinger, being apparently not aware of Fock's results, bypassed this issue altogether. b) It is of interest to recover the wavefunctions for hydrogen atom, Eq.(4.6a), using Eq.(4.35a).

The entire section 3 and a large part of Appendix F is devoted to the issues related to conformal invariance. Previously we mentioned the work by Makowski [53]. He took advantage of the fact that in 2 dimensions the Laplacian is conformally invariant. This fact enabled him to solve the 2-dimensional Sturmian Eq.(3.1b) for the fish-eye-type potentials exactly. In 3 dimensions the Laplacian no longer possess the property of conformal invariance. To bypass this difficulty the conformally invariant (Yamabe) Laplacian should be used instead. This fact created a lot of technical difficulties but allowed us to demonstrate in detail the full power of Hadamard-style calculations. These were only outlined in [3]. Such calculations are of significance not only for our work. To our knowledge, they have not been exposed in sufficient detail in physics literature, perhaps with exception of gravitation. But even in this field the alternative route of by-passing the issue of conformal invariance in 3 dimensions was not discussed also. It is associated with the systematic uses of the Hopf mapping. This mapping is not a novel item in physics literature [94]. Nevertheless, to our knowledge, the

²⁹Bertrand curved spaces of section 5 are 3 dimensional.

aspects we are willing to discuss are new. Because of this, they will be studied further in detail in Part II.

The idea of Hopf mapping is simple. Take the equation for the 3-sphere S^3 living in \mathbf{C}^2 , e.g. $|z_1|^2 + |z_2|^2 = 1$, $z_i = x_i + iy_i$, $i = 1, 2$, and look for the ratio $Z = \frac{z_1}{z_2}$ (or $Z = \frac{z_2}{z_1}$). While $S^3 \in \mathbf{C}^2$, $Z \in \mathbf{C}$. But $\mathbf{C} \cup \infty = S^2$! From this observation the Hopf map follows: $S^3 \rightarrow S^2$. Subsequently the map $S^2 \rightarrow S^3$ was also designed and used, e.g. read [95], etc. Above, using the pentaspherical coordinates we mapped the 3 dimensional Sturmian problem involving Eq.(3.19) into S^3 . We want now to reanalyze the 2 dimensional results by Makowski [53] using the notion of Hopf map. It should be noticed, though, that Makowski used the 2 dimensional plane \mathbf{C} in his work. By adding a point at infinity his results can be lifted to S^2 . E.g. look again at Eq.s(2.15)-(2.22) and (3.1a,b). The first thing we have to check: will his results reproduce known 2 dimensional results for the hydrogen atom on \mathbf{C} ? The answer depends upon our ability to solve the analogous problem on S^2 . On S^2 the potential, Eq.(3.19b), is constant. In fact, it is the same constant as we had calculated already: $\tilde{V}(x) = -\frac{(Ze^2)^2}{2|E_n|}$. The Laplacian restricted to S^2 is describing the rigid rotator. Therefore, we again run into the familiar equation: $\hat{\mathbf{L}}^2 Y_{lm}(\theta, \varphi) = \hbar^2 l(l+1) Y_{lm}(\theta, \varphi)$. The question remains: will this equation reproduce the 2-d hydrogen spectrum? Very fortunately, positive answer was found in [95]. These authors had adapted Pauli's dynamical group-theoretic method for solving hydrogen atom in 2 dimensions. By repeating the same steps, they indeed arrived at the rigid rotator equation. The Pauli-like group algebra enabled them to arrive at the equation for the spectrum

$$l(l+1) = -\left(\frac{1}{4} + \frac{1}{E}\right) \quad (4.39a)$$

leading to the known result:

$$E_l = -\frac{1}{(l + \frac{1}{2})^2}, \quad l = 0, 1, 2, \dots \quad (4.39b)$$

For the future reference, the 3-dimensional result taken from [84] and used in Eq.(4.38) is given as follows:

$$F(F+1) = -\left(\frac{1}{4} + \frac{1}{8E}\right), \quad F = 0, 1/2, 1, \dots \quad (4.40)$$

leading to $E_F = -\frac{1}{2n^2}$, $n = 2F + 1 = 1, 2, 3, \dots$

Both results can be made identical if one notices that Eq.(4.39a) uses only the angular momentum interpretation of the standard commutator algebra $[\hat{J}_j, \hat{J}_k] = i\epsilon_{jkl}\hat{J}_l$. In the case if the same commutator algebra is interpreted spin-theoretically, the results Eq.(4.39b) and (4.40) will coincide. This is becoming permissible in the case if the 2-dimensional results for S^2 are lifted to S^3 using the inverse Hopf map³⁰. We expect that the rest of 2-dimensional results to be discussed shortly below will follow the same trend.

³⁰Some technical details of this mapping are discussed at the end of section 5.

To prepare our readers for this discussion we recall that in mathematics the 3-dimensional rigid rotator eigenvalue problem is known under the name of eigenvalue equation problem for the associated Legendre polynomials $P_l^m(\cos \theta)$. In terms of the variable $x = \cos \theta$ the equation for these polynomials reads:

$$(1 - x^2) \frac{d^2}{dx^2} P_l^m(x) - 2x \frac{d}{dx} P_l^m(x) + [l(l+1) - \frac{m^2}{1-x^2}] P_l^m(x) = 0. \quad (4.41)$$

It should be compared with Eq.s(4.32)-(4.34). By analogy, we shall look for a solution of Eq.(4.41) in the form $P_l^m(x) = (1 - x^2)^{\frac{m}{2}} \bar{P}_l^m(x)$. Substitution of this ansatz into Eq.(4.41) leads to [97]

$$(1 - x^2) \frac{d^2}{dx^2} \bar{P}_l^m(x) - 2(m+1)x \frac{d}{dx} \bar{P}_l^m(x) + [l(l+1) - m(m+1)] \bar{P}_l^m(x) = 0. \quad (4.42)$$

But this is an equation for the Gegenbauer polynomials, Eq.(4.34)! In the simplest case, $m = 0$, we obtain: $\bar{P}_l^0(x) = C_n^{\frac{1}{2}}(x)$, $n = l$. For $m \neq 0$, we have to put $\lambda = m + 1/2$ in Eq.(4.34) and $l - m = n$ in Eq.(4.42) implying $\bar{P}_l^m(x) = C_n^{m+\frac{1}{2}}(x) = C_{l-m}^{m+\frac{1}{2}}(x)$. These results we would like to compare now against Eq.(9) of Makowski [53]. To do so, it is sufficient to make an identifications : Makowski's $\lambda \rightleftharpoons m$, also his $n_r + \lambda \rightleftharpoons n + m$. We are warning our readers not to confuse the factor λ in Gegenbauer's Eq.(4.34) and the factor λ in Makowski's paper [53]. Evidently, our n is the same as his n_r . Up to a complex factor $\exp(\pm im\varphi)$ the solution obtained by Makowski is given by

$$\psi_{n_r}(x, y) = \text{const}(1 - \xi^2)^{\frac{\lambda}{2}} C_{n_r}^{\lambda+\frac{1}{2}}(\xi), \quad \xi = \frac{1 - r^2}{1 + r^2} = \frac{1 - z^2}{1 + z^2}. \quad (4.43a)$$

Here we used our Eq.s (2.15)-(2.22) to write his Eq.(9) in our notations. This allows us to take into consideration that $\xi = \frac{1 - r^2}{1 + r^2} = Z = \cos \theta$ resulting in

$$\psi_{n_r}(\theta, \varphi) = N(\sin \theta)^m C_{n_r}^{m+\frac{1}{2}}(\cos \theta). \quad (4.43b)$$

Using Eq.(2.15), it is clear, that $x = \cos \theta \rightleftharpoons Z$. In addition, N is the normalization constant. We would like to compare Makowski's 2 dimensional result, Eq.(4.43a) (or Eq.(4.43b)), against that coming from the equation for the rigid rotator, Eq.(4.42). This equation is leading us to the solution

$$\psi_n(\theta, \varphi) = N'(\sin \theta)^m C_{l-m}^{m+\frac{1}{2}}(\cos \theta). \quad (4.44)$$

Since we just defined $l - m = n$ and $n = n_r$, this identification causes us to write $N' = N$. Makowski's result, Eq.(4.43b) was obtained using planar geometry while the result, Eq.(4.44), was obtained using the geometry/topology of S^2 . Use of stereographic projection connects these two geometries. When it comes to the connection between the respective differential equations -on \mathbf{C} and on S^2 -use of conformal invariance in Eq.(3.1b) is essential since the stereographic projection is a conformal mapping. Should the potential in the planar

Schrödinger equation be zero, the mapping $\mathbf{C} \rightleftharpoons S^2$ would transform the flat Laplacian to the Laplacian on S^2 . The Sturmian problem on \mathbf{C} is the eigenvalue problem for the Laplacian on S^2 . It is the familiar eigenvalue problem for the rigid rotator in quantum mechanics.

It is always possible to adapt the 3-dimensional Eq.(3.19) to S^2 as long as the conformal transformations are not involved. When they are involved, we only can lift Eq.(3.1b) to S^2 but we cannot make a reduction from 3 dimensions to 2 since in 3 and higher dimensions we are dealing with the conformal (Yamabe) Laplacian while in dimension 2- with the ordinary Laplacian which is conformally invariant. But we can use the stereographic projection from: from S^3 to \mathbf{R}^3 . Because of this, it is of interest to compare conformally modified planar results by Makowski and Peplowski [48] against the analogous 3-dimensional (actually the 4-dimensional, as demonstrated in previous subsection) results of Ref.[49]. Such a comparison

will give us an idea about the extent to which results on S^2 by Makowski [53], actually done on \mathbf{C} , change when they are lifted (via the Hopf map) to S^3 . It should be noted though that

we have not yet constructed rigorously the Hopf map. This is done in mathematical literature already and will be adapted for our needs in Part II.

To complete our calculation we need to reproduce Eq.s(4.39b). For this purpose we have to replace now the equation

$$\frac{\mathbf{L}^2}{\hbar^2} Y_{lm}(\theta, \varphi) = I_{lm} Y_{lm}(\theta, \varphi) \quad (4.45a)$$

by

$$\frac{\mathbf{L}^2}{\hbar^2} Y_{lm}(\theta, \varphi) = (I_{lm} - E) Y_{lm}(\theta, \varphi) \quad (4.45b)$$

where we have to choose, as before, $I_{lm} = -\frac{(Ze^2)^2}{2|E_n|}$ and select $-E = -\frac{1}{4}$. This choice follows from the identification

$$l(l+1) = -\frac{(Ze^2)^2}{2|E_n|} - \frac{1}{4} \quad (4.46a)$$

done with help of Eq.(4.42) and the property $l(l+1) + \frac{1}{4} = (l + \frac{1}{2})^2$. Using these results, we finally arrive at the equation for spectrum

$$E_l = -\frac{(Ze^2)^2}{(l + \frac{1}{2})^2}, l = 0, 1, 2, \dots \quad (4.46b)$$

Obtained result is in accord with Eq.(4.39b) as required. Furthermore, in the system of units $\hbar = 1, m = 1$, the obtained result exactly coincides with that reported in [98].

Now we want to discuss conformally deformed 2-dimensional results presented in [48]. They are immediately linked with results by the same authors just discussed. Instead of

duly reproducing the results of [48], we approach the whole problem differently. That is, first, in the absence of potential we lift the flat Laplacian from \mathbf{C} to S^2 . The conformal invariance is compatible with such a lift as we already know, e.g. see Eq.(3.1). Before performing any conformal transformation we notice that the potential $\tilde{V}(x)$ must be replaced by $l(l+1)$ in Eq.(4.45) in accord with theory of 2 dimensional rigid rotator. To study the effects of conformal transformations we analyze Eq.(5) of [48] by the same methods as we used in Eq.s(4.11)-(4.15). In the present case we start with

$$F_{nl}^{(\gamma)}(\xi) = N_{nl}^{(\gamma)}(1 - \xi^2)^{\frac{l}{2\gamma}} C_n^{l/\gamma+1/2}(\xi), \xi = \frac{1 - r^{2\gamma}}{1 + r^{2\gamma}}. \quad (4.47a)$$

To be compatible with our notations we replaced the k - factor in [48] by the γ factor defined in our Eq.(3.19). Before performing any conformal transformation we put $\gamma = 1$ in the above result to check it against Makowski's Eq.(4.43b) (see above). By comparing Eq.s (4.43) and (4.47a) we notice that l in Eq.(4.47a) must be replaced by m . If we treat $r^\gamma = \rho$ as independent variable, we still can use $\xi = \cos \theta = x$. Accordingly, we obtain:

$$F_{nm}^{(\gamma)}(\theta) = N_{nm}^{(\gamma)}(\sin \theta)^{\frac{m}{\gamma}} C_n^{m/\gamma+1/2}(\cos \theta), n = n_r. \quad (4.47b)$$

For $\gamma = 1$ Eq.s(4.43b) and (4.47b) coincide as required. Use of conformal transformations causes us to make a replacement $m \rightarrow m/\gamma$ in Eq.(4.43b) to reach an agreement with Eq.(4.47b). Such a replacement makes perfect physical sense because in this work the conformal transformation $f(z) = z^\gamma$ (e.g. see Eq.(2.22)) is used consistently. Since m is the azimuthal quantum number, the wave function boundary condition $\Psi(0) = \Psi(2\pi)$ prior to conformal mapping must be preserved after the conformal mapping is performed. This requirement is the cause of the transformation: $m \rightarrow m/\gamma$.

Now we are ready to analyze the 4-dimensional results of Ref.[49],page 34³¹. For $\gamma = 1$ this was done already in Eq.s (4.14),(4.15). Therefore, only the conformally deformed case is treated in this subsection. It is treated by analogy with that we just discussed when we considered the case of 2 dimensions. The additional help is coming from Eq.s(4.14),(4.15). Thus, now we obtain:

$$\cos \alpha = \frac{1 - \rho^{2\gamma}}{1 + \rho^{2\gamma}}; \rho^{2\gamma} = \frac{1 - \cos \alpha}{1 + \cos \alpha}; \rho^{2\gamma} + 1 = \frac{2}{1 + \cos \alpha}; \quad (4.48a)$$

$$\left(\frac{1 + \cos \alpha}{2}\right)^{\frac{2l+1}{2\gamma}} \left(\frac{1 - \cos \alpha}{1 + \cos \alpha}\right)^{\frac{l+1}{2\gamma}} = 2^{\frac{2l+1}{2\gamma}} \sin^{\frac{l}{\gamma}} \alpha \quad (4.48b)$$

Eq.(4.14) is replaced now by

$$\Psi(\alpha, \theta, \varphi) = \text{const}' (\sin^{\frac{l}{\gamma}} \alpha) C_{M - \frac{l}{\gamma} - 1}^{\frac{2l+1}{2\gamma} + \frac{1}{2}}(\cos \alpha) Y_{lm}(\theta, \varphi). \quad (4.49a)$$

³¹We would like to remind to our readers that the authors of Ref.[49] were not aware of 4 dimensional nature of their calculations.

Here, in accord with authors of [17] and [49], $M = n + (\gamma^{-1} - 1)l$. For $\gamma = 1$ Eq.(4.49a) coincides with Eq.s (4.14),(4.15) as required. Next, we compare this (4-dimensional) result against the 2-dimensional result, Eq.(4.47b), that is with

$$F_{nm}^{(\frac{1}{2})}(\theta) = N_{nm}^{(\frac{1}{2})}(\sin^{2m} \theta) C_{n_r}^{2m+1/2}(\cos \theta). \quad (4.47c)$$

In such a form we compare it now against Eq.(4.49a) in which we should put $\gamma = 1/2$:

$$\Psi(\alpha, \theta, \varphi) = \text{const}' (\sin^{2l} \alpha) C_{n_r}^{2l+\frac{3}{2}}(\cos \alpha) Y_{lm}(\theta, \varphi). \quad (4.49b)$$

This result is written in terms of notations used in [17] and [49]. In these references $M = n + (\gamma^{-1} - 1)l$ and, we have to write now $\gamma = 1/2$, and to use the standard result: $n = n_r + l + 1, n_r = 0, 1, 2, \dots$. The comparison implies that for $\gamma = 1/2$ the 4 dimensional wave function $\Psi(\alpha, \theta, \varphi)$ can be obtained with help of 2 dimensional $F_{n_r m}^{(\frac{1}{2})}(\theta)$ if we take into account that 3-dimensional $z = \cos \theta$ is equal to $t = \cos \alpha$ in 4 dimensions, e.g. see Eq.(4.7). Therefore we should make a replacement θ by $\theta \rightleftharpoons \alpha$ wherever this is appropriate. Next, we have to make a replacement $m \rightleftharpoons l$ and, finally, we have to use the addition formula [97],page121,

$$C_n^{\alpha+\beta}(x) = \sum_{m=0}^n C_m^{\alpha}(x) C_{n-m}^{\beta}(x). \quad (4.50)$$

After these replacements and use of the addition formula the final result should be multiplied by $Y_{lm}(\theta, \varphi)$. In the end, the obtained result must be properly normalized with help of the normalization condition for the Gegenbauer polynomials

$$\int_0^{\pi} d\alpha (\sin^{2p} \alpha) [C_l^p(\cos \alpha)]^2 = \frac{\pi \Gamma(2p+l)}{2^{2p-1} l! (l+p) \Gamma^2(p)}. \quad (4.51)$$

In writing the above results, following works by D-O and those who extended their works, we silently assumed that everything is fine with the result, Eq.(4.49a). But this is not at all the case. Mathematically correct results are presented below.

Calculation of the deformed spectrum is technically easy but, for the results to make physical sense, some prior discussion is absolutely essential. For readers convenience, needed background material is given in the Appendix G. In it we explain what is wrong with the existing "proofs" of the Madelung rule. We also describe the unexpected Hartree-Fock origins of the conformally deformed fish-eye potential, Eq.(3.19), with $\gamma = 1/2$. The description of this connection with the Hartree-Fock contribution to the effective potential $V_{eff}(\mathbf{r})$, Eq.(1.3), cannot be found among D-O papers cited thus far. Nevertheless, D-O did mention this connection in [99] and [100] but only in the context of the Thomas-Fermi (T-F) theory. The Appendix G does contain an information about the connection with the Hartree-Fock results. Yes, it is true that the contributions to the T-F theory made by Tietz resulted in the potential, Eq.s(G.8),(G.9), bearing his name. It is also true that, up to a constant factor, analytically

this potential is the same as the deformed fish-eye potential, Eq.(3.19) with $\gamma = 1/2$. But D-O have not cited the results by other authors in which it is demonstrated that *the combined use of the Coulombic and Hartree-Fock contributions to the effective potential $V_{eff}(\mathbf{r})$* , Eq.(1.2), *results in the deformed fish-eye potential with $\gamma = 1/2$.*

The problem of solving Eq.(1.2) had began with the work by Slater. In his 1932 paper [101] he writes " For any detailed calculations dealing with atomic or molecular structure, good approximations to the atomic wave functions are essential. The most satisfactory method, in general, for building up such functions to be by the use of one-electron functions which are solutions of the problem of an electron moving in a central field..." A noticeable advancement in this direction was made by Latter in 1955 [26]. He numerically solved Eq.(1.2) using the Thomas-Fermi-Dirac type potential for all elements with $Z = 1 \div 92$. Latter noticed that " The important error in this application of a fixed central field potential arises just from the use of the same potential for all configurations of the atom. This error clearly grows with increasing excitation of the atom above its ground state....The statistical potentials give acceptable values for ground state energies and, presumably, will therefore give semiquantitative predictions for states of low excitation." Results by Latter had been further analyzed by March. In his book [27] on p.76 March writes " In a Coulomb field we would, of course, fill the K, L, M shells successively. The closed shell of principal quantum number n holds $2n^2$ electrons and hence shells close, for a Coulomb field, for 2,10, 28, etc. electrons Now, in real atoms, the closed shell configurations occur of He ($Z=2$) and Ne ($Z=10$) which fit the Coulomb field scheme. But the next rare gas is Ar, with 18 electrons.... a screened Coulomb field such as given by Hartree or a Thomas-Fermi theory leads for an atomic number $Z=18$ to the filing of 3p sub shell which is rather far from the 3d level. Also, for larger atomic number, level crossing occurs. *The principal quantum number is no longer precise enough to group the energy levels: for a screened Coulomb field these depend upon n and the orbital angular momentum quantum number l* ". Thus, without using the word "Madelung rule", March recognized that the accurate calculations by Latter [26] provide the strongest possible justification of the validity of Madelung rule! It should be kept in mind that Latter compared the Thomas-Fermi and the Thomas-Fermi-Dirac solutions for energies of individual electrons against the available Hartree and Hartree-Fock results for respective atoms. All results by Latter are obtained numerically and are based on numerical solution of Eq.(1.2) with the Thomas-Fermi-Dirac type effective potential $V_{eff}(\mathbf{r})$. They demonstrate a very remarkable agreement with the Hartree and Hartree-Fock results. The solutions were obtained numerically because Latter was not using the Tietz potential. As discussed in the Appendix G, the Tietz potential is of Thomas-Fermi type but is of convenient analytical form and fits very nicely the Hartree-Fock data for the effective potential [25], p.664, Fig.10. Contrary to D-O descriptions of the Tietz work, Tietz recognized the qualities of his potential immediately after he discovered it. This caused him to apply the methodology developed by Kerner [102] in 1951 to the potential Tietz discovered later on. At the same time, Kerner, after consulting with Bethe and Feynman, obtained the exact analytical solution for the stationary Schrödinger equation having the potential $V_{eff}(\mathbf{r}) = -\frac{Ze}{r} \sqrt{\varphi(\frac{r}{a})}$. Here $\varphi(x)$ is the

same as in Eq.(G.8), except that Kerner was fiddling with the constant α (to get the best possible agreement with experiment) while in Eq.(G.8) this constant is fixed by the logic of the T-F calculations. Kerner tested his effective potential against the numerically obtained Hartree potential and found astonishingly good agreement as demonstrated in [102], Fig.1, page 71. Subsequently, Tietz adapted word-for-word Kerner's methodology for solving the Schrödinger's equation with the potential $V_{eff}(\mathbf{r}) = -\frac{Ze}{r}\varphi(\frac{r}{a})$. In [29] the latest attempt by Tietz at solving this type of Schrödinger equation is presented along with the list of all his previous attempts at exact solution. Unlike Latter, Kerner had not calculated the spectrum for all Z 's and for many l 's. He fixed his attention at the Mercury ($Z = 80$) and made a comparison of a single electron eigenvalues (for various n 's and l 's) calculated by his exact method against the relevant Hartree-type calculations. He obtained a good agreement with the Hartree results summarized in the Table II, page 74, of [102]. Kerner has no mention of the Madelung rule in his paper. Following into footsteps of Kerner, Tietz also selected Mercury for his study and compared his exact (Kerner-style) calculations against those published in Kerner's paper. Naturally, a good agreement with experimental data was also obtained.

D-O entered into research on this topic only in 1971-1972. Unlike other authors, they recognized the importance of restrictions caused by the Bertrand theorem and they reinterpreted the Tietz potential, without mentioning Tietz's name and his works, as the conformally deformed Maxwell's fish-eye potential. This is explained above, in section 2. This caused them to restrict their treatment [7],[17],[25] aimed at the proof of Madelung rule to studies of the Schrödinger equation (with Tietz potential) to the $E = 0$ case only. The rationale for doing so is discussed above, in sections 1–3. Already in section 2 we explained that the Schrödinger equation with the fish-eye potential **should** be used with $E = 0$ by design. This requirement has **nothing to do** with the energy spectrum for hydrogen atom though. It was calculated above, in this section. **But it has everything to do with the conformal invariance of the underlying Eq.(3.19) allowing us to conformally deform the Maxwell fish-eye potential thus allowing us to study the multielectron atoms with help of the same formalism.** Being unaware of just described facts, especially those, in the Appendix F, D-O studied only the $E = 0$ solution of the Schrödinger equation with the Tietz potential. At the same time, following the methodology developed by Kerner, Tietz obtained the exact solution of the same equation as studied by D-O in his 1956 paper [28] **without any restrictions on E .** The D-O method of solution of Eq.(3.19) differs from that by Tietz and, therefore, by Kerner. The analytical form of the wave function, Eq.(4.49b), is absent in the paper by Tietz.

Now, we are in the position to calculate the deformed spectrum using the above background information as well as results of Appendix C. We proceed by analogy with results just discussed. This brings us back to the deformed wave function Eq.(4.49b). The question emerges: Will this function be an eigenfunction of Eq.(4.31a)? A quick look at the analogous eigenfunction, Eq.(4.6a), and taking into account that $Y_{nlm}(\alpha, \theta, \varphi) = \Psi_{nl}(\alpha)Y_{lm}(\theta, \varphi)$, with $\Psi_{nl}(\alpha) = (1 - x^2)^{\frac{l}{2}}F_{nl}(\alpha)$, $x = \cos \alpha$ leads us to the following conclusions.

Using the factor $(1 - x^2)^{\frac{l}{2}} = \sin^l \alpha$ in $\Psi_{nl}(\alpha)$ and substitution of the ansatz $\Psi_{nl}(\alpha)Y_{lm}(\theta, \varphi)$

into Eq.(4.31a) allows us to get rid of the denominator $1/\sin^2 \alpha$ in Eq.(4.30b) (see also the analogous steps in Eq.s(4.41) and (4.42)). In view of Eq.(4.49b), in the deformed case the factor $(1-x^2)^{\frac{l}{2}}$ is replaced by the factor $(1-x^2)^l = \sin^{2l} \alpha$ in $\Psi_{n2l}(\alpha)$. When substituted back into Eq.(4.31a), calculation shows that only if we replace $l(l+1)$ by $2l(2l+1)$ in this equation the denominator $1/\sin^2 \alpha$ will disappear. Next, we need to adapt the fundamental Eq.(4.34) for the Gegenbauer polynomials to $C_{n_r}^{2l+\frac{3}{2}}(x)$ to our current needs. Such an adaptation produces:

$$(1-x^2)\frac{d^2}{dx^2}C_{n_r}^\lambda(x) - (4l+4)x\frac{d}{dx}C_{n_r}^\lambda(x) + n_r(n_r+4l+3)C_{n_r}^\lambda(x) = 0, \lambda = 2l + 3/2. \quad (4.52)$$

At the same time we should compare Eq.(4.52) with Eq.(4.33) in which we have to replace l by $2l$. Comparison indicates that the match is achieved only if we replace $C_{n_r}^{2l+\frac{3}{2}}(x)$ by $C_{n_r}^{\frac{l}{2}+1}(x)$ and take $\gamma = 1/2$. The combination $\left(\sin^{\frac{i}{\gamma}} \alpha\right) C_{n_r}^{\frac{l}{2}+1}(x)$ is fully consistent with two dimensional result, Eq.(4.47b). This is very plausible if we would like to apply the Hopf mapping to the two dimensional result, Eq.(4.47b), to get the result on S^3 . Furthermore, under such circumstances we can repeat word-for-word the discussion following Eq.(4.34), that is if $\lambda = 2l + 1$, then $2\lambda + 1 \rightleftharpoons 4l + 3$. Also, $n(n+2\lambda) \rightleftharpoons I_{nl} - 2l(2l+2)$ implying the relationship $n+2l+1 = 2F+1$. If, as before, we let $\hat{n} = n + l + 1$, then $2F+1 = \hat{n} + l$. By analogy with the derivation of the hydrogen spectrum, Eq.(4.38), we now obtain:

$$E_n = -\frac{Z^{\frac{7}{3}}e^4}{w(2F+1)^2}, F = 0, \frac{1}{2}, 1, \dots \text{ or, } E_n = -\frac{Z^{\frac{7}{3}}e^4}{w(\hat{n}+l)^2}, w = \text{const.} \quad (4.53)$$

In arriving at Eq.(4.53) we took into consideration the scaling analysis presented in Appendix G and in section 2. We also used the Remark 5.3. of the following section. The obtained result is fully consistent with the Madelung rule and group-theoretical analysis of periodic system [1], [103] done in close analogy with the Gell-Mann-Zweig theory of quark-hadron matter. It

should be noted though that the discussion of likely sources of anomalies of the Madelung rule given in [30] indicates that the group-theoretic methods are not capable of explaining these anomalies while the results presented thus far are fully compatible with the formalism describing the anomalies.

5. Bertrand spacetimes in the atomic physics and the Madelung rule

The title of our paper involves the phrase "modified Bertrand theorem". This terminology was introduced by Vler Perlick [23]. Perlick was interested in finding the relativistic-analogue of The Bertrand theorem. That is he wanted to find all spherically symmetric and static spacetimes (in the sense of general relativity) all of whose bounded trajectories are periodic. Although Perlick succeeded in finding such spacetimes (which he calls "Bertrand

spacetimes”), he was puzzled by the fact that the metric of such spacetimes is not reduced to the Schwartzshield metric used for description of spacetimes around massive objects, e.g. of our Sun. Such a metric leads to the precession of planetary trajectories rotating around the Sun. This result was explained by Einstein using his theory of general relativity. Classical Bertrand theorem forbids such a precession and, by design, its relativistic extension also forbids precession. 2 types of Bertrand metric found by Perlick caused Perlick to wonder about kind of a body (isolated from all other gravitational sources) that produces a metric in which all bounded trajectories are periodic. He concluded that ”such a body must be a rather exotic object”. The authors of [104] suggested that such a body should be identified with the dark matter. A quick introduction into this topic is given in our work [105], page 27. Further studies of dynamics in Bertrand spacetimes revealed [20] that dynamics in such spacetimes is superintegrable³² since the orbits are closed by design. Closure of orbits had prompted study of the quantum Bertrand systems [106], [107]. In these references no attempt was made to connect the obtained results with atomic physics. The purpose of this section is to demonstrate that the central object of our study, the D-O Eq.(3.19), justly belongs to the set of quantum Bertrand systems.

As before, this demonstration is essential to present in the proper context. Specifically, we begin with the Groenewold-van Hove (G-vH) theorem nicely explained in [108], [109]. Basically, it is telling us that only the harmonic oscillator-type dynamical systems can be unambiguously quantized. The Stäckel transform³³ connects the harmonic oscillator with the Kepler-Coulomb systems [106]. Only these two systems are permissible by the classical Bertrand theorem and, formally, can be quantized without violating the G-vH theorem. Extension of the Bertrand theorem to the curved spacetimes results in producing a class of superintegrable systems, including the Kepler-Coulomb -type [20], also quantizable [106], [107]. There are two independent ways to develop quantum mechanics without violating G-vH. The first one is using the Morse theory for development of the sophisticated asymptotic (semi-classical) methods of analysis of quantum mechanical problems [110]. The latest monograph [111] aimed at physically educated readers presents a detailed panoramic view of successful applications of semiclassical methods to all branches of quantum mechanics.

The idea of the second method belongs to Eisenhart. Although his results are summarized in section 4, because of their use in Perlick’s work [23], we need to add additional details at this point. These are coming from the recognition of the fact that the addition of time transformations into symplectomorphisms of classical mechanics makes it effectively general-relativistic [105]. Indeed, suppose that the Lagrangian \mathcal{L} of a dynamical system is given by

$$\mathcal{L} = T(p, q) - V(q), T(p, q) = \frac{1}{2}g_{ij}(q)v^i v^j. \quad (5.1)$$

The configurational space Q of this system is a Riemannian manifold whose metric $ds^2 = g_{ij}(q)dq^i dq^j$ is determined by the kinetic energy term in Eq.(5.1). The Hamiltonian $\mathcal{H}(q, p)$

³²E.g.read the Remark 4.5.

³³Definition 4.6 and Remark 4.7 .

is obtainable in a usual way as

$$\mathcal{H}(q, p) = \tilde{T}(p, q) + V(q), \tilde{T}(p, q) = \frac{1}{2}g^{ij}(q)p_i p_j, p_i = g_{ij}(q)v^j, g^{ij}g_{jk} = \delta_k^i. \quad (5.2)$$

By extending the configurational space Q to $\tilde{Q} = R \times Q$ and adding a new momentum, say, p_z leads to the replacement of $\mathcal{H}(q, p)$ by

$$\tilde{\mathcal{H}}(q, p) = \frac{1}{2}g^{ij}(q)p_i p_j + \frac{1}{2}V(q)p_z^2. \quad (5.3)$$

Accordingly, the metric ds^2 changes to

$$d\sigma^2 = g_{ij}(q)dq^i dq^j + \frac{dz^2}{V(q)}. \quad (5.4a)$$

If such an extension is caused by changes of time variable making it dynamical³⁴, we obtain the metric $d\sigma^2$ used in Perlick's paper [23], e.g.

$$d\sigma^2 = g_{ij}(q)dq^i dq^j - \frac{dt^2}{V(q)}. \quad (5.4b)$$

In section 4.3 we noticed that the Eisenhart lift of a dynamical system involves the condition $\tilde{\mathcal{H}}(q, p) = 0$ essential for recovery of the original dynamics. Such a condition leads to Eq.(4.3) associated with the equation for null geodesics. Eq.(4.3) is conformally invariant³⁵ Hamilton-Jacobi (H-J) equation. In spite of its simple look, the additive separation of variables³⁶ is highly nontrivial. The nontriviality of separation problem in classical H-J equation leads to the nontriviality of quantization problem associated with the conformal Laplacian, Eq.(3.12). References [69], [90] illustrate the essence of this problem. Results of these references should be considered as alternative to the G-vH theorem³⁷. In addition to the references just mentioned, we recommend to our readers excellent lectures by Benenti [112] as well as the thesis [113] by Bruce. Both are containing many practical examples. From Benenti lectures as well as from [90] and Appendix F it follows that analysis of the H-J equation can be done the same way even if $\tilde{\mathcal{H}}(q, p) = E \neq 0$, e.g. for Eq.(B.16). With such a background we are ready to demonstrate that, in accord with expectations of D-O [7], Eq.(3.19) indeed belongs to the set of equations describing quantum Bertrand systems in the sense of [106], [107].

We begin by rewriting Eq.(3,19) in the equivalent form

$$\left(\frac{r}{a}\right)^2 \left[(r/a)^{-\gamma} + (r/a)^{\gamma}\right]^2 \left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} + \frac{\partial^2}{\partial z^2}\right)\psi + n_0^2\psi = 0 \quad (5.5a)$$

³⁴Section 3 and Appendix B.

³⁵Theorem 3.1.

³⁶Eq.(4.4).

³⁷Perhaps, the Bertrand-Darboux theorem should be linked with the G-vH theorem in deciding which dynamical system is admitting quantization.

For $\gamma = 1$, that is for hydrogen atom, this equation coincides with Eq.(3.35) as required. Accounting for scaling arguments presented in Eq.(2.14), classical mechanics version of this equation coincides with Eq.(2.29). For arbitrary γ classical version of Eq.(5.5) reads ($\hbar = 1, m = 1$)

$$\left(\frac{r}{a}\right)^2 [(r/a)^{-\gamma} + (r/a)^\gamma]^2 |\mathbf{p}|^2 = n_0^2 \quad (5.5b)$$

or, in the case if $\gamma = 1/2$, we get

$$\left(\frac{r}{a}\right)^2 [1 + (r/a)]^2 |\mathbf{p}|^2 = n_0^2. \quad (5.5c)$$

In such a form (without applications to atomic physics) this result reads as Eq.(2.119), page 49, of the PhD thesis [108] on quantum Bertrand spaces. In Eq.(2.119) we have to put $K = -1$, $\beta = \gamma$, and $\frac{\mathbf{P}^2}{4} = |\mathbf{p}|^2$. It remains to demonstrate how this result emerges from Perlick's results [23].

We begin with some comments on two types of metric obtained in [23]. We describe only type I metric, that is the Kepler-Coulomb (KC) type. The type II metric, the oscillator-like, is obtainable from type I via Stäckel transform as demonstrated in [106]. The type I metric is described as

$$d\sigma_I^2 = \frac{dr^2}{\beta^2(1 + Kr^2)} + r^2(d\theta^2 + \sin^2\theta d\varphi^2) - \frac{dt^2}{G + \sqrt{r^{-2} + K}} \quad (5.6)$$

Here G and K are some constants while β is *rational* number. By comparing Eq.s (5.3), (5.4) and letting $G = K = 0$ and $\beta = 1$ we recover the KC Hamiltonian

$$\tilde{\mathcal{H}}_{CK} = p_r^2 + \frac{L^2}{r^2} - \frac{1}{r}. \quad (5.7)$$

In general, the associated Hamiltonian, Eq.(5.2), reads³⁸

$$\mathcal{H}_I(q, p) = \frac{\beta^2}{2}(1 + Kr^2)p_r^2 + \frac{L^2}{r^2} - \sqrt{r^{-2} + K} + G. \quad (5.8)$$

The task now is to relate Eq.(5.5b) to Eq.(5.8). This requires several steps. The Perlick metric, Eq.(5.4b), defines the kinetic energy realized on null geodesics of 3+1 dimensional space-time of Lorentzian signature. Alternatively, it can be used for description of the motion in 3 dimensional spherically symmetric space with the metric

$$d\Sigma_I^2 = \frac{dr^2}{\beta^2(1 + Kr^2)} + r^2(d\theta^2 + \sin^2\theta d\varphi^2) \quad (5.9)$$

describing the classical motion on 3 manifold with metric $d\Sigma_I^2$ in the presence of spherically symmetric potential $V(r)$. To move forward, following [114] we need to look at several alternative ways of writing $d\Sigma_I^2$. Specifically,

$$d\Sigma_I^2 = g(r)dr^2 + r^2d\Omega^2 = F^2(\rho)(d\rho^2 + d\Omega^2) = f(|\mathbf{q}|)d\mathbf{q}^2 = f^2(\tilde{r})(d\tilde{r}^2 + \tilde{r}^2d\Omega^2) \quad (5.10)$$

³⁸Since G is arbitrary constant thus far, we are free to change its sign.

Here $d\Omega^2 = d\theta^2 + \sin^2\theta d\varphi^2$, $d\mathbf{q}^2 = dq_1^2 + dq_2^2 + dq_3^2$, $q_1 = \tilde{r} \cos\theta$, $q_2 = \tilde{r} \sin\theta \cos\varphi$, $q_3 = \tilde{r} \sin\theta \sin\varphi$. $|\mathbf{q}| = \tilde{r}$. $\rho = \ln \tilde{r}$. Use of the metric written in the form $f(|\mathbf{r}|)d\mathbf{r}^2$ immediately brings us to Eq.(3.35). But we still can do more since our task is to relate Eq.(5.5b) to Eq.(5.8). Therefore, we have to take into account that $f(|\mathbf{r}|) = f(\tilde{r})$. To determine \tilde{r} and $f(\tilde{r})$ we have to take into account that : a) $F(\rho) = \tilde{r}f(\tilde{r})$, b) $\frac{d\rho}{dr} = r^{-1}g(r)$, c) $r = F(\rho)$. From Eq.(5.10) and just defined results we obtain:

$$a) \frac{r}{\tilde{r}} = f(\tilde{r}); \text{ b) } g(r) = \frac{1}{\beta} \frac{1}{\sqrt{1 + Kr^2}}; c) \frac{d\rho}{dr} = \frac{1}{\beta r \sqrt{1 + Kr^2}} \quad (5.11)$$

Integrating Eq.(5.11c)) yields

$$\rho = \frac{1}{\beta} \ln \left(\frac{r}{1 + \sqrt{1 + Kr^2}} \right). \quad (5.12)$$

Since $\rho = \ln \tilde{r}$ we obtain as well

$$\tilde{r}^\beta = \frac{r}{1 + \sqrt{1 + Kr^2}}. \quad (5.13)$$

Since according to Eq.(5.11a)) $\frac{r}{\tilde{r}} = f(\tilde{r})$, by resolving Eq. (5.13) with respect to r we obtain:

$$\left(\frac{r}{\tilde{r}} \right)^2 = f^2(\tilde{r}) = \frac{4}{\tilde{r}^2} \left(\frac{1}{\tilde{r}^{-\beta} - Kr^\beta} \right)^2. \quad (5.14)$$

Taking into account Eq.s (5.1)-(5.4), (5.6)-(5.10) (and omitting tildas) we obtain the (null) Hamiltonian

$$r^2(r^{-\beta} - Kr^\beta)^2 |\mathbf{p}|^2 + \alpha = 0 \quad (5.15)$$

Here K and α are some constants. They are fixed by comparing Eq.s (5.15) and (5.5c). This is achieved by selecting respectively $\beta = 1/2$, $K = -a^{-1}$ and $\alpha = -n_0^2$. Thus, we just established the desired correspondence between the D-O Eq.(3.19) and the generalized Bertrand problem.

By looking at Eq.s (5.6), (5.8) and (5.10) it remains to demonstrate that presence of the potential $-\sqrt{r^{-2} + K} + G$ in Perlick's Hamiltonian, Eq.(5.8), is harmless for the correspondence we just established. For this purpose we need to take into account that $\sqrt{r^{-2} + K} = \frac{1}{2}(\tilde{r}^{-\beta} + K\tilde{r}^\beta)$. This result is obtained on page 8 of [114]. In terms of these variables the Hamiltonian, Eq.(5.8), acquires the following form

$$\mathcal{H}_I(r, p) = \frac{1}{4}r^2(r^{-\beta} - Kr^\beta)^2 |\mathbf{p}|^2 - \frac{1}{2}(r^{-\beta} + Kr^\beta) + G. \quad (5.16)$$

Using the conformal Stäckel transform described in the Appendix F the Hamiltonian $\mathcal{H}_{II}(r, p)$ is obtainable from $\mathcal{H}_I(r, p)$ as follows. Relabel $\mathcal{H}_I(r, p)$ as $\mathcal{H}_{UI}(r, p)$, where $U = -\frac{1}{2}(r^{-\beta} + Kr^\beta) + G$. Let

$$\mathcal{H}(r, p) = r^2(r^{-\beta} - Kr^\beta)^2 |\mathbf{p}|^2 + \alpha. \quad (5.17a)$$

The Hamiltonian $\mathcal{H}_{\text{II}}(r, p)$ is defined now as

$$\mathcal{H}_{\text{II}}(r, p) = \frac{\mathcal{H}(r, p)}{U} = \frac{1}{2} \frac{r^2(r^{-\beta} - Kr^\beta)^2 |\mathbf{p}|^2}{r^{-\beta} + Kr^\beta - 2G} + \frac{2\alpha}{r^{-\beta} + Kr^\beta - 2G}. \quad (5.17b)$$

By introducing new variables r' and p' so that $r = \mathcal{F}(r')$, $p = F(p')$ with known³⁹ functions \mathcal{F} and F converts $\mathcal{H}_{\text{II}}(r, p)$ into the standard form. It is obtainable with help of Perlick's oscillator-type metric, Eq.(13) of [23], using Eq.s(5.3),(5.4). Thus, by using results of section 4, we just established the correspondence between the D-O Hamiltonian $\mathcal{H}(r, p)$ and that obtained with help of type II Bertrand metric. Since in the conformally flat spaces of constant curvature there is a Coulomb-oscillator duality [115], this duality does exist between the Bertrand spaces of types I and II. This result is supported by the following two theorems

Theorem 5.1. ⁴⁰ *Every nondegenerate second-order quantum superintegrable system on a 3d conformally flat space is Stäckel equivalent to a superintegrable system on constant curvature space*

Theorem 5.2. ⁴¹ *Every superintegrable system with nondegenerate potential on 3d conformally flat space is Stäckel equivalent to a superintegrable system on either 3d flat space or on S^3*

Remark 5.3. Theorem 5.2. provides justification of the result, Eq.(4.53), obtained in previous section.

For reader's convenience we notice that Perlick's spacetimes are, in fact, warped products. This means the following. Consider a static (spherically symmetric in Perlick's case) space-time $(\mathcal{M}, d\sigma^2)$ made of the product of spherically symmetric 3-manifold \mathcal{M} (with metric dg^2) and the line (time) thus making the Lorentzian warped product. The warping function $V(q)$ must be strictly positive. The projection of each timelike geodesic on a constant time leaf $M_0 = \mathcal{M} \times \{t_0\}$ is a trajectory of the dynamical system with Hamiltonian, Eq.(5.3), with $p_z^2 = 1$. Thus the trajectory in Perlick's spacetime $(\mathcal{M}, d\sigma^2)$ is just a projection of the timelike geodesic onto M_0 . For more details on warping products one should consult [118], [119].

In section 1 we mentioned work by Little [19]. In it he obtained a remarkable result. On S^2 there are only 3 distinct regular homotopy classes for oriented closed curves: a) those without self-intersections, b) those with just one self-intersection and, c) those with 2 self-intersections. Because the Bertrand spaces M_0 are spherically symmetric and, in view Theorems 5.1. and 5.2. and, of Hopf mapping, they are topologically either S^2 or S^3 . Based on these observations we anticipate Little's results to be fully enforced. This is indeed the case. In [25] Kirzhnits (with associates) studied Newtonian dynamic of a particle of unit mass

³⁹E.g. read page 49 of [106].

⁴⁰[116], Theorem 17.

⁴¹[116], Theorem 4.

moving in the presence of the Tietz potential⁴², Eq.(G.9), under standard initial conditions. Results were parametrized in terms of the angular momentum L of the particle, charge Z , and dimensionless distance $x = r/a$ defined with help of Eq.(G.7) of Appendix G. The obtained trajectory is described in polar coordinates with help of the equation

$$x + 1/x = \Delta + 1 + (\Delta - 1) \cos \varphi \quad (5.18a)$$

describing a closed curve with just one self-intersection. The observed period T is described as

$$T = \frac{\pi (\Delta + 1) (3\Delta - 1) a^2}{L}. \quad (5.18b)$$

Here $\Delta = \frac{Za}{L^2} - 1$. Exactly the same type of once self-intersecting closed (periodic) motion was obtained by Wheeler [14],[16] and his assistant Powers [15]. At the time of these studies the concept of Bertrand spacetime was nonexistent. Only in 2017 a systematic study of dynamical trajectories of type I (Kepler-Coulomb) Perlick systems with Hamiltonian $\mathcal{H}_1(q, p)$, Eq.(5.8), was performed by Kuru and associates [18]. In complete agreement with previously described results, for $\beta = 1/2$ the closed trajectories were obtained with either none or one self-intersection. These are displayed respectively in Fig.5 and Fig.7 on page 3362.

We conclude this section with some geometrical and topological remarks which would not be required should results of D-O papers be free of inaccuracies to be described momentarily. In atomic physics it is common to write $n = n_r + l + 1$. By referring to Appendix G and referring to Eq.(G.5) we recognize that it holds only if $n_r = 0$. From standard textbooks on quantum mechanics this result is looking suspicious. However, the results of section 4 indicate that, in fact, to use $n_r \neq 0$ is suspicious if calculations are made on S^2 and, via Hopf mapping on S^3 .

Below we explain why this is so using topological arguments. We begin by recalling the 2-dimensional results of section 4. Use of the conformal mapping permits us to solve the Schrödinger equation on \mathbf{R}^2 , to lift it to S^2 and, then to lift the obtained result to S^3 using Hopf mapping. In 2 dimensions the solution of the Kepler-Coulomb quantum problem is reduced to the solution of the standard rigid rotator equation as we have just demonstrated. By doing so known results were reproduced. Thus, we demonstrated that there is no room for n_r on S^2 in the 2d Kepler-Coulomb problem. Surely, it is possible to still use n_r on \mathbf{R}^2 as it was done by Makowski [53]. Since use of \mathbf{R}^2 (or S^2) allows us to perform the conformal transformations legitimately, it comes as no surprise that the result Eq.(4.49) for the deformed wave function in which n_r was used is not correct. The corrected result, Eq.(4.53), is analytical expression of the Madelung rule⁴³. In Appendix G this fact allows us to use in Eq.(G.5) $n = l + 1$ (instead of $n = n_r + l + 1$) as required.

The wave function $Y_{nlm}(\alpha, \theta, \varphi) = \Psi_{nl}(\alpha) Y_{lm}(\theta, \varphi)$ is not reflecting the underlying symmetry (geometry) and topology. This could be the likely cause for the D-O to use in the remnant

⁴²In earlier publication [120] by the same authors more focused exposition is given.

⁴³In Appendix G we demonstrate that this mathematically correct result is consistent with much earlier results by Enrico Fermi obtained in 1928 [121] !

of 3 dimensional calculations, the n_r . By using the above representation for $Y_{nlm}(\alpha, \theta, \varphi)$, the symmetry becomes broken. Its breakage is obscuring the relationship between the results on S^2 and S^3 . Group-theoretically, this issue is best understood following [87] and [123]. The Lie algebra $so(4)$ (working⁴⁴ on S^3) is a direct product of two $so(3)$ Lie algebras⁴⁵, each describing the rigid rotators with standard spectrum $j_i(j_i + 1)$, $i = 1 \div 2$. The Plücker embedding condition described in Appendix E leads to the requirement: $j_1 = j_2$ [122]. Thus, the Lie algebra $so(4) \simeq so(3) \oplus so(3)$. This result is achieved with help of the requirement $j_1 = j_2$. Topologically, S^3 is obtainable in two ways: a) either by gluing 2 solid tori to each other (Heegard splitting) or, b) by gluing two 3- balls pointwise to each other [123]. The last construction can be easily explained. Begin with S^2 . It can be made out of 2 discs glued to each other along the disc boundaries. Consider a stereographic projection- from the 2 sphere whose equator lies in the plane. Then, the projection can be made from the Northern pole or from the Southern pole. Thus, we obtain the correspondence between the half sphere -the disc- and the plane. It is possible then to map a plane onto the half plane and gluing of 2 half spheres -to make an S^2 - can be identified with gluing of 2 half planes along their common boundary. By extending these ideas to S^3 we have to glue together 2 half $(\mathbf{R}^3)'$ s along their common plane. This is exact topological equivalent of the group-theoretic condition: $j_1 = j_2$!

It is possible to still extend just obtained results. Following [88] the wavefunction $Y_{nlm}(\alpha, \theta, \varphi)$ can be replaced by its manifestly symmetric analog. For this, use of the manifestly symmetric 4 dimensional spherical coordinates is essential. These are given by

$$x_1 = R \cos \xi \cos \varphi, x_2 = R \cos \xi \sin \varphi, x_3 = R \sin \xi \cos \eta, x_4 = R \sin \xi \sin \eta \quad (5.19)$$

to be contrasted with those, given by Eq.(4.7). In terms of such coordinate parametrization the eigenvalue Eq.(4.31a) acquires the following look

$$\hat{\mathcal{L}}^2 Z_{nfg}(\xi, \varphi, \eta) = I_{nfg} Z_{nfg}(\xi, \varphi, \eta) \quad (5.20)$$

Here [87]

$$Z_{nfg}(\xi, \varphi, \eta) = (2\pi)^{-1} (2n)^{\frac{1}{2}} (-1)^{F+g} d_{gf}^F(2\xi) \exp\{i\varphi(f+g) + i\eta(f-g)\} \quad (5.21)$$

and, as before, $2F + 1 = n$. The detailed description of Wigner's coefficients $d_{gf}^F(2\xi)$ is given in [124].

Note added in proof: Existence of Bertrand spaces favors quantization of dynamics at all scales.

It also helps to understand and to describe the exceptions to Madelung's rule

Initially, the Bertrand spaces developed by Perlick [23] were **not** meant for description of microscopic systems. Nevertheless the results of this paper, e.g. sections 2 and 3, indicate

⁴⁴Surely, the $su(2)$ is working of S^3 . However, the $so(4)$ of is capable of working on S^3 very much like the rotation group $so(3)$ on S^2 .

⁴⁵E.g.read the Appendix E

that, by definition, superintegrable systems of physical interest do possess closed orbits irrespective to the actual size of these systems. If this is so, the immediate question emerges: Since quantum mechanically closure of orbits is linked with quantization, can one expect to encounter the macroscopic Bertrand dynamical systems which are quantum? Although we had initiated study of this topic previously [105], the results of this paper prompt us to make additional remarks.

First, the assumption that all planets of our solar systems lie in the same (Sun's equatorial) plane and move in the same direction coinciding with the direction of rotation of the Sun around its axis was used essentially by Henri Poincaré' in his "Les Methodes Nouvelles de la Mechanique Celeste" written between 1892 and 1899. Poincaré's assumptions are consistent with popular hypothesis⁴⁶ that our solar system was made out of pancake-like rotating cloud.

Second, this plausible hypothesis was confronted with the discovery (in 1898-1899)⁴⁷ of Phoebe-the ninth moon of Saturn rotating in the direction opposite to all other satellites of Saturn. Recall that the Jupiter, Saturn, Uranus-all have their own satellite systems analogous to the planetary system of our solar system. Therefore, apparently, the pancake models should be applicable to the satellite systems of heavy planets as well. But they are not! Since Phoebe's discovery many other Phoebe-like satellites were found. The satellites rotating in the "normal" direction had received the name "*prograde*" while those in the opposite direction had been called "*retrograde*". By 2018 in our solar system 125 retrograde satellites were found⁴⁸. Interestingly, the retrograde motion takes place in the same planes as prograde motion. Newton's mechanics does not imply whatsoever such a motion for retrogrades. Not only this fact breaks down the pancake model but it and other facts of satellite motion in solar system discussed in [105] point in favor of quantum mechanical Bertrand-type dynamical model of solar system dynamics.

Third, the description of multielectron quantum Bertrand models of multielectron atoms developed in this paper involves uses of conformal invariance. This fact makes the obtained results scale insensitive. However, general relativity together with classical mechanical Le Verrier's calculations⁴⁹ makes the assumption about the quantum nature of solar system dynamics very problematic. The observed planetary orbits are not closed, they precess. In atomic systems this fact was discovered by Arnold Sommerfeld in 1916 [30] immediately after seminal 1915 work by Einstein on general relativity. Unlike the case of celestial mechanics, Sommerfeld came up with ingenious way to by pass the issue of orbit nonclosure resulting in his result for the fine structure of hydrogen atom spectrum rediscovered by Dirac in 1928.

Forth, these facts are not detrimental, however, to the assumption that the Bertrand model spaces are capable of describing reality at all scales. This is so because of the following observations. In 1966 Greenberg⁵⁰ noticed that: a) the accidental degeneracy is equivalent to superintegrability, b) by applying perturbations the superintegrability can be removed. As an example Greenberg looked at the relativistic effects (e.g. velocity-dependent mass)

⁴⁶With lots of numerical simulations made in its support through years

⁴⁷[https://en.wikipedia.org/wiki/Phoebe_\(moon\)](https://en.wikipedia.org/wiki/Phoebe_(moon))

⁴⁸E.g. read https://en.wikipedia.org/wiki/Irregular_moon

⁴⁹https://en.wikipedia.org/wiki/Urbain_Le_Verrier

⁵⁰Accidental degeneracy *Am.J.Phys.*34 (1966) 1101-1109

in the Kepler-Coulomb problem. The same problem as studied by Sommerfeld! This effect is making Runge-Lenz vector nonconserved [30]. By considering the relativistic effects as small perturbations, the celebrated Einstein's general relativistic result for the precession of a planetary orbit was obtained. Much later, Riglioni (in the Appendix A of his PhD thesis [106]) had reobtained the same result as perturbation of the Bertrand-type motion caused by (incomplete) mapping of the Bertrand-like metric onto the Schwarzschild-like. These results surely require further study since, for instance, the exceptions to the Madelung rule⁵¹ are consequences of relativistic -type perturbations of superintegrable systems.

Apparently, these physics-inspired results are unknown to mathematicians thus far. At the same time, physics practitioners are not aware of the very important recent developments in mathematics related to the theory of perturbations of superintegrable systems. Extremely readable introduction into this topic can be found in papers by Francesco Fasso⁵² and Anthony Blaom⁵³. From both sources it follows that for the superintegrable systems theory of perturbations should be developed using Nekhoroshev's results rather than the KAM results⁵⁴. General principles of Nekhoroshev's theory are recently illustrated on the benchmark example of hydrogen atom in crossed electric and magnetic fields⁵⁵. This example was also used by Gutzwiller [12] as well as by many other authors without references to Nekhoroshev's results. Nevertheless, among these publications we notice the paper by Flöthman and Welge⁵⁶ in which the connection (isomorphism) between this quantum problem and the classical 3-body problem is developed in detail. These authors notice that such quantum problem is the benchmark problem for study of classical and quantum chaos. Rigorous study of manifestations of adiabatic classical and quantum chaos in perturbed superintegrable systems had just began⁵⁷.

Acknowledgements

Both authors are indebted to the members of the Department of Mathematics of the Novosibirsk State University for supplying us with unpublished lecture notes by L.Ovsiannikov. Work by Louis Kauffman's was supported in part by the Laboratory of Topology and Dynamics, Novosibirsk State University (contract no. 14.Y26.31.0025 with the Ministry of Education and Science of the Russian Federation).

⁵¹E.g. read https://en.wikipedia.org/wiki/Aufbau_principle

⁵²Notes on Finite dimensional integrable Hamiltonian systems (Padova: University of Padova, Department of Mathematics) 1999.

⁵³A geometric setting for Hamiltonian perturbational theory, AMS Memoir 153, 2001.

⁵⁴The book by Gutzwiller [12] and, with it all other books on classical and quantum chaos written in physics literature, does not contain any mention of Nekhoroshev's results.

⁵⁵Fasso F., Fontanari D. and Sadovskii D. , An application of Nekhoroshev theory to the study of the perturbed hydrogen atom *Math.Phys Anal Geom* 18 (1) (2015) Art.30

⁵⁶Crossed-field hydrogen atom and the three-body Sun-Earth-Moon problem, *Phys.Rev.A* 54 (1996) 1884-1888

⁵⁷Fontanari D. Quantum manifestations of the adiabatic chaos of perturbed superintegrable Hamiltonian systems (Padova: University of Padova, Department of Mathematics), 2016.

Appendix A. Contributions of Constantin Caratheodory to the design of absolute optical instruments

In this appendix we shall comment on the appropriate places in remarkable book by Caratheodory "Geometrical Optics" [22]. In it, he begins with Fermat principle, Eq.(2.1), which he rewrites as

$$\int_{\gamma'} L'(t', x'_i \frac{dx'_i}{dt'}) dt' = \int_{\gamma} L(t, x_i \frac{dx_i}{dt}) dt + \int_{\gamma} d\Psi, i = 1 \div 3. \quad (\text{A.1})$$

This equation is interpreted as follows. The difference between the optical lengths of two curve segments γ and γ' that correspond to each other by means of the stigmatic map is equal to the difference between the values of $\Psi(t, x_i)$ at the endpoints of the curve γ . This difference is then independent of the shape of the curves γ and γ' and depends only upon the position of their endpoints. Under such circumstances Eq.(A.1) implies that the relation

$$L'(t', x'_i \frac{dx'_i}{dt'}) dt' = L(t, x_i \frac{dx_i}{dt}) dt + \left(\frac{\partial}{\partial t} \Psi \right) dt + \left(\frac{\partial}{\partial x_i} \Psi \right) dx_i \quad (\text{A.2})$$

must be fulfilled identically when one replaces the old variables t, x_i by the new ones

$$t' = t'(t, x_i), x'_i = x'_i(t, x_i) \quad (\text{A.3})$$

so that,

$$dt' = \frac{\partial t'}{\partial t} dt + \frac{\partial t'}{\partial x_i} dx_i \text{ and } dx'_i = \frac{\partial x'_i}{\partial t} dt + \frac{\partial x'_i}{\partial x_j} dx_j. \quad (\text{A.4})$$

Caratheodory proves then that $\frac{\partial}{\partial t} \Psi \equiv \frac{\partial}{\partial x_i} \Psi \equiv 0$ causing Eq.(A.1) to be replaced by

$$\int_{\gamma'} L'(t', x'_i \frac{dx'_i}{dt'}) dt' = \int_{\gamma} L(t, x_i \frac{dx_i}{dt}) dt. \quad (\text{A.5})$$

We are not reproducing the proof since Eq.(A.5) was treated by Dirac whose arguments we are presenting in section 2 of the main text. Here we only notice that the map-from the object space to the image space, given by Eq.s(A.3), must be conformal. This fact causes us to bring some facts from our earlier work [3] superimposed with results of section 2. Caratheodory notices that "From a celebrated theorem of Liouville⁵⁸ in contrast to the planar conformal maps which depend upon infinitely many constants there is only restricted class of conformal maps of three-dimensional space. It can always be represented as a sequence of transformations through reciprocal radii of at most five spheres⁵⁹. *It follows then that the circles and lines in object space are transformed into curves in image space that will always*

⁵⁸Discuseed in our work [3], page 50, Theorem B.8.

⁵⁹Caratheodory is having in mind the Möbius and Lie sphere-type transformations discussed in our work [3] on pages 38-40 and the pentaspherical transformations discussed in section 4.

be either the circles or lines. Maxwell treated the simplest case of such a ray map (when one ignores a reflection) on occasion”⁶⁰. Furthermore, ” Maxwell imagined the entire space to be occupied by the medium whose index of refraction is described by Eq.(2.5) of section 2. The light rays themselves are then either circular or rectilinear. One confirms this result the most simply when one recalls the equation⁶¹

$$d\sigma = \frac{2ab}{b^2 + r^2} \sqrt{dx^2 + dy^2 + dz^2}, \quad r^2 = x^2 + y^2 + z^2. \quad (\text{A.6})$$

The differential $d\sigma$, which defines the optical length of a line element inside the interior of the Maxwell fish-eye, can be also interpreted as a line element in the three-dimensional boundary of a four-dimensional sphere of radius a that is projected stereographically onto a space of x, y, z that should be found at a distance b from the center of the projection.” Explicitly,

$$\xi = \frac{2abx}{b^2 + r^2}, \eta = \frac{2aby}{b^2 + r^2}, \zeta = \frac{2abz}{b^2 + r^2}, \tau = a \frac{b^2 - r^2}{b^2 + r^2} \quad (\text{A.7})$$

so that $\xi^2 + \eta^2 + \zeta^2 + \tau^2 = a^2$ and, reciprocally,

$$x = \frac{b\xi}{a + \tau}, y = \frac{b\eta}{a + \tau}, z = \frac{b\zeta}{a + \tau}, r^2 = \frac{b^2(a - \tau)}{a + \tau}. \quad (\text{A.8})$$

By differentiating either Eq.(A.7) or (A.8) we reobtain Eq.s(2.4a),(2.4b) of section 2. Let now x, y, z and, respectively, x', y', z' denote the stereographic projections of two opposite points ξ, η, ζ (respectively, $-\xi, -\eta, -\zeta$) on the four-sphere, then we obtain

$$x' = -\frac{b^2 x}{r^2}, y' = -\frac{b^2 y}{r^2}, z' = -\frac{b^2 z}{r^2} \quad (\text{A.9})$$

implying

$$r' r = b^2 \quad (\text{A.10})$$

easily recognizable as the main result defining the Möbius geometry. This is achieved when $b^2 = r^2 = x^2 + y^2 + z^2$ so that $x' = -x, y' = -y$ and $z' = -z$ and, finally, $\tau = 0$. With these results in our hands we are in the position to interpret them using either the language of Lie sphere geometry or, which is equivalent, of twistors⁶² By analogy with Eq.s (6.30a,b) of [3], page 30, and, in accord with Caratheodory, we obtain the great circles on the three sphere as intersection of two hyperplanes

$$A_k \xi + B_k \eta + C_k \zeta + b D_k \tau = 0, k = 1, 2. \quad (\text{A.11})$$

Their intersection, when projected into x, y, z space is described by

$$D_k(x^2 + y^2 + z^2 - b^2) - 2A_k x - 2B_k y - 2C_k z = 0, k = 1, 2. \quad (\text{A.12})$$

⁶⁰Here the emphasis is ours

⁶¹Eq.s (2.4a) and (2.4b) of the main text

⁶²E.g. read page 41 of [3].

These equations coincide exactly with Eq.s (7.16) of [3], page 41, as required. Recall, that these equations had been used in [3] to establish twistor- Lie sphere geometry correspondence. In the present case, following Caratheodory, the interpretation is as follows. Eq.s (A.12) are images of the great circles on the S^3 . Due to conformal property given by Eq.(A.10) the great circles on S^3 are in one-to one correspondence with great circles on S^2 ($S^2 : b^2 = x^2 + y^2 + z^2$) so that if for S^3 they intersect in diametrically opposite points, the same is happening in S^2 . Therefore, Eq.s.(A.1) and (A.5) can be interpreted as follows. The requirement $d\Psi = 0$ is coming from the equality of optical lengths of the corresponding curves. That is the spherical length of two diametrically opposite curve segments is the same for both curves. Having in mind additional applications mentioned in the main text, it is useful to think about just described results as follows. Suppose that there is a manifold M (Blashke manifold) with the property that for each point $P \in M$ all geodesics passing through P will meet again at the conjugated point P then, such a medium can be used as perfect optical instrument. Arthur Besse published a book entitled " Manifolds all of whose geodesics are closed" [125]. Blashke conjectured that the manifolds with such a property are S^n . The current status of this conjecture is described in [126]. For the latest optical applications of geometric and topological ideas of Besse's book, e.g. the geodesic lenses and similar devices, please, consult [127] and [128]. Results of Besse's book are further discussed in section 5 from the point of view of Bertrand spaces. Interesting enough, another construction of Blashke (e.g. Blashke products) had been used in studies of gravitational lensing [129]. Gravitational lensing can be connected with results of this Appendix as an elaboration on the discussion in section 2 of the main text. The topic of photon sphere known from general relativity falls also in the same category as that of Maxwell's fish -eye [130], [131].

Appendix B. Many uses of time changes in classical mechanics

Newtonian mechanics is invariant under Galilei transformations. Such transformations assume that time is nonchangeable parameter. However, this assumption breaks down in Newtonian mechanics as soon as mathematically rigorous treatment of collisions becomes of interest. Occurring mechanical singularities typically are being treated by the appropriate time changes. Short but excellent introduction to this topic is given in Ref.[132]. The need for variable time also emerges in theory of contact transformations. For a quick introduction, please, consult [94]. In mechanics the contact 1-form dS is given by

$$dS = \sum_{i=1}^n p_i dq_i - H(\{q_j\}, \{p_j\}) dt \quad (\text{B.1})$$

where $\{q_j\} = [q_1, \dots, q_n]$, etc. For the conservative system $E = H(\{q_j\}, \{p_j\})$. At least in that case it is possible to treat time t and energy E as canonically conjugate variables. So, we can make the following identifications: $t = q_{n+1}, -E = -H = p_{n+1}$. With this identification the action S can be written in the Maupertuis form

$$S = \int \sum_{i=1}^{n+1} p_i dq_i. \quad (\text{B.2})$$

We would like to connect this result with that discussed in the Appendix A. For this purpose consider an equality⁶³

$$dS' - p'_i dx'_i = \rho(S, \{x_j\}, \{p_j\})(dS - p_i dx_i) \quad (\text{B.3a})$$

where $\rho(S, \{x_j\}, \{p_j\}) \neq 0$ and $S' = \hat{S}(S, \{x_j\}, \{p_j\})$, $x'_i = x_i(S, \{x_j\}, \{p_j\})$, $p'_i = p_i(S, \{x_j\}, \{p_j\})$.

The equality, Eq.(B.3a), is defining a contactomorphism [94],[133],[134]. In contact geometry one always begins with the 1-form $\alpha = dS - p_i dx_i$ and studies groups of contactomorphisms. Clearly, the symplectic 2-form is readily obtainable via prescription: $\omega = d\alpha = dx_i \wedge dp_i$. These results create an impression that the contact and symplectic geometries are different in general since it is not a priori obvious that the contactomorphic transformations are isomorphic to symplectic. To shed some light on this issue, we multiply both sides of Eq.(B.3a) by new variable $\lambda \neq 0$. Then, Eq.(B.3a) acquires the following form

$$\frac{\lambda}{\rho} (dS' - p'_i dx'_i) = \lambda(dS - p_i dx_i). \quad (\text{B.3b})$$

By introducing new notations:

$$\begin{aligned} S &= x_0; \lambda = y_0, y_i = -\lambda p_i; \\ S' &= x'_0; \frac{\lambda}{\rho} = y'_0; y'_i = -\frac{\lambda p_i}{\rho} \end{aligned}$$

it is possible to rewrite Eq.(B.3.b) equivalently as

$$y'_0 dx'_0 + y'_i dx'_i = y_0 dx_0 + y_i dx_i. \quad (\text{B.4})$$

Using this result along with Eq.s(B.1),(B.2) it should be obvious that Eq.(B.4) is exactly the same as Eq.(A.2) (with $\Psi = 0$) implying Eq.(A.5) and, hence, the Dirac theory of constraints, and time changes discussed in section 2. We would like to do more in this appendix.

First, we want to emphasize the differences between the contact and symplectic geometries. The major difference between these two geometries takes place when they are confronted with the description of dynamical systems subject to the constraint $H = E$. At the quantum level this topic is discussed in the Appendix F.

As soon as we are having such a constraint, the choice between two geometries is broken in favor of contact geometry. The equation $H(\{q_j\}, \{p_j\}) = E$ defines the hypersurface Σ of dimension $2n - 1$ in $2n$ dimensional symplectic phase space \mathcal{M} . To relate the vector fields on Σ and \mathcal{M} the concept of Liouville vector field Y is essential. Given that the Lie derivative \mathcal{L}_Y is defined via

$$\mathcal{L}_Y = d \circ i_Y + i_Y \circ d, \quad (\text{B.5})$$

where d is the exterior differential and i_Y is the contraction (both applied to differential forms ω), the Liouville vector field Y is defined with help of the equation

$$\mathcal{L}_Y \omega = \omega. \quad (\text{B.6})$$

⁶³Summation over repeated indices is always being assumed

Based on this definition, the vector field Y can be defined, for example, as $Y = p_i \frac{\partial}{\partial p_i}$.

The Liouville vector field Y connects the symplectic and contact geometries with help of the equation

$$\alpha = i_Y \omega. \quad (\text{B.7})$$

At the same time, in symplectic geometry there is a Hamiltonian vector field X_H such that

$$i_{X_H} \omega = -dH \text{ or } \omega(X_H, -) = -dH(-). \quad (\text{B.8a})$$

Here $dH = \frac{\partial H}{\partial x_i} dx_i + \frac{\partial H}{\partial p_i} dp_i$ implying

$$X_H = \frac{\partial H}{\partial p_i} \partial_{x_i} - \frac{\partial H}{\partial x_i} \partial_{p_i} \quad (\text{B.8b})$$

since $\omega = dx_i \wedge dp_i$. With such definition of X_H we obtain:

$$\omega(X_H, X_H) = -dH(X_H) = 0 \quad (\text{B.8c})$$

as required. Consider now what will happen to $\mathcal{L}_{X_H} \omega$. Using Eq.(B.5) we obtain

$$\mathcal{L}_{X_H} \omega = d \circ i_{X_H} \omega + i_{X_H} \circ d\omega = 0 \quad (\text{B.9})$$

since $d\omega = 0$ and, because $i_{X_H} \omega = -dH$, we also have $d \circ i_{X_H} \omega = 0$. Thus, the Lie derivative of the Hamiltonian vector field X_H preserves the symplectic two-form ω . The question emerges: what vector field preserves the contact form α ? Let us denote this field as R_α , so that

$$\mathcal{L}_{R_\alpha} \alpha = d \circ i_{R_\alpha} \alpha + i_{R_\alpha} \circ d\alpha = 0. \quad (\text{B.10})$$

The Reeb vector field R_α is defined as the field that obeys two conditions:

$$i_{R_\alpha} \alpha = \text{const}, \quad i_{R_\alpha} \circ d\alpha = 0. \quad (\text{B.11})$$

Typically in the literature one encounters the condition, $\text{const} = 1$, but it is clear that one can choose as well the condition $\text{const} > 0$. The question of central importance is the following

Is there any connection between X_H and R_α ? and, if there is, how to describe such a connection?

We begin with the observation [94] that the very same manifold \mathcal{M} has both the symplectic and the Riemannian structure. The Riemannian structure is described in terms of the metric tensor 2-form: $g = g_{ij} dx^i \otimes dx^j$. Introduce now the (musical) \flat - operator converting the vector field $X = X^i \frac{\partial}{\partial x^i}$ into 1-form $X^\flat = i_X g = g(X, \cdot) = g_{ij} X^i dx^j$. Introduce also the inverse operator \sharp via $(X^\flat)^\sharp = g^{ij} X_i \frac{\partial}{\partial x^j} = X$. Suppose now that instead of Eq.(B.7) we can write

$$\alpha = i_X g. \quad (\text{B.12})$$

Suppose also that some vector field $X = \tilde{X}$ is Reeb-like, that is, looking at Eq.(B.11) we have to require

$$i_{\tilde{X}}\alpha = i_{\tilde{X}}i_{\tilde{X}}g = 1. \quad (\text{B.13a})$$

Explicitly,

$$i_{\tilde{X}}[g_{ij}\tilde{X}^i dx^j] = g_{ij}\tilde{X}^i \tilde{X}^j = 1. \quad (\text{B.13b})$$

But, since $\alpha = i_{\tilde{X}}g$ we also must have

$$\alpha = i_{\tilde{X}}g = i_Y\omega. \quad (\text{B.14})$$

In view of Eq.s(B5-B7) it should be clear that $d \circ i_Y\omega = \omega$. Accordingly, in view of Eq.(B.14) we also should have $d \circ i_{\tilde{X}}g = d\alpha$. If \tilde{X} is Reeb-like, then we have to have $i_{\tilde{X}} \circ d\alpha = 0$. But

$$i_{\tilde{X}} \circ d\alpha = \omega(\tilde{X}, -) = -dH(-) \quad (\text{B.15})$$

And this result becomes the same as Reeb's 2nd condition, Eq.(B.11), only when $dH \equiv 0$. Fortunately, this is the case. This follows from the observation that Eq.(B.13b) can be reinterpreted as an equation for the Jacobi-Maupertius Hamiltonian \tilde{H} for the fixed energy surface $H - E = 0$, [133]. That is

$$\tilde{H} = \frac{1}{2}\tilde{g}^{ij}(x)p_ip_j = 1, \quad (\text{B.16})$$

where

$$\tilde{g}^{ij}(x) = \frac{g^{ij}(x)}{m[E - U(x)]} \text{ or } \tilde{g}_{ij}(x) = m[E - U(x)]g_{ij}(x).$$

Alternatively, the results can be rewritten in the Lagrangian form as follows

$$\tilde{\mathcal{L}} = \frac{1}{2}\tilde{g}_{ij}(x)\xi^i\xi^j \equiv \frac{1}{2} < \xi^i, \xi^j > = 2 \quad (\text{B.17})$$

where $\xi^i = \frac{dx^i}{d\sigma}$ and $p_i = \tilde{g}_{ij}(x)\xi^j$. The Euler- Lagrange equations obtainable with Lagrangian $\tilde{\mathcal{L}}$ are equations for geodesics [134],[36] relative to the metric $\tilde{g}_{ij}(x)$. Equations for geodesics do not change if we replace $\tilde{\mathcal{L}}$ by $\bar{\mathcal{L}} = \sqrt{\tilde{\mathcal{L}}}$. But this Lagrangian leads to the reparametrization-invariant action and is homogenous of the first degree in ξ^i . That is $\bar{\mathcal{L}}(x, \lambda\xi) = \lambda\bar{\mathcal{L}}(x, \xi)$. Use of Euler's theorem produces then: $\xi^i \frac{\partial \bar{\mathcal{L}}}{\partial \xi^i} = \bar{\mathcal{L}}$. From here we obtain the Hamiltonian

$$H = \xi^i \frac{\partial \bar{\mathcal{L}}}{\partial \xi^i} - \bar{\mathcal{L}} \equiv 0.$$

Thus, the Reeb vector field \tilde{X} *always* describes the (null) geodesics of energy $E = 0$ in Riemannian space with metric $\tilde{g}_{ij}(x)$.

Obtained result is in agreement with that obtained by a very different set of arguments in [134], page 27. Now we are in the position to formulate the following (e.g. read [134], page

25):

Theorem B.1. *If a codimension 1 submanifold \mathcal{M} is both a hypersurface of contact type (that is with $\alpha = i_Y \omega$) and the level set of a Hamiltonian H , then the Reeb flow is reparametrization of the Hamiltonian flow.*

Proof: on one hand we have $i_{R_\alpha} \circ d\alpha|_{\Sigma} = 0$, on another hand, we have $i_{\tilde{X}} \circ d\alpha|_{\Sigma} = \omega(\tilde{X}, -)|_{\Sigma} = -dH(-)|_{\Sigma} = 0$.

Obtained results still requires some improvement. Let us return to the 1-form $\alpha = dS - p_i dx_i$ defined after Eq.(B.3a). A *contact structure* on a manifold M of odd dimension (say, $2n+1$) is a hyperplane field $\zeta = \ker \alpha$ such that $\alpha \wedge (d\alpha)^n \neq 0$. The pair (M, ζ) is called *contact manifold* [94],[134]. Now, notice that $\ker \alpha$ is known in mechanics as a statement: $p_i = \partial S / \partial x_i$. Thus, the action $S = S(\{x_i\})$ is a surface for which $\omega = dx_i \wedge dp_i = 0$. Such surfaces are called *Lagrangian surfaces*. On Lagrangian surfaces the Bohr-Sommerfeld quantization condition reads: $\oint p dx = 0$. The infinitesimal automorphisms of $\zeta = \ker \alpha$ are also described in terms of the vector fields. Suppose that \tilde{X} introduced after Eq.(B.12) is one of these fields. Then, Eq.(B.13b) can be equivalently rewritten as $\alpha(\tilde{X}) = 1$. But we already know from Eq.(B.16) that $\tilde{H} = 1$. Therefore we can write $\alpha(\tilde{X}) = \tilde{H}(\tilde{X})$ so that $\tilde{H}(\tilde{X})$ defines now the *contact Hamiltonian*. Unlike the Liouville vector field for which Eq.(B.6) holds, it makes sense to require

$$\mathcal{L}_{\tilde{X}} \alpha = \mu \alpha \quad (\text{B.18})$$

The physical meaning of μ multiplier is to be determined momentarily. For this purpose we follow [35], page 57. According to this reference for the Hamiltonian H we should discuss the dynamics under constraint: $H = c$. Suppose that initially such a Hamiltonian is used for description of dynamical system described by

$$\frac{dx}{dt} = H_y(x, y), \quad \frac{dy}{dt} = -H_x(x, y). \quad (\text{B.19a})$$

Changing time

$$s = \int_0^t d\tau \lambda(\phi^\tau(x, y)) \quad \text{or} \quad \frac{ds}{d\tau} = \lambda \quad (\text{B.19b})$$

converts the equations of motion into the following form:

$$\frac{dx}{dt} = \lambda^{-1}(x, y) H_y(x, y), \quad \frac{dy}{dt} = -\lambda^{-1}(x, y) H_x(x, y). \quad (\text{B.19c})$$

This causes the Hamiltonian Eq.s(B.19a) to lose their Hamiltonian form. Nevertheless the function $\tilde{H} = \lambda^{-1}(H - c)$ can be taken as new (actually contact) Hamiltonian on hypersurface $H = c$ where

$$d\tilde{H} = \lambda^{-1} dH \quad (\text{B.20})$$

Accordingly, for the vector fields on such hypersurface, $X_{\tilde{H}} = \lambda^{-1} X_H$. Earlier we obtained: $\alpha(\tilde{X}) = \tilde{H}(\tilde{X})$. Therefore, in view of Eq.(B.5), Eq.(B.18) can be rewritten now as

$$d\tilde{H} + i_{\tilde{X}} d\alpha = \mu\alpha. \quad (\text{B.21})$$

This result was obtained with help Eq.s(B.13) and (B.16). Taken the Reeb vector field R_α , this equation becomes: $d\tilde{H}(R_\alpha) = \mu$. By rewriting Eq.(B.21) as

$$d\tilde{H}(R_\alpha)\alpha - d\tilde{H}(\tilde{X}) = i_{\tilde{X}} d\alpha \quad (\text{B.22a})$$

and, since \tilde{X} is Reeb-like, we get $i_{\tilde{X}} d\alpha = 0$ causing us to arrive at the final (equivalent) form of Eq.(B.20), that is $d\tilde{H} = \mu\alpha$

Appendix C. Some auxiliary facts about conformal motions

Although there is a number of good sources on this topic, e.g. [62], [63], for the sake of uninterrupted reading, we present in this appendix some essentials.

Specifically, let x and x' be two different points on the Riemannian manifold \mathcal{M}_d of dimension d . Typically, the manifold \mathcal{M}_d is covered by a system of charts. To avoid complications, we shall only consider x and x' placed in just one chart. Then, the equation

$$g_{ij}(x) = \frac{\partial x'^k}{\partial x^i} \frac{\partial x'^l}{\partial x^j} g_{kl}(x') \quad (\text{C.1})$$

relates the components $g_{ij}(x)$ and $g_{kl}(x')$ of the metric tensor at points x and x' of \mathcal{M}_d . Suppose that for $\varepsilon \ll 1$ we have

$$x'^i = x^i + \varepsilon \xi^i(x), \quad (\text{C.2})$$

then the Killing vector field \mathbf{X} is defined by

$$\mathbf{X} = \xi^i(x) \frac{\partial}{\partial x^i}. \quad (\text{C.3})$$

Substituting of Eq.(C.2) into Eq.(C.1), Taylor expanding and keeping only the 1st order in ε terms results in the Killing Eq.(3.21) of the main text. Explicitly,

$$\nabla_i \xi_j + \nabla_j \xi_i = 0. \quad (\text{C.4})$$

Here $\xi_i = \xi^j g_{ji}$ and the covariant derivative $\nabla_i \xi_j$ is defined as usual by

$$\nabla_i \xi_j = \partial_i \xi_j - \Gamma_{ij}^k \xi_k. \quad (\text{C.5})$$

Familiar isometries are translations and rotations. For d -dimensional space these can be represented by the $d \times d$ matrices decomposable into diagonal d -dimensional matrix and skew symmetric $d \times d$ matrix having $\frac{d^2 - d}{2}$ components. Thus, in the case of isometries there are

$\frac{d^2 - d}{2} + d = \frac{d(d+1)}{2}$ isometry generators. There is a very important theorem by Eisenhart [65], Chr.6, which reads as follows

Theorem C.1. *Isometric group of motion for \mathcal{M}_d can have no more than $\frac{d(d+1)}{2}$ elements. The maximum number is achieved only for spaces of constant scalar curvature.*

For description of these spaces we follow the classical book by Petrov [52]. In chapter 1 of this reference we find the following

Theorem C.2. *In order for \mathcal{M}_d to be a space of constant curvature it is necessary and sufficient that the Riemannian curvature tensor R_{ijkl} can be represented in the form*

$$R_{ijkl} = K(g_{ik}g_{jl} - g_{jk}g_{il}) \quad (\text{C.6})$$

where K is constant.

Eq.(C.6) is satisfied if the metric can be presented in the form

$$ds^2 = \frac{1}{\sigma^2} \sum_{i=1}^d \varepsilon_i (dx^i)^2 \quad (\text{C.7})$$

where $\varepsilon_i = \pm 1$ are in accord with the signature of \mathcal{M}_d and, the conformal factor σ is determined based on Eq.(C.6) and known definition of R_{ijkl} via Christoffel's symbols. This leads to the following set of equations

$$\partial_{ij}\sigma = 0, \sigma[\varepsilon_i \partial_{jj}\sigma + \varepsilon_j \partial_{ii}\sigma] = \varepsilon_i \varepsilon_j [K + \sum_{i=1}^d \varepsilon_i (dx^i)^2], i \neq j. \quad (\text{C.8})$$

The solution of these equations is

$$\sigma = 1 + \frac{K}{4} \sum_{i=1}^d \varepsilon_i (dx^i)^2. \quad (\text{C.9})$$

Corollary C.3. In view of the Definition 3.6. and of Eq.s(C.7), (C.9), all spaces of constant curvature are conformally flat.

Appendix D. Calculation of Christoffel's symbols and related things

For the metric tensor in the form $g^{ij}(x) = \rho(x)\delta^{ij}$ (or $g_{ij}(x) = \frac{1}{\rho(x)}\delta_{ij}$) calculation of Christoffel's symbols proceeds as follows.

$$\begin{aligned}\Gamma_{kl}^i &= \frac{1}{2}\rho\delta^{ij}\left[\delta_{kj}\left(\frac{\partial}{\partial x^l}\frac{1}{\rho}\right) + \delta_{lj}\left(\frac{\partial}{\partial x^k}\frac{1}{\rho}\right) - \delta_{kl}\left(\frac{\partial}{\partial x^j}\frac{1}{\rho}\right)\right] \\ &= \frac{1}{2\rho}\left[\delta_k^i\left(\frac{\partial}{\partial x^l}\rho\right) + \delta_l^i\left(\frac{\partial}{\partial x^k}\rho\right) - \left(\frac{\partial}{\partial x^j}\rho\right)\delta^{ij}\delta_{kl}\right].\end{aligned}\quad (D.1)$$

From here, in view of Eq.(3.29a), we obtain:

$$g^{kl}\Gamma_{kl}^i = -\frac{1}{2}\left[\delta^{il}\left(\frac{\partial}{\partial x^l}\rho\right) + \delta^{ik}\left(\frac{\partial}{\partial x^k}\rho\right) - d\left(\frac{\partial}{\partial x^j}\rho\right)\delta^{ij}\right] = \frac{d-2}{2}\delta^{ij}\left(\frac{\partial}{\partial x^j}\rho\right) \quad (D.2)$$

Using Eq.(3.29a) and selecting sign "+" we obtain for *covariant* components of the vector b^i

$$b_i = a_i - \frac{d-2}{2}\frac{\partial}{\partial x^i}\ln\rho. \quad (D.3a)$$

In view of Eq.(3.44) we need to consider as well the sign "-" resulting in

$$b_i = a_i + \frac{d-2}{2}\frac{\partial}{\partial x^i}\ln\rho. \quad (D.3b)$$

But after Eq.(3.27b) we obtained for the contravariant components of b^i the following transformation law: $b'^i = b^i + 2g^{ij}\theta_j$. Written in the covariant form this relation looks as follows

$$b'_i = b_i + 2\theta_i. \quad (D.4)$$

Since the identically nonzero vector a_i always can be represented as

$$a_i = \frac{\partial}{\partial x^i}\psi \quad (D.5)$$

it is possible by combining Eq.s(D.3b) and (D.4) to represent vector b'_i as

$$b'_i = \frac{\partial}{\partial x^i}\left[\psi + \frac{d-2}{2}\ln\rho + 2\theta\right]. \quad (D.6)$$

From here, it follows that if in Eq.(3.27a) the transformation is performed with

$$\theta = -\frac{d-2}{4}\ln\rho - \frac{1}{2}\psi, \quad (D.7)$$

the transformed Eq.(3.27b) will have $b'^i = 0, i = 1, \dots, d$. In calculating of Eq.(3.37) we needed to evaluate Γ_{ki}^i . In view of Eq.(D.1), this can be done as follows.

$$\Gamma_{ki}^i = -\frac{1}{2\rho}(\rho_i\delta_k^i + \rho_k\delta_i^i - \rho_j\delta^{ij}\delta_{ki}) = -\frac{d}{2}\frac{\rho_k}{\rho}. \quad (D.8)$$

Appendix E. Solution of the constraint Eq.(4.29). SO(4) rotational group and Plücker matrices

In this appendix we are discussing the role of the constraint, Eq.(4.29), in finding solutions of Eq.(4.26b). This equation is obtained by mapping of the 3 dimensional Schrödinger's Eq.(3.19a) with potential Eq.(3.19b) ($\gamma = 1$) onto S^3 . We need to demonstrate that the constraint, Eq.(4.29), is enforcing the spherical topology of S^3 .

We begin with using some results from our work [3]. Specifically, in section 6 of [3] we defined the moment \mathbf{L} and the Laplace-Runge-Lenz vector \mathbf{A} for the hydrogen atom model Hamiltonian. These vectors are subject to the 3-dimensional constraint $\mathbf{L} \cdot \mathbf{A} = 0$. To connect this result with Fock's results on hydrogen atom [37], we found in [3] that it is convenient to look at this constraint from the point of view of the 4-dimensional space. Both in 3 and 4 dimensions this equation represents the Plücker embedding condition. Because of this, the constraint, Eq.(4.29), is of the same category as previously studied constraint $\mathbf{L} \cdot \mathbf{A} = 0$. Quantization of \mathbf{L} and \mathbf{A} leads to 2 decoupled $\mathfrak{so}(3)$ Lie algebras as demonstrated in [3] and [87]. From group theory [87] it is known that the Lie algebra $\mathfrak{so}(4) \simeq \mathfrak{so}(3) \oplus \mathfrak{so}(3)$. This observation is the starting point of our analysis.

Let the vector $\mathbf{X} \in \mathbf{R}^4$ be described as $\mathbf{X} = [U, V, W, T]^T$. Using this notation we consider all points $\mathbf{X} \in \mathbf{R}^4 \setminus \mathbf{0}$. They can be used for design of projective space \mathbf{P}^3 . It can be constructed out of equivalence classes made out of vectors \mathbf{X} and $\mathbf{Y} \in \mathbf{R}^4 \setminus \mathbf{0}$ such that $\mathbf{X} = \lambda \mathbf{Y}$, $\lambda \neq 0$. To distinguish between the \mathbf{R}^4 and \mathbf{P}^3 the *homogenous coordinates* in \mathbf{P}^3 should be introduced, e.g. via notation: $\mathbf{X} = [U:V:W:T]$. This helps us to identify the points of \mathbf{P}^3 with the equivalence classes. Clearly, the 3 dimensional vectors, e.g. \mathbf{v} , are obtained now as $\mathbf{v} = [\frac{U}{T}, \frac{V}{T}, \frac{W}{T}]^T$. Reciprocally, a homogenous presentation of the 3-vector $\mathbf{v} = [U, V, W]^T$ is given by $\mathbf{v} = [U:V:W:1]$. Thus, \mathbf{P}^3 is made out of all points $[U, V, W]^T$ of \mathbf{R}^3 plus the points $[U:V:W:0]$ at infinity comprising a projective plane \mathbf{P}^2 . Planes in \mathbf{P}^3 can be defined with help of the concept of duality defined with help of the scalar product \langle, \rangle in Euclidean space. Specifically, consider the equation

$$\langle \boldsymbol{\pi}, \mathbf{X} \rangle = \pi_1 X_1 + \pi_2 X_2 + \pi_3 X_3 + \pi_4 X_4 = 0. \quad (\text{E.1a})$$

It is looking exactly the same as Eq.(4.29). Its interpretation is the following. The point $\mathbf{X} = [X_1:X_2:X_3:X_4]$ lies on the hyperplane $\boldsymbol{\pi} = [\pi_1:\pi_2:\pi_3:\pi_4]$ if and only if Eq.(E.1a) is satisfied. Thus, in projective spaces the planes and the points are equivalent in view of the duality just defined. Clearly, in a more familiar 3 dimensional setting the same result is presented as

$$\pi_1 X + \pi_2 Y + \pi_3 Z + \pi_4 = 0, \quad (\text{E.1b})$$

where $X = X_1/X_4, Y = X_2/X_4, Z = X_3/X_4$. Eq.(E.1a) can be conveniently rewritten as

$$\boldsymbol{\pi}^T \mathbf{X} = 0. \quad (\text{E.1c})$$

Because of the duality, there is a compelling reason to look as well at the equation

$$\begin{bmatrix} \mathbf{X}_1^T \\ \mathbf{X}_2^T \\ \mathbf{X}_3^T \end{bmatrix} \boldsymbol{\pi} = 0. \quad (\text{E.2a})$$

This should be understood as follows. Three points $\mathbf{X}_1, \mathbf{X}_2, \mathbf{X}_3$ in general position define a plane. Each of these points is satisfying Eq.(E.1c), that is, $\boldsymbol{\pi}^T \mathbf{X}_i = 0, i = 1 \div 3$. Therefore, Eq.(E.2a) is defining a plane determined by 3 prescribed points in general position. By applying duality, we also get

$$\begin{bmatrix} \boldsymbol{\pi}_1^T \\ \boldsymbol{\pi}_2^T \\ \boldsymbol{\pi}_3^T \end{bmatrix} \mathbf{X} = 0. \quad (\text{E.2b})$$

To proceed, we need to define the notion of a line in \mathbf{P}^3 and to connect it with the results pertinent to the Lie algebra $so(4)$. In doing so, we follow the original Plücker ideas described in [3] and [94]. Some results from lecture notes by William Goldman [139] and those of [140] are also helpful. Specifically, we need to recall that the Lie algebra $so(4)$ is made out of the set of 4×4 skew symmetric matrices. These can be built as follows

$$\begin{aligned} \mathbf{R}^4 \times \mathbf{R}^4 &\rightarrow so(4) \text{ or, explicitly,} \\ (\mathbf{v}, \mathbf{w}) &\rightarrow \mathbf{v} \wedge \mathbf{w} := \mathbf{w}\mathbf{v}^T - \mathbf{v}\mathbf{w}^T := \mathbf{L}. \end{aligned} \quad (\text{E.3})$$

Here \mathbf{L} is defining the Plücker matrix describing the line l in \mathbf{P}^3 . The notion of Plücker matrix is absent in [3], [94] and [136]. But it can be found in [137]. It is essential, however, and requires some explanation.

To do so, we follow Plücker's ingenious idea that 3 dimensional lines can be described in terms of 3 dimensional vectors \mathbf{y} and \mathbf{x} , taken from the same origin \mathbf{o} , whose ends are located on the line l . In 3-space the direction for such a line is \mathbf{r} : $\mathbf{r} = \mathbf{y} - \mathbf{x}$. The shortest distance \mathbf{z} between l and \mathbf{o} , the cross product $\mathbf{n} = \mathbf{y} \times \mathbf{x}$ and \mathbf{r} are forming the right handed tripod. Any 2 of these 3 vectors are sufficient to determine the 3rd one. Typically the vectors \mathbf{r} and \mathbf{n} are chosen. With these results in our hands, following Plücker, we introduce

$$p_{01} = y_1 - x_1 ; p_{23} = x_2 y_3 - x_3 y_2, \quad (\text{E.4a})$$

$$p_{02} = y_2 - x_2 ; p_{31} = x_3 y_1 - x_1 y_3, \quad (\text{E.4b})$$

$$p_{03} = y_3 - x_3 ; p_{12} = x_1 y_2 - x_2 y_1. \quad (\text{E.4c})$$

From these results it follows that $\mathbf{x} = (x_1, x_2, x_3)^T, \mathbf{y} = (y_1, y_2, y_3)^T, p_{ij} = x_i y_j - x_j y_i, x_0 = y_0 = 1$. The Plücker relation, Eq.(6.18c), of [3] in terms of just defined notations acquires the form $\mathbf{n} \cdot \mathbf{r} = 0$, where

$$\mathbf{r} = \begin{bmatrix} p_{01} \\ p_{02} \\ p_{03} \end{bmatrix} \text{ and } \mathbf{n} = \begin{bmatrix} p_{23} \\ p_{31} \\ p_{12} \end{bmatrix}^T.$$

Explicitly,

$$p_{01}p_{23} + p_{02}p_{31} + p_{03}p_{12} = 0. \quad (\text{E.5})$$

Suppose now that \mathbf{q} is the point lying on the line l determined by \mathbf{n} and \mathbf{r} . Then, the condition that the point \mathbf{q} is indeed lying on the line l is given by

$$\mathbf{r} \times \mathbf{q} = \mathbf{n}. \quad (\text{E.6})$$

To prove this, we assume that $\mathbf{q} = \mathbf{y} + \lambda \mathbf{r}$ where λ is some nonzero arbitrary multiplier. By substituting this result into Eq.(E.6) we indeed obtain the identity. Next, we need to convert this result into manifestly 4 dimensional form. This can be achieved as follows. We begin with representation for the Lie algebra $\text{so}(3)$. A suitable basis is made out of matrices $(\mathcal{A}_i)_{jk} = -\epsilon_{ijk}$. Here i, j and k can take values $1 \div 3$. To generalize this result to $\text{so}(4)$, it is convenient to begin with

$$(\mathcal{A}_1)_{jk} = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, (\mathcal{A}_2)_{jk} = \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \\ -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, (\mathcal{A}_3)_{jk} = \begin{pmatrix} 0 & -1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} \quad (\text{E.7})$$

These matrices obey the $\text{so}(3)$ commutation relations $[\mathcal{A}_i, \mathcal{A}_j] = \epsilon_{ijk} \mathcal{A}_k$. Using just defined matrices, we can now define the \mathcal{B} -matrices also obeying the same commutator algebra $[\mathcal{B}_i, \mathcal{B}_j] = \epsilon_{ijk} \mathcal{B}_k$. Thus, we obtain $\text{so}(4) \simeq \text{so}(3) \oplus \text{so}(3)$. This is the Lie algebra of hydrogen atom [3], [87]. Explicitly, the \mathcal{B} matrices are given by

$$(\mathcal{B}_1)_{jk} = \begin{pmatrix} 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix}, (\mathcal{B}_2)_{jk} = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{pmatrix}, (\mathcal{B}_3)_{jk} = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 \\ 0 & 0 & 1 & 0 \end{pmatrix}. \quad (\text{E.8})$$

By looking at Eq.(E.3) we consider now the designing of a specific line- the X axis. It can be constructed using 2 reference points

$$\begin{bmatrix} 0 \\ 0 \\ 0 \\ 1 \end{bmatrix} = \mathbf{w} \text{ and } \begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \end{bmatrix} = \mathbf{v}$$

representing the origin of X-axis and the point at infinity for the X axis. Using Eq.(E.3), we obtain now for the X-line the following result:

$$\mathbf{L} = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 & 0 \end{bmatrix} - \begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \end{bmatrix} \begin{bmatrix} 0 & 0 & 0 & 1 \end{bmatrix} = \begin{pmatrix} 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix} = (\mathcal{B}_1)_{jk}. \quad (\text{E.9})$$

Just obtained result allows us to rewrite Eq.(E.6) in 4 dimensional form

$$\begin{pmatrix} 0 & p_{23} & -p_{13} & p_{12} \\ -p_{23} & 0 & p_{03} & -p_{02} \\ p_{13} & -p_{03} & 0 & p_{01} \\ -p_{12} & p_{02} & -p_{01} & 0 \end{pmatrix} \cdot \begin{pmatrix} q_0 \\ q_1 \\ q_2 \\ q_3 \end{pmatrix} = 0 \quad (\text{E.10})$$

The correctness of this result is easily checked following [140], page 201. From this reference we find

$$\mathbf{x} \times \mathbf{y} = \begin{pmatrix} 0 & -x_3 & x_2 \\ x_3 & 0 & -x_1 \\ -x_2 & x_1 & 0 \end{pmatrix} \cdot \begin{pmatrix} y_1 \\ y_2 \\ y_3 \end{pmatrix}. \quad (\text{E.11})$$

The 4 dimensional extension of this result is straightforward. Evidently, the result, Eq.(E.10), coincides with the constraint, Eq.(4.29). However from Eq.(4.26b) it follows that there are 4 momenta while we just got 6 vectors since every $\text{so}(4)$ matrix is decomposable in terms of the 6 skew-symmetric matrices \mathcal{A}_i and \mathcal{B}_i . To clarify this mismatch, we follow [3] and [139]. In \mathbf{R}^4 we are having $\{\mathbf{e}_0, \dots, \mathbf{e}_3\}$ as the basis. Using this basis we can construct the set of exterior products. To label these products we need to define the subsets $J_p = \{i_1, \dots, i_p\}, p \leq n$. There are exactly $n!/(n-p)!p!$ of such subsets. Change of the basis: $\{\mathbf{e}_0, \dots, \mathbf{e}_n\}$ to $\{\mathbf{e}'_0, \dots, \mathbf{e}'_n\}$ leads to the projective equivalence, e.g. to $\mathbf{e}'_{i_1} \wedge \dots \wedge \mathbf{e}'_{i_p} = \det(A_{ij}) \mathbf{e}_{i_1} \wedge \dots \wedge \mathbf{e}_{i_p}$. The Grassmannian $\text{Gr}_p(\mathbf{R}^n)$ is made out of all of just defined p -dimensional subspaces of \mathbf{R}^n . Let $n=4$ and consider all 2 dimensional subspaces of $\text{Gr}_2(\mathbf{R}^4)$. These are made of 2- planes, e.g. $\mathbf{e}_i \wedge \mathbf{e}_j$. There are $6=4!/2!2!$ planes which replace the 4 basis vectors $\{\mathbf{e}_0, \dots, \mathbf{e}_3\}$. Because of the projective equivalence of such planes, they are Plücker embedded into 5 dimensional projective space \mathbf{P}^5 . From Eq.s(E.4a-c) it is possible to establish 1-1 correspondence between the 6 planes set $\mathbf{e}_i \wedge \mathbf{e}_j$ and the 6 planes set p_{ij} . But p_{ij} are the momenta! The first of them, the \mathbf{p}_0 , is made out of planes $\mathbf{e}_0 \wedge \mathbf{e}_1$, $\mathbf{e}_0 \wedge \mathbf{e}_2$, $\mathbf{e}_0 \wedge \mathbf{e}_3$, There are 3 $\text{so}(4)$ matrices associated with these planes. The second- the \mathbf{p}_1 , is made of planes $\mathbf{e}_1 \wedge \mathbf{e}_2$ and $\mathbf{e}_1 \wedge \mathbf{e}_3$ and the associated with them $\text{so}(4)$ matrices and, the 3rd, the \mathbf{p}_2 , is made out of plane $\mathbf{e}_2 \wedge \mathbf{e}_3$ and, the associated with it, the $\text{so}(4)$ matrix. These arguments explain the mismatch. The 6 infinitesimal generators obtained with help of 6 $\text{so}(4)$ matrices are obtainable in exactly the same way as 3 generators for the 3 dimensional rigid rotator. Their explicit form is given in Appendix C of [87].

Appendix F. Essentials of the Sturmian problem and of the conformal Stäckel transform

In our exposition of this topic we follow [69],[90],[138] and [139]. In particular, following [138] we begin with brief discussion of the Sturmian problem. The hydrogen orbitals satisfy the 1-electron stationary Schrödinger equation (in the appropriately chosen system of units)

$$\left[-\frac{1}{2}\nabla^2 - \frac{Z}{r} + \frac{Z^2}{2n}\right]\psi_{nlm}(\mathbf{x}) = 0 \quad (\text{F.1a})$$

Denote $k = \frac{Z}{n}$, and rewrite Eq.(F.1a) using just introduced notation:

$$[-\frac{1}{2}\nabla^2 - \frac{nk}{r} + \frac{k^2}{2}]\tilde{\psi}_{nlm}(\mathbf{x}) = 0. \quad (\text{F.1b})$$

The eigenfunctions $\tilde{\psi}_{nlm}$ obey the *isoenergetic* Sturmian (Schrödinger) equation:

$$[-\frac{1}{2}\nabla^2 + \frac{k^2}{2}]\tilde{\psi}_{nlm}(\mathbf{x}) = \beta_n \frac{1}{r} \tilde{\psi}_{nlm}(\mathbf{x}). \quad (\text{F.1c})$$

In this equation, the energy $E = -\frac{k^2}{2}$ is fixed and the eigenvalue β_n is $\beta_n = nk$. If the eigenfunctions $\psi_{nlm}(\mathbf{x})$ are orthogonal, i.e.

$$\int d^3x \psi_{n'l'm'}^*(\mathbf{x}) \psi_{nlm}(\mathbf{x}) = \delta_{n'n} \delta_{l'l} \delta_{m'm} \quad (\text{F.2a})$$

, the eigenfunctions $\tilde{\psi}_{nlm}(\mathbf{x})$ are weighted orthogonal, i.e.

$$\int d^3x \tilde{\psi}_{n'l'm'}^*(\mathbf{x}) \left(\frac{n}{kr}\right) \tilde{\psi}_{nlm}(\mathbf{x}) = \delta_{n'n} \delta_{l'l} \delta_{m'm}. \quad (\text{F.2b})$$

The conformal Stäckel transform following [69],[90] can be defined as follows. Begin with the manifestly conformal (Laplace-type) equation

$$\mathcal{H}\Psi = [-\frac{1}{\lambda(\mathbf{x})}\nabla^2 + V(\mathbf{x})]\Psi = 0. \quad (\text{F.3})$$

In this equation let $V(\mathbf{x}) = W(\mathbf{x}) - U(\mathbf{x})E$. In accord with Eq.(F.1b) we shall consider the parameter E as fixed. The potential $U(\mathbf{x})$ defines the *conformal Stäckel transform*: from manifestly conformal Eq.(F.3) (e.g. read section 3) to the Helmholtz-type equation

$$\tilde{\mathcal{H}}\tilde{\Psi} = E\tilde{\Psi}, \quad (\text{F.4})$$

where

$$\tilde{\mathcal{H}} = -\frac{1}{\tilde{\lambda}(\mathbf{x})}\nabla^2 + \tilde{V}(\mathbf{x}). \quad (\text{F.5})$$

Here $\tilde{\lambda}(\mathbf{x}) = \lambda U$, $\tilde{V} = \frac{W(\mathbf{x})}{U(\mathbf{x})}$. The choice of the potential $U(\mathbf{x})$ is nontrivial. It is determined by the Bertrand-Darboux system of partial differential equations.

Theorem F.1. *Any second order conformal (Laplace-type) superintegrable system admitting nonconstant potential $U(\mathbf{x})$ can be Stäckel-transformed to a Helmholtz superintegrable system. This operation is invertible. The inverse mapping is taking true symmetries to conformal symmetries.*

Remark F.2. The nontriviality of the potential $U(\mathbf{x})$ is coming from the fact that it is obtainable as a solution of the system of second order (conformal) Bertrand-Darboux (B-D) partial differential equations (PDE).

Remark F.3. Solutions of the B-D system of PDE's resulting in obtaining of $U(\mathbf{x})$ is linked with solution of the Bertrand problem in curved spacetimes (e.g. read section 5). D-O by passed the problem of finding the $U(\mathbf{x})$ by solving the B-D system of PDE's. They simply guessed correctly the potential.

Remark F.4. Eq.(F.3) is exactly of the same type as Eq.s(3.14),(3.35). Therefore, methods described in section 3, also involving solutions of PDE's, should be considered as complementary to those in [69], [138], [139].

Remark F.5. There are three types of Laplace-type conformally superintegrable equations. These are:

a) Recasting a Helmholtz superintegrable system $H\Psi = E\Psi$ into the Laplace-type form $H'\Psi = 0$, $H' = H - E$. Here E is absorbed into potential as a parameter. Clearly, this is the case of Sturmian Eq.(F1.b).

b) Replacing a Helmholtz superintegrable system $H\Psi = E\Psi$ by the Laplace-type $H'\Psi = 0$ in which $H' = H - E_0$ with E_0 fixed once and for all eigenvalue. This is again the Sturmian-type problem.

c) Begin with a Helmholtz-type equation $H\Psi = E\Psi$ which is not superintegrable and convert it into the Laplace-type. The resulting equation will have the induced conformal symmetries.

Remark F.6. Just described classification justifies the transition from Eq.(1.2) to Eq.(1.3).

Appendix G. Thomas-Fermi treatment of multielectron atoms. Contributions of Fermi, Tietz and Klechkovski into the proof of Madelung's rule

This appendix is designed for chemists and atomic physicists. Mathematically inclined researchers can skip it. Although the Thomas-Fermi (T-F) method is only asymptotical (works well in the limit of high electron densities), because of its simplicity and reasonably good accuracy, we provide some facts about this method helpful for practical applications of the formalism of the main text to multielectron atoms. The T-F method and contributions by Tietz to this method used in section 4.5 are concisely summarized in the book by Flugge [24]. Belokolos [39] used the Tietz potential (Eq.s(G.8),(G.9) below) for the WKB-assisted "proof" of the Madelung rule. His results are neither exact not conclusive even though he is citing the paper by Wong [140] in which the Madelung rule is seemingly derived. In his derivation Wong used the Tietz potential. Wong's results are solely based on seminal works by Fermi [121], Tietz [24] and Klechkowski [141]. It is impossible to develop theory of the Madelung rule without acknowledging contributions of these authors. The strongest and

chronologically the earliest contribution to this topic was made by Enrico Fermi. Following publication of his ground breaking paper on the T-F method in 1927, in 1928 Fermi came up with the paper [121] describing applications of the T-F method to the periodic system of elements. In it he posed and solved the following problem:

Find analytically a relationship between the atomic number Z and the number of electrons N_l with azimuthal number l .

Knowledge of such a relationship allows then to solve the related problem of major significance:

Find a sequence of Z 's for which for the first time electrons with azimuthal numbers s, p, d, f, \dots emerge.

Fermi found a sequence : 1, 5, 21, 58. These are exactly the atomic numbers initiating sequential irregularities in filling of the periodic system. Subsequently, Tietz [142] simplified Fermi calculations with help of the potential bearing his name. The whole book by Klechkowski on "proving" the Madelung rule is effectively a discussion around these works by Tietz. The paper by Wong is a condensed summary of the results by Klechkowski summarized in his book [141]. As such, it suffers from the same drawbacks as Klechkowski's book. Specifically, in his book Klechkowski discusses from various perspectives the result by Tietz for N_l :

$$N_l = 2(2l + 1)(6Z)^{\frac{1}{3}} - 2(2l + 1)^2. \quad (\text{G.1})$$

Eq.(G.1) coincides with Eq.(14) of Wong's paper [120] where it was derived independently based directly on ideas of Enrico Fermi. Next, following Klechkowski's philosophy, Wong runs into myriad of the same kinds of problems. They began with an assumption

$$\frac{N_l}{2(2l + 1)} \simeq n - l - 1 = n_r. \quad (\text{G.2})$$

Here n and n_r are respectively the principal and radial quantum numbers. A comparison between Eq.s(G.1) and (G.2) then produces:

$$(6Z)^{\frac{1}{3}} - (2l + 1) = n - l - 1. \quad (\text{G.3a})$$

However, already from Eq.(G.1) it is evident that on the l.h.s the N_l should be a nonnegative integer while on the r.h.s. the term $(6Z)^{\frac{1}{3}}$ is making the r.h.s. noninteger for general Z 's. Since Eq.(G.3a) is incorrect for arbitrary nonnegative Z 's, the same is true for its corollary

$$n + l = (6Z)^{\frac{1}{3}}. \quad (\text{G.3b})$$

Wong claims, however, that this result is mathematically equivalent to the Madelung rule. Since $(6Z)^{\frac{1}{3}}$ is not always an integer, the rest of Klechkowski's book [141] is devoted to finding some corrections to Eq.(G.1) making $n + l$ an integer. Wong missed this issue (studied by Klechkowski) completely. His paper entitled "Theoretical justification of the Madelung rule" culminates in Eq.(G.3b). None of the corrections discussed in Klechkowski book are rigorous, systematic or convincing. Surprisingly, he made no attempts to use directly the results by

Fermi. Instead, he made a few comments on the paper by Ivanenko and Larin [143] which, in our opinion, deserves more attention than that given to it in Klechkowski's book. The authors of [143] used the Thomas-Fermi-Dirac method to recalculate the results by Fermi in a direct and simple way. The final result is expressible in compact form:

$$Z_l = \gamma(Z_l)(2l + 1)^3. \quad (\text{G.4})$$

The Z_l -dependent factor $\gamma(Z_l)$ is calculated iteratively for each Z_l . Fig.1 of [143] indicates that this variable factor can be safely replaced by the constant factor: $\gamma(Z_l) \simeq 0.169$. In such a case use of Eq.(G.4) reproduces (by rounding the numbers in Eq.(G.4)) the Fermi sequence of Z 's: $1(l = 0)$, $5(l = 1)$, $21(l = 2)$ and $58(l = 3)$ ⁶⁴. Strictly speaking, the minimization procedure involved in obtaining Eq.(G.4) is not extendable to the values of Z significantly different from those in the Fermi sequence. In his book Klechkowski missed this restriction completely. If however, in Klechkowski style, we do use Eq.(G.4) with unrestricted Z 's in Eq.(G.3b) in order to obtain

$$n + l = (6Z)^{\frac{1}{3}} = (6 \cdot 0.169)^{\frac{1}{3}}(2l + 1) = 1.004645(2l + 1) \simeq 2l + 1 \quad (\text{G.5})$$

then, since $n = n_r + 1 + 1$, Eq.(G.5) reduces to the identity, if $n_r = 0$. This requirement makes perfect physical sense as explained in section 5.

Now we are in the position to make a number of comments about the Tietz potential. In the book by Flugge [24] in sections devoted to the T-F theory our readers will find a nice discussion about the Tietz potential and how it fits the T-F theory. Typically, readers with standard physics background are left with an impression that the T-F theory is reasonably well describing multielectron atoms only when the number of electrons tend to infinity and is useless for description of, say, a hydrogen or helium atom. In the lecture notes by Roi Baer [144] it is demonstrated that, surprisingly, the T-F theory provides reasonable results even for the hydrogen atom. Additional details on T-F theory (without uses of the Tietz potential) are beautifully summarized in the book by March [27].

To develop our intuition the scaling analysis by Feynman, Metropolis and Teller [145] is the most helpful. Following these authors, we consider a similarity transformations for an atom. They are consisting of increasing the nuclear charge of an atom while simultaneously increasing the number of electrons so that the atom remains neutral. Evidently, such a transformation is describing a relation between solutions for various nuclear charges. It is assumed that the De Broglie length λ remains unchanged under scaling. Transformations consists of the following infinitesimal changes: $Z \rightarrow Z(1 + \zeta)$, $r \rightarrow r(1 + \rho)$, $\bar{E} \rightarrow \bar{E}(1 + \eta)$. Here \bar{E} represents any form of the energy (classical or quantum) per electron. At the same time, the electron density q is changed by $q(1 + \nu)$. The charge balance leads to the relation: $qr^3 \rightarrow q(1 + \nu)r^3(1 + \rho)^3 = Z(1 + \zeta)$ implying $(1 + \nu)(1 + \rho)^3 = 1 + \zeta$ or $\nu + 3\rho = \zeta$. Since the potential energy of an electron changes as $(1 + \zeta)/(1 + \rho) = (1 + \eta)$, this leads to: $\zeta - \rho = \eta$. In the T-F theory [24] $\bar{E} = \kappa q^{\frac{2}{3}}$ where κ is known constant. So, whenever this theory makes sense, we should have: $(1 + \nu)^{\frac{2}{3}} = (1 + \eta)$, implying $\frac{2}{3}\nu = \eta$. Elementary calculation based on these results

⁶⁴Evidently, Fermi also used some rounding.

produces: $\rho = -\frac{1}{3}\zeta$ and $\eta = \frac{4}{3}\zeta$. Thus, the energy \bar{E} per electron changes as $\bar{E}(1+\frac{4}{3}\zeta)$ and for neutral atom initially made of Z electrons it changes as $\bar{E}(1+\frac{4}{3}\zeta)Z(1+\zeta) = \bar{E}Z(1+\frac{7}{3}\zeta)$. Since in these arguments we assumed the validity of T-F relationship $\bar{E} = \kappa q^{\frac{2}{3}}$, the dimensionless T-F equation

$$\frac{d^2\varphi}{dx^2} = \frac{\varphi^{\frac{3}{2}}}{\sqrt{x}} \quad (\text{G.6})$$

connected with this relationship automatically establishes the characteristic length scale parameter

$$a = 0.8853a_0Z^{-\frac{1}{3}}, \quad (\text{G.7})$$

where a_0 is the Bohr radius, $a_0 = \frac{\hbar^2}{me^2}$. From here emerges the dimensionless length scale: $x = \frac{r}{a}$. As before, we shall use the system of units in which $\hbar = m = 1$. In view of just presented results, the potential energy of electron scales as $Z/a \simeq Z^{\frac{4}{3}}$ so that for Z atomic electrons the total energy scales as $Z \cdot Z^{\frac{4}{3}} = Z^{\frac{7}{3}}$. This result is consistent with the previously obtained scaling result $\bar{E}Z(1+\frac{7}{3}\zeta)$ for the total energy. The result $Z^{\frac{7}{3}}$ was first derived by Milne [27]. Schwinger [146] obtained a correction to just obtained scaling result. With it, such a correction gives good results for all Z 's. This fact, is of importance if one is interested in comparing much more technically cumbersome Hartree-Fock and T-F theories. Another example of this kind is related to the discussion related to the Tietz potential. The origins of this potential are connected with study of solutions of the T-F Eq.(G.6). This nonlinear equation does not admit exact analytical solution and, in addition, it requires the assignment of boundary conditions. For neutral atoms these are simple: $\varphi(0) = 1$ and $\varphi(\infty) = 0$. Because of their simplicity, Tietz suggested the analytical form for φ :

$$\varphi(x) = \frac{1}{(1 + \alpha x)^2}, \quad \alpha = 0.53625. \quad (\text{G.8})$$

The constant α is not a fitting parameter. It is determined by the electroneutrality normalization condition [24]. The Tietz potential $V(x)$ is obtained with help of $\varphi(x)$ as follows

$$V(r) = -\frac{Ze}{r}\varphi\left(\frac{r}{a}\right) \quad (\text{G.9})$$

For $r \rightarrow 0$, $V(r)$ tend to the usual attractive Coulombic potential while for larger r 's it represents $V_{eff}(r)$ introduced in Eq.(1.2). This fact was reported in [25] on p.664, Fig.10. From this figure it follows that the plot of $\varphi(x)$ vs. x obtained with help of the T-F method practically coincides with that obtained numerically with help of the Hartree-Fock method.

References

- [1] P.Thyssen, A. Ceulemans, Shattered Symmetry, Oxford University Press, New York, 2017.
- [2] H. Bethe, R. Jackiw, Intermediate Quantum Mechanics, CRC Press, New York, 2018.
- [3] A. Kholodenko, L. Kauffman, Huygens triviality of the time-independent Schrödinger equation. Applications to atomic and high energy physics, Ann.Phys. 390 (2018) 1-59.
- [4] P-O. Löwdin, Some comments on periodic system of elements, Int. J.Quant.Chemistry 3s, (1969) 331-334.
- [5] L.Allen,E. Knight, The Löwdin challenge. Origin of the $n + l, n$ (Madelung) rule for filling the orbital configurations, Int.J.Quantum Chemistry 90 (2002) 80-88.
- [6] E. Scerri, G. Restrepo, Mendeleev to Oganesson, Oxford University Press, New York, 2018.
- [7] Y. Demkov, V. Ostrovsky, Internal symmetry of the Maxwell "fish-eye" problem and the Fock group for hydrogen atom. Sov.Phys. JETP 13 (1971) 1083-1087.
- [8] Y. Kitagawara,A. Barut, Period doubling in the $n + l$ filling rule and dynamical symmetry of the Demkov-Ostrovsky atomic model, J.Phys.B 16 (1983) 3305-3322.
- [9] Y.Kitagawara, A. Barut, On the dynamical symmetry of the periodic table: II. Modified Demkov-Ostrovsky atomic model, J.Phys B 17 (1984) 4251-4259.
- [10] H.Goldstein, Ch. Poole, J. Safko, Classical Mechanics, Pearson Education Ltd, London, 2014.
- [11] M.Akila, D.Waltner, B. Gutkin, P. Braun,Th. Guhr, Semiclassical identification of periodic orbits in a quantum many-body system, PRL 118, (2017) 164101.
- [12] M. Gutzwiller, Chaos in Classical and Quantum Mechanic, Springer-Verlag, Berlin, 1990.
- [13] P.Cvitanovic, Chaos: Classical and Quantum, chaosbook.org
- [14] J.Wheeler, From Mendeleev's atom to the collapsing star, Atti del Convegno Mendeleeviano, pp 189-233, 1971.
- [15] R.Powers, Frequencies of radial oscillation and revolution as affected by features of a central potential, Atti del Convegno Mendeleeviano, pp 235-242, 1971.
- [16] J.Wheeler, Semi-classical analysis illuminates the connection between potential and bound states and scattering, in Studies in Mathematical Physics, pp 351-422. Princeton University Press, Princeton, 1976.
- [17] V.Ostrovsky, Dynamic symmetry of atomic potential. J.Phys. B 14 (1981) 4425-4439.
- [18] S.Kuru, J. Negro, O. Ragnisco, The Perlick system type I: From the algebra of symmetries to the geometry of the trajectories, Phys.Lett. A 381(2017) 3355-3363.

- [19] J. Little, Nondegenerate homotopies of curves on the unit 2-sphere.
J.Diff.Geom. 4 (1970) 339-34.
- [20] A.Ballesteros, A. Encio, F. Herranz, Hamiltonian systems admitting a
Runge–Lenz vector and an optimal extension of Bertrand’s theorem to curved
manifolds, Comm.Math.Phys. 290 (2009) 1033-1049.
- [21] R. Luneburg, Mathematical Theory of Optics, University of California Press,
Los Angeles, 1966.
- [22] C.Caratheodory, Optics, Julius Springer, Berlin, 1937.
- [23] V.Perlick, Bertrand spacetimes, Class.Quantum Grav. 9 (1992) 1009-1021.
- [24] S. Flugge, Practical Quantum Mechanics, Springer-Verlag, Berlin, 1999.
- [25] D. Kirzhnits, Y. Lozovik, G. Shpatkovskaya, Statistical model of matter.
Sov.Phys. Uspekhi 18 (1976) 649-672.
- [26] R. Latter, Atomic energy levels for the Thomas-Fermi and Thomas-Fermi-Dirac
potential, Phys.Rev. 99 (1955) 510-519.
- [27] N.March, Self-Consistent Fields in Atoms, Pergamon Press Ltd., London, 1975.
- [28] T.Tietz, Atomic energy levels for the approximate Thomas-Fermi Potential,
J.Chem.Phys. 25 (1956) 789-790.
- [29] T.Tietz, Contribution to the Thomas–Fermi Theory.
J.Chem.Phys. 49 (1968) 4391-4393.
- [30] A.Kholodenko, How the modified Bertrand theorem explains regularities
and anomalies of the periodic system of elements, arXiv.:2002.12128.
- [31] U. Leonhardt, Th. Philbin, Geometry and Light. Dover Publications,
New York, 2010.
- [32] H. Stephani, General Relativity, Cambridge University Press, Cambridge, 1990.
- [33] J.Milnor, On the Geometry of the Kepler Problem,
Am.Math.Monthly 90 (1983) 353-365.
- [34] B. Dubrovin, S. Novikov, A. Fomenko, Modern Geometry-Methods and
Applications, Vol.1, Springer-Verlag, Berlin, 1984.
- [35] J. Moser, E. Zehnder, Notes on Dynamical Systems,.
AMS Publishers, Providence, 2005.
- [36] J. Moser, Regularization of Kepler’s problem and the averaging method on
a manifold, Comm.Pure Appl.Math. 23(1970) 609-636.
- [37] S.Singer, Linearity, Symmetry and Prediction in the Hydrogen Atom,
Springer, Berlin, 2005.
- [38] A. Makowski, K.Gorska, Phys.Rev. A 79 (2009) 052116
- [39] E. Belokolos, Mendelev table: a proof of Madelung rule
and atomic Tietz potential, SIGMA 13 (2017) 038
- [40] P. Dirac, Lectures on Quantum Mechanics, Yeshiva University, New York, 1964.
- [41] V. Arnol’d, Mathematical Methods of Classical Mechanics,
Springer -Verlag, Berlin, 1989.
- [42] O. Jones, Analytical Mechanics for Relativity and Quantum Mechanics,
Oxford University Press, Oxford, 2011.

- [43] J. Struckmeier, Hamiltonian dynamics on the symplectic extended phase space for autonomous and non-autonomous systems, *J.Phys.A* 38 (2005) 1257-1278.
- [44] L. Prokhorov, S. Shabanov, *Hamiltonian Mechanics of Gauge Fields*, Cambridge University Press, Cambridge, 2011.
- [45] E. Anderson, *The problem of Time*, Springer International Publishing AG, Berlin, 2017.
- [46] K.Kuchar, Time and interpretations of quantum gravity, *Int.J.Mod.Phys.D* 20 (2011) 3-86.
- [47] K. Kuchar, Gravitation, geometry, and nonrelativistic quantum theory, *Phys.Rev.D* 22 (1980) 1285-1299.
- [48] A. Makowski, P. Peplowski, Zero energy wave packets which follow classical orbits, *Phys.Rev. A* 86 (2012) 042117.
- [49] H. Rosu, M. Reyers, K. Wolf, O.Obregon, Supersymmetry of Demkov-Ostrovsky effective potentials in the $R_0 = 0$ sector, *Phys.Lett. A* 208 (1995) 33-39.
- [50] A. Keane, R. Barrett, J. Simmons, The classical Kepler problem and geodesic motion on spaces of constant curvature, *JMP* 41 (2000) 8108-8116.
- [51] S. Chanda, G.Gibbons, P. Guha, Jacobi–Maupertuis metric and Kepler equation, *Int. J.of Geom.Methods in Mod.Physics* 14 (2017) 1730002.
- [52] A. Petrov, *Einstein Spaces*, Pergamon Press, London, 1969.
- [53] A. Makowski, Family of fish-eye-related models and their supersymmetric partners. *Phys. Rev. A* 81(2010) 052109.
- [54] R.Wald, *General Relativity*, The University of Chicago Press, Chicago, 1984.
- [55] M. Tsamparlis, A. Paliathanasis, Conformally related metrics and Lagrangians and their physical interpretation in cosmology, *Gen.Rel. Grav.* 45 (2013) 2003-2022.
- [56] A.Kholodenko, E. Ballard, From Ginzburg Landau to Hilbert Einstein via Yamabe, *Physica A* 380 (2007) 115-162.
- [57] A.Kholodenko, Towards physically motivated proofs of the Poincaré and geometrization conjectures, *Journal of Geometry and Physics* 58(2008) 259.
- [58] H.Urakawa, Geometry of Laplace-Beltrami operator on a complete Riemannian manifold, *Adv.Stud. Pure Math.* 22 (1993) 347-406
- [59] L.Ovsyannikov, *Group Analysis of Differential Equation*, Academic Press, New York,1982.
- [60] N.Ibragimov, *Transformation Groups Applied to Mathematical Physics*, D.Reidel Publ.Co, Boston, 1985.
- [61] K.Yano, *The Theory of Lie Derivatives and its Applications*, North -Holland, Amsterdam, 1957.
- [62] M.Obata, Conformal transformations of Riemannian manifolds, *J.Diff.Geom.* 4 (1970) 311-333.
- [63] H.Yamabe, On a deformation of Riemannian structures on compact manifolds, *Osaka Math.Journal* 12 (1960) 21-37.

- [64] T.Aubin, Some Nonlinear Problems in Riemannian Geometry, Springer-Verlag, Berlin, 1998.
- [65] L.Eisenhart, Riemannian Geometry, Princeton University Press, Princeton, 1926.
- [66] R. Courant, D. Hilbert, Methods of Mathematical Physics, Vol.2, Interscience Publishers, Amsterdam, 1966.
- [67] V. Ostrovsky, Group theory applied to the periodic table of the elements, in The Mathematics of Periodic Table. Nova Science Publishers, New York, pp. 265-311, 2006.
- [68] C. Boyer, E. Kalnins, W. Miller, Symmetry and separation of variables for the Helmholtz and Laplace equations, Nagoya Math. J. 60 (1976) 35-80.
- [69] E. Kalnins, J. Kress, W. Miller, Separation of Variables and Superintegrability. IOP Publishing, Bristol, 2018.
- [70] M. Chanachowicz, M. Chanu, R. McLenaghan, R-separation of variables for the conformally invariant Laplace equation, JGP 59 (2009) 876-884
- [71] N.Evans, Superintegrability in classical mechanics, Phys.Rev.A 41 (1990) 5666-5676.
- [72] T. Tuc, A. Danner, Absolute optical instruments, classical superintegrability, and separability of the Hamilton-Jacobi equation, Phys.Rev.A 96 (2017) 053838.
- [73] A. Linner, Periodic geodesics generator. Exp.Math 13 (2004) 199-206.
- [74] E.Kudryavtseva, D. Fedoseev, Superintegrable mechanical systems of Bertrand type, Modern Math and Its Applications 148, (2018) 37-57 (in Russian).
- [75] M.Cariglia, F. Alves, The Eisenhart lift: a didactical introduction of modern geometrical concepts from Hamiltonian dynamics, Eur. J. Phys. 36 (2015) 025018
- [76] J.Carinena, F. Herranz, M. Ranada, Superintegrable systems on 3-dimensional curved spaces: Eisenhart formalism and separability, J. Math.Phys. 58 (2017) 02270.
- [77] M. Cariglia, Null lifts and projective dynamics, Ann.Phys. 362 (2015) 642-658.
- [78] C.Boyer, E. Kalnins, W. Miller, Stäckel-equivalent integrable Hamiltonian systems, SIAM J.Math.Anal. 17 (1986) 778-797.
- [79] W.Miller, Multiseparability and superintegrability for classical and quantum systems, CRM Proc. Lecture Notes 26 (2000), 129-156.
- [80] A. Tsiganov, Transformation of the Stäckel matrices preserving superintegrability, JMP 60 (2019), 042701.
- [81] M.Berger, A Panoramic View of Riemannian Geometry, Springer-Verlag, Berlin, 2003.
- [82] A.Fordy, A. Galajinsky, Eisenhart lift of 2-dimensional mechanics. Eur.Phys.J. C 79 (2019) 301.
- [83] L.Landau, E. Lifshitz, Quantum Mechanics. Non-Relativistic Theory, Butterworth-Heinemann, London, 2003.
- [84] E. Kalnins, W. Miller, P. Winternitz, The group $O(4)$, separation of variables and the hydrogen atom, SIAM J.Appl.Math. 30 (1976) 630-664.
- [85] B.Adams, J. Cizek, L. Paldus, Representation Theory of $so(4,2)$ for the

- perturbation treatment of hydrogenic type hamiltonians by algebraic method, Int. J. Quantum Chem. 21 (1982) 153-171.
- [86] M. Gadella, J. Negro, L. Nieto, G. Pronko, M. Santander, Spectrum generating algebras for the free motion in S^3 , J.Math.Phys.52 (2011) 063509.
 - [87] M.Englefield, Group Theory and the Coulomb Problem, Wiley-Interscience, New York, 1972.
 - [88] A. Stone, Some Properties of Wigner coefficients and hyperspherical harmonics, Proc.Camb.Phil.Soc. 52 (1956) 424-430.
 - [89] B. Bransden, C. Joahain, Physics of Atoms and Molecules, Logman, Ltd., London, 1983.
 - [90] E. Kalnins, J. Kress, W. Miller, S. Post, Laplace-type equations as conformal superintegrable systems, Adv.Appl.Math. 46 (2011) 396-416.
 - [91] B. Kursunoglu, Modern Quantum Theory, W.H. Freeman and Co., New York, 1962.
 - [92] H. Bateman, A. Erdelyi, Higher Transcendental Functions. Vol.2., McGraw Hill Inc., New York, 1953.
 - [93] E. Schrödinger, A method of determining quantum-mechanical eigenvalues and eigenfunctions, Proc. Irish Acad, Section A 46 (1940) 9-16.
 - [94] A. Kholodenko, Applications of Contact Geometry and Topology in Physics, World Scientific, Singapore, 2013.
 - [95] P.Kustaanheimo, E. Steifel, Perturbation theory of Kepler motion based on spinor regularization, J. Reine Ang.Math.218 (1965) 204-219.
 - [96] D. Parfitt, D. Portnoi, The two-dimensional hydrogen atom revisited, J.Math.Phys. 43(2002) 4681-4691.
 - [97] B. George, S. Doman, The Classical Orthogonal Polynomials, World Scientific, Singapore, 2016.
 - [98] X. Yang, X. Guo, F. Chan, K. Wong, W. Ching, Analytic solution of a two-dimensional hydrogen atom. I. Nonrelativistic theory. Phys.Rev.A 43 (2001) 1186-1196.
 - [99] Y. Demkov, V. Ostrovsky, $n + l$ filling rule in the periodic system and focusing potentials, Sov. Phys. JETP 35(1972) 66-69.
 - [100] V. Ostrovsky, What and how physics contributes to understanding the periodic law, Found.Chemistry 3(2001) 145-181.
 - [101] J. Slater, Analytic Atomic Wave Functions, Phys.Rev.43 (1932) 33-43.
 - [102] E. Kerner, The solution of the Schrödinger equation for an approximate atomic field. Phys. Rev. 83, (1951) 71-75.
 - [103] A. Fet, Group Theory of the Chemical Elements, Walter de Gruyter, Boston, 2016.
 - [104] D. Dey, K. Bhattacharya, T. Sarkar, Astrophysics of Bertrand space-times. Phys.Rev.D 88 (2013) 083532.
 - [105] A. Kholodenko, Newtonian limit of Einsteinian gravity: from dynamics of Solar system to dynamics of stars in spiral galaxies, arXiv:1006.4650 v3
 - [106] D. Riglioni, Quantum Bertrand Systems,

- PhD Thesis. Roma III University, 2012.
- [107] D. Latini, Algebraic and Coalgebraic Aspects of Classical and Quantum Hamiltonian Systems, PhD Thesis. Roma III University, 2017.
 - [108] P. Woit, Quantum Theory, Groups and Representations, Springer, Berlin, 2017.
 - [109] G. Naber, Foundations of Quantum Mechanics,
<http://inspirehep.net/record/1411865?ln=en>
 - [110] M. Zworski, Semiclassical Analysis, AMS Publishers, Providence, 2012.
 - [111] E. Heller, The Semiclassical Way, Princeton University Press, Princeton, 2018.
 - [112] S. Benenti, Separation of variables in the geodesic Hamilton-Jacobi equation.
<http://www.sergiobenenti.it/cp/52.pdf>.
 - [113] A. Bruce, On the solution of the Hamilton-Jacobi equation by the method of separation of variables, PhD Thesis, University of Waterloo, 2000.
 - [114] A. Ballesteros, A. Enciso, F. Herranz, O. Ragnisco, Bertrand spacetimes as Kepler/oscillator potentials, *Class. Quantum Grav.* 25, (2008) 165005.
 - [115] E. Kalnins, W. Miller, G. Pogosyan, Coulomb-oscillator duality in spaces of constant curvature, *J. Math. Phys.* 41(2000) 2629-2657.
 - [116] E. Kalnins, J. Kress, W. Miller, Second-order superintegrable systems in conformally flat spaces. V. Two- and three-dimensional quantum systems, *J. Math. Phys.* 47 (2006) 093501.
 - [117] E. Kalnins, J. Kress, W. Miller, Second order superintegrable systems in conformally flat spaces. IV. The classical 3D Stäckel transform and 3D classification theory, *J. Math. Phys.* 47 (2006) 043514.
 - [118] P. Petersen, *Warped Products*.
<https://www.math.ucla.edu/~petersen/warpedproducts.pdf>.
 - [119] B-Y. Chen, Differential Geometry of Warped Product Manifolds and Submanifolds, World Scientific, Singapore, 2017.
 - [120] D. Kirzhnits, Y. Lozovik, Plasma oscillations of the electron shell of the atom, *Sov. Phys. Uspekhi* 9, (1966) 340-345.
 - [121] E. Fermi, Sulla deduzione statistica di alcune proprietà dell'atomo. Applicazione alla teoria del sistema periodico degli elementi, *Rend. Lincei* 7(1928) 342-346.
 - [122] M. Bander, C. Itzykson, Group theory and the hydrogen atom (I). *Rev. Mod. Phys.* 38(1966) 330-345.
 - [123] L. Kauffman, On Knots, Princeton University Press, Princeton, 1988.
 - [124] R. Albert, Computer-calculated explicit forms for representations of the three-dimensional pure rotation group, *Proc. Camb. Phil. Soc.* 65 (1969) 107-110.
 - [125] A. Besse, Manifolds all of Whose Geodesics are Closed, Springer-Verlag, Berlin, 1978.
 - [126] B. McKay, A summary of progress on the Blaschke conjecture. *Notices of the International Congress of Chinese Mathematics* 3(2015) 33-45.

- [127] L. Xu , X. Wang, T.Tyc, C. Sheng, S. Shu, H. Liu, H. Chen,
Light rays and waves on geodesic lenses. arXiv:1801.10438
- [128] M. Sarbort, Non–Euclidean geometry in optics. PhD Thesis.
Masaryk University, (2013)
www.researchgate.net/publication/287993246_Non-Euclidean_Geometry_in_Optics
- [129] L. Kuznia, E. Lundberg, Fixed points of conjugated Blaschke products with
applications to gravitational lensing, Computational methods and Function
Theory 9 (2009) 435-442.
- [130] A.Shoom, A.: Metamorphoses of a photon sphere,
Phys.Rev. D 96 (2017) 084056.
- [131] R.Nemiroff, Visual distortions near a neutron star and black hole
Am.J.Phys.61(1993) 619-632.
- [132] A.Celletti, From Order to Disorder in Gravitational
N-Body Dynamical Systems, Chaotic Worlds, pp. 203-226,
Springer, Dordrecht, 2006.
- [133] S.Chanda, G. Gibbons, P.Guha, Jacobi-Maupertuis metric and Kepler
equation, Int.J.of Geom.Methods in Mod.Physics 14(2016) 1730002.
- [134] H. Geiges, An Introduction to Contact Topology,
Cambridge University Press, Cambridge, 2008.
- [135] A. Kholodenko, Optical knots and contact geometry II. From Ranada dyons to
transverse and cosmetic knots, Ann.Phys. 371(2016) 77-124.
- [136] W.Goldmann, Lines in \mathbf{P}^3 , <http://terpconnect.umd.edu/~wmg/plucker.pdf>
- [137] W.Förstner,B.Wrobel, Photogrammetric Computer Vision,
Springer, Berlin, 2016.
- [138] C. Coletti, D. Calderini, V. Aquilanti, D-dimensional Kepler-Coulomb
sturmiens and hyperspherical harmonics as complete orthonormal atomic and
molecular orbitals, Adv.Quantum Chemistry 67 (2013) 73-127.
- [139] J. Avery, J. Avery,Jr, Hyperspherical Harmonics and Their Physical
Applications,World Scientific, Singapore, 2018.
- [140] D. Wong, Theoretical justification of Madelung’s rule,
J.Chem.Educ. 56 (1979) 714-717.
- [141] V.Klechkowski, Distribution of Atomic Electrons and the Rule of Sequential
Filling of the (n+l) Groups, Atomizdat, 1968, (in Russian)
- [142] T. Tietz, Über eine Approximation der Fermischen Verteilungsfunktion,
Ann.der Phys. 5 (1960) 237-240.
- [143] D. Ivanenko, S. Larin, Remarks about the periodic system of elements,
Sov. Physics Doklady 88 (1953) 45-47.
- [144] R.Baer, Electron Density Functional Theory,
The Fritz Haber Center for Molecular Dynamics, 2009.
- [145] R. Feynman, N. Metropolis, E. Teller, Equations of state of elements based
on the generalized Fermi-Thomas theory, Phys.Rev.75 (1949) 1561-1673.
- [146] J.Schwinger,Thomas-Fermi model: The leading correction,

Phys.Rev. A 22 (1980) 1827-1832.