GROUP GEOMETRICAL AXIOMS FOR MAGIC STATES OF QUANTUM COMPUTING

MICHEL PLANAT[†], RAYMOND ASCHHEIM[‡], MARCELO M. AMARAL[‡] AND KLEE IRWIN[‡]

ABSTRACT. Let H be a non trivial subgroup of index d of a free group G and N the normal closure of H in G. The coset organization in a subgroup H of G provides a group P of permutation gates whose common eigenstates are either stabilizer states of the Pauli group or magic states for universal quantum computing. A subset of magic states consists of MIC states associated to minimal informationally complete measurements. It is shown that, in most cases, the existence of a MIC state entails that the two conditions (i) N = G and (ii) no geometry (a triple of cosets cannot produce equal pairwise stabilizer subgroups), or that these conditions are both not satisfied. Our claim is verified by defining the low dimensional MIC states from subgroups of the fundamental group $G = \pi_1(M)$ of some manifolds encountered in our recent papers. e.g. the 3-manifolds attached to the trefoil knot and the figure-eight knot, and the 4-manifolds defined by 0-surgery of them. Exceptions to the aforementioned rule are classified in terms of geometric contextuality (which occurs when cosets on a line of the geometry do not all mutually commute).

MSC codes: 81P68, 51E12, 57M05, 81P50, 57M25, 57R65, 14H30

1. INTRODUCTION

Interpreting quantum theory is a long standing effort and not a single approach can exhaust all facets of this fascinating subject. Quantum information owes much to the concept of a (generalized) Pauli group for understanding quantum observables, their commutation, entanglement, contextuality and many other aspects, e.g. quantum computing. Quite recently, it has been shown that quantum commutation relies on some finite geometries such as generalized polygons and polar spaces [1]. Finite geometries connect to the classification of simple groups as understood by prominent researchers as Jacques Tits, Cohen Thas and many others [2, 3].

In the Atlas of finite group representations [4], one starts with a free group G with relations, then the finite group under investigation P is the permutation representation of the cosets of a subgroup of finite index d of G(obtained thanks to the Todd-Coxeter algorithm). As a way of illustrating this topic, one can refer to [3, Table 3] to observe that a certain subgroup of index 15 of the symplectic group $S'_4(2)$ corresponds to the 2QB (two-qubit) commutation of the 15 observables in terms of the generalized quadrangle of order two, denoted GQ(2,2) (alias the doily). For 3QB, a subgroup of index 63 in the symplectic group $S_6(2)$ does the job and the commutation relies on the symplectic polar space $W_5(2)$ [3, Table 7]. An alternative way to approach 3QB commutation is in terms of the generalized hexagon GH(2,2) (or its dual) which occurs from a subgroup of index 63 in the unitary group $U_3(3)$ [3, Table 8]. Similar geometries can be defined for multiple qudits (instead of qubits).

The straightforward relationship of quantum commutation to the appropriate symmetries and finite groups was made possible thanks to techniques developed by the first author (and coauthors) that we briefly summarize. This will be also useful at a later stage of the paper with the topic of magic state quantum computing.

The rest of this introduction recalls how the permutation group organizing the cosets leads to the finite geometries of quantum commutation (in Sec. 1.1) and how it allows the computation of magic states of universal quantum commutation (in Sec. 1.2). In this paper, it is shown that magic states themselves may be classified according to their coset geometry with two simple axioms (in Sec. 1.3).

1.1. Finite geometries from cosets [3, 6, 7]. One needs to define the rank r of a permutation group P. First it is expected that P acts faithfully and transitively on the set $\Omega = \{1, 2, \dots, n\}$ as a subgroup of the symmetric group S_n . The action of P on a pair of distinct elements of Ω is defined as $(\alpha,\beta)^p = (\alpha^p,\beta^p), p \in P, \alpha \neq \beta$. The orbits of P on $\Omega \times \Omega$ are called orbitals, and the number of orbits is called the rank r of P on Ω . The rank of P is at least two, and the two-transitive groups identify to the rank two permutation groups. Next, selecting a pair $(\alpha, \beta) \in \Omega \times \Omega$, $\alpha \neq \beta$, one introduces the two-point stabilizer subgroup $P_{(\alpha,\beta)} = \{p \in P | (\alpha,\beta)^p = (\alpha,\beta)\}$. There exist $1 < m \leq r$ such non-isomorphic (two-point stabilizer) subgroups S_m of P. Selecting one with $\alpha \neq \beta$, one defines a point/line incidence geometry \mathcal{G} whose points are the elements of Ω and whose lines are defined by the subsets of Ω sharing the same two-point stabilizer subgroup. Thus, two lines of \mathcal{G} are distinguished by their (isomorphic) stabilizers acting on distinct subsets of Ω . A non-trivial geometry arises from P as soon as the rank of the representation \mathcal{P} of P is r > 2, and simultaneously, the number of non isomorphic two-point stabilizers of \mathcal{P} is m > 2. Further, \mathcal{G} is said to be contextual (shows geometrical contextuality) if at least one of its lines/edges corresponds to a set/pair of vertices encoded by non-commuting cosets [7].

Figure 1 illustrates the application of the two-point stabilizer subgroup approach just described for the index 15 subgroup of the symplectic group is $S'_4(2) = A_6$ whose finite representation is

 $H = \langle a, b | a^2 = b^4 = (ab)^5 = (ab^2)^5 = 1 \rangle$. The finite geometry organizing the coset representatives is the generalized quadrangle GQ(2, 2). The other reresentation is in terms of the two-qubit Pauli operators, as first found in [1, 8]. It is easy to check that all lines not passing through the coset *e* contains some mutually not commuting cosets so that the GQ(2, 2) geometry is contextual. The embedded (3×3) -grid shown in bold (the so-called Mermin square) allows a 2QB proof of Kochen-Specker theorem [5].

1.2. Magic states in quantum computing. Now we recall our recent work about the relation of coset theory to the magic states of universal quantum computing. Bravyi & Kitaev introduced the principle of 'magic state distillation' [9]: universal quantum computation, the possibility of



FIGURE 1. The generalized quadrangle of order two GQ(2,2). The picture provides a representation in terms of the fifteen 2QB observables that are commuting by triples: the lines of the geometry. Bold lines are for an embedded 3×3 grid (also called Mermin square) that is a basic model of Kochen-Specker theorem (e.g. [3, Fig.1] or [5]). The second representation is in terms of the cosets of the permutation group arising from the index 15 subgroup of $G \cong A_6$ (the 6-letter alternating group).

getting an arbitrary quantum gate, may be realized thanks to stabilizer operations (Clifford group unitaries, preparations and measurements) and an appropriate single qubit non-stabilizer state, called a 'magic state'. Then, irrespective of the dimension of the Hilbert space where the quantum states live, a non-stabilizer pure state was called a magic state [10]. An improvement of this concept was carried out in [11] showing that a magic state could be at the same time a fiducial state for the construction of a minimal informationally complete positive operator-valued measure, or MIC, under the action on it of the Pauli group of the corresponding dimension. Thus UQC in this view happens to be relevant both to magic states and to MICs. In [11], a d-dimensional magic state is obtained from the permutation group that organizes the cosets of a subgroup H of index d of a two-generator free group G. This is due to the fact that a permutation may be realized as a permutation matrix/gate and that mutually commuting matrices share eigenstates - they are either of the stabilizer type (as elements of the Pauli group) or of the magic type. It is enough to keep magic states that are

MICHEL PLANAT[†], RAYMOND ASCHHEIM[‡], MARCELO M. AMARAL[‡] AND KLEE IRWIN[‡]

simultaneously fiducial states for a MIC because the other magic states may loose the information carried during the computation. A catalog of the magic states relevant to UQC and MICs can be obtained by selecting G as the two-letter representation of the modular group $\Gamma = PSL(2,\mathbb{Z})$ [12]. The next step, developed in [13, 14], is to relate the choice of the starting group G to three-dimensional topology. More precisely, G is taken as the fundamental group $\pi_1(S^3 \setminus L)$ of a 3-manifold M^3 defined as the complement of a knot or link L in the 3-sphere S^3 . A branched covering of degree d over the selected M^3 has a fundamental group corresponding to a subgroup of index d of $\pi_1(M^3)$ and may be identified as a sub-manifold of M^3 , the one leading to a MIC is a model of UQC. In the specific case of Γ , the knot involved is the left-handed trefoil knot $T_1 = 3^1$, as shown in [12] and [13, Sec. 2].

1.3. Coset geometry of magic states. The goal of the paper is to classify the magic states according to the coset geometry where they arise. We start from a non trivial subgroup H of index d of a free group G and we denote N the normal closure of H in G. As above, the coset organization in a subgroup H of G provides a group P of permutation gates whose common eigenstates are either stabilizer states of the Pauli group or magic states for universal quantum computing. A subset of magic states consists of MIC states associated to minimal informationally complete measurements.

It is shown in the paper that, in many cases, the existence of a MIC state entails that the two conditions (i) N = G and (ii) no geometry (a triple of cosets cannot produce equal pairwise stabilizer subgroups), or that these conditions are both not satisfied. Our claim is verified by defining the low dimensional MIC states from subgroups of the fundamental group $G = \pi_1(M)$ of manifolds encountered in our recent papers, e.g. the 3-manifolds attached to the trefoil knot and the figure-eight knot, and the 4-manifolds defined by 0-surgery of them.

Exceptions to the aforementioned rule are classified in terms of geometric contextuality (which occurs when cosets on a line of the geometry do not all mutually commute).

In section 2, one deals with the case of MIC states obtained from the subgroups of the fundamental group of Figure-of-Eight knot hyperbolic manifold and its 0-surgery. In section 3, the MIC states produced with the trefoil knot manifold and its 0-surgery are investigated.

2. MIC states pertaining to the Figure-of-Eight knot and its $0\text{-}\mathrm{surgery}$

We first investigate the relation of MIC states to the group geometrical axioms (i)-(ii) (or their negation) in the context of the Figure-of-Eight knot K4a1 (in Sec. 2.2) and its 0-surgery (in Sec. 2.1). The fundamental group of the complement of K4a1 in the 3-sphere $G = \pi_1(S^3 \setminus K4a1)$, and its connection to MICs, is first studied in [13, Table 2], below are new results and corrections.

2.1. Group geometrical axioms applied to the fundamental group $\pi_1(Y)$. The manifold Y defined by 0-surgery on the knot K4a1 is of special

interest as shown in [14, Sec 2] and references therein. The number of subgroups of index d of the fundamental group $\pi_1(Y)$ is as follows

 $\eta_d[\pi_1(Y)] = [1, 1, 1, 2, 2, 5, 1, 2, 2, 4, 3, 17, 1, 1, 2, 3, 1, 6, 3, 6, 1, 3, 1, 43, \ldots],$

where a bold number means that a MIC exists at the corresponding index.

In Table 1, one summarizes the check of our axioms (i) and (ii) applied to $\pi_1(Y)$. A triangle Δ means that a geometry does exist (corresponding to at least a triple of cosets with equal pairwise stabilizer subgroups), thus with (ii) is violated. According to our theory, for a MIC to exist, we should have (i) and (ii) satisfied, or both of them violated. The former case occurs for d = 9, 11 and 19. The latter case occurs for d = 6 where the geometry is that of the octahedron (with the 3-partite graph K(2, 2, 2)) and d = 20, where the geometry is encoded by the the complement of the line graph of the bipartite K(4, 5). In all of these five cases, a *pp*-valued MIC does exist.

For dimension 4, the bold triangle points out a violation since (i) is true and (ii) is false while the 2QB-MIC exists. In this case the geometry is the tetrahedron (with complete graph K_4) but not all cosets on a line/triangle are mutually commuting, a symptom of geometric contextuality, as shown in Fig. 2.



FIGURE 2. The contextual geometry associated to the 2QB-MIC and permutation group $P = A_4$ in Table 1. The line/triangle $\{a, ab, ab^{-1}\}$ is not made of mutually commuting cosets, thus geometric contextuality occurs.

2.2. Group geometrical axioms applied to the fundamental group $\pi_1(S^3 \setminus K4a1)$. The submanifolds obtained from the subgroups of index d of the fundamental group $\pi_1(S^3 \setminus K4a1)$ for the Figure-of-Eight knot complement are given in Table 2 (column 3), as identified in SnapPy [15] (this corrects a few mistakes of [13, Table 2]).

As for the subsection above, when axioms (i) and (ii) are simultaneously satisfied (or both are not satisfied), a MIC is created. Otherwise no MIC exist in the corresponding dimension, as expected.

There are three exceptions where (i) is true and a geometry does exist (when (ii) fails to be satisfied). This first occurs in dimension 4 with a 2QB MIC arising from the 3-manifold otet08₀₀₀₀₂, in this case geometric

MICHEL PLANAT[†], RAYMOND ASCHHEIM[‡], MARCELO M. AMARAL[‡] AND KLEE IRWIN[‡]

d	Р	(i)	pp	geometry
4	A_4	yes	2	2 QB MIC, Δ
5	10	yes		Δ
6	A_4	no	2	6-dit MIC, $K(2, 2, 2)$
9	$(36,9) \cong 3^2 \rtimes 4$	yes	2	2QT MIC
11	$(55,1) = 11 \rtimes 5, (\times 2)$	yes	3	11-dit MIC
16	$(48,3) \cong 4 \rtimes A_4$	yes		Δ
19	$(171,3) \cong 19 \rtimes 9$	yes	3	19-dit MIC
20	$(120,39) \cong 4 \rtimes (5 \rtimes (6,2))$	yes		$\overline{\mathcal{L}(K(4,5))}$

TABLE 1. Table of subgroups of the fundamental group $\pi_1[S^3 \setminus K4a1(0,1)]$ [with K4a1(0,1) the 0-surgery over the Figure-of-Eight knot]. The permutation group P organizing the cosets in column 2. If (i) is true, unless otherwise specified, the graph of cosets leading to a MIC is that of the d-simplex [and/or the condition (ii) is true: no geometry]. The symbol Δ means that (ii) fails to be satisfied. When there exists a MIC with (i) true and (ii) false, the geometry is shown in bold characters (here this occurs in dimension 4, see Fig. 2). If it exists, the MIC is pp-valued as given in column 4. In addition, K(2,2,2) is the binary tripartite graph (alias the octahedron) and $\overline{\mathcal{L}(K(4,5))}$ means the complement of the line graph of the bipartite graph K(4,5).

contextuality occurs in the cosets as in Fig. 2 of the previous subsection. Then, it occurs in dimension 7 (corresponding to 3-manifolds 00002 and 00014_{00035}) when the geometry of cosets is that of the Fano plane shown in Fig. 3a. Finally it occurs in dimension 10 when the geometry of cosets is that of a $[10_3]$ configuration shown in Fig. 3b [16, p 74].

In addition to the latter cases, false detection of a MIC may occur (this is denoted 'fd') in dimension 8 as shown in Table 2.



FIGURE 3. Contextual geometries associated to '(i) true and (ii) false' for the MICs of the Figure-of-Eight knot K4a1listed in Table 1: (a) the Fano plane related to the manifold otet14₀₀₀₀₂ at index 7, (b) the configuration [10₃] at index 10. The bold lines are for cosets that are not all mutually commuting. Each line corresponds to pair of cosets with the same stabilizer subgroup isomorphic to \mathbb{Z}_2^2 .

3. MIC STATES PERTAINING TO THE TREFOIL KNOT AND ITS 0-SURGERY

We now investigate the relation of MIC states to the group geometrical axioms (i)-(ii) (or their negation) in the context of the trefoil knot 3_1 (in Sec. 3.2) and its 0-sugery (in Sec. 3.1). The fundamental group of the complement of 3_1 in the 3-sphere $G = \pi_1(S^3 \setminus 3_1)$, and its connection to MICs, is studied in [13, Table 1] and below.

3.1. Group geometrical axioms applied to the fundamental group $\pi_1(E_8)$. The manifold \tilde{E}_8 is defined by 0-surgery on the trefoil knot 3_1 and is of special interest as shown in [14, Sec 3] and references therein. The number of subgroups of index d of the fundamental group $\pi_1(Y)$ is as follows

 $\eta_d[\tilde{E}_8] = [1, 1, 2, 2, 1, 5, 3, 2, 4, 1, 1, 12, 3, 3, 4, 3, 1, 17, 3, 2, 8, 1, 1, 27, 2, \ldots]$ where a bold number means that a MIC exists at the corresponding index.

Such cases are summarized in Table 3. As expected this occurs when the axioms (i) and (ii) are both true, or are both false. The latter case occurs at index 6 with geometry of the octahedron [and graph K(2, 2, 2)] and at index 15 with a geometry of graph K(5, 5, 5).

Exceptions to the rules are when a MIC exists with (i) true but not (ii). This occurs in dimension 3 (for the Hesse SIC) since the free group has a single generator (a trivial case), at index 4 (for the 2QB MIC) with a contextual geometry as in Fig. 2 and at index 21 with a contextual geometry (not shown) of graph K(3, 3, 3, 3, 3, 3, 3).

3.2. Group geometrical axioms applied to the fundamental group $\pi_1(S^3 \setminus 3_1)$. The characteristics of submanifolds obtained from the subgroups of index d of the fundamental group $\pi_1(S^3 \setminus 3_1)$ for the trefoil knot complement are given

d	ty	M^3 (or P)	ср	(i)	pp	geometry
2	cyc	$otet04_{00002}, m206$	1	no		
3	cyc	$otet 06_{00003}, s961$	1	no		
4	irr	otet $08_{00002}, L10n46, t_{12840}$	2	yes	2	2QB MIC, Δ
	cyc	$otet08_{00007}, t12839$	1	no		
5	cyc	$otet 10_{00019}$	1	no		
	irr	$otet 10_{00006}, L8a20$	3	yes		Δ
	$\operatorname{irr}(\times 2)$	$otet 10_{00026}$	2	yes	1	5-dit MIC
6	cyc	$otet 12_{00013}$	1	no		
	irr	$otet 12_{00039}$	1	no		
	$\operatorname{irr}(\times 2)$	$otet 12_{00038}$	1	yes	10	6-dit MIC
	irr	$otet 12_{00041}$	2	no		
	$\operatorname{irr}(\times 2)$	$otet 12_{00017}$	2	no		
	$\operatorname{irr}(\times 4)$	$otet 12_{00000}$	2	yes	2	6-dit MIC
7	cyc	$otet 14_{00019}$	1	no		
	$\operatorname{irr}(\times 4)$	$otet 14_{00002}, L14n55217$	3	yes	2	7-dit MIC, Δ : Fano
	$\operatorname{irr}(\times 4)$	$otet 14_{00035}$	1	yes	2	7-dit MIC, Δ : Fano
8	cyc	$otet 16_{00026}$	1	no		
	$\operatorname{irr}(\times 2)$	$0 tet 16_{00035}$	1	no		
	$\operatorname{irr}(\times 2)$	$0 tet 16_{00079}$	2	yes		fd
	$\operatorname{irr}(\times 2)$	$otet 16_{00016}$	2	yes		fd
	irr	$otet 16_{00092}$	2	no		
	irr	$otet 16_{00091}$	2	yes		16-cell
	irr	$otet 16_{00013}, L14n17678$	2	no		
9		$(36,9) \cong 3^2 \rtimes 4$		yes	2	2QT MIC
	$(\times 2)$	(504, 156) = PSL(2, 8)		yes	3	2QT MIC
	$(\times 2)$	$(216, 153) \cong 3^2 \rtimes (24, 3)$		yes	2	2QT MIC
10	$(\times 6)$	$(160, 234) \cong 2^4 \rtimes 10$		yes	5	10-dit MIC
	$(\times 2)$	$(120, 34) = S_5$		yes	4	10-dit MIC, $\boldsymbol{\Delta}: [\mathbf{10_3}]$
	$(\times 2)$	$(120, 34) = S_5$		no	7	10-dit MIC, 5-ortho
		$(360, 118) = A_6$		yes	5	10-dit MIC

MICHEL PLANAT[†], RAYMOND ASCHHEIM[‡], MARCELO M. AMARAL[‡] AND KLEE IRWIN[‡]

TABLE 2. Table of 3-manifolds M^3 found from subgroups of finite index d of the fundamental group $\pi_1(S^3 \setminus K4a1)$ (alias the d-fold coverings over the Figure-of-Eight knot 3-manifold). The covering type 'ty' in column 2, the manifold identification M^{3} , in column 3 and the number of cusps 'cp' in column 4 are from SnapPy [15]. For d = 9 and 10, SnapPy does not provide results so that we only identify the permutation group P = SmallGroup(o, k) (abbreviated as (o, k)), where o is the order and k is the k-th group of order o in the standard notation (that is used in Magma). If it exists, the MIC is 'pp'-valued. If (i) is true, unless otherwise specified, the graph of cosets leading to a MIC is that of the d-simplex [and/or the condition (ii) is true: no geometry]. The symbol Δ means that (ii) fails to be satisfied. When there exists a MIC with (i) true and (ii) false, the geometry is shown in bold characters. The symbol 'fd' means a false detection of a MIC when (i) and (ii) are satisfied simultaneously while a MIC does not exist. The abbreviations 'Fano', 'd-ortho' and ' $[10_3]$ ' are for the Fano plane, the d-orthoplex and the corresponding geometric configuration.

in Table 4 by using SnapPy [15] and Sage [17] for identifying the corresponding subgroup of the modular group Γ [12] (this improves [13, Table 1]).

GROUP GEOMETRICAL AXIOMS FOR MAGIC STATES OF QUANTUM COMPUTING

d	Р	(i)	pp	geometry
3	6	yes	1	Hesse SIC, Δ
4	A_4	yes	2	2QB MIC, $\boldsymbol{\Delta}$
6	A_4	no	2	6-dit MIC, $K(2, 2, 2)$
7	$(42,1) \cong 7 \rtimes (6,2)$	yes	2	7-dit MIC
9	$(54,5) \cong 3^2 \rtimes (18,3), (\times 2)$	yes		K(3,3,3)
12	$(72,44) \cong 2^2 \rtimes (18,3)$	yes		$\overline{\mathcal{L}(K(3,4))}$
13	$(78,1) \cong 13 \rtimes (6,2), (\times 2)$	yes	4	13-dit MIC
15	$(150, 6) \cong 5^2 \rtimes (6, 2), (\times 2)$	no	6	15-dit MIC, $K(5, 5, 5)$
16	$(96,72) \cong 2^3 \rtimes A_4$	yes		K(4, 4, 4, 4)
19	$(114,1) \cong 19 \rtimes (6,2)$	yes	3	19-dit MIC
21	$(126,9) \cong 7 \rtimes (18,3), (\times 2)$	yes	5	21-dit MIC, Δ: K(3,3,3,3,3,3,3)

TABLE 3. Table of subgroups of the fundamental group $\pi_1[S^3 \setminus 3_1(0,1)]$ [with $3_1(0,1)$ the 0-surgery over the trefoil knot] when the condition (i) is satisfied or when a MIC is missed. See the captions of Table 1 and 2 for the meaning of abbreviations.

As for the above sections, when axioms (i) and (ii) are simultaneously satisfied (or both are not satisfied), a MIC is created. Otherwise no MIC exist in the corresponding dimension, as one should expect.

There are a few exceptions where (i) is true and a geometry does exist (when (ii) fails to be satisfied). This first occurs in dimension 3 for the Hesse SIC where the free subgroup is trivial with a single generator. The next exceptions are for the 6-dit MIC related to the permutation group S_4 with the contextual geometry of the octahedron shown in Fig. 4a, in dimension 9 for the 2QT MIC related to the permutation group $3^3 \rtimes S_4$ with a contextual geometry consisting of three disjoint lines, and in dimension 10 for a 10-dit MIC related to the permutation group A_5 and the contextual geometry of the so-called Mermin pentagram. The latter geometry is known to allow a 3QB proof of the Kochen-Specker theorem [5].

4. Conclusion

Previous work about the relationship between quantum commutation and coset-generated finite geometries has been expanded here by establishing a connection between coset-generated magic states and coset-generated finite geometries. The magic states under question are those leading to MICs (with minimal complete quantum information in them). We found that, given an appropriate free group G, two axioms (i): the normal closure N of the subgroup of G generating the MIC is G itself and (ii): no coset-geometry should exist, or the negation of both axioms (i) and (ii), are almost enough to classify the MIC states. The few exceptions rely on configurations that admit geometric contextuality. We restricted the application of the theory to the fundamental group of the 3-manifolds defined from the Figure-of-Eight knot (an hyperbolic manifold) and from the trefoil knot, and to 4manifolds Y and \tilde{E}_8 obtained by 0-surgery on them. It is of importance to improve of knowledge of the magic states due to their application to quantum computing and we intend to pursue this research in future work.



FIGURE 4. Contextual geometries associated to '(i) true and (ii) false' for the MICs of the trefoil knot 3_1 listed in Table 2: (a) the octahedron, related to the subgroup $\Gamma_0(4)$ of Γ at index 6, (b) three disjoint lines K_3^3 at index 9, (c) The Mermin's pentagram at index 10. The bold lines are for cosets that are not all mutually commuting.

References

- M. Planat, Pauli graphs when the Hilbert space dimension contains a square: Why the Dedekind psi function? J. Phys. A Math. Theor. 2011, 44, 045301.
- [2] Thas, J.; van Maldeghem, H. Generalized polygons in finite projective spaces. In Distance-Regular Graphs and Finite Geometry, in Special Issue: Conference on Association Schemes, Codes and Designs, Proceedings of the 2004 Workshop on Distanceregular Graphs and Finite Geometry (Com 2 MaC 2004), Busan, Korea, 19–23 July 2004.
- [3] M. Planat and H. Zainuddin, Zoology of Atlas-Groups: Dessins D'enfants, Finite Geometries and Quantum Commutation, *Mathematics* (MDPI) 2017, 5, 6.
- [4] Wilson, R.; Walsh, P.; Tripp, J.; Suleiman, I.; Parker, R.; Norton, S.; Nickerson, S.; Linton, S.; Bray, J.; Abbott, R. ATLAS of Finite Group Representations, Version 3. Available online: http://brauer.maths.qmul.ac.uk/Atlas/v3/exc/TF42/ (accessed on June 2015).
- [5] Planat, M. On small proofs of the Bell-Kochen-Specker theorem for two, three and four qubits. *Eur. Phys. J. Plus* 2012, 127, 86.
- [6] M. Planat, A. Giorgetti, F. Holweck and M. Saniga, Quantum contextual finite geometries from dessins d'enfants, Int. J. Geom. Meth. in Mod. Phys. 12 1550067 (2015).
- [7] Planat, M. Geometry of contextuality from Grothendieck's coset space. Quantum Inf. Process. 2015, 14, 2563–2575.
- [8] M. Saniga and M. Planat, Multiple qubits as symplectic polar spaces of order two, Adv. Stud. Theor. Phys. 1, 1 (2007).
- [9] Sergey Bravyi and Alexei Kitaev, Universal quantum computation with ideal Clifford gates and noisy ancillas, *Phys. Rev.* A71 022316 (2005).
- [10] V. Veitch, S. A. Mousavian, D. Gottesman and J. Emerson, New J. of Phys. 16, Article ID 013009 (2014).
- [11] M. Planat and Z. Gedik, Magic informationally complete POVMs with permutations, R. Soc. open sci. 4 170387 (2017).

GROUP GEOMETRICAL AXIOMS FOR MAGIC STATES OF QUANTUM COMPUTING

- [12] M. Planat, The Poincaré half-plane for informationally complete POVMs, *Entropy* 20 16 (2018).
- [13] M. Planat, R. Aschheim, M. M. Amaral and K. Irwin, Universal quantum computing and three-manifolds, Universal quantum computing and three-manifolds Symmetry 10 773 (2018).
- [14] M. Planat, R. Aschheim, M. M. Amaral and K. Irwin, Quantum computing, Seifert surfaces and singular fibers, *Quantum Reports* 1 12-22 (2019).
- [15] M. Culler, N. M. Dunfield, M. Goerner, and J. R. Weeks, SnapPy, a computer program for studying the geometry and topology of 3-manifolds, http://snappy.computop.org.
- [16] B. Grünbaum, Configurations of points and lines (Graduate studies in mathematics, vol. 103, AMS, Providence, Rhode Island, 2009).
- [17] William A. Stein et al. Sage Mathematics Software (Version 8.6), The Sage Development Team, 2019, http://www.sagemath.org.

† UNIVERSITÉ DE BOURGOGNE/FRANCHE-COMTÉ, INSTITUT FEMTO-ST CNRS UMR
6174, 15 B AVENUE DES MONTBOUCONS, F-25044 BESANÇON, FRANCE.
E-mail address: michel.planat@femto-st.fr

‡ QUANTUM GRAVITY RESEARCH, LOS ANGELES, CA 90290, USA E-mail address: raymond@QuantumGravityResearch.org E-mail address: Klee@quantumgravityresearch.org E-mail address: Marcelo@quantumgravityresearch.org

d	ty	ср	Р	(i)	pp	type in Γ	geometry
2	cyc	1	$(2,1) \equiv 2$	no			
3	cyc	1	$(3,1) \equiv 3$	no			
	irr	2	$(6,1) \equiv 6$	yes	1	$\Gamma_0(2)$	Hesse SIC, Δ , L7n1
4	cyc	1	$(4,1) \equiv 4$	no			
	irr	2	$(12,3) = A_4$	yes	2	$\Gamma_0(3)$	2QB MIC, Δ , L6a3
	irr	1	$(24, 12) = S_4$	yes	2	$4A^0$	2QB MIC
5	cyc	2	$(5,1) \equiv 5$	no			
	irr	3	$(60,5) = A_5$	yes	1	$5A^0$	5-dit MIC
6	reg	3	$(6,1) \equiv 6$	no	2	$\Gamma(2)$	6-dit MIC, 6_3^3 [14]
	cyc	3	$(6,2) = 3 \times 2$	no		Γ'	
	irr	2	A_4	no	2	$3C^0$	6-dit MIC, $K(2,2,2)$
	irr	1	$(24,13) = 3 \rtimes 8$	no		$6B^0$	
	irr	1	$(18,3) \cong 3^2 \rtimes 2$	no		$6A^{0}$	
	irr	3	S_4	yes	2	$\Gamma_0(4)$	6-dit MIC, $\boldsymbol{\Delta}: \mathbf{octa}$
	irr	2	A_5	yes	2	$\Gamma_0(5)$	6-dit MIC
	irr	2	S_4	yes	2	$4C^{0}$	6-dit MIC, $\boldsymbol{\Delta}: \mathbf{octa}$
7	cyc	1	$(7,1)\equiv 7$	no			
	irr $(\times 2)$	2	$(42,1) \cong 7 \rtimes (6,2)$	yes	2	$NC(0, 6, 1, 1, [1^16^1])$	7-dit MIC
	irr $(\times 2)$	1	(168, 42) = PSL(2, 7)	yes	2	$7A^{0}$	7-dit MIC
	irr $(\times 2)$	2	S_7 (order 5040)	yes		$NC(0, 10, 1, 1, [2^15^1])$? 7-dit MIC
8	cyc	1	$(8,1) \equiv 8$	no			
	irr	2	(24, 13)	no		$6C^0$	
	irr	2	S_4	no		$4D^{0}$	
	irr $(\times 2)$	2	$(24,3) \cong 2.A_4$	yes			16-cell
	irr	2	PSL(2,7)	yes		$\Gamma_0(7)$	fd
	irr $(\times 2)$	1	SL(2,7)	yes		$NC(0, 8, 2, 2, [8^1])$	fd
	irr $(\times 2)$	2	$(48,29) \cong 2.(24,3)$	yes		$8A^0$	16-cell
9			$(9,1) \equiv 9$	no			
		2	(18,3)	no	7	$6D^0$	9-dit MIC, K(3,3,3)
		2	$(54,5) \cong 3^2 \rtimes (18,3)$	no	7	$NC(0, 6, 3, 0, [3^16^1])$	9-dit MIC, K(3,3,3)
		1	$(324, 160) \cong 3^3 \rtimes A_4$	no		$9A^0$	$K(3,3,3), K_3^3$
		3	(54,5)	yes	7	$NC(0, 6, 1, 0, [1^12^16^1])$	9-dit MIC
	$(\times 3)$		$(162,10) \cong 3^2 \rtimes 6$	yes			K(3,3,3)
	$(\times 2)$	1	(504, 156) = PSL(2, 8)	yes	3	$NC(1, 9, 1, 0, [9^1])$	2QT MIC
	$(\times 2)$	2	$(432,734) \cong 3^2 \rtimes (48,29)$	yes	2	$NC(0, 8, 3, 0, [8^11^1])$	2QT MIC
		3	$(648,703) \cong 3^3 \rtimes S_4$	yes	2	$NC(0, 12, 1, 0, [2^13^14^1])$	2QT MIC, $\boldsymbol{\Delta}: \boldsymbol{K_3^3}$
10		1	$(1\overline{20,35}) \cong 2 \rtimes \overline{A_5}$	no		$10A^{0}$	5-ortho
		2	A_5	yes	5	$5C^0$	10-dit MIC, Δ : MP
	$(\times 2)$	1	$(720,764) \cong A_6 \rtimes 2$	yes	9	$NC(0, 10, 0, 4, [10^1])\\$	10-dit MIC

MICHEL PLANAT†, RAYMOND ASCHHEIM‡, MARCELO M. AMARAL‡ AND KLEE IRWIN‡

TABLE 4. Subgroups of index d of the fundamental group $\pi_1(S^3 \setminus 3_1)$ (alias the d-fold coverings over the trefoil knot 3-manifold. The meaning of symbols is as in Table 2. When the subgroup in question is a subgroup of the modular group Γ , it is identified as a congruence subgroup or by its signature $NC(g, N, \nu_2, \nu_3, [c_i^{W_i}])$ (see [12] for the meaning of entries). The permutation group P =SmallGroup(o, k) is abbreviated as (o, k). As in Table 1, if (i) is true, unless otherwise specified, the graph of cosets leading to a MIC is the d-simplex [and/or the condition (ii) is true]. Exceptions (with geometry identified in bold characters) are for a MIC with (i) true and (ii) false. For index 9 and 10, some subgroups of large order could not be checked as leading to a MIC or not, they are not shown in the table. The abbreviation 'octa' is for the octahedron, 'MP' is for the Mermin pentagram and K_3^3 means three disjoint triangles.