

EXPANDING THE SAMPLE OF RADIO MINIHALOS IN GALAXY CLUSTERS

SIMONA GIACINTUCCI,¹ MAXIM MARKEVITCH,² ROSSELLA CASSANO,³ TIZIANA VENTURI,³
TRACY E. CLARKE,¹ RUTA KALE,⁴ AND VIRGINIA CUCITI³

¹*Naval Research Laboratory, 4555 Overlook Avenue SW, Code 7213, Washington, DC 20375, USA*

²*NASA/Goddard Space Flight Center, Greenbelt, MD 20771, USA*

³*INAF - Istituto di Radioastronomia, via Gobetti 101, I-40129 Bologna, Italy*

⁴*National Centre for Radio Astrophysics, Tata Institute of Fundamental Research, Pune University, Pune 411007, India*

(Received April 30, 2019; Revised June 12, 2019; Accepted June 12, 2019)

Submitted to ApJ

ABSTRACT

Radio minihalos are diffuse synchrotron sources of unknown origin found in the cool cores of some galaxy clusters. We use *GMRT* and *VLA* data to expand the sample of minihalos by reporting three new minihalo detections (A 2667, A 907 and PSZ1 G139.61+24.20) and confirming minihalos in five clusters (MACS J0159.8–0849, MACS J0329.6–0211, RXC J2129.6+0005, AS 780 and A 3444). With these new detections and confirmations, the sample now stands at 23, the largest sample to date. For consistency, we also reanalyze archival *VLA* 1.4 GHz observations of 7 known minihalos. We revisit possible correlations between the non-thermal emission and the thermal properties of their cluster hosts. Consistently with our earlier findings from a smaller sample, we find no strong relation between the minihalo radio luminosity and the total cluster mass. Instead, we find a strong positive correlation between the minihalo radio power and X-ray bolometric luminosity of the cool core ($r < 70$ kpc). This supplements our earlier result that most if not all cool cores in massive clusters contain a minihalo. Comparison of radio and *Chandra* X-ray images indicates that the minihalo emission is typically confined by concentric sloshing cold fronts in the cores of most of our clusters, supporting the hypothesis that minihalos arise from electron reacceleration by turbulence caused by core gas sloshing. Taken together, our findings suggest that the origin of minihalos should be closely related to the properties of thermal plasma in cluster cool cores.

Keywords: catalogs — galaxies: clusters: general — surveys — X-rays: galaxies: clusters — radio continuum: galaxies: clusters

1. INTRODUCTION

Galaxy clusters are filled with tenuous, hot X-ray emitting plasma, which is their dominant baryonic component. Over the cluster lifetime, it is heated by merger-generated shocks and turbulence, and cools via X-ray radiation, which occurs especially fast in the dense cluster cores. This plasma is permeated by tangled magnetic fields and relativistic particles, which contribute only a small fraction of the total energy density, but can strongly affect the plasma behavior as well as provide an interesting window into the cluster physics. The ultrarelativistic electrons in the intracluster medium (ICM) cool rapidly via synchrotron and inverse Compton radiation and have to be continuously energized. Thus, their synchrotron radio emission provides insight into the particle acceleration processes in clusters. Disregarding the radio galaxies often found in clusters, the diffuse radio emission that originates in the ICM comes as three broad phenomena, each apparently representing different acceleration mechanisms and energy sources (see, e.g., van Weeren et al. 2019 for a recent review). Peripheral radio relics are believed to be the result of acceleration at ICM shocks, while giant diffuse radio halos, typically found in disturbed clusters and extending over the whole cluster, are probably caused by electron reacceleration by merger-induced turbulence (e.g., Brunetti & Jones 2014).

The third phenomenon, radio minihalos, are confined to cluster cool cores (e.g., Giacintucci et al. 2017, hereafter G17). Minihalos are not directly powered by radio jets from the active galactic nucleus (AGN) harbored in the central cluster galaxy, nor they originate from diffusion, or other transport mechanisms, of relativistic electrons from the AGN. The radio-emitting electrons are instead generated *in situ* either as a by-product of hadronic collisions of cosmic-ray protons with thermal protons in the ICM, or reacceleration of seed relativistic elec-

trons by turbulence in the cool core (e.g., Gitti et al. 2002, Pfrommer & Ensslin 2004, Fujita et al. 2007, Keshet & Loeb 2010, Fujita & Ohira 2013, Zandanel et al. 2014, ZuHone et al. 2013, 2015; for a review see Brunetti & Jones 2014).

The distinction among these types of faint radio sources is not always clear in the radio data because of the finite instrument resolution and interferometric image quality, as well as projection effects and the presence of unrelated sources such as radio galaxies, but also because there is a continuum of source morphologies. A physically-motivated definition of a minihalo that we adopt is given in G17: it is an extended source in the cluster center that (1) does not consist of radio lobes or tails nor shows any morphological connection (such as jets) to the central cluster AGN, and (2) has a radius between 50 kpc and $0.2R_{500}$.¹ Smaller sources can plausibly be explained by diffusion from the central AGN (e.g., §6.4 in Giacintucci et al. 2014a) and $0.2R_{500}$ is the typical boundary between the cluster core, where non-gravitational processes (cooling, AGN and stellar feedback) become important, and the outer, simpler cluster region where density, temperature and pressure profiles of the ICM are self-similar (e.g., McDonald et al. 2017 and references therein).

Using a complete cluster sample, G17 showed that minihalos are rather exclusive to cool cores and that *most* massive ($M_{500} > 6 \times 10^{14} M_{\odot}$) clusters with cool cores possess minihalos. The cool cores often exhibit sloshing-driven X-ray cold fronts (e.g., Markevitch & Vikhlinin 2007). In sloshing cores with a minihalo, the radio emission is generally found to be co-spatial with the gas sloshing region (e.g., Mazzotta & Giacintucci 2008, Hlavacek-Larrondo et al. 2013, Giacintucci et al. 2014a,b, Gendron-Marsolais et

¹ R_{500} is the radius within which the cluster mean total density is 500 times the critical density at the cluster redshift.

al. 2017, this paper), suggesting a causal link between sloshing and minihalos – e.g., sloshing motions may amplify the magnetic fields (Keshet et al. 2010) and generate turbulence in the cool core, which in turn reaccelerates the relativistic electrons in minihalos (ZuHone et al. 2013). Gas turbulent motions, that are sufficiently strong for this scenario, have been measured by the *Hitomi* X-ray observatory in the cool core of the Perseus cluster (Hitomi Collaboration et al. 2016), which is host to a radio minihalo (Sijbring 1993, Burns et al. 1992, Gendron-Marsolais et al. 2017).

Even though most minihalos have active radio galaxies at their centers, it is unclear whether there is a relation between these AGNs and the surrounding diffuse radio emission. The lack of a morphological connection to the AGN (i.e., jets or lobes) in high-resolution radio images of minihalos, when available, suggests that the AGN does not directly replenish the minihalo with high-energy electrons. These AGNs could still be a source of electrons for the minihalos by providing seed particles (e.g., from aged, disrupted radio lobes) that are re-energized by turbulence (Cassano et al. 2008, ZuHone et al. 2013). Central AGNs are also a strong candidate source of the heating required to compensate for the otherwise catastrophic X-ray radiative cooling of the gas at the cluster centers (see McNamara & Nulsen 2007 for a review). The dissipation of the AGN-driven turbulence is one of the possible mechanisms by which energy of the AGN outbursts is transferred to the gas (Zhuravleva et al. 2014). Other proposed mechanisms include heating by cosmic rays (e.g., Guo & Oh 2008, Pfrommer 2013, Yang & Reynolds 2016, Ruszkowski et al. 2017) as well as gas sloshing, which can provide a net heat inflow by bringing the outer, higher-entropy gas into the cool core (ZuHone et al. 2010). If the radio minihalos are generated by turbulence and/or hadronic cosmic ray interactions, then a con-

nection between the non-thermal radio emission and the non-gravitational processes in cool cores (cooling and heating) may exist at a fundamental level (e.g., Fujita & Ohira 2013, Bravi et al. 2016, Jacob & Pfrommer 2017). This makes the minihalos a very interesting physical probe.

From an observational point of view, the detection and study of minihalos is complicated by a combination of their extent and low surface brightness and the presence of the often much brighter central radio galaxy. Sensitive, *low*-resolution radio observations with a good sampling of the interferometric *uv* plane, particularly at short antenna spacings (ideally, a full-synthesis single-dish imaging), are crucial to properly image the extended emission. At the same time, high-resolution images are needed to identify the emission associated with the central galaxy (as well as other possible radio galaxies in, or projected onto, the cluster core) and rule out the possibility of the large-scale diffuse emission being part of the radio galaxy. For most confirmed minihalos, radio images at different angular resolutions are available (e.g., Govoni et al. 2009, Giacintucci et al. 2014a,b). However, the existing images for a number of minihalo candidates do not sufficiently detail the morphology of the central galaxy and thus it is not possible to establish whether the AGN is connected to the surrounding extended emission.

In this paper, we present *Giant Metrewave Radio Telescope* (*GMRT*) images from our high-resolution follow-up of previously known candidate minihalos in MACS J0159.8-0849, MACS J0329.6-0211 and RXC J2129.6+0005 (Giacintucci et al. 2014a, henceforth G14a; Kale et al. 2015; hereafter K15). We present new *GMRT* and *Very Large Array* (*VLA*) images of the minihalos in AS 780 and A 3444 (Venturi et al. 2007, K15). Using *GMRT* data, we also report on the detection of new minihalos in the cool cores of A 2667, A 907 and PSZ1 G139.61+24.20. The latter is a newly

discovered *Planck* cluster, where a recent *LOFAR*² observation at 144 MHz has revealed faint ultra-steep spectrum radio emission, extending beyond the minihalo region (Savini et al. 2018). For consistency with our minihalo studies (G14a, G17, this paper), we also re-analyze archival, multi-configuration *VLA* observations at 1.4 GHz of 7 clusters, in which minihalos were reported in the literature — A1835, Ophiuchus, A2029, RBS797, RXJ1347.5–1145, MS1455.0+2232 and 2A0335+096 (Govoni et al. 2009, Gitti et al. 2006, 2007, Doria et al. 2012, Venturi et al. 2008, Mazzotta & Giacintucci 2008, Sarazin et al. 1995). Radio results presented here have been used for the G17 statistical sample study.

Using our larger sample of confirmed minihalos, we investigate correlations between the radio emission and the properties of the cluster

hosts, derived from *Chandra* X-ray data. Table 1 lists the minihalos used in our statistical analysis. We do not include clusters with a central extended radio source whose classification as a minihalo is still uncertain or unconfirmed (G14a, G17, van Weeren et al. 2019).

The paper is organized as follows: the radio observations are described in §2; our new radio images are presented in §3 and §4; the analysis of the *Chandra* archival X-ray data is described in §5, and a brief discussion of our results is given in §6. Details of our radio analysis of the previously known minihalos is presented in the Appendix.

We adopt a Λ CDM cosmology with $H_0=70$ km s⁻¹ Mpc⁻¹, $\Omega_m = 0.3$ and $\Omega_\Lambda = 0.7$. All errors are quoted at the 68% confidence level. For the radio spectral index α , we adopt $S_\nu \propto \nu^{-\alpha}$, where S_ν is the flux density at the frequency ν .

Table 1. List of minihalo clusters

| Cluster name | RA _{J2000} | DEC _{J2000} | z | scale |
|---------------------------------|---------------------|----------------------|-------|----------|
| | (h, m, s) | (° , ' , ") | | (kpc/'') |
| Clusters analyzed in this paper | | | | |
| MACSJ0159.8-0849 | 01 59 48.0 | −08 49 00 | 0.405 | 5.413 |
| MACSJ0329.6-0211 | 03 29 40.8 | −02 11 54 | 0.450 | 5.759 |
| RXCJ2129.6+0005 | 21 29 37.9 | +00 05 39 | 0.235 | 3.734 |
| AS780 | 14 59 29.3 | −18 11 13 | 0.236 | 3.746 |
| A3444 | 10 23 54.8 | −27 17 09 | 0.254 | 3.958 |
| A907 | 09 58 22.2 | −11 03 35 | 0.153 | 2.653 |
| A2667 | 23 51 40.7 | −26 05 01 | 0.230 | 3.674 |
| PSZ1G139.61+24.20 ^a | 06 22 13.9 | +74 41 39 | 0.267 | 4.102 |
| A1835 | 14 01 02.3 | +02 52 48 | 0.253 | 3.944 |
| Ophiuchus | 17 12 25.9 | −23 22 33 | 0.028 | 0.562 |

Table 1 continued

² Low Frequency Array, van Haarlem et al. (2013).

Table 1 (*continued*)

| Cluster name | RA _{J2000} (h, m, s) | DEC _{J2000} (° , ' , ") | z | scale (kpc/'') |
|--|----------------------------------|-------------------------------------|--------------------|-------------------|
| A 2029 | 15 10 55.0 | +05 43 12 | 0.077 | 1.451 |
| RBS 797 | 09 47 12.9 | +76 23 13 | 0.354 | 4.894 |
| RX J1347.5–1145 | 13 47 30.6 | −11 45 10 | 0.451 | 5.771 |
| MS 1455.0+2232 | 14 57 15.1 | +22 20 34 | 0.258 | 4.001 |
| 2A 0335+096 | 03 38 35.3 | +09 57 55 | 0.036 | 0.722 |
| Clusters with literature radio information | | | | |
| A 478 | 04 13 20.7 | +10 28 35 | 0.088 | 1.646 |
| ZwCl 3146 | 10 23 39.6 | +04 11 10 | 0.290 | 4.350 |
| RX J1532.9+3021 | 15 32 54.4 | +30 21 11 | 0.362 ^b | 5.048 |
| A 2204 | 16 32 45.7 | +05 34 43 | 0.152 | 2.643 |
| Perseus | 03 19 47.2 | +41 30 47 | 0.018 | 0.366 |
| RXC J1504.1–0248 | 15 04 07.5 | −02 48 16 | 0.215 | 3.494 |
| RX J1720.1+2638 | 17 20 08.9 | +26 38 06 | 0.164 | 2.814 |
| Phoenix ^c | 23 44 42.2 | −42 43 08 | 0.597 | 6.670 |

NOTE— Column 1: cluster name. Columns 2–4: cluster coordinates and redshift from the NASA/IPAC Extragalactic Database (NED). Column 5: angular to linear scale conversion. The literature radio information is from G14a, except for Perseus (Sijbring 1993), RXC J1504.1–0248 (Giacintucci et al. 2011), RX J1720.1+2638 (Giacintucci et al. 2014b) and Phoenix (van Weeren et al. 2014).

^a PSZ1 G139.61+24.20: Coordinates and redshift are from the *Planck* SZ cluster catalog (Planck Collaboration et al. 2014).

^b Crawford et al. (1999).

^c SPT-CL J2344-4243: coordinates and redshift are from McDonald et al. (2015).

2. RADIO OBSERVATIONS

In this section, we describe the new *GMRT* observations at 1.4 GHz and 1.3 GHz of the clusters MACS J0159.8-0849, MACS J0329.6-0211, RXC J2129.6+0005 and AS 780. For RXC J2129.6+0005 and AS 780, we complement these observations with higher-frequency data retrieved from the *VLA* archive. We also re-analyze archive observations of AS 780 (*GMRT*, 610 MHz), A 3444 (*VLA*, 1.4 GHz), A 907 (*GMRT*, 610 MHz) and A 2667 (*GMRT*, 610

MHz and 1.15 GHz). Finally, we present new *GMRT* data at 1.28 GHz and 610 MHz of the *Planck* cluster PSZ1 G139.61+24.20. Details on all these observations are provided in Table 2.

In Appendix A, we present our re-analysis of archival *VLA* observations at 1.4 GHz of A 1835, Ophiuchus, A 2029, RBS 797, RX J1347.5–1145, MS 1455.0+2232 and 2A 0335+096 (Table 8), known to possess a central minihalo at their center. New images and individual notes on these clusters are presented.

All data were reduced using the NRAO³ Astronomical Image Processing System (AIPS) package. A description of the *GMRT* and *VLA* data reduction is given in §2.1 and §2.2, respectively.

For all clusters, the flux densities of individual radio galaxies and minihalos are summarized in Tables 3 (Table 9 for the newly analyzed clusters with previously known minihalos). The position, size and flux density of the unresolved radio galaxies were determined by fitting the sources with a Gaussian model (JMFIT). For extended radio galaxies, we measured the total flux density by integration within the $+3\sigma$ surface brightness contour. As in G14a, the flux density of minihalos was obtained by measuring the total flux in circular regions of a radius that progressively increased from the radius encompassing the $+3\sigma$ isocontour until the integrated flux density reached saturation. The uncertainty on the minihalo flux density was estimated as described in G14a. For all minihalos, size (R_{MH} , measured as in G14a) and luminosity at 1.4 GHz are summarized in Table 4, which includes also 8 minihalos with literature information. The table also reports the 1.4 GHz luminosity of the central radio galaxy.

2.1. *GMRT* data reduction

The *GMRT* data were collected in spectral-line observing mode, using the software backend for all clusters (with the exception of A 907 and A 2667), which provides a total observing bandwidth of 33.3 MHz, subdivided in 256 channels. The observations of A 907 at 610 MHz and A 2667 at 1.15 GHz were taken with the old hardware correlator, using a 256-channel band of 16 MHz width at 1.15 GHz, and two 16 MHz-wide upper-side and lower-side bands (USB and LSB) at 610 MHz, each having 128 channels.

We accurately inspected the data and found that all observations were in part impacted by radio frequency interference (RFI). After an initial step of flux and bandpass calibration using standard calibration sources (3C48, 3C147 and 3C286), we used RFLAG to excise RFI-affected visibilities, followed by manual flagging to remove residual bad data. The flux and bandpass calibration was then recomputed and applied to the data. The flux densities of the primary calibrators were set in SETJY using the Perley & Butler (2013) coefficients. Phase calibrators, observed several times during the observation, were used to calibrate the data in phase. A number of phase-only self-calibration cycles and imaging were applied to the target visibilities to correct residual phase errors. To compensate for the non-complanarity of the array, we used wide-field imaging in each step of the self-calibration process and to produce the final images, decomposing the primary beam area into ~ 60 -80 smaller facets at 610 MHz and ~ 30 -50 facets at frequencies near 1 GHz. Finally, to improve the dynamic range of the final images, we used PEELR to “peel” problematic nearby bright sources, whose side lobes affected the area of the target. The final images were corrected for the *GMRT* primary beam response using PBCOR⁴. Residual amplitude errors are estimated to be within 5% at 1 GHz and 8% at 610 MHz (Chandra et al. 2004).

Table 2 provides the restoring beams and root mean square (rms) noise levels (1σ) of the final images, obtained using a “robustness” parameter (Briggs 1995) ROBUST=0 in IMAGR. For each cluster, we made sets of images with different angular resolution and weighting schemes, ranging from pure uniform weighting (ROBUST=−5) to natural weighting (ROBUST=+5), to enhance the extended emission. The higher-resolution images were

³ National Radio Astronomy Observatory.

⁴ www.ncra.tifr.res.in:8081/~ngk/primarybeam/beam.html

used to identify the radio source associated with the BCG and other possible radio galaxies in (or projected onto) the cluster core region. We also produced images by cutting the innermost region of the uv plane and using the remaining

long baselines ($\gtrsim 1 - 2 \text{ k}\lambda$) to better evaluate the contribution of the discrete sources. Low-resolution images were used to map the diffuse radio emission.

Table 2. Details of the radio observations

| Cluster | Array | Project | Frequency (GHz) | Bandwidth (MHz) | Date | Time (min) | FWHM, PA ("×", °) | rms ($\mu\text{Jy beam}^{-1}$) |
|--------------------|---------|---------|--------------------|--------------------|----------------|---------------|--------------------------|-------------------------------------|
| MACS J0159.8−0849 | GMRT | 28_077 | 1.39 | 33 | 2015 Aug 8 | 240 | 2.4×2.1 , 18 | 35 |
| MACS J0329.6−0211 | GMRT | 28_077 | 1.28 | 33 | 2015 Aug 7 | 123 | 2.7×2.1 , 61 | 23 |
| RXC J2129.6+0005 | GMRT | 28_077 | 1.30 | 33 | 2015 Jul 4 | 120 | 2.6×2.1 , 82 | 40 |
| | VLA−B | AH788 | 4.86 | 50 | 2002 Jul 07 | 60 | 1.3×1.2 , 10 | 22 |
| | VLA−A | AE177 | 8.46 | 50 | 1998 Apr 12 | 18 | 0.25×0.22 , −14 | 22 |
| AS 780 | GMRT | 07CRA01 | 0.61 | 32 | 2005 Jan 7 | 50 | 6.0×3.9 , 24 | 65 |
| | GMRT | 30_065 | 1.30 | 33 | 2016 Oct 1 | 76 | 3.2×1.8 , 33 | 83 |
| | VLA−A | AR517 | 8.46 | 50 | 2003 Jul 24 | 3 | 0.4×0.3 , −1 | 30 |
| A 3444 | VLA−BnA | AC696 | 1.44 | 50 | 2003 Oct 02 | 200 | 8.3×4.0 , 3 | 35 |
| | VLA−DnC | AC696 | 1.44 | 50 | 2004 May 30 | 260 | 42.5×30.2 , 58 | 50 |
| A 907 | GMRT | 15JHC01 | 0.61 | 32 | 2008 Dec 04 | 150 | 6.0×4.5 , −55 | 45 |
| A 2667 | GMRT | 10CRA01 | 0.61 | 16 ^a | 2006 Aug 27 | 80 | 7.3×4.3 , 2 | 50 |
| | GMRT | 13HBA01 | 1.15 | 16 | 2007 Nov 19,20 | 560 | 3.6×2.3 , 11 | 35 |
| PSZ1 G139.61+24.20 | GMRT | 28_077 | 0.61 | 33 | 2014 Oct 25 | 286 | 7.6×5.2 , 27 | 35 |
| | GMRT | 27_025 | 0.61 | 33 | 2015 Sept 4 | 280 | 6.3×3.7 , 44 | 70 |
| | GMRT | 30_065 | 1.28 | 33 | 2016 Aug 20 | 310 | 3.5×2.0 , 53 | 15 |

NOTE—Column 1: cluster name. Column 2: radio telescope (for the *VLA*, the array configuration is also reported). Column 3: project identifier. Columns 4–7: observation frequency, bandwidth, date and total time on source. Column 8: full-width half maximum (FWHM) and position angle of the array, obtained for ROBUST=0 in IMAGR. Column 9: image rms level (1σ).

^a The cluster was observed using both the USB and LSB, however, due to residual RFI and other errors affecting the LSB data, we chose to use only the USB data set.

2.2. *VLA data reduction*

The *VLA* data were calibrated and reduced following standard procedure. For each cluster, the observations in different array configurations were processed and imaged separately. Several loops of imaging and self-calibration were applied to each data set to reduce the effects of residual phase errors in the images. Tables 2 and 8 summarize radio beams and 1σ noise levels of the final images, made using a

ROBUST=0. When possible, data from different dates and configurations were combined together in the uv or image plane. As for the *GMRT* data, we used set of images with different angular resolution, weighting schemes and uv range to identify discrete radio galaxies and map the diffuse minihalos.

The flux density scale was set using standard calibration sources (3C48, 3C147 and 3C286) and the Perley & Butler (2013) coefficients. Residual amplitude errors are within 5%.

Table 3. Radio flux density and spectral index

| Cluster name | Source | ν (GHz) | S_ν (mJy) | α | Reference |
|-------------------|--------|----------------|------------------|------------------|-----------|
| MACS J0159.8–0849 | S1 | 1.39 | 34.0 ± 1.7 | -0.57 ± 0.04 | 1 |
| | | 1.40 | 35.0 ± 1.8 | | 2 |
| | | 8.46 | 94.8 ± 4.8 | | 2 |
| | MH | 1.39 | ~ 2 | \dots | 1 |
| | | 1.40 | 2.4 ± 0.2 | | 2 |
| | | | | | |
| MACS J0329.6–0211 | S1 | 1.30 | 3.9 ± 0.2 | \dots | 1 |
| | | 1.40 | 3.8 ± 0.2 | | 2 |
| | MH | 1.30 | 4.5 ± 0.4 | \dots | 1 |
| | | 1.40 | 3.8 ± 0.4 | | 2 |
| RXC J2129.6+0005 | S1 | 0.24 | 87 ± 5 | 0.77 ± 0.02 | 3 |
| | | 0.61 | 48.9 ± 2.5 | | 3 |
| | | 1.30 | 25.3 ± 1.3 | | 1 |
| | | 1.40 | 24 ± 1.2 | | 3 |
| | | 4.86 | 8.6 ± 0.3 | | 1 |
| | | 8.46 | 5.6 ± 0.2 | | 1 |
| | MH | 0.24 | 21.0 ± 1.6 | 1.2 ± 0.1 | 3 |
| | | 0.61 | 8.0 ± 0.7 | | 3 |
| | | 1.30 | 2.5 ± 0.2 | | 1 |
| | | 1.40 | 2.4 ± 0.2 | | 3 |
| AS 780 | S1 | 0.61 | 102 ± 8 | -0.17 ± 0.04 | 1 |
| | | 1.30 | 101 ± 5 | | 1 |
| | | 8.46 | 160 ± 8 | | 1 |
| | MH | 0.61 | 34 ± 2 | < 1.49 | 1 |
| | | 1.30 | > 11 | | 1 |
| A 3444 | S1 | 1.44 | 1.41 ± 0.07 | \dots | 1 |
| | MH | 1.44 | 12.1 ± 0.9 | \dots | 1 |
| A 907 | S1 | 0.61 | 117 ± 9 | \dots | 1 |

Table 3 continued

Table 3 (*continued*)

| Cluster name | Source | ν (GHz) | S_ν (mJy) | α | Reference |
|--------------------|--------|----------------|------------------|-----------------|-----------|
| | S2 | 0.61 | 42 ± 3 | ... | 1 |
| | MH | 0.61 | 34.9 ± 5.6 | ... | 1 |
| A 2667 | S1 | 0.61 | 20.1 ± 1.6 | 0.5 ± 0.1 | 1 |
| | | 1.15 | 14.8 ± 0.7 | | 1 |
| | S2 | 0.61 | 3.7 ± 0.3 | 0.7 ± 0.1 | 1 |
| | | 1.15 | 2.4 ± 0.1 | | 1 |
| | MH | 0.61 | 15.3 ± 1.2 | 1.0 ± 0.2 | 1 |
| | | 1.15 | 8.3 ± 0.7 | | 1 |
| PSZ1 G139.61+24.20 | S1 | 0.61 | 1.9 ± 0.2 | 1.05 ± 0.16 | 1 |
| | | 1.28 | 0.87 ± 0.05 | | 1 |
| | S2 | 0.61 | 0.25 ± 0.02 | 0.14 ± 0.02 | 1 |
| | | 1.28 | 0.18 ± 0.01 | | 1 |
| | MH | 0.14 | 12 ± 1.8 | 1.33 ± 0.08 | 4 |
| | | 0.61 | 2.3 ± 0.8 | | 1 |
| | | 1.28 | 0.65 ± 0.08 | | 1 |

NOTE—Column 1: cluster name. Column 2: radio source. Column 3: frequency. Column 4: flux density. Column 5: integrated spectral index, computed between the lowest and highest frequencies reported in the table. Column 6: reference for the radio flux. (1) this work. (2) G14a. (3) K15. (4) Savini et al. (2018).

Table 4. Properties of minihalos

| Cluster name | R_{MH} (kpc) | $P_{\text{MH}, 1.4 \text{ GHz}}$ ($10^{24} \text{ W Hz}^{-1}$) | $P_{\text{BCG}, 1.4 \text{ GHz}}$ ($10^{24} \text{ W Hz}^{-1}$) | Reference |
|---------------------------------|--------------------------|---|--|-----------|
| Clusters analyzed in this paper | | | | |
| MACS J0159.8–0849 | 90 | 1.40 ± 0.14 | 20.3 ± 0.1 | 2 |
| MACS J0329.6–0211 | 70 | 2.84 ± 0.30 | 2.8 ± 0.1 | 2 |
| PSZ1 G139.61+24.20 | 50 | $0.13 \pm 0.02^{\text{a}}$ | $< 0.007^{\text{a}}$ | 1 |
| A 907 | 65 | $0.9 \pm 0.2^{\text{b}}$ | $4.1 \pm 0.4^{\text{b}}$ | 1 |

Table 4 continued

Table 4 (*continued*)

| Cluster name | R_{MH} (kpc) | $P_{\text{MH}, 1.4 \text{ GHz}}$ ($10^{24} \text{ W Hz}^{-1}$) | $P_{\text{BCG}, 1.4 \text{ GHz}}$ ($10^{24} \text{ W Hz}^{-1}$) | Reference |
|--|--------------------------|---|--|-----------|
| AS 780 | 50 | $> 1.7^c$ | 17 ± 1^c | 1 |
| A 3444 | 120 | 2.4 ± 0.2 | 0.28 ± 0.01 | 1 |
| RX J2129.6+0005 | 80 | 0.40 ± 0.03 | 3.1 ± 0.2 | 6 |
| A 2667 | 70 | 1.1 ± 0.1^d | 2.1 ± 0.1^d | 1 |
| A 1835 | 240 | 1.19 ± 0.25 | 6.3 ± 0.3 | 1 |
| Ophiuchus | 250 | 0.11 ± 0.02 | 0.064 ± 0.003 | 1 |
| A 2029 | 270 | 0.28 ± 0.04 | 7.4 ± 0.4 | 1 |
| RBS 797 | 120 | 2.20 ± 0.24 | 7.6 ± 0.4 | 1 |
| RX J1347.5–1145 | 320 | 26.7 ± 2.1 | 22.7 ± 1.1 | 1 |
| MS 1455.0+2232 | 120 | 1.75 ± 0.23 | 0.96 ± 0.05 | 1 |
| 2A 0335+096 | 70 | 0.06 ± 0.01 | 0.058 ± 0.003 | 1 |
| Clusters with literature radio information | | | | |
| A 478 | 160 | 0.32 ± 0.06 | 0.60 ± 0.03 | 2 |
| ZwCl 3146 | 90 | 1.4 ± 0.3 | 0.8 ± 0.2 | 2 |
| RX J1532.9+3021 | 100 | 3.35 ± 0.17 | 7.0 ± 0.4 | 2 |
| A 2204 | 50 | 0.54 ± 0.05 | 3.7 ± 0.2 | 2 |
| Perseus | 130 | 2.18 ± 0.11 | 13.4 ± 0.1^e | 3 |
| RXC J1504.1–0248 | 140 | 2.70 ± 0.14 | 5.7 ± 0.3 | 4 |
| RX J1720.1+2638 | 140 | 5.33 ± 0.32 | 0.50 ± 0.02 | 5 |
| Phoenix | 230 | 9.6 ± 3.1^f | 73.9 ± 7.4^g | 7 |

NOTE—Column 1: cluster name. Column 2: average radius of the minihalo (estimated as in G14a). Column 3: radio luminosity of the minihalo at 1.4 GHz. Column 4: radio luminosity of the BCG at 1.4 GHz. Column 5: references. (1) this work. (2) G14a. (3) Sijbring (1993). (4) Giacintucci et al. (2011). (5) Giacintucci et al. (2014b). (6) K15. (7) van Weeren et al. (2014).

^a Extrapolated from 1.28 GHz (§4.3).

^b Extrapolated from 610 MHz (§4.1).

^c Extrapolated from 1.30 GHz (§3.4).

^d Extrapolated from 1.15 GHz (§4.2).

^e From NVSS.

^f Extrapolated from $S_{\text{MH}, 610 \text{ MHz}} = 17 \pm 5 \text{ mJy}$ assuming $\alpha = 1.0 \div 1.3$.

^g Extrapolated from $S_{\text{BCG}, 610 \text{ MHz}} = 87.9 \pm 9.0 \text{ mJy}$ assuming $\alpha = 0.6 \div 0.8$.

3. MINIHALO CONFIRMATION

In G14a, we suggested the presence of a minihalo in the clusters MACS J0329.6–0211 and

MACS J0159.8–0849, based on *VLA* 1.4 GHz images at $5''$ resolution. In K15, *GMRT* images at 235 MHz and 610 MHz revealed a central diffuse radio source in RXC J2129.6+0005, which was classified as a minihalo. A possible minihalo was reported at the center of AS 780 and A 3444 by Venturi et al. (2007) using *GMRT* observations at 610 MHz. However, due to the lack of images of all these clusters with a finer angular resolution, it was not possible to exclude the possibility that the observed extended emission is sustained directly by the central AGN through large-scale jets, rather than being an actual minihalo. Here, we present new high-resolution *GMRT* and *VLA* images that show no evidence of such jets (or other radio features on the minihalo scale directly connected to the AGN), thus confirming a minihalo in all these clusters.

3.1. MACS J0159.8–0849

MACS J0159.8–0849 is a relaxed, cool-core cluster at redshift $z = 0.405$, which is host to a candidate minihalo (G14a). Using the *GMRT* at 1.39 GHz, we imaged the cluster center at $2''$ resolution. Our image is shown in Figure 1. As a reference, the first contour of the *VLA* 1.4 GHz image of the minihalo from G14a is reported in magenta.

The radio source associated with the BCG (S1) is unresolved in our new image at 1.39 GHz. Its flux density is $S_{1.39\text{ GHz}} = 34.0 \pm 1.7$ mJy (Tab. 3), which is consistent with 35.0 ± 1.8 mJy measured by the *VLA* at 1.4 GHz. In the surroundings of the BCG, we detect only hints of the faint diffuse emission seen in the much higher sensitivity *VLA* image ($1\sigma = 15 \mu\text{Jy beam}^{-1}$). To enhance the extended emission, we made images at slightly lower resolution (down to $\sim 5''$, same as the *VLA* image), using values of ROBUST > 1 and tapering the *uv* data to weight down the long baselines. However, due to the higher noise in these images ($1\sigma \sim 40 \mu\text{Jy beam}^{-1}$), we were not able to map the dif-




Figure 1. MACS J0159.8–0849. *GMRT* image at 1.39 GHz (greyscale and black and white contours). The unresolved source S1 is associated with the BCG. The restoring beam is $2''.4 \times 2''.1$, in p.a. 18° and rms noise level is $1\sigma = 35 \mu\text{Jy beam}^{-1}$. Contours are 0.07, 0.14 (black), 0.28, 0.56, 1, 2, 4, 8, 16, and 32 (white) mJy beam^{-1} . Contours at $-0.07 \text{ mJy beam}^{-1}$ are shown as dashed. For a comparison, the first contour from the *VLA* 1.4 GHz image of G14a (their Figure 5) is reported in magenta. The contour level is $0.045 \text{ mJy beam}^{-1}$, for a $5''$ circular beam.

fuse emission around S1. On the other hand, integration over the area enclosed by the *VLA* isocontour in Figure 1 reveals an excess of ~ 2 mJy after subtraction of the flux of S1. Such flux excess is only $\sim 20\%$ less than the minihalo flux density measured on the *VLA* image ($2.4 \pm 0.2 \text{ mJy}$), thus confirming the presence of an extended component, that is not well imaged by our shallow *GMRT* observation.

Even though higher-sensitivity images are needed to adequately map the minihalo, the absence in Figure 1 of bright jets and/or lobes on the scale of the minihalo suggests that the central radio galaxy and outer extended emission are not morphologically connected. As a further note, the BCG is still unresolved at the sub-arcsecond resolution of a *VLA*-A config-




Figure 2. MACS J0329.6–0211. (a) *GMRT* contours at 1.28 GHz of the central unresolved radio galaxy (S1) and surrounding minihalo, overlaid on a *HST* ACS image (ID. 12452). The radio image has a restoring beam of $2''.7 \times 2''.1$, in p.a. 61° and rms noise level of $1\sigma = 23 \mu\text{Jy beam}^{-1}$. Contours are spaced by a factor of 2, starting from $+3\sigma$. Contours at -3σ are shown as dashed. (b) *GMRT* 1.28 GHz contours, same as in (a), overlaid on the *Chandra* combined X-ray image in the 0.5–4 keV band.

uration observation at 8.5 GHz (G14a). This indicates that any possible extended lobes/jets associated with S1 must be smaller than ~ 2 kpc.

3.2. MACS J0329.6–0211

MACS J0329.6–0211 is a distant ($z = 0.45$) relaxed cluster with a bright, low-entropy cool core (Cavagnolo et al. 2009). Figure 2 presents our new *GMRT* 1.28 GHz image at a resolution of $2''.7 \times 2''.1$, overlaid on the optical *Hubble Space Telescope* (*HST*) and X-ray *Chandra* images. The diffuse minihalo, first reported by G14a, is clearly detected in the cluster core region. It has a size of ~ 50 kpc in radius, which is comparable to its extent in the G14a *VLA* 1.4 GHz image at $5''$ resolution. No large-scale jets or plumes connect the diffuse emission to central radio galaxy (S1), which is unresolved, thus limiting its size to be less than 15 kpc. Observations at sub-arcsecond resolution are needed to reveal the details of its radio structure at smaller scales. We measure a flux density of $S_{1.28\text{GHz}} = 3.9 \pm 0.2$ mJy for the BCG

and $S_{1.28\text{GHz}} = 4.5 \pm 0.4$ mJy for the minihalo (Tab. 3).

3.3. RXC J2129.6+0005

RXC J2129.6+0005 is a relaxed cool-core cluster at $z = 0.235$. A central minihalo was reported by K15, based on a *GMRT* observation at 610 MHz. In Figure 3a, we present our new higher-resolution *GMRT* image at 1.30 GHz. The image detects only the innermost $r \sim 35$ kpc portion of the diffuse source seen at lower frequency (magenta contour). The extended emission encompasses a compact source associated with the BCG (S1). This latter is still point-like in a 4.86 GHz image at $1''$ resolution, obtained from reanalysis of archival *VLA* data (Table 2; image not shown here). The source structure is finally unveiled at 8.46 GHz (Figure 3b), which resolves the emission into a bright, compact component (possibly the core), coincident with the optical peak in the SDSS⁵ image,

⁵ Sloan Digital Sky Survey.




Figure 3. RXC J2129.6+0005. (a) *GMRT* image at 1.30 GHz (greyscale and black and grey contours) of the central radio galaxy (S1) and innermost part of the minihalo. The restoring beam is $2''.6 \times 2''.1$, in p.a. 82° and rms noise level is $1\sigma = 45 \mu\text{Jy beam}^{-1}$. Contours are spaced by a factor of 2 from $0.12 \text{ mJy beam}^{-1}$. Contours at $-0.12 \text{ mJy beam}^{-1}$ are shown as dashed. The first contour from the *GMRT* 610 MHz image of the minihalo from K15 (their Figure 2) is reported in magenta. The contour level is $0.33 \text{ mJy beam}^{-1}$, for a $11''.6 \times 10''.6$ beam. (b) *VLA* contours at 8.46 GHz of S1, overlaid on the r-band SDSS image. The radio image has a restoring beam of $0''.33 \times 0''.23$, in p.a. -9° . The rms noise level is $1\sigma = 22 \mu\text{Jy beam}^{-1}$. Contours are spaced by a factor of 2 from $0.06 \text{ mJy beam}^{-1}$. (c) *GMRT* 1.30 GHz contours, same as in (a), overlaid on the *Chandra* X-ray image in the 0.5–4 keV band.

and a fainter ~ 4 kpc jet pointing south-east. Hints of a much fainter counter jet are visible on the opposite side of the putative core. A comparison between the radio and X-ray *Chandra* images in Figure 3c shows that the diffuse emission detected at 1.30 GHz is centrally located and permeates the brightest part of the cluster core, as seen in other minihalo clusters.

The flux densities of S1 are summarized in Table 3, along with the measurements at lower frequencies by K15. Its spectral index is $\alpha = 0.77 \pm 0.02$ between 235 MHz and 8.46 GHz. For the minihalo, we measure a flux density of $S_{1.30\text{GHz}} = 2.5 \pm 0.2 \text{ mJy}$ in the region enclosed by the magenta contour in Figure 3a. A flux of $S_{1.4\text{GHz}} = 2.4 \pm 0.2 \text{ mJy}$, consistent with our measurement, was estimated by K15 from a comparison between the NVSS⁶ and FIRST⁷ images. The minihalo has an inferred spectral

index of $\alpha = 1.2 \pm 0.1$ between 235 MHz and 1.4 GHz.

3.4. AS 780

AS 780 is a relaxed cluster at $z = 0.236$ with a regular X-ray morphology on large scales. The X-ray surface brightness profile peaks at the BCG, and the gas temperature declines towards the center (e.g., G17), as typically observed in other cool-core clusters. Hints of diffuse radio emission around the central radio galaxy were reported at 610 MHz by Venturi et al. (2007). We have obtained new images at 610 MHz by re-analyzing the Venturi et al. data and compared them to higher-resolution images at 1.30 GHz and 8.46 GHz, as well as to optical (from Pan-STARRS-1⁸) and X-ray (from *Chandra*) images of the cluster central region (Figure 4). The 8.46 GHz image (white contours in panel b) shows an extended (~ 8 kpc) core-dominated and double-

⁶ NRAO VLA Sky Survey, Condon et al. (1998).

⁷ Faint Images of the Radio Sky at Twenty-cm, Becker et al. (1995).

⁸ Panoramic Survey Telescope and Rapid Response System, Chambers et al. (2016).




Figure 4. AS 780. (a) *GMRT* 1.30 GHz contours (cyan), overlaid on the *GMRT* image of the minihalo at 610 MHz (grey scale and black contours). The radio images have been restored with a $3'' \times 2''$ beam at 1.30 GHz and $6'' \times 4''$ at 610 MHz. Cyan contours start at $+3\sigma = 0.25 \text{ mJy beam}^{-1}$ and then scale by a factor of 2. Black contours are spaced by a factor of 2 between $3\sigma = 0.2 \text{ mJy beam}^{-1}$ and $1.6 \text{ mJy beam}^{-1}$. Contours at the -3σ level are shown as dashed. (b) *VLA*-A configuration contours at 8.46 GHz (white) of the cluster central radio galaxy S1 on the optical Pan-STARRS-1 *r* image. The angular resolution is $0''.4 \times 0''.3$. White contours start at $+3\sigma = 0.09 \text{ mJy beam}^{-1}$ and then scale by a factor of 2. Black contours show the radio emission detected by the *GMRT* at 1.30 GHz (a) and are spaced by a factor of 2 between $+3\sigma = 0.25 \text{ mJy beam}^{-1}$ and 16 mJy beam^{-1} . (c) *GMRT* 610 MHz contours from (a), superposed to the *Chandra* X-ray image in the 0.5-4 keV band. Radio contours start at $+3\sigma = 0.2 \text{ mJy beam}^{-1}$ and then scale by a factor of 2. (d) *Chandra* X-ray image, same as in (c). Arrows mark the position of two possible cold fronts.

jet radio source, coincident with the dominant optical galaxy and central X-ray point source in the *Chandra* image (panel *d*). The source is embedded in a region of diffuse emission at 1.30 GHz (black contours), which corresponds to the innermost part of the minihalo detected at 610 MHz (panel *a*). The size of the minihalo is ~ 50 kpc in radius at 610 MHz. As indicated by a comparison between the radio and X-ray emission (panels *c* and *d*), the minihalo is cospatial with the bright X-ray cool core and is edged by a pair of possible cold fronts, symmetric with respect to the cluster center.

The flux densities of the central radio galaxy S1 at all frequencies are summarized in Table 3. Its spectrum is inverted with $\alpha = -0.17 \pm 0.04$ between 610 MHz and 8.46 GHz. No high-resolution images at 1.4 GHz (e.g., from FIRST) are available for this source, therefore we estimate its radio luminosity at 1.4 GHz by extrapolating the 1.3 GHz flux density with $\alpha = -0.17 \pm 0.04$. We obtain $S_{\text{BCG}, 1.4 \text{ GHz}} = 102.3 \pm 0.3$ and $P_{\text{BCG}, 1.4 \text{ GHz}} = (17 \pm 1) \times 10^{24} \text{ W Hz}^{-1}$ (Table 4).

The minihalo has a flux of $S_{\text{MH}, 610 \text{ MHz}} = 34 \pm 2$ mJy. At 1.30 GHz, where only part of the minihalo is detected, we measure $S_{\text{MH}, 1.30 \text{ GHz}} = 11 \pm 1$ mJy, which has to be considered a lower limit to the actual flux of the minihalo at this frequency. This translates into a spectral index $\alpha < 1.45$ between 610 MHz and 1.30 GHz and an estimated flux density $S_{\text{MH}, 1.4 \text{ GHz}} > 10$ mJy, corresponding to a limit of $P_{\text{MH}, 1.4 \text{ GHz}} > 1.7 \times 10^{24} \text{ W Hz}^{-1}$ in radio power (Table 4).

3.5. *A 3444*

A 3444 is a relaxed cluster at $z = 0.254$ with a central, low-entropy cool core (G17). A possible central minihalo was reported in this cluster by Venturi et al. (2007), based on *GMRT* images at 610 MHz. In Figure 5, we present our *VLA* images at 1.4 GHz, compared to the cluster optical and X-ray emission. The BnA image (*a*) shows a central compact radio source (S1),

fully enveloped by a roundish minihalo containing no jets. The unresolved source is associated with the BCG and its size is < 20 kpc. In a higher-resolution image from recent *GMRT* observations at 1.28 GHz (Giacintucci et al. in preparation), the source is still pointlike, thus implying that its size must be even smaller (~ 8 kpc or less). The minihalo size reaches a radius of ~ 120 kpc in the combined DnC+BnA image in Figure 5*b*. The flux density of S1 is $S_{1.4 \text{ GHz}} = 1.41 \pm 0.07$ mJy and its radio power $P_{\text{BCG}, 1.4 \text{ GHz}} = (0.28 \pm 0.01) \times 10^{24} \text{ W Hz}^{-1}$. The surrounding minihalo has $S_{1.4 \text{ GHz}} = 12.1 \pm 0.9$ mJy and $P_{\text{MH}, 1.4 \text{ GHz}} = (2.38 \pm 0.17) \times 10^{24} \text{ W Hz}^{-1}$. A detailed spectral study of this minihalo, based on the *VLA* data presented here and new, deep *GMRT* observations at 1.28 GHz and 610 MHz will be presented in a future paper (Giacintucci et al. in preparation), where spatial correlation between the radio and X-ray emission is also investigated.

4. NEW MINIHALO DETECTIONS

Our analysis of *GMRT* data of the clusters A 907, A 2667 and PSZ1 G139.61+24.20 (Table 2) led to the detection of large-scale diffuse emission in their cool cores, which we classify as minihalos. In the following sections, we describe these newly discovered minihalos and summarize their properties.

4.1. *A 907*

We have reanalyzed an archival *GMRT* observation at 610 MHz (Table 2) of this relaxed, cool-core cluster at $z = 0.153$. In Figure 6*a*, we present a 610 MHz image, overlaid as contours an optical *r*-band image from Pan-STARRS1. The figure shows a radio point source (S1) at the center of a region of diffuse emission, about 70 kpc in radius, that we classify as a minihalo. The source S1 is identified with the cluster dominant galaxy. An extended head-tail source (S2) is also visible in the image. No redshift information is available for this object, however its radio




Figure 5. A3444. (a) VLA 1.4 GHz BnA-configuration image, restored with a $5''$ circular beam and overlaid on the optical Pan-STARRS-1 r image. The rms noise is $1\sigma = 35 \mu\text{Jy beam}^{-1}$. Contours start at $0.09 \text{ mJy beam}^{-1}$ and then scale by a factor of 2. Contours at $-0.09 \text{ mJy beam}^{-1}$ are shown as dashed. S1 is the central radio galaxy. (b) VLA 1.4 GHz combined DnC+BnA-configuration contours of the minihalo and central radio galaxy S1 (black cross), overlaid on the *Chandra* image in the 0.5-4 keV band. The radio image has been restored with a $11''$ circular beam. The rms noise is $1\sigma = 45 \mu\text{Jy beam}^{-1}$. Contours start at $0.13 \text{ mJy beam}^{-1}$ and then scale by a factor of 2. No contours at a level of $-0.13 \text{ mJy beam}^{-1}$ are present in the part of the image shown.

morphology, typical of dense environments, suggests that it is a cluster member. Panel *b* shows a comparison of the radio and *Chandra* X-ray emission. The minihalo fills a large fraction of the central cool core and appears to be bounded by a pair of apparent X-ray cold fronts, as it is often observed in minihalo clusters with similar sloshing features in their cores.

We measure a flux density of $S_{610 \text{ MHz}} = 117 \pm 9 \text{ mJy}$ for the central galaxy S1 and $S_{610 \text{ MHz}} = 42 \pm 3 \text{ mJy}$ for S2. The minihalo has $S_{610 \text{ MHz}} = 34.9 \pm 5.6 \text{ mJy}$ (Table 3). No high-resolution images at 1.4 GHz are available for this cluster. An estimate of the flux and radio luminosity at 1.4 GHz can be obtained by extrapolating the 610 MHz fluxes using a spectral index of $\alpha = 1.0 \div 1.3$ for the minihalo (e.g., G14a) and $\alpha = 0.6 \div 0.8$ for the central active radio galaxy (e.g., Condon 1992). We obtain $S_{1.4 \text{ GHz}} = 13.5 \pm 2.8 \text{ mJy}$ for the minihalo, which corresponds to a radio power of $P_{\text{MH}, 1.4 \text{ GHz}} = (0.9 \pm 0.2) \times 10^{24} \text{ W Hz}^{-1}$. For the BCG, we estimate $S_{1.4 \text{ GHz}} =$

$65 \pm 7 \text{ mJy}$ and $P_{\text{BCG}, 1.4 \text{ GHz}} = (4.1 \pm 0.4) \times 10^{24} \text{ W Hz}^{-1}$ (Table 4).

4.2. A 2667

A 2667 is a relaxed cluster at $z = 0.23$ with a bright, cool core (Cavagnolo et al. 2009). We imaged the central region of the cluster using two high-resolution observations at 1.15 GHz from the *GMRT* archive (Table 2). We also re-analyzed *GMRT* data at 610 MHz from Venturi et al. (2008). Our images are presented in Figure 7, compared with optical and X-ray images. Panel *a* shows an overlay of the radio contours from the combined 1.15 GHz data on the *HST* ACS image, revealing a bright unresolved radio source (S1), coincident with the cluster central galaxy, and a second point source (S2), associated with a member galaxy ($z = 0.234$; Covone et al. 2006). ATCA⁹ observations at 4.8 GHz and 8.6 GHz find S1 to be still unresolved at $\sim 1''$ resolution (Hogan et al. 2015), thus limit-

⁹ Australia Telescope Compact Array.




Figure 6. A 907. (a) *GMRT* 610 MHz contours, overlaid on the optical Pan-STARRS-1 *r*-band image (gray scale). The radio image has been restored with a $5''$ circular beam. The rms noise is $1\sigma = 40 \mu\text{Jy beam}^{-1}$ and contours are spaced by a factor of 2 starting from $0.12 \text{ mJy beam}^{-1}$. Contours at $-0.12 \text{ mJy beam}^{-1}$ are shown as dashed. S1 is associated with the cluster central galaxy and S2 is an extended, head-tail radio galaxy. (b) *GMRT* contours at 610 MHz, same as in (a), overlaid on the *Chandra* X-ray combined image in the 0.5-2 keV band. (c) *Chandra* X-ray combined image, same as in (b). Arrows mark the position of two possible X-ray cold fronts.




Figure 7. A 2667. (a) *GMRT* 1.15 GHz contours, overlaid on the optical *HST* ACS image (gray scale). The radio image has been restored with a circular $2''.5$ beam. The rms noise is $1\sigma = 35 \mu\text{Jy beam}^{-1}$. Contours are scaled by a factor of 2, starting from $0.12 \text{ mJy beam}^{-1}$. Contours at $-0.12 \text{ mJy beam}^{-1}$ are shown as dashed. Individual radio galaxies are labeled. (b) *GMRT* contours at 610 MHz, overlaid on the *Chandra* X-ray image in the 0.5-4 keV band. The radio image has been restored with a circular $7''$ beam. The rms noise is $1\sigma = 53 \mu\text{Jy beam}^{-1}$. Contours are scaled by a factor of 2, starting from $0.16 \text{ mJy beam}^{-1}$. Contours at $-0.16 \text{ mJy beam}^{-1}$ are shown as dashed. Individual radio galaxies are labeled. (c) *Chandra* X-ray image, same as in (b). White lines indicate two surface brightness edges, which may be cold fronts.

ing the spatial scale of any possible jets and/or lobes associated with the galaxy to be ~ 4 kpc or less. Diffuse radio emission, that we classify as a minihalo, is detected around S1. Panel *b* presents the radio contours at 610 MHz, overlaid on the *Chandra* image, which is also shown in panel *c* with arrows marking the position of two possible X-ray cold fronts. The minihalo covers an area of about 70 kpc in radius and appears to be contained within the sloshing region defined by the X-ray fronts.

The flux densities of S1 and S2 at 1.15 GHz and 610 MHz are summarized in Table 3. Their spectral indices are $\alpha = 0.5 \pm 0.1$ and $\alpha = 0.7 \pm 0.1$, respectively. We measure the minihalo flux density by subtracting the fluxes of S1 and S2 from the total emission. We obtain $S_{1.15\text{ GHz}} = 8.3 \pm 0.7$ mJy and $S_{610\text{ MHz}} = 15.3 \pm 1.2$ mJy. A spectral index $\alpha = 1.0 \pm 0.2$ is inferred for the minihalo. Using this spectral index, we estimate a minihalo flux density at 1.4 GHz of $S_{1.4\text{ GHz}} = 7.1 \pm 0.6$ mJy and a luminosity of $P_{\text{MH}, 1.4\text{ GHz}} = (1.1 \pm 0.1) \times 10^{24}$ W Hz $^{-1}$ (Table 4). For the central galaxy, we estimate $S_{1.4\text{ GHz}} = 13.4 \pm 0.7$ mJy and $P_{\text{BCG}, 1.4\text{ GHz}} = (2.1 \pm 0.1) \times 10^{24}$ W Hz $^{-1}$ using $\alpha = 0.5 \pm 0.1$.

4.3. PSZ1 G139.61+24.20

PSZ1 G139.61+24.20 is a massive, cool-core cluster at $z = 0.27$, that was discovered by *Planck* (Planck Collaboration et al. 2014). We observed the cluster with the *GMRT* at 1.28 GHz for approximately 5 hours and two times at 610 MHz for a total of 9.4 hours (Table 2). A high-resolution image at 1.28 GHz is shown as contours in Figure 8*a*, compared to an optical *r*-band image from Pan-STARRS1. A 610 MHz image at 7'' resolution is presented in Figure 8*b*. At 1.28 GHz, hints of extended emission are visible at the location of the central optical galaxy, which lacks a clear radio counterpart. An extended double radio source (S1) is detected at $\sim 15''$ from the center. The source has no visible optical counterpart in the Pan-

STARRS1 image, thus suggesting that it is a background, high-redshift radio galaxy. A second radio source (S2), located to the North-West, is associated with an optical galaxy with no redshift information.

At 610 MHz, the double source S1 blends with a central diffuse source, with a size of $r \sim 50$ kpc, that we classify as a minihalo (Figure 8*b*). An X-ray cold front, whose position is marked by arrows in panel *c*, was reported by Savini et al. (2018), indicating that the central cool gas is sloshing. The position of the cold front is roughly coincident with the northwestern boundary of the minihalo at 610 MHz, suggesting that its radio emission may be confined here by the front.

The minihalo has been also detected by Savini et al. (2018) at 610 MHz (using the same *GMRT* observations analyzed here) and at 144 MHz with LOFAR. The LOFAR image revealed the existence of a larger-scale diffuse component, which surrounds the cool core and its central and more compact minihalo. This second component is undetected at higher frequencies, implying a very steep radio spectrum ($\alpha > 1.7$), and has been interpreted as an ultra-steep spectrum radio halo, similar to those found in unrelaxed clusters without a central cool core (e.g., Brunetti et al. 2008, Macario et al. 2010). Savini et al. argued that a recent cluster collision with a smaller subcluster may have perturbed the cluster on the large scale, without destroying its central cool core, and generated a large scale radio halo. The same gravitational interaction may also have induced gas sloshing in the cool core and ensuing minihalo. This suggests that both a minihalo and a giant radio halo could at times co-exist in the same cluster (Storm et al. 2015, Venturi et al. 2017; Savini et al. 2018, 2019; van Weeren et al. 2019).

The flux densities of S1 and S2 are summarized in Table 3, along with their spectral indices. For the minihalo, we estimated a flux




Figure 8. PSZ1 G139.61+24.20. (a) *GMRT* 1.28 GHz contours at the resolution of $3'' \times 2''$, in p.a. 53° , overlaid on the *r*-band Pan-STARRS1 optical image. The rms noise of the radio image is $1\sigma = 15 \mu\text{Jy beam}^{-1}$. Contours are scaled by a factor of 2, starting from $0.05 \text{ mJy beam}^{-1}$. Contours at $-0.05 \text{ mJy beam}^{-1}$ are shown as dashed. S1 is an extended radio source with no optical identification on the POSS-II image. S2 is a radio source with an optical counterpart. (b) Contours from the combined image at 610 MHz, restored with a $7''$ circular beam, overlaid on a *Chandra* smoothed X-ray image in the 0.5–4 keV band. The rms noise of the radio image is $1\sigma = 30 \mu\text{Jy beam}^{-1}$. Contours are scaled by a factor of 2, starting from $0.08 \text{ mJy beam}^{-1}$. Crosses mark the position of the radio galaxies S1 and S2 (a). (c) *Chandra* X-ray image, same as in (b). White lines mark the position of an X-ray cold front (Savini et al. 2018).

density of $S_{610\text{MHz}} = 2.3 \pm 0.8 \text{ mJy}$. This value was obtained by subtracting the flux of S1 and S2, measured on the full $5''$ -resolution image (not shown here), from the total emission in Figure 8b. At 1.28 GHz, we used a $5''$ -resolution image (not shown here) to measure the minihalo flux and obtained $S_{1.28\text{GHz}} = 0.65 \pm 0.08 \text{ mJy}$. A flux density of 12 mJy (with 15% uncertainty) is reported at 144 MHz by Savini et al. (2018). A spectral index of $\alpha = 1.33 \pm 0.08$ is inferred between 144 MHz and 1.28 GHz, consistent with the index in the 144–610 MHz range (Savini et al. 2018).

The flux density of the minihalo at 1.4 GHz, extrapolated from the lower-frequency measurements with $\alpha = 1.33 \pm 0.08$, is $S_{1.4\text{GHz}} = 0.6 \pm 0.1 \text{ mJy}$. The corresponding luminosity at 1.4 GHz is $P_{\text{MH}, 1.4\text{GHz}} = (1.3 \pm 0.2) \times 10^{23} \text{ W Hz}^{-1}$ (Table 4). For the BCG, the non-detection at 1.28 GHz at a 3σ level of 0.035 mJy implies an upper limit to its 1.4 GHz radio power of $P_{\text{BCG}, 1.4\text{GHz}} < 7 \times 10^{21} \text{ W Hz}^{-1}$ (adopting $\alpha = 0.6 \div 0.8$). We note that, to date, this is

the only non-detection of radio emission from the BCG in a cluster that hosts a radio minihalo.

5. CHANDRA X-RAY ANALYSIS

We use high angular resolution *Chandra* observations to map the X-ray emitting gas in the cluster cores and compare its distribution to the radio minihalo images (Figs. 3–6, 10–16). In particular, we want to examine correlations of the minihalo radio luminosity at 1.4 GHz and the X-ray luminosity of the whole clusters and of their cores (§6). In the following sections, we describe the analysis of the *Chandra* observations that we have used. They are listed in Table 5 along with clean exposure times and the adopted Galactic absorption column density N_H (discussed below).

5.1. Data reduction

The Level-1 ACIS event files from the archive were reprocessed following the standard procedure (as described in Vikhlinin et al. 2005) us-

ing the *Chandra* Calibration Database (CALDB) 4.6.3. We excluded the background flare intervals and modeled the background as described in Markevitch et al. (2003). To model the detector + sky background, we used the CALDB blank-sky datasets appropriate for the date of each observation, projected to the sky and normalized using the ratio of the observed to blank-sky count rates in the source-free 9.5–12 keV energy band. Following Markevitch et al. (2000), we also subtracted the ACIS readout artifact (using the contributed tool `make_readout_bg`), particularly important in the presence of sharp X-ray brightness peaks in clusters that we study here. We used images in the 0.5–4 keV and 2–7 keV energy bands to detect the unrelated point sources and small-scale extended sources. These sources were masked out from the spectral analysis.

For each cluster, we obtained a background-subtracted, exposure-corrected image in the 0.5–4 keV band. For clusters with multiple ACIS observations, we first coadded the individual background-subtracted images and then divided the counts images by the sum of the corresponding exposure maps.

To derived the X-ray temperatures and luminosities, we extracted a spectrum of each cluster region of interest and generated the instrument responses (ARF and RMF) using calibration files version N0008 for the telescope effective area, N0006 for the CCD quantum efficiency, and N0009 for the ACIS time-dependent contamination model. Background spectra were

extracted for the same regions from the blank-sky datasets, normalized as described in §5.1. The cluster X-ray emission was modeled with an absorbed single-temperature APEC model in the the 0.7–7 keV energy band, with the heavy element abundance (same for all elements above He) relative to solar free to vary. The absorption column density N_H was initially allowed to vary; if the best-fit N_H value was found to be consistent with that from the Leiden/Argentine/Bonn (LAB) radio survey of Galactic HI (Kalberla et al. 2005), we then fixed it at the database value for subsequent analysis. Deviations from the radio survey values can be either real (because of nonuniformity of the column density on the cluster scale or additional molecular hydrogen) or caused by the residual inaccuracies of the ACIS contamination model. Using the best-fit N_H models such complications away and serves our purpose of deriving the cluster intrinsic X-ray luminosity and temperature. The choice of the lower energy boundary of 0.7 keV further limits the importance of such deviations. The adopted N_H values are listed in Table 5.

For several clusters, we combined multiple *Chandra* observations by fitting their spectra separately but simultaneously, with the X-ray model parameters tied together but using separate detector responses. For RXC J1504.1-0248, for which the observations had the same pointing position and roll angle and thus the same responses, we instead coadded the spectra.

Table 5. *Chandra* observations analyzed in this work

| Cluster name | Observation ID | Detector (ACIS) | Exposure (ksec) | N_H (10^{20} cm^{-2}) |
|--------------|-------------------|--------------------|--------------------|--|
| A 478 | 1669 | S | 36.3 | $31.03^{+0.18}_{-0.18}$ |

Table 5 continued

Table 5 (*continued*)

| Cluster name | Observation ID | Detector (ACIS) | Exposure (ksec) | N_H (10^{20} cm^{-2}) |
|--------------------|-------------------|--------------------|--------------------|--|
| MACS J0159.8–0849 | 6106 | I | 35.3 | 2.06* |
| MACS J0329.6–0211 | 3582 | I | 19.7 | 4.64* |
| ZwCl 3146 | 9371 | I | 38.1 | 2.46* |
| RX J1532.9+3021 | 1665,14009 | I,S | 9,88 | 2.30* |
| A 2204 | 7940 | I | 76.7 | $8.23^{+0.45}_{-0.39}$ |
| Perseus | 11715 | I | 70.5 | 13.6* |
| A 1835 | 6880 | I | 119 | 2.04* |
| Ophiuchus | 16627 | I | 35.0 | $35.7^{+0.26}_{-0.25}$ |
| A 2029 | 4977,6101 | S,I | 76.7,9.3 | 3.25* |
| RBS 797 | 2202,7902 | I,S | 11,7 38.4 | 2.27* |
| RXC J1504.1–0248 | 17197,17669,17670 | I | 29.8,28.5,44.6 | $14.35^{+4.6}_{-4.6}$ |
| RX J1347.5–1145 | 14407 | I | 60.1 | 4.60* |
| RXC J1720.1+2637 | 4361 | I | 22.0 | 3.36* |
| MS 1455.0+2232 | 4192 | I | 91.2 | 3.18* |
| 2A 0335+096 | 7939 | S | 49.8 | $23.18^{+0.27}_{-0.24}$ |
| A 907 | 3185,3205 | I | 47.7, 35.8 | 5.45* |
| A 2667 | 2214 | S | 9.3 | 1.73* |
| RX J2129.6+0005 | 9370 | I | 29.8 | 3.64* |
| AS 780 | 9428 | S | 39.4 | 7.39* |
| A 3444 | 9400 | S | 36.3 | 5.55* |
| PSZ1 G139.61+24.20 | 15139 | I | 17.9 | $13.11^{+1.7}_{-1.6}$ |
| Phoenix | 16545 | I | 59.9 | 1.52* |

NOTE—Column 1: cluster name; Column 2: observation identification number; Column 3: *Chandra* ACIS detector; Column 4: total clean exposure; Column 5: Galactic absorption column density adopted in this paper; values marked with * are from LAB (§5.1).

Table 6. Properties of cluster hosts

| Cluster name | M_{500} ($10^{14} M_{\odot}$) | R_{500} (Mpc) | $T_{X, R_{500}}$ (keV) | $T_{X, 0.15R_{500}}$ (keV) | $T_{X, 70 \text{ kpc}}$ (keV) | $L_{X, R_{500}}$ ($10^{44} \text{ erg s}^{-1}$) | $L_{X, 0.15R_{500}}$ ($10^{44} \text{ erg s}^{-1}$) | $L_{X, 70 \text{ kpc}}$ ($10^{44} \text{ erg s}^{-1}$) |
|-------------------|--------------------------------------|--------------------|---------------------------|-------------------------------|----------------------------------|--|--|---|
| ◇ † A 478 | $7.1^{+0.3}_{-0.4}$ | 1.3 | 6.4 ± 0.7 | 6.2 ± 0.6 | 5.2 ± 0.5 | 23.3 ± 3.0^a | 19.5 ± 2.3 | 7.8 ± 0.9 |
| MACS J0159.8–0849 | $6.9^{+0.9}_{-1.0}$ | 1.2 | 8.5 ± 0.5 | 7.5 ± 0.4 | 5.9 ± 0.3 | 40.6 ± 4.4 | 21.1 ± 2.2 | 9.6 ± 1.0 |

Table 6 continued

Table 6 (*continued*)

| Cluster name | M_{500} ($10^{14} M_{\odot}$) | R_{500} (Mpc) | $T_{X, R_{500}}$ (keV) | $T_{X, 0.15R_{500}}$ (keV) | $T_{X, 70 \text{ kpc}}$ (keV) | $L_{X, R_{500}}$ ($10^{44} \text{ erg s}^{-1}$) | $L_{X, 0.15R_{500}}$ ($10^{44} \text{ erg s}^{-1}$) | $L_{X, 70 \text{ kpc}}$ ($10^{44} \text{ erg s}^{-1}$) |
|----------------------------------|--------------------------------------|--------------------|---------------------------|-------------------------------|----------------------------------|--|--|---|
| ★ MACS J0329.6–0211 ^b | 4.9 ± 0.7 | 1.0 | 5.1 ± 0.3 | 5.1 ± 0.3 | 4.3 ± 0.3 | 27.2 ± 2.7 | 15.7 ± 1.7 | 9.5 ± 1.0 |
| ★ ZwCl 3146 | 6.7 ± 0.8 | 1.2 | 6.5 ± 0.1 | 5.9 ± 0.1 | 4.7 ± 0.1 | 45.6 ± 4.6 | 28.9 ± 3.0 | 13.8 ± 1.4 |
| ★ RX J1532.9+3021 | 4.7 ± 0.6 | 1.0 | 6.0 ± 0.3^c | 5.1 ± 0.1 | 4.5 ± 0.1 | 41.6 ± 4.5^c | 27.8 ± 2.8 | 15.9 ± 1.6 |
| ◇ † A 2204 | 8.0 ± 0.4 | 1.3 | 6.6 ± 0.2 | 6.3 ± 0.2 | 5.1 ± 0.1 | 35.0 ± 3.8 | 25.1 ± 2.7 | 13.1 ± 1.3 |
| ★ † Perseus | 6.1 ± 0.6 | 1.3 | 6.4 ± 0.1^d | 6.4 ± 0.2 | 4.1 ± 0.1 | 14.9 ± 1.5 | 10.4 ± 1.0^e | 4.4 ± 0.5 |
| A 1835 | $8.5^{+0.5}_{-0.6}$ | 1.3 | 7.7 ± 0.1 | 6.9 ± 0.1 | 5.6 ± 0.1 | 56.5 ± 5.7 | 34.5 ± 3.5 | 16.8 ± 1.7 |
| ★ † Ophiuchus | 12.4 ± 1.2 | 1.6 | 10.3 ± 0.2^d | 9.4 ± 0.1 | 8.8 ± 1.1 | 19.7 ± 2.0^f | 12.9 ± 1.3 | 3.0 ± 0.4 |
| ◇ † A 2029 | 6.8 ± 0.2 | 1.3 | 8.1 ± 0.2^g | 7.5 ± 0.1^h | 6.6 ± 0.1^h | 24.3 ± 2.4 | 15.0 ± 1.5^h | 6.3 ± 0.6^h |
| RBS 797 | $6.3^{+0.6}_{-0.7}$ | 1.2 | 6.7 ± 0.3^i | 6.1 ± 0.1^j | 5.3 ± 0.1^j | 41.9 ± 5.4^i | 32.2 ± 3.2^j | 18.9 ± 1.9^j |
| RXC J1504.1–0248 | 7.0 ± 0.6 | 1.3 | 5.9 ± 0.4 | 5.6 ± 0.4 | 4.9 ± 0.3 | 68.4 ± 7.0 | 52.1 ± 5.6 | 31.3 ± 3.3 |
| RX J1347.5–1145 | $10.6^{+0.7}_{-0.8}$ | 1.3 | 12.8 ± 0.5 | 12.7 ± 0.5 | 9.8 ± 0.4 | 147.3 ± 15.4 | 99.0 ± 10.0 | 40.9 ± 4.3 |
| ◇ † RX J1720.1+2638 | 6.3 ± 0.4 | 1.2 | 6.6 ± 0.1 | 6.1 ± 0.2 | 5.1 ± 0.1 | 17.2 ± 1.7 | 12.9 ± 1.3 | 6.2 ± 0.6 |
| ★ MS 1455.0+2232 | 3.5 ± 0.4 | 1.0 | 4.9 ± 0.1 | 4.8 ± 0.1 | 4.6 ± 0.1 | 21.1 ± 2.2 | 14.7 ± 1.5 | 8.6 ± 0.9 |
| † 2A 0335+096 | $2.3^{+0.2}_{-0.3}$ | 0.9 | 3.6 ± 0.1^k | 2.5 ± 0.1 | 2.3 ± 0.1 | 4.4 ± 0.5 | 3.4 ± 0.3 | 2.3 ± 0.2 |
| ◇ † A 907 | 5.2 ± 0.5 | 1.2 | 5.8 ± 0.1 | 5.4 ± 0.1 | 4.8 ± 0.1 | 12.1 ± 1.6 | 7.0 ± 0.7 | 2.9 ± 0.3 |
| ◇ † A 2667 | 6.8 ± 0.5 | 1.2 | 6.4 ± 0.2 | 5.5 ± 0.2 | 4.6 ± 0.2 | 28.8 ± 4.1 | 16.0 ± 1.7 | 6.2 ± 0.7 |
| ◇ † A 3444 | $7.6^{+0.5}_{-0.6}$ | 1.3 | 6.2 ± 0.2 | 5.7 ± 0.2 | 5.0 ± 0.2 | 28.3 ± 4.0 | 18.5 ± 1.9 | 8.9 ± 0.9 |
| RX J2129.6+0005 | 4.2 ± 0.6 | 1.1 | 6.4 ± 0.2 | 6.1 ± 0.2 | 5.1 ± 0.2 | 21.1 ± 2.1 | 11.2 ± 1.1 | 5.5 ± 0.5 |
| ◇ † AS 780 | 7.7 ± 0.6 | 1.3 | 5.2 ± 0.2 | 4.7 ± 0.1 | 3.9 ± 0.1 | 30.0 ± 6.2 | 10.9 ± 1.1 | 5.9 ± 0.6 |
| PSZ1 G139.61+24.20 | 7.1 ± 0.6 | 1.2 | 6.9 ± 0.5 | 7.2 ± 0.6 | 6.2 ± 0.5 | 25.2 ± 2.7 | 12.6 ± 1.4 | 5.6 ± 0.6 |
| Phoenix | 12.6 ± 2.0^l | 1.3 | 10.6 ± 0.4 | 10.0 ± 0.3 | 8.8 ± 0.4 | 143.7 ± 15.0 | 112.2 ± 11.3 | 62.6 ± 6.5 |

NOTE—Column 1: cluster name. Column 2: total mass within R_{500} from Planck Collaboration et al. (2014). For clusters marked with ★, M_{500} is from G17 and is estimated from the $M_{500} - T_X$ relation of Vikhlinin et al. (2009) using core-excised ($r > 70$ kpc) temperatures. Column 3: R_{500} , derived from M_{500} . Columns 4–6: cluster temperature measured within a radius corresponding to R_{500} , $0.15R_{500}$ and 70 kpc. For the clusters marked by a ◇, the total temperature was measured within radius smaller than R_{500} . A 478: $r = 315$ kpc; A 2204: $r = 360$ kpc; A 2029: $r = 390$ kpc; RX J1720.1+2638: $r = 860$ kpc; A 907: $r = 930$ kpc; A 2667: $r = 620$ kpc; A 3444: $r = 600$ kpc; AS 780: $r = 600$ kpc. Columns 7–9: bolometric X-ray luminosity measured within a radius corresponding to R_{500} , $0.15R_{500}$, and 70 kpc. For the clusters marked by a †, the R_{500} luminosity is from MCXC (see §??).

^a The MCXC luminosity was increased by factor 1.34 to account for the different N_H used in our *Chandra* analysis.

^b From ObsId 3582.

^c From ObsId 1665.

^d Ikebe et al. (2002).

^e See §??.

^f The MCXC luminosity was increased by factor 1.25 to account for the different N_H used in our *Chandra* analysis.

^g From ObsId 6101.

^h From ObsId 4977.

ⁱ From ObsId 2202.

^j From ObsId 7902.

^k David et al. (1993).

^l McDonald et al. (2015).

5.2. X-ray luminosities

For each cluster, we define 3 interesting regions: (1) the whole cluster, $r \leq R_{500}$, (2)

the core region, $r \leq 0.15R_{500}$ (which is in the $r = 140 - 240$ kpc range for our clusters), and (3) the central coolest core region, $r \leq 70$ kpc, which is within the cooling radius of all our clusters. R_{500} was derived from the total cluster masses M_{500} estimated from the Planck Sunyaev-Zeldovich (SZ) signal, which are available for 17 out of 23 clusters (Table 6). For the remaining clusters (marked by a \star), we used a mass estimate from G17, which is based on the $M_{500} - T_X$ relation of Vikhlinin et al. (2009), using X-ray measured, core-excised ($r > 70$ kpc) temperatures.

For these three regions, we first determined an X-ray luminosity in the 0.1–2.4 keV band for each cluster. For 12 clusters in the sample, *Chandra* covered R_{500} , and we determined the luminosity by fitting the spectrum in the respective region and converting its normalization to luminosity (in the XSPEC package). For 11 clusters marked with \dagger in Table 6, R_{500} was outside the *Chandra* field of view, so we adopted the 0.1–2.4 keV R_{500} luminosities from MCXC (Meta-Catalog of X-Ray Detected Clusters of Galaxies, Piffaretti et al. 2011), which is mostly based on the *ROSAT* All Sky Survey (RASS). For A478 and Ophiuchus, we corrected the MCXC luminosities by factors 1.34 and 1.25, respectively, to account for the different N_H derived from our *Chandra* fits, which were factors 2.0 and 1.8 higher than the database values adopted for the RASS catalogs. This makes the luminosities internally consistent with those for the central regions that were derived directly from *Chandra* observations. For Perseus, even the $0.15R_{500}$ radius is not fully covered by one *Chandra* ACIS-I observation, so we used the *Chandra* mosaic from Aharonian et al. (2016) to derive the luminosity within $0.15R_{500}$.

To convert the 0.1–2.4 keV luminosities to the more physically meaningful bolometric ones, we need gas temperatures, which we have derived from *Chandra* spectral fits in the same regions

for most clusters. The temperature fits are summarized in Table 6; their errors include the statistical uncertainty and a 10% uncertainty on the value of N_H . For Perseus, Ophiuchus and 2A 0335+096, *Chandra* coverage is too small to probe an average temperature within R_{500} , so we used average cluster temperatures from the literature instead, which come from instruments with a larger field of view (see Table 6). For Perseus and Ophiuchus, whose total mass is estimated from the $M_{500} - T_X$ relation (the 2A 0335+096 mass comes from Planck), this should result in slightly underestimated masses and R_{500} , because the cool core biases the average temperature low. This will not affect our conclusions in any significant way. For further 8 clusters, marked with \diamond in Table 6, the *Chandra* field of view is smaller than R_{500} but covers enough of it to derive a reliable temperature for the region, considering the steeply declining radial brightness profiles and the relatively small contribution of the missing emission to the spectrum. For these clusters, we use *Chandra* temperatures fitted in smaller regions given in the notes for Table 6. For Perseus, for simplicity, we used one ACIS-I pointing (Table 5) out of many available to derive a temperature for $r < 0.15R_{500}$, as it almost covers the whole region. The statistics of that data subset is more than sufficient. The above simplifications are adequate because we are using the temperatures only for the luminosity conversion.

For each cluster region, we converted the 0.1–2.4 keV luminosities to the bolometric luminosities (Table 6) using a conversion factor appropriate for the corresponding temperature (Boehringer et al. 2004). The luminosity errors in Table 6 include the uncertainty on the temperature and a 10% *ROSAT*–*Chandra* flux cross-calibration uncertainty, added in quadrature. The latter was evaluated by comparing the R_{500} 0.1–2.4 keV luminosities derived from *Chandra* and those determined from *ROSAT*

for the 10 clusters in our sample with both the *Chandra* and the MCXC measurement. We found that our *Chandra* luminosities are on average 6% higher than the RASS ones, with a scatter of $\sim 10\%$.

6. DISCUSSION

In this paper, we have presented new *GMRT* and *VLA* radio images of 8 cool-core clusters. Our images confirm the presence of radio minihalos in 3 of these clusters that had minihalo candidates, confirm previously-known minihalos in 2 clusters, and discover new minihalos in 3 clusters. For consistency with the present analysis and previous work (G14a, G17), we have also reanalyzed *VLA* 1.4 GHz archival observations of 7 clusters with known minihalos (Appendix A). Below, we briefly summarize our main radio results.

1. Our *GMRT* images of MACSJ0159.8-0849 (1.4 GHz), MACSJ0329.6-0211 and RXC J2129.6+0005 (1.3 GHz) detail the radio emission in the cluster center on angular scales $2''$ and up. This corresponds to a physical size of $\gtrsim 10$ kpc (7 kpc for RXC J2129.6+0005). We do not detect any well-defined jets, lobes or tails from the central AGN on this scale, which is considerably smaller than that of the surrounding diffuse minihalo seen in the lower-resolution images ($R_{\text{MH}} > 70$ kpc; G14a, K15). We thus confirm the presence of minihalos in these clusters (no morphological connection between the diffuse radio emission and the AGN).
2. We have presented new *GMRT* images at 1.28 GHz and 610 MHz of the minihalo in AS 780 (Venturi et al. 2017). The minihalo, which extends ~ 50 kpc in radius, appears to be confined within a pair of prominent sloshing cold fronts seen in the *Chandra* image. Our reanalysis of an archival *VLA* observation at 8.46 GHz

shows the cluster central galaxy to be a core-dominated, double-lobe radio source of 8 kpc in length with no morphological connection to the much larger minihalo.

3. We have presented *VLA* images at 1.4 GHz of the minihalo in A 3444, first reported by Venturi et al. (2007). The diffuse emission covers the central $r \sim 120$ kpc region of the cluster core and encloses a weak compact source ($\lesssim 8$ kpc) associated with the BCG. In a forthcoming paper (Giacintucci et al. in preparation), we will present new, deep *GMRT* observations at 1.28 GHz and 610 MHz, study the radio spectral properties of the minihalo and investigate possible spatial correlation between the radio and X-ray emission.
4. Using *GMRT* observations, we have detected new minihalos in the cores of A 2667, A 907 and PSZ1 G139.61+24.20. For the latter, a recent *LOFAR* observation at 144 MHz has shown the central minihalo to be surrounded by a larger-scale and fainter diffuse component, with a much steeper radio spectrum (Savini et al. 2018) — perhaps another example of minihalo-like emission inside a giant radio halo as possibly seen in A 2319 (Storm et al. 2015), A 2142 (Venturi et al. 2017) and RX J1720.1+2638 (Savini et al. 2019).

With the new minihalo detections and candidate verifications presented in this work, the total number of confirmed minihalos has now increased to 23 (Table 4). This allows us to revisit some correlations between non-thermal emission and thermal properties of the minihalo cluster hosts. We will consider some other correlations in the forthcoming work.

6.1. Radio minihalos and properties of the cluster hosts

A possible intrinsic relation between the non-thermal properties and global thermal properties of clusters with a central minihalo has been suggested by early works, based on a small number of minihalos ($\lesssim 10$, e.g., Cassano et al. 2008, Kale et al. 2013). More recently, K15 and Gitti et al. (2018) used a larger sample of minihalos to investigate the distribution of minihalo clusters in the $P_{\text{MH}, 1.4\text{ GHz}}-L_X$ plane and reported a positive correlation. The physical origin of this correlation might be a correlation of the radio power with the total cluster mass, as found for the giant radio halos (e.g., Cassano et al. 2013), because the cluster mass and X-ray luminosity correlate. However, in G14a, we did not find evidence for a correlation with the mass for 14 minihalos with available masses from the *Planck* SZ observations. We also did not find a strong correlation between the radio power and the cluster core-excised X-ray temperature, which is a better proxy for the cluster total mass than the X-ray luminosity (e.g., Vikhlinin et al. 2009).

With our new minihalo detections, we can explore these possible correlations further. Our statistical analysis will include 22 confirmed minihalos, excluding AS 780, which has only a lower limit for the minihalo 1.4 GHz luminosity, but we show it in the plots. To evaluate the presence of any possible correlation and its strength, we use the Spearman’s test, which computes a rank correlation coefficient, r_s , whose value can range from -1 (perfect anticorrelation) through 0 (no correlation) to 1 (perfect correlation). The test also returns the probability *probrs* to get the observed r_s value by chance under the null hypothesis of no correlation (false positive). In Table 7, we summarize the values of r_s and *probrs* found for the relations analyzed in the following paragraphs.

6.1.1. Radio luminosity–cluster mass

In Fig. 9a, we compare the radio luminosity (Table 4) and the total cluster mass within R_{500}

(Table 6) for our 23 minihalo clusters. The distribution in this plot is scattered and no obvious correlation is visible. The Spearman test gives a rank correlation coefficient of $r_s = 0.06$ and *probrs* = 0.79, which indicates lack of correlation with the cluster mass, consistently with the G14a finding from a smaller sample.

6.1.2. Luminosity relations

In line with previous studies (e.g., Gitti et al. 2018), we find a positive correlation ($r_s = 0.53$) between the minihalo radio power and cluster total X-ray bolometric luminosity measured within R_{500} (Fig. 9b). In cool-core clusters, such as the minihalo hosts, a large fraction of the total X-ray luminosity is found to arise from the central $0.15R_{500}$ region (e.g., Maughan et al. 2012), which is also the region where the minihalo is located. It is therefore interesting to check if the correlation seen in Fig. 9b is driven by a correlation with the X-ray luminosity of the cluster core. In Figs. 9c and d, we compare the minihalo radio power to the X-ray bolometric luminosity measured within the central $r = 0.15R_{500}$ (which is in the $r = 140 - 240$ kpc range for our clusters) and the smaller $r = 70$ kpc regions, respectively.

The plot with the $0.15R_{500}$ X-ray luminosity (c) is rather similar to the plot with the total X-ray luminosity (b), and we find a similar rank correlation coefficient, $r_s = 0.56$. This is not surprising, since for our sample of strongly peaked clusters, the core/total luminosity ratio, $L_{X, 0.15R_{500}}/L_{X, R_{500}}$, ranges from 0.4 (AS 780) to 0.8 (A 478), with an average value of 0.6. Thus, the central $r = 0.15R_{500}$ core region is the dominant contributor to the cluster total luminosity for most clusters in our sample, explaining the similarity of diagrams b and c.

In Fig. 9d, we cut an even smaller region of the core and plot the radio luminosity vs. the X-ray bolometric luminosity within the central $r = 70$ kpc, which is within the cooling radius for all our clusters. This correlation is tighter, with a




Figure 9. (a) Comparison between the 1.4 GHz radio luminosity of minihalos and total mass within R_{500} of their cluster hosts. (b,c,d) Comparison between the 1.4 GHz radio luminosity of minihalos and the cluster X-ray bolometric luminosity within R_{500} (a), $0.15R_{500}$ (b), and $r = 70 \text{ kpc}$ (c). Blue lines in (c) are the best-fit relations derived using the BCES bisector (solid line) and orthogonal (dashed line) regression algorithms (Eq. 6.1.2). (e) The minihalo 1.4 GHz luminosity is compared to the core-excised X-ray bolometric luminosity, a proxy for the cluster mass. (f) Comparison between the 1.4 GHz luminosity of minihalos and 1.4 GHz luminosity of the BCG.

Table 7. Spearman’s Rank Correlation Coefficient test

| Relation | r_s | $probs$ |
|---|-------|---------|
| $P_{\text{MH}, 1.4 \text{ GHz}} - M_{500}$ | +0.06 | 79% |
| $P_{\text{MH}, 1.4 \text{ GHz}} - L_{\text{X}, R_{500}}$ | +0.53 | 11% |
| $P_{\text{MH}, 1.4 \text{ GHz}} - L_{\text{X}, 0.15 R_{500}}$ | +0.56 | 6% |
| $P_{\text{MH}, 1.4 \text{ GHz}} - L_{\text{X}, 70 \text{ kpc}}$ | +0.67 | 0.7% |
| $P_{\text{MH}, 1.4 \text{ GHz}} - (L_{\text{X}, R_{500}} - L_{\text{X}, 70 \text{ kpc}})$ | +0.44 | 4% |
| * $P_{\text{MH}, 1.4 \text{ GHz}} - (L_{\text{X}, R_{500}} - L_{\text{X}, 0.15 R_{500}})$ | +0.43 | 4% |
| $P_{\text{MH}, 1.4 \text{ GHz}} - P_{\text{BCG}, 1.4 \text{ GHz}}$ | +0.43 | 5% |

NOTE. Column 1: relation (Fig. 9; * figure not shown). Column 2: Spearman’s rank correlation coefficient, a non-parametric measure of the statistical dependence between two variables. Column 3: probability of uncorrelated quantities producing datasets that have a Spearman correlation at least as extreme as the one computed from these datasets.

higher correlation coefficient, $r_s = 0.67$. The probability of the null hypothesis is less than 1%. For this relation, we also show the best-fit correlations in the form

$$\log(P_{\text{MH}, 1.4 \text{ GHz}}/10^{24}) = B \times \log(L_{\text{X}, 70 \text{ kpc}}/10^{44}) + A \quad (1)$$

derived by implementing the bivariate correlated error and intrinsic scatter (BCES) bisector (blue solid line) and orthogonal (blue dashed line) regression algorithms (Akritas & Bershady 1996). The best-fit parameters are $A = -1.60 \pm 0.23$ and $B = 1.71 \pm 0.21$ (bisector) and $A = -1.96 \pm 0.31$ and $B = 2.08 \pm 0.34$ (orthogonal).

In Fig. 9e we plot the radio luminosity as a function of the core *excised* ($r > 70 \text{ kpc}$) X-ray bolometric luminosity within R_{500} , which can be used as a proxy for the cluster total mass. This relation is weaker, with a correlation coefficient of $r_s = 0.44$, in agreement with the lack of

a strong scaling with the SZ-estimated cluster mass (a). A similar diagram (not shown here) is obtained if we excise the central $0.15 R_{500}$ region instead of the smaller $r = 70 \text{ kpc}$ region, with similar Spearman correlation coefficient and null-hypothesis probability (Table 7).

Finally, in Fig. 9f we compare the radio luminosity of the minihalo to the radio luminosity of the central BCG (Table 4). A possible trend is visible, in line with previous analysis by Govoni et al. (2009) and G14a, but it is relatively weak ($r_s = 0.43$, $probs = 5\%$ ¹⁰), indicating that the AGN activity and minihalo emission are not tightly correlated. Interestingly, this correlation is weaker than that with the $r < 70 \text{ kpc}$ core X-ray luminosity, suggesting that the properties of the core gas may have a more direct effect on the minihalos than the AGN output.

Our minihalo sample is not statistically complete. Analysis of selection effects is beyond the scope of this paper, however we checked for the possibility of the luminosity-luminosity correlations being dominated by effects originating from flux selection. The corresponding radio flux–X-ray flux diagrams show strong positive correlations reflecting the large range of distances, but do not show any obvious flux-selection effects. This does not affect our main result of a better correlation of the minihalo radio power with the cool-core luminosity compared to the total luminosity, which is derived from the same sample and thus not driven by selection effects.

Our finding of a significant correlation between the minihalo radio power and the X-ray luminosity of the cool core is consistent with earlier findings by Gitti et al. (2004, 2012) and

¹⁰ The test was done using 21 minihalos: in addition to AS 780, we also excluded PSZ1 G139.61+24.20, which has only a lower limit for the BCG luminosity at 1.4 GHz.

Bravi et al. (2016), who considered a correlation with the “cooling flow power” — which by their derivation is very close to the X-ray luminosity of the cooling region. Since we know that some physical process in the cluster cores compensates for X-ray radiative cooling (e.g., McNamara & Nulsen 2012), that process has to have the power very similar to the X-ray bolometric luminosity of the core, which is what cools the gas. Among the popular processes are the turbulence generated by AGN activity (Zhuravleva et al. 2014) and cosmic rays generated by the AGN (e.g., Guo & Oh 2008, Pfrommer 2013, Yang & Reynolds 2016, Ruszkowski et al. 2017). A further process is gas sloshing, which can provide a net heat inflow by bringing the outer, higher-entropy gas into the cool core (ZuHone et al. 2010). It is possible to make the conjecture that the same heating process is responsible for the radio minihalos. However, in addition to the minihalo power, any such hypothesis would need to explain other properties of the minihalos, in particular, their apparent close relation to the sloshing cold fronts (e.g., Mazzotta & Giacintucci 2008; ZuHone et al. 2013; Giacintucci et al. 2014b).

7. SUMMARY AND CONCLUSIONS

We have presented new *GMRT* and *VLA* images of 8 cool-core clusters. We have found new minihalos in A 2667, A 907 and PSZ1 G139.61+24.20, confirmed the candidates in MACS J0159.8–0849, MACS J0329.6–0211 and RXC J2129.6+0005, and presented new radio images of the previously-known minihalos AS 780 and A 3444. For consistency with our radio analysis, we have also reanalyzed archival

VLA 1.4 GHz observations of 7 cool-core clusters with known minihalos. Combining our new detections and confirmations with the previously known minihalos results in a total sample of 23 confirmed minihalos, the largest sample to date. We revisited possible correlations between non-thermal emission and thermal properties of their cluster hosts. We confirmed the lack of a strong relation between the minihalo radio power and the total cluster mass. Instead, we found a relatively strong positive correlation between the minihalo radio power and the X-ray bolometric luminosity of the cool core ($r < 70$ kpc). This supplements our earlier finding (G17) that most if not all cool cores in massive clusters contain radio minihalos. In many cases, we observe that the minihalo may be confined by the sloshing cold fronts in the core gas. Taken together, these findings suggest that the origin of the minihalos should be closely related to the properties of the cool-core gas.

Acknowledgements.

Basic research in radio astronomy at the Naval Research Laboratory is supported by 6.1 Base funding. RK acknowledges support through the DST-INSPIRE Faculty Award by the Government of India. We thank the staff of the GMRT that made these observations possible. GMRT is run by the National Centre for Astrophysics of the Tata Institute of Fundamental Research. The National Radio Astronomy Observatory is a facility of the National Science Foundation operated under cooperative agreement by Associated Universities, Inc.

REFERENCES

- | | |
|---|---|
| <p>Aharonian, F. A., Akamatsu, H., Akimoto, F., et al. 2017, <i>ApJL</i>, 837, L15</p> <p>Akritas, M. G., & Bershad, M. A. 1996, <i>ApJ</i>, 470, 706</p> | <p>Becker, R. H., White, R. L., & Helfand, D. J. 1995, <i>ApJ</i>, 450, 559</p> <p>Böhringer, H., Schuecker, P., Guzzo, L., et al. 2004, <i>A&A</i>, 425, 367</p> |
|---|---|

- Bravi, L., Gitti, M., & Brunetti, G. 2016, *MNRAS*, 455, L41
- Briggs, D. S. 1995, *Bulletin of the American Astronomical Society*, 27, 112.02
- Brunetti, G., Giacintucci, S., Cassano, R., et al. 2008, *Nature*, 455, 944
- Brunetti, G., & Jones, T. W. 2014, *International Journal of Modern Physics D*, 23, 1430007
- Burns, J. O., Sulkanen, M. E., Gisler, G. R., & Perley, R. A. 1992, *ApJL*, 388, L49
- Cassano, R., Gitti, M., & Brunetti, G. 2008, *A&A*, 486, L31
- Cassano, R., Ettori, S., Brunetti, G., et al. 2013, *ApJ*, 777, 141
- Cavagnolo, K. W., Donahue, M., Voit, G. M., & Sun, M. 2009, *ApJS*, 182, 12 (C09)
- Chandra, P., Ray, A., & Bhatnagar, S. 2004, *ApJ*, 612, 974
- K. C. Chambers, E. A. Magnier, N. Metcalfe, et al., 2016, arXiv:1612.05560
- Condon, J. J. 1992, *ARA&A*, 30, 575
- Condon, J. J., Cotton, W. D., Greisen, E. W., et al. 1998, *AJ*, 115, 1693
- Crawford, C. S., Allen, S. W., Ebeling, H., Edge, A. C., & Fabian, A. C. 1999, *MNRAS*, 306, 857
- David, L. P., Slyz, A., Jones, C., et al. 1993, *ApJ*, 412, 479
- Doria, A., Gitti, M., Ettori, S., et al. 2012, *ApJ*, 753, 47
- Fujita, Y., Kohri, K., Yamazaki, R., & Kino, M. 2007, *ApJL*, 663, L61
- Fujita, Y., & Ohira, Y. 2013, *MNRAS*, 428, 599
- Gendron-Marsolais, M., Hlavacek-Larrondo, J., van Weeren, R. J., et al. 2017, *MNRAS*, 469, 3872
- Giacintucci, S., Markevitch, M., Brunetti, G., Cassano, R., & Venturi, T. 2011, *A&A*, 525, L10
- Giacintucci, S., Markevitch, M., Venturi, T., et al. 2014a, *ApJ*, 781, 9 (G14a)
- Giacintucci, S., Markevitch, M., Brunetti, G., et al. 2014b, *ApJ*, 795, 73
- Giacintucci, S., Markevitch, M., Cassano, R., et al. 2017, *ApJ*, 841, 71 (G17)
- Giacintucci, S., Markevitch, M., Johnston-Hollitt, M., et al., 2019, *ApJ*, to be submitted
- Gitti, M., Brunetti, G., & Setti, G. 2002, *A&A*, 386, 456
- Gitti, M., Brunetti, G., Feretti, L., & Setti, G. 2004, *A&A*, 417, 1
- Gitti, M., Feretti, L., & Schindler, S. 2006, *A&A*, 448, 853
- Gitti, M., Ferrari, C., Domainko, W., Feretti, L., & Schindler, S. 2007, *A&A*, 470, L25
- Gitti, M., Tozzi, P., Brunetti, G., et al. 2015, *Advancing Astrophysics with the Square Kilometre Array (AASKA14)*, 76
- Govoni, F., Murgia, M., Markevitch, M., et al. 2009, *A&A*, 499, 371
- Guo, F., & Oh, S. P. 2008, *MNRAS*, 384, 251
- Hitomi Collaboration, Aharonian, F., Akamatsu, H., et al. 2016, *Nature*, 535, 117
- Hlavacek-Larrondo, J., Allen, S. W., Taylor, G. B., et al. 2013, *ApJ*, 777, 163
- Hogan, M. T., Edge, A. C., Hlavacek-Larrondo, J., et al. 2015, *MNRAS*, 453, 1201
- Ikebe, Y., Reiprich, T. H., Böhringer, H., Tanaka, Y., & Kitayama, T. 2002, *A&A*, 383, 773
- Jacob, S., & Pfrommer, C. 2017, *MNRAS*, 467, 1478
- Kalberla, P. M. W., Burton, W. B., Hartmann, D., et al. 2005, *A&A*, 440, 775
- Kale, R., Venturi, T., Giacintucci, S., et al. 2013, *A&A*, 557, A99
- Kale, R., Venturi, T., Giacintucci, S., et al. 2015, *A&A*, 579, A92 (K15)
- Keshet, U., Markevitch, M., Birnboim, Y., & Loeb, A. 2010, *ApJL*, 719, L74
- Keshet, U., & Loeb, A. 2010, *ApJ*, 722, 737
- Macario, G., Venturi, T., Brunetti, G., et al. 2010, *A&A*, 517, A43
- Markevitch, M., Ponman, T. J., Nulsen, P. E. J., et al. 2000, *ApJ*, 541, 542
- Markevitch, M., Mazzotta, P., Vikhlinin, A., et al. 2003, *ApJL*, 586, L19
- Markevitch, M., & Vikhlinin, A. 2007, *PhR*, 443, 1
- Maughan, B. J., Giles, P. A., Randall, S. W., Jones, C., & Forman, W. R. 2012, *MNRAS*, 421, 1583
- Mazzotta, P., Edge, A. C., & Markevitch, M. 2003, *ApJ*, 596, 190
- Mazzotta, P., & Giacintucci, S. 2008, *ApJL*, 675, L9
- McDonald, M., McNamara, B. R., van Weeren, R. J., et al. 2015, *ApJ*, 811, 111
- McDonald, M., Allen, S. W., Bayliss, M., et al. 2017, *ApJ*, 843, 28
- McNamara, B. R., & Nulsen, P. E. J. 2007, *ARA&A*, 45, 117

- McNamara, B. R., & Nulsen, P. E. J. 2012, *New Journal of Physics*, 14, 055023
- Murgia, M., Eckert, D., Govoni, F., et al. 2010, *A&A*, 514, A76
- Perley, R. A., & Butler, B. J. 2013, *ApJS*, 204, 19
- Piffaretti, R., Arnaud, M., Pratt, G. W., Pointecouteau, E., & Melin, J.-B. 2011, *A&A*, 534, A109
- Pfrommer, C., & Enßlin, T. A. 2004, *A&A*, 413, 17
- Pfrommer, C. 2013, *ApJ*, 779, 10
- Planck Collaboration, Ade, P. A. R., Aghanim, N., et al. 2014, *A&A*, 571, 29
- Ruszkowski, M., Yang, H.-Y. K., & Reynolds, C. S. 2017, *ApJ*, 844, 13
- Sanders, J. S., Fabian, A. C., & Taylor, G. B. 2009, *MNRAS*, 396, 1449
- Sarazin, C. L., Baum, S. A., & O’Dea, C. P. 1995, *ApJ*, 451, 125
- Savini, F., Bonafede, A., Brüggen, M., et al. 2018, *MNRAS*, 478, 2234
- Savini, F., Bonafede, A., Brüggen, M., et al. 2019, *A&A*, 622, A24
- Sijbring L.G., PhD Thesis, University of Groningen, A radio continuum and HI Line study of the Perseus cluster (1993)
- Storm, E., Jeltama, T. E., & Rudnick, L. 2015, *MNRAS*, 448, 2495
- Taylor, G. B., Barton, E. J., & Ge, J. 1994, *AJ*, 107, 1942
- van Haarlem, M. P., Wise, M. W., Gunst, A. W., et al. 2013, *A&A*, 556, A2
- van Weeren, R. J., Intema, H. T., Lal, D. V., et al. 2014, *ApJL*, 786, L17
- van Weeren, R. J., de Gasperin, F., Akamatsu, H., et al. 2019, *SSRv*, 215, 16
- Venturi, T., Giacintucci, S., Brunetti, G., et al. 2007, *A&A*, 463, 937
- Venturi, T., Giacintucci, S., Dallacasa, D., et al. 2008, *A&A*, 484, 327
- Venturi, T., Rossetti, M., Brunetti, G., et al. 2017, *A&A*, 603, A125
- Vikhlinin, A., Markevitch, M., Murray, S. S., et al. 2005, *ApJ*, 628, 655
- Vikhlinin, A., Burenin, R. A., Ebeling, H., et al. 2009, *ApJ*, 692, 1033
- Werner, N., Zhuravleva, I., Canning, R. E. A., et al. 2016, *MNRAS*, 460, 2752
- Yang, H.-Y. K., & Reynolds, C. S. 2016, *ApJ*, 829, 90
- Zandanel, F., Pfrommer, C., & Prada, F. 2014, *MNRAS*, 438, 124
- Zhuravleva, I., Churazov, E., Schekochihin, A. A., et al. 2014, *Nature*, 515, 85
- ZuHone, J. A., Markevitch, M., & Johnson, R. E. 2010, *ApJ*, 717, 908
- ZuHone, J. A., Markevitch, M., Brunetti, G., & Giacintucci, S. 2013, *ApJ*, 762, 78
- ZuHone, J. A., Brunetti, G., Giacintucci, S., & Markevitch, M. 2015, *ApJ*, 801, 146

Table 8. Newly analyzed 1.4 GHz *VLA* observations

| Cluster | Project | Configuration | Bandwidth (MHz) | Date | Time (min) | FWHM, PA ($'' \times ''$, $^\circ$) | rms ($\mu\text{Jy b}^{-1}$) |
|-----------------|---------|---------------|--------------------|----------------|---------------|---|----------------------------------|
| A 1835 | AI75 | B | 25 | 1998 Jul 13 | 443 | $5.2 \times 4.5, -74$ | 15 |
| | AG729 | C | 25 | 2006 Oct 23,28 | 325 | $16.2 \times 13.6, 37$ | 30 |
| | AG639 | D | 50 | 2003 Mar 11 | 260 | $50.5 \times 45.0, 4$ | 40 |
| Ophiuchus | AC261 | D | 25 | 1990 Jan 25 | 28 | $91.6 \times 39.2, -25$ | 100 |
| A 2029 | AL77 | B | 12.5 | 1984 Jan 15 | 69 | $4.5 \times 4.3, 11$ | 60 |
| | AG729 | C | 50 | 2006 Dec 26 | 355 | $15.2 \times 13.7, -15$ | 30 |
| | AG639 | D | 50 | 2003 Mar 09 | 293 | $57.3 \times 45.2, 0$ | 40 |
| RBS 797 | AS720 | A | 50 | 2002 Mar 27 | 257 | $1.6 \times 1.2, -11$ | 17 |
| | AS720 | B | 50 | 2002 Jul 27 | 45 | $5.2 \times 3.9, 10$ | 27 |
| RX J1347.5–1145 | AS632 | A | 25 | 1998 Apr 22 | 115 | $1.7 \times 1.3, -18$ | 35 |
| | AD511 | C | 50 | 2005 Sep 03 | 94 | $16.4 \times 13.0, -1$ | 40 |
| MS 1455.0+2232 | AL317 | A | 50 | 1994 Mar 12 | 33 | $1.5 \times 1.2, 33$ | 55 |
| | AS823 | C | 50 | 2006 Dec 12 | 31 | $16.5 \times 14.6, -7$ | 70 |
| 2A 0335+096 | AM828 | B | 6.3 | 2005 Apr 29 | 136 | $4.2 \times 3.9, -53$ | 60 |
| | AS465 | C | 50 | 1992 Mar 14 | 165 | $15.0 \times 14.3, -39$ | 38 |
| | AS465 | D | 50 | 1992 Jul 14 | 117 | $50.9 \times 45.2, 40$ | 50 |

Notes. Column 1: cluster name. Columns 2 and 3: *VLA* array configuration and project. Columns 4–6: observing bandwidth and date. Column 7: observing time. Column 8: radio beam (for ROBUST=0 in IMAGR). Column 9: image rms level (1σ).

APPENDIX

A. RADIO OBSERVATIONS AND IMAGES

Table 8 lists the *VLA* 1.4 GHz archival observations of 7 clusters with known minihalos, that we have reanalyzed in this work. The data were reduced as described in §2.2. The radio images are presented in Figs. 11–17, compared to the X-ray *Chandra* and *XMM-Newton* images. In Table 9, we summarize for each cluster the flux density of the discrete radio galaxies detected in the area occupied by the minihalo and the source-subtracted flux density of the minihalos, measured as described in §2.1. A brief description of each individual cluster is given below.

A.1. Notes on individual clusters

A 1835. A central minihalo was reported by Govoni et al. (2009). Our images are presented in Figure 10. The identification of the discrete sources embedded in the minihalo emission was done using the B-configuration image, shown in (a). We identified 17 discrete sources above a $5\sigma = 0.075$ mJy beam $^{-1}$ level (Table 9), accounting for a total of $S_{1.4\text{GHz}} = 44.8$ mJy. A global flux of $S_{\text{tot}, 1.4\text{GHz}} =$




Figure 10. A 1835. (a) *VLA* 1.4 GHz C-configuration contours at $20''$ resolution (black), overlaid on the 1.4 GHz B-configuration image (grey scale and red contours) at $5''$ resolution. The rms noise levels are $1\sigma = 30 \mu\text{Jy beam}^{-1}$ and $1\sigma = 15 \mu\text{Jy beam}^{-1}$, respectively. Black contours are -1 (dashed), $1, 2, 4, 8, 16, \dots \times 3\sigma$. Red contours are at $+5\sigma$. Individual discrete radio sources are labeled. (b) *VLA* 1.4 GHz D-configuration contours of the minihalo and embedded radio sources (yellow crosses; see panel a), overlaid on the *Chandra* X-ray image in the $0.5\text{--}4$ keV band. The radio image has a restoring beam of $51'' \times 50''$. The rms noise is $1\sigma = 40 \mu\text{Jy beam}^{-1}$. Contours are -1 (dashed), $1, 2, 4, 8, 16, \dots \times 2.5\sigma$.

50.6 ± 2.5 mJy is measured on the D-configuration image in the region occupied by the minihalo (Figure 10b). After subtraction of the flux in discrete sources, $S_{\text{MH}, 1.4\text{GHz}} = 6.1 \pm 1.3$ mJy remain for the minihalo, corresponding to a radio luminosity at 1.4 GHz of $P_{\text{MH}, 1.4\text{GHz}} = (1.19 \pm 0.25) \times 10^{24} \text{ W Hz}^{-1}$. Our flux value is in agreement with the flux density reported by Govoni et al. (2009) using the same C- and D-configuration observations analyzed here. The flux of the BCG is $S_{\text{BCG}, 1.4\text{GHz}} = 32.4 \pm 1.6$ mJy and its radio power $P_{\text{MH}, 1.4\text{GHz}} = (6.3 \pm 0.3) \times 10^{24} \text{ W Hz}^{-1}$.

Ophiuchus. A central minihalo was reported by Govoni et al. (2009) and Murgia et al. (2010). Our images are presented in Figure 11. We identified 10 discrete radio sources in the minihalo area (Table 9) using an image at $92'' \times 39''$ resolution, shown in (a). These sources account for a total flux of $S_{1.4\text{GHz}} = 733.8$ mJy. To measure the minihalo flux density, we used a lower resolution image, shown as contours in (b) on the *XMM-Newton* X-ray image. We subtracted the contribution from discrete sources. We also masked the region marked by a circle, which coincides with an X-ray cavity suggested by the *Chandra* and *XMM-Newton* images of the cluster core (Werner et al. 2016, Giacintucci et al. 2019). The cavity is filled by a very steep spectrum relic lobe, likely inflated by a previous outburst of central AGN, discovered in *GMRT* and *MWA* images at lower frequencies (Giacintucci et al. 2019). The relic lobe is only partially visible at 1.4 GHz (Figure 11b). We obtained a total flux of $S_{\text{MH}, 1.4\text{GHz}} = 62 \pm 9$ mJy for the minihalo ($P_{\text{MH}, 1.4\text{GHz}} = (0.11 \pm 0.02) \times 10^{24} \text{ W Hz}^{-1}$). A higher flux of $106.4^{+10.4}_{-8.9}$ mJy has been measured by Govoni et al. (2009) using an exponential model to fit the surface brightness profile of the minihalo. A flux density of 85 ± 3 mJy is reported by the same authors by integration within a radius of ~ 230 kpc. We note, however, that both these measurements include part of the emission from the newly detected relic lobe. The flux density of




Figure 11. Ophiuchus. (a) VLA 1.4 GHz D-configuration image. The restoring beam is $92'' \times 39''$ and rms noise is $1\sigma = 100 \mu\text{Jy beam}^{-1}$. The individual radio galaxies are labeled. (b) VLA 1.4 GHz D-configuration contours of the minihalo and radio galaxies (yellow and black crosses; see panel a), overlaid on the XMM-Newton X-ray image (OBS ID 0505150101). The radio image has a restoring beam of $110'' \times 60''$. The rms noise is $1\sigma = 120 \mu\text{Jy beam}^{-1}$. Contours are -1 (dashed), $1, 2, 4, 8, 16, \dots \times 2.5\sigma$. The region associated with the low-frequency relic lobe that fills a possible X-ray cavity (circle; Giacintucci et al. 2019) has been excluded to measure the flux density of the minihalo.

the BCG is $S_{\text{BCG}, 1.4\text{GHz}} = 30.6 \pm 1.5 \text{ mJy}$ and its radio power $P_{\text{MH}, 1.4\text{GHz}} = (0.064 \pm 0.003) \times 10^{24} \text{ W Hz}^{-1}$.

A2029. A central minihalo was reported by Govoni et al. (2009). Our images are presented in Figure 12. We identified 11 discrete radio sources (Table 9), including the central dominant wide angle tail (e.g., Taylor et al. 1994), using the B- and C-configuration images (a). The discrete source account for a total of $S_{1.4\text{GHz}} = 531.2 \text{ mJy}$. We used the D-configuration observation to map the minihalo (b) and measure its flux density. Our value of $S_{\text{MH}, 1.4\text{GHz}} = 19.5 \pm 2.5 \text{ mJy}$, after removal of the flux in discrete sources, is in agreement within the errors with $18.8 \pm 1.3 \text{ mJy}$ measured by Govoni et al. (2009). The 1.4 GHz radio luminosity of the minihalo is $P_{\text{MH}, 1.4\text{GHz}} = (0.28 \pm 0.04) \times 10^{24} \text{ W Hz}^{-1}$. The flux density of the BCG is $S_{\text{BCG}, 1.4\text{GHz}} = 513 \pm 26 \text{ mJy}$ and its radio power $P_{\text{BCG}, 1.4\text{GHz}} = (7.4 \pm 0.4) \times 10^{24} \text{ W Hz}^{-1}$.

RBS 797. A central minihalo was reported by Gitti et al. (2006) and Doria et al. (2012). Our images are presented in Figure 13. Using the A-configuration image (a), we measure a radio flux density of $S_{\text{BCG}, 1.4\text{GHz}} = 17.9 \pm 0.9 \text{ mJy}$ ($P_{\text{BCG}, 1.4\text{GHz}} = (7.6 \pm 0.4) \times 10^{24} \text{ W Hz}^{-1}$) for the BCG (S1), which includes the central compact component and the two radio lobes associated with a pair of X-ray cavities (circles in panel b; see Gitti et al. 2006, Doria et al. 2012). The flux density of whole radio emission detected by the B-configuration image (b) is $S_{\text{tot}, 1.4\text{GHz}} = 23.1 \pm 1.1 \text{ mJy}$, in agreement within the errors with the $24 \pm 0.3 \text{ mJy}$ reported by Doria et al. (2012). Once removed the BCG flux from the total emission, we find a residual flux of $S_{\text{MH}, 1.4\text{GHz}} = 5.2 \pm 0.6 \text{ mJy}$ for the minihalo (a flux density of $6.1 \pm 0.4 \text{ mJy}$ was estimated by Doria et al.), corresponding to $P_{\text{MH}, 1.4\text{GHz}} = (2.20 \pm 0.24) \times 10^{24} \text{ W Hz}^{-1}$.




Figure 12. A 2029. (a) *VLA* 1.4 GHz C-configuration contours, overlaid on the B-configuration image. The restoring beam and rms noise are $20'' \times 16''$ and $1\sigma = 30 \mu\text{Jy beam}^{-1}$ and $5'' \times 4''$ and $1\sigma = 50 \mu\text{Jy beam}^{-1}$, respectively. Contours are -0.1 (dashed), $0.1, 0.2, 0.4, 1.6, 6.4 \text{ mJy beam}^{-1}$. The individual radio galaxies are labeled. (b) *VLA* 1.4 GHz D-configuration contours of the minihalo and radio galaxies (yellow and black crosses; see panel a), overlaid on the *Chandra* image in the 0.5–4 keV band. The radio image has a restoring beam of $57'' \times 45''$. The rms noise is $1\sigma = 40 \mu\text{Jy beam}^{-1}$. Contours are -0.12 (dashed), $0.12, 0.24, 0.48, 0.96, 2, 4, 16, 64, 256 \text{ mJy beam}^{-1}$.




Figure 13. RBS 797. (a) *VLA* 1.4 GHz A-configuration image, restored with a $2'' \times 1''$ beam. The rms noise is $1\sigma = 17 \mu\text{Jy beam}^{-1}$. The central radio galaxy S1 and the radio lobes filling the X-ray cavities (white circles in panel b) are labeled. (b) *VLA* 1.4 GHz B-configuration contours overlaid on the *Chandra* image in the 0.5–4 keV band. The radio image has a restoring beam of $5'' \times 4''$. The rms noise is $1\sigma = 27 \mu\text{Jy beam}^{-1}$. Contours start at $70 \mu\text{Jy beam}^{-1}$ and then scale by a factor of 2. No levels at -3σ are present in the portion of the image shown. Circles mark the X-ray cavities filled by the radio lobes (a).




Figure 14. RX J1347.5–1145. (a) *VLA* 1.4 GHz C-configuration contours, overlaid on the 1.4 GHz A-configuration image. The restoring beam and rms noise are $17''$ and $1\sigma = 40 \mu\text{Jy beam}^{-1}$ and $1''.2$ and $1\sigma = 35 \mu\text{Jy beam}^{-1}$, respectively. Contours are -0.12 (dashed), 0.12 , 0.24 , 0.48 , 0.96 , 2 , 8 , 32 , $128 \text{ mJy beam}^{-1}$. Individual radio galaxies are labeled. (b) *VLA* 1.4 GHz C-configuration contours of the minihalo and radio galaxies (black and yellow crosses; see panel a), overlaid on the *Chandra* image in the 0.5–4 keV band.

RX J1347.5–1145. A central minihalo was reported by Gitti et al. (2007). Our images are presented in Figure 14. The central radio galaxy is a point source in the 1.4 GHz image from the A configuration (a). Its flux density is $S_{\text{BCG}, 1.4 \text{ GHz}} = 30.3 \pm 1.5 \text{ mJy}$ and its luminosity is $P_{\text{BCG}, 1.4 \text{ GHz}} = (22.7 \pm 1.1) \times 10^{24} \text{ W Hz}^{-1}$. A much fainter and extended source (S2) is also detected, with a flux of $S_{\text{S2}, 1.4 \text{ GHz}} = 2.0 \pm 0.1 \text{ mJy}$ within the 3σ isocontour. Its flux density increases to $4.9 \pm 0.2 \text{ mJy}$ in the lower-resolution image from the C configuration, where the source becomes unresolved. This flux difference is likely due to missing extended emission in the A-configuration image. Similar flux density values for S1 and S2 were measured by Gitti et al. (2007) using the same radio data analyzed here. The total flux in the C-configuration image is $S_{\text{tot}, 1.4 \text{ GHz}} = 69.3 \pm 3.5 \text{ mJy}$, measured out to a radius of $r = 80''$, where the integrated flux reaches saturation (§2.1). The total flux reported by Gitti et al. (2007) is $\sim 25\%$ lower than our measured flux. However, if we integrate the flux density within the 3σ isocontour of our C-configuration image, we obtain $60 \pm 3 \text{ mJy}$, which is consistent within 1.5σ with Gitti et al. (2007). After removal of S1 and S2 from the total flux, we estimate $S_{\text{MH}, 1.4 \text{ GHz}} = 35.6 \pm 2.9 \text{ mJy}$ for the minihalo. The error includes the uncertainty in the subtraction of S2 using its flux from the A- or C-configuration images. The minihalo luminosity at 1.4 GHz is $P_{\text{MH}, 1.4 \text{ GHz}} = (26.7 \pm 2.1) \times 10^{24} \text{ W Hz}^{-1}$.

MS 1455.0+2232 (Z7160). The minihalo, first detected by Venturi et al. (2008) with the *GMRT* at 610 MHz, is only marginally extended in the *VLA* C-configuration image at 1.4 GHz (Figure 15). The central compact radio galaxy has $S_{\text{BCG}, 1.4 \text{ GHz}} = 4.7 \pm 0.2 \text{ mJy}$ in the A-configuration image, corresponding to $P_{\text{BCG}, 1.4 \text{ GHz}} = (0.96 \pm 0.05) \times 10^{24} \text{ W Hz}^{-1}$. The total integrated flux density in the C-configuration image is $S_{\text{tot}, 1.4 \text{ GHz}} = 13.2 \pm 0.7 \text{ mJy}$, the minihalo accounting for $S_{\text{MH}, 1.4 \text{ GHz}} = 8.5 \pm 1.1 \text{ mJy}$ ($P_{\text{MH}, 1.4 \text{ GHz}} = (1.75 \pm 0.23) \times 10^{24} \text{ W Hz}^{-1}$). The BCG and minihalo flux densities at 610 MHz are $S_{\text{BCG}, 610 \text{ MHz}} = 9.5 \pm 0.8 \text{ mJy}$ and $S_{\text{MH}, 610 \text{ MHz}} = 28.7 \pm 3.7 \text{ mJy}$,




Figure 15. MS 1455.0+2232. (a) VLA 1.4 GHz C-configuration contours, overlaid on the 1.4 GHz A-configuration image. The restoring beam and rms noise are $16'' \times 15''$ and $1\sigma = 70 \mu\text{Jy beam}^{-1}$ and $1''.5 \times 1''.2$ and $1\sigma = 55 \mu\text{Jy beam}^{-1}$, respectively. Contours are -1 (dashed), 1, 2, 4, 8, 16, ... $\times 3\sigma$. S1 is the central radio galaxy. (b) VLA 1.4 GHz C-configuration contours of the minihalo and radio galaxy (black cross), overlaid on the *Chandra* image in the 0.5-4 keV band.

respectively (Mazzotta & Giacintucci 2008). Their spectral indices between 610 MHz and 1.4 GHz are $\alpha = 0.85 \pm 0.11$ and $\alpha = 1.46 \pm 0.22$.

2A 0335+096. A central minihalo was reported by Sarazin et al. (1995). Our 1.4 GHz images are presented in Figure 16. The central radio galaxy (S1) is a core-dominated, double-lobe source with a flux of $S_{\text{BCG}, 1.4\text{GHz}} = 16 \pm 1$ mJy in the B-configuration image (a). Its radio power is $P_{\text{BCG}, 1.4\text{GHz}} = (0.058 \pm 0.003) \times 10^{24} \text{ W Hz}^{-1}$. A nearby companion (S2) has $S_{\text{S2}, 1.4\text{GHz}} = 1.0 \pm 0.1$ mJy. Low-frequency *GMRT* images at 235 MHz and 610 MHz reveal the presence of two fossil radio lobes associated with a previous outburst of the central AGN, opposite with respect to S1 and aligned along a NorthWest-SouthEast axis (a paper is in preparation; see talk by Giacintucci at the Snowcluster 2018 conference¹¹ for an image at 610 MHz). Their position is marked by white circles in Figure 16b. The North-West lobe fills a prominent X-ray cavity seen in the *Chandra* image (Mazzotta et al. 2003, Sanders et al. 2009). A distinct patch of extended emission is detected at $\sim 25''$ (~ 18 kpc) from the cluster center (green circle). This feature becomes more prominent at lower frequencies due to its very steep radio spectrum and is interpreted as a fossil lobe from an even older AGN outburst. We measure a total flux of $S_{\text{tot}, 1.4\text{GHz}} = 48 \pm 2$ mJy in the combined C+D-configuration image (b). To estimate the 1.4 GHz flux density of the minihalo, we subtract the flux of S1 and S2 from the total emission as well as the flux in the fossil lobes (SE lobe: 4.1 ± 0.2 mJy, NW lobe: 2.7 ± 0.1 mJy, outer NW lobe: 3.1 ± 0.2 mJy). We obtain $S_{\text{MH}, 1.4\text{GHz}} = 21.1 \pm 2.1$ mJy and $P_{\text{MH}, 1.4\text{GHz}} = (0.06 \pm 0.01) \times 10^{24} \text{ W Hz}^{-1}$.

¹¹ <http://www.physics.utah.edu/snowcluster/archive/2018/talks/Giacintucci.pdf>.




Figure 16. 2A 0335+096. (a) *VLA* 1.4 GHz C-configuration contours, overlaid on the 1.4 GHz B-configuration image (grey scale and red contours at $+3, 6$, and 10σ). The restoring beam and rms noise are $15'' \times 14''$ and $1\sigma = 35 \text{ mJy beam}^{-1}$ and $4''$ and $1\sigma = 58 \mu\text{Jy beam}^{-1}$, respectively. Black contours are -1 (dashed), $1, 2, 4, 8, 16, \dots \times 3\sigma$. The central radio galaxy and nearby companion are labeled S1 and S2. Black circles mark the position of two X-ray cavities filled by fossil radio lobes (Giacintucci et al. in preparation). The green circle marks the location of another possible fossil radio lobe. (b) *VLA* 1.4 GHz C+D-configuration contours of the minihalo and radio galaxies (black crosses; see panel a), overlaid on the *Chandra* image in the 0.5-4 keV band. The radio image has a restoring beam of $30'' \times 27''$. The rms noise is $1\sigma = 40 \mu\text{Jy beam}^{-1}$. Contours are $1, 2, 4, 8, 16, \dots \times 3\sigma$. No levels at -3σ are present in the portion of the image shown. The region occupied by the X-ray cavities and fossil lobes (white and green circles) have been excluded to measure the flux density of the minihalo.