

Computational Many-Body Physics via \mathcal{M}_{2^q} Algebra

Emil Prodan

Department of Physics and Department of Mathematical Sciences, Yeshiva University, New York, USA*

The many-body Hamiltonians and other fermionic physical observables are expressed in terms of fermionic creation and annihilation operators, which, at an abstract level, form the algebra of Canonical Anti-commutation Relations. If the one-particle Hilbert space is q -dimensional, then this algebra is canonically isomorphic with the ordinary algebra \mathcal{M}_{2^q} of $2^q \times 2^q$ matrices with complex entries. In this work, we present a method that makes this isomorphism explicit. This supplies concrete matrix representations of various many-body operators without involving the traditional Fock space representation. The result is a steep simplification of the many-body exact diagonalization codes, which is a significant step towards the soft-coding of generic fermionic Hamiltonians. Pseudo-code implementing matrix representations of various many-body operators are supplied and Hubbard-type Hamiltonians are worked out explicitly.

I. INTRODUCTION

Local physical observables of fermionic systems are expressed as products and sums of creation a_n^* and annihilation a_n operators. The latter satisfy the canonical anti-commutation relations which automatically enforce Pauli's exclusion principle. The set of local fermionic physical observables can be closed to and given the structure of a C^* -algebra, called the canonical anti-commutation relations algebra, or in short CAR-algebra [1]. In the Heisenberg approach, one formulates the dynamics of fermions directly on the CAR-algebra and a many-body physical system is completely specified by a tuple (α, \mathcal{T}) , where α is a group homomorphism $\alpha : \mathbb{R} \rightarrow \text{Aut}(\text{CAR})$, specifying the time evolution of the physical observables, and \mathcal{T} is a state invariant w.r.t. the α -dynamics. In the Schroedinger picture, the dynamics of fermions is formulated on the anti-symmetric sector of the Fock space, which supplies a natural representation space for the CAR-algebra. Note that in Heisenberg's picture there is no place for Hilbert spaces and representations, and this observation is the starting point for our work. While we focus here at computational many-body aspects, this subtle but essential difference between the two pictures of quantum phenomena has conceptual consequences, as highlighted recently by Haldane in [2].

At the computational level, the difference manifests as follows: In the Heisenberg picture, one seeks a direct homomorphism that embeds the algebra of observables in a matrix algebra. In Schroedinger's picture, the matrix representations are generated by acting with the operators on the basis of the Hilbert space. To see the major difference, let us consider a generic system where the fermions populate a discrete set X of some physical space. We denote by q the cardinal of X . The generic many-body observables take the form:

$$A = \sum_{J, J' \subseteq X} a_{J, J'} \prod_{x \in J} a_x^* \prod_{x' \in J'} a_{x'} \quad (1)$$

Then in the Schroedinger picture, one will generate the $2^q \times 2^q$

matrix representation by looping over the occupation basis:

$$\langle n'_1 n'_2 \dots n'_q | A | n_1 n_2 \dots n_q \rangle, \quad n'_i, n_i \in \{0, 1\}, \quad (2)$$

using, for example, the action of the generators on the basis:

$$\begin{aligned} a_m^* |n_1 \dots n_m \dots n_q\rangle \\ = (-1)^\alpha (n_m \oplus 1) \text{mod } 2 |n_1 \dots n_m + 1 \dots n_q\rangle, \end{aligned} \quad (3)$$

where \oplus is addition mod 2. In a successful Heisenberg program, however, one will use the embedding homomorphism to explicitly specify the entire $2^q \times 2^q$ matrix in one step. When the many-body operator has a simple structure, the action of $\prod_{x \in J} a_x^* \prod_{x' \in J'} a_{x'}$ on the occupation basis can be computed by hand and then the result can be integrated in the computer codes. However, this is typically hard-coded and the entire task needs to be repeated when presented with a different operator. A soft-code, by definition, is one that can diagonalize any many-body observable A based on an input file that contains the subsets J and J' of X , as well as the associated coefficients $a_{J, J'}$. In the Schroedinger approach, the only way to achieve such soft-coding is to repeatedly apply the generators on the occupation basis but this leads to highly inefficient algorithms. This highlights one of the advantages of the Heisenberg approach, which become extremely useful when dealing with complicated Hamiltonians such as the ones often occurring in the research on topological phases of matters. For example, the Fidkowski-Kitaev Hamiltonians [3, 4] contains products of as many as 8 generators! The model Hamiltonians for higher fractional Hall sequences [5] present the same if not even higher level of complexity.

In this work, we exploit a well-know isomorphism between CAR and M_{2^∞} algebras [6] to derive matrix representations of generic products of creation and annihilation operators. Explicit analytic formulas are supplied for several key products of generators, which will enable one to analytically translate any many-fermion Hamiltonian into a matrix form. For the reader's convenience, we exemplify the algorithms with concrete pieces of code and we work out several interesting many-fermion eigen-problems.

In our opinion, the benefits of the proposed approach can materialize in two extreme settings. The first one, is that of small-scale computations involving complex Hamiltonians. For example, the search and characterization of topo-

* Financial support through an award from W. M. Keck Foundation is acknowledged.

logical boundary modes in correlated systems require precisely this type of computations, especially when the goal is to validate their robustness against arbitrary interaction potentials. The challenge for this type of research is that the one-particle Hilbert spaces and the many-body Hamiltonians can vary drastically from one application to another and it is precisely this challenge that is addressed by our approach. The second setting is that of large-scale computations with standard two-body potentials, such as the Coulomb potential. Since our approach supplies formal matrix representations of the Hamiltonians, one can estimate the sparseness of the matrices (see for example Fig. 1) and then decide more easily on the optimal linear-algebra package to be used. One can also estimate more accurately the numerical errors and speed-up of the computations can result from the analytically determined action of the whole Hamiltonian on vectors.

II. BACKGROUND

A. The Algebra of Canonical Anti-Commutation Relations

The algebra of canonical anti-commutation relations (CAR) is defined [1] by a linear map $a : \mathcal{H} \rightarrow \mathcal{B}(\mathcal{H}')$ from a Hilbert space \mathcal{H} onto the algebra of linear maps over another Hilbert space \mathcal{H}' , satisfying the following algebraic relations:

$$\begin{aligned} a(f)a(g) + a(g)a(f) &= 0 \\ a(f)^*a(g) + a(g)a(f)^* &= (g, f)1. \end{aligned} \quad (4)$$

for all $f, g \in \mathcal{H}$. Here and throughout, $(,)$ denotes the scalar product on \mathcal{H} . The CAR-algebra is the C^* -algebra generated by $\{a(f) : f \in \mathcal{H}\}$ modulo relations (4), endowed with the $*$ -operation and the C^* -norm borrowed from $\mathcal{B}(\mathcal{H}')$. Up to an isomorphism, this definition is completely independent of the concrete representations of the Hilbert spaces. In many-body physics, \mathcal{H} represents the one-particle Hilbert space and \mathcal{H}' is chosen as the Fock-space and one says that $a(f)^*$ creates a fermion in the quantum state f , while $a(f)$ destroys a fermion in quantum state f .

For condensed matter physicists, perhaps a more familiar representation of the CAR-algebra can be given in the following terms. Let $\{f_i\}_{i=0, \infty}$ denote an orthonormal basis on \mathcal{H} and let $a_i = a(f_i)$. Then the a_i 's satisfy the familiar anti-commutation relations:

$$a_i a_j + a_j a_i = a_i^* a_j^* + a_j^* a_i^* = 0, \quad a_i^* a_j + a_j a_i^* = \delta_{ij} 1. \quad (5)$$

If one prefers to maintain the liberty of choosing and changing the basis of the Hilbert space, the first representation in Eq. 4 is definitely more preferable.

We will denote the CAR-algebra over a finite dimensional Hilbert space $\dim \mathcal{H}_q = q < \infty$ by $CAR(q)$. Throughout our presentation, we will be consistent and enumerate the elements of the orthonormal basis starting from 0 and ending at $q - 1$. In other words, we will label the orthonormal basis of \mathcal{H}_q as f_0, f_1, \dots, f_{q-1} .

The CAR-algebra is a C^* -algebra, that is, it is closed under the addition, multiplication and the $*$ -transformation (or

dagger-operation). The CAR-algebra also comes equipped with a norm but, since we are mainly considering finite CAR-algebras, this norm will not play any special role here. If $a(f), a(g), \dots$ are some elements of $CAR(q)$, we will denote by $C^*(a(f), a(g), \dots)$ the sub-algebra generated by them. Henceforth, $C^*(a(f), a(g), \dots)$ contains all elements in $CAR(q)$ that can be formed through sums, multiplications and $*$ -transformations of $a(f), a(g), \dots$. In particular, let us point out that $CAR(q)$ can be naturally embedded in $CAR(q + 1)$ and this sets an inductive tower which enable one to define $CAR(\mathcal{H}_\infty)$ as its inductive limit.

B. The algebra \mathcal{M}_{2^∞}

Let \mathcal{M}_2 denote the algebra of 2×2 matrices with complex entries. Then $\mathcal{M}_{2^q} = \mathcal{M}_2 \otimes \mathcal{M}_2 \dots \otimes \mathcal{M}_2 = \mathcal{M}_2^{\otimes q}$, which is isomorphic to the algebra of $2^q \times 2^q$ matrices with complex entries. Note that \mathcal{M}_{2^q} can be embedded in $\mathcal{M}_{2^{q+1}}$ as $\begin{pmatrix} \mathcal{M}_{2^q} & 0 \\ 0 & \mathcal{M}_{2^q} \end{pmatrix}$ and, as such, one can set an inductive tower and define the UHF-algebra \mathcal{M}_{2^∞} as its inductive limit. The result is one of the most studied C^* -algebras in the mathematics literature. For example, its K-theory was worked out in [7] (see also [6]).

We now introduce notations and conventions for our exposition. For $A \in \mathcal{M}_2$ we choose to write $A = \begin{pmatrix} A_{00} & A_{01} \\ A_{10} & A_{11} \end{pmatrix}$. As a linear space, \mathcal{M}_{2^q} is generated by the system of units

$$\{E_{mn}^{(q)}\}_{m,n=0, \dots, 2^q-1}, \quad (6)$$

where $E_{mn}^{(q)}$ is the $2^q \times 2^q$ matrix with entry 1 at position (m, n) and 0 in rest. The system of units satisfies the usual algebraic relations:

$$E_{mn}^{(q)} E_{m'n'}^{(q)} = \delta_{nm'} E_{mn'}^{(q)}. \quad (7)$$

The system of units for \mathcal{M}_2 will be denoted by $\{e_{\alpha\beta}\}_{\alpha,\beta=0,1}$.

C. The link between the algebras

Theorem 1 *CAR(q) is isomorphic to \mathcal{M}_{2^q} for all $q \in \mathbb{N}$.*

Proof. A detailed proof can be found in Kenneth Davidson's monograph [6]. It will be, however, very instructive and helpful to present the proof in details once again here. Henceforth, let f_0, f_1, \dots, f_{q-1} be an orthonormal basis of \mathcal{H}_q and set $a_i = a(f_i)$. Then $CAR(q)$ is simply $C^*(a_0, a_1, \dots, a_{q-1})$. Our first task is to define a new set of generator that commute with each other rather than anti-commute. This can be accomplished via a Jordan-Wigner type transformation, whose main mechanism is recalled below.

Let f be a normalized vector from \mathcal{H} and let:

$$n_f := a(f)^* a(f). \quad (8)$$

Since $a(f)^2 = 0$ and $a(f)^* a(f) + a(f) a(f)^* = 1$, by multiplying the latter by $a(f) a(f)^*$, one obtains $n_f^2 = n_f$. In other words, n_f is an idempotent for any norm-one vector f from

\mathcal{H}_q . In fact, n_f is an orthogonal projector because $n_f^* = n_f$. Furthermore,

$$a(f)a(f)^* = 1 - a(f)^*a(f) = 1 - e(f), \quad (9)$$

hence $a(f)a(f)^*$ is the orthogonal complement of n_f :

$$n_f^\perp := a(f)a(f)^*, \quad n_f n_f^\perp = 0, \quad n_f + n_f^\perp = 1. \quad (10)$$

Consider now another vector g from \mathcal{H}_q which is orthogonal on f , $(f, g) = 0$. One can verify directly that $a(g)$ commutes with n_f (hence also with n_f^\perp) but of course, $a(g)$ does not commute with $a(f)$. This can be fixed as follows. Define:

$$v := n_f^\perp - n_f, \quad (11)$$

with the following obvious properties:

$$v^* = v, \quad v^2 = 1, \quad va(f) = -a(f)v, \quad va(f)^* = -a(f)^*v. \quad (12)$$

If $va(g)$ is considered instead of $a(g)$, then:

$$va(g)a(f) = -va(f)a(g) = a(f)va(g). \quad (13)$$

Similarly:

$$va(g)a(f)^* = -va(f)^*a(g) = a(f)^*va(g). \quad (14)$$

Hence, the substitution $a(g) \rightarrow va(g)$ made the operators commute. This is the essence of the Jordan-Wigner transformation.

Returning now to $C^*(a_0, a_1, \dots, a_{q-1})$, we can define a set of commuting generators by iterating the above construction. This leads us to the following substitutions:

$$a_i \rightarrow v_i a_i, \quad (15)$$

where $v_0 = 1$ and:

$$v_i = (n_{i-1}^\perp - n_{i-1})v_{i-1}, \quad v_i^* = v_i, \quad v_i^2 = 1, \quad i = 1, \dots, q-1, \quad (16)$$

with $n_i = a_i^* a_i$. It is important to keep in mind that n_i 's are all commuting orthogonal projections. The conclusion so far is that:

$$C^*(a_0, a_1, \dots, a_{q-1}) = C^*(v_0 a_0, v_1 a_1, \dots, v_{q-1} a_{q-1}), \quad (17)$$

and now all the generators commute with each other. This concludes the step of the proof that involves the Jordan-Wigner transformation.

The next step is to look at the sub-algebra generated by each of these generators. Because of the anti-commutation relations, one readily finds that $C^*(v_i a_i)$ coincides with the \mathbb{C} -linear span of just four operators:

$$C^*(v_i a_i) = \mathbb{C}\text{-Span}\{n_i, n_i^\perp, v_i a_i, v_i a_i^*\}. \quad (18)$$

Furthermore, if one sets:

$$e_{11}^{(i)} = n_i, \quad e_{00}^{(i)} = n_i^\perp, \quad e_{01}^{(i)} = v_i a_i, \quad e_{10}^{(i)} = v_i a_i^*, \quad (19)$$

then

$$e_{\alpha\beta}^{(i)} e_{\alpha'\beta'}^{(i)} = \delta_{\beta\alpha'} e_{\alpha\beta'}^{(i)} \quad (20)$$

which are exactly the algebraic relations satisfied by the generators of \mathcal{M}_2 . Hence, Eq. 20 defines an explicit isomorphic mapping of \mathcal{M}_2 into $C^*(v_i a_i)$.

The last step of the proof involves the following elements of $CAR(q)$:

$$E_{\varphi, \psi}^{(q)} = e_{\varphi(0)\psi(0)}^{(0)} e_{\varphi(1)\psi(1)}^{(1)} \cdots e_{\varphi(q-1)\psi(q-1)}^{(q-1)}, \quad (21)$$

where φ and ψ are two functions of the type:

$$\varphi, \psi : \{0, 1, \dots, q-1\} \rightarrow \{0, 1\}. \quad (22)$$

Note that there are exactly 2^q distinct such functions and one can verify explicitly that (21) span the entire $CAR(q)$ as well as that:

$$E_{\varphi, \psi}^{(q)} E_{\varphi', \psi'}^{(q)} = \delta_{\psi\varphi'} E_{\varphi, \psi'}^{(q)}, \quad (23)$$

which are precisely the algebraic relations (7) defining the generators of \mathcal{M}_{2^q} . The conclusion is that:

$$\{E_{\varphi, \psi}^{(q)} : \varphi, \psi : \{0, \dots, q-1\} \rightarrow \{0, 1\}\} \quad (24)$$

supply an explicit isomorphic mapping of \mathcal{M}_{2^q} into $CAR(q)$. Furthermore, this mapping respects the embedding of $CAR(q)$ into $CAR(q+1)$ and of \mathcal{M}_{2^q} into $\mathcal{M}_{2^{q+1}}$, hence the inductive towers are isomorphic and their limits are isomorphic as C^* -algebras [8]. \square

III. PRACTICAL REPRESENTATIONS

For practical applications, we need to devise an efficient way to account for all φ 's and ψ 's appearing in Eq. 24.

Proposition 1 *Let n be an integer between 0 and $2^q - 1$. Let:*

$$n = \alpha_0 \cdot 2^0 + \alpha_1 \cdot 2^1 + \dots + \alpha_{q-1} \cdot 2^{q-1}, \quad \alpha_i \in \{0, 1\}, \quad (25)$$

be its unique binary representations and define:

$$\mathfrak{b}_n : \{0, \dots, q-1\} \rightarrow \{0, 1\}, \quad \mathfrak{b}_n(i) = \alpha_i, \quad (26)$$

to be the function which outputs the binary digits of n . Then, when n is varied from 0 to $2^q - 1$, the \mathfrak{b}_n 's generate all the possible functions φ 's and ψ 's appearing in Eq. 24.

Remark 1 We introduce the following important conventions. Firstly, we will identify the elements $e_{\alpha\beta}^{(i)}$ of $CAR(q)$ introduced in (19) with the generators of \mathcal{M}_2 appearing at position i in the tensor product $\mathcal{M}_2 \otimes \dots \otimes \mathcal{M}_2$, tensored by the identity operators of the \mathcal{M}_2 's appearing at the other positions. Secondly, the system of units $E_{nm}^{(q)}$ generating \mathcal{M}_{2^q} and introduced in Eq. 6 will be identified with the elements of $CAR(q)$ via (23):

$$E_{nm}^{(q)} := E_{\mathfrak{b}_n \mathfrak{b}_m}^{(q)}, \quad (27)$$

and, as such, we will use the notations interchangeably. \diamond

The above proposition and Theorem 1 provides the following important Corollary.

Corollary 1 Let $\alpha_0\alpha_1 \dots \alpha_{q-1}$ and $\beta_0\beta_1 \dots \beta_{q-1}$ be two binary sequences of 1's and 0's. Then:

$$e_{\alpha_0\beta_0}^{(0)} e_{\alpha_1\beta_1}^{(1)} \dots e_{\alpha_{q-1}\beta_{q-1}}^{(q)} = E_{nm}^{(q)}, \quad (28)$$

where

$$n = \alpha_0 \cdot 2^0 + \alpha_1 \cdot 2^1 + \dots + \alpha_{q-1} \cdot 2^{q-1}, \quad (29)$$

$$m = \beta_0 \cdot 2^0 + \beta_1 \cdot 2^1 + \dots + \beta_{q-1} \cdot 2^{q-1}. \quad (30)$$

Conversely, for any m and n between 0 and $2^q - 1$ one has:

$$E_{nm}^{(q)} = e_{b_n(0), b_m(0)}^{(0)} e_{b_n(1), b_m(1)}^{(1)} \dots e_{b_n(q-1), b_m(q-1)}^{(q)}. \quad (31)$$

Computer Code 1 Below are code lines which performs the binary decomposition of an integer number $n \in \{0, 1, \dots, 2^q - 1\}$.

```

input n
bn(j) = 0, j = 0, ..., q - 1
p = 0
do j = 0, q - 1
  bn(j) = (n - p)/2j mod 2
  p = p + bn(j) * 2j
end do
return bn(j), j = 0, ..., q - 1.
```

Example 1 Let us compute $e_{12}^{(0)} e_{22}^{(1)} e_{12}^{(2)}$ from $CAR(3) \simeq \mathcal{M}_8$. We have successively:

$$\begin{aligned} e_{12}^{(0)} e_{22}^{(1)} e_{12}^{(2)} &= \begin{bmatrix} 0 & e_{22}^{(1)} e_{12}^{(2)} \\ 0 & 0 \end{bmatrix} \\ &= \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & e_{12}^{(2)} \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} = \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & \boxed{\begin{matrix} 0 & 1 \\ 0 & 0 \end{matrix}} \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} \end{aligned} \quad (33)$$

Above, all the 0's represent the null 2×2 matrix, excepting the 0's in the box, which are just ordinary 0's. On the other hand:

$$\begin{aligned} n &= 0 \times 2^0 + 1 \times 2^1 + 0 \times 2^2 = 2, \\ m &= 1 \times 2^0 + 1 \times 2^1 + 1 \times 2^2 = 7, \end{aligned} \quad (34)$$

hence Corollary 1 predicts:

$$e_{12}^{(0)} e_{22}^{(1)} e_{12}^{(2)} = E_{27}^{(3)}, \quad (35)$$

which is indeed the case (recall that we run the indices from 0 to 7). \diamond

IV. MATRIX REPRESENTATIONS OF MANY-FERMION OPERATORS

A. Matrix representations of the generators

As a model calculation, we derive first the matrix representations of a_i and a_i^* in \mathcal{M}_{2^q} . We start from:

$$a_i = v_i e_{01}^{(i)} = (n_0^\perp - n_0) \dots (n_{i-1}^\perp - n_{i-1}) e_{01}^{(i)}. \quad (36)$$

Using the definitions in Eq. 19 we obtain:

$$a_i = (e_{00}^{(0)} - e_{11}^{(0)}) \dots (e_{00}^{(i-1)} - e_{11}^{(i-1)}) e_{01}^{(i)}. \quad (37)$$

Corollary 1 gives matrix representations for products of e 's that contain exactly q terms. As such, we need to insert identity operators in Eq. 37 until we complete the products:

$$\begin{aligned} a_i &= (e_{00}^{(0)} - e_{11}^{(0)}) \dots (e_{00}^{(i-1)} - e_{11}^{(i-1)}) e_{01}^{(i)} \\ &\quad \times (e_{00}^{(i+1)} + e_{11}^{(i+1)}) \dots (e_{00}^{(q-1)} + e_{11}^{(q-1)}). \end{aligned} \quad (38)$$

Expanding:

$$a_i = \sum_{\alpha^i s} (-1)^{\alpha_0 + \dots + \alpha_{i-1}} e_{\alpha_0 \alpha_0}^{(0)} \dots e_{\alpha_{i-1} \alpha_{i-1}}^{(i-1)} e_{01}^{(i)} e_{\alpha_{i+1} \alpha_{i+1}}^{(i+1)} \dots e_{\alpha_{q-1} \alpha_{q-1}}^{(q-1)}. \quad (39)$$

The above sum is over the set of all binary sequences of the form

$$\alpha_0 \alpha_1 \dots \alpha_{i-1} 0 \alpha_{i+1} \dots \alpha_{q-1},$$

which coincides with the set of the binary expansions of $n \in \{0, \dots, 2^q - 1\}$ with $b_n(i) = 0$. Using Corollary 1 and accounting for α 's properly, we obtain a closed-form formula for a_i and, by applying the $*$ -operation, we also get a closed-form formula for a_i^* :

Proposition 2 In terms of the standard generators of M_{2^q} , we have:

$$a_i = \sum_{n=0}^{2^q-1} (-1)^{\sum_{s=0}^i b_n(s)} \delta_{b_n(i), 0} E_{n, n+2^i}^{(q)}, \quad (40)$$

$$a_i^* = \sum_{n=0}^{2^q-1} (-1)^{\sum_{s=0}^i b_n(s)} \delta_{b_n(i), 0} E_{n+2^i, n}^{(q)}. \quad (41)$$

Remark 2 We have verified analytically that the above matrices indeed satisfy the commutation relations (5). \diamond

B. Matrix representations of products of generators

We continue our computations with a derivation of the product $a_i^* a_j$, assuming for the beginning that $j > i$. Starting from (37), we have:

$$\begin{aligned} a_i^* a_j &= (e_{00}^{(0)} - e_{11}^{(0)}) \dots (e_{00}^{(i-1)} - e_{11}^{(i-1)}) e_{10}^{(i)} \\ &\quad \times (e_{00}^{(0)} - e_{11}^{(0)}) \dots (e_{00}^{(j-1)} - e_{11}^{(j-1)}) e_{01}^{(j)} \\ &= (e_{00}^{(0)} + e_{11}^{(0)}) \dots (e_{00}^{(i-1)} + e_{11}^{(i-1)}) e_{10}^{(i)} \\ &\quad \times (e_{00}^{(i+1)} - e_{11}^{(i+1)}) \dots (e_{00}^{(j-1)} - e_{11}^{(j-1)}) \\ &\quad \times e_{01}^{(j)} (e_{00}^{(j+1)} + e_{11}^{(j+1)}) \dots (e_{00}^{(q-1)} + e_{11}^{(q-1)}), \end{aligned} \quad (42)$$

where the middle line is missing if $j = i + 1$. Let us note that the case $i > j$ follows from the case treated above by applying the conjugation. Furthermore, we can straightforwardly

modify the above arguments to find that, for $i < j$:

$$\begin{aligned} a_i a_j^* &= -\left(e_{00}^{(0)} + e_{11}^{(0)}\right) \dots \left(e_{00}^{(i-1)} + e_{11}^{(i-1)}\right) e_{01}^{(i)} \\ &\quad \times \left(e_{00}^{(i+1)} - e_{11}^{(i+1)}\right) \dots \left(e_{00}^{(j-1)} - e_{11}^{(j-1)}\right) \\ &\quad \times e_{10}^{(j)} \left(e_{00}^{(j+1)} + e_{11}^{(j+1)}\right) \dots \left(e_{00}^{(q-1)} + e_{11}^{(q-1)}\right), \end{aligned} \quad (43)$$

and the reason for the minus sign is $e_{01}^{(i)} (e_{00}^{(i)} - e_{11}^{(i)}) = -e_{01}^{(i)}$, as opposed to $e_{10}^{(i)} (e_{00}^{(i)} - e_{11}^{(i)}) = +e_{10}^{(i)}$. After expanding and using Corollary 1, we obtained:

Proposition 3 *In terms of the standard generators of M_{2^q} , we have for $i \neq j$:*

$$\begin{aligned} a_i^* a_j &= \sum_{n=0}^{2^q-1} (-1)^{\sum_{s=\min(i,j)}^{\max(i,j)} b_n(s)} \delta_{b_n(i),0} \delta_{b_n(j),0} E_{n+2^i, n+2^j}^{(q)} \\ &= - \sum_{n=0}^{2^q-1} (-1)^{\sum_{s=\min(i,j)}^{\max(i,j)} b_n(s)} \delta_{b_n(i),1} \delta_{b_n(j),0} E_{n, n-2^i+2^j}^{(q)}, \end{aligned} \quad (44)$$

and $a_j a_i^* = -a_i^* a_j$.

If $i = j$, the calculations gives:

$$\begin{aligned} a_i^* a_i &= \left(e_{00}^{(0)} + e_{11}^{(0)}\right) \dots \left(e_{00}^{(i-1)} + e_{11}^{(i-1)}\right) e_{11}^{(i)} \\ &\quad \times \left(e_{00}^{(j+1)} + e_{11}^{(j+1)}\right) \dots \left(e_{00}^{(q-1)} + e_{11}^{(q-1)}\right), \end{aligned} \quad (45)$$

and:

$$\begin{aligned} a_i a_i^* &= \left(e_{00}^{(0)} + e_{11}^{(0)}\right) \dots \left(e_{00}^{(i-1)} + e_{11}^{(i-1)}\right) e_{00}^{(i)} \\ &\quad \times \left(e_{00}^{(j+1)} + e_{11}^{(j+1)}\right) \dots \left(e_{00}^{(q-1)} + e_{11}^{(q-1)}\right). \end{aligned} \quad (46)$$

The conclusion is:

Proposition 4 *In terms of the standard generators of M_{2^q} , we have:*

$$n_i = a_i^* a_i = \sum_{n=0}^{2^q-1} \delta_{b_n(i),1} E_{n,n}^{(q)}, \quad (47)$$

$$n_i^\perp = a_i a_i^* = \sum_{n=0}^{2^q-1} \delta_{b_n(i),0} E_{n,n}^{(q)}, \quad (48)$$

$$n_{i_1} n_{i_2} \dots n_{i_k} = \sum_{n=0}^{2^q-1} \delta_{b_n(i_1),1} \dots \delta_{b_n(i_k),1} E_{n,n}^{(q)}, \quad (49)$$

$$n_{i_1}^\perp n_{i_2}^\perp \dots n_{i_k}^\perp = \sum_{n=0}^{2^q-1} \delta_{b_n(i_1),0} \dots \delta_{b_n(i_k),0} E_{n,n}^{(q)}. \quad (50)$$

The products $a_i a_j$ and $a_i^* a_j^*$ can be treated similarly.

Proposition 5 *In terms of the standard generators of M_{2^q} , we have:*

$$\begin{aligned} a_i a_j &= \text{sgn}(i-j) \sum_{n=0}^{2^q-1} (-1)^{\sum_{s=\min(i,j)}^{\max(i,j)} b_n(s)} \\ &\quad \times \delta_{b_n(i),0} \delta_{b_n(j),0} E_{n, n+2^i+2^j}^{(q)}, \end{aligned} \quad (51)$$

and:

$$\begin{aligned} a_i^* a_j^* &= \text{sgn}(j-i) \sum_{n=0}^{2^q-1} (-1)^{\sum_{s=\min(i,j)}^{\max(i,j)} b_n(s)} \\ &\quad \times \delta_{b_n(i),0} \delta_{b_n(j),0} E_{n+2^i+2^j, n}^{(q)}, \end{aligned} \quad (52)$$

where we adopt the convention that $\text{sgn}(0) = 0$.

Remark 3 It will be convenient to introduce the notation:

$$N_{ij}(n) = \sum_{s=\min(i,j)}^{\max(i,j)} b_n(s), \quad i \neq j, \quad (53)$$

since the sign factors determined by these coefficients will appear often in the subsequent presentation. \diamond

A direct consequence of Proposition 5 is the following useful identity:

Corollary 2 *In terms of the standard generators of M_{2^q} , we have:*

$$\begin{aligned} a_i^* a_j^* a_k a_l &= \text{sgn}[(j-i)(k-l)] \sum_{n=0}^{2^q-1} (-1)^{N_{ij}(n)+N_{kl}(n)} \\ &\quad \times \delta_{b_n(i),0} \delta_{b_n(j),0} \delta_{b_n(k),0} \delta_{b_n(l),0} E_{n+2^i+2^j, n+2^k+2^l}^{(q)} \\ &= \text{sgn}[(j-i)(k-l)] \sum_{n=0}^{2^q-1} (-1)^{N_{ij}(n)+N_{kl}(n)} \\ &\quad \times \delta_{b_n(i),1} \delta_{b_n(j),1} \delta_{b_n(k),0} \delta_{b_n(l),0} E_{n, n-2^i-2^j+2^k+2^l}^{(q)}. \end{aligned} \quad (54)$$

Example 2 We derive the matrix representation of the following Hubbard-type Hamiltonian:

$$H = \sum_{i,j=0}^{q-1} \left(\delta_{ij} \epsilon_i n_i + (1-\delta_{ij})(t_{ij} a_i^* a_j + \bar{t}_{ij} a_j^* a_i) + u_{ij} n_i n_j \right), \quad (55)$$

where t_{ij} 's and u_{ij} 's and ϵ_i 's are some complex and real parameters, respectively. Browsing through the list of formulas supplied above, one can see that the matrix representation of H can be obtained automatically from Eqs. (44), (47) and (49):

$$\begin{aligned} H &= \sum_{n=0}^{2^q-1} \sum_{i,j=0}^{q-1} \left[(-1)^{N_{ij}(n)} \delta_{b_n(i),0} \delta_{b_n(j),0} (1-\delta_{ij}) \right. \\ &\quad \times \left(t_{ij} E_{n+2^i, n+2^j}^{(q)} + \bar{t}_{ij} E_{n+2^j, n+2^i}^{(q)} \right) \\ &\quad \left. + \delta_{b_n(i),1} \delta_{b_n(j),1} (u_{ij} + \delta_{ij} \epsilon_i) E_{n,n}^{(q)} \right]. \quad \diamond \end{aligned} \quad (56)$$

Computer Code 2 We provide here a basic piece of code which computes and stores the entire matrix of H from (56) in $\mathcal{M}_{2^{q+1}}$.

```

 $h_{n,n'} = 0, n, n' = 0, \dots, 2^q - 1$ 
do  $n = 0, 2^q - 1$ 
  Call Eq. 32
  do  $i = 0, q - 1$ 
    do  $j = 0, q - 1$ 
      if ( $b_n(i) = b_n(j) = 0$ ) then
         $\mathcal{N}_{ij} = \text{sum}[b_n(\min(i, j) : \max(i, j))]$ 
         $h_{n+2^i, n+2^j} = (-1)^{\mathcal{N}_{ij}} t_{ij}$ 
         $h_{n+2^j, n+2^i} = (-1)^{\mathcal{N}_{ij}} \bar{t}_{ij}$ 
      end if
      if ( $b_n(i) = b_n(j) = 1$ ) then
         $h_{n,n} = \delta_{ij} \epsilon_i + u_{ij}$ 
      end if
    end do end do
  end do
return  $h_{n,n'}, n, n' = 0, \dots, 2^q - 1$ .

```

Let us highlight the simplicity of the code. \diamond

Remark 4 Even though H conserves the number of particles, an issue to be addressed in the next section, there are cases where computing the full matrix of \hat{H} is still desirable, such as when H is perturbed with a potential that does not conserve the number of particles. \diamond

V. N-PARTICLES SECTORS

Our first goal is to give the spectral decomposition of the number of particles operator inside the algebra \mathcal{M}_{2^q} . We will then use its spectral sub-spaces to decompose the Hamiltonians in block diagonals.

A. Spectral resolution of the particle number operator

Let:

$$\hat{N} = \sum_{i=0}^{q-1} a_i^* a_i = \sum_{i=0}^{q-1} n_i = \sum_{i=0}^{q-1} e_{11}^{(i)} \quad (58)$$

be the classical particle-number operator. A direct way to generate its spectral decomposition inside \mathcal{M}_{2^q} will be to complete $e_{11}^{(i)}$'s to full product sequences and follow the steps above. We, however, proceed slightly differently.

Proposition 6 Let n be a number between 0 and $2^q - 1$. Then:

$$\hat{N} E_{n,n}^{(q)} = E_{n,n}^{(q)} \hat{N} = \mathcal{N}(n) E_{n,n}^{(q)}, \quad \mathcal{N}(n) = \sum_{s=0}^{q-1} b_n(s). \quad (59)$$

Proof. From Corollary 1:

$$E_{n,n}^{(q)} = e_{b_n(0), b_n(0)}^{(0)} e_{b_n(1), b_n(1)}^{(1)} \dots e_{b_n(q-1), b_n(q-1)}^{(q-1)}. \quad (60)$$

Since above all the e 's commute, we can separate the terms with $b_n(i) = 1$ to the left and the remaining terms with $b_n(i) = 0$ to the right. In this way, we obtain:

$$E_{n,n}^{(q)} = \prod_{b_n(i)=1} n_i \prod_{b_n(j)=0} n_j^\perp. \quad (61)$$

Then:

$$\begin{aligned} \hat{N} E_{n,n}^{(q)} &= \left(\sum_{k=0}^{q-1} n_k \right) \prod_{b_n(i)=1} n_i \prod_{b_n(j)=0} n_j^\perp \\ &= \sum_{k=0}^{q-1} (\delta_{b_n(k),0} + \delta_{b_n(k),1}) n_k \prod_{b_n(i)=1} n_i \prod_{b_n(j)=0} n_j^\perp, \end{aligned} \quad (62)$$

and since the n_i 's are projections, the last line can be written as:

$$\begin{aligned} &\sum_{k=0}^{q-1} \delta_{b_n(k),1} \prod_{b_n(i)=1} n_i \prod_{b_n(j)=0} n_j^\perp \\ &= \left(\sum_{k=0}^{q-1} b_n(k) \right) \prod_{b_n(i)=1} n_i \prod_{b_n(j)=0} n_j^\perp \end{aligned} \quad (63)$$

and the statement follows. \square

Corollary 3 The spectral decomposition of \hat{N} is:

$$\hat{N} = \sum_{N=0}^{q-1} N \left(\sum_{n=0}^{2^q-1} \delta_{\mathcal{N}(n),N} E_{n,n}^{(q)} \right). \quad (64)$$

Proof. The family of rank-one projections $E_{n,n}^{(q)}$, $n = 0, 2^q - 1$ gives a resolution of the identity in \mathcal{M}_{2^q} :

$$\sum_{n=0}^{2^q-1} E_{n,n}^{(q)} = I_{2^q \times 2^q}. \quad (65)$$

Hence, the range of the projections $E_{n,n}^{(q)}$ exhaust all the invariant Hilbert sub-spaces of \hat{N} when n is varied from 0 to $2^q - 1$, and the statement follows. \square .

Computer Code 3 We provide below lines of code that detect and re-label the original indices that belong to a specific N -particle sector. We call these new indices the N -compressed indices.

```

input  $N$ 
ind( $n$ ) = 0,  $n = 0, \dots, 2^q - 1$ ,
 $c = 0$ 
do  $n = 0, 2^q - 1$ 
  Call Eq. 32
   $\mathcal{N}(n) = \text{sum}(b_n)$ 
  if ( $\mathcal{N}(n) = N$ ) then
     $c = c + 1$ 
    ind( $n$ ) =  $c$ 
  end if
enddo
 $D_N = c$ 
return  $D_N, \text{ind}(n), n = 0, \dots, 2^q - 1$ .

```

These new indices will be used to generate, store and manipulate the diagonal blocks of the Hamiltonians corresponding to the N -particle sectors. Note that D_N is the dimension of the N -particle sector. \diamond

B. Elementary operators on N -particle sectors

Let $\Phi(a)$ be a product of a 's with equal number of creation and annihilation generators. Then $\Phi(a)$ commutes with N and the N -th block of the product can be computed from:

$$\begin{aligned} \Phi(a)_N &= \left(\sum_{n=0}^{2^q-1} \delta_{N(n),N} E_{n,n}^{(q)} \right) \Phi(a) \\ &= \Phi(a) \left(\sum_{n=0}^{2^q-1} \delta_{N(n),N} E_{n,n}^{(q)} \right). \end{aligned} \quad (67)$$

Applying this procedure on the products in Propositions 3 and 4 gives:

Proposition 7 *In terms of the standard generators of M_{2^q} , we have:*

$$\begin{aligned} (a_i^* a_j)_N &= -(a_j a_i^*)_N = \\ &= - \sum_{n=0}^{2^q-1} (-1)^{N_{ij}(n)} \delta_{N(n),N} \delta_{b_n(i),1} \delta_{b_n(j),0} E_{n,n-2^i+2^j}^{(q)}, \end{aligned} \quad (68)$$

$$(n_{i_1} n_{i_2} \dots n_{i_k})_N = \sum_{n=0}^{2^q-1} \delta_{N(n),N} \delta_{b_n(i_1),1} \dots \delta_{b_n(i_k),1} E_{n,n}^{(q)}. \quad (69)$$

$$(n_{i_1}^\dagger n_{i_2}^\dagger \dots n_{i_k}^\dagger)_N = \sum_{n=0}^{2^{q+1}-1} \delta_{N(n),N} \delta_{b_n(i_1),0} \dots \delta_{b_n(i_k),0} E_{n,n}^{(q)}. \quad (70)$$

$$\begin{aligned} (a_i^* a_j^* a_k a_l)_N &= \text{sgn}[(j-i)(k-l)] \sum_{n=0}^{2^q-1} (-1)^{N_{ij}(n)+N_{kl}(n)} \\ &\times \delta_{N(n),N} \delta_{b_n(i),1} \delta_{b_n(j),1} \delta_{b_n(k),0} \delta_{b_n(l),0} E_{n,n-2^j+2^k+2^l}^{(q)}. \end{aligned} \quad (71)$$

The particle number operator commutes with any product of generators which contains an equal number of creation and annihilation operators. In particular N commutes with the Hamiltonian defined in Example 2. Its block diagonals are worked out below.

Example 3 In the N -particle sector, the Hubbard model from Example 2 becomes:

$$\begin{aligned} H_N &= \sum_{i,j=0}^{q-1} \left((1 - \delta_{ij})(t_{ij} (a_i^* a_j)_N + \bar{t}_{ij} (a_j^* a_i)_N) \right. \\ &\quad \left. + (\delta_{ij} \epsilon_i + u_{ij})(n_i n_j)_N \right), \end{aligned} \quad (72)$$




FIG. 1. Rendering of $|h_{ab}|$ as function of the compressed indices (a, b) for the Hubbard-type model (72) with $t_{ij} = e^{-0.2|i-j|}$, $u_{ij} = 0.3e^{-0.5|i-j|}$ and $\epsilon_i = 0$. The plot was generated with (74), where the parameters were fixed at $q = 14$ and $N = 7$, in which case the dimension of the N -particle sector was $D_N = 3432$.

and its matrix form can be automatically generated from Proposition 7:

$$\begin{aligned} H_N &= \sum_{n=0}^{2^q-1} \delta_{N(n),N} \sum_{i,j=0}^{q-1} \left[(-1)^{N_{ij}(n)+1} \delta_{b_n(i),1} \delta_{b_n(j),0} (1 - \delta_{ij}) \right. \\ &\quad \times \left(t_{ij} E_{n,n-2^i+2^j}^{(q)} + \bar{t}_{ij} E_{n-2^i+2^j,n}^{(q)} \right) \\ &\quad \left. + \delta_{b_n(i),1} \delta_{b_n(j),1} (\delta_{ij} \epsilon_i + u_{ij}) E_{n,n}^{(q)} \right]. \diamond \end{aligned} \quad (73)$$

Remark 5 Comparing with Eq. (56), we see that the only change in (73) is a selective summation over n . However, when resolving over the particle number sectors, the computational challenge is two-fold: (a) determining the reduced form of the Hamiltonian, which (73) delivers, and (b) storing this reduced Hamiltonian using a minimal and natural set of indices. It is at this point where the indices introduced in (66) become useful, as we will see below. \diamond

Computer Code 4 We provide here code lines which compute and store the matrix of the Hamiltonian defined in Exam-

ple 2, this time in the N -th particle sector of \mathcal{M}_{2^q} .

```

input  $N$ 
Call Eq. (66)
 $h_{a,b} = 0, a, b = 1, \dots, D_N$ 
do  $n = 0, 2^q - 1$ 
  Call Eq. 32
  if ( $\text{sum}(b_n) = N$ ) then
    do  $i = 0, q - 1$ 
      do  $j = 0, q - 1$ 
        if ( $b_n(i) = 1$  &  $b_n(j) = 0$ ) then
           $\mathcal{N}_{ij} = \text{sum}(b_n(\min(i, j) : \max(i, j)))$ 
           $a = \text{ind}(n); b = \text{ind}(n - 2^i + 2^j)$ 
           $h_{a,b} = h_{a,b} - (-1)^{\mathcal{N}_{ij}}(1 - \delta_{ij})t_{ij}$ 
           $h_{b,a} = h_{b,a} - (-1)^{\mathcal{N}_{ij}}(1 - \delta_{ij})\bar{t}_{ij}$ 
        end if
        if ( $b_n(i) = b_n(j) = 1$ ) then
           $a = \text{ind}(n)$ 
           $h_{a,a} = h_{a,a} + \delta_{ij}\epsilon_i + u_{ij}$ 
        end if
      end do
    end do
  end if
end do
return  $h_{a,b}, a, b = 1, \dots, D_N$ .

```

An output of this algorithm is illustrated in Fig. 1. \diamond

VI. CONCLUSIONS

Although the examples we provided were all 1-dimensional, our analysis covers quite generic settings because, once a basis for the one-particle Hilbert space is chosen, Hamiltonians are all rendered using linear indices. To exemplify this point, let us consider a 2-dimensional $L \times L$

lattice $\mathcal{L}_L = \mathbb{Z}_L \times \mathbb{Z}_L$ ($\mathbb{Z}_L = \mathbb{Z}/L\mathbb{Z}$) with K quantum states per site, as well as a generic Hubbard-type Hamiltonian:

$$H = \sum_{i,j \in \mathcal{L}_L} \sum_{\alpha,\beta=0}^{K-1} \left[(1 - \delta_{ij} \delta_{\alpha\beta}) (t_{ij}^{\alpha\beta} a_{i,\alpha}^* a_{j,\beta} + \bar{t}_{ij}^{\alpha\beta} a_{j,\beta}^* a_{i,\alpha}) + (\delta_{ij} \delta_{\alpha\beta} \epsilon_i^\alpha + u_{ij}^{\alpha\beta}) n_{i,\alpha} n_{j,\beta} \right], \quad (75)$$

This model can be reduced identically to the Hamiltonian in Eq. (55), by creating a linear index i for the one-particle Hilbert space $\mathbb{C}^K \otimes \ell^2(\mathcal{L}_L)$ of the model. One way to achieve that is by applying the rule:

$$\begin{aligned} \mathcal{L}_L \times \mathbb{Z}_K \ni (\alpha, \mathbf{i}) &= (\alpha, i_1, i_2) \\ &\Downarrow \\ i &= \alpha + i_1 K + i_2 L \in \mathbb{Z}_q, \end{aligned} \quad (76)$$

with $q = KL^2$. Once we encode the information and re-write the Hamiltonian (75) using this linear index, which amounts to re-encoding the coefficients $t_{ij}^{\alpha\beta} \rightarrow t_{ij}$ and $u_{ij}^{\alpha\beta} \rightarrow u_{ij}$, there is nothing to be added to the previous analysis. Of course, not all basis set choices are the same and some can prove to be more optimal, in the sense that the coefficients t_{ij} are of shorter-range. This is an important issue which needs to be solved before the matrix-representation is attempted.

We also want to stress that the calculations can be straightforwardly expanded to cover higher order products of generators. This becomes quite apparent if the reader examines Eq. 42 and the manipulations after it. Specific applications taking advantage of these matrix representations will be reported in a future work.

ACKNOWLEDGMENTS

This work is supported by National Science Foundation through grant DMR-1823800.

-
- [1] O. Bratteli, D. W. Robinson, *Operator algebras and quantum statistical mechanics 2*, (Springer, Berlin, 2002).
 - [2] F. D. M. Haldane, *The origin of holomorphic states in Landau levels from non-commutative geometry, and a new formula for their overlaps on the torus*, J. Mathe. Phys. **59**, 081901 (2018).
 - [3] L. Fidkowski, A. Kitaev, *Effects of interactions on the topological classification of free fermion systems*, Phys. Rev. B **81**, 134509 (2010).
 - [4] Lukasz Fidkowski and Alexei Kitaev, *Topological phases of fermions in one dimension*, Phys. Rev. B **83**, 075103 (2011).
 - [5] E. Prodan, F.D.M. Haldane, *Mapping the braiding properties of the Moore-Read state*, Phys. Rev. B **80**, 11512 (2009).
 - [6] K. R. Davidson, *C*-Algebras by Example*, (AMS, Providence, 1996).
 - [7] J. Renault, *A groupoid approach to C*-algebras*, (Springer-Verlag, Berlin, 1980).
 - [8] E. G. Effros, *Dimensions and C*-algebras*, (AMS, Providence, 1981).