

CONFORMAL GEODESICS ON GRAVITATIONAL INSTANTONS

MACIEJ DUNAJSKI AND PAUL TOD

ABSTRACT. We study the integrability of the conformal geodesic flow (also known as the conformal circle flow) on the $SO(3)$ -invariant gravitational instantons. On a hyper-Kähler four-manifold the conformal geodesic equations reduce to geodesic equations of a charged particle moving in a constant self-dual magnetic field. In the case of the anti-self-dual Taub NUT instanton we integrate these equations completely by separating the Hamilton-Jacobi equations, and finding a commuting set of first integrals. This gives the first example of an integrable conformal geodesic flow on a four-manifold which is not a symmetric space. In the case of the Eguchi-Hanson we find all conformal geodesics which lie on the three-dimensional orbits of the isometry group. In the non-hyper-Kähler case of the Fubini-Study metric on $\mathbb{C}P^2$ we use the first integrals arising from the conformal Killing-Yano tensors to recover the known complete integrability of conformal geodesics.

1. INTRODUCTION

The geodesic flow on a (pseudo) Riemannian manifold (M, g) is integrable if the underlying metric admits a sufficient number of Killing vectors, or Killing tensors. The former correspond to first integrals linear in the velocity, and the latter give polynomial first integrals of higher order. For example the geodesic motion on a round two-sphere is integrable as any generator of the isometry group $SO(3)$ gives a Killing vector commuting with the Hamiltonian. On a tri-axial ellipsoid there are no Killing vectors, but the additional quadratic first integral required for integrability is given by a rank-two Killing tensor [17].

Let u be a unit tangent vector to a curve $\Gamma \subset M$, and let $a \equiv \nabla_u u$ be the acceleration of Γ . The geodesic condition $a = 0$ is not invariant under conformal rescalings of the metric $g \rightarrow \Omega^2 g$, but there is a different preferred set of curves on manifolds endowed with only conformal structures $(M, [g])$, where $[g] = \{\Omega^2 g, \Omega : M \rightarrow \mathbb{R}^+\}$. These *conformal geodesics* are also known as *conformal circles*, and arise as solutions to a system of third order ODEs on M . A conformal geodesic Γ is uniquely specified by a point, a tangent direction, and a perpendicular acceleration. If g is a representative metric in the conformal class, and ∇ is the Levi-Civita connection of g , then the conformal geodesic equations are

$$\nabla_u a = -(|a|^2 + L(u, u))u + L^\sharp(u), \quad (1.1)$$

where L is the Schouten tensor given in terms of the Ricci tensor by formula (2.5), and $L^\sharp : TM \rightarrow TM$ is the endomorphism corresponding to L . It can be demonstrated [3, 4, 24, 23] with varying degrees of sophistication, that the conformal geodesics only depend on the conformal class of $[g]$, and not on the choice of the representative metric.

Neither the Killing vectors nor the conformal Killing vectors of g give rise to first integrals of (1.1), and it is natural to ask whether there are examples of integrable conformal geodesic motions, and what geometric structures on $(M, [g])$ give rise to this integrability. In [24, 15] it was shown that the conformal Killing-Yano two-forms (CKY) give rise to first integrals of (1.1): $Y \in \Lambda^2(T^*M)$ is a CKY if $(\nabla Y)_0 \in \Gamma(\Lambda^3(T^*M))$, where T_0 denotes the trace-free part of T . The corresponding first integral is then

$$Q = Y(u, a) - \frac{1}{n-1} \operatorname{div}(Y)(u), \quad (1.2)$$

where $n = \dim(M)$, and $\operatorname{div} = *d*$ is the divergence. This was sufficient [24] to integrate (1.1) on a non-conformally flat **Nil** 3-manifold, and a squashed 3-sphere but already the

conformal class of the Schwarzschild metric proved too difficult to handle, as there do not exist sufficiently many CKYs.

In this paper we shall study the integrability of (1.1) on four-dimensional conformal structures corresponding to some gravitational instantons: solutions to Einstein equations on Riemannian four-manifolds which are compact or complete metrics which asymptotically look like flat space [12, 7].

We shall explore the existence of three CKYs corresponding to the underlying hyper-Kähler structure, and reduce (1.1) to a system of 2nd order ODEs corresponding to a forced geodesic equation in a constant magnetic field. Further progress can be made for the anti-self-dual (ASD) Taub-NUT and the Eguchi-Hanson instantons. In the ASD Taub-NUT case we shall establish complete integrability by separating the Hamilton-Jacobi equation.

Theorem 1.1. *The conformal geodesic equation on the anti-self-dual Taub-NUT manifold is completely integrable in the Arnold-Liouville sense: it reduces to a geodesic equation in a self-dual magnetic field which admits four first integrals in involution.*

The associated Hamilton-Jacobi equation is separable, and the conformal geodesic equations reduce to quadratures.

The Eguchi-Hanson metric admits an isometric and tri-holomorphic action of $SO(3)$ which preserves the magnetic field, and therefore gives rise to three charged linear first integrals of the Lorentz force equation. While full integrability cannot be established in this case, there is enough structure to reduce the conformal geodesic motion to quadratures under an additional assumption that the conformal geodesics lie on the 3-dimensional orbits of the $SO(3)$ subgroup of the isometry group.

The paper is organised as follows. In the next section we shall introduce the conformal geodesic equations (1.1), and focus on the special case where the underlying conformal structure admits an Einstein metric. We shall discuss the first-integrals of (1.1), and establish a link with the Lorentz force equations for hyper-Kähler four manifolds (Proposition 3.1). In §4 and §5 we shall study (1.1) on the ASD Taub-NUT metric, and the Eguchi-Hanson metric respectively. The ASD Taub-NUT case is completely integrable (Theorem 1.1 will follow from Proposition 4.1 and Proposition 4.2), and the Eguchi-Hanson case has a couple of integrable sub-cases (Proposition 5.1).

In §6 we shall establish the integrability of the conformal geodesic flow for the Fubini-Study metric on $\mathbb{C}\mathbb{P}^2$. We shall establish it by making use of nine CKYs on this space, and recover the results of [2], where all conformal geodesics of $\mathbb{C}\mathbb{P}^n$ have been characterised by their horizontal lifts to helices on the total space of the fibration $S^{2n+1} \rightarrow \mathbb{C}\mathbb{P}^n$. In Appendix A we shall rule out the existence of non-flat Riemannian Gibbons-Hawking metrics with three commuting vector fields, and in Appendix B we shall provide necessary and sufficient conditions for a Killing trajectory to be a conformal geodesic.

Acknowledgements. The work of MD has been partially supported by STFC consolidated grant no. ST/P000681/1.

2. CONFORMAL GEODESICS

In explicit computations involving conformal geodesics, and conformal Killing-Yano tensors it is convenient to adopt the abstract index notation [22]. Thus u^a , $a = 1, 2, \dots, \dim(M)$ denotes a vector, $u^a \nabla_a v^b$ denotes the directional covariant derivative $\nabla_u v$ of another vector v . In a presence of a Riemannian metric g an isomorphism between TM and T^*M is realised by $u_a = g_{ab} u^b$, where now u_a denotes a one-form, and the Einstein summation convention is used. Despite the presence of these indices, no choice of basis has been made.

We shall start–off by reformulating (1.1) in this notation. Let (M, g) be a Riemannian manifold. A curve Γ parametrised by an arc–length s , and with a unit tangent vector u is called a conformal geodesic, or a conformal circle if

$$u^c \nabla_c a^b = -(g_{ef} a^e a^f + L_{ef} u^e u^f) u^b + L^b_c u^c, \quad (2.3)$$

where

$$a^b = u^c \nabla_c u^b \quad \text{is the acceleration so that } g_{ef} a^e u^f = 0, \quad (2.4)$$

$$L_{ef} = \frac{1}{n-2} \left(R_{ef} - \frac{1}{2(n-1)} R g_{ef} \right) \quad (2.5)$$

is the Schouten tensor given in terms of the Ricci tensor R_{ef} and the Ricci scalar R of the metric g . If the metric g is Einstein, i.e. $R_{ab} = (R/n)g_{ab}$ then (2.3) reduces to

$$u^c \nabla_c a^b = -|a|^2 u^b, \quad \text{where } |a|^2 \equiv g_{ef} a^e a^f \geq 0 \text{ is a constant.} \quad (2.6)$$

The system of 3rd order equations (2.6) has long been studied in Riemannian geometry¹ [21, 2]. The terminology is motivated by the observation that a development of a circle Γ starting at a point $x \in M$ is an ordinary circle in $T_x M$. It is also the case that circles on the round sphere S^n are intersections of $S^n \subset \mathbb{R}^{n+1}$ with planes (not necessarily through the origin) in \mathbb{R}^{n+1} . Thus they are indeed circles.

2.1. First Integrals. If $Y \in \Lambda^2(M)$ satisfies the conformal Killing–Yano (CKY) equation

$$\nabla_a Y_{bc} = \nabla_{[a} Y_{bc]} - 2g_{a[b} K_{c]} \quad (2.7)$$

for some one–form $K \in \Lambda^1(M)$, then (if $\dim(M) = 4$)

$$Q = u^a a^b Y_{ab} - u^a K_a \quad (2.8)$$

is constant along the conformal geodesics. In the special case where Y_{ab} is a Kähler form, the one form K is zero and the linear term in (2.8) is not present. In this case the first integral (2.8) has been called the *complex torsion* (though it is real) in [2]. In four dimensions the condition $u^c \nabla_c Q = 0$ with Q given by (2.8) was established in [24]. In [15] it was put in the general context of parabolic geometries.

2.2. Examples.

2.2.1. Circles on \mathbb{R}^n . These are just ordinary circles. Equation (2.6) becomes $\ddot{\mathbf{x}} = -|a|^2 \dot{\mathbf{x}}$, and can be readily solved

$$\mathbf{x}(s) = \mathbf{x}(0) + |a|^{-1} \mathbf{v}(0) \sin(|a|s) + |a|^{-2} \mathbf{a}(0) (1 - \cos(|a|s)),$$

where $\mathbf{x}(0)$ is arbitrary, $\mathbf{v}(0)$ is a unit vector in \mathbb{R}^n , and the vector $\mathbf{a}(0)$ is orthogonal to $\mathbf{v}(0)$, and has squared norm $|a|^2$. We recognise these conformal geodesics as circles centered at $\mathbf{x}(0) + |a|^{-2} \mathbf{a}(0)$, and with Euclidean radius $|a|^{-1}$.

Conformal rescalings of the metric preserve conformal geodesics, so circles on S^n are also circles (intersections of $S^n \subset \mathbb{R}^{n+1}$ with planes in \mathbb{R}^{n+1} which do not necessarily pass through the origin). In particular all conformal geodesics on S^n are closed.

¹In these references equation (2.6) is written as $\nabla_s X_s = |a| Y_s$, $\nabla_s Y_s = -|a| X_s$ where X_s and Y_s are unit vector fields, X_s is the tangent vector to the curve Γ parametrised by the arc-length s , ∇_s is the covariant derivative of g along Γ and the positive constant $|a|^{-1}$ is the *radius* of the circle Γ .

2.2.2. *Magnetic motion on the upper half-plane.* Conformal geodesics on the hyperbolic space can also be obtained from those on \mathbb{R}^n . This time however they are not necessarily closed, and we shall consider this case separately to illustrate the significance of the first integrals in the qualitative behaviour of conformal geodesics.

Let \mathbb{H} be the upper half-plane with a constant curvature metric

$$g = \frac{dx^2 + dy^2}{y^2}, \quad y > 0$$

with Ricci scalar equal to -2 , and a magnetic field $F = B \operatorname{vol}_{\mathbb{H}}$ given by a constant $B > 0$ multiple of the parallel volume form on \mathbb{H} . We chose the potential $\Phi = By^{-1}dx$, so that $F = d\Phi$. The magnetic Lagrangian

$$L = \frac{1}{2} \frac{\dot{x}^2 + \dot{y}^2}{y^2} - B \frac{\dot{x}}{y}$$

admits a conserved energy integral which we set to 1. Taking u to be a unit tangent vector to an integral curve of the corresponding Euler–Lagrange equations we find

$$a \equiv \nabla_u u = BJ(u),$$

where J is the complex structure on \mathbb{H} defined by $g(v, J(w)) = \operatorname{vol}_{\mathbb{H}}(v, w)$. Therefore the circle equations (2.6) hold with $|a|^2 = B^2$, as

$$\nabla_u a = B \nabla_u J(u) = BJ(\nabla_u u) = B^2 J^2(u) = -B^2 u.$$

The two first integrals $|a|^2$, and Q given by (2.8) (where now $Y = F$) are therefore equal as

$$Q \equiv F(u, a) = B \operatorname{vol}_{\mathbb{H}}(u, BJ(u)) = B^2 g(u, u) = B^2.$$

The resulting trajectories are circles of radius $|B|^{-1}$. The behaviour of the circles depends on the value of B (Figure 1)




Figure 1. Geodesics (in blue) and circles on \mathbb{H} .

The circles are open (in the language of [6] the magnetic field is not strong enough to capture the particle) and unbounded if $0 < B < 1$, and closed if $B > 1$. The special case $B = 1$ corresponds to horocircles tangent to the $y = 0$ boundary of \mathbb{H} , or lines $y = \text{const}$ parallel to the x -axis.

3. CIRCLES ON HYPER-KÄHLER FOUR-MANIFOLDS AND THE LORENTZ FORCE

Let (M, g) be an oriented Riemannian–four manifold with a hyper–Kähler structure given by parallel two–forms $\Omega^1, \Omega^2, \Omega^3$. In particular (M, g) is Ricci–flat, so the conformal geodesic equations reduce to (2.6).

Proposition 3.1. *The conformal geodesic equations on a hyper-Kähler four-manifold (M, g) reduce to the Lorentz force equations*

$$u^a \nabla_a u^b = |a| F^b{}_c u^c, \quad (3.9)$$

where $F = \sum_j c^j \Omega^j$ is a constant self-dual Maxwell field, and (c^1, c^2, c^3) are constants such that $|\mathbf{c}| = 1$.

Proof. Each of the Kähler forms gives rise to a first integral (2.8), so there are three of those. These three integrals allow us to integrate the circle equations once, and reduce them, on a level set

$$c^i = \Omega^i(u, a), \quad i = 1, 2, 3. \quad (3.10)$$

to geodesic equations of a charged particle moving in a magnetic field.

If the orientation of (M, g) is chosen so that the Riemann tensor of g is anti-self-dual, then this magnetic field is self-dual. To see how the Lorentz force arises, chose an orthonormal frame such that

$$g = (e^1)^2 + (e^2)^2 + (e^3)^2 + (e^4)^2,$$

and

$$\Omega^1 = e^1 \wedge e^4 + e^2 \wedge e^3, \quad \Omega^2 = e^2 \wedge e^4 + e^3 \wedge e^1, \quad \Omega^3 = e^3 \wedge e^4 + e^1 \wedge e^2.$$

Let u^a and a^a be the components of u, a in the frame of vector-fields dual to e^a . The relations (3.10) can then be solved for a . Using the ordinary vector notation for the components $\mathbf{u} = (u^1, u^2, u^3), \mathbf{a} = (a^1, a^2, a^3)$ we can rewrite (3.10) as $\mathbf{c} = \mathbf{u}a^4 - \mathbf{a}u^4 + \mathbf{u} \wedge \mathbf{a}$, and find

$$\mathbf{a} = \mathbf{c} \wedge \mathbf{u} - u^4 \mathbf{c}, \quad a^4 = \mathbf{u} \cdot \mathbf{c}. \quad (3.11)$$

Note that

$$g(a, u) = \mathbf{a} \cdot \mathbf{u} + a^4 u^4 = 0, \quad g(a, a) = |\mathbf{c}|^2 \quad (3.12)$$

so the first integrals $g(a, a)$ and \mathbf{c} are not independent. The circle equations now reduce to (2.4), or (3.9) which is the Lorentz force equation for a unit mass particle with charge $e = |a|$ in a constant self-dual Maxwell field

$$F = \frac{1}{|a|} (c^1 \Omega^1 + c^2 \Omega^2 + c^3 \Omega^3).$$

In view of (3.12) we can redefine \mathbf{c} to be a unit-vector, and recover the statement of Proposition 3.1. With this interpretation metric geodesics correspond to uncharged particles. \square

To make further progress we need to seek first integrals of (3.9). Any Killing vector on (M, g) which is also tri-holomorphic, i.e. $\mathcal{L}_K \Omega^i = 0$, will give rise to a first integral: If $F = d\Phi$, and the potential one-form Φ is chosen such that $\mathcal{L}_K \Phi = 0$, then

$$\mathcal{K} := K^a (u_a + e\Phi_a) = K^a p_a \quad (3.13)$$

is conserved along conformal geodesics on a level set of the c^i . The momentum p_a is defined by

$$u^a = g^{ab} (p_b - e\Phi_b),$$

which is the usual convention for charged particles.

For (3.9) to be integrable we in particular need the geodesic equation to be integrable, and we need four constants of the motion in involution. One is the Hamiltonian (the unit mass) and the other three need to come from Killing vectors and Killing tensors. If at least two of them arise from Killing vectors then, to be involutive, these must commute and to give integrals for (3.9) they must Lie-drag Ω^i . Now necessarily there is a constant linear combination of them which has purely anti-self-dual covariant derivative (this is Theorem 2.3 in [14], credited to Hitchin by these authors) and this is the characterising property of

the Gibbons-Hawking metrics [12]. Consequently we henceforth restrict to these metrics, but there may be other integrable cases with two or more Killing tensors (see [16, 11] for other applications for Yano tensors to geodesic integrability).

3.1. The Gibbons-Hawking metrics. All hyper-Kähler four-manifolds which admit a tri-holomorphic Killing vector can be put in the Gibbons-Hawking form [12]

$$g = V \mathbf{dx} \cdot \mathbf{dx} + V^{-1} (d\tau + \omega)^2 \quad (3.14)$$

where (V, ω) is a function, and a one-form on \mathbb{R}^3 . Choosing a frame

$$e^4 = V^{-1/2} (d\tau + \omega), \quad e^i = V^{1/2} dx^i, \quad E_4 = V^{1/2} \partial_\tau, \quad E_i = V^{-1/2} (\partial_i - A_i \partial_\tau)$$

one verifies that the two-forms Ω^i are closed iff

$$d\omega = *dV,$$

which in particular implies that V is harmonic.

Consider the Killing vector $K = \partial_\tau$ of metric (3.14). This lowers with the metric to the one-form $K^\flat = V^{-1} (d\tau + \omega)$ and then its derivative is anti-self-dual (the characterising property of the Gibbons-Hawking metrics). It follows that K is tri-holomorphic, i. e.

$$\mathcal{L}_K \Omega^i = 0,$$

where \mathcal{L}_K is the Lie-derivative along K . It therefore follows that K gives rise to three moment maps: for each i there is a function x^i satisfying

$$K \lrcorner \Omega^i = dx^i \quad (3.15)$$

and these x^i are (up to scale) the flat coordinates (x, y, z) in (3.14).

The Killing vector K will give a constant of the motion (3.13) for (3.9). This is conserved for any V , and we will return to it when we consider examples below. It is however not sufficient for integrability.

To integrate (3.9) we need either two more Killing vectors, commuting with each other and with $K = \partial_\tau$, or another Killing vector commuting with K and a Killing tensor with suitable involutivity. The first case is not interesting (see the Appendix A) but for the second we can take motivation from the familiar fact that both the self-dual Taub-NUT metric and the Eguchi-Hanson metric admit Killing tensors [14] to concentrate on these two metrics is §4 and §5.

4. CIRCLES ON ANTI-SELF-DUAL TAUB-NUT

The ASD Taub-NUT metric [12] is a special case of (3.14) with $V = 1 + m/r$ where m is a non-negative constant, and (using spherical polar coordinates on \mathbb{R}^3) $\omega = m \cos \theta d\phi$

$$g = \left(1 + \frac{m}{r}\right) (dr^2 + r^2 (d\theta^2 + \sin^2 \theta d\phi^2)) + \left(1 + \frac{m}{r}\right)^{-1} (d\psi + m \cos \theta d\phi)^2$$

writing ψ for τ to emphasise the Bianchi IX $SU(2)$ symmetry. The singularity at $r = 0$ is removable. The infinity $r \rightarrow \infty$ has the topology of a one-monopole S^1 bundle over S^2 . This behaviour is referred to as asymptotic local flatness (ALF). Thus ASD Taub-NUT is regular everywhere, and is an example of a gravitational instanton.

The metric has $SU(2) \times U(1)$ as its group of isometries where the $U(1)$ action generated by $\partial/\partial\psi$ is tri-holomorphic, i. e.

$$\mathcal{L}_{\partial/\partial\psi} \Omega^i = 0, \quad i = 1, 2, 3$$

and so the self-dual derivative of the corresponding one-form $g(\partial/\partial\psi, \cdot) = V^{-1} (d\psi + \cos \theta d\phi)$ vanishes. The $SU(2)$ action is not tri-holomorphic, and the exterior derivatives of the corresponding Killing vectors do not have a definite duality (but their SD derivatives

are parallel). However for any unit vector $\mathbf{c} \in \mathbb{R}^3$ there exists a linear combination L of the isometric generators of $SU(2)$, such that L Lie-derives $\sum_i c^i \Omega^i$.

4.1. Conformal Killing–Yano tensors and Arnold–Liouville integrability. The ASD Taub–NUT metric admits a Killing–Yano 2–form Z , i. e. a two–form such that

$$\nabla_a Z_{bc} = \nabla_{[a} Z_{bc]}.$$

The self–dual, and anti–self–dual parts of Z are conformal Killing–Yano two–forms which satisfy² (2.7):

$$\begin{aligned} Z &= (d\psi + m \cos \theta d\phi) \wedge dr + (2r + m)(r + m)r\sigma_1 \wedge \sigma_2 \\ &= Y - W, \quad \text{where} \quad *Y = Y, \quad *W = -W \end{aligned}$$

and

$$Y = \frac{r^3}{m} d(V(d\psi + m \cos \theta d\phi)), \quad W = -\frac{(r + m)^3}{m} d(V^{-1}(d\psi + m \cos \theta d\phi)). \quad (4.17)$$

Both Y and W satisfy (2.7) with the same Killing vector $K = g(\partial/\partial\psi, \cdot)$. The CKY two–form Y was discovered in [9], where it was used to show that the ASD Taub–NUT metric is conformal to a scalar–flat Kähler metric with a non–constant conformal factor. Computing the first integral (2.8) corresponding to Y gives an expression which is functionally dependent with the first integrals arising from the Killing vectors $K = \partial/\partial\psi$ and $L = \partial/\partial\phi$, so it does not give anything useful. We instead turn to W , where the resulting first integral (2.8) is non–trivial, but it does not in general commute with the first integrals (3.13) arising from ∂_ψ and ∂_ϕ . To get around this problem we proceed as follows:

Given (3.9) with some F , we may first rotate the (x, y, z) coordinates until $F = -\Omega^3$ and then introduce the spherical polars: if this case is integrable then the general case is integrable. Now

$$\begin{aligned} F &= -\Omega^3 = d\Phi \quad \text{with} \\ \Phi &= -z(d\psi + \omega) + \left(\frac{m}{r} + \frac{1}{2}\right)(ydx - xdy) \\ &= -r \cos \theta d\psi - \left(mr + \frac{1}{2}r^2 \sin^2 \theta\right)d\phi \end{aligned} \quad (4.18)$$

or in polars and with the index raised

$$\Phi^\# = -zV\partial_\psi + \frac{1}{V}\partial_\phi.$$

In this form it is clear that $\Phi^\#$ commutes with the Killing vectors $K := \partial_\psi$ and $L := \partial_\phi$, which in turn commute with each other, leading to two constants of the motion for (3.9).

² The left invariant one–forms $\sigma_i, i = 1, 2, 3$ on the group manifold $SU(2)$ are such that

$$d\sigma_1 + \sigma_2 \wedge \sigma_3 = 0, \quad d\sigma_2 + \sigma_3 \wedge \sigma_1 = 0, \quad d\sigma_3 + \sigma_1 \wedge \sigma_2 = 0. \quad (4.16)$$

These one–forms can be represented in terms of Euler angles by

$$\sigma_1 + i\sigma_2 = e^{-i\psi}(d\theta + i \sin \theta d\phi), \quad \sigma_3 = d\psi + \cos \theta d\phi,$$

where to cover $SU(2) = S^3$ we require the ranges $0 \leq \theta \leq \pi, \quad 0 \leq \phi \leq 2\pi, \quad 0 \leq \psi \leq 4\pi$.

Up to a multiple of 4 the two–form Z coincides with the Killing–Yano two–form found by Gibbons and Ruback [14] (see formula (2.12) in this reference with $n = 1$, and make a coordinate transformation $t = m + 2r$).

This leads to four independent first integrals $\mathcal{I}_\alpha = (\mathcal{K}, \mathcal{L}, \mathcal{W}, \mathcal{H})$ two of which are linear in the velocity, and two of which are quadratic:

$$\begin{aligned}\mathcal{K} &= K^a(u_a + e\Phi_a), & \mathcal{L} &= L^a(u_a + e\Phi_a), \\ \mathcal{H} &= \frac{1}{2}g_{ab}u^a u^b, & \mathcal{W} &= eW_{ac}(F)^c_b u^a u^b - 2\mathcal{H} K^a u_a.\end{aligned}\tag{4.19}$$

To complete the calculation we shall use the Hamiltonian formalism. In the presence of a magnetic field there are two ways to proceed starting from the magnetic Lagrangian $\mathcal{G} = (1/2)g_{ab}u^a u^b + e\Phi_b u^b$. Either define the conjugate momentum

$$p_a = \frac{\partial \mathcal{G}}{\partial u^a} = u_a + e\Phi_a$$

and perform the Legendre transform to find four first integrals

$$\begin{aligned}\mathcal{K} &= p_\psi, & \mathcal{L} &= p_\phi, \\ \mathcal{H} &= \frac{1}{2}g^{ab}(p_a - e\Phi_a)(p_b - e\Phi_b), & \mathcal{W} &= eW^a_c F^{cb}(p_a - e\Phi_a)(p_b - e\Phi_b) - 2\mathcal{H}(\mathcal{K} - eK^a\Phi_a).\end{aligned}$$

These are in involution with respect to the standard symplectic structure on T^*M given by $dp_a \wedge dq^a$, where $q^a = (r, \phi, \theta, \psi)$ and $p_a = (p_r, p_\phi, p_\theta, p_\psi)$. Alternatively define a momentum by

$$P_a = p_a - e\Phi_a,$$

and consider the charged symplectic form

$$\Omega = dP_a \wedge dq^a + eF\tag{4.20}$$

on the 8-dimensional phase-space with coordinates $q^a = (r, \phi, \theta, \psi)$ and $P_a = (P_r, P_\phi, P_\theta, P_\psi)$. The set of first integrals is then $\mathcal{I}_\alpha = (\mathcal{K}, \mathcal{L}, \mathcal{H}, \mathcal{W})$, where

$$\mathcal{K} = P_\psi - er \cos \theta,\tag{4.21a}$$

$$\mathcal{L} = P_\phi - e\left(mr + \frac{1}{2}r^2 \sin^2 \theta\right)\tag{4.21b}$$

$$\mathcal{H} = \frac{1}{2}\left(\frac{r}{r+m}P_r^2 + \frac{1}{r(r+m)}P_\theta^2 + \frac{r+m}{r}P_\psi^2 + \frac{1}{r(r+m)\sin^2 \theta}(P_\phi - m \cos \theta P_\psi)^2\right)\tag{4.21c}$$

$$\begin{aligned}\mathcal{W} &= \frac{e \cos \theta}{r}(rP_r - \tan \theta P_\theta)^2 - \frac{e}{r \cos \theta}P_\theta^2 - \frac{e \cos \theta}{r \sin^2 \theta}(P_\phi^2 + (m^2 - r^2 \sin^2 \theta)P_\psi^2 - 2\frac{m+r \sin^2 \theta}{\cos \theta}P_\psi P_\phi) \\ &\quad - 2\mathcal{H}P_\psi.\end{aligned}\tag{4.21d}$$

These integrals are in involution

$$\{\mathcal{I}_\alpha, \mathcal{I}_\beta\} = 0, \quad \alpha, \beta = 1, \dots, 4$$

with respect to the charged symplectic structure (4.20). We have established

Proposition 4.1. *The conformal geodesic equation on the anti-self-dual Taub–NUT manifold is completely integrable in the sense of Arnold–Liouville.*

Rather than finding the action–angle variables explicitly, we follow [14] and seek to separate the Hamilton–Jacobi equation but for (3.9) rather than for the geodesic equation.

4.2. Separating the Hamilton-Jacobi equation for anti-self-dual Taub-NUT. Following Gibbons-Ruback [14], we introduce parabolic coordinates

$$\eta = r(1 + \cos \theta), \quad \xi = r(1 - \cos \theta), \quad \phi = \phi,$$

so that in the cylindrical polars obtained from (x, y, z) in (3.14)

$$z = \frac{1}{2}(\eta - \xi), \quad \rho = \sqrt{\eta\xi}, \quad \phi = \phi.$$

Now

$$z + i\rho = \frac{1}{2}(\sqrt{\eta} + i\sqrt{\xi})^2,$$

and

$$dz^2 + d\rho^2 = \frac{1}{4}(\eta + \xi)\left(\frac{d\eta^2}{\eta} + \frac{d\xi^2}{\xi}\right),$$

$$V = 1 + \frac{m}{r} = \frac{(\eta + \xi + 2m)}{(\eta + \xi)}, \quad \omega = m \cos \theta d\phi = m \frac{(\eta - \xi)}{(\eta + \xi)} d\phi,$$

so that the anti-self-dual Taub-NUT metric is

$$g = V \left(\frac{1}{4}(\eta + \xi)\left(\frac{d\eta^2}{\eta} + \frac{d\xi^2}{\xi}\right) + \eta\xi d\phi^2 \right) + \frac{1}{V} \left(d\psi + m \frac{(\eta - \xi)}{(\eta + \xi)} d\phi \right)^2.$$

For the geodesic equation first, we seek Hamilton's Principal Function S in the separated form

$$S = E\psi + J\phi + F(\xi) + G(\eta),$$

with constant E, J then the Hamilton-Jacobi equation is

$$\mu^2 = g^{ab} \nabla_a S \nabla_b S$$

where μ is the conserved mass which we will always take to be one.

Algebra leads to

$$4(\eta G_\eta^2 + \xi F_\xi^2) = \mu^2(\eta + \xi + 2m) - J^2\left(\frac{1}{\eta} + \frac{1}{\xi}\right) + 2JEm\left(-\frac{1}{\eta} + \frac{1}{\xi}\right) - E^2\left(\eta + \frac{m^2}{\eta} + \xi + \frac{m^2}{\xi} + 4m\right),$$

and, confirming [14], this separates with a new constant Q :

$$4\eta G_\eta^2 = Q + \mu^2(\eta + m) - \frac{J^2}{\eta} - \frac{2JEm}{\eta} - E^2\left(\eta + \frac{m^2}{\eta} + 2m\right),$$

$$4\xi F_\xi^2 = -Q + \mu^2(\xi + m) - \frac{J^2}{\xi} + \frac{2JEm}{\xi} - E^2\left(\xi + \frac{m^2}{\xi} + 2m\right).$$

The constant Q is quadratic in momenta and must be associated with a quadratic Killing tensor. To obtain the geodesics from S , solve

$$\dot{q}^a = g^{ab} p_b = g^{ab} \nabla_b S,$$

so that

$$\dot{\eta} = \frac{4\eta}{\eta + \xi + 2m} G_\eta, \quad \dot{\xi} = \frac{4\xi}{\eta + \xi + 2m} F_\xi, \quad p_\phi = J, \quad p_\psi = E.$$

We shall now modify this separation procedure to include a Maxwell field present for conformal geodesic motion.

Proposition 4.2. *The Hamilton-Jacobi equation for the conformal geodesics on anti-self-dual Taub-NUT manifold is separable.*

Proof. To include a Maxwell field, the general method (see e.g. [18]) is to suppose the potential is Φ_a , retain

$$\dot{q}^a = g^{ab}(p_b - e\Phi_b) \quad \text{and} \quad H = \frac{1}{2}g^{ab}(p_a - e\Phi_a)(p_b - e\Phi_b),$$

set $p_a = \nabla_a S$ and take the Hamilton-Jacobi equation to be

$$\mu^2 = g^{ab}(\nabla_a S - e\Phi_a)(\nabla_b S - e\Phi_b).$$

For us

$$\begin{aligned} \Phi &= -z(d\psi + \omega) - r^2 \sin^2 \theta \left(\frac{1}{2} + \frac{m}{r} \right) d\phi = -\frac{1}{2}(\eta - \xi)d\psi - \frac{1}{2}(\eta\xi + m(\eta + \xi))d\phi, \\ &= Xd\phi + Yd\psi, \text{ say.} \end{aligned}$$

The changes to the Hamilton-Jacobi equation are

$$J \rightarrow J - eX, \quad E \rightarrow E - eY,$$

leading to

$$\begin{aligned} 4(\eta G_\eta^2 + \xi F_\xi^2) &= \mu^2(\eta + \xi + 2m) - J^2\left(\frac{1}{\eta} + \frac{1}{\xi}\right) + 2JEm\left(-\frac{1}{\eta} + \frac{1}{\xi}\right) - E^2\left(\eta + \frac{m^2}{\eta} + \xi + \frac{m^2}{\xi} + 4m\right) \\ &\quad - eJ(\eta + \xi + 4m) - eE(\eta^2 - \xi^2 + 3m(\eta - \xi)) - \frac{e^2}{4}(\eta(\eta + 2m)^2 + \xi(\xi + 2m)^2), \end{aligned} \quad (4.22)$$

which still separates with a quadratic constant and implied Killing tensor. As before, for the particle paths solve

$$\dot{q}^a = g^{ab}p_b = g^{ab}\nabla_b S.$$

□

Thus in anti-self-dual Taub-NUT the conformal geodesic equation is completely integrable: it admits four first integrals in involution established in Proposition 4.1, and the associated Hamilton-Jacobi equation is separable which reduces the integration of the conformal geodesic equations to quadratures. This completes the proof of Theorem 1.1.

Separating (4.22) with a constant Q gives

$$\dot{\eta} = \frac{4\eta G_\eta}{\eta + \zeta + 2m}, \quad \dot{\zeta} = \frac{4\zeta F_\zeta}{\eta + \zeta + 2m} \quad (4.23)$$

with

$$4\eta G_\eta = U(\eta, E, Q), \quad 4\zeta F_\zeta = U(\zeta, -E, -Q)$$

where

$$\begin{aligned} U(x, E, Q) &= \sqrt{u_0 + u_1x + u_2x^2 + u_3x^3 + u_4x^4}, \\ u_0 &= -4(J + Em)^2, \quad u_1 = 4(Q + m(\mu^2 + 2Je - 2E^2)), \\ u_2 &= 4(\mu^2 - m^2e^2 - E^2 + Je + 3Eme), \quad u_3 = 4e(E - me), \quad u_4 = -e^2. \end{aligned} \quad (4.24)$$

Therefore (4.23) implies that the unparametrised conformal geodesic equations are solvable in terms of elliptic functions:

$$\int \frac{d\eta}{U(\eta, E, Q)} = \int \frac{d\zeta}{U(\zeta, -E, -Q)}.$$

5. CIRCLES ON EGUCHI–HANSON

The Eguchi–Hanson (EH) metric [10]

$$g = f^{-2}dr^2 + \frac{1}{4}r^2(\sigma_1^2 + \sigma_2^2) + \frac{1}{4}r^2f^2\sigma_3^2, \quad \text{where } f = \sqrt{1 - \frac{\alpha^4}{r^4}}, \quad \alpha = \text{const}$$

is another special case of (3.14). It corresponds to

$$V = \frac{1}{r_1} + \frac{1}{r_2}, \quad (5.25)$$

where r_1, r_2 are Euclidean distances from points P_1, P_2 which we can chose to be

$$P_1 = (0, 0, -\alpha), \quad P_2 = (0, 0, \alpha)$$

for constant positive α . To achieve regularity the period of the ψ coordinate in σ_3 should be 2π rather than 4π . Therefore the surfaces of constant r are real projective spaces, and at large r the metric looks like $\mathbb{R}^4/\mathbb{Z}_2$ rather than the Euclidean space. The Eguchi–Hanson is an example of the asymptotically locally Euclidean (ALE) manifold (see e.g. Chapter 9 of [7]).

The isometry group of EH is enhanced to $SO(3) \times U(1)$, but this time it is the $SO(3)$ which acts tri-holomorphically (so that EH can be put in the form (3.14) in many ways). The $U(1)$ action is not tri-holomorphic. The parallel basis of Λ^2_+ is given by

$$\Omega^i = e^i \wedge e^4 + \frac{1}{2}\varepsilon^{ijk}e^j \wedge e^k,$$

with

$$e^1 = \frac{1}{2}r\sigma_1, \quad e^2 = \frac{1}{2}r\sigma_2, \quad e^3 = \frac{1}{2}rf\sigma_3, \quad e^4 = f^{-1}dr. \quad (5.26)$$

Proposition 5.1. *All conformal geodesics on the $SO(3)$ orbits of constant r in the Eguchi–Hanson manifold are of the form*

$$(u^1)^2 + (u^2)^2 + (u^3)^2 = 1, \quad c_1u^1 + c_2u^2 = \frac{rf}{2(1-f^2)}(h^2 - 2(c_1^2 + c_2^2) - 2m_1),$$

$$\text{where } h = \frac{1-f^2}{fr}u^3 + \frac{m_0}{c_1^2 + c_2^2} \quad (5.27)$$

and (u^1, u^2, u^3) depend on (ϕ, ψ, θ) as in (5.33) where $(c_1, c_2, c_3, m_1, m_2, p_1, p_2, p_3)$ are constants of integration.

We shall first perform the general analysis of the circle equations using the Lorentz force system (3.9), and reduce the equations to a system of three 1st order ODEs (5.34a) for three unknown functions. We shall then show that this system is completely solvable under the additional assumption $r=\text{const}$. This will give the proof of Proposition 5.1.

With a determined in terms of u by (3.11) (note that now the components of a and u refer to the basis 5.26) the circle equations reduce to (2.4), which becomes

$$\dot{u}^1 = 2\frac{k^4}{r\sqrt{1-k^4}}u^2u^3 - \frac{\sqrt{1-k^4}}{r}u^1u^4 + c^2u^3 - c^3u^2 - c^1u^4, \quad (5.28a)$$

$$\dot{u}^2 = -2\frac{k^4}{r\sqrt{1-k^4}}u^1u^3 - \frac{\sqrt{1-k^4}}{r}u^2u^4 + c^3u^1 - c^1u^3 - c^2u^4, \quad (5.28b)$$

$$\dot{u}^3 = -\frac{1+k^4}{r\sqrt{1-k^4}}u^3u^4 + c^1u^2 - c^2u^1 - c^3u^4, \quad (5.28c)$$

$$\dot{u}^4 = \frac{\sqrt{1-k^4}}{r}(1 - (u^4)^2) + 2\frac{k^4}{r\sqrt{1-k^4}}(u^3)^2 + c^1u^1 + c^2u^2 + c^3u^3 \quad (5.28d)$$

where $k \equiv \alpha/r$. In this case we can also find the potentials of the triple of parallel two-forms

$$\Omega^1 = e^1 \wedge e^4 + e^2 \wedge e^3 = d\left(-\frac{rf}{2}e^1\right) \quad (5.29a)$$

$$\Omega^2 = e^2 \wedge e^4 + e^3 \wedge e^1 = d\left(-\frac{rf}{2}e^2\right) \quad (5.29b)$$

$$\Omega^3 = e^3 \wedge e^4 + e^1 \wedge e^2 = d\left(-\frac{r}{2f}e^3\right). \quad (5.29c)$$

The relation

$$u = \dot{\phi} \frac{\partial}{\partial \phi} + \dot{\theta} \frac{\partial}{\partial \theta} + \dot{\psi} \frac{\partial}{\partial \psi} + \dot{r} \frac{\partial}{\partial r} = u^a E_a$$

gives

$$u^1 = \frac{1}{2}r(\sin(\theta)\sin(\psi)\dot{\phi} + \cos(\psi)\dot{\theta}) \quad (5.30a)$$

$$u^2 = \frac{1}{2}r(\sin(\theta)\cos(\psi)(\dot{\phi}) - \sin(\psi)\dot{\theta}) \quad (5.30b)$$

$$u^3 = \frac{1}{2}rf(\dot{\psi} + \cos(\theta)\dot{\phi}) \quad (5.30c)$$

$$u^4 = \frac{1}{f}\dot{r}. \quad (5.30d)$$

5.1. Magnetic first integrals, and right invariant vector fields. The equations (5.28) correspond to a forced geodesic motion of a charged particle in a constant self-dual Maxwell field $F = c^1\Omega^1 + c^2\Omega^2 + c^3\Omega^3$

$$u^b \nabla_b u^a = F^a_b u^b. \quad (5.31)$$

Let Φ be a potential for $F = d\Phi$. Using (5.29) we find

$$\Phi = -\frac{1}{2}\left(c_1 r f e^1 + c_2 r f e^2 + c_3 \frac{r}{f} e^3\right).$$

For any tri-holomorphic Killing vector R , such that $\mathcal{L}_R \Phi = 0$ (this can always be arranged by adding a total derivative of a function to Φ), the function

$$p \equiv R^a (u_a + \kappa \Phi_a) \quad (5.32)$$

is a first integral of (5.31) (here κ is a numerical factor and the value of κ can be absorbed into constants c^i).

Given the left-invariant one-forms σ_i , let L_i be the dual basis of left-invariant vector fields, i. e. $L_i \lrcorner \sigma_j = \delta_{ij}$, and

$$[L_i, L_j] = \frac{1}{2} \epsilon_{ijk} L_k.$$

The tri-holomorphic and isometric $SO(3)$ action is generated by the right invariant vector fields R_j on $SO(3)$ such that

$$[L_i, R_j] = 0, \quad [R_i, R_j] = -\frac{1}{2} \epsilon_{ijk} R_k.$$

These vector fields can be expanded in terms of L_i s and therefore also in terms of the dual basis E_1, E_2, E_3 as

$$R_i = M^j_i E_j$$

where the components of the matrix M are explicit functions of (r, ψ, θ, ϕ) , and we have verified that $\mathcal{L}_{R_i} \Phi = 0$. Combining this with (5.32) leads to three first integrals

$$p_i \equiv R_i^a (u_a + \kappa \Phi_a) = M^j_i (u_j + \kappa \Phi_j).$$

These are three linear equations for \mathbf{u} which can be solved as $\mathbf{u} = M^{-1}\mathbf{p} - \kappa\Phi$, or (using the explicit form of M)

$$\begin{aligned} u^1 &= \frac{\kappa}{2}c_1rf + \frac{1}{r}\left(p_2(\cos\phi\cos\psi - \sin\phi\sin\psi\cos\theta) - p_3(\sin\phi\cos\psi - \cos\phi\sin\psi\cos\theta) + p_1\sin\theta\sin\psi\right) \\ u^2 &= \frac{\kappa}{2}c_2rf + \frac{1}{r}\left(-p_2(\cos\phi\sin\psi - \sin\phi\cos\psi\cos\theta) + p_3(\sin\phi\sin\psi - \cos\phi\cos\psi\cos\theta) + p_1\sin\theta\cos\psi\right) \\ u^3 &= \frac{\kappa}{2f}c_3r + \frac{1}{rf}(p_1\cos\theta + p_2\sin\phi\sin\theta + p_3\cos\phi\sin\theta). \end{aligned} \quad (5.33)$$

5.2. A final system of 1st order ODEs. Comparing the expressions (5.33) with (5.30) gives a system of three 1st order equations for three unknown functions $\theta(s), \psi(s), \phi(s)$

$$\dot{\phi} = \frac{1}{r^2}(2p_1 - 2\cot\theta(p_2\sin\phi + p_3\cos\phi)) + \kappa f \frac{c_1\sin\psi + c_2\cos\psi}{\sin\theta}, \quad (5.34a)$$

$$\dot{\theta} = \frac{1}{r^2}(p_2\cos\phi - p_3\sin\phi) + \kappa f(c_1\cos\psi - c_2\sin\psi), \quad (5.34b)$$

$$\begin{aligned} \dot{\psi} &= \kappa\left(f\cot\theta(c_1\sin\psi + c_2\cos\psi) + \frac{1}{f^2}c_3\right) + \frac{2}{r^2f^2\sin\theta}(p_2\sin\phi + p_3\cos\phi) \\ &\quad + \frac{2\alpha^4}{r^6f^2\sin\theta}(\cos\theta^2(p_2\sin\phi + p_3\cos\phi) - p_1\cos\theta\sin\theta). \end{aligned} \quad (5.34c)$$

The equation for \dot{r} can now be obtained from the 1st integral $g(u, u) = 1$.

5.3. An integrable case: circles on $SO(3)$ orbits. In this section we shall establish the complete integrability of the of the conformal geodesic equations under additional assumption that the conformal geodesics lie on surfaces of constant r .

Proof of Proposition 5.1. The circle equations can be reduced to a quadrature under the additional assumption that $\dot{r} = 0$, that is the circles are confined to the orbits of the isometric $SO(3)$ action which are surfaces of constant r . The condition $\dot{r} = 0$ is equivalent to $u^4 = 0$. Set

$$u^1 = \alpha(s), \quad u^2 = \beta(s), \quad u^3 = \gamma(s), \quad \frac{k^4}{r\sqrt{1-k^4}} = R = \text{const.}$$

Equations (5.28) take the form

$$\dot{\alpha} = R\beta\gamma + c_2\gamma - c_3\beta \quad (5.35a)$$

$$\dot{\beta} = -R\alpha\gamma + c_3\alpha - c_1\gamma \quad (5.35b)$$

$$\dot{\gamma} = c_1\beta - c_2\alpha. \quad (5.35c)$$

Algebraic manipulations with these equations give expressions for α and β in terms of γ and its derivatives

$$\begin{aligned} \alpha &= -\frac{c_1}{(c_1^2 + c_2^2)(R\gamma - c_3)}\ddot{\gamma} - \frac{c_2}{c_1^2 + c_2^2}\dot{\gamma} - \frac{c_1}{R\gamma - c_3}\gamma \\ \beta &= -\frac{c_2}{(c_1^2 + c_2^2)(R\gamma - c_3)}\ddot{\gamma} + \frac{c_1}{c_1^2 + c_2^2}\dot{\gamma} - \frac{c_2}{R\gamma - c_3}\gamma. \end{aligned} \quad (5.36)$$

The last equation in (5.35) now holds identically. The first two equations reduce to a single 3rd ODE

$$h\ddot{h} = (\ddot{h} - h^3 + m_0)\dot{h}, \quad \text{where } h(s) = R\gamma(s) - c_3, \quad \text{and } m_0 = c_3(c_1^2 + c_2^2). \quad (5.37)$$

This equation can be integrated in terms of elliptic functions

$$s - m_3 = 2 \int \frac{dh}{\sqrt{-h^4 + 4m_1h^2 - 8m_0h + 4m_2}}, \quad (5.38)$$

where m_0, m_1, m_2, m_3 are constants of integration. Inverting this formula for h , and substituting in the expressions above gives $\alpha(s), \beta(s)$. The elliptic functions are however not necessary to describe the conformal circles. To see this differentiate (5.38) twice, to obtain expressions for \dot{h} and \ddot{h} in terms of h . These expressions are then substituted into (5.36), and (remembering that $\gamma = R^{-1}(h(s) + c_3)$) yield the explicit parametrisation $\mathbf{u}(h)$ of u^1, u^2, u^3 in terms of a parameter h , and five integration constant (note that there is one constraint on the constants $m_0, m_1, m_2, m_3, c_1, c_2$ coming from $g(u, u) = 1$). The formulae for $\mathbf{u}(h)$ are now substituted into (5.33) which yields relations of the form

$$G_1(\phi, \psi, \theta, h) = 0, \quad G_2(\phi, \psi, \theta, h) = 0, \quad G_3(\phi, \psi, \theta, h) = 0$$

where the functions G_i can be read-off from (5.33), and depend on the 8 constants of integrations (five from the procedure above, and p_1, p_2, p_3). The parameter h may be eliminated between these three relations which gives unparametrised conformal circles on the $r = \text{const}$ orbits in the Eguchi–Hanson space. They are given by (5.27). □

5.4. Separability of the Hamilton–Jacobi equation. In this section we shall find another integrable subcase of the conformal geodesic equations on the Eguchi–Hanson background. This will be done by separating the associated Hamilton–Jacobi equation. Following [14] again, we introduce prolate spheroidal coordinates adapted to the potential (5.25)

$$\zeta = \frac{1}{2\alpha}(r_1 + r_2), \quad \lambda = \frac{1}{2\alpha}(r_1 - r_2),$$

so these are constant on confocal ellipses and confocal hyperbolae respectively. Introduce the angles θ_1, θ_2 as the angles supported at p_1, p_2 respectively w.r.t. the z -axis; now

$$r_1 = \alpha(\zeta + \lambda), \quad r_2 = \alpha(\zeta - \lambda), \quad \cos \theta_1 = \frac{\zeta\lambda + 1}{\zeta + \lambda}, \quad \cos \theta_2 = \frac{\zeta\lambda - 1}{\zeta - \lambda}.$$

We need V and ω in these coordinates: for the Eguchi–Hanson metric

$$V = \frac{1}{r_1} + \frac{1}{r_2} = \frac{2\zeta}{\zeta^2 - \lambda^2}, \quad \omega = \Omega d\phi \quad \text{with} \quad \Omega = -2 \frac{\lambda(\zeta^2 - 1)}{\zeta^2 - \lambda^2},$$

with conventions

$$\alpha(\zeta^2 - 1)V_\zeta = \Omega_\lambda, \quad -\alpha(1 - \lambda^2)V_\lambda = \Omega_\zeta.$$

Claim

$$d\rho^2 + dz^2 + \rho^2 d\phi^2 = \alpha^2(\zeta^2 - \lambda^2) \left(\frac{d\zeta^2}{\zeta^2 - 1} + \frac{d\lambda^2}{1 - \lambda^2} \right) + \alpha^2(\zeta^2 - 1)(1 - \lambda^2) d\phi^2$$

whence

$$g_{EH} = V \left(\alpha^2(\zeta^2 - \lambda^2) \left(\frac{d\zeta^2}{\zeta^2 - 1} + \frac{d\lambda^2}{1 - \lambda^2} \right) + \alpha^2(\zeta^2 - 1)(1 - \lambda^2) d\phi^2 \right) + V^{-1}(d\psi + \Omega d\phi)^2.$$

From the geodesic equations we obtain constants of the motion

$$p_\psi = V^{-1}(\dot{\psi} + \Omega \dot{\phi}) := E, \quad p_\phi = \alpha^2 V(\zeta^2 - 1)(1 - \lambda^2) \dot{\phi} + \Omega E := J,$$

from the Killing vectors $K := \partial_\psi$ and $L := \partial_\phi$, so that

$$p_\phi - \Omega p_\psi = \alpha^2 V(\zeta^2 - 1)(1 - \lambda^2) \dot{\phi}.$$

Seek a solution

$$S = E\psi + J\phi + F(\zeta) + G(\lambda)$$

for the Hamilton-Jacobi equation then

$$\mu^2 = \frac{\zeta^2 - 1}{\alpha^2 V(\zeta^2 - \lambda^2)} F_\zeta^2 + \frac{1 - \lambda^2}{\alpha^2 V(\zeta^2 - \lambda^2)} G_\lambda^2 + \frac{(J - \Omega E)^2}{\alpha^2 V(\zeta^2 - 1)(1 - \lambda^2)} + V E^2.$$

Multiply by $\alpha^2 V(\zeta^2 - \lambda^2)$ and simplify to obtain

$$2\alpha\zeta = (\zeta^2 - 1)F_\zeta^2 + (1 - \lambda^2)G_\lambda^2 + J^2\left(\frac{1}{\zeta^2 - 1} + \frac{1}{1 - \lambda^2}\right) + \frac{4JE\lambda}{1 - \lambda^2} + \frac{4E^2}{1 - \lambda^2}, \quad (5.39)$$

which separates.

Up to this point we are following [14]. To extend to the Lorentz force law, we need a Maxwell potential for F_{ab} which commutes with K and L . Now there is a problem: for anti-self-dual Taub-NUT there was no loss in generality in choosing $F = -\Omega^3$ but in Eguchi-Hanson there is, as K will Lie-drag any constant ϕ but L will not. Thus we can obtain integrability for a sub-class of conformal geodesics, those for which $F = -\Omega^3$, but not for all. With this restriction then we have

$$\Omega^3 = -(d\psi + \omega) \wedge dz + V dx \wedge dy.$$

If the potential for $-\Omega_3$ is

$$\Phi = -z(d\psi + \omega) + X d\phi = -z(d\psi + \Omega d\phi) + X d\phi,$$

then

$$dX \wedge d\phi = z d\Omega \wedge d\phi - V \rho d\rho \wedge d\phi.$$

This can be solved by

$$X = z\Omega + 2\alpha\zeta$$

when

$$\Phi = (X - z\Omega)d\phi - z d\psi = 2\alpha\zeta d\phi - \alpha\zeta \lambda d\psi.$$

To modify the Hamilton-Jacobi equation we make the replacements

$$J \rightarrow J + 2e\alpha\zeta, \quad E \rightarrow E - e\alpha\zeta\lambda$$

into (5.39). The added terms are

$$4e\alpha J \frac{\zeta^3}{\zeta^2 - 1} + 4e^2 \alpha^2 \frac{\zeta^4}{\zeta^2 - 1},$$

so this still separates, and this sub-class of the conformal geodesic equations is completely integrable.

6. CIRCLES ON \mathbb{CP}^2

Consider the Fubini–Study metric on \mathbb{CP}^2 . It is Einstein with the Ricci scalar equal to 24, has anti-self-dual (ASD) Weyl tensor, and is Kähler, but with the opposite orientation. That is to say the Kähler form is ASD. The local form of the metric is (see e.g. [13])

$$g_{\mathbb{CP}^2} = \frac{dr^2}{(1+r^2)^2} + \frac{1}{4} \frac{r^2 \sigma_3^2}{(1+r^2)^2} + \frac{1}{4} \frac{r^2}{1+r^2} (\sigma_1^2 + \sigma_2^2). \quad (6.40)$$

The metric is regular everywhere on \mathbb{CP}^2 , and the apparent singularity at $r = 0$ results from using spherical polars. The metric is also regular at $r = \infty$. To see it set $r = \rho^{-1}$. Fixing (ϕ, θ) now gives $g \sim d\rho^2 + (1/4)\rho^2 d\psi^2$ near $\rho = 0$. This is a removable bolt singularity (see [7]) if ψ is periodic with the period 2π . At $\rho = 0$ the three-dimensional orbits of $SU(2)$ collapse to a two-sphere of constant radius. We shall refer to this as the \mathbb{CP}^1 at infinity.

It was shown in [8] that in case of ASD Einstein manifolds with non-zero Ricci scalar there is a one-to-one correspondence between Killing vectors, and self-dual CKY tensors: If K is a Killing vector, then its self dual derivative

$$Y \equiv \frac{1}{2}(dK + *dK)$$

satisfies (2.7). The Fubini–Study metric is a symmetric space $M = SU(3)/U(2)$, so it has 8 Killing vectors generating the Lie algebra $\mathfrak{su}(3)$, and therefore 8 self-dual CKYs. This gives rise to 8 first integrals of the form (2.8). The 9th first integral is given by the parallel CKY which is the anti-self-dual Kähler form. In Section we aim to use these first integrals to find all conformal geodesics on \mathbb{CP}^2 . We shall first prove

Proposition 6.1. *All conformal geodesics on the Fubini–Study metric (6.40) on \mathbb{CP}^2 on the surfaces of constant r are trajectories of the Killing vector $\partial/\partial\psi$, or are of the form*

$$\chi = \operatorname{arccot}\left(\frac{\kappa(c_2 \cos(\kappa s) - \sin(\kappa s))}{Q(c_3 + c_2 \sin(\kappa s) + \cos(\kappa s))}\right), \quad \theta = \operatorname{arccot}\left(\frac{Q - \dot{\chi}}{P \sin \chi}\right), \quad \phi = P \int \frac{\sin \chi}{\sin \theta} ds \quad (6.41)$$

where

$$Q = 2\gamma/r - \gamma r, \quad P = \frac{2\sqrt{(1+r^2)(1-\gamma^2)}}{r}, \quad \kappa = \sqrt{P^2 + Q^2}$$

and r, γ, c_1, c_2, c_3 are constants.

and then deduce the the general form of conformal geodesics:

Theorem 6.2. *All conformal geodesics of the Fubini–Study metric on \mathbb{CP}^2 are of the form $\rho \circ \Gamma$ where Γ is some conformal geodesic on the $r=\text{const}$ surface given in Proposition 6.1, and ρ is an isometry of the Fubini–Study metric.*

Proof of Proposition 6.1. Pick a tetrad of one-forms

$$e^1 = \frac{r}{2\sqrt{1+r^2}}\sigma_1, \quad e^2 = \frac{r}{2\sqrt{1+r^2}}\sigma_2, \quad e^3 = \frac{r}{2(1+r^2)}\sigma_3, \quad e^4 = \frac{dr}{1+r^2},$$

so that the metric (6.40) is $g = \delta_{ab}e^a e^b$, and set

$$u = \alpha E_1 + \beta E_2 + \gamma E_3 + \delta E_4, \quad a = a^1 E_1 + a^2 E_2 + a^3 E_3 + a^4 E_4 \quad (6.42)$$

where $E_a, a = 1, 2, 3, 4$ is the dual tetrad of vector fields, and $\delta_{ab}u^a u^b = 1$. The conformal geodesics equations (2.6) become

$$a^1 = \dot{\alpha} + 2r\beta\gamma + \frac{1}{r}\alpha\delta \quad (6.43a)$$

$$a^2 = \dot{\beta} - 2r\gamma\alpha + \frac{1}{r}\beta\delta \quad (6.43b)$$

$$a^3 = \dot{\gamma} - r\gamma\delta + \frac{1}{r}\gamma\delta \quad (6.43c)$$

$$a^4 = \dot{\delta} + r\gamma^2 - \frac{1}{r}(1 - \delta^2) \quad (6.43d)$$

and

$$\dot{a}^1 = \frac{1}{r}(\beta a^3 - \alpha a^4 - \gamma a^2) - 2r\gamma a^2 - |a|^2 \alpha \quad (6.44a)$$

$$\dot{a}^2 = \frac{1}{r}(\gamma a^1 - \beta a^4 - \alpha a^3) + 2r\gamma a^1 - |a|^2 \beta \quad (6.44b)$$

$$\dot{a}^3 = \frac{1}{r}(\alpha a^2 - \gamma a^4 - \beta a^1) + r\gamma a^4 - |a|^2 \gamma \quad (6.44c)$$

$$\dot{a}^4 = \frac{1}{r}(\alpha a^1 + \beta a^2 + \gamma a^3) - r\gamma a^3 - |a|^2 \delta, \quad (6.44d)$$

together with

$$\delta_{ef} u^e u^f, \quad \delta_{ef} a^e a^f = |a|^2, \quad \text{where } u = (\alpha, \beta, \gamma, \delta).$$

Note that $\delta_{ef} u^e a^f = 0$ is satisfied identically if u^a is unit.

The CKY corresponding to the Killing vector $\partial/\partial\psi$ is

$$\frac{r^2}{r^2 + 1} (e^1 \wedge e^2 + e^3 \wedge e^4),$$

and gives rise to a first integral

$$C_Y := \frac{r^2}{r^2 + 1} (\alpha a^2 - \beta a^1 + \gamma a^4 - \delta a^3) + \frac{2r}{r^2 + 1} \gamma. \quad (6.45)$$

The ASD Kähler form is

$$e^1 \wedge e^2 - e^3 \wedge e^4. \quad (6.46)$$

This gives rise to a first integral

$$C_K := \alpha a^2 - \beta a^1 - \gamma a^4 + \delta a^3. \quad (6.47)$$

Combining the two gives

$$(r^2 + 1)C_Y - r^2 C_K = 2r^3 \gamma (\gamma^2 + \delta^2) + 2r^2 (\gamma \dot{\delta} - \delta \dot{\gamma}). \quad (6.48a)$$

$$(r^2 + 1)C_Y + r^2 C_K = 4r^3 \gamma (\gamma^2 + \delta^2) + 2r^2 (\alpha \dot{\beta} - \beta \dot{\alpha}) + 2r\gamma - 4r^3 \gamma. \quad (6.48b)$$

The first integral $|a|^2$ after some algebra simplifies to

$$\begin{aligned} |a|^2 &= \delta_{ef} \dot{u}^e \dot{u}^f + r^2 \gamma^2 (4 - 3(\gamma^2 + \delta^2)) + 2r\gamma (2\beta \dot{\alpha} - 2\alpha \dot{\beta} + \gamma \dot{\delta} - \delta \dot{\gamma}) \\ &\quad - 2\gamma^2 - \frac{2}{r} \dot{\delta} + \frac{1}{r^2} (1 - \delta^2). \end{aligned} \quad (6.49)$$

We now aim to find all circles such that $\delta = 0$, or equivalently all circles lying on surfaces of constant r . The vector field u is unit, so set $\beta = \sqrt{1 - \alpha^2 - \gamma^2}$. Equations (6.48a, 6.48b) and (6.49) give

$$0 = 2(r\gamma)^3 - r^2(C_Y - C_K) - C_Y \quad (6.50a)$$

$$0 = 2r^2(\alpha \dot{\beta} - \beta \dot{\alpha}) + 4(r\gamma)^3 - 4r^3 \gamma + 2r\gamma - r^2(C_K + C_Y) - C_Y \quad (6.50b)$$

$$A = \dot{\alpha}^2 + \dot{\beta}^2 + (r\gamma)^2 (4 - 3\gamma^2) + 4r\gamma(\beta \dot{\alpha} - \alpha \dot{\beta}) - 2\gamma^2 + \frac{1}{r^2} \quad (6.50c)$$

so that γ is a constant along circles (and this constant depends on r). We also get $a^4 = r\gamma^2 - 1/r$, which is a constant along circles, and $a^3 = 0$. Substituting $\beta = \sqrt{1 - \alpha^2 - \gamma^2}$ into (6.50b) yields a 1st order equation for $\alpha(s)$ which we can integrate to find

$$\alpha(s) = \sqrt{1 - \gamma^2} \sin(Bs + c_1), \quad \beta(s) = \sqrt{1 - \gamma^2} \cos(Bs + c_1), \quad \text{where} \quad (6.51)$$

$$B = \frac{4r^3(\gamma^3 - \gamma) - r^2(C_K + C_Y) + 2r\gamma - C_Y}{2r^2(1 - \gamma^2)}, \quad \text{where } \gamma \neq \pm 1$$

and γ is a root of (6.50a).

To find any additional constraints on B we substitute $u = (\alpha(s), \beta(s), \gamma, 0)$ into the conformal geodesic equations. We find that these equations hold iff

$$B = -3\gamma r, \quad \text{or} \quad \gamma^2 r^2 + \gamma B + 1 = 0. \quad (6.52)$$

The squared norm of acceleration (6.50c) is constant along conformal circles, and is given by

$$|a|^2 = B^2(1 - \gamma^2) + 4r\gamma B(1 - \gamma^2) + r^2\gamma^2(4 - 3\gamma^2) - 2\gamma^2 + \frac{1}{r^2}.$$

We can now compute the *complex torsion* $\tau = C_K/|a|$ and find that it can take any value between $(-1, 1)$ if $B = -3\gamma r$, but $\tau^2 = 1$ if $B = -\gamma r^2 - \gamma^{-1}$. We shall therefore take $B = -3\gamma r$ from now on, and return to this point in the proof of Theorem 6.2.

If $\gamma = \pm 1$, then the circle is a trajectory of the Killing vector $\partial/\partial\psi = r/(r^2 + 1)E_3$, and $\alpha = \beta = 0$ with the corresponding acceleration $a = (r^2 - 1)/rE_4$. In this case we find that $u = E_3$ and

$$\nabla_u \nabla_u u = -\frac{(r^2 - 1)^2}{r^2} u, \quad (6.53)$$

so the circle equations (2.6) hold³ with $|a|^2 = (r^2 - 1)^2/r^2$.

To complete the calculation for $\gamma \neq \pm 1$ we need to find the dependence of the coordinates (ϕ, θ, ψ, r) on the arc-length parameter s along the circles (6.51). Setting

$$u = \dot{\phi} \frac{\partial}{\partial\phi} + \dot{\theta} \frac{\partial}{\partial\theta} + \dot{\psi} \frac{\partial}{\partial\psi} + \dot{r} \frac{\partial}{\partial r},$$

equating it to u in (6.42) and substituting (6.51) yields a coupled system of three 1st order ODEs

$$\dot{\phi} = 2 \frac{\sqrt{1+r^2}}{r} \frac{\sqrt{1-\gamma^2}}{\sin\theta} \sin(\psi + Bs + c_1), \quad (6.54a)$$

$$\dot{\theta} = 2 \frac{\sqrt{1+r^2}}{r} \sqrt{1-\gamma^2} \cos(\psi + Bs + c_1), \quad (6.54b)$$

$$\dot{\psi} = 2 \frac{1+r^2}{r} \gamma - 2 \frac{\sqrt{1+r^2}}{r} \sqrt{1-\gamma^2} \cot\theta \sin(\psi + Bs + c_1), \quad (6.54c)$$

$$\dot{r} = 0. \quad (6.54d)$$

To simplify this slightly, set

$$\chi = \psi + Bs + c_1, \quad P = \frac{2\sqrt{(1+r^2)(1-\gamma^2)}}{r} = \text{const}, \quad Q = B + 2 \frac{1+r^2}{r} \gamma = \text{const} \quad (6.55)$$

so that

$$\dot{\phi} = P \frac{\sin\chi}{\sin\theta}, \quad \dot{\theta} = P \cos\chi, \quad \dot{\chi} = Q - P \cot\theta \sin\chi. \quad (6.56)$$

We claim that this system is solvable by quadratures. Given $\chi(s)$ we find

$$\theta = \text{arccot}\left(\frac{Q - \dot{\chi}}{P \sin\chi}\right), \quad \text{and then} \quad \phi = P \int \frac{\sin\chi}{\sin\theta} ds. \quad (6.57)$$

The final equation for χ arises from the middle equation in (6.56):

$$P^2 \cos(\chi)^3 - 2 \cos(\chi) \dot{\chi}^2 + 3Q \cos(\chi) \dot{\chi} - \cos(\chi)(P^2 + Q^2) + \sin(\chi) \ddot{\chi} = 0.$$

The general solution of this equation is given by

$$\chi = \text{arccot}\left(\frac{\kappa(c_2 \cos(\kappa s) - \sin(\kappa s))}{Q(c_3 + c_2 \sin(\kappa s) + \cos(\kappa s))}\right), \quad \text{where} \quad \kappa = \sqrt{P^2 + Q^2}. \quad (6.58)$$

³We also note that in general $\dot{r} = (1+r^2)\delta$, and use this to verify that $u = E_a$ ($a = 1, 2, 3, 4$) satisfies the circle eq. In the last case we get $|a|^2 = 0$.

This can now be substituted in (6.57) to find the explicit form of θ . It turns out that (according to MAPLE) even the ϕ integral can be computed in terms of elementary functions, but the answer is not illuminating. The integration will in any case introduce another constant - call it c_4 . Thus there exists a seven-dimensional family of conformal circles on surfaces of constant r . This family is parametrised by $c_1, c_2, c_3, c_4, C_Y, C_K, r$, and explicitly given by formulae (6.58, 6.57, 6.55), where B is given by (6.51), and γ is given by (6.50a).

□

We shall now use the conformal geodesics on the $r = \text{const}$ surfaces and conjugate them with the elements of the isometry group $SU(3)$ to find all conformal geodesics on \mathbb{CP}^2 . The fact that all conformal geodesics arise from this construction will follow from the results of [20] which we now recall.

We shall say that two circles Γ_1 and Γ_2 are congruent if there exists a holomorphic isometry (note - all isometries on \mathbb{CP}^2 are holomorphic) ρ such that $\Gamma_1 = \rho \circ \Gamma_2$.

In [20] (Theorem 5.1) it has been shown that two circles on \mathbb{CP}^2 with the same complex torsion

$$\tau = g_{\mathbb{CP}^2}(J(u), a/|a|) \quad (6.59)$$

and curvature $|a|^2$ are congruent. Here J is the complex structure of the ASD Kähler form on \mathbb{CP}^2 and we note that $\tau = C_K/|a|$, where C_K is the first integral (6.47). We also note $\tau^2 \leq |J(u)|^2|a|^2/|a|^2 = 1$, so that $\tau \in [-1, 1]$. Doing the counting, at a point on $r=\text{constant}$ there are 2 dimensions of (unit) velocity u and then 2 dimensions of acceleration orthogonal to velocity so with +3 for the dimension of $r=\text{constant}$ surface and -1 for the one-dimension of the curve that is 6 dimensions of conformal geodesics confined to $r=\text{const}$. But there are 4 dimensions of Killing vectors tangent to $r=\text{constant}$ to move them about, leaving 2 parameters. In the proof of Theorem 6.2 we shall show that these are $|a|$ and τ .

The general counting for unparametrised conformal geodesics on \mathbb{CP}^2 is as follows: There are 3 degrees of freedom of velocity, 3 of acceleration, 4 of position minus 1 for the dimension of the curve. This gives a 9 dimensional space of conformal geodesics on \mathbb{CP}^2 . There are 8 Killing vectors, but an orbit of $SU(3)$ has fixed $|a|$ and τ which is 7 dimensions in the space of conformal geodesics. Therefore there is always 1-dimensional isometry group stabilising a conformal geodesic i.e. every conformal geodesic $\Gamma(s) = \exp(sK) \Gamma(0)$, is a trajectory of some Killing vector K which confirms the results of [19]. This counting is correct as long as the Killing vector K is tangent to the conformal geodesic, rather than being vanishing on it. To examine this we look at fixed points of Killing vectors in \mathbb{CP}^2 . A one parameter subgroup of the isometry group comes from an $SU(3)$ matrix, and fixed points correspond to eigenvectors; if the eigenvalues are distinct then the Killing vector vanishes at isolated points (like ∂_ϕ in the coordinate system (6.40)) but if there is a repeat then the Killing vector vanishes on a \mathbb{CP}^1 in \mathbb{CP}^2 (like ∂_ψ which vanishes at $r = \infty$ - the \mathbb{CP}^1 at infinity). So if a conformal geodesic does not lie on a \mathbb{CP}^1 , then it can't lie in the zero set of a Killing so must be a Killing vector trajectory. If it does lie on a \mathbb{CP}^1 then there is a Killing vector that vanishes everywhere along the geodesic but it does not matter; take it to be the \mathbb{CP}^1 at infinity which is a metric sphere and extrinsically flat, so that the conformal geodesics are genuine circles and are Killing vector trajectories.

Proof of Theorem 6.2. In view of the discussion above, and the results of [20] it is sufficient to show that geodesics on constant r surfaces from Proposition 6.1 can have arbitrary values of the torsion τ , and the curvature $|a|^2$. We shall first consider the case when $\tau = \pm 1$. Then the trajectory of the Killing vector $\partial/\partial\psi$ is a conformal geodesic (6.53) with $|a|^2 = (r^2 - 1)^2/r^2$ which can take any value in $[0, \infty)$.

We shall therefore assume that $\tau^2 < 1$, and show that for any value of $(\tau, |a|)$ there is a corresponding value of (γ, r) . This will be achieved by finding two relations between $(\tau, |a|)$ and (γ, r) .

Recall from (6.52) that $B = -3\gamma r$, or $\gamma^2 r^2 + \gamma B + 1 = 0$. In the latter case we find $C_K^2 = |a|^2$, so that $\tau = \pm 1$, but we already have a conformal geodesics with these values of τ . We shall therefore assume that

$$B = -3\gamma r \quad (6.60)$$

in which case we find

$$|a|^2 = \gamma^2 r^2 - 2\gamma^2 + \frac{1}{r^2} \quad (6.61)$$

and

$$C_K = \frac{\gamma}{r}(1 + r^2 - 2\gamma^2 r^2). \quad (6.62)$$

As a check, calculate

$$1 - \tau^2 = 1 - \frac{C_K^2}{|a|^2} = \frac{(1 - \gamma^2)(1 - 2r^2\gamma^2)^2}{(1 - \gamma^2) + \gamma^2(r^2 - 1)^2} > 0,$$

as expected. Also (6.62) will give the sign of C_K/γ (note that, given a conformal geodesic, then traversing it in the opposite direction switches the signs on C_K and γ). Solving (6.61) for γ^2 gives

$$\gamma^2 = \frac{r^2|a|^2 - 1}{r^2(r^2 - 2)}, \quad (6.63)$$

(once we have r^2 we will need to check that this is less than 1.) This relation together with

$$C_K = -r^3\gamma^3 + r\gamma(1 + |a|^2)$$

(which follows from (6.47) gives a cubic

$$F(X) \equiv (XA - 1)(X - (2A + 1))^2 - TA(X - 2)^3 = 0, \quad \text{where } X \equiv r^2, A \equiv |a|^2, T \equiv \tau^2.$$

We will argue that this cubic has three positive roots which coalesce at $X = 2$ when $A = 1/2$.

Let us deal separately with $r^2 = 2$: from (6.61) for this case $|a|^2 = 1/r^2 = 1/2$ for any γ^2 ; then from (6.62)

$$\tau = \frac{C_K}{|a|} = \gamma(3 - 4\gamma^2),$$

and for any τ with $-1 < \tau < 1$, which is the currently allowed range, inspection of this cubic shows that there are three roots for γ in $(-1, 1)$.

Now we can suppose that $r^2 \neq 2$. Substitute for γ^2 into the square of (6.62):

$$\begin{aligned} C_K^2 &= \frac{\gamma^2}{r^2}(1 + r^2 - 2\gamma^2 r^2)^2 = \frac{r^2|a|^2 - 1}{r^4(r^2 - 2)} \left(1 + r^2 - 2r^2 \left(\frac{r^2|a|^2 - 1}{r^2(r^2 - 2)} \right) \right)^2 \\ &= \frac{(r^2|a|^2 - 1)}{(r^2 - 2)^3} (r^2 - (1 + 2|a|^2))^2. \end{aligned}$$

Rationalise:

$$0 = (r^2|a|^2 - 1)(r^2 - (1 + 2|a|^2))^2 - C_K^2(r^2 - 2)^3, \quad (6.64)$$

and put $r^2 = X$ to define the cubic

$$\begin{aligned} F(X) &:= X^3(|a|^2 - C_K^2) - X^2(1 + 2|a|^2 + 4|a|^4 - 6C_K^2) \\ &\quad + X(2 + 5|a|^2 + 4|a|^4 + 4|a|^6 - 12C_K^2) \\ &\quad - (1 + 4|a|^2 + 4|a|^4 - 8C_K^2), \end{aligned} \quad (6.65)$$

whose vanishing we want. Note that, in F :

- The coefficient of X^3 is

$$|a|^2 - C_K^2 = (1 - \tau^2)|a|^2,$$

which is strictly positive;

- the coefficient of X^2 is

$$-(1 + 2|a|^2(1 - 3\tau^2) + 4|a|^4) = -((1 - 2|a|^2)^2 + 6|a|^2(1 - \tau^2))$$

which is strictly negative;

- the coefficient of X is

$$\begin{aligned} 2 + 5|a|^2 + 4|a|^4 + 4|a|^6 - 12C_K^2 &> 2 - 7|a|^2 + 4|a|^4 + 4|a|^6 \\ &= (2|a|^2 - 1)^2(|a| + 2), \end{aligned}$$

where the first inequality uses $\tau^2 < 1$, so that this coefficient is strictly positive;

- and the constant term is

$$-(1 + 4|a|^2 + 4|a|^4 - 8C_K^2) = -(1 - 2|a|^2)^2 - 8|a|^2(1 - \tau^2),$$

which is strictly negative.

Thus $F(X)$ has no negative roots and at least one positive root. We will see below there are three positive roots, which coalesce at $r^2 = 2$ when $|a|^2 = 1/2$.

Next we note some particular values of F :

$$F(1/|a|^2) = \frac{\tau^2}{|a|^8}(2|a|^2 - 1)^3, \quad F(2) = (2|a|^2 - 1)^3 \quad (6.66)$$

and

$$F(1 + 2|a|^2) = -C_K^2(2|a|^2 - 1)^3. \quad (6.67)$$

Now we can analyse the roots of F

- If $|a|^2 = 1/2$ then $r^2 = 2$ is a root, and by inspection of (6.64) we see that it is three times repeated. By the discussion above, there is now a value of γ giving any value of τ .
- If $2|a|^2 - 1 < 0$ then the evaluations in (6.66) are both negative and that in (6.67) is positive, while

$$1 + 2|a|^2 < 2 < 1/|a|^2.$$

Therefore there is one root in each of the ranges $(0, 1 + 2|a|^2)$, $(1 + 2|a|^2, 2)$ and $(1/|a|^2, \infty)$. All three make γ^2 in (6.63) positive but we shall choose the largest root r_3^2 which has

$$r_3^2 > 1/|a|^2 > 2$$

so that there are positive δ, ϵ with $\delta < \epsilon$ and

$$r_3^2 = 1/|a|^2 + \delta = 2 + \epsilon.$$

Now from (6.63)

$$\gamma^2 = \frac{r_3^2|a|^2 - 1}{r_3^2(r_3^2 - 2)} = \frac{\delta|a|^2}{\epsilon r_3^2} < \frac{\delta}{2r_3^2\epsilon} < \frac{\delta}{4\epsilon} < \frac{1}{4} < 1,$$

so that this choice of r leads to a γ in the allowed range.

- If $2|a|^2 - 1 > 0$ then the evaluations in (6.66) are both positive and that in (6.67) is negative, while

$$1/|a|^2 < 2 < 1 + 2|a|^2.$$

Therefore there is one root in each of the ranges $(0, 1/|a|^2)$, $(2, 1 + 2|a|^2)$ and $(1 + 2|a|^2, \infty)$ and all make γ^2 in (6.63) positive. Again we choose the largest root r_3^2 which therefore has

$$r_3^2 > 1 + 2|a|^2 > 2.$$

We choose positive δ, ϵ with

$$|a|^2 = \frac{1}{2} + \epsilon, \quad r_3^2 = 2 + \delta,$$

so that also

$$r_3^2 = 2 + \delta > 1 + 2|a|^2 = 2 + 2\epsilon \text{ i.e. } \delta > 2\epsilon.$$

Now from (6.63)

$$\begin{aligned} \gamma^2 &= \frac{r_3^2|a|^2 - 1}{r_3^2(r_3^2 - 2)} = \frac{(1/2 + \epsilon)(2 + \delta) - 1}{\delta(2 + \delta)} = \frac{1}{4} \frac{\delta + 4\epsilon + 2\epsilon\delta}{\delta + \delta^2/2} \\ &= \frac{1}{4} \left(1 + \frac{4\epsilon + 2\epsilon^2 - (\delta - 2\epsilon)^2/2}{\delta + \delta^2/2} \right) < \frac{1}{4} \left(1 + \frac{4\epsilon + 2\epsilon^2}{\delta + \delta^2/2} \right) \end{aligned}$$

and with $\delta > 2\epsilon$:

$$< \frac{1}{4} \left(1 + \frac{4\epsilon + 2\epsilon^2}{2\epsilon + 2\epsilon^2} \right) < \frac{3}{4} < 1,$$

so again we have γ in the allowed range.

We have shown that any allowed $(C_K, |a|^2)$ can be obtained from an allowed (r, γ) . □

Remarks.

- A smooth curve Γ parametrised by an arc-length is called a helix of proper order d , if there exist orthonormal vector fields $\{V_1 \equiv \dot{\Gamma}, V_2, \dots, V_d\}$ such that

$$\nabla_s V_j = -\kappa_{j-1} V_{j-1} + \kappa_j V_{j+1}, \quad \text{where } j = 1, \dots, d$$

and $V_0 = V_d \equiv 0$, and the functions $\kappa_j = \kappa_j(s)$ (we set $\kappa_0 = \kappa_d \equiv 0$) where $j = 1, \dots, d-1$ are constant along Γ . Thus a helix of order 1 is a geodesic, and a helix of order 2 is a circle.

Let $\pi : S^5 \rightarrow \mathbb{C}\mathbb{P}^2$ be the Hopf fibration such that the round metric on S^5 is of the form

$$g_5 = (dt + \Theta)^2 + g_{\mathbb{C}\mathbb{P}^2},$$

where $d\Theta = \Omega$ is the Kähler form for $\mathbb{C}\mathbb{P}^2$ (this is also the derivative of $K = \partial/\partial t$ Killing vector). In [2] it is argued that horizontal (with respect to the $U(1)$ connection Θ) lifts of circles on $\mathbb{C}\mathbb{P}^2$ to S^5 are helices of order 2, 3 or 5. Moreover the order 2 occurs iff the first integral (6.47) vanishes. These are the conformal geodesics of S^5 for which velocity and acceleration are both orthogonal to K . A necessary condition is $J(u, a) = 0$ (to see it, differentiate $g_5(K, a)$ along a conformal geodesic on S^5) which we know can be satisfied, so the conformal geodesics of $\mathbb{C}\mathbb{P}^2$ with $J(u, a) = 0$ come from conformal geodesics of S^5 .

- There is another special case, where the conformal geodesic equations can be explicitly integrated with relative ease. Let J be the complex structure of the ASD Kähler form (6.46), and let λ be a non-zero constant. We shall make the ansatz $a = \lambda J(u)$ so that

$$a_1 = -\lambda\beta, \quad a_2 = \lambda\alpha, \quad a_3 = \lambda\delta, \quad a_4 = -\lambda\gamma.$$

Note that the first integral (6.47) is equal to λ , and

$$\nabla_u a = \lambda J^2(u) = -\lambda u, \quad \text{where } \lambda^2 = |a|^2, \quad \text{so that } \tau = 1.$$

The circle equations (2.6) hold identically as a consequence of the definition of a , and can be integrated explicitly. We will not reproduce this calculation here, as we know from Theorem 6.2, that the resulting conformal geodesics lie in the $SU(3)$ orbits of integral curves of E_4 .

APPENDIX A

In this Appendix we establish a non-existence result which narrows down the classification of integrable backgrounds for conformal geodesic equations.

Proposition .3. *There are no non-flat Riemannian Gibbons-Hawking metrics with three commuting Killing vectors.*

Proof. Suppose there are, and we have the Gibbons-Hawking metric (3.14) with two further Killing vectors, say L_1, L_2 , which commute with each other and with $K = \partial_\tau$. We find restrictions on the L_1, L_2 that force flatness. First note that for either L

$$L^a \partial_a (V^{-1}) = L^a \partial_a (K^b K_b) = 2L^a K^b \nabla_a K_b = 2K^a K^b \nabla_a L_b,$$

using $[K, L] = 0$, and this is manifestly zero, so that V is constant along each L . Decompose each L as

$$L^a \partial_a = L^0 \partial_\tau + L^i \partial_i,$$

then the vanishing of the commutator with K gives

$$\partial_\tau L^0 = 0 = \partial_\tau L^i.$$

Next with

$$h_{ab} := g_{ab} - V K_a K_b = V \delta_{ij},$$

where $\delta_{ij} = 0$ unless $i, j = 1, 2, 3$ when it is the usual Kronecker delta, we have

$$\mathcal{L}_L h_{ab} = 0,$$

so that $L^i \partial_i$ is a Killing vector of the flat (Riemannian) 3-metric δ_{ij} . Next, the (i, j) component of

$$0 = \mathcal{L}_L g_{ab} = L^a \partial_a g_{bc} + g_{ac} \partial_b L^a + g_{ba} \partial_c L^a$$

forces L^0 to be constant when it can be set to zero by subtracting a constant multiple of K from L . Finally from the $b = j$ component of

$$0 = \mathcal{L}_L K_b$$

we see that

$$0 = L^i \partial_i (V^{-1} \omega_j) + V^{-1} \omega_j \partial_i L^j,$$

so that L^i Lie-drags ω

Thus any further Killing vector is w.l.o.g. a symmetry of Riemannian flat 3-space, Lie-dragging both V and ω . These can be translations or rotations. Two independent rotations will not commute so L_1, L_2 can be either two independent translations, in which case we may suppose them orthogonal, or a translation and a rotation, in which case commutation requires them to be orthogonal. There are two cases to consider:

- Suppose $L_1 = \partial_x, L_2 = \partial_y$ then up to constants $V = z$ or $V = 1$. Now if $V \neq 1$

$$d\omega = *dV = dx \wedge dy,$$

and there is no ω preserved by both L_1 and L_2 . Thus $V = 1$ when $\omega = 0$ and this class of solutions is flat.

- Suppose $L_1 = \partial_z$ and $L_2 = \partial_\phi$ in cylindrical polars. Then V depends only on ρ and is harmonic so up to constants $V = \log \rho$ or $V = 1$. Now if $V \neq 1$

$$d\omega = \frac{*d\rho}{\rho} = dz \wedge d\phi,$$

and again there is no acceptable solution⁴, except constant V . □

APPENDIX B

In this appendix we shall give necessary and sufficient conditions for a trajectory of a Killing vector in a four-dimensional Einstein space to be a conformal geodesics

Proposition .4. *Let K be a non-null Killing vector on a (pseudo) Riemannian Einstein four manifold (M, g) . The trajectories of K are conformal geodesics if and only if*

$$d|K|^2 \wedge W = 0, \quad \text{where } W = \star_g(K \wedge dK). \quad (\text{B1})$$

Proof. Suppose the Killing vector is K with norm $V^2 = g(K, K)$ and claim that, for any Killing vector

$$\nabla_a K_b = \frac{2}{V} V_{[a} K_{b]} + \frac{1}{2V^2} \epsilon_{ab}{}^{cd} W_c K_d,$$

for some W_a orthogonal to K ; as a form $W = *(K \wedge dK)$. Now the velocity will be $u^a = V^{-1} K^a$ and the acceleration as above will be $a_b = -V^{-1} \nabla_b V$ so we need

$$\begin{aligned} -|a|^2 u_b &= u^c \nabla_c a_b = V^{-1} K^c \nabla_c (-V^{-1} V_b) = -V^{-2} K^c \nabla_b V_c = V^{-2} V^c \nabla_b K_c = \\ &= V^{-2} V^c \left(\frac{2}{V} V_{[b} K_{c]} + \frac{1}{2V^2} \epsilon_{bc}{}^{de} W_d K_e \right), \end{aligned}$$

i. e.

$$-V^{-3} |\nabla V|^2 K_b = -V^{-3} |\nabla V|^2 K_b + \frac{1}{2} V^{-4} \epsilon_b{}^{cde} V_c W_d K_e,$$

so the last term needs to vanish, which is equivalent to (B1). □

As a special case we recover the result of [24], where it was shown that if $W = 0$, then the Killing trajectories are conformal geodesics. We also find that in the case of the Fubini–Study metric (6.40) on \mathbb{CP}^2 the trajectories of $\partial/\partial\psi$ are conformal geodesics, but the trajectories of $K = \partial/\partial\phi$ (or any other Killing vector) are not.

⁴ Note that the claim is not true in the neutral signature: Consider the simplest ASD pp-wave

$$g = dw dx + dz dy + y^2 dw^2,$$

and complete the square to put it in the Gibbons–Hawking form

$$g = V \left(dY dz - \frac{1}{4} dx^2 \right) + V^{-1} \left(dw + \frac{1}{2} (3Y)^{-2/3} dx \right)^2,$$

where $Y = y^3/3$ and $V = (3Y)^{-2/3}$. This metric has a 9D isometry group [5], including the abelian 3D algebra spanned by $(\partial_x, \partial_w, \partial_z)$.

APPENDIX C

In this appendix we shall reformulate the Lorentz force equations, as well as some of the calculations underlying the conformal Killing–Yano tensors in terms of the two component spinor notation [22].

The Hodge $*$ operator of an oriented Riemannian manifold (M, g) is an involution on two-forms, and induces a decomposition

$$\Lambda^2(T^*M) = \Lambda_+^2(T^*M) \oplus \Lambda_-^2(T^*M)$$

of two-forms into self-dual (SD) and anti-self-dual (ASD) components. Locally there exist complex rank-two vector bundles \mathbb{S}, \mathbb{S}' over M equipped with covariantly constant symplectic structures ϵ, ϵ' such that

$$\mathbb{C} \otimes TM \cong \mathbb{S} \otimes \mathbb{S}', \quad g = \epsilon \otimes \epsilon',$$

and

$$\Lambda_+^2 \cong \mathbb{S}'^* \odot \mathbb{S}^*, \quad \Lambda_-^2 \cong \mathbb{S}^* \odot \mathbb{S}'.$$

Elements of \mathbb{S} are of the form $\kappa^A = (\kappa^0, \kappa^1)$, and the spinor indices $A, B, C, \dots = 0, 1$ are lowered using the symplectic form ϵ_{AB} with $\epsilon_{01} = 1$. Therefore any two-form admits a spinor decomposition

$$F_{ab} = f_{AB}\epsilon_{A'B'} + \tilde{f}_{A'B'}\epsilon_{AB},$$

where $f_{AB} = f_{(AB)}$, and $\tilde{f}_{A'B'} = \tilde{f}_{(A'B')}$. In particular the parallel basis $(\Omega^1, \Omega^2, \Omega^3)$ of Λ_+^2 on a hyper-Kähler four manifold decomposes as

$$\Omega^i{}_{ab} = \phi^i{}_{A'B'}\epsilon_{AB}, \quad i = 1, 2, 3.$$

Go back to the Lorentz force equation (3.9) with self-dual Maxwell field F , and set

$$F_{ab} = |a|\phi_{A'B'}\epsilon_{AB}$$

where $\phi_{A'B'} = \phi_{(A'B')}$ is a valence two symmetric spinor. The Lorentz force equation takes the form

$$u^{CC'}\nabla_{CC'}u_{AA'} = -|a|\phi_{A'B'}u^{B'}{}_A$$

and $\phi_{A'B'}\phi^{A'B'} = 2$. The ASD two-form W from (4.17) is

$$W_{ab} = \omega_{AB}\epsilon_{A'B'},$$

and the CKY equation (2.7) takes the form

$$\nabla_{AA'}\omega_{BC} = \epsilon_{A(B}K_{C)A'}, \quad \text{where } K = \partial/\partial\psi. \quad (\text{C1})$$

First note that ω_{AB} gives rise to three first integrals of the geodesic equations. To see it consider the quadratic in velocity $\omega_{AB}\phi^i{}_{A'B'}u^{AA'}u^{BB'}$. Then its rate of change is

$$u^{CC'}\nabla_{CC'}(\omega_{AB}\phi^i{}_{A'B'}u^{AA'}u^{BB'}) = \frac{1}{2}u^{AA'}\phi^i{}_{A'B'}K_A{}^{B'} = \frac{1}{2}u^{AA'}\nabla_{AA'}x^i,$$

by (C1) and (3.15) so that three constants of the motion for the geodesic equation are defined by

$$q^i := \omega_{AB}\phi^i{}_{A'B'}u^{AA'}u^{BB'} - \frac{1}{2}x^i.$$

It is known from [24] that a Killing spinor will give constants of the motion for the conformal geodesic equation: consider $\omega_{A'B'}u^{AA'}a_A{}^{B'}$ then its rate of change following (2.3)-(2.4) is

$$u^{CC'}\nabla_{CC'}(\omega_{AB}u^{AA'}a_{A'}{}^B) = -\frac{1}{2}K_b a^b = -\frac{1}{2}u^c\nabla_c(K_b u^b),$$

so that the following is a conserved quantity for the conformal geodesic equation:

$$\omega_{AB}u^{AA'}a_{A'}^B + \frac{1}{2}K_bu^b.$$

In the present case, when we have the relation

$$u_{AA'}a^A_{B'} = u_{A(A'}a^A_{B')} = -\frac{1}{2}|a|\phi_{A'B'}$$

for a real constant spinor $\phi_{A'B'}$ between velocity and acceleration, this constant becomes

$$-|a|u^{AA'}u^{BB'}\phi_{A'B'}\omega_{AB} + \frac{1}{2}K_bu^b. \quad (\text{C2})$$

This is the first integral \mathcal{W} in (4.19).

REFERENCES

- [1] Adachi, T. (1995) Kähler Magnetic Flows for a Manifold of Constant Holomorphic Sectional Curvature. *Tokyo J. Math.* **18**
- [2] Adachi, T. Maeda, S. and Udagawa, S. (1995) Circles in a complex projective space. *Osaka J. Math.* **32**, 709-719.
- [3] Bailey T. N. and Eastwood, M. G. (1990) Conformal circles and parametrizations of curves in conformal manifolds. *Proc. Amer. Math. Soc.*, **108** 215221.
- [4] Bailey, T. N., Eastwood, M. G. and Gover, A. R. (1994) Thomas's structure bundle for conformal, projective and related structures. *Rocky Mountain J. Math.* **24**, 1191-1217.
- [5] Casey, S., Dunajski, M. and Tod, K. P. (2013) Twistor geometry of a pair of second order ODEs *Comm. Math. Phys.* **321**, 681-701.
- [6] Comtet, A. (1987) On the Landau Levels on the Hyperbolic Plane , *Ann. of Phys. Solitons, Instantons, and Twistors*, Oxford University Press. **173**, 185-209.
- [7] Dunajski, M. (2009)
- [8] Dunajski, M. and Tod, K. P. (2010) Four Dimensional Metrics Conformal to Kahler, *Math. Proc. Camb. Phil. Soc.* **148**, 485.
- [9] Dunajski, M. and Tod, P. (2014) Self-Dual Conformal Gravity *Comm. Math. Phys.* **331**, 351373.
- [10] Eguchi, T. and Hanson, A. (1979) Selfdual solutions to Euclidean gravity. *Annals of Phys* **120**. 82105.
- [11] Frolov, Vv. P., Krtous, P., and Kubiznak, D. (2017) Black holes, hidden symmetries, and complete integrability *Living Reviews in Relativity*.
- [12] Gibbons, G. W. and Hawking, S. W. (1978) Gravitational Multi - Instantons. *Phys. Lett.* **B78** 430.
- [13] Gibbons, G. W. and Pope, C. (1978) $\mathbb{C}\mathbb{P}^2$ As a Gravitational Instanton. *Commun. Math. Phys.* **61** 239.
- [14] Gibbons, G. W. and Ruback, P. J. (1988) The hidden symmetries of multicentre metrics. *Comm. Math. Phys.* **115**, 267-300.
- [15] Gover, A. R., Snell, D. and Taghavi-Chabert, A. (2018) Distinguished curves and integrability in Riemannian, conformal, and projective geometry. [arXiv:1806.09830](https://arxiv.org/abs/1806.09830)
- [16] Houri, T., Takeshi, O. and Yukinori, Y. (2007) Closed conformal Killing-Yano tensor and geodesic integrability. *J. Phys.* **A41**.
- [17] Joachimsthal, F. (1843) Observationes de lineis brevissimis et curvis curvaturae in superficiebus secundi gradus, *JRAM* **26**, 155-171.
- [18] Landau, L. D. and Lifshitz (1962) *The Classical Theory of Fields*, Pergamon Press, 1962.
- [19] Maeda, S. and Adachi, T. (2001) Integral curves of Killing vector fields in a complex projective space. *Mem. Fac. Sci. Eng. Shimane Univ.* **B34**, 6185.
- [20] Maeda, S. and Ohnita, Y. (1989) Helical geodesic immersion into complex space forms, *Geom. Dedicata* **30**, 93-114.
- [21] Nomizu, K. and Yano, K. (1974) On circles and spheres in Riemannian geometry. *Mathematische Annalen* **210**, 163-170.
- [22] Penrose, R. & Rindler, W. (1987, 1988) *Spinors and space-time. Two-spinor calculus and relativistic fields*. Cambridge Monographs on Mathematical Physics. Cambridge University Press, Cambridge.
- [23] Sihlan, J. and Zadnik, V. (2018) Conformal theory of curves with tractors. [arXiv:1805.00422](https://arxiv.org/abs/1805.00422)
- [24] Tod, K. P. (2012) Some examples of the behaviour of conformal geodesics. *J. Geom. Phys.* **62** 1778-1792.

DEPARTMENT OF APPLIED MATHEMATICS AND THEORETICAL PHYSICS, UNIVERSITY OF CAMBRIDGE,
WILBERFORCE ROAD, CAMBRIDGE CB3 0WA, UK.

E-mail address: `m.dunajski@damtp.cam.ac.uk`

THE MATHEMATICAL INSTITUTE, OXFORD UNIVERSITY, WOODSTOCK ROAD, OXFORD OX2 6GG, UK.

E-mail address: `tod@maths.ox.ac.uk`