

On linear structure of non-commutative operator graphs

G.G.Amosov^{*1} and A.S. Mokeev^{†1,2}

¹Steklov Mathematical Institute of Russian Academy of Sciences, 8 Gubkina St., Moscow, 119991 Russia

²Chebyshev Laboratory of St. Petersburg State University, 14th Line V.O. 29B, St. Petersburg 199178, Russia

March 12, 2024

Abstract

We continue the study of non-commutative operator graphs generated by resolutions of identity covariant with respect to unitary actions of the circle group and the Heisenberg-Weyl group as well. It is shown that the graphs generated by the circle group has the system of unitary generators fulfilling permutations of basis vectors. For the graph generated by the Heisenberg-Weyl group the explicit formula for a dimension is given. Thus, we found a new description of the linear structure for the operator graphs introduced in our previous works.

Keywords non-commutative operator graphs, covariant resolution of identity, quantum antiques

1 Introduction

In this article we give a clarification of our current results on non-commutative operator graphs generated by covariant resolutions of identity [1, 2]. A linear subspace \mathcal{V} in the algebra of all bounded linear operators on the Hilbert space H is called a non-commutative operator graph if

$$A \in \mathcal{V} \Rightarrow A^* \in \mathcal{V}; \quad I \in \mathcal{V}.$$

^{*}gramos@mi-ras.ru

[†]alexandrmokeev@yandex.ru

These objects allow to determine a possibility of zero-error correction for a system of unitary operators representing errors in quantum information transmission (for explicit description of this theory see [3, 4]). Basically, we are interested only in graphs satisfying the Knill-Laflamme condition [5]. It is said that the graph \mathcal{V} satisfies the Knill-Laflamme condition, if there is an orthogonal projection P_K on the subspace $K \subset H$ such that $P_K \mathcal{V} P_K = \mathbb{C} P_K$, the projection P_K is called an anticlique and the subspace K is known as a quantum error correcting code.

Some properties and illustrating examples of non-commutative operators graphs could be given in terms of resolutions of identity covariant with respect to an action of locally-compact groups. Studying of non-commutative operator graphs generated by covariant resolution of identity was initiated in [6]. Let G be a locally compact group and \mathfrak{B} is a sigma-algebra generated by compact subsets of G , then the set of positive operators $\{M(B), B \in \mathfrak{B}\}$ is said to be a resolution of identity if

$$M(\emptyset) = 0, \quad M(G) = I,$$

$$M(\cup_j B_j) = \sum_j M(B_j), \quad B_k \cap B_l = \emptyset \text{ for } k \neq l, \quad B_j \in \mathfrak{B}.$$

Let $(U_g)_{g \in G}$ be a projective unitary representation of G . Define the action of this group on operator A as

$$\rho_g(A) = U_g A U_g^*.$$

Then $M(B)$ is said to be covariant with respect to this action if

$$\rho_g(M(B)) = M(gB).$$

Suppose that a graph \mathcal{V} is generated by $M(B)$ in the following sense

$$\mathcal{V} = \overline{\text{span}\{M(B), B \in \mathfrak{B}\}}.$$

Note that if H is finite dimensional there is no need to take a closure in the last equation.

2 Linear structure of \mathcal{V}

At first let us define the graph introduced in [1]. Consider the finite dimensional Hilbert space $H = \mathbb{C}^d \otimes \mathbb{C}^d$ and the orthonormal basis $|k\rangle$, $1 \leq k \leq d$ in \mathbb{C}^d . Then, vectors

$$|\psi_{sn}\rangle = \frac{1}{\sqrt{d}} \sum_{k=1}^d e^{\frac{2\pi i s k}{d}} |k \ k - n \bmod d\rangle, \quad 1 \leq s \leq d, \quad (1)$$

are known as the generalized Bell states. Let P_s , $1 \leq s \leq d$ be the projections on the subspaces

$$H_s = \text{span}\{|\psi_{sn}\rangle, 1 \leq n \leq d\}.$$

Define a unitary representation of the circle group $\mathbb{T} = [0, 2\pi]$ with the operation $+/mod(2\pi)$ by the formula

$$\varphi \rightarrow U_\varphi = \sum_{s=1}^d e^{i\varphi s} P_s, \quad \varphi \in \mathbb{T}.$$

Also consider a set of projections in \mathbb{C}^d of the form

$$Q_j = \sum_{k=1}^d |j \ j - k \ mod \ d\rangle \langle j \ j - k \ mod \ d|, \quad 1 \leq j \leq d. \quad (2)$$

The following corollary defines examples of the graphs mentioned above.

Corollary 1 [1]. *Given j , $1 \leq j \leq d$, the projection (2) generates the graph $\mathcal{V}_j = \text{span}\{U_\varphi Q_j U_\varphi^*, \varphi \in \mathbb{T}\}$ for which the projections $\{P_s, 1 \leq s \leq d\}$ are anticliques.*

Let us consider a set of unitary operators defined as follows

$$W_n = \sum_{l=1}^d \sum_{r=1}^d |\psi_{l+n \ mod \ d \ r}\rangle \langle \psi_{lr}|, \quad 0 \leq n \leq d-1. \quad (3)$$

The unitaries from (3) act as permutations on the set of vectors (1).

Below we clarify a structure of the graphs determined by Corollary 1.

Theorem 1. *All graphs $\mathcal{V}_j \equiv \mathcal{V}$ coincide for $1 \leq j \leq d$. Moreover, one can choose (3) as generators of \mathcal{V} .*

Proof.

For operators $U_\varphi Q_j U_\varphi^*$ we have such expansion in basis $|\psi_{kn}\rangle$

$$\begin{aligned} U_\varphi Q_j U_\varphi^* &= \sum_{s=1}^d \sum_{l=1}^d e^{i\varphi(s-l)} P_s Q_j P_l = \\ &= \sum_{s=1}^d \sum_{l=1}^d \sum_{r=1}^d e^{i\varphi(s-l)} P_s |j \ j - r \ mod \ d\rangle \langle j \ j - r \ mod \ d| P_l = \\ &= \sum_{s=1}^d \sum_{l=1}^d \sum_{r=1}^d \sum_{n=1}^d \sum_{h=1}^d e^{i\varphi(s-l)} |\psi_{sn}\rangle \langle \psi_{sn}| j \ j - r \ mod \ d\rangle \langle j \ j - r \ mod \ d| \psi_{lh}\rangle \langle \psi_{lh}| = \end{aligned}$$

$$\begin{aligned} & \frac{1}{d} \sum_{s=1}^d \sum_{l=1}^d \sum_{r=1}^d \sum_{n=1}^d \sum_{h=1}^d e^{i\varphi(s-l)} \delta_{nr} \delta_{hr} e^{-\frac{2\pi i j(s-l)}{d}} |\psi_{sn}\rangle \langle \psi_{lh}| = \\ & \frac{1}{d} \sum_{s=1}^d \sum_{l=1}^d e^{i\varphi(s-l)} e^{-\frac{2\pi i j(s-l)}{d}} \sum_{r=1}^d |\psi_{sr}\rangle \langle \psi_{lr}| = \frac{1}{d} \sum_{n=0}^{d-1} e^{i(\varphi - \frac{2\pi j}{d})n} W_n, \end{aligned} \quad (4)$$

where unitaries W_n are determined by (3). To complete the proof we need to show that $W_n \in \mathcal{V}_j$ for all j , $1 \leq j \leq d$. Substituting $\varphi = \varphi_{kj} = \frac{2\pi(k+j)}{d}$ into (4) we obtain

$$\frac{1}{d} \sum_{n=0}^{d-1} e^{i\frac{2\pi kn}{d}} W_n = U_{\varphi_{kj}} Q_j U_{\varphi_{kj}}^* \in \mathcal{V}_j$$

for all $0 \leq k \leq d-1$. The result follows. \square

3 Examples for Heisenberg-Weyl Group

Now we want to give an analogous clarification of linear structure for the example of the non-commutative operator graph generated by the same technique implied for the Heisenberg-Weyl group. Unlike the circle group this group is non-commutative.

At first we need to introduce the construction from [2]. Let \mathfrak{H} , $\dim \mathfrak{H} = d$ be a finite-dimensional Hilbert space with the fixed orthogonal basis $|j\rangle$, $0 \leq j \leq d-1$. Unitary operators S and M on \mathfrak{H} given by the formulae

$$S|j\rangle = |j+1 \bmod d\rangle, \quad M|j\rangle = e^{\frac{2\pi i}{d}j} |j\rangle$$

define the representation of Heisenberg-Weyl group G_d of rank d .

Consider $H = \mathfrak{H} \otimes \mathfrak{H}$ with the orthonormal basis $|kj\rangle$, $0 \leq k, j \leq d-1$. Also consider the orthogonal basis in H consisting of entangled vectors

$$h_k^0 = \frac{1}{\sqrt{d}} \sum_{j=0}^{d-1} e^{\frac{2\pi i k j}{d}} |jj\rangle,$$

$$h_k^j = (I \otimes S^j) h_k^0.$$

Define subspaces in H spanned by systems of the following entangled vectors

$$H_j = \text{span}\{h_k^j, 0 \leq k \leq d-1\}.$$

Let π be the map transmitting S and U to unitary operators in H as follows

$$\pi(S)h_k^j = h_{k+1 \bmod d}^j,$$

$$\pi(M)h_k^j = e^{\frac{2\pi i}{d}k}h_k^j.$$

It is shown in [2] that the map π can be extended to a unitary representation of the group G_d .

Also we need special sums of matrix units

$$y_{ml} = \sum_{k=0}^{d-1} |h_m^k\rangle \langle h_l^k|$$

and their linear combinations

$$h_0 = \sum_{m=0}^{d-1} y_{mm} \equiv I,$$

$$h_p = \sum_{m=0}^{d-1} y_{m+p \bmod d, m} + y_{m, m+p \bmod d}, \quad 1 \leq p \leq d-1.$$

Resulting linear structure of the operator graph generated by representation π was given in [2] as follows $\mathcal{V} = \text{span}\{h_p, 0 \leq p \leq d-1\}$. Here we give the explicit value for a dimension of this graph.

Theorem 2. *A dimension of the graph is given by the formula*

$$\begin{cases} \dim \mathcal{V} = \frac{d-1}{2} + 1 \text{ for odd } d \\ \dim \mathcal{V} = \frac{d}{2} + 1 \text{ for even } d \end{cases} \quad (5)$$

Proof.

Equation

$$h_p = h_{p'}$$

or equivalently

$$\sum_{m=0}^{d-1} y_{m+p \bmod d, m} + \sum_{m=0}^{d-1} y_{m, m+p \bmod d} = \sum_{m'=0}^{d-1} y_{m'+p' \bmod d, m'} + \sum_{m'=0}^{d-1} y_{m', m'+p' \bmod d}$$

holds true if

$$p' = d - p$$

because a first summand in the left hand side equals to a second summand in the right hand side and vice-versa. Thus, $\dim \mathcal{V}$ expresses by (5). \square

The next corollary follows from the proof of the theorem.

Corollary 2. *The graph \mathcal{V} has the system of generators*

$$\begin{cases} \mathcal{V} = \text{span}\{h_p, 0 \leq p \leq \frac{d-1}{2} + 1\} \text{ for odd } d \\ \mathcal{V} = \text{span}\{h_p, 0 \leq p \leq \frac{d}{2} + 1\} \text{ for even } d \end{cases}$$

4 Conclusion

We gave a clarification of the examples of the operator graphs given in [1]. It is shown that all these examples coincide and it is possible to choose unitary generators of a given graph, so we have the same situation as in [2]. For examples given in [2] we gave the explicit formula for dimensions of the graphs and specified their linear structure.

5 Acknowledgments

The work is supported by Russian Science Foundation under the grant no. 19-11-00086.

References

- [1] G.G. Amosov, A.S. Mokeev, “On non-commutative operator graphs generated by covariant resolutions of identity, ”Quantum Information Processing **17** (12), 325 (2018).
- [2] G.G. Amosov, A.S. Mokeev, “On non-commutative operator graphs generated by reducible unitary representation of the Heisenberg-Weyl group, ”International Journal of Theoretical Physics, doi: 10.1007/s10773-018-3963-4.
- [3] G.G. Amosov, A.S. Mokeev, “On construction of anticliques for non-commutative operator graphs, ”Zap. Nauchn. Sem. S.-Peterburg. Otdel. Mat. Inst. Steklov. (POMI) **456**, 5–15 (2017); J. Math. Sci. **234** (3), 269-275 (2018).
- [4] R. Duan, S. Severini, A. Winter, “Zero-error communication via quantum channels, noncommutative graphs and a quantum Lovasz theta function, ”IEEE Trans. Inf. Theory. **59** 1164-1174 (2013).
- [5] E. Knill, R. Laflamme, “Theory of quantum error-correcting codes, ”Phys. Rev. A **55** (2), 900 (1997).
- [6] G.G. Amosov, “On general properties of non-commutative operator graphs, ”Lobachevskii Journal of Mathematics **39** (3), 304-308 (2018).
- [7] N. Weaver, “A quantum Ramsey theorem for operator systems,”Proceedings of the American Mathematical Society **145** (11), 4595-4605 (2017).