

Bimetric interactions based on metric congruences

Mikica Kocic

*Department of Physics & The Oskar Klein Centre,
Stockholm University, AlbaNova University Centre, SE-106 91 Stockholm*

E-mail: mikica.kocic@fysik.su.se

ABSTRACT: In massive gravity and bigravity, spin-2 interactions are defined in terms of a square root matrix that involves two metrics. In this work, the interactions are constructed using a congruence matrix between the metrics. It is established that the primary square root matrix function is the only power series solution to the equations of motion for the congruence. Moreover, the shift vector redefinition that is used in the bimetric ghost-free proofs follows from the $N + 1$ form of the equations of motion. The analysis also gives an insight into the vielbein formulation of spin-2 interactions since the bimetric formulation in terms of a congruence is algebraically equivalent to the unconstrained vielbein formulation.

KEYWORDS: Modified gravity, Massive gravity, Bigravity, Ghost-free bimetric theory

Contents

1	Introduction	1
1.1	Metric congruences	4
1.2	Parametrization of orthogonal transformations	5
2	Congruence based bimetric scalar potential	6
3	Congruences in the $N+1$ formalism	9
3.1	The $N+1$ decomposition	9
3.2	Variation of the $N+1$ form of the potential	10
4	Discussion	12
	Appendices	13
A	Recovering indices	13
B	Detailed derivations	14
C	Bimetric actions with a scalar potential	17

1 Introduction

General relativity is the classical theory of nonlinear self-interactions for a massless spin-2 field governed by the Einstein–Hilbert action. The context of this paper are its extensions: de Rham–Gabadadze–Tolley (dRGT) massive gravity [1–3] and the Hassan–Rosen (HR) bimetric theory or bigravity [4–6]. Massive gravity is a nonlinear theory of a single massive spin-2 field, while bigravity is a nonlinear theory of two interacting spin-2 fields having both massless and massive modes. These theories have been studied extensively over the past years; for reviews, see [7–9].

Massive gravity and bigravity are classically consistent theories, free of instabilities such as the Boulware–Deser ghost [10]. This is not a coincidence, but due to the particular structure of the bimetric scalar potential proposed in [1] with a compact general form [3],

$$\mathcal{S}^{\text{int}}(g, f) = \sum_{n=0}^d \beta_n \int d^d x \sqrt{-g} e_n(\sqrt{g^{-1}f}). \quad (1.1)$$

The potential involves two metric fields g and f ; it is specifically constructed in terms of the square root matrix $(g^{-1}f)^{1/2}$ using the elementary symmetric polynomials e_n [11], where the interaction is parametrized by constants β_n . In massive gravity, the metric g carries dynamics while f is a reference nondynamical field. In bigravity, both g and f are dynamical, each having its own Einstein–Hilbert term.

The vielbein formulation of spin-2 interactions was constructed in [12]. The vielbein based potential has the following expression (given in [12], cf. [13]),

$$\mathcal{S}^{\text{int}}(g, f) = \sum_{n=0}^d \frac{\beta_n}{n!(d-n)!} \int \epsilon_{A_1 A_2 \dots A_d} L^{A_1} \wedge \dots \wedge L^{A_n} \wedge E^{A_{n+1}} \wedge \dots \wedge E^{A_d}. \quad (1.2)$$

Here, ϵ denotes the Levi-Civita symbol, and the vielbeins are represented by one-forms $E^A = E^A{}_{\mu} dx^{\mu}$ and $L^A = L^A{}_{\mu} dx^{\mu}$. The associated metrics are,

$$g = g_{\mu\nu} dx^{\mu} dx^{\nu} = \eta_{AB} E^A E^B, \quad f = f_{\mu\nu} dx^{\mu} dx^{\nu} = \eta_{AB} L^A L^B, \quad (1.3)$$

where $\eta = \text{diag}(-, +, \dots, +)$ is the metric of the local Lorentz frame. In the constrained version, $h = \eta_{AB} L^A E^B$ is symmetric, which is equivalent to having a real square root $(g^{-1}f)^{1/2} = E^{-1}L$ in matrix notation [14]. Consequently, the constrained vielbein formulation is equivalent to the metric formulation since (1.1) and (1.2) become equal. The metric formulation of the multivielbein theory was treated in [15].

Purpose of this work. We address several issues related to bimetric interactions. First, the question if there exists a more general bimetric formulation of the potential (1.1) yet *equivalent* to the vielbein formulation that is based the symmetric polynomials; then, if it exists, what governs the selection of the square root in the *constrained* case? Namely, the square root matrix $(g^{-1}f)^{1/2}$ must represent a well-defined tensor field. Notwithstanding, it may have multiple branches, and besides not being real, it can be solved ad hoc from the matrix equation $g^{-1}f = S^2$ which may have an infinite number of solutions (an example of such $g^{-1}f$ is shown in figure 1b). In the vielbein formulation, the square root selection is concealed inside the symmetrization condition giving no definite choice for S [14].

Another issue concerns the initial value problem for the unconstrained vielbein formulation. The ghost-free proof in [12] assumes the simultaneous $N+1$ decomposition with arbitrarily boosted vielbeins. This is in general not possible since one cannot ensure that an arbitrary vielbein L can simultaneously be triangularized with E already being in the triangular form. A typical example is shown in figure 1a, where the null cones of the two metrics doubly intersect; in this case the simultaneous $N+1$ decomposition of g & f , and the simultaneous triangularization of E & L , are not possible.




Figure 1. Null cones for the metric configurations which do not allow a boosted $N+1$ decomposition wherein both vielbeins are in the triangular form. Configuration (a) does not have a real square root, and (b) has an infinite number of nonprimary real square roots that are not tensor fields (for more details about possible metric configurations see [16]).

Summary of results. An overview of the paper with the key results is shown in figure 2. In section 2, we construct bimetric interactions in terms of a general congruence $f = S^\top g S$ between the metric fields g and f . This congruence based metric formulation is algebraically equivalent to the unconstrained vielbein formulation [12]. We solve the equation of motion for the congruence, which necessarily gives the primary square root $S = (g^{-1}f)^{1/2}$ as the congruence field. In section 3, the equations are solved using the $N+1$ decomposition. The obtained solution is the symmetrization of the spatial metrics together with the shift vector redefinition that was used in the bimetric ghost-free proofs [3, 4]. The rest of the introduction is devoted to a mathematical background on metric congruences.




Figure 2. Summary of results and paper structure. The starting point is the bimetric potential whose algebraic structure is based on the elementary symmetric polynomials $e_n(S)$ dependent on a general congruence S between the metrics $f = S^\top g S$. The on-shell conditions for S are obtained in two alternative ways: covariantly in section 2, and using the $N+1$ decomposition in section 3. Both approaches give the congruence $S = (g^{-1}f)^{1/2}$. The derivation in section 2 is more formal than in section 3, and it shows that the square root is necessarily a primary matrix function.

Notation. The metric signature is mostly positive $(-, +, \dots, +)$. The spacetime dimension is d . In the space-plus-time decomposition, $d = N+1$, hats over spacetime objects denote their spatial restrictions. The equations are mostly written in matrix notation.

Hence, the expressions are preferably stated using (1,1)-tensors with the default down-up contractions. Matrices do not naturally represent metrics and require transposes on the left side of the metric symbol in matrix notation. Examples of how to restore the indices are given in appendix A. Spacetime (world) indices are denoted by μ, ν, \dots and their spatial restrictions by i, j, \dots , while Lorentz (local frame) indices are denoted by A, B, \dots and their spatial restrictions by a, b, \dots .

1.1 Metric congruences

Here we review some basic properties of symmetric bilinear forms (metrics) and their isometries (congruent transformations or congruences); for more details see [17–19]. Two examples of congruences are orthogonal transformations and moving frames (vielbeins).

Let g be a nondegenerate symmetric bilinear form on a finite-dimensional real vector space V . The pair (V, g) is called a real symmetric bilinear space. Two symmetric bilinear spaces (V, g) and (\tilde{V}, f) are isometric iff there exists an invertible linear transformation $S : \tilde{V} \rightarrow V$ such that,

$$f(v, w) = g(S(v), S(w)), \quad \text{for all } v, w \in \tilde{V}. \quad (1.4)$$

That is, f is a precomposition of g with the map S (or the pullback of g by S). This corresponds to the matrix congruence $f = S^\top g S$ in some basis. In linear algebra, the linear transformation S is usually referred to as an isometry. To avoid possible confusion with the Killing symmetries of the metric fields, we adopt another frequently used name “congruence” or “congruent transformation.” By Sylvester’s law of inertia, the congruent transformations preserve the signature of the symmetric bilinear forms. They form a group of automorphisms on the same vector space $\tilde{V} = V$.

The point-wise notion of a congruence can be lifted from linear algebra to differential geometry (i.e., from a tensor to a tensor field). A possible obstruction is that the congruence field might not be defined as a global section of the fiber bundle. A typical example is a vielbein, which globally exists iff the manifold is parallelizable [20]. Nevertheless, one can always assume the local existence of sections, which we employ here.

Adjoint maps. Two linear transformations A and A' are adjoint with respect to g , iff

$$g(A(v), w) = g(v, A'(w)), \quad \text{for all } v, w \in V. \quad (1.5)$$

In matrix notation we have $A' = g^{-1} A^\top g$. This relation can be used to define the adjoint of A (having a regular g). A self-adjoint transformation A is such that $A' \equiv A$. An example of self-adjoint transformations with respect to g are the matrix functions of $g^{-1}f$.

Orthogonal group. An orthogonal transformation is a congruence X whose adjoint is equal its inverse, $X' \equiv X^{-1}$. Beware that “adjointness” is always stated with respect to some symmetric bilinear form (here g). The orthogonal group $O(V, g)$ comprise all the automorphisms which preserve g , that is, $g = X^\top g X$. Note that a congruence between (\tilde{V}, f) and (V, g) always contains excessive degrees of freedom because it can only be determined up to a residual orthogonal transformation of either g or f .

1.2 Parametrization of orthogonal transformations

A special orthogonal transformation X can be parametrized by a skew-symmetric bilinear form $\omega = -\omega^\top$ using the Cayley transform [21],

$$X := (g + \omega)^{-1}(g - \omega). \quad (1.6)$$

It is straightforward to verify that $g = X^\top g X = g^\top$ since,

$$\omega = X^\top \omega X = -\omega^\top, \quad g + \omega = X^\top (g + \omega) X. \quad (1.7)$$

The parametrization (1.6) is special since $\det X = 1$. In d -dimensions, X contains $d(d-1)/2$ degrees of freedom in the components of ω . Any X can be factored as a noncommutative product of orthogonal transformations $X = X_1 X_2 \cdots$ where each X_i contains partial degrees of freedom in the corresponding skew-symmetric ω_i .

The $N+1$ parametrization of boosts and rotations. Here we present a recursive definition of the orthogonal transformation, which is useful to triangularize vielbeins in their entire form. The same procedure is implicitly employed in the Cholesky decomposition to factor positive definite matrices into the product of a lower triangular matrix and its transpose [21]; see also the Cholesky–Banachiewicz and Cholesky–Crout algorithms [22].

Let η be a d -dimensional metric of either Lorentzian or Euclidean signature, where $\hat{\delta}$ is its $(d-1)$ -dimensional restriction,

$$\eta = \text{diag}(\sigma, 1, \dots, 1), \quad \sigma = \pm 1, \quad \hat{\delta} = \text{diag}(1, \dots, 1). \quad (1.8)$$

The orthogonal transformation $\eta = X^\top \eta X$ can be parametrized [19],

$$X := \begin{pmatrix} \varepsilon x & \varepsilon \sigma \xi' \\ \xi & \hat{X} \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & \hat{R} \end{pmatrix}, \quad x := \sqrt{1 - \sigma \xi' \xi}, \quad \hat{X} := \sqrt{\hat{I} - \sigma \xi \xi'}, \quad \xi' := \xi^\top \hat{\delta}, \quad (1.9)$$

where ξ is a $(d-1)$ -dimensional vector, $\hat{\delta} = \hat{R}^\top \hat{\delta} \hat{R}$ is an orthogonal transformation of the Euclidean restriction, \hat{I} is the identity map, and $\varepsilon = \pm 1$ denotes an optional reflection (ignored in the following). The parameter space is confined to $\sigma \xi' \xi \leq 1$, which is satisfied for any ξ in the Lorentzian case $\sigma = -1$.

The factorization (1.9) is recursive. For $d = 4$ and $\sigma = -1$, we have $X = X_1 X_2 X_3$ where X_1 comprises three-parameter boosts of the 4×4 Minkowski metric, X_2 contains two-parameter rotations of the 3×3 Euclidean metric, and X_3 is a one-parameter rotation of the 2×2 Euclidean metric. This effectively gives a polar decomposition of the orthogonal transformation X where the Lorentz boosts X_1 are parametrized by an arbitrary spatial Lorentz vector p (see Proposition 1.13 in [19]),

$$X_1 = \Lambda = \begin{pmatrix} \lambda & p' \\ p & \hat{\Lambda} \end{pmatrix}, \quad \lambda := \sqrt{1 + p' p}, \quad \hat{\Lambda} := \sqrt{\hat{I} + p p'}, \quad p' := p^\top \hat{\delta}. \quad (1.10)$$

The nonvanishing components of ω in the corresponding Cayley parametrization (1.6) are $\omega_{0a} = p^b \delta_{ba} / (\lambda + 1)$. Also note that,

$$\hat{\Lambda} = \sqrt{\hat{I} + p p'} = \hat{I} + \frac{1}{\lambda + 1} p p', \quad \hat{\Lambda}^{-1} = \hat{I} - \frac{1}{\lambda(\lambda + 1)} p p'. \quad (1.11)$$

The boosts can be reparametrized through $v := p/\lambda$ where,

$$p = \hat{\Lambda}v = \lambda v, \quad p' = v^\top \hat{\Lambda}^\top \hat{\delta} = \lambda v^\top \hat{\delta}, \quad v^\top \hat{\delta} v < 1. \quad (1.12)$$

Similar expressions hold for rotations. For instance, X_3 reads,

$$X_3 = \begin{pmatrix} \sqrt{1-\varrho^2} & -\varrho \\ \varrho & \sqrt{1-\varrho^2} \end{pmatrix}, \quad \varrho^2 \leq 1. \quad (1.13)$$

2 Congruence based bimetric scalar potential

We consider the scalar potential based on the elementary symmetric polynomials e_n ,

$$S^{\text{int}}(g, f) = \int d^d x \sqrt{-g} \mathcal{V}(S), \quad \mathcal{V}(S) := \sum_{n=0}^d \beta_n e_n(S), \quad (2.1)$$

where S is an arbitrary congruence between the metric fields g and f ,¹

$$f = S^\top g S, \quad f_{\mu\nu} = S^\rho{}_\mu S^\sigma{}_\nu g_{\rho\sigma}. \quad (2.2)$$

The congruence S is determined up to a local orthogonal transformation X of g ,

$$g = X^\top g X, \quad (2.3)$$

such that $S = X S_0$, where S_0 possibly depends on the degrees of freedom in g and f , but not on X . The additional components of the congruence are removed by on-shell conditions for X . All the fields are assumed to be regular since any singularity punctures the manifold.

The orthogonal transformation X can be parametrized by a skew-symmetric tensor field ω using the Cayley transform (1.6). The Einstein–Hilbert term which involves g is not affected by X . Hence, the equations of motion for S are obtained by varying the potential \mathcal{V} , which we summarize in the following.² The full derivation is in appendix B.

The variation of \mathcal{V} with respect to X reads,

$$\delta \mathcal{V} = \sum_{n=1}^d \beta_n \sum_{k=1}^n (-1)^k e_{n-k}(S) \text{Tr}(X^{-1} S^k \delta X), \quad (2.4)$$

where the further variation of X with respect to ω gives,

$$\delta X = -2(g + \omega)^{-1} \delta \omega (g + \omega)^{-1} g. \quad (2.5)$$

¹ Note that (2.2) is not the most general congruence between g and f since the metrics can be on different manifolds. Then, to write (2.2), we need a diffeomorphism between the manifolds (with the pullback of one of the metrics), which in turn introduces a diagonal group of common diffeomorphisms. This generalization, however, does not affect the presented analysis (see appendix C for more details).

² A similar variation was done in [15]; therein, however, S was a priori assumed to be the square root.

Substituting δX into $\delta\mathcal{V}$ yields,

$$\delta\mathcal{V} = -2 \sum_{n=1}^d \beta_n \sum_{k=1}^n (-1)^k e_{n-k}(S) \operatorname{Tr} \left[(g - \omega)^{-1} g S^k (g + \omega)^{-1} \delta\omega \right], \quad (2.6)$$

For the skew-symmetric ω , we have $\delta\omega^\top = -\delta\omega$ and it holds,

$$\partial \operatorname{Tr}(C\omega) / \partial\omega = (C - C^\top) / 2, \quad (2.7)$$

where C is an arbitrary (2,0)-tensor. Hence (see appendix B),

$$\frac{\partial\mathcal{V}}{\partial\omega} = - \sum_{n=1}^d \beta_n \sum_{k=1}^n (-1)^k e_{n-k}(S) \left\{ (g - \omega)^{-1} \left[g S^k - (g S^k)^\top \right] (g + \omega)^{-1} \right\}. \quad (2.8)$$

Having the nonsingular g (and $g \pm \omega$), the equations of motion $\partial\mathcal{V}/\partial\omega = 0$ become,

$$\sum_{n=1}^d \beta_n \sum_{k=1}^n (-1)^k e_{n-k}(S) \left[S^k - (g^{-1} S^\top g)^k \right] = 0. \quad (2.9)$$

The self-adjoint $S = g^{-1} S^\top g$ trivially solves (2.9), and since the β -parameters are arbitrary, this choice is unique for the independent g and f . When combined with $f = S^\top g S$, the self-adjoint S yields the equation $S^2 = g^{-1} f$, which is solved by a matrix function $S = \sqrt{g^{-1} f}$.

In the following we shall investigate in more detail the structure of all possible solutions to (2.9), showing that the primary square root is indeed the unique choice for the congruence field. Let us express (2.9) in terms of,

$$A := g^{-1} f. \quad (2.10)$$

Since $f = S^\top g S$ implies $g^{-1} S^\top g = A S^{-1}$, we have,

$$\sum_{n=1}^d \beta_n \sum_{k=1}^n (-1)^k e_{n-k}(S) \left[S^k - (A S^{-1})^k \right] = 0. \quad (2.11)$$

This is a nonlinear matrix equation with respect to S , depending only on the nonsingular A . Beside the powers of S , the equation contains the elementary symmetric polynomials of S . The elementary symmetric polynomials are the principal scalar invariants, which are part of the Cayley–Hamilton theorem. Subsequently, any analytic (power series) solution to the equation (2.11) is always a point-wise polynomial. Therefore, (2.11) possibly has three kinds of solutions:

- (i) $S = F(A)$ as a primary matrix function of A ,
- (ii) $S = F(A)$ as a nonprimary matrix function of A ,
- (iii) S is an isolated (incident) solution that is not a function of A .

In the first two cases, the solution is a matrix *function*. The matrix functions can be defined in many equivalent ways: by Jordan canonical form, polynomial interpolation, and Cauchy integral theorem [23]. All these definitions produce *primary* matrix functions.




Figure 3. The field of primary square roots where one point (or maybe some region) admits an infinite number of nonprimary solutions in addition to the primary ones. A well-defined interaction requires that different paths always agree on the chosen branch of the square root at the crossings.

A *nonprimary* matrix function is an “equation solving function” which cannot be expressed as a primary matrix function [23, 24]. An example of a nonprimary function is the square root of a matrix having the same eigenvalue a in different Jordan blocks where the chosen signs are not the same in $\pm\sqrt{a}$. Only if F is a primary function, $F(A)$ is a polynomial in A for all A . Nonprimary matrix functions do not allow perturbations [25].

Now, for any symmetric and nonsingular g and f , Corollary 1.34 from [24] asserts that $gF(g^{-1}f)$ and $fF(g^{-1}f)$ are also symmetric. This holds regardless of F being primary or nonprimary. Therefore, we necessarily have $gS = S^Tg$ and,

$$A = g^{-1}f = S^2, \quad F(A) = \sqrt{A}. \quad (2.12)$$

However, S is a well-defined tensor field only if $F(A)$ is a primary function (S can only then be expressed as a polynomial in A). This governs an unambiguous definition of the bimetric theory given in [16], which warrants the existence of a spacetime interpretation by singling out the principal square root whose eigenvalues lie in the open right complex half-plane, in which case the square root is unique.

An example of how a nonprimary solution can be encountered in the field of primary square roots is shown in figure 3. This happens whenever $g^{-1}f$ has the same eigenvalue in different Jordan blocks. A peculiar metric configuration that only has nonprimary real square roots (with no real primary roots) is shown in figure 1b.

As earlier noted, the orthogonal transformation X can be factored into several pieces by splitting the degrees of freedom, $X(\omega) = X(\omega_1)X(\omega_2) \cdots X(\omega_n)$. This will result in the equations of motion that form a *coupled system* for the parameters $\omega_1, \omega_2, \dots, \omega_n$. Such a case emerges in the following section where we employ the $N+1$ decomposition.

The presented analysis is covariant; no particular space-plus-time decomposition was used or assumed. Nevertheless, the on-shell congruence S (the *real* primary square root of $g^{-1}f$) enables the foliation of spacetime with common spacelike hypersurfaces [16]. The proper $N+1$ spacetime foliation is a prerequisite for the ghost-free proofs that are based on the canonical formalism in the metric [3–6] and the vielbein formulation [12].

3 Congruences in the $N+1$ formalism

In this section we derive the equations of motion for the congruence-based potential in the $N+1$ formalism. We shall see that the variation of this kind of potential gives the shift redefinition from [3, 4]. The results are also applicable to the equations of motion for the boost parameter p of the vielbein based potential from [12] when given in the $N+1$ form.

3.1 The $N+1$ decomposition

We first recall how the space-plus-time split [26–28] works for metrics and vielbeins. One can always find a coordinate patch where one of the metrics is properly $N+1$ decomposed, for instance g ,

$$g = \begin{pmatrix} -N^2 + \nu^\top \hat{g} \nu & \nu^\top \hat{g} \\ \hat{g} \nu & \hat{g} \end{pmatrix}. \quad (3.1)$$

Here, N is the lapse function, ν is the shift vector, and \hat{g} is the spatial projection of g . The lapse and the shift are already parts of the timelike vector in a vielbein. The spatial metric can further be factored $\hat{g} = e^\top \hat{\delta} e$, which fully defines g in terms of the vielbein E ,³

$$g = E^\top \eta E, \quad E = \begin{pmatrix} N & 0 \\ e \nu & e \end{pmatrix}, \quad E^{-1} = \begin{pmatrix} 1/N & 0 \\ -\nu/N & e^{-1} \end{pmatrix}. \quad (3.2)$$

On the other hand, the simultaneous $N+1$ split of f is not possible in general. Nevertheless, f can always be decomposed in an arbitrary non-null chart into the time restriction f^{00} of f^{-1} , the nonsingular space restriction \hat{f} of f (not necessarily positive definite), and the remaining shear μ (an apparent “shift” vector),

$$f = \begin{pmatrix} (f^{00})^{-1} + \mu^\top \hat{f} \mu & \mu^\top \hat{f} \\ \hat{f} \mu & \hat{f} \end{pmatrix}, \quad f^{-1} = \begin{pmatrix} f^{00} & -f^{00} \mu^\top \\ -f^{00} \mu & \hat{f}^{-1} + f^{00} \mu \mu^\top \end{pmatrix}. \quad (3.3)$$

The sign of f^{00} is arbitrary and depends on the chosen spacetime foliation, where f^{00} is negative if and only if \hat{f} is positive definite. In the same chart, f can be given in terms of a general vielbein L ,

$$f = L^\top \eta L, \quad L = \begin{pmatrix} u^0 + u^\top q & u^\top \\ \hat{L} q & \hat{L} \end{pmatrix}, \quad (3.4)$$

where \hat{L} , q , u , and u^0 are arbitrary, and \hat{L} is nonsingular. Equating (3.4) and (3.3) yields,

$$\hat{f} = -u u^\top + \hat{L}^\top \hat{\delta} \hat{L}, \quad \mu = q - u^0 \hat{f}^{-1} u. \quad (3.5)$$

Only in the case of the proper space-plus-time foliation, we have a real lapse function M ,

$$M^2 = -(f^{00})^{-1} > 0, \quad (3.6)$$

and the vielbein L can be put into a triangular form.

³ If we do not decompose \hat{g} , we end up with the D and Q variables introduced in [3]; this favors one metric (more precise, the metric f), and the expressions become “asymmetric.” Consequently, the duality between the metrics on the exchange $g \leftrightarrow f$ and $\beta_n \leftrightarrow \beta_{d-n}$ would not be explicit.

Boosted vielbeins. The formal claim is that a general vielbein can be triangularized by a local Lorentz transformation of the form (1.10) if and only if the apparent lapse of the associated metric is real in a given chart [29]. In other words, the condition on the coordinate system to be able to extract a real M from (3.3) is the same as to put L into the triangular form L_0 by Λ^{-1} so the triangular L_0 is boosted to $L = \Lambda L_0$,

$$\Lambda = \begin{pmatrix} \lambda & v^\top \hat{\delta} \hat{\Lambda} \\ \hat{\Lambda} v & \hat{\Lambda} \end{pmatrix}, \quad L_0 = \begin{pmatrix} M & 0 \\ m\mu & m \end{pmatrix}, \quad L = \begin{pmatrix} \lambda^{-1} M + \hat{\Lambda} m (M m^{-1} v + \mu) & v^\top \hat{\delta} \hat{\Lambda} m \\ \hat{\Lambda} m (M m^{-1} v + \mu) & \hat{\Lambda} m \end{pmatrix}. \quad (3.7)$$

Comparing (3.7) and (3.4), one concludes that only those general vielbeins that satisfy,

$$u^\top (\hat{L}^\top \hat{\delta} \hat{L})^{-1} u < 1, \quad (3.8)$$

can be triangularized by a Lorentz transformation since the boosts are restricted to an open ball $v^\top \hat{\delta} v < 1$. This is the same condition as for $f^{00} < 0$, that is, \hat{f} to be positive definite $\hat{f} > 0$ which follows from (3.5). For more details see Lemma 2 in [29].

3.2 Variation of the $N+1$ form of the potential

We again start from the bimetric potential (2.1) given in terms of a congruence $f = S^\top g S$, where S is now related to the vielbeins (3.2) and (3.4) by $S = E^{-1} L$. This form of the potential is algebraically equivalent to the unconstrained vielbein formulation [12].

The $N+1$ form of the potential reads (see appendix B for the derivation),

$$N e_n(S) = N e_n(\hat{B}) + u^0 e_{n-1}(\hat{B}) + \sum_{k=1}^n (-1)^{k-1} e_{n-k}(\hat{B}) u^\top \hat{B}^{k-1} (q - \nu), \quad (3.9)$$

where $\hat{B} := e^{-1} \hat{L}$ and $e_n(\hat{B}) = 0$ for $n > d - 1$. The potential is manifestly linear in N , ν , u^0 , and q . Note that \hat{B} is a congruence,

$$\hat{B}^\top \hat{g} \hat{B} = \hat{L}^\top \hat{\delta} \hat{L} = \hat{f} + u u^\top. \quad (3.10)$$

Let us consider a coordinate patch where both f and g admit the proper $N+1$ decomposition, that is, a patch where both E and L can be simultaneously triangularized. In this case, we have from (3.7) and (3.4),

$$u^0 = \lambda^{-1} M, \quad \text{boosted lapse,} \quad (3.11a)$$

$$q = M m^{-1} v + \mu, \quad \text{geometric mean shift,} \quad (3.11b)$$

$$u^\top = v^\top \hat{\delta} \hat{\Lambda} m = v^\top \hat{\delta} \hat{L}, \quad \text{space/time boost,} \quad (3.11c)$$

$$\hat{L} = \hat{\Lambda} m, \quad \text{boosted spatial vielbein.} \quad (3.11d)$$

Hence, $\hat{B} = e^{-1} \hat{\Lambda} m$ and,

$$\begin{aligned} N e_n(S) &= N e_n(\hat{B}) + \lambda^{-1} M e_{n-1}(\hat{B}) + \\ &+ \sum_{k=1}^n (-1)^{k-1} e_{n-k}(\hat{B}) v^\top \hat{\delta} \hat{L} \hat{B}^{k-1} (M m^{-1} v + \mu - \nu). \end{aligned} \quad (3.12)$$

This equation is linear in N , ν , M , and μ , which is a necessary condition for the ghost-free proofs [3–6, 12]. Note, however, that we still do not have a square root at this point; the velocity vector v and the residual spatial rotation of either e or m are not constrained.

To vary (3.12) with respect to v , the potential must be rewritten so the elementary symmetric polynomials conveniently depend only on the spatial vielbeins e and m . The derivation is lengthy and relegated to an ancillary Mathematica notebook (wherein the calculations are also verified). A final form that is suitable for variation reads,

$$\begin{aligned} N e_n(S) = N \left\{ e_n(m e^{-1}) + \sum_{k=1}^n (-1)^{k-1} e_{n-k}(m e^{-1}) p^\top \hat{\delta} (m e^{-1})^k e \left[\frac{1}{\lambda+1} e^{-1} p - N^{-1} \nu \right] \right\} \\ + M \left\{ e_{d-n}(e m^{-1}) + \sum_{k=1}^{d-n} (-1)^{k-1} e_{d-n-k}(e m^{-1}) p^\top \hat{\delta} (e m^{-1})^k m \left[\frac{1}{\lambda+1} m^{-1} p + M^{-1} \mu \right] \right\}. \end{aligned} \quad (3.13)$$

Introducing the derivatives of the elementary symmetric polynomials,

$$Y_n(X) := \sum_{k=0}^n (-1)^{n+k} e_k(X) X^{n-k} = \partial e_{n+1}(X) / \partial X^\top, \quad (3.14)$$

the equation (3.13) can be written,

$$\begin{aligned} N e_n(S) = N e_n(m e^{-1}) + M e_{n-1}(m e^{-1}) \\ + p^\top \hat{\delta} Y_{n-1}(m e^{-1}) m \left[\frac{1}{\lambda+1} (N e^{-1} + M m^{-1}) p + \mu - \nu \right]. \end{aligned} \quad (3.15)$$

Variation of the potential with respect to p gives (see appendix B),

$$\begin{aligned} N \partial e_n(S) / \partial p = \hat{\delta} Y_{n-1}(m e^{-1}) m \left[(N e^{-1} + M m^{-1}) v + \mu - \nu \right] \\ + \frac{1}{\lambda+1} \left[\hat{\Lambda}^\top \hat{\delta} Y_{n-1}(m e^{-1}) m (N e^{-1} + M m^{-1}) \hat{\Lambda}^{-1} \right]^\top v \\ - \frac{1}{\lambda+1} \left[\hat{\delta} Y_{n-1}(m e^{-1}) m (N e^{-1} + M m^{-1}) \right] v. \end{aligned} \quad (3.16)$$

Now, we need to address the residual spatial rotations \hat{R} of m (or e) because, together with $\partial \mathcal{V} / \partial p$, we have to vary \mathcal{V} with respect to \hat{R} . This can be done by parametrizing \hat{R} similarly to $\hat{\Lambda}$, then varying with respect to the parameters of \hat{R} . For instance, consider the 2×2 dimensional Euclidean metrics,

$$\tilde{g} = \begin{pmatrix} \tilde{N}^2 + \tilde{e}^2 \tilde{\nu}^2 & \tilde{e}^2 \tilde{\nu} \\ \tilde{e}^2 \tilde{\nu} & \tilde{e}^2 \end{pmatrix}, \quad \tilde{E} = \begin{pmatrix} \tilde{N} & 0 \\ \tilde{e} \tilde{\nu} & \tilde{e} \end{pmatrix}, \quad (3.17a)$$

$$\tilde{f} = \begin{pmatrix} \tilde{M}^2 + \tilde{m}^2 \tilde{\mu}^2 & \tilde{m}^2 \tilde{\mu} \\ \tilde{m}^2 \tilde{\mu} & \tilde{m}^2 \end{pmatrix}, \quad \tilde{L}_0 = \begin{pmatrix} \tilde{M} & 0 \\ \tilde{m} \tilde{\mu} & \tilde{m} \end{pmatrix}. \quad (3.17b)$$

Let \tilde{R} be the rotation of the triangular zweibein \tilde{L}_0 such that $\tilde{L} = \tilde{R} \tilde{L}_0$, where from (1.9),

$$\tilde{R}(\varrho) = \begin{pmatrix} \sqrt{1 - \varrho^2} & -\varrho \\ \varrho & \sqrt{1 - \varrho^2} \end{pmatrix}. \quad (3.18)$$

The variation of $e_1(\tilde{L}\tilde{E}^{-1})$ with respect to ϱ gives the equation of motion,

$$\tilde{\nu} - \tilde{\mu} = (\tilde{M}\tilde{m}^{-1} + N\tilde{e}^{-1})r, \quad r = \sqrt{1 - \varrho^2/\varrho} \in \mathbb{R}. \quad (3.19)$$

This is a ‘‘shift difference’’ condition which is similar to (3.23a) below. The solution to (3.19) symmetrizes the zweibeins from (3.17), which gives the square root $(\tilde{g}^{-1}\tilde{f})^{1/2} = \tilde{R}\tilde{L}_0\tilde{E}^{-1}$. A similar derivation, this time for the full rotations, yields the shift difference conditions for the symmetrization of the spatial part $e^\top\hat{\Delta}m$.

The symmetric $e^\top\hat{\Delta}m$ implies the symmetric $\hat{\Lambda}^\top\hat{\delta}Y_n(me^{-1})$, so (3.16) becomes,

$$N\partial e_n(S)/\partial v = \hat{\delta}Y_{n-1}(me^{-1})m \left[(Ne^{-1} + Mm^{-1})v + \mu - \nu \right]. \quad (3.20)$$

Hence, the equations of motion with respect to both p and \hat{R} are solved by,

$$\Delta n := (Ne^{-1} + Mm^{-1})v + \mu - \nu = 0, \quad (3.21)$$

together with $\hat{\delta}\hat{\Lambda}me^{-1} = (\hat{\delta}\hat{\Lambda}me^{-1})^\top$. These two conditions are equivalent to the symmetrization condition,

$$E^\top\eta L = (E^\top\eta L)^\top, \quad (3.22)$$

which gives the real square root $S = (g^{-1}f)^{1/2}$. Namely, a congruence S is self-adjoint $gS = (gS)^\top$ (it is a square root), if and only if [29, 30],

$$\nu - \mu = (Ne^{-1} + Mm^{-1})v, \quad (3.23a)$$

$$e^\top\hat{\delta}\hat{\Lambda}m = (e^\top\hat{\delta}\hat{\Lambda}m)^\top. \quad (3.23b)$$

In terms of the variables D and Q from [3, 4], the spatial symmetrization (3.23b) reads $\hat{f}D = (\hat{f}D)^\top$ where $DQD = \hat{g}^{-1}\hat{f}$, $DQ = \hat{B}$, and $Q = m^{-1}\hat{\Lambda}^2m$.

Finally, the coupled system (3.23) is equivalent to the condition on metrics to have intersecting null cones with a common timelike direction and a common spacelike hypersurface element [16]. The first equation controls the separation between the null cones, and the second ensures that the spatial shapes of the null cones properly intersect.

4 Discussion

The analysis in [3, 4] starts from the square root in the bimetric potential, which gives the required redefinition of the shift variable that is essential for the ghost-free proof in the $N+1$ formalism. We have shown that this shift redefinition and the spatial symmetrization come out from the equation of motion for a general congruence between the metrics. The primary square root naturally emerges as the on-shell condition in the covariant form, which further clarifies the square root branch and type selection in [16].

In earlier versions of [12], the authors proposed a method for dealing with the local spatial rotation invariance using constrained spatial vielbeins. This method is omitted in the most recent version of the paper. The authors justify the removal in a footnote, pointing out that it is not clear that such a method works since solving the constraint may

introduce dependence on the lapse or shift into the vielbeins. As shown in sec. 3, variation with respect to both p and the spatial rotation resolves this issue. In fact, it produces the square root $S = (g^{-1}f)^{1/2}$, which is compatible with the result from appendix C in [12].

Multiple spin-2 fields interactions in [12] and [31] rely on the vielbein formulation. In this work, we only treat bimetric interactions. An extension to multimetric congruences is possible using the generalized symmetric polynomials introduced in [12]. A similar remark holds for interactions of multiple spin-2 fields beyond pairwise couplings [31]. For a singled out metric g_I , the interaction given in [31] can be formulated as $\sqrt{-g_I} \det(\sum_{J=1}^N \beta^J S_{IJ})$ in terms of the congruences $g_J = S_{IJ}^\top g_I S_{IJ}$ (note the transitivity $S_{IJ} = S_{IK} S_{KJ}$).

Even though the vielbein formulation seems more fundamental, when it comes to partial differential equations and the causal propagation of the matter fields, one still needs the metric inverses and contracted covariant derivatives to write down the wave equation. This also holds for the fermionic fields. In general, a well-posed system of partial differential equations for arbitrary tensorial spacetimes carrying predictive, interpretable, and quantizable matter requires a bi-hyperbolic principal symbol [32–34]. For bimetric theory, the principal symbol is the totally symmetrized product $g^{(\mu\nu} f^{\rho\sigma)}$, which is bi-hyperbolic if and only if the real square root of $g^{\mu\rho} f_{\rho\nu}$ exists. A similar property still lacks for the multivielbein/multimetric formulation of spin-2 interactions.

Acknowledgments

I am grateful to Fawad Hassan for the valuable discussions and suggestion how to shorten the proof in section 2. I also thank Edvard Mörtsell, Fawad Hassan, Francesco Torsello, and Marcus Höggås for the comments and careful reading of the draft.

Appendix A Recovering indices

Below is the list of geometrical objects with their indices attached,

$$\begin{array}{cccccc}
S^\mu{}_\nu, & (S^\top)_{\nu}{}^\mu, & (S^{-1})^\nu{}_\mu, & (S^{-1,\top})_\mu{}^\nu, & \omega_{\mu\nu}, & X^\mu{}_\nu, \\
g_{\mu\nu}, & \hat{g}_{ij}, & \nu^i, & f_{\mu\nu}, & \hat{f}_{ij}, & \mu^i, \\
E^A{}_\mu, & (E^\top)_{\mu}{}^A, & (E^{-1})^\mu{}_A, & L^A{}_\mu, & (L^\top)_{\mu}{}^A, & (L^{-1})^\mu{}_A, \\
e^a{}_i, & (e^\top)_{i}{}^a, & (e^{-1})^i{}_a, & m^a{}_i, & (m^\top)_{i}{}^a, & (m^{-1})^i{}_a, \\
B^\mu{}_\nu, & \hat{B}^i{}_j, & & \hat{L}^a{}_i, & q^i, & u_i, \\
\eta_{AB}, & \hat{\delta}_{ab}, & p^a, & (p^\top)^a, & v^a, & (v^\top)^a, \\
(\eta^{-1})^{AB}, & (\hat{\delta}^{-1})^{ab}, & \hat{I}^a{}_b, & \hat{\Lambda}^a{}_b. & &
\end{array}$$

For example: $u^\top q \leftrightarrow u_i q^i$, $uu^\top \leftrightarrow u_i u_j$, $\hat{B} = e^{-1} \hat{L} \leftrightarrow \hat{B}^i{}_j = (e^{-1})^i{}_a \hat{L}^a{}_j$, and,

$$(Ne^{-1} + Mm^{-1})v + \mu - \nu \leftrightarrow \left[N(e^{-1})^i{}_a + M(m^{-1})^i{}_a \right] v^a + \mu^i - \nu^i. \quad (\text{A.1})$$

The adjoint of any A with respect to g reads $(A')^\mu{}_\nu = (g^{-1})^{\mu\rho} (A^\top)_\rho{}^\sigma g_{\sigma\nu}$. The functions Tr and e_n contract the first with the last index; hence, their argument must be a (1,1)-tensor.

Appendix B Detailed derivations

Detailed derivations for section 2

We start from (2.1) where S is not uniquely determined by g, f : we have additional degrees of freedom in an orthogonal transformation X of g , where $g = X^\top g X$, such that,

$$f = S^\top g S = S_0^\top X^\top g X S_0, \quad S = X S_0, \quad (\text{B.1})$$

where S_0 is a function only of g, f which does not depend on X .

We parametrize X using the Cayley transformation where,

$$X = (g + \omega)^{-1}(g - \omega), \quad X(\omega)^{-1} = X(-\omega), \quad X^\top = (g + \omega)(g - \omega)^{-1}, \quad (\text{B.2})$$

together with,

$$2(g + \omega)^{-1}g = I + X, \quad 2(g + \omega)^{-1}\omega = I - X. \quad (\text{B.3})$$

The variation of X in terms of ω is,

$$(g + \omega)X = g - \omega, \quad (\text{B.4})$$

$$(\delta\omega)X + (g + \omega)\delta X = -\delta\omega, \quad (\text{B.5})$$

$$\delta X = -(g + \omega)^{-1}\delta\omega(I + X) = -2(g + \omega)^{-1}\delta\omega(g + \omega)^{-1}g. \quad (\text{B.6})$$

On the other hand, the variation of potential in terms of X is,

$$\delta V(S) = \delta \left(\sum_{n=0}^d \beta_n e_n(S) \right) = \delta \left(\sum_{n=0}^d \frac{\beta_n}{n} \sum_{k=1}^n (-1)^k e_{n-k}(S) \text{Tr}(S^k) \right) \quad (\text{B.7})$$

$$= \sum_{n=1}^d \beta_n \sum_{k=1}^n (-1)^k e_{n-k}(S) \text{Tr}(S^{k-1} \delta S) \quad (\text{B.8})$$

$$= \sum_{n=1}^d \beta_n \sum_{k=1}^n (-1)^k e_{n-k}(S) \text{Tr}(X^{-1} S^k \delta X), \quad (\text{B.9})$$

where in the last two steps we used $\delta S = \delta(X S_0) = (\delta X) S_0 = \delta X X^{-1} X S_0 = \delta X X^{-1} S$ together with the cyclic property of the trace.

After substituting $X^\top =$ and δX into $\delta V(S)$, we get,

$$\delta V(S) = -2 \sum_{n=1}^d \beta_n \sum_{k=1}^n (-1)^k e_{n-k}(S) \text{Tr}(S^k (g + \omega)^{-1} \delta\omega (g + \omega)^{-1} g X^{-1}) \quad (\text{B.10})$$

$$= -2 \sum_{n=1}^d \beta_n \sum_{k=1}^n (-1)^k e_{n-k}(S) \text{Tr}(S^k (g + \omega)^{-1} \delta\omega (g + \omega)^{-1} X^\top g) \quad (\text{B.11})$$

$$= -2 \sum_{n=1}^d \beta_n \sum_{k=1}^n (-1)^k e_{n-k}(S) \text{Tr}(S^k (g + \omega)^{-1} \delta\omega (g - \omega)^{-1} g) \quad (\text{B.12})$$

$$= -2 \sum_{n=1}^d \beta_n \sum_{k=1}^n (-1)^k e_{n-k}(S) \text{Tr}((g - \omega)^{-1} g S^k (g + \omega)^{-1} \delta\omega). \quad (\text{B.13})$$

Since p is skew-symmetric, we have $\delta\omega^\top = -\delta\omega$ and,

$$\delta \operatorname{Tr}(C\omega) = \operatorname{Tr}(C \delta\omega) = \operatorname{Tr}(\delta\omega^\top C^\top) = -\operatorname{Tr}(C^\top \delta\omega), \quad (\text{B.14})$$

that is [35],

$$\delta \operatorname{Tr}(C\omega) = \frac{1}{2} \operatorname{Tr}[(C - C^\top)\delta\omega], \quad \rightarrow \quad \partial \operatorname{Tr}(C\omega)/\partial\omega = \frac{1}{2}(C - C^\top). \quad (\text{B.15})$$

Hence,

$$\begin{aligned} \partial V(S)/\partial\omega &= -\sum_{n=1}^d \beta_n \sum_{k=1}^n (-1)^k e_{n-k}(S) \times \\ &\times \left[\left((g - \omega)^{-1} g S^k (g + \omega)^{-1} - \left((g - \omega)^{-1} g S^k (g + \omega)^{-1} \right)^\top \right) \right] \end{aligned} \quad (\text{B.16})$$

$$= -\sum_{n=1}^d \beta_n \sum_{k=1}^n (-1)^k e_{n-k}(S) \left[(g - \omega)^{-1} (g S^k - (g S^k)^\top) (g + \omega)^{-1} \right]. \quad (\text{B.17})$$

The equations of motion for ω are $\partial V/\partial\omega = 0$, i.e., having a nonsingular g (and $g \pm \omega$),

$$\sum_{n=1}^d \beta_n \sum_{k=1}^n (-1)^k e_{n-k}(S) \left[S^k - (g^{-1} S^\top g)^k \right] = 0. \quad (\text{B.18})$$

Detailed derivations for section 3, part 1

Let us introduce the relation $\stackrel{e_n}{=}$ which indicates that the two expressions are equal after e_n is applied to both sides of the equation. Using the cyclic property of e_n , we obtain,

$$S \stackrel{e_n}{=} \frac{1}{N} \begin{pmatrix} u^0 + u^\top(q - \nu) \\ e^{-1} \hat{L}(q - \nu) \end{pmatrix} \begin{pmatrix} 1 \\ 0 \end{pmatrix}^\top + \begin{pmatrix} 0 & u^\top \\ 0 & e^{-1} \hat{L} \end{pmatrix} = \frac{1}{N} U W^\top + B, \quad (\text{B.19})$$

where,

$$U := \begin{pmatrix} u^0 \\ 0 \end{pmatrix} + B \begin{pmatrix} 0 \\ q - \nu \end{pmatrix}, \quad W := \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \quad B := \begin{pmatrix} 0 & u^\top \\ 0 & \hat{B} \end{pmatrix}, \quad \hat{B} := e^{-1} \hat{L}. \quad (\text{B.20})$$

Then,

$$S \stackrel{e_n}{=} \left[\frac{1}{N} \begin{pmatrix} 1 \\ -\nu \end{pmatrix} \begin{pmatrix} 1 \\ 0 \end{pmatrix}^\top + \begin{pmatrix} 0 & 0 \\ 0 & e^{-1} \end{pmatrix} \right] \begin{pmatrix} u^0 + u^\top q & u^\top \\ \hat{L} q & \hat{L} \end{pmatrix} \quad (\text{B.21})$$

$$\stackrel{e_n}{=} \begin{pmatrix} 0 & 0 \\ 0 & e^{-1} \end{pmatrix} \begin{pmatrix} u^0 + u^\top q & u^\top \\ \hat{L} q & \hat{L} \end{pmatrix} \left[\frac{1}{N} \begin{pmatrix} 1 \\ -\nu \end{pmatrix} \begin{pmatrix} 1 \\ 0 \end{pmatrix}^\top + \begin{pmatrix} 0 & 0 \\ 0 & \hat{I} \end{pmatrix} \right] \quad (\text{B.22})$$

$$= \begin{pmatrix} u^0 + u^\top q & u^\top \\ e^{-1} \hat{L} q & e^{-1} \hat{L} \end{pmatrix} \left[\frac{1}{N} \begin{pmatrix} 1 \\ -\nu \end{pmatrix} \begin{pmatrix} 1 \\ 0 \end{pmatrix}^\top + \begin{pmatrix} 0 & 0 \\ 0 & \hat{I} \end{pmatrix} \right]. \quad (\text{B.23})$$

Using the following identity (which holds for arbitrary B and vectors u, w),

$$e_n(B + u w^\top) = e_n(B) + \sum_{k=1}^n (-1)^{k-1} e_{n-k}(B) w^\top B^{k-1} u, \quad (\text{B.24})$$

we immediately have,

$$e_n(S) = e_n(B) + \frac{1}{N} \sum_{k=1}^n (-1)^{k-1} e_{n-k}(B) W^\top B^{k-1} U. \quad (\text{B.25})$$

Note that $e_n(B) = e_n(\hat{B})$ and $W^\top U = u^0$; thus,

$$N e_n(S) = N e_n(\hat{B}) + u^0 e_{n-1}(\hat{B}) + \sum_{k=1}^n (-1)^{k-1} e_{n-k}(\hat{B}) \begin{pmatrix} 1 \\ 0 \end{pmatrix}^\top B^k \begin{pmatrix} 0 \\ q - \nu \end{pmatrix}. \quad (\text{B.26})$$

Then from,

$$B^k = \begin{pmatrix} 0 & u^\top \hat{B}^{k-1} \\ 0 & \hat{B}^k \end{pmatrix}, \quad (\text{B.27})$$

it follows,

$$N e_n(S) = N e_n(\hat{B}) + u^0 e_{n-1}(\hat{B}) + \sum_{k=1}^n (-1)^{k-1} e_{n-k}(\hat{B}) u^\top \hat{B}^{k-1} (q - \nu). \quad (\text{B.28})$$

where $e_n(\hat{B}) = 0$ for $n > d - 1$.

Assuming that we are in a frame where both f and g admit proper decomposition (both E and L can be triangularized), we get,

$$\begin{aligned} N e_n(S) &= N e_n(\hat{B}) + \lambda^{-1} M e_{n-1}(\hat{B}) + \\ &+ \sum_{k=1}^n (-1)^{k-1} e_{n-k}(\hat{B}) u^\top \hat{B}^{k-1} (M m^{-1} v + \mu - \nu). \end{aligned} \quad (\text{B.29})$$

Note that $\hat{B} = e^{-1} \hat{L}$ corresponds to the congruence,

$$\hat{B}^\top \hat{g} \hat{B} = \hat{L}^\top \hat{\delta} \hat{L} = \hat{f} + uu^\top = \hat{f}(\hat{I} + \hat{f}^{-1}uu^\top) = \hat{f}Q = Q^\top \hat{f}, \quad Q := m^{-1} \hat{\Lambda}^2 m. \quad (\text{B.30})$$

Note on derivations for section 3, part 2

For the calculation of (3.13), see the derivation tracks 6 and 7 in the ancillary Mathematica notebook. The variation $\partial \mathcal{V} / \partial p$ in (3.16) is done using the following identities (for more details see Part 3 in the Mathematica notebook),

$$\partial(p^\top A) / \partial p = A^\top, \quad (\text{B.31})$$

and,

$$\partial\left(A \frac{p}{\lambda + 1}\right) / \partial p = \frac{1}{\lambda + 1} A \hat{\Lambda}^{-1}, \quad (\text{B.32})$$

where A is an arbitrary matrix that does not depend on p and,

$$\hat{\Lambda}^{-1} = \hat{I} - \frac{1}{\lambda(\lambda + 1)} pp^\top \hat{\delta}. \quad (\text{B.33})$$

Appendix C Bimetric actions with a scalar potential

Here we highlight GR-type actions where the interaction between metrics is given through a scalar potential. Consider the metric fields g and f where the dynamics of each metric is governed by the Einstein–Hilbert term. Each diffeomorphism group $\text{Diff}(g)$ and $\text{Diff}(f)$ acts separately on its own metric. In the interacting case where the interaction is given through a scalar potential $\mathcal{V}(g, f)$, the symmetry of the full action $(\phi_1, \phi_2) \in \text{Diff}(g) \times \text{Diff}(f)$ must be reduced to the diagonal group of common diffeomorphisms where $\phi_1 = \phi_2$ by the theorem in [36, 37]. This demands that the interaction term depends only on the scalars one can make with two metrics [38]. More precisely, the common diffeomorphism invariance restricts the scalar potential to depend only on the invariants of the (1,1) tensor field $g^{-1}\tilde{f}$ where φ is an overall diffeomorphism $\phi_1 = \varphi \circ \phi_2$ and $\tilde{f} = \varphi^* f$ is the pullback of f by φ ,

$$\tilde{f}_{\mu\nu} = \partial_\mu \varphi^{\bar{\alpha}} \partial_\nu \varphi^{\bar{\beta}} f_{\bar{\alpha}\bar{\beta}}. \quad (\text{C.1})$$

The map φ is part of the local trivialization of the tangent bundles. It gives rise to Stückelberg fields $\varphi^{\bar{\alpha}}$ that do not introduce new dynamics into the theory; hence, φ can be fixed to be the identity map in the unitary gauge where $\partial_\mu \varphi^{\bar{\alpha}} = \delta_\mu^{\bar{\alpha}}$, sloppily setting $\tilde{f} = f$.

The analysis in section 2 would not change if we had $\tilde{f} = \tilde{S}^\top g \tilde{S}$ and $\varphi^* f = S^\top g S$,

$$g^{-1} S^\top g = g^{-1} S^\top g S S^{-1} = g^{-1} \varphi^* \tilde{f} S^{-1} = g^{-1} f S^{-1} = A S^{-1}, \quad (\text{C.2})$$

since $\varphi^* \tilde{f} = \Phi^\top \tilde{S}^\top g \tilde{S} \Phi$ with $S = \tilde{S} \Phi$, that is, $S^\mu{}_\nu = \tilde{S}^\mu{}_{\bar{\alpha}} \partial_\nu \varphi^{\bar{\alpha}}$ where $\Phi^{\bar{\alpha}}{}_\mu = \partial_\mu \varphi^{\bar{\alpha}}$, and,

$$A = g^{-1} f = g^{-1} \Phi^\top \tilde{f} \Phi. \quad (\text{C.3})$$

Stückelberg trick. A congruence in the most general form reads,

$$f_{\bar{\alpha}\bar{\beta}} = S^\mu{}_{\bar{\alpha}} S^\nu{}_{\bar{\beta}} g_{\mu\nu}. \quad (\text{C.4})$$

where $S^\mu{}_{\bar{\alpha}} = \partial_{\bar{\alpha}} \varphi^\gamma X^\mu{}_\rho S_0^\rho{}_\gamma$ and φ is a diffeomorphism such that $\partial_{\bar{\alpha}} \varphi^\gamma$ is the pullback (differential map) that is “moving” f to the same tangent bundle where g lives ($\tilde{f} = \varphi^* f$). To be able to write down $e_n(S)$, we must use $S^\mu{}_\nu = S^\mu{}_{\bar{\alpha}} \partial_\nu \varphi^{\bar{\alpha}}$. The redundant gauge degrees of freedom in φ do not introduce new dynamics into the theory. The Einstein–Hilbert term is “blind” to both $\partial_{\bar{\alpha}} \varphi^\gamma$ and $X^\mu{}_\rho$. Also,

$$\frac{\delta}{\delta \varphi^{\bar{\alpha}}} \mathcal{V}(g, f) = 0. \quad (\text{C.5})$$

This equation will not give rise to any new dynamics since it is implied by the Bianchi constraint (for instance, see [9]),

$$\nabla^\mu V_{\mu\nu}(g, f) = 0, \quad \text{where } V_{\mu\nu}(g, f) := \frac{-2}{\sqrt{-g}} \frac{\delta}{\delta g^{\mu\nu}} [\sqrt{-g} \mathcal{V}(g, f)]. \quad (\text{C.6})$$

Indeed, a gauge transformation under the diagonal group of diffeomorphisms gives,

$$\nabla^\mu V_{\mu\nu}(g, f) = \frac{\delta \mathcal{V}}{\delta \varphi^{\bar{\alpha}}} \partial_\nu \varphi^{\bar{\alpha}}, \quad (\text{C.7})$$

where $\partial_\nu \varphi^{\bar{\alpha}}$ is nonsingular.

References

- [1] C. de Rham, G. Gabadadze and A. J. Tolley, *Resummation of Massive Gravity*, *Phys. Rev. Lett.* **106** (2011) 231101, [1011.1232].
- [2] S. F. Hassan and R. A. Rosen, *Resolving the Ghost Problem in non-Linear Massive Gravity*, *Phys. Rev. Lett.* **108** (2012) 041101, [1106.3344].
- [3] S. F. Hassan, R. A. Rosen and A. Schmidt-May, *Ghost-free Massive Gravity with a General Reference Metric*, *JHEP* **02** (2012) 026, [1109.3230].
- [4] S. F. Hassan and R. A. Rosen, *Bimetric Gravity from Ghost-free Massive Gravity*, *JHEP* **02** (2012) 126, [1109.3515].
- [5] S. F. Hassan and R. A. Rosen, *Confirmation of the Secondary Constraint and Absence of Ghost in Massive Gravity and Bimetric Gravity*, *JHEP* **04** (2012) 123, [1111.2070].
- [6] S. F. Hassan and A. Lundkvist, *Analysis of constraints and their algebra in bimetric theory*, *JHEP* **08** (2018) 182, [1802.07267].
- [7] C. de Rham, *Massive Gravity*, *Living Rev. Rel.* **17** (2014) 7, [1401.4173].
- [8] K. Hinterbichler, *Theoretical Aspects of Massive Gravity*, *Rev. Mod. Phys.* **84** (2012) 671–710, [1105.3735].
- [9] A. Schmidt-May and M. von Strauss, *Recent developments in bimetric theory*, *J. Phys.* **A49** (2016) 183001, [1512.00021].
- [10] D. G. Boulware and S. Deser, *Can gravitation have a finite range?*, *Phys. Rev.* **D6** (1972) 3368–3382.
- [11] I. G. Macdonald, *Symmetric functions and orthogonal polynomials*. AMS, 1998.
- [12] K. Hinterbichler and R. A. Rosen, *Interacting Spin-2 Fields*, *JHEP* **07** (2012) 047, [1203.5783].
- [13] B. Zumino, *Effective Lagrangians and Broken Symmetries*, in *Lectures on Elementary Particles and Quantum Field Theory vol.2*, Cambridge, Mass.: Brandeis Univ., pp. 437–500, 1970.
- [14] C. Deffayet, J. Mourad and G. Zahariade, *A note on ‘symmetric’ vielbeins in bimetric, massive, perturbative and non perturbative gravities*, *JHEP* **03** (2013) 086, [1208.4493].
- [15] S. F. Hassan, A. Schmidt-May and M. von Strauss, *Metric Formulation of Ghost-Free Multivielbein Theory*, 1204.5202.
- [16] S. F. Hassan and M. Kocic, *On the local structure of spacetime in ghost-free bimetric theory and massive gravity*, *JHEP* **05** (2018) 099, [1706.07806].
- [17] W. Scharlau, *Quadratic and hermitian forms*. Grundlehren der mathematischen Wissenschaften. Springer, 1985.
- [18] S. Lang, *Algebra*. Graduate Texts in Mathematics. Springer New York, 2002.
- [19] E. Meinrenken, *Clifford algebras and Lie theory*. Springer, 2013, 10.1007/978-3-642-36216-3.
- [20] S. Hawking and G. Ellis, *The Large Scale Structure of Space-Time*. Cambridge Monographs on Mathematical Physics. Cambridge University Press, 1973.
- [21] G. H. Golub and C. F. Van Loan, *Matrix Computations (3rd Ed.)*. Johns Hopkins University Press, Baltimore, MD, USA, 1996.

- [22] B. Obsieger, *Numerical Methods II - Roots and Equation Systems*. Numerical Methods. Boris Obsieger, 2015.
- [23] R. Horn and C. Johnson, *Topics in Matrix Analysis*. Cambridge University Press, 1994.
- [24] N. Higham, *Functions of Matrices: Theory and Computation*. SIAM, 2008.
- [25] M. Konstantinov, D. Gu, V. Mehrmann and P. Petkov, *Perturbation Theory for Matrix Equations*. Studies in Computational Mathematics. Elsevier Science, 2003.
- [26] É.ourgoulhon, *3+1 Formalism in General Relativity: Bases of Numerical Relativity*. Lecture Notes in Physics. Springer Berlin Heidelberg, 2012.
- [27] R. L. Arnowitt, S. Deser and C. W. Misner, *The dynamics of general relativity*, *Gen. Rel. Grav.* **40** (2008) 1997–2027, [gr-qc/0405109].
- [28] J. W. York, Jr., *Kinematics and Dynamics of General Relativity*, pp. 83–126.
- [29] M. Kocic, *Geometric mean of bimetric spacetimes*, 1803.09752.
- [30] S. F. Hassan, M. Kocic and A. Schmidt-May, *Absence of ghost in a new bimetric-matter coupling*, 1409.1909.
- [31] S. F. Hassan and A. Schmidt-May, *Interactions of multiple spin-2 fields beyond pairwise couplings*, 1804.09723.
- [32] D. Raetzl, S. Rivera and F. P. Schuller, *Geometry of physical dispersion relations*, *Phys. Rev.* **D83** (2011) 044047, [1010.1369].
- [33] F. P. Schuller and C. Witte, *How quantizable matter gravitates: A practitioner’s guide*, *Phys. Rev.* **D89** (2014) 104061, [1402.6548].
- [34] F. P. Schuller, N. Stritzelberger, F. Wolz and M. Düll, *Gravitational closure of matter field equations*, 1611.08878.
- [35] K. B. Petersen and M. S. Pedersen, *The matrix cookbook*, tech. rep., nov, 2012.
- [36] N. Boulanger, T. Damour, L. Gualtieri and M. Henneaux, *No consistent cross interactions for a collection of massless spin-2 fields*, in *Quantization, gauge theory, and strings. Proceedings, International Conference dedicated to the memory of Professor Efim Fradkin, Moscow, Russia, June 5-10, 2000. Vol. 1+2*, pp. 150–161, 2000, hep-th/0009109.
- [37] N. Boulanger, T. Damour, L. Gualtieri and M. Henneaux, *Inconsistency of interacting, multigraviton theories*, *Nucl. Phys.* **B597** (2001) 127–171, [hep-th/0007220].
- [38] T. Damour and I. I. Kogan, *Effective Lagrangians and universality classes of nonlinear bigravity*, *Phys. Rev.* **D66** (2002) 104024, [hep-th/0206042].