

DELAYED SPIN-UP AND PERSISTENT SHIFT PHENOMENA OF CRAB PULSAR GLITCHES: TWO SIDES OF THE SAME COIN?

WEI-HUA WANG¹ AND XIAO-PING ZHENG^{1,2}*Draft version January 20, 2022*

ABSTRACT

Pulsar glitches are sudden increase in their spin frequency, in most cases followed by the long timescale recovery process. As of this writing, about 546 glitches have been reported in 188 pulsars, the Crab pulsar is a special one with unique manifestations. This writing presents a statistic study on post-glitch observables of the Crab pulsar, especially the delayed spin-up in post-glitch phase and persistent shift in the slow-down rate of the star. By analyzing the radio data over 45 years, we find that two power law functions respectively fit the persistent shift and delayed spin-up timescales versus glitch size well, and we find a linear correlation between the persistent shift and delayed spin-up timescale from the consistency of the two fitting functions, probably indicating their same physical origin and may provide a new probe of interior physics of neutron stars.

Subject headings: stars: neutron — (stars:) pulsars: Crab: glitch — stars: statistics

1. INTRODUCTION

Glitch is a phenomenon that interrupts the monotonous spin down of pulsars due to electromagnetic braking (Taylor et al. 1993), it is characterized by a sudden increase in spin frequency, generally accompanied by a increase in spin down rate. The first glitch was discovered in the Vela pulsar in 1969 (Radhakrishnan & Manchester 1969; Reichely & Downs 1969), at present, about 546 glitches have been reported in 188 pulsars³, the famous Crab and Vela pulsars are both frequently glitch sources and are daily monitored. A series of models have been proposed ever since its first discovery, such as crustquake (Ruderman 1969), corequake (Pines et al. 1972), planetary perturbation (Michel 1970) and magnetospheric instabilities (Scargle & Pacini 1971), but none of these were convincing enough (Pines et al. 1974). In 1969, Baym et al. proposed the long timescale in the post-glitch recovery process of the Vela pulsar as a signature of neutron superfluid in inner neutron star (Baym et al. 1969). It should be noted that the absence of radiative and pulse profile changes in Vela glitches seems to support its internal origin. In 1975, Anderson and Itoh advanced the semina idea that glitches are triggered by sudden unpinning of superfluid vortices from neutron star crust (Anderson & Itoh 1975), resulting in a rapid transfer of angular momentum from the faster rotating superfluid component to the normal component, besides, as a small portion of moment of inertia decouples from the normal component while the external torque acting on the pulsar remains constant in short timescale, the observed spin down rate will thus increase temporarily. Alpar et al. further developed this into the vortex creep theory (Alpar et al. 1984), which

is now widely accepted as the standard scenario due to its success in explaining the post-glitch recovery process.

Within the framework of vortex creep theory, the pinning force and the friction between the crust and the superfluid component dominate the post-glitch relaxation process, thus in the relaxation timescale, the spin down rate will gradually go back to the value predicted by fitting to pre-glitch data. However, this is not the case for the young Crab pulsar. Despite its very low glitch activity (Fuentes et al. 2017), the Crab pulsar can not go back to predicted spin down rate even till three years later after glitches (Lyne et al. 2015), most evident in large Crab glitch recovery processes, this phenomenon is called the persistent shift. The persistent shift is accumulative if the time interval is less than three years and their effects can not be resolved. Besides, several large glitches in the Crab pulsar have experienced slow increase in spin frequency with timescales of days following the rapid rise, which mean day-long timescale positive or at least effective positive torques, this phenomenon is called delayed spin-up. Delayed spin-up was first discovered in the comparatively large glitch in 1989 (Lyne et al. 1992), and in two further glitches in 1996 (Wong et al. 2001) and 2017 (Shaw et al. 2018), Table(2) gives parameters of these three glitches. Remarkably, large Crab glitches are accompanied by both delayed spin-ups and persistent shifts, besides, larger glitch size corresponds to longer delayed spin-up timescale and larger persistent shift, from this point of view, it seems that delayed spin-up and persistent shift are tightly correlated. Other young neutron star also experience persistent shift, for instance, PSR B2334+61 (characteristic age $\tau_c \sim 41$ kyr) experienced a very large glitch (glitch size $\Delta\nu/\nu \sim 20.5 \times 10^{-6}$, much larger than Crab glitches) between MJDs 53608 and 53621, this glitch resulted in a large long-term persistent shift amounts to $\sim 1.1\%$ of the spin down rate at the time of the glitch (Yuan et al. 2010), but no delayed spin-up is reported, probably indicating a different physical origin.

The anomalous post-glitch behaviors of the Crab pulsar pose challenges to the standard vortex creep theory.

¹ Institute of Astrophysics, Central China Normal University, Wuhan 430079, China, wangweihua@mails.ccnuc.edu.cn, zhxp@phy.ccnuc.edu.cn

² Key Laboratory of Quark and Lepton Physics (Central China Normal University), Ministry of Education, Wuhan 430079, China

³ <http://www.jb.man.ac.uk/pulsar/glitches/gTable.html>, and <http://www.atnf.csiro.au/people/pulsar/psrcat/glitchTbl.html>.

Alpar et al. had explained this by combining the vortex creep and starquake (Alpar et al. 1994; Alpar et al. 1996). They proposed that starquake would result in vortex depletion region in the crust, when glitch is triggered and large amount of superfluid vortices move outward, part of the flowing vortices would transport inward and be trapped by vortex depletion region, resulting in the delayed spin-up. Besides, they interpreted the persistent shift as a decrease in effective moment of inertia through creation of new vortex depletion regions. The differences between the Crab and the Vela pulsar are understood from the view of evolutionary, as no new depletion region can be formed in the Vela pulsar because it is much older than the Crab pulsar. This theory phenomenologically explain the observations, but it depends strongly on the assumed notion of vortex depletion region that can not be verified. Besides, within this model, the delayed spin-up and persistent shift result from different physical origins, thus it is hard to build up any direct correlations between observables in these two phenomena. Haskell et al. emphasized the effect of vortex accumulation and proposed that vortex accumulation at certain part of the neutron star may account for the delayed spin-up that is seen as a extension of the fast spin-up (Haskell et al. 2018), but they provided no explanation for the physical origin of persistent shift phenomenon.

This letter aims at the data analysis to infer the possible correlations between observables of delayed spin-up and persistent shift phenomena from the view of statistics. We present the detailed statistics and analysis in Section 2, and the summary and discussion in Section 3.

2. STATISTICS AND ANALYSIS

All measured values of Crab pulsars glitches are presented in Table(1), data are taken from references Espinoza et al. 2011 and 2014, Lyne et al. 1992, Wong et al. 2001, Shaw et al. 2018 and from website <http://www.jb.man.ac.uk/pulsar/glitches/gTable.html>. The first and second columns correspond to time of the glitches, the third column is the fractional increase in spin frequency ($\Delta\nu/\nu$), the fourth column is the step increase in spin frequency ($\Delta\nu$), namely, glitch size, the fifth column is the fractional increase in spin down rate ($\Delta\dot{\nu}/\dot{\nu}$), and the last column is the persistent shift value ($\Delta\dot{\nu}_p$), X means unknown. Isolated glitches are separated from each other by lines in Table(1), but neighboring glitches whose effects are unresolved are not separated. Observables of three large glitches where both delayed spin-up and persistent shift occurred are listed in Table(2), $\Delta\nu_d$ is the frequency increase in the delayed spin-up process and τ_d is the timescale of delayed spin-up. Numbers in brackets represent error bars of the last significant digit.

Firstly, we analyze the relationship between $\Delta\dot{\nu}_p$ and $\Delta\nu$, as shown in Figure 1, 2 and 3. Figure 1 suggests two groups of $|\Delta\dot{\nu}_p|$ at cutoff $\Delta\nu \sim 1 \mu\text{Hz}$, in the following, glitches with $\Delta\nu > 1 \mu\text{Hz}$ are called large glitches, on the contrary, glitches with $\Delta\nu < 1 \mu\text{Hz}$ as small glitches. A linear fitting to five large glitches gives

$$|\Delta\dot{\nu}_p| = (24\Delta\nu + 90) \times 10^{-15} \text{ s}^{-2}, \quad (1)$$

where $\Delta\nu$ is in μHz throughout this writing. While linear




FIG. 1.— Persistent shift as a function of glitch size. Filled circles represent persistent shifts in isolated Crab glitches, blue and red unfilled circles represent cumulative persistent shifts after several neighboring glitches, but blue unfilled circles include a primary glitch which contributes most of the persistent shift, for example, Crab glitch MJD 53067.0780, red unfilled circles include several glitches with similar glitch sizes, for example, Crab glitch MJD 52498.257 and MJD 52587.20. Red and blue thick lines are our linear fitting to small and relatively large glitches separately. For comparison, Lyne's linear fitting is shown as the black thick line. For all unfilled circles, their glitch sizes are simply the sum of glitch size of several neighboring glitches.




FIG. 2.— Same with Figure 1 but persistent shift is fitted as power law functions of glitch size. Red thick and black dotted lines represent the fitting functions $|\Delta\dot{\nu}_p| = (a_1\Delta\nu^{b_1} + c_1) \times 10^{-15} \text{ s}^{-2}$ with $c_1 = 0$ set by hand and $|\Delta\dot{\nu}_p| = (a_2\Delta\nu^{b_2} + c_2) \times 10^{-15} \text{ s}^{-2}$ respectively.

fitting to small glitches gives

$$|\Delta\dot{\nu}_p| = 65 \times 10^{-15} \Delta\nu \text{ s}^{-2}, \quad (2)$$

Lyne et al. have also considered the inter-dependence of $\Delta\dot{\nu}_p$ and $\Delta\nu$ (Lyne et al. 2015), their linear fitting to all glitches gives,

$$|\Delta\dot{\nu}_p| = 70 \times 10^{-15} \Delta\nu \text{ s}^{-2}. \quad (3)$$

This result is very close to our fitting to small Crab glitches. A comparison between Figure 1 in this paper and Figure 5 in Lyne et al. 2015 shows clearly that, the fitting in logarithmic coordinate space seriously underestimated the contribution of large glitches.

Different linear relations between $|\Delta\dot{\nu}_p|$ and $\Delta\nu$ for large and small Crab glitches probably indicates their differences in physical origins, for example, large glitches may have the potential to influence internal structure

TABLE 1
OBSERVABLE OF ALL CRAB GLITCHES

Date	MJD (d)	$\Delta\nu/\nu$ (10^{-9})	$\Delta\nu$ (μHz)	$\Delta\dot{\nu}/\dot{\nu}$ (10^{-3})	$\Delta\dot{\nu}_p$ (10^{-15} s^{-2})
1969 September	40491.84(3)	7.2(4)	0.22(1)	0.44(4)	
1971 July	41161.98(4)	1.9(1)	0.057(4)	0.17(1)	
1971 October	41250.32(1)	2.1(1)	0.062(3)	0.11(1)	
1975 February	42447.26(4)	35.7(3)	1.08(1)	1.6(1)	−112(2)
1986 August	46663.69(3)	6.0	0.18(2)	0.5(1)	
1989 August	47767.504(3)	81.0(4)	2.43(1)	3.4(1)	−150(5)
1992 November	48945.6(1)	4.2(2)	0.13(1)	0.32(3)	
1995 October	50020.04(2)	2.1(1)	0.063(2)	0.20(1)	
1996 June	50260.031(4)	31.9(1)	0.953(4)	1.73(3)	
1997 January	50458.94(3)	6.1(4)	0.18(1)	1.1(1)	−116(5)
1997 December	50812.59(1)	6.2(2)	0.19(1)	0.62(4)	
1999 October	51452.02(1)	6.8(2)	0.20(1)	0.7(1)	−25(3)
2000 July	51740.656(2)	25.1(3)	0.75(1)	2.9(1)	
2000 September	51804.75(2)	3.5(1)	0.105(3)	0.53(3)	−53(3)
2001 June	52084.072(1)	22.6(1)	0.675(3)	2.07(3)	
2001 October	52146.7580(3)	8.87(5)	0.265(1)	0.57(1)	−70(10)
2002 August	52498.257(2)	3.4(1)	0.101(2)	0.70(2)	
2002 September	52587.20(1)	1.7(1)	0.050(3)	0.5(1)	−8(2)
2004 March	53067.0780(2)	214(1)	6.37(2)	6.2(2)	
2004 September	53254.109(2)	4.9(1)	0.145(3)	0.2(1)	
2004 November	53331.17(1)	2.8(2)	0.08(1)	0.7(1)	−250(20)
2006 August	53970.1900(3)	21.8(2)	0.65(1)	3.1(1)	−30(5)
2008 April	54580.38(1)	4.7(1)	0.140(4)	0.2(1)	
2011 November	55875.5(1)	49.2(3)	1.46(1)	X	−132(5)
2017 March	57839.92(6)	2.14(11)	0.064(3)	0.27(3)	
2017 November	58064.555(3)	516.37(10)	15.304(9)	6.969(21)	
2018 April	58237.357(5)	4.08(22)	0.122(6)	0.46(11)	

TABLE 2
OBSERVABLE OF THREE LARGE CRAB GLITCHES

MJD (d)	$\Delta\nu$ (μHz)	$\Delta\nu_d$ (μHz)	τ_d (days)	$\Delta\dot{\nu}_p$ (10^{-15} s^{-2})
47767.504(3)	2.43(1)	0.7	0.8	−150(5)
50260.031(4)	0.953(4)	0.31	0.5	−116(5)
58064.555(3)	15.304(9)	1.1	1.7	X

of neutron stars and result in relatively large persistent shifts, while effects of small glitches is limited and it seems impossible to change the structure, in this case, persistent shifts in small glitches may originate from some other unknown mechanism. However, conclusion that persistent shifts in small and large glitches arise from different physical processes seems to be unconvincing because of the absence of more data. Besides, distribution of points in the $|\Delta\dot{\nu}_p|$ versus $\Delta\nu$ plot also influence our judgement.

If the large and small glitches in the Crab pulsar do

have the same physical origin, it is reasonable to fit the whole range of data with one single function. Despite the linear fitting we discussed above, it is naturally to consider the power law fitting since persistent shifts of larger Crab pulsar glitches tend to be much larger. A fitting function in the form of $|\Delta\dot{\nu}_p| = a_1 \Delta\nu^{b_1} \times 10^{-15} \text{ s}^{-2}$ gives $a_1 = 87.24 \pm 1.288$, $b_1 = 0.7438 \pm 0.0186$, $c_1 = 0$ and $\chi^2/\text{dof} = 304.9/8$, while a fitting function in the form of $|\Delta\dot{\nu}_p| = (a_2 \Delta\nu^{b_2} + c_2) \times 10^{-15} \text{ s}^{-2}$ gives $a_2 = 151.5 \pm 13.1$, $b_2 = 0.4337 \pm 0.0444$, $c_2 = -59.08 \pm 12.33$ and $\chi^2/\text{dof} = 222.8/7$. The coefficient $c_1 = 0$ is obvi-




FIG. 3.— Same with Figure 1 but only persistent shifts in relatively isolated glitches are considered, which means glitches MJD 51804.75, MJD 52146.7580 and MJD 52587.20 are excluded. Black dotted and thick lines represent the fitting functions $|\dot{\Delta\nu}_p| = (a_2\Delta\nu^{b_2} + c_2) \times 10^{-15} \text{ s}^{-2}$ and $|\dot{\Delta\nu}_p| = (a_3\Delta\nu^{b_3} + c_3) \times 10^{-15} \text{ s}^{-2}$ respectively.




FIG. 4.— Delayed spin-up timescale as power law functions of glitch size. Black line represents the fitting function $\tau_d/(\text{days}) = a_4\Delta\nu^{b_4} + c_4$.




FIG. 5.— Application of the best fitting function of persistent shift versus glitch size to the delayed spin-up timescale with constant $K_1 = 201.98 \times 10^{-15} \text{ s}^{-2} \text{ days}^{-1}$.




FIG. 6.— Application of the best fitting function of delayed spin-up timescale versus glitch size to the persistent shift with constant $K_2 = 196.08 \times 10^{-15} \text{ s}^{-2} \text{ days}^{-1}$.

ously set by hand and the latter function gives a better fitting mathematically. Although the power law function seems to reconcile both large and small glitches, accumulative effect of persistent shift may contaminate the fitting. Therefore, we further fit persistent shifts of relatively isolated glitches after exclusion of glitch MJD 51804.75, MJD 52146.7580 and MJD 52587.20, a fitting function in the form of $|\dot{\Delta\nu}_p| = (a_3\Delta\nu^{b_3} + c_3) \times 10^{-15} \text{ s}^{-2}$ gives $a_3 = 243.1 \pm 60.34$, $b_3 = 0.2536 \pm 0.06654$, $c_3 = -141.9 \pm 59.51$ and $\chi^2/\text{dof} = 102.6/4$, this have improved the fitting mathematically. Comparison between $|\dot{\Delta\nu}_p| = (a_2\Delta\nu^{b_2} + c_2) \times 10^{-15} \text{ s}^{-2}$ (black dotted line in Figure 3) and $|\dot{\Delta\nu}_p| = (a_3\Delta\nu^{b_3} + c_3) \times 10^{-15} \text{ s}^{-2}$ (black thick line in Figure 3) shows that, $|\dot{\Delta\nu}_p| = (a_3\Delta\nu^{b_3} + c_3) \times 10^{-15} \text{ s}^{-2}$ fits the data better while $|\dot{\Delta\nu}_p| = (a_2\Delta\nu^{b_2} + c_2) \times 10^{-15} \text{ s}^{-2}$ underestimates contribution of large glitch MJD 53067.0780. Following this procedure, the function $|\dot{\Delta\nu}_p| = (a_3\Delta\nu^{b_3} + c_3) \times 10^{-15} \text{ s}^{-2}$ can be seen as the best fitting result.

We emphasize that, uncertainties of persistent shift in small glitches are relatively larger than that in large glitches, which means that persistent shift in large isolated glitches are more reliable, therefore, the best fitting should be close to points with large persistent shift. We thus removed the data points with serious accumulative effect so as not to affect the fitting. Besides, the best fitting function will inevitably bring in the cut-off glitch-size below which no persistent shift value is measured, physical meaning of this is unclear at present, probably related to some non-ideal effects, for example, nonspherically symmetric neutron star structure.

Secondly, the three large glitches in Table (2) brings another observable, the delayed spin-up timescale τ_d . As all three delayed spin-ups are observed in relatively large Crab glitches and larger glitch corresponds to longer delayed spin-up timescale, it is natural and meaningful to consider the link between τ_d and $\Delta\nu$. We perform the pure mathematical power law fitting as shown in Figure 4. A fitting function in the form of $\tau_d/(\text{days}) = a_4\Delta\nu^{b_4} + c_4$ (black line in Figure 4) gives $a_4 = 0.9467 \pm 0.022$, $b_4 = 0.2978 \pm 0.0046$, $c_4 = -0.4332 \pm 0.022$ with $\chi^2/\text{dof} = 4.81 \times 10^{-7}/0$. Though the fitting curve goes across the data points and seems to fit the data well, it is not reliable in principle as the data is much too less and any other functions may fit




FIG. 7.— Persistent shift as a function of fractional increase in spin down rate. If cumulative effect should be considered, we take the sum of fractional increase in spin down of several neighboring glitches as the abscissa.

these three points well. However, we noticed that the index $b_4 = 0.2978 \pm 0.0046$ is well within the uncertainty of $b_3 = 0.2536 \pm 0.06654$, it is probably that they are highly identical. Using $|\Delta\dot{\nu}_p| = (a_3\Delta\nu^{b_3} + c_3) \times 10^{-15} \text{ s}^{-2}$ and $\tau_d = (a_4\Delta\nu^{b_4} + c_4)$ days, we then try to fit τ_d versus $\Delta\nu$ by $\tau_d = |\Delta\dot{\nu}_p|/K_1$ through minimizing the χ^2 value, our calculations give the values $K_1 = 201.98 \times 10^{-15} \text{ s}^{-2} \text{ days}^{-1}$ and $\chi^2 = 0.00021$, as shown in Figure 5. On the other hand, we try to fit $|\Delta\dot{\nu}_p|$ versus $\Delta\nu$ by $|\Delta\dot{\nu}_p| = K_2\tau_d$ in the same way, our calculations give $K_2 = 196.08 \times 10^{-15} \text{ s}^{-2} \text{ days}^{-1}$ and $\chi^2 = 2698.87$, as shown in Figure 6. This cross check shows the possibility that the persistent shifts and delayed spin-up timescales versus glitch size follow the same power law distribution. Furthermore, ratio of absolute persistent shift to delayed spin-up timescale for glitch MJD 47767.504 is $|\Delta\dot{\nu}_p|/\tau_d \approx 187.5 \times 10^{-15} \text{ s}^{-2} \text{ days}^{-1}$, and for glitch MJD 50260.031 $|\Delta\dot{\nu}_p|/\tau_d \approx 232 \times 10^{-15} \text{ s}^{-2} \text{ days}^{-1}$, it is obvious that $187.5 \times 10^{-15} \text{ s}^{-2} \text{ days}^{-1} < K_1 \approx K_2 < 232 \times 10^{-15} \text{ s}^{-2} \text{ days}^{-1}$, which suggests a possible linear correlation between the persistent shift value and the delayed spin-up timescale, the possible linear relationship can be further tested by future measurement of the persistent shift of glitch MJD 58064.555.

Finally, we analyze the correlation between $\Delta\dot{\nu}_p$ and $\Delta\dot{\nu}/\nu$, as shown in Figure 7. The distribution is sparse and even worse if the only point in the top right corner of Figure 7 is not considered. The sparse distribution indicates weak correlation between $\Delta\dot{\nu}_p$ and $\Delta\dot{\nu}/\nu$, which means small possibility that persistent shift results from the decoupled moment of inertia that do not re-couple again. This is consistent with the absence of such persistent shift in the Vela pulsar and suggests some unknown physical difference between the Crab and Vela pulsars.

3. SUMMARY AND DISCUSSION

We have performed a statistic study on all measured values during the post-glitch recovery process in the Crab pulsar. Our pure mathematical fitting results show that, persistent shifts for the relatively large and small glitches may either have different linear dependence on glitch size or follow one single power law function.

Interestingly, the fact that the power law fitting also applies to delayed spin-up timescales demonstrates the

merit of a single power law fitting to persistent shift values. To overcome the drawback of too little delayed spin-up timescale data, we perform the cross check by applying the best fitting functions of persistent shift and delayed spin-up timescale to the other one respectively, at certain coefficients K_1 and K_2 which minimize the χ^2 values. The result that coefficients K_1 and K_2 are pretty close indicates a tight linear relationship between the persistent shift values and the delayed spin-up timescales. This strongly support the conclusion that they may have the same physical origin. As an explanation, we can expect a physical mechanism which can result in the extra angular momentum transfer $I\Delta\nu_d = \Delta I(\nu_s - \nu)$ and simultaneously the change in neutron star structure denoted by the fractional change of effective moment of inertia $\Delta I/I$ through the material transfer from the superfluid to normal component, where ν_s and ν are spin frequencies of the superfluid and normal components respectively, $\nu_s - \nu$ is the spin lag. The net spin-down rate of the star(the persistent shift) then arrives at $\Delta\dot{\nu}_p = \dot{\nu}\Delta I/I$ by angular momentum conservation, $\dot{\nu}$ is the spin down rate of the Crab pulsar, at present, $\nu = 29.63 \text{ Hz}$ and $\dot{\nu} = -3.68 \times 10^{-10} \text{ Hz s}^{-1}$. The Combination with the extra angular momentum transfer immediately gives a linear relationship $\Delta\dot{\nu}_p = \frac{\dot{\nu}\Delta\nu_d}{\nu_s - \nu}\tau_d$, the coefficient $\dot{\nu}\Delta\nu_d/(\nu_s - \nu)$ has the same unit with K_1 and K_2 . If we take the spin lag as $1.2 \times 10^{-3} \text{ s}^{-1}$ (Haskell & Melatos 2015), absolute value of this coefficient is about $200 \times 10^{-15} \text{ s}^{-2} \text{ days}^{-1}$, supporting the above fitting results. Monitoring of post-glitch evolution has been applied to constrain quantities such as the fractional moment of inertia involved in the re-coupling process and the mutual friction parameters which govern the re-coupling between the superfluid and normal components, however, requirement of self-consistency between the delayed spin-up and persistent shift phenomena may set more stringent constraints on these. A new window may be opened to probe the interior of neutron star, allowing stringently constraints on the vortex motion, even on the nuclear equation of state in high densities.

Future measurement of persistent shift of glitch MJD 58064.555 can serve as a test to the linear relationship between the persistent shift values and the delayed spin-up timescales. Lyne's linear fitting predicts $\Delta\dot{\nu}_p \sim -1000 \times 10^{-15} \text{ s}^{-2}$, while our linear fitting gives $\Delta\dot{\nu}_p \sim -457 \times 10^{-15} \text{ s}^{-2}$, our best power law fitting gives $\Delta\dot{\nu}_p \sim -(344 \pm 120) \times 10^{-15} \text{ s}^{-2}$. Zhang et al. observed this glitch in the 0.5-10 keV X-ray band with the *X-Ray Pulsar Navigation-I* (XPNAV-1) satellite, using the first 100 days-long post-glitch data, their fittings gave a persistent shift $\Delta\dot{\nu}_p \sim -(1040 \pm 150) \times 10^{-15} \text{ s}^{-2}$ (Zhang et al. 2018). However, the recovery process was not completed at that time, if the fitting function $\delta\dot{\nu} = |\Delta\dot{\nu}_p| \times (0.46 \times \exp(-t/320) - 1.0)$ (t is the time since the glitch epoch in units of days) is universal for all Crab glitches, the inferred final persistent shift should be $\Delta\dot{\nu}_p \sim -(1567 \pm 226) \times 10^{-15} \text{ s}^{-2}$.

It should be noticed that, fittings in section 2 are relatively rough at present because of (i)the lack of more data points, (ii)the effect of contamination of neighboring glitches, (iii)the non-uniform distribution of data points in glitch size. Thus, more persistent shift and delayed spin-up events are urgently needed for statistics and the-

oretical work.

The authors would like to thank Jin Wu from Key Laboratory of Quark and Lepton Physics (MOE) and Institute of Particle Physics, Central China Normal Univer-

sity for his help in data analysis. This work is supported by National Natural Science Foundation of China (Grant No. 11773011). Correspondence should be addressed to W. H. Wang (email: wangweihua@mails.ccnu.edu.cn) and X. P. Zheng (email: zhxp@phy.ccnu.edu.cn).

REFERENCES

- Taylor, J. H., Manchester, R. N., Lyne, A. G. 1993, *VizieR Online Data Catalog*, 7156
- Radhakrishnan, V., & Manchester, R. N. 1969, *Nature(London)*, 222, 228
- Reichely, P. E., & Downs, G. S. 1969, *Nature(London)*, 222, 229
- Ruderman, M. 1969, *Nature(London)*, 223, 597
- Pines, D., Shaham, J., Ruderman, M. 1972, *Nature Phys. Sci.*, 237, 83
- Michel, F. C. 1970, *ApJL*, 159, L25
- Scargle, J. D., & Pacini, F. 1971, *Nature Physical Science*, 232, 144
- Pines, D., Shaham, J., Ruderman, M., 1974, *IAU proceedings*, 53, 189
- Baym, G., Pethick, C., Pines, D., 1969, *Nature(London)*, 224, 673
- Anderson, P. W., & Itoh, N., 1975, *Nature(London)*, 256, 25
- Alpar, M. A., Anderson, P. W., Pines, D., & Shaham, J., 1984, *ApJ*, 276, 325
- Fuentes, J. R., Espinoza, C. M., Reisenegger, A., et al. 2017, *A&A*, 608, 131
- Lyne, A. G., Jordan, C. A., Graham-Smith, F., et al. 2015, *MNRAS*, 446, 857
- Lyne, A. G., Graham-Smith, F., & Pritchard, R. S., 1992, *Nature*, 359, 706
- Wong, T., Backer, D. C., Lyne, A. G., 2001, *ApJ*, 548, 447
- Shaw, B., Lyne, A. G., Stappers, B. W., et al. 2018, *MNRAS*, 478, 3832
- Yuan, J. P., Manchester, R. N., Wang, N., et al. 2010, *ApJL*, 719, L111
- Alpar, M. A., Chau, H. F., Cheng, K. S., et al. 1994, *ApJ*, 427, L29
- Espinoza, C. M., Lyne, A. G., Stappers, B. W., et al. 2011, *MNRAS*, 414, 1679
- Espinoza, C. M., Antonopoulou, D., Stappers, B. W., et al. 2014, *MNRAS*, 440, 2755
- Haskell, B., & Melatos, A., 2015, *International Journal of Modern Physics D*, 24, 1530008
- Zhang, X. Y., Shuai, P., Huang, L. W., et al. 2018, *ApJ*, 866, 82