

# A generalized Higgs potential with two degenerate minima for a dark QCD matter scenario

M.F.M. Lutz<sup>a,b</sup>, Yonggoo Heo<sup>c</sup>, Xiao-Yu Guo<sup>a</sup>

<sup>a</sup>*GSI Helmholtzzentrum für Schwerionenforschung GmbH,  
Planck Str. 1, 64291 Darmstadt, Germany*

<sup>b</sup>*Technische Universität Darmstadt, D-64289 Darmstadt, Germany*

<sup>c</sup>*Suranaree University of Technology, Nakhon Ratchasima, 30000, Thailand*

---

## Abstract

We consider the Higgs potential in generalizations of the Standard Model. The possibility of the potential to develop two almost degenerate minima is explored. This would imply that QCD matter at two distinct sets of quark masses is relevant for astrophysics and cosmology. If in the exotic minimum the QCD matter ground state is electromagnetically neutral, dark matter may consist of QCD matter and antimatter in bubbles of the Higgs field. We predict an abundance of  $\gamma$  rays in the few MeV region as messengers of dark matter regions in space. In addition the ratio of dark matter to normal matter is expected to show a time dependence.

---

## 1. Introduction

Dark-matter studies receive considerable attention in fundamental research (see e.g. [1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16]). Various scenarios proposed require new particles in extensions of the Standard Model (SM) (see e.g. [1, 6]).

The purpose of our Letter is to discuss a possible alternative of such scenarios based on exotic QCD matter. In a recent work the authors presented a detailed study suggesting that QCD matter depends crucially on the Higgs field [17, 18, 19]. Within the SM the quark masses in QCD are proportional to the Higgs field. As a consequence, changing its ground state value does change the quark masses in QCD, however, in a manner that keeps all quark-mass ratios fixed. In [19] a possible first order transition along the Higgs field




Figure 1: Isospin averaged baryon masses as a function of  $r = m_s/m_s^{\text{phys}}$  along the Higgs trajectory with  $2m_s/(m_u + m_d) = 26$  kept constant.

trajectory was discussed. It is compatible with current QCD lattice simulations of the baryon ground state masses, but should be scrutinized by further dedicated QCD lattice studies.

In Fig. 1 we show our prediction of the baryon masses along the Higgs trajectory [17, 18, 19]. The bands in the plot provide an estimate of uncertainties based on our Fit 1 and Fit 2 scenarios as discussed in [19]. At fixed ratio  $m_s/m = 26$  the masses are plotted as functions of the strange quark mass. The key observation is that within a critical region of the Higgs field, baryonic matter and antimatter are composed from  $\Lambda$  and  $\bar{\Lambda}$  particles rather than from nucleons and anti-nucleons. This follows from the relation  $M_\Lambda < M_N$ , which holds at a specific range of the strange quark mass. We point out that our dark-matter scenario does not rely necessarily on a first order transition. Since  $\Lambda$  particles are electromagnetically neutral such matter does not radiate and therefore appears dark. Since the Higgs sector of the SM drives a possible electroweak phase transition and underlies baryogenesis models (see e.g [20, 21, 22, 23]) it is important to explore exotic Higgs sector generalizations of the SM in more detail.

## 2. The Higgs potential

We consider the Higgs sector of the SM [24, 25, 26, 27, 28, 20, 29, 30, 23, 31, 32, 33]. At tree-level the Higgs potential in the SM can be expressed in terms of two parameters only

$$V(H) = \frac{M_h^2}{2v^2} \left( H^\dagger H - \frac{v^2}{2} \right)^2, \quad (1)$$

with the complex doublet Higgs field  $H$ , the Higgs mass parameter  $M_h \simeq 125.2$  GeV and the vacuum expectation value  $v \simeq 246.2$  GeV of the Higgs field in its physical vacuum state [34]. The value of  $v$  plays a decisive role in the QCD part of the SM since all quark masses are proportional to  $v$ . In this work we are interested in the Higgs potential at  $H^\dagger H \leq v^2$ , where it is known that even loop corrections in the SM are sizeable (see e.g. [24, 26, 28, 29, 35, 30, 36]). Since the Higgs potential will be affected in most extensions of the SM we follow here a phenomenological path where we explore the consequence of a fine-tuned potential with two degenerate minima. An effective field theory approach that implies two degenerate minima would require at least  $(H^\dagger H)^3$  and  $(H^\dagger H)^4$  operators. Consider the specific form

$$V(H) = \frac{2 M_h^2}{v^6 (1 - r^2)^2} \left( H^\dagger H - \frac{v^2}{2} \right)^2 \left( H^\dagger H - \frac{v_a^2}{2} \right)^2, \\ \text{and} \quad r = v_a/v = m_s/m_s^{\text{phys}}, \quad (2)$$

with  $v_a$  the vacuum expectation value of the Higgs field at the exotic minimum. By construction, the model potential (2) has two degenerate minima. At its physical one it recovers the empirical mass of the Higgs. The ratio  $r = v_a/v$  determines the strange quark mass in the exotic minimum.

The puzzle with (2) is that it may be unnatural in the size of its dimension-full operators. However, we may recast the problem by considering loop corrections (see e.g [24, 26, 29]). In the presence of multi-loop effects we may use the phenomenological ansatz

$$V(H) = \frac{M_h^2}{2v^2 [\log(\gamma + r^2) - \log(\gamma + 1)]^2} \left( H^\dagger H - \frac{v^2}{2} \right)^2 \\ \times \left( \log [\gamma + 2 H^\dagger H/v^2] - \log[\gamma + r^2] \right)^2, \quad (3)$$

where the particular form of the log term with the parameter  $\gamma$  is taken from [30]. There the value  $\gamma = 0.1$  is used. We note that the Higgs sector is the

least controlled part of the SM and therefore may be subject to significant model modifications.

According to Fit 1 and Fit 2 we expect dark QCD matter in the range  $0.39 < r < 0.57$  and  $0.39 < r < 0.54$  respectively. The critical values are close to those as derived in [19] on the unphysical trajectory where  $m_u + m_d$  is kept constant. In Fig. 2 we plot the effective potentials of (1-3) as a function of  $\sqrt{2 H^\dagger H}/v$  for the particular choice  $r = 0.45$ . The two degenerate minima are clearly visible for any of the three choices  $\gamma = 0.1$ ,  $\gamma = 0.2$  and  $\gamma = 0.3$ . With the parameter  $\gamma$  we can efficiently dial the magnitude of the Higgs potential close to the origin. In the vicinity of the two local minima we find a rather mild dependence on the form of our parametrization. The polynomial ansatz (2) or the log form (3) lead to almost indistinguishable results. We emphasize that both models are compatible with empirical constraints on the Higgs potential as discussed in [37]. For instance at  $r = 0.45$  we extract from (2) and (3) the range  $3.8 < \kappa_\lambda < 6$  for the three Higgs coupling constant  $\kappa_\lambda$ . This is well compatible with the empirical  $2\text{-}\sigma$  interval  $-5.0 < \kappa_\lambda < 12.1$  from ATLAS [37]. Our estimate excludes the SM value  $\kappa_\lambda = 1$ .

One may object to such a fine-tuned Higgs potential. However, we wish to recall that there are ample cases in physics in which a system is driven by fine-tuned dynamical assumptions. In particular the SM itself has various fine-tuning issues already. At this stage of the development we would not worry too much. Rather, we discuss in some detail the consequences of a possible dark QCD matter scenario.

Dark matter is believed to account for approximately 85% of the matter in the universe (see e.g. [38]). Within our model, matter freezes out from the Big Bang into the two available Higgs potential minima with roughly equal abundance. Thus we would start out with already 50% dark matter. This estimate is based on the assumption that the Higgs potential of Fig. 2 remains basically unchanged up to temperatures  $T \sim 200$  MeV, where matter constituents are quarks and gluons rather than hadrons. In a conventional Big Bang scenario, initially, a similar amount of matter and antimatter is produced. Sphaleron effects induce a small matter-antimatter asymmetry as a consequence of direct CP violation terms in the SM. However, the observed matter to antimatter ratio cannot be accommodated (see e.g. [39, 40, 41, 42, 30]). It would be of interest to investigate in the future in how far our exotic Higgs potential would affect this mechanism [30]. In the following we will consider the universe at times where it is already matter dominated.




Figure 2: A Higgs potential with two local minima as introduced in (2) and (3). It is compared with the tree-level potential of the SM (1).

If the exotic minimum in the Higgs potential is slightly metastable, we expect a scenario where the vacuum shows bubbles with dark QCD matter inside, but normal QCD matter outside. Inside the bubbles the matter or antimatter ground states consist of  $\Lambda$  or  $\bar{\Lambda}$  particles, however with exotic properties as shown by Fig. 1.

Let us explore the stability of a possible Higgs bubble. Since the boundary of such a bubble stores a significant amount of energy, there is a tendency that such a bubble shrinks or even collapses. From Fig. 2 we estimate the energy density

$$\epsilon_{\text{Higgs}} \simeq 1.1 \times 10^9 \text{ GeV}/\text{fm}^3 \quad (4)$$

from the Higgs potential taking in-between its two minima. We now assume a Higgs bubble with spherical geometry characterized by a radius  $R$  and a surface thickness  $d$ . That implies the total surface energy

$$E_{\text{surface}} = 4 \pi d R^2 \epsilon_{\text{Higgs}} + 4 \pi R^2 (\Delta v)^2/d, \quad (5)$$

where the second term in (5) follows from the kinetic term of the Higgs field. From Fig. 2 we can read off the change of the Higgs field across the

surface with  $(\Delta v)^2 \simeq v^2/8$ . We estimate the bulk energy by a free-Fermi gas approximation

$$E_{\text{bulk}} = \left( M_{\Lambda, \text{in}} + \frac{3}{10} k_F^2 / M_{\Lambda, \text{in}} \right) N + \frac{4\pi R^3}{3} \Delta\epsilon - \frac{3}{5} \frac{M_{\Lambda, \text{in}}^2}{R} G N^2, \\ N = \frac{4(R k_F)^3}{9\pi}, \quad (6)$$

with the gravitational constant  $G \simeq 6.709 \times 10^{-39} \text{ GeV}^{-2}$  and  $N$  the total number of  $\Lambda$ 's in the Higgs bubble. Their Fermi momentum is denoted by  $k_F$  with  $\rho = k_F^3 / (3\pi^2)$ , where  $\rho$  specifies the dark-matter density in the bubble. We parameterize a supposedly small difference in the vacuum energy densities at the two Higgs minima by  $\Delta\epsilon > 0$ , where we assume the dark-matter vacuum to be slightly disfavored.

A Higgs bubble can be stable provided that it encloses a sufficient amount of dark matter. We can make this more quantitative by a minimization of its energy  $E = E_{\text{bulk}} + E_{\text{surface}}$  with respect to the surface thickness  $d$  and the radius  $R$  at a fixed value of the total number of  $\Lambda$ 's in the bubble. From this we find the two relations,

$$d \simeq \sqrt{(\Delta v)^2 / \epsilon_{\text{Higgs}}} \simeq 6 \times 10^{-3} \text{ fm}, \\ \frac{1}{R} + \frac{R^2 G k_F^6 M_{\Lambda, \text{in}}^2}{135 \pi^2 d \epsilon_{\text{Higgs}}} = \frac{\Delta\epsilon}{4 d \epsilon_{\text{Higgs}}} \left( \frac{k_F^5}{15 \pi^2 M_{\Lambda, \text{in}} \Delta\epsilon} - 1 \right). \quad (7)$$

This implies that at given  $k_F$  it follows that  $\Delta\epsilon$  must be smaller than a critical value,

$$\Delta\epsilon < \Delta\epsilon_{\text{crit}} = \frac{k_F^5}{15 \pi^2 M_{\Lambda, \text{in}}}, \quad (8)$$

as to keep the dark-matter bubble stable. We checked that all second derivatives are positive so that with (7) we have at least a local minimum of the dark matter system (5, 6). In Fig. 3 we show the radius,  $R$ , and Fermi momentum,  $k_F$ , of the Higgs bubble as a function of  $N$  at various fixed values of  $\Delta\epsilon$ . Within the range  $10^{31} < N < 10^{57}$  the value of  $k_F < 250 \text{ MeV}$  is small enough to justify our free-Fermi gas approximation. We expect our results to hold at the qualitative level.




Figure 3: The radius and Fermi momentum as a function of  $N$  at fixed  $\Delta\epsilon$ . Note that  $N_{\text{Sun}} \simeq M_{\text{Sun}}/M_{\text{Nucleon}} \simeq 1.2 \times 10^{57}$ .

It is left to check whether such a dark matter Higgs bubble is stable with respect to a decay into a more conventional object consisting out of normal baryonic matter. A useful quantity to consider is the energy per particle in the bubble,  $(E_{\text{surface}} + E_{\text{bulk}})/N$ , with  $d$  and  $R$  as given in (7). In Fig. 4 we show such a dependence at various fixed values of  $\Delta\epsilon$ . The energy per particle is significantly smaller than the free nucleon mass and therefore, at least in the region  $10^{31} < N < 10^{57}$ , where the effects from gravity are not dominating the system yet, there is no phase-space available for such a decay.

As an example consider  $N = M_{\text{sun}}/M_N \simeq 1.2 \times 10^{57}$  for which it follows  $R \simeq 16$  km and  $k_F \simeq 250$  MeV at  $\Delta\epsilon = 0$ . The particle density with  $\rho \simeq 0.07 \text{ fm}^{-3}$  would be sufficiently small as to justify the application of the free-Fermi gas approximation. As we further increase  $N > N_{\text{Sun}}$  the dark matter will turn more dense and will be sensitive to the equation of state of the  $\Lambda$  particles in the Higgs bubble. It would be important to establish the latter from QCD and to derive mass limits for the dark matter bubbles. Here the Tolman-Oppenheimer-Volkoff equation has to be supplemented by a suitable boundary condition at the Higgs bubble surface. This may open the possibility for the existence of massive compact objects, with properties distinct to those of neutron stars and/or black holes of unconventionally small masses [43].




Figure 4: The energy per particle  $E_\Lambda/N$  as a function of  $N$  at fixed  $\Delta\epsilon$ . Note that  $N_{\text{Sun}} \simeq M_{\text{Sun}}/M_{\text{Nucleon}} \simeq 1.2 \times 10^{57}$ .

In a meson-exchange phenomenology, the repulsive omega-meson exchange process is expected to dominate the short-range interaction of  $\Lambda$  particles in the Higgs bubble. Due to the approximate isospin conservation, pion- and rho-meson exchange processes are suppressed. It remains the eta-meson exchange and the two-pion exchange contributions, which may bring in some weak intermediate-range attractive forces [44, 45, 46]. While there appears to be a rather weak net attraction at the physical point [46, 47, 48, 49] available studies suggest a sizeable quark-mass dependence thereof [44, 50, 48, 49]. We conclude that at the exotic Higgs minimum, that comes at much smaller up and down quark masses, there is little evidence to expect this weak attraction to survive. Whether and how massive dark-matter clusters form depends on the subtle balance of the gravitational force and the short-range strong interactions in the Higgs bubble.

We conclude that in any case the typical dark-matter density in a Higgs bubble should be significantly larger than the density of a cold interstellar medium, which is characterized by a baryon-number density smaller than about  $10^6/\text{cm}^3$ . In this context we discuss the so-called Bullet Cluster [51, 52]. While the radial velocity distributions of stars inside a galaxy or data on gravitational lensing effects (see e.g. [53, 38, 54, 8, 55]) put constraints on the dark matter distributions in and outside galaxies, more significant information on the possible nature of dark matter is set by the observation



of collisions of galaxy clusters [51, 52]. It is found that in such a collision there is no direct hint pointing at any sizeable interaction of dark matter with ordinary matter [52]. In this context we have to discuss how a Higgs bubble interacts with protons from the intergalactic hot gas. The relative velocity of the two colliding galaxies in [51] is of the order of 4500 km/s. An intergalactic gas of temperature  $T \simeq 6$  keV implies a typical proton velocity of about 1300 km/s. Thus most of the protons from the gas do not have sufficient kinetic energy to invade the bubble. In turn there will be no strong interaction effects visible. Second we need to consider the case where Higgs bubbles from the two galaxies collide. The chance that this happens depends on the typical size of such bubbles, which are not well constrained at this stage. They depend on the details of the Higgs potential, in particular the size of  $\Delta\epsilon$  term, and a cosmological model. The smaller the typical size of the Higgs bubbles, the smaller the likelihood that such a process turns relevant in a galaxy merger event. Even if two bubbles start to overlap, we would expect that the two bubbles merge into a larger one, since this reduces the energy stored in their surface. The residual interaction of the Lambda particles with kinetic energies of at most a few MeV should be dominated by elastic processes. In turn we do not see any reason to expect a strong visible effect of the dark matter component in such a galaxy collision event.

Last, we turn to a most interesting process where a sufficiently energetic cosmic proton tries to enter a dark-matter region in space with relative velocity,  $v_p$ . Note that, depending on the energy such a proton may be even trapped inside the dark-matter bubble and therefore the ratio of dark matter to normal matter is expected to show a time dependence in our dark-matter scenario [56]. According to Fig. 1 the nucleon mass inside the bubble is only up  $10 \text{ MeV} = \Delta M_N$  larger than its mass outside the bubble. Thus on the way into the bubble the nucleon has to either transfer momentum to the Higgs bubble and/or radiate photons. Such a Bremsstrahlung spectrum should be limited to  $\gamma$  rays with energies less than that 10 MeV. Here a crucial parameter is the acceleration,  $a \simeq (c^2/\gamma_p^2) \Delta M_N / (d M_N) \simeq 8 \times 10^{23} (c/\gamma_p^2) / s$ , of the proton across the Higgs bubble surface, since its total radiation power is proportional to  $a^2 \gamma_p^4$  with  $\gamma_p = (1 - v_p^2/c^2)^{-1/2}$ . To this extent our Higgs bubbles glim with a characteristic spectrum which depends on the details of the Higgs potential. It may be possible to identify such photons with satellite-based detectors like e-Astrogram or AMEGO [57, 58]. We note that so far there is basically no gamma ray data available in the 1-10 MeV region.

### 3. Summary and conclusions

We constructed a phenomenological Higgs potential with two degenerate local minima. It was argued that such a generalization of the SM may lead to dark QCD matter that lives in bubbles of the Higgs field, with normal QCD matter outside and dark QCD matter inside. Within the bubbles we expect exotic  $\Lambda$  and  $\bar{\Lambda}$  particles, that are formed by QCD at unconventionally small up, down and strange quark masses. We predict an abundance of  $\gamma$  rays in the few MeV region as messengers of dark matter regions in space. In addition the ratio of dark matter to normal matter is expected to show a time dependence.

It would be interesting to further scrutinize the dark QCD matter scenario proposed here. With current QCD lattice techniques it is possible to substantiate or rule out such a scenario by further studies of the strange quark-mass dependence of the nucleon and  $\Lambda$  baryon masses. It would be important to establish a more fundamental framework in which such an exotic Higgs potential is implied.

### Acknowledgments

Hans Feldmeier, Gregor Kasieczka, Thomas Mannel, Walter Schön, Madeleine Soyeur, Christian Sturm and David Urner are acknowledged for stimulating discussions. Y. H. received partial support from Suranaree University of Technology, Office of the Higher Education Commission under NRU project of Thailand (SUT-COE: High Energy Physics and Astrophysics) and SUT-CHE-NRU (Grant No. FtR.11/2561).

## References

- [1] G. Bertone, D. Hooper, J. Silk, Particle dark matter: Evidence, candidates and constraints, *Phys. Rept.* 405 (2005) 279–390. [arXiv:hep-ph/0404175](#), [doi:10.1016/j.physrep.2004.08.031](#).
- [2] M. Rocha, A. H. G. Peter, J. S. Bullock, M. Kaplinghat, S. Garrison-Kimmel, J. Onorbe, L. A. Moustakas, Cosmological Simulations with Self-Interacting Dark Matter I: Constant Density Cores and Substructure, *Mon. Not. Roy. Astron. Soc.* 430 (2013) 81–104. [arXiv:1208.3025](#), [doi:10.1093/mnras/sts514](#).
- [3] S. Tulin, H.-B. Yu, K. M. Zurek, Beyond Collisionless Dark Matter: Particle Physics Dynamics for Dark Matter Halo Structure, *Phys. Rev. D* 87 (11) (2013) 115007. [arXiv:1302.3898](#), [doi:10.1103/PhysRevD.87.115007](#).
- [4] K. Petraki, R. R. Volkas, Review of asymmetric dark matter, *Int. J. Mod. Phys. A* 28 (2013) 1330028. [arXiv:1305.4939](#), [doi:10.1142/S0217751X13300287](#).
- [5] C. Gordon, O. Macias, Dark Matter and Pulsar Model Constraints from Galactic Center Fermi-LAT Gamma Ray Observations, *Phys. Rev. D* 88 (8) (2013) 083521, [Erratum: *Phys. Rev. D* 89, no. 4, 049901 (2014)]. [arXiv:1306.5725](#), [doi:10.1103/PhysRevD.88.083521](#), [doi:10.1103/PhysRevD.89.049901](#).
- [6] L. Hui, J. P. Ostriker, S. Tremaine, E. Witten, Ultralight scalars as cosmological dark matter, *Phys. Rev. D* 95 (4) (2017) 043541. [arXiv:1610.08297](#), [doi:10.1103/PhysRevD.95.043541](#).
- [7] B. Carr, F. Kuhnel, M. Sandstad, Primordial Black Holes as Dark Matter, *Phys. Rev. D* 94 (8) (2016) 083504. [arXiv:1607.06077](#), [doi:10.1103/PhysRevD.94.083504](#).
- [8] P. F. de Salas, K. Malhan, K. Freese, K. Hattori, M. Valluri, On the estimation of the Local Dark Matter Density using the rotation curve of the Milky Way [arXiv:1906.06133](#).
- [9] P. Montero-Camacho, X. Fang, G. Vasquez, M. Silva, C. M. Hirata, Revisiting constraints on asteroid-mass primordial black holes as dark

- matter candidates, JCAP 1908 (2019) 031. [arXiv:1906.05950](#), [doi:10.1088/1475-7516/2019/08/031](#).
- [10] H. Grote, Y. V. Stadnik, Novel signatures of dark matter in laser-interferometric gravitational-wave detectors [arXiv:1906.06193](#).
- [11] Y. Bai, N. Orlofsky, Primordial Extremal Black Holes as Dark Matter [arXiv:1906.04858](#).
- [12] S. Savastano, L. Amendola, J. Rubio, C. Wetterich, Primordial dark matter halos from fifth-forces [arXiv:1906.05300](#).
- [13] C. E. Alvarez-Salazar, O. L. G. Peres, B. L. Sánchez-Vega, The dark matter puzzle in a class of models with gauge symmetry  $SU(3)_C \otimes SU(3)_L \otimes U(1)_N$ , Astron. Nachr. 340 (1-3) (2019) 135–138. [doi:10.1002/asna.201913577](#).
- [14] A. R. Romero Castellanos, C. E. Alvarez-Salazar, B. L. Sánchez-Vega, Constraints on axionic dark matter in the 3-3-1 model, Astron. Nachr. 340 (1-3) (2019) 131–134. [doi:10.1002/asna.201913576](#).
- [15] S. V. Ketov, Y. Aldabergenov, Inflation, dark energy, and dark matter in supergravity, Astron. Nachr. 340 (1-3) (2019) 126–130. [doi:10.1002/asna.201913575](#).
- [16] M. De Napoli, Production and Detection of Light Dark Matter at Jefferson Lab: The BDX Experiment, Universe 5 (5) (2019) 120. [doi:10.3390/universe5050120](#).
- [17] M. F. M. Lutz, R. Bavontaweepanya, C. Kobdaj, K. Schwarz, Finite volume effects in the chiral extrapolation of baryon masses, Phys. Rev. D90 (5) (2014) 054505. [arXiv:1401.7805](#), [doi:10.1103/PhysRevD.90.054505](#).
- [18] M. F. M. Lutz, Y. Heo, X.-Y. Guo, On the convergence of the chiral expansion for the baryon ground-state masses, Nucl. Phys. A977 (2018) 146–207. [arXiv:1801.06417](#), [doi:10.1016/j.nuclphysa.2018.05.007](#).
- [19] X.-Y. Guo, Y. Heo, M. F. M. Lutz, On a first order transition in QCD with up, down and strange quarks [arXiv:1907.00714](#).

- [20] A. Noble, M. Perelstein, Higgs self-coupling as a probe of electroweak phase transition, *Phys. Rev. D* 78 (2008) 063518. [arXiv:0711.3018](#), [doi:10.1103/PhysRevD.78.063518](#).
- [21] Q.-H. Cao, F. P. Huang, K.-P. Xie, X. Zhang, Testing the electroweak phase transition in scalar extension models at lepton colliders, *Chin. Phys. C* 42 (2) (2018) 023103. [arXiv:1708.04737](#), [doi:10.1088/1674-1137/42/2/023103](#).
- [22] A. Kobakhidze, L. Wu, J. Yue, Electroweak Baryogenesis with Anomalous Higgs Couplings, *JHEP* 04 (2016) 011. [arXiv:1512.08922](#), [doi:10.1007/JHEP04\(2016\)011](#).
- [23] M. Reichert, A. Eichhorn, H. Gies, J. M. Pawłowski, T. Plehn, M. M. Scherer, Probing baryogenesis through the Higgs boson self-coupling, *Phys. Rev. D* 97 (7) (2018) 075008. [arXiv:1711.00019](#), [doi:10.1103/PhysRevD.97.075008](#).
- [24] S. R. Coleman, E. J. Weinberg, Radiative Corrections as the Origin of Spontaneous Symmetry Breaking, *Phys. Rev. D* 7 (1973) 1888–1910. [doi:10.1103/PhysRevD.7.1888](#).
- [25] S. R. Coleman, The Fate of the False Vacuum. 1. Semiclassical Theory, *Phys. Rev. D* 15 (1977) 2929–2936, [Erratum: *Phys. Rev. D* 16,1248(1977)]. [doi:10.1103/PhysRevD.15.2929](#), [doi:10.1103/PhysRevD.16.1248](#).
- [26] S. P. Martin, A Supersymmetry primer, *Adv. Ser. Direct. High Energy Phys.* 18 (1998) 1–98. [arXiv:hep-ph/9709356](#), [doi:10.1142/9789812839657\\_0001](#), [doi:10.1142/9789814307505\\_0001](#).
- [27] G. Isidori, G. Ridolfi, A. Strumia, On the metastability of the standard model vacuum, *Nucl. Phys. B* 609 (2001) 387–409. [arXiv:hep-ph/0104016](#), [doi:10.1016/S0550-3213\(01\)00302-9](#).
- [28] C. Grojean, G. Servant, J. D. Wells, First-order electroweak phase transition in the standard model with a low cutoff, *Phys. Rev. D* 71 (2005) 036001. [arXiv:hep-ph/0407019](#), [doi:10.1103/PhysRevD.71.036001](#).
- [29] D. Buttazzo, G. Degrandi, P. P. Giardino, G. F. Giudice, F. Sala, A. Salvio, A. Strumia, Investigating the near-criticality of the Higgs

- boson, JHEP 12 (2013) 089. [arXiv:1307.3536](#), [doi:10.1007/JHEP12\(2013\)089](#).
- [30] M. Spannowsky, C. Tamarit, Sphalerons in composite and non-standard Higgs models, Phys. Rev. D95 (1) (2017) 015006. [arXiv:1611.05466](#), [doi:10.1103/PhysRevD.95.015006](#).
- [31] X. Gan, A. J. Long, L.-T. Wang, Electroweak sphaleron with dimension-six operators, Phys. Rev. D96 (11) (2017) 115018. [arXiv:1708.03061](#), [doi:10.1103/PhysRevD.96.115018](#).
- [32] B. Jain, S. J. Lee, M. Son, Validity of the effective potential and the precision of Higgs field self-couplings, Phys. Rev. D98 (7) (2018) 075002. [arXiv:1709.03232](#), [doi:10.1103/PhysRevD.98.075002](#).
- [33] J. Braathen, S. Kanemura, On two-loop corrections to the Higgs trilinear coupling in models with extended scalar sectors, Phys. Lett. B796 (2019) 38–46. [arXiv:1903.05417](#), [doi:10.1016/j.physletb.2019.07.021](#).
- [34] M. Tanabashi, et al., Review of Particle Physics, Phys. Rev. D98 (3) (2018) 030001. [doi:10.1103/PhysRevD.98.030001](#).
- [35] C. Tamarit, Higgs vacua with potential barriers, Phys. Rev. D90 (5) (2014) 055024. [arXiv:1404.7673](#), [doi:10.1103/PhysRevD.90.055024](#).
- [36] Y. Hamada, Higgs potential and naturalness after the Higgs discovery, Ph.D. thesis, Kyoto U. (2016).
- [37] M. Cepeda, et al., Higgs Physics at the HL-LHC and HE-LHC [arXiv:1902.00134](#).
- [38] J. de Swart, G. Bertone, J. van Dongen, How Dark Matter Came to Matter, Nature Astron. 1 (2017) 0059. [arXiv:1703.00013](#), [doi:10.1038/s41550017-0059](#), [10.1038/s41550-017-0059](#).
- [39] C. Jarlskog, Commutator of the Quark Mass Matrices in the Standard Electroweak Model and a Measure of Maximal CP Violation, Phys. Rev. Lett. 55 (1985) 1039. [doi:10.1103/PhysRevLett.55.1039](#).
- [40] M. E. Shaposhnikov, Possible Appearance of the Baryon Asymmetry of the Universe in an Electroweak Theory, JETP Lett. 44 (1986) 465–468, [Pisma Zh. Eksp. Teor. Fiz.44,364(1986)].

- [41] M. B. Gavela, P. Hernandez, J. Orloff, O. Pene, Standard model CP violation and baryon asymmetry, *Mod. Phys. Lett. A*9 (1994) 795–810. [arXiv:hep-ph/9312215](#), [doi:10.1142/S0217732394000629](#).
- [42] G. W. S. Hou, Source of CP Violation for the Baryon Asymmetry of the Universe, *Int. J. Mod. Phys. D*20 (2011) 1521–1532. [arXiv:1101.2161](#), [doi:10.1142/S0218271811019694](#).
- [43] T. A. Thompson, et al., A noninteracting low-mass black hole–giant star binary system, *Science* 366 (6465) (2019) 637–640. [doi:10.1126/science.aau4005](#).
- [44] K. Sasaki, E. Oset, M. J. Vicente Vacas, Scalar Lambda N and Lambda Lambda interaction in a chiral unitary approach, *Phys. Rev. C*74 (2006) 064002. [arXiv:nucl-th/0607068](#), [doi:10.1103/PhysRevC.74.064002](#).
- [45] H. Polinder, J. Haidenbauer, U. G. Meißner, Strangeness  $S = -2$  baryon-baryon interactions using chiral effective field theory, *Phys. Lett. B*653 (2007) 29–37. [arXiv:0705.3753](#), [doi:10.1016/j.physletb.2007.07.045](#).
- [46] E. Bauer, G. Garbarino, C. A. Rodríguez Peña, Nonmesonic weak decay of double- $\Lambda$  hypernuclei: A microscopic approach, *Phys. Rev. C*92 (1) (2015) 014301. [doi:10.1103/PhysRevC.92.014301](#).
- [47] J. Haidenbauer, U.-G. Meißner, S. Petschauer, Strangeness  $S = 2$  baryon-baryon interaction at next-to-leading order in chiral effective field theory, *Nucl. Phys. A*954 (2016) 273–293. [arXiv:1511.05859](#), [doi:10.1016/j.nuclphysa.2016.01.006](#).
- [48] K. Sasaki, S. Aoki, T. Doi, T. Hatsuda, Y. Ikeda, T. Inoue, N. Ishii, K. Murano, Coupled-channel approach to strangeness  $S = 2$  baryon-baryon interactions in lattice QCD, *PTEP* 2015 (11) (2015) 113B01. [arXiv:1504.01717](#), [doi:10.1093/ptep/ptv144](#).
- [49] K. Sasaki, S. Aoki, T. Doi, S. Gongyo, T. Hatsuda, Y. Ikeda, T. Inoue, T. Iritani, N. Ishii, T. Miyamoto, Lattice QCD studies on baryon interactions in the strangeness  $-2$  sector with physical quark masses, *EPJ Web Conf.* 175 (2018) 05010. [doi:10.1051/epjconf/201817505010](#).

- [50] S. R. Beane, et al., Present Constraints on the H-dibaryon at the Physical Point from Lattice QCD, *Mod. Phys. Lett. A* 26 (2011) 2587–2595. [arXiv:1103.2821](#), [doi:10.1142/S0217732311036978](#).
- [51] D. Clowe, A. Gonzalez, M. Markevitch, Weak-lensing mass reconstruction of the interacting cluster 1e 0657-558: Direct evidence for the existence of dark matter, *Astrophys. J.* 604 (2004) 596–603. [doi:10.1086/381970](#).
- [52] M. Markevitch, A. H. Gonzalez, D. Clowe, A. Vikhlinin, W. Forman, C. Jones, S. Murray, W. Tucker, Direct constraints on the dark matter self-interaction cross section from the merging galaxy cluster 1e 0657-56, *Astrophys. J.* 606 (2) (2004) 819–824. [doi:10.1086/383178](#).
- [53] A. B. Newman, T. Treu, R. S. Ellis, D. J. Sand, J. Richard, P. J. Marshall, P. Capak, S. Miyazaki, The Distribution of Dark Matter Over 3 Decades in Radius in the Lensing Cluster Abell 611, *Astrophys. J.* 706 (2009) 1078–1094. [arXiv:0909.3527](#), [doi:10.1088/0004-637X/706/2/1078](#).
- [54] Posti, Lorenzo, Helmi, Amina, [Mass and shape of the milky way’s dark matter halo with globular clusters from gaia and hubble](#), *A&A* 621 (2019) A56. [arXiv:1805.01408](#), [doi:10.1051/0004-6361/201833355](#). URL <https://doi.org/10.1051/0004-6361/201833355>
- [55] N. C. Relatores, A. B. Newman, J. D. Simon, R. Ellis, P. Truong, L. Blitz, A. Bolatto, C. Martin, P. Morrissey, The Dark Matter Distributions in Low-Mass Disk Galaxies. I. H $\alpha$  Observations Using the Palomar Cosmic Web Imager, *Astrophys. J.* 873 (1) (2019) 5. [arXiv:1902.09629](#), [doi:10.3847/1538-4357/ab0382](#).
- [56] R. Genzel, et al., Strongly baryon-dominated disk galaxies at the peak of galaxy formation ten billion years ago, *Nature* 543 (2017) 397. [arXiv:1703.04310](#), [doi:10.1038/nature21685](#).
- [57] R. Caputo, et al., All-sky Medium Energy Gamma-ray Observatory: Exploring the Extreme Multimessenger Universe [arXiv:1907.07558](#).
- [58] L. Oakes, et al., Combined Dark Matter searches towards dwarf spheroidal galaxies with Fermi-LAT, HAWC, HESS, MAGIC and VERITAS [arXiv:1909.06310](#).