

Some q -exponential formulas involving the double lowering operator ψ for a tridiagonal pair

Sarah Bockting-Conrad*

Abstract

Let \mathbb{K} denote an algebraically closed field and let V denote a vector space over \mathbb{K} with finite positive dimension. We consider an ordered pair of linear transformations $A : V \rightarrow V$ and $A^* : V \rightarrow V$ that satisfy the following four conditions: (i) Each of A, A^* is diagonalizable; (ii) there exists an ordering $\{V_i\}_{i=0}^d$ of the eigenspaces of A such that $A^*V_i \subseteq V_{i-1} + V_i + V_{i+1}$ for $0 \leq i \leq d$, where $V_{-1} = 0$ and $V_{d+1} = 0$; (iii) there exists an ordering $\{V_i^*\}_{i=0}^\delta$ of the eigenspaces of A^* such that $AV_i^* \subseteq V_{i-1}^* + V_i^* + V_{i+1}^*$ for $0 \leq i \leq \delta$, where $V_{-1}^* = 0$ and $V_{\delta+1}^* = 0$; (iv) there does not exist a subspace W of V such that $AW \subseteq W$, $A^*W \subseteq W$, $W \neq 0$, $W \neq V$. We call such a pair a *tridiagonal pair* on V . We assume that A, A^* belongs to a family of tridiagonal pairs said to have q -Racah type. Let $\{U_i\}_{i=0}^d$ and $\{U_i^\downarrow\}_{i=0}^d$ denote the first and second split decompositions of V . In an earlier paper we introduced a double lowering operator $\psi : V \rightarrow V$ with the notable feature that both $\psi U_i \subseteq U_{i-1}$ and $\psi U_i^\downarrow \subseteq U_{i-1}^\downarrow$ for $0 \leq i \leq d$, where $U_{-1} = 0$ and $U_{-1}^\downarrow = 0$. In the same paper, we showed that there exists a unique linear transformation $\Delta : V \rightarrow V$ such that $\Delta(U_i) \subseteq U_i^\downarrow$ and $(\Delta - I)U_i \subseteq U_0 + U_1 + \cdots + U_{i-1}$ for $0 \leq i \leq d$. In the present paper, we show that Δ can be expressed as a product of two linear transformations; one is a q -exponential in ψ and the other is a q^{-1} -exponential in ψ . We view Δ as a transition matrix from the first split decomposition of V to the second. Consequently, we view the q^{-1} -exponential in ψ as a transition matrix from the first split decomposition to a decomposition of V which we interpret as a kind of halfway point. This halfway point turns out to be the eigenspace decomposition of a certain linear transformation \mathcal{M} . We discuss the eigenspace decomposition of \mathcal{M} and give the actions of various operators on this decomposition.

Keywords. Tridiagonal pair, Leonard pair, quantum group $U_q(\mathfrak{sl}_2)$, quantum algebra, q -Racah polynomial, q -Serre relations, q -exponential function.

2010 Mathematics Subject Classification. Primary: 15A21. Secondary: 05E30.

1 Introduction

Throughout this paper, \mathbb{K} denotes an algebraically closed field. We begin by recalling the notion of a tridiagonal pair. We will use the following terms. Let V denote a vector space

*This research was partially supported by a grant from the College of Science and Health at DePaul University.

over \mathbb{K} with finite positive dimension. For a linear transformation $A : V \rightarrow V$ and a subspace $W \subseteq V$, we say that W is an *eigenspace* of A whenever $W \neq 0$ and there exists $\theta \in \mathbb{K}$ such that $W = \{v \in V \mid Av = \theta v\}$. In this case, θ is called the *eigenvalue* of A associated with W . We say that A is *diagonalizable* whenever V is spanned by the eigenspaces of A .

Definition 1.1. [9, Definition 1.1]. Let V denote a vector space over \mathbb{K} with finite positive dimension. By a *tridiagonal pair* (or *TD pair*) on V we mean an ordered pair of linear transformations $A : V \rightarrow V$ and $A^* : V \rightarrow V$ that satisfy the following four conditions.

(i) Each of A, A^* is diagonalizable.

(ii) There exists an ordering $\{V_i\}_{i=0}^d$ of the eigenspaces of A such that

$$A^*V_i \subseteq V_{i-1} + V_i + V_{i+1} \quad (0 \leq i \leq d), \quad (1)$$

where $V_{-1} = 0$ and $V_{d+1} = 0$.

(iii) There exists an ordering $\{V_i^*\}_{i=0}^\delta$ of the eigenspaces of A^* such that

$$AV_i^* \subseteq V_{i-1}^* + V_i^* + V_{i+1}^* \quad (0 \leq i \leq \delta), \quad (2)$$

where $V_{-1}^* = 0$ and $V_{\delta+1}^* = 0$.

(iv) There does not exist a subspace W of V such that $AW \subseteq W$, $A^*W \subseteq W$, $W \neq 0$, $W \neq V$.

We say the pair A, A^* is *over* \mathbb{K} .

Note 1.2. According to a common notational convention A^* denotes the conjugate-transpose of A . We are not using this convention. In a TD pair A, A^* the linear transformations A and A^* are arbitrary subject to (i)–(iv) above.

Referring to the TD pair in Definition 1.1, by [9, Lemma 4.5] the scalars d and δ are equal. We call this common value the *diameter* of A, A^* . To avoid trivialities, throughout this paper we assume that the diameter is at least one.

TD pairs first arose in the study of Q -polynomial distance-regular graphs and provided a way to study the irreducible modules of the Terwilliger algebra associated with such a graph. Since their introduction, TD pairs have been found to appear naturally in a variety of other contexts including representation theory [1, 7, 11, 10, 12, 14, 15, 25], orthogonal polynomials [24, 23], partially ordered sets [22], statistical mechanical models [3, 6, 19], and other areas of physics [16, 18]. As a result, TD pairs have become an area of interest in their own right. Among the above papers on representation theory, there are several works that connect TD pairs to quantum groups [1, 5, 7, 11, 12]. These papers consider certain special classes of TD pairs. We call particular attention to [5], in which the present author describes a new relationship between TD pairs in the q -Racah class and quantum groups. The present paper builds off of this work.

In the present paper, we give a new relationship between the maps $\Delta, \psi : V \rightarrow V$ introduced in [4], as well as describe a new decomposition of the underlying vector space that, in some sense, lies between the first and second split decompositions associated with a TD pair. In order to motivate our results, we now recall some basic facts concerning TD pairs. For the rest of this section, let A, A^* denote a TD pair on V , as in Definition 1.1. Fix an ordering $\{V_i\}_{i=0}^d$ (resp. $\{V_i^*\}_{i=0}^d$) of the eigenspaces of A (resp. A^*) which satisfies (1) (resp. (2)). For $0 \leq i \leq d$ let θ_i (resp. θ_i^*) denote the eigenvalue of A (resp. A^*) corresponding to V_i (resp. V_i^*). By [9, Theorem 11.1] the ratios

$$\frac{\theta_{i-2} - \theta_{i+1}}{\theta_{i-1} - \theta_i}, \quad \frac{\theta_{i-2}^* - \theta_{i+1}^*}{\theta_{i-1}^* - \theta_i^*}$$

are equal and independent of i for $2 \leq i \leq d-1$. This gives two recurrence relations, whose solutions can be written in closed form. There are several cases [9, Theorem 11.2]. The most general case is called the q -Racah case [12, Section 1]. We will discuss this case shortly.

We now recall the split decompositions of V [9]. For $0 \leq i \leq d$ define

$$\begin{aligned} U_i &= (V_0^* + V_1^* + \cdots + V_i^*) \cap (V_i + V_{i+1} + \cdots + V_d), \\ U_i^\Downarrow &= (V_0^* + V_1^* + \cdots + V_i^*) \cap (V_0 + V_1 + \cdots + V_{d-i}). \end{aligned}$$

By [9, Theorem 4.6], both the sums $V = \sum_{i=0}^d U_i$ and $V = \sum_{i=0}^d U_i^\Downarrow$ are direct. We call $\{U_i\}_{i=0}^d$ (resp. $\{U_i^\Downarrow\}_{i=0}^d$) the first split decomposition (resp. second split decomposition) of V . In [9], the authors showed that A, A^* act on the first and second split decomposition in a particularly attractive way. This will be described in more detail in Section 3.

We now describe the q -Racah case. We say that the TD pair A, A^* has q -Racah type whenever there exist nonzero scalars $q, a, b \in \mathbb{K}$ such that $q^4 \neq 1$ and

$$\theta_i = aq^{d-2i} + a^{-1}q^{2i-d}, \quad \theta_i^* = bq^{d-2i} + b^{-1}q^{2i-d}$$

for $0 \leq i \leq d$. For the rest of this section assume that A, A^* has q -Racah type.

We recall the maps K and B [13, Section 1.1]. Let $K : V \rightarrow V$ denote the linear transformation such that for $0 \leq i \leq d$, U_i is an eigenspace of K with eigenvalue q^{d-2i} . Let $B : V \rightarrow V$ denote the linear transformation such that for $0 \leq i \leq d$, U_i^\Downarrow is an eigenspace of B with eigenvalue q^{d-2i} . The relationship between K and B is discussed in considerable detail in [5].

We now bring in the linear transformation $\Psi : V \rightarrow V$ [4, Lemma 11.1]. As in [5], we work with the normalization $\psi = (q - q^{-1})(q^d - q^{-d})\Psi$. A key feature of ψ is that by [4, Lemma 11.2, Corollary 15.3],

$$\psi U_i \subseteq U_{i-1}, \quad \psi U_i^\Downarrow \subseteq U_{i-1}^\Downarrow$$

for $1 \leq i \leq d$ and both $\psi U_0 = 0$ and $\psi U_0^\Downarrow = 0$. In [5], it is shown how ψ is related to several maps, including the maps K, B , as well as the map Δ which we now recall. By [4, Lemma

9.5], there exists a unique linear transformation $\Delta : V \rightarrow V$ such that

$$\begin{aligned}\Delta U_i &\subseteq U_i^\Downarrow & (0 \leq i \leq d), \\ (\Delta - I)U_i &\subseteq U_0 + U_1 + \cdots + U_{i-1} & (0 \leq i \leq d).\end{aligned}$$

In [4, Theorem 17.1], the present author showed that both

$$\Delta = \sum_{i=0}^d \left(\prod_{j=1}^i \frac{aq^{j-1} - a^{-1}q^{1-j}}{q^j - q^{-j}} \right) \psi^i, \quad \Delta^{-1} = \sum_{i=0}^d \left(\prod_{j=1}^i \frac{a^{-1}q^{j-1} - aq^{1-j}}{q^j - q^{-j}} \right) \psi^i.$$

The primary goal of this paper is to provide factorizations of these power series in ψ and to investigate the consequences of these factorizations. We accomplish this goal using a linear transformation $\mathcal{M} : V \rightarrow V$ given by

$$\mathcal{M} = \frac{aK - a^{-1}B}{a - a^{-1}}.$$

By construction, $\mathcal{M}^\Downarrow = \mathcal{M}$. One can quickly check that \mathcal{M} is invertible. We show that the map \mathcal{M} is equal to each of

$$(I - a^{-1}q\psi)^{-1}K, \quad K(I - a^{-1}q^{-1}\psi)^{-1}, \quad (I - aq\psi)^{-1}B, \quad B(I - aq^{-1}\psi)^{-1}.$$

We give a number of different relations involving the maps \mathcal{M}, K, B, ψ , the most significant of which are the following:

$$\begin{aligned}K \exp_q \left(\frac{a^{-1}}{q - q^{-1}} \psi \right) &= \exp_q \left(\frac{a^{-1}}{q - q^{-1}} \psi \right) \mathcal{M}, \\ B \exp_q \left(\frac{a}{q - q^{-1}} \psi \right) &= \exp_q \left(\frac{a}{q - q^{-1}} \psi \right) \mathcal{M}.\end{aligned}$$

Using these equations, we obtain our main result which is that both

$$\begin{aligned}\Delta &= \exp_q \left(\frac{a}{q - q^{-1}} \psi \right) \exp_{q^{-1}} \left(-\frac{a^{-1}}{q - q^{-1}} \psi \right), \\ \Delta^{-1} &= \exp_q \left(\frac{a^{-1}}{q - q^{-1}} \psi \right) \exp_{q^{-1}} \left(-\frac{a}{q - q^{-1}} \psi \right).\end{aligned}$$

Due to its important role in the factorization of Δ , we explore the map \mathcal{M} further. We show that \mathcal{M} is diagonalizable with eigenvalues $q^d, q^{d-2}, q^{d-4}, \dots, q^{-d}$. For $0 \leq i \leq d$, let W_i denote the eigenspace of \mathcal{M} corresponding to the eigenvalue q^{d-2i} . We show that for $0 \leq i \leq d$,

$$\begin{aligned}U_i &= \exp_q \left(\frac{a^{-1}}{q - q^{-1}} \psi \right) W_i, & U_i^\Downarrow &= \exp_q \left(\frac{a}{q - q^{-1}} \psi \right) W_i, \\ W_i &= \exp_{q^{-1}} \left(-\frac{a^{-1}}{q - q^{-1}} \psi \right) U_i, & W_i &= \exp_{q^{-1}} \left(-\frac{a}{q - q^{-1}} \psi \right) U_i^\Downarrow.\end{aligned}$$

In light of this result, we interpret the decomposition $\{W_i\}_{i=0}^d$ as a sort of halfway point between the first and second split decompositions. We explore this decomposition further and give the actions of $\psi, K, B, \Delta, A, A^*$ on $\{W_i\}_{i=0}^d$. We then give the actions of $\mathcal{M}^{\pm 1}$ on $\{U_i\}_{i=0}^d, \{U_i^\downarrow\}_{i=0}^d, \{V_i\}_{i=0}^d, \{V_i^*\}_{i=0}^d$. We conclude the paper with a discussion of the special case when A, A^* is a Leonard pair.

The present paper is organized as follows. In Section 2 we discuss some preliminary facts concerning TD pairs and TD systems. In Sections 3 we discuss the split decompositions of V as well as the maps K and B . In Section 4 we discuss the map ψ . In Section 5 we recall the map Δ and give Δ as a power series in ψ . In Section 6 we introduce the map \mathcal{M} and describe its relationship with A, K, B, ψ . In Section 7 we express Δ as a product of two linear transformations; one is a q -exponential in ψ and the other is a q^{-1} -exponential in ψ . In Section 8 we describe the eigenvalues and eigenspaces of \mathcal{M} and discuss how the eigenspace decomposition of \mathcal{M} is related to the first and second split decompositions. In Section 9 we discuss the actions of $\psi, K, B, \Delta, A, A^*$ on the eigenspace decomposition of \mathcal{M} . In Section 10 we describe the action of \mathcal{M} on the first and second split decompositions of V , as well as on the eigenspace decompositions of A, A^* . In Section 11 we consider the case when A, A^* is a Leonard pair.

2 Preliminaries

When working with a tridiagonal pair, it is useful to consider a closely related object called a tridiagonal system. In order to define this object, we first recall some facts from elementary linear algebra [9, Section 2].

We use the following conventions. When we discuss an algebra, we mean a unital associative algebra. When we discuss a subalgebra, we assume that it has the same unit as the parent algebra.

Let V denote a vector space over \mathbb{K} with finite positive dimension. By a *decomposition* of V , we mean a sequence of nonzero subspaces whose direct sum is V . Let $\text{End}(V)$ denote the \mathbb{K} -algebra consisting of all linear transformations from V to V . Let A denote a diagonalizable element in $\text{End}(V)$. Let $\{V_i\}_{i=0}^d$ denote an ordering of the eigenspaces of A . For $0 \leq i \leq d$ let θ_i be the eigenvalue of A corresponding to V_i . Define $E_i \in \text{End}(V)$ by $(E_i - I)V_i = 0$ and $E_i V_j = 0$ if $j \neq i$ ($0 \leq j \leq d$). In other words, E_i is the projection map from V onto V_i . We refer to E_i as the *primitive idempotent* of A associated with θ_i . By elementary linear algebra, (i) $AE_i = E_i A = \theta_i E_i$ ($0 \leq i \leq d$); (ii) $E_i E_j = \delta_{ij} E_i$ ($0 \leq i, j \leq d$); (iii) $V_i = E_i V$ ($0 \leq i \leq d$); (iv) $I = \sum_{i=0}^d E_i$. Moreover

$$E_i = \prod_{\substack{0 \leq j \leq d \\ j \neq i}} \frac{A - \theta_j I}{\theta_i - \theta_j} \quad (0 \leq i \leq d).$$

Let M denote the subalgebra of $\text{End}(V)$ generated by A . Note that each of $\{A^i\}_{i=0}^d, \{E_i\}_{i=0}^d$ is a basis for the \mathbb{K} -vector space M .

Let A, A^* denote a TD pair on V . An ordering of the eigenspaces of A (resp. A^*) is said to be *standard* whenever it satisfies (1) (resp. (2)). Let $\{V_i\}_{i=0}^d$ denote a standard ordering of the eigenspaces of A . By [9, Lemma 2.4], the ordering $\{V_{d-i}\}_{i=0}^d$ is standard and no further ordering of the eigenspaces of A is standard. A similar result holds for the eigenspaces of A^* . An ordering of the primitive idempotents of A (resp. A^*) is said to be *standard* whenever the corresponding ordering of the eigenspaces of A (resp. A^*) is standard.

Definition 2.1. [17, Definition 2.1]. Let V denote a vector space over \mathbb{K} with finite positive dimension. By a *tridiagonal system* (or *TD system*) on V , we mean a sequence

$$\Phi = (A; \{E_i\}_{i=0}^d; A^*; \{E_i^*\}_{i=0}^d)$$

that satisfies (i)–(iii) below.

- (i) A, A^* is a tridiagonal pair on V .
- (ii) $\{E_i\}_{i=0}^d$ is a standard ordering of the primitive idempotents of A .
- (iii) $\{E_i^*\}_{i=0}^d$ is a standard ordering of the primitive idempotents of A^* .

We call d the *diameter* of Φ , and say Φ is *over* \mathbb{K} . For notational convenience, set $E_{-1} = 0$, $E_{d+1} = 0$, $E_{-1}^* = 0$, $E_{d+1}^* = 0$.

In Definition 2.1 we do not assume that the primitive idempotents $\{E_i\}_{i=0}^d, \{E_i^*\}_{i=0}^d$ all have rank 1. A TD system for which each of these primitive idempotents has rank 1 is called a Leonard system [20]. The Leonard systems are classified up to isomorphism [20, Theorem 1.9].

For the rest of this paper, fix a TD system Φ on V as in Definition 2.1. Our TD system Φ can be modified in a number of ways to get a new TD system [9, Section 3]. For example, the sequence

$$\Phi^\Downarrow = (A; \{E_{d-i}\}_{i=0}^d; A^*; \{E_i^*\}_{i=0}^d)$$

is a TD system on V . Following [9, Section 3], we call Φ^\Downarrow the *second inversion* of Φ . When discussing Φ^\Downarrow , we use the following notational convention. For any object f associated with Φ , let f^\Downarrow denote the corresponding object associated with Φ^\Downarrow .

Definition 2.2. For $0 \leq i \leq d$ let θ_i (resp. θ_i^*) denote the eigenvalue of A (resp. A^*) associated with E_i (resp. E_i^*). We refer to $\{\theta_i\}_{i=0}^d$ (resp. $\{\theta_i^*\}_{i=0}^d$) as the *eigenvalue sequence* (resp. *dual eigenvalue sequence*) of Φ .

By construction $\{\theta_i\}_{i=0}^d$ are mutually distinct and $\{\theta_i^*\}_{i=0}^d$ are mutually distinct. By [9, Theorem 11.1], the scalars

$$\frac{\theta_{i-2} - \theta_{i+1}}{\theta_{i-1} - \theta_i}, \quad \frac{\theta_{i-2}^* - \theta_{i+1}^*}{\theta_{i-1}^* - \theta_i^*}$$

are equal and independent of i for $2 \leq i \leq d-1$. For this restriction, the solutions have been found in closed form [9, Theorem 11.2]. The most general solution is called q -Racah [12, Section 1]. This solution is described as follows.

Definition 2.3. Let Φ denote a TD system on V as in Definition 2.1. We say that Φ has *q-Racah type* whenever there exist nonzero scalars $q, a, b \in \mathbb{K}$ such that $q^4 \neq 1$ and

$$\theta_i = aq^{d-2i} + a^{-1}q^{2i-d}, \quad \theta_i^* = bq^{d-2i} + b^{-1}q^{2i-d} \quad (3)$$

for $0 \leq i \leq d$.

Note 2.4. Referring to Definition 2.3, the scalars q, a, b are not uniquely defined by Φ . If q, a, b is one solution, then their inverses give another solution.

For the rest of the paper, we make the following assumption.

Assumption 2.5. We assume that our TD system Φ has *q-Racah type*. We fix q, a, b as in Definition 2.3.

Lemma 2.6. [5, Lemma 2.4]. *With reference to Assumption 2.5, the following hold.*

- (i) Neither of a^2, b^2 is among $q^{2d-2}, q^{2d-4}, \dots, q^{2-2d}$.
- (ii) $q^{2i} \neq 1$ for $1 \leq i \leq d$.

Proof: The result follows from the comment below Definition 2.2. □

3 The first and second split decomposition of V

Recall the TD system Φ from Assumption 2.5. In this section we consider two decompositions of V associated with Φ , called the first and second split decomposition.

For $0 \leq i \leq d$ define

$$U_i = (E_0^*V + E_1^*V + \dots + E_i^*V) \cap (E_iV + E_{i+1}V + \dots + E_dV).$$

For notational convenience, define $U_{-1} = 0$ and $U_{d+1} = 0$. Note that for $0 \leq i \leq d$,

$$U_i^\Downarrow = (E_0^*V + E_1^*V + \dots + E_i^*V) \cap (E_0V + E_1V + \dots + E_{d-i}V).$$

By [9, Theorem 4.6], the sequence $\{U_i\}_{i=0}^d$ (resp. $\{U_i^\Downarrow\}_{i=0}^d$) is a decomposition of V . Following [9], we refer to $\{U_i\}_{i=0}^d$ (resp. $\{U_i^\Downarrow\}_{i=0}^d$) as the *first split decomposition* (resp. *second split decomposition*) of V with respect to Φ . By [9, Corollary 5.7], for $0 \leq i \leq d$ the dimensions of $E_iV, E_i^*V, U_i, U_i^\Downarrow$ coincide; we denote the common dimension by ρ_i . By [9, Theorem 4.6],

$$E_iV + E_{i+1}V + \dots + E_dV = U_i + U_{i+1} + \dots + U_d, \quad (4)$$

$$E_0V + E_1V + \dots + E_iV = U_{d-i}^\Downarrow + U_{d-i+1}^\Downarrow + \dots + U_d^\Downarrow, \quad (5)$$

$$E_0^*V + E_1^*V + \dots + E_i^*V = U_0 + U_1 + \dots + U_i = U_0^\Downarrow + U_1^\Downarrow + \dots + U_i^\Downarrow. \quad (6)$$

By [9, Theorem 4.6], A and A^* act on the first split decomposition in the following way:

$$\begin{aligned} (A - \theta_i I)U_i &\subseteq U_{i+1} & (0 \leq i \leq d-1), & & (A - \theta_d I)U_d &= 0, \\ (A^* - \theta_i^* I)U_i &\subseteq U_{i-1} & (1 \leq i \leq d), & & (A^* - \theta_0^* I)U_0 &= 0. \end{aligned}$$

By [9, Theorem 4.6], A and A^* act on the second split decomposition in the following way:

$$\begin{aligned} (A - \theta_{d-i}I)U_i^\downarrow &\subseteq U_{i+1}^\downarrow & (0 \leq i \leq d-1), & (A - \theta_0I)U_d^\downarrow = 0, \\ (A^* - \theta_i^*I)U_i^\downarrow &\subseteq U_{i-1}^\downarrow & (1 \leq i \leq d), & (A^* - \theta_0^*I)U_0^\downarrow = 0. \end{aligned}$$

Definition 3.1. [5, Definitions 3.1 and 3.2]. Define $K, B \in \text{End}(V)$ such that for $0 \leq i \leq d$, U_i (resp. U_i^\downarrow) is the eigenspace of K (resp. B) with eigenvalue q^{d-2i} . In other words,

$$(K - q^{d-2i}I)U_i = 0, \quad (B - q^{d-2i}I)U_i^\downarrow = 0 \quad (0 \leq i \leq d). \quad (7)$$

Observe that $B = K^\downarrow$.

By construction each of K, B is invertible and diagonalizable on V .

We now describe how K and B act on the eigenspaces of the other one.

Lemma 3.2. [5, Lemma 3.3]. *For $0 \leq i \leq d$,*

$$(B - q^{d-2i}I)U_i \subseteq U_0 + U_1 + \cdots + U_{i-1}, \quad (8)$$

$$(K - q^{d-2i}I)U_i^\downarrow \subseteq U_0^\downarrow + U_1^\downarrow + \cdots + U_{i-1}^\downarrow. \quad (9)$$

Next we describe how A, K, B are related.

Lemma 3.3. [13, Section 1.1]. *Both*

$$\frac{qKA - q^{-1}AK}{q - q^{-1}} = aK^2 + a^{-1}I, \quad \frac{qBA - q^{-1}AB}{q - q^{-1}} = a^{-1}B^2 + aI. \quad (10)$$

Lemma 3.4. [5, Theorem 9.9]. *We have*

$$aK^2 - \frac{a^{-1}q - aq^{-1}}{q - q^{-1}} KB - \frac{aq - a^{-1}q^{-1}}{q - q^{-1}} BK + a^{-1}B^2 = 0. \quad (11)$$

4 The linear transformation ψ

We continue to discuss the situation of Assumption 2.5. In [4, Section 11] we introduced an element $\Psi \in \text{End}(V)$. In [5] we used the normalization $\psi = (q - q^{-1})(q^d - q^{-d})\Psi$. In [5, Theorem 9.8], we showed that ψ is equal to some rational expressions involving K, B . We now recall this result. We start with a comment.

Lemma 4.1. [5, Lemma 9.7]. *Each of the following is invertible:*

$$aI - a^{-1}BK^{-1}, \quad a^{-1}I - aKB^{-1}, \quad (12)$$

$$aI - a^{-1}K^{-1}B, \quad a^{-1}I - aB^{-1}K. \quad (13)$$

Lemma 4.2. [5, Theorem 9.8]. *The following four expressions coincide:*

$$\frac{I - BK^{-1}}{q(aI - a^{-1}BK^{-1})}, \quad \frac{I - KB^{-1}}{q(a^{-1}I - aKB^{-1})}, \quad (14)$$

$$\frac{q(I - K^{-1}B)}{aI - a^{-1}K^{-1}B}, \quad \frac{q(I - B^{-1}K)}{a^{-1}I - aB^{-1}K}. \quad (15)$$

In (14), (15) the denominators are invertible by Lemma 4.1.

Definition 4.3. Define $\psi \in \text{End}(V)$ to be the common value of the four expressions in Lemma 4.2.

We now recall some facts concerning ψ .

Lemma 4.4. [5, Lemma 5.4]. *Both*

$$K\psi = q^2\psi K, \quad B\psi = q^2\psi B. \quad (16)$$

Lemma 4.5. [4, Lemma 11.2, Corollary 15.3]. *We have*

$$\psi U_i \subseteq U_{i-1}, \quad \psi U_i^\Downarrow \subseteq U_{i-1}^\Downarrow \quad (1 \leq i \leq d) \quad (17)$$

and also $\psi U_0 = 0$ and $\psi U_0^\Downarrow = 0$. Moreover $\psi^{d+1} = 0$.

In Lemma 4.2 we obtained ψ as a rational expression in BK^{-1} or $K^{-1}B$. Next we solve for BK^{-1} and $K^{-1}B$ as a rational function in ψ . In order to state the answer, we will need the following result.

Lemma 4.6. [5, Lemma 9.2]. *Each of the following is invertible:*

$$I - aq\psi, \quad I - a^{-1}q\psi, \quad I - aq^{-1}\psi, \quad I - a^{-1}q^{-1}\psi. \quad (18)$$

Their inverses are as follows:

$$(I - aq\psi)^{-1} = \sum_{i=0}^d a^i q^i \psi^i, \quad (I - a^{-1}q\psi)^{-1} = \sum_{i=0}^d a^{-i} q^i \psi^i, \quad (19)$$

$$(I - aq^{-1}\psi)^{-1} = \sum_{i=0}^d a^i q^{-i} \psi^i, \quad (I - a^{-1}q^{-1}\psi)^{-1} = \sum_{i=0}^d a^{-i} q^{-i} \psi^i. \quad (20)$$

The next result is an immediate consequence of Lemma 4.2, Definition 4.3, and Lemma 4.6.

Theorem 4.7. [5, Theorem 9.4]. *The following hold:*

$$BK^{-1} = \frac{I - aq\psi}{I - a^{-1}q\psi}, \quad KB^{-1} = \frac{I - a^{-1}q\psi}{I - aq\psi}, \quad (21)$$

$$K^{-1}B = \frac{I - aq^{-1}\psi}{I - a^{-1}q^{-1}\psi}, \quad B^{-1}K = \frac{I - a^{-1}q^{-1}\psi}{I - aq^{-1}\psi}. \quad (22)$$

In (21), (22) the denominators are invertible by Lemma 4.6.

Lemma 4.8. [5, Equation (22)]. *We have*

$$\frac{\psi A - A\psi}{q - q^{-1}} = (I - aq\psi) K - (I - a^{-1}q^{-1}\psi) K^{-1}. \quad (23)$$

Proof: This result is a reformulation of [5, Equation (22)] using [5, Equation (14)]. \square

5 The linear transformation Δ

We continue to discuss the situation of Assumption 2.5. In [4, Section 9] we introduced an invertible element $\Delta \in \text{End}(V)$. In [4] we showed that Δ, ψ commute and in fact both Δ, Δ^{-1} are power series in ψ . These power series will be the central focus of this paper. We will show that each of those power series factors as a product of two power series, each of which is a quantum exponential in ψ .

Lemma 5.1. [4, Lemma 9.5]. *There exists a unique $\Delta \in \text{End}(V)$ such that*

$$\Delta U_i \subseteq U_i^\natural \quad (0 \leq i \leq d), \quad (24)$$

$$(\Delta - I)U_i \subseteq U_0 + U_1 + \cdots + U_{i-1} \quad (0 \leq i \leq d). \quad (25)$$

Lemma 5.2. [4, Lemmas 9.3 and 9.6]. *The map Δ is invertible. Moreover $\Delta^{-1} = \Delta^\natural$ and*

$$(\Delta^{-1} - I)U_i \subseteq U_0 + U_1 + \cdots + U_{i-1} \quad (0 \leq i \leq d). \quad (26)$$

Lemma 5.3. *The map $\Delta - I$ is nilpotent. Moreover $\Delta K = B\Delta$.*

Proof: The first assertion follows from (25). The last assertion follows from (24) and Definition 3.1. \square

The map Δ is characterized as follows.

Lemma 5.4. [4, Lemma 9.8]. *The map Δ is the unique element of $\text{End}(V)$ such that*

$$(\Delta - I)E_i^*V \subseteq E_0^*V + E_1^*V + \cdots + E_{i-1}^*V \quad (0 \leq i \leq d), \quad (27)$$

$$\Delta(E_iV + E_{i+1}V + \cdots + E_dV) = E_0V + E_1V + \cdots + E_{d-i}V \quad (0 \leq i \leq d). \quad (28)$$

Theorem 5.5. [4, Theorem 17.1]. *Both*

$$\Delta = \sum_{i=0}^d \left(\prod_{j=1}^i \frac{aq^{j-1} - a^{-1}q^{1-j}}{q^j - q^{-j}} \right) \psi^i, \quad (29)$$

$$\Delta^{-1} = \sum_{i=0}^d \left(\prod_{j=1}^i \frac{a^{-1}q^{j-1} - aq^{1-j}}{q^j - q^{-j}} \right) \psi^i. \quad (30)$$

In (29) and (30), the elements Δ, Δ^{-1} are expressed as a power series in ψ . In the present paper, we factor these power series and interpret the results. This interpretation will involve a linear transformation \mathcal{M} . We introduce \mathcal{M} in the next section.

6 The linear transformation \mathcal{M}

We continue to discuss the situation of Assumption 2.5. In this section we introduce an element $\mathcal{M} \in \text{End}(V)$. We explain how \mathcal{M} is related to K, B, ψ, A .

Definition 6.1. Define $\mathcal{M} \in \text{End}(V)$ by

$$\mathcal{M} = \frac{aK - a^{-1}B}{a - a^{-1}}. \quad (31)$$

By construction, $\mathcal{M}^\sharp = \mathcal{M}$. Evaluating (31) using Lemma 4.1, we see that \mathcal{M} is invertible.

Lemma 6.2. *The map \mathcal{M} is equal to each of:*

$$(I - a^{-1}q\psi)^{-1}K, \quad K(I - a^{-1}q^{-1}\psi)^{-1}, \quad (I - aq\psi)^{-1}B, \quad B(I - aq^{-1}\psi)^{-1}.$$

Proof: We first show that $\mathcal{M} = (I - a^{-1}q\psi)^{-1}K$. By Definition 6.1,

$$(a - a^{-1})\mathcal{M}K^{-1} = aI - a^{-1}BK^{-1}.$$

The result follows from this fact along with the equation on the left in (21).

The remaining assertions follow from Theorem 4.7. \square

Lemma 6.2 can be reformulated as follows.

Lemma 6.3. *We have*

$$K = (I - a^{-1}q\psi)\mathcal{M}, \quad K = \mathcal{M}(I - a^{-1}q^{-1}\psi), \quad (32)$$

$$B = (I - aq\psi)\mathcal{M}, \quad B = \mathcal{M}(I - aq^{-1}\psi). \quad (33)$$

For later use, we give several descriptions of $\mathcal{M}^{\pm 1}$.

Lemma 6.4. *The map \mathcal{M}^{-1} is equal to each of:*

$$K^{-1}(I - a^{-1}q\psi), \quad (I - a^{-1}q^{-1}\psi)K^{-1}, \quad B^{-1}(I - aq\psi), \quad (I - aq^{-1}\psi)B^{-1}.$$

Proof: Immediate from Lemma 6.2. \square

Lemma 6.5. *The map \mathcal{M} is equal to each of:*

$$K \sum_{n=0}^d a^{-n}q^{-n}\psi^n, \quad \sum_{n=0}^d a^{-n}q^n\psi^n K, \quad B \sum_{n=0}^d a^nq^{-n}\psi^n, \quad \sum_{n=0}^d a^nq^n\psi^n B \quad (34)$$

Proof: Use Lemma 4.6 and Lemma 6.2. \square

We now give some attractive equations that show how \mathcal{M} is related to ψ, K, B, A .

Lemma 6.6. *We have*

$$\mathcal{M}\psi = q^2\psi\mathcal{M}. \quad (35)$$

Proof: Use Lemma 4.4 and Definition 6.1. \square

Lemma 6.7. *We have*

$$\frac{q\mathcal{M}^{-1}K - q^{-1}K\mathcal{M}^{-1}}{q - q^{-1}} = I, \quad \frac{q\mathcal{M}^{-1}B - q^{-1}B\mathcal{M}^{-1}}{q - q^{-1}} = I. \quad (36)$$

Proof: Use Lemma 6.4. □

Lemma 6.8. *We have*

$$\frac{qA\mathcal{M}^{-1} - q^{-1}\mathcal{M}^{-1}A}{q - q^{-1}} = (a + a^{-1})I - (q + q^{-1})\psi. \quad (37)$$

Proof: Use Lemma 3.3, Lemma 4.4, Lemma 4.8, and Lemma 6.4. □

Lemma 6.9. *We have*

$$\mathcal{M}^{-2}A - (q^2 + q^{-2})\mathcal{M}^{-1}A\mathcal{M}^{-1} + A\mathcal{M}^{-2} = -(q - q^{-1})^2(a + a^{-1})\mathcal{M}^{-1}. \quad (38)$$

Proof: Use Lemma 6.6 and Lemma 6.8. □

7 A factorization of Δ

We continue to discuss the situation of Assumption 2.5. We now bring in the q -exponential function [8]. In [4, Theorem 17.1] we expressed Δ as a power series in ψ . In this section we strengthen this result in the following way. We express Δ as a product of two linear transformations; one is a q -exponential in ψ and the other is a q^{-1} -exponential in ψ .

For an integer n , define

$$[n]_q = \frac{q^n - q^{-n}}{q - q^{-1}} \quad (39)$$

and for $n \geq 0$, define

$$[n]_q! = [n]_q[n-1]_q \cdots [1]_q. \quad (40)$$

We interpret $[0]_q! = 1$.

We now recall the q -exponential function [8]. For a nilpotent $T \in \text{End}(V)$,

$$\exp_q(T) = \sum_{n=0}^{\infty} \frac{q^{\binom{n}{2}}}{[n]_q!} T^n. \quad (41)$$

The map $\exp_q(T)$ is invertible. Its inverse is given by

$$\exp_{q^{-1}}(-T) = \sum_{n=0}^{\infty} \frac{(-1)^n q^{-\binom{n}{2}}}{[n]_q!} T^n. \quad (42)$$

Using (41) we obtain

$$(I - (q^2 - 1)T) \exp_q(q^2T) = \exp_q(T). \quad (43)$$

For $S \in \text{End}(V)$ such that $ST = q^2TS$, we have

$$S \exp_q(T) S^{-1} = \exp_q(STS^{-1}) = \exp_q(q^2T).$$

Consequently

$$S \exp_q(T) = \exp_q(q^2T)S. \quad (44)$$

Combining (43) and (44),

$$(I - (q^2 - 1)T)S \exp_q(T) = \exp_q(T)S. \quad (45)$$

We return our attention to K, B, ψ, \mathcal{M} .

Proposition 7.1. *Both*

$$K \exp_q\left(\frac{a^{-1}}{q - q^{-1}}\psi\right) = \exp_q\left(\frac{a^{-1}}{q - q^{-1}}\psi\right) \mathcal{M}, \quad (46)$$

$$B \exp_q\left(\frac{a}{q - q^{-1}}\psi\right) = \exp_q\left(\frac{a}{q - q^{-1}}\psi\right) \mathcal{M}. \quad (47)$$

Proof: Recall from Lemma 6.6 that $\mathcal{M}\psi = q^2\psi\mathcal{M}$. We first obtain (46). To do this, in (45) take $S = \mathcal{M}$ and $T = \frac{a^{-1}}{q - q^{-1}}\psi$. Evaluate the result using the equation $\mathcal{M} = (I - a^{-1}q\psi)^{-1}K$ from Lemma 6.2.

Next we obtain (47). To do this, in (45) take $S = \mathcal{M}$ and $T = \frac{a}{q - q^{-1}}\psi$. Evaluate the result using the equation $\mathcal{M} = (I - aq\psi)^{-1}B$ from Lemma 6.2. \square

The following is our main result.

Theorem 7.2. *Both*

$$\Delta = \exp_q\left(\frac{a}{q - q^{-1}}\psi\right) \exp_{q^{-1}}\left(-\frac{a^{-1}}{q - q^{-1}}\psi\right), \quad (48)$$

$$\Delta^{-1} = \exp_q\left(\frac{a^{-1}}{q - q^{-1}}\psi\right) \exp_{q^{-1}}\left(-\frac{a}{q - q^{-1}}\psi\right). \quad (49)$$

Proof: We first show (48). Let $\tilde{\Delta}$ denote the expression on the right in (48). Combining (46) and (47), we see that $\tilde{\Delta}K = B\tilde{\Delta}$. Therefore $\tilde{\Delta}U_i = U_i^\Downarrow$ for $0 \leq i \leq d$. Observe that $\tilde{\Delta} - I$ is a polynomial in ψ with zero constant term. By Lemma 4.5, $(\tilde{\Delta} - I)U_i \subseteq U_0 + U_1 + \cdots + U_{i-1}$ for $0 \leq i \leq d$. By Lemma 5.1, $\tilde{\Delta} = \Delta$.

To obtain (49) from (48), use (42). \square

Corollary 7.3. *We have*

$$\begin{aligned} \exp_q \left(\frac{a}{q - q^{-1}} \psi \right) \exp_{q^{-1}} \left(-\frac{a^{-1}}{q - q^{-1}} \psi \right) &= \sum_{i=0}^d \left(\prod_{j=1}^i \frac{aq^{j-1} - a^{-1}q^{1-j}}{q^j - q^{-j}} \right) \psi^i, \\ \exp_q \left(\frac{a^{-1}}{q - q^{-1}} \psi \right) \exp_{q^{-1}} \left(-\frac{a}{q - q^{-1}} \psi \right) &= \sum_{i=0}^d \left(\prod_{j=1}^i \frac{a^{-1}q^{j-1} - aq^{1-j}}{q^j - q^{-j}} \right) \psi^i. \end{aligned}$$

Proof: Combine Theorem 5.5 and Theorem 7.2. The equations can also be obtained directly by expanding their left-hand sides using (41) and (42), and evaluating the results using the q -binomial theorem [2, Theorem 10.2.1]. \square

8 The eigenvalues and eigenspaces of \mathcal{M}

We continue to discuss the situation of Assumption 2.5. In Section 6 we introduced the linear transformation \mathcal{M} . Proposition 7.1 indicates the role of \mathcal{M} in the factorization of Δ in Theorem 7.2. In this section we show that \mathcal{M} is diagonalizable. We describe the eigenvalues and eigenspaces of \mathcal{M} . We also explain how the eigenspace decomposition for \mathcal{M} is related to the first and second split decompositions.

Lemma 8.1. *The map \mathcal{M} is diagonalizable with eigenvalues $q^d, q^{d-2}, q^{d-4}, \dots, q^{-d}$.*

Proof: Let $E = \exp_q \left(\frac{a^{-1}}{q - q^{-1}} \psi \right)$. By (46), $\mathcal{M} = E^{-1}KE$. By construction K is diagonalizable with eigenvalues $q^d, q^{d-2}, q^{d-4}, \dots, q^{-d}$. The result follows. \square

Definition 8.2. For $0 \leq i \leq d$, let W_i denote the eigenspace of \mathcal{M} corresponding to the eigenvalue q^{d-2i} . Note that $\{W_i\}_{i=0}^d$ is a decomposition of V , and that $W_i^\Downarrow = W_i$ for $0 \leq i \leq d$. For notational convenience, let $W_{-1} = 0$ and $W_{d+1} = 0$.

Proposition 8.3. *For $0 \leq i \leq d$,*

$$U_i = \exp_q \left(\frac{a^{-1}}{q - q^{-1}} \psi \right) W_i, \quad U_i^\Downarrow = \exp_q \left(\frac{a}{q - q^{-1}} \psi \right) W_i, \quad (50)$$

$$W_i = \exp_{q^{-1}} \left(-\frac{a^{-1}}{q - q^{-1}} \psi \right) U_i, \quad W_i = \exp_{q^{-1}} \left(-\frac{a}{q - q^{-1}} \psi \right) U_i^\Downarrow. \quad (51)$$

Proof: Define E as in the proof of Lemma 8.1. We show that $U_i = EW_i$. By (46), $KE = E\mathcal{M}$. Recall that U_i (resp. W_i) is the eigenspace of K (resp. \mathcal{M}) corresponding to the eigenvalue q^{d-2i} . By these comments $U_i = EW_i$.

Define $F = \exp_q \left(\frac{a}{q - q^{-1}} \psi \right)$. We show $U_i^\Downarrow = FW_i$. By (47), $BF = F\mathcal{M}$. Recall that U_i^\Downarrow (resp. W_i) is the eigenspace of B (resp. \mathcal{M}) corresponding to the eigenvalue q^{d-2i} . By these comments $U_i^\Downarrow = FW_i$.

To obtain (51) from (50), use (42). \square

Lemma 8.4. For $0 \leq i \leq d$, the dimension of W_i is ρ_i .

Proof: This follows from Proposition 8.3 and the fact that U_i, U_i^\Downarrow have dimension ρ_i . \square

Recall from (6) that

$$\sum_{h=0}^i E_h^* V = \sum_{h=0}^i U_h = \sum_{h=0}^i U_h^\Downarrow \quad (52)$$

for $0 \leq i \leq d$.

Lemma 8.5. For $0 \leq i \leq d$, the sum $\sum_{h=0}^i W_h$ is equal to the common value of (52).

Proof: Define $W = \sum_{h=0}^i W_h$ and let U denote the common value of (52). We show that $W = U$. By Lemma 4.5 and the equation on the left in (51), $W \subseteq U$. By Lemma 8.4, W and U have the same dimension. Thus $W = U$. \square

9 The actions of $\psi, K, B, \Delta, A, A^*$ on $\{W_i\}_{i=0}^d$

We continue to discuss the situation of Assumption 2.5. Recall the eigenspace decomposition $\{W_i\}_{i=0}^d$ for \mathcal{M} . In this section, we discuss the actions of $\psi, K, B, \Delta, A, A^*$ on $\{W_i\}_{i=0}^d$.

Lemma 9.1. For $0 \leq i \leq d$,

$$\psi W_i \subseteq W_{i-1}. \quad (53)$$

Proof: Use Lemma 6.6. \square

Lemma 9.2. For $0 \leq i \leq d$,

$$(K - q^{d-2i}I)W_i \subseteq W_{i-1}, \quad (B - q^{d-2i}I)W_i \subseteq W_{i-1}. \quad (54)$$

Proof: Use Lemma 6.3 and Lemma 9.1. \square

Lemma 9.3. For $0 \leq i \leq d$,

$$(\Delta - I)W_i \subseteq W_0 + W_1 + \cdots + W_{i-1}, \quad (55)$$

$$(\Delta^{-1} - I)W_i \subseteq W_0 + W_1 + \cdots + W_{i-1}. \quad (56)$$

Proof: To show (55), use (25) and Lemma 8.5.

To show (56), use (26) and Lemma 8.5. \square

Lemma 9.4. For $0 \leq i \leq d$,

$$(A - (a + a^{-1})q^{d-2i}I)W_i \subseteq W_{i-1} + W_{i+1}. \quad (57)$$

Proof: By Lemma 6.9, the expression

$$(\mathcal{M}^{-1} - q^{2i+2-d}I)(\mathcal{M}^{-1} - q^{2i-2-d}I)(A - (a + a^{-1})q^{d-2i}I)$$

vanishes on W_i . Therefore $(\mathcal{M}^{-1} - q^{2i+2-d}I)(\mathcal{M}^{-1} - q^{2i-2-d}I)$ vanishes on $(A - (a + a^{-1})q^{d-2i}I)W_i$. The result follows. \square

Lemma 9.5. For $0 \leq i \leq d$,

$$(A^* - \theta_i^*I)W_i \subseteq W_0 + W_1 + \cdots + W_{i-1}. \quad (58)$$

Proof: Use $(A^* - \theta_i^*I)E_i^*V = 0$ together with (25) and Lemma 8.5. \square

10 The actions of $\mathcal{M}^{\pm 1}$ on $\{U_i\}_{i=0}^d, \{U_i^\downarrow\}_{i=0}^d, \{E_iV\}_{i=0}^d, \{E_i^*V\}_{i=0}^d$

We continue to discuss the situation of Assumption 2.5. In Section 8 we saw how various operators act on the decomposition $\{W_i\}_{i=0}^d$. In this section we investigate the action of \mathcal{M} on the first and second split decompositions of V , as well as on the eigenspace decompositions of A, A^* .

Lemma 10.1. For $0 \leq i \leq d$,

$$(\mathcal{M} - q^{d-2i}I)U_i \subseteq U_0 + U_1 + \cdots + U_{i-1}, \quad (59)$$

$$(\mathcal{M} - q^{d-2i}I)U_i^\downarrow \subseteq U_0^\downarrow + U_1^\downarrow + \cdots + U_{i-1}^\downarrow. \quad (60)$$

Proof: To show (59), use Definition 3.1, Lemma 3.2, and Definition 6.1.

To show (60), use (59) applied to Φ^\downarrow , along with $\mathcal{M}^\downarrow = \mathcal{M}$. \square

Lemma 10.2. For $0 \leq i \leq d$,

$$(\mathcal{M}^{-1} - q^{2i-d}I)U_i \subseteq U_{i-1}, \quad (\mathcal{M}^{-1} - q^{2i-d}I)U_i^\downarrow \subseteq U_{i-1}^\downarrow. \quad (61)$$

Proof: We first show the equation on the left in (61). By Lemma 6.4,

$$\mathcal{M}^{-1} = (I - a^{-1}q^{-1}\psi)K^{-1}. \quad (62)$$

From this and Definition 3.1, it follows that on U_i ,

$$\mathcal{M}^{-1} - q^{2i-d}I = a^{-1}q^{2i-d-1}\psi. \quad (63)$$

The result follows from this along with Lemma 4.5.

The proof of the equation on the right in (61) follows from the equation on the left in (61) applied to Φ^\downarrow , along with the fact that $\mathcal{M}^\downarrow = \mathcal{M}$. \square

Lemma 10.3. For $0 \leq i \leq d$,

$$\mathcal{M}^{-1}E_iV \subseteq E_{i-1}V + E_iV + E_{i+1}V. \quad (64)$$

Proof: We first show that $\mathcal{M}^{-1}E_iV \subseteq \sum_{h=0}^{i+1} E_hV$. Recall from (5) that $E_iV \subseteq \sum_{h=d-i}^d U_h^\downarrow$. By this, Lemma 10.2, and (5), we obtain $\mathcal{M}^{-1}E_iV \subseteq \sum_{h=0}^{i+1} E_hV$.

We now show that $\mathcal{M}^{-1}E_iV \subseteq \sum_{h=i-1}^d E_hV$. Recall from (4) that $E_iV \subseteq \sum_{h=i}^d U_h$. By this, Lemma 10.2, and (4), we obtain $\mathcal{M}^{-1}E_iV \subseteq \sum_{h=i-1}^d E_hV$.

Thus $\mathcal{M}^{-1}E_iV$ is contained in the intersection of $\sum_{h=0}^{i+1} E_hV$ and $\sum_{h=i-1}^d E_hV$, which is $E_{i-1}V + E_iV + E_{i+1}V$. \square

Lemma 10.4. For $0 \leq i \leq d$,

$$\begin{aligned} (\mathcal{M} - q^{d-2i}I)E_i^*V &\subseteq E_0^*V + E_1^*V + \cdots + E_{i-1}^*V, \\ (\mathcal{M}^{-1} - q^{2i-d}I)E_i^*V &\subseteq E_0^*V + E_1^*V + \cdots + E_{i-1}^*V. \end{aligned}$$

Proof: Note that $E_i^*V \subseteq E_0^*V + E_1^*V + \cdots + E_i^*V = W_0 + W_1 + \cdots + W_i$ by Lemma 8.5. The result follows from this fact along with Definition 8.2. \square

11 When Φ is a Leonard system

We continue to discuss the situation of Assumption 2.5. For the rest of the paper we assume $\rho_i = 1$ for $0 \leq i \leq d$. In this case Φ is called a Leonard system.

We use the following notational convention. Let $\{v_i\}_{i=0}^d$ denote a basis for V . The sequence of subspaces $\{\mathbb{K}v_i\}_{i=0}^d$ is a decomposition of V said, to be *induced* by the basis $\{v_i\}_{i=0}^d$.

We display a basis $\{u_i\}_{i=0}^d$ (resp. $\{u_i^\downarrow\}_{i=0}^d$) (resp. $\{w_i\}_{i=0}^d$) that induces the decomposition $\{U_i\}_{i=0}^d$ (resp. $\{U_i^\downarrow\}_{i=0}^d$) (resp. $\{W_i\}_{i=0}^d$). We find the actions of $\psi, K, B, \Delta^{\pm 1}, A$ on these bases. We also display the transition matrices between these bases.

For the rest of this section fix $0 \neq u_0 \in U_0$. Let M denote the subalgebra of $\text{End}(V)$ generated by A . By [21, Lemma 5.1], the map $M \rightarrow V, X \mapsto Xu_0$ is an isomorphism of vector spaces. Consequently, the vectors $\{A^i u_0\}_{i=0}^d$ form a basis for V .

We now define a basis $\{u_i\}_{i=0}^d$ of V that induces $\{U_i\}_{i=0}^d$. For $0 \leq i \leq d$, define

$$u_i = \left(\prod_{j=0}^{i-1} (A - \theta_j I) \right) u_0. \quad (65)$$

Observe that $u_i \neq 0$. By [9, Theorem 4.6], $u_i \in U_i$. So u_i is a basis for U_i . Consequently, $\{u_i\}_{i=0}^d$ is a basis for V that induces $\{U_i\}_{i=0}^d$.

Next we define a basis $\{u_i^\downarrow\}_{i=0}^d$ of V that induces $\{U_i^\downarrow\}_{i=0}^d$. For $0 \leq i \leq d$, define

$$u_i^\downarrow = \left(\prod_{j=0}^{i-1} (A - \theta_{d-j}I) \right) u_0. \quad (66)$$

Observe that $u_i^\downarrow \neq 0$. By Lemma 5.1, $u_i^\downarrow \in U_i^\downarrow$. So u_i^\downarrow is a basis for U_i^\downarrow . Consequently, $\{u_i^\downarrow\}_{i=0}^d$ is a basis for V that induces $\{U_i^\downarrow\}_{i=0}^d$.

Lemma 11.1. For $0 \leq i \leq d$,

$$u_i^\downarrow = \Delta u_i. \quad (67)$$

Proof: By Lemma 5.1, $\Delta U_i = U_i^\downarrow$. So there exists $0 \neq \lambda \in \mathbb{K}$ such that $\Delta u_i = \lambda u_i^\downarrow$. We show that $\lambda = 1$. By [4, Lemma 7.3] and (25), $\Delta u_i - A^i u$ is a linear combination of $\{A^j u\}_{j=0}^{i-1}$. Also, $u_i^\downarrow - A^i u$ is a linear combination of $\{A^j u\}_{j=0}^{i-1}$. The vectors $\{A^j u\}_{j=0}^{i-1}$ are linearly independent. By these comments $\lambda = 1$. \square

We next define a basis $\{w_i\}_{i=0}^d$ of V that induces $\{W_i\}_{i=0}^d$. For $0 \leq i \leq d$, define

$$w_i = \exp_{q^{-1}} \left(-\frac{a^{-1}}{q - q^{-1}} \psi \right) u_i. \quad (68)$$

Since $\{u_i\}_{i=0}^d$ is a basis of V and $\exp_{q^{-1}} \left(-\frac{a^{-1}}{q - q^{-1}} \psi \right)$ is invertible, w_i is a basis for W_i . Consequently, $\{w_i\}_{i=0}^d$ is a basis for V that induces $\{W_i\}_{i=0}^d$.

Lemma 11.2. For $0 \leq i \leq d$,

$$u_i = \exp_q \left(\frac{a^{-1}}{q - q^{-1}} \psi \right) w_i, \quad u_i^\downarrow = \exp_q \left(\frac{a}{q - q^{-1}} \psi \right) w_i, \quad (69)$$

$$w_i = \exp_{q^{-1}} \left(-\frac{a^{-1}}{q - q^{-1}} \psi \right) u_i, \quad w_i = \exp_{q^{-1}} \left(-\frac{a}{q - q^{-1}} \psi \right) u_i^\downarrow. \quad (70)$$

Proof: Use (68) to obtain the equations on the left in (69),(70). To obtain the equations on the right in (69),(70), use Theorem 7.2, Lemma 11.1, and (68). \square

We now describe the actions of $\psi, K, B, \mathcal{M}, \Delta, A$ on the bases $\{u_i\}_{i=0}^d, \{u_i^\downarrow\}_{i=0}^d, \{w_i\}_{i=0}^d$. First we recall a notion from linear algebra. Let $\text{Mat}_{d+1}(\mathbb{K})$ denote the \mathbb{K} -algebra of $(d+1) \times (d+1)$ matrices that have all entries in \mathbb{K} . We index the rows and columns by $0, 1, \dots, d$. Let $\{v_i\}_{i=0}^d$ denote a basis of V . For $T \in \text{End}(V)$ and $X \in \text{Mat}_{d+1}(\mathbb{K})$, we say that X represents T with respect to $\{v_i\}_{i=0}^d$ whenever $Tv_j = \sum_{i=0}^d X_{ij}v_i$ for $0 \leq j \leq d$.

By (65) and (66), the matrices that represent A with respect to $\{u_i\}_{i=0}^d$ and $\{u_i^\downarrow\}_{i=0}^d$ are, respectively,

$$\begin{pmatrix} \theta_0 & & & \mathbf{0} \\ 1 & \theta_1 & & \\ & \ddots & \ddots & \\ \mathbf{0} & & 1 & \theta_d \end{pmatrix}, \quad \begin{pmatrix} \theta_d & & & \mathbf{0} \\ 1 & \theta_{d-1} & & \\ & \ddots & \ddots & \\ \mathbf{0} & & 1 & \theta_0 \end{pmatrix}. \quad (71)$$

By construction, the matrix $\text{diag}(q^d, q^{d-2}, \dots, q^{-d})$ represents K with respect to $\{u_i\}_{i=0}^d$, and B with respect to $\{u_i^\downarrow\}_{i=0}^d$, and \mathcal{M} with respect to $\{w_i\}_{i=0}^d$.

Definition 11.3. We define a matrix $\widehat{\psi} \in \text{Mat}_{d+1}(\mathbb{K})$. For $1 \leq i \leq d$, the $(i-1, i)$ -entry is $(q^i - q^{-i})(q^{d-i+1} - q^{i-d-1})$. All other entries are 0.

Proposition 11.4. The matrix $\widehat{\psi}$ represents ψ with respect to each of the bases $\{u_i\}_{i=0}^d$, $\{u_i^\downarrow\}_{i=0}^d$, $\{w_i\}_{i=0}^d$.

Proof: By [5, Line (23)], $\widehat{\psi}$ represents ψ with respect to $\{u_i\}_{i=0}^d$. The remaining assertions follow from Lemma 11.2. \square

Next we give the matrices that represent $\mathcal{M}^{\pm 1}$ with respect to the bases $\{u_i\}_{i=0}^d$, $\{u_i^\downarrow\}_{i=0}^d$.

Lemma 11.5. We give the matrix in $\text{Mat}_{d+1}(\mathbb{K})$ that represents \mathcal{M} with respect to $\{u_i\}_{i=0}^d$. This matrix is upper triangular. For $0 \leq i \leq j \leq d$, the (i, j) -entry is

$$a^{i-j} q^{d-j-i} (q - q^{-1})^{2(j-i)} \frac{[j]_q! [d-i]_q!}{[i]_q! [d-j]_q!}. \quad (72)$$

Proof: The matrix $\text{diag}(q^d, q^{d-2}, \dots, q^{-d})$ represents K with respect to $\{u_i\}_{i=0}^d$. Use this fact along with Lemma 6.5 and Proposition 11.4. \square

Lemma 11.6. We give the matrix in $\text{Mat}_{d+1}(\mathbb{K})$ that represents \mathcal{M}^{-1} with respect to $\{u_i\}_{i=0}^d$. For $0 \leq i \leq d$, the (i, i) -entry is q^{2i-d} . For $1 \leq i \leq d$, the $(i-1, i)$ -entry is

$$-a^{-1} q^{2i-d-1} (q^i - q^{-i}) (q^{d-i+1} - q^{i-d-1}).$$

All other entries are zero.

Proof: The matrix $\text{diag}(q^{-d}, q^{2-d}, \dots, q^d)$ represents K^{-1} with respect to $\{u_i\}_{i=0}^d$. Use this fact along with Lemma 6.4 and Proposition 11.4. \square

Lemma 11.7. We give the matrix in $\text{Mat}_{d+1}(\mathbb{K})$ that represents \mathcal{M} with respect to $\{u_i^\downarrow\}_{i=0}^d$. This matrix is upper triangular. For $0 \leq i \leq j \leq d$, the (i, j) -entry is

$$a^{j-i} q^{d-j-i} (q - q^{-1})^{2(j-i)} \frac{[j]_q! [d-i]_q!}{[i]_q! [d-j]_q!}. \quad (73)$$

Proof: The matrix $\text{diag}(q^d, q^{d-2}, \dots, q^{-d})$ represents B with respect to $\{u_i^\downarrow\}_{i=0}^d$. Use this fact along with Lemma 6.5 and Proposition 11.4. \square

Lemma 11.8. We give the matrix in $\text{Mat}_{d+1}(\mathbb{K})$ that represents \mathcal{M}^{-1} with respect to $\{u_i^\downarrow\}_{i=0}^d$. For $0 \leq i \leq d$, the (i, i) -entry is q^{2i-d} . For $1 \leq i \leq d$, the $(i-1, i)$ -entry is

$$-a q^{2i-d-1} (q^i - q^{-i}) (q^{d-i+1} - q^{i-d-1}).$$

All other entries are zero.

Proof: The matrix $\text{diag}(q^{-d}, q^{2-d}, \dots, q^d)$ represents B^{-1} with respect to $\{u_i^\downarrow\}_{i=0}^d$. Use this fact along with Lemma 6.4 and Proposition 11.4. \square

Next we give the matrices that represent K with respect to the bases $\{u_i^\downarrow\}_{i=0}^d, \{w_i\}_{i=0}^d$.

Lemma 11.9. *We give the matrix in $\text{Mat}_{d+1}(\mathbb{K})$ that represents K with respect to $\{u_i^\downarrow\}_{i=0}^d$. For $0 \leq i \leq d$, the (i, i) -entry is q^{d-2i} . For $0 \leq i < j \leq d$, the (i, j) -entry is*

$$(1 - a^{-2}) a^{j-i} q^{d-j-i} (q - q^{-1})^{2(j-i)} \frac{[j]_q! [d-i]_q!}{[i]_q! [d-j]_q!}. \quad (74)$$

All other entries are zero.

Proof: Evaluating the equation on the right in (14) using the equation on the left in (12) we get

$$K = \left(a^{-2}I + (1 - a^{-2}) \sum_{n=0}^d a^n q^n \psi^n \right) B. \quad (75)$$

The result follows from this along with Proposition 11.4 and the fact that the matrix $\text{diag}(q^d, q^{d-2}, \dots, q^{-d})$ represents B with respect to $\{u_i^\downarrow\}_{i=0}^d$. \square

Lemma 11.10. *We give the matrix in $\text{Mat}_{d+1}(\mathbb{K})$ that represents K with respect to $\{w_i\}_{i=0}^d$. For $0 \leq i \leq d$, the (i, i) -entry is q^{d-2i} . For $1 \leq i \leq d$, the $(i-1, i)$ -entry is*

$$-a^{-1} q^{d-2i+1} (q^i - q^{-i}) (q^{d-i+1} - q^{i-d-1}).$$

All other entries are zero.

Proof: The matrix $\text{diag}(q^d, q^{d-2}, \dots, q^{-d})$ represents \mathcal{M} with respect to $\{w_i\}_{i=0}^d$. Use this fact along with Proposition 11.4 and the equation on the left in (32). \square

Next we give the matrices that represent B with respect to the bases $\{u_i\}_{i=0}^d, \{w_i\}_{i=0}^d$.

Lemma 11.11. *We give the matrix in $\text{Mat}_{d+1}(\mathbb{K})$ that represents B with respect to $\{u_i\}_{i=0}^d$. For $0 \leq i \leq d$, the (i, i) -entry is q^{d-2i} . For $0 \leq i < j \leq d$, the (i, j) -entry is*

$$(1 - a^2) a^{i-j} q^{d-j-i} (q - q^{-1})^{2(j-i)} \frac{[j]_q! [d-i]_q!}{[i]_q! [d-j]_q!}. \quad (76)$$

All other entries are zero.

Proof: Evaluating the equation on the left in (14) using the equation on the right in (12) we get

$$B = \left(a^2I + (1 - a^2) \sum_{n=0}^d a^{-n} q^n \psi^n \right) K. \quad (77)$$

The result follows from this along with Proposition 11.4 and the fact that the matrix $\text{diag}(q^d, q^{d-2}, \dots, q^{-d})$ represents K with respect to $\{u_i\}_{i=0}^d$. \square

Lemma 11.12. *We give the matrix in $\text{Mat}_{d+1}(\mathbb{K})$ that represents B with respect to $\{w_i\}_{i=0}^d$. For $0 \leq i \leq d$, the (i, i) -entry is q^{d-2i} . For $1 \leq i \leq d$, the $(i-1, i)$ -entry is*

$$-aq^{d-2i+1}(q^i - q^{-i})(q^{d-i+1} - q^{i-d-1}).$$

All other entries are zero.

Proof: The matrix $\text{diag}(q^d, q^{d-2}, \dots, q^{-d})$ represents \mathcal{M} with respect to $\{w_i\}_{i=0}^d$. Use this fact along with Proposition 11.4 and the equation on the left in (33). \square

Next we consider the matrices

$$\exp_q \left(\frac{a}{q - q^{-1}} \widehat{\psi} \right), \quad \exp_q \left(\frac{a^{-1}}{q - q^{-1}} \widehat{\psi} \right). \quad (78)$$

Their inverses are

$$\exp_{q^{-1}} \left(-\frac{a}{q - q^{-1}} \widehat{\psi} \right), \quad \exp_{q^{-1}} \left(-\frac{a^{-1}}{q - q^{-1}} \widehat{\psi} \right) \quad (79)$$

respectively. The matrices in (78),(79) are upper triangular. We now consider the entries of (78),(79).

Lemma 11.13. *For $0 \neq x \in \mathbb{K}$, the matrix $\exp_q(x\widehat{\psi})$ is upper triangular. For $0 \leq i \leq j \leq d$, the (i, j) -entry is*

$$x^{j-i} q^{\binom{j-i}{2}} (q - q^{-1})^{2(j-i)} \cdot \frac{[j]_q! [d-i]_q!}{[i]_q! [j-i]_q! [d-j]_q!}. \quad (80)$$

The matrix $\exp_{q^{-1}}(x\widehat{\psi})$ is upper triangular. For $0 \leq i \leq j \leq d$, the (i, j) -entry is

$$x^{j-i} q^{-\binom{j-i}{2}} (q - q^{-1})^{2(j-i)} \cdot \frac{[j]_q! [d-i]_q!}{[i]_q! [j-i]_q! [d-j]_q!}. \quad (81)$$

Lemma 11.14. *The transition matrices between the basis $\{w_i\}_{i=0}^d$ and the bases $\{u_i\}_{i=0}^d$, $\{u_i^\downarrow\}_{i=0}^d$ are given in the table below.*

<i>From</i>	<i>To</i>	<i>Transition Matrix</i>
$\{u_i\}_{i=0}^d$	$\{w_i\}_{i=0}^d$	$\exp_{q^{-1}} \left(-\frac{a^{-1}}{q - q^{-1}} \widehat{\psi} \right)$
$\{w_i\}_{i=0}^d$	$\{u_i\}_{i=0}^d$	$\exp_q \left(\frac{a^{-1}}{q - q^{-1}} \widehat{\psi} \right)$
$\{u_i^\downarrow\}_{i=0}^d$	$\{w_i\}_{i=0}^d$	$\exp_{q^{-1}} \left(-\frac{a}{q - q^{-1}} \widehat{\psi} \right)$
$\{w_i\}_{i=0}^d$	$\{u_i^\downarrow\}_{i=0}^d$	$\exp_q \left(\frac{a}{q - q^{-1}} \widehat{\psi} \right)$

Proof: Use Lemma 11.2 and Proposition 11.4. \square

We next consider the product

$$\exp_q \left(\frac{a}{q - q^{-1}} \widehat{\psi} \right) \exp_{q^{-1}} \left(-\frac{a^{-1}}{q - q^{-1}} \widehat{\psi} \right). \quad (82)$$

The inverse of (82) is

$$\exp_q \left(\frac{a^{-1}}{q - q^{-1}} \widehat{\psi} \right) \exp_{q^{-1}} \left(-\frac{a}{q - q^{-1}} \widehat{\psi} \right). \quad (83)$$

The matrices in (82),(83) are upper triangular.

Lemma 11.15. *The transition matrices between the bases $\{u_i\}_{i=0}^d$, $\{u_i^\downarrow\}_{i=0}^d$ are given in the table below.*

From	To	Transition Matrix
$\{u_i\}_{i=0}^d$	$\{u_i^\downarrow\}_{i=0}^d$	$\exp_q \left(\frac{a}{q - q^{-1}} \widehat{\psi} \right) \exp_{q^{-1}} \left(-\frac{a^{-1}}{q - q^{-1}} \widehat{\psi} \right)$
$\{u_i^\downarrow\}_{i=0}^d$	$\{u_i\}_{i=0}^d$	$\exp_q \left(\frac{a^{-1}}{q - q^{-1}} \widehat{\psi} \right) \exp_{q^{-1}} \left(-\frac{a}{q - q^{-1}} \widehat{\psi} \right)$

Proof: Use Lemma 11.14. □

Lemma 11.16. *With respect to each of the bases $\{u_i\}_{i=0}^d$, $\{u_i^\downarrow\}_{i=0}^d$, $\{w_i\}_{i=0}^d$, the matrices that represent Δ and Δ^{-1} are $\exp_q \left(\frac{a}{q - q^{-1}} \widehat{\psi} \right) \exp_{q^{-1}} \left(-\frac{a^{-1}}{q - q^{-1}} \widehat{\psi} \right)$ and $\exp_q \left(\frac{a^{-1}}{q - q^{-1}} \widehat{\psi} \right) \exp_{q^{-1}} \left(-\frac{a}{q - q^{-1}} \widehat{\psi} \right)$ respectively.*

Proof: Use Theorem 5.5 and Proposition 11.4. □

We give the entries of the matrices representing Δ, Δ^{-1} in the following lemma.

Lemma 11.17. *The matrix in (82) is upper triangular. For $0 \leq i \leq j \leq d$, the (i, j) -entry of (82) is*

$$\frac{(q - q^{-1})^{j-i} [j]_q! [d - i]_q!}{[i]_q! [j - i]_q! [d - j]_q!} \prod_{n=1}^{j-i} (aq^{n-1} - a^{-1}q^{1-n}). \quad (84)$$

The matrix in (83) is upper triangular. For $0 \leq i \leq j \leq d$, the (i, j) -entry of (83) is

$$\frac{(q - q^{-1})^{j-i} [j]_q! [d - i]_q!}{[i]_q! [j - i]_q! [d - j]_q!} \prod_{n=1}^{j-i} (a^{-1}q^{n-1} - aq^{1-n}). \quad (85)$$

Proof: Use Corollary 7.3 and Proposition 11.4. □

We finish the paper by giving the matrix that represents A with respect to $\{w_i\}_{i=0}^d$.

Lemma 11.18. *We give the matrix in $\text{Mat}_{d+1}(\mathbb{K})$ that represents A with respect to $\{w_i\}_{i=0}^d$. For $1 \leq i \leq d$, the $(i, i-1)$ -entry is 1. For $0 \leq i \leq d$, the (i, i) -entry is $(a + a^{-1})q^{d-2i}$. For $1 \leq i \leq d$, the $(i-1, i)$ -entry is*

$$-q^{d-2i+1}(q^i - q^{-i})(q^{d-i+1} - q^{i-d-1}).$$

All other entries are zero.

Proof: Let \mathcal{A} denote the matrix that represents A with respect to $\{w_i\}_{i=0}^d$. By Lemma 9.4, \mathcal{A} is tridiagonal with (i, i) -entry given by $(a + a^{-1})q^{d-2i}$ for $0 \leq i \leq d$.

We now show that the subdiagonal entries of \mathcal{A} are all 1. Let \mathcal{A}' denote the matrix that represents A with respect to $\{u_i\}_{i=0}^d$. Recall that this matrix is displayed on the left in (71). Observe that \mathcal{A} is equal to $\exp_{q^{-1}}(-\frac{a^{-1}}{q-q^{-1}}\widehat{\psi})\mathcal{A}'\exp_q(\frac{a^{-1}}{q-q^{-1}}\widehat{\psi})$. It follows from this fact that the subdiagonal entries of \mathcal{A} are all 1.

We next obtain the superdiagonal entries of \mathcal{A} . Let $0 \leq i \leq d$. Apply both sides of (37) to w_i . Evaluate the result using Proposition 11.4 and the fact that the w_i is an eigenvector for \mathcal{M} with eigenvalue q^{2i-d} . Analyze the result in light of the above comments concerning the entries of \mathcal{A} to obtain the desired result. \square

References

- [1] H. Alnajjar and B. Curtin, *A family of tridiagonal pairs related to the quantum affine algebra $U_q(\widehat{\mathfrak{sl}}_2)$* , Electron. J. Linear Algebra **13** (2005), 1–9.
- [2] G. Andrews, R. Askey, and R. Roy, *Special functions*, Encyclopedia of Mathematics and its Applications, Cambridge University Press, Cambridge, 1999.
- [3] P. Baseilhac, *A family of tridiagonal pairs and related symmetric functions*, J. Phys. A **39** (2006), 11773–11791.
- [4] S. Bockting-Conrad, *Two commuting operators associated with a tridiagonal pair*, Linear Algebra Appl. **437** (2012), 242–270.
- [5] ———, *Tridiagonal pairs of q -Racah type, the double lowering operator ψ , and the quantum algebra $U_q(\mathfrak{sl}_2)$* , Linear Algebra Appl. **445** (2014), 256–279.
- [6] L. Dolan and M. Grady, *Conserved charges from self-duality*, Phys. Rev. D **25** (1982), 1587–1604.
- [7] D. Funk-Neubauer, *Tridiagonal pairs and the q -tetrahedron algebra*, Linear Algebra Appl. **431** (2009), 903–925.
- [8] G. Gasper and M. Rahman, *Basic hypergeometric series*, second ed. ed., Cambridge University Press, New York, 2003.

- [9] T. Ito, K. Tanabe, and P. Terwilliger, *Some algebra related to P- and Q-polynomial association schemes*, Codes and Association Schemes (Piscataway NJ, 1999) (Providence RI), Amer. Math. Soc., 2001, pp. 167–192.
- [10] T. Ito and P. Terwilliger, *The q-tetrahedron algebra and its finite-dimensional irreducible modules*, Comm. Algebra **35** (2007), 3415–3439.
- [11] ———, *Tridiagonal pairs and the quantum affine algebra $U_q(\widehat{\mathfrak{sl}}_2)$* , Ramanujan J. **13** (2007), 39–62.
- [12] ———, *Tridiagonal pairs of q-Racah type*, J. Algebra **322** (2009), 68–93.
- [13] ———, *The augmented tridiagonal algebra*, Kyushu J. Math **64** (2010), 81–144.
- [14] T.H. Koornwinder, *The relationship between Zhedanov’s algebra $AW(3)$ and the double affine Hecke algebra in the rank one case*, SIGMA **3** (2007), 063, 15 pages.
- [15] ———, *Zhedanov’s algebra $AW(3)$ and the double affine Hecke algebra in the rank one case. II. The spherical subalgebra*, SIGMA **4** (2008), 052, 17 pages.
- [16] A. Korovnichenko and A. S. Zhedanov, *“Leonard pairs” in classical mechanics*, J. Phys. A **35** (2002), 5767–5780.
- [17] K. Nomura and P. Terwilliger, *Towards a classification of the tridiagonal pairs*, Linear Algebra Appl. **429** (2008), 503–518.
- [18] S. Odake and R. Sasaki, *Orthogonal polynomials from hermitian matrices*, J. Math. Phys. **49** (2008), 053503, 43 pages.
- [19] L. Onsager, *Crystal statistics. I. A two-dimensional model with an order-disorder transition*, Phys. Rev. **65** (1944), 117–149.
- [20] P. Terwilliger, *Two linear transformations each tridiagonal with respect to an eigenbasis of the other*, Linear Algebra Appl. **330** (2001), 149–203.
- [21] ———, *Leonard pairs from 24 points of view*, Rocky Mountain J. Math **32** (2002), 827–888.
- [22] ———, *Introduction to Leonard pairs*, OPSFA Rome 2001, J. Comput. Appl. Math. **153** (2003), 463–475.
- [23] ———, *Two linear transformations each tridiagonal with respect to an eigenbasis of the other; comments on the parameter array*, Des. Codes Cryptogr. **34** (2005), 307–332.
- [24] ———, *An algebraic approach to the Askey scheme of orthogonal polynomials*, Orthogonal polynomials and special functions, Lecture Notes in Math., 1883 (Berlin), Springer, 2006, pp. 255–330.
- [25] P. Terwilliger and R. Vidunas, *Leonard pairs and the Askey-Wilson relations*, J. Algebra Appl. **3** (2004), 411–426.

Sarah Bockting-Conrad
Department of Mathematical Sciences
DePaul University
2320 N. Kenmore Avenue
Chicago, IL 60614 USA
email: sarah.bockting@depaul.edu