Alternative Renormalizable SO(10) GUTs and Data Fitting

Takeshi Fukuyama^{a,1}, Nobuchika Okada^{b,2} and Hieu Minh Tran^{c,3}

 ^a Research Center for Nuclear Physics (RCNP), Osaka University, Ibaraki, Osaka, 567-0047, Japan
 ^b Department of Physics and Astronomy, University of Alabama,

Tuscaloosa, Alabama 35487, USA

^cHanoi University of Science and Technology, 1 Dai Co Viet Road, Hanoi, Vietnam

Abstract

The alternative renormalizable minimal SO(10) model is composed of the Yukawa couplings with $10 \oplus 120$ Higgs fields, whereas the right-handed (RH) neutrino Majorana masses are generated via the Witten mechanism. The gauge coupling unification is achieved together with a unique pattern of the fermion masses and mixing at the grand unification scale due to additional contributions of vector-like quarks to the standard model renormalization group equations. We perform the fitting of the model to the experimental data of charged fermion masses and the CKM matrix. The best fit point is obtained with large pulls for m_c, m_s, m_b , and m_{τ} . For the modifications to the minimal model by adding either 10' or 120', a large deviation for the tau mass rules out all these models. In the case with the bottom and vectorlike quark mixing, the mass matrices are well fitted the charged fermions but the bound on the light neutrino mass scale excludes this scenario. To ameliorate this deficit, we consider the two-step symmetry breaking scenario, $SO(10) \rightarrow SU(5) \rightarrow SU(3)_C \times SU(2)_L \times U(1)_Y$, with the SO(10) breaking at the Planck scale leading to the radiatively generated RH neutrino Majorana masses being at the ordinary seesaw scale. For this case, we find the best fit point with $\chi^2 = 7.8$ consistent with experimental results including the neutrino sector. The largest deviation is 2.3σ corresponding to the strange quark mass. Hence, a more precise determination of the strange quark mass can test this model. For the best fit point, we find the effective Majorana neutrino mass of $m_{\beta\beta} = 0.22$ meV and the sum of light neutrino masses as $\Sigma = 0.078$ eV, which are consistent with the current constraints from the search for the neutrinoless double beta decay and the CMB anisotropy measurement.

¹E-mail: fukuyama@se.ritsumei.ac.jp

²E-mail: okadan@ua.edu

³E-mail: hieu.tranminh@hust.edu.vn

1 Introduction

The grand unified theory (GUT) with the underlined symmetry of an SO(10) group is a beautiful and convincing picture for the unification of interactions beyond the standard model (SM). It is not only the unification of the SM gauge symmetries, but also that of the SM fermions of each generation into a single **16**-dimensional multiplet. In the SM, it is well known that the gauge couplings do not unify at high energy scales. This issue can be resolved in scenarios where physics at intermediate scales is introduced for the deformation of the renormalization group (RG) trajectories. A popular direction motivated by the gauge hierarchy problem is the assumption of supersymetry (SUSY) with the SUSY breaking scale of about O(1-10) TeV. The SUSY GUTs have been investigated in many aspects.

Since the fermion masses in SO(10) models originate from the Yukawa couplings between Higgs fields and the tensor product of two matter multiplets [1],

$$\mathbf{16} \otimes \mathbf{16} = \mathbf{10}_s \oplus \mathbf{120}_a \oplus \mathbf{126}_s, \qquad (1)$$

the construction of the Higgs sector can be varied. In the minimal SO(10) models with $\mathbf{10} \oplus \overline{\mathbf{126}}$ Higgs fields [2], the heavy right-handed (RH) neutrinos obtain their Majorana masses from the vacuum expectation value (VEV) of ($\overline{\mathbf{10}}, \mathbf{1}, \mathbf{3}$) under the subgroup $SU(4)_C \otimes SU(2)_L \otimes SU(2)_R (G_{422})$ at the tree level, and the light left-handed (LH) neutrino masses are generated via a canonical seesaw mechanism [3]. This minimal model is supplemented with **120** Higgs as a general renormalizable model which has served to correct minor mismatching with data in both SUSY and non-SUSY cases. So there arise interesting issues whether SUSY or non-SUSY and whether the minimal SO(10) is the best SO(10) model or the best GUT model in terms of reproducing experimental data.

If we respect the renormalizability and minimality, a model with $10 \oplus 120$ Higgs fields (we call this model the alternative minimal model) is alternative to the ordinary minimal model with $10 \oplus \overline{126}$ Higgs fields. Unlike the case of $\overline{126}$, 120 Higgs has no VEV which directly generates the RH-neutrino Majorana (and LH-neutrino Majorana) masses. The RH-neutrinos with vanishing masses at the tree level become massive by virtue of the Witten mechanism [4]. It gives an effective **126** coupling with **16**-plets of matters via quantum corrections at the 2-loop level. Thus, the seesaw scale in this scenario is relatively low because of the 2-loop suppression [5, 6]. Due to the non-renormalization theorem of SUSY theories, the Witten mechanism is peculiar and applied only in the non-SUSY framework. Similar to the ordinary minimal SO(10) model, the theoretical predictions of this model on the particle mass spectrum, the Cabibbo-Kobayashi-Maskawa (CKM) and Pontecorvo-Maki-Nakagawa-Sakata (PMNS) mixing matrices should be checked if they are all in agreement with experimental results. Beside the minimal alternative SO(10) model, we consider other two simple extensions with the Higgs sectors, respectively, comprised of $10 \oplus 10' \oplus 120$ and $10 \oplus 120 \oplus 120'$. Although the former is simpler in terms of SO(10) representation, it has more degrees of freedom in the Yukawa couplings. Meanwhile, the latter with larger SO(10)representation is more predictive thanks to its smaller number of free inputs. In all three considered models, we assume a single-scale symmetry breaking pattern, $SO(10) \rightarrow SU(3)_C \otimes SU(2)_L \otimes U(1)_Y$, at the GUT scale (M_G) for simplicity.

Assuming the minimal SUSY extension of the SM (the MSSM) as an effective theory below the GUT scale, the SO(10) model with $10 \oplus 120$ Higgs fields was investigated in Refs. [7, 8, 9, 10]. According to that, the result of data fitting without considering the neutrino sector was not

satisfactory with much larger χ^2 than the minimal model with $10 + \overline{126}$ where $\chi^2 \leq O(1)$ for SUSY [11, 12] and $\chi^2 = 14.7$ for non-SUSY [13] including the neutrino sector fitting. Here, we are interested in the scenario where the successful gauge coupling unification is achieved by the contribution of additional vector-like quarks to the beta function coefficients [14]. The vector-like quarks change the RGE trajectories of the Yukawa couplings compared to those in the SM and the MSSM, resulting in a distinctive pattern of fermion masses and mixing at the GUT scale. We will also investigate the two-step symmetry breaking scenario where the GUT symmetry breaking chain $SO(10) \rightarrow SU(5) \rightarrow SU(3)_C \times SU(2)_L \times U(1)_Y$ happens at two different scales. The matching of the alternative SO(10) models to such an effective non-SUSY model at low energies has not been investigated so far, and will show new fitting results. This analysis is essential to quantify how well the above model setups are compatible with the updated experimental results, for which the answer could not be derived from previous works. The existence of vector-like quarks is also well motivated by the stability of the electroweak vacuum below the GUT scale [15], and could play the role in the observed experimental anomalies [16]. In this paper, we employ the following procedure to fit the model parameters to the measured observables including those in the neutrino sector. Firstly, the experimental values and errors of the fermion masses and mixing parameters are evolved to the GUT scale, taking into account the effects of the additional vector-like quarks. Using the χ^2 function as a measure, we then look for the best fit points of these models.

The paper is organized as follows. In the next section, the alternative minimal SO(10) model is briefly reviewed, and its two simple extensions are introduced. In Section 3, the experimental observables are evolved to the GUT scale by solving the RG equations. In Section 4, we perform the data fittings for each alternative renormalizable SO(10) models and show the results. In Section 5, we introduce the mixing between the bottom and vector-like quarks mixing in the minimal alternative model and present the data fitting results for this scenario. The last section is devoted to the conclusion.

2 Alternative renormalizable *SO*(10) models

2.1 The minimal model

When an SO(10) model includes the **126** Higgs field, a VEV of the (**10**, **1**, **3**) component under G_{422} generates the Majorana masses for RH neutrinos (N_R 's) at the tree level. However, even without the **126** Higgs field, we can generate these Majorana masses when the model includes a **16**-plet Higgs field (H_{16}), since a bilinear product of H_{16}^{\dagger} can play a role of the **126** Higgs field. An effective operator relevant to this mass generation is given by

$$\mathcal{L} \supset \frac{1}{M} \mathbf{16}_i \mathbf{16}_j H_{16}^{\dagger} H_{16}^{\dagger}, \tag{2}$$

where M is a mass parameter. Although we cannot introduce such a higher dimensional term by hand in a renormalizable model, it can be induced through quantum corrections at the 2-loop level as pointed out by Witten [4]. This is very interesting since the loop corrections suppress the seesaw scale.

In the simplest model discussed in Ref. [4], the matter fermions couple directly only to the 10-plet Higgs and the SO(10) gauge field of the 45 representation. The basic idea is that 126



Figure 1: The construction of the RH neutrino mass from $\mathbf{10} \otimes \mathbf{45} \otimes \mathbf{45}$ via a 2-loop diagram. Shown in parenthesis are the SO(10) and the broken subgroup $(SU(5) \text{ or } G_{422})$ representations, which are summarized in Table 2. Two blobs represent the insertion of $\langle H_{16} \rangle = M_G$. The crossed diagram should be added.

	(a)	(b)	(c)	(d)	(e)	(f)	(g)	(h)
SU(5)	1	$\overline{5}$	10	5	10	10	1	5
G_{422}	$\overline{4}, 1, 2$	${f 4, 2, 1}$	${f 4, 2, 1}$	1, 2, 2	15, 1, 1	${\bf 6, 2, 2}$	$\overline{4}, 1, 2$	${f 4, 2, 1}$

Table 1: Representations of particles in the 2-loop diagram under the SO(10) subgroups.

representation is a 5th rank tensor which can be constructed by the product $\mathbf{10} \otimes \mathbf{45} \otimes \mathbf{45}$ with a vector $\mathbf{10}$ and a 2nd rank tensor $\mathbf{45}$. In fact, the N_R mass is generated by quantum corrections at the 2-loop level as shown in Fig. 1, when H_{16} develops its vacuum expectation value (VEV) $(\mathbf{1}, -5)$ for the subgroup $SU(5) \otimes U(1)_X$ or a VEV $(\mathbf{\overline{4}}, \mathbf{1}, \mathbf{2})$ for the subgroup $SU(4) \otimes SU(2)_L \otimes SU(2)_R$. Here, note that a triple scalar coupling among the Higgs fields of $\mathbf{10}$ -plet and $\mathbf{16}$ -plets also plays a crucial role:

$$\mathcal{L} \supset \lambda_{10} M_G H_{10} H_{16} H_{16},\tag{3}$$

where we have parametrize the triple scalar coupling with M_G and a dimensionless coupling constant λ_{10} . The resultant N_R mass is estimated as

$$M_R = \left(\frac{m_q}{M_W}\right) \epsilon_{10} \left(\frac{\alpha_G}{\pi}\right)^2 M_G. \tag{4}$$

Here, we have used a relation, $Y_{10} \sim m_q/M_W$ between the Yukawa coupling of H_{10} and an up-type quark mass m_q , and ϵ_{10} represents a mixing angle between H_{10} and H_{16} induced by their coupling in Eq. (3) with a VEV of H_{16} . Note that the M_R scale is much lower than the usual seesaw scale of the model $\sim M_G$. In the present SO(10) model with only one Yukawa coupling Y_{10} , the Dirac neutrino mass matrix is the same as the up-type quark mass matrix. Therefore, the light neutrino mass m_{ν_L} due to the type I seesaw mechanism,

$$m_{\nu_L} = -M_D^T M_R^{-1} M_D, (5)$$

is given by

$$m_{\nu_L} = m_q \left[\epsilon_{10} \left(\frac{\alpha_G}{\pi} \right)^2 \right]^{-1} \frac{M_W}{M_G}.$$
 (6)

As in Ref. [4], we estimate $m_{\nu_L} = 10^{-7} m_q$ by using $(\alpha_G/\pi)^2 = 10^{-5}$, $M_G = 10^{15}$ GeV and $\epsilon_{10} = 0.1$. Clearly, the light neutrino mass spectrum predicted by this formula is unrealistic. For example, the heaviest light neutrino mass is $10^{-7} m_t \sim 20$ keV, where $m_t = 173$ GeV is the top quark mass. Thus, the light neutrino masses from the type I seesaw with the 2-loop induced M_R are too heavy. This is due to the quark mass m_q insertions in Eqs. (4) and (6), which originate from the single Yukawa coupling Y_{10} . A simple way to ameliorate the problem is to add one more Yukawa coupling with **120** Higgs. **120** Higgs field includes two pairs of $SU(2)_L$ Higgs doublets (**15**, **2**, **2**) and (**1**, **2**, **2**) unlike the **126** Higgs field. Their general renormalizable mass formula was given in [8]. **120** Higgs does not involve (**10**, **3**, **1**) \oplus (**10**, **1**, **3**) under the subgroup G_{422} . Hence, the Majorana neutrino masses are not generated at the tree-level but via the two-loop correction a la Witten mechanism. With the **120** Higgs field, the Yukawa interactions are given by

$$\mathcal{L}_Y = Y_{10}^{ij} \mathbf{16}_i \mathbf{10}_H \mathbf{16}_j + Y_{120}^{ij} \mathbf{16}_i \mathbf{120}_H \mathbf{16}_j.$$
(7)

With the VEVs of three pairs of Higgs doublets (one in $\mathbf{10}_H$ and two in $\mathbf{120}_H$), the fermion mass matrices at M_G are described as [8]

$$M_u = c_{10} M_{10} + c_{120} m_{120} + c'_{120} m'_{120}, \qquad (8)$$

$$M_d = M_{10} + m_{120} + m'_{120}, (9)$$

$$M_D = c_{10} M_{10} + c_{120} m_{120} - 3 c'_{120} m'_{120}, \qquad (10)$$

$$M_e = M_{10} + m_{120} - 3 m'_{120}, \tag{11}$$

where M_u , M_d , M_D , and M_e denote the up-type quark, down-type quark, Dirac neutrino, and charged-lepton, respectively. Here, the mass matrices m_{120} and m'_{120} come from (1, 2, 2) and (15, 2, 2) of 120, respectively. The mass matrices of the right-hand sides are defined as $M_{10} =$ $Y_{10} \alpha_d^1 v \cos \beta$, $m_{120} = Y_{120} \alpha_d^2 v \cos \beta$, and $m'_{120} = Y_{120} \alpha_d^3 v \cos \beta$ respectively, and the coefficients are defined as $c_{10} = (\alpha_u^1/\alpha_d^1) \tan \beta$, $c_{120} = (\alpha_u^2/\alpha_d^2) \tan \beta$, $c'_{120} = (\alpha_u^3/\alpha_d^3) \tan \beta$. $\alpha_{u,d}^i$ are the complex elements of the unitary matrices which make the light pair of Higgs doublets H_u and H_d from many doublets. These complex values depend on the Higgs potential. See [8] for details. Here it is sufficient to recognize that the coefficients cs are independent complex numbers. These mass matrices are directly connected with low-energy observations and are resumed as

$$M_u = c_{10}M_{10} + c_{120}^u M_{120}, (12)$$

$$M_d = M_{10} + M_{120}, (13)$$

$$M_D = c_{10}M_{10} + c_{120}^n M_{120}, (14)$$

$$M_e = M_{10} + c_{120}^e M_{120}. (15)$$

Here, M_{120} is

$$M_{120} = m_{120} + m'_{120} \,, \tag{16}$$

and the complex coefficients $c_{120}^{u,n,e}$ are

$$c_{120}^{u} = \frac{\alpha_u^2 + \alpha_u^3}{\alpha_d^2 + \alpha_d^3} \tan\beta, \qquad (17)$$

$$c_{120}^n = \frac{\alpha_u^2 - 3\alpha_u^3}{\alpha_d^2 + \alpha_d^3} \tan\beta, \qquad (18)$$

$$c_{120}^e = \frac{\alpha_d^2 - 3\alpha_d^3}{\alpha_d^2 + \alpha_d^3} \tan \beta.$$
(19)

The mass matrices M_{10} and M_{120} are respectively complex symmetric and antisymmetric. Since the neutrino Dirac mass matrix M_D is not equal to M_u , hence the problem appeared in Eq. (6) does not occur. Here, the light LH neutrino mass can be estimated as

$$m_{\nu_L} \sim \frac{(M_u + (c_{120}^n - c_{120}^u)M_{120})^2}{M_u - c_{120}^u M_{120}} \times \left[\epsilon_{10} \left(\frac{\alpha_G}{\pi}\right)^2\right]^{-1} \frac{M_W}{M_G}.$$
 (20)

With appropriate values of the parameters c_{120}^n , c_{120}^u , and M_{120} , the light LH neutrino mass can be of the right order. For example, in the case $c_{120}^u M_{120} = \mathcal{O}(M_u)$, we can set $c_{120}^n M_{120}$ so as to almost cancel the numerator such that the light LH neutrino mass can be $m_{\nu_L} = \mathcal{O}(0.1)$ eV. In this estimation, the hierarchy in m_{ν_L} is similar to that in M_u .

Given the experimental data at that time, the system with $10 \oplus 120$ Higgs fields was shown to be consistent with the realistic charged fermion mass spectra [7] when the Yukawa coupling matrices of 10 and 120 are respectively assumed to be real and pure imaginary for simplicity. The fitting was not exhausted with full data. Afterward, more elaborate data fittings of this model have been performed in the Ref. [9, 10] for the SUSY case. We perform the data fitting of our models in Section 4.

2.2 Two simple extensions

Beside the alternative minimal SO(10) model, we also consider some extensions of it. As simple possibilities, we introduce one more Higgs multiplet of either 10'-plet or 120'-plet. Although the 10'-plet is simpler than the 120'-plet in terms of the field content, the Yukawa sector of the former case has more independent parameters than that of the latter one.

In the extension with the Higgs sector $10 \oplus 120 \oplus 120'$, the Yukawa sector is given as

$$\mathcal{L}_{Y}^{1} = Y_{10}^{ij} \mathbf{16}_{i} \mathbf{10}_{H} \mathbf{16}_{j} + Y_{120}^{ij} \mathbf{16}_{i} \mathbf{120}_{H} \mathbf{16}_{j} + Y_{120'}^{ij} \mathbf{16}_{i} \mathbf{120'}_{H} \mathbf{16}_{j}.$$
(21)

There are five pairs of Higgs doublets developing VEVs in this model (one in $\mathbf{10}_H$, and two in each representation of $\mathbf{120}_H$ and $\mathbf{120}'_H$) where only one pair of their linear combination (H_u, H_d) are assumed to be light while the other four are heavy [9]. Once the light Higgs doublets develop

their VEVs (v_u, v_d) , the SM fermion masses are generated via the Yukawa couplings at M_G :

$$M_u = c_{10}M_{10} + c_{120}^u m_{120} + c_{120}'^u m_{120}', \qquad (22)$$

$$M_d = M_{10} + m_{120} + m'_{120}, (23)$$

$$M_{D} = c_{10}M_{10} + c_{120}^{n}m_{120} + c_{120}^{\prime n}m_{120}^{\prime}, \qquad (24)$$

$$M_{D} = M_{10} + c_{120}^{n}m_{120} + c_{120}^{\prime n}m_{120}^{\prime}, \qquad (24)$$

$$M_e = M_{10} + c_{120}^e m_{120} + c_{120}^c m_{120}^\prime.$$
⁽²⁵⁾

After a re-parameterization,

$$M_{120} = m_{120} + m'_{120}, (26)$$

$$M'_{120} = c^u_{120} m_{120} + c'^u_{120} m'_{120}, \qquad (27)$$

$$d_{120}^{n} = \frac{c_{120}^{n} c_{120}^{\prime u} - c_{120}^{\prime n} c_{120}^{u}}{c_{120}^{\prime u} - c_{120}^{u}},$$
(28)

$$d'_{120}^{n} = \frac{c'_{120}^{n} - c_{120}^{n}}{c'_{120}^{u} - c_{120}^{u}},$$
(29)

$$d_{120}^{e} = \frac{c_{120}^{e} c_{120}^{\prime u} - c_{120}^{\prime e} c_{120}^{u}}{c_{120}^{\prime u} - c_{120}^{u}},$$
(30)

$$d'^{e}_{120} = \frac{c'^{e}_{120} - c^{e}_{120}}{c'^{u}_{120} - c^{u}_{120}},$$
(31)

the fermion mass matrices at the GUT scale in this model are determined by the GUT parameters as

$$M_u = c_{10}M_{10} + M'_{120}, (32)$$

$$M_d = M_{10} + M_{120}, (33)$$

$$M_D = c_{10}M_{10} + d_{120}^n M_{120} + d'_{120}^n M'_{120}, \qquad (34)$$

$$M_e = M_{10} + d_{120}^e M_{120} + d_{120}^{\prime e} M_{120}^{\prime}, \qquad (35)$$

where the coefficients c_{10} , d_{120}^n , d_{120}^n , d_{120}^e , and d'_{120}^e are complex. The matrix M_{10} is complex symmetric, and the matrices M_{120} and M'_{120} are both complex antisymmetric.

In the other extension, the Higgs sector consists of $10 \oplus 10' \oplus 120$ representations. The Yukawa sector is given as

$$\mathcal{L}_{Y}^{2} = Y_{10}^{ij} \mathbf{16}_{i} \mathbf{10}_{H} \mathbf{16}_{j} + Y_{10'}^{ij} \mathbf{16}_{i} \mathbf{10}_{H}' \mathbf{16}_{j} + Y_{120}^{ij} \mathbf{16}_{i} \mathbf{120}_{H} \mathbf{16}_{j}.$$
(36)

Similar to the case of $10 \oplus 120 \oplus 120'$ Higgs fields, in this case the fermion mass matrices at the GUT scale are determined by the GUT parameters as

$$M_u = M'_{10} + c^u_{120} M_{120}, (37)$$

$$M_d = M_{10} + M_{120}, (38)$$

$$M_D = M'_{10} + c^n_{120} M_{120}, (39)$$

$$M_e = M_{10} + c_{120}^e M_{120}, (40)$$

where the coefficients $c_{120}^{u,n,e}$ are complex, the matrices M_{10} and M'_{10} are both complex symmetric, and the matrix M_{120} is complex antisymmetric.

To fit the model parameters to the experimental data, we firstly evaluate the experimental values of observables at the GUT scale by solving the RG equations (RGEs). The theoretical prediction of these observables at the GUT scale to be compared with the experimental values are determined for each point of the free parameter space. Then, we scan over the parameter space to find the best fit point.

3 Observables at the GUT scale

In the SM, it is well-known that the three gauge couplings are not successfully unified at high energy scales. To resolve this shortcoming, we assume that below the GUT scale there are two pairs of vector-like quarks (Q_L, Q_R, D_L, D_R) carrying the SM charges [14] as given in Table 2.

	$SU(3)_C$	$SU(2)_L$	$U(1)_Y$
$Q_{L,R}$	3	2	$\frac{1}{6}$
$D_{L,R}$	3	1	$\frac{1}{3}$

Table 2: The SM charges of the additional vector-like quarks

Similar to the case of a SU(5) model [17], in the framework of the SO(10) GUT, these vectorlike quarks are embedded in two pairs of vector-like SO(10) representation, $F_{16}^Q + \bar{F}_{16}^Q$ and $F_{16}^D + \bar{F}_{16}^D$. The Lagrangian for them is written as follows:

$$\mathcal{L}_{SO(10)} \supset \operatorname{Tr} \left[\bar{F}_{16}^{Q} (Y_{16}^{Q} \Sigma - M_{16}^{Q}) F_{16}^{Q} \right] + \operatorname{Tr} \left[\bar{F}_{16}^{D} (Y_{16}^{D} \Sigma - M_{16}^{D}) F_{16}^{D} \right],$$
(41)

where $Y_{16}^{Q,D}$ are Yukawa couplings, $M_{16}^{Q,D}$ are vector-like fermion masses, and Σ is an SO(10)adjoint Higgs field of **45**-plet whose vacuum expectation value with $\langle H_{16} \rangle$ breaks SO(10) to $SU(3)_C \times SU(2)_L \times U(1)_Y$. By tuning $Y_{16}^{Q,D}$ and $M_{16}^{Q,D}$ according to the method of generating doublet-triplet mass splitting, the vector-like quarks (Q_L, Q_R, D_L, D_R) can be as light as $\mathcal{O}(\text{TeV})$, while other components are super-heavy with masses around the GUT scale.

At low energies, the Lagrangian of the effective theory includes that of the SM and the mass terms for the vector-like quarks:

$$\mathcal{L}_{\text{mass}} \supset m_Q \bar{Q}_L Q_R + m_D \bar{D}_L D_R + \text{h.c.}$$
(42)

In this analysis, we assume that $m_Q = m_D = M_F = \mathcal{O}(\text{TeV})$ for simplicity.

From the electroweak scale to the vector-like quark scale M_F , the gauge coupling running follows the SM RGEs. At the 2-loop level, they are given by [18]

$$\frac{dg_i}{d\ln\mu} = \frac{b_i}{16\pi^2}g_i^3 + \frac{g_i^3}{(16\pi^2)^2} \left(\sum_{j=1}^3 B_{ij}g_j^2 - \operatorname{Tr}(C_{i,u}Y_u^{\dagger}Y_u + C_{i,d}Y_d^{\dagger}Y_d + C_{i,e}Y_e^{\dagger}Y_e)\right), \quad (43)$$

where the values of the index i = 1, 2, 3 correspond to the SM gauge groups $U(1)_Y, SU(2)_L$, and $SU(3)_C$. The other coefficients are as follows

$$b_i = \left(\frac{41}{10}, -\frac{19}{6}, -7\right), \quad (i = 1, 2, 3),$$
(44)

$$B_{ij} = \begin{pmatrix} \frac{199}{50} & \frac{27}{10} & \frac{44}{5} \\ \frac{9}{10} & \frac{35}{6} & 12 \\ \frac{11}{10} & \frac{9}{2} & -26 \end{pmatrix},$$
(45)

$$C_{i,f} = \begin{pmatrix} \frac{17}{10} & \frac{1}{2} & \frac{3}{2} \\ \frac{3}{2} & \frac{3}{2} & \frac{1}{2} \\ 2 & 2 & 0 \end{pmatrix}, \qquad (f = u, d, e).$$
(46)

With the existence of vector-like quarks between M_F and the GUT scale M_G , the beta function coefficients of the RGEs in this energy range for the SM gauge couplings are modified by additional contributions from these fermions [14]:

$$b'_i = \left(\frac{2}{5}, 2, 2\right), \qquad (i = 1, 2, 3),$$
(47)

$$B'_{ij} = \begin{pmatrix} \frac{3}{50} & \frac{3}{10} & \frac{8}{5} \\ \frac{1}{10} & \frac{49}{2} & 8 \\ \frac{1}{5} & 3 & \frac{114}{3} \end{pmatrix}.$$
 (48)

The RGEs for other parameters in this energy range are the same as those for the SM parameters. The vector-like quarks affect these RGEs indirectly via those of the gauge couplings.

By varying the vector-like quark mass scale, we find that a successful gauge coupling unification is achieved at $M_G = 1.5 \times 10^{16}$ GeV for $M_F = 2$ TeV. The RG evolution of three gauge couplings in this case is demonstrated in Figure 2.

In order to evaluate the fermion masses and mixing at a high energy scale (like the GUT scale), all the relevant quantities must be determined at that scale for consistency [19]. To determine the VEV at the GUT scale we used the RGE of the VEV in the paper by Arason *et al* (Phys. Rev. D 46, 3945 (1992)) where the VEV running originates from the Higgs doublets wavefunction renormalization.⁴ In our analysis, we solve the RGEs for the Yukawa coupling matrices and the VEV at the two-loop level [18] from m_Z to M_G with the boundary conditions given at the electroweak (EW) scale. The center values and the corresponding errors⁵ of charged fermion

 $^{^{4}}$ Some of the previous works neglected the evolution of the VEV for simplicity, see for example Ref. [20]. We thank the referee for commenting on this point.

⁵ It is worth noting that observables with large errors (like m_u and m_d) have negligible contributions to the RGEs, while the observable having significant contributions to the RGEs have a very small relative error (e.g m_t). We have checked that the correlations among errors are negligibly small and, in a good approximation, the upper/lower bound of an observable at the EW scale corresponds to its upper/lower (or lower/upper) bound at the GUT scale.



Figure 2: The RG runnings of three SM gauge couplings when the vector-like quark mass is $M_F = 2$ TeV. In this case, the gauge couplings unify at $M_G = 1.5 \times 10^{16}$ GeV.

masses and mixing at the EW scale are taken from Table 1 of Ref. [21] where the results in [22] were invoked. For the neutrino oscillation data at low energies, we use the values in Table 1 of Ref. [23].

The charged fermion mass matrices at the GUT scale M_G are calculated as

$$M_f(M_G) = Y_f(M_G) \frac{v(M_G)}{\sqrt{2}}, \qquad (f = u, d, e).$$
 (49)

The CKM mixing matrix at M_G is determined from the rotation matrices that diagonalize the matrices $Y_u^{\dagger}Y_u$ and $Y_d^{\dagger}Y_d$ as follows:

$$U_{\rm CKM}(M_G) = V_{uL}(M_G) \cdot V_{dL}^{\dagger}(M_G).$$
⁽⁵⁰⁾

In Table 3, we present the mean values and errors of the fermion masses and mixings at the GUT scale. Here, $m_{u,c,t}$, $m_{d,s,b}$ and $m_{e,\mu,\tau}$ are, respectively, the singular values of the mass matrices M_u, M_d , and M_e in Eq. (49), while $\lambda, A, \bar{\rho}, \bar{\eta}$ are the Wolfenstein parameters of the CKM mixing matrix in Eq. (50). Δm_{21}^2 and Δm_{31}^2 are the neutrino squared mass differences, while $\theta_{12}, \theta_{23}, \theta_{13}$, and δ_{CP} are the parameters of the PMNS mixing matrix.

4 Data fitting for alternative models

To measure how well a point of the parameter space can predict the experimental data, we use the χ^2 function defined as

$$\chi^2 = \sum_i p_i^2 = \sum_i \left(\frac{x_i - \bar{x}_i}{\sigma_i}\right)^2,\tag{51}$$

Observables	\bar{x}_i	σ_i
m_u	0.40×10^{-3}	0.20×10^{-3}
m_c	0.199	0.031
m_t	60.4	1.7
m_d	0.88×10^{-3}	0.46×10^{-3}
m_s	0.018	0.006
m_b	0.829	0.026
m_e	0.440×10^{-3}	0.022×10^{-3}
m_{μ}	0.093	0.005
$m_{ au}$	1.583	0.079
λ	0.22469	$0.39 imes 10^{-3}$
A	0.932	0.012
$\bar{ ho}$	0.140	0.016
$ar\eta$	0.356	0.010
Δm_{21}^2	1.112×10^{-4}	0.032×10^{-4}
Δm_{31}^2	3.798×10^{-3}	0.048×10^{-3}
θ_{12}	0.573	0.013
θ_{23}	0.867	0.017
θ_{13}	0.1503	0.0023
$\delta_{ m CP}$	-2.50	0.59

Table 3: The charged fermion masses (in GeV), the Wolfenstein parameters of the CKM mixing matrix, the neutrino squared mass differences (in eV^2), and the parameters of the PMNS matrix at the GUT scale M_G . The \bar{x}_i and σ_i columns correspond to the mean values and the uncertainties.

where x_i indicate the theoretical values of the observables, \bar{x}_i and σ_i are the mean value and the uncertainties of the observables, and p_i are the pulls to the corresponding observables. The sum is taken over all 13 observables in Table 3. The best fit point of the model that is the global minimum of the χ^2 function is found by performing two sequential steps. Firstly, we randomly scan over the parameter space with the package MultiNest version 3.11 [24] where the nested sampling algorithm is employed. The ranges of the free inputs are chosen to be [-100,100] for those relevant to the coefficients (c_{10}, c_{120}, d_{120}), and [$10^{-20}, 100$] (GeV) for those relevant to the mass matrices ($M_{10}, M_{120}, M'_{10}, M'_{120}$) with positive and negative signs allowed. This scan is carried out many times to avoid falling into some local minimum. Secondly, the smallest minimum found in the first step is refined further by using the Nelder-Mead simplex algorithm [25] with a high precision.

4.1 The minimal model with $10 \oplus 120$ Higgs multiplets

In the SO(10) model with $10 \oplus 120$ Higgs multiplets, we consider a general case where the coefficients $(c_{120}^{u,n,e})$ and the matrices (M_{10}, M_{120}) in Eqs. (12)-(15) are complex. Inherited from the properties of the corresponding Yukawa couplings, the mass matrices M_{10} and M_{120} are symmetric and antisymmetric, respectively. With an appropriate choice of the SO(10) fermion basis, we can set M_{10} to be real and diagonal without loss of generality. By rephasing u-type quarks, c_{10} can be real. Thus, relevant to the data fitting for the charged fermions masses and mixing there are totally 14 free real parameters including four parameters for the complex coefficients $c_{120}^{u,e}$, one parameter for the real coefficient c_{10} , three parameters for the real diagonal matrix M_{10} , and six parameters for the complex antisymmetric matrix M_{120} .

The best fit point that we have found in the numerical analysis is

$$c_{10} = 67.126,$$
 (52)

$$c_{120}^u = 146.235 + 64.985i, (53)$$

$$c_{120}^e = 9.8535 + 5.3155i, (54)$$

$$M_{10} = \begin{pmatrix} 0.000516765 & 0 & 0 \\ 0 & 0.00432434 & 0 \\ 0 & 0 & 0.893733 \end{pmatrix},$$
(55)

(56)

$$M_{120} =$$

$$\left(\begin{array}{cccc} 0 & 0.00034554 + 0.00042528i & 0.0061402 - 0.0059386i \\ -0.00034554 - 0.00042528i & 0 & 0.013011 + 0.023147i \\ -0.0061402 + 0.0059386i & -0.013011 - 0.023147i & 0 \end{array}\right).$$

In Table 4, the predicted values and the pulls of the charged fermion masses and the Wolfenstein parameters of the CKM mixing matrix at the GUT scale for the best fit point are presented. We see that the total χ^2 value is 69.5079 which is not good. It is mostly due the large pulls toward

Observables	x_i	p_i
m_u	0.40089×10^{-3}	0.0096
m_c	0.20216	0.096
m_t	60.3245	0.018
m_d	0.49026×10^{-3}	0.87
m_s	0.004064	2.5
m_b	0.89461	2.5
m_e	0.44180×10^{-3}	0.061
m_{μ}	0.095153	0.43
$m_{ au}$	0.99239	7.4
λ	0.22469	0.0055
A	0.93283	0.031
$\bar{ ho}$	0.13985	0.020
$ar\eta$	0.34626	0.22×10^{-3}
Total χ^2	69.50	79

Table 4: The best fit point values for the charged fermion masses (in GeV) and the Wolfenstein parameters of the CKM mixing matrix in the $\mathbf{10} \oplus \mathbf{120}$ model at the GUT scale. The x_i and p_i columns correspond to the theoretical predicted values and the pulls.

opposite sides of the mean values for the bottom and tau masses:

$$m_b^{\text{best-fit}} = \overline{m}_b + 2.5 \,\sigma_b, \tag{57}$$

$$m_{\tau}^{\text{best-fit}} = \overline{m}_{\tau} - 7.3 \,\sigma_{\tau}. \tag{58}$$

We can also see that there is a mild pull for the strange quark mass:

$$m_s^{\text{best-fit}} = \overline{m}_s - 2.5 \,\sigma_s. \tag{59}$$

4.2 The model with $10 \oplus 120 \oplus 120'$ Higgs sector

The number of free real parameter of this model is 20 including one parameter for the real coefficient c_{10} , four parameters for the complex coefficients d_{120}^e and d'_{120}^e three parameters for the real diagonal matrix M_{10} (in an appropriate basis), twelve parameters for the complex antisymmetric matrices M_{120} , and M'_{120} . The data fitting procedure for the $\mathbf{10} \oplus \mathbf{120} \oplus \mathbf{120'}$ model results in the best fit point determined by

$$c_{10} = 66.335,$$
 (60)

$$d_{120}^e = -0.93895 + 0.39900i, (61)$$

$$d'_{120}^{e} = -0.016416 - 0.0047637i, (62)$$

$$M_{10} = \begin{pmatrix} 0.00047530 & 0 & 0\\ 0 & 0.050943 & 0\\ 0 & 0 & 0.86104 \end{pmatrix},$$
(63)

Observables	x_i	p_i
m_u	0.39931×10^{-3}	0.17×10^{-2}
m_c	0.19919	7.7×10^{-4}
m_t	60.3537	0.10×10^{-2}
m_d	0.86175×10^{-3}	0.042
m_s	0.020710	0.50
m_b	0.89166	2.4
m_e	0.44049×10^{-3}	0.18×10^{-2}
m_{μ}	0.095107	0.42
$m_{ au}$	1.00705	7.3
λ	0.22469	0.37×10^{-2}
A	0.93248	0.17×10^{-2}
$\bar{ ho}$	0.13952	8.4×10^{-7}
$ar\eta$	0.34626	3.2×10^{-4}
Total χ^2	59.30	01

Table 5: The best fit point values for the charged fermion masses (in GeV) and the Wolfenstein parameters of the CKM mixing matrix in the $\mathbf{10} \oplus \mathbf{120} \oplus \mathbf{120'}$ model at the GUT scale. The x_i and p_i columns correspond to the theoretical predicted values and the pulls.

$$M_{120} = \begin{pmatrix} 0 & 0.0021268 - 3.5349 \times 10^{-5}i & -0.0081535 - 0.00049252i \\ -0.0021268 + 3.5349 \times 10^{-5}i & 0 & -3.5746 \times 10^{-5} + 0.16025i \\ 0.0081535 + 0.00049252i & 3.5746 \times 10^{-5} - 0.16025i & 0 \end{pmatrix},$$

$$M'_{120} = \begin{pmatrix} 0 & 0.012290 + 0.079333i & 0.19358 - 0.11446i \\ -0.012290 - 0.079333i & 0 & 0.046671 + 13.5779i \\ -0.19358 + 0.11446i & -0.046671 - 13.5779i & 0 \end{pmatrix}.$$
(65)

In Table 5, the predicted values and the pulls for the observables of this model at the GUT scale are shown. We see that there is no more tension on the second generation fermion masses (m_c, m_s) due to the model's flexibility with more degrees of freedom than that of the minimal one. However, the tension in the third generation between the the pulls for the bottom and tau masses persists with smaller deviation for the tau mass:

$$m_b^{\text{best-fit}} = \overline{m}_b + 2.4 \,\sigma_b, \tag{66}$$

$$m_{\tau}^{\text{best-fit}} = \overline{m}_{\tau} - 7.3 \,\sigma_{\tau}. \tag{67}$$

4.3 The model with $10 \oplus 10' \oplus 120$ Higgs sector

In this extended model, the number of free real parameter is 24 including one parameter for the real coefficient c_{120}^u , two parameters for the complex coefficient c_{120}^e , three parameters for the real diagonal matrix M_{10} (in an appropriate basis), twelve parameters for the complex symmetric matrix M'_{10} , and six parameters for the complex antisymmetric matrices M_{120} . The best fit point for the $\mathbf{10} \oplus \mathbf{10}' \oplus \mathbf{120}$ model is found to be

$$c_{120}^u = -20.3343, (68)$$

$$c_{120}^e = 3.08017 - 5.24577 \times 10^{-6} i,$$
 (69)

$$M_{10} = \begin{pmatrix} -0.000717021 & 0 & 0 \\ 0 & 0.0362958 & 0 \\ 0 & 0 & 0.875936 \end{pmatrix},$$
(70)

$$M_{10}' = \begin{pmatrix} 0.0059128 + 0.0054818i & -0.04779 + 0.013064i & 0.11143 + 0.13764i \\ -0.04779 + 0.013064i & -0.082199 + 0.20885i & 4.37003 - 0.61857i \\ 0.11143 + 0.13764i & 4.37003 - 0.61857i & -34.4169 - 49.0727i \end{pmatrix}, (71)$$

$$M_{120} =$$

$$\begin{pmatrix} 0 & 1.5193 \times 10^{-5} - 0.0013945i & 5.0427 \times 10^{-5} + 0.0053721i \\ -1.5193 \times 10^{-5} + 0.0013945i & 0 & 9.7925 \times 10^{-7} - 0.11571i \\ -5.0427 \times 10^{-5} - 0.0053721i & -9.7925 \times 10^{-7} + 0.11571i & 0 \end{pmatrix}.$$
(72)

In Table 6, we present the predicted values and the pulls for the observables of this model at the GUT scale for the best fit point of this model. Here, although the pull for the bottom mass is slightly smaller than that in the previous model due to a larger number of degrees of freedom, the large pull for the tau mass is still approximately the same:

$$m_b^{\text{best-fit}} = \overline{m}_b + 2.4 \,\sigma_b, \tag{73}$$

$$m_{\tau}^{\text{best-fit}} = \overline{m}_{\tau} - 7.3 \,\sigma_{\tau}. \tag{74}$$

Meanwhile, as expected the fittings for other observables are much improved thanks to the large number of degrees of freedom in this model.

In the alternative minimal model with $\mathbf{10} \oplus \mathbf{120}$ Higgs, both M_{10} and M_{120} that are used to fit down-type quark and charged lepton masses are also involved in the fitting u-type quark masses. Therefore, the tension is severe leading to a larger total χ^2 value of 69.5079 for the best fit point. In the extended models, one of these two matrices $(M_{120} \text{ in the } \mathbf{10} \oplus \mathbf{120} \oplus \mathbf{120'} \text{ model})$, M'_{10} in the $\mathbf{10} \oplus \mathbf{10'} \oplus \mathbf{120} \text{ model})$ is relaxed and does not involve in fitting u-type quark masses. Therefore, the mild pull for m_s of the minimal model (Eq. (59)) disappear, the tension between the pulls for m_b and m_{τ} is slightly reduced, and other fittings are much improved in these two

Observables	x_i	p_i
m_u	0.398968×10^{-3}	2.4×10^{-5}
m_c	0.199171	5.8×10^{-5}
m_t	60.3555	3.9×10^{-6}
m_d	0.861726×10^{-3}	0.042
m_s	0.0207547	0.51
m_b	0.891622	2.4
m_e	0.440494×10^{-3}	0.20×10^{-2}
m_{μ}	0.0951064	0.42
$m_{ au}$	1.00706	7.3
λ	0.224686	2.7×10^{-4}
A	0.932457	3.2×10^{-5}
$\bar{ ho}$	0.139518	5.2×10^{-5}
$ar\eta$	0.34626	4.7×10^{-5}
Total χ^2	59.298	34

Table 6: The best fit point values for the charged fermion masses (in GeV) and the Wolfenstein parameters of the CKM mixing matrix in the $\mathbf{10} \oplus \mathbf{10'} \oplus \mathbf{120}$ model at the GUT scale. The x_i and p_i columns correspond to the theoretical predicted values and the pulls.

extension. As a consequence, the best total χ^2 values are 59.3001 and 59.2984 in the two extended models respectively. In all these considered models, there is always a tension between the pulls for bottom and tau masses. It is due to the fact that down-type quark and charged lepton masses (M_d, M_e) in these three models are determined mainly by the symmetric matrix M_{10} while the antisymmetric matrices (M_{120}, M'_{120}) are not enough to generate the adequate corrections. Thus, large pulls of opposite sides persist for all three models. Because of this property, even if more additional **10**-plets or **120**-plets Higgs fields of the same kind are introduced, the situation will not be improved.

5 The minimal alternative model with bottom and vectorlike quark mixing

5.1 Bottom and vector-like quark mixing

From the results of data fitting to the alternative models, we can see the tension in the fitting of the bottom and tau masses. In this section we consider a Yukawa interaction between the 3rd generation fermions, 16_3 , and the 16_D representation introduced in the minimal alternative model⁶:

$$\mathcal{L}_{SO(10)} \supset Y_m \mathbf{16}_3 \mathbf{10}_H \mathbf{16}_D. \tag{75}$$

⁶ In general, the vector-like quarks have Yukawa interactions with all three generations of the SM quarks. In this analysis, we assume, for simplicity, that the dominant term is the Yukawa coupling with the third generation while other terms are negligible. This hierarchy is inspired from that in the SM. Such hierarchical picture should be governed by some underlying symmetry and deserves further study in the future.

Assuming that this Yukawa coupling is adequately small, we can neglect its RGE effects below the GUT scale. After the SO(10) symmetry breaking at the GUT scale, the new interaction in Eq. (75) results in the mixing mass term between the bottom quark and the vector-like quark D_L while other heavy states are neglected:

$$\mathcal{L}_{effective} \supset Y_m \alpha_d^1 v_d \bar{b}_R D_L. \tag{76}$$

Due to such mixing, the bottom mass is deviated from that in the non-mixing case by a small amount while other charged fermion masses remain intact.

We parameterize the GUT mass relation for down-type quarks as follows:

$$M_d = M_{10} + M_{120} + M_{mix}, (77)$$

where

$$M_{mix} = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & m_{vl} \end{pmatrix}$$
(78)

is the contribution from the above mass mixing with the vector-like quark.

5.2 Data fitting

Considering the mixing between the bottom and the vector-like quark D_L , we have performed the fitting to the charged fermion masses and mixing. Relevant to this sector, this scenario has 16 free real parameters including 14 free inputs as those in the above minimal alternative model and 2 additional free inputs from the complex number m_{vl} . The best fit point in this case is found to be

$$c_{10} = 40.3765,$$
 (79)

$$c_{120}^u = -31.5752 - 40.1781i, (80)$$

$$c_{120}^e = -3.0042 - 2.67699i, \tag{81}$$

$$M_{10} = \begin{pmatrix} -0.000557118 & 0 & 0 \\ 0 & 0.00866987 & 0 \\ 0 & 0 & -1.48365 \end{pmatrix},$$
(82)

$$M_{120} = \begin{pmatrix} 0 & 0.00032580 - 0.0016479i & 0.0049921 - 0.023637i \\ -0.00032580 + 0.0016479i & 0 & 0.054465 - 0.078530i \\ -0.0049921 + 0.023637i & -0.054465 + 0.078530i & 0 \end{pmatrix},$$
(83)

$$m_{vl} = 2.16619 - 0.450918i. \tag{84}$$

Note that, for the above best fit point, $|m_{vl}| \ll m_t$ implies that the condition $|Y_m| \ll Y_t$ is satisfied (given that $|\alpha_1^d|v_d = \mathcal{O}(100)$ GeV). This ensures the consistency of our assumption that the RGE effects of Y_m on the considered observables are negligible.

In Table 7, the best fit values of the charged fermion masses and the Wolfenstein parameters of the CKM matrix are shown together with the corresponding pulls. In this case, we have obtained a very good value for the total χ^2 that is 0.217. Therefore, the tension in fitting between the bottom and tau masses is resolved.

Observables	x_i	p_i
m_u	0.400616×10^{-3}	8.2×10^{-3}
m_c	0.201931	0.089
m_t	60.3277	0.016
m_d	0.747056×10^{-3}	0.30
m_s	0.016069	0.34
m_b	0.829718	0.017
m_e	0.44133×10^{-3}	0.040
m_{μ}	0.0931455	2.0×10^{-3}
$m_{ au}$	1.58347	8.9×10^{-3}
λ	0.224687	3.0×10^{-3}
A	0.932497	3.3×10^{-3}
$\bar{ ho}$	0.139571	3.3×10^{-3}
$ar\eta$	0.346291	3.1×10^{-3}
Total χ^2	0.216997	

Table 7: The best fit point values for the charged fermion masses (in GeV) and the Wolfenstein parameters of the CKM mixing matrix in the $10 \oplus 120$ model with the Yukawa interaction between the 3rd generation and the vector-like quark D (see Eq. (75)) at the GUT scale. The x_i and p_i columns correspond to the theoretical predicted values and the pulls.

With the good fitting result for the charged fermion sector, it is necessary to verify if the model is capable to predict the realistic neutrino mass scale as obtained from cosmological observation [26]. We find that the Dirac neutrino mass scale given by Eq. (14) is of $\mathcal{O}(m_t)$ or larger for any value of the coefficient c_{120}^n . On the other hand, the Majorana RH-neutrino mass scale is suppressed by a 2-loop factor of about $10^{-5} - 10^{-6}$ compared to the GUT scale due to the Witten mechanism. Therefore, it is impossible to get the correct mass scale of the left handed neutrinos of $\mathcal{O}(0.1)$ eV [26] by the seesaw type I mechanism.

5.3 Two-step symmetry breaking

To ameliorate the problem that seesaw scale through the Witten mechanism is too low, we consider that the GUT symmetry breaking chain occurs at two distinct scales with $SO(10) \rightarrow SU(5)$ at $M_{SO(10)}$, and $SU(5) \rightarrow SU(3)_C \times SU(2)_L \times U(1)_Y$ at $M_G = 1.5 \times 10^{16} \text{ GeV} < M_{SO(10)}$. The SO(10) GUT symmetry breaking scale is a free parameter bounded from above by the (reduced) Planck scale, $M_{SO(10)} \lesssim M_{\text{Planck}} = 2.4 \times 10^{18}$ GeV. In this case, the RH neutrino Majorana mass through the Witten mechanism is found to be

$$M_R = \left(\frac{m_q}{M_W}\right) \epsilon_{10} \left(\frac{\alpha_G}{\pi}\right)^2 M_{SO(10)}.$$
(85)

Note that for $M_{SO(10)} \sim M_{\text{Planck}}$, the RH neutrino scale becomes of $\mathcal{O}(10^{13})$ GeV, resulting in an acceptable LH neutrino mass scale of about $\mathcal{O}(0.1)$ eV.

We repeat the data fitting analysis to the masses and mixings of all fermions including those of the neutrinos, but with $M_{SO(10)}$ being a new free parameter. Since M_G is close to $M_{SO(10)}$, we have neglected the RGE evolutions of the fermion masses and mixings from M_G to $M_{SO(10)}$, and have used the SO(10) GUT mass relations, Eqs. (12)-(15), to match with experimental data at M_G . In our analysis, the radiatively generated RH-neutrino Majorana mass is parameterized as follows

$$M_R = c_{10}^R M_{10}. ag{86}$$

We have obtained the best fit point with the χ^2 value of 7.8198 that is experimentally viable. The parameters corresponding to this point are determined as

$$c_{10} = 40.6978, (87)$$

$$c_{120}^u = -17.6178 - 10.7482i, (88)$$

$$c_{120}^e = -3.151 + 6.33661i, (89)$$

$$c_{120}^n = 47.9034 - 38.7087i, (90)$$

$$c_{10}^R = -3.81633 \times 10^{12} + 3.22151 \times 10^{11} i, \tag{91}$$

$$M_{10} = \begin{pmatrix} -3.9037 \times 10^{-5} & 0 & 0 \\ 0 & 0.0045238 & 0 \\ 0 & 0 & -1.48135 \end{pmatrix},$$
(92)

$$M_{120} = \begin{pmatrix} 0 & 0.0007942 - 0.0005567i & 0.001138 - 1.6085 \times 10^{-6}i \\ -0.0007942 + 0.0005567i & 0 & 0.042495 + 0.034654i \\ -0.001138 + 1.6085 \times 10^{-6}i & -0.042495 - 0.034654i & 0 \end{pmatrix},$$

$$m_{vl} = 2.20452 - 0.402702i. \tag{94}$$

For this best fit point, the RH neutrino Majorana mass is $M_R \approx 5.6 \times 10^{12}$ GeV. Taking $\epsilon_{10} = 0.11$ as an example, the SO(10) GUT breaking scale is found to be $M_{SO(10)} \approx 2.35 \times 10^{18}$ GeV, which is smaller than M_{Planck} , and we neglect gravity effects in our analysis.

Observables	x_i	p_i
m_u	0.412643×10^{-3}	0.068
m_c	0.203016	0.12
m_t	60.309	0.027
m_d	0.20426×10^{-3}	1.5
m_s	0.504091×10^{-2}	2.3
m_b	0.831351	0.080
m_e	0.441872×10^{-3}	0.065
m_{μ}	0.0932523	0.021
$m_{ au}$	1.57706	0.072
λ	0.224687	0.19×10^{-2}
A	0.932724	0.022
$ar{ ho}$	0.139633	0.71×10^{-2}
$ar\eta$	0.346093	0.017
Δm_{21}^2	1.1119×10^{-4}	0.78×10^{-2}
Δm_{31}^2	3.79811×10^{-3}	0.35×10^{-2}
θ_{12}	0.573177	0.026
θ_{23}	0.866723	0.041
$ heta_{13}$	0.150271	2.8×10^{-4}
$\delta_{ m CP}$	-2.49584	2.9×10^{-5}
Total χ^2	7.819	8
$m_{\beta\beta}$	0.22 meV	
Σ	$0.078 { m eV}$	

Table 8: The best fit point values at M_G for the charged fermion masses (in GeV), the Wolfenstein parameters of the CKM mixing matrix, neutrino mass squared differences (in eV^2), and the parameters of the PMNS mixing matrix in the two-step symmetry breaking scenario of the $10\oplus120$ model with the Yukawa interaction between the 3rd generation and the vector-like quark D (see Eq. (75)). The x_i and p_i columns correspond to the theoretical predicted values and the pulls. In the last three rows, the total χ^2 value, the predicted values of the effective Majorana neutrino mass, $m_{\beta\beta}$, and the sum of light neutrino masses, Σ , for the best fit point are presented.

The predicted observables and their pulls for the best fit point are given in Table 8. The largest deviation is 2.3σ corresponding to the strange quark mass. Therefore, a precise determination of the strange quark mass in the near future can test the validity of this model.

For consistency, we take into account the current constraints on the neutrino sector from nuclear physics and astrophysics. The Majorana nature of light neutrinos can be revealed by the neutrinoless double beta decay. The current constraint on the effective Majorana mass reads [27]

$$m_{\beta\beta} \equiv \left| \sum_{i=1}^{3} m_{i}^{\nu} U_{ei}^{2} e^{i\varphi_{i}} \right| \lesssim \mathcal{O}(100) \text{ meV}, \tag{95}$$

where m_i^{ν}, U_{ei} , and φ_i are the light neutrino mass eigenvalues, the PMNS matrix elements, and the Majorana phases, respectively. The effective Majorana mass predicted by our best fit point is found to be $m_{\beta\beta} = 0.22$ meV, that is well below the current upper limit. On the other hand, the sum of light neutrino masses is constrained by the measurement of the Cosmic Microwave Background (CMB) anisotropy [26]:

$$\Sigma = m_1^{\nu} + m_2^{\nu} + m_3^{\nu} \lesssim 0.12 \text{ eV}.$$
(96)

For the best fit point in our scenario, Σ is found to be 0.078 eV, which is consistent with the upper bound.

6 Conclusion

In the alternative SO(10) GUTs, only 10-plets and 120-plets of Higgs fields are involved in generating masses for the SM charged fermions. The RH-neutrinos with vanishing Majorana masses at tree level obtain nonzero ones at quantum two-loop level via the Witten mechanism. The LH-neutrinos acquire tiny masses in charge of neutrino oscillations via the seesaw mechanism. The success of the gauge coupling unification is achieved by introducing the vector-like quarks with the mass scale of $\mathcal{O}(\text{TeV})$. The existence of these vector-like fermions deflects the SM RGE trajectories, resulting in a unique pattern of fermion masses and mixing at the GUT scale. The matching between such a particular effective model at low energies and the alternative SO(10)models has been investigated for the first time in this paper. The experimental data for the charged fermion masses and the CKM matrix are evolved to the GUT scale and have been used to fit the model parameters. In the alternative minimal model with $10 \oplus 120$ Higgs fields, we have found the best fit point among the space of 14 free parameters. The total χ^2 value of this point is 69.5079. Large pulls are observed corresponding to m_s , m_b , and m_{τ} . In two simple extensions to the minimal models with $10 \oplus 120 \oplus 120'$ and $10 \oplus 10' \oplus 120$ of Higgs fields, there are more degrees of freedom with 20 and 24 free parameters, respectively. The results of data fitting are improved with the best χ^2 equals 59.3001 for the former and 59.2984 for the latter. However, due to the GUT mass relation among down-type quarks and charged leptons, the severe tension between the pulls for m_b and m_{τ} still exists. Especially, the tau mass always suffers from large pulls of about 7.3 that rule out all the three models.

We have shown that the tension in the fitting to the bottom and tau masses in the minimal alternative model can be resolved by introducing the new Yukawa interaction between the 3rd generation fermions, $\mathbf{10}_3$, the vector-like fermions, $\mathbf{10}_D$, and the $\mathbf{10}$ -plet of Higgs fields. Neglecting all the heavy fields, this interaction results in the mixing between the bottom quark and the vectorlike quark D. We have found that the best χ^2 value in the data fitting to the charged fermion sector is 0.217. However, the Dirac neutrino mass scale is too large in this scenario such that the 2-loop suppressed Majorana RH-neutrino mass scale could not lead to the LH-neutrino mass scale compatible with its cosmological bound. To ameliorate this problem of the radiatively generated seesaw scale being too small, we have considered the two-step symmetry breaking scenario. Here, the GUT symmetry breaking chain occurs at two distinct scales with $SO(10) \rightarrow SU(5)$ at $M_{SO(10)}$, and $SU(5) \rightarrow SU(3)_C \times SU(2)_L \times U(1)_Y$ at $M_G = 1.5 \times 10^{16}$ GeV. Performing the data fitting analysis in this case, we have found the best fit point with $\chi^2_{\text{best}} = 7.8198$ that is experimentally allowed. For this point, the largest pull of 2.3 comes from the strange quark mass. Thus, a more precise determination of the strange quark mass can test this scenario. We have found the effective Majorana neutrino mass of $m_{\beta\beta} = 0.22$ meV and the sum of light neutrino masses as $\Sigma = 0.078$ eV, which are consistent with the current constraints from the search for the neutrinoless double beta decay and the CMB anisotropy measurement, respectively.

To summarize, in this work, almost all possibilities have been exhausted for the alternative models with $\mathbf{10}_H$ and $\mathbf{120}_H$. The vector-like quarks turn out to be important not only for the success of the gauge coupling unification, but also for the fitting of charged fermion masses and mixing. In the context of the Witten mechanism, the two-step symmetry breaking is necessary for the prediction of observables in both charged and neutral fermion sectors consistent with experimental results.

Acknowledgments

This work is supported in part by JSPS Kakenhi No. 17H01133 (T.F.), the U.S. Department of Energy No. DE-SE0012447 (N.O.), and Vietnam National Foundation for Science and Technology Development (NAFOSTED) No. 103.01-2017.301 (H.M.T.).

References

- [1] R. Slansky, Phys. Rept. **79**, 1 (1981).
- [2] K. S. Babu and R. N. Mohapatra, Phys. Rev. Lett. **70**, 2845 (1993) [hep-ph/9209215];
 D. G. Lee and R. N. Mohapatra, Phys. Rev. D **51**, 1353 (1995) [hep-ph/9406328]; See for a review of the minimal SO(10) model, for instance, T. Fukuyama, Int. J. Mod. Phys. A **28**, 1330008 (2013) [arXiv:1212.3407 [hep-ph]].
- [3] P. Minkowski, Phys. Lett. 67B, 421 (1977); T. Yanagida, Conf. Proc. C 7902131, 95 (1979), KEK-79-18-95. M. Gell-Mann, P. Ramond and R. Slansky, Conf. Proc. C 790927, 315 (1979) [arXiv:1306.4669 [hep-th]]; S. L. Glashow, NATO Sci. Ser. B 61, 687 (1980); R. N. Mohapatra and G. Senjanovic, Phys. Rev. Lett. 44, 912 (1980); R. N. Mohapatra and G. Senjanovic, Phys. Rev. D 23, 165 (1981).
- [4] E. Witten, Phys. Lett. **91B**, 81 (1980).

- [5] T. Fukuyama and N. Okada, Mod. Phys. Lett. A 33, no. 29, 1850167 (2018) [arXiv:1802.06530 [hep-ph]].
- [6] B. Bajc and G. Senjanovic, Phys. Lett. B 610, 80 (2005) [hep-ph/0411193]; B. Bajc and G. Senjanovic, Phys. Rev. Lett. 95, 261804 (2005) [hep-ph/0507169].
- [7] K. Matsuda, Y. Koide and T. Fukuyama, Phys. Rev. D 64, 053015 (2001) [hep-ph/0010026].
- [8] T. Fukuyama, A. Ilakovac, T. Kikuchi, S. Meljanac, N. Okada, J.Math.Phys. 46, 033505 (2005) [hep-ph/0405300].
- [9] D. Chang, T. Fukuyama, Y. Y. Keum, T. Kikuchi and N. Okada, Phys. Rev. D 71, 095002 (2005) [hep-ph/0412011].
- [10] L. Lavoura, H. Kuhbock and W. Grimus, Nucl. Phys. B **754**, 1 (2006) [hep-ph/0603259].
- [11] T. Fukuyama, K. Ichikawa, and Y. Mimura, Phys. Rev. D94 (2016) 075018.
- [12] K. S. Babu, T. Fukuyama, S. Khan and S. Saad, JHEP **1906**, 045 (2019) [arXiv:1812.11695 [hep-ph]].
- [13] T. Ohlsson and M. Pernow, JHEP **1906**, 085 (2019) [arXiv:1903.08241 [hep-ph]].
- [14] I. Gogoladze, B. He and Q. Shafi, Phys. Lett. B 690, 495 (2010) [arXiv:1004.4217 [hep-ph]].
- [15] H. Y. Chen, I. Gogoladze, S. Hu, T. Li and L. Wu, Eur. Phys. J. C 78, no. 1, 26 (2018) [arXiv:1703.07542 [hep-ph]].
- [16] J. M. Cline, Phys. Rev. D 97, no. 1, 015013 (2018) [arXiv:1710.02140 [hep-ph]]; L. Calibbi, A. Crivellin and T. Li, Phys. Rev. D 98, no. 11, 115002 (2018) [arXiv:1709.00692 [hep-ph]]; F. Z. Xu, W. Zhang, J. Li and T. Li, Phys. Rev. D 98, no. 11, 115033 (2018) [arXiv:1809.01472 [hep-ph]]; N. Assad, B. Fornal and B. Grinstein, Phys. Lett. B 777, 324 (2018) [arXiv:1708.06350 [hep-ph]]; B. Fornal, S. A. Gadam and B. Grinstein, Phys. Rev. D 99, no. 5, 055025 (2019) [arXiv:1812.01603 [hep-ph]].
- [17] N. Okada, S. Okada and D. Raut, Phys. Lett. B 780, 422 (2018) [arXiv:1712.05290 [hep-ph]].
- [18] M. E. Machacek and M. T. Vaughn, Nucl. Phys. B 222, 83 (1983); M. E. Machacek and M. T. Vaughn, Nucl. Phys. B 236, 221 (1984); M. E. Machacek and M. T. Vaughn, Nucl. Phys. B 249, 70 (1985); C. Ford, I. Jack and D. R. T. Jones, Nucl. Phys. B 387, 373 (1992) Erratum: [Nucl. Phys. B 504, 551 (1997)] [hep-ph/0111190]; H. Arason, D. J. Castano, B. Keszthelyi, S. Mikaelian, E. J. Piard, P. Ramond and B. D. Wright, Phys. Rev. D 46, 3945 (1992); V. D. Barger, M. S. Berger and P. Ohmann, Phys. Rev. D 47, 1093 (1993) [hep-ph/9209232]; M. x. Luo and Y. Xiao, Phys. Rev. Lett. 90, 011601 (2003) [hep-ph/0207271]; S. Antusch, J. Kersten, M. Lindner and M. Ratz, Nucl. Phys. B 674, 401 (2003) [hep-ph/0305273].
- [19] See for example C. R. Das and M. K. Parida, Eur. Phys. J. C 20, 121 (2001) [hep-ph/0010004], and references therein.
- [20] A. Dueck and W. Rodejohann, JHEP **1309**, 024 (2013) [arXiv:1306.4468 [hep-ph]].

- [21] T. Ohlsson and M. Pernow, JHEP 1811, 028 (2018) [arXiv:1804.04560 [hep-ph]].
- [22] Z. z. Xing, H. Zhang and S. Zhou, Phys. Rev. D 77, 113016 (2008) [arXiv:0712.1419 [hep-ph]];
 Z. z. Xing, H. Zhang and S. Zhou, Phys. Rev. D 86, 013013 (2012) [arXiv:1112.3112 [hep-ph]];
 D. Meloni, T. Ohlsson and S. Riad, JHEP 1412, 052 (2014) [arXiv:1409.3730 [hep-ph]];
 D. Meloni, T. Ohlsson and S. Riad, JHEP 1703, 045 (2017) [arXiv:1612.07973 [hep-ph]];
 I. Esteban, M. C. Gonzalez-Garcia, M. Maltoni, I. Martinez-Soler and T. Schwetz, JHEP 1701, 087 (2017) [arXiv:1611.01514 [hep-ph]]; C. Patrignani et al. [Particle Data Group], Chin. Phys. C 40, no. 10, 100001 (2016); NuFIT3.2, http://www.nu-fit.org/?q=node/166
- [23] I. Esteban, M. C. Gonzalez-Garcia, A. Hernandez-Cabezudo, M. Maltoni and T. Schwetz, JHEP 1901, 106 (2019) [arXiv:1811.05487 [hep-ph]].
- [24] F. Feroz and M. P. Hobson, Mon. Not. Roy. Astron. Soc. 384, 449 (2008) [arXiv:0704.3704 [astro-ph]]; F. Feroz, M. P. Hobson and M. Bridges, Mon. Not. Roy. Astron. Soc. 398, 1601 (2009) [arXiv:0809.3437 [astro-ph]]; F. Feroz, M. P. Hobson, E. Cameron and A. N. Pettitt, arXiv:1306.2144 [astro-ph.IM].
- [25] J. A. Nelder, R. Mead, A Simplex Method for Function Minimization, The Computer Journal, Volume 7, Issue 4, 1 January 1965, Pages 308-313; O'Neill, R. Journal of the Royal Statistical Society, Series C (Applied Statistics) 20, no. 3 (1971), 338-345.
- [26] N. Palanque-Delabrouille et al., JCAP 1511, 011 (2015) [arXiv:1506.05976 [astro-ph.CO]];
 E. Di Valentino, E. Giusarma, O. Mena, A. Melchiorri and J. Silk, Phys. Rev. D 93, no. 8, 083527 (2016) [arXiv:1511.00975 [astro-ph.CO]]; X. Zhang, Phys. Rev. D 93, no. 8, 083011 (2016) [arXiv:1511.02651 [astro-ph.CO]]; A. J. Cuesta, V. Niro and L. Verde, Phys. Dark Univ. 13, 77 (2016) [arXiv:1511.05983 [astro-ph.CO]]; E. Giusarma, M. Gerbino, O. Mena, S. Vagnozzi, S. Ho and K. Freese, Phys. Rev. D 94, no. 8, 083522 (2016) [arXiv:1605.04320 [astro-ph.CO]]; C. Yche, N. Palanque-Delabrouille, J. Baur and H. du Mas des Bourboux, JCAP 1706, 047 (2017) [arXiv:1702.03314 [astro-ph.CO]]; N. Aghanim et al. [Planck Collaboration], arXiv:1807.06209 [astro-ph.CO]; S. Roy Choudhury and S. Choubey, JCAP 1809, no. 09, 017 (2018) [arXiv:1806.10832 [astro-ph.CO]].
- [27] S. I. Alvis *et al.* [Majorana Collaboration], Phys. Rev. C 100, no. 2, 025501 (2019) [arXiv:1902.02299 [nucl-ex]]; M. Agostini *et al.* [GERDA Collaboration], Phys. Rev. Lett. 120, no. 13, 132503 (2018) [arXiv:1803.11100 [nucl-ex]]; C. Alduino *et al.* [CUORE Collaboration], Phys. Rev. Lett. 120, no. 13, 132501 (2018) [arXiv:1710.07988 [nucl-ex]]; G. Anton *et al.* [EXO-200 Collaboration], arXiv:1906.02723 [hep-ex]; A. Gando *et al.* [KamLAND-Zen Collaboration], Phys. Rev. Lett. 117, no. 8, 082503 (2016) Addendum: [Phys. Rev. Lett. 117, no. 10, 109903 (2016)] [arXiv:1605.02889 [hep-ex]].