

Generalised reversible transformations and the inhomogeneous nonlinear Schrödinger equation hierarchy

Sudipta Nandy* and Abhijit Barthakur
Cotton University, Panbazar, Guwahati -781001, India

March 1, 2022

Abstract

Under investigation is the nonlinear Schrödinger equation hierarchies and the reversible transformations. We propose a generalized reversible transformation between the the generalized NLSE hierarchy with focussing and defocussing nonlinearity and the NLSE hierarchy forced with a linear potential term. The corresponding extended concept of classical dark and bright solitons of the forced hierarchy, accelerating due to linear potential as well as due to the dispersion are obtained directly without resolving the nonisospectral inverse scattering problem. We have identified a set of new constraints among the dispersion and the nonlinear coefficients in the inhomogeneous NLSE hierarchy, which are preserved after the transformations. The reversible transformations allow us to encompass inhomogeneous NLS, HNLS and higher order equations belonging to the class of nonisospectral family of inverse scattering problems to the isospectral NLS class of equations and study them under a general mathematical framework. We hope that our analysis provides a mathematical platform to study inhomogeneous NLSEs as well as open up the possibility of new applications in physics.

05.45.Yv, 42.81Dp, 42.65.Tg

Introduction

Possibility of optical soliton in fiber and in bulk medium was first proposed by Hasegawa and Tappert [1]. Two of the most discussed optical solitons are, spatial soliton, that do

*sudipta.nandy@cottonuniversity.ac.in

not spread spatially due to diffraction and temporal soliton, that do not spread temporally due to dispersion[2, 3, 4]. The dynamics of both spatial soliton and temporal soliton are described by the same nonlinear Schrödinger equation (NLSE) except that the spatial parameters are replaced by their temporal counterparts. The applications of NLSE have spread into many other branches of sciences over the past few decades. Reports of new applications of NLSE soliton are coming in every month. NLSE soliton are among the most widely studied concept in nonlinear optics. Temporal soliton, due to its remarkable dispersion-less property is suitable for all optical communication system[37]. On the other hand, in case of spatial soliton the variety of nonlinearity accessible is much broader. A detail account of the spatial soliton and various types of nonlinearity are found in [35, 36, 5]. Langmuir wave soliton in plasma physics is another example, where coherent nonlinear structures associated with electron plasma can be described with the NLSE[6, 7, 8]. Matter wave soliton in Bose-Einstein condensate is yet another example, where the behaviour of large number of particles at near absolute zero temperature, known as Bose-Einstein condensate can be described by Gross-Pitaevski equation [9, 10, 11] which in $1 - D$ is the NLSE. The list continues.

Drawing an analogy with the time independent Schrödinger equation, solitons of the integrable NLSE are sometimes referred to as *autonomous* soliton, that is a soliton solution of NLSE in the absence of an external potential. The NLSE in presence of external potential is an inhomogeneous nonlinear equation and is not integrable in general. With certain potential however, the equation is found to be integrable and admit soliton solution. The soliton solution in presence of such potential is called *nonautonomous*. In 1976 Chen and Liu in [6] introduced inhomogeneity in terms of a "gravitational" like potential to the integrable NLSE, to describe the behaviour of the Langmuir wave in a linearly inhomogeneous plasma. In the same communication Chen and Liu also introduced a symmetry transformation called the *Tappert transformation*, which transform NLSE with linear potential to NLSE without the potential and vice versa. The transformation is very much similar to the transformation introduced by Husimi and Taniuti[12] to transform Schrodinger equation with a linear potential into the free Schrödinger equation. It would be interesting to check whether such symmetries can also be extended to NLSE hierarchies.

The spectral problem associated with the NLSE is well known[13], where the eigen values are time independent(isospectral). The introduction of inhomogeneity in NLSE causes a major change in the spectral problem. In presence of inhomogeneity the eigen values are no longer time independent and they vary linearly with time. Consequently one has to solve the nontrivial nonisospectral inverse scattering problem. However, with the Tappert transformation [6] the NLSE with linear potential can be transformed into NLSE without potential and vice versa. Consequently one may obtain the soliton solution of the nonisospectral integrable models by the reversible transformation from the soliton solutions of corresponding isospectral NLSE, without going through the

rigorous procedure of solving the nonisospectral IST problem.

The transformations are however, nontrivial in nature and hence are applied to only a handful of inhomogeneous problems. Recently in [14] authors considered a "gravitational" like potential to the integrable higher order nonlinear Schrödinger equations (HNLSE) in order to take in to account: (1) the sliding filter method of the noise separation from femto-second pulse train, where the peak frequency of the sliding frequency filter is assumed to be a linear function of time variable; (2) the Raman induced self scattering shift. The resulting forced HNLSE though exactly solvable but is a case of nontrivial nonisospectral inverse scattering problem. They used reversible gauge transformations and obtained the soliton solution from the corresponding soliton solution of isospectral HNLSEs. It is well known that NLSE shares many of the mathematical properties with its hierarchy, infinite conserved quantities for instance, is one of them. The symmetry transformations for NLSE and HNLSE suggest that there might be a general symmetry transformation between force free NLSE hierarchy and forced NLSE hierarchy. The usefulness and potential of such transformations are unquestionable as they provide a bridge between the more nontrivial nonisospectral integrable models and the isospectral integrable models. One obtains the solutions without solving the IST problem.

In this paper we explore a generalized reversible transformation between the the generalized NLSE hierarchy with focussing and defocussing nonlinearity and the generalized NLSE hierarchy forced with a linear potential. We start with the transformation between force free NLSE and forced NLSE in the following section, and then obtain the transformations for the second equation and the third equation of the hierarchy and then propose a general transformation relation between the equations of forced and force free hierarchies. We also present a discussion about the soliton properties, considering bright and dark soliton in the inhomogeneous regime. We conclude with the summary of our results and future possibilities in the subsequent section.

Reversible transformations

Consider the inhomogeneous equation:

$$\mathbf{i}q_t + \sum_{i=1}^{\infty} S_{2i}\{q\} + \mathbf{i}P_{2i+1}\{q\} - V(x, t)q = 0 \quad (1)$$

where, $S_{2i}\{q\}$ and $P_{2i+1}\{q\}$ are respectively the even and odd operators of the NLSE hierarchy. For instance, $S_2\{q\} = \frac{1}{2}D_{20}q_{xx} + \sigma R_{20}(|q|^2q)$ is the NLSE operator. The subscripts (x, t) denote the partial derivatives with respect to x and t respectively. The inhomogeneous term namely, $V(x, t) = 2\lambda(t)x$ is a gravitational like potential [33].

$\lambda(t)$ is the time derivative of the spectral parameter. Note that the role of x and t in NLSE are exchanged when applied to optics. For the spectral parameter, linear in t , $\lambda(t) = \lambda$, is a constant. In absence of the inhomogeneous term the equation represents the NLSE hierarchy generalized for focusing and defocusing nonlinearity. In absence of the last term in eq. 1 the S_{2i} and P_{2i+1} are determined from the infinite conserved charges with appropriate Poisson bracket [37, 16]. It may be noted here that NLSE hierarchy equations with $P_{2i+1} = 0$ are different from the equations with $S_{2i} = 0$ at least from one aspect, that is in the second case the equations are the real modified KdV equation hierarchy. The distinction is expected to appear also in the soliton solutions of forced equations. With $S_2 = \frac{D_{20}}{2} q_{xx} + \sigma R_{20} |q|^2 q$ and $S_{2i}|_{i>1} = 0$, $P_{2i+1} = 0$, eq. 1 reduces to the integrable inhomogeneous NLSE [6] for focusing or defocusing nonlinearity, according to the sign of σ .

$$iq_t + \frac{D_{20}}{2} q_{xx} + \sigma R_{20} |q|^2 q - 2\lambda(t)qx = 0 \quad (2)$$

The ”+ sign” for σ accounts for the focusing nonlinearity and the ”– sign” accounts for the defocusing nonlinearity. In optical applications (x, t) represent the dimensionless time in a comoving frame and the normalized propagation distance respectively. The forced eq. 2 is a case of nonisospectral IST problem and is exactly solvable. An easier alternative approach to obtain the solution is to use Tappert transformation[6]:

$$\begin{aligned} E(T(x, t), Z) &= q(x, t)e^{i\phi(x, t)} \\ \phi(x, t) &= 2\lambda xt + \frac{2}{3}D_{20}\lambda^2 t^3 \\ T(x, t) &= x + D_{20}\lambda t^2 \\ Z &= t \end{aligned} \quad (3)$$

and convert eq. 2 into free NLSE:

$$iE_Z + \frac{1}{2}D_{20}E_{TT} + \sigma R_{20}|E|^2 E = 0 \quad (4)$$

The well known NLSE soliton solution may be obtained through several methods, namely IST method for isospectral problems [13], Hirota method [3], Bačklund transformations [4] etc. The solutions for focusing ($\sigma = +1$) and defocusing ($\sigma = -1$) nonlinearity respectively are:

$$E = \begin{cases} 2\eta\sqrt{\frac{D_{20}}{R_{20}}} \text{sech}(2\eta T + 4D_{20}\eta\kappa Z)e^{(-i2\kappa T + i2D_{20}(\eta^2 - \kappa^2)Z)}; & \text{bright soliton}(\sigma = 1) \\ 2\eta\sqrt{\frac{D_{20}}{R_{20}}} \tanh(2\eta T + 4D_{20}\eta\kappa Z)e^{(-i2\kappa T - i2D_{20}(2\eta^2 + \kappa^2)Z)}; & \text{dark soliton}(\sigma = -1) \end{cases} \quad (5)$$

Finally through reverse transformation:

$$\begin{aligned} q(x(T, Z), t) &= E(T, Z)e^{-i\psi(T, Z)} \\ \psi(T, Z) &= 2\lambda T Z - \frac{4}{3}D_{20}\lambda^2 Z^3 \\ x &= T - D_{20}\lambda Z^2 \\ t &= Z \end{aligned} \quad (6)$$

the soliton solutions of eq. 2 are obtained. The solutions are:

$$q = \begin{cases} 2\eta\sqrt{\frac{D_{20}}{R_{20}}} & sech(2\eta x + 2\eta D_{20}\lambda t^2 + 4D_{20}\eta\kappa t)e^{-i2\kappa x - i2\kappa D_{20}\lambda t^2 + i2D_{20}(\eta^2 - \kappa^2)t} \\ & \times e^{-i(2\lambda x t + \frac{2}{3}D_{20}\lambda^2 t^3)}; \quad \text{bright soliton}(\sigma = 1) \\ 2\eta\sqrt{\frac{D_{20}}{R_{20}}} & tanh(2\eta x + 2\eta D_{20}\lambda t^2 + 4D_{20}\eta\kappa t)e^{-i2\kappa x - i2\kappa D_{20}\lambda t^2 - i2D_{20}(2\eta^2 + \kappa^2)t} \\ & \times e^{-i(2\lambda x t + \frac{2}{3}D_{20}\lambda^2 t^3)}; \quad \text{dark soliton}(\sigma = -1) \end{cases} \quad (7)$$

It may be noted that the reversible transformations(3, 6) do not depend on parameter σ . That is Tappart transformation is independent of type of nonlinearity, focusing or defocusing. The inhomogeneity in the equation introduces many changes in the soliton dynamics, such as in soliton's velocity and phase, which are two of the principle quantities controlling the dynamics of soliton. For instance a dark NLSE soliton with group velocity " $V_g = 2D_{20}\kappa$ " and phase velocity " $V_{ph} = \frac{1}{\kappa}D_{20}(2\eta^2 + \kappa^2)$ " are changed to $2D_{20}\kappa + D_{20}\lambda t$ and $\frac{-D_{20}(2\eta^2 + \kappa^2) + 2D_{20}\kappa\lambda t + D_{20}\lambda^2 t^2 + \lambda x}{\kappa + \lambda t}$ respectively in the inhomogeneous regime. The group velocity and the phase velocity are now linear function of 't', that is the motion of soliton is "accelerated" in the presence of external potential. Further note that the soliton also changes it's colour during motion as it's frequency changes quadratically with time.

The same approach may also be adopted to higher order NLSE with the linear external potential. Next let us consider the second equation of the NLSE hierarchy, and rewrite the eq. 1 with $S_2 = \frac{D_{20}}{2}q_{xx} + \sigma R_{20}|q|^2 q$ and $P_3 = D_{30}q_{xxx} + \sigma R_{30}6|q|^2 q_x + \beta\sigma R_{30}(|q|^2)_x q$,

$$iq_t + \frac{1}{2}D_{20}q_{xx} + \sigma R_{20}(|q|^2 q) + i(D_{30}q_{xxx} + \sigma R_{30}6|q|^2 q_x + \beta\sigma R_{30}(|q|^2)_x q) - 2\lambda qx = 0 \quad (8)$$

We shall refer eq. 8 as FHNLSSE for future reference. In absence of the inhomogeneous term eq. 8 represents three known cases of the hierarchy, namely the complex modified KdV equation(CMKdVE) ($D_{20} = R_{20} = 0$; $\beta = 0$ or 3)[18]; Hirota equation(HE) ($\beta = 0$) [17] and Sasa Satsuma equation(SSE) ($\beta = 3$)[18] provided the parameters D_{20} , D_{30} , R_{20} , R_{30} satisfy the Hirota condition, namely $D_{20}R_{30} = D_{30}R_{20}$. Eq. 8 is important for the description of ultrashort pulse propagation in fiber in presence of the effects such as, third order dispersion, self steepening and stimulated Raman Scattering in an inhomogeneous medium. The potential term, $V(x, t) = 2\lambda(t)x$ in the equation may account for the sliding frequency method of the noise separation from a soliton [15, 19]. Secondly, the inhomogeneous term also describe the change in the frequency of the pulse due to Raman induced delayed nonlinear response [26, 27]. The parameter σ has the same significance as it has in the NLSE (eq. 2).

The transformations of the FHNLSSE (eq. 8) for $\sigma(= +1)$ is given in [14] for three different cases for HNLSE. Here we rewrite the reversible transformations in a generalised form, which would represent transformations for both focusing and defocusing cases

($\sigma = \pm 1$) and would transform eq. 8 into the corresponding force free HNLSE. The transformations are:

$$\begin{aligned} E(T(x, t), Z) &= q(x, t)e^{i\phi(x, t)} \\ \phi(x, t) &= 2\lambda xt + \frac{2}{3}D_{20}\lambda^2 t^3 + 2D_{30}\lambda^3 t^4 \\ T(x, t) &= x + D_{20}\lambda t^2 + 4D_{30}\lambda^2 t^3 \\ Z &= t \end{aligned} \quad (9)$$

and the converted force free equation is:

$$\mathbf{i}E_Z + \frac{1}{2}D_2(Z)E_{TT} + \sigma R_2(z)(|E|^2 E) + \mathbf{i}(D_{30}E_{TTT} + \sigma R_{30}6|E|^2 E_T + \beta\sigma R_{30}(|E|^2)_T E) = 0 \quad (10)$$

where, $D_2(Z) = D_{20} + 12D_{30}\lambda Z$ and $R_2(Z) = R_{20} + 12R_{30}\lambda Z$. Eq. 10 corresponds to free Hirota equation, for $\beta = 0$; free Sasa-Satsuma equation, for $\beta = 3$ and free CMKDV equation for $D_{20} = R_{20} = 0$. Notice that after the transformation the Hirota condition namely, $D_2(Z)R_{30} = D_{30}R_2(Z)$ is still satisfied. At this point it may be worthwhile to mention that the force free Hirota equation and the Sasa Satsuma equation can be converted into CMKDV equation with gauge transformations[17, 18]. Let us consider soliton solution of one of these cases, Hirota equation for instance,

$$E(T, Z) = \begin{cases} 2\eta\sqrt{\frac{D_{30}}{R_{30}}} \text{sech}(2\eta T + 4D_{20}\eta\kappa Z + 8\eta D_{30}(3\kappa^2 - \eta^2)Z + 24D_{30}\eta\kappa\lambda Z^2) \\ \times e^{-2i\kappa T + 2iD_{20}(\eta^2 - \kappa^2)Z - 8i\kappa D_{30}(\kappa^2 - 3\eta^2)Z + 12iD_{30}(\eta^2 - \kappa^2)\lambda Z^2}; \\ \text{bright soliton } (\sigma = 1) \\ 2\eta\sqrt{\frac{D_{30}}{R_{30}}} \tanh(2\eta T + 4D_{20}\eta\kappa Z + 8\eta D_{30}(3\kappa^2 + 2\eta^2)Z + 24D_{30}\eta\kappa\lambda Z^2) \\ \times e^{-i(2\kappa T + 2D_{20}(2\eta^2 + \kappa^2)Z + 8D_{30}\kappa(\kappa^2 + 6\eta^2)Z + 12D_{30}(\kappa^2 + 2\eta^2)\lambda Z^2)}; \\ \text{dark soliton } (\sigma = -1) \end{cases} \quad (11)$$

Now with the following reverse transformations:

$$\begin{aligned} q(x(T, Z), t) &= E(T, Z)e^{i\psi(T, Z)} \\ \psi(T, Z) &= -2\lambda TZ + \frac{4}{3}D_{20}\lambda^2 Z^3 + 6D_{30}\lambda^3 Z^4 \\ x(T, Z) &= T - D_{20}\lambda Z^2 - 4D_{30}\lambda^2 Z^3 \\ t(Z) &= Z \end{aligned} \quad (12)$$

the 1-soliton solution of the forced HNLS equation form eq. 8 is obtained,

$$q(x, t) = \begin{cases} 2\eta\sqrt{\frac{D_{30}}{R_{30}}} \text{sech}(2\eta x + 4D_{20}\eta\kappa t + 8\eta D_{30}(3\kappa^2 - \eta^2)t + 24D_{30}\eta\kappa\lambda t^2 \\ + 2\eta D_{20}\lambda t^2 + 8\eta D_{30}\lambda^2 t^3) e^{-i(2\kappa x + 2\kappa D_{20}\lambda t^2 + 8\kappa D_{30}\lambda^2 t^3 + 8D_{30}\kappa(\kappa^2 - 3\eta^2)t)} \\ \times e^{-i(-2D_{20}(\eta^2 - \kappa^2)t - 12D_{30}(\eta^2 - \kappa^2)\lambda t^2 + 2\lambda x t + \frac{2}{3}D_{20}\lambda^2 t^3 + 2D_{30}\lambda^3 t^4)} \\ \text{bright soliton}(\sigma = 1); \\ 2\eta\sqrt{\frac{D_{30}}{R_{30}}} \tanh(2\eta x + 4D_{20}\eta\kappa t + 8\eta D_{30}(3\kappa^2 + 2\eta^2)t + 24\eta D_{30}\kappa\lambda t^2 \\ + 2\eta D_{20}\lambda t^2 + 8\eta D_{30}\lambda^2 t^3) e^{-i(2\kappa x + 2\kappa D_{20}\lambda t^2 + 8\kappa D_{30}\lambda^2 t^3 + 8\eta D_{30}\kappa(\kappa^2 + 6\eta^2)t)} \\ \times e^{-i(2D_{20}(2\eta^2 + \kappa^2)t + 12D_{30}(\kappa^2 + 2\eta^2)\lambda t^2 + 2\lambda x t + \frac{2}{3}D_{20}\lambda^2 t^3 + 2D_{30}\lambda^3 t^4)} \\ \text{dark soliton}(\sigma = -1); \end{cases} \quad (13)$$

The soliton solutions pair (eq. 13) is the nonautonomous soliton solution of eq. 8 and is interpreted as a coloured (multi wavelength) soliton [25] because its frequency is changing with 't' during propagation. The group velocity of nonautonomous NLS soliton the nonautonomous HNLS solitons are fundamentally different in the sense that in the former case soliton's velocity is reversed after reaching a minimum and in the later case soliton's velocity increases in the same direction after reaching a minimum. Secondly the third order dispersion(TOD) causes an asymmetric broadening of the soliton pulse that is the higher amplitude soliton moves faster, a phenomenon also exhibited by KdV soliton[32] which is because of the presence of TOD term in the equation. The consequence of dependence of the soliton velocity on the amplitude is that the noise(having lower amplitudes) can be filtered out from the ultrashort soliton pulses.

The transformation discussed above may also be extended to NLSE with higher order dispersion and nonlinear terms. In the literature study of such equations is rare, other than a few reported theoretical models [20, 23, 34, 33]. However, recent development of extremely short pulse (attosecond), analysis of higher order terms is gaining momentum again. Effects of the individual terms such as forth order dispersion (FOD) [27], quintic nonlinearity [28, 5, 30] are studied separately in different physical contexts. For instance, in dispersion flattened fibers, it is found that FOD is significant[27] compared to TOD, such equation is not integrable but admit "autosoliton" solution. Recent reports suggest that $CDS_x Se_{1-x}$ doped glasses shows fifth order susceptibility and parabolic form of refractive index[29]. NLSE in presence of either FOD or quintic nonlinearity is also not integrable but there exist solitary waves solutions for these equations too. Theoretical and experimental investigation on higher order derivatives of nonlinear terms are not available in the literature. This is due to the fact the detail analysis on the effects of these terms as well as the collective behaviour of all the terms simultaneously is extremely difficult task at present and require further advancement of both theory and experimental technology. However, to start with theoretically it is always convenient to consider an integrable model, which contains these terms.

Next in our analysis we consider equations having quintic nonlinearity and fourth order dispersion term along with the inhomogeneous term. That is along with S_2 and P_3 , we also consider the term $S_4 = D_{40}q_{xxxx} + R_{40}(6|q|^4q + 2\sigma q^2q_{xx}^* + 4\sigma|q_x|^2q + 6\sigma q_x^2q^* + 8\sigma|q|^2q_{xx}) - 2\lambda(t)qx$ and S_{2i+2} , $P_{2i+1} = 0$ for $i \geq 2$, then eq. 1 reduces to

$$\begin{aligned} & \mathbf{i}q_t + \frac{1}{2}D_{20}q_{xx} + \sigma R_{20}|q|^2q + \mathbf{i}(D_{30}q_{xxx} + \sigma R_{30}|q|^2q_x) + D_{40}q_{xxxx} \\ & + R_{40}(6|q|^4q + 2\sigma q^2q_{xx}^* + 4\sigma|q_x|^2q + 6\sigma q_x^2q^* + 8\sigma|q|^2q_{xx}) - 2\lambda(t)qx = 0 \end{aligned} \quad (14)$$

With $\sigma = 1$, eq.14 with constant coefficients is the Lakshmanan-Porsezian-Daniel equation [20, 21, 22], attached with the force term. In future we refer eq.14 as forced Lakshmanan-Porsezian-Daniel equation (FLPDE). Notice that if $D_{20} = D_{30} = R_{20} = R_{30} = 0$, then FLPDE reduces to the following equation:

$$\begin{aligned} & \mathbf{i}q_t + D_{40}q_{xxxx} \\ & + R_{40}(6|q|^4q + 2\sigma q^2q_{xx}^* + 4\sigma|q_x|^2q + 6\sigma q_x^2q^* + 8\sigma|q|^2q_{xx}) - 2\lambda(t)qx = 0 \end{aligned} \quad (15)$$

Note that this equation is also referred as the Lakshmanan-Porsezian-Daniel equation and is a hierarchy of the CMKdVE. We consider additional two equations, which admit soliton solution. First one is FLPDE (eq. 14) in absence of the NLS operator (S_2):

$$\begin{aligned} & \mathbf{i}q_t + \mathbf{i}(D_{30}q_{xxx} + \sigma R_{30}|q|^2q_x) + D_{40}q_{xxxx} \\ & + R_{40}(6|q|^4q + 2\sigma q^2q_{xx}^* + 4\sigma|q_x|^2q + 6\sigma q_x^2q^* + 8\sigma|q|^2q_{xx}) - 2\lambda(t)qx = 0 \end{aligned} \quad (16)$$

and the second one is FLPDE (eq. 14) in absence of the Hirota operator (P_3):

$$\begin{aligned} & \mathbf{i}q_t + \frac{1}{2}D_{20}q_{xx} + \sigma R_{20}|q|^2q + D_{40}q_{xxxx} \\ & + R_{40}(6|q|^4q + 2\sigma q^2q_{xx}^* + 4\sigma|q_x|^2q + 6\sigma q_x^2q^* + 8\sigma|q|^2q_{xx}) - 2\lambda(t)qx = 0 \end{aligned} \quad (17)$$

Note that the structures of integrable hierarchies are highly rigid. Each of the hierarchies of the NLSE contains dispersion and nonlinear terms in a fixed ratio. However, the nice mathematical symmetries among the hierarchy makes it convenient to study the complex higher order equations.

Let us introduce the following gauge transformation and transformation of variables:

$$\begin{aligned} E(x(T, Z), t) &= q(x, t)e^{i\phi(x, t)} \\ \phi(x, t) &= 2\lambda xt + \frac{2}{3}D_{20}\lambda^2t^3 + 2D_{30}\lambda^3t^4 - \frac{16}{5}D_{40}\lambda^4t^5 \\ T(x, t) &= x + D_{20}\lambda t^2 + 4D_{30}\lambda^2t^3 - 8D_{40}\lambda^3t^4 \\ Z(t) &= t \end{aligned} \quad (18)$$

which converts the forced eq. 14 into the following a free higher order NLSEs with variable coefficients,

$$\begin{aligned} \mathbf{i}E_Z + D_{40}E_{TTTT} + \sigma R_{40}(6|E|^4E + |E_T|^2E + 6E_T^2E^* + 2E^2E_{TT}^* + 8|E|^2E_{TT}) \\ + \mathbf{i}(D_3(Z)E_{TTT} + 6\sigma R_3(Z)|E|^2E) + \frac{D_2(Z)}{2}E_{TT} + R_2(Z)\sigma|E|^2E = 0 \end{aligned} \quad (19)$$

where $D_2(Z) = D_{20} + 12D_{30}\lambda Z - 48D_{40}\lambda^2 Z^2$, $D_3(Z) = D_{30} - D_{40}8\lambda Z$, $R_2(Z) = R_{20} + 12D_{30}\lambda Z - D_{40}48\lambda^2 Z^2$, and $R_3(Z) = R_{30} - D_{40}8\lambda Z$.

Note that the transformation eqs.18 with $D_{20} = D_{30} = R_{20} = R_{30} = 0$, that is,

$$\begin{aligned} E(x(T, Z), t) &= q(x, t)e^{i\phi(x, t)} \\ \phi(T, Z) &= 2\lambda x t - \frac{16}{5}D_{40}\lambda^4 t^5 \\ T(x, t) &= x - 8D_{40}\lambda^3 t^4 \\ Z(t) &= t \end{aligned} \quad (20)$$

convert eq. 15 into eq. 19 with $D_2(Z) = -D_{40}48\lambda^2 Z^2$, $D_3(Z) = -D_{40}8\lambda Z$, $R_2(Z) = -48D_{40}\lambda^2 Z^2$, and $R_3(Z) = -D_{40}8\lambda Z$.

The transformations from eq. 16 to eq. 19 is accomplished from eq. 18 with $D_{20} = 0$. Consequently in the transformed equation, $D_2(Z) = 12D_{30}\lambda Z - 48D_{40}\lambda^2 Z^2$, $D_3(Z) = D_{30} - 8D_{40}\lambda Z$, $R_2(Z) = 12R_{30}\lambda Z - 48D_{40}\lambda^2 Z^2$, and $R_3(Z) = R_{30} - 8D_{40}\lambda Z$.

similarly the transformations from eq. 17 to eq. 19 is accomplished by using eqs. 18 with $D_{30} = 0$. Consequently, $D_2(Z) = D_{20} - 48D_{40}\lambda^2 Z^2$, $D_3(Z) = -8D_{40}\lambda Z$, $R_2(Z) = R_{20} - 48R_{40}\lambda^2 Z^2$, and $R_3(Z) = -8R_{40}\lambda Z$.

Further note that similar to the Hirota constraint [24] in HNLS equations, we identify a set of new constraints among the dispersion and nonlinear coefficients, namely, $D_{20}R_{40} = D_{40}R_{20}$; $D_{30}R_{40} = D_{40}R_{30}$ and $D_{20}R_{30} = D_{30}R_{20}$ which are preserved under the transformations eqs. 18. Similar constraints are expected from the equations as one goes higher in the hierarchy. Number of constraints increases as nC_2 , where n is the order of the equation in the NLSE hierarchy.

Let us consider 1- soliton solutions of the force free eq. 19 :

$$E(T, Z) = \begin{cases} 2\eta\sqrt{\frac{D_4}{R_4}}\text{sech}(2\eta T + 4D_{20}\eta\kappa Z + 8D_{30}\eta(3\kappa^2 - \eta^2)Z + 24D_{30}\eta\kappa\lambda Z^2 \\ -64D_{40}\eta\kappa(\kappa^2 - \eta^2)Z - 32D_{40}\eta(3\kappa^2 - \eta^2)\lambda Z^2 - 64D_{40}\eta\kappa\lambda^2 Z^3) \\ \times e^{-i2\kappa T + i2D_{20}(\eta^2 - \kappa^2)Z - i8D_{30}\kappa(\kappa^2 - 3\eta^2)Z + i12D_{30}(\eta^2 - \kappa^2)\lambda Z^2} \\ \times e^{i16D_{40}(\kappa^4 - 6\kappa^2\eta^2 + \eta^4)Z - i32D_{40}(\eta^2 - \kappa^2)\lambda^2 Z^3 + i32D_{40}\kappa(\kappa^2 - 3\eta^2)\lambda Z^2}; \\ \text{bright soliton } (\sigma = 1) \\ 2\eta\sqrt{\frac{D_4}{R_4}}\tanh(2\eta T + 4D_{20}\eta\kappa Z + 8D_{30}\eta(3\kappa^2 + 2\eta^2)Z + 24D_{30}\eta\kappa\lambda Z^2 \\ -64D_{40}\eta\kappa\lambda^2 Z^3 - 64D_{40}\eta\kappa(\kappa^2 + 2\eta^2)Z - 32D_{40}\eta(3\kappa^2 + 2\eta^2)\lambda Z^2) \\ \times e^{-i2\kappa T - i2D_{20}(2\eta^2 + \kappa^2)Z - i8D_{30}\kappa(\kappa^2 + 6\eta^2)Z + i12D_{30}(2\eta^2 + \kappa^2)\lambda Z^2} \\ \times e^{i16D_{40}(\kappa^4 + 12\kappa^2\eta^2 + 6\eta^4)Z + i32D_{40}(2\eta^2 + \kappa^2)\lambda^2 Z^3 + i32D_{40}\kappa(\kappa^2 + 6\eta^2)\lambda Z^2} \\ \text{dark soliton } (\sigma = -1); \end{cases} \quad (21)$$

we shall obtain the soliton solution of one of the solutions of eq. (14-17). For instance, the soliton solution of the first of these equations and the reverse transformations for this equation is:

$$\begin{aligned} q(x(T, Z), t) &= E(T, Z)e^{-i\phi(T, Z)} \\ \phi(T, Z) &= 2\lambda TZ - \frac{4}{3}D_{20}\lambda^2 Z^3 - 4D_{30}\lambda^3 Z^4 + \frac{64}{5}D_{40}\lambda^4 Z^5 \\ x(T, Z) &= T - D_{20}\lambda Z^2 - 4D_{30}\lambda^2 Z^3 + 8D_{40}\lambda^3 Z^4 \\ t(Z) &= Z \end{aligned} \quad (22)$$

Thus the soliton solution of the forced eq. 14 is :

$$q(x, t) = \begin{cases} 2\eta\sqrt{\frac{D_4}{R_4}}\text{sech}(2\eta x + 4D_{20}\eta\kappa t + 2\eta D_{20}\lambda t^2 + 8D_{30}\eta(3\kappa^2 - \eta^2)t \\ +8\eta D_{30}\lambda^2 t^3 + 24D_{30}\eta\kappa\lambda t^2 - 64D_{40}\eta\kappa(\kappa^2 - \eta^2)t - 64D_{40}\eta\kappa\lambda^2 t^3 \\ -32D_{40}\eta(3\kappa^2 - \eta^2)\lambda t^2 - 16\eta D_{40}\lambda^3 t^4)e^{-i2\kappa x - i2D_{20}\kappa\lambda t^2 + i2D_{20}(\eta^2 - \kappa^2)t} \\ \times e^{-i\frac{2}{3}D_{20}\lambda^2 t^3 - i8D_{30}\kappa(\kappa^2 - 3\eta^2)t - 4iD_{30}\lambda^3 t^4 - 8iD_{30}\kappa\lambda^2 t^3 + i16D_{40}(\kappa^4 - 6\kappa^2\eta^2 + \eta^4)t} \\ \times e^{i16D_{40}\kappa\lambda^3 t^4 - i32D_{40}(\eta^2 - \kappa^2)\lambda^2 t^3 + i32D_{40}\kappa(\kappa^2 - 3\eta^2)\lambda t^2 + i\frac{16}{5}D_{40}\lambda^4 t^5 - i2\lambda x t}; \\ \text{bright soliton } (\sigma = 1) \\ 2\eta\sqrt{\frac{D_4}{R_4}}\tanh(2\eta x + 4D_{20}\eta\kappa t + 2D_{20}\eta\lambda t^2 + 8D_{30}\eta(3\kappa^2 + 2\eta^2)t \\ +8D_{30}\eta\lambda^2 t^3 + 24D_{30}\eta\kappa\lambda t^2 - 64D_{40}\eta\kappa(\kappa^2 + 2\eta^2)t - 64D_{40}\eta\kappa\lambda^2 t^3 \\ -32D_{40}\eta(3\kappa^2 + 2\eta^2)\lambda t^2 - 16\eta D_{40}\lambda^3 t^4) \\ \times e^{-i2\kappa x - i\frac{2}{3}D_{20}\lambda^2 t^3 - i2D_{20}(2\eta^2 + \kappa^2)t} \\ \times e^{-i2D_{20}\kappa\lambda t^2 - i8D_{30}\kappa(\kappa^2 + 6\eta^2)t - i8D_{30}\kappa\lambda^2 t^3 - 4iD_{30}\lambda^3 t^4 + i16D_{40}(\kappa^4 + 12\kappa^2\eta^2 + 6\eta^4)t} \\ \times e^{i32D_{40}(2\eta^2 + \kappa^2)\lambda^2 t^3 + i16D_{40}\kappa\lambda^3 t^4 + i32D_{40}\kappa(\kappa^2 + 6\eta^2)\lambda t^2 - i2\lambda x t + i\frac{16}{5}D_{40}\lambda^4 t^5}; \\ \text{dark soliton } (\sigma = -1); \end{cases} \quad (23)$$

Solution of the rest of the eqs. (15-17) may similarly be obtained with the corresponding transformations discussed above. The analysis can be carried forward to even higher equations, such as equations in [23, 31] along with the potential term. We notice a few observations in the above analysis. First, solitons of odd order hierarchy have different space-time profile in general than the soliton of even order hierarchy, an observation also noticed in [16] but for force free NLSE hierarchy. In the odd order hierarchy soliton the *Sech* argument contains even order polynomial in λt where as the even order hierarchy the the argument contains an odd order polynomial in λt . Secondly, as $\lambda \rightarrow 0$ the nonautonomous solitons reduces to their autonomous counterparts. Thirdly each of the higher order equations are reducible to the lower order equations by removing the higher order parameters, for instance, the FLPDE reduces to forced HNLSE with $D_{40} = R_{40} = 0$ and the corresponding nonautonomous solitons reduce to the autonomous soliton of HNLSE. Similarly forced HNLSE reduces to the forced NLSE with $D_{30} = R_{30} = 0$. Above observations show that our analysis is consistent with the earlier analysis, with force free NLSE hierarchy. The simple form of reversible transformations suggests that the generalized reversible transformations between the force free NLSE hierarchy and the forced NLSE hierarchy follow a general form:

$$E(T(x, t), Z(t)) = q(x, t)e^{i\phi(x, t)} \quad (24)$$

Where the phase $\phi(x, t)$ and the variable $T(x, t)$ are expressed as a series:

$$\begin{aligned} \phi &= 2t(\lambda x + a_0(\lambda t)^2 + a_1(\lambda t)^3 + \dots); \\ T &= x + v(t) t; \\ v(t) &= b_0(\lambda t) + b_1(\lambda t)^2 + \dots; \\ Z &= t \end{aligned} \quad (25)$$

where $a_1, a_2, a_3, \dots, b_0, b_1, b_2, \dots$ are constants, to be determined.

For instance for forced NLSE:

$a_0 = \frac{1}{3}D_{20}$, $b_0 = D_{20}$ and $a_i, b_i = 0$ for $i \geq 1$;

for forced CMKdVE:

$a_1 = D_{30}$, $b_1 = 4D_{30}$ and $a_i, b_i = 0$ for $i \neq 1$;

for forced Hirota equation and Sasa Satsuma equation:

$a_0 = \frac{1}{3}D_{20}$, $a_1 = D_{30}$, $b_0 = D_{20}$, $b_1 = 4D_{30}$, $a_i, b_i = 0$ for $i \geq 2$;

for FLPDE without NLS and Hirota operator:

$a_2 = \frac{-8}{5}D_{40}$, $b_2 = -8D_{40}$ and $a_i, b_i = 0$ for $i \neq 2$;

forced HNLSE with cubic, quintic nonlinearities and fourth order dispersion, FLPDE eq. (14)

$a_0 = \frac{1}{3}D_{20}$, $a_1 = D_{30}$, $a_{20} = \frac{-8}{5}D_{40}$, $b_0 = D_{20}$, $b_1 = 4D_{30}$, $b_2 = -8D_{40}$, and $a_i, b_i = 0$ for $i \geq 2$;

HNLSE with cubic, quintic nonlinearities and fifth order dispersion

$a_0 = \frac{1}{3}D_{20}$, $a_1 = D_{30}$, $a_2 = \frac{-8}{5}D_{40}$, $a_3 = \frac{-16}{5}D_{50}$, $b_0 = D_{20}$, $b_1 = 4D_{30}$, $b_2 = -8D_{40}$, $b_{30} = -16D_{50}$ and $a_i, b_i = 0$ for $i \geq 3$;

and so on.




Figure 1: Evolution of an autonomous NLSE bright(a) and dark(b) solitons at constant velocity with $D_{20} = R_{20} = 1$, $\lambda = 0$ and soliton parameters $\eta = 0.5$ and $\kappa = 0.9$




Figure 2: Evolution of a nonautonomous NLSE bright(a) and dark(b) solitons with $D_{20} = R_{20} = 1$, $\lambda = 1.2$ and soliton parameters $\eta = 0.5$ and $\kappa = 0.9$ (13)




Figure 3: Evolution of a nonautonomous HNLSE bright(a) and dark(b) solitons with $D_{20} = R_{20} = 1$, $D_{30} = R_{30} = 1$, $\lambda = 1.2$ and soliton parameters $\eta = 0.5$ and $\kappa = 0.9$




Figure 4: Evolution of a nonautonomous FLPDE bright(a) and dark(b) solitons with $D_{20} = 3$, $R_{20} = 6$, $D_{30} = 1$, $R_{30} = 2$, $D_{40} = 2$, $R_{40} = 4$, $\lambda = 1.2$ and soliton parameters $\eta = 0.5$ and $\kappa = 0.9$

Figures 1 shows the evolution of typical constant coefficient NLSE autonomous bright and dark soliton having a constant velocity. Velocity of solitons of rest of the equations in the hierarchy also remain constant with time as the soliton evolves. Figures (2 - 4) present evolution of a bright soliton (a) and a dark soliton (b) of first three equations of the NLSE hierarchy in presence of a linear potential. In the plot absolute value of the soliton's amplitude are plotted, suppressing the phase part. Notice that in presence of the potential, NLSE nonautonomous solitons are "accelerated" (Fig. 2), that is the soliton group velocity is a function of t , $v_g = 2D_{20}\kappa + D_{20}\lambda t$. In this case the GVD parameter along with the linear potential term account for the acceleration. On the other hand nonautonomous HNLSE solitons are subjected to "accelerations" proportional to t^2 and t^3 , appearing in the argument of envelope functions *sech* and *tanh*. Moving one up in the hierarchy, the figure shows that the nonautonomous solitons are subjected to "accelerations" proportional to t^2 , t^3 and t^4 . Thus the argument of envelope functions turn out to be a polynomial in t , the degree of the which increases with the increase in the equation order in the hierarchy. The "accelerations" also become a polynomial in t . and its degree increases as the order of the hierarchy. Note that the group velocity of nonautonomous soliton of even order NLSE hierarchy are fundamentally different from the nonautonomous soliton of odd order NLSE hierarchy, that is in the former case soliton's velocity is reversed after reaching the minima and in the later case soliton's velocity increases in the same direction after reaching the minima. The number of minima increases with the order of hierarchy, see Figures (2 - 4). The inhomogeneous term in the NLSE hierarchy makes frequency, phase velocity also dependent on the time.

1 Conclusion

We propose a generalized reversible transformation between the the generalized NLSE hierarchy and the generalized forced NLSE hierarchy with a linear potential term. The reversible transformations allow to encompass inhomogeneous NLS, HNLS and higher order equations belonging to the class of nonisospectral family of inverse scattering problems into the isospectral NLS class of equations and study them under the general mathematical framework. The corresponding extended concept of classical dark and bright solitons, accelerating in the linear potential of the forced hierarchy are obtained directly without resolving the nonisospectral inverse scattering problem. The transformation is not restricted to the case of dark and bright soliton only and may also be extended to breather and other types of solitons. Soliton properties of the forced higher order NLSE family of equations are changed and the parameters such as phase velocity, group velocity, frequency change with propagation and depends on time polynomially. We introduce a set of constraints for the FLPDE which are preserved after the transformations. Similar constraints are expected as one consider higher order equations. It has been noticed that the the odd order inhomogeneous NLSE hierarchy

are fundamentally different from the even order inhomogeneous NLSE hierarchy. It would be interesting to extend the analysis to the coupled systems, that is Manakov equation hierarchy, which would be our future investigation. We hope that our analysis open up the possibility of new applications in physics.

References

- [1] Hasegawa A, Tappert F. Appl Phys Lett 1973;23:142.
- [2] Zakharov VE, Shabat AB. Sov Phys JEPT 1972;34:62.
- [3] Hietarinta J. J Math Phys 1988;29:628.
- [4] Matveev B, Salle MA. Darboux transformation and solitons. Springer-Verlag Berlin, Heidelberg; 1991.
- [5] Biswas A, Konar S. Introduction to nonkerr law optical solitons, Chapman & Hall/CRC Applied Mathematics & Nonlinear Science; 2006.
- [6] Chen HH, Liu CS. Phys. Rev. Lett. 1976;37:693.
- [7] Antipov SV, Nezlin MV, Snezhkin EN, Trubnikov AS. Sov Phys JETP 1978;47:506.
- [8] Zhen HL, Tian B, Wang YF, Liu DY. Phys Plasmas 2015;22: 032307.
- [9] Serkin VN, Hasegawa A, Belayeva TL. Phys Rev Lett 2004;92:199401.
- [10] Raju TS, Panigrahi PK, Porsezian K. Phys Rev E 2005; 71:026608;
- [11] Atre R, Panigrahi PK, Agarwal GS. Phys Rev E 2006;73:056611.
- [12] Husimi K. Misellanea in elementary quantum mechanics I Prog Theor Phys 1953;9:238.
- [13] Ablowitz MJ, Kaup DJ, Newell AC, Segur H. Phys Rev Lett 1973;31:125
- [14] Serkin VN. Belyaeva TL. Optik 2018;158:1-1584.
- [15] Hasegawa A, Kodama Y. Soliton in optical communications, Oxford University Press, New York, 1995.
- [16] Chawdury A, Krolkowski W. Phys. Rev. E 2017;95:062226.
- [17] Hirota R. J Math Phys 1973;14:805.

-
- [18] Sasa N, Satsuma J. 1991;60:409.
- [19] Burtsev S, Kaup DJ. J Opt Soc Am B 1997;14:627.
- [20] Porsezian, Phys Rev E 1997;55:3785.
- [21] Ankiewicz A, Wang Y, Wabnitz S, Akhmediev N. Phys Rev E 2014;89:012907.
- [22] Ankiewicz A, Akhmediev N. Phys. Lett A 2014;378:358.
- [23] Yang C, Liu W, Zhou Q, Mihalache Q, Malomed BA. <https://doi.org/10.1007/s11071-018-4569-3>
- [24] Dianov EM, Karasik AYa, Mamyshev PV, Prokhorov AM, Serkin VN, Stel'Makh MF, Fomichev AA. JEPT Lett 1985; 41:294.
- [25] Maimistov AI, Basharov AM. Nonlinear optical waves, Kluwer Academic Publishers, Dordrecht; 1999.
- [26] Serkin VN, Hasegawa A, Belyaeva TL. Phys Rev Lett 2007;98:074102.
- [27] Agrawal GP, Nonlinear Fiber Optics. Academic Press, London; 1995.
- [28] Akhmediev N, Soto-Crespo JM, Town G. Phys Rev E 2001; 63: 056602.
- [29] A. Biswas. Optics Commun 2003;216:427.
- [30] Komarov A, Leblond H, Sanchez F. Phys Rev E 2005; 72: 2, 025604(R).
- [31] Chen SS, Tian B, Chai J. Wu XY, Du Z. Waves in Random and Complex Media DOI:10.1080/17455030.2018.1516053.
- [32] Das A. Integrable Models. World Scientific, Singapore; 1989.
- [33] Serkin VN. Balyaeva TL. Optik 2018;175:17.
- [34] Dai CQ, Wang YY, Fan Y, Yu DG. Nonlinear Dyn <https://doi.org/10.1007/s11071-018-4130-4>.
- [35] Segev M. Optical and Quantum Electronics. 1998;30:503.
- [36] Kivshar YS. Agrawal GP. Optical Solitons. Academic Press, London 2003.
- [37] Hasegawa A, Matsumoto M. Optical Solitons in Fibers. Springer-Verlag Berlin 2003.