## **ON HIGHER** $\{g_n, h_n\}$ -DERIVATIONS

AMIN HOSSEINI\* AND NADEEM UR REHMAN

ABSTRACT. In this article, we introduce the concepts of higher  $\{g_n, h_n\}$ -derivation and Jordan higher  $\{g_n, h_n\}$ -derivation, and then we give a characterization of higher  $\{g_n, h_n\}$ -derivations in terms of  $\{g, h\}$ -derivations. Using this result, we prove that every Jordan higher  $\{g_n, h_n\}$ -derivation on a semiprime algebra is a higher  $\{g_n, h_n\}$ -derivation.

## 1. INTRODUCTION AND PRELIMINARIES

Recently, in 2016 Brešar [1] introduced the notion of  $\{g, h\}$ -derivation. Let  $\mathcal{A}$  be an algebra over a field  $\mathbb{F}$  with  $char(\mathbb{F}) \neq 2$ , and let  $f, g, h : \mathcal{A} \to \mathcal{A}$  be linear maps. We say that f is a  $\{g, h\}$ -derivation if f(ab) = g(a)b + ah(b) = h(a)b + ag(b) for all  $a, b \in \mathcal{A}$ , and f is called a Jordan  $\{g, h\}$ -derivation if  $f(a \circ b) = g(a) \circ b + a \circ h(b)$  for all  $a, b \in \mathcal{A}$ , where  $a \circ b = ab + ba$ . We call  $a \circ b$  the Jordan product of a and b. It is evident that  $a \circ b = b \circ a$  for all  $a, b \in \mathcal{A}$ . The notion of a Jordan  $\{g, h\}$ -derivation is a generalization of what is called a Jordan generalized derivation in [8]. Recall that a linear mapping  $f : \mathcal{A} \to \mathcal{A}$  is called a Jordan generalized derivation if there exists a linear mapping  $d : \mathcal{A} \to \mathcal{A}$  such that  $f(a \circ b) = f(a) \circ b + a \circ d(b)$  for all  $a, b \in \mathcal{A}$ , where d is called an associated linear map of f. It is clear that  $f(a \circ b) = d(a) \circ b + a \circ f(b)$  for all  $a, b \in \mathcal{A}$ . Obviously, the definition of a Jordan generalized derivation. For more details in this regard, see [1, 8] and references therein.

As an important result, Brešar [1, Theorem 4.3] established that every Jordan  $\{g, h\}$ -derivation of a semiprime algebra  $\mathcal{A}$  is a  $\{g, h\}$ -derivation. He also showed that every Jordan  $\{g, h\}$ -derivation of the tensor product of a semiprime and a commutative algebra is a  $\{g, h\}$ -derivation. Obviously, every  $\{g, h\}$ -derivation is a Jordan  $\{g, h\}$ -derivation, but the converse is in general not true. For instance in this regard see [1, Example 2.1].

In this study, we introduce the concept of a higher  $\{g_n, h_n\}$ -derivation, and then we characterize it on algebras. Throughout this paper,  $\mathcal{A}$  denotes an algebra over a field of characteristic zero, and I denotes the identity mapping on  $\mathcal{A}$ . Let f be a  $\{g, h\}$ -derivation on an algebra  $\mathcal{A}$ . An easy induction argument implies that  $f^n(ab) = \sum_{k=0}^n \binom{n}{k} g^{n-k}(a)h^k(b) = \sum_{k=0}^n \binom{n}{k} h^{n-k}(a)g^k(b)$  (Leibniz rule) for each

Date: Received: xxxxx; Revised: yyyyyy; Accepted: zzzzz.

<sup>\*</sup> Corresponding author.

<sup>2010</sup> Mathematics Subject Classification. 47B47; 16W10.

Key words and phrases. derivation;  $\{g,h\}$ -derivation; higher  $\{g_n,h_n\}$ -derivation; Jordan higher  $\{g_n,h_n\}$ -derivation.

 $a, b \in \mathcal{A}$  and each non-negative integer n, where  $f^0 = g^0 = h^0 = I$ . If we define the sequences  $\{f_n\}, \{g_n\}$  and  $\{h_n\}$  of linear mappings on  $\mathcal{A}$  by  $f_0 = g_0 = h_0 = I$ , and  $f_n = \frac{f^n}{n!}, g_n = \frac{g^n}{n!}$  and  $h_n = \frac{h^n}{n!}$ , then it follows from the Leibniz rule that  $f_n$ 's,  $g_n$ 's and  $h_n$ 's satisfy

$$f_n(ab) = \sum_{k=0}^n g_{n-k}(a)h_k(b) = \sum_{k=0}^n h_{n-k}(a)g_k(b),$$
(1.1)

for each  $a, b \in \mathcal{A}$  and each non-negative integer n. This is our motivation to consider the sequences  $\{f_n\}$ ,  $\{g_n\}$  and  $\{h_n\}$  of linear mappings on an algebra  $\mathcal{A}$  satisfying (1.1). A sequence  $\{f_n\}$  of linear mappings on  $\mathcal{A}$  is called a higher  $\{g_n, h_n\}$ -derivation if there exist two sequences  $\{g_n\}$  and  $\{h_n\}$  of linear mappings on  $\mathcal{A}$  satisfying  $f_n(ab) = \sum_{k=0}^n g_{n-k}(a)h_k(b) = \sum_{k=0}^n h_{n-k}(a)g_k(b)$  for any  $a, b \in \mathcal{A}$ and any non-negative integer n. Additionally, a sequence  $\{f_n\}$  of linear mappings on  $\mathcal{A}$  is called a Jordan higher  $\{g_n, h_n\}$ -derivation if

$$f_n(a \circ b) = \sum_{k=0}^n g_{n-k}(a) \circ h_k(b),$$
(1.2)

for each  $a, b \in \mathcal{A}$  and each non-negative integer n. Notice that if  $\{f_n\}$  is a higher  $\{f_n, f_n\}$ -derivation (resp. Jordan higher  $\{f_n, f_n\}$ -derivation), then it is an ordinary higher derivation (resp. Jordan higher derivation). We know that if f is a  $\{g, h\}$ -derivation, then  $\{f_n = \frac{f^n}{n!}\}$  is a higher  $\{g_n, h_n\}$ -derivation, where  $g_n = \frac{g^n}{n!}$ ,  $h_n = \frac{h^n}{n!}$  and  $f_0 = g_0 = h_0 = I$ . We call this kind of higher  $\{g_n, h_n\}$ -derivation an ordinary higher  $\{g_n, h_n\}$ -derivation, but this is not the only example of a higher  $\{g_n, h_n\}$ -derivation.

In 2010, Miravaziri [9] characterized all higher derivations on an algebra  $\mathcal{A}$  in terms of the derivations on  $\mathcal{A}$ . In this article, by getting idea from [9], our aim is to characterize all higher  $\{g_n, h_n\}$ -derivations on an algebra  $\mathcal{A}$  in terms of the  $\{g, h\}$ -derivations on  $\mathcal{A}$ . Indeed, we show that each higher  $\{g_n, h_n\}$ -derivation is a combination of compositions of  $\{g, h\}$ -derivations. As the main result of this article, we prove that if  $\{f_n\}$  is a higher  $\{g_n, h_n\}$ -derivation on an algebra  $\mathcal{A}$  with  $f_0 = g_0 = h_0 = I$ , then there is a sequence  $\{F_n\}$  of  $\{G_n, H_n\}$ -derivations on  $\mathcal{A}$  such that

$$\begin{cases} f_n = \sum_{i=1}^n \left( \sum_{\sum_{j=1}^i r_j = n} \left( \prod_{j=1}^i \frac{1}{r_j + \ldots + r_i} \right) F_{r_1} \ldots F_{r_i} \right), \\ g_n = \sum_{i=1}^n \left( \sum_{\sum_{j=1}^i r_j = n} \left( \prod_{j=1}^i \frac{1}{r_j + \ldots + r_i} \right) G_{r_1} \ldots G_{r_i} \right), \\ h_n = \sum_{i=1}^n \left( \sum_{\sum_{j=1}^i r_j = n} \left( \prod_{j=1}^i \frac{1}{r_j + \ldots + r_i} \right) H_{r_1} \ldots H_{r_i} \right), \end{cases}$$

where the inner summation is taken over all positive integers  $r_j$  with  $\sum_{i=1}^{i} r_j = n$ . The importance of this result is to transfer the problems such as the characterization of Jordan higher  $\{g_n, h_n\}$ -derivations on semiprime algebras and automatic continuity of higher  $\{g_n, h_n\}$ -derivations into the same problems concerning  $\{g,h\}$ -derivations. Let  $\mathcal{A}$  be an algebra, D be the set of all higher derivations  $\{d_n\}_{n=0,1,\dots}$  on  $\mathcal{A}$  with  $d_0 = I$  and  $\Delta$  be the set of all sequences  $\{\delta_n\}_{n=0,1,\dots}$ of derivations on  $\mathcal{A}$  with  $\delta_0 = 0$ . As an application of the main result of this article, we investigate Jordan higher  $\{g_n, h_n\}$ -derivations on algebras. It is a classical question in which algebras (and rings) a Jordan derivation is necessarily a derivation. Let us give a brief background in this issue. In 1957, Herstein [7] achieved a result which asserts any Jordan derivation on a prime ring of characteristic different from two is a derivation. A brief proof of Herstein's result can be found in [3]. In 1975, Cusack [5] generalized Herstein's result to 2-torsion free semiprime rings (see also [2] for an alternative proof). Moreover, Vukman [12] investigated generalized Jordan derivations on semiprime rings and he proved that every generalized Jordan derivation of a 2-torsion free semiprime ring is a generalized derivation. Recently, the first name author along with Ajda Fošner [6] have studied the same problem for  $(\sigma, \tau)$ -derivations from a C<sup>\*</sup>-algebra  $\mathcal{A}$  into a Banach  $\mathcal{A}$ -module  $\mathcal{M}$ . In this paper, we show that if  $\{f_n\}$  is a Jordan higher  $\{g_n, h_n\}$ -derivation of a semiprime algebra  $\mathcal{A}$  with  $f_0 = g_0 = h_0 = I$ , then it is a higher  $\{g_n, h_n\}$ -derivation.

## 2. CHARACTERIZATION OF HIGHER $\{g_n, h_n\}$ -DERIVATIONS ON ALGEBRAS

Throughout the article,  $\mathcal{A}$  denotes an algebra over a field of characteristic zero, and I is the identity mapping on  $\mathcal{A}$ . Let  $f, g, h : \mathcal{A} \to \mathcal{A}$  be linear maps. We say that f is a  $\{g, h\}$ -derivation if f(ab) = g(a)b + ah(b) = h(a)b + ag(b) for all  $a, b \in \mathcal{A}$ , and f is called a Jordan  $\{g, h\}$ -derivation if  $f(a \circ b) = g(a) \circ b + a \circ h(b)$ for all  $a, b \in \mathcal{A}$ , where  $a \circ b = ab + ba$ .

We begin with the following definition.

**Definition 2.1.** A sequence  $\{f_n\}$  of linear mappings on  $\mathcal{A}$  is called a higher  $\{g_n, h_n\}$ -derivation if there are two sequences  $\{g_n\}$  and  $\{h_n\}$  of linear mappings on  $\mathcal{A}$  such that  $f_n(ab) = \sum_{k=0}^n g_{n-k}(a)h_k(b) = \sum_{k=0}^n h_{n-k}(a)g_k(b)$  for each  $a, b \in \mathcal{A}$  and each non-negative integer n.

We begin our results with the following lemma which will be used extensively to prove the main theorem of this article. The following lemma has been motivated by [9].

**Lemma 2.2.** Let  $\{f_n\}$  be a higher  $\{g_n, h_n\}$ -derivation on an algebra  $\mathcal{A}$  with  $f_0 = g_0 = h_0 = I$ . Then there is a sequence  $\{F_n\}$  of  $\{G_n, H_n\}$ -derivations on  $\mathcal{A}$  such

that

$$\begin{cases} (n+1)f_{n+1} = \sum_{k=0}^{n} F_{k+1}f_{n-k}, \\ (n+1)g_{n+1} = \sum_{k=0}^{n} G_{k+1}g_{n-k}, \\ (n+1)h_{n+1} = \sum_{k=0}^{n} H_{k+1}h_{n-k} \end{cases}$$

for each non-negative integer n.

*Proof.* Using induction on n, we prove this lemma. Let n = 0. We know that  $f_1(ab) = g_1(a)b + ah_1(b) = h_1(a)b + ag_1(b)$  for all  $a, b \in \mathcal{A}$ . Thus, if  $F_1 = f_1, G_1 = g_1$  and  $H_1 = h_1$ , then  $F_1$  is a  $\{G_1, H_1\}$ -derivation on  $\mathcal{A}$  and further,  $(0+1)f_{0+1} = \sum_{k=0}^{0} F_{k+1}f_{0-k}, (0+1)g_{0+1} = \sum_{k=0}^{0} G_{k+1}g_{0-k}$  and  $(0+1)h_{0+1} = \sum_{k=0}^{0} H_{k+1}h_{0-k}$ . As induction assumption, suppose that  $F_k$  is a  $\{G_k, H_k\}$ -derivation for any  $k \leq n$  and further

$$\begin{cases} (r+1)f_{r+1} = \sum_{k=0}^{r} F_{k+1}f_{r-k}, \\ (r+1)g_{r+1} = \sum_{k=0}^{r} G_{k+1}g_{r-k}, \\ (r+1)h_{r+1} = \sum_{k=0}^{r} H_{k+1}h_{r-k} \end{cases}$$

for r = 0, 1, ..., n - 1. Put  $F_{n+1} = (n+1)f_{n+1} - \sum_{k=0}^{n-1} F_{k+1}f_{n-k}$ ,  $G_{n+1} = (n+1)g_{n+1} - \sum_{k=0}^{n-1} G_{k+1}g_{n-k}$  and  $H_{n+1} = (n+1)h_{n+1} - \sum_{k=0}^{n-1} H_{k+1}h_{n-k}$ . Our next task is to show that  $F_{n+1}$  is a  $\{G_{n+1}, H_{n+1}\}$ -derivation on  $\mathcal{A}$ . For  $a, b \in \mathcal{A}$ , we have

$$F_{n+1}(ab) = (n+1)f_{n+1}(ab) - \sum_{k=0}^{n-1} F_{k+1}f_{n-k}(ab)$$
$$= (n+1)\sum_{k=0}^{n+1} g_k(a)h_{n+1-k}(b) - \sum_{k=0}^{n-1} F_{k+1}\Big(\sum_{l=0}^{n-k} g_l(a)h_{n-k-l}(b)\Big).$$

So, we have

$$F_{n+1}(ab) = \sum_{k=0}^{n+1} (n+1)g_k(a)h_{n+1-k}(b) - \sum_{k=0}^{n-1} F_{k+1}\Big(\sum_{l=0}^{n-k} g_l(a)h_{n-k-l}(b)\Big)$$
$$= \sum_{k=0}^{n+1} (k+n+1-k)g_k(a)h_{n+1-k}(b) - \sum_{k=0}^{n-1} F_{k+1}\Big(\sum_{l=0}^{n-k} g_l(a)h_{n-k-l}(b)\Big).$$

Since  $F_k$  is a  $\{G_k, H_k\}$ -derivation for each k = 1, 2, ..., n,

$$F_{n+1}(ab) = \sum_{k=0}^{n+1} kg_k(a)h_{n+1-k}(b) + \sum_{k=0}^{n+1} g_k(a)(n+1-k)h_{n+1-k}(b) - \sum_{k=0}^{n-1} \sum_{l=0}^{n-k} \left[ G_{k+1}(g_l(a))h_{n-k-l}(b) + g_l(a)H_{k+1}(h_{n-k-l}(b)) \right].$$

4

Letting

$$G = \sum_{k=0}^{n+1} kg_k(a)h_{n+1-k}(b) - \sum_{k=0}^{n-1} \sum_{l=0}^{n-k} G_{k+1}(g_l(a))h_{n-k-l}(b),$$

$$H = \sum_{k=0}^{n+1} g_k(a)(n+1-k)h_{n+1-k}(b) - \sum_{k=0}^{n-1} \sum_{l=0}^{n-k} g_l(a)H_{k+l}(h_{n-k-l}(b)),$$

we have  $F_{n+1}(ab) = G + H$ . Here, we compute G and H. In the summation  $\sum_{k=0}^{n-1} \sum_{l=0}^{n-k}$ , we have  $0 \le k+l \le n$  and  $k \ne n$ . Thus if we put r = k+l, then we can write it as the form  $\sum_{r=0}^{n} \sum_{k+l=r,k\ne n}$ . Putting l = r - k, we find that

$$G = \sum_{k=0}^{n+1} kg_k(a)h_{n+1-k}(b) - \sum_{r=0}^n \sum_{0 \le k \le r, k \ne n} G_{k+1}(g_{r-k}(a))h_{n-r}(b)$$
  
=  $\sum_{k=0}^{n+1} kg_k(a)h_{n+1-k}(b) - \sum_{r=0}^{n-1} \sum_{k=0}^r G_{k+1}(g_{r-k}(a))h_{n-r}(b) - \sum_{k=0}^{n-1} G_{k+1}(g_{n-k}(a))b.$ 

It means that

$$G + \sum_{k=0}^{n-1} G_{k+1}(g_{n-k}(a))b = \sum_{k=0}^{n+1} kg_k(a)h_{n+1-k}(b) - \sum_{r=0}^{n-1} \sum_{k=0}^r G_{k+1}(g_{r-k}(a))h_{n-r}(b).$$

Putting r + 1 instead of k in the first summation of above, we have

$$G + \sum_{k=0}^{n-1} G_{k+1}(g_{n-k}(a))b$$
  
=  $\sum_{r=0}^{n} (r+1)g_{r+1}(a)h_{n-r}(b) - \sum_{r=0}^{n-1} \sum_{k=0}^{r} G_{k+1}(g_{r-k}(a))h_{n-r}(b)$   
=  $\sum_{r=0}^{n-1} \left[ (r+1)g_{r+1}(a) - \sum_{k=0}^{r} G_{k+1}(g_{r-k}(a)) \right] h_{n-r}(b) + (n+1)g_{n+1}(a)b.$ 

According to the induction hypothesis,  $(r+1)g_{r+1}(a) = \sum_{k=0}^{r} G_{k+1}(g_{r-k}(a))$  for r=0,...,n-1. So, it is obtained that

$$G = \left[ (n+1)g_{n+1}(a) - \sum_{k=0}^{n-1} G_{k+1}(g_{n-k}(a)) \right] b = G_{n+1}(a)b.$$

Like above, we achieve that

$$H = a \Big[ (n+1)h_{n+1}(b) - \sum_{k=0}^{n-1} H_{k+1} \big( h_{n-k}(b) \big) \Big] = a H_{n+1}(b).$$

Therefore, we have  $F_{n+1}(ab) = G + H = G_{n+1}(a)b + aH_{n+1}(b)$ . In the next step, we will show that  $F_{n+1}(ab) = H_{n+1}(a)b + aG_{n+1}(b)$ . We have

$$F_{n+1}(ab) = (n+1)f_{n+1}(ab) - \sum_{k=0}^{n-1} F_{k+1}f_{n-k}(ab)$$
$$= (n+1)\sum_{k=0}^{n+1} h_k(a)g_{n+1-k}(b) - \sum_{k=0}^{n-1} F_{k+1}\Big(\sum_{l=0}^{n-k} h_l(a)g_{n-k-l}(b)\Big).$$

So,

$$F_{n+1}(ab) = \sum_{k=0}^{n+1} (n+1)h_k(a)g_{n+1-k}(b) - \sum_{k=0}^{n-1} F_{k+1}\left(\sum_{l=0}^{n-k} h_l(a)g_{n-k-l}(b)\right)$$
$$= \sum_{k=0}^{n+1} (k+n+1-k)h_k(a)g_{n+1-k}(b) - \sum_{k=0}^{n-1} F_{k+1}\left(\sum_{l=0}^{n-k} h_l(a)g_{n-k-l}(b)\right).$$

Since  $F_k$  is a  $\{G_k, H_k\}$ -derivation for each k = 1, 2, ..., n,

$$F_{n+1}(ab) = \sum_{k=0}^{n+1} kh_k(a)g_{n+1-k}(b) + \sum_{k=0}^{n+1} h_k(a)(n+1-k)g_{n+1-k}(b) - \sum_{k=0}^{n-1} \sum_{l=0}^{n-k} \Big[ H_{k+1}(h_l(a))g_{n-k-l}(b) + h_l(a)G_{k+1}(g_{n-k-l}(b)) \Big].$$

Letting

$$\mathfrak{G} = \sum_{k=0}^{n+1} kh_k(a)g_{n+1-k}(b) - \sum_{k=0}^{n-1} \sum_{l=0}^{n-k} H_{k+1}(h_l(a))g_{n-k-l}(b),$$

$$\mathfrak{H} = \sum_{k=0}^{n+1} h_k(a)(n+1-k)g_{n+1-k}(b) - \sum_{k=0}^{n-1} \sum_{l=0}^{n-k} h_l(a)G_{k+l}(g_{n-k-l}(b)),$$

we have  $F_{n+1}(ab) = \mathfrak{G} + \mathfrak{H}$ . The next step is to compute  $\mathfrak{G}$  and  $\mathfrak{H}$ . In the summation  $\sum_{k=0}^{n-1} \sum_{l=0}^{n-k}$ , we have  $0 \leq k+l \leq n$  and  $k \neq n$ . Thus if we put r = k+l, then we can write it as the form  $\sum_{r=0}^{n} \sum_{k+l=r,k\neq n}$ . Putting l = r-k, we find that

$$\mathfrak{G} = \sum_{k=0}^{n+1} kh_k(a)g_{n+1-k}(b) - \sum_{r=0}^n \sum_{0 \le k \le r, k \ne n} H_{k+1}(h_{r-k}(a))g_{n-r}(b)$$
$$= \sum_{k=0}^{n+1} kh_k(a)g_{n+1-k}(b) - \sum_{r=0}^{n-1} \sum_{k=0}^r H_{k+1}(h_{r-k}(a))g_{n-r}(b) - \sum_{k=0}^{n-1} H_{k+1}(h_{n-k}(a))b.$$

It means that

$$\mathfrak{G} + \sum_{k=0}^{n-1} H_{k+1}(h_{n-k}(a))b = \sum_{k=0}^{n+1} kh_k(a)g_{n+1-k}(b) - \sum_{r=0}^{n-1} \sum_{k=0}^r H_{k+1}(h_{r-k}(a))g_{n-r}(b).$$

Putting r + 1 instead of k in the first summation of above, we have

$$\mathfrak{G} + \sum_{k=0}^{n-1} H_{k+1}(h_{n-k}(a))b$$
  
=  $\sum_{r=0}^{n} (r+1)h_{r+1}(a)g_{n-r}(b) - \sum_{r=0}^{n-1} \sum_{k=0}^{r} H_{k+1}(h_{r-k}(a))g_{n-r}(b)$   
=  $\sum_{r=0}^{n-1} \left[ (r+1)h_{r+1}(a) - \sum_{k=0}^{r} H_{k+1}(h_{r-k}(a)) \right] g_{n-r}(b) + (n+1)h_{n+1}(a)b.$ 

According to the induction hypothesis,  $(r+1)h_{r+1}(a) = \sum_{k=0}^{r} H_{k+1}(h_{r-k}(a))$  for r = 0, ..., n-1. So, it is obtained that

$$\mathfrak{G} = \left[ (n+1)h_{n+1}(a) - \sum_{k=0}^{n-1} H_{k+1}(h_{n-k}(a)) \right] b = H_{n+1}(a)b.$$

By a reasoning like above, we get that

$$\mathfrak{H} = a \Big[ (n+1)g_{n+1}(b) - \sum_{k=0}^{n-1} G_{k+1} \big( g_{n-k}(b) \big) \Big] = a G_{n+1}(b).$$

Therefore, we have  $F_{n+1}(ab) = \mathfrak{G} + \mathfrak{H} = H_{n+1}(a)b + aG_{n+1}(b)$ . Consequently,  $F_{n+1}$  is a  $\{G_{n+1}, H_{n+1}\}$ -derivation and the proof is complete.  $\Box$ 

**Example 2.3.** Using Lemma 2.2, the first five terms of a higher  $\{g_n, h_n\}$ -derivation  $\{f_n\}$  are

$$\begin{split} f_0 &= I, \\ f_1 &= F_1, \\ 2f_2 &= F_1 f_1 + F_2 f_0 = F_1 F_1 + F_2, \\ f_2 &= \frac{1}{2} F_1^2 + \frac{1}{2} F_2, \\ 3f_3 &= F_1 f_2 + F_2 f_1 + F_3 f_0 = F_1 (\frac{1}{2} F_1^2 + \frac{1}{2} F_2) + F_2 F_1 + F_3, \\ f_3 &= \frac{1}{6} F_1^3 + \frac{1}{6} F_1 F_2 + \frac{1}{3} F_2 F_1 + \frac{1}{3} F_3, \\ 4f_4 &= F_1 f_3 + F_2 f_2 + F_3 f_1 + F_4 f_0 \\ &= F_1 (\frac{1}{6} F_1^3 + \frac{1}{6} F_1 F_2 + \frac{1}{3} F_2 F_1 + \frac{1}{3} F_3) + F_2 (\frac{1}{2} F_1^2 + \frac{1}{2} F_2) + F_3 F_1 + F_4, \\ f_4 &= \frac{1}{24} F_1^4 + \frac{1}{24} F_1^2 F_2 + \frac{1}{12} F_1 F_2 F_1 + \frac{1}{12} F_1 F_3 + \frac{1}{8} F_2 F_1^2 + \frac{1}{8} F_2^2 + \frac{1}{4} F_3 F_1 + \frac{1}{4} F_4. \end{split}$$

Now the main result of this paper reads as follows:

**Theorem 2.4.** Let  $\{f_n\}$  be a higher  $\{g_n, h_n\}$ -derivation on an algebra  $\mathcal{A}$  with  $f_0 = g_0 = h_0 = I$ . Then there is a sequence  $\{F_n\}$  of  $\{G_n, H_n\}$ -derivations on  $\mathcal{A}$  such that

$$\begin{cases} f_n = \sum_{i=1}^n \left( \sum_{\sum_{j=1}^i r_j = n} \left( \prod_{j=1}^i \frac{1}{r_j + \dots + r_i} \right) F_{r_1} \dots F_{r_i} \right), \\\\ g_n = \sum_{i=1}^n \left( \sum_{\sum_{j=1}^i r_j = n} \left( \prod_{j=1}^i \frac{1}{r_j + \dots + r_i} \right) G_{r_1} \dots G_{r_i} \right), \\\\ h_n = \sum_{i=1}^n \left( \sum_{\sum_{j=1}^i r_j = n} \left( \prod_{j=1}^i \frac{1}{r_j + \dots + r_i} \right) H_{r_1} \dots H_{r_i} \right), \end{cases}$$

where the inner summation is taken over all positive integers  $r_j$  with  $\sum_{j=1}^{i} r_j = n$ .

*Proof.* First, we show that if  $f_n$ ,  $g_n$  and  $h_n$  are of the above form, then they satisfy the recursive relations of Lemma 2.2. Simplifying the notation, we put  $a_{r_1,\ldots,r_i} = \prod_{j=1}^{i} \frac{1}{r_j+\ldots+r_i}$ . Note that if  $r_1 + \ldots + r_i = n+1$ , then  $(n+1)a_{r_1,\ldots,r_i} = a_{r_2,\ldots,r_i}$ . Furthermore,  $a_{n+1} = \frac{1}{n+1}$ . According to the aforementioned assumptions, we have

$$f_{n+1} = \sum_{i=2}^{n+1} \left( \sum_{\substack{\sum_{j=1}^{i} r_j = n+1 \\ \sum_{j=1}^{i} r_j = n+1}} a_{r_1,\dots,r_i} F_{r_1} \dots F_{r_i} \right) + a_{n+1} F_{n+1}$$
$$= \sum_{i=2}^{n+1} \left( \sum_{\substack{\sum_{j=1}^{i} r_j = n+1 \\ \sum_{j=1}^{i} r_j = n+1}} a_{r_1,\dots,r_i} F_{r_1} \dots F_{r_i} \right) + \frac{F_{n+1}}{n+1}.$$

So,

$$(n+1)f_{n+1} = \sum_{i=2}^{n+1} \left( \sum_{\substack{\sum_{j=1}^{i} r_j = n+1 \\ \sum_{j=1}^{i} r_j = n+1}} (n+1)a_{r_1,\dots,r_i}F_{r_1}\dots F_{r_i} \right) + F_{n+1}$$

$$= \sum_{i=2}^{n+1} \left( \sum_{\substack{\sum_{j=1}^{i} r_j = n+1 \\ \sum_{j=2}^{i} r_j = n+1}} F_{r_1} \sum_{\substack{\sum_{j=2}^{i} r_j = n+1-r_1}} a_{r_2,\dots,r_i}F_{r_2}\dots F_{r_i} \right) + F_{n+1}$$

$$= \sum_{r_1=1}^{n} F_{r_1} \sum_{i=2}^{n-(r_1-1)} \left( \sum_{\substack{\sum_{j=2}^{i} r_j = n-(r_1-1) \\ \sum_{j=2}^{i} r_j = n-(r_1-1)}} a_{r_2,\dots,r_i}F_{r_2}\dots F_{r_i} \right) + F_{n+1}$$

$$= \sum_{r_1=1}^{n} F_{r_1}f_{n-(r_1-1)} + F_{n+1}$$

$$= \sum_{k=0}^{n} F_{k+1}f_{n-k}.$$

Using a reasoning like above, we get that

$$\begin{cases} (n+1)g_{n+1} = \sum_{k=0}^{n} G_{k+1}g_{n-k}, \\ (n+1)h_{n+1} = \sum_{k=0}^{n} H_{k+1}h_{n-k}, \end{cases}$$

for each non-negative integer *n*. Putting n + 1 = m, we find that  $mf_m = \sum_{k=0}^{m-1} F_{k+1}f_{m-1-k} = \sum_{k=0}^{m-2} F_{k+1}f_{m-1-k} + F_m$ , and consequently

$$F_m = mf_m - \sum_{k=0}^{m-2} F_{k+1}f_{m-1-k}.$$

Similarly, we have

$$\begin{cases} G_m = mg_m - \sum_{k=0}^{m-2} G_{k+1}g_{m-1-k}, \\ H_m = mh_m - \sum_{k=0}^{m-2} H_{k+1}h_{m-1-k}. \end{cases}$$

Therefore, we can define  $F_n, G_n, H_n : \mathcal{A} \to \mathcal{A}$  by  $F_0 = G_0 = H_0 = 0$  and

$$F_n = nf_n - \sum_{k=0}^{n-2} F_{k+1}f_{n-1-k},$$

$$G_n = ng_n - \sum_{k=0}^{n-2} G_{k+1}g_{n-1-k},$$

$$H_n = nh_n - \sum_{k=0}^{n-2} H_{k+1}h_{n-1-k},$$

for each positive integer n. It follows from Lemma 2.2 that  $\{F_n\}$  is a sequence of  $\{G_n, H_n\}$ -derivations. In addition, we prove that if  $f_n$ ,  $g_n$  and  $h_n$  are of the form

$$\begin{cases} (n+1)f_{n+1} = \sum_{k=0}^{n} F_{k+1}f_{n-k}, \\ (n+1)g_{n+1} = \sum_{k=0}^{n} G_{k+1}g_{n-k}, \\ (n+1)h_{n+1} = \sum_{k=0}^{n} H_{k+1}h_{n-k}, \end{cases}$$

where  $\{F_n\}$  is a sequence of  $\{G_n, H_n\}$ -derivations, then  $\{f_n\}$  is a higher  $\{g_n, h_n\}$ derivation on  $\mathcal{A}$  with  $f_0 = g_0 = h_0 = I$ . To see this, we use induction on n. For n = 0, we have  $f_0(ab) = ab = g_0(a)h_0(b) = h_0(a)g_0(b)$ . As the inductive hypothesis, assume that

$$f_k(ab) = \sum_{i=0}^k g_i(a)h_{k-i}(b) = \sum_{i=0}^k h_i(a)g_{k-i}(b) \text{ for } k \le n.$$

Therefore, we have

$$(n+1)f_{n+1}(ab) = \sum_{k=0}^{n} F_{k+1}f_{n-k}(ab)$$
  
=  $\sum_{k=0}^{n} F_{k+1} \sum_{i=0}^{n-k} g_i(a)h_{n-k-i}(b)$   
=  $\sum_{i=0}^{n} \left(\sum_{k=0}^{n-i} G_{k+1}g_{n-k-i}(a)\right)h_i(b) + \sum_{i=0}^{n} g_i(a)\left(\sum_{k=0}^{n-i} H_{k+1}h_{n-k-i}(b)\right).$ 

According to the above-mentioned recursive relations, we continue the previous expressions as follows:

$$(n+1)f_{n+1}(ab) = \sum_{i=0}^{n} (n-i+1)g_{n-i+1}(a)h_i(b) + \sum_{i=0}^{n} g_i(a)(n-i+1)h_{n-i+1}(b)$$
$$= \sum_{i=1}^{n+1} ig_i(a)h_{n+1-i}(b) + \sum_{i=0}^{n} (n-i+1)g_i(a)h_{n+1-i}(b)$$
$$= (n+1)\sum_{i=0}^{n+1} g_i(a)h_{n+1-i}(b),$$

which means that  $f_{n+1}(ab) = \sum_{i=0}^{n+1} g_i(a)h_{n+1-i}(b)$ . By an argument like above, we can obtain that  $f_{n+1}(ab) = \sum_{i=0}^{n+1} h_i(a)g_{n+1-i}(b)$ . Thus,  $\{f_n\}$  is a higher  $\{g_n, h_n\}$ -derivation on  $\mathcal{A}$  which is characterized by the sequence  $\{F_n\}$  of  $\{G_n, H_n\}$ derivations. This completes the proof.  $\Box$ 

In the next example, using the above theorem, we characterize term  $f_4$  of a higher  $\{g_n, h_n\}$ -derivation  $\{f_n\}$ .

**Example 2.5.** We compute the coefficients  $a_{r_1,...,r_i}$  for the case n = 4. First, note that 4 = 1 + 3 = 3 + 1 = 2 + 2 = 1 + 1 + 2 = 1 + 2 + 1 = 2 + 1 + 1 = 1 + 1 + 1 + 1.

10

Based on the definition of  $a_{r_1,\ldots,r_i}$  we have

$$a_{4} = \frac{1}{4},$$

$$a_{1,3} = \frac{1}{1+3} \cdot \frac{1}{3} = \frac{1}{12},$$

$$a_{3,1} = \frac{1}{3+1} \cdot \frac{1}{1} = \frac{1}{4},$$

$$a_{2,2} = \frac{1}{2+2} \cdot \frac{1}{2} = \frac{1}{8},$$

$$a_{1,1,2} = \frac{1}{1+1+2} \cdot \frac{1}{1+2} \cdot \frac{1}{2} = \frac{1}{24},$$

$$a_{1,2,1} = \frac{1}{1+2+1} \cdot \frac{1}{2+1} \cdot \frac{1}{1} = \frac{1}{12},$$

$$a_{2,1,1} = \frac{1}{2+1+1} \cdot \frac{1}{1+1} \cdot \frac{1}{1} = \frac{1}{8},$$

$$a_{1,1,1,1} = \frac{1}{1+1+1+1} \cdot \frac{1}{1+1+1} \cdot \frac{1}{1+1+1} \cdot \frac{1}{1} = \frac{1}{24}.$$

Therefore,  $f_4$  is characterized as follows:

$$\begin{split} f_4 = & \frac{1}{4}F_4 + \frac{1}{12}F_1F_3 + \frac{1}{4}F_3F_1 + \frac{1}{8}F_2F_2 + \frac{1}{24}F_1F_1F_2 + \frac{1}{12}F_1F_2F_1 \\ & + \frac{1}{8}F_2F_1F_1 + \frac{1}{24}F_1F_1F_1F_1. \end{split}$$

In the following there are some immediate consequences of the above theorem. Before it, recall that a sequence  $\{f_n\}$  of linear mappings on  $\mathcal{A}$  is called a Jordan higher  $\{g_n, h_n\}$ -derivation if there exist two sequences  $\{g_n\}$  and  $\{h_n\}$  of linear mappings on  $\mathcal{A}$  such that  $f_n(a \circ b) = \sum_{k=0}^n g_{n-k}(a) \circ h_k(b)$  holds for each  $a, b \in \mathcal{A}$ and each non-negative integer n. Since the Jordan product is commutative, we have

$$f_n(a \circ b) = f_n(b \circ a) = \sum_{k=0}^n g_{n-k}(b) \circ h_k(a)$$
$$= \sum_{k=0}^n g_k(b) \circ h_{n-k}(a)$$
$$= \sum_{k=0}^n h_{n-k}(a) \circ g_k(b).$$

So, it is observed that if  $\{f_n\}$  is a Jordan higher  $\{g_n, h_n\}$ -derivation, then

$$f_n(a \circ b) = \sum_{k=0}^n g_{n-k}(a) \circ h_k(b) = \sum_{k=0}^n h_{n-k}(a) \circ g_k(b),$$

for all  $a, b \in \mathcal{A}$ .

**Corollary 2.6.** Let  $\{f_n\}$  be a Jordan higher  $\{g_n, h_n\}$ -derivation on a semiprime algebra  $\mathcal{A}$  with  $f_0 = g_0 = h_0 = I$ . Then  $\{f_n\}$  is a higher  $\{g_n, h_n\}$ -derivation.

*Proof.* Using the proof of Theorem 2.4, we can show that if  $\{f_n\}$  is a Jordan higher  $\{g_n, h_n\}$ -derivation on an algebra  $\mathcal{A}$  with  $f_0 = g_0 = h_0 = I$ , then there exists a sequence  $\{F_n\}$  of Jordan  $\{G_n, H_n\}$ -derivations on  $\mathcal{A}$  such that

$$f_n = \sum_{i=1}^n \Big(\sum_{\sum_{j=1}^i r_j = n} \Big(\prod_{j=1}^i \frac{1}{r_j + \dots + r_i}\Big)F_{r_1}\dots F_{r_i}\Big),$$

where the inner summation is taken over all positive integers  $r_j$  with  $\sum_{j=1}^{i} r_j = n$ . Since  $\mathcal{A}$  is a semiprime algebra, [1, Theorem 4.3] proves the corollary.  $\Box$ 

Remark 2.7. We know that the notion of a Jordan  $\{g, h\}$ -derivation is a generalization of a Jordan generalized derivation (see Introduction). A sequence  $\{f_n\}$  of linear mappings on an algebra  $\mathcal{A}$  is called a *Jordan higher generalized* derivation if there exists a sequence  $\{d_n\}$  of linear mappings on  $\mathcal{A}$  such that  $f_n(a \circ b) = \sum_{k=0}^n f_{n-k}(a) \circ d_k(b)$  for all  $a, b \in \mathcal{A}$ . Obviously, if  $\{f_n\}$  is a Jordan higher generalized derivation associated with a sequence  $\{d_n\}$  of linear mappings on  $\mathcal{A}$ , then it is a Jordan higher  $\{f_n, d_n\}$ -derivation. So, Corollary 2.6 is also valid for Jordan higher generalized derivations.

**Theorem 2.8.** If  $\{f_n\}_{n=0,1,\ldots}$  is a higher  $\{g_n, h_n\}$ -derivation on  $\mathcal{A}$  with  $f_0 = g_0 = h_0 = I$ , then there is a sequence  $\{F_n\}_{n=0,1,\ldots}$  of  $\{G_n, H_n\}$ -derivations with  $F_0 = G_0 = H_0 = 0$  characterizing the higher  $\{g_n, h_n\}$ -derivation  $\{f_n\}_{n=0,1,\ldots}$ . As well as if  $\{F_n\}_{n=0,1,\ldots}$  is a sequence of  $\{G_n, H_n\}$ -derivations with  $F_0 = G_0 = H_0 = 0$ , then there exists a higher  $\{g_n, h_n\}$ -derivation  $\{f_n\}$  with  $f_0 = g_0 = h_0 = I$  which is characterized by the sequence  $\{F_n\}_{n=0,1,\ldots}$ .

Proof. Let  $\{f_n\}$  be a higher  $\{g_n, h_n\}$ -derivation on  $\mathcal{A}$  with  $f_0 = g_0 = h_0 = I$ . We are going to obtain a sequence  $\{F_n\}_{n=0,1,\dots}$  of  $\{G_n, H_n\}$ -derivations with  $F_0 = G_0 = H_0 = 0$  that characterizes the higher  $\{g_n, h_n\}$ -derivation  $\{f_n\}$ . Define  $F_n, G_n, H_n : \mathcal{A} \to \mathcal{A}$  by  $F_0 = G_0 = H_0 = 0$  and

$$\begin{cases} F_n = nf_n - \sum_{k=0}^{n-2} F_{k+1}f_{n-1-k}, \\ G_n = ng_n - \sum_{k=0}^{n-2} G_{k+1}g_{n-1-k}, \\ H_n = nh_n - \sum_{k=0}^{n-2} H_{k+1}h_{n-1-k}, \end{cases}$$

for each positive integer n. Then it follows from Lemma 2.2 that  $\{F_n\}$  is a sequence of  $\{G_n, H_n\}$ -derivations characterizing the higher  $\{g_n, h_n\}$ -derivation  $\{f_n\}$ . Now suppose that  $\{F_n\}_{n=0,1,\ldots}$  is a sequence of  $\{G_n, H_n\}$ -derivations with  $F_0 = G_0 = H_0 = 0$ . We will show that there exists a higher  $\{g_n, h_n\}$ -derivation  $\{f_n\}$  with  $f_0 = g_0 = h_0 = I$  which is characterized by the sequence  $\{F_n\}_{n=0,1,\ldots}$ .

We define  $f_n, g_n, h_n : \mathcal{A} \to \mathcal{A}$  by  $f_0 = g_0 = h_0 = I$  and

$$\begin{cases} f_n = \sum_{i=1}^n \left( \sum_{\sum_{j=1}^i r_j = n} \left( \prod_{j=1}^i \frac{1}{r_j + \dots + r_i} \right) F_{r_1} \dots F_{r_i} \right), \\ g_n = \sum_{i=1}^n \left( \sum_{\sum_{j=1}^i r_j = n} \left( \prod_{j=1}^i \frac{1}{r_j + \dots + r_i} \right) G_{r_1} \dots G_{r_i} \right), \\ h_n = \sum_{i=1}^n \left( \sum_{\sum_{j=1}^i r_j = n} \left( \prod_{j=1}^i \frac{1}{r_j + \dots + r_i} \right) H_{r_1} \dots H_{r_i} \right). \end{cases}$$

By Theorem 2.4,  $f_n$ ,  $g_n$  and  $h_n$  satisfy the following recursive relations:

$$\begin{cases} (n+1)f_{n+1} = \sum_{k=0}^{n} F_{k+1}f_{n-k}, \\ (n+1)g_{n+1} = \sum_{k=0}^{n} G_{k+1}g_{n-k}, \\ (n+1)h_{n+1} = \sum_{k=0}^{n} H_{k+1}h_{n-k}. \end{cases}$$

Based on the last part of the proof of Theorem 2.4,  $\{f_n\}$  is a higher  $\{g_n, h_n\}$ -derivation on  $\mathcal{A}$  with  $f_0 = g_0 = h_0 = I$ . This yields the desired result.

An immediate corollary of the precede theorem reads as follows:

**Corollary 2.9.** If  $\{d_n\}_{n=0,1,\ldots}$  with  $d_0 = I$  is a higher derivation on  $\mathcal{A}$ , then there is a sequence  $\{\delta_n\}_{n=0,1,\ldots}$  of derivations with  $\delta_0 = 0$  characterizing the higher derivation  $\{d_n\}_{n=0,1,\ldots}$ . As well as if  $\{\delta_n\}_{n=0,1,\ldots}$  is a sequence of derivations with  $\delta_0 = 0$ , then there exists a higher derivation  $\{d_n\}$  with  $d_0 = I$  which is characterized by the sequence  $\{\delta_n\}_{n=0,1,\ldots}$ .

## References

- M. Brešar, Jordan {g, h}-derivations on tensor products of algebras, Linear Multilinear Algebra. 64 (2016), no. 11, 2199-2207.
- M. Brešar, Jordan derivations on semiprime rings, Proc. Amer. Math. Soc. 140 (1988), no. 4, 1003–1006.
- M. Brešar and J. Vukman, Jordan derivations on prime rings, Bull. Austral. Math. Soc. 37 (1988), 321–322.
- M. Brešar, Characterizations of derivations on some normed algebras with involution, J. Algebra 152 (1992), 454–462.
- 5. J. Cusack, Jordan derivations on rings, Proc. Amer. Math. Soc. 53 (1975), 1104–1110.
- A. Hosseini and A. Fošner, Identities related to (σ, τ)-Lie derivations and (σ, τ)-derivations, Boll. Unione Mat. Ital. DOI 10.1007/s40574-017-0141-1, (to appear).
- I. N. Herstein, Jordan derivations of prime rings, Proc.Amer. Math. Soc. 8 (1957), 1104– 1110.
- Y. Li and D. Benkovič. Jordan generalized derivations on triangular algebras. Linear Multilinear Algebra. 59 (2011), 841849.
- M. Mirzavaziri, Characterization of higher derivations on algebras, Communications in algebra. 38 (2010), no. 3, 981–987.
- 10. G. J. Murphy, C<sup>\*</sup>-Algebras and Operator Theory. Boston Academic Press, (1990).

- 11. M. Takesaki, *Theory of Operator Algebras*, Springer-Verlag Berlin Heidelberg New York, 2001.
- J. Vukman, A note on generalized derivations of semiprime rings, Taiwanese J. Math. 11 (2007), 367-370.

Amin Hosseini, Department of Mathematics, Kashmar Higher Education Institute-Kashmar- Iran

*E-mail address*: hosseini.amin820gmail.com, a.hosseini@kashmar.ac.ir

NADEEM UR REHMAN, DEPARTMENT OF MATHEMATICS, ALIGARH MUSLIM UNIVERSITY, ALIGARH-202002 INDIA

*E-mail address*: rehman100@gmail.com

14