Hodge-theoretic analysis on manifolds with boundary, heatable currents, and Onsager's conjecture in fluid dynamics

Khang Manh Huynh

Abstract

We use Hodge theory and functional analysis to develop a clean approach to heat flows and Onsager's conjecture on Riemannian manifolds with boundary, where the weak solution lies in the trace-critical Besov space $B_{3,1}^{\frac{1}{3}}$. We also introduce heatable currents as the natural analogue to tempered distributions and justify their importance in Hodge theory.

Acknowledgments

The author thanks his advisor Terence Tao for the invaluable guidance and patience as the project quickly outgrew its original scale. The author is also grateful to Jochen Glück for recommending Stephan Fackler's insightful PhD thesis [Fac15].

Contents

1	Inti	roduction	2
	1.1	Onsager's conjecture	2
	1.2	Modularity	4
	1.3	Motivation behind the approach	4
	1.4	Blackboxes	5
	1.5	For the specialists	6
2	Cor	nmon notation	7
3	Ons	sager's conjecture	8
	3.1	Summary of preliminaries	8
	3.2	Searching for the proper formulation	12
	3.3	Justification of formulation	13
	3.4	Heating the nonlinear term	14
	3.5	Proof of Onsager's conjecture	15
4	Fun	actional analysis	20
	4.1	Common tools	20
	4.2	Interpolation theory	20
	4.3	Stein extrapolation of analyticity of semigroups	23
		4.3.1 Semigroup definitions	24
		4.3.2 Simple extrapolation (with core)	25
		4.3.3 Coreless version	26

5 Scalar function spaces	80	
5.1 On \mathbb{R}^n	30	
5.2 On domains \ldots	31	
5.3 Holder & Zygmund spaces	33	
5.4 Interpolation & embedding	34	
5.5 Strip decay	35	
6 Hodge theory	88	
6.1 The setting \ldots	38	
6.1.1 Vector bundles	38	
6.1.2 Compatibility with scalar function spaces	39	
6.1.3 Complexification issue	11	
6.2 Differential forms & boundary	41	
6.3 Boundary conditions and potential theory	14	
6.4 Hodge decomposition	17	
6.5 An easy mistake	52	
7 Heat flow	52	
7.1 L^2 -analyticity	52	
7.2 L^p -analyticity	53	
7.3 $W^{1,p}$ -analyticity	55	
7.4 Distributions and adjoints	57	
7.5 Square root \ldots	30	
7.6 Some trace-zero results	31	
8 Results related to the Euler equation 6	52	
8.1 Hodge-Sobolev spaces	32	
8.2 Calculating the pressure	34	
8.3 On an interpolation identity \ldots \ldots \ldots \ldots \ldots \ldots \ldots \ldots \ldots	34	
A Complexification	57	
Nomenclature		

Introduction 1

Onsager's conjecture 1.1

Recall the incompressible Euler equation in fluid dynamics:

$$\begin{cases} \partial_t V + \operatorname{div} \left(V \otimes V \right) &= -\operatorname{grad} p & \operatorname{in} M \\ \operatorname{div} V &= 0 & \operatorname{in} M \\ \langle V, \nu \rangle &= 0 & \operatorname{on} \partial M \end{cases}$$
(1)

 $\begin{array}{l} \text{where} \left\{ \begin{array}{l} (M,g) \text{ is an oriented, compact smooth Riemannian manifold with smooth boundary, dimension} \geq 2 \\ \nu \text{ is the outwards unit normal vector field on } \partial M. \\ I \subset \mathbb{R} \text{ is an open interval, } V: I \rightarrow \mathfrak{X}M, \ p: I \times M \rightarrow \mathbb{R}. \end{array} \right. \end{array}$

Observe that the **Neumann condition** $\langle V, \nu \rangle = 0$ means $V \in \mathfrak{X}_N$, where \mathfrak{X}_N is the set of vector fields on M which are tangent to the boundary. Note that when V is not smooth, we need the trace theorem to define the condition (see Subsection 5.2).

Roughly speaking, Onsager's conjecture says that the energy $||V(t, \cdot)||_{L^2}$ is a.e. constant in time when V is a weak solution whose regularity is at least $\frac{1}{3}$. Making that statement precise is part of the challenge.

In the boundaryless case, the "positive direction" (conservation when regularity is at least $\frac{1}{3}$) has been known for a long time [Eyi94; CET94; Che+08]. The "negative direction" (failure of energy conservation when regularity is less than $\frac{1}{3}$) is substantially harder [LS12; LJ14], and was finally settled by Isett in his seminal paper [Ise18] (see the survey in [LS19] for more details and references).

Since then more attention has been directed towards the case with boundary, and its effects in the generation of turbulence. In [BT17], the "positive direction" was proven in the case M is a bounded domain in \mathbb{R}^n and $V \in L^3_t C^{0,\alpha} \mathfrak{X}_N$ ($\alpha > \frac{1}{3}$). The result was then improved in various ways [DN18; Bar+18; BTW19]. In [NN18], the conjecture was proven for V in $L^3_t B^{\alpha}_{3,\infty} \mathfrak{X}$ ($\alpha > \frac{1}{3}$) along with some "strip decay" conditions for V and p near the boundary (more details in Subsection 3.2). Most recently, the conjecture was proven as part of a more general conservation of entropy law in [Bar+19], where M is a domain in \mathbb{R}^n , $V \in L^3_t \underline{B}^{1/3}_{3,\text{VMO}} \mathfrak{X}$ (where $\underline{B}^{1/3}_{3,\text{VMO}} \mathfrak{X}$ is a VMO-type subspace of $B^{1/3}_{3,\infty} \mathfrak{X}$), along with a "strip decay" condition involving both V and p near the boundary (see Subsection 3.2).

Much less is known about the conjecture on general Riemannian manifolds. The key arguments on flat spaces rely on the nice properties of convolution, such as div $(T * \phi_{\varepsilon}) = \operatorname{div}(T) * \phi_{\varepsilon}$ where T is a tensor field and $\phi_{\varepsilon} \xrightarrow{\varepsilon \downarrow 0} \delta_0$ is a mollifier, or that mollification is essentially local. This "local approach" by convolution does not generalize well to Riemannian manifolds. In [IO15] – the main inspiration for this paper – Isett and Oh used the heat flow to prove the conjecture on compact Riemannian manifolds without boundary, for $V \in L_t^3 B_{3,c(\mathbb{N})}^{\frac{1}{3}} \mathfrak{X}$ (where $B_{3,c(\mathbb{N})}^{\frac{1}{3}} \mathfrak{X}$ is the $B_{3,\infty}^{\frac{1}{3}}$ -closure of compactly supported smooth vector fields). The situation becomes more complicated when the boundary is involved. Most notably, the covariant derivative behaves badly on the boundary (e.g. the second fundamental form), and it is difficult to avoid boundary terms that come from integration by parts. Even applying the heat flow to a distribution might no longer be well-defined. This requires a finer understanding of analysis involving the boundary, as well as the properties of the heat flow.

In this paper, we will see how we can resolve these issues, and that the conjecture still holds true with the boundary:

Fact. Assuming M as in Equation (1), conservation of energy is true when (V, p) is a weak solution with $V \in L^3_t B^{\frac{1}{3}}_{3,1} \mathfrak{X}_N$.

It is not a coincidence that this is also the lowest regularity where the trace theorem holds. We also note a very curious fact that no "strip decay" condition involving p (which is present in different forms for the results on flat spaces) seems to be necessary, and we only need $p \in L^1_{loc} (I \times M)$ (see Subsection 3.3 for details). One way to explain this minor improvement is that the "strip decay" condition involving Vnaturally originates from the trace theorem (see Subsection 3.3), and is therefore included in the condition $V \in L^3_t B^{\frac{1}{3}}_{3,1} \mathfrak{X}_N$, while the presence of p is more of a technical artifact arising from localization (see [Bar+19, Section 4]), which typically does not respect the Leray projection. By using the trace theorem and the heat flow, our approach becomes global in nature, and thus avoids the artifact. Another approach is to formulate the conjecture in terms of Leray weak solutions like in [RRS18], without mentioning p at all, and we justify how this is possible in Subsection 3.3.

A more local approach, where we assume $V \in L^3_t B^{\frac{1}{3}}_{3,c(\mathbb{N})} \mathfrak{X}$ as in [IO15], and the "strip decay" condition as in [Bar+19, Equation 4.9], would be a good topic for another paper. Nevertheless, $B^{\frac{1}{3}}_{3,1}\mathfrak{X}_N$ is an interesting

space with its own unique results, which keep the exposition simple and allow the boundary condition to be natural.

1.2 Modularity

The paper is intended to be modular: the part dealing with Onsager's conjecture (Section 3) is relatively short, while the rest is to detail the tools for harmonic analysis on manifolds we will need (and more). As we will summarize the tools in Section 3, they can be read independently.

1.3 Motivation behind the approach

Riemannian manifolds (and their semi-Riemannian counterparts) are among the most important natural settings for modern geometric PDEs and physics, where the objects for analysis are often vector bundles and differential forms. The two fundamental tools for a harmonic analyst – mollification and Littlewood-Paley projection via the Fourier transform – do not straightforwardly carry over to this setting, especially when the boundary is involved. Even in the case of scalar functions on bounded domains in \mathbb{R}^n , mollification arguments often need to stay away from the boundary, which can present a problem when the trace is nonzero. Consider, however, the idea of a special kind of Littlewood-Paley projection which preserves the boundary conditions and commutes with important operators such as divergence and the Leray projection, or using the principles of harmonic analysis without translation invariance. It is one among a vast constellation of ideas which have steadily become more popular over the years, with various approaches proposed (and we can not hope to fully recount here).

For our discussion, the starting point of interest is perhaps [Str83], in which Strichartz introduced to analysts what had long been known to geometers, the rich setting of complete Riemannian manifolds, where harmonic analysis (and the **Riesz transform** in particular) can be done via the Laplacian and the **heat semigroup** $e^{t\Delta}$, constructed by **dissipative operators** and Yau's lemma. Then in [KR06], Klainerman and Rodnianski defined the L^2 -heat flow by the **spectral theorem** and used it to get the Littlewood-Paley projection on compact 2-surfaces. In [IO15], Isett and Oh successfully tackled Onsager's conjecture on Riemannian manifolds without boundary by using Strichartz's heat flow. These results hint at the central importance of the heat flow for analysis on manifolds. But it is not enough to settle the case with boundary, especially when derivatives are involved. Some pieces of the puzzle are still missing.

To paraphrase James Arthur (in his introduction to the trace formula and the Langlands program), there is an intimate link between geometric objects and "spectral" phenomena, much like how the shape of a drum affects its sounds. For a Riemannian manifold, that link is better known as the Laplacian – the generator of the heat flow – and **Hodge theory** is the study of how the Laplacian governs the cohomology of a Riemannian manifold. An oversimplified description of Fourier analysis on \mathbb{R}^n would be "the spectral theory of the Laplacian" [Str89], where the heat kernel is the Gaussian function, invariant under the Fourier transform and a possible choice of mollifier. Additionally, the Helmholtz decomposition, originally discovered in a hydrodynamic context, turned out to be a part of Hodge theory. It should therefore be no surprise that Hodge theory is the natural framework in which we formulate harmonic analysis on manifolds, heat flows and Onsager's conjecture. Wherever there is the Laplacian, there is harmonic analysis. Historically, Milgram managed to establish a subset of Hodge theory by heat flow methods [MR51]. Here, however, we will establish Hodge theory by standard elliptic estimates, from which we develop analysis on manifolds and construct the heat flow. Most notably, Hodge theory greatly simplifies some crucial approximation steps involving the boundary (Corollary 70), and helps predict some key results Onsager's conjecture would require (Theorem 15, Subsection 7.4, Subsection 8.3). That such leaps of faith turn out to be true only further underscore how well-made the conjecture is in its anticipation of undiscovered mathematics.

For those familiar with the smoothing properties of Littlewood-Paley projection as well as **Bernstein** inequalities [Tao06, Appendix A], the rough picture is that $e^{t\Delta} \approx P_{\leq \frac{1}{\sqrt{t}}}$. While the introduction of curvature necessitates the change of constants in estimates, and the boundary requires its own considerations, it is remarkable how far we can go with this analogy. Regarding the properties we will need for Onsager's conjecture, there is a satisfying explanation: the theory of sectorial operators in functional analysis. This, together with Hodge theory, the theory of Besov spaces and interpolation theory, allows us to build a basic foundation for global analysis on Riemannian manifolds in general, which will be more than enough to handle Onsager's conjecture.

Hodge theory and sectorial operators, in their various forms, have been used in fluid dynamics for a long time by Fujita, Kato, Giga, Miyakawa *et al.* (cf. [FK64; Miy80; Gig81; GM85; BAE16] and their references). Although we will not use them for this paper, we also ought to mention the results regarding bisectorial operators, H^{∞} functional calculus, and Hodge theory on rough domains developed by Alan McIntosh, Marius Mitrea, Sylvie Monniaux *et al.* (cf. [McI86; DM96; FMM98; AM04; MM08; MM09a; MM09b; GMM10; She12; MM18] and their references), which generalize many Hodge-theoretic results in this paper. Alternative formalizations of Littlewood-Paley theory also exist (cf. [HMY08; KP14; FFP16; KW16; BBD18; Tan18] and their references). Here, we are mainly focused on the analogy between the heat flow and the Littlewood-Paley projection on L^p spaces of differential forms (over manifolds with boundary), as well as the interplay with Hodge theory.

Lastly, we also introduce **heatable currents** – the largest space on which the heat flow can be profitably defined – as the analogue to tempered distributions on manifolds (Subsection 7.4). In doing so, we will realize that the energy-conserving weak solution in Onsager's conjecture solves the Euler equation in the sense of heatable currents. This is an elegant insight that helps show how interconnected these subjects are. Much like how learning the language of measure theory can shed light on problems in calculus and familiarity with differential geometry simplifies many calculations in fluid dynamics, the cost of learning ostensibly complicated formalism is often dwarfed by the benefits in clarity it brings. That being said, accessibility is also important, and besides providing a gentle introduction to the theory with copious references, this paper also hopes to convince the reader of the naturality behind the formalism.

1.4 Blackboxes

Since we draw upon many areas, the paper is intended to be as self-contained as possible, but we will assume familiarity with basic elements of functional analysis, harmonic analysis and complex analysis. Some familiarity with differential and Riemannian geometry is certainly needed (cf. [Lee09; Cha06]), as well as **Penrose notation** (cf. [Wal84, Section 2.4]). In addition, a number of blackbox theorems will be borrowed from the following sources:

- 1. For interpolation theory: Interpolation Spaces [BL76] and "Abstract Stein Interpolation" [Voi92]
- 2. For harmonic analysis and elements of functional analysis:
 - Singular Integrals and Differentiability Properties of Functions. (PMS-30) [Ste71]
 - Partial Differential Equations I [Tay11a]
 - Recent developments in the Navier-Stokes problem (Chapman & Hall/CRC Research Notes in Mathematics Series) [Lem02]
- 3. For Besov spaces: Theory of Function Spaces; Theory of Function Spaces II [Tri10; Tri92]
- 4. For Hodge theory: Hodge Decomposition—A Method for Solving Boundary Value Problems [Sch95]

5. For semigroups and sectorial operators: One-parameter semigroups for linear evolution equations [Eng00] and Vector-valued Laplace Transforms and Cauchy Problems: Second Edition (Monographs in Mathematics) [Are+11]

The first three categories should be familiar with harmonic analysts.

1.5 For the specialists

Some noteworthy characteristics of our approach:

- An alternative development of the (absolute Neumann) heat flow. In particular, the extrapolation of analyticity to L^p spaces does not involve establishing the resolvent estimate in Yosida's half-plane criterion (Theorem 39), either via "Agmon's trick" [Agm62] as done in [Miy80] or manual estimates as in [BAE16]. Instead, by abstract Stein interpolation, we only need the local boundedness of the heat flow on L^p , which can follow cleanly from Gronwall and integration by parts (Theorem 71). In short, functional analysis does the heavy lifting. We also managed to attain $W^{1,p}$ -analyticity assuming the Neumann condition (Subsection 7.3), and $B_{p,1}^{\frac{1}{p}}$ -analyticity via the Leray projection (Subsection 8.3).
- We do not focus on the **Stokes operator** in this paper, but our results (Subsection 7.3, Subsection 8.3) do contain the case of the Stokes operator corresponding to the "Navier-type" / "free" boundary condition, as discussed in [Miy80; Gig82; MM09a; MM09b; BAE16] and others. This should not be confused with the Stokes operator corresponding to the "no-slip" boundary condition, as discussed in [FK64; GM85; MM08] and others. See [HS18] for more references.
- For simplicity, we stay within the smooth and compact setting, which, as Hilbert would say, is that special case containing all the germs of generality. An effort has also been made to keep the material concrete (as opposed to, for instance, using Hilbert complexes).
- Heatable currents are introduced as the analogue to tempered distributions, and we show how they naturally appear in the characterization of the adjoints of d and δ (Subsection 7.4).
- A refinement of a special case of the fractional Leibniz rule, with the supports of functions taken into account, is given in Theorem 54.
- For the proof of Onsager's conjecture, there are some subtle, but substantial differences with [IO15]:
 - In [IO15], Besov spaces are *defined* by the heat flow, and compatibility with the usual scalar Besov spaces is proven when M is \mathbb{R}^n or \mathbb{T}^n . Here we will use the standard scalar Besov spaces as defined by Triebel in [Tri10; Tri92], and prove the appropriate estimates for the heat flow by interpolation.
 - The heat flow used by Isett & Oh (constructed by Strichartz using dissipative operators) is generated by the **Hodge Laplacian**, which is self-adjoint in the no-boundary case. In the case with boundary, there are four different self-adjoint versions for the Hodge Laplacian (see Theorem 61), and we choose the **absolute Neumann** version. There are also heat flows generated by the **connection Laplacian**, but we do not use them in this paper since the connection Laplacian does not commute with the **exterior derivative** and the Leray projection etc. The theory of dissipative operators is also not sufficient to establish L^p -analyticity and $W^{1,p}$ -analyticity for all $p \in (1, \infty)$, so we instead use the theory of sectorial operators, which is made for this purpose.

- The commutator we will use is a bit different from that in [IO15]. This will help us eliminate some boundary terms. We will also avoid the explicit formula and computations in [IO15, Lemma 4.4], as they also lead to various boundary terms. Generally speaking, the covariant derivative behaves badly on the boundary.
- A calculation of the pressure by negative-order Hodge-Sobolev spaces (Subsection 8.2).
- More results will be proven for analysis on manifolds than needed for Onsager's conjecture, as they are of independent interest. For the sake of accessibility, we will also review most of the relevant background material, with the assumption that the reader is a harmonic analyst who knows some differential geometry.

It is hard to overstate our indebtedness to all the mathematicians whose work our theory will build upon, from harmonic analysis to Hodge theory and sectorial operators, and yet hopefully each will be able to find within this paper something new and interesting.

2 Common notation

It might not be an exaggeration to say the main difficulty in reading a paper dealing with Hodge theory is understanding the notation, and an effort has been made to keep our notation as standard and selfexplanatory as possible.

Some common notation we use:

- $A \leq_{x,\neg y} B$ means $A \leq CB$ where C > 0 depends on x and not y. Similarly, $A \sim_{x,\neg y} B$ means $A \leq_{x,\neg y} B$ and $B \leq_{x,\neg y} A$. When the dependencies are obvious by context, we do not need to make them explicit.
- $\mathbb{N}_0, \mathbb{N}_1$: the set of natural numbers, starting with 0 and 1 respectively.
- DCT: dominated convergence theorem, FTC: fundamental theorem of calculus, PTAS: passing to a subsequence, WLOG: without loss of generality.
- TVS: topological vector space, NVS: normed vector space, SOT: strong operator topology.
- For TVS $X, Y \leq X$ means Y is a subspace of X.
- $\mathcal{L}(X, Y)$: the space of continuous linear maps from TVS X to Y. Also $\mathcal{L}(X) = \mathcal{L}(X, X)$.
- $C^0(S \to Y)$: the space of bounded, continuous functions from metric space S to normed vector space Y. Not to be confused with $C^0_{\text{loc}}(S \to Y)$, which is the space of locally bounded, continuous functions.
- $||x||_{D(A)} = ||x||_X + ||Ax||_X$ and $||x||_{D(A)}^* = ||Ax||_X$ where A is an unbounded operator on (real/complex) Banach space X and $x \in D(A)$. Note that $||\cdot||_{D(A)}^*$ is not always a norm. Also define $D(A^{\infty}) = \bigcap_{k \in \mathbb{N}_1} D(A^k)$.
- For $\delta \in (0, \pi]$, define the open sector $\Sigma_{\delta}^+ = \{z \in \mathbb{C} \setminus \{0\} : |\arg z| < \delta\}$, $\Sigma_{\delta}^- = -\Sigma_{\delta}^+$, $\mathbb{D} = \{z \in \mathbb{C} : |z| < 1\}$. Also define $\Sigma_0^+ = (0, \infty)$ and $\Sigma_0^- = -\Sigma_0^+$.
- B(x,r): the open ball of radius r centered at x in a metric space.
- $\mathcal{S}(\mathbb{R}^n)$: the space of Schwartz functions on \mathbb{R}^n , $\mathcal{S}(\overline{\Omega})$: restrictions of Schwartz functions to the domain $\Omega \subset \mathbb{R}^n$.

There is also a list of other symbols we will use at the end of the paper.

3 Onsager's conjecture

3.1 Summary of preliminaries

At the cost of some slight duplication of exposition, we will quickly summarize the key tools we need for the proof, and leave the development of such tools for the rest of the paper. Alternatively, the reader can read the theory first and come back to this section later.

Definition 1. For the rest of the paper, unless otherwise stated, let M be a compact, smooth, Riemannian n-dimensional manifold, with no or smooth boundary. We also let $I \subset \mathbb{R}$ be an open time interval. We write $M_{\leq r} = \{x \in M : \operatorname{dist}(x, \partial M) < r\}$ for r > 0 small. Similarly define $M_{\geq r}, M_{\leq r}, M_{[r_1, r_2]}$ etc. Let \mathring{M} denote the interior of M.

By musical isomorphism, we can consider $\mathfrak{X}M$ (the space of **smooth vector fields**) mostly the same as $\Omega^1(M)$ (the space of **smooth 1-forms**), *mutatis mutandis*. We note that $\mathfrak{X}M$, $\mathfrak{X}(\partial M)$ and $\mathfrak{X}M|_{\partial M}$ are different. Unless otherwise stated, let the implicit domain be M, so \mathfrak{X} stands for $\mathfrak{X}M$, and similarly Ω^k for $\Omega^k M$. For $X \in \mathfrak{X}$, we write X^{\flat} as its dual 1-form. For $\omega \in \Omega^1$, we write ω^{\sharp} as its dual vector field.

Let $\mathfrak{X}_{00}(M)$ denote the set of smooth vector fields of compact support in \tilde{M} . Define $\Omega_{00}^{k}(M)$ similarly (smooth differential forms with compact support in \tilde{M}).

Let ν denote the outwards unit normal vector field on ∂M . ν can be extended via geodesics to a smooth vector field $\tilde{\nu}$ which is of unit length near the boundary (and cut off at some point away from the boundary).

For $X \in \mathfrak{X}M$, define $\mathbf{n}X = \langle X, \nu \rangle \nu \in \mathfrak{X}M|_{\partial M}$ (the **normal part**) and $\mathbf{t}X = X|_{\partial M} - \mathbf{n}X$ (the **tangential part**). We note that $\mathbf{t}X$ and $\mathbf{n}X$ only depend on $X|_{\partial M}$, so \mathbf{t} and \mathbf{n} can be defined on $\mathfrak{X}M|_{\partial M}$, and $\mathbf{t}(\mathfrak{X}M|_{\partial M}) \xrightarrow{\sim} \mathfrak{X}(\partial M)$.

For $\omega \in \Omega^{k}(M)$, define $\mathbf{t}\omega$ and $\mathbf{n}\omega$ by

$$\mathbf{t}\omega(X_1,...,X_k) := \omega(\mathbf{t}X_1,...,\mathbf{t}X_k) \ \forall X_j \in \mathfrak{X}M, j = 1,...,k$$

and $\mathbf{n}\omega = \omega|_{\partial M} - \mathbf{t}\omega$. Note that $(\mathbf{n}X)^{\flat} = \mathbf{n}X^{\flat} \ \forall X \in \mathfrak{X}$.

Let ∇ denote the **Levi-Civita connection**, d the **exterior derivative**, δ the **codifferential**, and $\Delta = -(d\delta + \delta d)$ the **Hodge-Laplacian**, which is defined on vector fields by the musical isomorphism.

Familiar scalar function spaces such as $L^p, W^{m,p}$ (Lebesgue-Sobolev spaces), $B^s_{p,q}$ (Besov spaces), $C^{0,\alpha}$ (Holder spaces) (see Section 5 for precise definitions) can be defined on M by partitions of unity and given a unique topology (Subsection 5.2, Subsection 6.1.2). Similarly, we define such function spaces for tensor fields and differential forms on M by partitions of unity and local coordinates (see subsection 6.1). For instance, we can define $L^2\mathfrak{X}$ or $B_{3,1}^{\frac{1}{3}}\mathfrak{X}$.

Fact 2. $\forall \alpha \in \left(\frac{1}{3}, 1\right), \forall p \in (1, \infty) : W^{1,p} \mathfrak{X} \hookrightarrow B_{p,1}^{\frac{1}{p}} \mathfrak{X} \hookrightarrow L^{p} \mathfrak{X} \text{ and } C^{0,\alpha} \mathfrak{X} = B_{\infty,\infty}^{\alpha} \mathfrak{X} \hookrightarrow B_{3,\infty}^{\alpha} \mathfrak{X} \hookrightarrow B_{3,1}^{\frac{1}{3}} \mathfrak{X} \text{ (cf. Subsection 5.2, Subsection 5.4)}$

Definition 3. We write $\langle \cdot, \cdot \rangle$ to denote the **Riemannian fiber metric** for tensor fields on M. We also define the dot product

$$\langle \langle \sigma, \theta \rangle \rangle = \int_M \langle \sigma, \theta \rangle$$
 vol

where σ and θ are tensor fields of the same type, while vol is the **Riemannian volume form**. When there is no possible confusion, we will omit writing vol.

We define $\mathfrak{X}_N = \{X \in \mathfrak{X} : \mathbf{n}X = 0\}$ (Neumann condition). Similarly, we can define Ω_N^k . In order to define the Neumann condition for less regular vector fields (and differential forms), we need to use the trace theorem.

Fact 4. (Subsection 5.2, Subsection 6.1.2) Let $p \in [1, \infty)$. Then

- $B_{p,1}^{\frac{1}{p}}(M) \twoheadrightarrow L^{p}(\partial M)$ and $B_{p,1}^{\frac{1}{p}}\mathfrak{X}M \twoheadrightarrow L^{p}\mathfrak{X}M\Big|_{\partial M}$ are continuous surjections.
- $\forall m \in \mathbb{N}_1 : B_{p,1}^{m+\frac{1}{p}} \mathfrak{X}M \twoheadrightarrow B_{p,1}^m \mathfrak{X}M \big|_{\partial M} \hookrightarrow W^{m,p} \mathfrak{X}M \big|_{\partial M}$ is continuous.

Also closely related is the **coarea formula**:

Fact 5. (Theorem 53) Let $p \in [1, \infty)$, r > 0 be small and f be in $B_{p,1}^{\frac{1}{p}}(M)$:

- 1. $([0,r) \to \mathbb{R}, \rho \mapsto \|f\|_{L^p(\partial M_{>\rho})})$ is continuous and bounded by $C \|f\|_{B^{\frac{1}{p}}_{p,1}}$ for some C > 0.
- 2. $|M_{\leq r}| \sim_{M,\neg r} |\partial M| r$ and $||f||_{L^p(M_{\leq r})} \sim_{\neg r} \left\| ||f||_{L^p(\partial M_{> \rho})} \right\|_{L^p_{\rho}((0,r))}$.
- 3. $||f||_{L^p(M_{\leq r}, \operatorname{avg})} \lesssim_{\neg r} ||f||_{B^{\frac{1}{p}}_{p,1}(M)}$ and $||f||_{L^p(M_{\leq r}, \operatorname{avg})} \xrightarrow{r\downarrow 0} ||f||_{L^p(\partial M, \operatorname{avg})}$, where avg means normalizing the measure to make it a probability measure.
- $4. Let \mathfrak{f} \in L^p(I \to B_{p,1}^{\frac{1}{p}}(M)), then \|\mathfrak{f}\|_{L_t^p B_{p,1}^{\frac{1}{p}}(M)} \gtrsim_{\neg r} \|\mathfrak{f}\|_{L_t^p L^p(M_{\leq r}, \operatorname{avg})} \xrightarrow{r \downarrow 0} \|\mathfrak{f}\|_{L_t^p L^p(\partial M, \operatorname{avg})}.$

Analogous results hold if $f \in B_{p,1}^{\frac{1}{p}} \mathfrak{X}$. (Subsection 6.1.2)

Therefore, we can define spaces such as $B_{3,1}^{\frac{1}{3}}\mathfrak{X}_N = \{X \in B_{3,1}^{\frac{1}{3}}\mathfrak{X} : \mathbf{n}X = 0\}$ and $W^{1,3}\mathfrak{X}_N$. However, something like $L^2\mathfrak{X}_N$ would not make sense since the trace map does not continuously extend to $L^2\mathfrak{X}$.

Definition 6. We define \mathbb{P} as the **Leray projection** (constructed in Theorem 68), which projects \mathfrak{X} onto Ker $(\operatorname{div}|_{\mathfrak{X}_N})$. Note that the Neumann condition is enforced by \mathbb{P} .

Fact 7. $\forall m \in \mathbb{N}_0, \forall p \in (1, \infty), \mathbb{P} \text{ is continuous on } W^{m,p}\mathfrak{X} \text{ and } \mathbb{P}(W^{m,p}\mathfrak{X}) = W^{m,p}\text{-cl}\left(\operatorname{Ker}\left(\operatorname{div}_{\mathfrak{X}_N}\right)\right)$ (closure in the $W^{m,p}$ -topology). (Subsection 6.4)

We collect some results regarding our heat flow in one place:

Fact 8 (Absolute Neumann heat flow). There exists a semigroup of operators $(S(t))_{t\geq 0}$ acting on $\cup_{p\in(1,\infty)}L^p\mathfrak{X}$ such that

- 1. $S(t_1) S(t_2) = S(t_1 + t_2) \ \forall t_1, t_2 \ge 0 \ and \ S(0) = 1.$
- 2. (Subsection 7.2) $\forall p \in (1,\infty), \forall X \in L^p \mathfrak{X}$:
 - (a) $S(t)X \in \mathfrak{X}_N$ and $\partial_t (S(t)X) = \Delta S(t)X \ \forall t > 0.$
 - (b) $S(t)X \xrightarrow{C^{\infty}} S(t_0)X \quad \forall t_0 > 0.$
 - (c) $\|S(t)X\|_{W^{m,p}} \lesssim_{m,p} \left(\frac{1}{t}\right)^{\frac{m}{2}} \|X\|_{L^p} \ \forall m \in \mathbb{N}_0, \forall t \in (0,1).$
 - (d) $S(t)X \xrightarrow{L^p} X$.
- 3. (Subsection 7.3) $\forall p \in (1, \infty), \forall X \in W^{1,p} \mathfrak{X}_N$:
 - (a) $||S(t)X||_{W^{m+1,p}} \lesssim_{m,p} \left(\frac{1}{t}\right)^{\frac{m}{2}} ||X||_{W^{1,p}} \ \forall m \in \mathbb{N}_0, \forall t \in (0,1).$

- (b) $S(t)X \xrightarrow[t \to 0]{W^{1,p}} X.$
- 4. (Theorem 76) $S(t) \mathbb{P} = \mathbb{P}S(t)$ on $W^{m,p}\mathfrak{X} \ \forall m \in \mathbb{N}_0, \forall p \in (1,\infty), \forall t \ge 0.$
- 5. (Subsection 7.2) $\langle \langle S(t)X,Y \rangle \rangle = \langle \langle X,S(t)Y \rangle \rangle \forall t \ge 0, \forall p \in (1,\infty), \forall X \in L^p \mathfrak{X}, \forall Y \in L^{p'} \mathfrak{X}.$

These estimates precisely fit the analogy $e^{t\Delta} \approx P_{\leq \frac{1}{\sqrt{t}}}$ where P is the **Littlewood-Paley projection**. We also stress that the heat flow preserves the space of tangential, divergence-free vector fields (the range of \mathbb{P}), and is intrinsic (with no dependence on choices of local coordinates).

Analogous results hold for scalar functions and differential forms (Section 7). We also have commutativity with the exterior derivative and codifferential in the case of differential forms (Theorem 73). Loosely speaking, this allows the heat flow to preserve the overall Hodge structure on the manifold. All these properties would not be possible under standard mollification via partitions of unity.

Note that for $X \in \mathfrak{X}$, $X \otimes X$ is not dual to a differential form. As our heat flow is generated by the Hodge Laplacian, it is less useful in mollifying general tensor fields (for which the connection Laplacian is better suited). Fortunately, we will never actually have to do so in this paper.

We observe some basic identities (cf. Theorem 58):

- Using **Penrose abstract index notation** (see Subsection 6.2), for any smooth tensors $T_{a_1...a_k}$, we define $(\nabla T)_{ia_1...a_k} = \nabla_i T_{a_1...a_k}$ and div $T = \nabla^i T_{ia_2...a_k}$.
- For all smooth tensors $T_{a_1...a_k}$ and $Q_{a_1...a_{k+1}}$:

$$\int_{M} \nabla_i \left(T_{a_1 \dots a_k} Q^{ia_1 \dots a_k} \right) = \int_{M} \nabla_i T_{a_1 \dots a_k} Q^{ia_1 \dots a_k} + \int_{M} T_{a_1 \dots a_k} \nabla_i Q^{ia_1 \dots a_k} = \int_{\partial M} \nu_i T_{a_1 \dots a_k} Q^{ia_1 \dots a_k}$$

• For $X \in \mathfrak{X}_N, Y \in \mathfrak{X}, f \in C^{\infty}(M)$:

1.
$$\int_{M} Xf = \int_{M} \operatorname{div} (fX) - \int_{M} f \operatorname{div} (X) = \int_{\partial M} \langle fX, \nu \rangle - \int_{M} f \operatorname{div} X = -\int_{M} f \operatorname{div} X$$

2.
$$\int_{M} \langle \operatorname{div} (X \otimes X), Y \rangle = -\int_{M} \langle X \otimes X, \nabla Y \rangle$$

• $(\nabla_a \nabla_b - \nabla_b \nabla_a) T^{ij}{}_{kl} = -R_{ab\sigma}{}^i T^{\sigma j}{}_{kl} - R_{ab\sigma}{}^j T^{i\sigma}{}_{kl} + R_{abk}{}^{\sigma} T^{ij}{}_{\sigma l} + R_{abl}{}^{\sigma} T^{ij}{}_{k\sigma}$ for any tensor $T^{ij}{}_{kl}$, where R is the **Riemann curvature tensor**. Similar identities hold for other types of tensors. When we do not care about the exact indices and how they contract, we can just write the **schematic identity** $(\nabla_a \nabla_b - \nabla_b \nabla_a) T^{ij}{}_{kl} = R * T$. As R is bounded on compact M, interchanging derivatives is a zeroth-order operation on M. In particular, we have the **Weitzenbock formula**:

$$\Delta X = \nabla_i \nabla^i X + R * X \ \forall X \in \mathfrak{X} M \tag{2}$$

- For $X \in \mathbb{P}L^2 \mathfrak{X}, Y \in \mathfrak{X}, Z \in \mathfrak{X}, f \in C^{\infty}(M)$:
 - 1. $\int_M Xf = 0$ 2. $\int_M \langle \nabla_X Y, Z \rangle = - \int_M \langle Y, \nabla_X Z \rangle$.

There is an elementary lemma which is useful for convergence (the proof is straightforward and omitted):

Lemma 9 (Dense convergence). Let X, Y be (real/complex) Banach spaces and $X_0 \leq X$ be norm-dense. Let $(T_j)_{j \in \mathbb{N}}$ be bounded in $\mathcal{L}(X, Y)$ and $T \in \mathcal{L}(X, Y)$.

If $T_j x_0 \to T x_0 \ \forall x_0 \in X_0$ then $T_j x \to T x \ \forall x \in X$.

Definition 10 (Heatable currents). As the heat flow does not preserve compact supports in \mathring{M} , it is not defined on distributions. This inspires the formulation of **heatable currents**. Define:

- $\mathscr{D}\Omega^k = \Omega_{00}^k = \operatorname{colim}\{(\Omega_{00}^k(K), C^\infty \operatorname{topo}) : K \subset \mathring{M} \operatorname{compact}\}$ as the space of **test** *k*-forms with **Schwartz's topology**¹ (colimit in the category of locally convex TVS).
- $\mathscr{D}'\Omega^k = (\mathscr{D}\Omega^k)^*$ as the space of k-currents (or distributional k-forms), equipped with the weak* topology.
- $\mathscr{D}_N \Omega^k = \{ \omega \in \Omega^k : \mathbf{n} \Delta^m \omega = 0, \mathbf{n} d \Delta^m \omega = 0 \ \forall m \in \mathbb{N}_0 \}$ as the space of **heated** k-forms with the Frechet C^{∞} topology and $\mathscr{D}'_N \Omega^k = (\mathscr{D}_N \Omega^k)^*$ as the space of **heatable** k-currents (or **heatable** distributional k-forms) with the weak* topology.
- Spacetime test forms: $\mathscr{D}(I, \Omega^k) = C_c^{\infty}(I, \Omega_{00}^k) = \operatorname{colim}\{(C_c^{\infty}(I_1, \Omega_{00}^k(K)), C^{\infty} \operatorname{topo}) : I_1 \times K \subset I \times \mathring{M} \operatorname{compact}\}$ and $\mathscr{D}_N(I, \Omega^k) = \operatorname{colim}\{(C_c^{\infty}(I_1, \mathscr{D}_N \Omega^k), C^{\infty} \operatorname{topo}) : I_1 \subset I \operatorname{compact}\}.$
- Spacetime distributions $\mathscr{D}'(I,\Omega^k) = \mathscr{D}(I,\Omega^k)^*, \ \mathscr{D}'_N(I,\Omega^k) = \mathscr{D}_N(I,\Omega^k)^*.$

In particular, $\mathscr{D}_N \mathfrak{X}$ is defined from $\mathscr{D}_N \Omega^1$ by the musical isomorphism, and it is invariant under our heat flow (much like how the space of Schwartz functions $\mathcal{S}(\mathbb{R}^n)$ is invariant under the Littlewood-Paley projection). By that analogy, heatable currents are tempered distributions on manifolds, and we can write

$$\langle\langle S(t)\Lambda, X\rangle\rangle = \langle\langle\Lambda, S(t)X\rangle\rangle \ \forall\Lambda \in \mathscr{D}'_N \mathfrak{X}, \forall X \in \mathscr{D}_N \mathfrak{X}, \forall t \ge 0$$

where the dot product $\langle \langle \cdot, \cdot \rangle \rangle$ is simply abuse of notation.

Fact 11. Some basic properties of $\mathscr{D}_N \mathfrak{X}$ and $\mathscr{D}'_N \mathfrak{X}$:

- $\langle \langle \Delta X, Y \rangle \rangle = \langle \langle X, \Delta Y \rangle \rangle \ \forall X, Y \in \mathscr{D}_N \mathfrak{X}.$ (Theorem 58)
- $S(t)\Lambda \in \mathscr{D}_N \mathfrak{X} \ \forall t > 0, \forall \Lambda \in \mathscr{D}'_N \mathfrak{X}$. (Subsection 7.4, a heatable current becomes heated once the heat flow is applied)
- $\mathfrak{X}_{00} \subset \mathscr{D}_N \mathfrak{X}$ and is dense in $L^p \mathfrak{X} \forall p \in [1, \infty)$. Also, $L^p \mathfrak{X} \hookrightarrow \mathscr{D}'_N \mathfrak{X}$ is continuous $\forall p \in [1, \infty]$.
- $\mathbb{P}B_{3,1}^{\frac{1}{3}}\mathfrak{X} = \mathbb{P}B_{3,1}^{\frac{1}{3}}\mathfrak{X}_N$, $\mathbb{P}W^{1,p}\mathfrak{X} = \mathbb{P}W^{1,p}\mathfrak{X}_N$ and $\mathbb{P}\mathscr{D}_N\mathfrak{X} \leq \mathscr{D}_N\mathfrak{X}$. (Subsection 6.4)
- $W^{1,p}$ -cl $(\mathscr{D}_N\mathfrak{X}) = W^{1,p}\mathfrak{X}_N \ \forall p \in (1,\infty) \ (Subsection \ 7.3), \ B_{3,1}^{\frac{1}{3}}$ -cl $(\mathbb{P}\mathscr{D}_N\mathfrak{X}) = \mathbb{P}B_{3,1}^{\frac{1}{3}}\mathfrak{X}_N \ (Subsection \ 8.3)$
- $\forall X \in \mathscr{D}_N \mathfrak{X} : S(t)X \xrightarrow{C^{\infty}} t \downarrow 0$ $X \text{ and } \partial_t (S(t)X) = \Delta S(t)X = S(t)\Delta X \ \forall t \ge 0.$ (Theorem 32, Subsection 7.2)
- (Subsection 7.2, Subsection 8.3) $\forall t \in (0,1), \forall m, m' \in \mathbb{N}_0, \forall p \in (1,\infty), \forall X \in \mathscr{D}_N \mathfrak{X}$:

$$1. \|S(t)X\|_{W^{m+m',p}} \lesssim \left(\frac{1}{t}\right)^{\frac{m'}{2}} \|X\|_{W^{m,p}}.$$

$$2. \|S(t)X\|_{B^{m+m'+\frac{1}{p}}_{p,1}} \lesssim \left(\frac{1}{t}\right)^{\frac{1}{2p}+\frac{m'}{2}} \|X\|_{W^{m,p}}.$$

$$3. t^{\frac{1}{2}(m-\frac{1}{p})} \|S(t)X\|_{W^{m,p}} + \|S(t)X\|_{B^{\frac{1}{p}}_{p,1}} \lesssim \|X\|_{B^{\frac{1}{p}}_{p,1}} \text{ when } m \ge 1 \text{ and } X \in \mathbb{P}\mathscr{D}_{N}\mathfrak{X}$$
By dense convergence (Lemma 9), this means $S(t)X \xrightarrow{B^{\frac{1}{3}}_{3,1}}{t+0} X \ \forall X \in \mathbb{P}B^{\frac{1}{3}}_{3,1}\mathfrak{X}_{N}.$

 $^{^1{\}rm Confusingly\ enough},$ "Schwartz's topology" refers to the topology on the space of distributions, not the topology for Schwartz functions.

Corollary 12 (Vanishing). $\forall X \in \mathbb{P}B_{3,1}^{\frac{1}{3}}\mathfrak{X}_N : s^{\frac{1}{3}} \|S(s)X\|_{W^{1,3}} \xrightarrow{s\downarrow 0} 0.$

Remark. So, for $\mathcal{U} \in L^3_t \mathbb{P}B^{\frac{1}{3}}_{3,1} \mathfrak{X}_N$: $\|\mathcal{U}(t)\|_{L^3_t B^{\frac{1}{3}}_{3,1}} \gtrsim \left\| \left\| \sigma^{\frac{1}{3}} \|S(\sigma)\mathcal{U}(t)\|_{W^{1,3}} \right\|_{L^{\infty}_{\sigma}([0,s])} \right\|_{L^3_t} \xrightarrow{s\downarrow 0} 0.$

This pointwise vanishing property becomes important for the commutator estimate in Onsager's conjecture at the critical regularity level $\frac{1}{3}$, while higher regularity levels have enough room for vanishing in norm (which is better).

Proof. For
$$Y \in \mathbb{P}\mathscr{D}_N \mathfrak{X}$$
, as $s > 0$ small: $s^{\frac{1}{3}} \|S(s)Y\|_{W^{1,3}} \lesssim s^{\frac{1}{3}} \|Y\|_{W^{1,3}} \xrightarrow{s\downarrow 0} 0$. Then note $s^{\frac{1}{3}} \|S(s)X\|_{W^{1,3}} \lesssim \|X\|_{B^{\frac{1}{3}}_{3,1}} \forall X \in \mathbb{P}B^{\frac{1}{3}}_{3,1} \mathfrak{X}_N$, so we can apply dense convergence (Lemma 9). \Box

3.2 Searching for the proper formulation

Onsager's conjecture states that energy is conserved when \mathcal{V} has enough regularity, with appropriate conditions near the boundary. But making this statement precise is half of the challenge.

Definition 13. We say $(\mathcal{V}, \mathfrak{p})$ is a weak solution to the Euler equation when

- $\mathcal{V} \in L^2_{\text{loc}}(I, \mathbb{P}L^2\mathfrak{X}), \mathfrak{p} \in L^1_{\text{loc}}(I \times M)$
- $\forall \mathcal{X} \in C_c^{\infty}(I, \mathfrak{X}_{00}) : \iint_{I \times M} \langle \mathcal{V}, \partial_t \mathcal{X} \rangle + \langle \mathcal{V} \otimes \mathcal{V}, \nabla \mathcal{X} \rangle + \mathfrak{p} \operatorname{div} \mathcal{X} = 0.$

The last condition means $\partial_t \mathcal{V} + \operatorname{div}(\mathcal{V} \otimes \mathcal{V}) + \operatorname{grad} \mathfrak{p} = 0$ as spacetime distributions. Note that $\mathcal{V} \otimes \mathcal{V} \in L^1_{\operatorname{loc}}(I, L^1\mathfrak{X})$ so it is a distribution.

The keen reader should notice we use a different font for time-dependent vector fields.

There is not enough time-regularity for FTC, and we cannot say

$$\langle \langle \mathcal{V}(t_1), X \rangle \rangle - \langle \langle \mathcal{V}(t_0), X \rangle \rangle = \int_{t_0}^{t_1} \langle \langle \mathcal{V} \otimes \mathcal{V}, \nabla X \rangle \rangle + \int_{t_0}^{t_1} \int_M \mathfrak{p} \operatorname{div} X \,\forall X \in \mathfrak{X}_{00}$$

But we can still use approximation to the identity (in the time variable) near t_0, t_1 , as well as Lebesgue differentiation to get something similar for a.e. t_0, t_1 . By using dense convergence (Lemma 9) and modifying I into $I_0 \subset I$ such that $|I \setminus I_0| = 0$, we can say $\mathcal{V} \in C^0_{\text{loc}} \left(I_0, \left(L^2 \mathfrak{X}, \text{weak} \right) \right) \leq L^{\infty}_{\text{loc}} \left(I, L^2 \mathfrak{X} \right)$.

We do not have $\mathcal{V} \in C^0_{\text{loc}}(I, L^2\mathfrak{X})$, so energy conservation only means $\partial_t \left(\|\mathcal{V}(t)\|^2_{L^2\mathfrak{X}} \right) = 0$ as a distribution. In other words, the goal is to show

$$\int_{I} \eta'(t) \left\langle \left\langle \mathcal{V}(t), \mathcal{V}(t) \right\rangle \right\rangle \mathrm{d}t = 0 \; \forall \eta \in C_{c}^{\infty}(I)$$

Next, having the test vector field $\mathcal{X} \in C_c^{\infty}(I, \mathfrak{X}_{00})$ can be quite restrictive, since the heat flow (much like the Littlewood-Paley projection) does not preserve compact supports in \mathring{M} . We need a notion that is more in tune with our theory.

Definition 14. We say $(\mathcal{V}, \mathfrak{p})$ is a **Hodge weak solution** to the Euler equation when $\mathcal{V} \in L^2_{\text{loc}}(I, \mathbb{P}L^2\mathfrak{X})$, $\mathfrak{p} \in L^1_{\text{loc}}(I \times M)$ and

$$\forall \mathcal{X} \in C_c^{\infty}\left(I, \mathfrak{X}_N\right) : \iint_{I \times M} \left\langle \mathcal{V}, \partial_t \mathcal{X} \right\rangle + \left\langle \mathcal{V} \otimes \mathcal{V}, \nabla \mathcal{X} \right\rangle + \mathfrak{p} \operatorname{div} \mathcal{X} = 0$$

Now this looks better, since \mathfrak{X}_N is invariant under the heat flow. However, this is a leap of faith we will need to justify later (cf. Subsection 3.3).

As $\mathbb{P}\mathfrak{X} \leq \mathfrak{X}_N$, we can go further and say \mathcal{V} is a **Hodge-Leray weak solution** to the Euler equation when $\mathcal{V} \in L^2_{loc}(I, \mathbb{P}L^2\mathfrak{X})$ and

$$\forall \mathcal{X} \in C_c^{\infty}\left(I, \mathbb{P}\mathfrak{X}\right) : \iint_{I \times M} \left\langle \mathcal{V}, \partial_t \mathcal{X} \right\rangle + \left\langle \mathcal{V} \otimes \mathcal{V}, \nabla \mathcal{X} \right\rangle = 0$$

This would help give a formulation of Onsager's conjecture that does not depend on the pressure, similar to [RRS18].

Next, we look at the conditions for \mathcal{V} and \mathfrak{p} near ∂M . In [BT17], they assumed $\mathcal{V} \in L^3_t C^{0,\alpha} \mathfrak{X}_N$ with $\alpha \in (\frac{1}{3}, 1)$. In [NN18], they assumed $\mathcal{V} \in L^3_t B^{\alpha}_{3,\infty} \mathfrak{X}$ ($\alpha \in (\frac{1}{3}, 1)$) with a more general "strip decay" condition:

- $\|\mathcal{V}\|_{L^3_t L^3(M_{< r}, \operatorname{avg})}^2 \|\langle \mathcal{V}, \widetilde{\nu} \rangle\|_{L^3_t L^3(M_{< r}, \operatorname{avg})} \xrightarrow{r\downarrow 0} 0$
- $\|\mathfrak{p}\|_{L^{\frac{3}{2}}_{t}L^{\frac{3}{2}}(M_{< r}, \operatorname{avg})} \|\langle \mathcal{V}, \widetilde{\nu} \rangle\|_{L^{3}_{t}L^{3}(M_{< r}, \operatorname{avg})} \xrightarrow{r\downarrow 0} 0.$

In [Bar+19] (the most recent result), they assumed $\mathcal{V} \in L^3_t \underline{B}^{1/3}_{3,\text{VMO}} \mathfrak{X}$ (see the paper for the full definition), along with a minor relaxation for the "strip decay" condition:

$$\left\| \left(\frac{\left| \mathcal{V} \right|^2}{2} + \mathfrak{p} \right) \left\langle \mathcal{V}, \widetilde{\nu} \right\rangle \right\|_{L^1_t L^1 \left(M_{[\frac{r}{4}, \frac{r}{2}]}, \operatorname{avg} \right)} \xrightarrow{r \downarrow 0} 0$$

When $\mathcal{V} \in L^3_t B^{\frac{1}{3}}_{3,1} \mathfrak{X}$, $\|\langle \mathcal{V}, \widetilde{\nu} \rangle\|_{L^3_t L^3(M_{< r}, \operatorname{avg})} \xrightarrow{r\downarrow 0} \|\langle \mathcal{V}, \nu \rangle\|_{L^3_t L^3(\partial M, \operatorname{avg})}$ by Fact 5. This motivates our formulation later in Subsection 3.5, where we put $\mathcal{V} \in L^3_t \mathbb{P} B^{\frac{1}{3}}_{3,1} \mathfrak{X}_N$.

3.3 Justification of formulation

We define the cutoffs

$$\psi_r(x) = \Psi_r\left(\operatorname{dist}\left(x, \partial M\right)\right) \tag{3}$$

where r > 0 small, $\Psi_r \in C^{\infty}([0,\infty), [0,\infty))$ such that $\mathbf{1}_{[0,\frac{3}{4}r)} \ge \Psi_r \ge \mathbf{1}_{[0,\frac{r}{2}]}$ and $\|\Psi'_r\|_{\infty} \lesssim \frac{1}{r}$. Then $\nabla \psi_r(x) = f_r(x)\widetilde{\nu}(x)$ where $|f_r(x)| \lesssim \frac{1}{r}$ and $\operatorname{supp} \psi_r \subset M_{< r}$.

Let $(\mathcal{V}, \mathfrak{p})$ be a weak solution to the Euler equation and $\alpha \in (\frac{1}{3}, 1)$. Define different conditions:

- 1. $\mathcal{V} \in L^3_t C^{0,\alpha} \mathfrak{X}_N$.
- $2. \ \mathcal{V} \in L^3_t B^\alpha_{3,\infty} \mathfrak{X} \text{ and } \|\mathcal{V}\|^2_{L^3_t L^3(M_{< r}, \mathrm{avg})} \|\langle \mathcal{V}, \widetilde{\nu} \rangle\|_{L^3_t L^3(M_{< r}, \mathrm{avg})} \xrightarrow{r \downarrow 0} 0.$
- 3. $\mathcal{V} \in L^3_t B^{\frac{1}{3}}_{3,1} \mathfrak{X}_N.$
- 4. $(\mathcal{V}, \mathfrak{p})$ is a Hodge weak solution.
- 5. \mathcal{V} is a Hodge-Leray weak solution.

Theorem 15. We have $(1) \implies (2) \implies (3) \implies (4) \implies (5)$.

Proof. By Fact 2,
$$C^{0,\alpha}\mathfrak{X}_N = B^{\alpha}_{\infty,\infty}\mathfrak{X}_N \hookrightarrow B^{\alpha}_{3,\infty}\mathfrak{X}_N \hookrightarrow B^{\frac{1}{3}}_{3,1}\mathfrak{X}_N$$
. Then by the coarea formula,
 $\|\langle \mathcal{V}, \widetilde{\nu} \rangle\|^3_{L^3_t L^3(M_{< r}, \operatorname{avg})} \lesssim \|\mathcal{V}\|^2_{L^3_t L^3(M_{< r}, \operatorname{avg})} \|\langle \mathcal{V}, \widetilde{\nu} \rangle\|_{L^3_t L^3(M_{< r}, \operatorname{avg})} \lesssim \|\mathcal{V}\|^2_{L^3_t B^{\frac{1}{3}}_{3,1}\mathfrak{X}} \|\langle \mathcal{V}, \widetilde{\nu} \rangle\|_{L^3_t L^3(M_{< r}, \operatorname{avg})}$

So for $\mathcal{V} \in L^3_t B^{\frac{1}{3}}_{3,1} \mathfrak{X}$: $\|\mathcal{V}\|^2_{L^3_t L^3(M_{< r}, \operatorname{avg})} \|\langle \mathcal{V}, \widetilde{\nu} \rangle\|_{L^3_t L^3(M_{< r}, \operatorname{avg})} \xrightarrow{r\downarrow 0} 0 \iff \|\langle \mathcal{V}, \nu \rangle\|_{L^3_t L^3(\partial M)} = 0 \iff \mathbf{n} \mathcal{V} = 0.$

As (4) \implies (5) is obvious, the only thing left is to show (3) \implies (4). Recall the cutoffs ψ_r from Equation (3).

Let $I_1 \subset I$ be bounded and $\mathcal{X} \in C_c^{\infty}(I_1, \mathfrak{X}_N)$, then $(1 - \psi_r) \mathcal{X} \in C_c^{\infty}(I, \mathfrak{X}_{00})$, and so by the definition of weak solution:

$$0 = \iint_{I \times M} (1 - \psi_r) \langle \mathcal{V}, \partial_t \mathcal{X} \rangle + \langle \mathcal{V}, \nabla_{\mathcal{V}} ((1 - \psi_r) \mathcal{X}) \rangle + \mathfrak{p} \operatorname{div} ((1 - \psi_r) \mathcal{X})$$
$$= \iint_{I \times M} (1 - \psi_r) (\langle \mathcal{V}, \partial_t \mathcal{X} \rangle + \langle \mathcal{V}, \nabla_{\mathcal{V}} \mathcal{X} \rangle + \mathfrak{p} \operatorname{div} \mathcal{X}) - \iint_{I \times M} (\langle \mathcal{V}, \nabla \psi_r \rangle \langle \mathcal{V}, \mathcal{X} \rangle + \mathfrak{p} \langle \mathcal{X}, \nabla \psi_r \rangle)$$

We are done if the first term goes to zero as $r \downarrow 0$. So we only need to show the second term goes to zero. Since $\nabla \psi_r = f_r \tilde{\nu}$ and $\operatorname{supp} \psi_r \subset M_{< r}$, we only need to bound

$$\begin{split} & \left| \iint_{I_1 \times M_{< r}} f_r \left\langle \mathcal{V}, \widetilde{\nu} \right\rangle \left\langle \mathcal{V}, \mathcal{X} \right\rangle + \mathfrak{p} f_r \left\langle \mathcal{X}, \widetilde{\nu} \right\rangle \right| \\ \lesssim & \frac{1}{r} \left\| \mathcal{V} \right\|_{L^3_t L^3(M_{< r})} \left\| \left\langle \mathcal{V}, \widetilde{\nu} \right\rangle \right\|_{L^3_t L^3(M_{< r})} \left\| \mathcal{X} \right\|_{L^3_t L^3(M_{< r})} + \frac{1}{r} \left\| \mathfrak{p} \right\|_{L^1(I_1 \times M_{< r})} \left\| \left\langle \mathcal{X}, \widetilde{\nu} \right\rangle \right\|_{L^\infty_t L^\infty(M_{< r})} \\ \lesssim & \left\| \mathcal{V} \right\|_{L^3_t L^3(M_{< r}, \operatorname{avg})} \left\| \left\langle \mathcal{V}, \widetilde{\nu} \right\rangle \right\|_{L^3_t L^3(M_{< r}, \operatorname{avg})} \left\| \mathcal{X} \right\|_{L^3_t L^3(M_{< r}, \operatorname{avg})} + \left\| \mathfrak{p} \right\|_{L^1(I_1 \times M_{< r})} \left\| \left\langle \mathcal{X}, \widetilde{\nu} \right\rangle \right\|_{L^\infty_t C^{0,1}(M_{< r})} \\ \lesssim & \left\| \mathcal{V} \right\|_{L^3_t B^{\frac{1}{3}}_{3,1} \mathfrak{X}} \left\| \left\langle \mathcal{V}, \widetilde{\nu} \right\rangle \right\|_{L^3_t L^3(M_{< r}, \operatorname{avg})} \left\| \mathcal{X} \right\|_{L^3_t B^{\frac{1}{3}}_{3,1} \mathfrak{X}} + \left\| \mathfrak{p} \right\|_{L^1(I_1 \times M_{< r})} \left\| \left\langle \mathcal{X}, \widetilde{\nu} \right\rangle \right\|_{L^\infty_t C^{0,1}(M_{< r})} \qquad \xrightarrow{r \downarrow 0} 0 \\ \text{We used the estimate } & \left\| \left\langle \mathcal{X}, \widetilde{\nu} \right\rangle \right\|_{L^\infty(M_{< r})} \lesssim r \left\| \left\langle \mathcal{X}, \widetilde{\nu} \right\rangle \right\|_{C^{0,1}(M_{< r})} \text{ since } \left\langle \mathcal{X}, \nu \right\rangle = 0 \text{ on } \partial M. \qquad \Box \end{split}$$

Remark. Interestingly, as Subsection 3.5 will show, no "strip decay" condition involving \mathfrak{p} seems to be necessary. See the end of Subsection 1.1 for a discussion of this minor improvement.

We briefly note that when $\partial M = \emptyset$, it is customary to set dist $(x, \partial M) = \infty$, and $\psi_r = 0$, $M_{>r} = M = M$, $M_{< r} = \emptyset$, and $\mathscr{D}_N \mathfrak{X} M = \mathscr{D} \mathfrak{X} M = \mathfrak{X} M$.

3.4 Heating the nonlinear term

Let $U, V \in B_{3,1}^{\frac{1}{3}} \mathfrak{X}$. Then $U \otimes V \in L^1 \mathfrak{X}$ and div $(U \otimes V)$ is defined as a distribution. To apply the heat flow to div $(U \otimes V)$, we need to define $(\operatorname{div} (U \otimes V))^{\flat}$ so that it is heatable. Recall integration by parts:

$$\left\langle \left\langle \operatorname{div}\left(Y\otimes Z\right),X\right\rangle \right\rangle = -\left\langle \left\langle Y\otimes Z,\nabla X\right\rangle \right\rangle + \int_{\partial M}\left\langle \nu,Y\right\rangle \left\langle Z,X\right\rangle \ \forall X,Y,Z\in\mathfrak{X}\left(M\right)$$

Observe that for $X \in \mathfrak{X}$, even though $\langle \langle \operatorname{div}(U \otimes V), X \rangle \rangle$ is not defined, $\int_{\partial M} \langle \nu, U \rangle \langle V, X \rangle - \langle \langle U \otimes V, \nabla X \rangle \rangle$ is well-defined by the trace theorem. So we will define the heatable 1-current $(\operatorname{div}(U \otimes V))^{\flat}$ by

$$\langle \langle \operatorname{div} (U \otimes V), X \rangle \rangle = - \langle \langle U \otimes V, \nabla X \rangle \rangle + \int_{\partial M} \langle \nu, U \rangle \langle V, X \rangle \ \forall X \in \mathscr{D}_N \mathfrak{X} \ (X \text{ is heated})$$

It is continuous on $\mathscr{D}_N \mathfrak{X}$ since $|\langle \langle \operatorname{div}(U \otimes V), X \rangle \rangle| \lesssim ||U||_{B_{3,1}^{\frac{1}{3}}} ||V||_{B_{3,1}^{\frac{1}{3}}} ||X||_{B_{3,1}^{\frac{1}{3}}} + ||U||_{L^3} ||V||_{L^3} ||\nabla X||_{L^3}$. By the same formula and reasoning, we see that $(\operatorname{div}(U \otimes V))^{\flat}$ is not just heatable, but also a continuous linear functional on $(\mathfrak{X}(M), C^{\infty}$ topo).

On the other hand, we can get away with less regularity by assuming $U \in \mathbb{P}L^2 \mathfrak{X}$. Then we simply need to define $\langle \langle \operatorname{div}(U \otimes V), X \rangle \rangle = - \langle \langle U \otimes V, \nabla X \rangle \rangle \quad \forall X \in \mathfrak{X}.$

In short, $(\operatorname{div}(U \otimes V))^{\flat}$ is heatable when $U \in \mathbb{P}L^2 \mathfrak{X}$ and $V \in L^2 \mathfrak{X}$. Consequently, by Theorem 15, when $(\mathcal{V}, \mathfrak{p})$ is a weak solution to the Euler equation and $\mathcal{V} \in L^3_t B^{\frac{1}{3}}_{3,1} \mathfrak{X}_N$: $(\mathcal{V}, \mathfrak{p})$ is a Hodge weak solution and

$$\partial_t \mathcal{V} + \operatorname{div}(\mathcal{V} \otimes \mathcal{V}) + \operatorname{grad} \mathfrak{p} = 0 \text{ in } \mathscr{D}'_N(I, \mathfrak{X}).$$
 (4)

3.5 Proof of Onsager's conjecture

For the rest of the proof, we will write $e^{t\Delta}$ for S(t), as we will not need another heat flow. For $\varepsilon > 0$ and vector field X, we will write X^{ε} for $e^{\varepsilon \Delta}X$.

We opt to formulate the conjecture without mentioning the pressure (see Subsection 3.3 for the justification).

Theorem 16 (Onsager's conjecture). Let M be a compact, oriented Riemannian manifold with no or smooth boundary. Let $\mathcal{V} \in L^3_t \mathbb{P}B^{\frac{1}{3}}_{3,1} \mathfrak{X}_N$ such that $\forall \mathcal{X} \in C^{\infty}_c(I, \mathbb{P}\mathfrak{X}) : \iint_{I \times M} \langle \mathcal{V}, \partial_t \mathcal{X} \rangle + \langle \mathcal{V} \otimes \mathcal{V}, \nabla \mathcal{X} \rangle = 0$ (Hodge-Leray weak solution).

Then we can show

$$\int_{I} \eta'(t) \left\langle \left\langle \mathcal{V}(t), \mathcal{V}(t) \right\rangle \right\rangle \mathrm{d}t = 0 \; \forall \eta \in C_{c}^{\infty}(I)$$

Consequently, $\langle \langle \mathcal{V}(t), \mathcal{V}(t) \rangle \rangle$ is constant for a.e. $t \in I$.

As usual, there is a **commutator estimate** which we will leave for later:

$$\int_{I} \eta \left\langle \left\langle \operatorname{div} \left(\mathcal{U} \otimes \mathcal{U} \right)^{2\varepsilon}, \mathcal{U}^{2\varepsilon} \right\rangle \right\rangle - \int_{I} \eta \left\langle \left\langle \operatorname{div} \left(\mathcal{U}^{2\varepsilon} \otimes \mathcal{U}^{2\varepsilon} \right), \mathcal{U}^{2\varepsilon} \right\rangle \right\rangle \\
= \int_{I} \eta \left\langle \left\langle \operatorname{div} \left(\mathcal{U} \otimes \mathcal{U} \right)^{3\varepsilon}, \mathcal{U}^{\varepsilon} \right\rangle \right\rangle - \int_{I} \eta \left\langle \left\langle \operatorname{div} \left(\mathcal{U}^{2\varepsilon} \otimes \mathcal{U}^{2\varepsilon} \right)^{\varepsilon}, \mathcal{U}^{\varepsilon} \right\rangle \right\rangle \xrightarrow{\varepsilon \downarrow 0} 0 \tag{5}$$

for all $\mathcal{U} \in L^3_t \mathbb{P}B^{\frac{1}{3}}_{3,1}\mathfrak{X}_N, \eta \in C^{\infty}_c(I).$

Notation: we write div $(\mathcal{U} \otimes \mathcal{U})^{\varepsilon}$ for $(\operatorname{div} (\mathcal{U} \otimes \mathcal{U}))^{\varepsilon}$ and $\nabla \mathcal{U}^{\varepsilon}$ for $\nabla (\mathcal{U}^{\varepsilon})$ (recall that the heat flow does not work on tensors $\mathcal{U} \otimes \mathcal{U}$ and $\nabla \mathcal{U}$). Compared with [IO15], our commutator estimate looks a bit different, to ease some integration by parts procedures down the line.

Remark. For any U in $\mathbb{P}L^2\mathfrak{X}$, div $(U \otimes U)^{\flat}$ is a heatable 1-current (see Subsection 3.4). In particular, for $\varepsilon > 0$, div $(U \otimes U)^{\varepsilon}$ is smooth and

$$\langle \langle \operatorname{div} \left(U \otimes U \right)^{\varepsilon}, Y \rangle \rangle = - \langle \langle U \otimes U, \nabla \left(Y^{\varepsilon} \right) \rangle \rangle \quad \forall Y \in \mathfrak{X}$$

$$(6)$$

Consequently, Equation (5) is well-defined.

Theorem 17 (Onsager). Assume Equation (5) is true. Then $\int_{I} \eta'(t) \langle \langle \mathcal{V}(t), \mathcal{V}(t) \rangle \rangle dt = 0$.

Proof. Let $\Phi \in C_c^{\infty}(\mathbb{R})$ and $\Phi_{\tau} \xrightarrow{\tau \downarrow 0} \delta_0$ be a radially symmetric mollifier. Write $\mathcal{V}^{\varepsilon}$ for $e^{\varepsilon \Delta} \mathcal{V}$ (spatial mollification) and \mathcal{V}_{τ} for $\Phi_{\tau} * \mathcal{V}$ (temporal mollification). First, we mollify in time and space

$$\frac{1}{2} \int_{I} \eta' \left\langle \left\langle \mathcal{V}, \mathcal{V} \right\rangle \right\rangle \stackrel{\text{DCT}}{=} \liminf_{\varepsilon \downarrow 0} \lim_{\tau \downarrow 0} \frac{1}{2} \int_{I} \eta' \left\langle \left\langle \mathcal{V}_{\tau}^{\varepsilon}, \mathcal{V}_{\tau}^{\varepsilon} \right\rangle \right\rangle$$

Then we want to get rid of the time derivative:

$$\frac{1}{2}\int_{I}\eta'\left\langle\left\langle\mathcal{V}_{\tau}^{\varepsilon},\mathcal{V}_{\tau}^{\varepsilon}\right\rangle\right\rangle = -\int_{I}\eta\left\langle\left\langle\partial_{t}\mathcal{V}_{\tau}^{\varepsilon},\mathcal{V}_{\tau}^{\varepsilon}\right\rangle\right\rangle = -\int_{I}\left\langle\left\langle\partial_{t}\left(\eta\mathcal{V}_{\tau}^{\varepsilon}\right),\mathcal{V}_{\tau}^{\varepsilon}\right\rangle\right\rangle + \int_{I}\eta'\left\langle\left\langle\mathcal{V}_{\tau}^{\varepsilon},\mathcal{V}_{\tau}^{\varepsilon}\right\rangle\right\rangle$$

Then we use the definition of Hodge-Leray weak solution, and exploit the commutativity between spatial and temporal operators:

$$\begin{split} \frac{1}{2} \int_{I} \eta' \left\langle \left\langle \mathcal{V}_{\tau}^{\varepsilon}, \mathcal{V}_{\tau}^{\varepsilon} \right\rangle \right\rangle &= \int_{I} \left\langle \left\langle \partial_{t} \left(\eta \mathcal{V}_{\tau}^{\varepsilon} \right), \mathcal{V}_{\tau}^{\varepsilon} \right\rangle \right\rangle \\ &= - \int_{I} \left\langle \left\langle \nabla \left[\left(\eta \mathcal{V}_{\tau}^{2\varepsilon} \right)_{\tau} \right], \mathcal{V} \otimes \mathcal{V} \right\rangle \right\rangle \\ &= - \int_{I} \left\langle \left\langle \left[\eta \left(\nabla \mathcal{V}_{\tau}^{2\varepsilon} \right)_{\tau} \right], \mathcal{V} \otimes \mathcal{V} \right\rangle \right\rangle \\ &= - \int_{I} \left\langle \left\langle \left[\eta \left(\nabla \mathcal{V}_{\tau}^{2\varepsilon} \right)_{\tau}, \mathcal{V} \otimes \mathcal{V} \right\rangle \right\rangle \\ &= - \int_{I} \eta \left\langle \left\langle \left(\nabla \mathcal{V}^{2\varepsilon} \right)_{\tau}, (\mathcal{V} \otimes \mathcal{V})_{\tau} \right\rangle \right\rangle \end{split}$$

As there is no longer a time derivative on \mathcal{V} , we get rid of τ by letting $\tau \downarrow 0$ (fine as \mathcal{V} is L^3 in time). Recall Equation (6):

$$\begin{split} \frac{1}{2} \int_{I} \eta' \left\langle \left\langle \mathcal{V}^{\varepsilon}, \mathcal{V}^{\varepsilon} \right\rangle \right\rangle &= -\int_{I} \eta \left\langle \left\langle \nabla \left(\mathcal{V}^{2\varepsilon} \right), \mathcal{V} \otimes \mathcal{V} \right\rangle \right\rangle = \int_{I} \eta \left\langle \left\langle \mathcal{V}^{\varepsilon}, \operatorname{div} \left(\mathcal{V} \otimes \mathcal{V} \right)^{\varepsilon} \right\rangle \right\rangle \\ &= \int_{I} \eta \left\langle \left\langle \mathcal{V}^{\varepsilon}, \operatorname{div} \left(\mathcal{V}^{\varepsilon} \otimes \mathcal{V}^{\varepsilon} \right) \right\rangle \right\rangle + o_{\varepsilon}(1) \\ &= \int_{I} \eta \left\langle \left\langle \mathcal{V}^{\varepsilon}, \nabla_{\mathcal{V}^{\varepsilon}} \mathcal{V}^{\varepsilon} \right\rangle \right\rangle + o_{\varepsilon}(1) = \int_{I} \eta \int_{M} \mathcal{V}^{\varepsilon} \left(\frac{|\mathcal{V}^{\varepsilon}|^{2}}{2} \right) + o_{\varepsilon}(1) = o_{\varepsilon}(1) \\ & \text{ as } \mathcal{V}^{\varepsilon} \in \mathbb{P}\mathfrak{X} \end{split}$$

So
$$\frac{1}{2} \int_{I} \eta' \langle \langle \mathcal{V}, \mathcal{V} \rangle \rangle = \lim_{\varepsilon \downarrow 0} \lim_{\tau \downarrow 0} \frac{1}{2} \int_{I} \eta' \langle \langle \mathcal{V}_{\tau}^{\varepsilon}, \mathcal{V}_{\tau}^{\varepsilon} \rangle \rangle = \lim_{\varepsilon \downarrow 0} \frac{1}{2} \int_{I} \eta' \langle \langle \mathcal{V}^{\varepsilon}, \mathcal{V}^{\varepsilon} \rangle \rangle = 0.$$

The proof is short and did not much use the Besov regularity of \mathcal{V} . It is the commutator estimate that presents the main difficulty. We proceed similarly as in [IO15].

Let $\mathcal{U} \in L^3_t \mathbb{P}B^{\frac{1}{3}}_{3,1}\mathfrak{X}_N$. By setting $\mathcal{U}(t)$ to 0 for t in a null set, WLOG $\mathcal{U}(t) \in \mathbb{P}B^{\frac{1}{3}}_{3,1}\mathfrak{X}_N \ \forall t \in I$. Define the commutator

$$\mathcal{W}(t,s) = \operatorname{div} \left(\mathcal{U}(t) \otimes \mathcal{U}(t) \right)^{3s} - \operatorname{div} \left(\mathcal{U}(t)^{2s} \otimes \mathcal{U}(t)^{2s} \right)$$

When t and s are implicitly understood, we will not write them. As div $(\mathcal{U}(t) \otimes \mathcal{U}(t))^{3s}$ solves $(\partial_s - 3\Delta) \mathcal{X} = 0$, we define $\mathcal{N} = (\partial_s - 3\Delta) \mathcal{W}$. Then \mathcal{W} and \mathcal{N} obey the Duhamel formula:

Lemma 18 (Duhamel formulas).

- 1. $\mathcal{W}(t,s) \xrightarrow{s\downarrow 0} 0$ in $\mathscr{D}'_N \mathfrak{X}$ and therefore in $\mathscr{D}'\mathfrak{X}$. Furthermore, $\mathcal{W}(\cdot,s) \xrightarrow{s\downarrow 0} 0$ in $\mathscr{D}'_N(I,\mathfrak{X})$ and therefore in $\mathscr{D}'(I,\mathfrak{X})$ (spacetime distribution).
- 2. For fixed $t_0 \in I$ and s > 0: $\int_{\varepsilon}^{s} \mathcal{N}(t_0, \sigma)^{3(s-\sigma)} d\sigma \xrightarrow{\varepsilon \downarrow 0} \mathcal{W}(t_0, s)$ in $\mathscr{D}'_N \mathfrak{X}$.

Proof.

1. Let $X \in \mathscr{D}_N \mathfrak{X}, \mathcal{X} \in C_c^{\infty}(I, \mathscr{D}_N \mathfrak{X})$. It is trivial to check (with DCT)

$$\left\langle \left\langle \mathcal{U}(t) \otimes \mathcal{U}(t), \nabla \left(X^{3s} \right) \right\rangle \right\rangle - \left\langle \left\langle \mathcal{U}(t)^{2s} \otimes \mathcal{U}(t)^{2s}, \nabla \left(X^{s} \right) \right\rangle \right\rangle \xrightarrow{s\downarrow 0} 0$$

$$\int_{I} \left\langle \left\langle \mathcal{U} \otimes \mathcal{U}, \nabla \left(\mathcal{X}^{3s} \right) \right\rangle \right\rangle - \int_{I} \left\langle \left\langle \mathcal{U}^{2s} \otimes \mathcal{U}^{2s}, \nabla \left(\mathcal{X}^{s} \right) \right\rangle \right\rangle \xrightarrow{s\downarrow 0} 0$$

2. Let $\varepsilon > 0$. By the smoothing effect of $e^{s\Delta}$, $\mathcal{W}(t_0, \cdot)$ and $\mathcal{N}(t_0, \cdot)$ are in $C^0_{\text{loc}}((0, 1], \mathscr{D}_N \mathfrak{X})$. As $(e^{s\Delta})_{s>0}$ is a C_0 semigroup on $(H^m - \operatorname{cl}(\mathscr{D}_N \mathfrak{X}), \|\cdot\|_{H^m}) \ \forall m \in \mathbb{N}_0$, and a semigroup basically corresponds to an ODE (cf. [Tay11a, Appendix A, Proposition 9.10 & 9.11]), from $\partial_s W =$ $3\Delta \mathcal{W} + \mathcal{N}$ for $s \geq \varepsilon$ we get the Duhamel formula

$$\forall s > \varepsilon : \mathcal{W}(t_0, s) = \mathcal{W}(t_0, \varepsilon)^{3(s-\varepsilon)} + \int_{\varepsilon}^{s} \mathcal{N}(t_0, \sigma)^{3(s-\sigma)} \,\mathrm{d}\sigma$$

So we only need to show $\mathcal{W}(t_0,\varepsilon)^{3(s-\varepsilon)} \xrightarrow{\mathscr{D}'_N\mathfrak{X}} 0$. Let $X \in \mathscr{D}_N\mathfrak{X}$.

$$\left\langle \left\langle X, \mathcal{W}\left(t_{0},\varepsilon\right)^{3(s-\varepsilon)}\right\rangle \right\rangle = \left\langle \left\langle X^{3(s-\varepsilon)}, \operatorname{div}\left(\mathcal{U}\left(t_{0}\right)\otimes\mathcal{U}\left(t_{0}\right)\right)^{3\varepsilon}\right\rangle \right\rangle - \left\langle \left\langle X^{3(s-\varepsilon)}, \operatorname{div}\left(\mathcal{U}\left(t_{0}\right)^{2\varepsilon}\otimes\mathcal{U}\left(t_{0}\right)^{2\varepsilon}\right)^{\varepsilon}\right\rangle \right\rangle$$
$$= -\left\langle \left\langle \nabla\left(X^{3s}\right), \mathcal{U}\left(t_{0}\right)\otimes\mathcal{U}\left(t_{0}\right)\right\rangle \right\rangle + \left\langle \left\langle \nabla\left(X^{3s-2\varepsilon}\right), \mathcal{U}\left(t_{0}\right)^{2\varepsilon}\otimes\mathcal{U}\left(t_{0}\right)^{2\varepsilon}\right\rangle \right\rangle \xrightarrow{\varepsilon\downarrow 0} 0.$$

that $\left\|\mathcal{U}\left(t\right)^{s}\right\|_{B^{2+\frac{1}{3}}_{3,1}} \lesssim$

From now on, we write \int_{0+}^{s} for $\lim_{\varepsilon \downarrow 0} \int_{\varepsilon}^{s}$. Then

$$\int_{I} \mathrm{d}t \ \eta(t) \left\langle \left\langle \mathcal{W}(t,s), \mathcal{U}(t)^{s} \right\rangle \right\rangle = \int_{I} \mathrm{d}t \ \eta(t) \int_{0+}^{s} \mathrm{d}\sigma \left\langle \left\langle \mathcal{N}(t,\sigma)^{3(s-\sigma)}, \mathcal{U}(t)^{s} \right\rangle \right\rangle$$

To clean up the algebra, we will classify the terms that are going to appear but are actually negligible in the end. The following estimates lie at the heart of the problem, showing why the regularity needs to be at least $\frac{1}{3}$, and that our argument barely holds thanks to the pointwise vanishing property (Corollary 12).

Lemma 19 (3 error estimates). Define the k-jet fiber norm $|X|_{J^k} = \left(\sum_{i=0}^k |\nabla^{(i)}X|^2\right)^{\frac{1}{2}} \quad \forall X \in \mathfrak{X} \text{ (more }$ details in Subsection 6.1.1). Then we have

Now we can prove the error estimates go to 0:

$$\begin{split} 1. \\ & \int_{I} |\eta| \int_{0+}^{s} \mathrm{d}\sigma \int_{M} \left| \mathcal{U}^{2\sigma} \right|_{J^{1}}^{2} \left| \mathcal{U}^{4s-2\sigma} \right|_{J^{1}} \lesssim \int_{I} |\eta| \int_{0+}^{s} \mathrm{d}\sigma \left\| \mathcal{U}^{2\sigma} \right\|_{W^{1,3}}^{2} \left\| \mathcal{U}^{4s-2\sigma} \right\|_{W^{1,3}} \\ & \lesssim \int_{I} \mathrm{d}t \left| \eta(t) \right| \int_{0+}^{s} \mathrm{d}\sigma \left(\frac{1}{\sigma} \right)^{\frac{2}{3}} \left(\frac{1}{2s-\sigma} \right)^{\frac{1}{3}} A(t, 2\sigma)^{2} A(t, 4s-2\sigma) \\ & \stackrel{\sigma \mapsto s\sigma}{=} \int_{I} \mathrm{d}t \left| \eta(t) \right| \int_{0+}^{1} \mathrm{d}\sigma \ \left(\frac{1}{\sigma} \right)^{\frac{2}{3}} \left(\frac{1}{2-\sigma} \right)^{\frac{1}{3}} A(t, 2s\sigma)^{2} A(t, 4s-2s\sigma) \lesssim \int_{I} \mathrm{d}t \ |\eta(t)| \left\| A(t, \sigma) \right\|_{L^{\infty}_{\sigma \leq 4s}}^{3} \xrightarrow{s\downarrow 0} 0. \end{split}$$

$$2.$$

$$\begin{split} & \int_{I} |\eta| \int_{0+}^{s} \mathrm{d}\sigma \int_{\partial M} \left| \mathcal{U}^{2\sigma} \right|^{2} \left| \mathcal{U}^{4s-2\sigma} \right|_{J^{2}} \lesssim \int_{I} |\eta| \int_{0+}^{s} \mathrm{d}\sigma \left\| \mathcal{U}^{2\sigma} \right\|_{L^{3}\mathfrak{X}M|_{\partial M}}^{2} \left\| \mathcal{U}^{4s-2\sigma} \right\|_{W^{2,3}\mathfrak{X}M|_{\partial M}} \\ \stackrel{\mathrm{Trace}}{\lesssim} \int_{I} |\eta| \int_{0+}^{s} \mathrm{d}\sigma \left\| \mathcal{U}^{2\sigma} \right\|_{B^{\frac{1}{3},1}_{3,1}\mathfrak{X}M}^{2} \left\| \mathcal{U}^{4s-2\sigma} \right\|_{B^{2+\frac{1}{3}}_{3,1}\mathfrak{X}M} \\ \lesssim \int_{I} \mathrm{d}t \ |\eta(t)| \left\| \mathcal{U}(t) \right\|_{B^{\frac{1}{3},1}_{3,1}\mathfrak{X}M}^{2} \int_{0+}^{s} \mathrm{d}\sigma \left(\frac{1}{2s-\sigma} \right) A\left(t, 4s-2\sigma \right) \\ \stackrel{\sigma \mapsto s\sigma}{=} \int_{I} \mathrm{d}t \ |\eta(t)| \left\| \mathcal{U}(t) \right\|_{B^{\frac{1}{3},1}_{3,1}\mathfrak{X}M}^{2} \int_{0+}^{1} \mathrm{d}\sigma \left(\frac{1}{2-\sigma} \right) A\left(t, 4s-2s\sigma \right) \lesssim \int_{I} \mathrm{d}t \ |\eta(t)| \left\| \mathcal{U}(t) \right\|_{B^{\frac{1}{3},1}_{3,1}\mathfrak{X}M}^{2} \left\| A(t,\sigma) \right\|_{L^{\infty}_{\sigma \leq 4s}} \\ \lesssim \left\| \mathcal{U} \right\|_{L^{3}_{t}B^{\frac{1}{3},1}_{3,1}(M)}^{2} \left\| \left\| A(t,\sigma) \right\|_{L^{\infty}_{\sigma \leq 4s}} \left\| L^{3}_{t} \right\|_{T^{3}}^{s\downarrow 0} 0 \end{split}$$

$$\begin{split} & \int_{I} |\eta| \int_{0+}^{s} \mathrm{d}\sigma \int_{\partial M} \left| \mathcal{U}^{2\sigma} \right| \left| \mathcal{U}^{2\sigma} \right|_{J^{1}} \left| \mathcal{U}^{4s-2\sigma} \right|_{J^{1}} \\ & \overset{\mathrm{Trace}}{\lesssim} \int_{I} |\eta| \int_{0+}^{s} \mathrm{d}\sigma \left\| \mathcal{U}^{2\sigma} \right\|_{B^{\frac{1}{3}}_{3,1} \mathfrak{X}M} \left\| \mathcal{U}^{2\sigma} \right\|_{B^{1+\frac{1}{3}}_{3,1} \mathfrak{X}M} \left\| \mathcal{U}^{4s-2\sigma} \right\|_{B^{1+\frac{1}{3}}_{3,1} \mathfrak{X}M} \\ & \lesssim \int_{I} \mathrm{d}t \, |\eta(t)| \left\| \mathcal{U}(t) \right\|_{B^{\frac{1}{3}}_{3,1}} \int_{0+}^{s} \mathrm{d}\sigma \left(\frac{1}{\sigma} \right)^{\frac{1}{2}} \left(\frac{1}{2s-\sigma} \right)^{\frac{1}{2}} A(t,2\sigma) \, A(t,4s-2\sigma) \\ & \overset{\sigma \mapsto s\sigma}{=} \int_{I} \mathrm{d}t \, |\eta(t)| \left\| \mathcal{U}(t) \right\|_{B^{\frac{1}{3}}_{3,1}} \int_{0+}^{1} \mathrm{d}\sigma \left(\frac{1}{\sigma} \right)^{\frac{1}{2}} \left(\frac{1}{2-\sigma} \right)^{\frac{1}{2}} A(t,2s\sigma) \, A(t,4s-2s\sigma) \\ & \lesssim \int_{I} \mathrm{d}t \, |\eta(t)| \left\| \mathcal{U}(t) \right\|_{B^{\frac{1}{3}}_{3,1}} \left\| A(t,\sigma) \right\|_{L^{\infty}_{\sigma \leq 4s}}^{2} \lesssim \left\| \mathcal{U} \right\|_{L^{3}_{t} B^{\frac{1}{3}}_{3,1}} \left\| \| A(t,\sigma) \right\|_{L^{\infty}_{\sigma \leq 4s}} \left\| \frac{2}{L^{3}_{t}} \frac{s \downarrow 0}{\to} 0 \\ & \Box \end{split}$$

Note that

 $\mathcal{N}(t,\sigma) = (\partial_{\sigma} - 3\Delta) \left(-\operatorname{div} \left(\mathcal{U}^{2\sigma} \otimes \mathcal{U}^{2\sigma} \right)^{\sigma} \right) = -2 \operatorname{div} \left(\Delta \mathcal{U}^{2\sigma} \otimes \mathcal{U}^{2\sigma} \right)^{\sigma} - 2 \operatorname{div} \left(\mathcal{U}^{2\sigma} \otimes \Delta \mathcal{U}^{2\sigma} \right)^{\sigma} + 2\Delta \operatorname{div} \left(\mathcal{U}^{2\sigma} \otimes \mathcal{U}^{2\sigma} \right)^{\sigma}$ Finally, we will do not

Finally, we will show

$$\int_{I} \eta \left\langle \left\langle \mathcal{W}(s), \mathcal{U}^{s} \right\rangle \right\rangle = \int_{I} \mathrm{d}t \; \eta \left(t \right) \left\langle \left\langle \mathcal{W}(t, s), \mathcal{U} \left(t \right)^{s} \right\rangle \right\rangle \xrightarrow{s \downarrow 0} 0$$

Proof. Integrate by parts into 3 components:

$$\begin{split} \int_{I} \eta \left\langle \left\langle \mathcal{W}(s), \mathcal{U}^{s} \right\rangle \right\rangle &= \int_{I} \mathrm{d}t \ \eta \left(t \right) \int_{0+}^{s} \mathrm{d}\sigma \left\langle \left\langle \mathcal{N}\left(t, \sigma \right)^{3(s-\sigma)}, \mathcal{U}\left(t \right)^{s} \right\rangle \right\rangle \\ &= 2 \int_{I} \eta \int_{0+}^{s} \mathrm{d}\sigma \left\langle \left\langle \Delta \mathcal{U}^{2\sigma} \otimes \mathcal{U}^{2\sigma}, \nabla \left(\mathcal{U}^{4s-2\sigma} \right) \right\rangle \right\rangle + 2 \int_{I} \eta \int_{0+}^{s} \mathrm{d}\sigma \left\langle \left\langle \mathcal{U}^{2\sigma} \otimes \Delta \mathcal{U}^{2\sigma}, \nabla \left(\mathcal{U}^{4s-2\sigma} \right) \right\rangle \right\rangle \\ &- 2 \int_{I} \eta \int_{0+}^{s} \mathrm{d}\sigma \left\langle \left\langle \mathcal{U}^{2\sigma} \otimes \mathcal{U}^{2\sigma}, \nabla \left(\Delta \mathcal{U}^{4s-2\sigma} \right) \right\rangle \right\rangle \end{split}$$

Note that for the third component, we used some properties from Fact 11 to move the Laplacian. It also explains our choice of \mathcal{W} .

We now use Penrose notation to estimate the 3 components. To clean up the notation, we only focus on the integral on M, with the other integrals $2 \int_{I} \eta \int_{0+}^{s} d\sigma(\cdot)$ in variables t and σ implicitly understood. We also use **schematic identities** for linear combinations of similar-looking tensor terms where we do not care how the indices contract (recall Equation (2)). By the error estimates above, all the terms with R or ν will be negligible as $s \downarrow 0$, and interchanging derivatives will be a free action. We write \approx to throw the negligible error terms away. Also, when we write $(\nabla_j \mathcal{U}_l)^{4s-2\sigma}$, we mean the heat flow is applied to \mathcal{U} , not $\nabla \mathcal{U}$ (which is not possible anyway).

First component:

$$\int_{M} \left\langle \Delta \mathcal{U}^{2\sigma} \otimes \mathcal{U}^{2\sigma}, \nabla \left(\mathcal{U}^{4s-2\sigma} \right) \right\rangle = \int_{M} \frac{R * \mathcal{U}^{2\sigma} * \mathcal{U}^{2\sigma} * \nabla \left(\mathcal{U}^{4s-2\sigma} \right)}{\left(\mathcal{U}^{i} \nabla^{i} \mathcal{U}^{j} \right)^{2\sigma} \left(\mathcal{U}^{l} \right)^{2\sigma} \left(\nabla_{j} \mathcal{U}_{l} \right)^{4s-2\sigma}} - \int_{M} \frac{\left(\nabla^{i} \mathcal{U}^{j} \right)^{2\sigma} \left(\nabla_{i} \mathcal{U}^{l} \right)^{2\sigma} \left(\nabla_{j} \mathcal{U}_{l} \right)^{4s-2\sigma}}{\left(\nabla_{i} \mathcal{U}^{j} \right)^{2\sigma} \left(\nabla_{i} \mathcal{U}_{l} \right)^{2\sigma} \left(\nabla_{i} \nabla_{j} \mathcal{U}_{l} \right)^{4s-2\sigma}} - \int_{M} \frac{\left(\nabla^{i} \mathcal{U}^{j} \right)^{2\sigma} \left(\nabla_{i} \mathcal{U}_{l} \right)^{2\sigma} \left(\nabla_{j} \mathcal{U}_{l} \right)^{4s-2\sigma}}{\left(\nabla_{i} \mathcal{U}^{j} \right)^{2\sigma} \left(\nabla_{i} \nabla_{j} \mathcal{U}_{l} \right)^{4s-2\sigma}}$$

Second component:

$$\begin{split} &\int_{M} \left\langle \mathcal{U}^{2\sigma} \otimes \Delta \mathcal{U}^{2\sigma}, \nabla \left(\mathcal{U}^{4s-2\sigma} \right) \right\rangle = \underbrace{\int_{\mathcal{M}} \mathcal{U}^{2\sigma} * R * \mathcal{U}^{2\sigma} * \nabla \left(\mathcal{U}^{4s-2\sigma} \right)}_{\mathcal{M}} + \int_{M} \left(\mathcal{U}^{j} \right)^{2\sigma} \left(\nabla_{i} \nabla^{i} \mathcal{U}^{l} \right)^{2\sigma} \left(\nabla_{j} \mathcal{U}_{l} \right)^{4s-2\sigma} \\ \approx \underbrace{\int_{\partial \mathcal{M}} \left(\mathcal{U}^{j} \right)^{2\sigma} \left(\nabla_{i} \mathcal{U}^{l} \right)^{2\sigma} \left(\nabla_{j} \mathcal{U}_{l} \right)^{4s-2\sigma}}_{-\int_{M} \left(\mathcal{U}^{j} \right)^{2\sigma} \left(\nabla^{i} \mathcal{U}^{l} \right)^{2\sigma} \left(\nabla_{i} \mathcal{U}_{j} \right)^{4s-2\sigma}} \\ = \int_{M} \left(\mathcal{U}^{j} \right)^{2\sigma} \left(\nabla^{i} \mathcal{U}^{l} \right)^{2\sigma} \left(\nabla_{i} \nabla_{j} \mathcal{U}_{l} \right)^{4s-2\sigma} \end{split}$$

For the third component, note $\nabla (R * U) = \nabla R * U + R * \nabla U$

$$-\int_{M} \left\langle \mathcal{U}^{2\sigma} \otimes \mathcal{U}^{2\sigma}, \nabla \left(\Delta \mathcal{U}^{4s-2\sigma} \right) \right\rangle = -\int_{M} \mathcal{U}^{2\sigma} * \mathcal{U}^{2\sigma} * \nabla \left(R * \mathcal{U}^{4s-2\sigma} \right) - \int_{M} \left(\mathcal{U}^{j} \right)^{2\sigma} \left(\mathcal{U}^{l} \right)^{2\sigma} \left(\nabla_{j} \nabla^{i} \nabla_{i} \mathcal{U}_{l} \right)^{4s-2\sigma} \\ \approx -\int_{M} \left(\mathcal{U}^{j} \right)^{2\sigma} \left(\mathcal{U}^{l} \right)^{2\sigma} \left(\underline{R} * \mathcal{U}^{4s-2\sigma} \right) + \nabla^{i} \nabla_{j} \nabla_{i} \mathcal{U}_{l}^{4s-2\sigma} \right) \\ \approx -\int_{M} \left(\mathcal{U}^{j} \right)^{2\sigma} \left(\mathcal{U}^{l} \right)^{2\sigma} \left(\underline{\nabla} \left(\underline{R} * \mathcal{U}^{4s-2\sigma} \right) + \nabla^{i} \nabla_{i} \nabla_{j} \mathcal{U}_{l}^{4s-2\sigma} \right) \\ \approx -\int_{\partial M} \left(\mathcal{U}^{j} \right)^{2\sigma} \left(\mathcal{U}^{l} \right)^{2\sigma} \left(\nabla^{i} \nabla_{i} \nabla_{j} \mathcal{U}_{l} \right)^{4s-2\sigma} + \int_{M} \left(\nabla^{i} \mathcal{U}^{j} \right)^{2\sigma} \left(\nabla_{i} \nabla_{j} \mathcal{U}_{l} \right)^{4s-2\sigma}$$

$$+ \int_{M} \left(\mathcal{U}^{j} \right)^{2\sigma} \left(\nabla^{i} \mathcal{U}^{l} \right)^{2\sigma} \left(\nabla_{i} \nabla_{j} \mathcal{U}_{l} \right)^{4s-2\sigma}$$

Add them up, and we get 0 as $2 \int_{I} \eta \int_{0+}^{s} \mathrm{d}\sigma \left(\cdot \right) \xrightarrow{s \downarrow 0} 0.$

So we are done and the rest of the paper is to develop the tools we have borrowed for the proof.

4 Functional analysis

4.1 Common tools

We note a useful inequality:

Theorem 20 (Ehrling's inequality). Let X, Y, \widetilde{X} be (real/complex) Banach spaces such that X is reflexive and $X \hookrightarrow \widetilde{X}$ is a continuous injection. Let $T: X \to Y$ be a linear compact operator. Then $\forall \varepsilon > 0, \exists C_{\varepsilon} > 0$:

$$\|Tx\|_{Y} \le \varepsilon \, \|x\|_{X} + C_{\varepsilon} \, \|x\|_{\widetilde{X}} \, \, \forall x \in X$$

Remark. Usually, X is some higher-regularity space than \widetilde{X} (e.g. H^1 and L^2). The inequality is useful when the higher-regularity norm is expensive. We will need this for the L^p -analyticity of the heat flow (Theorem 71).

Proof. Proof by contradiction: Assume $\varepsilon > 0$ and there is $(x_j)_{j \in \mathbb{N}}$ such that $||x_j||_X = 1$ and $||Tx_j||_Y > \varepsilon + j ||x_j||_{\widetilde{X}}$. Since X is reflexive, by Banach-Alaoglu and PTAS, WLOG assume $x_j \xrightarrow{X} x_{\infty}$. Then $Tx_j \xrightarrow{Y} Tx_{\infty}$ and $x_j \xrightarrow{\widetilde{X}} x_{\infty}$. As T is compact, PTAS, WLOG $Tx_j \to Tx_{\infty}$. So $||Tx_{\infty}||_Y \ge \limsup_{j\to\infty} (\varepsilon + j ||x_j||_{\widetilde{X}}) > 0$ and $x_j \xrightarrow{\widetilde{X}} 0$. Then $x_j \xrightarrow{\widetilde{X}} 0$ and $x_{\infty} = 0$, contradicting $||Tx_{\infty}||_Y > 0$.

Definition 21 (Banach-valued holomorphic functions). Let $\Omega \subset \mathbb{C}$ be an open set and X be a complex Banach space. Then a function $f : \Omega \to X$ is said to be **holomorphic** (or **analytic**) when $\forall z \in \Omega$: $f'(z) := \lim_{|h|\to 0} \frac{f(z+h)-f(z)}{h}$ exists. The words "holomorphic" and "analytic" are mostly interchangeable, but "analytic" stresses the existence of power series expansion and can also describe functions on \mathbb{R} for which analytic continuation into the complex plane exists.

Theorem 22 (Identity theorem). Let X be a complex Banach space and $X_0 \leq X$ closed. Let $\Omega \subset \mathbb{C}$ be connected, open and $f : \Omega \to X$ holomorphic. Assume there is a sequence $(z_j)_{j \in \mathbb{N}}$ such that $z_j \to z \in \Omega$ and $f(z_j) \in X_0 \forall j$. Then $f(\Omega) \subset X_0$.

Proof. Let $\Lambda \in X^*$ such that $\Lambda(X_0) = 0$. Reduce this to the scalar version in complex analysis. \Box

In fact, many theorems from scalar complex analysis similarly carry over via linear functionals (cf. [Rud91, Theorem 3.31]).

4.2 Interpolation theory

We will quickly review the theory of complex and real interpolation, and state the abstract Stein interpolation theorem. Interpolation theory can be seen as vast generalizations of the Marcinkiewicz and Riesz-Thorin interpolation theorems. **Definition 23.** An interpolation couple of (real/complex) Banach spaces is a pair (X_0, X_1) of Banach spaces with a Hausdorff TVS \mathcal{X} such that $X_0 \hookrightarrow \mathcal{X}, X_1 \hookrightarrow \mathcal{X}$ are continuous injections. Then $X_0 \cap X_1$ and $X_0 + X_1$ are Banach spaces under the norms

$$\|x\|_{X_0 \cap X_1} = \max\left(\|x\|_{X_0}, \|x\|_{X_1}\right) \text{ and } \|x\|_{X_0 + X_1} = \inf_{x = x_0 + x_1, x_j \in X_j} \|x_0\|_{X_0} + \|x_1\|_{X_1}$$

Let (Y_0, Y_1) be another interpolation couple. We say $T : (X_0, X_1) \to (Y_0, Y_1)$ is a **morphism** when $T \in \mathcal{L}(X_0 + X_1, Y_0 + Y_1)$ and $T \in \mathcal{L}(X_j, Y_j)$ for j = 0, 1 under domain restriction. That implies $T \in \mathcal{L}(X_0 \cap X_1, Y_0 \cap Y_1)$ and we write $T \in \mathcal{L}((X_0, X_1), (Y_0, Y_1))$. We also write $\mathcal{L}((X_0, X_1)) = \mathcal{L}((X_0, X_1), (X_0, X_1))$.

Let $P \in \mathcal{L}((X_0, X_1))$ such that $P^2 = P$. Then we call P a **projection** on the interpolation couple (X_0, X_1) .

Definition 24. Let (X_0, X_1) be an interpolation couple of (real/complex) Banach spaces. Then define the *J*-functional:

$$J: \quad (0,\infty) \times X_0 \cap X_1 \quad \longrightarrow \mathbb{R}$$
$$(t,x) \qquad \qquad \longmapsto \|x\|_{X_0} + t \, \|x\|_{X_1}$$

For $\theta \in (0, 1), q \in [1, \infty]$, define the real interpolation space

$$(X_0, X_1)_{\theta, q} = \left\{ \sum_{j \in \mathbb{Z}} u_j : u_j \in X_0 \cap X_1, \left(2^{-j\theta} J(2^j, u_j) \right)_{j \in \mathbb{Z}} \in l_j^q(\mathbb{Z}) \right\}$$

which is Banach under the norm $||x||_{(X_0,X_1)_{\theta,q}} = \inf_{\substack{x=\sum_{j\in\mathbb{Z}}u_j\\j\in\mathbb{Z}}u_j} ||2^{-j\theta}J(2^j,u_j)||_{l_j^q}$. Note that $\sum_{j\in\mathbb{Z}}u_j$ denotes a series that converges in $X_0 + X_1$.

• When $q \in [1, \infty]$ and $x \in X_0 \cap X_1$, note that $\forall j \in \mathbb{Z} : x = \sum_{k \in \mathbb{Z}} \delta_{kj} x$ and

$$\|x\|_{(X_0,X_1)_{\theta,q}} \le \inf_{j\in\mathbb{Z}} \left|2^{-j\theta} J(2^j,x)\right| = \inf_{j\in\mathbb{Z}} \left|2^{-j\theta} \|x\|_{X_0} + 2^{j(1-\theta)} \|x\|_{X_1} \right| \sim_{\neg\theta,\neg q} \|x\|_{X_0}^{1-\theta} \|x\|_{X_1}^{\theta}$$

The last estimate comes from AM-GM and shifting j so that $||x||_{X_0} \sim 2^j ||x||_{X_1}$. Note that the implied constants do not depend on θ and q.

- By considering the finite partial sums $\sum_{|j| < j_0} u_j$, we conclude that $X_0 \cap X_1$ is dense in $(X_0, X_1)_{\theta,q}$ when $q \in [1, \infty)$.
- Let (Y_0, Y_1) be another interpolation couple and $T \in \mathcal{L}((X_0, X_1), (Y_0, Y_1))$. For $\theta \in (0, 1), q \in [1, \infty]$, define $X_{\theta,q} = (X_0, X_1)_{\theta,q}, Y_{\theta,q} = (Y_0, Y_1)_{\theta,q}$. Then $T \in \mathcal{L}(X_{\theta,q}, Y_{\theta,q})$ and

$$\|T\|_{\mathcal{L}(X_{\theta,q},Y_{\theta,q})} \lesssim_{\neg\theta,\neg q,\neg T} \|T\|_{\mathcal{L}(X_0,Y_0)}^{1-\theta} \|T\|_{\mathcal{L}(X_1,Y_1)}^{\theta}$$

where the implied constant does not depend on θ and q. This can be proved by a simple shifting argument.

• If P is a projection on (X_0, X_1) then $(PX_0, PX_1)_{\theta,q} = P(X_0, X_1)_{\theta,q}$.

Remark. There is also an equivalent characterization by the K-functional, which we shall omit. This theory can also be extended to quasi-Banach spaces. We refer to [BL76; Tri10] for more details.

Definition 25. Let (X_0, X_1) be an interpolation couple of complex Banach spaces.

Let $\Omega = \{z \in \mathbb{C} : 0 < \operatorname{Re} z < 1\}$. We then define the Banach space of vector-valued holomorphic/analytic functions on the strip:

$$\mathcal{F}_{X_0,X_1} = \{ f \in C^0(\overline{\Omega} \to X_0 + X_1) : f \text{ holomorphic in } \Omega, \ \|f(it)\|_{X_0} + \|f(1+it)\|_{X_1} \xrightarrow{|t| \to \infty} 0 \}$$

with the norm $\|f\|_{\mathcal{F}_{X_0,X_1}} = \max\left(\sup_{t\in\mathbb{R}} \|f(it)\|_{X_0}, \sup_{t\in\mathbb{R}} \|f(1+it)\|_{X_1}\right)$. For $\theta \in [0,1]$, define the **complex interpolation space** $[X_0, X_1]_{\theta} = \{f(\theta) : f \in \mathcal{F}_{X_0,X_1}\}$, which is Banach under the norm

$$\|x\|_{[X_0,X_1]_{\theta}} = \inf_{\substack{f \in \mathcal{F}_{X_0,X_1} \\ f(\theta) = x}} \|f\|_{\mathcal{F}_{X_0,X_1}}$$

• When $x \in X_0 \cap X_1 \setminus \{0\}$, $\theta \in [0,1]$, $\varepsilon > 0$, define $f_{\varepsilon}(z) = e^{\varepsilon(z^2 - \theta^2)} \frac{x}{\|x\|_{X_0}^{1-z} \|x\|_{X_1}^{z}}$. By the freedom in choosing ε , we conclude

$$\|x\|_{[X_0,X_1]_{\theta}} \le \inf_{\varepsilon > 0} \|f_{\varepsilon}\|_{\mathcal{F}_{X_0,X_1}} \|x\|_{X_0}^{1-\theta} \|x\|_{X_1}^{\theta} \le \inf_{\varepsilon > 0} \max\left(e^{\varepsilon(1-\theta^2)}, e^{-\epsilon\theta^2}\right) \|x\|_{X_0}^{1-\theta} \|x\|_{X_1}^{\theta} = \|x\|_{X_0}^{1-\theta} \|x\|_{X_1}^{\theta}$$

• When $\theta \in [0, 1]$, by Poisson summation and Fourier series, we can prove that

$$\mathcal{F}^{0}_{X_{0},X_{1}} = \{ e^{Cz^{2}} \sum_{j=1}^{N} e^{\lambda_{j}z} x_{j} : N \in \mathbb{N}, C > 0, \lambda_{j} \in \mathbb{R}, x_{j} \in X_{0} \cap X_{1} \}$$

is dense in \mathcal{F}_{X_0,X_1} (cf. [BL76, Lemma 4.2.3]). This implies $X_0 \cap X_1$ is dense in $[X_0,X_1]_{\theta}$. There is a simple extension of the above density result. Let U be dense in $X_0 \cap X_1$ and define $A(\Omega) = \{ \phi \in C^0(\overline{\Omega} \to \mathbb{C}) : \phi \text{ holomorphic in } \Omega \}.$ Then

$$\mathcal{F}_{X_0,X_1}^U = \{ e^{Cz^2} \sum_{j=1}^N \phi_j(z) u_j : N \in \mathbb{N}, C > 0, \phi_j \in A(\Omega), u_j \in U \}$$

is dense in \mathcal{F}_{X_0,X_1} . This will lead to the abstract Stein interpolation theorem.

• Let (Y_0, Y_1) be another interpolation couple and $T \in \mathcal{L}((X_0, X_1), (Y_0, Y_1))$. Then for $\theta \in [0, 1]$, almost by the definitions, we conclude

$$\|T\|_{\mathcal{L}([X_0,X_1]_{\theta},[Y_0,Y_1]_{\theta})} \le \|T\|_{\mathcal{L}(X_0,Y_0)}^{1-\theta} \|T\|_{\mathcal{L}(X_1,Y_1)}^{\theta}$$

• If P is a projection on (X_0, X_1) then $[PX_0, PX_1]_{\theta} = P[X_0, X_1]_{\theta}$

Remark. A keen reader would notice that we use square brackets for complex interpolation, and parentheses for real interpolation. One reason is that the real interpolation methods easily extend to quasi-Banach spaces, while the complex interpolation method does not. There is a version of complex interpolation for special quasi-Banach spaces, which is denoted by parentheses (cf. [Tri10, Section 2.4.4]), but we shall omit it for simplicity.

Blackbox 26 (Abstract Stein interpolation). Let (X_0, X_1) and (Y_0, Y_1) be interpolation couples of complex Banach spaces and U dense in $X_0 \cap X_1$. Let $\Omega = \{z \in \mathbb{C} : 0 < \text{Re } z < 1\}$ and $(T(z))_{z \in \overline{\Omega}}$ be a family of linear mappings $T(z): U \to Y_0 + Y_1$ such that

1. $\forall u \in U : (\overline{\Omega} \to Y_0 + Y_1, z \mapsto T(z)u)$ is continuous, bounded and analytic in Ω .

2. For j = 0, 1 and $u \in U$: $(\mathbb{R} \to Y_j, t \mapsto T(j + it)u)$ is continuous and bounded by $M_j ||u||_{X_j}$ for some $M_j > 0$.

Then for $\theta \in [0,1]$, we can conclude

$$\|T(\theta)u\|_{[Y_0,Y_1]_{\theta}} \le M_0^{1-\theta} M_1^{\theta} \|u\|_{[X_0,X_1]_{\theta}} \quad \forall u \in U$$

Consequently, by unique extension, we have $T(\theta) \in \mathcal{L}([X_0, X_1]_{\theta}, [Y_0, Y_1]_{\theta})$.

Proof. See [Voi92], which is a very short read.

Remark. We will only use Stein interpolation in Subsection 4.3.

4.3 Stein extrapolation of analyticity of semigroups

We are inspired by [Fac15, Theorem 3.1.1] (Stein extrapolation) and [Fac15, Theorem 3.1.10] (Kato-Beurling extrapolation), and wish to create variants for our own use. We will focus on Stein extrapolation, since it is simpler to deal with.

There exists a subtle, but very important criterion to establish analyticity/holomorphicity:

Blackbox 27 (Holo on total). Let $\Omega \subset \mathbb{C}$ be open and X complex Banach. Let $f : \Omega \to X$ be a function. Assume $N \leq X^*$ is **total** (separating points) and f is locally bounded. Then f is analytic iff Λf is analytic $\forall \Lambda \in N$.

Proof. This is a consequence of Krein-Smulian and the Vitali holomorphic convergence theorem, and we refer to [Are+11, Theorem A.7]. \Box

Remark. It will quickly become obvious how crucial this criterion is for the rest of the paper. Let us briefly note that an improvement has just been discovered by Arendt *et al.* [ABK19] (the author thanks Stephan Fackler for bringing this news).

Corollary 28 (Inheritance of analyticity). Let $\Omega \subset \mathbb{C}$ be open and X, Y be complex Banach spaces where $j: X \hookrightarrow Y$ is a continuous injection. Let $f: \Omega \to X$ be locally bounded. Then f is analytic iff $j \circ f$ is analytic.

Proof. $\text{Im}(j^*)$ is weak*-dense, therefore total.

Corollary 29 (Evaluation on dense set). Let X, Y be complex Banach spaces with $X_0 \leq X$. Let $\Omega \subset \mathbb{C}$ be open and $f : \Omega \to \mathcal{L}(X, Y)$ be a function. Assume $X_0 \leq X$ is weakly dense and f is locally bounded. Then f is analytic $\iff \forall x_0 \in X_0, f(\cdot)x_0 : \Omega \to Y$ is analytic.

Proof. Consider $N_{X_0} = \operatorname{span}\{y^* \circ \operatorname{ev}_{x_0} : x_0 \in X_0, y^* \in Y^*\} \leq \mathcal{L}(X, Y)^*$. It is total as X_0 is weakly dense. Use Blackbox 27.

4.3.1 Semigroup definitions

As mentioned before, we assume the reader is familiar with basic elements of functional analysis, including semigroup theory as covered in [Tay11a, Appendix A.9].

Unfortunately, definitions vary depending on the authors, so we need to be careful about which ones we are using.

Definition 30. For $\delta \in (0, \pi]$, define $\Sigma_{\delta}^+ = \{z \in \mathbb{C} \setminus \{0\} : |\arg z| < \delta\}, \Sigma_{\delta}^- = -\Sigma_{\delta}^+, \mathbb{D} = \{z \in \mathbb{C} : |z| < 1\}$. Also define $\Sigma_0^+ = (0, \infty)$ and $\Sigma_0^- = -\Sigma_0^+$.

Let X be a complex Banach space.

 $(T(t))_{t\geq 0} \subset \mathcal{L}(X)$ is called:

- a semigroup when $T: [0,\infty) \to \mathcal{L}(X)$ is a monoid homomorphism $(T(0) = 1, T(t_1 + t_2) = T(t_1)T(t_2))$
- degenerate when $T: (0, \infty) \to \mathcal{L}(X)$ is continuous in the SOT (strong operator topology).
- immediately norm-continuous when $T: (0, \infty) \to \mathcal{L}(X)$ is norm-continuous.
- C_0 (strongly continuous) when $T: [0, \infty) \to \mathcal{L}(X)$ is continuous in the SOT.
- bounded when $T([0,\infty))$ is bounded in $\mathcal{L}(X)$, and locally bounded when T(K) is bounded $\forall K \subset [0,\infty)$ bounded. (so C_0 implies local boundedness by Banach-Steinhaus, and the semigroup property implies we just need to test $K \subset [0,1)$)

 $(T(z))_{z\in\Sigma^+_{s}\cup\{0\}}\subset\mathcal{L}(X)$ is called

- a semigroup when $T: \Sigma_{\delta}^+ \cup \{0\} \to \mathcal{L}(X)$ is a monoid homomorphism.
- C_0 when $\forall \delta' \in (0, \delta), T : \Sigma_{\delta'}^+ \cup \{0\} \to \mathcal{L}(X)$ is continuous in the SOT.
- bounded when $T(\Sigma_{\delta'}^+)$ is bounded $\forall \delta' \in (0, \delta)$ and locally bounded when T(K) is bounded $\forall K \subset \Sigma_{\delta'}^+$ bounded. (so C_0 implies local boundedness, and the semigroup property implies we just need to test $K \subset \mathbb{D} \cap \Sigma_{\delta'}^+$)
- analytic when $T: \Sigma_{\delta}^+ \to \mathcal{L}(X)$ is analytic

We say $(T(t))_{t\geq 0}$ is **analytic** of angle $\delta \in (0, \frac{\pi}{2}]$ if there is an extension $(T(z))_{z\in\Sigma_{\delta}^{+}\cup\{0\}} \subset \mathcal{L}(X)$ which is analytic and <u>locally bounded</u>. If furthermore $(T(z))_{z\in\Sigma_{\delta}^{+}\cup\{0\}}$ is bounded, we say $(T(t))_{t\geq 0}$ is **boundedly analytic** of angle δ .

Remark. A subtle problem is that when $(T(t))_{t\geq 0}$ is bounded and analytic, we cannot conclude $(T(t))_{t\geq 0}$ is boundedly analytic (cf. [Are+11, Definition 3.7.3]).

Blackbox 31. If $(T(t))_{t\geq 0}$ is a C_0 semigroup which is (boundedly) analytic of angle $\delta \in (0, \frac{\pi}{2}]$, then $(T(z))_{z\in \Sigma_{k}^{+}\cup\{0\}}$ is a C_0 , (bounded) semigroup.

Proof. The semigroup property comes from the identity theorem, and C_0 comes from the Vitali holomorphic convergence theorem. We refer to [Are+11, Proposition 3.7.2].

Theorem 32 (Sobolev tower). Let $(e^{tA})_{t\geq 0}$ be a C_0 semigroup on a (real/complex) Banach space X with generator A (implying A is closed and densely defined). Then $\forall m \in \mathbb{N}_1$, $D(A^m)$ is a Banach space under the norm $\|x\|_{D(A^m)} = \|x\|_X + \sum_{k=1}^m \|A^k x\|_X$, and $D(A^m)$ is dense in X.

As e^{tA} and A commute on D(A), we conclude that $(e^{tA})_{t>0}$, after domain restriction, is also a C_0 semigroup on $D(A^m)$ and $\|e^{tA}\|_{\mathcal{L}(D(A^m))} \leq \|e^{tA}\|_{\mathcal{L}(X)} \quad \forall t \geq 0.$ Lastly, if X is a complex Banach space and $(e^{tA})_{t\geq 0}$ is (boundedly) analytic on X, $(e^{tA})_{t\geq 0}$ is also

(boundedly) analytic on $D(A^m)$ after domain restriction.

Proof. Most are just the basics of semigroup theory (cf. [Tay11a, Appendix A.9]). We only prove the last assertion. All we need is commutativity: if $(e^{tA})_{t>0}$ is extended to $(e^{zA})_{z\in\Sigma_{t}^{+}\cup\{0\}}$, we want to show $e^{zA}A = Ae^{zA}$ on D(A).

By Blackbox 31, $(e^{zA})_{z \in \Sigma_{+}^{+} \cup \{0\}}$ is a C_0 semigroup. Therefore $\forall x \in D(A), \forall z \in \Sigma_{\delta}^{+}$:

$$e^{zA}Ax = e^{zA}\left(X - \lim_{t \downarrow 0} \frac{e^{tA} - 1}{t}x\right) = X - \lim_{t \downarrow 0} e^{zA} \frac{e^{tA} - 1}{t}x = X - \lim_{t \downarrow 0} \frac{e^{tA} - 1}{t}e^{zA}x$$

The last term implies $e^{zA}x \in D(A)$ and $e^{zA}Ax = Ae^{zA}x$. Then use Corollary 28 and Corollary 29 to get analyticity.

4.3.2 Simple extrapolation (with core)

Lemma 33. Let U, X be complex Banach spaces and $U \hookrightarrow X$ be a continuous injection with dense image.

- 1. Let $(T(t))_{t\geq 0} \subset \mathcal{L}(X)$ be locally bounded and $T(t)U \leq U \ \forall t \geq 0$. Assume $(T(t))_{t\geq 0}$ is a C_0 semigroup on U. Then $(T(t))_{t>0}$ on X is also a C_0 semigroup.
- 2. Let $(T(z))_{z \in \Sigma_{\delta}^+ \cup \{0\}} \subset \mathcal{L}(X)$ (where $\delta \in (0, \frac{\pi}{2}]$) be locally bounded and $T(z)U \leq U \ \forall z \in \Sigma_{\delta}^+$. Assume $(T(z))_{z \in \Sigma^+_{+} \cup \{0\}}$ is a C_0 , analytic semigroup on U. Then $(T(z))_{z \in \Sigma^+_{+} \cup \{0\}}$ on X is also a C_0 , analytic semigroup.

Remark. The assumption of local boundedness on X is important. We will also use this result in Subsection 7.3 to establish the $W^{1,p}$ -analyticity of the heat flow.

Proof. The semigroup property comes from the density of U in X. To get C_0 on X, use the local boundedness on X and dense convergence (Lemma 9). For analyticity in (2), use Corollary 29.

Lemma 34 (Core). Let A be an unbounded linear operator on a (real/complex) Banach space X and $E \leq$ D(A). E is called a core when E is dense in $(D(A), \|\cdot\|_{D(A)})$.

If A is the generator of a C_0 semigroup on X, E is dense in X and $e^{tA}E \leq E$, then E is a core.

Proof. Let $x \in D(A)$. Then there is $(x_j)_{j \in \mathbb{N}}$ in E such that $x_j \xrightarrow{X} x$. It is trivial to check

$$\frac{1}{t} \int_0^t e^{sA} x_j \, \mathrm{d}s \xrightarrow[j \to \infty]{\|\cdot\|_{D(A)}} \frac{1}{t} \int_0^t e^{sA} x \, \mathrm{d}s \xrightarrow[t \downarrow 0]{\|\cdot\|_{D(A)}} x$$

as $\left(e^{sA}\Big|_{D(A)}\right)_{s>0}$ is $\|\cdot\|_{D(A)}$ -continuous. Note that $\int_0^t e^{sA}x_j \, ds$ is in the $\|\cdot\|_{D(A)}$ -closure of E by the Riemann integral. **Theorem 35** (Simple extrapolation with core). Let (X_0, X_1) be an interpolation couple of complex Banach spaces and $X_{\theta} = [X_0, X_1]_{\theta}$ for $\theta \in (0, 1]$.

Let $(T(t))_{t\geq 0} \subset \mathcal{L}((X_0, X_1))$. Assume that on $X_0, (T(t))_{t\geq 0}$ is bounded.

Assume that on X_1 , $(T(t))_{t\geq 0}$ is a C_0 semigroup, boundedly analytic of angle $\delta \in (0, \frac{\pi}{2}]$ with generator A_1 .

Assume $\exists m \in \mathbb{N}_1 : \left(D(A_1^m), \|\cdot\|_{D(A_1^m)} \right) \hookrightarrow \left(X_0 \cap X_1, \|\cdot\|_{X_0 \cap X_1} \right) \hookrightarrow X_0$ are continuous injections with dense images.

Then on X_{θ} , $(T(t))_{t>0}$ is a C_0 semigroup, and boundedly analytic of angle $\theta\delta$.

Remark. The existence of a convenient core like $D(A_1^m)$ is usually a trivial consequence of Sobolev embedding. We can replace bounded analyticity on X_1 and X_{θ} with analyticity, and boundedness on X_0 with local boundedness via the usual rescaling argument $(\forall \delta' \in (0, \delta) \subset (0, \frac{\pi}{2}), \exists C_{\delta'} > 0 : \|e^{-C_{\delta'}z}T(z)\|_{\mathcal{L}(X_1)} \lesssim_{\delta'} 1 \forall z \in \Sigma_{\delta'}^+).$

The existence of a core allows conditions on X_0 and X_1 to be more general than those in [Fac15, Theorem 3.1.1] (which requires immediate norm-continuity on X_0), and actually be equivalent to those in [Fac15, Theorem 3.1.10] (though Kato-Beurling covers more than just complex interpolation). Once again, the assumption of (local) boundedness on X_0 is important.

We will use this result to establish the L^p -analyticity of the heat flow in Subsection 7.2.

Proof. Let $U = D(A_1^m)$. Then U is Banach as A_1 is closed. Obviously $U \hookrightarrow X_{\theta}$ is a continuous injection with dense image, and $(T(z))_{z \in \Sigma_{\delta}^+ \cup \{0\}}$ is a C_0 , bounded, analytic semigroup on U (via Sobolev tower).

By Lemma 33, $(T(t))_{t\geq 0}$ is a C_0 , bounded semigroup on X_0 . Also by Lemma 33, to get the desired conclusion, we only need to show $(T(z))_{z\in \Sigma_{as}^+\cup\{0\}}$ is locally bounded in $\mathcal{L}(X_{\theta})$.

Fix $\delta' \in (0, \delta)$. We use abstract Stein interpolation. Define the strip $\Omega = \{0 < \text{Re} < 1\}$. Let $\alpha \in (-\delta', \delta'), \rho > 0, u \in U$ and

$$L(z) = T(\rho e^{i\alpha z})u \; \forall z \in \overline{\Omega}$$

Note that $U \leq X_0 \cap X_1$ is dense. We check the other conditions for interpolation:

- As $U \hookrightarrow X_0$ and $U \hookrightarrow X_1$ are continuous, $(\overline{\Omega} \to X_0 + X_1, z \mapsto L(z)u)$ is continuous, bounded on $\overline{\Omega}$ and analytic on Ω (as $L(z)u \in X_1 \hookrightarrow X_0 + X_1$).
- For $j = 0, 1 \ (\mathbb{R} \to X_j, s \mapsto L(j+is)u)$ is
 - continuous since $U \hookrightarrow X_j$ is continuous.
 - bounded by $C_{j,T} \|u\|_{X_j}$ for some $C_{j,T} > 0$ since $(T(t))_{t \ge 0}$ is bounded on X_0 and $(T(te^{i\alpha}))_{t \ge 0}$ is bounded on X_1 .

Then by Stein interpolation, we conclude $\{T(\rho e^{i\theta\alpha}) : \rho > 0, \alpha \in (-\delta', \delta')\} = T(\Sigma_{\theta\delta'}^+) \subset \mathcal{L}(X_{\theta})$ is bounded.

4.3.3 Coreless version

There is an alternative version which we will not use, but is of independent interest:

Theorem 36 (Coreless extrapolation). Let (X_0, X_1) be an interpolation couple of complex Banach spaces and $X_{\theta} = [X_0, X_1]_{\theta}$ for $\theta \in (0, 1]$.

Let $(T(t))_{t>0} \subset \mathcal{L}((X_0, X_1))$ be <u>a semigroup</u>. Assume that on $X_0, (T(t))_{t>0}$ is bounded and <u>degenerate</u>.

Assume that on X_1 , $(T(t))_{t\geq 0}$ is a C_0 semigroup, boundedly analytic of angle $\delta \in (0, \frac{\pi}{2}]$ with generator A_1 .

Then on X_{θ} , $(T(t))_{t>0}$ is a C_0 semigroup, boundedly analytic of angle $\theta\delta$.

Remark. The differences with the previous version are underlined. Again, via rescaling we can replace bounded analyticity on X_1 and X_{θ} with analyticity, and boundedness on X_0 with local boundedness. The conditions on X_0 and X_1 are still a bit more general than those in [Fac15, Theorem 3.1.1], which requires immediate norm-continuity on X_0 . In practice local boundedness on X_0 can usually come from global analysis, while degeneracy can come from Sobolev embedding and dense convergence (Lemma 9). Immediate norm-continuity is harder to establish.

Note that Theorem 36 is not as general as [Fac15, Theorem 3.1.10] (which removes the need for degeneracy and covers more than just complex interpolation), though it is markedly easier to prove.

Proof. By interpolation, $(T(t))_{t\geq 0}$ is a bounded semigroup on X_{θ} . Let $U = X_0 \cap X_1$. Obviously $(T(t))_{t>0}$ is a bounded semigroup on U.

Then observe that $\forall u \in U, \forall t, t_0 \ge 0$:

$$\left\| \left(T(t) - T(t_0) \right) u \right\|_{X_{\theta}} \le \left\| \left(T(t) - T(t_0) \right) u \right\|_{X_0}^{1-\theta} \left\| \left(T(t) - T(t_0) \right) u \right\|_{X_1}^{\theta} \le \left\| \left(T(t) - T(t_0) \right) u \right\|_{X_0}^{\theta} \le \| \left(T(t) - T(t_0) \right) u \|_{X_0}^{\theta} \le \| \left(T(t) - T(t_0) \right) u \|_{X_0}^{\theta} \le \| \left(T(t) - T(t_0) \right) u \|_{X_0}^{\theta} \le \| \left(T(t) - T(t_0) \right) u \|_{X_0}^{\theta} \le \| \left(T(t) - T(t_0) \right) u \|_{X_0}^{\theta} \le \| \left(T(t) - T(t_0) \right) u \|_{X_0}^{\theta} \le \| \left(T(t) - T(t_0) \right) u \|_{X_0}^{\theta} \le \| \left(T(t) - T(t_0) \right) u \|_{X_0}^{\theta} \le \| \left(T(t) - T(t_0) \right) u \|_{X_0}^{\theta} \le \| \left(T(t) - T(t_0) \right) u \|_{X_0}^{\theta} \le \| \left(T(t) - T(t_0) \right) u \|_{X_0}^{\theta} \le \| \left(T(t) - T(t_0) \right) u \|_{X_0}^{\theta} \le \| \left(T(t) - T(t_0) \right) u \|_{X_0}^{\theta} \le \| \left(T(t) - T(t_0) \right) u \|_{X_0}^{\theta} \le \| \left(T(t) - T(t_0) \right) u \|_{X_0}^{\theta} \le \| \left(T(t) - T(t_0) \right) u \|_{X_0}^{\theta} \le \| \left(T(t) - T(t_0) \right) u \|_{X_0}^{\theta} \le \| \left(T(t) - T(t_0) \right) u \|_{X_0}^{\theta} \le \| \left(T(t) - T(t_0) \right) u \|_{X_0}^{\theta} \le \| \left(T(t) - T(t_0) \right) u \|_{X_0}^{\theta} \le \| \left(T(t) - T(t_0) \right) u \|_{X_0}^{\theta} \le \| \left(T(t) - T(t_0) \right) u \|_{X_0}^{\theta} \le \| \left(T(t) - T(t_0) \right) u \|_{X_0}^{\theta} \le \| \left(T(t) - T(t_0) \right) u \|_{X_0}^{\theta} \le \| \left(T(t) - T(t_0) \right) u \|_{X_0}^{\theta} \le \| \left(T(t) - T(t_0) \right) u \|_{X_0}^{\theta} \le \| \left(T(t) - T(t_0) \right) u \|_{X_0}^{\theta} \le \| \left(T(t) - T(t_0) \right) u \|_{X_0}^{\theta} \le \| \left(T(t) - T(t_0) \right) u \|_{X_0}^{\theta} \le \| \left(T(t) - T(t_0) \right) u \|_{X_0}^{\theta} \le \| \left(T(t) - T(t_0) \right) u \|_{X_0}^{\theta} \le \| \left(T(t) - T(t_0) \right) u \|_{X_0}^{\theta} \le \| \left(T(t) - T(t_0) \right) u \|_{X_0}^{\theta} \le \| \left(T(t) - T(t_0) \right) u \|_{X_0}^{\theta} \le \| \left(T(t) - T(t_0) \right) u \|_{X_0}^{\theta} \le \| \left(T(t) - T(t_0) \right) u \|_{X_0}^{\theta} \le \| \left(T(t) - T(t_0) \right) u \|_{X_0}^{\theta} \le \| \left(T(t) - T(t_0) \right) u \|_{X_0}^{\theta} \le \| \left(T(t) - T(t_0) \right) u \|_{X_0}^{\theta} \le \| \left(T(t) - T(t_0) \right) u \|_{X_0}^{\theta} \le \| \left(T(t) - T(t_0) \right) u \|_{X_0}^{\theta} \le \| \left(T(t) - T(t_0) \right) u \|_{X_0}^{\theta} \le \| \left(T(t) - T(t_0) \right) u \|_{X_0}^{\theta} \le \| \left(T(t) - T(t_0) \right) u \|_{X_0}^{\theta} \le \| \left(T(t) - T(t_0) \right) u \|_{X_0}^{\theta} \le \| \left(T(t) - T(t_0) \right) u \|_{X_0}^{\theta} \le \| \left(T(t) - T(t_0) \right) u \|_{X_0}^{\theta} \le \| \left(T(t) - T(t_0) \right) u \|_{X_0}^{\theta} \le \| \left(T(t) - T(t_0) \right) u \|_{X_0}^{\theta} \le \| \left(T(t) - T(t_0) \right) u$$

Since $\theta \neq 0$, we have $T(t)u \xrightarrow[t \to t_0]{X_{\theta}} T(t_0)u$. As $(T(t))_{t \geq 0}$ is bounded on X_{θ} and U is dense in X_{θ} , we conclude $(T(t))_{t>0}$ is C_0 on X_{θ} by dense convergence (Lemma 9).

Fix $\delta' \in (0, \delta)$. We use abstract Stein interpolation. Define the strip $\Omega = \{0 < \text{Re} < 1\}$. Let $\alpha \in (-\delta', \delta'), \rho > 0, u \in U$ and

$$L(z) = T(\rho e^{i\alpha z})u \; \forall z \in \overline{\Omega}$$

Note that $U = X_0 \cap X_1$. We check the other conditions for interpolation:

- As $U \hookrightarrow X_0$ and $U \hookrightarrow X_1$ are continuous, $(\overline{\Omega} \to X_0 + X_1, z \mapsto L(z)u)$ is continuous, bounded on $\overline{\Omega}$ and analytic on Ω (as $L(z)u \in X_1 \hookrightarrow X_0 + X_1$).
- For $j = 0, 1 \ (\mathbb{R} \to X_j, s \mapsto L(j + is)u)$ is
 - continuous since $(T(t))_{t>0}$ is degenerate on X_0 and $(T(te^{i\alpha}))_{t>0}$ is C_0 on X_1 .
 - bounded by $C_{j,T} \|u\|_{X_j}$ for some $C_{j,T} > 0$ since $(T(t))_{t \ge 0}$ is bounded on X_0 and $(T(te^{i\alpha}))_{t \ge 0}$ is bounded on X_1 .
- By Stein interpolation, $\{T(\rho e^{i\theta\alpha}) : \rho > 0, \alpha \in (-\delta', \delta')\} = T(\Sigma_{\theta\delta'}^+) \subset \mathcal{L}(X_{\theta})$ is bounded. Finally, we just need to show $(T(z))_{z \in \Sigma_{\sigma s}^+ \cup \{0\}}$ is analytic on X_{θ} . Let $u \in U$. Then

$$\left(\Sigma_{\delta}^{+} \to X_{1} \hookrightarrow X_{0} + X_{1}, z \mapsto T(z)u\right)$$

is analytic. Therefore $(\Sigma_{\theta\delta}^+ \to X_\theta \hookrightarrow X_0 + X_1, z \mapsto T(z)u)$ is analytic, while $(\Sigma_{\theta\delta}^+ \to X_\theta, z \mapsto T(z)u)$ is locally bounded, so we can use Corollary 28 to conclude $(\Sigma_{\theta\delta}^+ \to X_\theta, z \mapsto T(z)u)$ is analytic. As U is dense in X_θ , by corollary 29, we conclude $(\Sigma_{\theta\delta}^+ \to \mathcal{L}(X_\theta), z \mapsto T(z))$ is analytic. \Box

4.4 Sectorial operators

Recall that if $(T(t))_{t\geq 0}$ is a C_0 semigroup on a complex Banach space X, then it has a closed, densely defined generator A, and $T(t) = e^{tA}$ is exponentially bounded: $||e^{tA}|| \lesssim_{\neg t} e^{Ct}$ for some C > 0. Then

 $\forall \zeta \in \{ \operatorname{Re} > C \} : \zeta \in \rho(A) \text{ and }$

$$\frac{1}{\zeta - A}x = \int_0^\infty e^{-\zeta t} e^{tA}x \, \mathrm{d}t \, \forall x \in X$$

(cf. [Tay11a, Appendix A, Proposition 9.2])

This means that the resolvent $\frac{1}{\zeta - A}$ is the Laplace transform of the semigroup e^{tA} . This naturally leads to the question when we can perform the inverse Laplace transform, to recover the semigroup from the resolvent. This motivates the definition of sectorial operators, which includes the Laplacian.

Unfortunately, there are wildly different definitions currently in use by authors. The reader should study the definitions closely whenever they consult any literature on sectorial operators (e.g. [Lun95; Haa06; Are+11; Eng00]).

Definition 37. Let A be an unbounded operator on a complex Banach space X. For $\theta \in [0, \pi)$, we say A is

- sectorial of angle θ $(A \in \text{Sect}(\theta))$ when $\begin{cases} \sigma(A) \subset \overline{\Sigma_{\theta}^{-}} \\ \forall \omega \in [0, \pi \theta) : M(A, \omega) := \sup_{\lambda \in \Sigma_{\omega}^{+}} \left\| \frac{\lambda}{\lambda A} \right\| < \infty \end{cases}$
- quasi-sectorial when $\exists a \in \mathbb{R} : A a$ is sectorial.
- acutely sectorial when $A \in \text{Sect}(\theta)$ for some $\theta \in [0, \frac{\pi}{2})$
- acutely quasi-sectorial when $\exists a \in \mathbb{R} : A a$ is acutely sectorial.

For $r > 0, \eta \in (\frac{\pi}{2}, \pi)$, we define the (counterclockwise-oriented) Mellin curve

$$\gamma_{r,\eta} = e^{i\eta}[r,\infty) \cup e^{-i\eta}[r,\infty) \cup re^{i[-\eta,\eta]}$$

Remark. Depending on the author, "sectorial" can mean any of those four, and that is not taking sign conventions into account (some authors want $-\Delta$ to be sectorial), as well as whether A should be densely defined. The term "quasi-sectorial" is taken from [Haa06].

In particular, letting the spectrum be in the left half-plane means we agree with [Eng00; Lun95] and disagree with [Are+11; McI86; Haa06]. This is simply a personal preference, of being able to say "the Laplacian is sectorial", or "generators of C_0 analytic semigroups are acutely sectorial". Also, for bounded holomorphic calculus, $e^{t\Delta}$ morally comes from $(e^{tz})_{z\in\sigma(\Delta)}$ which is bounded in the left half-plane.

In keeping with tradition, here is the usual visualization:



Figure 1: Acutely sectorial operators

Blackbox 38. A generates a C_0 , boundedly analytic semigroup on complex Banach space X if and only if A is densely defined and acutely sectorial.

When that happens, $\exists \delta \in (0, \frac{\pi}{2})$ and $\eta \in (\frac{\pi}{2}, \pi)$ such that $(e^{tA})_{t>0}$ extends to $(e^{\zeta A})_{\zeta \in \Sigma^+_{t} \cup \{0\}}$ and

$$e^{\zeta A} = \frac{1}{2\pi i} \int_{\gamma_{r,\eta}} e^{\zeta z} \frac{1}{z - A} \, dz \quad \forall \zeta \in \Sigma_{\delta}^+, \forall r > 0$$

Also $\forall t > 0, \forall k \in \mathbb{N}_1 : e^{tA}(X) \le D(A^\infty), \left\| A^k e^{tA} \right\| \lesssim_{\neg t, \neg k} \frac{k^k}{t^k} \text{ and } \partial_t^k(e^{tA}x) = A^k e^{tA}x \ \forall x \in X.$

Remark. This is the aforementioned inverse Laplace transform. The Mellin curve and the resolvent estimate in the definition of sectoriality ensure sufficient decay for the integral to make sense. As it is a complex line integral and the resolvent is analytic, the semigroup becomes analytic.

A trivial consequence is that $D(A^{\infty})$ is dense in X and therefore a core.

When A is densely defined and acutely quasi-sectorial, a simple rescaling $e^{t(A-a)} = e^{-ta}e^{tA}$ implies $(e^{tA})_{t>0}$ is a C_0 , analytic semigroup.

Proof. See [Eng00, Section II.4.a]. The curious figure
$$\frac{k^k}{t^k}$$
 comes from $A^k e^{tA} = \left(Ae^{\frac{t}{k}A}\right)^k$.

Theorem 39 (Yosida's half-plane criterion). A is acutely quasi-sectorial if and only if $\exists C > 0$ such that

- {Re > C} $\subset \rho(A)$
- $\bullet \ \sup_{\lambda \in \{\operatorname{Re} > C\}} \left\| \tfrac{\lambda}{\lambda A} \right\| < \infty$

Remark. This is how the L^p -analyticity of the heat flow is traditionally established. Yet proving the resolvent estimate is nontrivial, as it is quite a refinement of elliptic estimates, so we choose not to do so. Interestingly, we will instead use this for the $B_{3,1}^{\frac{1}{3}}$ -analyticity of the heat flow in Subsection 8.3, though that case is especially easy since we already have analyticity at the two endpoints L^3 and $W^{1,3}$.

Proof. We only need to prove \Leftarrow . Recall the proof of how $\rho(A)$ is open: $\forall \lambda \in \rho(A), B\left(\lambda, \left\|\frac{1}{\lambda-A}\right\|^{-1}\right) \subset \rho(A)$. Applying this allows us to open up {Re > C} and get $C + \Sigma_{\eta}^+ \subset \rho(A)$ for some $\eta \in (\frac{\pi}{2}, \pi)$. By choosing η near $\frac{\pi}{2}$, the resolvent estimate is retained.

Definition 40. Let A be an unbounded operator on a Hilbert space X. Then A is called

- symmetric when $\langle Ax, y \rangle = \langle x, Ay \rangle \quad \forall x, y \in D(A)$, or equivalently, $A \subset A^*$ (where A and A^* are identified with their graphs).
- self-adjoint when $A = A^*$. This implies $\sigma(A) \subset \mathbb{R}$ (cf. [Tay11a, Appendix A, Proposition 8.5]).
- dissipative when $\operatorname{Re} \langle Ax, x \rangle \leq 0 \ \forall x \in D(A)$.

When A is dissipative, $\forall \lambda \in \{\operatorname{Re} > 0\}, \forall x \in D(A) : \operatorname{Re} \langle (\lambda - A) x, x \rangle \ge \operatorname{Re} \langle \lambda x, x \rangle$ so $\|(\lambda - A) x\| \ge \operatorname{Re} \lambda \|x\|$. Recall how $\rho(A)$ is proved to be open: $\forall \lambda \in \rho(A), B\left(\lambda, \left\|\frac{1}{\lambda - A}\right\|^{-1}\right) \subset \rho(A)$. Consequently, if A is dissipative and $\exists \lambda_0 \in \{\operatorname{Re} > 0\} \cap \rho(A)$, we can conclude $\{\operatorname{Re} > 0\} \subset \rho(A)$.

Theorem 41 (Dissipative sectoriality). Assume X is a complex Hilbert space and A is an unbounded, self-adjoint, dissipative operator on X. Then A is acutely sectorial of angle 0.

Remark. Though standard, this might be the most elegant theorem in the theory, and later on will instantly imply the L^2 -analyticity of the heat flow in Subsection 7.1. The theorem can also be proved by Euclidean geometry. When X is separable, we can also use the spectral theorem for unbounded operators.

Proof. As A is self-adjoint, $\mathbb{C}\setminus\mathbb{R} \subset \rho(A)$. By dissipativity, we conclude $\sigma(A) \subset (-\infty, 0]$. Also by self-adjointness, Re $\langle Ax, x \rangle = \langle Ax, x \rangle \leq 0 \ \forall x \in D(A)$.

Arbitrarily pick $\theta \in (\frac{\pi}{2}, \pi)$. We want to show $\left\| \frac{z}{z-A} \right\| \lesssim_{\theta} 1 \ \forall z \in \Sigma_{\theta}^+$.

Let $x \in X$ and $u = \frac{1}{z-A}x$. As $|\langle u, x \rangle| \le ||u||_X ||x||_X$, we want to show $||u||_X^2 \lesssim_{\theta} \left|\frac{1}{z} \langle u, x \rangle\right|$. Note that

$$\frac{1}{z} \langle u, x \rangle = \frac{1}{z} \langle u, (z - A)u \rangle = \langle u, u \rangle - \frac{1}{z} \langle Au, u \rangle$$

WLOG assume $||u||_X = 1$. Then we want $1 \leq_{\theta} |1 - \frac{1}{z} \langle Au, u \rangle|$. Note that $-\langle Au, u \rangle \geq 0$ and $-\frac{1}{z} \langle Au, u \rangle \in \Sigma_{\theta}^+$. Then we are done since

$$\left|1 - \frac{1}{z} \langle Au, u \rangle \right| \ge \operatorname{dist}(0, 1 + \Sigma_{\theta}^{+}) > 0.$$

By Euclidean geometry, we can even calculate dist $(0, 1 + \Sigma_{\theta}^{+})$. We will not need it though. \Box

5 Scalar function spaces

Throughout this section, we work with complex-valued functions.

5.1 On \mathbb{R}^n

Definition 42. Here we recall the various (inhomogeneous) function spaces which are particularly suitable for interpolation. They are defined as subspaces of $\mathcal{S}'(\mathbb{R}^n)$ with certain norms being finite:

- 1. Lebesgue-Sobolev spaces: for $m \in \mathbb{N}_0, p \in [1, \infty]$: $\|f\|_{W^{m,p}(\mathbb{R}^n)} \sim \sum_{k=0}^m \|\nabla^k f\|_p$ where $\nabla^k f \in L^p$ are tensors defined by distributions. It is customary to write H^m for $W^{m,2}$.
- 2. Bessel potential spaces: for $s \in \mathbb{R}, p \in [1, \infty]$: $||f||_{H^{s,p}(\mathbb{R}^n)} \sim ||\langle \nabla \rangle^s f||_p$ where $\langle \nabla \rangle^s = (1 \Delta)^{\frac{s}{2}}$ is the Bessel potential.
- 3. Besov spaces: for $s \in \mathbb{R}, p \in [1, \infty], q \in [1, \infty]$: $||f||_{B^s_{p,q}(\mathbb{R}^n)} \sim ||P_{\leq 1}f||_p + ||N^s||P_Nf||_p ||_{l^q_{N>1}}$ where P_N and $P_{\leq N}$ (for $N \in 2^{\mathbb{Z}}$) are the standard Littlewood-Paley projections (cf. [Tao06, Appendix A]).
- 4. Triebel-Lizorkin spaces: for $s \in \mathbb{R}, p \in [1, \infty), q \in [1, \infty]$: $\|f\|_{F^s_{p,q}(\mathbb{R}^n)} \sim \|P_{\leq 1}f\|_p + \|N^s\|P_Nf\|_{l^q_{N \leq 1}}\|_{L^q}$

Remark. As there are multiple characterizations for the same spaces, we only define up to equivalent norms. Of course, the topologies induced by equivalent norms are the same.

In the literature, "Fractional Sobolev spaces" like $W^{s,p}$ could either refer to $B_{p,p}^{s}$ (Sobolev–Slobodeckij spaces) or $H^{s,p}$. We shall avoid using the term at all. There are also some delicate issues with $F_{\infty,q}^{s}$ which we do not need to discuss here (cf. [Tri10, Section 2.3.4]).

Blackbox 43. Recall from harmonic analysis (cf. [Tri10, Section 2.5.6, 2.3.3, 2.11.2] and [Lem02, Part 1, Chapter 3.1]):

- $W^{m,p}(\mathbb{R}^n) = H^{m,p}(\mathbb{R}^n)$ for $m \in \mathbb{N}_0, p \in (1,\infty)$.
- $F_{p,2}^s(\mathbb{R}^n) = H^{s,p}(\mathbb{R}^n)$ for $s \in \mathbb{R}, p \in (1,\infty)$.
- $B_{p,1}^m(\mathbb{R}^n) \hookrightarrow W^{m,p}(\mathbb{R}^n) \hookrightarrow B_{p,\infty}^m(\mathbb{R}^n)$ for $m \in \mathbb{N}_0, p \in [1,\infty]$.
- $\mathcal{S}(\mathbb{R}^n)$ is dense in $W^{m,p}(\mathbb{R}^n)$, $B^s_{p,q}(\mathbb{R}^n)$ and $F^s_{p,q}(\mathbb{R}^n)$ for $m \in \mathbb{N}_0$, $s \in \mathbb{R}$, $p \in [1,\infty)$, $q \in [1,\infty)$.
- $B^s_{p,\min(p,q)}(\mathbb{R}^n) \hookrightarrow F^s_{p,q}(\mathbb{R}^n) \hookrightarrow B^s_{p,\max(p,q)}(\mathbb{R}^n) \text{ for } s \in \mathbb{R}, p \in [1,\infty), q \in [1,\infty].$
- $\left(B_{p,q}^s\left(\mathbb{R}^n\right)\right)^* = B_{p',q'}^{-s}\left(\mathbb{R}^n\right)$ for $s \in \mathbb{R}, p \in [1,\infty), q \in [1,\infty)$. $\left(F_{p,q}^s\left(\mathbb{R}^n\right)\right)^* = F_{p',q'}^{-s}\left(\mathbb{R}^n\right)$ for $s \in \mathbb{R}, p \in (1,\infty), q \in (1,\infty)$.

5.2 On domains

Definition 44. A C^{∞} domain Ω in \mathbb{R}^n is defined as an open subset of \mathbb{R}^n with smooth boundary, and scalar function spaces are then defined on Ω . If $\Omega \subset S \subset \overline{\Omega}$, let function spaces on S implicitly refer to function spaces on Ω . This will make it possible to discuss function spaces on, for example, $\overline{\mathbb{R}^n_+} \cap B_{\mathbb{R}^n}(0,1)$, or compact Riemannian manifolds with boundary.

Obviously, Sobolev spaces are still defined on domains by distributions. The big question is finding a good characterization for $B_{p,q}^s$ and $F_{p,q}^s$ on domains, when the Fourier transform is no longer available. This is among the main topics of Triebel's seminal books. Let us review the results:

Definition 45. Let Ω be either \mathbb{R}^n , or the half-space \mathbb{R}^n_+ , or a bounded C^{∞} domain in \mathbb{R}^n .

Then $B_{p,q}^s(\Omega)$ and $F_{p,q}^s(\Omega)$ can simply be defined as the restrictions of $B_{p,q}^s(\mathbb{R}^n)$ and $F_{p,q}^s(\mathbb{R}^n)$ to Ω and

$$\|f\|_{B^{s}_{p,q}(\Omega)} = \inf\{\|F\|_{B^{s}_{p,q}(\mathbb{R}^{n})} : F \in B^{s}_{p,q}(\mathbb{R}^{n}), F|_{\Omega} = f\} \text{ for } s \in \mathbb{R}; p, q \in [1,\infty]$$

$$\|f\|_{F^{s}_{p,q}(\Omega)} = \inf\{\|F\|_{F^{s}_{p,q}(\mathbb{R}^{n})} : F \in F^{s}_{p,q}(\mathbb{R}^{n}), \, F|_{\Omega} = f\} \ \text{ for } s \in \mathbb{R}, p \in [1,\infty), q \in [1,\infty]$$

A more useful characterization is via BMD (**ball mean difference**). Let $\tau_h f(x) = f(x+h)$ be the translation operator and $\Delta_h f = \tau_h f - f$ be the difference operator. Then for $m \in \mathbb{N}_1$, we can define $\Delta_h^m = (\Delta_h)^m$ as the *m*-th difference operator. As we need to stay on the domain Ω , define

$$V^m(x,t) = \frac{1}{m} \left(B(x,mt) \cap \Omega - x \right) \text{ for } x \in \Omega, t > 0, m \in \mathbb{N}_1$$

So $V^m(x,t) \subset B(0,t)$, $x + mV^m(x,t) \subset \Omega$ and $\Delta_h^l f(x)$ is well-defined when $h \in V^m(x,t)$. Also note for $t \in (0,1)$: $|V^m(x,t)| \sim_{\Omega,m} t^n$. Then by [Tri92, Section 3.5.3, 5.2.2]:

1. For $m \in \mathbb{N}_1, p \in [1, \infty], q \in [1, \infty], s \in (0, m), r \in [1, p]$:

$$\|f\|_{B^{s}_{p,q}(\Omega)} \sim \|f\|_{p} + \left\|t^{-s} \left\| \|\Delta^{m}_{h}f(x)\|_{L^{r}_{h}\left(\frac{1}{t^{n}}\mathrm{d}h, V^{m}(x,t)\right)} \right\|_{L^{p}_{x}(\Omega)} \right\|_{L^{q}\left(\frac{1}{t}\mathrm{d}t, (0,1)\right)}$$
(7)

We carefully note here that m > s (the difference operator must be strictly higher-order than the regularity), and that the variable t is small, which will play a big role in Theorem 54. We also note that this is different from the classical characterization via differences ([Tri10, Section 3.4.2], [Tri92, Section 1.10.3]) which analysts might be more familiar with:

$$\|f\|_{B^{s}_{p,q}(\Omega)} \sim \|f\|_{p} + \left\||h|^{-s} \left\|\Delta^{m}_{h,\Omega}f(x)\right\|_{L^{p}_{x}(\Omega)}\right\|_{L^{q}\left(\frac{dh}{|h|^{n}}, B(0,1)\right)}$$

where $\Delta_{h,\Omega}^m f(x)$ is the same as $\Delta_h^m f(x)$, but zero wherever undefined, and $m \in \mathbb{N}_1, p \in (1,\infty), q \in [1,\infty], s \in (0,m)$.

2. For $m \in \mathbb{N}_1, p \in [1, \infty), q \in [1, \infty], s \in (0, m), r \in [1, p]$:

$$\|f\|_{F^{s}_{p,q}(\Omega)} \sim \|f\|_{p} + \left\|t^{-s} \left\| \|\Delta_{h}^{m} f(x)\|_{L^{r}_{h}\left(\frac{\mathrm{d}h}{t^{h}}, V^{m}(x,t)\right)} \right\|_{L^{q}\left(\frac{\mathrm{d}t}{t}, (0,1)\right)} \right\|_{L^{p}_{x}(\Omega)}$$

Blackbox 46 (Diffeomorphisms and smooth multipliers). Every diffeomorphism on \mathbb{R}^n preserves (under pullback) the topology of

- $W^{k,p}(\mathbb{R}^n)$ for $k \in \mathbb{N}_0, p \in [1,\infty]$
- $B^s_{p,q}(\mathbb{R}^n)$ for $s \in \mathbb{R}, p \in [1,\infty], q \in [1,\infty]$
- $F_{p,q}^s(\mathbb{R}^n)$ for $s \in \mathbb{R}, p \in [1,\infty), q \in [1,\infty]$

Also on the same spaces, for $\phi \in C^\infty_c(\mathbb{R}^n)$, $f \mapsto \phi f$ is a bounded linear map .

Remark. This allows us to trivially define function spaces on compact Riemannian manifolds with boundary via partitions of unity and give them unique topologies.

Proof. For $W^{k,p}$ it is trivial. For $B^s_{p,q}$ and $F^s_{p,q}$, see [Tri92, Section 4.3, 4.2.2] and [Tri10, Section 2.8.2].

Blackbox 47 (Extension and trace). Let Ω be either the half-space \mathbb{R}^n_{\perp} or a bounded C^{∞} domain in \mathbb{R}^n .

1. Stein extension: There exists a common (continuous linear) extension operator \mathfrak{E} : $W^{k,p}(\Omega) \hookrightarrow W^{k,p}(\mathbb{R}^n)$ for all $k \in \mathbb{N}_0, p \in [1,\infty]$

- 2. **Triebel extension**: For any $N \in \mathbb{N}_1$, there exists a common (continuous linear) extension operator \mathfrak{E}^N such that
 - $(a) \ \mathfrak{E}^N: B^s_{p,q}(\Omega) \hookrightarrow B^s_{p,q}(\mathbb{R}^n) \ for \ all \ |s| < N, p \in [1,\infty], q \in [1,\infty]$
 - (b) $\mathfrak{E}^N : F^s_{p,q}(\Omega) \hookrightarrow F^s_{p,q}(\mathbb{R}^n)$ for all $|s| < N, p \in [1,\infty), q \in [1,\infty]$
- 3. Trace theorems: Let $n \geq 2$.
 - (a) For $p \in [1,\infty], q \in [1,\infty], s > \frac{1}{p} : B_{p,q}^s(\Omega) \twoheadrightarrow B_{p,q}^{s-\frac{1}{p}}(\partial\Omega)$ is a retraction (continuous surjection with a bounded linear section as a right inverse).
 - (b) For $p \in [1,\infty), q \in [1,\infty], s > \frac{1}{p} : F_{p,q}^s(\Omega) \twoheadrightarrow B_{p,p}^{s-\frac{1}{p}}(\partial\Omega)$ is a retraction.
 - (c) (Limiting case) For $p \in [1,\infty)$, $B_{p,1}^{\frac{1}{p}}(\Omega) \rightarrow L^{p}(\partial\Omega)$ and $W^{1,1}(\Omega) \rightarrow L^{1}(\partial\Omega)$ are continuous surjections.

Remark. It is important to note that we do not have the trace theorem for, say, $B_{3,2}^{\frac{1}{3}}(\Omega)$ (cf. [Sch11, Section 3])

Proof.

- 1. See [Ste71, Section VI.3].
- 2. See [Tri92, Section 4.5, 5.1.3].
- 3. See [Tri10, Section 2.7.2, 3.3.3] and the remarks.

- **Corollary 48.** Let Ω be either the half-space \mathbb{R}^n_+ or a bounded C^{∞} domain in \mathbb{R}^n .
 - $F_{p,2}^m(\Omega) = W^{m,p}(\Omega)$ for $m \in \mathbb{N}_0, p \in (1,\infty)$.
 - $B_{p,1}^m(\Omega) \hookrightarrow W^{m,p}(\Omega) \hookrightarrow B_{p,\infty}^m(\Omega)$ for $m \in \mathbb{N}_0, p \in [1,\infty]$.
 - $\mathcal{S}(\overline{\Omega})$ is dense in $W^{m,p}(\Omega)$, $F^s_{p,q}(\Omega)$ and $B^s_{p,q}(\Omega)$ for $m \in \mathbb{N}_0$, $s \in \mathbb{R}$, $p \in [1,\infty)$, $q \in [1,\infty)$.
 - $B^s_{p,\min(p,q)}(\Omega) \hookrightarrow F^s_{p,q}(\Omega) \hookrightarrow B^s_{p,\max(p,q)}(\Omega) \text{ for } s \in \mathbb{R}, p \in [1,\infty), q \in [1,\infty]$

Remark. When Ω is a bounded C^{∞} domain, $\mathcal{S}(\overline{\Omega}) = C^{\infty}(\overline{\Omega})$.

Proof. Use Triebel and Stein extensions.

5.3 Holder & Zygmund spaces

Definition 49. Let Ω be either \mathbb{R}^n , the half-space \mathbb{R}^n_+ or a bounded C^{∞} domain in \mathbb{R}^n . Recall some L^{∞} type spaces:

• Holder spaces: for $k \in \mathbb{N}_0, \alpha \in (0, 1]$,

$$\|f\|_{C^{k,\alpha}(\Omega)} = \|f\|_{C^{k}(\Omega)} + \max_{|\beta|=k} \left[D^{\beta}f\right]_{C^{0,\alpha}(\Omega)}$$

where $[g]_{C^{0,\alpha}(\Omega)} = \sup_{x \neq y} \frac{|g(x) - g(y)|}{|x - y|^{\alpha}}$

• **Zygmund spaces**: for s > 0, define $\mathfrak{C}^s(\Omega) = B^s_{\infty,\infty}(\Omega)$. Then for $m \in \mathbb{N}_1, s \in (0, m)$:

$$\|f\|_{\mathfrak{C}^{s}(\Omega)} \sim \|f\|_{L^{\infty}(\Omega)} + \left\|t^{-s} \left\|\|\Delta_{h}^{m}f(x)\|_{L^{\infty}_{h}(V^{m}(x,t))}\right\|_{L^{\infty}_{x}(\Omega)}\right\|_{L^{\infty}_{t}((0,1))} \sim \sup|f| + \sup_{0 < |h| \le 1, x \in \Omega} \frac{\left|\Delta_{h,\Omega}^{m}f(x)\right|}{|h|^{s}}$$

It is well-known (cf. [Tri10, Section 2.2.2, 2.5.7, 2.5.12, 2.8.3]) that

- $\|f\|_{\mathfrak{C}^{k+\alpha}(\Omega)} \sim \|f\|_{C^k} + \max_{|\beta|=k} \|D^{\beta}f\|_{\mathfrak{C}^{\alpha}(\Omega)}$ for $k \in \mathbb{N}_0, \alpha \in (0,1]$
- $||f||_{\mathfrak{C}^{k+\alpha}(\Omega)} \sim ||f||_{C^{k,\alpha}}$ for $k \in \mathbb{N}_0, \alpha \in (0,1)$.
- $||fg||_{\mathfrak{C}^s(\Omega)} \lesssim ||f||_{\mathfrak{C}^s} ||g||_{\mathfrak{C}^s}$ for s > 0.

Note that $C^{0,1}, C^1$ and \mathfrak{C}^1 are different.

5.4 Interpolation & embedding

Blackbox 50 (Interpolation). Let Ω be either \mathbb{R}^n , the half-space \mathbb{R}^n_+ or a bounded C^{∞} domain in \mathbb{R}^n . Throughout the theorem, always assume $\theta \in (0,1), s_{\theta} = (1-\theta)s_0 + \theta s_1$.

1. $(B_{p,q_0}^{s_0}(\Omega), B_{p,q_1}^{s_1}(\Omega))_{\theta,q} = B_{p,q}^{s_{\theta}}(\Omega) \text{ for } s_0 \neq s_1, s_j \in \mathbb{R}, p \in [1,\infty], q_j, q \in [1,\infty].$

$$\left(F_{p,q_0}^{s_0}(\Omega), F_{p,q_1}^{s_1}(\Omega)\right)_{\theta,q} = B_{p,q}^{s_\theta}(\Omega) \text{ for } s_0 \neq s_1, s_j \in \mathbb{R}, p \in [1,\infty), q_j, q \in [1,\infty]$$

- 2. $\left(B_{p_0,q_0}^{s_0}(\Omega), B_{p_1,q_1}^{s_1}(\Omega)\right)_{\theta,p_{\theta}} = B_{p_{\theta},p_{\theta}}^{s_{\theta}}(\Omega) \text{ for } s_0 \neq s_1, s_j \in \mathbb{R}, p_j \in [1,\infty], q_j \in [1,\infty], \frac{1}{p_{\theta}} = \frac{1-\theta}{p_0} + \frac{\theta}{p_1} = \frac{1-\theta}{q_0} + \frac{\theta}{q_1}$
- $3. \ \left[B_{p_0,q_0}^{s_0}(\Omega), B_{p_1,q_1}^{s_1}(\Omega)\right]_{\theta} = B_{p_{\theta},q_{\theta}}^{s_{\theta}}(\Omega) \ and \ \left[F_{p_0,q_0}^{s_0}(\Omega), F_{p_1,q_1}^{s_1}(\Omega)\right]_{\theta} = F_{p_{\theta},q_{\theta}}^{s_{\theta}}(\Omega)$

for
$$s_j \in \mathbb{R}, p_j \in (1, \infty), q_j \in (1, \infty), \frac{1}{p_\theta} = \frac{1-\theta}{p_0} + \frac{\theta}{p_1}, \frac{1}{q_\theta} = \frac{1-\theta}{q_0} + \frac{\theta}{q_1}$$

4.
$$[L^{p_0}(\Omega), L^{p_1}(\Omega)]_{\theta} = L^{p_{\theta}}(\Omega) \text{ for } p_j \in [1, \infty], \frac{1}{p_{\theta}} = \frac{1-\theta}{p_0} + \frac{\theta}{p_1}.$$

5. $(W^{m_0,p}(\Omega), W^{m_1,p}(\Omega))_{\theta,q} = B^{m_\theta}_{p,q}(\Omega)$ for $m_j \in \mathbb{N}_0, m_0 \neq m_1, p \in [1,\infty], q \in [1,\infty], m_\theta = (1-\theta)m_0 + \theta m_1.$

Proof.

- 1. Extension operators and [Tri10, Section 2.4.2].
- 2. Extension operators and [BL76, Theorem 6.4.5].
- 3. Extension operators and [Tri10, Section 2.4.7].
- 4. Extension by zero and [BL76, Section 5.1.1]

5. Recall
$$B_{p,1}^m(\Omega) \hookrightarrow W^{m,p}(\Omega) \hookrightarrow B_{p,\infty}^m(\Omega)$$
 for $m \in \mathbb{N}_0, p \in [1,\infty]$. Then apply 1.

Blackbox 51 (Embedding). Let Ω be a bounded C^{∞} domain in \mathbb{R}^n . Assume $\infty > s_0 > s_1 > -\infty$. Then

- 1. $B_{p_0,q_0}^{s_0}(\Omega) \hookrightarrow B_{p_1,q_1}^{s_1}(\Omega)$ is compact for $p_j \in [1,\infty], q_j \in [1,\infty], \frac{1}{p_1} > \frac{1}{p_0} \frac{s_0 s_1}{n}$ $F_{p_0,q_0}^{s_0}(\Omega) \hookrightarrow F_{p_1,q_1}^{s_1}(\Omega)$ is compact for $p_j \in [1,\infty), q_j \in [1,\infty], \frac{1}{p_1} > \frac{1}{p_0} - \frac{s_0-s_1}{n}$
- 2. $B_{p_0,q}^{s_0}(\Omega) \hookrightarrow B_{p_1,q}^{s_1}(\Omega)$ is continuous for $p_j \in [1,\infty], q \in [1,\infty], \frac{1}{p_1} = \frac{1}{p_0} \frac{s_0 s_1}{n}$

$$F_{p_0,q_0}^{s_0}(\Omega) \hookrightarrow F_{p_1,q_1}^{s_1}(\Omega)$$
 is continuous for $p_j \in [1,\infty), q_j \in [1,\infty], \frac{1}{p_1} = \frac{1}{p_0} - \frac{s_0 - s_1}{n}$

Proof.

- See [Tri10, Section 4.3.2, Remark 1] and [Tri10, Section 3.3.1].
 See [Tri10, Section 3.3.1].

Corollary 52. Let Ω be a bounded C^{∞} domain in \mathbb{R}^n . Then

- 1. For $m_j \in \mathbb{N}_0, m_0 > m_1, p_j \in [1, \infty], \frac{1}{p_1} > \frac{1}{p_0} \frac{m_0 m_1}{n}$: $W^{m_0, p_0}(\Omega) \hookrightarrow B^{m_0}_{p_0, \infty}(\Omega) \hookrightarrow B^{m_1}_{p_1, 1}(\Omega) \hookrightarrow W^{m_1, p_1}(\Omega) \text{ is compact.}$
- 2. For $m_j \in \mathbb{N}_0, m_0 > m_1, p_0 \in [1, \infty], \alpha \in (0, 1), 0 > \frac{1}{p_0} \frac{m_0 (m_1 + \alpha)}{n}$: $W^{m_0, p_0}(\Omega) \hookrightarrow B^{m_0}_{p_0, \infty}(\Omega) \hookrightarrow B^{m_1 + \alpha}_{\infty, \infty}(\Omega) = C^{m_1, \alpha}(\overline{\Omega}) \text{ is compact.}$
- 3. For $m \in \mathbb{N}_1, p \in (1,\infty) : W^{m,p}(\Omega) \hookrightarrow B^m_{p,\infty}(\Omega) \hookrightarrow B^{\frac{1}{p}}_{p,1}(\Omega) \twoheadrightarrow L^p(\partial\Omega)$ is compact.

Remark. These include the Rellich-Kondrachov embeddings found in [Ada03, Theorem 6.3], so the Besov embeddings generalize Sobolev embeddings.

5.5Strip decay

Some notation first: let Ω be a C^{∞} domain in \mathbb{R}^n or a compact Riemannian manifold with or without boundary. Define $\Omega_{>r} = \{x \in \Omega : \operatorname{dist}(x, \partial \Omega) > r\}$ where $\operatorname{dist}(x, \partial \Omega) = \infty$ if $\partial \Omega = \emptyset$. Similarly define $\Omega_{\geq r}, \Omega_{< r}, \Omega_{[r_1, r_2]}.$

When $|\Omega| < \infty$ and $p \in [1, \infty)$, we write

$$\|f\|_{L^{p}(\Omega, \operatorname{avg})} = \|f(x)\|_{L^{p}_{x}(\frac{\mathrm{d}x}{|\Omega|}, \Omega)} = \left(\int_{\Omega} |f|^{p}\right)^{\frac{1}{p}} = \frac{1}{|\Omega|^{1/p}} \left(\int_{\Omega} |f|^{p}\right)^{\frac{1}{p}}$$

By convention, we set $\|f\|_{L^{\infty}(\Omega, \operatorname{avg})} = \|f(x)\|_{L^{\infty}_{x}(\overline{|\Omega|}, \Omega)} = \|f\|_{L^{\infty}(\Omega)}$. The implicit measure is of course the Riemannian measure. In such mean integrals, the domain becomes a probability space.

Theorem 53 (Coarea formula).

1. For any $h \in \mathbb{R}^n$, the translation semigroup $(\tau_{th})_{t\geq 0}$ is a C_0 semigroup on $W^{m,p}(\mathbb{R}^n)$, $B^s_{p,q}(\mathbb{R}^n)$ and $F_{p,q}^s(\mathbb{R}^n)$ for $m \in \mathbb{N}_0$, $s \in \mathbb{R}$, $p \in [1,\infty)$, $q \in [1,\infty)$. Consequently, for $p \in [1,\infty)$ and $f \in B_{p,1}^{\frac{1}{p}}(\mathbb{R}^n)$,

$$([0,\infty) \to L^p(\mathbb{R}^{n-1}), t \mapsto \tau_{th} f|_{\mathbb{R}^{n-1}})$$

is continuous and bounded by $C \|f\|_{B^{\frac{1}{p}}_{n-1}(\mathbb{R}^n)}$ for some C > 0.

- 2. Let Ω be a bounded C^{∞} domain in \mathbb{R}^n (or a compact Riemannian manifold with boundary). Let $p \in [1, \infty)$. Then for $f \in B_{p,1}^{\frac{1}{p}}(\Omega)$ and r > 0 small:
 - (a) $([0,r) \to \mathbb{R}, \rho \mapsto \|f\|_{L^p(\partial\Omega_{>\rho})})$ is continuous and bounded by $C \|f\|_{B^{\frac{1}{p}}_{p,1}(\Omega)}$ for some C > 0.

(b)
$$\|f\|_{L^{p}(\Omega_{\leq r})} \sim_{\neg r} \|\|f\|_{L^{p}(\partial\Omega_{>\rho})}\|_{L^{p}_{\rho}((0,r))}$$

- (c) $\|f\|_{L^p(\Omega_{\leq r}, \operatorname{avg})} \lesssim_{\neg r} \|f\|_{B^{\frac{1}{p}}_{p,1}(\Omega)}$ and $\|f\|_{L^p(\Omega_{\leq r}, \operatorname{avg})} \xrightarrow{r \downarrow 0} \|f\|_{L^p(\partial\Omega, \operatorname{avg})}.$
- (d) Let $I \subset \mathbb{R}$ be an open interval and $\mathfrak{f} \in L^p(I \to B_{p,1}^{\frac{1}{p}}(\Omega))$, then $\|\mathfrak{f}\|_{L^p_t B^{\frac{1}{p}}_{p,1}(\Omega)} \gtrsim_{\neg r} \|\mathfrak{f}\|_{L^p_t L^p(\Omega_{\leq r}, \operatorname{avg})} \xrightarrow{r\downarrow 0} \|\mathfrak{f}\|_{L^p_t L^p(\partial\Omega, \operatorname{avg})}$.

3. Let $p \in [1,\infty), f \in W^{1,p}(\Omega)$, show that $\|f\|_{L^p(\Omega_{< r})} \lesssim_{\neg r} r \|f\|_{W^{1,p}(\Omega_{< r})} + r^{\frac{1}{p}} \|f\|_{L^p(\partial\Omega)}$ for r > 0 small.

Proof.

1. Use the density of $\mathcal{S}(\mathbb{R}^n)$ and Lemma 9.

2.

- (a) By partition of unity, geodesic normals, diffeomorphisms and the smallness of r, reduce the problem to the half-space case, which is just 1).
- (b) Approximate f in $B_{p,1}^{\frac{1}{p}}$ by $C^{\infty}(\overline{\Omega})$ functions. This is the well-known coarea formula, which corresponds to Fubini's theorem in the half-space case. Note that $||f||_{L^{p}(\partial\Omega_{>\rho})}$ is defined by the trace theorem. See [Cha06, Section III.5] for more details.
- (c) For r small, $|\Omega_{< r}| \sim |\partial \Omega| r$ and $|\partial \Omega_{> r}| \sim |\partial \Omega|$, so

$$\|f\|_{L^p(\Omega_{\leq r}, \operatorname{avg})} \sim \left\|\|f\|_{L^p(\partial\Omega_{>\rho}, \operatorname{avg})}\right\|_{L^p_\rho((0, r), \operatorname{avg})} \leq \sup_{\rho < r} \|f\|_{L^p(\partial\Omega_{>\rho}, \operatorname{avg})}$$

and
$$\left\| \|f\|_{L^p(\partial\Omega_{>\rho},\mathrm{avg})} \right\|_{L^p_\rho((0,r),\mathrm{avg})} \xrightarrow{r\downarrow 0} \|f\|_{L^p(\partial\Omega,\mathrm{avg})}$$
 by continuity in a)

- (d) Dominated convergence.
- 3. By the trace theorem, WLOG $f \in C^{\infty}(\overline{\Omega})$. By partition of unity and diffeomorphisms, WLOG $\overline{\Omega} = \overline{\mathbb{R}^n_+} = \{(\mathbf{x}, y) : \mathbf{x} \in \mathbb{R}^{n-1}, y \ge 0\}$. Then

$$\begin{split} \|f\|_{L^{p}(\Omega_{< r})} &\sim \left\|\|f(\mathbf{x}, y)\|_{L^{p}_{y}([0, r])}\right\|_{L^{p}_{\mathbf{x}}} \lesssim \left\|\left\|\|\partial_{y}f(\mathbf{x}, \rho)\|_{L^{1}_{\rho}([0, y])} + |f(\mathbf{x}, 0)|\right\|_{L^{p}_{y}([0, r])}\right\|_{L^{p}_{\mathbf{x}}} \\ &\lesssim \left\|\left\|\|\partial_{y}f(\mathbf{x}, \rho)\|_{L^{p}_{\rho}([0, y])} \left|y\right|^{\frac{1}{p'}}\right\|_{L^{p}_{y}([0, r])}\right\|_{L^{p}_{\mathbf{x}}} + r^{\frac{1}{p}} \|f(\mathbf{x}, 0)\|_{L^{p}_{\mathbf{x}}} \\ \end{split}$$
The first term $\lesssim \left\|\|\partial_{y}f(\mathbf{x}, \rho)\|_{L^{p}_{\rho}([0, r])} \left\||y|^{\frac{1}{p'}}\right\|_{L^{p}_{y}([0, r])}\right\|_{L^{p}_{\mathbf{x}}} \lesssim r \|\partial_{y}f\|_{L^{p}(\Omega_{< r})}.$ So we are done.

Theorem 54 (Product estimate). Let M be a bounded C^{∞} domain in \mathbb{R}^n (or a compact Riemannian manifold with boundary). Assume r > 0 small, $f_r \in C^{\infty}(\overline{M})$ with support in $M_{\leq r}$. Then for $p \in (1, \infty)$,
$$g \in B_{p,1}^{\frac{1}{p}}(M):$$

$$\|f_r g\|_{B_{p,1}^{1/p}(M)} \lesssim_{M,\neg r} \|f_r\|_{B_{\infty,1}^{1/p}(M)} \|g\|_{L^p(M_{\leq 4r})} + \|f_r\|_{L^{\infty}(M_{\leq r})} \|g\|_{B_{p,1}^{1/p}(M)}$$

Remark. The theory of product and commutator estimates (Kato-Ponce, Coifman-Meyer etc.) has a long and rich history which we will not recount here (cf. [KP88; Tao07; GO14; NT19]). However, for our intended application, f_r has very small support and we want to use $||g||_{L^p(M < 4r)}$ instead of $||g||_{L^p(M)}$ to control the product. Unfortunately there does not seem to be much, if at all, literature on this issue. This theorem will only be used for Theorem 86, and is not necessary for Onsager's conjecture.

Proof. By diffeomorphisms, partition of unity, and geodesic normals, WLOG assume $M = \overline{\mathbb{R}^n_+}$ with $M_{\leq r} = \{x \in \mathbb{R}^n : 0 \leq x_n < r\}.$

Recall $\|g\|_{L^p(x_n=a)} \lesssim \|g\|_{B^{1/p}_{p,1}(\mathbb{R}^n_+)}$ $\forall 0 \le a < \infty$ where $\|g\|_{L^p(x_n=a)} := \|g\|_{L^p(\{x \in \mathbb{R}^n : x_n=a\})}$ is defined by the trace theorem.

WLOG, assume $||f_r||_{\infty} \leq 1$. Recall the characterization of Besov spaces by ball mean difference (BMD) and write V(x,t) for $V^1(x,t)$ (see Equation (7)). Then

$$\|f_r g\|_{B^{1/p}_{p,1}(M)} \sim \|f_r g\|_{L^p(M)} + \left\| t^{-\frac{1}{p}-n} \left\| \|\Delta_h(f_r g)(x)\|_{L^1_h(V(x,t))} \right\|_{L^p_x(M)} \right\|_{L^1_t(\frac{\mathrm{d}t}{t},(0,1))}$$

The term $||f_rg||_{L^p(M)}$ is easily bounded and thrown away. For the remaining term, we use the identity $\Delta_h(f_rg) = \Delta_h f_rg + \tau_h f_r \Delta_h g$ to bound it by

$$\left\| t^{-\frac{1}{p}-n} \left\| \|\Delta_h f_r(x)\|_{L^1_h(V(x,t))} g(x) \right\|_{L^p_x(M)} \right\|_{L^1(\frac{\mathrm{d}t}{t},(0,1))} + \left\| t^{-\frac{1}{p}-n} \left\| \|f_r\|_{\infty} \left\|\Delta_h g(x)\|_{L^1_h(V(x,t))} \right\|_{L^p_x(M)} \right\|_{L^1(\frac{\mathrm{d}t}{t},(0,1))} + \left\| t^{-\frac{1}{p}-n} \left\| \|f_r\|_{\infty} \left\|\Delta_h g(x)\|_{L^1_h(V(x,t))} \right\|_{L^p_x(M)} \right\|_{L^1(\frac{\mathrm{d}t}{t},(0,1))} + \left\| t^{-\frac{1}{p}-n} \left\| \|f_r\|_{\infty} \left\|\Delta_h g(x)\|_{L^1_h(V(x,t))} \right\|_{L^p_x(M)} \right\|_{L^1(\frac{\mathrm{d}t}{t},(0,1))} + \left\| t^{-\frac{1}{p}-n} \left\| \|f_r\|_{\infty} \left\|\Delta_h g(x)\|_{L^1_h(V(x,t))} \right\|_{L^p_x(M)} \right\|_{L^1(\frac{\mathrm{d}t}{t},(0,1))} + \left\| t^{-\frac{1}{p}-n} \left\| \|f_r\|_{\infty} \left\|\Delta_h g(x)\|_{L^1_h(V(x,t))} \right\|_{L^p_x(M)} \right\|_{L^1(\frac{\mathrm{d}t}{t},(0,1))} + \left\| t^{-\frac{1}{p}-n} \left\| \|f_r\|_{\infty} \left\|\Delta_h g(x)\|_{L^1_h(V(x,t))} \right\|_{L^p_x(M)} \right\|_{L^1(\frac{\mathrm{d}t}{t},(0,1))} + \left\| t^{-\frac{1}{p}-n} \left\| \|f_r\|_{\infty} \left\|\Delta_h g(x)\|_{L^1_h(V(x,t))} \right\|_{L^p_x(M)} \right\|_{L^1(\frac{\mathrm{d}t}{t},(0,1))} + \left\| t^{-\frac{1}{p}-n} \left\| \|f_r\|_{\infty} \left\|\Delta_h g(x)\|_{L^1_h(V(x,t))} \right\|_{L^p_x(M)} \right\|_{L^1(\frac{\mathrm{d}t}{t},(0,1))} + \left\| t^{-\frac{1}{p}-n} \left\| \|f_r\|_{\infty} \left\|\Delta_h g(x)\|_{L^1(\frac{\mathrm{d}t}{t},(0,1))} \right\|_{L^p_x(M)} \right\|_{L^1(\frac{\mathrm{d}t}{t},(0,1))} + \left\| t^{-\frac{1}{p}-n} \left\| \|f_r\|_{\infty} \left\|\Delta_h g(x)\|_{L^1(\frac{\mathrm{d}t}{t},(0,1))} \right\|_{L^p_x(M)} \right\|_{L^p_x(M)} + \left\| t^{-\frac{1}{p}-n} \left\| \|f_r\|_{\infty} \left\|\Delta_h g(x)\|_{L^p_x(M)} \right\|_{L^p_x(M)} \right\|_{L^p_x(M)} + \left\| t^{-\frac{1}{p}-n} \left\| \|f_r\|_{\infty} \right\|_{L^p_x(M)} + \left\| t^{-\frac{1}{p}-n} \right\|_{L^p_x(M)} + \left\| t^{-\frac{1}{p}-n} \right\|_{L^p_x(M)} + \left\| t^{-\frac{1}{p}-n} \right\|_{L^p_x(M)} + \left\| t^{-\frac{1}{p}-n} \right\|$$

The second term here is just $||f_r||_{L^{\infty}} ||g||_{B^{1/p}_{p,1}(M)}$, so throw it away. For the remaining term, by using $||\cdot||_{L^p(M)} \lesssim ||\cdot||_{L^p(M_{\leq 4r})} + ||\cdot||_{L^p(M_{\geq 4r})}$ and

$$\left\| \|\Delta_h f_r(x)\|_{L^1_h(V(x,t))} g(x) \right\|_{L^p_x(M_{<4r})} \lesssim \left\| \|\Delta_h f_r(x)\|_{L^1_h(V(x,t))} \right\|_{L^\infty_x(M_{<4r})} \|g(x)\|_{L^p_x(M_{<4r})}$$

we are left with

$$\|f_r\|_{B^{1/p}_{\infty,1}(M)}\|g\|_{L^p(M_{\leq 4r})} + \left\|t^{-\frac{1}{p}-n} \left\|\|\Delta_h f_r(x)\|_{L^1_h(V(x,t))}g(x)\right\|_{L^p_x(M_{\geq 4r})}\right\|_{L^1(\frac{\mathrm{d}t}{t},(0,1))}$$

Throwing away the first term, we have arrived at the important estimate: what happens on $M_{>4r}$. It will turn out that the values of g on $M_{>4r}$ are well-controlled by $\|g\|_{B^{1/p}_{p,1}(M)}$. To begin, recall f_r is supported on $M_{< r}$ and use the crude geometric estimate

$$t^{-n} \|\Delta_h f_r(x)\|_{L^1_h(V(x,t))} = t^{-n} \|f_r(x+h)\|_{L^1_h(V(x,t))} \lesssim \frac{|B(x,t) \cap M_{\leq r}|}{|B(x,t)|} \lesssim \frac{r}{x_n} \mathbf{1}_{t > x_n - r} \ \forall x \in M_{>4r}, \forall t \in (3r,1)$$

Note that t > 3r comes from $t > x_n - r > 4r - r$. So we have used the "room" from 4r to get an O(r)-lower bound for t. By $x_n < r + t$, we now only need to bound

$$\left\| t^{-\frac{1}{p}} \left\| g(x) \frac{r}{x_n} \right\|_{L^p_x(M_{[4r,r+t]})} \right\|_{L^1(\frac{\mathrm{d}t}{t},(3r,1))}$$

Obviously, we will integrate g on x_n -slices (using p > 1):

$$\left\|g(x)\frac{r}{x_n}\right\|_{L^p_x(M_{[4r,r+t]})} = r \left\|\frac{1}{\rho} \left\|g\right\|_{L^p(x_n=\rho)}\right\|_{L^p_\rho([4r,r+t])} \lesssim \left\|g\right\|_{B^{1/p}_{p,1}(M)} r \left\|\frac{1}{\rho}\right\|_{L^p_\rho([4r,\infty))} \lesssim r^{\frac{1}{p}} \|g\|_{B^{1/p}_{p,1}(M)}$$

Then we are done (using $p < \infty$):

$$r^{\frac{1}{p}} \left\| t^{-\frac{1}{p}} \right\|_{L^{1}(\frac{dt}{t},(3r,1))} = \left\| \left(\frac{r}{t} \right)^{\frac{1}{p}} \right\|_{L^{1}(\frac{dt}{t},(3r,1))} = \left\| \left(\frac{1}{t} \right)^{\frac{1}{p}} \right\|_{L^{1}(\frac{dt}{t},(3,\frac{1}{r}))} \lesssim_{\neg r} 1$$

6 Hodge theory

We stick closely to the terminology and symbols of [Sch95], with some careful exceptions.

6.1 The setting

Definition 55. Define a ∂ -manifold as a paracompact, Hausdorff, metric-complete, oriented, smooth manifold, with no or smooth boundary.

Note that this means $B_{\mathbb{R}^n}(0,1)$ is not a ∂ -manifold (as it is not complete), but $\overline{B_{\mathbb{R}^n}(0,1)}$ is.

For the rest of this paper, unless mentioned otherwise, we work on M which is a compact Riemannian *n*-dimensional ∂ -manifold (where $n \geq 2$), and use ν to denote the outwards unit normal vector field on ∂M . As before, define $M_{>r} = \{x \in M : \operatorname{dist}(x, \partial M) > r\}$, and similarly for $M_{>r}, M_{< r}, M_{[r_1, r_2]}$ etc.

For r > 0 small, the map $(\partial M \times [0, r) \to M_{< r}, (x, t) \mapsto \exp_x(-t\nu))$ is a diffeomorphism, which we call a **Riemannian collar**. Then ν can be extended via geodesics to a smooth vector field $\tilde{\nu}$ which is of unit length near the boundary (cut off at some point away from the boundary, but we only care about the area near the boundary).

Let vol stand for the Riemannian volume form orienting M and vol ∂ for that of ∂M . Let $j: \partial M \hookrightarrow M$ be the smooth inclusion map and ι stand for interior product (contraction) of differential forms. Note that for a smooth differential form $\omega, j^*\omega$ only depends on $\omega|_{\partial M}$, so by abuse of notation, we can write

$$\operatorname{vol}_{\partial} = j^*(\iota_{\nu} \operatorname{vol})$$

where ι_{ν} vol $\in \Omega^{n-1}(M) \mid_{\partial M}$. Additionally, the Stokes theorem reads $\int_{M} d\omega = \int_{\partial M} j^* \omega$ for $\omega \in \Omega^{n-1}(M)$.

6.1.1 Vector bundles

Let \mathbb{F} be a real vector bundle over M with a Riemannian fiber metric $\langle \cdot, \cdot \rangle_{\mathbb{F}}$.

Define

- $\Gamma(\mathbb{F})$: the space of smooth sections of \mathbb{F}
- $\Gamma_c(\mathbb{F})$: smooth sections with compact support (so $\Gamma_c(\mathbb{F}) = \Gamma(\mathbb{F})$ since M is compact)
- $\Gamma_{00}(\mathbb{F})$: smooth sections with compact support in \mathring{M} (the interior of M).

Remark. We are following [Sch95], where Hodge theory is also formulated for non-compact M. In the book, $\Gamma_0 \mathbb{F}$ is used instead of $\Gamma_{00} \mathbb{F}$ to denote compact support in \mathring{M} . As that can be confused with having zero trace, we opt to write $\Gamma_{00} \mathbb{F}$ instead.

Then on $\Gamma_c(\mathbb{F})$, define the dot product

$$\langle \langle \sigma, \theta \rangle \rangle = \int_M \langle \sigma, \theta \rangle_{\mathbb{F}} \operatorname{vol}$$

and $|\sigma|_{\mathbb{F}} = \sqrt{\langle \sigma, \sigma \rangle_{\mathbb{F}}}$. Then for $p \in [1, \infty)$, $L^p \Gamma(\mathbb{F})$ is the completion of $\Gamma_c(\mathbb{F})$ under the norm

$$\|\sigma\|_{L^p\Gamma(\mathbb{F})} = \||\sigma|_{\mathbb{F}}\|_{L^p(M)}$$

Let $\nabla^{\mathbb{F}}$ be a connection on \mathbb{F} . Then for $\sigma \in \Gamma(\mathbb{F}), \nabla^{\mathbb{F}}\sigma \in \Gamma(T^*M \otimes \mathbb{F})$ and we can define the fiber metric

$$\langle \alpha \otimes \sigma, \beta \otimes \theta \rangle_{T^*M \otimes \mathbb{F}} = \langle \alpha, \beta \rangle_{T^*M} \left\langle \sigma, \theta \right\rangle_{\mathbb{F}}$$

In local coordinates (Einstein notation):

$$\left\langle \nabla^{\mathbb{F}}\sigma, \nabla^{\mathbb{F}}\theta \right\rangle_{T^{*}M \otimes \mathbb{F}} = \left\langle dx^{i} \otimes \nabla^{\mathbb{F}}_{i}\sigma, dx^{j} \otimes \nabla^{\mathbb{F}}_{j}\sigma \right\rangle_{\mathbb{F}} = \left\langle dx^{i}, dx^{j} \right\rangle_{T^{*}M} \left\langle \nabla^{\mathbb{F}}_{i}\sigma, \nabla^{\mathbb{F}}_{j}\theta \right\rangle_{\mathbb{F}} = g^{ij} \left\langle \nabla^{\mathbb{F}}_{i}\sigma, \nabla^{\mathbb{F}}_{j}\theta \right\rangle_{\mathbb{F}}$$

For higher derivatives, define the k-jet fiber metric

$$\langle \sigma, \theta \rangle_{J^k \mathbb{F}} = \sum_{0 \le j \le k} \left\langle \left(\nabla^{\mathbb{F}} \right)^{(j)} \sigma, \left(\nabla^{\mathbb{F}} \right)^{(j)} \theta \right\rangle_{\left(\bigotimes^j T^* M \right) \otimes \mathbb{F}}$$

and $|\sigma|_{J^k\mathbb{F}} = \sqrt{\langle \sigma, \sigma \rangle_{J^k\mathbb{F}}}$. Then we have Cauchy-Schwarz: $|\langle \sigma, \theta \rangle_{J^k\mathbb{F}}| \le |\sigma|_{J^k\mathbb{F}} |\theta|_{J^k\mathbb{F}}$.

Then for $m \in \mathbb{N}_0, p \in [1, \infty)$, we define the Sobolev space $W^{m,p}\Gamma(\mathbb{F})$ as the completion of $\Gamma_c(\mathbb{F})$ under the norm

$$\|\sigma\|_{W^{m,p}\Gamma(\mathbb{F})} = \||\sigma|_{J^m\mathbb{F}}\|_{L^p(M)}$$

It is worth noting that $|\sigma|_{J^m\mathbb{F}}$, up to some constants, does not depend on $\nabla^{\mathbb{F}}$. Indeed, assume there is another connection $\widetilde{\nabla}^{\mathbb{F}}$, then $\nabla^{\mathbb{F}} - \widetilde{\nabla}^{\mathbb{F}}$ is tensorial:

$$\left(\nabla_X^{\mathbb{F}} - \widetilde{\nabla}_X^{\mathbb{F}}\right)(f\sigma) = f\left(\nabla_X^{\mathbb{F}} - \widetilde{\nabla}_X^{\mathbb{F}}\right)(\sigma) = \left(\nabla_{fX}^{\mathbb{F}} - \widetilde{\nabla}_{fX}^{\mathbb{F}}\right)(\sigma) \text{ for } f \in C^{\infty}(M), \sigma \in \Gamma(\mathbb{F}), X \in \mathfrak{X}M$$

So there is a $C^{\infty}(M)$ -multilinear map $A : \mathfrak{X}M \otimes_{C^{\infty}(M)} \Gamma(\mathbb{F}) \to \Gamma(\mathbb{F})$ such that $\left(\nabla_{X}^{\mathbb{F}} - \widetilde{\nabla}_{X}^{\mathbb{F}}\right)(\sigma) = A(X, \sigma)$. By the compactness of M and the boundedness of A, we conclude $|\sigma|_{J^m\mathbb{F},\nabla^{\mathbb{F}}} \sim |\sigma|_{J^m\mathbb{F},\widetilde{\nabla}^{\mathbb{F}}}$. Therefore the topology of $W^{m,p}\Gamma(\mathbb{F})$ is uniquely defined.

Definition 56 (Distributions). Set $\mathscr{D}\Gamma(\mathbb{F}) = \Gamma_{00}(\mathbb{F})$ as the space of **test sections** and $\mathscr{D}'\Gamma(\mathbb{F}) = (\mathscr{D}\Gamma(\mathbb{F}))^*$ the space of **distributional sections**. As usual, in the category of locally convex TVS, $\mathscr{D}\Gamma(\mathbb{F})$ is given **Schwartz's topology** as the colimit of $\{\Gamma(\mathbb{F})_K : K \subset \mathring{M} \text{ compact}\}$, where $\Gamma(\mathbb{F})_K := \{\sigma \in \Gamma(\mathbb{F}) : \text{supp } \sigma \subset K\}$ has the **Frechet** C^{∞} topology.

6.1.2 Compatibility with scalar function spaces

We aim to show that the global definitions of Sobolev spaces in Subsection 6.1.1 are compatible with the definitions of Sobolev spaces by local coordinates.

Let $(\psi_{\alpha}, U_{\alpha})_{\alpha}$ be a finite partition of unity, where U_{α} is open in M and ψ_{α} is supported in U_{α} . Normally in differential geometry, U_{α} is diffeomorphic to either $\overline{\mathbb{R}^n_+} \cap B_{\mathbb{R}^n}(0, 1)$ or $B_{\mathbb{R}^n}(0, 1)$. However, it is problematic that the half-ball does not have C^{∞} boundary, so we use some piecewise-linear functions and mollification to create a bounded C^{∞} domain.



Figure 2: Smoothing the corners

So WLOG, $\overline{U_{\alpha}}$ is diffeomorphic to the closure of a bounded C^{∞} domain in \mathbb{R}^n , and scalar function spaces are well-defined on U_{α} (recall Definition 44). Note that supp ψ_{α} might intersect with ∂M .

For U_{α} chosen small enough, the bundle \mathbb{F} on U_{α} is diffeomorphic to $U_{\alpha} \times F$ (where F is the typical fiber of \mathbb{F}).

Let $(e^{\alpha}_{\beta})_{\beta}$ be the coordinate sections on $\operatorname{supp} \psi_{\alpha}$, and cut off such that $\operatorname{supp} \psi_{\alpha} \subset (\operatorname{supp} e^{\alpha}_{\beta}) \subset \operatorname{supp} e^{\alpha}_{\beta} \subset U_{\alpha}$. Let $\sigma \in \Gamma(\mathbb{F})$. Then there exist $c^{\alpha}_{\beta}(\sigma) \in C^{\infty}_{c}(U_{\alpha})$ such that $\operatorname{supp} c^{\alpha}_{\beta}(\sigma) \subset \operatorname{supp} \psi_{\alpha}, \ \psi_{\alpha}\sigma = \sum_{\beta} c^{\alpha}_{\beta}(\sigma) e^{\alpha}_{\beta}$ and

$$\sigma = \sum_{\alpha,\beta} c^{\alpha}_{\beta}(\sigma) e^{\alpha}_{\beta}$$

Now, observe that $|\sigma|_{\mathbb{F}} \sim \sum_{\alpha} |\psi_{\alpha}\sigma|_{\mathbb{F}}$ and

$$\left|\psi_{\alpha}\sigma\right|_{\mathbb{F}} = \left(\sum_{\beta,\beta'} c_{\beta}^{\alpha}(\sigma)c_{\beta'}^{\alpha}(\sigma)\left\langle e_{\beta}^{\alpha}, e_{\beta'}^{\alpha}\right\rangle_{\mathbb{F}}\right)^{\frac{1}{2}} \sim \left(\sum_{\beta} \left|c_{\beta}^{\alpha}(\sigma)\right|^{2}\right)^{\frac{1}{2}}$$

To see this, let $x \in \operatorname{supp} \psi_{\alpha}$ and $\left\langle e_{\beta}^{\alpha}, e_{\beta'}^{\alpha} \right\rangle_{\mathbb{F}}(x) = B_{\beta\beta'}(x)$. Then $B_x(u, v) := \sum_{\beta,\beta'} u_{\beta}v_{\beta'}B_{\beta\beta'}(x)$ is a positive-definite inner product, which induces a norm on a finite-dimensional vector space, where all norms are equivalent. Then simply note $B_x(u, u)$ is continuous in variable $x \in \operatorname{supp} \psi_{\alpha}$.

Also, in local coordinates, there are $s_{i\beta}^{\gamma} \in C_c^{\infty}(U_{\alpha})$ such that $\nabla_i^{\mathbb{F}} e_{\beta}^{\alpha} = \sum_{\gamma} s_{i\beta}^{\gamma} e_{\gamma}^{\alpha}$ on $\operatorname{supp} \psi_{\alpha}$. Then

$$\nabla_{i}^{\mathbb{F}}(\psi_{\alpha}\sigma) = \sum_{\beta} \partial_{i}c_{\beta}^{\alpha}(\sigma)e_{\beta}^{\alpha} + \sum_{\beta,\gamma} c_{\beta}^{\alpha}(\sigma)s_{i\beta}^{\gamma}e_{\gamma}^{\alpha} = \sum_{\beta} d_{i\beta}^{\alpha}(\sigma)e_{\beta}^{\alpha} \quad \text{where } d_{i\beta}^{\alpha}(\sigma) = \partial_{i}c_{\beta}^{\alpha}(\sigma) + \sum_{\gamma} c_{\gamma}^{\alpha}(\sigma)s_{i\gamma}^{\beta}e_{\gamma}^{\alpha}$$

So $|\sigma|_{J^{1}\mathbb{F}} \sim \sum_{\alpha,\beta} |c_{\beta}^{\alpha}(\sigma)| + \sum_{\alpha,\beta,i} |d_{i\beta}^{\alpha}(\sigma)| \sim \sum_{\alpha,\beta} |c_{\beta}^{\alpha}(\sigma)| + \sum_{\alpha,\beta,i} |\partial_{i}c_{\beta}^{\alpha}(\sigma)|.$ Similarly $|\sigma|_{J^{m}\mathbb{F}} \sim \sum_{\alpha,\beta} \sum_{k \leq m} |\nabla^{(k)}c_{\beta}^{\alpha}(\sigma)|.$ So for $m \in \mathbb{N}_{0}, p \in [1, \infty),$ $\|\sigma\|_{W^{m,p}} \sim \sum_{\alpha,\beta} \|c_{\beta}^{\alpha}(\sigma)\|_{W^{m,p}(U_{\alpha},\mathbb{R})}$

Now define $S\sigma = (c^{\alpha}_{\beta}(\sigma))_{\alpha,\beta}$ and $R(c^{\alpha}_{\beta})_{\alpha,\beta} = \sum_{\alpha,\beta} c^{\alpha}_{\beta} e^{\alpha}_{\beta}$. Then RS = 1 on $\Gamma(\mathbb{F})$ and P := SR is a projection on $\prod_{\alpha,\beta} C^{\infty}(\overline{U_{\alpha}})$. Note that P depends on the choice of partition of unity. By looking into the definitions of R and S, we can extend this to have P = SR as a continuous projection on $\prod_{\alpha,\beta} L^{1}(U_{\alpha})$ and

$$\left\| P\left(c_{\beta}^{\alpha}\right)_{\alpha,\beta} \right\|_{\prod_{\alpha,\beta} W^{m,p}(U_{\alpha})} \lesssim \sum_{\alpha,\beta} \left\| c_{\beta}^{\alpha} \right\|_{W^{m,p}(U_{\alpha})} \text{ for } m \in \mathbb{N}_{0}, p \in [1,\infty], c_{\beta}^{\alpha} \in W^{m,p}(U_{\alpha})$$

The keen reader should have noticed we never mentioned the case $p = \infty$ in Subsection 6.1.1 as we defined $W^{m,p}\Gamma(\mathbb{F})$ by the completion of smooth sections, and $C^{\infty}(M)$ is not dense in $W^{m,\infty}(M)$. Now,

however, by using local coordinates, we are justified in defining $W^{m,p}\Gamma(\mathbb{F}) = \{\sum_{\alpha,\beta} c_{\beta}^{\alpha} e_{\beta}^{\alpha} : c_{\beta}^{\alpha} \in W^{m,p}(U_{\alpha})\}$ for $m \in \mathbb{N}_0, p \in [1, \infty]$ with the norm defined (up to equivalent norms) as

$$\left\|\sum_{\alpha,\beta} c^{\alpha}_{\beta} e^{\alpha}_{\beta}\right\|_{W^{m,p}\Gamma\mathbb{F}} := \left\|S\sum_{\alpha,\beta} c^{\alpha}_{\beta} e^{\alpha}_{\beta}\right\|_{\prod_{\alpha,\beta} W^{m,p}(U_{\alpha})}$$

Then $B_{p,q}^s\Gamma(\mathbb{F})$ and $F_{p,q}^s\Gamma(\mathbb{F})$ can be defined similarly. In other words, for $m \in \mathbb{N}_0, p \in [1,\infty], q \in [1,\infty], s \ge 0$:

- $W^{m,p}\Gamma(\mathbb{F}) \simeq P \prod_{\alpha,\beta} W^{m,p}(U_{\alpha})$
- $B_{p,q}^{s}\Gamma(\mathbb{F}) \simeq P \prod_{\alpha,\beta} B_{p,q}^{s}(U_{\alpha})$
- $F_{p,q}^s \Gamma(\mathbb{F}) \simeq P \prod_{\alpha,\beta} F_{p,q}^s(U_\alpha), \ p \neq \infty$

By using Blackbox 46, we can show the Banach topologies of these spaces are uniquely defined (independent of the choices of $\psi_{\alpha}, U_{\alpha}$). For convenience (such as working with Holder's inequality), we still use the Sobolev norms $W^{m,p}$ ($m \in \mathbb{N}_0, p \in [1, \infty)$) defined globally in Subsection 6.1.1.

All theorems from section 5 that worked on bounded C^{∞} domains carry over to our setting on M, mutatis mutandis. For instance, $B_{3,1}^{\frac{1}{3}}\Gamma(\mathbb{F}) \twoheadrightarrow L^3 \Gamma(\mathbb{F})|_{\partial M}$ is a continuous surjection and

$$B_{3,1}^{\frac{1}{3}}\Gamma(\mathbb{F}) = \left(L^{3}\Gamma(\mathbb{F}), W^{1,3}\Gamma(\mathbb{F})\right)_{\frac{1}{2},1}$$

Moreover, for $p \in (1, \infty)$, $L^p \Gamma(\mathbb{F})$ is reflexive. By Holder's inequality, $(L^p \Gamma(\mathbb{F}))^* = L^{p'} \Gamma(\mathbb{F})$ for $p \in (1, \infty)$.

6.1.3 Complexification issue

A small step which we omitted is complexification. As \mathbb{F} is a real vector bundle, the previous definitions only give $W^{m,p}\Gamma(\mathbb{F}) \simeq P \prod_{\alpha,\beta} W^{m,p}(U_{\alpha},\mathbb{R})$ for $m \in \mathbb{N}_0, p \in [1,\infty]$. In working with real manifolds, differential forms/tensors and their dot products, we always assume real-valued coefficients for sections, but whenever we need to use theorems involving complex Banach spaces or the theory of function spaces, we assume an implicit complexification step. Fortunately, no complications arise from complexification (see Section A for the full reasoning), so for the rest of the paper we can ignore this detail. When we want to be explicit, we will specify the scalars we are using, e.g. $\mathbb{R}W^{m,p}\Gamma(\mathbb{F})$ versus $\mathbb{C}W^{m,p}\Gamma(\mathbb{F})$.

6.2 Differential forms & boundary

Unless mentioned otherwise, the metric is the Riemannian metric, and the connection is the **Levi-Civita** connection.

For $X \in \mathfrak{X}M$, define $\mathbf{n}X = \langle X, \nu \rangle \nu \in \mathfrak{X}M|_{\partial M}$ (the **normal part**) and $\mathbf{t}X = X|_{\partial M} - \mathbf{n}X$ (the **tangential part**). We note that $\mathbf{t}X$ and $\mathbf{n}X$ only depend on $X|_{\partial M}$, so \mathbf{t} and \mathbf{n} can be defined on $\mathfrak{X}M|_{\partial M}$, and by abuse of notation, $\mathbf{t}(\mathfrak{X}M|_{\partial M}) \xrightarrow{\sim} \mathfrak{X}(\partial M)$.

For $\omega \in \Omega^{k}(M)$, define $\mathbf{t}\omega$ and $\mathbf{n}\omega$ by

$$\mathbf{t}\omega(X_1,...,X_k) := \omega(\mathbf{t}X_1,...,\mathbf{t}X_k) \ \forall X_j \in \mathfrak{X}M, j = 1,...,k$$

and $\mathbf{n}\omega = \omega|_{\partial M} - \mathbf{t}\omega$. By abuse of notation, we similarly observe that $\mathbf{t}\left(\Omega^{k}(M)\Big|_{\partial M}\right) \xrightarrow{\sim} \Omega^{k}(\partial M) = j^{*}\left(\Omega^{k}(M)\Big|_{\partial M}\right) = j^{*}\left(\Omega^{k}(M)\Big|_{\partial M}\right) = j^{*}\left(\Omega^{k}(M)\right).$

Recall the **musical isomorphism**: $X_p^{\flat}(Y_p) = \langle X_p, Y_p \rangle$ and $\langle \omega_p^{\sharp}, Y_p \rangle = \omega_p(Y_p)$ for $p \in M, \omega_p \in T_p^*M, X_p \in T_pM, Y_p \in T_pM$.

Recall the usual Hodge star operator $\star : \Omega^k(M) \xrightarrow{\sim} \Omega^{n-k}(M)$, exterior derivative $d : \Omega^k(M) \rightarrow \Omega^{k+1}(M)$, codifferential $\delta : \Omega^k(M) \rightarrow \Omega^{k-1}(M)$, and Hodge Laplacian $\Delta = -(d\delta + \delta d)$ (cf. [Tayl1a, Section 2.10] and [Sch95, Definition 1.2.2]).

We will often use **Penrose abstract index notation** (cf. [Wal84, Section 2.4]), which should not be confused with the similar-looking **Einstein notation** for local coordinates, or the similar-sounding **Penrose graphical notation**. In Penrose notation, we collect the usual identities in differential geometry (cf. [Lee09]):

- For any tensor $T_{a_1...a_k}$, define $(\nabla T)_{ia_1...a_k} = \nabla_i T_{a_1...a_k}$ and div $T = \nabla^i T_{ia_2...a_k}$.
- $(d\omega)_{ba_1...a_k} = (k+1)\widetilde{\nabla}_{[b}\omega_{a_1...a_k]} \quad \forall \omega \in \Omega^k(M)$ where $\widetilde{\nabla}$ is any torsion-free connection.
- $\bullet \ (\delta \omega)_{a_1 \ldots a_{k-1}} = \nabla^b \omega_{b a_1 \ldots a_{k-1}} = (\operatorname{div} w)_{a_1 \ldots a_{k-1}} \forall \omega \in \Omega^k(M)$
- $(\nabla_a \nabla_b \nabla_b \nabla_a) T^{ij}{}_{kl} = -R_{ab\sigma}{}^i T^{\sigma j}{}_{kl} R_{ab\sigma}{}^j T^{i\sigma}{}_{kl} + R_{abk}{}^{\sigma} T^{ij}{}_{\sigma l} + R_{abl}{}^{\sigma} T^{ij}{}_{k\sigma}$ for any tensor $T^{ij}{}_{kl}$, where R is the **Riemann curvature tensor** and ∇ the Levi-Civita connection. Similar identities hold for other types of tensors. When we do not care about the exact indices and how they contract, we can just write the **schematic identity** $(\nabla_a \nabla_b - \nabla_b \nabla_a) T^{ij}{}_{kl} = R * T$. As R is bounded on compact M, interchanging derivatives is a zeroth-order operation on M.
- For tensor $T_{a_1...a_k}$, define the Weitzenbock curvature operator

$$\operatorname{Ric}(T)_{a_1...a_k} = 2\sum_{j=1}^k \nabla_{[i} \nabla_{a_j]} T_{a_1...a_{j-1}}{}^i{}_{a_{j+1}...a_k} = \sum_j R_{a_j}{}^{\sigma} T_{a_1...a_{j-1}\sigma a_{j+1}...a_k} - \sum_{j \neq l} R_{a_j}{}^{\mu}{}_{a_l}{}^{\sigma} T_{a_1...\sigma ...\mu ...a_k}$$

where $R_{ab} = R_{a\sigma b}{}^{\sigma}$ is the **Ricci tensor**. The invariant form is

$$\operatorname{Ric}(T)(X_1,...X_k) = \sum_{a} \left(R(\partial_i, X_a) T \right) (X_1, ..., X_{a-1}, \partial^i, X_{a+1}, ..., X_k) \; \forall X_j \in \mathfrak{X} M$$

where $\partial^i = g^{ij}\partial_j$ and $R(\partial_i, \partial_j) = \nabla_i \nabla_j - \nabla_j \nabla_i$ (Penrose notation). Note that $\langle R(\partial_a, \partial_b)\partial_d, \partial_c \rangle = R_{abcd}$. Special cases include $\operatorname{Ric}(f) = 0 \ \forall f \in C^{\infty}(M)$ and $\operatorname{Ric}(X)_a = R_a{}^{\sigma}X_{\sigma} \ \forall X \in \mathfrak{X}M$ (justifying the notation Ric).

In local coordinates

$$\operatorname{Ric}\left(\omega\right) = dx^{j} \wedge \left(R(\partial_{i}, \partial_{j})\omega \cdot \partial^{i}\right) \ \forall \omega \in \Omega^{k}(M),$$

where \cdot stands for contraction (interior product). Then we have the **Weitzenbock formula**:

$$\Delta \omega = \nabla_i \nabla^i \omega - \operatorname{Ric}(\omega) \; \forall \omega \in \Omega^k(M)$$

where $\nabla_i \nabla^i \omega = \operatorname{tr}(\nabla^2 \omega)$ is also called the **connection Laplacian**, which differs from the Hodge Laplacian by a zeroth-order term. The geometry of M and differential forms are more easily handled by the Hodge Laplacian, while the connection Laplacian is more useful in calculations with tensors and the Penrose notation.

• For tensors $T_{a_1...a_k}$ and $Q_{a_1...a_k}$, the **tensor inner product** is $\langle T, Q \rangle = T_{a_1...a_k}Q^{a_1...a_k}$. But for $\omega, \eta \in \Omega^k(M)$, there is another dot product, called the **Hodge inner product**, where

$$\left< \omega, \eta \right>_{\Lambda} = \frac{1}{k!} \left< \omega, \eta \right>$$

So $|\omega|_{\Lambda} = \sqrt{\frac{1}{k!}} |\omega|$. Then we define $\langle \langle \omega, \eta \rangle \rangle = \int_M \langle \omega, \eta \rangle$ vol and $\langle \langle \omega, \eta \rangle \rangle_{\Lambda} = \int_M \langle \omega, \eta \rangle_{\Lambda}$ vol. Recall that $\omega \wedge \star \eta = \langle \omega, \eta \rangle_{\Lambda}$ vol $\forall \omega \in \Omega^k(M), \forall \eta \in \Omega^k(M)$. Also

$$\left\langle \left\langle d\omega,\eta\right\rangle \right\rangle _{\Lambda}=\left\langle \left\langle \omega,\delta\eta\right\rangle \right\rangle _{\Lambda}\ \forall\omega\in\Omega_{00}^{k}(M),\forall\eta\in\Omega_{00}^{k+1}(M)$$

So $\langle \cdot, \cdot \rangle_{\Lambda}$ is more convenient for integration by parts and the Hodge star. Nevertheless, as they only differ up to a constant factor, we can still define $W^{m,p}\Omega^k(M)$ $(m \in \mathbb{N}_0, p \in [1,\infty))$ by $\langle \cdot, \cdot \rangle$ as in Subsection 6.1. Finally, by the Weitzenbock formula and Penrose notation, we easily get the **Bochner formula**:

$$\frac{1}{2}\Delta\left(|\omega|^{2}\right) = \frac{1}{2}\nabla_{i}\nabla^{i}\left(\langle\omega,\omega\rangle\right) = \langle\Delta\omega,\omega\rangle + |\nabla\omega|^{2} + \langle\operatorname{Ric}\left(\omega\right),\omega\rangle$$

Remark. In [Sch95], the conventions are a bit different, with $\Delta = (d\delta + \delta d)$, $\Delta^{\Lambda} = -\nabla_i \nabla^i$, $R^W = -$ Ric and \mathcal{N} the inwards unit normal vector field. Also the difference between $\langle \langle \cdot, \cdot \rangle \rangle$ and $\langle \langle \cdot, \cdot \rangle \rangle_{\Lambda}$ is not made explicit in the book. We will not use such notation.

Lemma 57. Some basic identities:

$$\begin{array}{l} 1. \ \forall \omega \in \Omega^{k}\left(M\right): \mathbf{t}\omega = 0 \iff j^{*}\omega = 0 \ . \ Similarly, \ \mathbf{n}\omega = 0 \iff \iota_{\nu}\omega = 0.\\ 2. \ \left(\mathbf{t}X\right)^{\flat} = \mathbf{t}(X^{\flat}) \ \forall X \in \mathfrak{X}M\\ 3. \ j^{*}\mathbf{t}\omega = j^{*}\omega, \ \mathbf{t}\omega = \iota_{\nu}(\nu^{\flat} \wedge \omega), \ \mathbf{n}\omega = \nu^{\flat} \wedge \iota_{\nu}\omega, \ \mathbf{t}(\omega \wedge \eta) = \mathbf{t}\omega \wedge \mathbf{t}\eta \ \forall \omega \in \Omega^{k}(M), \forall \eta \in \Omega^{l}(M)\\ 4. \ \left\langle\langle \mathbf{t}\omega, \eta \right\rangle\rangle_{\Lambda} = \left\langle\langle \mathbf{t}\omega, \mathbf{t}\eta \right\rangle\rangle_{\Lambda} = \left\langle\langle \mathbf{t}\omega, \mathbf{t}\eta \right\rangle\rangle_{\Lambda} \ \forall \omega, \eta \in \Omega^{k}(M)\\ 5. \ \mathbf{t} \ (\star\omega) = \star \left(\mathbf{n}\omega\right), \ \mathbf{n} \ (\star\omega) = \star \left(\mathbf{t}\omega\right), \ \star d\omega = (-1)^{k+1} \ \delta \star \omega, \ \star \delta\omega = (-1)^{k} \ d \star \omega, \ \star \Delta\omega = \Delta \star \omega \ \forall \omega \in \Omega^{k}(M)\\ 6. \ j^{*}\mathbf{t}d\omega = j^{*}d\omega = d^{\partial M}j^{*}\omega = d^{\partial M}j^{*}\mathbf{t}\omega \ \forall \omega \in \Omega^{k}(M)\\ 7. \ Let \ \omega \in \Omega^{k}(M). \ If \ \mathbf{t}\omega = 0 \ then \ \mathbf{t}d\omega = 0. \ If \ \mathbf{n}\omega = 0 \ then \ \mathbf{n}\delta\omega = 0.\\ 8. \ \iota_{\nu}\omega = \mathbf{t} \ (\iota_{\nu}\omega) = \iota_{\nu}\mathbf{n}\omega \ \forall \omega \in \Omega^{k}(M)\\ 9. \ j^{*} \ (\omega \wedge \star \eta) = \left\langle j^{*}\omega, j^{*}\iota_{\nu}\eta \right\rangle_{\Lambda} \ \mathrm{vol}_{\partial} \ \forall \omega \in \Omega^{k}(M), \ \forall \eta \in \Omega^{k+1}(M)\\ \\ \end{array}$$

$$\nu^{\flat} \wedge \mathbf{t}\omega \wedge \mathbf{t} (\star \eta) = \nu^{\flat} \wedge \mathbf{t}\omega \wedge \star \mathbf{n}\eta = \left\langle \nu^{\flat} \wedge \mathbf{t}\omega, \mathbf{n}\eta \right\rangle_{\Lambda} \text{vol} = \left\langle \nu^{\flat} \wedge \mathbf{t}\omega, \nu^{\flat} \wedge \iota_{\nu}\eta \right\rangle_{\Lambda} \text{vol} = \left\langle \mathbf{t}\omega, \iota_{\nu}\eta \right\rangle_{\Lambda} \text{vol}$$

Theorem 58 (Integration of tensors and forms by parts).

1. For tensors $T_{a_1...a_k}$ and $Q_{a_1...a_{k+1}}$,

$$\int_{M} \nabla_i \left(T_{a_1 \dots a_k} Q^{ia_1 \dots a_k} \right) = \int_{M} \nabla_i T_{a_1 \dots a_k} Q^{ia_1 \dots a_k} + \int_{M} T_{a_1 \dots a_k} \nabla_i Q^{ia_1 \dots a_k} = \int_{\partial M} \nu_i T_{a_1 \dots a_k} Q^{ia_1 \dots a_k}$$
(8)

In other words, $\int_M \langle \nabla T, Q \rangle \operatorname{vol} + \int_M \langle T, \operatorname{div} Q \rangle \operatorname{vol} = \int_{\partial M} \langle \nu \otimes T, Q \rangle \operatorname{vol}_\partial$.

2. For $p \in (1, \infty), \omega \in \mathbb{R}W^{1, p}\Omega^k, \eta \in \mathbb{R}W^{1, p'}\Omega^{k+1}$:

$$\langle\langle d\omega, \eta \rangle\rangle_{\Lambda} = \langle\langle \omega, \delta\eta \rangle\rangle_{\Lambda} + \langle\langle j^*\omega, j^*\iota_{\nu}\eta \rangle\rangle_{\Lambda}$$
(9)

where $\langle \langle j^* \omega, j^* \iota_{\nu} \eta \rangle \rangle_{\Lambda} = \int_{\partial M} \langle j^* \omega, j^* \iota_{\nu} \eta \rangle_{\Lambda} \operatorname{vol}_{\partial}.$

3. For $p \in (1, \infty), \omega \in \mathbb{R}W^{2, p}\Omega^k(M), \eta \in \mathbb{R}W^{1, p'}\Omega^k(M)$:

$$\mathcal{D}(\omega,\eta) = \left\langle \left\langle -\Delta\omega,\eta\right\rangle \right\rangle_{\Lambda} + \left\langle \left\langle j^*\iota_{\nu}d\omega,j^*\eta\right\rangle \right\rangle_{\Lambda} - \left\langle \left\langle j^*\delta\omega,j^*\iota_{\nu}\eta\right\rangle \right\rangle_{\Lambda} \tag{10}$$

where $\mathcal{D}(\omega,\eta) := \langle \langle d\omega, d\eta \rangle \rangle_{\Lambda} + \langle \langle \delta\omega, \delta\eta \rangle \rangle_{\Lambda}$ is called the **Dirichlet integral**.

Proof.

- 1. Let $X^i = T_{a_1...a_k} Q^{ia_1...a_k}$. Then it is just the divergence theorem.
- 2. By approximation, it is enough to prove the smooth case.

$$\int_{\partial M} \langle j^* \omega, j^* \iota_{\nu} \eta \rangle_{\Lambda} \operatorname{vol}_{\partial} = \int_{\partial M} j^* (\omega \wedge \star \eta) = \int_M d (\omega \wedge \star \eta)$$
$$= \int_M d\omega \wedge \star \eta + (-1)^k \int_M \omega \wedge d \star \eta = \langle \langle d\omega, \eta \rangle \rangle_{\Lambda} - \langle \langle \omega, \delta \eta \rangle \rangle_{\Lambda}$$

3. Trivial.

6.3 Boundary conditions and potential theory

Definition 59. We define:

- $\Omega_D^k(M) = \{\omega \in \Omega^k(M) : \mathbf{t}\omega = 0\}$ (Dirichlet boundary condition)
- $\Omega^k_{\text{hom}D}(M) = \{\omega \in \Omega^k(M) : \mathbf{t}\omega = 0, \mathbf{t}\delta\omega = 0\}$ (relative Dirichlet boundary condition)
- $\Omega_N^k(M) = \{\omega \in \Omega^k(M) : \mathbf{n}\omega = 0\}$ (Neumann boundary condition)
- $\Omega^k_{\hom N}(M) = \{\omega \in \Omega^k(M) : \mathbf{n}\omega = 0, \mathbf{n}d\omega = 0\}$ (absolute Neumann boundary condition)
- $\Omega_{0}^{k}(M) = \Omega_{D}^{k}(M) \cap \Omega_{N}^{k}(M)$ (trace-zero boundary condition)
- $\mathcal{H}^k(M) = \{\omega \in \Omega^k(M) : d\omega = 0, \delta\omega = 0\}$ (harmonic fields)
- $\mathcal{H}^k_D(M) = \mathcal{H}^k(M) \cap \Omega^k_D(M)$ (Dirichlet fields)
- $\mathcal{H}^k_N(M) = \mathcal{H}^k(M) \cap \Omega^k_N(M)$ (Neumann fields)

Remark. In writing the function spaces, we omit M when there is no possible confusion. Note that Ω_{00}^k (compact support in \mathring{M}) is different from Ω_0^k .

We can readily extend these definitions to less regular spaces by replacing $\omega \in \Omega^k$ with, for example, $\omega \in B_{3,1}^{\frac{1}{3}}\Omega^k$. Boundary conditions are defined via the trace theorem, and therefore require some regularity.

For example, $B_{3,1}^{\frac{1}{3}}\Omega_N^k$ makes sense, while $L^2\Omega_N^k$ and $H^1\Omega_{\text{hom}N}^k$ do not make sense. Observe that L^2 -cl (Ω_N^k) (closure in the L^2 norm) is just $L^2\Omega^k$ since Ω_{00}^k is dense in $L^2\Omega^k$. Most of these symbols come from [Sch95]. Note that in [Sch95], the difference between L^2X and L^2 -cl(X) (where X is some space) is not made explicit.

Function spaces of type $p = \infty$ are problematic since the smooth members are not dense (see Corollary 48). For instance, $W^{m,\infty}\Omega^k \neq W^{m,\infty}$ -cl (Ω^k) in general.

A special case is when k = 0: $\Omega_N^0(M) = \Omega^0(M) = C^{\infty}(M)$ and $\Omega_{\text{hom}D}^0(M) = \Omega_D^0(M)$. Indeed, the conditions for $\Omega_{\text{hom}D}^0$ and $\Omega_{\text{hom}N}^0$ are what analysts often call "Dirichlet" and "Neumann" boundary conditions respectively.

In fluid dynamics, the condition for Ω_N^1 is also called "impermeable", while Ω_0^1 is "no-slip". On the other hand, $\Omega_{\text{hom}N}^1$ is often given various names, such as "Navier-type", "free boundary" or "Hodge" [MM09a; Mon13; BAE16]. The consensus, however, seems to be that $\Omega_{\text{hom}N}^1$ should be called the "absolute boundary condition" [Wu91; Hsu02; COQ09; Bau17; Ouy17], which explains our choice of naming.

Lemma 60. We have Hodge duality:

- $\star : \Omega^k_D(M) \xrightarrow{\sim} \Omega^{n-k}_N(M), \star : \Omega^k_{\hom D}(M) \xrightarrow{\sim} \Omega^{n-k}_{\hom N}(M), \star : \mathcal{H}^k_D(M) \xrightarrow{\sim} \mathcal{H}^{n-k}_N(M).$
- $\nabla_X(\star\omega) = \star (\nabla_X \omega), \ |\star\omega|_{\Lambda} = |\omega|_{\Lambda} \text{ for } \omega \in \Omega^k, X \in \mathfrak{X}M.$
- For $m \in \mathbb{N}_0, p \in [1, \infty)$, we have $\star : W^{m,p}\Omega_D^k(M) \xrightarrow{\sim} W^{m,p}\Omega_N^{n-k}(M)$, $\star : W^{m,p}\Omega_{\hom D}^k(M) \xrightarrow{\sim} W^{m,p}\Omega_{\hom N}^{n-k}(M)$.

We stress that harmonic fields are **harmonic forms**, i.e. $\Delta \omega = 0$, but the converse is not true in general.

Theorem 61 (4 versions). Let $\omega \in \Omega^k(M)$ be a harmonic form. Then ω is a harmonic field if either

1.
$$\mathbf{t}\omega = 0, \mathbf{n}\omega = 0$$
 (trace-zero)

- 2. $\mathbf{t}\omega = 0, \mathbf{t}\delta\omega = 0$ (relative Dirichlet)
- 3. $\mathbf{n}\omega = 0, \mathbf{n}d\omega = 0$ (absolute Neumann)

4.
$$\mathbf{t}\delta\omega = 0, \mathbf{n}d\omega = 0$$
 (Gaffney)

Proof. Trivial to show $\mathcal{D}(\omega, \omega) = 0$ via integration by parts.

Remark. The four conditions correspond to four different versions of the Poisson equation $\Delta \omega = \eta$ (cf. [Sch95, Section 3.4]), and four ways we can make Δ self-adjoint. In this paper, we will just focus on the absolute Neumann Laplacian and the absolute Neumann heat flow.

Gaffney, one of the earliest figures in the field, showed that the Laplacian corresponding to the 4th boundary condition is self-adjoint and called it the "Neumann problem" (cf. [Gaf54; Con54]). We, however, feel the name "Neumann" should only be used when its Hodge dual is Dirichlet-related (for instance, the Dirichlet potential vs the Neumann potential, to be introduced shortly). Therefore, absent a better rationalization or convention, we see no reason not to honor the name of the mathematician.

In the same vein, some authors consider the 1st condition to be the "Dirichlet boundary condition" (following the intuition from the scalar case, where the trace and the tangential part coincide). By the same reasoning as above, we choose not to do so in this paper.

Blackbox 62 (Dirichlet/Neumann fields). $\mathcal{H}_D^k(M)$ and $\mathcal{H}_N^k(M)$ are finite-dimensional, and therefore complemented in $\mathbb{R}W^{m,p}\Omega^k(M) \ \forall m \in \mathbb{N}_0, p \in [1, \infty].$

Remark. All norms on \mathcal{H}_N^k are equivalent, so we do not need to specify which norm on \mathcal{H}_N^k we are using at any time.

These are very nice spaces, yet they often prevent uniqueness for boundary value problems. We almost always want to work on their orthogonal complements, where Hodge theory truly shines.

Proof. See [Sch95, Theorem 2.2.6].

Corollary 63. $\forall m \in \mathbb{N}_0, p \in [1, \infty]$, there is a continuous projection $P_{m,p} : \mathbb{R}W^{m,p}\Omega^k \twoheadrightarrow \mathcal{H}_N^k$ such that

- it is compatible across different Sobolev spaces, i.e. $P_{m_0,p_0}(\omega) = P_{m_1,p_1}(\omega)$ if $\omega \in W^{m_0,p_0}\Omega^k \cap W^{m_1,p_1}\Omega^k$.
- $1 P_{m,p} : \mathbb{R}W^{m,p}\Omega^k \twoheadrightarrow W^{m,p} (\mathcal{H}_N^k)^{\perp} := \{ \omega \in W^{m,p}\Omega^k : \langle \langle \omega, \phi \rangle \rangle_{\Lambda} = 0 \ \forall \phi \in \mathcal{H}_N^k \} \text{ is also a compatible projection.}$

Proof. Define the continuous linear map $\mathcal{I}_{m,p}: W^{m,p}\Omega^k \to (\mathcal{H}_N^k)^*$ where

$$\mathcal{I}_{m,p}\omega(\phi) = \left\langle \left\langle \omega, \phi \right\rangle \right\rangle_{\Lambda} \forall \phi \in \mathcal{H}_{N}^{k}, \forall \omega \in W^{m,p}\Omega^{k}$$

Then note that $(\phi_1, \phi_2) \mapsto \langle \langle \phi_1, \phi_2 \rangle \rangle_{\Lambda}$ is a positive-definite inner product on \mathcal{H}_N^k , so $\mathcal{I}_{m,p}|_{\mathcal{H}_N^k}$: $\mathcal{H}_N^k \xrightarrow{\sim} (\mathcal{H}_N^k)^*$. We also observe that $\mathcal{I}_{m,p}|_{\mathcal{H}_N^k}$ does not depend on m, p, so we can define the continuous inverse $\mathcal{J} : (\mathcal{H}_N^k)^* \xrightarrow{\sim} \mathcal{H}_N^k$. Then we can just set $P_{m,p} = \mathcal{J} \circ \mathcal{I}_{m,p}$. As we defined $\mathcal{I}_{m,p}$ by $\langle \langle \cdot, \cdot \rangle \rangle_{\Lambda}$, $P_{m,p}$ is compatible across different m, p.

Remark. From now on, for $\omega \in W^{m,p}\Omega^k$, we can decompose $\omega = \mathcal{P}^N \omega + \mathcal{P}^{N\perp} \omega$ where $\mathcal{P}^N \omega = \omega_{\mathcal{H}_N^k} \in \mathcal{H}_N^k$ and $\mathcal{P}^{N\perp} \omega = \omega_{(\mathcal{H}_N^k)^{\perp}} \in W^{m,p} (\mathcal{H}_N^k)^{\perp}$. The decomposition is **natural**, i.e. continuous and compatible across different Sobolev spaces. By Hodge duality, similarly define \mathcal{P}^D and $\mathcal{P}^{D\perp}$. Note $\mathcal{P}^{N\perp}W^{1,p}\Omega_N^k \leq W^{1,p}\Omega_N^k$ and $\mathcal{P}^{N\perp}W^{2,p}\Omega_{\hom M}^k \leq W^{2,p}\Omega_{\hom M}^k$.

Blackbox 64 (Potential theory). For $m \in \mathbb{N}_0$, $p \in (1, \infty)$, we define the injective Neumann Laplacian

 $\Delta_N: \mathcal{P}^{N\perp} W^{m+2,p} \Omega^k_{\hom N} \to \mathcal{P}^{N\perp} W^{m,p} \Omega^k$

as simply Δ under domain restriction. Then $(-\Delta_N)^{-1}$ is called the **Neumann potential**, which is bounded (and actually a Banach isomorphism). Δ_N can also be thought of as an unbounded operator on $\mathcal{P}^{N\perp}W^{m,p}\Omega^k$. By Hodge duality, we also define the Dirichlet counterparts Δ_D and $(-\Delta_D)^{-1}$.

 $\underline{D}g$ from $\underline{D}g$ is a set of a $\underline{D}g$ into the $\underline{D}g$ is the interval of $\underline{D}g$ into $\underline{$

Remark. Because duality is involved, we stay away from $p \in \{1, \infty\}$. Amazingly enough, this is the only elliptic estimate we will need for the rest of the paper. One could say the whole theory is a functional analytic consequence of elliptic regularity (much like how the Nash embedding theorem is a consequence of Schauder estimates, following Günther's approach [Tao16]).

There are many identities which might seem complicated, but are actually trivial to check and helpful for grasping the intuition behind routine operations in Hodge theory, as well as its rich algebraic structure.

Definition. We write d_c as d restricted to $W^{1,p}\Omega_D^k$ and δ_c as δ restricted to $W^{1,p}\Omega_N^k$ for $p \in (1,\infty)$. We will prove in Subsection 7.4 that they are essentially adjoints of δ and d. Let us note that $\Delta_N = -(d\delta_c + \delta_c d)$ on $\mathcal{P}^{N\perp}W^{2,p}\Omega_{\text{hom }N}^k$.

Corollary 65. Let $p \in (1, \infty)$. Some basic properties:

1. $\mathcal{P}^{D\perp}\delta = \delta$ and $\mathcal{P}^{N\perp}d = d$ on $W^{1,p}\Omega^k$. $\mathcal{P}^{N\perp}\delta_c = \delta_c$ on $W^{1,p}\Omega^k_N$ and $\mathcal{P}^{D\perp}d_c = d_c$ on $W^{1,p}\Omega^k_D$. 2.
$$(-\Delta_D)^{-1} \delta = \delta (-\Delta_D)^{-1}$$
 on $\mathcal{P}^{D\perp} W^{1,p} \Omega^k$ and $(-\Delta_N)^{-1} d = d (-\Delta_N)^{-1}$ on $\mathcal{P}^{N\perp} W^{1,p} \Omega^k$.
 $(-\Delta_N)^{-1} \delta_c = \delta_c (-\Delta_N)^{-1}$ on $\mathcal{P}^{N\perp} W^{1,p} \Omega^k_N$ and $(-\Delta_D)^{-1} d_c = d_c (-\Delta_D)^{-1}$ on $\mathcal{P}^{D\perp} W^{1,p} \Omega^k_D$.
3. $\delta = \delta \mathcal{P}^{D\perp} = \delta \mathcal{P}^{N\perp}$ and $d = d\mathcal{P}^{D\perp} = d\mathcal{P}^{N\perp}$ on $W^{1,p} \Omega^k$.
4. $d\delta d = d (\delta d + d\delta) = d (-\Delta)$.
 $\delta d\delta (-\Delta_D)^{-1} = \delta$ on $\mathcal{P}^{D\perp} W^{1,p} \Omega^k$ and $d\delta d (-\Delta_N)^{-1} = d$ on $\mathcal{P}^{N\perp} W^{1,p} \Omega^k$.
5. $d (W^{2,p} \Omega^k_{\hom N}) = d (W^{2,p} \Omega^k_N) \cap W^{1,p} \Omega^{k+1}_N, \delta (W^{2,p} \Omega^k_{\hom D}) = \delta (W^{2,p} \Omega^k_D) \cap W^{1,p} \Omega^{k-1}_D.$

Remark. A good mnemonic device is that Δ_N is formed by d and δ_c , so $(-\Delta_N)^{-1}$ commutes with d and δ_c .

Proof.

- 1. Integration by parts.
- 2. Just check that the expressions are defined by using 1).

The rest is trivial.

6.4 Hodge decomposition

We proceed differently from [Sch95], by using a more algebraic approach in order to derive some results not found in the book. There will be a lot of identities gathered through experience, so their appearances can seem unmotivated at first. Hence, as motivation, let's look at an example of a problem we will need Hodge theory for: is it true that $W^{2,p}\Omega^k_{\text{hom}N}$ is dense in $W^{1,p}\Omega^k_N$ for $p \in (1,\infty)$? The problem is more subtle than it seems, and it is true that the heat flow, once constructed, will imply the answer is yes. But we do not yet have the heat flow, and it turns out this problem is needed for the $W^{1,p}$ -analyticity of the heat flow itself. This foundational approximation of boundary conditions can be done easily once we understand Hodge theory and the myriad connections between different boundary conditions.

Let $\omega \in W^{m,p}\Omega^k$ $(m \in \mathbb{N}_0, p \in (1,\infty))$. In one line, the **Hodge-Morrey decomposition algorithm** is

$$\omega = d_c \delta \left(-\Delta_D \right)^{-1} \mathcal{P}^{D\perp} \omega + \delta_c d \left(-\Delta_N \right)^{-1} \mathcal{P}^{N\perp} \omega + \omega_{\mathcal{H}^k}$$

where $\mathcal{P}^{D\perp}\omega = \omega_{(\mathcal{H}_D^k)^{\perp}}, \mathcal{P}^{N\perp}\omega = \omega_{(\mathcal{H}_N^k)^{\perp}}$ are defined as in Corollary 63, and $\omega_{\mathcal{H}^k}$ is simply defined by subtraction. This is the heart of the matter, and the rest is arguably just bookkeeping.

Note that if $\omega \in W^{1,p}\Omega^k$, $d\omega = d\delta d (-\Delta_N)^{-1} \mathcal{P}^{N\perp} \omega + d\omega_{\mathcal{H}^k} = d\mathcal{P}^{N\perp} \omega + d\omega_{\mathcal{H}^k} = d\omega + d\omega_{\mathcal{H}^k}$. So $d\omega_{\mathcal{H}^k} = 0$ and similarly $\delta\omega_{\mathcal{H}^k} = 0$, justifying the notation. A mild warning is that we do not yet have $W^{1,p}\mathcal{H}^k = W^{1,p}$ -cl (\mathcal{H}^k) .

As we will keep referring to this decomposition, let us define

- $\mathcal{P}_1 = d_c \delta \left(-\Delta_D\right)^{-1} \mathcal{P}^{D\perp}$. Then $\mathcal{P}_1 = d_c \left(-\Delta_D\right)^{-1} \delta \mathcal{P}^{D\perp} = d_c \left(-\Delta_D\right)^{-1} \delta$ on $W^{1,p} \Omega^k$. • $\mathcal{P}_2 = \delta_c d \left(-\Delta_N\right)^{-1} \mathcal{P}^{N\perp}$ Then $\mathcal{P}_2 = \delta_c \left(-\Delta_N\right)^{-1} d$ on $W^{1,p} \Omega^k$.
- $\mathcal{P}_3 = 1 \mathcal{P}_1 \mathcal{P}_2$.

We observe that the decomposition $1 = \mathcal{P}_1 + \mathcal{P}_2 + \mathcal{P}_3$ is natural (continuous and compatible across different Sobolev spaces) since all the operations are natural. In particular, \mathcal{P}_j (for $j \in \{1, 2, 3\}$) is a zeroth-order operator, and if ω is smooth, so is $\mathcal{P}_j \omega$ by Sobolev embedding. Recall that $\mathbf{t}\omega = 0$ implies $\mathbf{t}d\omega = 0$, while $\mathbf{n}\omega = 0$ implies $\mathbf{n}\delta\omega = 0$ (Lemma 57). **Theorem 66** (Smooth decomposition). Some basic properties of \mathcal{P}_j on Ω^k :

- 1. $\mathcal{P}_1 \delta = 0 \text{ on } \Omega^{k+1} \text{ and } \mathcal{P}_2 d = 0 \text{ on } \Omega^{k-1}.$ $\mathcal{P}_1 = \mathcal{P}_2 = 0 \text{ on } \mathcal{H}^k.$ 2. $\mathcal{P}_3 \delta_c = 0 \text{ on } \Omega_N^{k+1} \text{ and } \mathcal{P}_3 d_c = 0 \text{ on } \Omega_D^{k-1}.$
- 3. $\mathcal{P}_{j}\mathcal{P}_{i} = \delta_{ij}\mathcal{P}_{i}$. Therefore $\Omega^{k} = \bigoplus_{j=1}^{3} \mathcal{P}_{j}(\Omega^{k})$.
- 4. $\mathcal{P}_{1}\left(\Omega^{k}\right) = d_{c}\left(\Omega_{D}^{k-1}\right) = d_{c}\mathcal{P}^{D\perp}\left(\Omega_{D}^{k-1}\right) = d_{c}\delta\mathcal{P}^{D\perp}\left(\Omega_{\hom D}^{k}\right) \leq \Omega_{D}^{k}.$ $\mathcal{P}_{2}\left(\Omega^{k}\right) = \delta_{c}\left(\Omega_{N}^{k+1}\right) = \delta_{c}\mathcal{P}^{N\perp}\left(\Omega_{N}^{k+1}\right) = \delta_{c}d\mathcal{P}^{N\perp}\left(\Omega_{\hom N}^{k}\right) \leq \Omega_{N}^{k}.$ $\mathcal{P}_{3}\left(\Omega^{k}\right) = \mathcal{H}^{k}.$
- 5. $\Omega^{k} = \bigoplus_{j=1}^{3} \mathcal{P}_{j}(\Omega^{k})$ is $\langle \langle \cdot, \cdot \rangle \rangle_{\Lambda}$ -orthogonal decomposition.

Proof.

- 1. On Ω^{k+1} , $\mathcal{P}_1 \delta = d_c (-\Delta_D)^{-1} \delta \delta = 0$. Let $\eta \in \mathcal{H}^k$. Then $\mathcal{P}_1 \eta = d_c (-\Delta_D)^{-1} \delta \eta = 0$.
- 2. We just need $\mathcal{P}_2 \delta_c = \delta_c$ on Ω_N^{k+1} . Indeed, $\mathcal{P}_2 \delta_c = \delta_c d (-\Delta_N)^{-1} \delta_c \mathcal{P}^{N\perp} = \delta_c d \delta_c (-\Delta_N)^{-1} \mathcal{P}^{N\perp} = \delta_c \mathcal{P}^{N\perp} = \delta_c$.
- 3. By 1), $\mathcal{P}_2\mathcal{P}_1 = \mathcal{P}_1\mathcal{P}_2 = \mathcal{P}_1\mathcal{P}_3 = \mathcal{P}_2\mathcal{P}_3 = 0$. By 2), $\mathcal{P}_3\mathcal{P}_2 = \mathcal{P}_3\mathcal{P}_1 = 0$. Then observe $\mathcal{P}_2 = (\mathcal{P}_1 + \mathcal{P}_2 + \mathcal{P}_3)\mathcal{P}_2 = \mathcal{P}_2^2$. Similarly, $\mathcal{P}_1^2 = \mathcal{P}_1$ and $\mathcal{P}_3^2 = \mathcal{P}_3$.
- 4. Recall $\mathcal{P}_3(\Omega^k) \leq \mathcal{H}^k$. It becomes an equality since $\mathcal{P}_2(\mathcal{H}^k) = \mathcal{P}_1(\mathcal{H}^k) = 0$. Similarly, obviously $\mathcal{P}_1(\Omega^k) = d_c \delta \mathcal{P}^{D\perp}(\Omega^k_{\text{hom }D}) \leq d_c(\Omega^{k-1}_D)$. It becomes an equality since $\mathcal{P}_2 d = 0$ and $\mathcal{P}_3 d_c = 0$.
- 5. Trivial.

		ı
		I
		I
н		I

To extend this to Sobolev spaces, we will need to use distributions and duality.

Corollary 67 (Sobolev version). Some basic properties of \mathcal{P}_j on $W^{m,p}\Omega^k$ $(m \in \mathbb{N}_0, p \in (1, \infty))$:

- $1. \ \left\langle \left\langle \mathcal{P}_{j}\omega,\phi\right\rangle \right\rangle _{\Lambda}=\left\langle \left\langle \omega,\mathcal{P}_{j}\phi\right\rangle \right\rangle _{\Lambda} \ \forall \omega\in W^{m,p}\Omega^{k}, \forall\phi\in\Omega_{00}^{k}, j=1,2,3$
- 2. $\mathcal{P}_1 \delta = 0 \text{ on } W^{m+1,p} \Omega^{k+1} \text{ and } \mathcal{P}_2 d = 0 \text{ on } W^{m+1,p} \Omega^{k-1}.$
- 3. $\mathcal{P}_1 = \mathcal{P}_2 = 0$ on $W^{m+1,p}\mathcal{H}^k$ and $W^{m,p}$ -cl (\mathcal{H}^k) .
- 4. $\mathcal{P}_{3}\delta_{c} = 0 \text{ on } W^{m+1,p}\Omega_{N}^{k+1} \text{ and } \mathcal{P}_{3}d_{c} = 0 \text{ on } W^{m+1,p}\Omega_{D}^{k-1}.$
- 5. $\mathcal{P}_{j}\mathcal{P}_{i} = \delta_{ij}\mathcal{P}_{i}$. Therefore $W^{m,p}\Omega^{k} = \bigoplus_{j=1}^{3} \mathcal{P}_{j}(W^{m,p}\Omega^{k})$.
- 6. $\mathcal{P}_{3}\left(W^{m,p}\Omega^{k}\right) = W^{m,p}\mathcal{H}^{k}$ for $m \geq 1$ and $W^{m,p}\operatorname{-cl}\left(\mathcal{H}^{k}\right)$ for $m \geq 0$. $\mathcal{P}_{2}\left(W^{m,p}\Omega^{k}\right) = \delta_{c}\left(W^{m+1,p}\Omega^{k+1}_{N}\right) = \delta_{c}d\mathcal{P}^{N\perp}\left(W^{m+2,p}\Omega^{k}_{\hom M}\right)$. $\mathcal{P}_{1}\left(W^{m,p}\Omega^{k}\right) = d_{c}\left(W^{m+1,p}\Omega^{k-1}_{D}\right) = d_{c}\delta\mathcal{P}^{D\perp}\left(W^{m+2,p}\Omega^{k}_{\hom D}\right)$.
- 7. $\mathbf{t}\mathcal{P}_1 = 0$ and $\mathbf{n}\mathcal{P}_2 = 0$ on $W^{m+1,p}\Omega^k$.
- 8. For $p \geq 2$, $W^{m,p}\Omega^k = \bigoplus_{j=1}^3 \mathcal{P}_j(W^{m,p}\Omega^k)$ is $\langle \langle \cdot, \cdot \rangle \rangle_{\Lambda}$ -orthogonal decomposition.

- 9. $W^{m,p}$ -cl $\left(d_c\left(\Omega_D^{k-1}\right)\right) = d_c\left(W^{m+1,p}\Omega_D^{k-1}\right)$ and $W^{m,p}$ -cl $\left(\delta_c\left(\Omega_N^{k+1}\right)\right) = \delta_c\left(W^{m+1,p}\Omega_N^{k+1}\right)$. $W^{m+1,p}$ -cl $\left(\mathcal{H}^k\right) = W^{m+1,p}\mathcal{H}^k$.
- 10. $d = d\left(\mathcal{P}_1 + \mathcal{P}_2 + \mathcal{P}_3\right) = d\mathcal{P}_2 = d\mathcal{P}^{N\perp} = \mathcal{P}^{N\perp} d \text{ on } W^{m+1,p}\Omega^k.$ Consequently, $\mathbf{n}d\mathcal{P}_2\left(W^{m+2,p}\Omega^k_{\hom N}\right) = \mathbf{n}d\left(W^{m+2,p}\Omega^k_{\hom N}\right) = 0, \text{ and } \mathcal{P}_2\left(W^{m+2,p}\Omega^k_{\hom N}\right) \leq W^{m+2,p}\Omega^k_{\hom N}.$ We also have

$$d\left(W^{m+1,p}\Omega^{k-1}\right) = d\mathcal{P}_2\left(W^{m+1,p}\Omega^{k-1}\right) = d\left(W^{m+1,p}\Omega_N^{k-1}\right) = d\mathcal{P}^{N\perp}\left(W^{m+1,p}\Omega_N^{k-1}\right)$$

11. $\delta_c = \mathcal{P}_2 \delta_c$ on $W^{m+1,p} \Omega_N^k$ and

$$\mathcal{P}_2\left(W^{m+1,p}\Omega^k\right) = \delta_c\left(W^{m+2,p}\Omega_N^{k+1}\right) = \delta_c d\mathcal{P}^{N\perp}\left(W^{m+3,p}\Omega_{\hom N}^k\right) = \delta_c \mathcal{P}^{N\perp}\left(W^{m+2,p}\Omega_{\hom N}^{k+1}\right)$$

Remark. Note that L^p -cl (\mathcal{H}^k) $(p \in (1, \infty))$ is defined, while $L^p \mathcal{H}^k$ is not.

Proof.

- 1. Observe $\mathcal{P}_1 \omega \in d_c \left(W^{m+1,p} \Omega_D^{k-1} \right), \mathcal{P}_2 \omega \in \delta_c \left(W^{m+1,p} \Omega_N^{k+1} \right), \mathcal{P}_3 \omega \in W^{m,p}\text{-cl}\left(\mathcal{H}^k\right)$. Simply show $d_c \left(W^{m+1,p} \Omega_D^{k-1} \right) \perp \delta_c \left(\Omega_N^{k+1} \right), W^{m,p}\text{-cl}\left(\mathcal{H}^k\right) \perp d_c \left(\Omega_D^{k-1} \right)$, and so forth via integration by parts.
- 2. $W^{m+1,p}\Omega^{k+1} = W^{m+1,p}$ -cl (Ω^{k+1}) .
- 3. The case $W^{m,p}$ -cl (\mathcal{H}^k) is trivial. For $\omega \in W^{m+1,p}\mathcal{H}^k$, $\langle\langle \mathcal{P}_1\omega, \phi \rangle\rangle_{\Lambda} = \langle\langle \omega, \mathcal{P}_1\phi \rangle\rangle_{\Lambda} = 0 \ \forall \phi \in \Omega_{00}^k$ since $W^{m+1,p}\mathcal{H}^k \perp d_c(\Omega_D^{k-1})$ (integration by parts).
- 4. Let $\omega \in W^{m+1,p}\Omega_N^{k+1}$. Then $\langle \langle \mathcal{P}_3 \delta_c \omega, \phi \rangle \rangle_{\Lambda} = \langle \langle \delta_c \omega, \mathcal{P}_3 \phi \rangle \rangle_{\Lambda} = 0 \ \forall \phi \in \Omega_{00}^k \text{ since } \delta_c \left(W^{m+1,p}\Omega_N^{k+1} \right) \perp \mathcal{H}^k$.

The rest is trivial.

To connect Hodge decomposition to fluid dynamics, we will need the Friedrichs decomposition:

$$\mathcal{P}_3 = \left(\mathcal{P}^N + \mathcal{P}^{N\perp}\right)\mathcal{P}_3 = \mathcal{P}_3^N + \mathcal{P}_3^{\mathrm{ex}}$$

where

P₃^N := P^NP₃ = P^N = P₃P^N (as P^{N⊥}P₁ = P₁ and P^{N⊥}P₂ = P₂)
 P₃^{ex} := P^{N⊥}P₃ = P₃P^{N⊥}

We similarly define $\mathcal{P}_3^D, \mathcal{P}_3^{co}$ via Hodge duality. Note that ex and co stand for "exact" and "coexact" (and we will see why shortly).

Then we define $\mathbb{P} := \mathcal{P}_3^N + \mathcal{P}_2$ as the **Leray projection**. Then $1 = (\mathcal{P}_3^{ex} + \mathcal{P}_1) + (\mathcal{P}_3^N + \mathcal{P}_2) = (\mathcal{P}_3^{ex} + \mathcal{P}_1) + \mathbb{P}$ is called the **Helmholtz decomposition**.

Theorem 68 (Friedrichs decomposition). Basic properties of \mathcal{P}_3^N , $\mathcal{P}_3^{\text{ex}}$ on $W^{m,p}\Omega^k$ $(m \in \mathbb{N}_0, p \in (1, \infty))$:

1. $\mathcal{P}_{3}^{\mathrm{ex}} = d\delta(-\Delta_{N})^{-1}\mathcal{P}_{3}^{\mathrm{ex}} \text{ on } W^{m,p}\Omega^{k}.$ 2. $\mathcal{P}_{3}^{\mathrm{ex}}\left(W^{m,p}\Omega^{k}\right) = W^{m,p}\text{-cl}\left(\mathcal{H}^{k}\right) \cap d\left(W^{m+1,p}\Omega^{k-1}\right).$ 3. $\left(\mathcal{P}_{3}^{\mathrm{ex}} + \mathcal{P}_{1}\right)\left(W^{m,p}\Omega^{k}\right) = d\left(W^{m+1,p}\Omega^{k-1}\right) = d\left(W^{m+1,p}\Omega^{k-1}_{N}\right) = d\mathcal{P}^{N\perp}\left(W^{m+1,p}\Omega^{k-1}_{N}\right).$

- 4. $\mathbb{P}(W^{m,p}\Omega^k) = (\mathcal{P}_3^N + \mathcal{P}_2)(W^{m,p}\Omega^k) = \operatorname{Ker}\left(\delta_c\big|_{W^{m,q}\Omega_N^k}\right)$ when $m \ge 1$ and $W^{m,p}\operatorname{-cl}\left(\operatorname{Ker}\left(\delta_c\big|_{\Omega_N^k}\right)\right)$ when $m \ge 0$.
- 5. $(\mathcal{P}_3 + \mathcal{P}_2) \left(W^{m,p} \Omega^k \right) = \operatorname{Ker} \left(\delta \big|_{W^{m,q} \Omega^k} \right)$ when $m \ge 1$ and $W^{m,p}$ -cl $\left(\operatorname{Ker} \left(\delta \big|_{\Omega^k} \right) \right)$ when $m \ge 0$.
- $\begin{aligned} 6. \ \mathcal{P}^{N\perp}\mathbb{P} &= \mathcal{P}_2 = \mathbb{P}\mathcal{P}^{N\perp} \ on \ W^{m,p}\Omega^k. \\ Therefore \ d\mathbb{P} &= d\mathcal{P}^{N\perp}\mathbb{P} = d\mathcal{P}_2 = d = d\mathcal{P}^{N\perp} = \mathcal{P}^{N\perp}d \ on \ W^{m+1,p}\Omega^k. \end{aligned}$
- 7. $\mathbb{P}\left(W^{m+2,p}\Omega_{\hom N}^{k}\right) \leq \mathcal{P}_{2}\left(W^{m+2,p}\Omega_{\hom N}^{k}\right) \oplus \mathcal{H}_{N}^{k} \leq W^{m+2,p}\Omega_{\hom N}^{k}.$

Proof.

- 1. On Ω^k : $\delta d(-\Delta_N)^{-1}\mathcal{P}_3^{\text{ex}} = \delta(-\Delta_N)^{-1}d\mathcal{P}_3^{\text{ex}} = 0$, so $\mathcal{P}_3^{\text{ex}} = (-\Delta)(-\Delta_N)^{-1}\mathcal{P}_3^{\text{ex}} = d\delta(-\Delta_N)^{-1}\mathcal{P}_3^{\text{ex}}$. Then we are done by density.
- 2. $\mathcal{P}_3^N d = \mathcal{P}_3 \mathcal{P}^N d = 0$ as $\mathcal{P}^{N\perp} d = d$.
- 3. $\mathcal{P}_2 d = 0$ and $\mathcal{P}_3^N d = 0$.

4. We first prove the smooth version. Let ω ∈ Ker (δ_c|_{Ω^k_N}). Then ⟨⟨P₁ω, P₁ω⟩⟩_Λ = ⟨⟨P₁ω, ω⟩⟩_Λ = 0 as Ker (δ_c|_{Ω^k_N}) ⊥ d (Ω^{k-1}), so P₁ω = 0. Similarly, P^{ex}₃ω = 0. Then (P^N₃ + P₂) Ω^k = Ker (δ_c|_{Ω^k_N}).
 For W^{m,p}Ω^k, the case W^{m,p}-cl (Ker (δ_c|_{Ω^k_N})) is trivial. Then assume m ≥ 1 and ω ∈ Ker (δ_c|_{W^{m,q}Ω^k_N}). We can show P₁ω = P^{ex}₃ω = 0 as distributions since Ker (δ_c|_{W^{m,q}Ω^k_N}) ⊥ d(Ω^{k-1}).
 Just note that Ker (δ|_{W^{m,q}Ω^k}) ⊥ d_c(Ω^{k-1}_D) and argue similarly.
 Easy to check that P^{N⊥}P^N₃ = P^N₃P^{N⊥} = 0 and P^{N⊥}P₂ = P₂P^{N⊥} = P₂.

7. Trivial.

-	-	1	
		L	
		L	

Remark. Similar results for $\mathcal{P}_3^D, \mathcal{P}_3^{co}$ hold by Hodge duality. When M has no boundary, $\mathcal{H}^k = \mathcal{H}_D^k = \mathcal{H}_N^k$ so $\mathcal{P}_3 = \mathcal{P}_3^N = \mathcal{P}_3^D$.

A simple consequence of the Hodge-Helmholtz decomposition is that

$$\frac{\operatorname{Ker}\left(\delta_{c}\big|_{\Omega_{N}^{k}}\right)}{\delta_{c}\left(\Omega_{N}^{k+1}\right)} = \frac{\left(\mathcal{P}_{3}^{N} + \mathcal{P}_{2}\right)\left(\Omega^{k}\right)}{\mathcal{P}_{2}\left(\Omega^{k}\right)} = \mathcal{P}_{3}^{N}\left(\Omega^{k}\right) = \frac{\left(\mathcal{P}_{3} + \mathcal{P}_{1}\right)\left(\Omega^{k}\right)}{\left(\mathcal{P}_{3}^{\mathrm{ex}} + \mathcal{P}_{1}\right)\left(\Omega^{k}\right)} = \frac{\operatorname{Ker}\left(d\big|_{\Omega^{k}}\right)}{d\left(\Omega^{k-1}\right)}$$

This can be rewritten as $\mathbb{H}_{a}^{k}(M) = \mathcal{H}_{N}^{k}(M) = \mathbb{H}_{dR}^{k}(M,d)$ (Hodge isomorphism theorem) where $\mathbb{H}_{dR}^{k}(M,d) := \frac{\operatorname{Ker}\left(d|_{\Omega^{k}}\right)}{d(\Omega^{k-1})}$ is called the *k*-th de Rham cohomology group, and $\mathbb{H}_{a}^{k}(M) := \frac{\operatorname{Ker}\left(\delta_{c}|_{\Omega^{k}_{N}}\right)}{\delta_{c}(\Omega^{k+1}_{N})}$ is called the *k*-th absolute de Rham cohomology group. In particular, $\beta^{k}(M) := \dim \mathcal{H}_{N}^{k}(M) = \dim \mathbb{H}_{dR}^{k}(M,d)$ is called the *k*-th relative de Rham cohomology group.

We can also define right inverses (*potentials*) for d, δ, δ_c, d_c (see Subsection 8.1).

In many ways, Hodge theory reduces otherwise complicated boundary value problems into purely algebraic calculations. A standard Hodge-theoretic calculation related to the Euler equation is given later in Subsection 8.2. We can also derive a general form of the Poincare inequality:

Corollary 69 (Poincare-Hodge-Dirac inequality). Let $\omega \in \mathcal{P}^{N\perp}W^{m+1,p}\Omega_N^k$ $(m \in \mathbb{N}_0, p \in (1,\infty))$. Then

$$\|\omega\|_{W^{m+1,p}} \sim \|d\omega\|_{W^{m,p}} + \|\delta_c \omega\|_{W^{m,p}}$$

and we have a bijection

$$\mathcal{P}^{N\perp}W^{m+1,p}\Omega_N^k \xrightarrow{d\oplus\delta_c} d\left(W^{m+1,p}\Omega^k\right) \oplus \delta_c\left(W^{m+1,p}\Omega_N^k\right) = \left(\mathcal{P}_1 + \mathcal{P}_3^{ex}\right)\left(W^{m,p}\Omega^{k+1}\right) \oplus \mathcal{P}_2(W^{m,p}\Omega^{k-1})$$

In particular,
$$\boxed{\left(d\oplus\delta_c\right)^{-1}\left(d\eta,\delta_c\upsilon\right) = \mathcal{P}_2\left(\eta-\upsilon\right) + \upsilon\;\forall\eta,\upsilon\in\mathcal{P}^{N\perp}W^{m+1,p}\Omega_N^k}.$$

Proof. Observe that

- $\mathcal{P}^{N\perp}W^{m+1,p}\Omega_N^k \xrightarrow{d \oplus \delta_c} d\mathcal{P}^{N\perp} \left(W^{m+1,p}\Omega_N^k \right) \oplus \delta_c \mathcal{P}^{N\perp} \left(W^{m+1,p}\Omega_N^k \right)$ is a continuous injection.
- $d\mathcal{P}^{N\perp}\left(W^{m+1,p}\Omega_N^k\right) = d\left(W^{m+1,p}\Omega^k\right) = (\mathcal{P}_1 + \mathcal{P}_3^{\mathrm{ex}})\left(W^{m,p}\Omega^{k+1}\right)$ by Corollary 67.
- $\delta_c \mathcal{P}^{N\perp} \left(W^{m+1,p} \Omega_N^k \right) = \delta_c \left(W^{m+1,p} \Omega_N^k \right) = \mathcal{P}_2(W^{m,p} \Omega^{k-1})$ by Corollary 65 and 67.

By open mapping, we only need to prove $d \oplus \delta_c$ (the **injective Hodge-Dirac operator**) is surjective: let $\eta, \upsilon \in \mathcal{P}^{N\perp}W^{m+1,p}\Omega_N^k$. We want to find $\omega \in \mathcal{P}^{N\perp}W^{m+1,p}\Omega_N^k$ such that $d\omega = d\eta, \delta_c \omega = \delta_c \upsilon$. By the restriction $\delta_c \omega = \delta_c \upsilon$, the freedom is in choosing

$$\vartheta := \omega - \upsilon \in \mathcal{P}^{N\perp} \operatorname{Ker} \left(\delta_c \big|_{W^{m+1,p}\Omega_N^k} \right) = \mathcal{P}^{N\perp} \mathbb{P}(W^{m+1,p}\Omega^k) = \mathcal{P}_2(W^{m+1,p}\Omega^k)$$

such that $d\omega = dv + d\vartheta = d\eta$. In other words, we want ϑ such that $d\vartheta = d(\eta - v)$ and $\mathcal{P}_2 \vartheta = \vartheta$. Then we are done by setting $\vartheta = \mathcal{P}_2(\eta - v)$.

Remark. We note that a less general version of the Poincare inequality was used in [Sch95] to establish the potential estimates in Blackbox 64 as well as Blackbox 62. A more general version [Sch95, Lemma 2.4.10] deals with the case $p \ge 2$. Our version here only requires $p \in (1, \infty)$.

Among other things, the inequality allows the following approximation of boundary conditions, which will play a crucial role for the $W^{1,p}$ -analyticity of the heat flow in Subsection 7.3.

Corollary 70. Let $p \in (1, \infty)$.

1.
$$W^{1,p}\Omega_N^k = d\left(W^{2,p}\Omega_{\mathrm{hom}N}^{k-1}\right) \oplus \operatorname{Ker}\left(\delta_c\big|_{W^{1,p}\Omega_N^k}\right) and \Omega_N^k = d\left(\Omega_{\mathrm{hom}N}^{k-1}\right) \oplus \operatorname{Ker}\left(\delta_c\big|_{\Omega_N^k}\right)$$

2.
$$L^{p}$$
-cl $\left(d\left(\Omega_{\text{hom}N}^{k}\right)\right) = d\left(W^{1,p}\Omega_{N}^{k}\right) = d\left(W^{1,p}\Omega^{k}\right)$

3. $W^{1,p}$ -cl $\left(W^{2,p}\Omega^k_{\mathrm{hom}N}\right) = W^{1,p}\Omega^k_N$.

Proof.

1. Because $\mathbb{P}W^{1,p}\Omega^k \leq W^{1,p}\Omega^k_N$, we conclude $\mathbb{P}W^{1,p}\Omega^k = \mathbb{P}W^{1,p}\Omega^k_N$. Meanwhile, $(\mathcal{P}_1 + \mathcal{P}_3^{ex})W^{1,p}\Omega^k_N = (1-\mathbb{P})W^{1,p}\Omega^k_N \leq W^{1,p}\Omega^k_N$, so $(\mathcal{P}_1 + \mathcal{P}_3^{ex})W^{1,p}\Omega^k_N \leq d(W^{2,p}\Omega^{k-1}_N) \cap W^{1,p}\Omega^k_N = d(W^{2,p}\Omega^{k-1}_{\mathrm{hom}N})$.

2.
$$L^p$$
-cl $\left(\left(\mathcal{P}_1 + \mathcal{P}_3^{\text{ex}}\right)\Omega_N^{k+1}\right) = \left(\mathcal{P}_1 + \mathcal{P}_3^{\text{ex}}\right)L^p$ -cl $\left(\Omega_N^{k+1}\right) = \left(\mathcal{P}_1 + \mathcal{P}_3^{\text{ex}}\right)L^p\Omega^{k+1}$.

3. We are done if $W^{1,p}$ -cl $(\mathcal{P}^{N\perp}\Omega_{\text{hom}N}^k) = \mathcal{P}^{N\perp}W^{1,p}\Omega_N^k$. Recall $\mathcal{P}_2(\Omega_{\text{hom}N}^k) \leq \Omega_{\text{hom}N}^k$ and $\delta_c(\Omega_N^k) = \delta_c \mathcal{P}^{N\perp}(\Omega_{\text{hom}N}^k)$ by Corollary 67, so by the formula of $(d \oplus \delta_c)^{-1}$ from Corollary 69:

$$(d \oplus \delta_c)^{-1} \left[d \left(\Omega_{\hom N}^k \right) \oplus \delta_c \left(\Omega_N^k \right) \right] = (d \oplus \delta_c)^{-1} \left[d \mathcal{P}^{N\perp} \left(\Omega_{\hom N}^k \right) \oplus \delta_c \mathcal{P}^{N\perp} \left(\Omega_{\hom N}^k \right) \right] = \mathcal{P}^{N\perp} \Omega_{\hom N}^k$$

So

$$W^{1,p}\text{-}\mathrm{cl}\left(\mathcal{P}^{N\perp}\Omega_{\mathrm{hom}N}^{k}\right) = \left(d \oplus \delta_{c}\right)^{-1} \left[L^{p}\text{-}\mathrm{cl}\left(d\left(\Omega_{\mathrm{hom}N}^{k}\right)\right) \oplus L^{p}\text{-}\mathrm{cl}\left(\delta_{c}\left(\Omega_{N}^{k}\right)\right)\right]$$
$$= \left(d \oplus \delta_{c}\right)^{-1} \left[d\left(W^{1,p}\Omega^{k}\right) \oplus \delta_{c}\left(W^{1,p}\Omega_{N}^{k}\right)\right]$$
$$= \mathcal{P}^{N\perp}W^{1,p}\Omega_{N}^{k}$$

6.5 An easy mistake

Let $p \in (1, \infty)$, $\omega \in \Omega^k_{\text{hom}N}$. In other words, $\mathbf{n}\omega = 0$ and $\mathbf{n}d\omega = 0$. Using intuition from Euclidean space, it is tempting to conclude $\nabla_{\nu}\omega = 0$, but this is not true in general.

We will not use Penrose notation but work in local coordinates on ∂M , with $\partial_1, ..., \partial_{n-1}$ for directions on ∂M and ∂_n for the direction of $\tilde{\nu}$. Let $\{a_1, ..., a_k\} \subset \{1, ..., n-1\}$. Observe that $\mathbf{n}d\omega = 0$ implies

$$0 = (d\omega)_{na_1\dots a_k} = \partial_n \omega_{a_1\dots a_k} + \sum_i (\pm 1) \partial_{a_i} \omega_{na_1\dots \widehat{a_i}\dots a_k} = \partial_n \omega_{a_1\dots a_k}$$

since $\omega_{na_1...\hat{a_i}...a_k} = 0$ on ∂M . Then recall $\partial_n \omega_{a_1...a_k} = (\nabla_n \omega)_{a_1...a_k} + \Gamma * \omega$ where $\Gamma * \omega$ is schematic for some terms with the Christoffel symbols. As Γ is bounded on M, we conclude $|\mathbf{t}\nabla_{\nu}\omega| \leq |\omega|$ and $|\mathbf{t}\nabla_{\nu}\omega|_{\Lambda} \leq |\omega|_{\Lambda}$ on ∂M . Then

$$\iota_{\nu}d\left(|\omega|^{2}\right) = \nabla_{\nu}\left\langle\omega,\omega\right\rangle = 2\left\langle\nabla_{\nu}\omega,\omega\right\rangle = 2\left\langle\mathbf{t}\nabla_{\nu}\omega,\omega\right\rangle$$

so $|\nabla_{\nu}(|\omega|^2)| \leq |\omega|^2$ on ∂M . This will be important in establishing the L^p -analyticity of the heat flow in Subsection 7.2.

7 Heat flow

As promised, we now obtain a simple construction of the heat flow. We still work on the same setting as in Subsection 6.1.

7.1 L^2 -analyticity

Recall that Δ_N is an unbounded operator on $\mathbb{R}\mathcal{P}^{N\perp}L^2\Omega^k$ and $(-\Delta_N)^{-1}$ is bounded. It is trivial to check that $(-\Delta_N)^{-1}$ is symmetric, therefore self-adjoint. Then Δ_N is also self-adjoint. Then for $\omega \in D(\Delta_N) = \mathcal{P}^{N\perp}H^2\Omega_{\text{hom}N}^k$: $\langle\langle\Delta_N\omega,\omega\rangle\rangle_{\Lambda} = -\mathcal{D}(\omega,\omega) \leq 0$. So Δ_N is dissipative. Therefore, by a complexification argument, $\Delta_N^{\mathbb{C}}$ is acutely sectorial of angle 0 by Theorem 41 and $\left(e^{t\Delta_N^{\mathbb{C}}}\right)_{t\geq 0}$ is a C_0 , analytic semigroup on $\mathbb{C}\mathcal{P}^{N\perp}L^2\Omega^k$. By Blackbox 38, we can derive some basic facts about $e^{t\Delta_N}$:

• For $m \in \mathbb{N}_1$, $D(\Delta_N^m) \leq \mathcal{P}^{N\perp} H^{2m} \Omega^k$ and $\|\Delta_N^m \omega\|_{L^2} \sim \|\omega\|_{H^{2m}} \sim \|\omega\|_{D(\Delta_N^m)} \quad \forall \omega \in D(\Delta_N^m)$ by potential estimates. Recall that $(e^{t\Delta_N})_{t>0}$ on $(D(\Delta_N^m), \|\cdot\|_{H^{2m}})$ is also a C_0 semigroup by Sobolev tower

(Theorem 32).

- For t > 0, by either the spectral theorem (with a complexification step) or semigroup theory, $e^{t\Delta_N}$ is a self-adjoint contraction on $\mathbb{R}\mathcal{P}^{N\perp}L^2\Omega^k$, with image in $D(\Delta_N^\infty) \leq \mathcal{P}^{N\perp}\Omega^k$ by the analyticity of $\left(e^{s\Delta_N^{\mathbb{C}}}\right)_{s\geq 0}$.
- $\forall \omega \in \mathcal{P}^{N\perp}L^2\Omega^k$, $((0,\infty) \to \mathcal{P}^{N\perp}\Omega^k, t \mapsto e^{t\Delta_N}\omega)$ is C^{∞} -continuous by Sobolev tower. Let $m \in \mathbb{N}_1$, then $\partial_t^m \left(e^{t\Delta_N}\omega\right) = \Delta_N^m e^{t\Delta_N}\omega$ and $\left\|e^{t\Delta_N}\omega\right\|_{H^{2m}} \sim \left\|\Delta_N^m e^{t\Delta_N}\omega\right\|_{L^2} \lesssim_{\neg m, \neg t} \frac{m^m}{t^m} \|\omega\|_{L^2}$

Next we define the **non-injective Neumann Laplacian** $\widetilde{\Delta_N}$ as an unbounded operator on $L^2\Omega^k$ with $D(\widetilde{\Delta_N}^m) = D(\Delta_N^m) \oplus \mathcal{H}_N^k$ and $\widetilde{\Delta_N}^m = \Delta_N^m \oplus 0 \ \forall m \in \mathbb{N}_1$. By using either the spectral theorem or checking the definitions manually, $\widetilde{\Delta_N}$ is also a self-adjoint, dissipative operator. Then we also get an analytic heat flow, and $\widetilde{\Delta_N} = \Delta_N \oplus 0_{\mathcal{H}_N^k}$ with $e^{t\widetilde{\Delta_N}} = e^{t\Delta_N} \oplus \mathrm{Id}_{\mathcal{H}_N^k}$.

Recall that for $m \in \mathbb{N}_0, p \in (1, \infty), \omega \in W^{m, p} \Omega^k : \|\omega\|_{W^{m, p}} \sim \|\mathcal{P}^{N\perp}\omega\|_{W^{m, p}} + \|\mathcal{P}^N\omega\|_{\mathcal{H}^k_N}$ where we do not need to specify the norm on \mathcal{H}^k_N as they're all equivalent. Then the previous results for Δ_N can easily be extended to $\widetilde{\Delta_N}$:

- For $m \in \mathbb{N}_1$, $D(\widetilde{\Delta_N}^m) \leq H^{2m}\Omega^k$ and $\forall \omega \in D(\widetilde{\Delta_N}^m)$: $\left\|\widetilde{\Delta_N}^m \omega\right\|_{L^2} \sim \left\|\mathcal{P}^{N\perp}\omega\right\|_{H^{2m}}$ and $\|\omega\|_{D(\widetilde{\Delta_N}^m)} \sim \|\omega\|_{H^{2m}}$. Recall $\left(e^{t\widetilde{\Delta_N}}\right)_{t\geq 0}$ on $D(\widetilde{\Delta_N}^m)$ is also an a C_0 semigroup. (Sobolev tower)
- For t > 0, by either the spectral theorem (with a complexification step) or semigroup theory, $e^{t\Delta_N}$ is a self-adjoint contraction on $\mathbb{R}L^2\Omega^k$, with image in $D\left(\widetilde{\Delta_N}^\infty\right) \leq \Omega^k$.
- $\forall \omega \in L^2 \Omega^k$, $((0,\infty) \to \Omega^k, t \mapsto e^{t\widetilde{\Delta_N}} \omega)$ is C^{∞} -continuous by Sobolev tower. Let $m \in \mathbb{N}_1$, then $\partial_t^m \left(e^{t\widetilde{\Delta_N}} \omega \right) = \widetilde{\Delta_N}^m e^{t\widetilde{\Delta_N}} \omega$ and

$$\left\|e^{t\widetilde{\Delta_N}}\omega\right\|_{H^{2m}} \sim \left\|e^{t\widetilde{\Delta_N}}\mathcal{P}^{N\perp}\omega\right\|_{H^{2m}} + \left\|\mathcal{P}^N\omega\right\|_{\mathcal{H}^k_N} \lesssim_{\neg m,\neg t} \frac{m^m}{t^m} \left\|\mathcal{P}^{N\perp}\omega\right\|_{L^2} + \left\|\mathcal{P}^N\omega\right\|_{\mathcal{H}^k_N}$$

By these estimates, we conclude that $e^{t\widetilde{\Delta_N}} \xrightarrow{t\to\infty} \mathcal{P}^N$ in $\mathcal{L}(L^2\Omega^k)$ (Kodaira projection). In fact, this is how Hodge decomposition was done historically.

7.2 L^p -analyticity

Though we could use the same symbols Δ_N and $\widetilde{\Delta_N}$ for the Neumann Laplacian on L^p , that can create confusion regarding the domains. Let them still refer to the unbounded operators on $\mathbb{R}\mathcal{P}^{N\perp}L^2\Omega^k$ and $\mathbb{R}L^2\Omega^k$ as before. However, $e^{t\Delta_N}$ and $e^{t\widetilde{\Delta_N}}$ are compatible across all L^p spaces (as we will see).

First we note that $\Omega_{00}^k \leq D\left(\widetilde{\Delta_N}^\infty\right)$ so $D\left(\widetilde{\Delta_N}^\infty\right)$ is dense in $L^p \ \forall p \in (1,\infty)$.

Then for L^p -analyticity, we make a Gronwall-type argument (adapted from [IO15, Appendix A] to handle the boundary).

Theorem 71 (Local boundedness). For $p \in (1, \infty)$, $s \in (0, 1)$ and $u \in D\left(\widetilde{\Delta_N}^{\infty}\right)$:

$$\left\|e^{s\Delta_N}u\right\|_p \lesssim_p \|u\|_p$$

Proof. By duality and the density of $D\left(\widetilde{\Delta_N}^{\infty}\right)$ in $L^2 \cap L^p$, WLOG assume $p \ge 2$. By complex interpolation (with a complexification step), WLOG assume p = 4K where K is a large natural number.

Let $U(s) = e^{s\Delta_N}u$, so $\partial_s U = \Delta U$ and

$$\partial_s \left(|U|^{4K} \right) = 2K |U|^{4K-2} \left\langle 2\Delta U, U \right\rangle \stackrel{\text{Bochner}}{=} 2K |U|^{4K-2} \left(\Delta \left(|U|^2 \right) - 2 \left| \nabla U \right|^2 - 2 \left\langle \text{Ric} \left(U \right), U \right\rangle \right)$$

 So

$$\partial_s \int_M |U|^{4K} \le 2K \int_M |U|^{4K-2} \Delta\left(|U|^2\right) + \mathcal{O}_{M,K}\left(\int_M |U|^{4K}\right)$$

Let $f = |U|^2$. As $U \in D\left(\widetilde{\Delta_N}^{\infty}\right) \leq \Omega_{\hom N}^k$, $|\nabla_{\nu} f| \lesssim f$ on ∂M by Subsection 6.5. By Gronwall, we just need $\int_M f^{2K-1} \Delta f \lesssim \int_M f^{2K}$ (pseudo-dissipativity). Simply integrate by parts:

$$\begin{split} \left\langle \left\langle \Delta f, f^{2K-1} \right\rangle \right\rangle &= -\left\langle \left\langle df, d\left(f^{2K-1}\right) \right\rangle \right\rangle + \left\langle \left\langle \nabla_{\nu} f, f^{2K-1} \right\rangle \right\rangle \\ &= -(2K-1) \int_{M} |df|^{2} f^{2K-2} + \mathcal{O}_{M} \left(\int_{\partial M} f^{2K} \right) \\ &= -\frac{2K-1}{K^{2}} \int_{M} |d\left(f^{K}\right)|^{2} + \mathcal{O}_{M} \left(\int_{\partial M} f^{2K} \right) \end{split}$$

Let $F = |f|^K$. So for any $\varepsilon > 0$, we want $C_{\varepsilon} > 0$ such that $\int_{\partial M} F^2 \leq \varepsilon \int_M |dF|^2 + C_{\varepsilon} \int_M F^2$. This follows from Ehrling's inequality, and the fact that $H^1(M) \to L^2(\partial M)$ is compact.

So $(e^{t\widetilde{\Delta_N}})_{t\geq 0}$ can be uniquely extended by density to $L^2\Omega^k + L^p\Omega^k$ and $e^{t\widetilde{\Delta_N}}\Big|_{L^p\Omega^k} \in \mathcal{L}(L^p\Omega^k)$. With a complexification step and an appropriate core chosen by Sobolev embedding, local boundedness on L^p implies L^p -analyticity for all $p \in (1, \infty)$ by Theorem 35.

Let A_p be the generator of $\left(e^{t\widetilde{\Delta_N}}\right)_{t\geq 0}$ on $L^p\Omega^k$. By the definition of generator, $A_p = \widetilde{\Delta_N}$ on $D\left(\widetilde{\Delta_N}^{\infty}\right)$. In our terminology, $A_p^{\mathbb{C}}$ is acutely quasi-sectorial. But we want a more concrete description of $D(A_p)$.

Lemma 72. Let $p \in (1, \infty)$. Then $\left(D(A_p), \|\cdot\|_{D(A_p)}\right) \sim \left(W^{2,p}\Omega^k_{\hom N}, \|\cdot\|_{W^{2,p}}\right)$ and $W^{2,p}\operatorname{-cl}\left(D\left(\widetilde{\Delta_N}^{\infty}\right)\right) = W^{2,p}\Omega^k_{\hom N}$.

Proof. Observe that
$$\forall u \in D\left(\widetilde{\Delta_N}^{\infty}\right) : \mathcal{P}^{N\perp} u \in D(\Delta_N^{\infty})$$
 and
 $\|u\|_{D(A_p)} = \|u\|_p + \left\|\widetilde{\Delta_N} u\right\|_p \sim \left\|\mathcal{P}^N u\right\|_{\mathcal{H}_N^k} + \left\|\mathcal{P}^{N\perp} u\right\|_p + \left\|\Delta_N \mathcal{P}^{N\perp} u\right\|_p \sim \left\|\mathcal{P}^N u\right\|_{\mathcal{H}_N^k} + \left\|\mathcal{P}^{N\perp} u\right\|_{W^{2,p}} \sim \|u\|_{W^{2,p}}$
Then $\|\cdot\|_{D(A_p)} \sim \|\cdot\|_{W^{2,p}}$ since $D\left(\widetilde{\Delta_N}^{\infty}\right)$ is a dense core in $\left(D(A_p), \|\cdot\|_{D(A_p)}\right)$ (see Lemma 34).
This also implies $D(A_p) = W^{2,p}$ -cl $\left(D\left(\widetilde{\Delta_N}^{\infty}\right)\right) = W^{2,p}$ -cl $\left(D\left(\Delta_N^{\infty}\right)\right) \oplus \mathcal{H}_N^k$.
Recall that $D\left(\Delta_N^{\infty}\right) \leq \left(\mathcal{P}^{N\perp} W^{2,p} \Omega_{\hom N}^k, \|\cdot\|_{W^{2,p}}\right) \xrightarrow{\Delta_N} \left(\mathcal{P}^{N\perp} L^p \Omega^k, \|\cdot\|_{L^p}\right)$. Since L^p -cl $\left(\Delta_N D\left(\Delta_N^{\infty}\right)\right) = L^p$ -cl $\left(D\left(\Delta_N^{\infty}\right)\right) = \mathcal{P}^{N\perp} L^p \Omega^k$, we conclude $W^{2,p}$ -cl $\left(D\left(\Delta_N^{\infty}\right)\right) = \left(-\Delta_N\right)^{-1} \left(\mathcal{P}^{N\perp} L^p \Omega^k\right) = \mathcal{P}^{N\perp} W^{2,p} \Omega_{\hom N}^k$

So for $p \in (1, \infty)$, $s \in (0, 1)$ and $u \in L^p \Omega^k$: $\left\| e^{s\widetilde{\Delta_N}} u \right\|_{W^{2,p}} \lesssim \frac{1}{s} \|u\|_p$. That implies $\left\| e^{s\widetilde{\Delta_N}} u \right\|_{W^{1,p}} \lesssim \frac{1}{\sqrt{s}} \|u\|_p$ by complex interpolation (with complexification), using $\left[\mathbb{C}L^p, \mathbb{C}W^{2,p} \right]_{\frac{1}{2}} = \left[\mathbb{C}F_{p,2}^0, \mathbb{C}F_{p,2}^2 \right]_{\frac{1}{2}} = \mathbb{C}F_{p,2}^1$.

Obviously, $D(A_p^{\infty}) = \{\omega \in \Omega_{\hom N}^k : \Delta^m \omega \in W^{2,p} \Omega_{\hom N}^k \ \forall m \in \mathbb{N}_0\} = D(\widetilde{\Delta_N}^{\infty})$ by Sobolev embedding. Additionally, by the density of $D(A_p^{\infty})$ in L^p , we can show by approximation that

$$\left\langle \left\langle e^{t\widetilde{\Delta_{N}}}\omega,\eta\right\rangle \right\rangle _{\Lambda}=\left\langle \left\langle \omega,e^{t\widetilde{\Delta_{N}}}\eta\right\rangle \right\rangle _{\Lambda}\ \forall\omega\in L^{p}\Omega^{k},\eta\in L^{p'}\Omega^{k},p\in\left(1,\infty\right),t\geq0$$

This implies that $e^{t\widetilde{\Delta_N}}\mathcal{P}^{N\perp} = \mathcal{P}^{N\perp}e^{t\widetilde{\Delta_N}}$ on $W^{m,p}\Omega^k \ \forall m \in \mathbb{N}_0, \forall p \in (1,\infty).$

7.3 $W^{1,p}$ -analyticity

We first observe that $W^{1,p}-\operatorname{cl}\left(D\left(\widetilde{\Delta_N}^{\infty}\right)\right) = W^{1,p}-\operatorname{cl}\left(W^{2,p}-\operatorname{cl}\left(D\left(\widetilde{\Delta_N}^{\infty}\right)\right)\right) = W^{1,p}-\operatorname{cl}\left(W^{2,p}\Omega_{\hom N}^k\right) = W^{1,p}\Omega_N^k$ by Corollary 70 and Lemma 72.

Because we will soon be dealing with differential forms of different degrees, define $\Omega(M) = \bigoplus_{k=0}^{n} \Omega^{k}(M)$ as the **graded algebra** of differential forms where multiplication is the wedge product. We simply define $W^{m,p}\Omega(M) = \bigoplus_{k=0}^{n} W^{m,p}\Omega^{k}(M)$, and similarly for $B_{p,q}^{s}, F_{p,q}^{s}$ spaces. Spaces like $\Omega_{D}(M)$, $\Omega_{00}(M)$ or $W^{m,p}\Omega_{\text{hom}N}$ are also defined by direct sums. The dot products $\langle \cdot, \cdot \rangle_{\Lambda}$ and $\langle \langle \cdot, \cdot \rangle \rangle_{\Lambda}$ are also definable as the sum from each degree. Also define $\mathcal{H}(M) = \bigoplus_{k=0}^{n} \mathcal{H}^{k}(M)$.

As an example, $\omega \in L^2\Omega(M)$ and $\eta \in L^2\Omega(M)$ would imply $\omega \wedge \eta \in L^1\Omega(M)$. We also recover integration by parts:

$$\left\langle \left\langle d\omega,\eta\right\rangle \right\rangle _{\Lambda}=\left\langle \left\langle \omega,\delta\eta\right\rangle \right\rangle _{\Lambda}+\left\langle \left\langle j^{*}\omega,j^{*}\iota_{\nu}\eta\right\rangle \right\rangle _{\Lambda}\ \forall\omega\in\mathbb{R}W^{1,p}\Omega\left(M\right),\forall\eta\in\mathbb{R}W^{1,p'}\Omega\left(M\right),p\in\left(1,\infty\right)$$

Then we can set $D\left(\widetilde{\Delta_N}\right) = H^2 \Omega_{\text{hom }N}$ and $D(A_p) = W^{2,p} \Omega_{\text{hom }N}$ $(p \in (1, \infty))$, and previous results such as sectoriality or the Poincare inequality still hold true in this new degree-independent framework, *mutatis mutandis*.

Theorem 73 (Commuting with derivatives I). Let $p \in (1, \infty)$.

1.
$$\delta_c\left(D\left(A_p^{\infty}\right)\right) \leq D\left(A_p^{\infty}\right) \text{ and } d\left(D\left(A_p^{\infty}\right)\right) \leq D\left(A_p^{\infty}\right)$$

2. Let $\omega \in D(A_p) = W^{2,p}\Omega_{\text{hom }N}$ and $\mathfrak{D} \in \{d, \delta_c, \delta_c d, d\delta_c\}$. Then for t > 0: $\mathfrak{D}e^{t\widetilde{\Delta_N}}\omega = e^{t\widetilde{\Delta_N}}\mathfrak{D}\omega$.

Proof.

- 1. Let $\eta \in D(A_p^{\infty})$. Obviously $d\eta \in W^{2,p}\Omega_{\hom N}$, so $d\Delta^m \eta \in W^{2,p}\Omega_{\hom N} \,\forall m \in \mathbb{N}_0$. Observe that $\mathbf{n}\eta = 0$ implies $\mathbf{n}\delta\eta = 0$, and $\mathbf{n}d\eta = 0$ implies $\mathbf{n}\delta d\eta = 0$. But $\mathbf{n}\Delta\eta = 0$ so $\mathbf{n}d\delta\eta = 0$ and we conclude $\delta_c \eta \in W^{2,p}\Omega_{\hom N}$. Similarly, $\delta_c \Delta^m \eta \in W^{2,p}\Omega_{\hom N} \,\forall m \in \mathbb{N}_0$.
- 2. Let t > 0. Note that $\mathfrak{D}e^{t\widetilde{\Delta_N}}\omega \in D(A_p^{\infty})$. Then $\frac{e^{h\widetilde{\Delta_N}}-1}{h}e^{t\widetilde{\Delta_N}}\omega \xrightarrow{C^{\infty}}{h\downarrow 0}\widetilde{\Delta_N}e^{t\widetilde{\Delta_N}}\omega$ so $\partial_t\left(\mathfrak{D}e^{t\widetilde{\Delta_N}}\omega\right) = \mathfrak{D}\widetilde{\Delta_N}e^{t\widetilde{\Delta_N}}\omega = \widetilde{\Delta_N}\mathfrak{D}e^{t\widetilde{\Delta_N}}\omega$. Therefore

$$e^{h\widetilde{\Delta_N}}\mathfrak{D}e^{t\widetilde{\Delta_N}}\omega=\mathfrak{D}e^{(t+h)\widetilde{\Delta_N}}\omega\;\forall t>0,\forall h>0$$

Note that $\mathfrak{D}e^{(t+h)\widetilde{\Delta_N}}\omega \xrightarrow{L^p}{t\downarrow 0} \mathfrak{D}e^{h\widetilde{\Delta_N}}\omega$ since $e^{(t+h)\widetilde{\Delta_N}}\omega \xrightarrow{C^{\infty}}{t\downarrow 0} e^{h\widetilde{\Delta_N}}\omega$. On the other hand, $e^{h\widetilde{\Delta_N}}\mathfrak{D}e^{t\widetilde{\Delta_N}}\omega \xrightarrow{L^p}{t\downarrow 0} e^{h\widetilde{\Delta_N}}\mathfrak{D}\omega$ as $e^{t\widetilde{\Delta_N}}\omega \xrightarrow{W^{2,p}}{t\downarrow 0}\omega$ (why we need $\omega \in D(A_p)$). So $\mathfrak{D}e^{h\widetilde{\Delta_N}}\omega = e^{h\widetilde{\Delta_N}}\mathfrak{D}\omega \ \forall h > 0$. We can extend this via complexification. For $\omega \in \mathbb{C}W^{2,p}\Omega_{\text{hom}N}$, $\mathfrak{D}^{\mathbb{C}}e^{t\widetilde{\Delta_N^{\mathbb{C}}}}\omega = e^{t\widetilde{\Delta_N^{\mathbb{C}}}}\mathfrak{D}^{\mathbb{C}}\omega \ \forall t > 0$.

By L^p -analyticity, $\exists \alpha = \alpha(p) > 0$ such that $\left(e^{z\widetilde{\Delta_N^c}}\right)_{z\in\Sigma_\alpha^+\cup\{0\}}$ is a C_0 , locally bounded, analytic semigroup on $\mathbb{C}L^p\Omega$. Then by the identity theorem, $\mathfrak{D}^{\mathbb{C}}e^{z\widetilde{\Delta_N^c}}\omega = e^{z\widetilde{\Delta_N^c}}\mathfrak{D}^{\mathbb{C}}\omega \ \forall z\in\Sigma_\alpha^+$.

Theorem 74 ($W^{1,p}$ -analyticity). $\left(e^{z\widetilde{\Delta_N^{\mathbb{C}}}}\right)_{z\in\Sigma_{\alpha}^+\cup\{0\}}$ is a C_0 , analytic semigroup on $\mathbb{C}W^{1,p}\Omega_N$.

Proof. Note that $(D(A_p^{\mathbb{C}}), \|\cdot\|_{W^{2,p}})$ is dense in $(\mathbb{C}W^{1,p}\Omega_N, \|\cdot\|_{W^{1,p}})$ by Corollary 70.

So by Lemma 33, we just need to show $\left(e^{z\overline{\Delta_N^{\mathbb{C}}}}\right)_{z\in\Sigma_{\alpha}^+\cup\{0\}} \subset \mathcal{L}\left(\mathbb{C}W^{1,p}\Omega_N\right)$ and is locally bounded. So it is enough to show

$$\left\| e^{z \widetilde{\Delta_{N}^{\mathbb{C}}}} u \right\|_{W^{1,p}} \lesssim \|u\|_{W^{1,p}} \,\,\forall u \in D\left(A_{p}^{\mathbb{C}}\right), \forall z \in \mathbb{D} \cap \Sigma_{\alpha}^{+}$$

Consider $\mathcal{P}^{N\perp}u$, then we only need $\left\|e^{z\widetilde{\Delta_N^{\mathbb{C}}}}u\right\|_{W^{1,p}} \lesssim \|u\|_{W^{1,p}} \quad \forall u \in \mathcal{P}^{N\perp}D\left(A_p^{\mathbb{C}}\right), \forall z \in \mathbb{D} \cap \Sigma_{\alpha}^+.$ Recall $e^{t\widetilde{\Delta_N}}\mathcal{P}^{N\perp} = \mathcal{P}^{N\perp}e^{t\widetilde{\Delta_N}}$ from Subsection 7.2. By the Poincare inequality (Corollary 69):

$$\begin{aligned} \left\| e^{z\widetilde{\Delta_{N}^{\mathbb{C}}}} u \right\|_{W^{1,p}} &\sim & \left\| d^{\mathbb{C}} e^{z\widetilde{\Delta_{N}^{\mathbb{C}}}} u \right\|_{p} + \left\| \delta_{c}^{\mathbb{C}} e^{z\widetilde{\Delta_{N}^{\mathbb{C}}}} u \right\|_{p} = \left\| e^{z\widetilde{\Delta_{N}^{\mathbb{C}}}} d^{\mathbb{C}} u \right\|_{p} + \left\| e^{z\widetilde{\Delta_{N}^{\mathbb{C}}}} \delta_{c}^{\mathbb{C}} u \right\|_{p} \\ &\lesssim & \left\| d^{\mathbb{C}} u \right\|_{p} + \left\| \delta_{c}^{\mathbb{C}} u \right\|_{p} \sim \left\| u \right\|_{W^{1,p}} \ \forall u \in \mathcal{P}^{N\perp} D\left(A_{p}^{\mathbb{C}}\right), \forall z \in \mathbb{D} \cap \Sigma_{\alpha}^{+} \end{aligned}$$

Corollary 75. Let $\omega \in W^{1,p}\Omega_N$ and $\mathfrak{D} \in \{d, \delta_c\}$. Then for t > 0: $\mathfrak{D}e^{t\widetilde{\Delta_N}}\omega = e^{t\widetilde{\Delta_N}}\mathfrak{D}\omega$.

Proof. Same as before, but with
$$e^{t \widetilde{\Delta_N}} \omega \xrightarrow[t \downarrow 0]{W^{1,p}} \omega$$
.

Let $A_{1,p}$ be the generator of $(e^{t\widetilde{\Delta_N}})_{t\geq 0}$ on $W^{1,p}\Omega_N$. Then $A_{1,p}$ and A_p agree on $D(A_p^2)$ by the definition of generators, so $A_{1,p} = \widetilde{\Delta_N}$ on $D(\widetilde{\Delta_N}^{\infty})$. By potential estimates, $\|\cdot\|_{D(A_{1,p})} \sim \|\cdot\|_{W^{3,p}}$ on $D(\widetilde{\Delta_N}^{\infty})$ and therefore on $\|\cdot\|_{W^{3,p}}$ -cl $(D(\widetilde{\Delta_N}^{\infty})) = D(A_{1,p})$. By the same argument as in Lemma 72, $D(A_{1,p}) = (-\Delta_N)^{-1} (\mathcal{P}^{N\perp}W^{1,p}\Omega_N) \oplus \mathcal{H}_N \ge D(\widetilde{\Delta_N}^{\infty})$.

Theorem 76 (Compatibility with Hodge-Helmholtz). Let $m \in \mathbb{N}_0$, $p \in (1, \infty)$, t > 0. By Corollary 75 and Corollary 67:

•
$$e^{t\widetilde{\Delta_N}}d\left(W^{m+1,p}\Omega_N\right) = d\left(e^{t\widetilde{\Delta_N}}W^{m+1,p}\Omega_N\right) \le d\left(\Omega_N\right) = d\left(\Omega\right).$$

• $e^{t\widetilde{\Delta_N}}\delta_c\left(W^{m+1,p}\Omega_N\right) = \delta_c\left(e^{t\widetilde{\Delta_N}}W^{m+1,p}\Omega_N\right) \le \delta_c\left(\Omega_N\right) = \delta_c\left(\Omega_{\text{hom}N}\right).$

As $e^{t\widetilde{\Delta_N}} = 1$ on \mathcal{H}_N , we finally conclude $e^{t\widetilde{\Delta_N}}(\mathcal{P}_3^{ex} + \mathcal{P}_1) = (\mathcal{P}_3^{ex} + \mathcal{P}_1)e^{t\widetilde{\Delta_N}}, e^{t\widetilde{\Delta_N}}\mathcal{P}_2 = \mathcal{P}_2 e^{t\widetilde{\Delta_N}}$ and $e^{t\widetilde{\Delta_N}}\mathcal{P}_3^N = \mathcal{P}_3^N e^{t\widetilde{\Delta_N}} = \mathcal{P}_3^N$ on $W^{m,p}\Omega(M)$. Also, $e^{t\widetilde{\Delta_N}}\mathbb{P} = \mathbb{P}e^{t\widetilde{\Delta_N}}$ on $W^{m,p}\Omega(M)$ where \mathbb{P} is the Leray projection.

By the definition of generators,

$$\widetilde{\Delta_N}\left(\mathcal{P}_3^{\text{ex}} + \mathcal{P}_1\right) = \left(\mathcal{P}_3^{\text{ex}} + \mathcal{P}_1\right)\widetilde{\Delta_N}, \mathcal{P}_3^N\widetilde{\Delta_N} = \widetilde{\Delta_N}\mathcal{P}_3^N = 0, \mathcal{P}_2\widetilde{\Delta_N} = \widetilde{\Delta_N}\mathcal{P}_2 = \widetilde{\Delta_N}\mathbb{P} = \mathbb{P}\widetilde{\Delta_N}$$

on $D(A_p) = W^{2,p}\Omega_{\text{hom}N}$.

We briefly note that in the no-boundary case, we have $\Omega = \Omega_N = \Omega_{\text{hom}N}$, $\widetilde{\Delta_N} = \widetilde{\Delta_D} = \Delta$, $e^{t\Delta}\mathcal{P}_1 = \mathcal{P}_1 e^{t\Delta}$ on $W^{m,p}\Omega$, $\mathcal{P}_1\Delta = \Delta\mathcal{P}_1$ on $W^{2,p}\Omega$.

Remark. The operator $\mathbb{P}\widetilde{\Delta_N}$, with the domain $\mathbb{P}D(A_p)$, is a well-defined unbounded operator on $\mathbb{P}L^p\Omega$. By our arguments, its complexification is acutely sectorial, and $\mathbb{P}\widetilde{\Delta_N} = \widetilde{\Delta_N}, e^{t\mathbb{P}\widetilde{\Delta_N}} = e^{t\widetilde{\Delta_N}}$ on $\mathbb{P}L^p\Omega$. Other authors call it the **Stokes operator** corresponding to the "Navier-type" / "free" boundary condition [Miy80; Gig82; MM09a; MM09b; BAE16].

7.4 Distributions and adjoints

Like the Littlewood-Paley projection, the heat flow does not preserve compact supports in \check{M} . So applying the heat flow to a distribution is not well-defined. This can be a problem as we will need to heat up the nonlinear term in the Euler equation for Onsager's conjecture. For the Littlewood-Paley projection, we fixed it by introducing tempered distributions. That in turn motivates the following definition.

Definition 77. Let $I \subset \mathbb{R}$ be an open interval. Define

- $\mathscr{D}\Omega^{k} = \Omega_{00}^{k} = \operatorname{colim}\{(\Omega_{00}^{k}(K), C^{\infty} \operatorname{topo}) : K \subset \mathring{M} \operatorname{compact}\}$ as the space of **test** *k*-forms with Schwartz's topology (colimit in the category of locally convex TVS).
- $\mathscr{D}'\Omega^k = (\mathscr{D}\Omega^k)^*$ as the space of k-currents (or distributional k-forms), equipped with the weak* topology.
- $\mathscr{D}_N \Omega^k = D\left(\widetilde{\Delta_N}^\infty\right)$ as the space of heated *k*-forms with the Frechet C^∞ topology and $\mathscr{D}'_N \Omega^k = (\mathscr{D}_N \Omega^k)^*$ as the space of heatable *k*-currents (or heatable distributional *k*-forms) with the weak* topology.
- Spacetime test forms: $\mathscr{D}(I, \Omega^k) = C_c^{\infty}(I, \Omega_{00}^k) = \operatorname{colim}\{(C_c^{\infty}(I_1, \Omega_{00}^k(K)), C^{\infty} \operatorname{topo}) : I_1 \times K \subset I \times \mathring{M} \operatorname{compact}\}$ and $\mathscr{D}_N(I, \Omega^k) = \operatorname{colim}\{(C_c^{\infty}(I_1, \mathscr{D}_N \Omega^k), C^{\infty} \operatorname{topo}) : I_1 \subset I \operatorname{compact}\}.$
- Spacetime distributions $\mathscr{D}'\left(I,\Omega^k\right) = \mathscr{D}\left(I,\Omega^k\right)^*, \ \mathscr{D}'_N\left(I,\Omega^k\right) = \mathscr{D}_N\left(I,\Omega^k\right)^*.$

Obviously $\mathscr{D}\Omega^k \xrightarrow{i} \mathscr{D}_N \Omega^k$, so there is an adjoint $\mathscr{D}'_N \Omega^k \xrightarrow{i^*} \mathscr{D}' \Omega^k$. Unfortunately, $\operatorname{Im}(i)$ is not dense so i^* is not injective. Nevertheless, we will make i^* the implicit canonical map from \mathscr{D}'_N to \mathscr{D}' . In particular, $\omega_j \xrightarrow{\mathscr{D}'_N} 0$ implies $\omega_j \xrightarrow{\mathscr{D}'} 0$. Similarly, $\mathscr{D}(I, \Omega^k) \hookrightarrow \mathscr{D}_N(I, \Omega^k)$ and $\mathscr{D}'_N(I, \Omega^k) \to \mathscr{D}'(I, \Omega^k)$. By Sobolev tower (Theorem 32), we observe that $e^{t\widetilde{\Delta_N}}\phi \xrightarrow{C^{\infty}}_{t \mapsto 0} \phi \ \forall \phi \in \mathscr{D}_N \Omega^k$.

For $\Lambda \in \mathscr{D}'_N \Omega^k$, $t \ge 0$ and $\phi \in \mathscr{D}_N \Omega^k$, we define $e^{t\widetilde{\Delta_N}} \Lambda(\phi) = \Lambda\left(e^{t\widetilde{\Delta_N}}\phi\right)$. As Λ is continuous, $\exists m_0, m_1 \in \mathbb{N}_0$ such that $|\Lambda(\phi)| \lesssim \|\phi\|_{C^{m_0}} \lesssim \|\phi\|_{H^{m_1}}$. Then for t > 0 and $\phi \in \mathscr{D}_N \Omega^k$: $\left|e^{t\widetilde{\Delta_N}} \Lambda(\phi)\right| \lesssim \left\|e^{t\widetilde{\Delta_N}}\phi\right\|_{H^{m_1}} \lesssim_{t,m_1} \|\phi\|_{L^2} \implies e^{t\widetilde{\Delta_N}} \Lambda \in L^2 \Omega^k$ and $e^{t\widetilde{\Delta_N}} \Lambda = e^{\frac{t}{2}\widetilde{\Delta_N}} \Lambda \in \mathscr{D}_N \Omega^k$.

Also, for $p \in (1, \infty)$ and $\omega \in L^p \Omega^k$, $e^{t \widetilde{\Delta_N}} \omega$ is the same in $L^p \Omega^k$ and $\mathscr{D}'_N \Omega^k$.

Remark. We note an important limitation: though heated forms are closed under d and δ by Theorem 73, because of integration by parts, we cannot naively define δ or Δ on heatable currents.

Analogous concepts such as \mathscr{D}_D and \mathscr{D}'_D can be defined via Hodge duality for the relative Dirichlet heat flow.

Recall the graded algebra $\Omega(M) = \bigoplus_{k=0}^{n} \Omega^{k}(M)$ from Subsection 7.3. We can easily define $\mathscr{D}\Omega, \mathscr{D}_{N}\Omega$ etc. by direct sums.

For $\Lambda \in \mathscr{D}'_N \Omega$ and $\phi \in \mathscr{D}_N \Omega$, we can define $\delta_c^{\mathscr{D}'_N} \Lambda(\phi) = \Lambda(d\phi)$ and $d^{\mathscr{D}'_N} \Lambda(\phi) = \Lambda(\delta_c \phi)$. These will be consistent with the smooth versions, though we take care to note that

$$\left\langle \left\langle \delta_{c}^{\mathscr{D}'_{N}}\omega,\phi\right\rangle \right\rangle _{\Lambda}=\left\langle \left\langle \omega,d\phi\right\rangle \right\rangle _{\Lambda}=\left\langle \left\langle \delta\omega,\phi\right\rangle \right\rangle _{\Lambda}+\left\langle \left\langle j^{*}\iota_{\nu}\omega,j^{*}\phi\right\rangle \right\rangle _{\Lambda}\ \forall\omega\in W^{1,p}\Omega,\phi\in\mathscr{D}_{N}\Omega,p\in(1,\infty)$$

$$\tag{11}$$

So $\delta_c^{\mathscr{D}'_N}$ agrees with δ_c on $W^{1,p}\Omega_N$ as defined previously. In particular, $\widetilde{\Delta_N}^{\mathscr{D}'_N} = -\left(d^{\mathscr{D}'_N}\delta_c^{\mathscr{D}'_N} + \delta_c^{\mathscr{D}'_N}d^{\mathscr{D}'_N}\right)$ is well-defined on $\mathscr{D}'_N\Omega$.

Note that $\delta^{\mathscr{D}'_N}\Lambda$ cannot be defined since there is $\phi \in \mathscr{D}_N\Omega$ such that $d_c\phi$ is not defined.

For convenience, we also write $\Lambda(\phi) = \langle \langle \Lambda, \phi \rangle \rangle_{\Lambda}$ (abuse of notation) and $\Lambda^{\varepsilon} = e^{\varepsilon \Delta_N} \Lambda$ for $\varepsilon > 0$. Observe that for all $\Lambda \in \mathscr{D}'_N \Omega, \phi \in \mathscr{D}_N \Omega$:

$$\begin{split} \langle \langle d\left(\Lambda^{\varepsilon}\right),\phi\rangle \rangle_{\Lambda} &= \langle \langle\Lambda^{\varepsilon},\delta_{c}\phi\rangle \rangle_{\Lambda} = \langle \langle\Lambda,(\delta_{c}\phi)^{\varepsilon}\rangle \rangle_{\Lambda} = \langle \langle\Lambda,\delta_{c}\left(\phi^{\varepsilon}\right)\rangle \rangle_{\Lambda} = \left\langle \left\langle \left(d^{\mathscr{D}'_{N}}\Lambda\right)^{\varepsilon},\phi\right\rangle \right\rangle_{\Lambda} \end{split}$$

Then $d\left(\Lambda^{\varepsilon}\right) &= \left(d^{\mathscr{D}'_{N}}\Lambda\right)^{\varepsilon}$ and similarly $\delta_{c}\left(\Lambda^{\varepsilon}\right) = \left(\delta_{c}^{\mathscr{D}'_{N}}\Lambda\right)^{\varepsilon} \forall \Lambda \in \mathscr{D}'_{N}\Omega.$

Problem (Consistency problem). For $p \in (1, \infty)$, we have $L^p\Omega \hookrightarrow \mathscr{D}'_N\Omega$ and $L^p\Omega \hookrightarrow \mathscr{D}'\Omega$, and we can identify $\mathscr{D}'_N\Omega \cap L^p\Omega = \mathscr{D}'\Omega \cap L^p\Omega = L^p\Omega$. Let $d^{\mathscr{D}'}$ and $d^{\mathscr{D}'_N}$ be d defined on \mathscr{D}' and \mathscr{D}'_N respectively. For $\omega \in L^p\Omega$, if $d^{\mathscr{D}'}\omega \in \mathscr{D}'\Omega \cap L^p\Omega$, the question is whether we can say $d^{\mathscr{D}'_N}\omega \in \mathscr{D}'_N\Omega \cap L^p\Omega$.

More explicitly, if $\alpha, \omega \in L^p\Omega$ and $\langle \langle \alpha, \phi_0 \rangle \rangle_{\Lambda} = \langle \langle \omega, \delta_c \phi_0 \rangle \rangle_{\Lambda} \quad \forall \phi_0 \in \mathscr{D}\Omega$, can we say $\langle \langle \alpha, \phi \rangle \rangle_{\Lambda} = \langle \langle \omega, \delta_c \phi \rangle \rangle_{\Lambda} \quad \forall \phi \in \mathscr{D}_N \Omega$? The answer is yes, and the method is analogous to some key steps in Subsection 3.3 and Subsection 8.3.

Recall the cutoffs ψ_r from Equation (3).

Lemma 78. Let
$$p \in (1,\infty)$$
 and $\phi \in W^{1,p}\Omega_N^k$. Then $(1-\psi_r)\phi \xrightarrow[r\downarrow 0]{L^p} \phi$ and $\delta_c ((1-\psi_r)\phi) \xrightarrow[r\downarrow 0]{L^p} \delta_c \phi$.

Proof. In Penrose notation,

$$\delta_{c} \left((1 - \psi_{r}) \phi \right)_{a_{1} \dots a_{k-1}} = -\nabla^{i} \left((1 - \psi_{r}) \phi \right)_{ia_{1} \dots a_{k-1}} = \nabla^{i} \psi_{r} \phi_{ia_{1} \dots a_{k-1}} - (1 - \psi_{r}) \nabla^{i} \phi_{ia_{1} \dots a_{k-1}} \\ \implies \delta_{c} \left((1 - \psi_{r}) \phi \right) = \iota_{\nabla \psi_{r}} \phi + (1 - \psi_{r}) \delta_{c} \phi = f_{r} \iota_{\widetilde{\nu}} \phi + (1 - \psi_{r}) \delta_{c} \phi$$

Then we only need $f_r \iota_{\widetilde{\nu}} \phi \xrightarrow[r \downarrow 0]{L^p} 0$. As $\iota_{\widetilde{\nu}} \phi = 0$ on ∂M , by Theorem 53, $\|f_r \iota_{\widetilde{\nu}} \phi\|_{L^p} \lesssim \frac{1}{r} \|\iota_{\widetilde{\nu}} \phi\|_{L^p(M_{< r})} \lesssim \|\iota_{\widetilde{\nu}} \phi\|_{W^{1,p}(M_{< r})} \xrightarrow{r \downarrow 0} 0$.

Then we can conclude $\{\omega \in L^p\Omega(M) : d^{\mathscr{D}'_N}\omega \in L^p\} = \{\omega \in L^p\Omega(M) : d^{\mathscr{D}'}\omega \in L^p\}.$

Recall that for an unbounded operator A, we write (A, D(A)) to specify its domain.

Theorem 79 (Adjoints of d, δ). For $p \in (1, \infty)$, the closure of $(d, \Omega(M))$ as well as $(d, \mathscr{D}_N \Omega(M))$ on $L^p\Omega(M)$ is d_{L^p} where $D(d_{L^p}) = \{\omega \in L^p\Omega(M) : d^{\mathscr{D}'_N}\omega \in L^p\} = \{\omega \in L^p\Omega(M) : d^{\mathscr{D}'}\omega \in L^p\}.$

By Hodge duality, the closure of $(\delta, \Omega(M))$ as well as $(\delta, \mathscr{D}_D \Omega(M))$ on $L^p \Omega(M)$ is δ_{L^p} where $D(\delta_{L^p}) = \{\omega \in L^p \Omega(M) : \delta^{\mathscr{D}'_D} \omega \in L^p\} = \{\omega \in L^p \Omega(M) : \delta^{\mathscr{D}'} \omega \in L^p\}.$

 $Define \boxed{\delta_{c,L^p} = d_{L^{p'}}^*} and \boxed{d_{c,L^p} = \delta_{L^{p'}}^*}. Then \, \delta_{c,L^p} \text{ is the closure of } (\delta, \mathscr{D}_N \Omega(M)) \text{ as well as } (\delta, \mathscr{D}\Omega(M)).$ $Also, \ D(\delta_{c,L^p}) = \{ \omega \in L^p \Omega(M) : \delta_c^{\mathscr{D}'_N} \omega \in L^p \}.$

Similarly, d_{c,L^p} is the closure of $(d, \mathscr{D}_D\Omega(M))$ and $(d, \mathscr{D}\Omega(M))$. Also, $D(d_{c,L^p}) = \{\omega \in L^p\Omega(M) : d_c^{\mathscr{D}'_D}\omega \in L^p\}.$

Proof. Firstly, it is trivial to check d_{L^p} is closed and $(d, \Omega(M))$ is closable $(\omega_j \xrightarrow{L^p} 0 \text{ and } d\omega_j \xrightarrow{L^p} \eta$ would imply $\eta = 0$ since $d\omega_j \xrightarrow{\mathscr{D}'} 0$. Then let $\omega \in D(d_{L^p})$. We can conclude $(\omega^{\varepsilon}, d(\omega^{\varepsilon})) = (\omega^{\varepsilon}, (d^{\mathscr{D}'_N}\omega)^{\varepsilon}) \xrightarrow{L^p \oplus L^p} (\omega, d^{\mathscr{D}'_N}\omega)$. This also gives the closure of $(d, \mathscr{D}_N\Omega(M))$. Then let $G(\delta_{c,L^p}) \leq L^p\Omega \oplus L^p\Omega$ be the graph of δ_{c,L^p} . Similarly for $G(d_{L^{p'}}) \leq L^{p'}\Omega \oplus L^{p'}\Omega$. Write $\mathfrak{J}(x, y) = (-y, x)$. By the definition of adjoints, $\mathfrak{J}(G(\delta_{c,L^p})) = G(d_{L^{p'}})^{\perp}$. Then observe that

$$\begin{split} \left((L^p \oplus L^p) \operatorname{-cl} \left\{ (-\delta_c \phi, \phi) : \phi \in \mathscr{D}\Omega \right\} \right)^{\perp} &= \left\{ (\omega_1, \omega_2) \in L^{p'} \oplus L^{p'} : \langle \langle \omega_1, \delta_c \phi \rangle \rangle_{\Lambda} = \langle \langle \omega_2, \phi \rangle \rangle_{\Lambda} \ \forall \phi \in \mathscr{D}\Omega \right\} \\ &= \left\{ (\omega_1, \omega_2) \in L^{p'} \oplus L^{p'} : \omega_2 = d^{\mathscr{D}'} \omega_1 \right\} = G\left(d_{L^{p'}} \right) \end{split}$$

Then $G(\delta_{c,L^p}) = (L^p \oplus L^p) \operatorname{-cl} \{(\phi, \delta_c \phi) : \phi \in \mathscr{D}\Omega\}$. Do the same for $\phi \in \mathscr{D}_N \Omega$. Finally, by the definition of adjoints:

$$D\left(\delta_{c,L^{p}}\right) = \left\{\omega \in L^{p}\Omega(M) : \left|\langle\langle\omega, d_{L^{p'}}\phi\rangle\rangle_{\Lambda}\right| \lesssim \|\phi\|_{L^{p'}} \ \forall\phi \in D\left(d_{L^{p'}}\right)\right\}$$
$$= \left\{\omega \in L^{p}\Omega(M) : \left|\delta_{c}^{\mathscr{D}'_{N}}\omega\left(\phi^{\varepsilon}\right)\right| = \left|\langle\langle\omega, d\phi^{\varepsilon}\rangle\rangle_{\Lambda}\right| = \left|\langle\langle\omega, (d_{L^{p'}}\phi)^{\varepsilon}\rangle\rangle_{\Lambda}\right| \lesssim \|\phi^{\varepsilon}\|_{L^{p'}} \ \forall\phi \in D\left(d_{L^{p'}}\right), \forall\varepsilon > 0\right\}$$
$$= \left\{\omega \in L^{p}\Omega(M) : \left|\delta_{c}^{\mathscr{D}'_{N}}\omega\left(\phi\right)\right| \lesssim \|\phi\|_{L^{p'}} \ \forall\phi \in \mathscr{D}_{N}\Omega\right\} = \left\{\omega \in L^{p}\Omega(M) : \delta_{c}^{\mathscr{D}'_{N}}\omega \in L^{p}\right\}$$

For the third equal sign, we implicitly used the fact that $e^{t\widetilde{\Delta_N}}\phi \xrightarrow[t\downarrow 0]{C^{\infty}} \phi \ \forall \phi \in \mathscr{D}_N \Omega^k$.

In particular, $W^{1,p}\Omega_N = W^{1,p}\operatorname{-cl}(\mathscr{D}_N\Omega) \leq D(\delta_{c,L^p})$. Similarly, $W^{1,p}\Omega_D \leq D(d_{c,L^p})$. This makes our choice of notation consistent.

Interestingly, a literature search yields a similar result regarding the adjoints of d and δ in [AM04, Proposition 4.3], where the authors used Lie flows on the domain M which is bounded in \mathbb{R}^n , as well as zero extensions to \mathbb{R}^n to characterize $D(d_{L^p})$ and $D(d_{L^p}^*)$. In [MM09a, Equation 2.12], for $\eta \in D(\delta_{L^p})$, the authors defined $\nu \lor \eta \in B_{p,p}^{-\frac{1}{p}}\Omega(\partial M) = \left(B_{p',p'}^{\frac{1}{p}}\Omega(\partial M)\right)^*$ $(p \in (1,\infty))$ by

$$\left\langle \left\langle \nu \vee \eta, j^* \omega \right\rangle \right\rangle_{\Lambda} = \left\langle \left\langle \eta, d \omega \right\rangle \right\rangle_{\Lambda} - \left\langle \left\langle \delta^{\mathscr{D}'} \eta, \omega \right\rangle \right\rangle_{\Lambda} \, \forall \omega \in \Omega\left(M\right)$$

which is reminiscent of Equation (11). Note that $\langle \langle \nu \lor \eta, j^* \omega \rangle \rangle_{\Lambda}$ is abuse of notation (referring to the natural pairing via duality). Recall from Blackbox 47 that $W^{1,p'}\Omega(M) = F^1_{p',2}\Omega(M) \xrightarrow{\text{Trace}} B^{\frac{1}{p}}_{p',p'}\Omega(M) |_{\partial M}$ has a bounded linear section Ext, so it is possible to choose ω such that $\|j^*\omega\|_{B^{\frac{1}{p}}_{p',p'}} \sim \|\omega\|_{W^{1,p'}}$ and therefore $\nu \lor \eta$ is well-defined with

$$\|\nu \vee \eta\|_{B^{-\frac{1}{p}}_{p,p}} \sim \sup_{\substack{\omega \in W^{1,p'}\Omega(M)\\ \|j^*\omega\|_{B^{1/p}_{p',p'}} = 1}} |\langle \langle \nu \vee \eta, j^*\omega \rangle \rangle_{\Lambda}| \lesssim \|\eta\|_{L^p} + \left\|\delta^{\mathscr{D}'}\eta\right\|_{L^p}$$

Of course, for $\eta \in W^{1,p}\Omega$, $\nu \lor \eta = j^* \iota_{\nu} \eta$. We can now show an alternative description of $D(\delta_{c,L^p})$:

Theorem 80. For $p \in (1,\infty)$, $D(\delta_{c,L^p}) = \{\eta \in L^p\Omega(M) : \delta_c^{\mathscr{D}'_N}\eta \in L^p\} = \{\eta \in L^p\Omega(M) : \delta^{\mathscr{D}'}\eta \in L^p \text{ and } \nu \lor \eta = 0\}.$

Proof. Assume $\eta \in L^p\Omega(M)$ and $\delta_c^{\mathscr{D}'_N}\eta \in L^p$. Then $\exists \alpha \in L^p\Omega(M) : \alpha = \delta_c^{\mathscr{D}'_N}\eta = \delta^{\mathscr{D}'}\eta$. By the definition of $\nu \lor \eta$, $\langle \langle \alpha, \omega \rangle \rangle_{\Lambda} + \langle \langle \nu \lor \eta, j^* \omega \rangle \rangle_{\Lambda} = \langle \langle \eta, d\omega \rangle \rangle_{\Lambda} \ \forall \omega \in \Omega(M)$. By the definition of $\delta_c^{\mathscr{D}'_N}\eta$, $\langle \langle \alpha, \omega \rangle \rangle_{\Lambda} = \langle \langle \eta, d\omega \rangle \rangle_{\Lambda} \ \forall \omega \in \mathscr{D}_N\Omega$. So $\langle \langle \nu \lor \eta, j^* \omega \rangle \rangle_{\Lambda} = 0 \ \forall \omega \in \mathscr{D}_N\Omega$. Recall that Ext (the right inverse of Trace) is bounded, so $B_{p',p'}^{\frac{1}{p}}$ -cl $(j^*(\mathscr{D}_N\Omega)) \stackrel{\text{Ext}}{=} j^*(W^{1,p}\text{-cl}(\mathscr{D}_N\Omega)) = j^*(W^{1,p}\Omega_N(M)) \stackrel{\text{Ext}}{=} j^*(W^{1,p}\Omega(M)) = B_{p',p'}^{\frac{1}{p}}\Omega(\partial M)$. Therefore $\nu \lor \eta = 0$. Conversely, now assume $\eta \in L^p\Omega(M)$, $\delta^{\mathscr{D}'}\eta = \alpha \in L^p$ and $\nu \lor \eta = 0$. Then by the definition of

Conversely, now assume $\eta \in L^{p}\Omega(M)$, $\delta^{\mathscr{D}} \eta = \alpha \in L^{p}$ and $\nu \vee \eta = 0$. Then by the definition of $\nu \vee \eta$ for $\eta \in D(\delta_{L^{p}})$, $\langle \langle \alpha, \omega \rangle \rangle_{\Lambda} = \langle \langle \eta, d\omega \rangle \rangle_{\Lambda} \quad \forall \omega \in \Omega(M)$. The formula also holds for $\omega \in \mathscr{D}_{N}\Omega$, and therefore $\delta_{c}^{\mathscr{D}'_{N}} \eta = \alpha \in L^{p}$.

This result agrees with [MM09a, Equation 2.17]. Our characterization of the adjoints of d and δ further highlights how heatable currents are truly natural objects in Hodge theory, independent of the theory of heat flows.

In particular, it is trivial to show $\mathbb{P}L^p\Omega = L^p$ -cl Ker $\left(\delta_c|_{\Omega_N}\right) = \{\eta \in D\left(\delta_{c,L^p}\right) : \delta_c^{\mathscr{D}'_N}\eta = 0\}$ for $p \in (1,\infty)$.

Remark. The name "heatable current" simply refers to the largest topological vector space of differential forms (and hence vector fields) for which the heat equation can be solved (i.e. *heatable*), and once we apply the heat flow a heatable current becomes heated. The name "current" for distributional forms was introduced by Georges de Rham [Rha84], likely with its physical equivalents in mind, and has since become standard in various areas of mathematics such as geometric measure theory and complex manifolds.

It is not easy to search for literature dealing with the subject and how it relates to Hodge theory. They are mentioned in a couple of papers [BB97; Tro09] dealing with "tempered currents" or "temperate currents" on \mathbb{R}^n – differential forms with tempered-distributional coefficients. Yet the notion of "tempered" – not growing too fast – does not make sense on a compact manifold with boundary. Arguably, it is the ability to facilitate the heat flow, or the Littlewood-Paley projection, that most characterizes tempered distributions and makes them ideal for harmonic analysis. For scalar functions, much more is known (cf. [KP14; BBD18; Tan18] and their references). In the same vein, various results from harmonic analysis should also hold for heatable currents.

7.5 Square root

We will not need this for the rest of the paper, but a popular question is the characterization of the square root of the Laplacian.

By the Poincare inequality, $\mathcal{P}^{N\perp}H^1\Omega_N^k$ is a Hilbert space where the H^1 -inner product can be replaced by $(\omega, \eta) \mapsto \mathcal{D}(\omega, \eta)$ (the Dirichlet integral). The space is dense in $\mathcal{P}^{N\perp}L^2\Omega^k$. Define \mathcal{A} as an unbounded operator on $\mathcal{P}^{N\perp}L^2\Omega^k$ where

$$D\left(\mathcal{A}\right) = \left\{ \omega \in \mathcal{P}^{N\perp} H^1 \Omega_N^k : \left| \mathcal{D}(\omega, \eta) \right| \lesssim_\omega \left\| \eta \right\|_2 \, \forall \eta \in \mathcal{P}^{N\perp} H^1 \Omega_N^k \right\}$$

and $\langle\langle \mathcal{A}\omega,\eta\rangle\rangle_{\Lambda} = \mathcal{D}(\omega,\eta) \ \forall \omega \in D(\mathcal{A}), \forall \eta \in \mathcal{P}^{N\perp}H^{1}\Omega_{N}^{k}$. Easy to check that $\langle\langle \mathcal{A}\omega,\eta\rangle\rangle_{\Lambda} = \mathcal{D}\left(\left(-\Delta_{N}\right)^{-1}\mathcal{A}\omega,\eta\right)$ $\forall \eta \in \mathcal{P}^{N\perp}H^{1}\Omega_{N}^{k}$. Therefore $\omega = (-\Delta_{N})^{-1}\mathcal{A}\omega \in \mathcal{P}^{N\perp}H^{2}\Omega_{\text{hom}N}^{k}$ and $\mathcal{A}\omega = (-\Delta_{N})\omega \ \forall \omega \in D(\mathcal{A})$, so $\mathcal{A} \subset -\Delta_{N}$. It is trivial to check $D(-\Delta_{N}) \leq D(\mathcal{A})$, so $\mathcal{A} = -\Delta_{N}$. By Friedrichs extension (cf. [Tay11a, Appendix A, Proposition 8.7], [Tay11b, Section 8, Proposition 2.2]), we conclude that

$$\begin{split} \mathbb{C}\mathcal{P}^{N\perp}H^{1}\Omega_{N}^{k} &= \left[\mathbb{C}\mathcal{P}^{N\perp}L^{2}\Omega^{k}, \left(D\left(\Delta_{N}^{\mathbb{C}}\right), \|\cdot\|_{D\left(\Delta_{N}^{\mathbb{C}}\right)}\right)\right]_{\frac{1}{2}} = \left[\mathbb{C}\mathcal{P}^{N\perp}L^{2}\Omega^{k}, \mathbb{C}\mathcal{P}^{N\perp}H^{2}\Omega_{\hom N}^{k}\right]_{\frac{1}{2}} \\ &= \left(D\left(\sqrt{-\Delta_{N}^{\mathbb{C}}}\right), \|\cdot\|_{D\left(\sqrt{-\Delta_{N}^{\mathbb{C}}}\right)}\right) \end{split}$$

By direct summing, we can extend the result to $\widetilde{\Delta_N}$ to get

$$\mathbb{C}H^{1}\Omega_{N}^{k} = \left[\mathbb{C}L^{2}\Omega^{k}, \mathbb{C}H^{2}\Omega_{\hom N}^{k}\right]_{\frac{1}{2}} = \left(D\left(\sqrt{-\widetilde{\Delta_{N}^{\mathbb{C}}}}\right), \left\|\cdot\right\|_{D\left(\sqrt{-\widetilde{\Delta_{N}^{\mathbb{C}}}}\right)\right)$$

We note that the norms are only defined up to equivalent norms, and $\|\cdot\|_{D(\mathcal{A})}$ is not the same as $\|\cdot\|_{D(\mathcal{A})}^*$ (see Section 2). This difference is not always made explicit in [Tay11a; Tay11b].

7.6 Some trace-zero results

Although we will not need them for the rest of the paper, let us briefly delineate some results regarding the trace-zero Laplacian (cf. Theorem 61) which are similar to those obtained above for the absolute Neumann Laplacian. We begin by retracing our steps from Corollary 63.

Define $\mathcal{H}_0^k(M) = \mathcal{H}_N^k(M) \cap \mathcal{H}_D^k(M)$. Obviously, $\mathcal{H}_0^k(M)$ is finite-dimensional and we can define \mathcal{P}^0 and $\mathcal{P}^{0\perp}$ the same way we did for \mathcal{P}^N and $\mathcal{P}^{N\perp}$ in Corollary 63. When M has no boundary, $\mathcal{P}^{0\perp} = \mathcal{P}^{N\perp}$ and $\mathcal{P}^0 = \mathcal{P}^N = \mathcal{P}_3$.

It is a celebrated theorem, following from the **Aronszajn continuation theorem** [AKS62], that $\mathcal{H}_0^k(M) = 0$ when every connected component of M has nonempty boundary (cf. [Sch95, Theorem 3.4.4]). When that happens, $\mathcal{P}^{0\perp} = 1$ and $\mathcal{P}^0 = 0$.

Blackbox 81 (Potential theory). For $m \in \mathbb{N}_0, p \in (1, \infty)$, we define the injective trace-zero Laplacian

$$\Delta_0: \mathcal{P}^{0\perp} W^{m+2,p} \Omega_0^k \to \mathcal{P}^{0\perp} W^{m,p} \Omega^k$$

as simply Δ under domain restriction. Then $(-\Delta_0)^{-1}$ is called the **trace-zero potential**, which is bounded. Δ_0 can also be thought of as an unbounded operator on $\mathcal{P}^{0\perp}W^{m,p}\Omega_0^k$.

Proof. We only need to prove the theorem on each connected component of M. So WLOG, M is connected. If $\partial M = \emptyset$, we are back to the absolute Neumann case in Blackbox 64. When $\partial M \neq \emptyset$, $\mathcal{P}^{0\perp} = 1$ and we only need to show the trace-zero Poisson problem $(\Delta \omega, \omega|_{\partial M}) = (\eta, 0)$ is uniquely solvable for each $\eta \in W^{m,p}\Omega^k$. This is [Sch95, Theorem 3.4.10].

Consequently, we have a trivial decomposition

$$\omega = \mathcal{P}^{0\perp}\omega + \mathcal{P}^{0}\omega = d\delta \left(-\Delta_0\right)^{-1} \mathcal{P}^{0\perp}\omega + \delta d \left(-\Delta_0\right)^{-1} \mathcal{P}^{0\perp}\omega + \mathcal{P}^{0}\omega$$

for $\omega \in W^{m,p}\Omega^k$, $m \in \mathbb{N}_0$, $p \in (1, \infty)$. This decomposition is not as useful as the Hodge-Morrey decomposition (Subsection 6.4) since the first two terms are not orthogonal. However, it does mean that, when $\mathcal{P}^0 = 0$, every differential form is a sum of exact and coexact forms.

For $\omega \in \mathcal{P}^{0\perp}W^{m+2,p}\Omega_0^k$, $m \in \mathbb{N}_0$, $p \in (1,\infty)$, we also have $\omega = (-\Delta_0)^{-1}(-\Delta_0)\omega = (-\Delta_0)^{-1}(d\delta\omega + \delta d\omega)$, so $\|\omega\|_{W^{m+2,p}} \sim \|\delta\omega\|_{W^{m+1,p}} + \|d\omega\|_{W^{m+1,p}}$. This trick is not enough to get the full Poincare inequality $\|\omega\|_{W^{1,p}} \sim \|\delta\omega\|_p + \|d\omega\|_p$, and therefore [Sch95, Lemma 2.4.10.iv] might be wrong.

As $(-\Delta_0)^{-1}$ is symmetric and bounded on $\mathcal{P}^{0\perp}L^2\Omega^k$, we conclude Δ_0 is a self-adjoint and dissipative operator on $\mathcal{P}^{0\perp}L^2\Omega^k$, with the domain $D(\Delta_0) = \mathcal{P}^{0\perp}H^2\Omega_0^k$. This means $\Delta_0^{\mathbb{C}}$ is acutely sectorial on $\mathbb{C}\mathcal{P}^{0\perp}L^2\Omega^k$.

Next we define the **non-injective trace-zero Laplacian** $\widetilde{\Delta_0}$ as an unbounded operator on $L^2\Omega^k$ with $D\left(\widetilde{\Delta_0}^m\right) = D\left(\Delta_0^m\right) \oplus \mathcal{H}_0^k$ and $\widetilde{\Delta_0}^m = \Delta_0^m \oplus 0 \ \forall m \in \mathbb{N}_1$. Again, $\widetilde{\Delta_0^{\mathbb{C}}}$ is acutely sectorial on $\mathbb{C}L^2\Omega^k$ and $\|\omega\|_{D\left(\widetilde{\Delta_0}^m\right)} \sim \|\omega\|_{H^{2m}} \ \forall \omega \in D\left(\widetilde{\Delta_0}^m\right), \forall m \in \mathbb{N}_1$. In particular, $D\left(\widetilde{\Delta_0}\right) = \mathcal{P}^{0\perp}H^2\Omega_0^k \oplus \mathcal{H}_0^k = H^2\Omega_0^k$.

For L^p -analyticity, observe that on ∂M : $|\nabla_{\nu} (|\omega|^2)| = 2 |\langle \nabla_{\nu} \omega, \omega \rangle| = 0 \leq |\omega|^2 \ \forall \omega \in W^{2,p}\Omega_0^k, \forall p \in (1,\infty).$ So we argue as in Theorem 71, and L^p -analyticity follows.

Remark. The operator $\mathbb{P}\Delta_0$, with the domain $H^2\Omega_0^k \cap \mathbb{P}L^2\Omega^k$, is a well-defined unbounded operator on $\mathbb{P}L^2\Omega^k$. It is called the **Stokes operator** corresponding to the trace-zero/no-slip boundary condition, as discussed in [FK64; GM85; MM08] and others. It lies outside the scope of this paper. For more information, see [HS18] and its references.

8 Results related to the Euler equation

8.1 Hodge-Sobolev spaces

We will have need of negative-order Sobolev spaces when we calculate the pressure in the Euler equation.

Recall the space of heatable currents $\mathscr{D}'_N\Omega$ (defined in Subsection 7.4). Note that $\mathcal{P}^{N\perp}$ is well-defined on $\mathscr{D}'_N\Omega$ by $\langle\langle \mathcal{P}^{N\perp}\Lambda,\phi\rangle\rangle_{\Lambda} = \langle\langle\Lambda,\mathcal{P}^{N\perp}\phi\rangle\rangle \ \forall\Lambda\in\mathscr{D}'_N\Omega, \forall\phi\in\mathscr{D}_N\Omega$. Same for \mathcal{P}^N , and we can uniquely identify $\mathcal{P}^N\Lambda\in\mathcal{H}_N\ \forall\Lambda\in\mathscr{D}'_N\Omega$.

Similarly, $\mathbb{P}(\mathscr{D}_N \Omega) \leq \mathscr{D}_N \Omega$ (use Theorem 76 and Theorem 68), so $\mathbb{P}, 1-\mathbb{P} = (\mathcal{P}_1 + \mathcal{P}_3^{ex})$ and $\mathcal{P}_2 = \mathbb{P} - \mathcal{P}^N$ are well-defined on $\mathscr{D}'_N \Omega$.

For all $p \in (1, \infty)$, define $D_N = d^{\mathscr{D}'_N} + \delta_c^{\mathscr{D}'_N}$ on $\mathcal{P}^{N\perp} \mathscr{D}'_N \Omega$ and $\widetilde{D_N} = d^{\mathscr{D}'_N} + \delta_c^{\mathscr{D}'_N}$ on $\mathscr{D}'_N \Omega$ as the **injective** and **non-injective** (Neumann) Hodge-Dirac operators.

By the Poincare inequality (Corollary 69), it is easy to check that $D_N|_{\mathcal{P}^{N\perp}\mathscr{D}_N\Omega} : \mathcal{P}^{N\perp}\mathscr{D}_N\Omega \to \mathcal{P}^{N\perp}\mathscr{D}_N\Omega$ is bijective. Consequently, so is D_N on $\mathcal{P}^{N\perp}\mathscr{D}'_N\Omega$.

Observe that $\forall m \in \mathbb{N}_0, \forall p \in (1, \infty), \forall \alpha \in \mathcal{P}^{N \perp} W^{m, p} \Omega(M), \exists ! \beta = (D_N)^{-1} \alpha \in \mathcal{P}^{N \perp} W^{m+1, p} \Omega_N$ and

$$\|\beta\|_{W^{m+1,p}} \sim \|\alpha\|_{W^{m,p}} = \|d\beta + \delta_c\beta\|_{W^{m,p}} \sim \|d\beta\|_{W^{m,p}} + \|\delta_c\beta\|_{W^{m,p}}$$
(12)

because $\mathcal{P}^{N\perp}W^{m,p}\Omega = d\left(W^{m+1,p}\Omega\right) \oplus \delta_c\left(W^{m+1,p}\Omega_N\right)$ is a direct sum of closed subspaces (corresponding to $\mathcal{P}_1 + \mathcal{P}_3^{\text{ex}}$ and \mathcal{P}_2).

Note that we do not have $d^{\mathscr{D}'_N}D_N = D_N d^{\mathscr{D}'_N}$, but $d^{\mathscr{D}'_N}D_N^2 = D_N^2 d^{\mathscr{D}'_N} = -\Delta_N^{\mathscr{D}'_N} d^{\mathscr{D}'_N}$ is true.

Definition 82. For $m \in \mathbb{Z}, p \in (1, \infty)$, let $W^{m,p}(D_N) := (D_N)^{-m} (\mathcal{P}^{N\perp} L^p \Omega) = \{ \alpha \in \mathcal{P}^{N\perp} \mathscr{D}'_N \Omega : (D_N)^m \alpha \in L^p \Omega \}$ and $W^{m,p} (\widetilde{D_N}) := W^{m,p} (D_N) \oplus \mathcal{H}_N$. They are Banach spaces under the norms $\|\alpha\|_{W^{m,p}(D_N)} := \|(D_N)^m \alpha\|_{L^p \Omega}$ and $\|\beta\|_{W^{m,p}(\widetilde{D_N})} := \|\mathcal{P}^{N\perp}\beta\|_{W^{m,p}(D_N)} + \|\mathcal{P}^N\beta\|_{\mathcal{H}_N}$.

In a sense, these are comparable to homogeneous and inhomogeneous Bessel potential spaces. We can extend the definitions to fractional powers, but that is outside the scope of this paper.

It is trivial to check that $\|\alpha\|_{W^{m,p}(\widetilde{D_N})} \sim \|\alpha\|_{W^{m,p}\Omega} \ \forall \alpha \in \mathscr{D}_N\Omega, \forall m \in \mathbb{N}_0, \forall p \in (1,\infty).$

Theorem 83. Some basic properties of $W^{m,p}\left(\widetilde{D_N}\right)$:

- 1. $\mathscr{D}_N \Omega$ is dense in $W^{m,p}\left(\widetilde{D_N}\right) \, \forall m \in \mathbb{Z}, \forall p \in (1,\infty).$
- 2. $W^{m,p}\left(\widetilde{D_N}\right) = W^{m,p} \operatorname{-cl}\left(\mathscr{D}_N\Omega\right) \ \forall m \in \mathbb{N}_0, \forall p \in (1,\infty).$
- 3. $\left\| d^{\mathscr{D}'_{N}} \beta \right\|_{W^{m,p}(\widetilde{D_{N}})} + \left\| \delta_{c}^{\mathscr{D}'_{N}} \beta \right\|_{W^{m,p}(\widetilde{D_{N}})} \lesssim \left\| \beta \right\|_{W^{m+1,p}(\widetilde{D_{N}})} \, \forall \beta \in W^{m+1,p}\left(\widetilde{D_{N}}\right), \forall m \in \mathbb{Z}, \forall p \in (1,\infty)$ $Then \ \mathcal{P}_{2} = \delta_{c}^{\mathscr{D}'_{N}} d^{\mathscr{D}'_{N}} \left(-\Delta_{N}^{\mathscr{D}'_{N}} \right)^{-1} \mathcal{P}^{N\perp} = \delta_{c}^{\mathscr{D}'_{N}} \left(-\Delta_{N}^{\mathscr{D}'_{N}} \right)^{-1} d^{\mathscr{D}'_{N}} and \ \mathbb{P} = \mathcal{P}_{2} + \mathcal{P}^{N} are of order 0 on$ $W^{m,p}\left(\widetilde{D_{N}}\right).$
- $4. \left(W^{m,p}\left(\widetilde{D_N}\right) \right)^* = W^{-m,p'}\left(\widetilde{D_N}\right) \forall m \in \mathbb{Z}, \forall p \in (1,\infty) \text{ via the pairing } \langle \alpha, \phi \rangle_{W^{-m,p}\left(\widetilde{D_N}\right),W^{m,p'}\left(\widetilde{D_N}\right)} = \left\langle \left\langle D_N^{-m}\mathcal{P}^{N\perp}\alpha, D_N^m\mathcal{P}^{N\perp}\phi \right\rangle \right\rangle_{\Lambda} + \left\langle \left\langle \mathcal{P}^N\alpha, \mathcal{P}^N\phi \right\rangle \right\rangle_{\Lambda}$

Proof.

- 1. Because $D_N^m \left(\mathcal{P}^{N\perp} \mathscr{D}_N \Omega \right) = \mathcal{P}^{N\perp} \mathscr{D}_N \Omega$ is dense in $\mathcal{P}^{N\perp} L^p \Omega$.
- 2. We only need $W^{m,p}(D_N) = \mathcal{P}^{N\perp} W^{m,p}\left(\widetilde{D_N}\right) \leq W^{m,p}\text{-cl}\left(\mathcal{P}^{N\perp}\mathscr{D}_N\Omega\right)$. Let $\alpha \in \mathcal{P}^{N\perp} W^{m,p}\left(\widetilde{D_N}\right)$ and $\alpha^{\varepsilon} = e^{\varepsilon \widetilde{\Delta_N}} \alpha$ as usual. Then $D_N^m(\alpha^{\varepsilon}) = (D_N^m \alpha)^{\varepsilon} \xrightarrow[\varepsilon \downarrow 0]{} D_N^m \alpha$. So $D_N^{-m} D_N^m(\alpha^{\varepsilon}) = \alpha^{\varepsilon} \xrightarrow[\varepsilon \downarrow 0]{} \alpha$ by Equation (12).
- 3. Let $D_N^{m+1} \mathcal{P}^{N\perp} \beta \in L^p$. Then $D_N^m \mathcal{P}^{N\perp} \beta \in \mathcal{P}^{N\perp} W^{1,p} \Omega_N$ by Equation (12).

When $m = 2k \ (k \in \mathbb{Z})$: $\left\| dD_N^{2k} \mathcal{P}^{N\perp} \beta \right\|_{L^p} + \left\| \delta_c D_N^{2k} \mathcal{P}^{N\perp} \beta \right\|_{L^p} \sim \left\| dD_N^{2k} \mathcal{P}^{N\perp} \beta + \delta_c D_N^{2k} \mathcal{P}^{N\perp} \beta \right\|_{L^p} = \left\| D_N^{2k+1} \mathcal{P}^{N\perp} \beta \right\|_{L^p}.$

When
$$m = 2k + 1$$
 $(k \in \mathbb{Z})$: $D_N^{2k} \mathcal{P}^{N\perp} \beta \in \mathcal{P}^{N\perp} W^{2,p} \Omega_{\hom N}$ and

$$\begin{split} \left\| D_N dD_N^{2k} \mathcal{P}^{N\perp} \beta \right\|_{L^p} + \left\| D_N \delta_c D_N^{2k} \mathcal{P}^{N\perp} \beta \right\|_{L^p} &= \left\| \delta_c dD_N^{2k} \mathcal{P}^{N\perp} \beta \right\|_{L^p} + \left\| d\delta_c D_N^{2k} \mathcal{P}^{N\perp} \beta \right\|_{L^p} \\ &\sim \left\| \delta_c dD_N^{2k} \mathcal{P}^{N\perp} \beta + d\delta_c D_N^{2k} \mathcal{P}^{N\perp} \beta \right\|_{L^p} = \left\| D_N^{2k+2} \mathcal{P}^{N\perp} \beta \right\|_{L^p} \end{split}$$

4. Simply observe that $(W^{m,p}(D_N))^* = W^{-m,p'}(D_N)$ via the isomorphisms $W^{m,p}(D_N) \xrightarrow{D_N^m} \mathcal{P}^{N\perp}L^p\Omega$ and $W^{-m,p'}(D_N) \xrightarrow{D_N^{-m}} \mathcal{P}^{N\perp}L^{p'}\Omega$.

Remark. We briefly note that D_N with the domain $\mathcal{P}^{N\perp}H^1\Omega_N$ is self-adjoint on $\mathcal{P}^{N\perp}L^2\Omega$ and its complexification is therefore "bisectorial". For more on this, see [McI86; McI10; MM18].

Corollary 84. Assume $U \in \mathbb{P}L^2\mathfrak{X}$. Define $\operatorname{div}(U \otimes U) \in \mathscr{D}'_N\mathfrak{X}$ by $\langle \langle \operatorname{div}(U \otimes U), X \rangle \rangle_{\Lambda} := - \langle \langle U \otimes U, \nabla X \rangle \rangle$ $\forall X \in \mathscr{D}_N\mathfrak{X}$.

If
$$p \in (1,\infty)$$
 and $U \otimes U \in L^p \Gamma(TM \otimes TM)$, then $\|\operatorname{div}(U \otimes U)^{\flat}\|_{W^{-1,p}(\widetilde{D_N})} \lesssim \|U \otimes U\|_{L^p}$.

Proof. For
$$\eta \in \Omega(M)$$
, write η_k for the part of η in Ω^k . Let $\phi \in \mathscr{D}_N \Omega$, then
 $\left|\left\langle \left\langle D_N^{-1} \mathcal{P}^{N\perp} \operatorname{div}(U \otimes U)^{\flat}, \phi \right\rangle \right\rangle_{\Lambda} \right| = \left|\left\langle \left\langle U \otimes U, \nabla \left(D_N^{-1} \mathcal{P}^{N\perp} \phi \right)_1^{\sharp} \right\rangle \right\rangle \right| \lesssim \|U \otimes U\|_{L^p} \|\phi\|_{L^{p'}}$

This implies $\operatorname{div}(U \otimes U)^{\flat} \in W^{-1,p}\left(\widetilde{D_N}\right)$. Then observe $\left|\left\langle\left\langle \operatorname{div}(U \otimes U)^{\flat}, \phi\right\rangle\right\rangle_{\Lambda}\right| = \left|\left\langle\left\langle U \otimes U, \nabla\left(\phi\right)_1^{\sharp}\right\rangle\right\rangle\right| \lesssim \|U \otimes U\|_{L^p} \|\phi\|_{W^{1,p'}(\widetilde{D_N})}$.

8.2 Calculating the pressure

In this subsection, we assume that $\partial_t \mathcal{V} + \operatorname{div}(\mathcal{V} \otimes \mathcal{V}) + \operatorname{grad} \mathfrak{p} \xrightarrow{\mathscr{D}'_N(I,\mathfrak{X})} 0, \mathcal{V} \in L^2_{\operatorname{loc}}(I, \mathbb{P}L^2\mathfrak{X}), \mathfrak{p} \in L^1_{\operatorname{loc}}(I \times M).$ This is true, for instance, in the case of Onsager's conjecture (see Subsection 3.3 and Subsection 3.4).

We first note that $\mathcal{H}_N^0 = \mathcal{H}^0 = \{\text{locally constant functions}\}$. Then we can show \mathcal{V} uniquely determines \mathfrak{p} by a formula, up to a difference in \mathcal{H}_N^0 ($d\mathfrak{p}$ is always unique). It is no loss of generality to set $\mathfrak{p} = \mathcal{P}^{N\perp}\mathfrak{p}$ (implying $\int_M \mathfrak{p} = 0$).

1. Assume $\mathcal{V} \otimes \mathcal{V} \in L^q_t W^{m+1,p} \Gamma(TM \otimes TM)$ for some $m \in \mathbb{N}_0, p \in (1,\infty), q \in [1,\infty]$.

Let $\omega = \operatorname{div}(\mathcal{V} \otimes \mathcal{V})^{\flat}$. Then $d^{\mathscr{D}'_N} \mathfrak{p} \stackrel{\mathscr{D}'_N(I,\mathfrak{X})}{=} (\mathbb{P}-1) \omega \in L^q_t W^{m,p} \Omega^1$. By the Poincare inequality (Corollary 69), there is a unique $\mathfrak{f} \in L^q_t \mathcal{P}^{N\perp} W^{m+1,p} \Omega^0$ such that $d\mathfrak{f} = (\mathbb{P}-1) \omega \stackrel{\mathscr{D}'_N(I,\mathfrak{X})}{=} d^{\mathscr{D}'_N} \mathfrak{p}$. An explicit formula is $\mathfrak{f} = -R_d \omega$ where $R_d := \mathcal{P}^{N\perp} \delta (-\Delta_D)^{-1} \mathcal{P}^{D\perp} + \mathcal{P}^{N\perp} \delta (-\Delta_N)^{-1} \mathcal{P}^{\mathfrak{s}}_3$ is the potential for d. We aim to show $\mathfrak{f} = \mathfrak{p}$. Let $\psi \in C^{\infty}(I, \mathscr{Q}, \Omega^0)$. Then because $\Omega^0 = \mathcal{P}_{\mathfrak{s}}(\Omega^0) \oplus \mathcal{P}^N(\Omega^0)$, we conclude

We aim to show $\mathfrak{f} = \mathfrak{p}$. Let $\psi \in C_c^{\infty}(I, \mathscr{D}_N \Omega^0)$. Then because $\Omega^0 = \mathcal{P}_2(\Omega^0) \oplus \mathcal{P}_3^N(\Omega^0)$, we conclude $\mathcal{P}^{N\perp}\psi = \delta_c \phi$ where $\phi := d(-\Delta_N)^{-1} \mathcal{P}^{N\perp}\psi \in C_c^{\infty}(I, \mathscr{D}_N \Omega^1)$ and

$$\int_{I} \langle \langle \mathfrak{f}, \psi \rangle \rangle_{\Lambda} = \int_{I} \left\langle \left\langle \mathfrak{f}, \mathcal{P}^{N\perp}\psi \right\rangle \right\rangle_{\Lambda} = \int_{I} \left\langle \left\langle \mathfrak{f}, \delta_{c}\phi \right\rangle \right\rangle_{\Lambda} = \int_{I} \left\langle \left\langle d\mathfrak{f}, \phi \right\rangle \right\rangle_{\Lambda} = \int_{I} \left\langle \left\langle d\mathfrak{f}, \psi \right\rangle \right\rangle_{\Lambda} = \int_{I} \left\langle \left\langle \mathfrak{g}, \psi \right\rangle \right\rangle_{\Lambda} = \int_{I} \left\langle \left\langle \mathfrak{g}$$

Therefore $\mathfrak{p} = \mathfrak{f}$ and $\|\mathfrak{p}\|_{L^q_t W^{m+1,p}} \lesssim \|\omega\|_{L^q_t W^{m,p}} \lesssim \|\mathcal{V} \otimes \mathcal{V}\|_{L^q_t W^{m+1,p}}$

2. Assume $\mathcal{V} \otimes \mathcal{V} \in L^q_t L^p \Gamma(TM \otimes TM)$ for some $p \in (1, \infty), q \in [1, \infty]$.

Let $\omega = \operatorname{div}(\mathcal{V} \otimes \mathcal{V})^{\flat}$. Then $d^{\mathscr{D}'_{N}} \mathfrak{p} \stackrel{\mathscr{D}'_{N}(I,\mathfrak{X})}{=} (\mathbb{P}-1) \omega \in L^{q}_{t} W^{-1,p}\left(\widetilde{D_{N}}\right)$ by Corollary 84 and Theorem 83. Then $-\delta^{\mathscr{D}'_{N}}_{c} d^{\mathscr{D}'_{N}} \mathfrak{p} \stackrel{\mathscr{D}'_{N}(I,\mathfrak{X})}{=} \delta^{\mathscr{D}'_{N}}_{c} (1-\mathbb{P}) \omega = \delta^{\mathscr{D}'_{N}}_{c} \omega \in L^{q}_{t} W^{-2,p}\left(\widetilde{D_{N}}\right)$ and $\mathfrak{p} = -D^{-2}_{N} \delta^{\mathscr{D}'_{N}}_{c} \omega$, so $\|\mathfrak{p}\|_{L^{q}_{t}L^{p}} \lesssim \|\delta^{\mathscr{D}'_{N}}_{c} \omega\|_{L^{q}_{t} W^{-2,p}(\widetilde{D_{N}})} \lesssim \|\omega\|_{L^{q}_{t} W^{-1,p}(\widetilde{D_{N}})} \lesssim \|\mathcal{V} \otimes \mathcal{V}\|_{L^{q}_{t}L^{p}}.$

Remark. It is also possible to define $R_{\delta_c} := d^{\mathscr{D}'_N} \left(-\Delta_N^{\mathscr{D}'_N} \right)^{-1} \mathcal{P}^{N\perp}$ on $\mathscr{D}'_N \Omega$ and have $R_d = \left(D_N^{-1} - R_{\delta_c} \right) \mathcal{P}^{N\perp}$ on $\mathscr{D}'_N \Omega$. This would then imply $\|R_d \alpha\|_{W^{m+1,p}(\widetilde{D_N})} \lesssim \|\alpha\|_{W^{m,p}(\widetilde{D_N})} \quad \forall \alpha \in W^{m,p}(\widetilde{D_N}), \forall m \in \mathbb{Z}, \forall p \in (1,\infty).$

8.3 On an interpolation identity

Let $p \in (1, \infty)$. We are faced with the difficulty of finding a good interpolation characterization for $B_{p,1}^{\frac{1}{p}}\Omega_N$. We do have $B_{p,1}^{\frac{1}{p}}\Omega = (L^p\Omega, W^{1,p}\Omega)_{\frac{1}{p},1}$ (complexification, then projection onto the real part), but our heat flow is not analytic on $\mathbb{C}W^{1,p}\Omega$. The hope is that $B_{p,1}^{\frac{1}{p}}\Omega_N = (L^p\Omega, W^{1,p}\Omega_N)_{\frac{1}{p},1}$, and our first guess is to try to find some kind of projection. Indeed, the Leray projection yields

$$\mathbb{P}B_{p,1}^{\frac{1}{p}}\Omega = \left(\mathbb{P}L^{p}\Omega, \mathbb{P}W^{1,p}\Omega\right)_{\frac{1}{p},1}$$
(13)

and the heat flow is well-behaved on $\mathbb{P}W^{1,p}\Omega = \mathbb{P}W^{1,p}\Omega_N$ (Theorem 68, Theorem 76). By interpolation, \mathbb{P} is $B_{p,1}^{\frac{1}{p}}$ -continuous, so $\mathbf{n}\mathbb{P}: B_{p,1}^{\frac{1}{p}}\Omega \to L^p\Omega|_{\partial M}$ is continuous and $\mathbb{P}B_{p,1}^{\frac{1}{p}}\Omega = \mathbb{P}B_{p,1}^{\frac{1}{p}}\Omega_N$. This is enough to get all the Besov estimates we will need for Onsager's conjecture.

Additionally, it is true that the heat semigroup is also C_0 and analytic on $\mathbb{CP}B_{p,1}^{\frac{1}{p}}\Omega_N$ by Yosida's halfplane criterion (Theorem 39). Unlike the L^p -analyticity case, here we already have analyticity on the 2 endpoints, so the criterion simply follows by interpolation. Alternatively, observe that there exists C > 0such that $\sup_{t>0} \left\| t\left(\widetilde{\Delta_N^{\mathbb{C}}} - C\right) e^{t\left(\widetilde{\Delta_N^{\mathbb{C}}} - C\right)} \right\|_{\mathcal{L}(V)} < \infty \text{ for } V \in \{\mathbb{CP}L^p\Omega, \mathbb{CP}W^{1,p}\Omega_N\}.$ Therefore it also holds for $V = \mathbb{CPB}_{p,1}^{\frac{1}{p}} \Omega_N$ by interpolation, and that is another criterion for analyticity ([Eng00, Section II, Theorem 4.6.c]).

Unfortunately, this does not tell us about the relationship between $(L^p\Omega, W^{1,p}\Omega_N)_{1,1}$ and $B_{p,1}^{\frac{1}{p}}\Omega_N$. Obviously $(L^p\Omega, W^{1,p}\Omega_N)_{\frac{1}{2}} \hookrightarrow B_{p,1}^{\frac{1}{p}}\Omega_N$ by the density of $W^{1,p}\Omega_N$. The other direction is more delicate. Interpolation involving boundary conditions is often nontrivial. The reader can see [Gui91; Lof92; Ama19] to get an idea of the challenges involved, especially at the critical regularity levels $\mathbb{N} + \frac{1}{n}$.

Nevertheless, there are a few interesting things we can say about these spaces.

Definition 85 (Neumann condition on strip). For vector field X and r > 0 small, with ψ_r as in Equation (3), define

$$\mathbf{n}_r X = \psi_r \langle X, \widetilde{\nu} \rangle \widetilde{\nu}$$
 and $\mathbf{t}_r X = X - \mathbf{n}_r X$

Then define $\mathfrak{X}_{N,r} = \{X \in \mathfrak{X} : \langle X, \widetilde{\nu} \rangle = 0 \text{ on } M_{\leq r}\}$. Similarly we can define $W^{m,p}\mathfrak{X}_{N,r}$ and $B^s_{p,q}\mathfrak{X}_{N,r}$ by setting $\|\langle X, \tilde{\nu} \rangle\|_{L^1(M_{\leq r})} = 0$. We note that $L^3 \mathfrak{X}_{N,r}$ makes sense since the definition does not require the trace theorem, unlike $L^3 \mathfrak{X}_N$ which is ill-defined.

Some basic facts:

- 1. $\mathbf{t}_r \mathfrak{X} \leq \mathfrak{X}_{N, \frac{r}{2}}$
- 2. $\mathbf{t}_r = 1$ and $\mathbf{n}_r = 0$ on $\mathfrak{X}_{N,r}$
- 3. $\mathbf{t}_{\frac{r}{2}}\mathbf{t}_r = \mathbf{t}_r$
- 4. $\|\mathbf{t}_{T}X\|_{W^{m,p}} \lesssim_{r,m,p} \|X\|_{W^{m,p}}$ for $m \in \mathbb{N}_{0}, p \in [1,\infty]$
- 5. $W^{m,p}\mathfrak{X}_{N,r}$ and $B^s_{n,q}\mathfrak{X}_{N,r}$ are Banach for $m \in \mathbb{N}_0, p \in [1,\infty], s \ge 0, q \in [1,\infty]$
- 6. $B_{p,q}^{m_{\theta}}\mathfrak{X}_{N,r} \xrightarrow{\mathbf{t}_{r}=1} (W^{m_{0},p}\mathfrak{X}_{N,\frac{r}{2}}, W^{m_{1},p}\mathfrak{X}_{N,\frac{r}{2}})_{\theta,q} \hookrightarrow B_{p,q}^{m_{\theta}}\mathfrak{X}_{N,\frac{r}{2}} \text{ for } \theta \in (0,1), m_{j} \in \mathbb{N}_{0}, m_{0} \neq m_{1}, p \in \mathbb{N}_{0}, m_{0} \neq m_{0}, m_{0} \neq m_{0} \neq m_{0}, m_{0} \neq$ $[1, \infty], q \in [1, \infty], m_{\theta} = (1 - \theta) m_0 + \theta m_1.$

Remark. The last assertion is proven by the definition of the J-method, and it works like partial interpolation. The reader can notice the similarity with the Littlewood-Paley projection $(P_{\leq N}P_{<\frac{N}{2}} = P_{<\frac{N}{2}})$. The hope is that $\mathbf{t}_r X \xrightarrow{t \downarrow 0} X$ in a good way for $X \in \mathfrak{X}_N$.

A subtle issue is that for $X \in B_{p,q}^{m_{\theta}}\mathfrak{X}_{N,r}, \|X\|_{\left(W^{m_{0},p}\mathfrak{X}_{N,\frac{r}{2}},W^{m_{1},p}\mathfrak{X}_{N,\frac{r}{2}}\right)_{\theta,r}} \lesssim_{r} \|X\|_{B_{p,q}^{m_{\theta}}\mathfrak{X}_{N,r}}.$ The implicit constant which depends on r can blow up as $r \downarrow 0$.

Define $B_{p,q}^s \mathfrak{X}_{N,0+} = B_{p,q}^s \operatorname{-cl}\left(\bigcup_{r>0 \text{ small}} B_{p,q}^s \mathfrak{X}_{N,r}\right)$ and $W^{m,p} \mathfrak{X}_{N,0+} = W^{m,p} \operatorname{-cl}\left(\bigcup_{r>0 \text{ small}} W^{m,p} \mathfrak{X}_{N,r}\right)$. Then we recover the usual spaces by results from Subsection 5.5:

Theorem 86. Let $p \in (1, \infty)$:

$$\begin{array}{l} l.\ L^{p}\mathbf{X}_{N,0,1} = L^{p}\mathbf{X},\ W^{1,p}\mathbf{X}_{N,0,1} = W^{1,p}\mathbf{X}_{N}. \\ \\ 2.\ B_{p,1}^{1}\mathbf{X}_{N,0+} = B_{p,1}^{1}\mathbf{X}_{N}. \\ \\ \hline Proof. \\ 1.\ Let\ X \in L^{p}\mathbf{X}.\ Then\ \mathbf{n}_{r}X\ \frac{L^{p}}{r_{10}}\ 0\ w shrinking support.\ If\ X \in W^{1,p}\mathbf{X}_{N},\ then\ by\ Theorem 53 \\ & \|\mathbf{n}_{r}X\|_{W^{1,p}} = \|\psi_{r}\langle X,\bar{\nu}\rangle\|_{W^{1,p}(M_{$$

.

These results hold not just for vector fields, but also for differential forms once we perform the proper modifications: for differential form ω , define $\mathbf{n}_r \omega = \psi_r \tilde{\nu}^{\flat} \wedge (\iota_{\tilde{\nu}} \omega)$, $\mathbf{t}_r \omega = \omega - \mathbf{n}_r \omega$, $W^{m,p} \Omega_r^k = \{\omega \in W^{m,p} \Omega^k :$

 $\iota_{\widetilde{\nu}}\omega = 0 \text{ on } M_{< r}\}$, replace $\langle X, \widetilde{\nu} \rangle$ with $\iota_{\widetilde{\nu}}\omega$ in the proofs etc. In particular, $B_{p,1}^{\frac{1}{p}}\Omega_{N,0+}^k = B_{p,1}^{\frac{1}{p}}\Omega_N^k$ for $p \in (1,\infty)$.

A Complexification

Throughout this small subsection, the overline always stands for conjugation, and not topological closure.

Let $\mathbb{R}X$ be a real NVS, then a **complexification** of $\mathbb{R}X$ is a tuple $\left(\mathbb{C}X, \mathbb{R}X \stackrel{\phi}{\to} \mathbb{C}X\right)$ such that

- 1. $\mathbb{C}X$ is a complex NVS.
- 2. ϕ is a linear, continuous injection and $\phi(\mathbb{R}X) \oplus i\phi(\mathbb{R}X) = \mathbb{C}X$.
- 3. $\|\phi(x)\|_{\mathbb{C}X} = \|x\|_{\mathbb{R}X}$ and $\|\phi(x) + i\phi(y)\|_{\mathbb{C}X} = \|\phi(x) i\phi(y)\|_{\mathbb{C}X} \quad \forall x, y \in \mathbb{R}X.$

The last property says $\|\cdot\|_{\mathbb{C}X}$ is a **complexification norm**. By treating $\phi(\mathbb{R}X)$ as the real part, $\forall z \in \mathbb{C}X$, we can define $\Re z$, $\Im z$ as the real and imaginary parts respectively, so $z = \Re z + i\Im z$. Then define $\overline{z} = \Re z - i\Im z$. So $\overline{\lambda z} = \overline{\lambda \overline{z}} \ \forall z \in \mathbb{C}X, \forall \lambda \in \mathbb{C}$.

Construction A standard construction of such a complexification is $\mathbb{C}X = \mathbb{R}X \otimes_{\mathbb{R}} \mathbb{C}$. As $\mathbb{R}X$ is a flat and free \mathbb{R} -module, $0 \to \mathbb{R} \hookrightarrow \mathbb{C} \xrightarrow{\mathfrak{I}} \mathbb{R} \to 0$ induces $0 \to \mathbb{R}X \xrightarrow{\phi} \mathbb{C}X \xrightarrow{\mathfrak{I}} \mathbb{R}X \to 0$ as a split short exact sequence and $\mathbb{C}X = \phi(\mathbb{R}X) \oplus i\phi(\mathbb{R}X)$. Then we can make ϕ implicit and not write it again. The representation z = x + iy = (x, y) is unique. Easy to see that any two complexifications of $\mathbb{R}X$ must be isomorphic as \mathbb{C} -modules.

We define the **minimal complexification norm** (also called **Taylor norm**)

$$\|x+iy\|_T := \sup_{\theta \in [0,2\pi]} \|x\cos\theta - y\sin\theta\|_{\mathbb{R}X} = \sup_{\theta \in [0,2\pi]} \left\|\Re e^{i\theta} \left(x+iy\right)\right\|_{\mathbb{R}X} \,\,\forall x, y \in \mathbb{R}X$$

Any other complexification norm is equivalent to $\|\cdot\|_T$.

Proof. Let $\|\cdot\|_B$ be another complexification norm. Then $\|\Re e^{i\theta}(x+iy)\|_{\mathbb{R}X} = \|\Re e^{i\theta}(x+iy)\|_B \leq \|x+iy\|_B$ (minimal) and $\|x+iy\|_B \leq \|x\|_{\mathbb{R}X} + \|y\|_{\mathbb{R}X} = \|\Re(x+iy)\|_{\mathbb{R}X} + \|\Re(-i(x+iy))\|_{\mathbb{R}X} \leq 2\|x+iy\|_T$.

So the topology of $\mathbb{C}X$ is unique. It is more convenient, however, to set $||x + iy||_{\mathbb{C}X} = ||(x,y)||_{\mathbb{R}X \oplus \mathbb{R}X} = (||x||_{\mathbb{R}X}^2 + ||y||_{\mathbb{R}X}^2)^{\frac{1}{2}} \quad \forall x, y \in \mathbb{R}X$. Easy to see that any two complexifications of $\mathbb{R}X$ must be isomorphic as complex NVS, so we write $\mathbb{C}X = \mathbb{R}X \otimes_{\mathbb{R}} \mathbb{C}$ from this point on, and if $\mathbb{R}X$ is normed, so is $\mathbb{C}X$. Obviously, if $\mathbb{R}X$ is Banach, so is $\mathbb{C}X$, and when that happens, we call $(\mathbb{R}X, \mathbb{C}X)$ a **Banach complexification couple**.

Real operators Let $(\mathbb{R}X, \mathbb{C}X)$ and $(\mathbb{R}Y, \mathbb{C}Y)$ be 2 Banach complexification couples.

- An operator $A : D(A) \leq \mathbb{C}X \to \mathbb{C}Y$ is called a **real operator** when $D(A) = \mathbb{C}\Re D(A)$ and $A\Re (D(A)) \leq \mathbb{R}Y$. In particular, $A(x, y) = (Ax, Ay) \ \forall x, y \in \mathbb{R}X$.
- An unbounded \mathbb{R} -linear operator $T: D(T) \leq \mathbb{R}X \to \mathbb{R}Y$ has a natural complexified version $T^{\mathbb{C}} = T \otimes_{\mathbb{R}} 1_{\mathbb{C}} : \mathbb{C}X \to \mathbb{C}Y$ where $D(T^{\mathbb{C}}) = \mathbb{C}D(T)$. Obviously $T^{\mathbb{C}}$ is a real operator and we write $(\mathbb{R}X, \mathbb{C}X) \xrightarrow{(T,T^{\mathbb{C}})} (\mathbb{R}Y, \mathbb{C}Y)$.

- $-\overline{D(T^{\mathbb{C}})} = D(T^{\mathbb{C}}) \text{ and } T^{\mathbb{C}}\overline{z} = \overline{T^{\mathbb{C}}z} \ \forall z \in \mathbb{C}X.$
- -T is closed $\iff T^{\mathbb{C}}$ is closed. Same for bounded, compact, densely defined.
- For any unbounded \mathbb{C} -linear operator $A: D(A) \leq \mathbb{C}X \to \mathbb{C}Y$ such that $D(A) = \mathbb{C}\Re(D(A))$, define 2 real operators $\begin{cases} \Re A = \left(\Re \circ \left(A \right|_{\Re D(A)} \right) \right)^{\mathbb{C}} \\ \Im A = \left(\Im \circ \left(A \right|_{\Re D(A)} \right) \right)^{\mathbb{C}} \\ \text{Then } A = \Re A + i \Im A. \text{ We can see that } A \text{ is real } \iff \Re A = A \iff \Im A = 0. \text{ Also, } A \text{ is bounded} \end{cases}$

 $\iff \Re A, \Im A$ are bounded.

Spectrum For $(\mathbb{R}X, \mathbb{C}X) \xrightarrow{(T,T^{\mathbb{C}})} (\mathbb{R}Y, \mathbb{C}Y)$, define

- $\rho(T) := \rho(T^{\mathbb{C}}), \sigma(T) := \sigma(T^{\mathbb{C}}).$
- $\rho_{\mathbb{R}}(T) := \{\lambda \in \mathbb{R} : \lambda T \text{ is boundedly invertible}\}\$ and $\sigma_{\mathbb{R}}(T) := \mathbb{R} \setminus \rho_{\mathbb{R}}(T).$

If $\zeta \in \mathbb{C}$ and $\zeta - T^{\mathbb{C}}$ is boundedly invertible, so is $\overline{\zeta} - T^{\mathbb{C}}$. So $\overline{\sigma(T)} = \sigma(T)$ and $\overline{\rho(T)} = \rho(T)$. For $\lambda \in \mathbb{R}$, $\lambda - T^{\mathbb{C}}$ is boundedly invertible $\iff \lambda - T$ is boundedly invertible. So $\rho_{\mathbb{R}}(T) = \rho(T) \cap \mathbb{R}$ and $\sigma_{\mathbb{R}}\left(T\right) = \sigma\left(T\right) \cap \mathbb{R}.$

Semigroup T generates an \mathbb{R} -linear C_0 semigroup $\iff T^{\mathbb{C}}$ generates a \mathbb{C} -linear C_0 semigroup. When that happens, $(e^{tT})^{\mathbb{C}} = e^{tT^{\mathbb{C}}}$.

Proof. When either happens, T and $T^{\mathbb{C}}$ are densely defined. Also, T - j and $T^{\mathbb{C}} - j$ are boundedly invertible for $j \in \mathbb{N}$ large enough, so T and $T^{\mathbb{C}}$ are closed. Easy to use Hille-Yosida to show both T and $T^{\mathbb{C}}$ must generate C_0 semigroups.

As in the proof of Hille-Yosida, define the Yosida approximations $T_j = T \frac{1}{1 - \frac{1}{j}T}, T_j^{\mathbb{C}} = T^{\mathbb{C}} \frac{1}{1 - \frac{1}{j}T^{\mathbb{C}}} = (T_j)^{\mathbb{C}}$. As T_j and $T_j^{\mathbb{C}}$ are bounded, $(e^{tT_j})^{\mathbb{C}} = e^{tT_j^{\mathbb{C}}}$ by power series expansion. Then $(e^{tT})^{\mathbb{C}} = e^{tT^{\mathbb{C}}}$ as $e^{tT} = \lim_{j \to \infty} e^{tT_j}$ pointwise.

Hilbert spaces Let $\mathbb{R}H$ be a real Hilbert space with inner product $\langle \cdot, \cdot \rangle$. Then $\mathbb{C}H$ is also Hilbert with the inner product

$$\langle x_1 + iy_1, x_2 + iy_2 \rangle_{\mathbb{C}H} := \langle x_1, x_2 \rangle + \langle y_1, y_2 \rangle + i\left(\langle y_1, x_2 \rangle - \langle x_1, y_2 \rangle \right) \ \forall x_j, y_j \in \mathbb{R}H$$

Then $||x + iy||_{\mathbb{C}H} = \left(||x||_{\mathbb{R}H}^2 + ||y||_{\mathbb{R}H}^2 \right)^{\frac{1}{2}} \forall x, y \in \mathbb{R}H$, consistent with our previously chosen norm. Also, $\langle z_1, z_2 \rangle_{\mathbb{C}H} = \langle z_2, z_1 \rangle_{\mathbb{C}H} \ \forall z_1, z_2 \in \mathbb{C}H.$ Let $(A, A^{\mathbb{C}}) : (\mathbb{R}H, \mathbb{C}H) \to (\mathbb{R}H, \mathbb{C}H)$ be unbounded.

- A is symmetric $\iff A^{\mathbb{C}}$ is symmetric. When that happens, $\langle Ax + iAy, x + iy \rangle_{\mathbb{C}H} = \langle Ax, x \rangle +$ $\langle Ay, y \rangle \ \forall x, y \in \mathbb{R}H.$
- $\mathbb{C}(\mathbb{R}H \oplus \mathbb{R}H) = \mathbb{C}H \oplus \mathbb{C}H$ and $G(A^{\mathbb{C}}) = \mathbb{C}G(A)$ (graphs). Also $\mathbb{C}(G(A)^{\perp}) = G(A^{\mathbb{C}})^{\perp}$.
- A is self-adjoint $\iff A^{\mathbb{C}}$ is self-adjoint. When this happens, $\sigma(A) = \sigma(A^{\mathbb{C}}) \subset \mathbb{R}$.
- A is dissipative $\iff A^{\mathbb{C}}$ is dissipative.

For more information on complexification, see [Glü17, Appendix C].

Nomenclature

- ψ_r, f_r cutoffs on M living near the boundary, page 13
- $e^{t\Delta}$ the absolute Neumann heat flow, defined for the proof of Onsager's conjecture, page 14
- $\mathcal{L}((X_0, X_0), (Y_0, Y_1))$ morphisms between interpolation couples, page 20

 $(X_0, X_1)_{\theta,q}$ real interpolation, page 21

 $[X_0,X_1]_{\theta}\,$ complex interpolation, page 21

- $W^{m,p}$ Sobolev spaces, page 30
- $B_{p,q}^s$ Besov spaces, page 30
- $F_{p,q}^s$ Triebel-Lizorkin spaces, page 30
- $\mathfrak{C}^{s}(\Omega)$ Zygmund spaces, page 33

 $\Omega_{< r} \quad \{ x \in \Omega : \operatorname{dist}(x, \partial \Omega) < r \}$, page 34

 $\|f\|_{L^p(\Omega, avg)}$ integration on probability space, page 34

 ν outwards unit normal vector field on ∂M , page 37

- $\tilde{\nu}$ extension of ν near ∂M , page 37
- $j \qquad j: \partial M \hookrightarrow M$ is the smooth inclusion map, page 37
- ι interior product (contraction) of differential forms, page 37

 $\operatorname{vol}_{\partial}$ volume form of ∂M , page 37

 $\Gamma(\mathbb{F}), \Gamma_c(\mathbb{F}), \Gamma_{00}(\mathbb{F})$ the space of smooth sections of \mathbb{F} with different support conditions, page 38

 $\langle \langle \sigma, \theta \rangle \rangle$ dot product on $\Gamma(\mathbb{F})$, page 38

 $\mathbb{R}W^{m,p},\ \mathbb{C}W^{m,p}$ real and complexified versions of function space, page 40

 $\mathfrak{X}M$ set of smooth vector fields on M, page 40

t tangential part, page 40

 \mathbf{n} normal part, page 40

 $\Omega^{k}(M)$ set of smooth differential forms on M, page 41

 $X_p^{\flat}, \ \omega_p^{\sharp}$ musical isomorphism, page 41

 \star Hodge star, page 41

 δ codifferential, page 41

- Δ Hodge Laplacian, page 41
- R_{abcd} Riemann curvature tensor, page 41
- Ric Weitzenbock curvature operator, page 41

 $\langle T, Q \rangle$ tensor inner product, page 42

 $\langle \omega, \eta \rangle_{\Lambda}, \langle \langle \omega, \eta \rangle \rangle_{\Lambda}$ Hodge inner product, page 42

 $\mathcal{D}(\omega, \eta)$ Dirichlet integral, page 43

 $\Omega^k_D, \Omega^k_{\mathrm{hom}D}$ different Dirichlet conditions for differential forms, page 44

 $\Omega_N^k, \Omega_{\text{hom }N}^k$ different Neumann conditions for differential forms, page 44

 $\mathcal{H}^k, \mathcal{H}^k_D, \mathcal{H}^k_N$ harmonic fields, then with Dirichlet and Neumann conditions, page 44

 $L^2\text{-}\mathrm{cl}\left(\cdot\right)\,$ closure under L^2 norm, page 44

 $\mathcal{P}^N,~\mathcal{P}^{N\perp},~\mathcal{P}^D,~\mathcal{P}^{D\perp}$ natural orthogonal decomposition, page 45

 Δ_N injective Neumann Laplacian, page 45

 $(-\Delta_N)^{-1}$, $(-\Delta_D)^{-1}$ Neumann and Dirichlet potentials, page 45

 δ_c, d_c adjoints of d and δ , page 46

 $\mathcal{P}_1\omega$, $\mathcal{P}_2\omega$, $\mathcal{P}_3\omega$ the component projections in Hodge decomposition, page 47

 $\mathcal{P}_3^N, \mathcal{P}_3^{\text{ex}}, \mathcal{P}_3^D, \mathcal{P}_3^{\text{co}}$ Friedrichs decomposition, page 49

- \mathbb{P} Leray projection, page 49
- $\widetilde{\Delta_N}$ non-injective Neumann Laplacian, page 52
- A_p generator of heat flow on L^p , page 53
- $A_{1,p}$ generator of heat flow on $W^{1,p}$, page 55
- $\mathscr{D}_N\Omega^k, \mathscr{D}'_N\Omega^k$ heated forms and heatable currents, page 56

 $D_N, \widetilde{D_N}$ the injective and non-injective (Neumann) Hodge-Dirac operators, page 65

 $W^{m,p}(D_N), W^{m,p}\left(\widetilde{D_N}\right)$ Hodge-Sobolev spaces, page 65

 $\mathbb{C}Y, T^{\mathbb{C}}$ complexification of spaces and operators , page 68

References

- [ABK19] Wolfgang Arendt, Manuel Bernhard, and Marcel Kreuter. *Elliptic problems and holomorphic functions in Banach spaces.* 2019. eprint: arXiv:1904.03088.
- [Ada03] R. A. Adams. Sobolev spaces. Amsterdam Boston: Academic Press, 2003. ISBN: 9780120441433.
- [Agm62] Shmuel Agmon. "On the eigenfunctions and on the eigenvalues of general elliptic boundary value problems". In: Communications on Pure and Applied Mathematics 15.2 (1962), pp. 119–147. DOI: 10.1002/cpa.3160150203. URL: https://doi.org/10.1002/cpa.3160150203.
- [AKS62] N. Aronszajn, A. Krzywicki, and J. Szarski. "A unique continuation theorem for exterior differential forms on Riemannian manifolds". In: Arkiv för Matematik 4.5 (Aug. 1962), pp. 417–453.
 DOI: 10.1007/bf02591624. URL: https://doi.org/10.1007/bf02591624.

- [AM04] Andreas Axelsson and Alan McIntosh. "Hodge Decompositions on Weakly Lipschitz Domains". In: Advances in Analysis and Geometry. Birkhäuser Basel, 2004, pp. 3–29. DOI: 10.1007/978– 3-0348-7838-8_1. URL: https://doi.org/10.1007/978-3-0348-7838-8_1.
- [Ama19] Herbert Amann. Linear and Quasilinear Parabolic Problems. Springer International Publishing, 2019. DOI: 10.1007/978-3-030-11763-4. URL: https://doi.org/10.1007/978-3-030-11763-4.
- [Are+11] Wolfgang Arendt et al. Vector-valued Laplace Transforms and Cauchy Problems: Second Edition (Monographs in Mathematics). Birkhäuser, 2011. ISBN: 303480086X.
- [BAE16] Hind Baba, Chérif Amrouche, and Miguel Escobedo. Analyticity of the semi-group generated by the Stokes operator with Navier-type boundary conditions on Lp-spaces. 2016. DOI: 10.1090/ conm/666/13337. URL: https://doi.org/10.1090/conm/666/13337.
- [Bar+18] Claude Bardos et al. "On the Extension of Onsager's Conjecture for General Conservation Laws".
 In: Journal of Nonlinear Science 29.2 (Sept. 2018), pp. 501-510. DOI: 10.1007/s00332-018-9496-4.
 URL: https://doi.org/10.1007%2Fs00332-018-9496-4.
- [Bar+19] Claude Bardos et al. Onsager's conjecture in bounded domains for the conservation of entropy and other companion laws. 2019. eprint: arXiv:1902.07120.
- [Bau17] Fabrice Baudoin. Stochastic analysis and related topics : a festschrift in honor of Rodrigo Bañuelos. Cham: Birkhäuser, 2017. ISBN: 978-3-319-59670-9.
- [BB97] Alexander Braverman and Maxim Braverman. Tempered currents and the cohomology of the remote fiber of a real polynomial map. 1997. eprint: arXiv:alg-geom/9704001.
- [BBD18] Huy-Qui Bui, The Anh Bui, and Xuan Thinh Duong. Weighted Besov and Triebel-Lizorkin spaces associated to operators. 2018. eprint: arXiv:1809.02795.
- [BL76] Jöran Bergh and Jörgen Löfström. Interpolation Spaces. Springer Berlin Heidelberg, 1976. DOI: 10.1007/978-3-642-66451-9. URL: https://doi.org/10.1007/978-3-642-66451-9.
- [BT17] Claude Bardos and Edriss S. Titi. "Onsager's Conjecture for the Incompressible Euler Equations in Bounded Domains". In: Archive for Rational Mechanics and Analysis 228.1 (Nov. 2017), pp. 197-207. DOI: 10.1007/s00205-017-1189-x. URL: https://doi.org/10.1007/s00205-017-1189-x.
- [BTW19] Claude Bardos, Edriss S. Titi, and Emil Wiedemann. "Onsager's Conjecture with Physical Boundaries and an Application to the Vanishing Viscosity Limit". In: Communications in Mathematical Physics (July 2019). DOI: 10.1007/s00220-019-03493-6. URL: https://doi.org/ 10.1007%2Fs00220-019-03493-6.
- [CET94] Peter Constantin, Weinan E, and Edriss S. Titi. "Onsager's conjecture on the energy conservation for solutions of Euler's equation". In: Comm. Math. Phys. 165.1 (1994), pp. 207–209. URL: https: //projecteuclid.org:443/euclid.cmp/1104271041.
- [Cha06] Isaac Chavel. *Riemannian geometry : a modern introduction*. New York: Cambridge University Press, 2006. ISBN: 978-0521619547.
- [Che+08] A Cheskidov et al. "Energy conservation and Onsager's conjecture for the Euler equations".
 In: Nonlinearity 21.6 (Apr. 2008), pp. 1233–1252. DOI: 10.1088/0951-7715/21/6/005. URL: https://doi.org/10.1088/0951-7715/21/6/005.
- [Con54] P. E. Conner. "The Green's and Neumann's Problems for Differential Forms on Riemannian Manifolds". In: Proceedings of the National Academy of Sciences 40.12 (Dec. 1954), pp. 1151– 1155. DOI: 10.1073/pnas.40.12.1151. URL: https://doi.org/10.1073/pnas.40.12.1151.

- [COQ09] Gui-Qiang Chen, Dan Osborne, and Zhongmin Qian. "The Navier-Stokes equations with the kinematic and vorticity boundary conditions on non-flat boundaries". In: Acta Mathematica Scientia 29.4 (July 2009), pp. 919–948. DOI: 10.1016/s0252-9602(09)60078-3. URL: https: //doi.org/10.1016%2Fs0252-9602%2809%2960078-3.
- [DM96] Xuan Thinh Duong and Alan McIntosh. "Functional calculi of second-order elliptic partial differential operators with bounded measurable coefficients". In: *The Journal of Geometric Analysis* 6.2 (June 1996), pp. 181–205. DOI: 10.1007/bf02921599. URL: https://doi.org/10.1007/bf02921599.
- [DN18] Theodore D. Drivas and Huy Q. Nguyen. "Onsager's Conjecture and Anomalous Dissipation on Domains with Boundary". In: SIAM Journal on Mathematical Analysis 50.5 (Jan. 2018), pp. 4785–4811. DOI: 10.1137/18m1178864. URL: https://doi.org/10.1137%2F18m1178864.
- [Eng00] Engel. One-parameter semigroups for linear evolution equations. New York: Springer, 2000. ISBN: 978-0-387-98463-6.
- [Eyi94] Gregory L. Eyink. "Energy dissipation without viscosity in ideal hydrodynamics I. Fourier analysis and local energy transfer". In: *Physica D: Nonlinear Phenomena* 78.3-4 (Nov. 1994), pp. 222–240. DOI: 10.1016/0167-2789(94)90117-1. URL: https://doi.org/10.1016/0167-2789(94)90117-1.
- [Fac15] Stephan Fackler. "Regularity properties of sectorial operators: extrapolation, counterexamples and generic classes". en. PhD thesis. 2015. DOI: 10.18725/oparu-3268. URL: https://oparu. uni-ulm.de/xmlui/handle/123456789/3295.
- [FFP16] Hans G. Feichtinger, Hartmut Führ, and Isaac Z. Pesenson. "Geometric Space-Frequency Analysis on Manifolds". In: Journal of Fourier Analysis and Applications 22.6 (Jan. 2016), pp. 1294– 1355. DOI: 10.1007/s00041-015-9457-3. URL: https://doi.org/10.1007%2Fs00041-015-9457-3.
- [FK64] Hiroshi Fujita and Tosio Kato. "On the Navier-Stokes initial value problem. I". In: Archive for Rational Mechanics and Analysis 16.4 (1964), pp. 269–315. DOI: 10.1007/bf00276188. URL: https://doi.org/10.1007/bf00276188.
- [FMM98] Eugene Fabes, Osvaldo Mendez, and Marius Mitrea. "Boundary Layers on Sobolev-Besov Spaces and Poisson's Equation for the Laplacian in Lipschitz Domains". In: Journal of Functional Analysis 159.2 (Nov. 1998), pp. 323-368. DOI: 10.1006/jfan.1998.3316. URL: https://doi.org/ 10.1006/jfan.1998.3316.
- [Gaf54] Matthew P. Gaffney. "The Heat Equation Method of Milgram and Rosenbloom for Open Riemannian Manifolds". In: *The Annals of Mathematics* 60.3 (Nov. 1954), p. 458. DOI: 10.2307/1969846. URL: https://doi.org/10.2307/1969846.
- [Gig81] Yoshikazu Giga. "Analyticity of the semigroup generated by the Stokes operator inL r spaces".
 In: Mathematische Zeitschrift 178.3 (Sept. 1981), pp. 297–329. DOI: 10.1007/bf01214869. URL: https://doi.org/10.1007/bf01214869.
- [Gig82] Yoshikazu Giga. "The nonstationary Navier-Stokes system with some first order boundary condition". In: Proceedings of the Japan Academy, Series A, Mathematical Sciences 58.3 (1982), pp. 101-104. DOI: 10.3792/pjaa.58.101. URL: https://doi.org/10.3792/pjaa.58.101.
- [Glü17] Jochen Glück. "Invariant sets and long time behaviour of operator semigroups". en. PhD thesis. 2017. DOI: 10.18725/oparu-4238. URL: https://oparu.uni-ulm.de/xmlui/handle/ 123456789/4277.
- [GM85] Yoshikazu Giga and Tetsuro Miyakawa. "Solutions in Lr of the Navier-Stokes initial value problem". In: Archive for Rational Mechanics and Analysis 89.3 (1985), pp. 267–281. DOI: 10.1007/ bf00276875. URL: https://doi.org/10.1007/bf00276875.
- [GMM10] V. Gol'dshtein, I. Mitrea, and M. Mitrea. "Hodge decompositions with mixed boundary conditions and applications to partial differential equations on lipschitz manifolds". In: Journal of Mathematical Sciences 172.3 (Dec. 2010), pp. 347-400. DOI: 10.1007/s10958-010-0200-y. URL: https://doi.org/10.1007%2Fs10958-010-0200-y.
- [GO14] Loukas Grafakos and Seungly Oh. "The Kato-Ponce Inequality". In: Communications in Partial Differential Equations 39.6 (May 2014), pp. 1128–1157. DOI: 10.1080/03605302.2013.822885.
 URL: https://doi.org/10.1080/03605302.2013.822885.
- [Gui91] Davide Guidetti. "On interpolation with boundary conditions". In: Mathematische Zeitschrift 207.1 (May 1991), pp. 439–460. DOI: 10.1007/bf02571401. URL: https://doi.org/10.1007/bf02571401.
- [Haa06] Markus Haase. The Functional Calculus for Sectorial Operators. Birkhäuser Basel, 2006. DOI: 10.1007/3-7643-7698-8. URL: https://doi.org/10.1007/3-7643-7698-8.
- [HMY08] Yongsheng Han, Detlef Müller, and Dachun Yang. "A Theory of Besov and Triebel-Lizorkin Spaces on Metric Measure Spaces Modeled on Carnot-Carathéodory Spaces". In: Abstract and Applied Analysis 2008 (2008), pp. 1–250. DOI: 10.1155/2008/893409. URL: https://doi.org/ 10.1155%2F2008%2F893409.
- [HS18] Matthias Hieber and Jürgen Saal. "The Stokes Equation in the Lp-Setting: Well-Posedness and Regularity Properties". In: Handbook of Mathematical Analysis in Mechanics of Viscous Fluids. Springer International Publishing, 2018, pp. 117–206. DOI: 10.1007/978-3-319-13344-7_3. URL: https://doi.org/10.1007/978-3-319-13344-7_3.
- [Hsu02] Elton Hsu. "Multiplicative functional for the heat equation on manifolds with boundary". In: The Michigan Mathematical Journal 50.2 (Aug. 2002), pp. 351-367. DOI: 10.1307/mmj/1028575738.
 URL: https://doi.org/10.1307%2Fmmj%2F1028575738.
- [IO15] Philip Isett and Sung-Jin Oh. "A heat flow approach to Onsager's conjecture for the Euler equations on manifolds". In: Transactions of the American Mathematical Society 368.9 (Nov. 2015), pp. 6519–6537. DOI: 10.1090/tran/6549. URL: https://doi.org/10.1090/tran/6549.
- [Ise18] Philip Isett. "A proof of Onsager's conjecture". In: Annals of Mathematics 188.3 (2018), p. 871.
 DOI: 10.4007/annals.2018.188.3.4. URL: https://doi.org/10.4007/annals.2018.188.3.
 4.
- [KP14] Gerard Kerkyacharian and Pencho Petrushev. "Heat kernel based decomposition of spaces of distributions in the framework of Dirichlet spaces". In: Transactions of the American Mathematical Society 367.1 (June 2014), pp. 121–189. DOI: 10.1090/s0002-9947-2014-05993-x. URL: https://doi.org/10.1090%2Fs0002-9947-2014-05993-x.
- [KP88] Tosio Kato and Gustavo Ponce. "Commutator estimates and the euler and navier-stokes equations". In: Communications on Pure and Applied Mathematics 41.7 (Oct. 1988), pp. 891–907. DOI: 10.1002/cpa.3160410704. URL: https://doi.org/10.1002/cpa.3160410704.
- [KR06] S. Klainerman and I. Rodnianski. "A geometric approach to the Littlewood-Paley theory". In: GAFA Geometric And Functional Analysis 16.1 (Feb. 2006), pp. 126–163. DOI: 10.1007/s00039-006-0551-1. URL: https://doi.org/10.1007/s00039-006-0551-1.

- [KW16] Christoph Kriegler and Lutz Weis. "Paley-Littlewood decomposition for sectorial operators and interpolation spaces". In: *Mathematische Nachrichten* 289.11-12 (Feb. 2016), pp. 1488–1525. DOI: 10.1002/mana.201400223. URL: https://doi.org/10.1002%2Fmana.201400223.
- [Lee09] Jeffrey Lee. Manifolds and differential geometry. Providence, R.I: American Mathematical Society, 2009. ISBN: 978-0821848159.
- [Lem02] Pierre Gilles Lemarie-Rieusset. Recent developments in the Navier-Stokes problem (Chapman & Hall/CRC Research Notes in Mathematics Series). CRC Press, Apr. 2002. ISBN: 1584882204. URL: https://www.xarg.org/ref/a/1584882204/.
- [LJ14] Camillo De Lellis and László Székelyhidi Jr. "Dissipative Euler flows and Onsager's conjecture".
 In: Journal of the European Mathematical Society 16.7 (2014), pp. 1467–1505. DOI: 10.4171/ jems/466. URL: https://doi.org/10.4171/jems/466.
- [Lof92] Jorgen Lofstrom. "Interpolation of Boundary Value Problems of Neumann Type on Smooth Domains". In: Journal of the London Mathematical Society s2-46.3 (Dec. 1992), pp. 499-516.
 DOI: 10.1112/jlms/s2-46.3.499. URL: https://doi.org/10.1112/jlms/s2-46.3.499.
- [LS12] Camillo De Lellis and László Székelyhidi. "Dissipative continuous Euler flows". In: Inventiones mathematicae 193.2 (Oct. 2012), pp. 377–407. DOI: 10.1007/s00222-012-0429-9. URL: https: //doi.org/10.1007/s00222-012-0429-9.
- [LS19] Camillo De Lellis and László Székelyhidi. "On Turbulence and Geometry: from Nash to Onsager". In: Notices of the American Mathematical Society ??05 (May 2019), p. 1. DOI: 10.1090/ noti1868. URL: https://doi.org/10.1090/noti1868.
- [Lun95] Alessandra Lunardi. Analytic semigroups and optimal regularity in parabolic problems. Basel: Birkhäuser, 1995. ISBN: 978-3-0348-0556-8.
- [McI10] Alan McIntosh. "Operator theory-Spectra and functional calculi". In: (2010).
- [McI86] Alan McIntosh. Operators which have an Hinf functional calculus. English. Operator theory and partial differential equations, Miniconf. Ryde/Aust. 1986, Proc. Cent. Math. Anal. Aust. Natl. Univ. 14, 210-231 (1986). 1986.
- [Miy80] Tetsuro Miyakawa. "The Lp approach to the Navier-Stokes equations with the Neumann boundary condition". In: *Hiroshima Mathematical Journal* 10.3 (1980), pp. 517–537. DOI: 10.32917/ hmj/1206134338. URL: https://doi.org/10.32917/hmj/1206134338.
- [MM08] Marius Mitrea and Sylvie Monniaux. "The regularity of the Stokes operator and the Fujita-Kato approach to the Navier-Stokes initial value problem in Lipschitz domains". In: Journal of Functional Analysis 254.6 (Mar. 2008), pp. 1522–1574. DOI: 10.1016/j.jfa.2007.11.021. URL: https://doi.org/10.1016/j.jfa.2007.11.021.
- [MM09a] Marius Mitrea and Sylvie Monniaux. "On the analyticity of the semigroup generated by the Stokes operator with Neumann-type boundary conditions on Lipschitz subdomains of Riemannian manifolds". In: Transactions of the American Mathematical Society 361.06 (June 2009), pp. 3125–3125. DOI: 10.1090/s0002-9947-08-04827-7. URL: https://doi.org/10.1090/s0002-9947-08-04827-7.
- [MM09b] Marius Mitrea and Sylvie Monniaux. "The nonlinear Hodge-Navier-Stokes equations in Lipschitz domains". In: *Differential Integral Equations* 22.3/4 (Mar. 2009), pp. 339–356. URL: https://projecteuclid.org:443/euclid.die/1356019778.
- [MM18] Alan Mcintosh and Sylvie Monniaux. "Hodge-Dirac, Hodge-Laplacian and Hodge-Stokes operators in Lp spaces on Lipschitz domains". In: *Revista Matemática Iberoamericana* 34.4 (2018), pp. 1711–1753. URL: https://hal.archives-ouvertes.fr/hal-01351604.

- [Mon13] Sylvie Monniaux. "Various boundary conditions for Navier-Stokes equations in bounded Lipschitz domains". In: Discrete and Continuous Dynamical Systems Series S 6.5 (Mar. 2013), pp. 1355–1369. DOI: 10.3934/dcdss.2013.6.1355. URL: https://doi.org/10.3934%2Fdcdss.2013.6.1355.
- [MR51] A. N. Milgram and P. C. Rosenbloom. "Harmonic Forms and Heat Conduction: I: Closed Riemannian Manifolds". In: Proceedings of the National Academy of Sciences 37.3 (Mar. 1951), pp. 180–184. DOI: 10.1073/pnas.37.3.180. URL: https://doi.org/10.1073/pnas.37.3.180.
- [NN18] Quoc-Hung Nguyen and Phuoc-Tai Nguyen. "Onsager's Conjecture on the Energy Conservation for Solutions of Euler Equations in Bounded Domains". In: Journal of Nonlinear Science 29.1 (July 2018), pp. 207–213. DOI: 10.1007/s00332-018-9483-9. URL: https://doi.org/10. 1007/s00332-018-9483-9.
- [NT19] Virginia Naibo and Alexander Thomson. "Coifman-Meyer multipliers: Leibniz-type rules and applications to scattering of solutions to PDEs". In: *Transactions of the American Mathematical Society* (Apr. 2019), p. 1. DOI: 10.1090/tran/7866. URL: https://doi.org/10.1090/tran/7866.
- [Ouy17] Cheng Ouyang. "Multiplicative functional for the heat equation on manifolds with boundary". In: Stochastic Analysis and Related Topics. Springer, 2017, pp. 67–83.
- [Rha84] Rham. Differentiable manifolds : forms, currents, harmonic forms. Berlin, Heidelberg: Springer-Verlag, 1984. ISBN: 978-3-642-61754-6.
- [RRS18] James C. Robinson, José L. Rodrigo, and Jack W.D. Skipper. "Energy conservation for the Euler equations on T2xR+ for weak solutions defined without reference to the pressure". In: Asymptotic Analysis 110.3-4 (Nov. 2018), pp. 185–202. ISSN: 1875-8576. DOI: 10.3233/ASY-181482. URL: http://doi.org/10.3233/ASY-181482.
- [Rud91] Walter Rudin. Functional Analysis. McGraw-Hill Science/Engineering/Math, 1991. ISBN: 0070542368.
- [Sch11] Cornelia Schneider. "Traces of Besov and Triebel-Lizorkin spaces on domains". In: Mathematische Nachrichten 284.5-6 (Mar. 2011), pp. 572–586. DOI: 10.1002/mana.201010052. URL: https: //doi.org/10.1002/mana.201010052.
- [Sch95] Günter Schwarz. Hodge Decomposition—A Method for Solving Boundary Value Problems. Springer Berlin Heidelberg, 1995. DOI: 10.1007/bfb0095978. URL: https://doi.org/10.1007/ bfb0095978.
- [She12] Zhongwei Shen. "Resolvent Estimates in L p for the Stokes Operator in Lipschitz Domains". In: Archive for Rational Mechanics and Analysis 205.2 (Mar. 2012), pp. 395–424. DOI: 10.1007/ s00205-012-0506-7. URL: https://doi.org/10.1007/s00205-012-0506-7.
- [Ste71] Elias M. Stein. Singular Integrals and Differentiability Properties of Functions. (PMS-30). Princeton University Press, Feb. 1971. ISBN: 0691080798. URL: https://www.xarg.org/ref/a/ 0691080798/.
- [Str83] Robert S Strichartz. "Analysis of the Laplacian on the complete Riemannian manifold". In: Journal of Functional Analysis 52.1 (June 1983), pp. 48–79. DOI: 10.1016/0022-1236(83) 90090-3. URL: https://doi.org/10.1016/0022-1236(83)90090-3.
- [Str89] Robert S. Strichartz. "Harmonic analysis as spectral theory of Laplacians". In: Journal of Functional Analysis 87.1 (Nov. 1989), pp. 51–148. DOI: 10.1016/0022-1236(89)90004-9. URL: https://doi.org/10.1016%2F0022-1236%2889%2990004-9.

- [Tan18] Koichi Taniguchi. "Besov spaces generated by the Neumann Laplacian". In: European Journal of Mathematics 4.4 (Mar. 2018), pp. 1521–1548. DOI: 10.1007/s40879-018-0224-2. URL: https://doi.org/10.1007%2Fs40879-018-0224-2.
- [Tao06] Terence Tao. Nonlinear dispersive equations : local and global analysis. Providence, R.I: American Mathematical Society, 2006. ISBN: 978-0-8218-4143-3.
- [Tao07] Terence Tao. LECTURE NOTES 6 FOR 247B. 2007.
- [Tao16] Terrence Tao. Notes on the Nash embedding theorem. May 2016. URL: https://terrytao. wordpress.com/2016/05/11/notes-on-the-nash-embedding-theorem/.
- [Tay11a] Michael E. Taylor. Partial Differential Equations I. Springer New York, 2011. DOI: 10.1007/978-1-4419-7055-8. URL: https://doi.org/10.1007/978-1-4419-7055-8.
- [Tay11b] Michael E. Taylor. Partial Differential Equations II. Springer New York, 2011. DOI: 10.1007/ 978-1-4419-7052-7. URL: https://doi.org/10.1007/978-1-4419-7052-7.
- [Tri10] H. Triebel. *Theory of Function Spaces*. Modern Birkhäuser Classics. Springer Basel, 2010. ISBN: 9783034604154. URL: https://books.google.com/books?id=fMUM16iLMGwC.
- [Tri92] H. Triebel. *Theory of Function Spaces II*. Monographs in Mathematics. Birkhäuser Verlag, 1992. ISBN: 9780817626396. URL: https://books.google.com/books?id=-0vvAAAMAAJ.
- [Tro09] M. Troyanov. "On the Hodge Decomposition in Rn". In: Moscow Mathematical Journal 9.4 (2009), pp. 899–926. DOI: 10.17323/1609-4514-2009-9-4-899-926. URL: https://doi.org/10.17323%2F1609-4514-2009-9-4-899-926.
- [Voi92] Jürgen Voigt. "Abstract Stein Interpolation". In: Mathematische Nachrichten 157.1 (1992), pp. 197-199. DOI: 10.1002/mana.19921570115. URL: https://doi.org/10.1002/mana. 19921570115.
- [Wal84] Robert Wald. General relativity. Chicago: University of Chicago Press, 1984. ISBN: 978-0226870335.
- [Wu91] Siye Wu. "Topological quantum field theories on manifolds with a boundary". In: Comm. Math. Phys. 136.1 (1991), pp. 157–168. URL: https://projecteuclid.org:443/euclid.cmp/ 1104202263.