

On Peterson's open problem and representations of the general linear groups

Đặng Võ Phúc

*Faculty of Education Studies, University of Khanh Hoa,
01 Nguyen Chanh, Nha Trang, Khanh Hoa, Viet Nam*

Abstract

Fix $\mathbb{Z}/2$ is the prime field of two elements and write $\mathcal{A}_2$ for the mod 2 Steenrod algebra. Denote by $GL_d := GL(d, \mathbb{Z}/2)$ the general linear group of rank d over $\mathbb{Z}/2$ and by $\mathcal{P}_d$ the polynomial algebra $\mathbb{Z}/2[x_1, x_2, \dots, x_d]$ as a connected unstable left $\mathcal{A}_2$ -module on d generators of degree one. We study the *Peterson hit problem* of finding a minimal set of $\mathcal{A}_2$ -generators for $\mathcal{P}_d$. Equivalently, we need to determine a basis for the $\mathbb{Z}/2$ -vector space

$$Q\mathcal{P}_d := \mathbb{Z}/2 \otimes_{\mathcal{A}_2} \mathcal{P}_d \cong \mathcal{P}_d / \mathcal{A}_2^+ \mathcal{P}_d$$

in each degree $n \geq 1$. Note that this space is a representation of GL_d over $\mathbb{Z}/2$. The problem for $d = 5$ is not yet completely solved, and unknown in general.

In this work, we give an explicit solution to the hit problem of five variables in the generic degree $n = r(2^t - 1) + 2^t s$ with $r = d = 5$, $s = 8$ and t an arbitrary non-negative integer. An application of this study to the cases $t = 0$ and $t = 1$ shows that the Singer algebraic transfer of rank 5 is an isomorphism in bidegrees $(5, 5 + 8)$ and $(5, 5 + 21)$. Here, the Singer transfer of rank d is a $\mathbb{Z}/2$ -algebra homomorphism from GL_d -coinvariants of certain subspaces of $Q\mathcal{P}_d$ to the cohomology groups of the Steenrod algebra, $\text{Ext}_{\mathcal{A}_2}^{d, d+*}(\mathbb{Z}/2, \mathbb{Z}/2)$. It is one of the useful tools for describing these cohomology groups.

Keywords:

Adams spectral sequences; Steenrod algebra; Peterson hit problem; Algebraic transfer

2010 MSC: Primary 55Q45, 55S10; Secondary 55S05, 55T15.

Contents

1	Introduction and statement of results	2
2	Preliminaries	7
2.1	Kameko's squaring operation	7
2.2	On the Peterson "hit" problem	8
3	Generators of the $\mathcal{A}_2$-module $\mathcal{P}_5$ in the generic degree $13 \cdot 2^t - 5$	10
3.1	Some homomorphisms	10
3.2	Some examples on $\phi_{(k; \mathcal{K})}$	11
3.3	Singer's criterion on the $\mathcal{A}_2$ -decomposable and Sum's conjecture	12
3.4	Proof of Theorem 1.1	14
3.4.1	The case $t = 1$	14
3.4.2	The case $t = 2$	18

4	An application of Theorem 1.1	34
4.1	Computation of $(Q\mathcal{P}_5)_8^{GL_5}$	35
4.2	Computation of $(\text{Ker}(Sq_*^0)_{(5,21)})^{S_5}$	37
4.3	Proof of Theorem 1.2	40
5	Proof of Theorem 1.3	40
6	Appendix	41
6.1	$\mathcal{A}_2$ -generators for $\mathcal{P}_5$ in degree 8	41
6.2	$\mathcal{A}_2$ -generators for $\mathcal{P}_5^0$ in degree 21	42
6.3	$\mathcal{A}_2$ -generators for $\mathcal{P}_5^+$ in degree 21	45
6.4	$\mathcal{A}_2$ -generators for $\mathcal{P}_5^0$ in degree 22	46
6.5	$\mathcal{A}_2$ -generators for $\mathcal{P}_5^+$ in degree 22	49
6.6	$\mathcal{A}_2$ -generators for $\mathcal{P}_5^0$ in degree 47	52
6.7	$\mathcal{A}_2$ -generators for $\mathcal{P}_5^+$ in degree 47	55

1. Introduction and statement of results

Throughout this article, we shall work only at the prime 2. Let $Sq^k : H^*(\mathbb{X}) \rightarrow H^{k+*}(\mathbb{X})$ be the stable cohomology operation of degree $k \geq 0$, which is introduced by Steenrod in 1947 (see [45]). Here $H^*(\mathbb{X})$ is the singular cohomology group of the topological space $\mathbb{X}$ with coefficients in $\mathbb{Z}/2$. The $\mathbb{Z}/2$ -graded algebra $\mathcal{A}_2$ generated by the operations Sq^k is called *the mod 2 Steenrod algebra* and acts in a natural way on the cohomology of any space $\mathbb{X}$. For d a natural number, we denote by $B(\mathbb{Z}/2)^{\times d}$ the classifying space of elementary abelian 2-group $(\mathbb{Z}/2)^{\times d}$ of rank d and by $\mathcal{P}_d = \mathbb{Z}/2[x_1, x_2, \dots, x_d]$ the polynomial algebra on d variables of degree 1. This algebra is isomorphic to the cohomology with $\mathbb{Z}/2$ -coefficients of $B(\mathbb{Z}/2)^{\times d}$, hence $\mathcal{P}_d$ has an unstable left $\mathcal{A}_2$ -module structure. The left action of $\mathcal{A}_2$ on $\mathcal{P}_d$ is determined by the unstable condition $Sq^1(x_i) = x_i^2$, $Sq^k(x_i) = 0$ for $k > 1$ and Cartan's formula $Sq^k(x \cdot y) = \sum_{1 \leq j \leq k} Sq^{k-j}(x) \cdot Sq^j(y)$ (see [45]).

The investigation of the homotopy classification of topological spaces leads us to the study the (mod 2) cohomology groups of the Steenrod algebra, $\text{Ext}_{\mathcal{A}_2}^{d,d+*}(\mathbb{Z}/2, \mathbb{Z}/2)$. It has been thoroughly studied for $d \leq 5$ (see [6, 17, 52]). However, for d higher, the calculations seem to be difficult. Moreover, it has a deep connection with the "hit" problem of our interest in determining a minimal set of $\mathcal{A}_2$ -generators for $\mathcal{P}_d$. Equivalently, we need to find a basis for the $\mathbb{Z}/2$ -graded vector space

$$Q\mathcal{P}_d := \mathbb{Z}/2 \otimes_{\mathcal{A}_2} \mathcal{P}_d \cong \mathcal{P}_d / \mathcal{A}_2^+ \mathcal{P}_d,$$

where $\mathcal{A}_2^+$ denotes the *augmentation ideal* of $\mathcal{A}_2$ and $\mathbb{Z}/2$ is viewed as a right $\mathcal{A}_2$ -module concentrated in grading 0. This problem was posed by Peterson [32] in 1987. However, it remains open for $d \geq 5$.

For a non-negative integer n , let $(\mathcal{P}_d)_n$ be the subspace of $\mathcal{P}_d$ consisting of all the homogeneous polynomials of degree n in $\mathcal{P}_d$. Denote by $(Q\mathcal{P}_d)_n$ the subspace of $Q\mathcal{P}_d$ consisting of all the classes represented by the homogeneous polynomials in $(\mathcal{P}_d)_n$. As well known, the general linear group $GL_d := GL(d, \mathbb{Z}/2)$ acts regularly on $\mathcal{P}_d$ by matrix substitution. Further, the two actions of $\mathcal{A}_2$ and GL_d upon $\mathcal{P}_d$ commute with each other; hence there is an inherited action of GL_d on $Q\mathcal{P}_d$. From this event, the hit problem becomes one of the main tools for studying representations of the general linear groups over $\mathbb{Z}/2$. Therefrom the hit problem has attracted great interest of many algebraic topologists (see Ault and Singer [3], Crabb and Hubbuck [8], Crossley [9], Kameko [15], Mothebe and his collaborators [21]-[24], Nam [25], Pengelley and Williams [28, 30], Priddy [39], Silverman and Singer [41], Singer [43, 44], Peterson [33], the present author [34]-[38], Sum [46]-[51], Walker and Wood [58, 59, 60], Wood [61, 62] and others).

Several other aspects of the hit problems were then studied by many authors. For instance, the hit problem for the symmetric of polynomials $\mathcal{P}_d^{S_d}$ as an $\mathcal{A}_2$ -submodule of $\mathcal{P}_d$, has been of interest in [14], where S_d is the symmetric group on d letters acting on the right of $\mathcal{P}_d$, and $\mathcal{P}_d^{S_d}$ is isomorphic to the cohomology of the classifying space $BO(d)$ of the orthogonal group $O(d)$. In [11, 12], Hưng and his collaborators have studied the hit problem for the rank d Dickson algebra, $\mathcal{P}_d^{GL_d}$ (the algebra of invariants, which is also an unstable $\mathcal{A}_2$ -module). In addition, the Dickson algebra is related to the dual of the Dyer-Lashof algebra (see Madsen [19]). The relationship between Kudo-Araki-May algebra and the hit problem has been investigated by Pengelley and Williams [27, 29, 31], and by Singer [44]. Recently, Zare [63] has used geometric methods to study the hit problem of Peterson [32] as well as the symmetric hit problem of Janfada and Wood [14].

Let $P_{\mathcal{A}_2}H_*(B(\mathbb{Z}/2)^{\times d})$ be the subspace of $H_*(B(\mathbb{Z}/2)^{\times d})$ consisting of all elements that are $\mathcal{A}_2^+$ -annihilated. With the idea of studying the cohomology groups of the Steenrod algebra by means of modular representations of the general linear groups, William Singer [42] constructed a transfer homomorphism of rank d from GL_d -coinvariants of the $\mathcal{A}_2^+$ -annihilated elements of the dual of $\mathcal{P}_d$ to the cohomology of the Steenrod algebra:

$$Tr_d : \mathbb{Z}/2 \otimes_{GL_d} P_{\mathcal{A}_2}H_*(B(\mathbb{Z}/2)^{\times d}) \rightarrow \text{Ext}_{\mathcal{A}_2}^{d,d+*}(\mathbb{Z}/2, \mathbb{Z}/2)$$

which is related to the geometrical transfer $tr_d : \pi_*^{\mathbb{S}}(B(\mathbb{Z}/2)_+^{\times d}) \rightarrow \pi_*^{\mathbb{S}}(\mathbb{S}^0)$ of the stable homotopy of spheres. More explicitly, tr_d induces Tr_d at the E_2 -term of the Adams spectral sequence [1]. Many mathematicians then investigated this algebraic transfer (see Boardman [4], Bruner and his collaborators [5], Chơn and Hà [7], Crossley [10], M.H. Lê [16], Hưng [13], Minami [20], Nam [26], the present author [36, 37, 38], Sum [48, 49, 51] and others). Singer [42] has indicated the importance of this transfer by proving that Tr_d is an isomorphism with $d \leq 2$ and at some degrees with $d \in \{3, 4\}$. In 1993, Boardman [4] confirmed this again by using a basis consisting of the all the classes represented by certain polynomials in $\mathcal{P}_3$ to show that Tr_3 is also an isomorphism. Therefore, the Singer transfer is a useful tool for studying the cohomology groups of the Steenrod algebra, $\text{Ext}_{\mathcal{A}_2}^{d,*}(\mathbb{Z}/2, \mathbb{Z}/2)$.

In [42], by using the invariant theory, Singer proved that Tr_5 is not an epimorphism in bidegree $(5, 5 + 9)$ and conjectured that Tr_d is a monomorphism for any positive integer d . By Boardman [4] and Singer [42], the conjecture is true for $d \leq 3$. However, it is still open in general. One of the extremely useful tools for computing the hit problem and studying Singer's transfer is the Kameko squaring operation [15] $\widetilde{Sq}_*^0 = (\widetilde{Sq}_*^0)_{(d, 2n+d)} : (Q\mathcal{P}_d)_{2n+d} \rightarrow (Q\mathcal{P}_d)_n$. This operation is an $\mathbb{Z}/2GL_d$ -homomorphism induced by the $\mathbb{Z}/2$ -linear map $\delta : (\mathcal{P}_d)_{2n+d} \rightarrow (\mathcal{P}_d)_n$, given by $\delta(x = x_1^{t_1} x_2^{t_2} \dots x_d^{t_d}) = x_1^{\frac{t_1-1}{2}} x_2^{\frac{t_2-1}{2}} \dots x_d^{\frac{t_d-1}{2}}$, if $t_1, t_2, \dots, t_d$ odd while $\delta(x) = 0$, if any of the exponents t_i are even.

In [37], we have proved that the hit problem is reduced to the case of the degree n of the form

$$(1.1) \quad n = r(2^t - 1) + 2^t s,$$

with r, t, s non-negative integers such that $1 \leq r = \mu(n) \leq d$ and $\mu(s) < r$. Here, $\mu(n)$ is the smallest number u such that $\alpha(n + u) \leq u$, where $\alpha(k)$ denotes the number of 1's in the dyadic expansion of the positive integer k . We note that if $\mu(2n + d) = d$ then Kameko's operation $\widetilde{Sq}_*^0$ is an isomorphism of $\mathbb{Z}/2(GL_d)$ -modules (see Sect.2). The hit problem was solved explicitly for the cases $d \leq 4$ (see [15, 32, 46, 47]).

For $r = d - 1$, the problem was surveyed by Crabb-Hubbuck [8], Nam [25], Repka-Selick [40], and Sum [46, 47] with $s > 0$, by Mothebe [21] and by us [34, 35] with $s = 0$. For $r = d - 2 = 3$, the problem was studied by the present author [37] with $s = 1$ and by Sum [50] with $s = 2^{m+u} + 2^m - 2$, $m \geq 0$, $u > 0$, $t \geq 6$. For $r = d = 5$, it was explicitly determined by Tín [54, 55, 56], with $s = 2, 3, 7$, by Sum [49, 51] with $s = 5, 10$, by the present author [36, 38] with $s = 6$, and by Walker-Wood [58] with $s = 26$, $t = 0$. Note that in [58], they showed that in any minimal

generating set for the $\mathcal{A}_2$ -module $\mathcal{P}_d$, there are $2^{\binom{d}{2}}$ elements in degree $2^d - d - 1$. For $d = 5$, we see that $2^5 - 6 = 26 = 5(2^0 - 1) + 26 \cdot 2^0$ and $\dim((Q\mathcal{P}_5)_{5(2^0-1)+26 \cdot 2^0}) = 2^{\binom{5}{2}} = 1024$. Further, it is easy to see that $\mu(5(2^t - 1) + 26 \cdot 2^t) = 5$ for $t > 0$. This leads to the iterated Kameko operation

$$((\widetilde{Sq}_*^0)_{(5,5(2^t-1)+26 \cdot 2^t)})^t : (Q\mathcal{P}_5)_{5(2^t-1)+26 \cdot 2^t} \rightarrow (Q\mathcal{P}_5)_{5(2^0-1)+26 \cdot 2^0}$$

is an isomorphism for every $t \geq 0$. So, one gets

$$\dim((Q\mathcal{P}_5)_{5(2^t-1)+26 \cdot 2^t}) = \dim((Q\mathcal{P}_5)_{5(2^0-1)+26 \cdot 2^0}) = 1024$$

for all $t \geq 0$. From this event, we have the following remark for the dimension of $(Q\mathcal{P}_d)_{5(2^t-1)+26 \cdot 2^t}$ in the case $d = 6$: We see that $\mu(26) = 4$ and $\alpha(26 + \mu(26)) = 4 = \mu(26)$. Then, using a result in Sum [47, Theorem 1.3], we conclude

$$\dim((Q\mathcal{P}_6)_{5(2^t-1)+26 \cdot 2^t}) = (2^6 - 1) \dim((Q\mathcal{P}_5)_{26}) = 64512 \text{ for any } t \geq 5.$$

The above event has also been studied in [57]. In addition, we also have a similar result for the degree $5(2^t - 1) + 42 \cdot 2^t$. More precisely, since $\mu(n) = 5$ for $t > 0$, the Kameko operation

$$((\widetilde{Sq}_*^0)_{(5,5(2^t-1)+42 \cdot 2^t)})^t : (Q\mathcal{P}_5)_{5(2^t-1)+42 \cdot 2^t} \rightarrow (Q\mathcal{P}_5)_{42}$$

is an isomorphism for each $t \geq 0$. This implies $\dim((Q\mathcal{P}_5)_{5(2^t-1)+42 \cdot 2^t}) = \dim((Q\mathcal{P}_5)_{42})$ for $t \geq 0$. By using a computer program of Robert R. Bruner written in MAGMA, we have $\dim((Q\mathcal{P}_5)_{42}) = 2520$. The direct proofs of this result will be published in detail elsewhere. Thus, we see that $\dim((Q\mathcal{P}_5)_{5(2^t-1)+42 \cdot 2^t}) = 2520$ for any $t \geq 0$. On the other hand, since $\mu(42) = 4$ and $\alpha(42 + \mu(42)) = 4 = \mu(42)$, by Sum [47, Theorem 1.3], we get

$$\dim((Q\mathcal{P}_6)_{5(2^t-1)+42 \cdot 2^t}) = (2^6 - 1) \dim((Q\mathcal{P}_5)_{42}) = 158760 \text{ for all } t \geq 5.$$

It is currently difficult to solve the hit problem in the general case.

In this article, based on our works in [34]-[38], we continue our study of the hit problem of five variables and by using these computational techniques, we survey Singer's algebraic transfer for the case $d = 5$ and the generic degree n of the form (1.1). More precisely, we explicitly determine a basis of $Q\mathcal{P}_d$ for $d = 5$ and the degree n of the form (1.1) with $r = d = 5$, $s = 8$ and t an arbitrary non-negative integer. A basis of this space is a set consisting of all the classes represent by the admissible monomials of degree n in $\mathcal{P}_d$. We refer to Sect.2 for the definition of the admissible monomial. Using this result combining with the computations of Chen [6], Lin [17], Tangora [52] on the cohomology groups $\text{Ext}_{\mathcal{A}_2}^{5,13 \cdot 2^t}(\mathbb{Z}/2, \mathbb{Z}/2)$, and a representation in the $\mathbb{Z}/2$ -algebra of Tr_5 , we show that the fifth Singer transfer is an isomorphism in bidegree $(5, (13 \cdot 2^t - 5) + 5)$ for $t \in \{0, 1\}$. (See the below information about the lambda algebra.) This gives a quite efficient method to study the isomorphism of the fifth transfer in some internal degrees of the form (1.1) and it is different from the ones of Boardman [4], and of Singer [42]. The following is our first main result.

Theorem 1.1. *Let $n = 13 \cdot 2^t - 5 = 5(2^t - 1) + 8 \cdot 2^t$ with t a non-negative integer. Then, we have*

$$\dim(Q\mathcal{P}_5)_{13 \cdot 2^t - 5} = \begin{cases} 174 & \text{if } t = 0, \\ 840 & \text{if } t = 1, \\ 1894 & \text{if } t \geq 2. \end{cases}$$

We prove the theorem by determining all the admissible monomials of degree n in the $\mathcal{A}_2$ -module $\mathcal{P}_5$. The first idea for the proof of the theorem is from Kameko's squaring operation [15]. More clearly, since $13 \cdot 2^t - 5 = 2^{t+3} + 2^{t+2} + 2^{t-1} + 2^{t-2} + 2^{t-2} - 5$, $\mu(13 \cdot 2^t - 5) = 5$, for $t > 2$. This leads to the iterated Kameko operation $(\widetilde{Sq}_*^0)_{(5,13 \cdot 2^t - 5)}^{t-2} : (Q\mathcal{P}_5)_{13 \cdot 2^t - 5} \rightarrow (Q\mathcal{P}_5)_{13 \cdot 2^2 - 5}$ is an isomorphism of $\mathbb{Z}/2(GL_5)$ -modules for any $t \geq 2$. From this we need only to study the structure

of $(Q\mathcal{P}_5)_{13,2^t-5}$ for $t = 0, 1, 2$. The case $t = 0$ has been computed by Tín [53]. For $t \in \{1, 2\}$, we see that $(\widetilde{Sq}_*^0)_{(5,13,2^t-5)} : (Q\mathcal{P}_5)_{13,2^t-5} \rightarrow (Q\mathcal{P}_5)_{13,2^t-1-5}$ is an epimorphism. In these cases, we need only to determine the kernel of $(\widetilde{Sq}_*^0)_{(5,13,2^t-5)}$. To study this space, we combine our recent results in [34] with previous results by Kameko [15], Mothebe [22, 24], Singer [43], Sum [47], and Tín [53, 55].

Recently, in [49], Sum has proved some properties on the $\mathcal{A}_2$ -generators for $\mathcal{P}_d$. Then, he made a conjecture on the relation between the admissible monomials for the polynomial algebras (see Sect.3.3). The conjecture helps us to reduce remarkably in computing the hit problem. From the results of Peterson [32], Kameko [15] and Sum [47], this conjecture is true for $d \leq 4$. Sum proved in [49] that the conjecture is true for $d = 5$ and the degree n of the form (1.1) with $(r; s) = (5; 10)$ and $t \geq 0$. Based on the proof of Theorem 1.1 in Sect.3.4 and our results in [34]–[37] and Sum [50], the conjecture is also true for $d = 5$ and the degrees of the form (1.1) with $(r; s) = (4; 0), (5; 6), (5; 8)$ and $(3; s)$, for $s = 1, t > 0, s = 2^{m+u} + 2^m - 2, m \geq 0, u > 0, t \geq 6$.

For $t = 1$, we have $n = 13 \cdot 2^1 - 5 = 21$. Applying Theorem 1.1, we get the following, which is our second main result.

Theorem 1.2. *There exists uniquely a non-zero class in $(Q\mathcal{P}_5)_{21}$ invariant under the action of the group GL_5 .*

Note that the Kameko operation $(\widetilde{Sq}_*^0)_{(5,21)} : (Q\mathcal{P}_5)_{21} \rightarrow (Q\mathcal{P}_5)_8$ is an epimorphism of $\mathbb{Z}/2GL_5$ -modules. So, in order to prove Theorem 1.2, we describe the $\mathbb{Z}/2$ -vector space structure of $(Q\mathcal{P}_5)_8^{GL_5}$. Combining this with a basis of $(QP_5)_{21}$ in Theorem 1.1, we explicitly compute all GL_5 -invariants of $Q\mathcal{P}_5$ in degree 21.

The cohomology of Steenrod algebra, $\text{Ext}_{\mathcal{A}_2}^{*,*}(\mathbb{Z}/2, \mathbb{Z}/2)$, is described in term of the mod 2 lambda algebra Λ (see Lin [17]). Recall that Λ is a bigraded differential algebra over $\mathbb{Z}/2$ generated by $\lambda_j \in \Lambda^{1,j}, j \geq 0$, with the Adem relations $\lambda_j \lambda_{2j+1+m} = \sum_{\nu \geq 0} \binom{m-\nu-1}{\nu} \lambda_{j+m-\nu} \lambda_{2j+1+\nu}$, for $m \geq 0$, and the differential $\partial(\lambda_s) = \sum_{\nu \geq 0} \binom{s-\nu-1}{\nu+1} \lambda_{s-\nu-1} \lambda_\nu$, for $s > 0$. Further, $H^{s,t}(\Lambda, \partial) \cong \text{Ext}_{\mathcal{A}_2}^{s,t}(\mathbb{Z}/2, \mathbb{Z}/2)$. As well known, the dual of $\mathcal{P}_d = \mathbb{Z}/2[x_1, \dots, x_d]$ is isomorphic to $\Gamma(a_1, \dots, a_d)$, the divided power algebra generated by $a_1, \dots, a_d$, where a_t is dual to $x_t \in \mathcal{P}_d$ with respect to the basis of $\mathcal{P}_d$ consisting of all monomials in $x_1, \dots, x_d$. So, $H_*(B(\mathbb{Z}/2)^{\times d}) \cong \Gamma(a_1, \dots, a_d)$. The algebra $H_*(B(\mathbb{Z}/2)^{\times d})$ has a right $\mathcal{A}_2$ -module structure. The right action of $\mathcal{A}_2$ on $H_*(B(\mathbb{Z}/2)^{\times d})$ is given by $(a_t^{(s)})Sq^n = \binom{s-n}{n} a_t^{(s-n)}$ and Cartan's formula. In [7], Chơn and Hà have constructed the $\mathbb{Z}/2$ -linear map $\psi_d : H_*(B(\mathbb{Z}/2)^{\times d}) \longrightarrow \Lambda_d$, which is determined by the following inductive formula:

$$\psi_d(a^{(I,s)}) = \begin{cases} \lambda_s & \text{if } d-1 = \ell(I) = 0, \\ \sum_{t \geq s} \psi_{d-1}(a^I Sq^{t-s}) \lambda_t & \text{if } d-1 = \ell(I) > 0, \end{cases}$$

for any $a^{(I,s)} = a_1^{(n_1)} \dots a_{d-1}^{(n_{d-1})} a_d^{(s)} \in H_*(B(\mathbb{Z}/2)^{\times d})$ and $I = (n_1, n_2, \dots, n_{d-1})$. This map is a presentation in the $\mathbb{Z}/2$ -lambda algebra of the transfer homomorphism Tr_d . Note that ψ_d is not an $\mathcal{A}_2$ -homomorphism. Then, we have the homomorphism of algebras

$$\psi = \oplus_{d \geq 1} \psi_d : \oplus_{d \geq 1} H_*(B(\mathbb{Z}/2)^{\times d}) \longrightarrow \oplus_{d \geq 1} \Lambda_d = \Lambda,$$

which induces the Singer transfer. Chơn and Hà showed in [7] that if $u \in P_{\mathcal{A}_2} H_*(B(\mathbb{Z}/2)^{\times d})$, then $\psi_d(u)$ is a cycle in Λ and $Tr_d([u]) = [\psi_d(u)]$. Combining this with Theorem 1.2, and by a direct computation using a representation in the $\mathbb{Z}/2$ -lambda algebra of Tr_5 , we obtain the following theorem, which is our third main result.

Theorem 1.3. *The fifth transfer homomorphism*

$$Tr_5 : \mathbb{Z}/2 \otimes_{GL_5} P_{\mathcal{A}_2} H_{21}(B(\mathbb{Z}/2)^{\times 5}) \rightarrow \text{Ext}_{\mathcal{A}_2}^{5,5+21}(\mathbb{Z}/2, \mathbb{Z}/2)$$

is an isomorphism.

We know that the classical squaring operations

$$Sq^0 : \text{Ext}_{\mathcal{A}_2}^{d,d+*}(\mathbb{Z}/2, \mathbb{Z}/2) \rightarrow \text{Ext}_{\mathcal{A}_2}^{d,2(d+*)}(\mathbb{Z}/2, \mathbb{Z}/2)$$

acting on the cohomology groups of $\mathcal{A}_2$ (see [18]), but they are not the identity maps. As above mentioned, the structure of the cohomology groups $\text{Ext}_{\mathcal{A}_2}^{d,*}(\mathbb{Z}/2, \mathbb{Z}/2)$ has intensively been studied for $d \leq 5$, but remains very mysterious in general. A family $\{a_t : t \geq 0\} \subset \text{Ext}_{\mathcal{A}_2}^{d,*}(\mathbb{Z}/2, \mathbb{Z}/2)$ is called a Sq^0 -family if $a_t = (Sq^0)^t(a_0)$ for $t \geq 0$. According to Lin [17] and Chen [6], we have

$$\text{Ext}_{\mathcal{A}_2}^{5,13 \cdot 2^t}(\mathbb{Z}/2, \mathbb{Z}/2) = \begin{cases} 0 & \text{if } t = 0, \\ \langle h_t g_t \rangle & \text{if } t \geq 1, \end{cases}$$

and $h_t g_t = h_{t+1} f_{t-1} \neq 0$, where h_t is the Adams element in $\text{Ext}_{\mathcal{A}_2}^{1,2^t}(\mathbb{Z}/2, \mathbb{Z}/2)$, and $g_t = (Sq^0)^{t-1}(g_1)$, $f_{t-1} = (Sq^0)^{t-1}(f_0)$ are non-zero in $\text{Ext}_{\mathcal{A}_2}^{4,12 \cdot 2^t}(\mathbb{Z}/2, \mathbb{Z}/2)$ and $\text{Ext}_{\mathcal{A}_2}^{4,11 \cdot 2^t}(\mathbb{Z}/2, \mathbb{Z}/2)$, respectively, for any $t \geq 1$. Now, for $t = 0$, we have $n = 13 \cdot 2^t - 5 = 8$. Based on the computations of the GL_5 -invariants of $(Q\mathcal{P}_5)_8$ (see Sect. 4.1), we see that the space $\mathbb{Z}/2 \otimes_{GL_5} P_{\mathcal{A}_2} H_8(B(\mathbb{Z}/2)^{\times 5})$ is trivial. From this and $\text{Ext}_{\mathcal{A}_2}^{5,13}(\mathbb{Z}/2, \mathbb{Z}/2) = 0$, we conclude that the fifth transfer

$$Tr_5 : \mathbb{Z}/2 \otimes_{GL_5} P_{\mathcal{A}_2} H_8(B(\mathbb{Z}/2)^{\times 5}) \rightarrow \text{Ext}_{\mathcal{A}_2}^{5,5+8}(\mathbb{Z}/2, \mathbb{Z}/2)$$

is also an isomorphism. As it is known, the transfer Tr_1 detects the family $\{h_t | t \geq 1\}$ (see Singer [42]). In [26], Nam proved that Tr_4 detects the family $\{f_{t-1} | t \geq 1\}$. Note that the family $\{g_t | t \geq 1\}$ was not detected by Tr_4 (see Bruner-Hà-Hung [5]). Combining these data with an event that $Tr = \bigoplus_{d \geq 0} Tr_d$ is the homomorphism of $\mathbb{Z}/2$ -algebras (see [42]), we conclude that $h_t g_t$ is in the image of Tr_5 . In Sect. 5, we give another direct proof of this event for the case $t = 1$. More specifically, we proved $h_1 g_1 \in \text{Im}(Tr_5)$ by a direct computation using Theorem 1.2 and a representation in the $\mathbb{Z}/2$ -lambda algebra of Tr_5 .

From Theorem 1.3 and the above data, we have immediately

Corollary 1.4. *Singer's conjecture for Tr_5 is true in bidegrees $(5, 5 + 8)$ and $(5, 5 + 21)$.*

We end this introduction with a short discussion on Singer's conjecture for Tr_5 in the bidegree $(5, (13 \cdot 2^t - 5) + 5)$ with $t \geq 2$. Recall that the iterated Kameko operation $((\widetilde{Sq}_*^0)_{(5,13 \cdot 2^t - 5)})^{t-2} : (Q\mathcal{P}_5)_{13 \cdot 2^t - 5} \rightarrow (Q\mathcal{P}_5)_{47=13 \cdot 2^2 - 5}$ is an isomorphism of GL_5 -modules for every $t \geq 2$. So, to verify Singer's conjecture for Tr_5 in the above bidegree, we need only to determine all GL_5 -invariants of $(Q\mathcal{P}_5)_{13 \cdot 2^t - 5}$ for $t = 2$.

It is well known that Kameko's operation $(\widetilde{Sq}_*^0)_{(5,47)} : (Q\mathcal{P}_5)_{47} \rightarrow (Q\mathcal{P}_5)_{21}$ is an epimorphism of GL_5 -modules and the element $h_2 g_2 = h_3 f_1 \in \text{Im}(Tr_5)$. So, by Theorem 1.2, we get

$$1 \leq \dim \left(\mathbb{Z}/2 \otimes_{GL_5} P_{\mathcal{A}_2} H_{47}(B(\mathbb{Z}/2)^{\times 5}) \right) \leq \dim(\text{Ker}(\widetilde{Sq}_*^0)_{(5,47)})^{GL_5} + 1.$$

Furthermore, all elements of $\mathbb{Z}/2 \otimes_{GL_5} P_{\mathcal{A}_2} H_{47}(B(\mathbb{Z}/2)^{\times 5})$ are of the form $(\gamma[\varphi(u_0)] + [v])^*$, where $\gamma \in \mathbb{Z}/2$; $\varphi : \mathcal{P}_5 \rightarrow \mathcal{P}_5$, which is determined by $\varphi(u) = x_1 x_2 x_3 x_4 x_5 u^2$ for any $u \in \mathcal{P}_5$; $f \in (\mathcal{P}_5)_{47}$ such that $[f] \in \text{Ker}(\widetilde{Sq}_*^0)_{(5,47)}$, and $u_0 \in (\mathcal{P}_5)_{21}$. By Theorem 1.2, $[u_0]$ is the only non-zero element in $(Q\mathcal{P}_5)_{21}^{GL_5}$. Direct calculating the elements $(\gamma[\varphi(u_0)] + [v])^*$ is a hard work. By using techniques of the hit problem of five variables, we will describe explicitly all these elements in the near future.

From the above events and $h_t g_t \in \text{Im}(Tr_5)$ for $t \geq 1$, we see that if $(\text{Ker}(\widetilde{Sq}_*^0)_{(5,47)})^{GL_5} = 0$, then Singer's conjecture for Tr_5 is also true at the bidegree $(5, (13 \cdot 2^t - 5) + 5)$ with $t \geq 2$. However, it will be much more interesting if $(\text{Ker}(\widetilde{Sq}_*^0)_{(5,47)})^{GL_5} \neq 0$ and $\dim(Q\mathcal{P}_5)_{47}^{GL_5} \neq 1$.

Our work is organized as follows: First, some information about Kameko's squaring operation $\widetilde{Sq}_*^0$, and the "hit" problem of Peterson is given in the section of preliminaries (see Sect. 2). In the

next section, the generators of the $\mathcal{A}_2$ -module $\mathcal{P}_5$ in degree $n = 13 \cdot 2^t - 5$ are described explicitly by proving Theorem 1.1 (see Sect.3). The proof of Theorem 1.2 is presented in Sect.4 by using the results in Sect.3. In Sect.5, we prove Theorem 1.3. Finally, the Appendix describes all the admissible monomials of degrees 8, 21, 22, and 47 in the $\mathcal{A}_2$ -module $\mathcal{P}_5$.

Acknowledgments. This work is supported financially by the National Foundation for Science and Technology Development (NAFOSTED) of Viet Nam under Grant No. 101.04-2017.05.

The author thanks the anonymous referees for their valuable remarks which were incorporated into this revised version. I would also like to thank Robert R. Bruner, Phan H. Chon, Mbakiso F. Mothebe for helpful conversations related to this work, Hadi Zare for sending a copy of [63], and Le M. Ha for pointing out an error in the original manuscript.

Finally, the author would like to give my deepest sincere thanks to Professor Nguyen Sum, my PhD thesis advisor, for introducing me to the wonderful subject of the hit problem, for encouraging me to think about the relationship between the hit problem and the algebraic transfer, and for many helpful discussions.

2. Preliminaries

This section starts with a recollection of the Kameko homomorphism and some results related to the Peterson hit problem.

2.1. Kameko's squaring operation

Denote by $GL_d := GL(d, \mathbb{Z}/2)$ the general linear group of rank d over the field $\mathbb{Z}/2$. This GL_d left acts on $\mathcal{P}_d$ according to the following standard action

$$(w(f))(x_1, x_2, \dots, x_d) = f(w(x_1), w(x_2), \dots, w(x_d)),$$

where $w = (w_{ij}) \in GL_d$ and $w(x_j) = \sum_{i=1}^d w_{ij}x_i$, $1 \leq j \leq d$. Then, $\mathcal{P}_d$ has a left GL_d -module structure. Since the two actions of $\mathcal{A}_2$ and GL_d upon $\mathcal{P}_d$ commute with each other, there is an inherited action of GL_d on $Q\mathcal{P}_d$.

For n a positive integer, let $(Q\mathcal{P}_d)_n^{GL_d}$ be the subspace of $Q\mathcal{P}_d$ generated all GL_d -invariants of degree n . Then, we have the Kameko squaring operation

$$\widetilde{Sq}^0 : \mathbb{Z}/2 \otimes_{GL_d} P_{\mathcal{A}_2} H_n(B(\mathbb{Z}/2)^{\times d}) \rightarrow \mathbb{Z}/2 \otimes_{GL_d} P_{\mathcal{A}_2} H_{d+2n}(B(\mathbb{Z}/2)^{\times d}).$$

It is well known that this operation commutes with the classical squaring operation

$$Sq^0 : \text{Ext}_{\mathcal{A}_2}^{d, d+n}(\mathbb{Z}/2, \mathbb{Z}/2) \rightarrow \text{Ext}_{\mathcal{A}_2}^{d, 2d+2n}(\mathbb{Z}/2, \mathbb{Z}/2)$$

through the Singer transfer Tr_d (see [2, 4, 20]). This means that the following diagram is commutative:

$$\begin{array}{ccc} \mathbb{Z}/2 \otimes_{GL_d} P_{\mathcal{A}_2} H_n(B(\mathbb{Z}/2)^{\times d}) & \xrightarrow{Tr_d} & \text{Ext}_{\mathcal{A}_2}^{d, d+n}(\mathbb{Z}/2, \mathbb{Z}/2) \\ \downarrow \widetilde{Sq}^0 & & \downarrow Sq^0 \\ \mathbb{Z}/2 \otimes_{GL_d} P_{\mathcal{A}_2} H_{d+2n}(B(\mathbb{Z}/2)^{\times d}) & \xrightarrow{Tr_d} & \text{Ext}_{\mathcal{A}_2}^{d, 2d+2n}(\mathbb{Z}/2, \mathbb{Z}/2). \end{array}$$

Note that Sq^0 is completely different from the identity map. Recall that the map

$$\widetilde{Sq}_*^0 : (Q\mathcal{P}_d)_{d+2n}^{GL_d} \rightarrow (Q\mathcal{P}_d)_n^{GL_d}$$

is dual to $\widetilde{Sq^0}$, and it is induced by the homomorphism $(\widetilde{Sq_*^0})_{(d,d+2n)} : (Q\mathcal{P}_d)_{d+2n} \rightarrow (Q\mathcal{P}_d)_n$. The latter is given by the $\mathbb{Z}/2$ -linear map $\delta : (\mathcal{P}_d)_{d+2n} \rightarrow (\mathcal{P}_d)_n$, determined by

$$\delta(x) = \begin{cases} u & \text{if } u = x_1 x_2 \dots x_d u^2, \\ 0 & \text{otherwise,} \end{cases}$$

for any $x \in (\mathcal{P}_d)_{d+2n}$. Thus, we see that $(\widetilde{Sq_*^0})_{(d,d+2n)}$ is an epimorphism of $\mathbb{Z}/2(GL_d)$ -modules.

For a positive integer n , denote by $\alpha(n)$ the number of 1's in dyadic expansion of n . We consider the following arithmetic function (see [51]):

$$\begin{aligned} \mu(n) &= \min\{u \in \mathbb{N} : \alpha(n+u) \leq u\} \\ &= \min\{u \in \mathbb{N} : n = \sum_{1 \leq j \leq u} (2^{d_j} - 1), d_j > 0, 1 \leq j \leq u\}. \end{aligned}$$

By Kameko [15], we have the following result:

Theorem 2.1.1 (see [15]). *If $\mu(d+2n) = d$, then the homomorphism*

$$(\widetilde{Sq_*^0})_{(d,d+2n)} : (Q\mathcal{P}_d)_{d+2n} \rightarrow (Q\mathcal{P}_d)_n$$

is an isomorphism of $\mathbb{Z}/2(GL_d)$ -modules.

According to Hưng [13], if Singer's conjecture that Tr_d is a monomorphism for every d is true, then the algebraic transfer does not detect $\text{Ker}(\widetilde{Sq^0})$ in degrees n such that $\mu(n) = d$. This leads us to the study of the kernel of $\widetilde{Sq^0}$ and its dual. Obviously, the map $\widetilde{Sq^0} : P_{A_2} H_n(B(\mathbb{Z}/2)^{\times d}) \rightarrow P_{A_2} H_{2d+n}(B(\mathbb{Z}/2)^{\times d})$ is injective. However, the Kameko squaring operation $\widetilde{Sq^0} = id_{\mathbb{Z}/2} \otimes_{GL_d} \widetilde{Sq^0}$ is not a monomorphism in general (see Hưng [13]). Thereafter, Hưng [13] conjectured that $\widetilde{Sq^0}$ is a monomorphism in positive degrees of $\mathbb{Z}/2 \otimes_{GL_d} P_{A_2} H_n(B(\mathbb{Z}/2)^{\times d})$ if and only if $d \leq 4$. It can be verified by using the dual of $\widetilde{Sq^0}$ and the results on the hit problem in [15, 32, 49].

2.2. On the Peterson "hit" problem

To study the hit problem, we need to some relevant notations and definitions. For a natural number n , denote by $\alpha_t(n)$ the t -th coefficients in dyadic expansion of n . This means $\alpha(n) = \sum_{t \geq 0} \alpha_t(n)$. Further, n can be represented as follows: $n = \sum_{t \geq 0} \alpha_t(n) 2^t$, where $\alpha_t(n) \in \{0, 1\}$, $t = 0, 1, \dots$. Let $x = x_1^{u_1} x_2^{u_2} \dots x_d^{u_d}$ be a monomial in $\mathcal{P}_d$. We define two sequences associated with x by $\omega(x) = (\omega_1(x), \omega_2(x), \dots, \omega_t(x), \dots)$ and $\xi(x) = (u_1, u_2, \dots, u_d)$, where $\omega_t(x) = \sum_{1 \leq j \leq d} \alpha_{t-1}(u_j)$. The sequences $\omega(x)$ and $\xi(x)$ are called the *weight vector* and the *exponent vector* of x , respectively.

Let $\omega = (\omega_1, \omega_2, \dots, \omega_t, \dots)$ be a sequence of non-negative integers. The sequence ω are called the weight vector if $\omega_t = 0$ for $t \gg 0$. The sets of all the weight vectors and the exponent vectors are given the left lexicographical order.

For a weight vector ω , we define $\deg \omega = \sum_{t \geq 1} 2^{t-1} \omega_t$. Denote by $\mathcal{P}_d(\omega)$ the subspace of $\mathcal{P}_d$ spanned by all monomials $x \in \mathcal{P}_d$ such that $\deg x = \deg \omega$, $\omega(x) \leq \omega$, and by $\mathcal{P}_d^-(\omega)$ the subspace of $\mathcal{P}_d$ spanned by all monomials $x \in \mathcal{P}_d$ such that $\deg x = \deg \omega$, $\omega(x) < \omega$.

Recall that a homogeneous element f of degree n in $\mathcal{P}_d$ is A_2 -decomposable (or "hit") if there is a finite sum $f = \sum_{k > 0} Sq^k(f_k)$, where the homogeneous elements $f_k \in \mathcal{P}_d$ have grading strictly less than n . This means that f belong to $A_2^+ \mathcal{P}_d$.

The equivalence relations on $\mathcal{P}_d$ (see [15, 37]). Let ω be a weight vector and f, g two homogeneous polynomials of the same degree in $\mathcal{P}_d$. We define the binary relations " $\equiv$ " and " $\equiv_\omega$ " on $\mathcal{P}_d$.

- (i) $f \equiv g$ if and only if $f = g$ modulo $(A_2^+ \mathcal{P}_d)$. If $f \equiv 0$ then f is A_2 -decomposable.
- (ii) $f \equiv_\omega g$ if and only if $f, g \in \mathcal{P}_d(\omega)$ and $f = g$ modulo $((A_2^+ \mathcal{P}_d \cap \mathcal{P}_d(\omega)) + \mathcal{P}_d^-(\omega))$.

These binary relations are equivalence ones. Denote by $Q\mathcal{P}_d(\omega)$ the quotient of $\mathcal{P}_d(\omega)$ by the equivalence relation " $\equiv_\omega$ ". Then, we have the $\mathbb{Z}/2$ -quotient space

$$Q\mathcal{P}_d(\omega) = \mathcal{P}_d(\omega) / ((\mathcal{A}_2^+ \mathcal{P}_d \cap \mathcal{P}_d(\omega)) + \mathcal{P}_d^-(\omega)),$$

By Sum [51], $Q\mathcal{P}_d(\omega)$ is also an GL_d -module. The following events are shown in [37]. However, to make the paper self-contained, we will present again them in detail.

$$\dim((Q\mathcal{P}_d)_n) = \sum_{\deg \omega = n} \dim(Q\mathcal{P}_d(\omega)), \quad \dim((Q\mathcal{P}_d)_n^{GL_d}) \leq \sum_{\deg(\omega) = n} \dim(Q\mathcal{P}_d(\omega)^{GL_d}).$$

Indeed, by Walker and Wood [60], we have a filtration of $Q\mathcal{P}_d$:

$$\{0\} \subseteq \cdots \subseteq \mathcal{P}_d^-(\omega) / ((\mathcal{A}_2^+ \mathcal{P}_d \cap \mathcal{P}_d^-(\omega))) \subseteq \mathcal{P}_d(\omega) / ((\mathcal{A}_2^+ \mathcal{P}_d \cap \mathcal{P}_d(\omega))) \subseteq \cdots \subseteq \mathcal{P}_d / (\mathcal{A}_2^+ \mathcal{P}_d) = Q\mathcal{P}_d.$$

This is not only a filtration of $Q\mathcal{P}_d$ as a vector space, but also as a GL_d -module. The inclusion of $\mathcal{P}_d^-(\omega)$ into $\mathcal{P}_d(\omega)$ induces the monomorphism

$$\mathcal{P}_d^-(\omega) / ((\mathcal{A}_2^+ \mathcal{P}_d \cap \mathcal{P}_d^-(\omega))) \rightarrow \mathcal{P}_d(\omega) / ((\mathcal{A}_2^+ \mathcal{P}_d \cap \mathcal{P}_d(\omega)))$$

and the following sequence is short exact:

$$0 \rightarrow \mathcal{P}_d^-(\omega) / ((\mathcal{A}_2^+ \mathcal{P}_d \cap \mathcal{P}_d^-(\omega))) \rightarrow \mathcal{P}_d(\omega) / ((\mathcal{A}_2^+ \mathcal{P}_d \cap \mathcal{P}_d(\omega))) \rightarrow \mathcal{P}_d(\omega) / (((\mathcal{A}_2^+ \mathcal{P}_d \cap \mathcal{P}_d(\omega)) + \mathcal{P}_d^-(\omega))) \rightarrow 0.$$

From this, we get

$$Q\mathcal{P}_d(\omega) \cong (\mathcal{P}_d(\omega) / ((\mathcal{A}_2^+ \mathcal{P}_d \cap \mathcal{P}_d(\omega))) / (\mathcal{P}_d^-(\omega) / ((\mathcal{A}_2^+ \mathcal{P}_d \cap \mathcal{P}_d^-(\omega)))).$$

This isomorphism is also an isomorphism of GL_d -modules. Combining this and the filtration of $Q\mathcal{P}_d$, we conclude the above statements.

The linear order on $\mathcal{P}_d$ (see [15]). Let u, v be monomials of the same degree in $\mathcal{P}_d$. We say that $u < v$ if and only if one of the following holds:

- (i) $\omega(u) < \omega(v)$;
- (ii) $\omega(u) = \omega(v)$ and $\xi(u) < \xi(v)$.

The inadmissible monomial (see [15]). A monomial u is said to be *inadmissible* if there exist monomials $x_1, x_2, \dots, x_k$ such that $x_j < u$ for $1 \leq j \leq k$ and $u \equiv \sum_{1 \leq i \leq k} x_i$. Then, u is said to be *admissible* if it is not inadmissible.

Note that the set of all the admissible monomials of degree n in $\mathcal{P}_d$ is a *minimal set of $\mathcal{A}_2$ -generators for $\mathcal{P}_d$ in degree n* .

The strictly inadmissible monomial (see [15]). A monomial u is said to be *strictly inadmissible* if and only if there exist monomials $x_1, x_2, \dots, x_k$ such that $x_j < u$ for $1 \leq j \leq k$ and $u = \sum_{1 \leq j \leq k} x_j + \sum_{1 \leq t \leq 2^s - 1} Sq^t(y_t)$ with $s = \max\{i \in \mathbb{Z} : \omega_i(u) > 0\}$ and suitable polynomials $y_t \in \mathcal{P}_d$.

We see that if a monomial is strictly inadmissible, then it is inadmissible. The following result is used to study $Q\mathcal{P}_5$ in the next section.

Theorem 2.2.1 (see [15]). *Let x, y and u be monomials in $\mathcal{P}_d$ such that $\omega_i(x) = 0$ for $i > r > 0$, $\omega_t(u) \neq 0$ and $\omega_i(u) = 0$ for $i > t > 0$. Then, if u is inadmissible, then xu^{2^r} is also inadmissible. Furthermore, if u is strictly inadmissible, then uy^{2^t} is also strictly inadmissible.*

For a natural number d , we now denote

$$\begin{aligned} \mathcal{P}_d^0 &= \langle x_1^{t_1} x_2^{t_2} \dots x_d^{t_d} \in \mathcal{P}_d \mid t_1 t_2 \dots t_d = 0 \rangle, \\ \mathcal{P}_d^+ &= \langle x_1^{t_1} x_2^{t_2} \dots x_d^{t_d} \in \mathcal{P}_d \mid t_1 t_2 \dots t_d > 0 \rangle. \end{aligned}$$

Then $\mathcal{P}_d^0$ and $\mathcal{P}_d^+$ are the $\mathcal{A}_2$ -submodules of $\mathcal{P}_d$. Furthermore, we have a direct summand decomposition of the $\mathbb{Z}/2$ -vector spaces

$$Q\mathcal{P}_d = Q\mathcal{P}_d^0 \oplus Q\mathcal{P}_d^+,$$

where $Q\mathcal{P}_d^0 := \mathbb{Z}/2 \otimes_{\mathcal{A}_2} \mathcal{P}_d^0$ and $Q\mathcal{P}_d^+ := \mathbb{Z}/2 \otimes_{\mathcal{A}_2} \mathcal{P}_d^+$.

We end this section by setting up a formula on the dimension of $Q\mathcal{P}_d^0$ in degree n , which will be used in the next section. Note that this formula is similar to the one of Mothebe-Kaelo-Ramatebele [24].

Let $\mathcal{I} = (j_1, j_2, \dots, j_r)$, with $1 \leq j_1 < \dots < j_r \leq d$, $1 \leq r \leq d-1$, and let $r := \ell(\mathcal{I})$ be the length of $\mathcal{I}$. We denote $\mathcal{P}_{\mathcal{I}} = \langle \{x_{j_1}^{t_1} x_{j_2}^{t_2} \dots x_{j_r}^{t_r} \mid t_s \in \mathbb{N}, s = 1, 2, \dots, r\} \rangle \subset \mathcal{P}_d$. Then, $\mathcal{P}_{\mathcal{I}}$ is $\mathcal{A}_2$ -submodule of $\mathcal{P}_d$. Further, it is isomorphic to $\mathcal{P}_r$. By a simple computation, one gets

$$Q\mathcal{P}_d^0 = \bigoplus_{1 \leq r \leq d-1} \bigoplus_{\ell(\mathcal{I})=r} Q\mathcal{P}_{\mathcal{I}}^+,$$

where $\mathcal{P}_{\mathcal{I}}^+ = \langle \{x_{j_1}^{t_1} x_{j_2}^{t_2} \dots x_{j_r}^{t_r} \in \mathcal{P}_{\mathcal{I}} \mid t_1 t_2 \dots t_r > 0, 1 \leq r \leq d-1\} \rangle$. It can be seen that $\dim(Q\mathcal{P}_{\mathcal{I}}^+)_n = \dim(Q\mathcal{P}_r^+)_n$ for all n , and that $\binom{d}{r}$ is the number of the sequences $\mathcal{I}$ of length r . Therefore, we get

$$\dim(Q\mathcal{P}_d^0)_n = \sum_{1 \leq r \leq d-1} \binom{d}{r} \dim(Q\mathcal{P}_r^+)_n.$$

3. Generators of the $\mathcal{A}_2$ -module $\mathcal{P}_5$ in the generic degree $13.2^t - 5$

In this section, we prove Theorem 1.1. More precisely, we explicitly compute the admissible monomials of degree $13.2^t - 5$ in the $\mathcal{A}_2$ -module $\mathcal{P}_5$. We first review some homomorphisms and related results, Sum's conjecture [49] and Singer's criterion on the $\mathcal{A}_2$ -decomposable [42].

3.1. Some homomorphisms

For $1 \leq k \leq d$, define the homomorphism $\rho_{(k,d)} : \mathcal{P}_{d-1} \rightarrow \mathcal{P}_d$ of algebras by substituting

$$\rho_{(k,d)}(x_j) = \begin{cases} x_j & \text{if } 1 \leq j < k, \\ x_{j+1} & \text{if } k \leq j < d. \end{cases}$$

We now denote

$$\mathcal{N}_d := \{(k; \mathcal{K}) \mid \mathcal{K} = (k_1, k_2, \dots, k_r), 1 \leq k < k_1 < k_2 < \dots < k_r \leq d, 0 \leq r < d\}.$$

Here, by convention, $\mathcal{K} = \emptyset$, if $r = 0$. Denote by $r = \ell(\mathcal{K})$ the length of $\mathcal{K}$.

For any $(k; \mathcal{K}) \in \mathcal{N}_d$, we define the homomorphism $\pi_{(k; \mathcal{K})} : \mathcal{P}_d \rightarrow \mathcal{P}_{d-1}$ of algebras by substituting

$$\pi_{(k; \mathcal{K})}(x_j) = \begin{cases} x_j & \text{if } 1 \leq j < k, \\ \sum_{p \in \mathcal{K}} x_{p-1} & \text{if } j = k, \\ x_{j-1} & \text{if } k < j \leq d. \end{cases}$$

Note that $\rho_{(k,d)}$ and $\pi_{(k; \mathcal{K})}$ are also the homomorphisms of the $\mathcal{A}_2$ -modules. In particular, we have $\pi_{(k; \emptyset)}(x_k) = 0$ for $1 \leq k \leq d$ and $\pi_{(k; \mathcal{K})}(\rho_{(k,d)}(u)) = u$ for any $u \in \mathcal{P}_{d-1}$.

Proposition 3.1.1 (see [34]). *If x is a monomial in $\mathcal{P}_d$, then $\pi_{(k; \mathcal{K})}(x) \in \mathcal{P}_{d-1}(\omega(x))$.*

This result implies that if ω is a weight vector and $x \in \mathcal{P}_d(\omega(x))$, then $\pi_{(k; \mathcal{K})}(x) \in \mathcal{P}_{d-1}(\omega)$. Furthermore, $\pi_{(k; \mathcal{K})}$ passes to a homomorphism from $Q\mathcal{P}_d(\omega)$ to $Q\mathcal{P}_{d-1}(\omega)$.

From now on, we use the following notations: For a natural number d , we set

$$\begin{aligned} \Gamma_d &= \{1, 2, \dots, d\}, \\ X_{(S,d)} &= X_{(\{s_1, s_2, \dots, s_r\}, d)} = \prod_{s \in \Gamma_d \setminus S} x_s, \quad S = \{s_1, s_2, \dots, s_r\} \subseteq \Gamma_d. \end{aligned}$$

In particular,

$$\begin{aligned} X_{\Gamma_d} &= 1, \quad X_{(\emptyset, d)} = x_1 x_2 \dots x_d, \\ X_{(\{s\}, d)} &= x_1 \dots \hat{x}_s \dots x_d, \quad 1 \leq s \leq d. \end{aligned}$$

Let $X = x_1^{t_1} x_2^{t_2} \dots x_d^{t_d} \in \mathcal{P}_d$. Set $S_j(X) = \{s \in \Gamma_d : \alpha_j(t_s) = 0\}$, for $j \geq 0$. Then we have

$$X = \prod_{j \geq 0} X_{(S_j(X), d)}^{2^j}.$$

Definition 3.1.2 (see [47]). Let $(k; \mathcal{K}) \in \mathcal{N}_d$, $0 < r < d$. Denote

$$x_{(\mathcal{K}, u)} = x_{k_u}^{2^{r-1} + 2^{r-2} + \dots + 2^{r-u}} \prod_{u < m \leq r} x_{k_m}^{2^{r-m}}$$

for $1 \leq u \leq r$, $x_{(\emptyset, 1)} = 1$. For a monomial $x = x_1^{t_1} x_2^{t_2} \dots x_{d-1}^{t_{d-1}} \in \mathcal{P}_{d-1}$, we define the monomial $\phi_{(k; \mathcal{K})}(x)$ in $\mathcal{P}_d$ by setting

$$\phi_{(k; \mathcal{K})}(x) = \begin{cases} \frac{x_k^{2^r-1} \rho_{(k, d)}(x)}{x_{(\mathcal{K}, u)}} & \text{if there exist } u \text{ such that,} \\ & t_{k_1-1} = t_{k_2-1} = \dots = t_{k_{(u-1)}-1} = 2^r - 1, \\ & t_{k_u-1} > 2^r - 1, \\ & \alpha_{r-m}(t_{k_u-1}) = 1, \quad \forall m, \quad 1 \leq m \leq u, \\ & \alpha_{r-m}(t_{k_m-1}) = 1, \quad \forall m, \quad u < m \leq r, \\ & \text{otherwise.} \\ 0 \end{cases}$$

Then we have the $\mathbb{Z}/2$ -linear map $\phi_{(k; \mathcal{K})} : \mathcal{P}_{d-1} \rightarrow \mathcal{P}_d$. In particular, $\phi_{(k; \emptyset)} = \rho_{(k, d)}$. Note that $\phi_{(k; \mathcal{K})}$ is not an $\mathcal{A}_2$ -homomorphism. Moreover, we see that for each $x \in \mathcal{P}_{d-1}$, if $\phi_{(k; \mathcal{K})}(x) \neq 0$, then it is not a spike and $\omega(\phi_{(k; \mathcal{K})}(x)) = \omega(x)$.

3.2. Some examples on $\phi_{(k; \mathcal{K})}$

The following examples can be found in [47]. However, we present them in more detail.

Let $\mathcal{K} = (j)$ and $1 \leq k < j \leq d$. Then, for any the monomial $x = x_1^{a_1} x_2^{a_2} \dots x_{d-1}^{a_{d-1}} \in \mathcal{P}_{d-1}$ and $\alpha_0(a_{j-1}) = 1$, we get $\phi_{(k; \mathcal{K})}(x) = \frac{x_k \rho_{(k, d)}(x)}{x_j}$.

Let m be a positive integer and let $x = Y^{2^m-1} y^{2^m}$ with $y = x_1^{b_1} x_2^{b_2} \dots x_{d-1}^{b_{d-1}}$ and $Y = X_{(\{d\}, d)} = x_1 x_2 \dots x_{d-1} \in \mathcal{P}_{d-1}$. Then if $m > r = \ell(\mathcal{K})$ and $u = 1$ then

$$\phi_{(k; \mathcal{K})}(x) = \phi_{(k; \mathcal{K})}(Y^{2^m-1})(\rho_{(k, d)}(y))^{2^m} = x_k^{2^r-1} \prod_{1 \leq t \leq r} x_{k_t}^{2^m-2^{r-t}-1} X_{(\{k, k_1, \dots, k_r\}, d)}^{2^m-1} (\rho_{(k, d)}(y))^{2^m}.$$

Indeed, since $\rho_{(k, d)}$ is an $\mathbb{Z}/2$ -algebras homomorphism,

$$\rho_{(k, d)}(x) = \rho_{(k, d)}(X^{2^m-1})(\rho_{(k, d)}(y))^{2^m}, \quad 1 \leq k \leq d.$$

Since $Y^{2^m-1} = x_1^{2^m-1} \dots x_{d-1}^{2^m-1}$ and $2^d - 1 > 2^r - 1$, for each $(k; \mathcal{K})$, $\mathcal{K} = (k_1, k_2, \dots, k_r)$ and $u = 1$, we have

$$\begin{aligned} \phi_{(k; \mathcal{K})}(Y^{2^m-1}) &= \frac{x_k^{2^r-1} x_2^{2^m-1} \dots x_{k_1}^{2^m-1} \dots x_{k_r}^{2^m-1} \dots x_d^{2^m-1}}{x_k^{2^r-1} x_{k_1}^{2^r-2} x_{k_2}^{2^r-2} \dots x_{k_r}^{2^r-r}} \\ &= x_k^{2^r-1} x_{k_1}^{2^m-2^{r-1}-1} \dots x_{k_2}^{2^m-2^{r-2}-1} \dots x_{k_r}^{2^m-2^{r-r}-1} X_{(\{k, k_1, \dots, k_r\}, d)}^{2^m-1} \\ &= x_k^{2^r-1} \prod_{1 \leq t \leq r} x_{k_t}^{2^m-2^{r-t}-1} X_{(\{k, k_1, \dots, k_r\}, d)}^{2^m-1}. \end{aligned}$$

Then, one gets

$$\begin{aligned}
\phi_{(k;\mathcal{K})}(x) &= \phi_{(k;\mathcal{K})}(Y^{2^m-1}y^{2^m}) \\
&= \frac{x_k^{2^r-1}\rho_{(k,d)}(X^{2^m-1}y^{2^m})}{x_{(\mathcal{K};1)}} = \left(\frac{x_k^{2^r-1}\rho_{(k,d)}(Y^{2^m-1})}{x_{(\mathcal{K};1)}} \right) (\rho_{(k,d)}(y))^{2^m} \\
&= \phi_{(k;\mathcal{K})}(X^{2^m-1})(\rho_{(k,d)}(y))^{2^m} \\
&= x_k^{2^r-1} \prod_{1 \leq t \leq r} x_{k_t}^{2^m-2^{r-t}-1} X_{(\{k,k_1,\dots,k_r\},d)}^{2^m-1} (\rho_{(k,d)}(y))^{2^m}.
\end{aligned}$$

Now, if $m = r$, $b_{j-1} = 0$, $j = k_1, k_2, \dots, k_{u-1}$ and $b_{k_u-1} > 0$ then for each $(k; \mathcal{K})$ and $1 \leq u \leq r = m$, we have

$$\phi_{(k;\mathcal{K})}(x) = \phi_{(k_u;\{k_{u+1},\dots,k_m\})}(Y^{2^m-1})(\rho_{(k,d)}(y))^{2^m}.$$

Indeed, we have

$$\begin{aligned}
Y^{2^m-1}y^{2^m} &= (x_1 \dots x_{d-1})^{2^m-1} (x_1^{b_1} \dots \hat{x}_{k_1}^{b_{k_1}} \dots \hat{x}_{k_{(u-1)-1}}^{b_{k_{(u-1)-1}}} x_{k_u-1}^{b_{k_u-1}} \dots x_{d-1}^{b_{d-1}})^{2^m} \\
&= (x_1 \dots x_{k_1} \dots x_{k_{u-1}-1} x_{k_u-1} \dots x_{d-1})^{2^m-1} (x_1^{b_1} \dots \hat{x}_{k_1}^{b_{k_1}} \dots \hat{x}_{k_{(u-1)-1}}^{b_{k_{(u-1)-1}}} x_{k_u-1}^{b_{k_u-1}} \dots x_{d-1}^{b_{d-1}})^{2^m}.
\end{aligned}$$

Then, we get

$$\begin{aligned}
\phi_{(k;\mathcal{K})}(x) &= \frac{x_k^{2^m-1}\rho_{(k,d)}(Y^{2^m-1})}{x_{k_u}^{2^{m-1}+2^{m-2}+\dots+2^{m-u}} \prod_{\substack{u < t \leq r \\ 1 \leq t \leq m}} x_{k_t}^{2^m-t}} (\rho_{(k,d)}(y))^{2^m}. \\
&= \frac{x_k^{2^m-1} x_2^{2^m-1} \dots x_{k_{u-1}}^{2^m-1} x_{k_u}^{2^m-1} \prod_{u+1 \leq t \leq m} x_{k_t}^{2^m-1} \dots x_d^{2^m-1}}{x_{k_u}^{2^{m-1}+2^{m-2}+\dots+2^{m-u}} \prod_{\substack{u < t \leq m \\ 1 \leq t \leq m}} x_{k_t}^{2^m-t}} (\rho_{(k,d)}(y))^{2^m} \\
&= x_k^{2^m-1} x_2^{2^m-1} \dots x_{k_{u-1}}^{2^m-1} x_{k_u}^{2^{m-(u+1)}+\dots+2^{m-m}} \dots x_d^{2^m-1} \prod_{u+1 \leq t \leq m} x_{k_t}^{2^m-2^{m-t}-1} \dots x_d^{2^m-1} (\rho_{(k,d)}(y))^{2^m} \\
&= \left(x_{k_u}^{2^{m-(u+1)}+\dots+2^{m-m}} \prod_{u+1 \leq t \leq m} x_{k_t}^{2^m-2^{m-t}-1} X_{(\{k_u,k_{u+1},\dots,k_m\},d)}^{2^m-1} \right) (\rho_{(k,d)}(y))^{2^m}. \\
&= \phi_{(k_u;\{k_{u+1},\dots,k_m\})}(Y^{2^m-1})(\rho_{(k,d)}(y))^{2^m}.
\end{aligned}$$

3.3. Singer's criterion on the $\mathcal{A}_2$ -decomposable and Sum's conjecture

For a subset $\mathcal{U} \subset \mathcal{P}_{d-1}$, we denote

$$\begin{aligned}
\overline{\Phi}^0(\mathcal{U}) &= \bigcup_{1 \leq k \leq d} \phi_{(k;\emptyset)}(\mathcal{U}) = \bigcup_{1 \leq k \leq d} \rho_{(k,d)}(\mathcal{U}), \\
\overline{\Phi}^+(\mathcal{U}) &= \bigcup_{(k;\mathcal{K}) \in \mathcal{N}_d, 0 < \ell(\mathcal{K}) < d} \phi_{(k;\mathcal{K})}(\mathcal{U}) \setminus \mathcal{P}_d^0, \\
\overline{\Phi}(\mathcal{U}) &= \overline{\Phi}^0(\mathcal{U}) \cup \overline{\Phi}^+(\mathcal{U}).
\end{aligned}$$

Since $\rho_{(k,d)}$ is a homomorphism of the $\mathcal{A}_2$ -modules, if $\mathcal{U}$ is a minimal set of generators for the $\mathcal{A}_2$ -module $\mathcal{P}_{d-1}$ in degree n , then $\overline{\Phi}^0(\mathcal{U})$ is also a minimal set of generators for the $\mathcal{A}_2$ -module $\mathcal{P}_d^0$ in degree n .

Now, for a polynomial $f \in \mathcal{P}_d$, we denote by $[f]$ the classes in $Q\mathcal{P}_d$ represented by f . If ω is a weight vector and $f \in \mathcal{P}_d(\omega)$, then denote by $[f]_\omega$ the classes in $Q\mathcal{P}_d(\omega)$ represented by f .

Denote by $\mathcal{B}_d(n)$ the set of all admissible monomials of degree n in $\mathcal{P}_d$. Set

$$\mathcal{B}_d^0(n) := \mathcal{B}_d(n) \cap (\mathcal{P}_d^0)_n, \quad \mathcal{B}_d^+(n) := \mathcal{B}_d(n) \cap (\mathcal{P}_d^+)_n.$$

If ω is a weight vector of degree n , we set

$$\mathcal{B}_d(\omega) := \mathcal{B}_d(n) \cap \mathcal{P}_d(\omega), \quad \mathcal{B}_d^0(\omega) := \mathcal{B}_d(\omega) \cap (\mathcal{P}_d^0)_n, \quad \mathcal{B}_d^+(\omega) := \mathcal{B}_d(\omega) \cap (\mathcal{P}_d^+)_n.$$

Then, $[\mathcal{B}_d(\omega)]_\omega$, $[\mathcal{B}_d^0(\omega)]_\omega$ and $[\mathcal{B}_d^+(\omega)]_\omega$ are respectively the bases of the $\mathbb{Z}/2$ -vector spaces

$$Q\mathcal{P}_d(\omega), \quad Q\mathcal{P}_d^0(\omega) := Q\mathcal{P}_d(\omega) \cap (Q\mathcal{P}_d^0)_n, \quad \text{and} \quad Q\mathcal{P}_d^+(\omega) := Q\mathcal{P}_d(\omega) \cap (Q\mathcal{P}_d^+)_n.$$

For a subset $\mathcal{B} \subset \mathcal{P}_d$, we denote $[\mathcal{B}] = \{[f] : f \in \mathcal{B}\}$. If $\mathcal{B} \subset \mathcal{P}_d(\omega)$, then we set $[\mathcal{B}]_\omega = \{[f]_\omega : f \in \mathcal{B}\}$. Throughout this paper, to prove a certain subset of $Q\mathcal{P}_d$ is linearly independent, we use Singer's criterion on the $\mathcal{A}_2$ -decomposable (see Theorem 3.3.5 below), Proposition 3.1.1, and a result in Sum [47]. More precisely, let $\mathcal{B}$ be a finite subset of $\mathcal{P}_d$ consisting of some monomials of degree n . Denote by $|\mathcal{B}|$ the cardinal of $\mathcal{B}$. To prove the set $[\mathcal{B}]$ is linearly independent in $(Q\mathcal{P}_d)_n$, we denote the elements of $\mathcal{B}$ by $\mathcal{Y}_{n,i}$, $1 \leq i \leq m = |\mathcal{B}|$ and assume that there is a linear relation

$$\mathcal{S} = \sum_{1 \leq i \leq m} \gamma_i \mathcal{Y}_{n,i} = 0 \text{ modulo } (\mathcal{A}_2^+ \mathcal{P}_d + \mathcal{P}_d^-(\omega)),$$

with $\gamma_i \in \mathbb{Z}/2$ for all i , $1 \leq i \leq m$. For $(k; \mathcal{K}) \in \mathcal{N}_d$, we explicitly compute $\pi_{(k; \mathcal{K})}(\mathcal{S})$ in terms of the admissible monomials in $\mathcal{P}_{d-1}$ (modulo $(\mathcal{A}_2^+ \mathcal{P}_{d-1} + \mathcal{P}_{d-1}^-(\omega))$). Computing from some relations $\pi_{(k; \mathcal{K})}(\mathcal{S}) = 0$ modulo $(\mathcal{A}_2^+ \mathcal{P}_{d-1} + \mathcal{P}_{d-1}^-(\omega))$ with $(k; \mathcal{K}) \in \mathcal{N}_d$, we obtain $\gamma_i = 0$ for all i .

In [49], Sum made the following conjecture on the relation between the admissible monomials for the polynomial algebras.

Conjecture 3.3.1 (see [49]). *If ω is a weight vector, then $\overline{\Phi}(\mathcal{B}_{d-1}(\omega)) \subseteq \mathcal{B}_d(\omega)$.*

We see that if this conjecture is true, then $\overline{\Phi}(\mathcal{B}_{d-1}(n)) \subseteq \mathcal{B}_d(n)$ for any positive integer n . In other words, if $x \in \mathcal{B}_{d-1}(n)$, then $\phi_{(k; \mathcal{K})}(x) \in \mathcal{B}_d(n)$. Based on the results of Peterson [32], Kameko [15] and Sum [47], the conjecture is true for $d \leq 4$. In particular, we have the following remark.

Remark 3.3.2. Consider the spike monomial $Y = X_{(\{d\}, d)} = x_1 x_2 \dots x_{d-1} \in \mathcal{P}_{d-1}$. Let m be a positive integer such that $m > r = \ell(\mathcal{K})$. Then, according to Sect. 3.2, we have

$$\phi_{(k; \mathcal{K})}(Y^{2^m-1}) = x_k^{2^r-1} \prod_{1 \leq t \leq r} x_{i_t}^{2^m-2^{r-t}-1} X_{(\{k, k_1, \dots, k_r\}, d)}^{2^m-1}.$$

It is easy to see that $\overline{\omega} := \omega(Y) = \underbrace{(d-1, d-1, \dots, d-1)}_{m \text{ times of } (d-1)}$. Based on the results in [35, 50], the set

$\{\phi_{(k; \mathcal{K})}(Y^{2^m-1}) : (k; \mathcal{K}) \in \mathcal{N}_d\}$ is a basis of $Q\mathcal{P}_d(\overline{\omega})$. Note that this also holds true for $m \leq r$ (see Sum [50]). Since Y^{2^m-1} is a spike monomial, by Proposition 3.1.1, Y^{2^m-1} is admissible. Combining these data, Sum's conjecture is true for the weight vector $\overline{\omega}$, where d is an arbitrary positive integer.

In [49], Sum showed that Conjecture 3.3.1 is true for $d = 5$ and any the weight vector ω of the degree $15 \cdot 2^t - 5$ for t an arbitrary non-negative integer. In Sect. 3.4, we will show that this conjecture is also true for $d = 5$ and in degree $13 \cdot 2^t - 5$ for any positive integer t .

To end this section, we review Singer's criterion on the $\mathcal{A}_2$ -decomposable.

Definition 3.3.3. A monomial $z = x_1^{t_1} x_2^{t_2} \dots x_d^{t_d}$ in $\mathcal{P}_d$ is called a *spike* if $t_j = 2^{a_j} - 1$ for a_j a non-negative integer and $1 \leq j \leq d$. If z is a spike with $a_1 > a_2 > \dots > a_{r-1} \geq a_r > 0$ and $a_j = 0$ for $j > r$, then it is called a *minimal spike*.

Lemma 3.3.4 (see [34, 37]). *All the spikes in $\mathcal{P}_d$ are admissible and their weight vectors are weakly decreasing. Furthermore, if a weight vector $\omega = (\omega_1, \omega_2, \dots)$ is weakly decreasing and $\omega_1 \leq d$, then there is a spike z in $\mathcal{P}_d$ such that $\omega(z) = \omega$.*

We refer the reader to [37] for the detailed proofs of the lemma. Singer showed in [43] that if $\mu(n) \leq d$, then there exists uniquely a minimal spike of degree n in $\mathcal{P}_d$. Further, we have the following.

Theorem 3.3.5 (see [43]). *Suppose that $X \in \mathcal{P}_d$ is a monomial of degree n , where $\mu(X) \leq d$. Let z be the minimal spike of degree n in $\mathcal{P}_d$. If $\omega(X) < \omega(z)$, then X is $\mathcal{A}_2$ -decomposable.*

This standard is also a useful tool to solve the hit problem.

3.4. Proof of Theorem 1.1

As shown in Sect.1, the iterated Kameko squaring operation

$$(\widetilde{Sq_*^0})_{(5,13,2^t-5)}^{t-2} : (Q\mathcal{P}_5)_{13,2^t-5} \rightarrow (Q\mathcal{P}_5)_{13,2^2-5}$$

is an isomorphism of $\mathbb{Z}/2GL_5$ -modules for every $t \geq 2$. So, we need only to study the structure of $(Q\mathcal{P}_5)_{13,2^t-5}$ for $t = 0, 1, 2$. Note that the case $t = 0$ has been determined by T  n [53].

3.4.1. The case $t = 1$

Now, for $t = 1$, we have $n = 13 \cdot 2^1 - 5 = 21$. Consider Kameko's squaring operation

$$(\widetilde{Sq_*^0})_{(5,21)} : (Q\mathcal{P}_5)_{21} \longrightarrow (Q\mathcal{P}_5)_8.$$

Since $(\widetilde{Sq_*^0})_{(5,21)}$ is an epimorphism of $\mathbb{Z}/2$ -vector spaces, one gets

$$(Q\mathcal{P}_5)_{21} \cong \text{Ker}(\widetilde{Sq_*^0})_{(5,21)} \bigoplus (Q\mathcal{P}_5)_8 \cong (Q\mathcal{P}_5^0)_{21} \bigoplus (\text{Ker}(\widetilde{Sq_*^0})_{(5,21)} \cap (Q\mathcal{P}_5^+)_{21}) \bigoplus (Q\mathcal{P}_5)_8.$$

By T  n [53], $\dim(Q\mathcal{P}_5)_8 = 174$. According to the results in [32], [15] and [47], we have

$$|\mathcal{B}_1^+(21)| = 0, |\mathcal{B}_2^+(21)| = 0, |\mathcal{B}_3^+(21)| = 7, \text{ and } |\mathcal{B}_4^+(21)| = 66.$$

Note that $(Q\mathcal{P}_3)_{21} \cong (Q\mathcal{P}_3)_3$ and $\mathcal{B}_3(21) = \mathcal{B}_3^+(21) = \tilde{\varphi}^2(\mathcal{B}_3(3))$ with the $\mathbb{Z}/2$ -linear map $\tilde{\varphi} : \mathcal{P}_3 \rightarrow \mathcal{P}_3$, given by $\tilde{\varphi}(u) = X_{(\emptyset,3)}u^2$, $\forall u \in \mathcal{P}_3$. Since $(Q\mathcal{P}_5^0)_{21} = \bigoplus_{1 \leq r \leq 4} \bigoplus_{1 \leq j \leq \binom{5}{r}} (Q\mathcal{P}_r^+)_{21}$, one gets $\dim(Q\mathcal{P}_5^0)_{21} = \binom{5}{3} \cdot 7 + \binom{5}{4} \cdot 66 = 400$, and a direct computation shows that $\mathcal{B}_5^0(21) = \overline{\Phi}^0(\mathcal{B}_4(21)) = \{\mathcal{Y}_{21,i} : 1 \leq i \leq 400\}$, where the monomials $\mathcal{Y}_{21,i}$, $1 \leq i \leq 400$, are listed in Sect.6.2.

Proposition 3.4.1. *The set $\{[\mathcal{Y}_{21,i}] : 401 \leq i \leq 666\}$ is the basis of the $\mathbb{Z}/2$ -vector space $\text{Ker}(\widetilde{Sq_*^0})_{(5,21)} \cap (Q\mathcal{P}_5^+)_{21}$. Here the monomials $\mathcal{Y}_i := \mathcal{Y}_{21,i}$, $401 \leq i \leq 666$, which are determined in Sect.6.3.*

Combining this result and the above data, we get $\dim(Q\mathcal{P}_5)_{21} = 840$.

We prepare some lemmas for the proof of Proposition 3.4.1.

Lemma 3.4.2. *If u is an admissible monomial of degree 21 in $\mathcal{P}_5$ and $[u]$ belong to $\text{Ker}(\widetilde{Sq_*^0})_{(5,21)}$, then we have either $\omega(u) = (3, 3, 1, 1)$ or $\omega(u) = (3, 3, 3)$.*

Proof. Observe $\mathcal{Z} = x_1^{15}x_2^3x_3^3$ is the minimal spike of degree 21 in $\mathcal{P}_5$ and $\omega(\mathcal{Z}) = (3, 3, 1, 1)$. Since $[u] \neq [0]$, by Theorem 3.3.5, we get either $\omega_1(u) = 3$ or $\omega_1(u) = 5$. If $\omega_1(u) = 5$ then $u = X_{(\emptyset,5)}y^2$ with y a monomial of degree 8 in $\mathcal{P}_5$. Since u is admissible, by Theorem 2.2.1, one gets $y \in \mathcal{B}_5(8)$. So $(\widetilde{Sq_*^0})_{(5,21)}([u]) = [y] \neq [0]$. This contradicts the fact that $[u] \in \text{Ker}(\widetilde{Sq_*^0})_{(5,21)}$. Hence, $\omega_1(u) = 3$. Then, we have $u = X_{(\{i,j\},5)}y_1^2$ with $1 \leq i < j \leq 5$ and $y_1 \in \mathcal{B}_5(9)$. Since y_1 is admissible, according to a result in [55], we have either $\omega(y_1) = (3, 1, 1)$ or $\omega(y_1) = (3, 3)$. The lemma is proved. $\square$

Based on Lemma 3.4.2, we conclude

$$\text{Ker}((\widetilde{Sq_*^0})_{(5,21)}) \cap (Q\mathcal{P}_5^+)_{21} \cong Q\mathcal{P}_5^+(3, 3, 1, 1) \bigoplus Q\mathcal{P}_5^+(3, 3, 3).$$

The following lemma is an immediate corollary from a result in [47].

Lemma 3.4.3. *The following monomials are strictly inadmissible:*

- (i) $x_i^2 x_j x_k^3 x_\ell^7, x_i^6 x_j x_k^3 x_\ell^3, x_i^2 x_j^5 x_k^3 x_\ell^3, x_i^2 x_j x_k^3 x_\ell^3, 1 \leq i < j \leq 5, 1 \leq k, \ell \leq 5, k \neq \ell, k, \ell \neq i, j$;
- (ii) $x_i^3 x_j^4 x_k^3 x_\ell^3, 1 \leq i < j < k < \ell \leq 5$;
- (iii) $\rho_{(k,5)}(X), 1 \leq k \leq 5$, where X is one of the following monomials:
 $x_1^3 x_2^4 x_3^7 x_4^7, x_1^3 x_2^7 x_3^4 x_4^7, x_1^3 x_2^7 x_3^7 x_4^4, x_1^7 x_2^3 x_3^4 x_4^7,$
 $x_1^7 x_2^3 x_3^7 x_4^4, x_1^7 x_2^7 x_3^3 x_4^4, x_1^7 x_2^8 x_3^3 x_4^3.$

Lemma 3.4.4. *The following monomials are strictly inadmissible:*

- (i) $x_i^2 x_j^2 x_k x_\ell x_m^7, x_i^2 x_j^6 x_k x_\ell x_m^3, x_i^6 x_j^2 x_k x_\ell x_m^3, x_i^2 x_j^2 x_k^5 x_\ell x_m^3, x_i^2 x_j^4 x_k x_\ell^3 x_m^3,$
 $x_i^4 x_j^2 x_k x_\ell^3 x_m^3, x_i^2 x_j x_k^4 x_\ell^3 x_m^3$, where (i, j, k, ℓ, m) is a permutation of $(1, 2, 3, 4, 5)$;
- (ii) $x_1^3 x_2^2 x_3 x_4 x_5^6, x_1^3 x_2^6 x_3 x_4 x_5^2, x_1^3 x_2^2 x_3^5 x_4 x_5, x_1^3 x_2^2 x_3 x_4^5 x_5^2,$
 $x_1^3 x_2^2 x_3^5 x_4 x_5^2, x_1^3 x_2^5 x_3^2 x_4 x_5, x_1^3 x_2^2 x_3^2 x_4 x_5, x_1^3 x_2^2 x_3 x_4^2 x_5,$
 $x_1^3 x_2^2 x_3 x_4 x_5^2.$

Proof. Consider the monomials $X = x_i^2 x_j^2 x_k x_\ell x_m^7$ and $Y = x_1^3 x_2^2 x_3 x_4 x_5^6$. We prove that these monomials are strictly inadmissible. The others can be proved by a similar computation. Obviously, $\omega(X) = \omega(Y) = (3, 3, 1)$. By a direct computation using the Cartan formula, we obtain

$$\begin{aligned} X &= Sq^2(x_i x_j x_k x_\ell x_m^7) + Sq^4(x_i x_j x_k x_\ell x_m^5) + x_i x_j x_k^2 x_\ell^2 x_m^7 \\ &\quad + x_i x_j^2 x_k x_\ell^2 x_m^7 + x_i x_j^2 x_k^2 x_\ell x_m^7 \text{ modulo } (\mathcal{P}_5^-(3, 3, 1)); \\ Y &= Sq^1(x_1^3 x_2 x_3 x_4 x_5^6) + x_1^3 x_2 x_3^2 x_4 x_5^6 + x_1^3 x_2 x_3 x_4^2 x_5^6 \text{ modulo } (\mathcal{P}_5^-(3, 3, 1)). \end{aligned}$$

These equalities show that X and Y are strictly inadmissible. The lemma follows. $\square$

Lemma 3.4.5. *The following monomials are strictly inadmissible:*

$$\begin{aligned} X_1 &= x_1 x_2^6 x_3^8 x_4^3 x_5^3, & X_2 &= x_1 x_2^3 x_3^6 x_4^6 x_5^5, & X_3 &= x_1^3 x_2^5 x_3^2 x_4^6 x_5^5, & X_4 &= x_1 x_2^6 x_3^7 x_4^6 x_5, \\ X_5 &= x_1 x_2^7 x_3^6 x_4^6 x_5, & X_6 &= x_1^7 x_2 x_3^6 x_4^6 x_5, & X_7 &= x_1 x_2^2 x_3^6 x_4^5 x_5^7, & X_8 &= x_1 x_2^2 x_3^6 x_4^7 x_5^5, \\ X_9 &= x_1 x_2^6 x_3^2 x_4^5 x_5^7, & X_{10} &= x_1 x_2^6 x_3^2 x_4^7 x_5^5, & X_{11} &= x_1 x_2^2 x_3^7 x_4^6 x_5^5, & X_{12} &= x_1 x_2^2 x_3^7 x_4^2 x_5^5, \\ X_{13} &= x_1 x_2^7 x_3^2 x_4^6 x_5^5, & X_{14} &= x_1 x_2^7 x_3^6 x_4^2 x_5^5, & X_{15} &= x_1^7 x_2 x_3^2 x_4^6 x_5^5, & X_{16} &= x_1^7 x_2 x_3^6 x_4^2 x_5^5, \\ X_{17} &= x_1 x_2^6 x_3^3 x_4^4 x_5^7, & X_{18} &= x_1 x_2^6 x_3^3 x_4^7 x_5^4, & X_{19} &= x_1 x_2^6 x_3^7 x_4^3 x_5^4, & X_{20} &= x_1 x_2^7 x_3^6 x_4^3 x_5^4, \\ X_{21} &= x_1^3 x_2^4 x_3 x_4^6 x_5^7, & X_{22} &= x_1^3 x_2^4 x_3 x_4^7 x_5^6, & X_{23} &= x_1^3 x_2^4 x_3^7 x_4^6 x_5^6, & X_{24} &= x_1^3 x_2^7 x_3^4 x_4^6 x_5^6, \\ X_{25} &= x_1^7 x_2 x_3^6 x_4^3 x_5^4, & X_{26} &= x_1^7 x_2 x_3^4 x_4^6 x_5^6, & X_{27} &= x_1 x_2^6 x_3^6 x_4^3 x_5^5, & X_{28} &= x_1 x_2^6 x_3^3 x_4^6 x_5^5, \\ X_{29} &= x_1 x_2^6 x_3^6 x_4^7 x_5, & X_{30} &= x_1^3 x_2 x_3^6 x_4^6 x_5^5, & X_{31} &= x_1^3 x_2^5 x_3^6 x_4^6 x_5^5, & X_{32} &= x_1^3 x_2^5 x_3^5 x_4^2 x_5^7, \\ X_{33} &= x_1^3 x_2^4 x_3^7 x_4^2 x_5^2, & X_{34} &= x_1^3 x_2^4 x_3^7 x_4^2 x_5^2, & X_{35} &= x_1^3 x_2^5 x_3^4 x_4^2 x_5^7, & X_{36} &= x_1^3 x_2^5 x_3^4 x_4^2 x_5^2, \\ X_{37} &= x_1^3 x_2^7 x_3^4 x_5^2 x_5^2, & X_{38} &= x_1^3 x_2^5 x_3^7 x_4^2 x_5^2, & X_{39} &= x_1^3 x_2^7 x_3^5 x_4^2 x_5^2, & X_{40} &= x_1^7 x_2^3 x_3^4 x_5^2 x_5^2, \\ X_{41} &= x_1^7 x_2^3 x_3^5 x_4^2 x_5^2, & X_{42} &= x_1 x_2^6 x_3^6 x_4^7 x_5^7, & X_{43} &= x_1^3 x_2^5 x_3^6 x_4^2 x_5^5, & X_{44} &= x_1^3 x_2^5 x_3^6 x_4^5 x_5^2, \\ X_{45} &= x_1^3 x_2^5 x_3^5 x_4^2 x_5^6, & X_{46} &= x_1^3 x_2^5 x_3^5 x_4^2 x_5^2, & X_{47} &= x_1^3 x_2^4 x_3^4 x_4^7 x_5^7, & X_{48} &= x_1^3 x_2^4 x_3^4 x_4^7 x_5^3, \\ X_{49} &= x_1^3 x_2^4 x_3^3 x_4^4 x_5^7, & X_{50} &= x_1^3 x_2^4 x_3^7 x_4^4 x_5^3, & X_{51} &= x_1^3 x_2^4 x_3^3 x_4^7 x_5^4, & X_{52} &= x_1^3 x_2^4 x_3^7 x_4^3 x_5^4, \\ X_{53} &= x_1^3 x_2^7 x_3^4 x_4^4 x_5^3, & X_{54} &= x_1^3 x_2^7 x_3^4 x_4^3 x_5^4, & X_{55} &= x_1^7 x_2^3 x_4^4 x_4^3 x_5^3, & X_{56} &= x_1^7 x_2^3 x_4^3 x_4^3 x_5^4, \\ X_{57} &= x_1^3 x_2^4 x_3^5 x_4^6 x_5^3, & X_{58} &= x_1^3 x_2^4 x_3^5 x_4^6 x_5^3, & X_{59} &= x_1^3 x_2^5 x_3^4 x_4^6 x_5^3, & X_{60} &= x_1^3 x_2^5 x_3^4 x_4^6 x_5^3, \\ X_{61} &= x_1^3 x_2^5 x_3^4 x_4^6 x_5^3. \end{aligned}$$

Proof. It is easy to see that $\omega(X_1) = (3, 3, 1, 1)$ and $\omega(X_j) = (3, 3, 3)$ for $j = 2, 3, \dots, 61$. We prove the lemma for the monomials $X_1 = x_1 x_2^6 x_3^8 x_4^3 x_5^3$, $X_2 = x_1^3 x_2^5 x_3^2 x_4^6 x_5^5$ and $X_3 = x_1 x_2^6 x_3^7 x_4^6 x_5$. The others can be proven by a similar computation. By a direct computation, we have

$$\begin{aligned} X_1 &= x_1 x_2^3 x_3^3 x_4^6 x_5^8 + x_1 x_2^3 x_3^3 x_4^8 x_5^6 + x_1 x_2^3 x_3^6 x_4^3 x_5^8 + x_1 x_2^3 x_3^6 x_4^8 x_5^3 + x_1 x_2^3 x_3^8 x_4^3 x_5^6 \\ &\quad + x_1 x_2^3 x_3^8 x_4^6 x_5^3 + x_1 x_2^4 x_3^3 x_4^3 x_5^{10} + x_1 x_2^4 x_3^3 x_4^{10} x_5^3 + x_1 x_2^4 x_3^{10} x_4^3 x_5^3 \\ &\quad + x_1 x_2^6 x_3^3 x_4^3 x_5^8 + x_1 x_2^6 x_3^3 x_4^8 x_5^3 + Sq^1(A_1) + Sq^2(A_2) + Sq^4(A_4) \text{ modulo } (\mathcal{P}_5^-(3, 3, 1, 1)), \end{aligned}$$

where

$$\begin{aligned} A_1 &= x_1^2 x_2^3 x_3^5 x_4^5 x_5^5 + x_1^2 x_2^5 x_3^3 x_4^5 x_5^5 + x_1^2 x_2^5 x_3^5 x_4^3 x_5^5 + x_1^2 x_2^5 x_3^5 x_4^5 x_5^3, \\ A_2 &= x_1 x_2^3 x_3^3 x_4^6 x_5^6 + x_1 x_2^3 x_3^5 x_4^5 x_5^5 + x_1 x_2^3 x_3^6 x_4^3 x_5^6 + x_1 x_2^3 x_3^6 x_4^5 x_5^3 + x_1 x_2^5 x_3^3 x_4^5 x_5^5 \\ &\quad + x_1 x_2^5 x_3^5 x_4^3 x_5^5 + x_1 x_2^5 x_3^5 x_4^5 x_5^3 + x_1 x_2^6 x_3^3 x_4^6 x_5^6 + x_1 x_2^6 x_3^3 x_4^6 x_5^3 + x_1 x_2^6 x_3^6 x_4^3 x_5^3, \\ A_4 &= x_1 x_2^4 x_3^3 x_4^3 x_5^6 + x_1 x_2^4 x_3^3 x_4^6 x_5^3 + x_1 x_2^4 x_3^6 x_4^3 x_5^3. \end{aligned}$$

This relation implies that X_1 is strictly inadmissible. By a similar technique, we obtain

$$X_2 = Sq^1(B_1) + Sq^2(B_2) + Sq^8(B_8) + x_1 x_2^3 x_3^5 x_4^6 x_5^6 + x_1 x_2^3 x_3^6 x_4^5 x_5^6 \text{ modulo } (\mathcal{P}_5^-(3, 3, 3)),$$

where

$$\begin{aligned} B_1 &= x_1^2 x_2^5 x_3^3 x_4^5 x_5^5 + x_1^2 x_2^5 x_3^5 x_4^3 x_5^5 + x_1^2 x_2^5 x_3^5 x_4^5 x_5^3, \\ B_2 &= x_1 x_2^3 x_3^3 x_4^6 x_5^6 + x_1 x_2^3 x_3^5 x_4^5 x_5^5 + x_1 x_2^3 x_3^6 x_4^3 x_5^6 + x_1 x_2^6 x_3^3 x_4^3 x_5^6 + x_1 x_2^6 x_3^3 x_4^6 x_5^3 + x_1 x_2^6 x_3^6 x_4^3 x_5^3, \\ B_8 &= x_1 x_2^3 x_3^3 x_4^3 x_5^3, \end{aligned}$$

$$X_3 = Sq^1(C_1) + Sq^2(C_2) + Sq^8(C_8) + x_1^3 x_2^5 x_3 x_4^6 x_5^6 + x_1^3 x_2^5 x_3^2 x_4^5 x_5^6 \text{ modulo } (\mathcal{P}_5^-(3, 3, 3)),$$

where

$$\begin{aligned} C_1 &= x_1^3 x_2^5 x_3 x_4^5 x_5^6 + x_1^3 x_2^5 x_3 x_4^6 x_5^5 + x_1^5 x_2^3 x_3^2 x_4^5 x_5^5 + x_1^5 x_2^5 x_3^2 x_4^3 x_5^5 + x_1^5 x_2^5 x_3^2 x_4^5 x_5^3, \\ C_2 &= x_1^3 x_2^3 x_3 x_4^6 x_5^6 + x_1^3 x_2^5 x_3 x_4^5 x_5^5 + x_1^3 x_2^6 x_3 x_4^3 x_5^6 + x_1^3 x_2^6 x_3 x_4^6 x_5^3 + x_1^6 x_2^3 x_3 x_4^6 x_5^6 \\ &\quad + x_1^6 x_2^3 x_3 x_4^3 x_5^3 + x_1^6 x_2^6 x_3 x_4^3 x_5^3, \\ C_8 &= x_1^3 x_2^3 x_3 x_4^3 x_5^3. \end{aligned}$$

The lemma is proved. $\square$

Now we denote by $\mathcal{C}$ the set of the following monomials:

$$\begin{array}{cccc} x_1^3 x_2^{12} x_3 x_4^2 x_5^3 & x_1^3 x_2^{12} x_3 x_4^3 x_5^2 & x_1^3 x_2^{12} x_3^2 x_4 x_5^2 & x_1^3 x_2^4 x_3 x_4^2 x_5^{11}, \\ x_1^3 x_2^4 x_3 x_4^{11} x_5^2 & x_1^3 x_2^4 x_3^{11} x_4 x_5^2 & x_1^7 x_2^8 x_3 x_4^2 x_5^3 & x_1^7 x_2^8 x_3 x_4^3 x_5^2, \\ x_1^7 x_2^8 x_3^2 x_4 x_5^2 & x_1^3 x_2^4 x_3 x_4^{10} x_5^3 & x_1^3 x_2^4 x_3 x_4^{10} x_5^3 & x_1^3 x_2^4 x_3^2 x_4 x_5^{10}, \\ x_1^3 x_2^4 x_3^2 x_4^9 x_5^2 & x_1^3 x_2^4 x_3^2 x_4^9 x_5^2 & x_1^3 x_2^4 x_3^2 x_4^9 x_5^2 & \end{array}$$

A direct computation shows that $\overline{\Phi}^+(\mathcal{B}_4(3, 3, 1, 1)) \cup \mathcal{C}$ is the set of 196 monomials: $\mathcal{Y}_j := \mathcal{Y}_{21,j}$, $401 \leq j \leq 596$ (see Sect.6.3.)

Proposition 3.4.6. *Under the above notations, the $\mathbb{Z}/2$ -vector space $Q\mathcal{P}_5^+(3, 3, 1, 1)$ is spanned by the set*

$$[\overline{\Phi}^+(\mathcal{B}_4(3, 3, 1, 1)) \cup \mathcal{C}].$$

Proof. Let X be an admissible monomial in $\mathcal{P}_5$ such that $\omega(X) = (3, 3, 1, 1)$. Then $X = X_{(\{k,\ell\},5)} Y^2$ with $1 \leq k < \ell \leq 5$ and Y a monomial of degree 9 in $\mathcal{P}_5$. Since X is admissible, according to Theorem 2.2.1, $Y \in \mathcal{B}_5(3, 1, 1)$.

A direct computation shows that if $z \in \mathcal{B}_5^+(3, 1, 1)$, $1 \leq k < \ell \leq 5$, and $X_{(\{k,\ell\},5)} z^2 \neq \mathcal{Y}_j$, $\forall j$, $401 \leq j \leq 596$ then there exists a monomial w which is given in one of Lemmas 3.4.3-3.4.5 such that $X_{(\{k,\ell\},5)} z^2 = w z_1^{2^a}$ with a monomial $z_1 \in \mathcal{P}_5$ and $a = \max\{m \in \mathbb{Z} : \omega_m(w) > 0\}$. By Theorem 2.2.1, $X_{(\{k,\ell\},5)} z^2$ is inadmissible. Since $X = X_{(\{k,\ell\},5)} Y^2$ with $Y \in \mathcal{B}_5(3, 1, 1)$ and X is admissible, one can see that $X = \mathcal{Y}_j$, $401 \leq j \leq 596$. The lemma follows. $\square$

By a direct computation, we see that

$$\overline{\Phi}^+(\mathcal{B}_4(3, 3, 3)) \cup \{x_1^3 x_2^4 x_3 x_4^5 x_5^6, x_1^3 x_2^5 x_3 x_4^6 x_5^4, x_1^3 x_2^5 x_3^2 x_4^3 x_5^4\}$$

is the set consisting of 70 monomials: $\mathcal{Y}_t := \mathcal{Y}_{21,t}$, $597 \leq t \leq 666$ (see Sect.6.3.)

Proposition 3.4.7. *The $\mathbb{Z}/2$ -vector space $Q\mathcal{P}_5^+(3, 3, 3)$ is spanned by the set $\{[\mathcal{Y}_t] : 597 \leq t \leq 666\}$.*

Proof. Let u be an admissible monomial in $\mathcal{P}_5$ such that $\omega(u) = (3, 3, 3)$. Then $u = x_i x_j x_\ell y^2$ with $1 \leq i < j < \ell \leq 5$ and $y \in \mathcal{B}_5(3, 3)$.

By a direct computation, we can verify that for any $X \in \mathcal{B}_5(3, 3)$, $1 \leq i < j < \ell \leq 5$, such that $x_i x_j x_\ell X^2 \neq \mathcal{Y}_t$, $\forall t$, $597 \leq t \leq 666$, there is a monomial z which is given in one of Lemmas 3.4.3-3.4.5 such that $x_i x_j x_\ell X^2 = zw^{2^b}$ with suitable monomial $w \in \mathcal{P}_5$ and $b = \max\{r \in \mathbb{Z} : \omega_r(z) > 0\}$. Then, according to Theorem 2.2.1, $x_i x_j x_\ell X^2$ nadmissible. Since $u = x_i x_j x_\ell y^2$ is admissible and $y \in \mathcal{B}_5(3, 3)$, one gets $u = \mathcal{Y}_t$, for some t . This proves the proposition. $\square$

Proof of Proposition 3.4.1. From Propositions 3.4.6 and 3.4.7, the space $\text{Ker}((\widetilde{Sq}_*^0)_{(5,21)}) \cap (Q\mathcal{P}_5^+)_{21}$ is spanned by the set $\{[\mathcal{Y}_i := \mathcal{Y}_{21,i}] : 401 \leq i \leq 666\}$. Futhermore, this set is linearly independent in $(Q\mathcal{P}_5)_{21}$. Indeed, suppose there is a linear relation

$$\mathcal{S} = \sum_{401 \leq i \leq 666} \gamma_i \mathcal{Y}_i = 0 \text{ modulo } (\mathcal{A}_2^+ \mathcal{P}_5),$$

where $\gamma_i \in \mathbb{Z}/2$. Let $\mathcal{H}$ be a sequence of non-negative integers and $\gamma_h \in \mathbb{Z}/2$ for $h \in \mathcal{H}$. Denote $\gamma_{\mathcal{H}} = \sum_{h \in \mathcal{H}} \gamma_h \in \mathbb{Z}/2$. Based on Theorem 3.3.5 and Proposition 3.1.1, for $(k; \mathcal{K}) \in \mathcal{N}_5$, we explicitly compute $\pi_{(k; \mathcal{K})}(\mathcal{S})$ in terms of a given minimal set of $\mathcal{A}_2$ -generators in $\mathcal{P}_4$ (modulo $(\mathcal{A}_2^+ \mathcal{P}_4)$). Computing directly from the relations $\pi_{(k; p)}(\mathcal{S}) \equiv 0$, $1 \leq k < p \leq 5$, one gets $\gamma_h = 0$, $h \in \mathcal{H}$, where the set $\mathcal{H} = \{401, 402, \dots, 425, 434, 435, 439, 441,$

442, 443, 446, 451, 453, 454, 457, 458, 460, 461, 462, 463, 465, 467, 470, 471, $\dots$, 477, 479, 483, 487, 489, 497, 498, 500, 501, 502, 504, 505, 506, 507, 510, 513, 515, 517, 519, 520, 523, 527, 532, 533, 540, 542, 543, 560, 561, 562, 565, 566, 567, 568, 570, 573, 575, 577, 578, 579, 580, 581, 585, 597, 598, 599, 600, 601, 602, 604, 606, 607, 609, 610, 613, 615, 616, 617, 620, 622, 629, 630, 632, 633, $\dots$, 647, 650, 651, 652, 653, 655, 656, 657, 658, 659, 660, 662, 663}, and

$$\left\{ \begin{array}{l} \gamma_{430} = \gamma_{593}, \\ \gamma_{431} = \gamma_{594}, \gamma_{536} = \gamma_{541}, \\ \gamma_{603} = \gamma_{664}, \gamma_h = \gamma_{426}, h \in \mathbb{I}_1, \\ \gamma_{608} = \gamma_{627}, \gamma_h = \gamma_{427}, h \in \mathbb{I}_2, \\ \gamma_{619} = \gamma_{626}, \gamma_h = \gamma_{432}, h = 468, 491, 518, 595, \\ \gamma_{621} = \gamma_{665}, \gamma_h = \gamma_{433}, h = 596, 623, 666, \\ \gamma_h = \gamma_{436}, h = 437, 438, 469, 492, 495, 496, 499, 534, 539, 545, 614, 654, \\ \gamma_{\{450, 595, 661\}} = \gamma_{\{522, 648, 661\}} = \gamma_{\{531, 648, 661\}} = \gamma_{\{549, 654, 661\}} = 0, \\ \gamma_{\{555, 595, 654\}} = \gamma_{\{618, 626, 628\}} = 0, \\ \gamma_{\{582, 654, 661\}} = \gamma_{\{583, 654, 661\}} = \gamma_{\{584, 654, 661\}} = 0, \\ \gamma_{\{429, 648, 595, 661\}} = \gamma_{\{455, 595, 654, 661\}} = \gamma_{\{456, 654, 661, 666\}} = \gamma_{\{486, 648, 594, 654\}} = 0, \\ \gamma_{\{493, 664, 665, 594\}} = \gamma_{\{503, 654, 552, 666\}} = \gamma_{\{512, 648, 595, 661\}} = \gamma_{\{529, 664, 665, 594\}} = 0, \\ \gamma_{\{544, 654, 661, 595\}} = \gamma_{\{547, 654, 661, 595\}} = \gamma_{\{548, 654, 661, 666\}} = \gamma_{\{550, 654, 661, 595\}} = 0, \\ \gamma_{\{551, 648, 594, 654\}} = \gamma_{\{557, 654, 661, 666\}} = \gamma_{\{558, 654, 661, 648\}} = \gamma_{\{592, 648, 595, 661\}} = 0, \\ \gamma_{\{481, 648, 664, 665, 594\}} = \gamma_{\{526, 648, 664, 665, 594\}} = \gamma_{\{554, 654, 628, 595, 626\}} = 0, \\ \gamma_{\{624, 626, 628, 627, 664\}} = \gamma_{\{459, 628, 648, 595, 626, 666\}} = 0, \\ \gamma_{\{488, 628, 595, 626, 648, 594\}} = \gamma_{\{535, 627, 654, 595, 661, 626\}} = 0, \\ \gamma_{\{546, 661, 666, 648, 594, 552\}} = \gamma_{\{612, 628, 627, 665, 666, 664\}} = 0, \\ \gamma_{\{485, 627, 654, 648, 595, 661, 626\}} = \gamma_{\{490, 626, 628, 627, 665, 666, 664\}} = 0, \\ \gamma_{\{509, 666, 552, 654, 628, 595, 626\}} = \gamma_{\{524, 666, 552, 654, 628, 595, 626\}} = 0, \\ \gamma_{\{538, 627, 654, 648, 595, 661, 626\}} = \gamma_{\{556, 654, 628, 648, 595, 661, 626\}} = 0, \\ \gamma_{\{559, 648, 626, 628, 627, 666, 661\}} = \gamma_{\{625, 626, 628, 627, 665, 666, 664\}} = 0, \\ \gamma_{\{521, 654, 666, 648, 594, 552, 628, 664, 665\}} = \gamma_{\{553, 654, 648, 594, 626, 628, 627, 666, 664\}} = 0. \end{array} \right.$$

Here $\mathbb{I}_1 = \{440, 447, 448, 464, 466, 478, 484, 508, 514, 516, 586, 605, , 648\}$,

$$\mathbb{I}_2 = \left\{ 428, 444, 445, 449, 452, 480, 482, 494, 511, 525, 528, 530, 537, 563, \right. \\ \left. 564, 569, 571, 572, 574, 576, 587, 588, 589, 590, 591, 611, 631, 649, 661 \right\},$$

Combining the above computations and the relations $\pi_{(1,(2;j))}(\mathcal{S}) \equiv 0$, $j = 3, 4, 5$, $\pi_{(1,(3;4))}(\mathcal{S}) \equiv 0$, we obtain $\gamma_i = 0$, $\forall i$, $401 \leq i \leq 666$. This finishes the proof. $\square$

3.4.2. The case $t = 2$

For $t = 2$, we have $13 \cdot 2^t - 5 = 47$ and $\mu(47) = 3 < 5$. Since Kameko's operation

$$(\widetilde{Sq}_*^0)_{(5,47)} : (Q\mathcal{P}_5)_{47} \rightarrow (Q\mathcal{P}_5)_{21}$$

is an epimorphism of the $\mathbb{Z}/2GL_5$ -modules, hence $(Q\mathcal{P}_5)_{47} = \text{Ker}((\widetilde{Sq}_*^0)_{(5,47)}) \oplus (Q\mathcal{P}_5)_{21}$. Hence, we need to compute $\text{Ker}((\widetilde{Sq}_*^0)_{(5,47)})$.

Remark 3.4.8. If $Y \in \mathcal{B}_5(47)$ and $[Y] \in \text{Ker}((\widetilde{Sq}_*^0)_{(5,47)})$, then $\omega_1(Y) = 3$.

Indeed, we see that $\mathcal{Z} = x_1^{31}x_2^{15}x_3$ is the minimal spike of degree 47 in $\mathcal{P}_5$. By Lemma 3.3.4, $\mathcal{Z} \in \mathcal{B}_5(47)$. Since $[Y] \neq 0$, by Theorem 3.3.5, either $\omega_1(Y) = 3$ or $\omega_1(Y) = 5$. If $\omega_1(Y) = 5$, then $Y = X_{(\emptyset,5)}Z^2$ with Z a monomial of degree 21 in $\mathcal{P}_5$. Since Y is admissible, by Theorem 2.2.1, $Z \in \mathcal{B}_5(21)$. So, we have $(\widetilde{Sq}_*^0)_{(5,47)}([Y]) = [Z] \neq [0]$. This contradicts the fact that $[Y] \in \text{Ker}((\widetilde{Sq}_*^0)_{(5,47)})$; hence we get $\omega_1(Y) = 3$.

From Remark 3.4.8, we have $Y = x_kx_\ell x_m g^2$ with $1 \leq k < \ell < m \leq 5$ and $g \in \mathcal{B}_5(22)$. Thus, to determine $\text{Ker}((\widetilde{Sq}_*^0)_{(5,47)})$, we need to compute all the admissible monomials of degree 22 in the A_2 -module $\mathcal{P}_5$.

Computation of $(Q\mathcal{P}_5)_{22}$

We consider the weight vectors

$$\omega_{(1)} = (2, 2, 2, 1), \omega_{(2)} = (2, 4, 1, 1), \omega_{(3)} = (2, 4, 3), \omega_{(4)} = (4, 3, 1, 1), \omega_{(5)} = (4, 3, 3).$$

We have $\deg \omega_{(i)} = 22$, $1 \leq i \leq 5$. It is easy to see that $\mathcal{Z} = x_1^{15}x_2^7$ is the minimal spike of degree 22 in $\mathcal{P}_5$ and $\omega(z) = \omega_{(1)}$. Let u be an admissible monomial of degree 22 in $\mathcal{P}_5$. Then $[u] \neq [0]$ and by Theorem 3.3.5, either $\omega_1(u) = 2$ or $\omega_1(u) = 4$. Since $u \in \mathcal{B}_5(22)$, by Theorem 2.2.1, if $\omega_1(u) = 2$, then $u = X_{(\{i,j,k\},5)}y^2$ with $y \in \mathcal{B}_5(10)$ and $1 \leq i < j < k \leq 5$. According to T  n [54], $\omega(y)$ is one of the following sequences:

$$(2, 2, 1), (4, 1, 1), (4, 3).$$

If $\omega_1(u) = 4$, then $u = X_{(\{i\},5)}y_1^2$ with y_1 a monomial of degree 9 in $\mathcal{P}_5$ and $1 \leq i \leq 5$. By T  n [54], either $\omega(y_1) = (3, 1, 1)$ or $\omega(y_1) = (3, 3)$. Hence, one gets the following.

Remark 3.4.9. If $u \in \mathcal{B}_5(22)$, then $\omega(u)$ is one of the sequences $\omega_{(t)}$, $1 \leq t \leq 5$.

As it is known,

$$(Q\mathcal{P}_5)_{22} = (Q\mathcal{P}_5^0)_{22} \oplus (Q\mathcal{P}_5^+)_{22}.$$

By Sum [47], we have $|\mathcal{B}_4^+(22)| = 72$. Then, combining the results in [32] and [15] with fact that $(Q\mathcal{P}_5^0)_{22} = \bigoplus_{1 \leq s \leq 4} \bigoplus_{1 \leq u \leq \binom{5}{u}} (Q\mathcal{P}_s^+)_{22}$, we get

$$\dim(Q\mathcal{P}_5^0)_{22} = \binom{5}{2} \cdot 2 + \binom{5}{3} \cdot 8 + \binom{5}{4} \cdot 72 = 460,$$

and $\mathcal{B}_5^0(22) = \overline{\Phi}^0(\mathcal{B}_4(22)) = \{\mathcal{Y}_{22,t} : 1 \leq t \leq 460\}$, where the monomials $\mathcal{Y}_{22,t}$, $1 \leq t \leq 460$, are determined in Sect.6.4.

Next, we determine $(Q\mathcal{P}_5^+)_{22}$. We need some notations.

Notation 3.4.10. For n a natural number, consider the set

$$\mathcal{U}(5, n) := \{r \in \mathbb{N} : \alpha(n - (2^r - 1) + 4) \leq 4\}.$$

For $r \in \mathcal{U}(5, n)$, let $X = x_1^{a_1} \dots x_4^{a_4}$ be an admissible monomial of degree $m = n - (2^r - 1)$ in $\mathcal{A}_2$ -module $\mathcal{P}_4$. We set

$$\mathcal{B}(5, n) := \bigcup_{r \in \mathcal{U}(5, 22)} \left\langle \{x_1^{a_1} \dots x_{j-1}^{a_{j-1}} x_j^{2^r-1} x_{j+1}^{a_{j+1}} \dots x_5^{a_4} \in (\mathcal{P}_5)_{n=m+2^r-1} : 1 \leq j \leq 5\} \right\rangle.$$

Then, according to Mothebe [22], $\mathcal{B}(5, n) \subseteq \mathcal{B}_5(n)$. If ω is a weight vector of degree n , then we write

$$\mathcal{B}(5, \omega) := \mathcal{B}(5, n) \cap \mathcal{P}_5(\omega), \quad \mathcal{B}^+(5, \omega) := \mathcal{B}(5, \omega) \cap (\mathcal{P}_5^+)_n.$$

By a simple computation, we see that $\overline{\Phi}^+(\mathcal{B}_4(\omega_{(1)}) \cup \mathcal{B}^+(5, \omega_{(1)}))$ is the set of 31 admissible monomials: $\mathcal{Y}_{22,i}$, $461 \leq i \leq 491$ (see Sect.6.5.)

Denote by $\mathcal{D}$ is the set of the following monomials:

$$\begin{aligned} \mathcal{Y}_{22,492} &= x_1 x_2 x_3^6 x_4^8 x_5^8, & \mathcal{Y}_{22,493} &= x_1 x_2 x_3^6 x_4^{10} x_5^4, & \mathcal{Y}_{22,494} &= x_1 x_2^2 x_3^3 x_4^4 x_5^{12}, \\ \mathcal{Y}_{22,495} &= x_1 x_2^2 x_3^3 x_4^{12} x_5^4, & \mathcal{Y}_{22,496} &= x_1 x_2^2 x_3^4 x_4^9 x_5^6, & \mathcal{Y}_{22,497} &= x_1 x_2^2 x_3^5 x_4^8 x_5^6, \\ \mathcal{Y}_{22,498} &= x_1 x_2^3 x_3^2 x_4^4 x_5^{12}, & \mathcal{Y}_{22,499} &= x_1 x_2^3 x_3^2 x_4^{12} x_5^4, & \mathcal{Y}_{22,500} &= x_1 x_2^3 x_3^4 x_4^8 x_5^6, \\ \mathcal{Y}_{22,501} &= x_1 x_2^3 x_3^6 x_4^4 x_5^8, & \mathcal{Y}_{22,502} &= x_1 x_2^3 x_3^6 x_4^8 x_5^4, & \mathcal{Y}_{22,503} &= x_1^3 x_2 x_3^2 x_4^4 x_5^{12}, \\ \mathcal{Y}_{22,504} &= x_1^3 x_2 x_3^2 x_4^{12} x_5^4, & \mathcal{Y}_{22,505} &= x_1^3 x_2 x_3^4 x_4^8 x_5^6, & \mathcal{Y}_{22,506} &= x_1^3 x_2 x_3^6 x_4^4 x_5^8, \\ \mathcal{Y}_{22,507} &= x_1^3 x_2 x_3^6 x_4^8 x_5^4, & \mathcal{Y}_{22,508} &= x_1^3 x_2^5 x_3^2 x_4^4 x_5^8, & \mathcal{Y}_{22,509} &= x_1^3 x_2^5 x_3^2 x_4^8 x_5^4, \\ \mathcal{Y}_{22,510} &= x_1^3 x_2^5 x_3^8 x_4^2 x_5^4. \end{aligned}$$

Proposition 3.4.11. $\mathcal{B}_5^+(\omega_{(1)}) = \overline{\Phi}^+(\mathcal{B}_4(\omega_{(1)}) \cup \mathcal{B}^+(5, \omega_{(1)}) \cup \mathcal{D})$.

In order to prove the proposition, we need some lemmas.

Lemma 3.4.12. *The following monomials are strictly inadmissible:*

(i) $x_1^2 x_j x_k^2 x_l^6 x_m^4$, $x_1^6 x_j x_k x_l^2 x_m^4$, $l < m$, $x_1^2 x_j x_k x_l^4 x_m^6$.

Here (j, k, l, m) is a permutation of $(2, 3, 4, 5)$;

(ii) $x_1 x_2^2 x_3^4 x_4^6 x_5$, $x_1 x_2^2 x_3^6 x_4^4 x_5^4$, $x_1 x_2^2 x_3^6 x_4^4 x_5$, $x_1 x_2^6 x_3^2 x_4 x_5^4$,
 $x_1 x_2^6 x_3^4 x_4^2 x_5^2$, $x_1^3 x_2^3 x_3^4 x_4^2 x_5^2$, $x_1^3 x_2^4 x_3^2 x_4^2 x_5^3$, $x_1^3 x_2^4 x_3^2 x_4^2 x_5^2$,
 $x_1^3 x_2^4 x_3^2 x_4^2 x_5^6$, $x_1^3 x_2^5 x_3^2 x_4^2 x_5^2$, $x_1^3 x_2^5 x_3^2 x_4^2 x_5^6$, $x_1^3 x_2^6 x_3^2 x_4^2 x_5^2$,
 $x_1^3 x_2^6 x_3^2 x_4^2 x_5^6$, $x_1^3 x_2^6 x_3^2 x_4^2 x_5^2$, $x_1^3 x_2^6 x_3^2 x_4^2 x_5^6$, $x_1^3 x_2^6 x_3^2 x_4^2 x_5^2$,
 $x_1^3 x_2^6 x_3^2 x_4^2 x_5^6$, $x_1^3 x_2^6 x_3^2 x_4^2 x_5^2$, $x_1^3 x_2^6 x_3^2 x_4^2 x_5^6$, $x_1^3 x_2^6 x_3^2 x_4^2 x_5^2$.

Proof. We prove the lemma for the monomials $u = x_1^2 x_j x_k^2 x_l^6 x_m^4$, and $v = x_1 x_2^2 x_3^4 x_4^6 x_5$. The others can be proved by a similar computation. We have $\omega(u) = (2, 4, 1)$ and $\omega(v) = (2, 2, 2)$. By a simple computation, one gets

$$\begin{aligned} u &= x_1 x_j^2 x_k^2 x_l^3 x_m^6 + Sq^1(x_1 x_j x_k^2 x_l^3 x_m^6) \text{ modulo } (\mathcal{P}_5^-(2, 4, 1)), \\ v &= x_1 x_2 x_3^4 x_4^6 x_5^2 + x_1 x_2^2 x_3^4 x_4^5 x_5^2 + Sq^1(f_1) + Sq^2(f_2) \text{ modulo } (\mathcal{P}_5^-(2, 2, 2)), \end{aligned}$$

where $f_1 = x_1^2 x_2 x_3^4 x_4^5 x_5$ and $f_2 = x_1 x_2 x_3^4 x_4^5 x_5$. Hence, u and v are strictly inadmissible. The lemma follows. $\square$

The following lemma can easily be proved by a direct computation.

Lemma 3.4.13. *If (i, j, k, l, m) is a permutation of $(1, 2, 3, 4, 5)$, then the following monomials are strictly inadmissible:*

- (i) $x_i^6 x_j x_k^7, x_i^2 x_j^5 x_k^7, x_i^3 x_j^4 x_k^7, x_i^2 x_j x_k^2 x_l^2 x_m^7,$
 $x_i^2 x_j^5 x_k^2 x_l^2 x_m^3, x_i^2 x_j^4 x_k^2 x_l^2 x_m^3, x_i^2 x_j x_k^2 x_l^2 x_m^3, i < j;$
- (ii) $x_i x_j^6 x_k^3 x_l^2 x_m^2, j < k, x_i^3 x_j^6 x_k^5, x_i^6 x_j^3 x_k^5, x_i^2 x_j x_k x_l^3 x_m^3, x_i^2 x_j x_k x_l^2, i < j < k;$
- (iii) $x_i^2 x_j x_k^4 x_l^3 x_m^4, x_i^2 x_j^4 x_k x_l^3 x_m^4, i < j < k, l < m.$

Lemma 3.4.14. *The following monomials are strictly inadmissible:*

$$\begin{array}{cccc}
x_1 x_2^2 x_3^2 x_4^7 x_5^{10}, & x_1 x_2^2 x_3^4 x_4^3 x_5^{12}, & x_1 x_2^2 x_3^4 x_4^{11} x_5^4, & x_1 x_2^2 x_3^7 x_4^2 x_5^{10}, \\
x_1 x_2^2 x_3^7 x_4^8 x_5^4, & x_1 x_2^2 x_3^7 x_4^{10} x_5^2, & x_1 x_2^2 x_3^{12} x_4^3 x_5^4, & x_1 x_2^6 x_3 x_4^6 x_5^8, \\
x_1 x_2^6 x_3 x_4^{10} x_5^4, & x_1 x_2^6 x_3^3 x_4^6 x_5^6, & x_1 x_2^6 x_3^6 x_4^3 x_5^6, & x_1 x_2^6 x_3^6 x_4^6 x_5^3, \\
x_1 x_2^6 x_3^9 x_4^2 x_5^4, & x_1 x_2^7 x_3^2 x_4^2 x_5^{10}, & x_1 x_2^7 x_3^2 x_4^8 x_5^4, & x_1 x_2^7 x_3^2 x_4^{10} x_5^2, \\
x_1 x_2^7 x_3^8 x_4^2 x_5^4, & x_1 x_2^7 x_3^{10} x_4^2 x_5^2, & x_1^3 x_2^3 x_4^4 x_5^8, & x_1^3 x_2^3 x_4^8 x_5^4, \\
x_1^3 x_2^4 x_3^2 x_4^{12} x_5^4, & x_1^3 x_2^4 x_3^4 x_4^{10} x_5^4, & x_1^3 x_2^4 x_3^4 x_4^7 x_5^6, & x_1^3 x_2^4 x_3^4 x_4^5 x_5^6, \\
x_1^3 x_2^4 x_3^4 x_4^7 x_5^4, & x_1^3 x_2^4 x_3^5 x_4^4 x_5^6, & x_1^3 x_2^4 x_3^5 x_4^6 x_5^4, & x_1^3 x_2^4 x_3^7 x_4^4 x_5^4, \\
x_1^3 x_2^4 x_3^9 x_4^2 x_5^4, & x_1^3 x_2^5 x_3^4 x_4^4 x_5^6, & x_1^3 x_2^5 x_3^4 x_6^4 x_5^4, & x_1^3 x_2^5 x_3^6 x_4^4 x_5^4, \\
x_1^3 x_2^7 x_3^4 x_4^4 x_5^4, & x_1^3 x_2^{12} x_3 x_4^2 x_5^4, & x_1^7 x_2 x_3^2 x_4^{10} x_5^2, & x_1^7 x_2 x_3^2 x_4^8 x_5^4, \\
x_1^7 x_2 x_3^2 x_4^{10} x_5^2, & x_1^7 x_2 x_3^8 x_4^2 x_5^4, & x_1^7 x_2 x_3^{10} x_4^2 x_5^2, & x_1^7 x_2^3 x_4^4 x_5^4, \\
x_1^7 x_2^8 x_3^2 x_4^4 x_5^4, & x_1^7 x_2^9 x_3^2 x_4^2 x_5^2. & &
\end{array}$$

Proof. We prove the lemma for the monomials $x = x_1 x_2^2 x_3^4 x_4^3 x_5^{12}$, and $y = x_1^3 x_2^3 x_4^4 x_5^8$. The others can be proved by a similar computation. By a direct computation using the Cartan formula, we have

$$\begin{aligned}
x &= x_1 x_2^2 x_3^4 x_4^3 x_5^{12} + x_1 x_2^2 x_3^4 x_4^2 x_5^{13} + x_1 x_2^2 x_3^2 x_4^4 x_5^{13} + x_1 x_2 x_3^6 x_4^6 x_5^8 + x_1 x_2 x_3^6 x_4^4 x_5^{10} \\
&\quad + x_1 x_2 x_3^4 x_4^6 x_5^{10} + Sq^1(f_1) + Sq^2(f_2) + Sq^4(f_4) \text{ modulo } (\mathcal{P}_5^-(\omega_{(1)})), \text{ where} \\
f_1 &= x_1 x_2^4 x_3^4 x_4^9 x_5^9 + x_1 x_2^4 x_3^3 x_4^5 x_5^8 + x_1 x_2^4 x_3^4 x_4^3 x_5^9 + x_1 x_2^4 x_3^4 x_4^5 x_5^7 \\
&\quad + x_1 x_2^4 x_3^5 x_4^3 x_5^8 + x_1 x_2^4 x_3^5 x_4^4 x_5^7 + x_1^2 x_2 x_3^4 x_4^5 x_5^9 + x_1^2 x_2 x_3^5 x_4^4 x_5^9 \\
&\quad + x_1^2 x_2 x_3^5 x_4^5 x_5^8 + x_1^4 x_2^4 x_3^3 x_4^3 x_5^7, \\
f_2 &= x_1 x_2 x_3^4 x_4^5 x_5^9 + x_1 x_2 x_3^5 x_4^4 x_5^9 + x_1 x_2 x_3^5 x_4^5 x_5^8 + x_1 x_2^2 x_3^3 x_4^4 x_5^{10} \\
&\quad + x_1 x_2^2 x_3^3 x_4^6 x_5^8 + x_1 x_2^2 x_3^4 x_4^3 x_5^{10} + x_1 x_2^2 x_3^4 x_4^6 x_5^7 + x_1 x_2^2 x_3^6 x_4^3 x_5^8 \\
&\quad + x_1 x_2^2 x_3^6 x_4^4 x_5^7 + x_1 x_2^4 x_3^2 x_4^2 x_5^{11} + x_1^2 x_2^4 x_3^3 x_4^3 x_5^8 + x_1^2 x_2^4 x_3^3 x_4^4 x_5^7 \\
&\quad + x_1^2 x_2^4 x_3^4 x_4^3 x_5^7, \\
f_4 &= x_1 x_2^2 x_3^2 x_4^2 x_5^{11} + x_1 x_2^2 x_3^4 x_4^4 x_5^7.
\end{aligned}$$

The above equalities show that x is strictly inadmissible. By a similar computation, we obtain

$$\begin{aligned}
y &= x_1^2 x_2 x_3^2 x_4^{13} x_5^4 + x_1^2 x_2 x_3^3 x_4^{12} x_5^4 + x_1^2 x_2 x_3^4 x_4^7 x_5^8 + x_1^2 x_2 x_3^4 x_4^{13} x_5^2 + x_1^2 x_2 x_3^5 x_4^6 x_5^8 \\
&\quad + x_1^2 x_2 x_3^6 x_4^9 x_5^4 + x_1^2 x_2 x_3^8 x_4^7 x_5^4 + x_1^2 x_2 x_3^{12} x_4^5 x_5^2 + x_1^2 x_3^5 x_4^8 x_5^4 + x_1^2 x_3^8 x_4^5 x_5^4 \\
&\quad + x_1^2 x_2^5 x_3^2 x_4^5 x_5^8 + x_1^2 x_2^5 x_3^2 x_4^9 x_5^4 + x_1^2 x_2^5 x_3^3 x_4^8 x_5^4 + x_1^2 x_2^5 x_3^4 x_4^3 x_5^8 + x_1^2 x_2^5 x_3^4 x_4^9 x_5^2 \\
&\quad + x_1^2 x_2^5 x_3^8 x_4^5 x_5^2 + x_1^3 x_2^2 x_3^2 x_4^{12} x_5^4 + x_1^3 x_2^2 x_3^4 x_4^6 x_5^8 + x_1^3 x_2^2 x_3^4 x_4^{10} x_5^4 + x_1^3 x_2^2 x_3^4 x_4^{12} x_5^2 \\
&\quad + x_1^3 x_2^2 x_3^4 x_4^{12} x_5^2 + x_1^3 x_2^2 x_3^4 x_4^9 x_5^4 + x_1^3 x_2^2 x_3^8 x_4^5 x_5^4 \\
&\quad + Sq^1(g_1) + Sq^2(g_2) + Sq^4(g_4) + Sq^8(x_1^3 x_2^2 x_3^2 x_4^4 x_5^2) \text{ modulo } (\mathcal{P}_5^-(\omega_{(1)})), \text{ where} \\
g_1 &= x_1^3 x_2^3 x_3^2 x_4^9 x_5^4 + x_1^3 x_2^3 x_3^4 x_4^9 x_5^2 + x_1^3 x_2^3 x_3^2 x_4^5 x_5^8 + x_1^3 x_2^3 x_3^8 x_4^5 x_5^2 + x_1^5 x_2 x_3^2 x_4^9 x_5^4 \\
&\quad + x_1^5 x_2 x_3^4 x_4^9 x_5^4 + x_1^5 x_2 x_3^4 x_4^9 x_5^2 + x_1^5 x_2 x_3^8 x_4^5 x_5^4 + x_1^5 x_2 x_3^6 x_4^5 x_5^4 \\
&\quad + x_1^5 x_2^4 x_3^5 x_4^5 x_5^4 + x_1^5 x_2^5 x_3^6 x_4^4 x_5^4 + x_1^5 x_2^5 x_3^8 x_4^4 x_5^4 + x_1^5 x_2^5 x_3^4 x_4^5 x_5^8 \\
&\quad + x_1^5 x_2^4 x_3^4 x_4^3 x_5^8 + x_1^3 x_2^3 x_3^4 x_4^3 x_5^8,
\end{aligned}$$

$$\begin{aligned}
g_2 &= x_1^5 x_2^3 x_3^2 x_4^8 x_5^2 + x_1^5 x_2^3 x_3^2 x_4^6 x_5^4 + x_1^5 x_2^3 x_3^4 x_4^6 x_5^2 + x_1^2 x_2^3 x_3^2 x_4^9 x_5^4 + x_1^3 x_2^2 x_3^2 x_4^9 x_5^4 \\
&\quad + x_1^3 x_2 x_3^2 x_4^{10} x_5^4 + x_1^6 x_2 x_3^2 x_4^7 x_5^4 + x_1^2 x_2^3 x_3^4 x_4^9 x_5^2 + x_1^3 x_2^2 x_3^4 x_4^9 x_5^2 + x_1^3 x_2 x_3^4 x_4^{10} x_5^2 \\
&\quad + x_1^6 x_2 x_3^3 x_4^7 x_5^2 + x_1^2 x_2^3 x_3^8 x_4^5 x_5^2 + x_1^2 x_2^3 x_3^2 x_4^5 x_5^8 + x_1^3 x_2^2 x_3^8 x_4^5 x_5^2 + x_1^3 x_2^2 x_3^2 x_4^5 x_5^8 \\
&\quad + x_1^3 x_2 x_3^8 x_4^5 x_5^2 + x_1^6 x_2 x_3^6 x_4^5 x_5^2 + x_1^3 x_2^2 x_3^6 x_4^5 x_5^4 + x_1^3 x_2^2 x_3^5 x_4^6 x_5^4 + x_1^3 x_2^2 x_3^3 x_4^8 x_5^4 \\
&\quad + x_1^2 x_2^3 x_3^8 x_4^5 x_5^4 + x_1^6 x_2 x_3^2 x_4^8 x_5^4 + x_1^3 x_2 x_3^2 x_4^6 x_5^8 + x_1^3 x_2^2 x_3^4 x_4^8 x_5^4 + x_1^6 x_2 x_3^3 x_4^6 x_5^4 \\
&\quad + x_1^2 x_2^3 x_4^8 x_5^4 + x_1^2 x_2 x_3^{10} x_4^5 x_5^2 + x_1^2 x_2 x_3^2 x_4^7 x_5^8 + x_1^2 x_2 x_3^2 x_4^{11} x_5^4 + x_1^2 x_2 x_3^3 x_4^6 x_5^8 \\
&\quad + x_1^2 x_2 x_3^3 x_4^{10} x_5^4 + x_1^2 x_2 x_3^4 x_4^{11} x_5^2 + x_1^2 x_2 x_3^6 x_4^9 x_5^2 + x_1^2 x_2 x_3^8 x_4^7 x_5^2, \\
g_4 &= x_1^3 x_2^3 x_3^2 x_4^6 x_5^4 + x_1^3 x_2^3 x_3^4 x_4^6 x_5^2 + x_1^4 x_2 x_3^2 x_4^7 x_5^4 + x_1^4 x_2 x_3^4 x_4^7 x_5^2 + x_1^4 x_2 x_3^6 x_4^5 x_5^2 + x_1^4 x_2 x_3^3 x_4^6 x_5^4.
\end{aligned}$$

The above relations imply that y is also strictly inadmissible. The lemma is proved. $\square$

Proof of Proposition 3.4.11. We denote by $\mathcal{Y}_t := \mathcal{Y}_{22,t}$, $461 \leq t \leq 510$ the admissible monomials in $\mathcal{B}_5^+(\omega_{(1)})$ (see Sect.6.5.) For $x \in \mathcal{B}_5^+(\omega_{(1)})$, we have $x = X_{\{i,j,k\}} y^2$ with y a monomial of degree 10 in $\mathcal{P}_5$, and $1 \leq i < j < k \leq 5$. Since x is admissible, by Theorem 2.2.1, $y \in \mathcal{B}_5(2, 2, 1)$.

Let $z \in \mathcal{B}_5(2, 2, 1)$ such that $X_{\{i,j,k\},5} y_1^2 \in \mathcal{P}_5^+$. By a direct computation using a result in [55], we see that if $X_{\{i,j,k\},5} y_1^2 \neq \mathcal{Y}_t$, for all t , $461 \leq t \leq 510$, then there is a monomial w which is given in one of Lemmas 3.4.13 - 3.4.14 such that $X_{\{i,j,k\},5} y_1^2 = w z^{2^u}$ with suitable monomial $z \in \mathcal{P}_5$ and $u = \max\{j \in \mathbb{Z} : \omega_j(w) > 0\}$. By Theorem 2.2.1, $X_{\{i,j,k\},5} y_1^2$ is inadmissible. Since $x = X_{\{i,j,k\},5} y^2$ and x is admissible, one gets $x = \mathcal{Y}_t$. This implies $Q\mathcal{P}_5^+(\omega_{(1)})$ is spanned by the set $\{[\mathcal{Y}_t := \mathcal{Y}_{22,t}]_{\omega_{(1)}} : 461 \leq t \leq 510\}$.

We now prove the set $\{[\mathcal{Y}_t]_{\omega_{(1)}} : 461 \leq t \leq 510\}$ is linearly independent in $Q\mathcal{P}_5(\omega_{(1)})$. Suppose there is a linear relation

$$(3.2) \quad \mathcal{S} = \sum_{461 \leq t \leq 510} \gamma_t \mathcal{Y}_t \equiv_{\omega_{(1)}} 0,$$

where $\gamma_t \in \mathbb{Z}/2$. From a result in [47], $\dim Q\mathcal{P}_4^+(\omega_{(1)}) = 26$, with the basis $\{[u_j]_{\omega_{(1)}} : 1 \leq j \leq 26\}$, where

$$\begin{aligned}
&u_1. x_1 x_2 x_3^6 x_4^{14}, & u_2. x_1 x_2 x_3^{14} x_4^6, & u_3. x_1 x_2^2 x_3^4 x_4^{15}, & u_4. x_1 x_2^2 x_3^5 x_4^{14}, & u_5. x_1 x_2^2 x_3^7 x_4^{12}, \\
&u_6. x_1 x_2^2 x_3^{12} x_4^7, & u_7. x_1 x_2^2 x_3^{13} x_4^6, & u_8. x_1 x_2^2 x_3^{15} x_4^4, & u_9. x_1 x_2^3 x_3^4 x_4^{14}, & u_{10}. x_1 x_2^3 x_3^6 x_4^{12}, \\
&u_{11}. x_1 x_2^3 x_3^{12} x_4^6, & u_{12}. x_1 x_2^3 x_3^{14} x_4^4, & u_{13}. x_1 x_2^6 x_3^3 x_4^{14}, & u_{14}. x_1 x_2^7 x_3^2 x_4^{12}, & u_{15}. x_1 x_2^{14} x_3 x_4^6, \\
&u_{16}. x_1 x_2^{15} x_3^2 x_4^4, & u_{17}. x_1^3 x_2 x_3^4 x_4^{14}, & u_{18}. x_1^3 x_2 x_3^6 x_4^{12}, & u_{19}. x_1^3 x_2 x_3^{12} x_4^6, & u_{20}. x_1^3 x_2 x_3^{14} x_4^4, \\
&u_{21}. x_1^3 x_2^5 x_3^2 x_4^{12}, & u_{22}. x_1^3 x_2^5 x_3^6 x_4^8, & u_{23}. x_1^3 x_2^5 x_3^{10} x_4^4, & u_{24}. x_1^3 x_2^{13} x_3^2 x_4^4, & u_{25}. x_1^7 x_2 x_3^2 x_4^{12}, \\
&u_{26}. x_1^{15} x_2 x_3^2 x_4^4.
\end{aligned}$$

Consider the homomorphism $\pi_{(1;2)} : \mathcal{P}_5 \rightarrow \mathcal{P}_4$. By a direct computation using Theorem 3.3.5 and Proposition 3.1.1, we have

$$\begin{aligned}
\pi_{(1;2)}(\mathcal{S}) &\equiv_{\omega_{(1)}} (\gamma_{473} + \gamma_{475} + \gamma_{481} + \gamma_{483})u_1 + \gamma_{477}u_6 + \gamma_{478}u_7 \\
&\quad + (\gamma_{476} + \gamma_{477} + \gamma_{478})u_4 + \gamma_{483}(u_9 + u_{14}) + \gamma_{481}(u_{10} + u_{11}) \\
&\quad + (\gamma_{475} + \gamma_{476} + \gamma_{477} + \gamma_{478})u_{13} + (\gamma_{470} + \gamma_{481} + \gamma_{484})u_{18} \\
&\quad + (\gamma_{471} + \gamma_{478} + \gamma_{481} + \gamma_{484})u_{19} + \gamma_{472}u_{20} + (\gamma_{479} + \gamma_{484})u_{21} \\
&\quad + (\gamma_{480} + \gamma_{481} + \gamma_{484})u_{22} + \gamma_{482}u_{23} + \gamma_{485}u_{24} + \gamma_{495}u_{25} + \gamma_{497}u_{26} \\
&\quad + (\gamma_{474} + \gamma_{478} + \gamma_{481} + \gamma_{484})u_2 + \gamma_{476}u_5 + (\gamma_{469} + \gamma_{475})u_{17} + \gamma_{484}u_{15} \equiv_{\omega_{(1)}} 0.
\end{aligned}$$

This relation implies

$$\begin{aligned}
(3.3) \quad \gamma_{469} &= \gamma_{470} = \gamma_{471} = \gamma_{472} = \gamma_{473} = \gamma_{474} = \gamma_{475} = \gamma_{476} \\
&= \gamma_{477} = \gamma_{478} = \gamma_{479} = \gamma_{480} = \gamma_{481} = \gamma_{482} = \gamma_{483} \\
&= \gamma_{484} = \gamma_{485} = \gamma_{495} = \gamma_{497} = 0.
\end{aligned}$$

Substituting (3.3) into the relation (3.2), we have

$$(3.4) \quad \sum_{461 \leq t \leq 468} \gamma_t \mathcal{Y}_t + \sum_{486 \leq t \leq 494} \gamma_t \mathcal{Y}_t + \gamma_{496} \mathcal{Y}_{36} + \sum_{498 \leq t \leq 510} \gamma_t \mathcal{Y}_t \equiv_{\omega_{(1)}} 0.$$

Applying the homomorphisms $\pi_{(1;3)}, \pi_{(1;4)} : \mathcal{P}_5 \rightarrow \mathcal{P}_4$ to (3.4), we get

$$(3.5) \quad \begin{cases} \gamma_t = 0, \quad t \in \mathbb{J}, \\ \gamma_{464} = \gamma_{487} = \gamma_{499} = \gamma_{508}, \gamma_{468} = \gamma_{494} = \gamma_{506} = \gamma_{509}, \gamma_{498} = \gamma_{502}, \\ \gamma_{467} + \gamma_{468} + \gamma_{503} = \gamma_{467} + \gamma_{468} + \gamma_{505} = 0, \\ \gamma_{462} + \gamma_{468} + \gamma_{502} + \gamma_{508} = \gamma_{463} + \gamma_{468} + \gamma_{502} + \gamma_{508} = 0, \\ \gamma_{466} + \gamma_{501} + \gamma_{502} + \gamma_{508} = \gamma_{466} + \gamma_{468} + \gamma_{504} + \gamma_{507} + \gamma_{508} = 0. \end{cases}$$

Here $\mathbb{J} = \{461, 465, 486, 488, 489, 490, 491, 492, 493, 496, 500\}$. Then, combining (3.3), (3.5), and the relation $\pi_{(1;5)}(\mathcal{S}) \equiv_{\omega_{(1)}} 0$, we obtain $\gamma_t = 0$ for $461 \leq t \leq 510$. The proposition is proved. $\square$

Using a similar technique as mentioned in the proof of Proposition 3.4.11, we obtain

Proposition 3.4.15.

(I) $\mathcal{B}_5^+(\omega_{(2)}) = \mathcal{B}^+(5, \omega_{(2)}) \cup \mathcal{E}$, where $\mathcal{E}$ is the set of the following monomials:

$$\begin{aligned} & x_1 x_2^2 x_3^2 x_4^3 x_5^{14}, \quad x_1 x_2^2 x_3^3 x_4^2 x_5^{14}, \quad x_1 x_2^2 x_3^3 x_4^6 x_5^{10}, \quad x_1 x_2^2 x_3^3 x_4^{14} x_5^2, \quad x_1 x_2^3 x_3^2 x_4^2 x_5^{14}, \\ & x_1 x_2^3 x_3^2 x_4^6 x_5^{10}, \quad x_1 x_2^3 x_3^2 x_4^{14} x_5^2, \quad x_1 x_2^3 x_3^6 x_4^2 x_5^{10}, \quad x_1 x_2^3 x_3^6 x_4^{10} x_5^2, \quad x_1 x_2^3 x_3^{14} x_4^2 x_5^2, \\ & x_1^3 x_2 x_3^2 x_4^2 x_5^{14}, \quad x_1^3 x_2 x_3^2 x_4^6 x_5^{10}, \quad x_1^3 x_2 x_3^2 x_4^{14} x_5^2, \quad x_1^3 x_2 x_3^6 x_4^2 x_5^{10}, \quad x_1^3 x_2 x_3^6 x_4^{10} x_5^2, \\ & x_1^3 x_2 x_3^{14} x_4^2 x_5^2, \quad x_1^3 x_2^5 x_3^2 x_4^2 x_5^{10}, \quad x_1^3 x_2^5 x_3^2 x_4^{10} x_5^2, \quad x_1^3 x_2^5 x_3^{10} x_4^2 x_5^2, \quad x_1^3 x_2^{13} x_3^2 x_4^2 x_5^2, \end{aligned}$$

(II) $\mathcal{B}_5^+(\omega_{(3)}) = \{x_1 x_2^3 x_3^6 x_4^6 x_5^6, x_1^3 x_2 x_3^6 x_4^6 x_5^6, x_1^3 x_2^5 x_3^6 x_4^6 x_5^6, x_1^3 x_2^5 x_3^6 x_4^2 x_5^6, x_1^3 x_2^5 x_3^6 x_4^6 x_5^2\}$,

(III) $\mathcal{B}_5^+(\omega_{(4)}) = \overline{\Phi}^+(\mathcal{B}_4(\omega_{(4)}) \cup \mathcal{B}^+(5, \omega_{(4)}))$,

(IV) $\mathcal{B}_5^+(\omega_{(5)}) = \overline{\Phi}^+(\mathcal{B}_4(\omega_{(5)}) \cup \mathcal{B}^+(5, \omega_{(5)}))$.

A direct computation shows: $|\mathcal{B}_5^+(\omega_{(2)})| = 25$, $|\mathcal{B}_5^+(\omega_{(4)})| = 300$ and $|\mathcal{B}_5^+(\omega_{(5)})| = 125$ (see Sect. 6.5.) On the other hand, by Remark 3.4.9, we have $(Q\mathcal{P}_5)_{22}^+ \cong \bigoplus_{1 \leq j \leq 5} Q\mathcal{P}_5^+(\omega_{(j)})$. Combining this with the above results, we obtain

Corollary 3.4.16. $(Q\mathcal{P}_5^+)_{22}$ is the $\mathbb{Z}/2$ -vector space of dimension 505 with a basis consisting of all the classes represented by the monomials $\mathcal{Y}_{22,t}$, $1 \leq t \leq 505$, which are determined in Sect. 6.5.

Computation of $\text{Ker}((\widetilde{Sq}_*^0)_{(5,47)})$

We consider the weight vectors

$$\overline{\omega}_{(1)} = (3, 2, 2, 2, 1), \quad \overline{\omega}_{(2)} = (3, 2, 4, 1, 1), \quad \overline{\omega}_{(3)} = (3, 2, 4, 3),$$

$$\overline{\omega}_{(4)} = (3, 4, 3, 1, 1), \quad \overline{\omega}_{(5)} = (3, 4, 3, 3).$$

It is easy to see that $\deg \overline{\omega}_{(1)} = \deg \overline{\omega}_{(2)} = \dots = \deg \overline{\omega}_{(5)} = 47$. Then, from Remarks 3.4.8 and 3.4.9, one gets the following.

Lemma 3.4.17. If X is an admissible monomial of degree 47 in $\mathcal{P}_5$ and $[X] \in \text{Ker}((\widetilde{Sq}_*^0)_{(5,47)})$, then $\omega(X)$ is one of the sequences $\overline{\omega}_{(k)}$, $1 \leq k \leq 5$.

From this lemma, we have

$$\begin{aligned} \text{Ker}((\widetilde{Sq}_*^0)_{(5,47)}) & \cong (Q\mathcal{P}_5^0)_{47} \oplus (\text{Ker}((\widetilde{Sq}_*^0)_{(5,47)}) \cap (Q\mathcal{P}_5^+)_{47}) \\ & \cong (Q\mathcal{P}_5^0)_{47} \oplus (\bigoplus_{1 \leq i \leq 5} Q\mathcal{P}_5^+(\overline{\omega}_{(i)})). \end{aligned}$$

Recall that $(Q\mathcal{P}_5^0)_{47} = \bigoplus_{1 \leq t \leq 4} \bigoplus_{1 \leq \ell \leq \binom{5}{t}} (Q\mathcal{P}_t^+)_{47}$. Then, by a direct computation using a result in [15], [32] and [47], we conclude

$$\dim(Q\mathcal{P}_5^0)_{47} = \binom{5}{3} \cdot 14 + \binom{5}{4} \cdot 84 = 560.$$

The set $\mathcal{B}_5^0(47) = \mathcal{B}_5^0(\overline{\omega}_{(1)}) = \overline{\Phi}^0(\mathcal{B}_4(47)) = \{\mathcal{Y}_{47,i} : 1 \leq i \leq 560\}$ is described in Sect. 6.6. We now determine $Q\mathcal{P}_5^+(\overline{\omega}_{(j)})$ for $1 \leq j \leq 5$.

Lemma 3.4.18. *The following monomials are strictly inadmissible:*

- I) $X_1 = x_1^3 x_2^{12} x_k x_\ell^3 x_m^{12}$, $X_2 = x_1^3 x_2^4 x_k^3 x_\ell^8 x_m^{13}$, $X_3 = x_1 x_2^{14} x_k^2 x_\ell x_m^{13}$,
 $X_5 = x_1^3 x_2^{14} x_k^{12} x_\ell x_m^{13}$, $X_6 = x_1^7 x_2^{10} x_k^{12} x_\ell x_m^{13}$, $X_7 = x_1^3 x_2^2 x_k^{12} x_\ell x_m^{13}$,
 $X_8 = x_1^3 x_2^{12} x_k^2 x_\ell x_m^{13}$, $X_9 = x_1^{15} x_2^2 x_k x_\ell^4 x_m^9$, $X_{10} = x_1^{15} x_2^2 x_k x_\ell^5 x_m^8$,
 $X_{11} = x_1^7 x_2^2 x_k x_\ell^8 x_m^{13}$, $X_{12} = x_1^7 x_2^2 x_k x_\ell^9 x_m^{12}$, $X_{13} = x_1^{15} x_2^2 x_k^{12} x_\ell x_m^{13}$,
 $X_{14} = x_1^3 x_2^4 x_k x_\ell^8 x_m^{15}$, $X_{15} = x_1^3 x_2^4 x_k x_\ell^9 x_m^{14}$, $X_{16} = x_1^3 x_2^{14} x_k x_\ell^4 x_m^9$,
 $X_{17} = x_1^3 x_2^5 x_k x_\ell^8 x_m^{14}$, $X_{18} = x_1^3 x_2^6 x_k x_\ell^8 x_m^{13}$, $X_{19} = x_1^3 x_2^6 x_k x_\ell^9 x_m^{12}$,
 $X_{20} = x_1^7 x_2^{10} x_k x_\ell^4 x_m^9$, $X_{21} = x_1^7 x_2^{10} x_k x_\ell^5 x_m^8$, $X_{22} = x_1^3 x_2^2 x_k^5 x_\ell^8 x_m^{13}$,
 $X_{23} = x_1^3 x_2^2 x_k^5 x_\ell^9 x_m^{12}$, $X_{24} = x_1^7 x_2^2 x_k^5 x_\ell^8 x_m^9$, $X_{25} = x_1 x_2^2 x_k^{14} x_\ell x_m^{13}$,
 $X_{26} = x_1^3 x_2^2 x_k^5 x_\ell^8 x_m^9$, $X_{27} = x_1^3 x_2^{12} x_k^{14} x_\ell x_m^{13}$. Here (k, ℓ, m) is a permutation of $(3, 4, 5)$;
II) $X_{27} = x_i^3 x_j^2 x_k^{13} x_\ell^4 x_m^9$, $j < k$, $X_{28} = x_i x_j^2 x_k^6 x_\ell^9 x_m^{13}$, $X_{29} = x_i^3 x_j^4 x_k^6 x_\ell^9 x_m^9$, where (i, j, k, ℓ, m) is a permutation of $(1, 2, 3, 4, 5)$.

Proof. It is easy to see that $\omega(X_t) = \omega^* := (3, 2, 2, 2)$ for $1 \leq t \leq 29$. We prove the lemma for the monomials $X_1 = x_1^3 x_2^{12} x_k x_\ell^3 x_m^{12}$ and $X_2 = x_1^3 x_2^4 x_k^3 x_\ell^8 x_m^{13}$, where (k, ℓ, m) is a permutation of $(3, 4, 5)$. The others can be proved by a similar technique. We have

$$\begin{aligned} X_1 &= x_1^2 x_2^{11} x_k x_\ell^5 x_m^{12} + x_1^2 x_2^{11} x_k^4 x_\ell^5 x_m^9 + x_1^2 x_2^{11} x_k^8 x_\ell^5 x_m^5 + x_1^2 x_2^{13} x_k x_\ell^3 x_m^{12} \\ &\quad + x_1^2 x_2^{13} x_k^4 x_\ell^3 x_m^9 + x_1^2 x_2^{13} x_k^8 x_\ell^3 x_m^5 + x_1^3 x_2^3 x_k^8 x_\ell^5 x_m^{12} + x_1^3 x_2^5 x_k^8 x_\ell^5 x_m^{10} \\ &\quad + x_1^3 x_2^5 x_k^8 x_\ell^6 x_m^9 + x_1^3 x_2^7 x_k^4 x_\ell^8 x_m^9 + x_1^3 x_2^7 x_k^8 x_\ell^5 x_m^8 + x_1^3 x_2^7 x_k^8 x_\ell^8 x_m^5 \\ &\quad + x_1^3 x_2^9 x_k^2 x_\ell^5 x_m^{12} + x_1^3 x_2^9 x_k^4 x_\ell^3 x_m^{12} + x_1^3 x_2^9 x_k^4 x_\ell^5 x_m^{10} + x_1^3 x_2^9 x_k^4 x_\ell^6 x_m^9 \\ &\quad + x_1^3 x_2^9 x_k^8 x_\ell^5 x_m^6 + x_1^3 x_2^9 x_k^8 x_\ell^6 x_m^5 + x_1^3 x_2^{11} x_k x_\ell^4 x_m^{12} \\ &\quad + Sq^1(g_1) + Sq^2(g_2) + Sq^4(g_4) + Sq^8(x_1^6 x_2^5 x_k^4 x_\ell^3 x_m^5) \text{ modulo } (\mathcal{P}_5^-(\omega^*)), \text{ where} \\ g_1 &= x_1^3 x_2^7 x_k x_\ell^3 x_m^{16} + x_1^3 x_2^{11} x_k x_\ell^3 x_m^{12} + x_1^5 x_2^3 x_k^8 x_\ell^5 x_m^9 + x_1^5 x_2^7 x_k^4 x_\ell^5 x_m^9 + x_1^5 x_2^7 x_k^8 x_\ell^5 x_m^5, \\ g_2 &= x_1^2 x_2^{11} x_k x_\ell^3 x_m^{12} + x_1^2 x_2^{11} x_k^4 x_\ell^3 x_m^9 + x_1^2 x_2^{11} x_k^8 x_\ell^3 x_m^5 + x_1^3 x_2^3 x_k^8 x_\ell^5 x_m^{10} + x_1^3 x_2^3 x_k^8 x_\ell^6 x_m^9 \\ &\quad + x_1^3 x_2^7 x_k^4 x_\ell^5 x_m^{12} + x_1^3 x_2^7 x_k^4 x_\ell^6 x_m^9 + x_1^3 x_2^7 x_k^8 x_\ell^5 x_m^6 + x_1^3 x_2^7 x_k^8 x_\ell^6 x_m^5 + x_1^5 x_2^7 x_k^2 x_\ell^3 x_m^{12} \\ &\quad + x_1^6 x_2^3 x_k^8 x_\ell^3 x_m^9 + x_1^6 x_2^7 x_k^4 x_\ell^3 x_m^9 + x_1^6 x_2^7 x_k^8 x_\ell^3 x_m^5, \\ g_4 &= x_1^3 x_2^7 x_k^2 x_\ell^3 x_m^{12} + x_1^4 x_2^7 x_k^4 x_\ell^3 x_m^9 + x_1^4 x_2^7 x_k^8 x_\ell^3 x_m^5 + x_1^{10} x_2^5 x_k^4 x_\ell^3 x_m^5. \end{aligned}$$

This equality implies that X_1 is strictly inadmissible.

Next, we show that X_2 is also strictly inadmissible. Indeed, using Cartan's formula, we obtain

$$\begin{aligned} X_2 &= x_1^2 x_2 x_k^3 x_\ell^{12} x_m^{13} + x_1^2 x_2 x_k^5 x_\ell^9 x_m^{14} + x_1^2 x_2 x_k^5 x_\ell^{10} x_m^{13} + x_1^2 x_2 x_k^6 x_\ell^9 x_m^{13} + x_1^2 x_2 x_k^{10} x_\ell^5 x_m^{13} \\ &\quad + x_1^2 x_2 x_k^{12} x_\ell^5 x_m^{11} + x_1^2 x_2^5 x_k x_\ell^9 x_m^{14} + x_1^2 x_2^5 x_k x_\ell^{10} x_m^{13} + x_1^2 x_2^5 x_k^8 x_\ell^3 x_m^{13} + x_1^2 x_2^5 x_k^8 x_\ell^5 x_m^{11} \\ &\quad + x_1^3 x_2 x_k^4 x_\ell^9 x_m^{14} + x_1^3 x_2 x_k^4 x_\ell^{10} x_m^{13} + x_1^3 x_2 x_k^6 x_\ell^8 x_m^{13} + x_1^3 x_2 x_k^8 x_\ell^6 x_m^{13} + x_1^3 x_2 x_k^4 x_\ell^9 x_m^{13} \\ &\quad + x_1^3 x_2^2 x_k^5 x_\ell^8 x_m^{13} + x_1^3 x_2^2 x_k^8 x_\ell^4 x_m^{13} + x_1^3 x_2^2 x_k^8 x_\ell^5 x_m^{12} + x_1^3 x_2^4 x_k x_\ell^9 x_m^{14} + x_1^3 x_2^4 x_k x_\ell^{10} x_m^{13} \\ &\quad + x_1^3 x_2^4 x_k^2 x_\ell^9 x_m^{13} \\ &\quad + Sq^1(Z_1) + Sq^2(Z_2) + Sq^4(Z_4) + Sq^8(x_1^3 x_2^4 x_k^4 x_\ell^5 x_m^7) \text{ modulo } (\mathcal{P}_5^-(\omega^*)), \end{aligned}$$

where

$$\begin{aligned} Z_1 &= x_1^3 x_2 x_k^3 x_\ell^5 x_m^{18} + x_1^3 x_2 x_k^3 x_\ell^6 x_m^{17} + x_1^3 x_2 x_k^3 x_\ell^9 x_m^{14} + x_1^3 x_2 x_k^3 x_\ell^{10} x_m^{13} \\ &\quad + x_1^3 x_2 x_k^6 x_\ell^9 x_m^{11} + x_1^3 x_2 x_k^{10} x_\ell^5 x_m^{11} + x_1^3 x_2^3 x_k x_\ell^6 x_m^{17} + x_1^3 x_2^3 x_k x_\ell^{10} x_m^{13} \\ &\quad + x_1^3 x_2^3 x_k x_\ell^{12} x_m^{11} + x_1^3 x_2^3 x_k x_\ell^{16} x_m^7 + x_1^3 x_2^3 x_k^4 x_\ell^3 x_m^{17} + x_1^3 x_2^3 x_k^8 x_\ell^3 x_m^{13} \\ &\quad + x_1^3 x_2^3 x_k^8 x_\ell^5 x_m^{11} + x_1^3 x_2^3 x_k^8 x_\ell^9 x_m^7 + x_1^3 x_2^4 x_k^5 x_\ell^5 x_m^{13} + x_1^5 x_2^2 x_k x_\ell^5 x_m^{14}, \\ Z_2 &= x_1^2 x_2 x_k^3 x_\ell^9 x_m^{14} + x_1^2 x_2 x_k^3 x_\ell^{10} x_m^{13} + x_1^2 x_2 x_k^6 x_\ell^9 x_m^{11} + x_1^2 x_2 x_k^{10} x_\ell^5 x_m^{11} \\ &\quad + x_1^2 x_2^3 x_k x_\ell^9 x_m^{14} + x_1^2 x_2^3 x_k x_\ell^{10} x_m^{13} + x_1^2 x_2^3 x_k x_\ell^{12} x_m^{11} + x_1^2 x_2^3 x_k^8 x_\ell^3 x_m^{13} \\ &\quad + x_1^2 x_2^3 x_k^8 x_\ell^5 x_m^{11} + x_1^2 x_2^3 x_k^8 x_\ell^9 x_m^7 + x_1^2 x_2^5 x_k x_\ell^6 x_m^{14} + x_1^2 x_2^5 x_k^2 x_\ell^5 x_m^{14} \\ &\quad + x_1^3 x_2^6 x_k x_\ell^5 x_m^{14} + x_1^3 x_2^6 x_k^6 x_\ell^{11} + x_1^5 x_2^2 x_k^2 x_\ell^9 x_m^{11} + x_1^5 x_2^2 x_k^3 x_\ell^5 x_m^{14} \\ &\quad + x_1^5 x_2^2 x_k^3 x_\ell^6 x_m^{13} + x_1^5 x_2^2 x_k^6 x_\ell^5 x_m^{11} + x_1^5 x_2^2 x_k^8 x_\ell^3 x_m^{11} + x_1^5 x_2^3 x_k x_\ell^6 x_m^{14} \\ &\quad + x_1^5 x_2^3 x_k^2 x_\ell^6 x_m^{13} + x_1^5 x_2^3 x_k^2 x_\ell^8 x_m^{11} + x_1^5 x_2^3 x_k^2 x_\ell^{10} x_m^9 + x_1^5 x_2^3 x_k^2 x_\ell^{12} x_m^7 \\ &\quad + x_1^5 x_2^3 x_k^4 x_\ell^3 x_m^{14} + x_1^5 x_2^3 x_k^4 x_\ell^6 x_m^{11} + x_1^5 x_2^3 x_k^4 x_\ell^{10} x_m^7 + x_1^6 x_2^3 x_k x_\ell^5 x_m^{14}, \end{aligned}$$

$$\begin{aligned}
Z_4 = & x_1^3 x_2 x_k^6 x_\ell^5 x_m^{12} + x_1^3 x_2 x_k^6 x_\ell^6 x_m^{11} + x_1^3 x_2^2 x_k^2 x_\ell^9 x_m^{11} + x_1^3 x_2^2 x_k^3 x_\ell^6 x_m^{13} + x_1^3 x_2^2 x_k^6 x_\ell^5 x_m^{11} \\
& + x_1^3 x_2^2 x_k^8 x_\ell^3 x_m^{11} + x_1^3 x_2^3 x_k x_\ell^6 x_m^{14} + x_1^3 x_2^3 x_k^2 x_\ell^6 x_m^{13} + x_1^3 x_2^3 x_k^2 x_\ell^8 x_m^{11} + x_1^3 x_2^3 x_k^2 x_\ell^{10} x_m^9 \\
& + x_1^3 x_2^3 x_k^2 x_\ell^{12} x_m^7 + x_1^3 x_2^3 x_k^4 x_\ell^3 x_m^{14} + x_1^3 x_2^3 x_k^4 x_\ell^6 x_m^{11} + x_1^3 x_2^3 x_k^4 x_\ell^{10} x_m^7 + x_1^3 x_2^4 x_k x_\ell^5 x_m^{14} \\
& + x_1^3 x_2^8 x_k^4 x_\ell^5 x_m^7 + x_1^4 x_2^3 x_k x_\ell^5 x_m^{14}.
\end{aligned}$$

The above relations imply that X_2 is strictly inadmissible. The lemma follows. $\square$

Lemma 3.4.19. *The following monomials are strictly inadmissible:*

$$\begin{aligned}
Y_1 &= x_1^3 x_2^4 x_3^7 x_4^8 x_5^9, & Y_2 &= x_1^3 x_2^7 x_3^8 x_4^5 x_5^8, & Y_3 &= x_1^6 x_2^3 x_3^{10} x_4 x_5^{13}, & Y_4 &= x_1 x_2^6 x_3^{10} x_4^{13} x_5, \\
Y_5 &= x_1 x_2^6 x_3^{11} x_4^{12} x_5, & Y_6 &= x_1 x_2^7 x_3^{10} x_4^{12} x_5, & Y_7 &= x_1^7 x_2 x_3^{10} x_4^{12} x_5, & Y_8 &= x_1 x_2^2 x_3^2 x_4^{13} x_5^{13}, \\
Y_9 &= x_1 x_2^2 x_3^{14} x_4^5 x_5^9, & Y_{10} &= x_1 x_2^{14} x_3^2 x_4^5 x_5^9, & Y_{11} &= x_1^3 x_2^3 x_3^{12} x_4^{12} x_5, & Y_{12} &= x_1^3 x_2^{15} x_3^4 x_4^8 x_5, \\
Y_{13} &= x_1^3 x_2^{15} x_3^4 x_4^8 x_5, & Y_{14} &= x_1^{15} x_2^3 x_3^4 x_4^8 x_5, & Y_{15} &= x_1^{15} x_2^3 x_3^4 x_4^8 x_5, & Y_{16} &= x_1 x_2^{14} x_3^3 x_4^4 x_5^9, \\
Y_{17} &= x_1^3 x_2^4 x_3^{10} x_4 x_5^{13}, & Y_{18} &= x_1^3 x_2^4 x_3^{10} x_4^{13} x_5, & Y_{19} &= x_1^3 x_2^4 x_3^{11} x_4^{12} x_5, & Y_{20} &= x_1 x_2^{14} x_3^3 x_4^5 x_5^8, \\
Y_{21} &= x_1^3 x_2^{14} x_3^5 x_4^8 x_5, & Y_{22} &= x_1^3 x_2^{14} x_3^5 x_4^8 x_5, & Y_{23} &= x_1^3 x_2^{14} x_3^5 x_4^8 x_5, & Y_{24} &= x_1 x_2^6 x_3^3 x_4^{13} x_5^8, \\
Y_{25} &= x_1 x_2^{14} x_3^{14} x_4 x_5, & Y_{26} &= x_1^3 x_2^7 x_3^{12} x_4^{12} x_5, & Y_{27} &= x_1^3 x_2^7 x_3^{12} x_4^8 x_5, & Y_{28} &= x_1^3 x_2^7 x_3^{12} x_4^8 x_5, \\
Y_{29} &= x_1^7 x_2^3 x_3^8 x_4^{12} x_5, & Y_{30} &= x_1^7 x_2^3 x_3^{12} x_4^8 x_5, & Y_{31} &= x_1^7 x_2^3 x_3^{12} x_4^8 x_5, & Y_{32} &= x_1 x_2^6 x_3^{11} x_4^4 x_5^9, \\
Y_{33} &= x_1^7 x_2^{11} x_3^4 x_4^8 x_5, & Y_{34} &= x_1^7 x_2^{11} x_3^4 x_4^8 x_5, & Y_{35} &= x_1 x_2^6 x_3^{11} x_4^5 x_5^8, & Y_{36} &= x_1 x_2^6 x_3^{10} x_4^5 x_5^9, \\
Y_{37} &= x_1^3 x_2^5 x_3^8 x_4^2 x_5^{13}, & Y_{38} &= x_1^3 x_2^{12} x_3^2 x_4^5 x_5^9, & Y_{39} &= x_1^3 x_2^5 x_3^9 x_4^2 x_5^{12}, & Y_{40} &= x_1^7 x_2^2 x_3^4 x_4^9 x_5^9, \\
Y_{41} &= x_1^7 x_2^2 x_3^9 x_4^4 x_5^9, & Y_{42} &= x_1^3 x_2^4 x_3^3 x_4^{12} x_5^9, & Y_{43} &= x_1^3 x_2^{12} x_3^3 x_4^4 x_5^9, & Y_{44} &= x_1^3 x_2^4 x_3^9 x_4^{12} x_5^{12}, \\
Y_{45} &= x_1^3 x_2^{12} x_3^3 x_4^5 x_5^8, & Y_{46} &= x_1^3 x_2^5 x_3^8 x_4^3 x_5^{12}, & Y_{47} &= x_1^3 x_2^4 x_3^{11} x_4^4 x_5^9, & Y_{48} &= x_1^3 x_2^4 x_3^{11} x_4^5 x_5^8, \\
Y_{49} &= x_1^3 x_2^4 x_3^{10} x_4^5 x_5^9, & Y_{50} &= x_1 x_2^6 x_3^3 x_4^{12} x_5^9, & Y_{51} &= x_1^3 x_2^4 x_3^7 x_4^9 x_5^8, & Y_{52} &= x_1^3 x_2^7 x_3^8 x_4^4 x_5^9, \\
Y_{53} &= x_1^3 x_2^3 x_3^8 x_4^4 x_5^9, & Y_{54} &= x_1 x_2^{14} x_3^3 x_4^{12} x_5, & Y_{55} &= x_1^7 x_2^3 x_3^8 x_4^5 x_5^8.
\end{aligned}$$

Proof. Note that $\omega(Y_j) = \omega^*$, $j = 1, 2, \dots, 55$. We prove this lemma for the monomials $Y_1 = x_1^3 x_2^4 x_3^7 x_4^8 x_5^9$, and $Y_2 = x_1^3 x_2^7 x_3^8 x_4^5 x_5^8$. The others can be proved by a similar computation. A direct computation shows:

$$\begin{aligned}
Y_1 &= x_1^2 x_2 x_3^7 x_4^8 x_5^{13} + x_1^2 x_2 x_3^{13} x_4 x_5^{14} + x_1^2 x_2 x_3^{13} x_4^8 x_5^7 + x_1^2 x_3^9 x_4^4 x_5^{13} + x_1^2 x_3^{12} x_4 x_5^{13} \\
&+ x_1^2 x_3^9 x_4^4 x_5^{11} + x_1^2 x_3^9 x_4^8 x_5^{11} + x_1^2 x_3^{12} x_4 x_5^{11} + x_1^3 x_2^2 x_3^5 x_4^8 x_5^{14} + x_1^3 x_2^2 x_3^7 x_4^8 x_5^{12} \\
&+ x_1^3 x_2^2 x_3^{12} x_4 x_5^{14} + x_1^3 x_2^2 x_3^{12} x_4^4 x_5^{11} + x_1^3 x_2^2 x_3^9 x_4^4 x_5^{13} + x_1^3 x_2^2 x_3^9 x_4^4 x_5^{12} + x_1^3 x_2^2 x_3^{12} x_4 x_5^{12} \\
&+ x_1^3 x_2^3 x_3^{12} x_4^4 x_5^9 + Sq^1(u_1) + Sq^2(u_2) + Sq^4(u_4) + Sq^8(u_8) \text{ modulo } (\mathcal{P}_5^-(\omega^*)), \text{ where} \\
u_1 &= x_1^3 x_2 x_3^5 x_4^8 x_5^{13} + x_1^3 x_2 x_3^7 x_4 x_5^{18} + x_1^3 x_2 x_3^9 x_4^4 x_5^{13} + x_1^3 x_2 x_3^{11} x_4 x_5^{14} \\
&+ x_1^3 x_2^3 x_3^9 x_4^4 x_5^{11} + x_1^3 x_2^3 x_3^9 x_4^8 x_5^7 + x_1^3 x_2^3 x_3^{12} x_4 x_5^{11} + x_1^3 x_2^3 x_3^{16} x_4 x_5^7 \\
&+ x_1^3 x_2^4 x_3^9 x_4^4 x_5^{13} + x_1^5 x_2 x_3^5 x_4^8 x_5^{11} + x_1^5 x_2 x_3^7 x_4^8 x_5^9, \\
u_2 &= x_1^2 x_2 x_3^7 x_4^8 x_5^{11} + x_1^2 x_2 x_3^{11} x_4 x_5^{14} + x_1^2 x_2 x_3^{11} x_4^8 x_5^7 + x_1^2 x_3^9 x_4^4 x_5^{11} \\
&+ x_1^2 x_3^9 x_4^8 x_5^{11} + x_1^2 x_3^{12} x_4 x_5^{11} + x_1^3 x_2^2 x_3^6 x_4^8 x_5^{11} + x_1^3 x_2^2 x_3^7 x_4^8 x_5^{10} \\
&+ x_1^3 x_2^2 x_3^{10} x_4 x_5^{11} + x_1^3 x_2^2 x_3^5 x_4^8 x_5^{11} + x_1^3 x_2^2 x_3^7 x_4^8 x_5^9 + x_1^3 x_2^2 x_3^9 x_4^4 x_5^{11} \\
&+ x_1^5 x_2 x_3^7 x_4^2 x_5^{14} + x_1^5 x_2 x_3^7 x_4^2 x_5^{14} + x_1^5 x_2^2 x_3^9 x_4^2 x_5^{11} + x_1^5 x_2^2 x_3^{10} x_4^2 x_5^9 \\
&+ x_1^5 x_2^3 x_3^{10} x_4 x_5^7 + x_1^5 x_2^3 x_3^{12} x_4^2 x_5^7 + x_1^6 x_2 x_3^7 x_4^8 x_5^7, \\
u_4 &= x_1^3 x_2 x_3^7 x_4^2 x_5^{14} + x_1^3 x_2^2 x_3^7 x_4 x_5^{14} + x_1^3 x_2^2 x_3^9 x_4^2 x_5^{11} + x_1^3 x_2^2 x_3^{10} x_4 x_5^{11} \\
&+ x_1^3 x_2^3 x_3^{10} x_4^2 x_5^9 + x_1^3 x_2^3 x_3^{10} x_4^4 x_5^7 + x_1^3 x_2^3 x_3^{12} x_4^2 x_5^7 + x_1^3 x_2^8 x_3^5 x_4^4 x_5^7 \\
&+ x_1^4 x_2 x_3^7 x_4^8 x_5^7 + x_1^{10} x_2 x_3^5 x_4^4 x_5^7, \\
u_8 &= x_1^3 x_2^4 x_3^5 x_4^4 x_5^7 + x_1^6 x_2 x_3^5 x_4^4 x_5^7.
\end{aligned}$$

By a similar computation, we have

$$\begin{aligned}
Y_2 &= x_1^2 x_2^7 x_3^8 x_4^5 x_5^9 + x_1^2 x_2^{11} x_3 x_4^5 x_5^{12} + x_1^2 x_2^{11} x_3^8 x_4^5 x_5^5 + x_1^2 x_2^{13} x_3 x_4^5 x_5^{10} + x_1^2 x_2^{13} x_3 x_4^9 x_5^6 \\
&\quad + x_1^2 x_2^{13} x_3^8 x_4^5 x_5^5 + x_1^3 x_2^3 x_3 x_4^{12} x_5^{12} + x_1^3 x_2^3 x_3^4 x_4^9 x_5^{12} + x_1^3 x_2^3 x_3^8 x_4^5 x_5^{12} + x_1^3 x_2^5 x_3^4 x_4^9 x_5^{10} \\
&\quad + x_1^3 x_2^5 x_3^4 x_4^{10} x_5^9 + x_1^3 x_2^5 x_3^8 x_4^5 x_5^{10} + x_1^3 x_2^5 x_3^8 x_4^6 x_5^9 + x_1^3 x_2^5 x_3^8 x_4^9 x_5^6 + x_1^3 x_2^5 x_3^8 x_4^{10} x_5^5 \\
&\quad + x_1^3 x_2^7 x_3 x_4^8 x_5^{12} + x_1^3 x_2^7 x_3 x_4^{12} x_5^8 + x_1^3 x_2^7 x_3^4 x_4^9 x_5^8 + x_1^3 x_2^7 x_3^8 x_4^5 x_5^9 \\
&\quad + Sq^1(v_1) + Sq^2(v_2) + Sq^4(v_4) + Sq^8(v_8) \text{ modulo } (\mathcal{P}_5^-(\omega^*)), \text{ where} \\
v_1 &= x_1^5 x_2^3 x_3 x_4^9 x_5^{12} + x_1^5 x_2^3 x_3^8 x_4^5 x_5^9 + x_1^5 x_2^7 x_3 x_4^5 x_5^{12} \\
&\quad + x_1^5 x_2^7 x_3 x_4^9 x_5^8 + x_1^5 x_2^7 x_3^4 x_4^5 x_5^9 + x_1^5 x_2^7 x_3^8 x_4^5 x_5^5, \\
v_2 &= x_1^2 x_2^7 x_3^8 x_4^5 x_5^9 + x_1^2 x_2^{11} x_3 x_4^5 x_5^{10} + x_1^2 x_2^{11} x_3 x_4^9 x_5^6 + x_1^2 x_2^{11} x_3^8 x_4^5 x_5^5 + x_1^2 x_2^{13} x_3 x_4^{10} x_5^{12} \\
&\quad + x_1^3 x_2^3 x_3^2 x_4^9 x_5^{12} + x_1^3 x_2^3 x_3^8 x_4^5 x_5^{10} + x_1^3 x_2^3 x_3^8 x_4^6 x_5^9 + x_1^3 x_2^7 x_3 x_4^6 x_5^{12} + x_1^3 x_2^7 x_3 x_4^{10} x_5^8 \\
&\quad + x_1^3 x_2^7 x_3^2 x_4^5 x_5^{12} + x_1^3 x_2^7 x_3^2 x_4^9 x_5^8 + x_1^3 x_2^7 x_3^4 x_4^5 x_5^{10} + x_1^3 x_2^7 x_3^4 x_4^6 x_5^9 + x_1^3 x_2^7 x_3^8 x_4^5 x_5^6 \\
&\quad + x_1^3 x_2^7 x_3^8 x_4^6 x_5^5 + x_1^6 x_2^3 x_3 x_4^9 x_5^{10} + x_1^6 x_2^3 x_3^8 x_4^5 x_5^9 + x_1^6 x_2^7 x_3 x_4^5 x_5^{10} + x_1^6 x_2^7 x_3 x_4^9 x_5^6 \\
&\quad + x_1^6 x_2^7 x_3^4 x_4^5 x_5^9 + x_1^6 x_2^7 x_3^8 x_4^5 x_5^5, \\
v_4 &= x_1^3 x_2^5 x_3 x_4^6 x_5^{12} + x_1^3 x_2^5 x_3^2 x_4^5 x_5^{12} + x_1^3 x_2^5 x_3^4 x_4^5 x_5^{10} + x_1^3 x_2^5 x_3^4 x_4^6 x_5^9 + x_1^3 x_2^5 x_3^8 x_4^5 x_5^6 \\
&\quad + x_1^3 x_2^5 x_3^8 x_4^6 x_5^5 + x_1^3 x_2^{11} x_3^4 x_4^5 x_5^5 + x_1^4 x_2^7 x_3 x_4^5 x_5^{10} + x_1^4 x_2^7 x_3 x_4^9 x_5^6 + x_1^4 x_2^7 x_3^8 x_4^5 x_5^5 \\
&\quad + x_1^{10} x_2^5 x_3^4 x_4^5 x_5^6 + x_1^{10} x_2^5 x_3^4 x_4^5 x_5^5, \\
v_8 &= x_1^3 x_2^7 x_3^4 x_4^5 x_5^5 + x_1^6 x_2^5 x_3^4 x_4^5 x_5^6 + x_1^6 x_2^5 x_3^4 x_4^5 x_5^5.
\end{aligned}$$

From the above equalities, we see that Y_1 and Y_2 are strictly inadmissible. The lemma is proved. $\square$

Proposition 3.4.20. *We have $Q\mathcal{P}_5^+(\overline{\omega}_{(1)}) = \langle [\overline{\Phi}^+(\mathcal{B}_4(\overline{\omega}_{(1)})) \cup \mathcal{B}^+(5, \overline{\omega}_{(1)}) \cup \mathcal{F}]_{\overline{\omega}_{(1)}} \rangle$, where*

$$\overline{\Phi}^+(\mathcal{B}_4(\overline{\omega}_{(1)})) \cup \mathcal{B}^+(5, \overline{\omega}_{(1)}) \cup \mathcal{F} = \{\mathcal{Y}_{47,j} : 1 \leq j \leq 370\}$$

with the monomials $\mathcal{Y}_{47,j} : 1 \leq j \leq 370$, are listed in Sect. 6.7.

The proof of the proposition is based on Lemmas 3.4.18, 3.4.19 and some results below. The following lemma is an immediate consequence of a result in Sum [47].

Lemma 3.4.21. *The following monomials are strictly inadmissible:*

- I) $x_i^{15} x_j^2 x_k x_\ell^{13}$, $x_i^3 x_j^{12} x_k x_\ell^{15}$, $x_i^3 x_j^{14} x_k x_\ell^{13}$,
 $x_i^7 x_j^{10} x_k x_\ell^{13}$, $x_i^3 x_j^{13} x_k^8 x_\ell^7$, $x_i^3 x_j^{12} x_k^9 x_\ell^7$, $1 \leq j < k \leq 5$, $1 \leq i, \ell \leq 5$, $i \neq \ell$, $i, \ell \neq j, k$;
- II) $x_i^{15} x_j^{14} x_k x_\ell$, $x_i^3 x_j^2 x_k^{13} x_\ell^{13}$, $x_i^{15} x_j^2 x_k^5 x_\ell^9$, $x_i^{13} x_j^2 x_k^7 x_\ell^9$, $x_i^{13} x_j^7 x_k^2 x_\ell^9$, $x_i^{15} x_j^3 x_k^4 x_\ell^9$, $x_i^{15} x_j^3 x_k^5 x_\ell^8$,
 $x_i^{14} x_j^3 x_k^5 x_\ell^9$, $x_i^{13} x_j^3 x_k^6 x_\ell^9$, $x_i^{13} x_j^6 x_k^3 x_\ell^9$, $1 \leq j < k < \ell \leq 5$, $1 \leq i \leq 5$, $i \neq j, k, \ell$;
- III) $\rho_{(1,5)}(x_1^2 x_2^5 x_3^{11} x_4^{13})$, $\rho_{(1,5)}(x_1^6 x_2^{11} x_3^5 x_4^9)$, $\rho_{(1,5)}(x_1^6 x_2 x_3^{11} x_4^{13})$,
 $\rho_{(1,5)}(x_1^6 x_2^{11} x_3 x_4^{13})$, $\rho_{(1,5)}(x_1^6 x_2^{11} x_3^5 x_4^9)$.

Lemma 3.4.22. *The following monomials are strictly inadmissible:*

$$\begin{aligned}
Z_1 &= x_1^3 x_2^3 x_3^{12} x_4^5 x_5^{24}, & Z_2 &= x_1^7 x_2 x_3^{11} x_4^{20} x_5^8, & Z_3 &= x_1^7 x_2 x_3^2 x_4^{28} x_5^9, & Z_4 &= x_1 x_2^2 x_3^7 x_4^{12} x_5^{25}, \\
Z_5 &= x_1 x_2^7 x_3^{12} x_4^{25}, & Z_6 &= x_1^7 x_2 x_3^2 x_4^{12} x_5^{25}, & Z_7 &= x_1 x_2^6 x_3^3 x_4^8 x_5^{29}, & Z_8 &= x_1 x_2^6 x_3^3 x_4^{24} x_5^{13}, \\
Z_9 &= x_1^3 x_2^7 x_3^8 x_4^5 x_5^{28}, & Z_{10} &= x_1^7 x_2^3 x_3^8 x_4^5 x_5^{28}, & Z_{11} &= x_1^3 x_2^7 x_3^{24} x_4^5 x_5^{12}, & Z_{12} &= x_1^7 x_3^3 x_3^{24} x_4^5 x_5^{12}, \\
Z_{13} &= x_1 x_2^7 x_3^{26} x_4^5 x_5^9, & Z_{14} &= x_1^7 x_2 x_3^{26} x_4^5 x_5^9, & Z_{15} &= x_1 x_2^7 x_3^{10} x_4^5 x_5^{25}, & Z_{16} &= x_1^7 x_2 x_3^{10} x_4^5 x_5^{25}, \\
Z_{17} &= x_1 x_2^7 x_3^{26} x_4^5 x_5^8, & Z_{18} &= x_1^7 x_2 x_3^{26} x_4^5 x_5^8, & Z_{19} &= x_1 x_2^7 x_3^{10} x_4^5 x_5^{24}, & Z_{20} &= x_1^7 x_2 x_3^{10} x_4^5 x_5^{24}, \\
Z_{21} &= x_1 x_2^6 x_3^{11} x_4^{13} x_5^{16}, & Z_{22} &= x_1 x_2^7 x_3^{10} x_4^{21} x_5^8, & Z_{23} &= x_1^7 x_2 x_3^{10} x_4^{21} x_5^8, & Z_{24} &= x_1 x_2^7 x_3^{11} x_4^{20} x_5^8, \\
Z_{25} &= x_1 x_2^2 x_3^7 x_4^{28} x_5^9, & Z_{26} &= x_1^7 x_2^{11} x_3 x_4^{20} x_5^8, & Z_{27} &= x_1 x_2^7 x_3^{10} x_4^{20} x_5^9, & Z_{28} &= x_1^7 x_2 x_3^{10} x_4^{20} x_5^9, \\
Z_{29} &= x_1 x_2^7 x_3^{10} x_4^{13} x_5^{16}, & Z_{30} &= x_1^7 x_2 x_3^{10} x_4^{13} x_5^{16}, & Z_{31} &= x_1 x_2^7 x_3^{11} x_4^{12} x_5^{16}, & Z_{32} &= x_1^7 x_2 x_3^{11} x_4^{12} x_5^{16}, \\
Z_{33} &= x_1^7 x_2^{11} x_3 x_4^{12} x_5^{16}, & Z_{34} &= x_1^3 x_2^3 x_3^4 x_4^{28} x_5^9, & Z_{35} &= x_1^3 x_2^3 x_3^{28} x_4^5 x_5^9, & Z_{36} &= x_1^3 x_2^3 x_3^4 x_4^{12} x_5^{25}, \\
Z_{37} &= x_1^3 x_2^3 x_3^{12} x_4^5 x_5^{25}, & Z_{38} &= x_1^3 x_2^3 x_3^{28} x_4^5 x_5^8, & Z_{39} &= x_1 x_2^7 x_3^2 x_4^{28} x_5^9, & Z_{40} &= x_1^3 x_2^3 x_3^{12} x_4^{21} x_5^8.
\end{aligned}$$

Proof. We have $\omega(Z_u) = \overline{\omega}_{(1)}$, $\forall u$, $1 \leq u \leq 40$. Consider the monomials $Z_1 = x_1^3 x_2^3 x_3^{12} x_4^5 x_5^{24}$ and $Z_2 = x_1^7 x_2 x_3^{11} x_4^{20} x_5^8$. We prove that these monomials are strictly inadmissible. The others can be

proved by a similar technique. Computing the monomials Z_1, Z_2 is long and technical. Indeed, by using Cartan's formula, we get

$$Z_1 = \sum X + Sq^1(\sum \sigma_1) + Sq^2(\sum \sigma_2) + Sq^4(\sum \sigma_4) + Sq^8(\sum \sigma_8) \text{ modulo } (\mathcal{P}_5^-(\bar{\omega}_{(1)})), \text{ where}$$

$$\begin{aligned} \sum X &= x_1^2 x_2 x_3^3 x_4^{12} x_5^{29} + x_1^2 x_2 x_3^3 x_4^{13} x_5^{28} + x_1^2 x_2 x_3^5 x_4^{12} x_5^{27} + x_1^2 x_2 x_3^5 x_4^{13} x_5^{26} + x_1^2 x_2 x_3^5 x_4^{24} x_5^{15} \\ &+ x_1^2 x_2 x_3^5 x_4^{25} x_5^{14} + x_1^2 x_2 x_3^7 x_4^9 x_5^{28} + x_1^2 x_2 x_3^7 x_4^{13} x_5^{24} + x_1^2 x_2 x_3^{12} x_4^3 x_5^{29} + x_1^2 x_2 x_3^{12} x_4^5 x_5^{27} \\ &+ x_1^2 x_2 x_3^{13} x_4^7 x_5^{24} + x_1^2 x_2 x_3^{13} x_4^9 x_5^{22} + x_1^2 x_2 x_3^{13} x_4^{16} x_5^{15} + x_1^2 x_2 x_3^{13} x_4^{17} x_5^{14} + x_1^2 x_2 x_3^{16} x_4^{13} x_5^{15} \\ &+ x_1^2 x_2 x_3^{24} x_4^5 x_5^{15} + x_1^2 x_2^3 x_3^5 x_4^9 x_5^{28} + x_1^2 x_2^3 x_3^5 x_4^{12} x_5^{25} + x_1^2 x_2^3 x_3^9 x_4^5 x_5^{28} + x_1^2 x_2^3 x_3^{12} x_4^5 x_5^{25} \\ &+ x_1^2 x_2^5 x_3^3 x_4^9 x_5^{28} + x_1^2 x_2^5 x_3^3 x_4^{12} x_5^{25} + x_1^2 x_2^5 x_3^9 x_4^5 x_5^{26} + x_1^2 x_2^5 x_3^{12} x_4^5 x_5^{25} + x_1^3 x_2 x_3^3 x_4^{12} x_5^{28} \\ &+ x_1^3 x_2 x_3^4 x_4^{12} x_5^{27} + x_1^3 x_2 x_3^4 x_4^{13} x_5^{26} + x_1^3 x_2 x_3^4 x_4^{24} x_5^{15} + x_1^3 x_2 x_3^4 x_4^{25} x_5^{14} + x_1^3 x_2 x_3^7 x_4^{12} x_5^{24} \\ &+ x_1^3 x_2 x_3^7 x_4^{12} x_5^{24} + x_1^3 x_2 x_3^9 x_4^{12} x_5^{22} + x_1^3 x_2 x_3^{12} x_4^3 x_5^{28} + x_1^3 x_2 x_3^{12} x_4^5 x_5^{27} + x_1^3 x_2 x_3^{12} x_4^7 x_5^{24} \\ &+ x_1^3 x_2 x_3^{12} x_4^9 x_5^{22} + x_1^3 x_2 x_3^{12} x_4^{16} x_5^{15} + x_1^3 x_2 x_3^{12} x_4^{17} x_5^{14} + x_1^3 x_2 x_3^{16} x_4^{12} x_5^{15} + x_1^3 x_2 x_3^{24} x_4^4 x_5^{15} \\ &+ x_1^3 x_2^2 x_3^3 x_4^9 x_5^{28} + x_1^3 x_2^2 x_3^5 x_4^{12} x_5^{25} + x_1^3 x_2^2 x_3^9 x_4^{12} x_5^{21} + x_1^3 x_2^2 x_3^{12} x_4^5 x_5^{25} + x_1^3 x_2^2 x_3^{12} x_4^9 x_5^{21} \\ &+ x_1^3 x_2^3 x_3^4 x_4^9 x_5^{28} + x_1^3 x_2^3 x_3^4 x_4^{12} x_5^{25} + x_1^3 x_2^3 x_3^5 x_4^8 x_5^{28} + x_1^3 x_2^3 x_3^5 x_4^{12} x_5^{24} \end{aligned}$$

$$\begin{aligned} \sum \sigma_1 &= x_1^3 x_2 x_3^3 x_4^{12} x_5^{27} + x_1^3 x_2 x_3^3 x_4^{13} x_5^{26} + x_1^3 x_2 x_3^3 x_4^{16} x_5^{23} + x_1^3 x_2 x_3^3 x_4^{17} x_5^{22} + x_1^3 x_2 x_3^3 x_4^{20} x_5^{19} \\ &+ x_1^3 x_2 x_3^3 x_4^{21} x_5^{18} + x_1^3 x_2 x_3^3 x_4^{24} x_5^{15} + x_1^3 x_2 x_3^3 x_4^{25} x_5^{14} + x_1^3 x_2 x_3^7 x_4^9 x_5^{26} + x_1^3 x_2 x_3^7 x_4^{11} x_5^{24} \\ &+ x_1^3 x_2 x_3^7 x_4^{16} x_5^{19} + x_1^3 x_2 x_3^7 x_4^{17} x_5^{18} + x_1^3 x_2 x_3^9 x_4^{13} x_5^{20} + x_1^3 x_2 x_3^{11} x_4^7 x_5^{24} + x_1^3 x_2 x_3^{11} x_4^9 x_5^{22} \\ &+ x_1^3 x_2 x_3^{11} x_4^{16} x_5^{15} + x_1^3 x_2 x_3^{11} x_4^{17} x_5^{14} + x_1^3 x_2 x_3^{12} x_4^3 x_5^{27} + x_1^3 x_2 x_3^{16} x_4^3 x_5^{23} + x_1^3 x_2 x_3^{16} x_4^7 x_5^{19} \\ &+ x_1^3 x_2 x_3^{16} x_4^{11} x_5^{15} + x_1^3 x_2 x_3^{20} x_4^3 x_5^{15} + x_1^3 x_2^2 x_3^3 x_4^5 x_5^{32} + x_1^3 x_2^2 x_3^3 x_4^9 x_5^{28} \\ &+ x_1^3 x_2^3 x_3^3 x_4^{12} x_5^{25} + x_1^3 x_2^3 x_3^3 x_4^{16} x_5^{21} + x_1^3 x_2^3 x_3^9 x_4^5 x_5^{26} + x_1^3 x_2^3 x_3^9 x_4^9 x_5^{22} + x_1^3 x_2^3 x_3^{12} x_4^3 x_5^{25} \\ &+ x_1^3 x_2^3 x_3^{16} x_4^3 x_5^{21} + x_1^3 x_2^4 x_3^9 x_4^9 x_5^{21}, \end{aligned}$$

$$\begin{aligned} \sum \sigma_2 &= x_1^2 x_2 x_3^3 x_4^{12} x_5^{27} + x_1^2 x_2 x_3^3 x_4^{13} x_5^{26} + x_1^2 x_2 x_3^3 x_4^{24} x_5^{15} + x_1^2 x_2 x_3^3 x_4^{25} x_5^{14} + x_1^2 x_2 x_3^7 x_4^9 x_5^{26} \\ &+ x_1^2 x_2 x_3^7 x_4^{11} x_5^{24} + x_1^2 x_2 x_3^{11} x_4^7 x_5^{24} + x_1^2 x_2 x_3^{11} x_4^9 x_5^{22} + x_1^2 x_2 x_3^{11} x_4^{16} x_5^{15} + x_1^2 x_2 x_3^{11} x_4^{17} x_5^{14} \\ &+ x_1^2 x_2 x_3^{12} x_4^3 x_5^{27} + x_1^2 x_2 x_3^{16} x_4^{11} x_5^{15} + x_1^2 x_2 x_3^{24} x_4^3 x_5^{15} + x_1^2 x_2^3 x_3^3 x_4^9 x_5^{28} + x_1^2 x_2^3 x_3^3 x_4^{12} x_5^{25} \\ &+ x_1^2 x_2^3 x_3^9 x_4^5 x_5^{26} + x_1^2 x_2^3 x_3^9 x_4^9 x_5^{22} + x_1^2 x_2^3 x_3^{12} x_4^3 x_5^{25} + x_1^2 x_2^3 x_3^{14} x_4^5 x_5^{22} + x_1^2 x_2^3 x_3^{22} x_4^5 x_5^{14} \\ &+ x_1^5 x_2 x_3^7 x_4^{10} x_5^{22} + x_1^5 x_2 x_3^7 x_4^{18} x_5^{14} + x_1^5 x_2^2 x_3^3 x_4^9 x_5^{26} + x_1^5 x_2^2 x_3^3 x_4^{11} x_5^{24} + x_1^5 x_2^2 x_3^3 x_4^{12} x_5^{23} \\ &+ x_1^5 x_2^2 x_3^3 x_4^{13} x_5^{22} + x_1^5 x_2^2 x_3^3 x_4^{20} x_5^{15} + x_1^5 x_2^2 x_3^3 x_4^{21} x_5^{14} + x_1^5 x_2^2 x_3^7 x_4^7 x_5^{24} + x_1^5 x_2^2 x_3^7 x_4^9 x_5^{22} \\ &+ x_1^5 x_2^2 x_3^7 x_4^{16} x_5^{15} + x_1^5 x_2^2 x_3^7 x_4^{17} x_5^{14} + x_1^5 x_2^2 x_3^9 x_4^{10} x_5^{19} + x_1^5 x_2^2 x_3^9 x_4^{11} x_5^{18} + x_1^5 x_2^2 x_3^{10} x_4^9 x_5^{19} \\ &+ x_1^5 x_2^2 x_3^{12} x_4^3 x_5^{23} + x_1^5 x_2^2 x_3^{16} x_4^7 x_5^{15} + x_1^5 x_2^2 x_3^{20} x_4^3 x_5^{15} + x_1^5 x_2^3 x_3^3 x_4^6 x_5^{28} + x_1^5 x_2^3 x_3^3 x_4^8 x_5^{26} \\ &+ x_1^5 x_2^3 x_3^3 x_4^{10} x_5^{24} + x_1^5 x_2^3 x_3^3 x_4^{12} x_5^{22} + x_1^5 x_2^3 x_3^5 x_4^6 x_5^{26} + x_1^5 x_2^3 x_3^5 x_4^{10} x_5^{22} + x_1^5 x_2^3 x_3^{10} x_4^3 x_5^{24} \\ &+ x_1^5 x_2^3 x_3^{10} x_4^5 x_5^{22} + x_1^5 x_2^3 x_3^{12} x_4^3 x_5^{22}, \end{aligned}$$

$$\begin{aligned} \sum \sigma_4 &= x_1^3 x_2 x_3^3 x_4^{14} x_5^{22} + x_1^3 x_2 x_3^3 x_4^{22} x_5^{14} + x_1^3 x_2 x_3^5 x_4^{14} x_5^{20} + x_1^3 x_2 x_3^5 x_4^{20} x_5^{14} + x_1^3 x_2 x_3^6 x_4^{13} x_5^{20} \\ &+ x_1^3 x_2 x_3^7 x_4^{10} x_5^{22} + x_1^3 x_2 x_3^7 x_4^{18} x_5^{14} + x_1^3 x_2^2 x_3^3 x_4^9 x_5^{26} + x_1^3 x_2^2 x_3^3 x_4^{11} x_5^{24} + x_1^3 x_2^2 x_3^3 x_4^{12} x_5^{23} \\ &+ x_1^3 x_2^2 x_3^3 x_4^{13} x_5^{22} + x_1^3 x_2^2 x_3^3 x_4^{20} x_5^{15} + x_1^3 x_2^2 x_3^3 x_4^{21} x_5^{14} + x_1^3 x_2^2 x_3^7 x_4^7 x_5^{24} + x_1^3 x_2^2 x_3^7 x_4^9 x_5^{22} \\ &+ x_1^3 x_2^2 x_3^7 x_4^{16} x_5^{15} + x_1^3 x_2^2 x_3^7 x_4^{17} x_5^{14} + x_1^3 x_2^2 x_3^9 x_4^{10} x_5^{19} + x_1^3 x_2^2 x_3^9 x_4^{11} x_5^{18} + x_1^3 x_2^2 x_3^{10} x_4^9 x_5^{19} \\ &+ x_1^3 x_2^2 x_3^{12} x_4^3 x_5^{23} + x_1^3 x_2^2 x_3^{16} x_4^7 x_5^{15} + x_1^3 x_2^2 x_3^{20} x_4^3 x_5^{15} + x_1^3 x_2^3 x_3^3 x_4^6 x_5^{28} + x_1^3 x_2^3 x_3^3 x_4^8 x_5^{26} \\ &+ x_1^3 x_2^3 x_3^3 x_4^{10} x_5^{24} + x_1^3 x_2^3 x_3^3 x_4^{12} x_5^{22} + x_1^3 x_2^3 x_3^5 x_4^6 x_5^{26} + x_1^3 x_2^3 x_3^5 x_4^{10} x_5^{22} + x_1^3 x_2^3 x_3^{10} x_4^3 x_5^{24} \\ &+ x_1^3 x_2^3 x_3^{10} x_4^5 x_5^{22} + x_1^3 x_2^3 x_3^{12} x_4^3 x_5^{22} + x_1^3 x_2^8 x_3^5 x_4^5 x_5^{22} + x_1^3 x_2^8 x_3^5 x_4^7 x_5^{20} + x_1^3 x_2^8 x_3^5 x_4^{12} x_5^{15} \\ &+ x_1^3 x_2^8 x_3^5 x_4^{13} x_5^{14} + x_1^3 x_2^8 x_3^{12} x_4^5 x_5^{15}, \end{aligned}$$

$$\sum \sigma_8 = x_1^3 x_2^4 x_3^5 x_4^5 x_5^{22} + x_1^3 x_2^4 x_3^5 x_4^7 x_5^{20} + x_1^3 x_2^4 x_3^5 x_4^{12} x_5^{15} + x_1^3 x_2^4 x_3^5 x_4^{13} x_5^{14} + x_1^3 x_2^4 x_3^{12} x_4^5 x_5^{15},$$

The above relations imply that Z_1 is strictly inadmissible.

Next we prove that Z_2 is also strictly inadmissible. A direct computation shows that

$$Z_2 = \sum Y + Sq^1(\sum \beta_1) + Sq^2(\sum \beta_2) + Sq^4(\sum \beta_4) + Sq^8(\sum \beta_8) \text{ modulo } (\mathcal{P}_5^-(\bar{\omega}_{(1)})),$$

$$\begin{aligned}
\sum Y &= x^3_1 x_2 x_3 x_4^{12} x_5^{24} + x^3_1 x_2 x_3 x_4^{28} x_5^8 + x^3_1 x_2 x_3 x_4^{20} x_5^{14} + x^3_1 x_2 x_3 x_4^{28} x_5^6 + x^3_1 x_2 x_3^{11} x_4^{20} x_5^{12} \\
&+ x^3_1 x_2 x_3^{12} x_4^{17} x_5^{14} + x^3_1 x_2 x_3^{12} x_4^{19} x_5^{12} + x^3_1 x_2 x_3^{12} x_4^{21} x_5^{10} + x^3_1 x_2 x_3^{14} x_4^{17} x_5^{12} + x^3_1 x_2 x_3^{17} x_4^{12} x_5^{14} \\
&+ x^3_1 x_2 x_3^{20} x_4^9 x_5^{14} + x^3_1 x_2 x_3^{13} x_4^{13} x_5^{16} + x^3_1 x_2 x_3^{13} x_4^{21} x_5^8 + x^3_1 x_2 x_3^{17} x_4^{25} x_5^8 + x^3_1 x_2 x_3^{19} x_4^{11} x_5^{20} \\
&+ x^3_1 x_2 x_3^4 x_4^{21} x_5^{10} + x^3_1 x_2 x_3^4 x_4^{25} x_5^6 + x^3_1 x_2 x_3^{11} x_4^{12} x_5^{17} + x^3_1 x_2 x_3^{11} x_4^{20} x_5^9 + x^3_1 x_2 x_3^{12} x_4^{11} x_5^{17} \\
&+ x^3_1 x_2 x_3^{12} x_4^{19} x_5^9 + x^3_1 x_2 x_3^{12} x_4^{24} x_5^{14} + x^3_1 x_2 x_3^{11} x_4^{17} x_5^{14} + x^3_1 x_2 x_3^{11} x_4^{19} x_5^{12} + x^3_1 x_2 x_3^{11} x_4^{21} x_5^{10} \\
&+ x^3_1 x_2 x_3^{11} x_4^{19} x_5^{14} + x^3_1 x_2 x_3^{11} x_4^{25} x_5^9 + x^3_1 x_2 x_3^{11} x_4^{13} x_5^{17} + x^3_1 x_2 x_3^{11} x_4^{21} x_5^9 + x^3_1 x_2 x_3^{11} x_4^{11} x_5^{17} \\
&+ x^3_1 x_2 x_3^{11} x_4^{19} x_5^9 + x^3_1 x_2 x_3^{11} x_4^{24} x_5^{10} + x^3_1 x_2 x_3^{11} x_4^{12} x_5^{18} + x^3_1 x_2 x_3^{11} x_4^{14} x_5^{16} + x^3_1 x_2 x_3^{11} x_4^{20} x_5^{10} \\
&+ x^3_1 x_2 x_3^{24} x_4^7 x_5^{10} + x^3_1 x_2 x_3^{24} x_4^{11} x_5^6 + x^3_1 x_2 x_3^{24} x_4^9 x_5^{10} + x^3_1 x_2 x_3^{11} x_4^{12} x_5^{17} + x^3_1 x_2 x_3^{11} x_4^{20} x_5^9 \\
&+ x^3_1 x_2 x_3^8 x_4^{17} x_5^{10} + x^3_1 x_2 x_3^{11} x_4^{17} x_5^6 + x^3_1 x_2 x_3^8 x_4^{13} x_5^{18} + x^3_1 x_2 x_3^8 x_4^{14} x_5^{17} + x^3_1 x_2 x_3^8 x_4^{21} x_5^{10} \\
&+ x^3_1 x_2 x_3^8 x_4^{22} x_5^9 + x^3_1 x_2 x_3^9 x_4^{12} x_5^{18} + x^3_1 x_2 x_3^9 x_4^{20} x_5^{10} + x^3_1 x_2 x_3^{10} x_4^9 x_5^{20} + x^3_1 x_2 x_3^{10} x_4^{12} x_5^{17} \\
&+ x^3_1 x_2 x_3^{10} x_4^{13} x_5^{16} + x^3_1 x_2 x_3^{10} x_4^{17} x_5^{12} + x^3_1 x_2 x_3^{10} x_4^{20} x_5^9 + x^3_1 x_2 x_3^{10} x_4^{21} x_5^8 + x^3_1 x_2 x_3^{11} x_4^{16} x_5^{12}, \\
\sum \beta_1 &= x^3_1 x_2 x_3 x_4^{11} x_5^{24} + x^3_1 x_2 x_3^{11} x_4^{17} x_5^{14} + x^3_1 x_2 x_3^{11} x_4^{19} x_5^{12} + x^3_1 x_2 x_3^{11} x_4^{21} x_5^{10} + x^3_1 x_2 x_3^{17} x_4^{11} x_5^{14} \\
&+ x^3_1 x_2 x_3^{17} x_4^{19} x_5^6 + x^3_1 x_2 x_3^{11} x_4^{11} x_5^{17} + x^3_1 x_2 x_3^{11} x_4^{19} x_5^9 + x^3_1 x_2 x_3^9 x_4^{17} x_5^{12} + x^3_1 x_2 x_3^{11} x_4^{17} x_5^{10} \\
&+ x^3_1 x_2 x_3^{13} x_4^{17} x_5^8 + x^3_1 x_2 x_3^4 x_4^{13} x_5^{17} + x^3_1 x_2 x_3^4 x_4^{21} x_5^9 + x^3_1 x_2 x_3^9 x_4^{11} x_5^{17} + x^3_1 x_2 x_3^9 x_4^{19} x_5^9 \\
&+ x^3_1 x_2 x_3^8 x_4^{13} x_5^{17} + x^3_1 x_2 x_3^8 x_4^{21} x_5^9 + x^3_1 x_2 x_3^{11} x_4^{17} x_5^{17} + x^3_1 x_2 x_3^{11} x_4^{19} x_5^9, \\
\sum \beta_2 &= x^3_1 x_2 x_3^{14} x_4^{17} x_5^{10} + x^3_1 x_2 x_3^7 x_4^{13} x_5^{20} + x^3_1 x_2 x_3^7 x_4^{21} x_5^{12} + x^3_1 x_2 x_3^{11} x_4^{11} x_5^{18} + x^3_1 x_2 x_3^{11} x_4^{19} x_5^{10} \\
&+ x^3_1 x_2 x_3^7 x_4^{18} x_5^{14} + x^3_1 x_2 x_3^7 x_4^{26} x_5^6 + x^3_1 x_2 x_3^{18} x_4^7 x_5^{14} + x^3_1 x_2 x_3^{11} x_4^{20} x_5^{20} + x^3_1 x_2 x_3^{21} x_4^{10} x_5^{10} \\
&+ x^3_1 x_2 x_3^7 x_4^{25} x_5^6 + x^3_1 x_2 x_3^{11} x_4^{11} x_5^{16} + x^3_1 x_2 x_3^{11} x_4^{13} x_5^{14} + x^3_1 x_2 x_3^{11} x_4^{19} x_5^8 + x^3_1 x_2 x_3^{11} x_4^{21} x_5^6 \\
&+ x^3_1 x_2 x_3^{13} x_4^{11} x_5^{14} + x^3_1 x_2 x_3^{13} x_4^{19} x_5^6 + x^3_1 x_2 x_3^4 x_4^{11} x_5^{18} + x^3_1 x_2 x_3^7 x_4^{19} x_5^{10} + x^3_1 x_2 x_3^7 x_4^{13} x_5^{17} \\
&+ x^3_1 x_2 x_3^7 x_4^{20} x_5^{10} + x^3_1 x_2 x_3^7 x_4^{21} x_5^9 + x^3_1 x_2 x_3^9 x_4^{11} x_5^{17} + x^3_1 x_2 x_3^9 x_4^{19} x_5^9 + x^3_1 x_2 x_3^{10} x_4^9 x_5^{18} \\
&+ x^3_1 x_2 x_3^{10} x_4^{11} x_5^{16} + x^3_1 x_2 x_3^{10} x_4^{19} x_5^8 + x^3_1 x_2 x_3^{11} x_4^{18} x_5^8 + x^3_1 x_2 x_3^{13} x_4^{14} x_5^{10} + x^3_1 x_2 x_3^{14} x_4^{17} x_5^6 \\
&+ x^3_1 x_2 x_3^{18} x_4^7 x_5^{12} + x^3_1 x_2 x_3^{18} x_4^{13} x_5^6 + x^3_1 x_2 x_3^7 x_4^{17} x_5^{12} + x^3_1 x_2 x_3^9 x_4^{17} x_5^{10} + x^3_1 x_2 x_3^{11} x_4^{17} x_5^8 \\
&+ x^3_1 x_2 x_3^{13} x_4^{17} x_5^6 + x^3_1 x_2 x_3^{11} x_4^{17} x_5^{17} + x^3_1 x_2 x_3^7 x_4^{19} x_5^9, \\
\sum \beta_4 &= x^3_1 x_2 x_3 x_4^{18} x_5^{14} + x^3_1 x_2 x_3^7 x_4^{22} x_5^{10} + x^3_1 x_2 x_3^{10} x_4^{11} x_5^{18} + x^3_1 x_2 x_3^{10} x_4^{19} x_5^{10} + x^3_1 x_2 x_3^{11} x_4^{10} x_5^{18} \\
&+ x^3_1 x_2 x_3^{11} x_4^{22} x_5^6 + x^3_1 x_2 x_3^{18} x_4^7 x_5^{14} + x^3_1 x_2 x_3^{11} x_4^{20} x_5^{20} + x^3_1 x_2 x_3^{21} x_4^{10} x_5^{10} + x^3_1 x_2 x_3^{25} x_4^6 x_5^6 \\
&+ x^3_1 x_2 x_3^{11} x_4^{11} x_5^{16} + x^3_1 x_2 x_3^{11} x_4^{13} x_5^{14} + x^3_1 x_2 x_3^{11} x_4^{19} x_5^8 + x^3_1 x_2 x_3^{11} x_4^{21} x_5^6 + x^3_1 x_2 x_3^{13} x_4^{11} x_5^{14} \\
&+ x^3_1 x_2 x_3^{13} x_4^{19} x_5^6 + x^3_1 x_2 x_3^7 x_4^{13} x_5^{17} + x^3_1 x_2 x_3^7 x_4^{21} x_5^9 + x^3_1 x_2 x_3^7 x_4^{12} x_5^{18} + x^3_1 x_2 x_3^7 x_4^{18} x_5^{12} \\
&+ x^3_1 x_2 x_3^7 x_4^{20} x_5^{10} + x^3_1 x_2 x_3^{11} x_4^{14} x_5^{12} + x^3_1 x_2 x_3^{14} x_4^{17} x_5^6 + x^3_1 x_2 x_3^{20} x_4^7 x_5^{10} + x^3_1 x_2 x_3^{20} x_4^{11} x_5^6 \\
&+ x^3_1 x_2 x_3^4 x_4^{11} x_5^{18} + x^3_1 x_2 x_3$$

$$\begin{aligned}
d_1 &= x_1^3 x_2^3 x_3^{12} x_4^{13} x_5^{16}, & d_2 &= x_1^3 x_2^3 x_3^{13} x_4^{12} x_5^{16}, & d_3 &= x_1^3 x_2^3 x_3^{13} x_4^{20} x_5^8, & d_4 &= x_1^3 x_2^3 x_3^{12} x_4^{20} x_5^9, \\
d_5 &= x_1^3 x_2^7 x_3^8 x_4^{25}, & d_6 &= x_1^3 x_2^7 x_3^4 x_4^{25} x_5^8, & d_7 &= x_1^7 x_2^3 x_3^4 x_4^8 x_5^{25}, & d_8 &= x_1^7 x_2^3 x_3^4 x_4^{25} x_5^8, \\
d_9 &= x_1^3 x_2^7 x_3^4 x_4^{24} x_5^9, & d_{10} &= x_1^3 x_2^7 x_3^4 x_4^9 x_5^{24}, & d_{11} &= x_1^7 x_2^3 x_3^4 x_4^{24} x_5^9, & d_{12} &= x_1^7 x_2^3 x_3^4 x_4^9 x_5^{24}, \\
d_{13} &= x_1^3 x_2^3 x_3^{11} x_4^{17} x_5^{12}, & d_{14} &= x_1^3 x_2^3 x_3^{11} x_4^{16} x_5^{13}, & d_{15} &= x_1^3 x_2^3 x_3^{11} x_4^{13} x_5^{16}, & d_{16} &= x_1^3 x_2^3 x_3^6 x_4^{25}, \\
d_{17} &= x_1^3 x_2^5 x_3^6 x_4^{25} x_5^8, & d_{18} &= x_1^3 x_2^5 x_3^6 x_4^{24} x_5^9, & d_{19} &= x_1^3 x_2^5 x_3^6 x_4^{24} x_5^{24}, & d_{20} &= x_1^3 x_2^5 x_3^8 x_4^{24}, \\
d_{21} &= x_1^3 x_2^5 x_3^7 x_4^{24} x_5^8, & d_{22} &= x_1^3 x_2^5 x_3^8 x_4^{24} x_5^{24}, & d_{23} &= x_1^3 x_2^5 x_3^{24} x_4^{24} x_5^8, & d_{24} &= x_1^7 x_2^3 x_3^8 x_4^{24}, \\
d_{25} &= x_1^7 x_2^3 x_3^5 x_4^{24} x_5^8, & d_{26} &= x_1^3 x_2^5 x_3^{10} x_4^{17} x_5^{12}, & d_{27} &= x_1^3 x_2^5 x_3^{10} x_4^{16} x_5^{13}, & d_{28} &= x_1^3 x_2^5 x_3^{11} x_4^{16} x_5^{12}, \\
d_{29} &= x_1^3 x_2^7 x_3^8 x_4^{17} x_5^{12}, & d_{30} &= x_1^7 x_2^3 x_3^8 x_4^{17} x_5^{12}, & d_{31} &= x_1^3 x_2^7 x_3^8 x_4^{16} x_5^{13}, & d_{32} &= x_1^3 x_2^7 x_3^8 x_4^{13} x_5^{16}, \\
d_{33} &= x_1^7 x_2^3 x_3^8 x_4^{16} x_5^{13}, & d_{34} &= x_1^7 x_2^3 x_3^8 x_4^{13} x_5^{16}, & d_{35} &= x_1^3 x_2^7 x_3^9 x_4^{16} x_5^{12}, & d_{36} &= x_1^7 x_2^3 x_3^9 x_4^{16} x_5^{12}, \\
d_{37} &= x_1^7 x_2^{11} x_3^{17} x_4^8 x_5.
\end{aligned}$$

27

Lemma 3.4.22. By a direct computation, we have

$$\begin{aligned}
d_1 &= x_1^2 x_2 x_3^3 x_4^{13} x_5^{28} + x_1^2 x_2 x_3^3 x_4^{28} x_5^{13} + x_1^2 x_2 x_3^5 x_4^{13} x_5^{26} + x_1^2 x_2 x_3^5 x_4^{14} x_5^{25} + x_1^2 x_2 x_3^5 x_4^{25} x_5^{14} \\
&\quad + x_1^2 x_2 x_3^5 x_4^{26} x_5^{13} + x_1^2 x_2 x_3^{10} x_4^{13} x_5^{21} + x_1^2 x_2 x_3^{12} x_4^{11} x_5^{25} + x_1^2 x_2 x_3^{12} x_4^{11} x_5^{21} + x_1^2 x_2 x_3^{13} x_4^{14} x_5^{17} \\
&\quad + x_1^2 x_2 x_3^{13} x_4^{17} x_5^{14} + x_1^2 x_2 x_3^{18} x_4^{13} x_5^{13} + x_1^2 x_2 x_3^{20} x_4^{11} x_5^{13} + x_1^2 x_2 x_3^5 x_4^{13} x_5^{24} + x_1^2 x_2 x_3^5 x_4^{24} x_5^{13} \\
&\quad + x_1^2 x_2 x_3^9 x_4^{13} x_5^{20} + x_1^2 x_2 x_3^{12} x_4^{13} x_5^{17} + x_1^2 x_2 x_3^{16} x_4^{13} x_5^{13} + x_1^2 x_2 x_3^5 x_4^{13} x_5^{24} + x_1^2 x_2 x_3^5 x_4^{24} x_5^{13} \\
&\quad + x_1^2 x_2 x_3^9 x_4^{13} x_5^{18} + x_1^2 x_2 x_3^9 x_4^{17} x_5^{14} + x_1^2 x_2 x_3^{12} x_4^{11} x_5^{17} + x_1^2 x_2 x_3^{16} x_4^{11} x_5^{13} + x_1^3 x_2 x_3^4 x_4^{13} x_5^{26} \\
&\quad + x_1^3 x_2 x_3^4 x_4^{14} x_5^{25} + x_1^3 x_2 x_3^4 x_4^{25} x_5^{14} + x_1^3 x_2 x_3^4 x_4^{26} x_5^{13} + x_1^3 x_2 x_3^9 x_4^{14} x_5^{20} + x_1^3 x_2 x_3^9 x_4^{20} x_5^{14} \\
&\quad + x_1^3 x_2 x_3^{10} x_4^{12} x_5^{21} + x_1^3 x_2 x_3^{12} x_4^{17} x_5^{24} + x_1^3 x_2 x_3^{12} x_4^{17} x_5^{22} + x_1^3 x_2 x_3^{12} x_4^{14} x_5^{17} + x_1^3 x_2 x_3^{12} x_4^{17} x_5^{14} \\
&\quad + x_1^3 x_2 x_3^{18} x_4^{12} x_5^{13} + x_1^3 x_2 x_3^{20} x_4^{9} x_5^{14} + x_1^3 x_2 x_3^5 x_4^{13} x_5^{24} + x_1^3 x_2 x_3^5 x_4^{24} x_5^{13} + x_1^3 x_2 x_3^{12} x_4^{17} x_5^{13} \\
&\quad + x_1^3 x_2 x_3^4 x_4^{13} x_5^{24} + x_1^3 x_2 x_3^8 x_4^{12} x_5^{21} + x_1^3 x_2 x_3^{12} x_4^{12} x_5^{17} \\
&\quad + Sq^1(\sum X) + Sq^2(\sum Y) + Sq^4(\sum Z) + Sq^8(\sum T) \text{ modulo } (\mathcal{P}_5^-(\bar{\omega}_{(1)})), \text{ where} \\
\sum X &= x_1^3 x_2 x_3^3 x_4^{13} x_5^{26} + x_1^3 x_2 x_3^3 x_4^{14} x_5^{25} + x_1^3 x_2 x_3^3 x_4^{17} x_5^{22} + x_1^3 x_2 x_3^3 x_4^{18} x_5^{21} + x_1^3 x_2 x_3^3 x_4^{21} x_5^{18} \\
&\quad + x_1^3 x_2 x_3^3 x_4^{22} x_5^{17} + x_1^3 x_2 x_3^3 x_4^{25} x_5^{14} + x_1^3 x_2 x_3^3 x_4^{26} x_5^{13} + x_1^3 x_2 x_3^7 x_4^{17} x_5^{18} + x_1^3 x_2 x_3^7 x_4^{18} x_5^{17} \\
&\quad + x_1^3 x_2 x_3^{10} x_4^{11} x_5^{25} + x_1^3 x_2 x_3^{10} x_4^{11} x_5^{21} + x_1^3 x_2 x_3^{11} x_4^{14} x_5^{17} + x_1^3 x_2 x_3^{11} x_4^{17} x_5^{14} + x_1^3 x_2 x_3^{18} x_4^{17} x_5^{13} \\
&\quad + x_1^3 x_2 x_3^{18} x_4^{11} x_5^{13} + x_1^3 x_2 x_3^{18} x_4^{13} x_5^{24} + x_1^3 x_2 x_3^{17} x_4^{20} x_5^{17} + x_1^3 x_2 x_3^{20} x_4^{17} x_5^{13} + x_1^3 x_2 x_3^9 x_4^{13} x_5^{18} \\
&\quad + x_1^3 x_2 x_3^9 x_4^{17} x_5^{14} + x_1^3 x_2 x_3^{12} x_4^{11} x_5^{17} + x_1^3 x_2 x_3^{16} x_4^{11} x_5^{13} + x_1^3 x_2 x_3^4 x_4^{13} x_5^{17} x_5^{13}, \\
\sum Y &= x_1^2 x_2 x_3^3 x_4^{13} x_5^{26} + x_1^2 x_2 x_3^3 x_4^{14} x_5^{25} + x_1^2 x_2 x_3^3 x_4^{25} x_5^{14} + x_1^2 x_2 x_3^3 x_4^{26} x_5^{13} + x_1^2 x_2 x_3^{10} x_4^{11} x_5^{25} \\
&\quad + x_1^2 x_2 x_3^{10} x_4^{11} x_5^{21} + x_1^2 x_2 x_3^{11} x_4^{14} x_5^{17} + x_1^2 x_2 x_3^{11} x_4^{17} x_5^{14} + x_1^2 x_2 x_3^{18} x_4^{17} x_5^{13} + x_1^2 x_2 x_3^3 x_4^{13} x_5^{24} \\
&\quad + x_1^2 x_2 x_3^9 x_4^{13} x_5^{18} + x_1^2 x_2 x_3^9 x_4^{17} x_5^{14} + x_1^2 x_2 x_3^{10} x_4^{11} x_5^{22} + x_1^2 x_2 x_3^{18} x_4^{17} x_5^{14} + x_1^2 x_2 x_3^{16} x_4^{13} x_5^{22} \\
&\quad + x_1^2 x_2 x_3^3 x_4^{14} x_5^{21} + x_1^2 x_2 x_3^3 x_4^{21} x_5^{14} + x_1^2 x_2 x_3^3 x_4^{22} x_5^{13} + x_1^2 x_2 x_3^7 x_4^{14} x_5^{17} + x_1^2 x_2 x_3^7 x_4^{17} x_5^{14} \\
&\quad + x_1^2 x_2 x_3^{10} x_4^{11} x_5^{21} + x_1^2 x_2 x_3^{18} x_4^{17} x_5^{13} + x_1^2 x_2 x_3^3 x_4^{14} x_5^{20} + x_1^2 x_2 x_3^3 x_4^{20} x_5^{14} + x_1^2 x_2 x_3^5 x_4^{14} x_5^{18} \\
&\quad + x_1^2 x_2 x_3^5 x_4^{18} x_5^{14} + x_1^2 x_2 x_3^{10} x_4^{11} x_5^{16} + x_1^2 x_2 x_3^{10} x_4^{13} x_5^{14} + x_1^2 x_2 x_3^{12} x_4^{11} x_5^{14}, \\
\sum Z &= x_1^3 x_2 x_3^7 x_4^{14} x_5^{18} + x_1^3 x_2 x_3^7 x_4^{18} x_5^{14} + x_1^3 x_2 x_3^{18} x_4^{17} x_5^{14} + x_1^3 x_2 x_3^3 x_4^{13} x_5^{22} + x_1^3 x_2 x_3^3 x_4^{14} x_5^{21} \\
&\quad + x_1^3 x_2 x_3^3 x_4^{21} x_5^{14} + x_1^3 x_2 x_3^3 x_4^{22} x_5^{13} + x_1^3 x_2 x_3^7 x_4^{14} x_5^{17} + x_1^3 x_2 x_3^7 x_4^{17} x_5^{14} + x_1^3 x_2 x_3^{10} x_4^{17} x_5^{21} \\
&\quad + x_1^3 x_2 x_3^{18} x_4^{17} x_5^{13} + x_1^3 x_2 x_3^3 x_4^{14} x_5^{20} + x_1^3 x_2 x_3^3 x_4^{20} x_5^{14} + x_1^3 x_2 x_3^4 x_4^{20} x_5^{13} + x_1^3 x_2 x_3^5 x_4^{14} x_5^{18} \\
&\quad + x_1^3 x_2 x_3^5 x_4^{18} x_5^{14} + x_1^3 x_2 x_3^{10} x_4^{11} x_5^{16} + x_1^3 x_2 x_3^{10} x_4^{13} x_5^{14} + x_1^3 x_2 x_3^{12} x_4^{11} x_5^{14} + x_1^3 x_2 x_3^5 x_4^{13} x_5^{14} \\
&\quad + x_1^3 x_2 x_3^8 x_4^{12} x_5^{13} + x_1^3 x_2 x_3^7 x_4^{13} x_5^{13}, \\
\sum T &= x_1^3 x_2 x_3^{12} x_4^9 x_5^{13} + x_1^3 x_2 x_3^8 x_4^{12} x_5^{13} + x_1^3 x_2 x_3^4 x_4^{13} x_5^{14} + x_1^3 x_2 x_3^4 x_4^{14} x_5^{13} + x_1^3 x_2 x_3^{10} x_4^9 x_5^{13} \\
&\quad + x_1^3 x_2 x_3^4 x_4^{12} x_5^{13}.
\end{aligned}$$

The above relations show that d_1 is strictly inadmissible. The lemma follows. $\square$

Proof of Proposition 3.4.20. Let a be an admissible monomial in $\mathcal{P}_5^+$ such that $\omega(a) = \bar{\omega}_{(1)}$. Then $a = x_t x_k x_m u^2$ with $1 \leq t < k < m \leq 5$ and $u \in (\mathcal{P}_5^+)_{22}$. Since a is admissible, by Theorem 2.2.1, u is also admissible. Further, $u \in \mathcal{B}_5^+(\omega_{(1)})$.

Let $X \in \mathcal{B}_5^+(\omega_{(1)})$ such that $x_t x_k x_m X^2 \in (\mathcal{P}_5^+)_{47}$. By a direct computation using Proposition 3.4.11, we see that if $x_t x_k x_m X^2 \neq \mathcal{Y}_{47,j}$, $1 \leq j \leq 370$, then there is a monomial b which is given in one of Lemmas 3.4.18 - 3.4.23 such that $x_t x_k x_m X^2 = b Y^{2^r}$ with suitable monomial $Y \in \mathcal{P}_5$, and $r = \max\{s \in \mathbb{Z} : \omega_s(b) > 0\}$. By Theorem 2.2.1, $x_t x_k x_m X^2$ is inadmissible. On the other hand, we have $a = x_t x_k x_m u^2$ and a is admissible, hence $a = \mathcal{Y}_j$ for some $j = 1, 2, \dots, 370$. This completes the proof of the proposition. $\square$

Remark 3.4.24. For $1 \leq j \leq 370$, we have $[\mathcal{Y}_j := \mathcal{Y}_{47,j}]_{\bar{\omega}_{(1)}} \neq [0]$. Indeed, suppose that there is a linear relation

$$\mathcal{S} = \sum_{1 \leq j \leq 370} \gamma_j \mathcal{Y}_j \equiv_{\bar{\omega}_{(1)}} 0,$$

where $\gamma_j \in \mathbb{Z}/2$, for all j . Based on Theorem 3.3.5 and Proposition 3.1.1, for $(k; \mathcal{K}) \in \mathcal{N}_5$, we explicitly compute $\pi_{(k; \mathcal{K})}(\mathcal{S})$ in terms of a given minimal set of $\mathcal{A}_2$ -generators in $\mathcal{P}_4$ (modulo $(\mathcal{A}_2^+ \mathcal{P}_4)$). By computing from the relations $\pi_{(k; \mathcal{K})}(\mathcal{S}) \equiv_{\bar{\omega}_{(1)}} 0$, $\ell(\mathcal{K}) = 1$, one gets $\gamma_j = 0$, $\forall j$, $1 \leq j \leq 370$.

Note that these computations are similar to the proof of Propositions 3.4.1 and 3.4.11. Combining this and Proposition 3.4.20, we have a direct corollary.

Corollary 3.4.25. *The set $[\overline{\Phi}^+(\mathcal{B}_4(\overline{\omega}_{(1)})) \cup \mathcal{B}^+(5, \overline{\omega}_{(1)}) \cup \mathcal{F}]_{\overline{\omega}_{(1)}}$ is a basis of the $\mathbb{Z}/2$ -vector space $Q\mathcal{P}_5^+(\overline{\omega}_{(1)})$. This implies $\dim(Q\mathcal{P}_5^+(\overline{\omega}_{(1)})) = 370$.*

Remark 3.4.26. Consider the weight vector $\omega^{**} = (3, 2, 4)$ with $\deg \omega^{**} = 23$. By using a result in [55], we see that the following monomials are strictly inadmissible:

$$\begin{aligned} e_1 &= x_1^7 x_2^2 x_3^4 x_4^5 x_5^5, & e_2 &= x_1^7 x_2^2 x_3^5 x_4^4 x_5^5, & e_3 &= x_1^7 x_2^2 x_3^5 x_4^5 x_5^4, & e_4 &= x_1^3 x_2^7 x_3^4 x_4^4 x_5^5, \\ e_5 &= x_1^3 x_2^7 x_3^4 x_4^5 x_5^4, & e_6 &= x_1^3 x_2^7 x_3^5 x_4^4 x_5^4, & e_7 &= x_1^7 x_2^3 x_3^4 x_4^4 x_5^5, & e_8 &= x_1^7 x_2^3 x_3^4 x_4^5 x_5^4, \\ e_9 &= x_1^7 x_2^3 x_3^4 x_4^5 x_5^4, & e_{10} &= x_1^3 x_2^4 x_3^4 x_4^5 x_5^7, & e_{11} &= x_1^3 x_2^4 x_3^4 x_4^5 x_5^7, & e_{12} &= x_1^3 x_2^4 x_3^5 x_4^4 x_5^7, \\ e_{13} &= x_1^3 x_2^4 x_3^5 x_4^4 x_5^7, & e_{14} &= x_1^3 x_2^4 x_3^5 x_4^5 x_5^4, & e_{15} &= x_1^3 x_2^4 x_3^7 x_4^5 x_5^4, & e_{16} &= x_1^3 x_2^5 x_3^4 x_4^4 x_5^7, \\ e_{17} &= x_1^3 x_2^5 x_3^4 x_4^5 x_5^7, & e_{18} &= x_1^3 x_2^5 x_3^4 x_4^5 x_5^7, & e_{19} &= x_1^3 x_2^6 x_3^4 x_4^5 x_5^5, & e_{20} &= x_1^3 x_2^6 x_3^5 x_4^4 x_5^5, \\ e_{21} &= x_1^3 x_2^6 x_3^5 x_4^4 x_5^5, & e_{22} &= x_1^3 x_2^4 x_3^6 x_4^5 x_5^5, & e_{23} &= x_1^3 x_2^4 x_3^6 x_4^5 x_5^5, & e_{24} &= x_1^3 x_2^4 x_3^5 x_4^5 x_5^6, \\ e_{25} &= x_1^3 x_2^5 x_3^4 x_4^6 x_5^5, & e_{26} &= x_1^3 x_2^5 x_3^6 x_4^4 x_5^5, & e_{27} &= x_1^3 x_2^5 x_3^4 x_4^5 x_5^6, & e_{28} &= x_1^3 x_2^5 x_3^6 x_4^5 x_5^4, \\ e_{29} &= x_1^3 x_2^5 x_3^5 x_4^4 x_5^6, & e_{30} &= x_1^3 x_2^5 x_3^5 x_4^6 x_5^4. \end{aligned}$$

Note that $\omega(e_k) = \omega^{**}$, $k = 1, 2, \dots, 30$. Let $X \in \mathcal{B}_5^+(47)$ such that either $\omega(X) = \overline{\omega}_{(2)}$ or $\omega(X) = \overline{\omega}_{(3)}$. Then $X = X_{(\{\ell, m\}, 5)} y^2$ with $1 \leq \ell < m \leq 5$, and $y \in (\mathcal{P}_5^+)_{22}$. By a direct computation using Theorem 2.2.1, and Proposition 3.4.15(I), (II), we see that either $y \in \mathcal{B}_5^+(\omega_{(2)})$ or $y \in \mathcal{B}_5^+(\omega_{(3)})$ and there is a monomial $Z = e_i$ for some i , $1 \leq i \leq 30$ such that $X = X_{(\{\ell, m\}, 5)} y^2 = Zu^{2^t}$, $1 \leq \ell < m \leq 5$, with suitable monomial $u \in \mathcal{P}_5$, and $t = \max\{a \in \mathbb{Z} : \omega_a(Z) > 0\}$. By Theorem 2.2.1, we get either $[X] = [0]_{\overline{\omega}_{(2)}}$ or $[X] = [0]_{\overline{\omega}_{(3)}}$. As a consequence, we get the following.

Proposition 3.4.27. *The $\mathbb{Z}/2$ -vector spaces $Q\mathcal{P}_5^+(\overline{\omega}_{(2)})$ and $Q\mathcal{P}_5^+(\overline{\omega}_{(3)})$ are trivial.*

Lemma 3.4.28. *The set $[\{\mathcal{Y}_{47,j} : 371 \leq j \leq 480\}]_{\overline{\omega}_{(4)}}$ is a minimal system of generators for $Q\mathcal{P}_5^+(\overline{\omega}_{(4)})$, where the monomials $\mathcal{Y}_{47,j} : 371 \leq j \leq 480$, are determined in Sect. 6.7.*

In order to prove the lemma, we need to use some results. As an immediate consequence of a result in [47], we obtain the following.

Lemma 3.4.29. *The following monomials are strictly inadmissible:*

$$\begin{aligned} I) & x_a^7 x_b^2 x_c^{15} x_d^7, x_a^7 x_b^{15} x_c^2 x_d^7, x_a^7 x_b^6 x_c^3 x_d^{15}, x_a^7 x_b^3 x_c^6 x_d^{15}, x_a^7 x_b^3 x_c^{14} x_d^7, \\ & x_a^7 x_b^{14} x_c^3 x_d^7, x_a^7 x_b^6 x_c^{11} x_d^7, x_a^7 x_b^{11} x_c^6 x_d^7, \quad 1 \leq b < c \leq 5, \quad 1 \leq a, d \leq 5, \quad a \neq d, \quad a, d \neq b, c; \end{aligned}$$

II) $\rho_{(k,5)}(v)$, $1 \leq k \leq 5$, where v is one of the following monomials:

$$x_1^7 x_2^7 x_3^7 x_4^{10}, x_1^7 x_2^7 x_3^{10} x_4^7, x_1^7 x_2^{10} x_3^7 x_4^7.$$

We now consider the weight vector $\omega^{***} = (3, 4, 3, 1)$ with $\deg \omega^{***} = 31$. The following lemma can be easily proved by a direct computation.

Lemma 3.4.30. *All permutations of the following monomials are strictly inadmissible:*

$$\begin{aligned} & x_1 x_2^2 x_3^6 x_4^7 x_5^{15}, x_1 x_2^2 x_3^4 x_4^7 x_5^7, x_1 x_2^6 x_3^7 x_4^7 x_5^{10}, x_1^7 x_2^2 x_3^2 x_4^5 x_5^{15}, x_1^7 x_2^2 x_3^2 x_4^7 x_5^{13}, x_1^3 x_2^2 x_3^4 x_4^7 x_5^{15}, \\ & x_1^3 x_2^2 x_3^7 x_4^7 x_5^{12}, x_1^7 x_2^2 x_3^4 x_4^7 x_5^{11}, x_1^7 x_2^2 x_3^5 x_4^7 x_5^{10}, x_1^7 x_2^2 x_3^6 x_4^7 x_5^9, x_1^7 x_2^2 x_3^7 x_4^7 x_5^8, x_1^3 x_2^3 x_3^4 x_4^6 x_5^{15}, \\ & x_1^3 x_2^3 x_3^4 x_4^7 x_5^{14}, x_1^3 x_2^3 x_3^6 x_4^7 x_5^{12}, x_1^3 x_2^4 x_3^6 x_4^7 x_5^{11}, x_1^3 x_2^4 x_3^7 x_4^7 x_5^{10}, x_1^3 x_2^6 x_3^7 x_4^7 x_5^8. \end{aligned}$$

Lemma 3.4.31. *If (m, n, p, q, r) is a permutation of $(1, 2, 3, 4, 5)$, then the following monomials are strictly inadmissible:*

$$\begin{aligned}
x_m^3 x_n^{13} x_p^6 x_q^6 x_r^6 &\neq X = x_1^3 x_2^3 x_3^{13} x_4^6 x_5^6, \quad x_1^3 x_2^{13} x_3^6 x_4^6 x_5^6, \\
x_m^7 x_n^2 x_p^5 x_q^6 x_r^{11} &\neq Y = x_1^7 x_2^{11} x_3^5 x_4^6 x_5^6, \quad x_1^7 x_2^{11} x_3^5 x_4^6 x_5^2, \\
x_m^3 x_n^5 x_p^6 x_q^6 x_r^{11} &\neq Z = x_1^3 x_2^5 x_3^6 x_4^6 x_5^{11}, \quad x_1^3 x_2^5 x_3^{11} x_4^6 x_5^6, \\
x_m^3 x_n^7 x_p^9 x_q^6 x_r^6 &\neq G = x_1^3 x_2^7 x_3^9 x_4^6 x_5^6, \quad x_1^7 x_2^3 x_3^9 x_4^6 x_5^6, \quad x_1^7 x_2^9 x_3^6 x_4^6 x_5^6, \\
x_m^3 x_n^2 x_p^6 x_q^7 x_r^{13} &\neq H = x_1^3 x_2^7 x_3^{13} x_4^2 x_5^6, \quad x_1^3 x_2^7 x_3^{13} x_4^6 x_5^2, \quad x_1^7 x_2^3 x_3^{13} x_4^2 x_5^6, \quad x_1^7 x_2^3 x_3^{13} x_4^6 x_5^2, \\
x_m^3 x_n^5 x_p^6 x_q^7 x_r^{10} &\neq F = x_1^3 x_2^5 x_3^6 x_4^7 x_5^{10}, \quad x_1^3 x_2^5 x_3^7 x_4^6 x_5^{10}, \quad x_1^3 x_2^5 x_3^7 x_4^{10} x_5^6, \quad x_1^3 x_2^7 x_3^5 x_4^6 x_5^{10}, \\
&\quad x_1^3 x_2^7 x_3^5 x_4^{10} x_5^6, \quad x_1^7 x_2^3 x_3^5 x_4^6 x_5^{10}, \quad x_1^7 x_2^3 x_3^5 x_4^{10} x_5^6, \\
x_m^3 x_n^5 x_p^{14} x_q^3 x_r^6 &\neq T = x_1^3 x_2^3 x_3^5 x_4^6 x_5^{14}, \quad x_1^3 x_2^3 x_3^5 x_4^{14} x_5^6, \quad x_1^3 x_2^5 x_3^3 x_4^6 x_5^{14}, \\
&\quad x_1^3 x_2^5 x_3^3 x_4^{14} x_5^6, \quad x_1^3 x_2^5 x_3^6 x_4^3 x_5^{14}, \quad x_1^3 x_2^5 x_3^6 x_4^{14} x_5^3.
\end{aligned}$$

Proof. It is easy to see that the weight vector of these monomials is ω^{***} . Note that the monomials X, Y, Z, G, H, F , and T are admissible. We now prove the lemma for the monomials $f = x_m^3 x_n^{13} x_p^6 x_q^6 x_r^6$ and $g = x_m^3 x_n^5 x_p^{14} x_q^3 x_r^6$. The others can be proved by an argument similar to the proofs of Lemmas 3.4.18 and 3.4.19. Applying the Cartan formula, we get

$$f = x_m^3 x_n^{13} x_p^6 x_q^6 x_r^6 + Sq^1(\sum \bar{A}_1) + Sq^2(\sum \bar{B}_1) + Sq^4(\sum \bar{C}_1) \text{ modulo } (\mathcal{P}_5^-(\omega^{***})), \text{ where}$$

$$\begin{aligned}
\sum \bar{A}_1 &= x_m^3 x_n^7 x_p^5 x_q^5 x_r^{10} + x_m^3 x_n^7 x_p^5 x_q^6 x_r^9 + x_m^3 x_n^7 x_p^6 x_q^5 x_r^9 + x_m^3 x_n^7 x_p^6 x_q^9 x_r^5 + x_m^3 x_n^7 x_p^9 x_q^6 x_r^5 \\
&\quad + x_m^3 x_n^{11} x_p^5 x_q^6 x_r^5 + x_m^3 x_n^{14} x_p^3 x_q^5 x_r^5 + x_m^6 x_n^{11} x_p^5 x_q^3 x_r^5, \\
\sum \bar{B}_1 &= x_m^3 x_n^{11} x_p^5 x_q^5 x_r^5 + x_m^3 x_n^{11} x_p^6 x_q^3 x_r^6 + x_m^3 x_n^{13} x_p^5 x_q^3 x_r^5 + x_m^5 x_n^7 x_p^3 x_q^5 x_r^9 + x_m^5 x_n^7 x_p^3 x_q^9 x_r^5 \\
&\quad + x_m^5 x_n^7 x_p^5 x_q^3 x_r^9 + x_m^5 x_n^7 x_p^9 x_q^3 x_r^5 + x_m^5 x_n^9 x_p^3 x_q^3 x_r^9 + x_m^5 x_n^{11} x_p^3 x_q^5 x_r^5, \\
\sum \bar{C}_1 &= x_m^3 x_n^7 x_p^3 x_q^5 x_r^9 + x_m^3 x_n^7 x_p^3 x_q^9 x_r^5 + x_m^3 x_n^7 x_p^5 x_q^3 x_r^9 + x_m^3 x_n^7 x_p^9 x_q^3 x_r^5 + x_m^3 x_n^9 x_p^3 x_q^3 x_r^9 \\
&\quad + x_m^3 x_n^9 x_p^5 x_q^5 x_r^5 + x_m^3 x_n^{11} x_p^3 x_q^5 x_r^5 + x_m^3 x_n^{11} x_p^5 x_q^3 x_r^5,
\end{aligned}$$

Since $x_m^3 x_n^{13} x_p^6 x_q^6 x_r^6 < f$, the monomial f is strictly inadmissible. Next, by a direct computation, we have

$$g = \sum \mathcal{L} + Sq^1(\sum \bar{A}_2) + Sq^2(\sum \bar{B}_2) + Sq^4(\sum \bar{C}_2) \text{ modulo } (\mathcal{P}_5^-(\omega^{***})), \text{ where}$$

$$\begin{aligned}
\sum \mathcal{L} &= x_m^3 x_n^3 x_p^{14} x_q^5 x_r^6 + x_m^3 x_n^5 x_p^7 x_q^6 x_r^{10} + x_m^3 x_n^5 x_p^{11} x_q^6 x_r^6, \\
\sum \bar{A}_2 &= x_m^3 x_n^3 x_p^{13} x_q^5 x_r^6 + x_m^3 x_n^3 x_p^{14} x_q^5 x_r^5 + x_m^3 x_n^5 x_p^{11} x_q^5 x_r^6 + x_m^3 x_n^5 x_p^{13} x_q^3 x_r^6 + x_m^3 x_n^6 x_p^7 x_q^9 x_r^5 \\
&\quad + x_m^3 x_n^6 x_p^{11} x_q^5 x_r^5 + x_m^3 x_n^{10} x_p^7 x_q^5 x_r^5 + x_m^6 x_n^5 x_p^{11} x_q^3 x_r^5, \\
\sum \bar{B}_2 &= x_m^3 x_n^5 x_p^{11} x_q^5 x_r^5 + x_m^3 x_n^5 x_p^{13} x_q^3 x_r^5 + x_m^3 x_n^6 x_p^{11} x_q^3 x_r^6 + x_m^5 x_n^3 x_p^7 x_q^5 x_r^9 + x_m^5 x_n^3 x_p^7 x_q^9 x_r^5 \\
&\quad + x_m^5 x_n^3 x_p^9 x_q^3 x_r^9 + x_m^5 x_n^3 x_p^{11} x_q^5 x_r^5 + x_m^5 x_n^5 x_p^7 x_q^3 x_r^9 + x_m^5 x_n^9 x_p^7 x_q^3 x_r^5, \\
\sum \bar{C}_2 &= x_m^3 x_n^3 x_p^7 x_q^5 x_r^9 + x_m^3 x_n^3 x_p^7 x_q^9 x_r^5 + x_m^3 x_n^3 x_p^9 x_q^3 x_r^9 + x_m^3 x_n^3 x_p^{11} x_q^5 x_r^5 + x_m^3 x_n^5 x_p^7 x_q^3 x_r^9 \\
&\quad + x_m^3 x_n^5 x_p^7 x_q^6 x_r^6 + x_m^3 x_n^5 x_p^9 x_q^5 x_r^5 + x_m^3 x_n^5 x_p^{11} x_q^3 x_r^5 + x_m^3 x_n^9 x_p^7 x_q^3 x_r^5.
\end{aligned}$$

These relations show that g is strictly inadmissible. The lemmas follows. $\square$

The proof of the following lemmas is analogous to the proofs of Lemmas 3.4.18, 3.4.19, 3.4.30 and 3.4.31.

Lemma 3.4.32. *If (p, q, r) is a permutation of $(3, 4, 5)$, then the following monomials are strictly inadmissible:*

$$\begin{aligned}
u_1 &= x_1 x_2^6 x_p^6 x_q^3 x_r^{15}, \quad u_2 = x_1^3 x_2^6 x_p^6 x_q x_r^{15}, \quad u_3 = x_1^{16} x_2^6 x_p^6 x_q x_r^3, \quad u_4 = x_1 x_2^6 x_p^3 x_q^7 x_r^{14}, \\
u_5 &= x_1 x_2^{14} x_p^3 x_q^6 x_r^7, \quad u_6 = x_1^3 x_2^6 x_p x_q^7 x_r^{14}, \quad u_7 = x_1^3 x_2^{14} x_p x_q^6 x_r^7, \quad u_8 = x_1^7 x_2^6 x_p x_q^3 x_r^{14}, \\
u_9 &= x_1^7 x_2^{14} x_p x_q^3 x_r^6, \quad u_{10} = x_1 x_2^6 x_p^6 x_q^7 x_r^{11}, \quad u_{11} = x_1^7 x_2^6 x_p x_q^6 x_r^{11}, \quad u_{12} = x_1^3 x_2^2 x_p^5 x_q^6 x_r^{15}, \\
u_{13} &= x_1^3 x_2^6 x_p^2 x_q^5 x_r^{15}, \quad u_{14} = x_1^{15} x_2^2 x_p^3 x_q^5 x_r^6, \quad u_{15} = x_1^{15} x_2^6 x_p^2 x_q^3 x_r^5, \quad u_{16} = x_1^3 x_2^2 x_p^5 x_q^7 x_r^{14}, \\
u_{17} &= x_1^3 x_2^{14} x_p^2 x_q^5 x_r^7, \quad u_{18} = x_1^7 x_2^2 x_p^{14} x_q^3 x_r^5, \quad u_{19} = x_1^7 x_2^{14} x_p^2 x_q^3 x_r^5.
\end{aligned}$$

Lemma 3.4.33. *The following monomials are strictly inadmissible:*

$$\begin{aligned}
I) \quad & u_{20} = x_1 x_2^{15} x_3^6 x_q^3 x_r^6, \quad u_{21} = x_1^3 x_2^{15} x_3^6 x_q x_r^6, \quad u_{22} = x_1^{15} x_2 x_3^6 x_q^3 x_r^6, \quad u_{23} = x_1^{15} x_2^3 x_3^6 x_q x_r^6, \\
& u_{24} = x_1^7 x_2 x_3^6 x_q^3 x_r^{14}, \quad u_{25} = x_1^7 x_2 x_3^{14} x_q^3 x_r^6, \quad u_{26} = x_1^7 x_2^3 x_3^6 x_q x_r^{14}, \quad u_{27} = x_1^7 x_2^3 x_3^{14} x_q x_r^6, \\
& u_{28} = x_1 x_2^3 x_3^{14} x_q^6 x_r^7, \quad u_{29} = x_1 x_2^7 x_3^6 x_q^3 x_r^{14}, \quad u_{30} = x_1 x_2^7 x_3^{14} x_q^3 x_r^6, \quad u_{31} = x_1^3 x_2 x_3^{14} x_q^6 x_r^7, \\
& u_{32} = x_1^3 x_2^7 x_3^6 x_q x_r^{14}, \quad u_{33} = x_1^3 x_2^7 x_3^{14} x_q x_r^6, \quad u_{34} = x_1 x_2^7 x_3^6 x_q^{11}, \quad u_{35} = x_1^7 x_2 x_3^6 x_q^{11}, \\
& u_{36} = x_1^7 x_2^{11} x_3^6 x_q x_r^6, \quad u_{37} = x_1^3 x_2^{15} x_3^2 x_q^5 x_r^6, \quad u_{38} = x_1^3 x_2^{15} x_3^6 x_q^2 x_r^5, \quad u_{39} = x_1^{15} x_2^3 x_3^2 x_q^5 x_r^6, \\
& u_{40} = x_1^{15} x_2^3 x_3^6 x_q^2 x_r^5, \quad u_{41} = x_1^3 x_2^5 x_3^{14} x_q^2 x_r^7, \quad u_{42} = x_1^3 x_2^5 x_3^{14} x_q^2 x_r^7, \quad u_{43} = x_1^3 x_2^7 x_3^2 x_q^5 x_r^{14}, \\
& u_{44} = x_1^3 x_2^7 x_3^{14} x_q^2 x_r^5, \quad u_{45} = x_1^3 x_2^3 x_3^2 x_q^5 x_r^{14}, \quad u_{46} = x_1^7 x_2^3 x_3^{14} x_q^2 x_r^5, \quad \text{where } q, r = 4, 5, \\
& q \neq r.
\end{aligned}$$

$$II) \quad u_{47} = x_1 x_2^3 x_3^6 x_4^{14} x_5^7, \quad u_{48} = x_1^3 x_2 x_3^6 x_4^{14} x_5^7, \quad u_{49} = x_1^3 x_2^5 x_3^2 x_4^{14} x_5^7.$$

Note that $\omega(u_t) = \omega^{***}$ for $t = 1, 2, \dots, 49$. Next we consider the following lemma.

Lemma 3.4.34. *The following monomials are strictly inadmissible:*

$$\begin{aligned}
A_1 &= x_1 x_2^3 x_3^6 x_4^{15} x_5^{22}, & A_2 &= x_1 x_2^3 x_3^{15} x_4^6 x_5^{22}, & A_3 &= x_1 x_2^3 x_3^{15} x_4^{22} x_5^6, & A_4 &= x_1 x_2^{15} x_3^3 x_4^6 x_5^{22}, \\
A_5 &= x_1 x_2^{15} x_3^3 x_4^{22} x_5^6, & A_6 &= x_1 x_2^{15} x_3^{19} x_4^6 x_5^6, & A_7 &= x_1^3 x_2 x_3^6 x_4^{15} x_5^{22}, & A_8 &= x_1^3 x_2^5 x_3^2 x_4^{15} x_5^{22}, \\
A_9 &= x_1^3 x_2^5 x_3^6 x_4^{15} x_5^{18}, & A_{10} &= x_1^3 x_2 x_3^{15} x_4^6 x_5^{22}, & A_{11} &= x_1^3 x_2 x_3^{15} x_4^{22} x_5^6, & A_{12} &= x_1^3 x_2^5 x_3^{15} x_4^2 x_5^{22}, \\
A_{13} &= x_1^3 x_2^5 x_3^{15} x_4^6 x_5^{18}, & A_{14} &= x_1^3 x_2^5 x_3^{15} x_4^{18} x_5^6, & A_{15} &= x_1^3 x_2^5 x_3^{15} x_4^{22} x_5^2, & A_{16} &= x_1^3 x_2^{13} x_3^3 x_4^6 x_5^{22}, \\
A_{17} &= x_1^3 x_2^{13} x_3^3 x_4^{22} x_5^6, & A_{18} &= x_1^3 x_2^{13} x_3^{19} x_4^6 x_5^6, & A_{19} &= x_1^3 x_2^{15} x_3^6 x_4^{22} x_5^2, & A_{20} &= x_1^3 x_2^{15} x_3^6 x_4^2 x_5^6, \\
A_{21} &= x_1^3 x_2^{15} x_3^5 x_4^{22} x_5^2, & A_{22} &= x_1^3 x_2^{15} x_3^5 x_4^{18} x_5^6, & A_{23} &= x_1^3 x_2^{15} x_3^5 x_4^{18} x_5^6, & A_{24} &= x_1^3 x_2^{15} x_3^5 x_4^2 x_5^2, \\
A_{25} &= x_1^3 x_2^{15} x_3^{17} x_4^6 x_5^6, & A_{26} &= x_1^3 x_2^{15} x_3^{21} x_4^6 x_5^2, & A_{27} &= x_1^3 x_2^{15} x_3^{21} x_4^6 x_5^2, & A_{28} &= x_1^7 x_2^3 x_3^3 x_4^6 x_5^{22}, \\
A_{29} &= x_1^7 x_2^3 x_3^3 x_4^{22} x_5^6, & A_{30} &= x_1^7 x_2^3 x_3^{19} x_4^6 x_5^6, & A_{31} &= x_1^7 x_2^{11} x_3^{17} x_4^6 x_5^6, & A_{32} &= x_1^7 x_2^{25} x_3^3 x_4^6 x_5^6, \\
A_{33} &= x_1^{15} x_2 x_3^3 x_4^6 x_5^{22}, & A_{34} &= x_1^{15} x_2 x_3^3 x_4^{22} x_5^6, & A_{35} &= x_1^{15} x_2 x_3^{19} x_4^6 x_5^6, & A_{36} &= x_1^{15} x_2^3 x_3^6 x_4^6 x_5^{22}, \\
A_{37} &= x_1^{15} x_2 x_3^3 x_4^{22} x_5^6, & A_{38} &= x_1^{15} x_2^3 x_3^5 x_4^2 x_5^{22}, & A_{39} &= x_1^{15} x_2^3 x_3^5 x_4^6 x_5^{18}, & A_{40} &= x_1^{15} x_2^3 x_3^5 x_4^{18} x_5^6, \\
A_{41} &= x_1^{15} x_2^3 x_3^5 x_4^{22} x_5^2, & A_{42} &= x_1^{15} x_2^3 x_3^{17} x_4^6 x_5^6, & A_{43} &= x_1^{15} x_2^3 x_3^{21} x_4^2 x_5^6, & A_{44} &= x_1^{15} x_2^3 x_3^{21} x_4^6 x_5^2, \\
A_{45} &= x_1^{15} x_2^{17} x_3^3 x_4^6 x_5^6, & A_{46} &= x_1^{15} x_2^{19} x_3^6 x_4^6 x_5^6, & A_{47} &= x_1^{15} x_2^{19} x_3^5 x_4^6 x_5^6, & A_{48} &= x_1^{15} x_2^{19} x_3^5 x_4^6 x_5^2, \\
A_{49} &= x_1^7 x_2^{11} x_3^5 x_4^{18} x_5^6, & A_{50} &= x_1^7 x_2^{11} x_3^5 x_4^{18} x_5^6.
\end{aligned}$$

Proof. We have $\omega(A_i) = \bar{\omega}_{(4)}$, $1 \leq i \leq 50$. We prove the lemma for the monomials $A_{23} = x_1^3 x_2^{15} x_3^5 x_4^{18} x_5^6$ and $A_{29} = x_1^7 x_2^3 x_3^3 x_4^{22} x_5^6$. The others can be proved by using a similar technique as in Lemmas 3.4.22 and 3.4.23. Direct computing from Cartan's formula, we get

$$\begin{aligned}
A_{23} &= \sum_{1 \leq i \leq 3} b_i + Sq^1(\sum f) + Sq^2(\sum g) + Sq^4(\sum h) + Sq^8(\sum p) \text{ modulo } (\mathcal{P}_5^-(\bar{\omega}_{(4)})), \text{ where} \\
b_1 &= x_1^3 x_2^{11} x_3^5 x_4^{22} x_5^6, \quad b_2 = x_1^3 x_2^{13} x_3^3 x_4^{22} x_5^6, \quad b_3 = x_1^3 x_2^{13} x_3^6 x_4^{19} x_5^6, \\
\sum f &= x_1^3 x_2^7 x_3^6 x_4^{21} x_5^9 + x_1^3 x_2^7 x_3^6 x_4^{25} x_5^5 + x_1^3 x_2^{11} x_3^6 x_4^{21} x_5^5 + x_1^3 x_2^{14} x_3^3 x_4^{21} x_5^5 + x_1^3 x_2^{15} x_3^5 x_4^{14} x_5^9 \\
&\quad + x_1^3 x_2^{15} x_3^5 x_4^{17} x_5^6 + x_1^3 x_2^{15} x_3^9 x_4^{14} x_5^5 + x_1^3 x_2^{19} x_3^5 x_4^{14} x_5^5 + x_1^3 x_2^{22} x_3^5 x_4^{11} x_5^5, \\
\sum g &= x_1^5 x_2^7 x_3^3 x_4^{21} x_5^9 + x_1^5 x_2^7 x_3^3 x_4^{25} x_5^5 + x_1^5 x_2^9 x_3^3 x_4^{19} x_5^3 + x_1^5 x_2^{11} x_3^3 x_4^{17} x_5^9 + x_1^5 x_2^{15} x_3^5 x_4^{11} x_5^9 \\
&\quad + x_1^5 x_2^{15} x_3^9 x_4^{11} x_5^5 + x_1^5 x_2^{17} x_3^3 x_4^{17} x_5^3 + x_1^5 x_2^{17} x_3^3 x_4^{11} x_5^3 + x_1^5 x_2^{19} x_3^5 x_4^{11} x_5^5, \\
\sum h &= x_1^3 x_2^7 x_3^3 x_4^{21} x_5^9 + x_1^3 x_2^7 x_3^3 x_4^{25} x_5^5 + x_1^3 x_2^9 x_3^5 x_4^{21} x_5^5 + x_1^3 x_2^9 x_3^9 x_4^{19} x_5^3 + x_1^3 x_2^{11} x_3^3 x_4^{17} x_5^9 \\
&\quad + x_1^3 x_2^{11} x_3^3 x_4^{21} x_5^5 + x_1^3 x_2^{13} x_3^9 x_4^{13} x_5^5 + x_1^3 x_2^{15} x_3^5 x_4^{11} x_5^9 + x_1^3 x_2^{15} x_3^6 x_4^{13} x_5^5 + x_1^3 x_2^{15} x_3^5 x_4^{11} x_5^5 \\
&\quad + x_1^3 x_2^{17} x_3^3 x_4^{17} x_5^3 + x_1^3 x_2^{17} x_3^3 x_4^{11} x_5^3 + x_1^3 x_2^{19} x_3^5 x_4^{11} x_5^5, \\
\sum p &= x_1^3 x_2^{13} x_3^5 x_4^{13} x_5^5 + x_1^3 x_2^{13} x_3^6 x_4^{11} x_5^6.
\end{aligned}$$

Since $b_i < A_1$, $1 \leq i \leq 3$, A_1 is strictly inadmissible. By a similar computation, we obtain

$$A_{29} = \sum Z + Sq^1(\sum \bar{f}) + Sq^2(\sum \bar{g}) + Sq^4(\sum \bar{h}) + Sq^8(x_1^7 x_2^5 x_3^6 x_4^{15} x_5^6) \text{ modulo } (\mathcal{P}_5^-(\bar{\omega}_{(4)})),$$

where the polynomials $\sum Z$, $\sum \bar{f}$, $\sum \bar{g}$, and $\sum \bar{h}$ are as follows:

$$\begin{aligned}
\sum Z &= x_1^5 x_2^3 x_3^6 x_4^{23} x_5^{10} + x_1^5 x_2^3 x_3^6 x_4^{27} x_5^6 + x_1^5 x_2^3 x_3^{10} x_4^{23} x_5^6 + x_1^5 x_2^7 x_3^3 x_4^{22} x_5^{10} + x_1^5 x_2^7 x_3^3 x_4^{26} x_5^6 \\
&\quad + x_1^5 x_2^7 x_3^6 x_4^{19} x_5^{10} + x_1^5 x_2^7 x_3^{10} x_4^{19} x_5^6 + x_1^5 x_2^{11} x_3^3 x_4^{22} x_5^6 + x_1^5 x_2^{11} x_3^6 x_4^{19} x_5^6 + x_1^7 x_2^3 x_3^5 x_4^{22} x_5^{10} \\
&\quad + x_1^7 x_2^3 x_3^6 x_4^{23} x_5^8 + x_1^7 x_2^3 x_3^6 x_4^{25} x_5^6 + x_1^7 x_2^3 x_3^8 x_4^{23} x_5^6 + x_1^7 x_2^3 x_3^9 x_4^{22} x_5^6 + x_1^7 x_2^7 x_3^3 x_4^{22} x_5^8 \\
&\quad + x_1^7 x_2^7 x_3^3 x_4^{24} x_5^6, \\
\sum \bar{f} &= x_1^7 x_2^3 x_3^5 x_4^{19} x_5^{12} + x_1^7 x_2^3 x_3^5 x_4^{21} x_5^{10} + x_1^7 x_2^3 x_3^9 x_4^{21} x_5^6 + x_1^7 x_2^3 x_3^{12} x_4^{19} x_5^5 + x_1^7 x_2^6 x_3^9 x_4^{15} x_5^9 \\
&\quad + x_1^7 x_2^7 x_3^5 x_4^{19} x_5^8 + x_1^7 x_2^7 x_3^5 x_4^{21} x_5^6 + x_1^7 x_2^7 x_3^8 x_4^{19} x_5^5 + x_1^7 x_2^{10} x_3^5 x_4^{15} x_5^9 \\
&\quad + x_1^7 x_2^{10} x_3^9 x_4^{15} x_5^5, \\
\sum \bar{g} &= x_1^7 x_2^3 x_3^6 x_4^{19} x_5^{10} + x_1^7 x_2^3 x_3^6 x_4^{23} x_5^6 + x_1^7 x_2^3 x_3^9 x_4^{17} x_5^9 + x_1^7 x_2^3 x_3^{10} x_4^{19} x_5^6 + x_1^7 x_2^5 x_3^9 x_4^{15} x_5^9 \\
&\quad + x_1^7 x_2^7 x_3^3 x_4^{22} x_5^6 + x_1^7 x_2^7 x_3^6 x_4^{19} x_5^6 + x_1^7 x_2^9 x_3^3 x_4^{17} x_5^9 + x_1^7 x_2^9 x_3^5 x_4^{15} x_5^9 + x_1^7 x_2^9 x_3^9 x_4^{15} x_5^5 \\
&\quad + x_1^7 x_2^9 x_3^9 x_4^{17} x_5^3 + x_1^9 x_2^3 x_3^9 x_4^{15} x_5^9 + x_1^9 x_2^9 x_3^3 x_4^{15} x_5^9 + x_1^9 x_2^9 x_3^9 x_4^{15} x_5^3, \\
\sum \bar{h} &= x_1^5 x_2^3 x_3^6 x_4^{23} x_5^6 + x_1^5 x_2^7 x_3^3 x_4^{22} x_5^6 + x_1^5 x_2^7 x_3^6 x_4^{19} x_5^6.
\end{aligned}$$

The above equalities imply that A_2 is strictly inadmissible. The lemma is proved. $\square$

Proof of Lemma 3.4.28. Let b be an admissible monomial in $(\mathcal{P}_5^+)_{47}$ such that $\omega(b) = \bar{\omega}_{(4)}$. Then $\omega_1(b) = 3$ and $b = X_{(\{t,k\},5)} Y^2$ with $1 \leq t < k \leq 5$ and Y a monomial of degree 22 in $\mathcal{P}_5$. Since b is admissible, according to Theorem 2.2.1, $Y \in \mathcal{B}_5(\omega_{(4)})$.

Using Proposition 3.4.15(III) and a simple computation shows that if $Z \in \mathcal{B}_5(\omega_{(4)})$, $1 \leq t < k \leq 5$, and $X_{(\{t,k\},5)} Z^2 \neq \mathcal{Y}_{47,j}$, $\forall j$, $371 \leq j \leq 480$, then there is a monomial u which is given in one of Lemmas 3.4.29 - 3.4.34 such that $X_{(\{t,k\},5)} Z^2 = u g^{2^s}$ with a monomial $g \in \mathcal{P}_5$, and $s = \max\{\ell \in \mathbb{Z} : \omega_\ell(u) > 0\}$. Then, By Theorem 2.2.1, $X_{(\{t,k\},5)} Z^2$ is inadmissible. Finally, we see that $b = X_{(\{t,k\},5)} Y^2$ is admissible with $Y \in \mathcal{B}_5(\omega_{(4)})$; hence $b = \mathcal{Y}_{47,j}$ for some $j = 371, \dots, 480$. This implies $\mathcal{B}_5^+(\bar{\omega}_{(4)}) \subseteq \{\mathcal{Y}_{47,j} : 371 \leq j \leq 480\}$. $\square$

Proposition 3.4.35. $Q\mathcal{P}_5^+(\bar{\omega}_{(4)})$ is the $\mathbb{Z}/2$ -vector space of dimension 110 with a basis consisting of all the classes represented by the monomials $\mathcal{Y}_{47,j}$, $371 \leq j \leq 480$.

Proof. First, we show that the set $[V := \{\mathcal{Y}_{47,j} : 371 \leq j \leq 480\}]_{\bar{\omega}_{(4)}}$ is linearly independent in the space $Q\mathcal{P}_5^+(\bar{\omega}_{(4)})$. Indeed, suppose there is a linear relation $\mathcal{S} = \sum_{371 \leq j \leq 482} \gamma_j \mathcal{Y}_j \equiv_{\bar{\omega}_{(4)}} 0$ with $\gamma_j \in \mathbb{Z}/2$ and $\mathcal{Y}_{47,j} \in V$. By using Theorem 3.3.5 and Proposition 3.1.1, we determine explicitly $\pi_{(k;\mathcal{K})}(\mathcal{S})$ in terms of the admissible monomials in $(\mathcal{P}_4^+)_{47}$. From the relations $\pi_{(k;\mathcal{K})}(\mathcal{S}) \equiv_{\bar{\omega}_{(4)}} 0$ with $\ell(\mathcal{K}) > 0$, one gets $\gamma_j = 0$ for all $371 \leq j \leq 480$.

Now, by Lemma 3.4.28, to prove $[V]_{\bar{\omega}_{(4)}}$ is a basis of $Q\mathcal{P}_5^+(\bar{\omega}_{(4)})$ we need to show that $[\mathcal{Y}_{47,j}]_{\bar{\omega}_{(4)}} \neq [0]$ for all $\mathcal{Y}_{47,j} \in V$. By a similar argument as given in the proof of Propositions 3.4.1 and 3.4.11, we can prove that the set $[\mathcal{B}_5^+(\bar{\omega}_{(1)}) \cup V]$ is linearly independent in $(Q\mathcal{P}_5^+)_{47}$. This fact shows that $[\mathcal{Y}_{47,j}]_{\bar{\omega}_{(4)}} \neq [0]$ for all $\mathcal{Y}_{47,j}$. The proposition is proved. $\square$

Proposition 3.4.36. There exist exactly 14 admissible monomials in $(\mathcal{P}_5^+)_{47}$ such that their weight vectors are $\bar{\omega}_{(5)}$. Consequently $\dim(Q\mathcal{P}_5^+(\bar{\omega}_{(5)})) = 14$.

We prove the proposition by showing that

$$\mathcal{B}_5^+(\bar{\omega}_{(5)}) = \{\mathcal{Y}_{47,m} : 481 \leq m \leq 494\},$$

where the monomials $\mathcal{Y}_{47,m} : 481 \leq m \leq 494$, are described in Sect.6.7. We need some lemmas for the proof of this proposition.

The following result is proved by using a result in [55].

Lemma 3.4.37. *The following monomials are strictly inadmissible:*

- a) $x_u x_v^2 x_m^6 x_n^7 x_p^7$, $x_u^7 x_v^2 x_m^2 x_n^5 x_p^7$, $x_u^3 x_v^2 x_m^4 x_n^7 x_p^7$, $x_u^3 x_v^4 x_m^6 x_n^3 x_p^7$, $x_u^3 x_v^6 x_m^6 x_n^3 x_p^5$,
where (u, v, m, n, p) is a permutation of $(1, 2, 3, 4, 5)$;
- b) $x_1 x_2^6 x_q^6 x_r^3 x_t^7$, $x_1^3 x_2^6 x_q^6 x_r^7 x_t^3$, $x_1^3 x_2^2 x_q^6 x_r^5 x_t^7$, $x_1^3 x_2^6 x_q^2 x_r^5 x_t^7$, $x_1^7 x_2^2 x_q^6 x_r^3 x_t^5$,
where (q, r, t) is a permutation of $(3, 4, 5)$;
- c) $x_1 x_2^7 x_3^6 x_4^6 x_5^3$, $x_1 x_2^7 x_3^6 x_4^3 x_5^6$, $x_1^3 x_2^7 x_3^6 x_4^6 x_5^3$, $x_1^3 x_2^7 x_3^6 x_4^3 x_5^6$,
 $x_1^7 x_2^6 x_3^6 x_4^3 x_5^3$, $x_1^7 x_2^6 x_3^6 x_4^6 x_5^3$, $x_1^7 x_2^6 x_3^3 x_4^6 x_5^6$, $x_1^7 x_2^3 x_3^6 x_4^6 x_5^6$,
 $x_1^3 x_2^7 x_3^6 x_4^6 x_5^3$, $x_1^3 x_2^7 x_3^6 x_4^3 x_5^6$, $x_1^3 x_2^7 x_3^3 x_4^6 x_5^6$, $x_1^3 x_2^7 x_3^6 x_4^6 x_5^3$,
 $x_1^7 x_2^6 x_3^6 x_4^3 x_5^3$, $x_1^7 x_2^6 x_3^6 x_4^6 x_5^3$, $x_1^7 x_2^6 x_3^3 x_4^6 x_5^6$, $x_1^7 x_2^3 x_3^6 x_4^6 x_5^6$;
- d) $\rho_{(k,5)}(Z)$, $1 \leq k \leq 5$, where Z is one of the following monomials:
 $x_1^2 x_2^7 x_3^7 x_4^7$, $x_1^7 x_2^2 x_3^7 x_4^7$, $x_1^7 x_2^7 x_3^2 x_4^7$, $x_1^7 x_2^7 x_3^7 x_4^2$, $x_1^6 x_2^3 x_3^7 x_4^7$, $x_1^6 x_2^7 x_3^3 x_4^7$,
 $x_1^6 x_2^7 x_3^7 x_4^3$, $x_1^3 x_2^6 x_3^7 x_4^7$, $x_1^7 x_2^6 x_3^3 x_4^7$, $x_1^7 x_2^6 x_3^7 x_4^3$, $x_1^3 x_2^7 x_3^6 x_4^7$, $x_1^7 x_2^3 x_3^6 x_4^7$,
 $x_1^7 x_2^7 x_3^3 x_4^6$, $x_1^3 x_2^7 x_3^6 x_4^6$, $x_1^7 x_2^3 x_3^6 x_4^6$, $x_1^7 x_2^7 x_3^3 x_4^6$.

Lemma 3.4.38. *The following monomials are strictly inadmissible:*

$$\begin{aligned}
T_1 &= x_1 x_2^3 x_3^{14} x_4^{14} x_5^{15}, & T_2 &= x_1 x_2^3 x_3^{14} x_4^{15} x_5^{14}, & T_3 &= x_1 x_2^3 x_3^{15} x_4^{14} x_5^{14}, & T_4 &= x_1 x_2^{15} x_3^3 x_4^{14} x_5^{14}, \\
T_5 &= x_1^3 x_2^3 x_3^{14} x_4^{14} x_5^{15}, & T_6 &= x_1^3 x_2^3 x_3^{14} x_4^{15} x_5^{14}, & T_7 &= x_1^3 x_2^3 x_3^{15} x_4^{14} x_5^{14}, & T_8 &= x_1^3 x_2^{15} x_3^3 x_4^{14} x_5^{14}, \\
T_9 &= x_1^{15} x_2^3 x_3^3 x_4^{14} x_5^{14}, & T_{10} &= x_1^{15} x_2^3 x_3^3 x_4^{14} x_5^{14}, & T_{11} &= x_1^3 x_2^{13} x_3^2 x_4^{14} x_5^{15}, & T_{12} &= x_1^3 x_2^{13} x_3^2 x_4^{15} x_5^{14}, \\
T_{13} &= x_1^3 x_2^{13} x_3^{14} x_4^{14} x_5^{15}, & T_{14} &= x_1^3 x_2^{13} x_3^{14} x_4^{15} x_5^{14}, & T_{15} &= x_1^3 x_2^{13} x_3^{15} x_4^{14} x_5^{14}, & T_{16} &= x_1^3 x_2^{13} x_3^{15} x_4^{14} x_5^{14}, \\
T_{17} &= x_1^3 x_2^{15} x_3^{13} x_4^{14} x_5^{14}, & T_{18} &= x_1^3 x_2^{15} x_3^{13} x_4^{14} x_5^{14}, & T_{19} &= x_1^{15} x_2^3 x_3^{13} x_4^{14} x_5^{14}, & T_{20} &= x_1^{15} x_2^3 x_3^{13} x_4^{14} x_5^{14}, \\
T_{21} &= x_1^3 x_2^{13} x_3^{13} x_4^{14} x_5^{14}, & T_{22} &= x_1^3 x_2^{13} x_3^{13} x_4^{14} x_5^{14}, & T_{23} &= x_1^3 x_2^{13} x_3^{14} x_4^{14} x_5^{14}, & T_{24} &= x_1^3 x_2^{13} x_3^{14} x_4^{14} x_5^{14}, \\
T_{25} &= x_1^3 x_2^5 x_3^{10} x_4^{14} x_5^{15}, & T_{26} &= x_1^3 x_2^5 x_3^{10} x_4^{15} x_5^{14}, & T_{27} &= x_1^3 x_2^5 x_3^{14} x_4^{10} x_5^{15}, & T_{28} &= x_1^3 x_2^5 x_3^{14} x_4^{15} x_5^{10}, \\
T_{29} &= x_1^3 x_2^5 x_3^{15} x_4^{10} x_5^{14}, & T_{30} &= x_1^3 x_2^5 x_3^{15} x_4^{10} x_5^{14}, & T_{31} &= x_1^3 x_2^{15} x_3^5 x_4^{10} x_5^{14}, & T_{32} &= x_1^3 x_2^{15} x_3^5 x_4^{10} x_5^{14}, \\
T_{33} &= x_1^{15} x_2^3 x_3^5 x_4^{10} x_5^{14}, & T_{34} &= x_1^{15} x_2^3 x_3^5 x_4^{10} x_5^{14}, & T_{35} &= x_1^3 x_2^5 x_3^{11} x_4^{14} x_5^{14}, & T_{36} &= x_1^3 x_2^5 x_3^{14} x_4^{11} x_5^{14}, \\
T_{37} &= x_1^3 x_2^5 x_3^{14} x_4^{14} x_5^{11}, & T_{38} &= x_1^3 x_2^{13} x_3^6 x_4^{10} x_5^{15}, & T_{39} &= x_1^3 x_2^{13} x_3^6 x_4^{15} x_5^{10}, & T_{40} &= x_1^3 x_2^{13} x_3^{15} x_4^6 x_5^{10}, \\
T_{41} &= x_1^3 x_2^{15} x_3^{13} x_4^6 x_5^{10}, & T_{42} &= x_1^{15} x_2^3 x_3^6 x_4^{10} x_5^{10}, & T_{43} &= x_1^3 x_2^{13} x_3^6 x_4^{11} x_5^{14}, & T_{44} &= x_1^3 x_2^{13} x_3^6 x_4^{14} x_5^{11}, \\
T_{45} &= x_1^3 x_2^{13} x_3^6 x_4^{11} x_5^{14}, & T_{46} &= x_1^7 x_2^3 x_3^{14} x_4^{14} x_5^{14}, & T_{47} &= x_1^3 x_2^{13} x_3^7 x_4^{10} x_5^{14}, & T_{48} &= x_1^3 x_2^{13} x_3^7 x_4^{14} x_5^{10}, \\
T_{49} &= x_1^3 x_2^{13} x_3^{14} x_4^7 x_5^{10}.
\end{aligned}$$

Proof. It is easy to see that $\omega(T_s) = \bar{\omega}_{(5)}$ for $s = 1, 2, \dots, 49$. We prove the lemma for the monomials $T_{35} = x_1^3 x_2^5 x_3^{14} x_4^{11} x_5^{14}$ and $T_{42} = x_1^{15} x_2^3 x_3^6 x_4^{10} x_5^{10}$. The others can be proved by using a similar technique as in Lemmas 3.4.22, 3.4.23 and 3.4.34. Computing from the Cartan formula, we get

$$\begin{aligned}
T_{35} &= Sq^1 \left(x_1^3 x_2^6 x_3^{11} x_4^{13} x_5^{13} + x_1^3 x_2^6 x_3^{13} x_4^{11} x_5^{13} \right) + Sq^2 \left(x_1^5 x_2^3 x_3^7 x_4^{13} x_5^{17} + x_1^5 x_2^3 x_3^7 x_4^{17} x_5^{13} \right. \\
&\quad + x_1^5 x_2^3 x_3^9 x_4^{11} x_5^{17} + x_1^5 x_2^3 x_3^9 x_4^{17} x_5^{11} + x_1^5 x_2^3 x_3^{11} x_4^9 x_5^{17} + x_1^5 x_2^3 x_3^{11} x_4^{13} x_5^{13} \\
&\quad \left. + x_1^5 x_2^3 x_3^{13} x_4^7 x_5^{17} + x_1^5 x_2^3 x_3^{13} x_4^{11} x_5^{13} + x_1^5 x_2^3 x_3^{17} x_4^7 x_5^{13} + x_1^5 x_2^3 x_3^{17} x_4^9 x_5^{11} \right) \\
&\quad + Sq^4 \left(x_1^3 x_2^3 x_3^7 x_4^{13} x_5^{17} + x_1^3 x_2^3 x_3^7 x_4^{17} x_5^{13} + x_1^3 x_2^3 x_3^9 x_4^{11} x_5^{17} + x_1^3 x_2^3 x_3^9 x_4^{17} x_5^{11} \right. \\
&\quad \left. + x_1^3 x_2^3 x_3^{11} x_4^9 x_5^{17} x_1^3 x_2^3 x_3^{11} x_4^{13} x_5^{13} + x_1^3 x_2^3 x_3^{13} x_4^7 x_5^{17} + x_1^3 x_2^3 x_3^{13} x_4^{11} x_5^{13} \right) \\
&\quad \left. + x_1^3 x_2^3 x_3^{17} x_4^7 x_5^{13} + x_1^3 x_2^3 x_3^{17} x_4^9 x_5^{11} + x_1^3 x_2^5 x_3^{11} x_4^{14} x_5^{14} \text{ modulo } (\mathcal{P}_5^-(\bar{\omega}_{(5)})) \right).
\end{aligned}$$

We have $x_1^3 x_2^5 x_3^{11} x_4^{14} x_5^{14} < T_{35}$. This implies that T_{35} is strictly inadmissible. By a similar computation, we obtain

$$T_{42} = (p_1 + p_2) \text{ modulo } (\mathcal{A}_2^+ \mathcal{P}_5 + \mathcal{P}_5^-(\bar{\omega}_{(5)})),$$

where $p_1 = x_1^3 x_2^7 x_3^{10} x_4^{13} x_5^{14}$, $p_2 = x_1^3 x_2^{13} x_3^3 x_4^{14} x_5^{14}$. Since $p_i < T_{42}$, $i = 1, 2$, the monomial T_{42} is strictly inadmissible. The lemma follows. $\square$

Proof of Proposition 3.4.36. Suppose that X is a admissible monomial in $\mathcal{P}_5^+$ and $\omega(X) = \bar{\omega}_{(5)}$. Then $\omega_1(X) = 3$ and $X = x_a x_b x_c u^2$ with $1 \leq a < b < c \leq 5$ and $u \in (\mathcal{P}_5^+)_{22}$. Then, by Theorem 2.2.1, u is admissible and $u \in \mathcal{B}_5^+(\omega_{(5)})$.

Let $Z \in B_5^+(\omega_{(5)})$ such that $x_a x_b x_c Z^2 \in (\mathcal{P}_5^+)_{47}$ with $1 \leq a < b < c \leq 5$. Denote by $\bar{V}$ the set of all the monomials as given in Proposition 3.4.36. By a direct computation using Proposition 3.4.15(IV), we see that if $x_a x_b x_c Z^2 \notin \bar{V}$ then there is a monomial X_1 which is given in Lemmas 3.4.37 and 3.4.38 such that $x_a x_b x_c Z^2 = X_1 Z_1^{2^t}$ with suitable monomial $Z_1 \in \mathcal{P}_5$ and $t = \max\{q \in \mathbb{Z} : \omega_q(X_1) > 0\}$. Based on Theorem 2.2.1, we conclude that $x_a x_b x_c Z^2$ is inadmissible. Combining this with the above data, one gets $X \in \bar{V}$. This means $\mathcal{B}_5^+(\bar{\omega}_{(5)}) \subseteq \bar{V}$.

Next, we show that the set $[\bar{V}]_{\bar{\omega}_{(5)}}$ is linearly independent in the space $Q\mathcal{P}_5^+(\bar{\omega}_{(5)})$. Indeed, suppose there is a linear relation $\mathcal{S} = \sum_{481 \leq m \leq 494} \gamma_m \mathcal{Y}_{47,m} \equiv_{\bar{\omega}_{(5)}} 0$, where with $\gamma_m \in \mathbb{Z}/2$, $m = 481, \dots, 494$ and $\mathcal{Y}_{47,m} \in \bar{V}$. By combining Theorem 3.3.5 and Proposition 3.1.1, we explicitly calculate $\pi_{(k;\mathcal{K})}(\mathcal{S})$ in terms of a given minimal set of $\mathcal{A}_2$ -generators in $\mathcal{P}_4$ (modulo $(\mathcal{A}_2^+ \mathcal{P}_4)$). From the relations $\pi_{(k;\mathcal{K})}(\mathcal{S}) \equiv_{\bar{\omega}_{(5)}} 0$ with $\ell(\mathcal{K}) \leq 2$, we get $\gamma_m = 0$ for all m .

To prove $[\bar{V}]_{\bar{\omega}_{(5)}}$ is a basis of $Q\mathcal{P}_5^+(\bar{\omega}_{(5)})$ we need to show that $[\mathcal{Y}_{47,m}]_{\bar{\omega}_{(5)}} \neq [0]$ for all $\mathcal{Y}_{47,m} \in \bar{V}$. Denote by $\bar{V}$ the set of all the monomials as given in Proposition 3.4.36. By a similar argument as given in the proof of Propositions 3.4.1 and 3.4.11, we can prove that the set $[\mathcal{B}_5^+(\bar{\omega}_{(1)}) \cup V \cup \bar{V}]$ is linearly independent in $(Q\mathcal{P}_5^+)_{47}$, where V the set of all the admissible monomials as given in Proposition 3.4.35. This implies $[\mathcal{Y}_{47,m}]_{\bar{\omega}_{(5)}} \neq [0]$ for all $\mathcal{Y}_{47,m}$. The proposition is proved. $\square$

Now, since $\dim(Q\mathcal{P}_5)_{47} = 560$ and $\dim(Q\mathcal{P}_5^0)_{21} = 460$, by Corollaries 3.4.16, 3.4.25 and Propositions 3.4.27, 3.4.35, 3.4.36, we conclude $\dim(Q\mathcal{P}_5)_{47} = 1894$. The proof of Theorem 1.1 is completed.

Final remarks. Since the homomorphism $(\widetilde{Sq_*^0})_{(5,13.2^t)} : (Q\mathcal{P}_5)_{13.2^t-5} \longrightarrow (Q\mathcal{P}_5)_{13.2^{t-1}-5}$ is an epimorphism of $\mathbb{Z}/2(GL_5)$ -modules, $(Q\mathcal{P}_5)_{13.2^t-5} \cong \text{Ker}(\widetilde{Sq_*^0})_{(5,13.2^t-5)} \oplus (Q\mathcal{P}_5)_{13.2^{t-1}-5}$.

According to Lemma 3.4.2, we have

$$\text{Ker}(\widetilde{Sq_*^0})_{(5,13.2^1-5=21)} \cong Q\mathcal{P}_5(3, 3, 1, 1) \bigoplus Q\mathcal{P}_5(3, 3, 3).$$

By a simple computation, we conclude $\mathcal{B}_5^0(21) = \bar{\Phi}(\mathcal{B}_4(21)) = \mathcal{B}_5^0(3, 3, 1, 1) \cup \mathcal{B}_5^0(3, 3, 3)$. Combining this with Propositions 3.4.6 and 3.4.7, we get

$$\bar{\Phi}(\mathcal{B}_4(3, 3, 1, 1)) \subset \mathcal{B}_5(3, 3, 1, 1), \quad \bar{\Phi}(\mathcal{B}_4(3, 3, 3)) \subset \mathcal{B}_5(3, 3, 3).$$

On the other hand, $(Q\mathcal{P}_5)_{13.2^t-5} \cong (Q\mathcal{P}_5)_{13.2^2-5=47}$ for any $t > 2$. By Lemma 3.4.17, $\text{Ker}(\widetilde{Sq_*^0})_{(5,47)} \cong \bigoplus_{1 \leq j \leq 5} Q\mathcal{P}_5(\bar{\omega}_{(j)})$. From the above computations, we obtain $\mathcal{B}_5^0(\bar{\omega}_{(1)}) = \mathcal{B}_5^0(47) = \bar{\Phi}(\mathcal{B}_4(47)) = \bar{\Phi}(\mathcal{B}_4(\bar{\omega}_{(1)}))$. Then, by Corollary 3.4.25, and Propositions 3.4.27, 3.4.35, 3.4.36, one gets

$$\bar{\Phi}(\mathcal{B}_4(\bar{\omega}_{(1)})) \subset \mathcal{B}_5(\bar{\omega}_{(1)}).$$

If ω is a weight vector of degree 47 and $\omega \neq \bar{\omega}_{(1)}$, then $\mathcal{B}_4(\omega) = \emptyset$. Furthermore, if $t > 2$, then $\mathcal{B}_4(13.2^t - 5) = \emptyset$.

From the above remarks, we see that Conjecture 3.3.1 is true in case of five variables and the generic degree $n = 13.2^t - 5$ with t an arbitrary positive integer.

4. An application of Theorem 1.1

The goal of this section is to prove Theorem 1.2. More precisely, by using our results in Sect.3 and a result in [53], we describe the $\mathbb{Z}/2GL_d$ -modules structure of $(Q\mathcal{P}_d)_n$ for $d = 5$ and $n = 8, 21$. Then, we explicitly determine all GL_5 -invariants of these spaces.

Before coming to the proof of the theorem, we introduce some notations and homomorphisms. We note that $(\mathbb{Z}/2)^{\times d}$ regarded as a $\mathbb{Z}/2$ -vector space of dimension d and

$$(\mathbb{Z}/2)^{\times d} \cong \langle x_1, \dots, x_d \rangle \subset \mathcal{P}_d.$$

For $1 \leq t \leq d$, define the $\mathbb{Z}/2$ -linear map $\tau_t : (\mathbb{Z}/2)^{\times d} \rightarrow (\mathbb{Z}/2)^{\times d}$, which is determined by $\tau_t(x_t) = x_{t+1}$, $\tau_t(x_{t+1}) = x_t$, $\tau_t(x_m) = x_m$ for $m \neq t, t+1$, $1 \leq t \leq d-1$ and $\tau_d(x_1) = x_1 + x_2$, $\tau_d(x_m) = x_m$ for $m > 1$.

Denote by S_d the symmetric group of degree d . Then, S_d is generated by τ_t , $1 \leq t \leq d-1$. For each permutation in S_d , consider corresponding permutation matrix; these form a group of matrices isomorphic to S_d . So, the general linear group $GL_d = GL(d, \mathbb{Z}/2) \cong GL((\mathbb{Z}/2)^{\times d})$ is generated by S_d and τ_d . Let $X = x_1^{a_1} x_2^{a_2} \dots x_d^{a_d}$ be a monomial in $\mathcal{P}_d$. Then, the weight vector $\omega(X)$ is invariant under the permutation of the generators x_j , $j = 1, 2, \dots, d$; hence $Q\mathcal{P}_d(\omega)$ also has a S_d -module structure. The homomorphism τ_t induces a homomorphism of $\mathcal{A}_2$ -algebras which is also denoted by $\tau_t : \mathcal{P}_d \rightarrow \mathcal{P}_d$. Hence, a class $[u]_\omega \in Q\mathcal{P}_d(\omega)$ is an GL_d -invariant if and only if $\tau_t(u) + u = 0$ modulo $(\mathcal{A}_2^+ \mathcal{P}_5 + \mathcal{P}_5^-(\omega))$ for $1 \leq t \leq d$. If $\tau_t(u) + u = 0$ modulo $(\mathcal{A}_2^+ \mathcal{P}_5 + \mathcal{P}_5^-(\omega))$ for $1 \leq t \leq d-1$, then $[u]_\omega$ is an S_d -invariant. Note that $\dim((QP_d)_n^{GL_d}) \leq \sum_{\deg(\omega)=n} \dim(QP_d(\omega)^{GL_d})$.

Let ω be a weight vector of degree n and the monomials $\mathcal{Y}_1, \mathcal{Y}_2, \dots, \mathcal{Y}_s \in \mathcal{P}_d(\omega)$, with $s \geq 1$. For a subgroup $L \subseteq GL_d$, we denote

$$\begin{aligned} L(\mathcal{Y}_1, \mathcal{Y}_2, \dots, \mathcal{Y}_s) &= \{\sigma(\mathcal{Y}_j) : \sigma \in L, 1 \leq j \leq s\} \subset \mathcal{P}_d(\omega), \\ [\mathcal{B}(\mathcal{Y}_1, \mathcal{Y}_2, \dots, \mathcal{Y}_s)]_\omega &= [\mathcal{B}_d(\omega)]_\omega \cap \langle [L(\mathcal{Y}_1, \mathcal{Y}_2, \dots, \mathcal{Y}_s)]_\omega \rangle, \\ \theta(\mathcal{Y}_j) &= \sum_{x \in \mathcal{B}_d(n) \cap L(\mathcal{Y}_j)} x, \end{aligned}$$

We note that $\langle [L(\mathcal{Y}_1, \mathcal{Y}_2, \dots, \mathcal{Y}_s)]_\omega \rangle$ is the L -submodule of $Q\mathcal{P}_d(\omega)$ generated by the set

$$\{[\mathcal{Y}_1]_\omega, [\mathcal{Y}_2]_\omega, \dots, [\mathcal{Y}_s]_\omega\}.$$

Now, recall that the squaring operation $(\widetilde{Sq}_*^0)_{(5,21)} : (Q\mathcal{P}_5)_{21} \rightarrow (Q\mathcal{P}_5)_8$ is an epimorphism of GL_5 -modules. So, to prove Theorem 1.2, we need to compute all GL_5 -invariants of $(Q\mathcal{P}_5)_8$.

4.1. Computation of $(Q\mathcal{P}_5)_8^{GL_5}$

According to T  n [53], the $\mathbb{Z}/2$ -vector space $(Q\mathcal{P}_5)_8$ has the basis $[\{\mathcal{Y}_{8,i} : 1 \leq i \leq 174\}]$, where the monomials $\mathcal{Y}_i := \mathcal{Y}_{8,i}$, $1 \leq i \leq 174$, as given in Sect. 6.1.

Proposition 4.1.1. *The space $(Q\mathcal{P}_5)_8^{GL_5}$ is trivial in degree 8.*

We prepare some lemmas for the proof of the proposition. We have

$$(Q\mathcal{P}_5)_8 = Q\mathcal{P}_5^0(\tilde{\omega}_{(1)}) \oplus Q\mathcal{P}_5(\tilde{\omega}_{(2)}) \oplus Q\mathcal{P}_5(\tilde{\omega}_{(3)}),$$

where $\tilde{\omega}_{(1)} = (2, 1, 1)$, $\tilde{\omega}_{(2)} = (2, 3)$, $\tilde{\omega}_{(3)} = (4, 2)$. We see that $\dim Q\mathcal{P}_5^0(\tilde{\omega}_{(1)}) = 105$ with the basis $\bigcup_{1 \leq i \leq 6} [\mathcal{B}(\mathcal{Y}_i)]_{\tilde{\omega}_{(1)}}$, where

$$\begin{aligned} \mathcal{Y}_1 &= x_4 x_5^7, \quad \mathcal{Y}_2 = x_4^3 x_5^5, \quad \mathcal{Y}_3 = x_3 x_4 x_5^6, \\ \mathcal{Y}_4 &= x_3 x_4^2 x_5^5, \quad \mathcal{Y}_5 = x_3 x_4^3 x_5^4, \quad \mathcal{Y}_6 = x_2 x_3 x_4^2 x_5^4. \end{aligned}$$

Observe that $\tilde{\omega}_{(1)}$ is the weight vector of the minimal spike $x_1^7 x_2$, so $[X]_{\tilde{\omega}_{(1)}} = [X]$ for all $X \in \mathcal{P}_5$. A direct computation shows that

$$\begin{aligned} [S_5(\mathcal{Y}_1)] &= \langle [\mathcal{Y}_i] : 1 \leq i \leq 20 \rangle, \\ [S_5(\mathcal{Y}_2)] &= \langle [\mathcal{Y}_i] : 21 \leq i \leq 30 \rangle, \\ [S_5(\mathcal{Y}_3, \mathcal{Y}_4, \mathcal{Y}_5)] &= \langle [\mathcal{Y}_i] : 31 \leq i \leq 90 \rangle, \\ [S_5(\mathcal{Y}_6)] &= \langle [\mathcal{Y}_i] : 91 \leq i \leq 105 \rangle \end{aligned}$$

are S_5 -submodules of $Q\mathcal{P}_5^0(\tilde{\omega}_{(1)})$. Hence, we have a direct summand decomposition of S_5 -modules: $QP_5^0(\tilde{\omega}_{(1)}) = [S_5(\mathcal{Y}_1)] \oplus [S_5(\mathcal{Y}_2)] \oplus [S_5(\mathcal{Y}_3, \mathcal{Y}_4, \mathcal{Y}_5)] \oplus [S_5(\mathcal{Y}_6)]$.

Lemma 4.1.2. $Q\mathcal{P}_5^0(\tilde{\omega}_{(1)})^{S_5}$ has dimension 4.

Proof. We prove the following:

$$\begin{aligned} [S_5(\mathcal{Y}_j)]^{S_5} &= \langle [\theta(\mathcal{Y}_j)] \rangle, \quad j = 1, 2, 6 \\ [S_5(\mathcal{Y}_3, \mathcal{Y}_4, \mathcal{Y}_5)]^{S_5} &= \langle [q := \mathcal{Y}_{31} + \mathcal{Y}_{32} + \cdots + \mathcal{Y}_{70}] \rangle. \end{aligned}$$

Indeed, we compute $[S_5(\mathcal{Y}_j)]^{S_5}$ for $j = 2$ and $[S_5(\mathcal{Y}_3, \mathcal{Y}_4, \mathcal{Y}_5)]^{S_5}$. The others can be proved by a similar technique. Note that $\dim[S_5(\mathcal{Y}_2)] = 10$ with a basis consisting of all the classes represented by the monomials $\mathcal{Y}_i : 21 \leq i \leq 30$. Suppose that $a = \sum_{21 \leq i \leq 30} \gamma_i \mathcal{Y}_i$ with $\gamma_i \in \mathbb{Z}/2$ and $[a] \in [S_5(\mathcal{Y}_2)]^{S_5}$. By a direct computation using Theorem 3.3.5, we have

$$\begin{aligned} \tau_1(a) + a &= \sum X \text{ modulo } (\mathcal{A}_2^+ \mathcal{P}_5) = 0 \text{ modulo } (\mathcal{A}_2^+ \mathcal{P}_5), \\ \tau_2(a) + a &= \sum Y \text{ modulo } (\mathcal{A}_2^+ \mathcal{P}_5) = 0 \text{ modulo } (\mathcal{A}_2^+ \mathcal{P}_5), \\ \tau_3(a) + a &= \sum Z \text{ modulo } (\mathcal{A}_2^+ \mathcal{P}_5) = 0 \text{ modulo } (\mathcal{A}_2^+ \mathcal{P}_5), \\ \tau_4(a) + a &= \sum W \text{ modulo } (\mathcal{A}_2^+ \mathcal{P}_5) = 0 \text{ modulo } (\mathcal{A}_2^+ \mathcal{P}_5), \end{aligned}$$

where

$$\begin{aligned} \sum X &= (\gamma_{24} + \gamma_{27})(\mathcal{Y}_{24} + \mathcal{Y}_{27}) + (\gamma_{25} + \gamma_{28})(\mathcal{Y}_{25} + \mathcal{Y}_{28}) + (\gamma_{26} + \gamma_{29})(\mathcal{Y}_{26} + \mathcal{Y}_{29}), \\ \sum Y &= (\gamma_{22} + \gamma_{24})(\mathcal{Y}_{22} + \mathcal{Y}_{24}) + (\gamma_{23} + \gamma_{25})(\mathcal{Y}_{23} + \mathcal{Y}_{25}) + (\gamma_{29} + \gamma_{30})(\mathcal{Y}_{29} + \mathcal{Y}_{30}), \\ \sum Z &= (\gamma_{21} + \gamma_{22})(\mathcal{Y}_{21} + \mathcal{Y}_{22}) + (\gamma_{25} + \gamma_{26})(\mathcal{Y}_{25} + \mathcal{Y}_{26}) + (\gamma_{28} + \gamma_{29})(\mathcal{Y}_{28} + \mathcal{Y}_{29}), \\ \sum W &= (\gamma_{22} + \gamma_{23})(\mathcal{Y}_{22} + \mathcal{Y}_{23}) + (\gamma_{24} + \gamma_{25})(\mathcal{Y}_{24} + \mathcal{Y}_{25}) + (\gamma_{27} + \gamma_{28})(\mathcal{Y}_{27} + \mathcal{Y}_{28}). \end{aligned}$$

These relations imply that $\gamma_i = \gamma_{21}$ for $i = 22, \dots, 30$. Hence, we get $a = \theta(\mathcal{Y}_2) \text{ modulo } (\mathcal{A}_2^+ \mathcal{P}_5)$ with $\theta(\mathcal{Y}_2) = \sum_{21 \leq j \leq 30} \mathcal{Y}_j$.

Now, we have the set $\{[\mathcal{Y}_i] : 31 \leq i \leq 90\}$ is a basis of $[S_5(\mathcal{Y}_3, \mathcal{Y}_4, \mathcal{Y}_5)]$. Suppose that $b = \sum_{31 \leq i \leq 90} \gamma_i \mathcal{Y}_i$ with $\gamma_i \in \mathbb{Z}/2$ and $[b] \in [S_5(\mathcal{Y}_3, \mathcal{Y}_4, \mathcal{Y}_5)]^{S_5}$. By a similar computation from the relations $\tau_t(b) + b = 0 \text{ modulo } (\mathcal{A}_2^+ \mathcal{P}_5)$, $t = 1, 2, 3, 4$, one gets $\gamma_i = 0$ for $71 \leq i \leq 90$ and $\gamma_i = \gamma_{31}$, $i = 32, 33, \dots, 70$. This means $b = q \text{ modulo } (\mathcal{A}_2^+ \mathcal{P}_5)$ with $q = \sum_{31 \leq i \leq 70} \mathcal{Y}_i$. The lemma is proved. $\square$

Lemma 4.1.3. The subspace $(Q\mathcal{P}_5(\tilde{\omega}_{(3)}))^{GL_5}$ is trivial.

Proof. Using a result in [53], we see that $Q\mathcal{P}_5(\tilde{\omega}_{(3)})$ is the $\mathbb{Z}/2$ -vector space of dimension 45 with the basis $[\mathcal{B}(\mathcal{Y}_{130} = x_2 x_3 x_4^3 x_5^3)]_{\tilde{\omega}_{(3)}} \cup [\mathcal{B}(\mathcal{Y}_{160} = x_1 x_2 x_3 x_4^2 x_5^3)]_{\tilde{\omega}_{(3)}}$. Furthermore, $[S_5(\mathcal{Y}_{130})]$ and $[S_5(\mathcal{Y}_{160})]$ are S_5 -submodules of $Q\mathcal{P}_5^0(\tilde{\omega}_{(3)})$, where $[S_5(\mathcal{Y}_{130})]_{\tilde{\omega}_{(3)}} = \langle [\mathcal{Y}_i : 130 \leq i \leq 159]_{\tilde{\omega}_{(3)}} \rangle$ and $[S_5(\mathcal{Y}_{160})]_{\tilde{\omega}_{(3)}} = \langle [\mathcal{Y}_i : 160 \leq i \leq 174]_{\tilde{\omega}_{(3)}} \rangle$. So, we have a direct summand decomposition of S_5 -modules: $Q\mathcal{P}_5(\tilde{\omega}_{(3)}) = [S_5(\mathcal{Y}_{130})]_{\tilde{\omega}_{(3)}} \oplus [S_5(\mathcal{Y}_{160})]_{\tilde{\omega}_{(3)}}$. The set $[B(\mathcal{Y}_{130})]_{\tilde{\omega}_{(3)}}$ is a basis of $[S_5(\mathcal{Y}_{130})]_{\tilde{\omega}_{(3)}}$. The action of S_5 on $Q\mathcal{P}_5$ induces the one of it on $[B(\mathcal{Y}_{130})]_{\tilde{\omega}_{(3)}}$. On the other hand, this action is transitive, hence if $a = \sum_{130 \leq i \leq 159} \gamma_i \mathcal{Y}_i$ with $\gamma_i \in \mathbb{Z}/2$ and $[a] \in [S_5(\mathcal{Y}_{130})]^{S_5}$, then from the relations $\tau_t(a) + a = 0 \text{ modulo } (\mathcal{A}_2^+ \mathcal{P}_5 + \mathcal{P}_5^-(\tilde{\omega}_{(3)}))$, $1 \leq t \leq 4$, we get $\gamma_i = \gamma_{130}$, $\forall i$, $131 \leq i \leq 159$. In other words, $[S_5(\mathcal{Y}_j)]_{\tilde{\omega}_{(3)}}^{S_5} = \langle [\theta(\mathcal{Y}_{130})]_{\tilde{\omega}_{(3)}} \rangle$ with $\theta(\mathcal{Y}_{130}) = \sum_{130 \leq i \leq 159} \mathcal{Y}_i$.

Next, we have $\dim[S_5(\mathcal{Y}_{160})]_{\tilde{\omega}_{(3)}} = 15$ with the basis $[B(\mathcal{Y}_{160})]_{\tilde{\omega}_{(3)}}$. Suppose $b = \sum_{160 \leq i \leq 174} \gamma_i \mathcal{Y}_i$

with $\gamma_j \in \mathbb{Z}/2$ and $[b] \in [S_5(\mathcal{Y}_{160})]^{S_5}$. A direct computation shows:

$$\begin{aligned}
\tau_1(b) + b &\equiv_{\tilde{\omega}_{(3)}} \gamma_{169}(\mathcal{Y}_{160} + \mathcal{Y}_{162}) + \gamma_{170}(\mathcal{Y}_{161} + \mathcal{Y}_{163}) + \gamma_{171}(\mathcal{Y}_{164} + \mathcal{Y}_{165}) \\
&\quad + (\gamma_{166} + \gamma_{172})(\mathcal{Y}_{166} + \mathcal{Y}_{172}) + (\gamma_{167} + \gamma_{173})(\mathcal{Y}_{167} + \mathcal{Y}_{173}) \\
&\quad + (\gamma_{168} + \gamma_{174})(\mathcal{Y}_{168} + \mathcal{Y}_{174}) \\
\tau_2(b) + b &\equiv_{\tilde{\omega}_{(3)}} (\gamma_{162} + \gamma_{169})(\mathcal{Y}_{162} + \mathcal{Y}_{169}) + (\gamma_{163} + \gamma_{170})(\mathcal{Y}_{163} + \mathcal{Y}_{170}) \\
&\quad + (\gamma_{164} + \gamma_{166})(\mathcal{Y}_{164} + \mathcal{Y}_{166}) + (\gamma_{165} + \gamma_{167})(\mathcal{Y}_{165} + \mathcal{Y}_{167}) \\
&\quad + (\gamma_{168} + \gamma_{171})(\mathcal{Y}_{168} + \mathcal{Y}_{171}) + (\gamma_{165} + \gamma_{167})(\mathcal{Y}_{165} + \mathcal{Y}_{167}) \\
&\quad + \gamma_{174}(\mathcal{Y}_{172} + \mathcal{Y}_{173}), \\
\tau_3(b) + b &\equiv_{\tilde{\omega}_{(3)}} (\gamma_{160} + \gamma_{162})(\mathcal{Y}_{160} + \mathcal{Y}_{162}) + (\gamma_{161} + \gamma_{164})(\mathcal{Y}_{161} + \mathcal{Y}_{164}) \\
&\quad + (\gamma_{163} + \gamma_{165})(\mathcal{Y}_{163} + \mathcal{Y}_{165}) + (\gamma_{167} + \gamma_{168})(\mathcal{Y}_{167} + \mathcal{Y}_{168}) \\
&\quad + (\gamma_{170} + \gamma_{171})(\mathcal{Y}_{170} + \mathcal{Y}_{171}) + (\gamma_{173} + \gamma_{174})(\mathcal{Y}_{173} + \mathcal{Y}_{174}), \\
\tau_4(b) + b &\equiv_{\tilde{\omega}_{(3)}} (\gamma_{160} + \gamma_{161})(\mathcal{Y}_{160} + \mathcal{Y}_{161}) + (\gamma_{162} + \gamma_{163})(\mathcal{Y}_{162} + \mathcal{Y}_{163}) \\
&\quad + (\gamma_{164} + \gamma_{165})(\mathcal{Y}_{164} + \mathcal{Y}_{165}) + (\gamma_{166} + \gamma_{167})(\mathcal{Y}_{166} + \mathcal{Y}_{167}) \\
&\quad + (\gamma_{169} + \gamma_{170})(\mathcal{Y}_{169} + \mathcal{Y}_{170}) + (\gamma_{172} + \gamma_{173})(\mathcal{Y}_{172} + \mathcal{Y}_{173}).
\end{aligned}$$

Then, from the relations $\tau_t(b) + b = 0 \pmod{(\mathcal{A}_2^+ \mathcal{P}_5 + \mathcal{P}_5^-(\tilde{\omega}_{(3)}))}$, we obtain $\gamma_j = 0, \forall j$.

Now, let $[X]_{\tilde{\omega}_{(3)}} \in (Q\mathcal{P}_5(\tilde{\omega}_{(3)}))^{GL_5}$ with $X \in \mathcal{P}_5(\tilde{\omega}_{(3)})$, then $[X]_{\tilde{\omega}_{(3)}} \in (Q\mathcal{P}_5(\tilde{\omega}_{(3)}))^{S_5}$. So, we have $X = \gamma\theta(\mathcal{Y}_{130}) \pmod{(\mathcal{A}_2^+ \mathcal{P}_5 + \mathcal{P}_5^-(\tilde{\omega}_{(3)}))}$ with $\gamma \in \mathbb{Z}/2$. By a direct computation, we have

$$\tau_5(X) + X = \gamma\mathcal{Y}_{130} + \text{other terms} \pmod{(\mathcal{A}_2^+ \mathcal{P}_5 + \mathcal{P}_5^-(\tilde{\omega}_{(3)}))} = 0 \pmod{(\mathcal{A}_2^+ \mathcal{P}_5 + \mathcal{P}_5^-(\tilde{\omega}_{(3)}))}.$$

This implies $\gamma = 0$. The proposition follows. $\square$

By a simple computation using the techniques as given in the proof of Lemmas 4.1.2 and 4.1.3, we get the following.

Lemma 4.1.4.

i) We have a direct summand decomposition of the S_5 -modules:

$$Q\mathcal{P}_5(\tilde{\omega}_{(2)}) = [S_5(\mathcal{Y}_{106})]_{\tilde{\omega}_{(2)}} \bigoplus [S_5(\mathcal{Y}_{126})]_{\tilde{\omega}_{(2)}}.$$

ii) The subspace $(Q\mathcal{P}_5(\tilde{\omega}_{(2)}))^{GL_5}$ is trivial.

Proof of Proposition 4.1.1. Let $[X] \in (Q\mathcal{P}_5)^{GL_5}$. Then, from Lemmas 4.1.2 - 4.1.4, we have

$$X = \beta_1\theta(\mathcal{Y}_1) + \beta_2\theta(\mathcal{Y}_2) + \beta_3q + \beta_4\theta(\mathcal{Y}_3) \pmod{(\mathcal{A}_2^+ \mathcal{P}_5)},$$

with $\beta_t \in \mathbb{Z}/2, 1 \leq t \leq 4$. By using Theorem 3.3.5 and computing $\tau_5(X) + X$ in terms of the admissible monomials $(\pmod{(\mathcal{A}_2^+ \mathcal{P}_5)})$, we conclude

$$\tau_5(X) + X \equiv \beta_1\mathcal{Y}_7 + (\beta_1 + \beta_2)\mathcal{Y}_{16} + \beta_3\mathcal{Y}_{33} + \beta_4(\mathcal{Y}_{91} + \mathcal{Y}_{92} + \mathcal{Y}_{93}) + \text{other terms} \equiv 0.$$

This relation shows $\beta_1 = \beta_2 = \beta_3 = \beta_4 = 0$. The proposition is proved. $\square$

4.2. Computation of $(\text{Ker}(\widetilde{Sq}_*^0)_{(5,21)})^{S_5}$

From the results in Sect. 3.4.1, we see that $\dim \text{Ker}(\widetilde{Sq}_*^0)_{(5,21)} = 666$ with the basis $\{\mathcal{Y}_{21,t} : 1 \leq t \leq 666\}$. Here, the admissible monomials $\mathcal{Y}_t := \mathcal{Y}_{21,t}, 1 \leq t \leq 666$, as given in Sect. 6.6 and Sect. 6.7. Recall that $(\widetilde{Sq}_*^0)_{(5,21)}$ is an epimorphism of GL_5 -modules. Combining this and the results in Subsection 4.1, we get $(Q\mathcal{P}_5)_{21}^{GL_5} \subseteq (\text{Ker}(\widetilde{Sq}_*^0)_{(5,21)})^{GL_5}$. By Lemma 3.4.2, we have a direct summand decomposition of the S_5 -modules:

$$\text{Ker}(\widetilde{Sq}_*^0)_{(5,21)} = Q\mathcal{P}_5(3, 3, 1, 1) \bigoplus Q\mathcal{P}_5(3, 3, 3).$$

For $\omega = (3, 3, 1, 1)$, according to the results in Sect. 3.4.1, we get $Q\mathcal{P}_5(\omega) = Q\mathcal{P}_5^0(\omega) \oplus Q\mathcal{P}_5^+(\omega)$ with $\dim Q\mathcal{P}_5^0(\omega) = 340$ and $\dim Q\mathcal{P}_5^+(\omega) = 196$. Note that $\mathcal{Z} = x_1^{15}x_2^3x_3^3$ is the minimal spike monomial in $(\mathcal{P}_5)_{21}$ and $\omega(\mathcal{Z}) = \omega$. So, $[x]_\omega = [x]$ for any $x \in (\mathcal{P}_5)_{21}$. By using the results in Sect. 3.4.1, we see that there is a direct summand decomposition of the S_5 -modules:

$$Q\mathcal{P}_5^0(\omega) = \langle [S_5(\mathcal{Y}_1)] \rangle \oplus \langle [S_5(\mathcal{Y}_{31})] \rangle \oplus \langle [S_5(\mathcal{Y}_{61})] \rangle \oplus \langle [S_5(\mathcal{Y}_{121})] \rangle \oplus \langle \mathbb{V}_1 \rangle,$$

where

$$\begin{aligned} \mathcal{B}(\mathcal{Y}_1) &= \{\mathcal{Y}_t : 1 \leq t \leq 30\}, & \mathcal{B}(\mathcal{Y}_{31}) &= \{\mathcal{Y}_t : 31 \leq t \leq 60\}, \\ \mathcal{B}(\mathcal{Y}_{61}) &= \{\mathcal{Y}_t : 61 \leq t \leq 120\}, & \mathcal{B}(\mathcal{Y}_{121}) &= \{\mathcal{Y}_t : 121 \leq t \leq 150\}, \\ \mathbb{V}_1 &= \mathcal{B}(\mathcal{Y}_{151}, \mathcal{Y}_{181}, \mathcal{Y}_{201}, \mathcal{Y}_{241}, \mathcal{Y}_{256}, \mathcal{Y}_{266}, \mathcal{Y}_{286}, \mathcal{Y}_{296}, \mathcal{Y}_{301}, \mathcal{Y}_{316}) \\ &= \{\mathcal{Y}_t : 151 \leq t \leq 340\}, \end{aligned}$$

Lemma 4.2.1.

a) $\langle [S_5(\mathcal{Y}_j)] \rangle^{S_5} = \langle [\theta(\mathcal{Y}_j)] \rangle$ for $j = 1, 31, 61, 121$.

b) The subspace $\langle [S_5(\mathbb{V}_1)] \rangle^{S_5}$ is trivial.

Outline of the proof. For $j = 1$, let $[f_1] \in \langle [S_5(\mathcal{Y}_1)] \rangle^{S_5}$. Then, we have

$$\tau_m(f_1) = \sum_{X \in \mathcal{B}(\mathcal{Y}_1)} \beta_X \cdot X \text{ modulo } (\mathcal{A}_2^+ \mathcal{P}_5), \quad 0 < m < 5$$

with $f_1 = \sum_{X \in \mathcal{B}(\mathcal{Y}_1)} \beta_X \cdot X$ and $\beta_X \in \mathbb{Z}/2$. By a direct computation, we can see that the action of the symmetric group S_5 on $Q\mathcal{P}_5$ induces the one of it on the set $[\mathcal{B}(\mathcal{Y}_j)]$ and this action is transitive. So, we get $\beta_X = \beta_{X'} = \beta \in \mathbb{Z}/2$ for all $X, X' \in \mathcal{B}(\mathcal{Y}_j)$. This means $f_1 = \theta(\mathcal{Y}_1) \text{ modulo } (\mathcal{A}_2^+ \mathcal{P}_5)$. For $j = 31, 61, 121$, we determine $\tau_m(f_j) + f_j$ in terms of $\mathcal{Y}_j$. Then, by a simple computation using the relations $\tau_m(f_j) = f_j \text{ modulo } (\mathcal{A}_2^+ \mathcal{P}_5)$, we conclude $f_j = \theta(\mathcal{Y}_j) \text{ modulo } (\mathcal{A}_2^+ \mathcal{P}_5)$.

Next, we have $\dim \langle \mathbb{V}_1 \rangle = 190$ with the basis $\{[\mathcal{Y}_t] : 151 \leq t \leq 340\}$. Assume that $g = \sum_{u \in \mathbb{V}_1} \gamma_u \cdot u$ with $\gamma_u \in \mathbb{Z}/2$ and $[g] \in \langle [\mathbb{V}_1] \rangle^{S_5}$. By using Theorem 3.3.5 and a similar computation as given in the proof of Lemmas 4.1.2 and 4.1.3, we obtain $\gamma_u = 0$ for all $u \in \mathbb{V}_1$. This implies that g is $\mathcal{A}_2$ -decomposable. The lemma follows. $\square$

Lemma 4.2.2. $Q\mathcal{P}_5^+(\omega)^{S_5} = \langle [p := \mathcal{Y}_{411} + \mathcal{Y}_{412} + \cdots + \mathcal{Y}_{419}] \rangle$.

Proof. From Proposition 3.4.6, we see that the sets $[S_5(\mathcal{Y}_{401})] = \langle [\mathcal{Y}_t] : 401 \leq t \leq 410 \rangle$ and $\mathbb{V}_2 = \langle [\mathcal{Y}_t] : 411 \leq t \leq 596 \rangle$, are S_5 -submodules of $Q\mathcal{P}_5^+(\omega)$. Hence, we have a direct summand decomposition of the S_5 -modules:

$$Q\mathcal{P}_5^+(\omega) = [S_5(\mathcal{Y}_{401})] \oplus \mathbb{V}_2.$$

The set $[\mathcal{B}(\mathcal{Y}_{401})]$ is a basis of $[S_5(\mathcal{Y}_{401})]$. Assume that Z is a polynomial such that $[Z] \in [S_5(\mathcal{Y}_{401})]^{S_5}$ and $Z = \sum_{401 \leq t \leq 410} \ell_t \mathcal{Y}_t$ with $\ell_t \in \mathbb{Z}/2$. For $1 \leq j \leq 4$, we explicitly compute $\tau_j(Z) + Z$ in the terms of the admissible monomials $\mathcal{Y}_t$, $401 \leq t \leq 410$. By a direct computation using Theorem 3.3.5, we get

$$\begin{aligned} \tau_1(Z) + Z &\equiv \ell_{401} \mathcal{Y}_{404} + \ell_{402} \mathcal{Y}_{405} + \ell_{403} \mathcal{Y}_{407} + (\ell_{406} + \ell_{409})(\mathcal{Y}_{406} + \mathcal{Y}_{409}) \\ &\quad + (\ell_{408} + \ell_{410})(\mathcal{Y}_{408} + \mathcal{Y}_{410}), \\ \tau_2(Z) + Z &\equiv (\ell_{401} + \ell_{404})(\mathcal{Y}_{401} + \mathcal{Y}_{404}) + (\ell_{402} + \ell_{405})(\mathcal{Y}_{402} + \mathcal{Y}_{405}) \\ &\quad + (\ell_{403} + \ell_{406})(\mathcal{Y}_{403} + \mathcal{Y}_{406}) + (\ell_{407} + \ell_{408})(\mathcal{Y}_{407} + \mathcal{Y}_{408}) + \ell_{409} \mathcal{Y}_{410}, \\ \tau_3(Z) + Z &\equiv \ell_{401} \mathcal{Y}_{404} + (\ell_{402} + \ell_{403})(\mathcal{Y}_{402} + \mathcal{Y}_{403}) + (\ell_{405} + \ell_{407})(\mathcal{Y}_{405} + \mathcal{Y}_{407}), \\ &\quad + (\ell_{406} + \ell_{408})(\mathcal{Y}_{406} + \mathcal{Y}_{408}) + (\ell_{409} + \ell_{410})(\mathcal{Y}_{409} + \mathcal{Y}_{410}), \\ \tau_4(Z) + Z &\equiv (\ell_{401} + \ell_{402})(\mathcal{Y}_{401} + \mathcal{Y}_{402}) + \ell_{403} \mathcal{Y}_{407} + (\ell_{404} + \ell_{405})(\mathcal{Y}_{404} + \mathcal{Y}_{405}) \\ &\quad + \ell_{406} \mathcal{Y}_{408} + \ell_{409} \mathcal{Y}_{410}. \end{aligned}$$

Then, by the relations $\tau_j(Z) = Z \text{ modulo } (\mathcal{A}_2^+ \mathcal{P}_5)$, $j = 1, 2, 3, 4$, one gets $\ell_t = 0$, $\forall t$. This implies that Z is $\mathcal{A}_2$ -decomposable.

Note that $\dim \mathbb{V}_2 = 186$ with the basis $\{[\mathcal{Y}_j] : 411 \leq j \leq 596\}$. Then if the polynomial $g = \sum_{u \in \mathbb{V}_2} \sigma_u \cdot u$, $\sigma_u \in \mathbb{Z}/2$ such that $[g] \in \mathbb{V}_2^{S_5}$, then by a similar argument as given above, we obtain $g = p \text{ modulo } (\mathcal{A}_2^+ \mathcal{P}_5)$. This completes the proof of the lemma. $\square$

Proposition 4.2.3. *For $\bar{\omega} = (3, 3, 3)$, we have $Q\mathcal{P}_5(\bar{\omega})^{GL_5} = \langle [q_2]_{\bar{\omega}} \rangle$, where*

$$\begin{aligned} q_2 &= \mathcal{Y}_{652} + \mathcal{Y}_{653} + \mathcal{Y}_{654} + \mathcal{Y}_{656} + \mathcal{Y}_{657} + \mathcal{Y}_{658} + \sum_{661 \leq t \leq 666} \mathcal{Y}_t \\ &= x_1 x_2^6 x_3^3 x_4^5 x_5^6 + x_1 x_2^3 x_3^6 x_4^5 x_5^6 + x_1 x_2^3 x_3^5 x_4^6 x_5^6 + x_1^3 x_2 x_3^5 x_4^6 x_5^6 \\ &\quad + x_1^3 x_2^5 x_3 x_4^6 x_5^6 + x_1^3 x_2^5 x_3^6 x_4 x_5^6 + x_1^3 x_2^3 x_3^5 x_4^4 x_5^6 + x_1^3 x_2^3 x_3^4 x_4^5 x_5^6 \\ &\quad + x_1^3 x_2^3 x_3^5 x_4^6 x_5^4 + x_1^3 x_2^4 x_3^3 x_4^5 x_5^6 + x_1^3 x_2^5 x_3^3 x_4^6 x_5^4 + x_1^3 x_2^5 x_3^6 x_4^3 x_5^4. \end{aligned}$$

Based on the results in Sect. 3.4, we have

$$\dim Q\mathcal{P}_5(\bar{\omega}) = \dim Q\mathcal{P}_5^0(\bar{\omega}) + \dim Q\mathcal{P}_5^+(\bar{\omega}) = 60 + 70 = 130.$$

Consider the following monomials:

$$\begin{aligned} \mathcal{Y}_{341} &= x_3^7 x_4^7 x_5^7, & \mathcal{Y}_{351} &= x_2 x_3^6 x_4^7 x_5^7, & \mathcal{Y}_{381} &= x_2^3 x_3^5 x_4^6 x_5^7, \\ \mathcal{Y}_{597} &= x_1 x_2^2 x_3^4 x_4^7 x_5^7, & \mathcal{Y}_{607} &= x_1 x_2^6 x_3 x_4^6 x_5^7, & \mathcal{Y}_{617} &= x_1 x_2^2 x_3^5 x_4^6 x_5^7, \\ \mathcal{Y}_{622} &= x_1 x_2^3 x_3^4 x_4^6 x_5^7, & \mathcal{Y}_{642} &= x_1^3 x_2^5 x_3^2 x_4^4 x_5^7, & \mathcal{Y}_{647} &= x_1^3 x_2^3 x_3^4 x_4^4 x_5^7, \\ \mathcal{Y}_{652} &= x_1 x_2^6 x_3^3 x_4^5 x_5^6, & \mathcal{Y}_{659} &= x_1^3 x_2^5 x_3^2 x_4^5 x_5^6, & \mathcal{Y}_{660} &= x_1^3 x_2^5 x_3^3 x_4^4 x_5^6. \end{aligned}$$

The following lemma can be easily proved by a direct computation.

Lemma 4.2.4.

i) *The following subspaces are S_5 -submodules of $Q\mathcal{P}_5(\bar{\omega})$:*

$$\begin{aligned} \langle [S_5(\mathcal{Y}_a)]_{\bar{\omega}} \rangle, \quad a &= 341, 351, 381, 597, \quad \mathbb{V}_3 := \langle [S_5(\mathcal{Y}_{607}, \mathcal{Y}_{617}, \mathcal{Y}_{622}, \mathcal{Y}_{642}, \mathcal{Y}_{647})]_{\bar{\omega}} \rangle, \\ \mathbb{V}_4 &:= \langle [S_5(\mathcal{Y}_{652}, \mathcal{Y}_{659}, \mathcal{Y}_{660})]_{\bar{\omega}} \rangle. \end{aligned}$$

ii) *We have a direct summand decomposition of the S_5 -modules:*

$$\begin{aligned} Q\mathcal{P}_5(\bar{\omega}) &= \langle [S_5(\mathcal{Y}_{341})]_{\bar{\omega}} \rangle \oplus \langle [S_5(\mathcal{Y}_{351})]_{\bar{\omega}} \rangle \oplus \langle [S_5(\mathcal{Y}_{381})]_{\bar{\omega}} \rangle \\ &\quad \oplus \langle [S_5(\mathcal{Y}_{597})]_{\bar{\omega}} \rangle \oplus \mathbb{V}_3 \oplus \mathbb{V}_4. \end{aligned}$$

Lemma 4.2.5.

- i) $\langle [S_5(\mathcal{Y}_a)]_{\bar{\omega}} \rangle^{S_5} = \langle [\theta(\mathcal{Y}_a)]_{\bar{\omega}} \rangle$, for $a = 341, 351, 381, 597$.
- ii) $\langle [S_5(\mathbb{V}_3)]_{\bar{\omega}} \rangle^{S_5} = \langle [q_1 := \mathcal{Y}_{610} + \mathcal{Y}_{611} + \mathcal{Y}_{613} + \mathcal{Y}_{614} + \mathcal{Y}_{616} + \sum_{647 \leq t \leq 651} \mathcal{Y}_t]_{\bar{\omega}} \rangle$.
- iii) $\langle [S_5(\mathbb{V}_4)]_{\bar{\omega}} \rangle^{S_5} = \langle [q_2 := \mathcal{Y}_{652} + \mathcal{Y}_{653} + \mathcal{Y}_{654} + \mathcal{Y}_{656} + \mathcal{Y}_{657} + \mathcal{Y}_{658} + \sum_{661 \leq t \leq 666} \mathcal{Y}_t]_{\bar{\omega}} \rangle$.

The proof of the lemma is straightforward.

Proof of Proposition 4.2.3. By Lemmas 4.2.4 and 4.2.5, we get

$$Q\mathcal{P}_5(\bar{\omega})^{S_5} = \langle [\theta(\mathcal{Y}_{341})]_{\bar{\omega}}, [\theta(\mathcal{Y}_{351})]_{\bar{\omega}}, [\theta(\mathcal{Y}_{381})]_{\bar{\omega}}, [\theta(\mathcal{Y}_{597})]_{\bar{\omega}}, [q_1]_{\bar{\omega}}, [q_2]_{\bar{\omega}} \rangle.$$

Let X be a polynomial in $\mathcal{P}_5(\bar{\omega})$ such that $[X]_{\bar{\omega}} \in Q\mathcal{P}_5(\bar{\omega})^{GL_5}$. Then, we have

$$X = v_1 \theta(\mathcal{Y}_{341}) + v_2 \theta(\mathcal{Y}_{351}) + v_3 \theta(\mathcal{Y}_{381}) + v_4 \theta(\mathcal{Y}_{597}) + v_5 q_1 + v_6 q_2 \text{ modulo } (\mathcal{A}_2^+ \mathcal{P}_5 + \mathcal{P}_5^-(\bar{\omega})),$$

with $v_j \in \mathbb{Z}/2$ for $1 \leq j \leq 6$. We explicitly compute $\tau_5(X)$ in terms of the admissible monomials $\mathcal{Y}_t$ with $t = 341, 342, \dots, 400, 597, 598, \dots, 666$. By a direct computation, one gets

$$\begin{aligned} \tau_5(X) + X &= (v_1 + v_2) \mathcal{Y}_{342} + (v_2 + v_4) \mathcal{Y}_{351} + v_2 \mathcal{Y}_{354} + v_3 \mathcal{Y}_{383} \\ &\quad + v_5 \mathcal{Y}_{650} + \text{other terms modulo } (\mathcal{A}_2^+ \mathcal{P}_5 + \mathcal{P}_5^-(\bar{\omega})). \end{aligned}$$

Since $[X]_{\bar{\omega}} \in Q\mathcal{P}_5(\bar{\omega})^{GL_5}$, $v_j = 0$, $1 \leq j \leq 5$. The proposition is proved. $\square$

Combining Lemmas 4.2.1, 4.2.2 and 4.2.5, one gets the following.

Corollary 4.2.6. $\text{Ker}(\widetilde{Sq_*^0})_{(5,21)}^{S_5}$ has dimension 11.

4.3. Proof of Theorem 1.2

Suppose that $[T] \in (Q\mathcal{P}_5)_{21}^{GL_5}$ with T is a polynomial in $(\mathcal{P}_5)_{21}$. From Proposition 4.2.3, we have $T = T^* + \zeta_6 q_2$ modulo $(\mathcal{A}_2^+ \mathcal{P}_5)$ with $T^* \in \mathcal{P}_5^-(\overline{\omega})$ and $\zeta_6 \in \mathbb{Z}/2$. By a simple computation, we see that $[q_2] \in (Q\mathcal{P}_5)_{21}^{S_5}$. This implies that $[T^*]$ is an S_5 -invariant. On the other hand, $[\mathcal{P}_5^-(\overline{\omega})] = Q\mathcal{P}_5(\omega)$. Hence, by Lemmas 4.2.1 and 4.2.2, we obtain

$$T^* = \zeta_1 \theta(\mathcal{Y}_1) + \zeta_2 \theta(\mathcal{Y}_{31}) + \zeta_3 \theta(\mathcal{Y}_{61}) + \zeta_4 \theta(\mathcal{Y}_{121}) + \zeta_5 p \text{ modulo } (\mathcal{A}_2^+ \mathcal{P}_5),$$

where $\zeta_i \in \mathbb{Z}/2$. Using Theorem 1.1 and computing $\tau_5(T) + T$ in terms of the admissible monomials $\mathcal{Y}_t$, $1 \leq t \leq 666$, we conclude

$$\tau_5(T) + T \equiv (\zeta_1 + \zeta_3) \mathcal{Y}_4 + (\zeta_2 + \zeta_4) \mathcal{Y}_{35} + \zeta_3 \mathcal{Y}_{61} + (\zeta_3 + \zeta_4) \mathcal{Y}_{107} + \zeta_5 \mathcal{Y}_{153} + \text{other terms}.$$

By the relation $\tau_5(T) = T$ modulo $(\mathcal{A}_2^+ \mathcal{P}_5)$, one gets $\zeta_i = 0$, $1 \leq i \leq 5$. This shows $T = \zeta_6 q_2$ modulo $(\mathcal{A}_2^+ \mathcal{P}_5)$. The proof of the theorem is completed.

5. Proof of Theorem 1.3

We see that $\lambda_3 \in \Lambda^{1,3}$, and $\overline{f}_0 = \lambda_3 \lambda_5 \lambda_6 \lambda_4 + \lambda_3^2 \lambda_7 \lambda_5 + \lambda_7 \lambda_5 \lambda_2 \lambda_3^2 + \lambda_7 \lambda_5 \lambda_4 \lambda_2 \in \Lambda^{4,18}$ are the cycles in the lambda algebra Λ . By Lin [17], we have $\text{Ext}_{\mathcal{A}_2}^{5,26}(\mathbb{Z}/2, \mathbb{Z}/2) = \langle h_2 f_0 \rangle$, with $h_2 = [\lambda_3] \in \text{Ext}_{\mathcal{A}_2}^{1,4}(\mathbb{Z}/2, \mathbb{Z}/2)$ and $f_0 = [\overline{f}_0] \in \text{Ext}_{\mathcal{A}_2}^{4,22}(\mathbb{Z}/2, \mathbb{Z}/2)$. Note that $h_2 f_0 = h_1 g_1$ with $h_1 \in \text{Ext}_{\mathcal{A}_2}^{1,2}(\mathbb{Z}/2, \mathbb{Z}/2)$ and $g_1 \in \text{Ext}_{\mathcal{A}_2}^{4,24}(\mathbb{Z}/2, \mathbb{Z}/2)$. We consider the following element in $H_{21}(B(\mathbb{Z}/2)^{\times 5})$:

$$\begin{aligned} Z = & a_1^{(3)} a_2^{(3)} a_3^{(5)} a_4^{(1)} a_5^{(9)} + a_1^{(3)} a_2^{(3)} a_3^{(5)} a_4^{(2)} a_5^{(8)} + a_1^{(3)} a_2^{(3)} a_3^{(6)} a_4^{(1)} a_5^{(8)} + a_1^{(3)} a_2^{(3)} a_3^{(6)} a_4^{(2)} a_5^{(7)} \\ & + a_1^{(3)} a_2^{(3)} a_3^{(5)} a_4^{(4)} a_5^{(6)} + a_1^{(3)} a_2^{(3)} a_3^{(6)} a_4^{(3)} a_5^{(6)} + a_1^{(3)} a_2^{(5)} a_3^{(6)} a_4^{(1)} a_5^{(6)} + a_1^{(3)} a_2^{(3)} a_3^{(5)} a_4^{(5)} a_5^{(5)} \\ & + a_1^{(3)} a_2^{(3)} a_3^{(6)} a_4^{(4)} a_5^{(5)} + a_1^{(3)} a_2^{(5)} a_3^{(6)} a_4^{(2)} a_5^{(5)} + a_1^{(3)} a_2^{(3)} a_3^{(9)} a_4^{(1)} a_5^{(5)} + a_1^{(3)} a_2^{(5)} a_3^{(7)} a_4^{(1)} a_5^{(5)} \\ & + a_1^{(3)} a_2^{(3)} a_3^{(9)} a_4^{(2)} a_5^{(4)} + a_1^{(3)} a_2^{(5)} a_3^{(7)} a_4^{(2)} a_5^{(4)} + a_1^{(3)} a_2^{(3)} a_3^{(10)} a_4^{(1)} a_5^{(4)} + a_1^{(3)} a_2^{(6)} a_3^{(7)} a_4^{(1)} a_5^{(4)} \\ & + a_1^{(3)} a_2^{(5)} a_3^{(6)} a_4^{(4)} a_5^{(3)} + a_1^{(3)} a_2^{(6)} a_3^{(7)} a_4^{(2)} a_5^{(3)} + a_1^{(3)} a_2^{(3)} a_3^{(10)} a_4^{(2)} a_5^{(3)} + a_1^{(3)} a_2^{(3)} a_3^{(11)} a_4^{(2)} a_5^{(2)} \\ & + a_1^{(3)} a_2^{(5)} a_3^{(9)} a_4^{(2)} a_5^{(2)} + a_1^{(3)} a_2^{(6)} a_3^{(10)} a_4^{(1)} a_5^{(1)} + a_1^{(3)} a_2^{(3)} a_3^{(5)} a_4^{(9)} a_5^{(1)} + a_1^{(3)} a_2^{(3)} a_3^{(5)} a_4^{(8)} a_5^{(2)} \\ & + a_1^{(3)} a_2^{(3)} a_3^{(6)} a_4^{(8)} a_5^{(1)} + a_1^{(3)} a_2^{(3)} a_3^{(6)} a_4^{(7)} a_5^{(2)} + a_1^{(3)} a_2^{(3)} a_3^{(5)} a_4^{(6)} a_5^{(4)} + a_1^{(3)} a_2^{(3)} a_3^{(6)} a_4^{(6)} a_5^{(3)} \\ & + a_1^{(3)} a_2^{(5)} a_3^{(6)} a_4^{(6)} a_5^{(1)} + a_1^{(3)} a_2^{(3)} a_3^{(5)} a_4^{(5)} a_5^{(5)} + a_1^{(3)} a_2^{(3)} a_3^{(6)} a_4^{(5)} a_5^{(4)} + a_1^{(3)} a_2^{(5)} a_3^{(6)} a_4^{(5)} a_5^{(2)} \\ & + a_1^{(3)} a_2^{(3)} a_3^{(9)} a_4^{(5)} a_5^{(1)} + a_1^{(3)} a_2^{(5)} a_3^{(7)} a_4^{(5)} a_5^{(1)} + a_1^{(3)} a_2^{(3)} a_3^{(9)} a_4^{(4)} a_5^{(2)} + a_1^{(3)} a_2^{(5)} a_3^{(7)} a_4^{(4)} a_5^{(2)} \\ & + a_1^{(3)} a_2^{(3)} a_3^{(10)} a_4^{(4)} a_5^{(1)} + a_1^{(3)} a_2^{(6)} a_3^{(7)} a_4^{(4)} a_5^{(1)} + a_1^{(3)} a_2^{(5)} a_3^{(6)} a_4^{(3)} a_5^{(4)} + a_1^{(3)} a_2^{(6)} a_3^{(7)} a_4^{(3)} a_5^{(2)} \\ & + a_1^{(3)} a_2^{(3)} a_3^{(10)} a_4^{(3)} a_5^{(2)} + a_1^{(3)} a_2^{(3)} a_3^{(11)} a_4^{(2)} a_5^{(2)} + a_1^{(3)} a_2^{(5)} a_3^{(9)} a_4^{(2)} a_5^{(2)} + a_1^{(3)} a_2^{(6)} a_3^{(10)} a_4^{(1)} a_5^{(1)} \\ & + a_1^{(3)} a_2^{(3)} a_3^{(12)} a_4^{(1)} a_5^{(2)} + a_1^{(3)} a_2^{(7)} a_3^{(8)} a_4^{(1)} a_5^{(2)} + a_1^{(3)} a_2^{(11)} a_3^{(4)} a_4^{(1)} a_5^{(2)} + a_1^{(3)} a_2^{(13)} a_3^{(2)} a_4^{(1)} a_5^{(2)} \\ & + a_1^{(3)} a_2^{(14)} a_3^{(1)} a_4^{(1)} a_5^{(2)} + a_1^{(3)} a_2^{(12)} a_3^{(3)} a_4^{(1)} a_5^{(2)} + a_1^{(3)} a_2^{(8)} a_3^{(7)} a_4^{(1)} a_5^{(2)} + a_1^{(3)} a_2^{(4)} a_3^{(11)} a_4^{(1)} a_5^{(2)} \\ & + a_1^{(3)} a_2^{(2)} a_3^{(13)} a_4^{(1)} a_5^{(2)} + a_1^{(3)} a_2^{(1)} a_3^{(14)} a_4^{(1)} a_5^{(2)} + a_1^{(3)} a_2^{(6)} a_3^{(6)} a_4^{(3)} a_5^{(3)} + a_1^{(3)} a_2^{(5)} a_3^{(5)} a_4^{(5)} a_5^{(3)} \\ & + a_1^{(3)} a_2^{(3)} a_3^{(3)} a_4^{(9)} a_5^{(3)} + a_1^{(3)} a_2^{(5)} a_3^{(3)} a_4^{(7)} a_5^{(3)} + a_1^{(3)} a_2^{(7)} a_3^{(7)} a_4^{(2)} a_5^{(2)} + a_1^{(3)} a_2^{(6)} a_3^{(9)} a_4^{(1)} a_5^{(2)} \\ & + a_1^{(3)} a_2^{(9)} a_3^{(6)} a_4^{(1)} a_5^{(2)} + a_1^{(3)} a_2^{(10)} a_3^{(5)} a_4^{(1)} a_5^{(2)} + a_1^{(3)} a_2^{(5)} a_3^{(10)} a_4^{(2)} a_5^{(1)} + a_1^{(3)} a_2^{(13)} a_3^{(3)} a_4^{(1)} a_5^{(1)} \\ & + a_1^{(3)} a_2^{(5)} a_3^{(11)} a_4^{(1)} a_5^{(1)} + a_1^{(3)} a_2^{(9)} a_3^{(7)} a_4^{(1)} a_5^{(1)}. \end{aligned}$$

From the proof of Theorem 1.2 (see Sect. 4.3), $\{[q_2]\}$ is a basis of $(Q\mathcal{P}_5)_{21}^{GL_5}$. Obviously, $\langle Z, q_2 \rangle = 1$. Then, by a routine computation, we can verify that $Z \in P_{\mathcal{A}_2} H_{21}(B(\mathbb{Z}/2)^{\times 5})$. This implies $[Z] = ([q_2])^*$. By a direct computation, we have the representation in the $\mathbb{Z}/2$ -lambda algebra of Tr_5 :

$$a^{(n_1, \dots, n_5)} \longmapsto \sum_{\substack{i, j, h, m, k_1+k_2+k_3+k_4=i-n_1, \\ \ell_1+\ell_2+\ell_3=j-n_2+k_4, \\ u_1+u_2=h-n_3+k_3+\ell_3}} b_1.b_2.b_3.b_4.b_5,$$

where

$$\begin{aligned} b_1 &= \binom{n_5-k_1}{k_1} \binom{n_4-k_2}{k_2} \binom{n_3-k_3}{k_3} \binom{n_2-k_4}{k_4}, \\ b_2 &= \binom{n_5-k_1-\ell_1}{\ell_1} \binom{n_4-k_2-\ell_2}{\ell_2} \binom{n_3-k_3-\ell_3}{\ell_3}, \\ b_3 &= \binom{n_5-k_1-\ell_1-u_1}{u_1} \binom{n_4-k_2-\ell_2-u_2}{u_2}, \\ b_4 &= \binom{n_1+n_2+n_3+n_4+n_5-i-j-h-m}{m-n_4+k_2+\ell_2+u_2}, \\ b_5 &= \lambda_{i, j, h, m, n_1+n_2+n_3+n_4+n_5-i-j-h-m}, \end{aligned}$$

with the binomial factors (modulo 2). Using this presentation, we get

$$Z \longmapsto \lambda_3 \bar{f}_0 + \lambda_3^2 \lambda_5 \lambda_3 \lambda_7 = \lambda_3 \bar{f}_0 + \partial(\lambda_3^2 \lambda_5 \lambda_{11}).$$

Combining this with a previous result that $\psi_5(Z)$ is a cycle in the lambda algebra Λ and $Tr_5([Z]) = [\psi_5(Z)]$ (see [7]), we see that $h_2 f_0$ is in the image of Tr_5 . By Theorem 1.2 and the results of Lin [17] and Chen [6], $\mathbb{Z}/2 \otimes_{GL_5} P_{\mathcal{A}_2} H_{21}(B(\mathbb{Z}/2)^{\times 5})$ and $\text{Ext}_{\mathcal{A}_2}^{5,26}(\mathbb{Z}/2, \mathbb{Z}/2)$ that have the same dimensions are 1. So, from these data, we see that Tr_5 is an isomorphism at the bidegree $(5, 26)$. Theorem 1.3 is proved.

6. Appendix

In this section, we describe all the admissible monomials in $\mathcal{B}_5(n)$ for $n = 8, 21, 22, 47$. These monomials are mentioned in Sects. 3 and 4.

6.1. $\mathcal{A}_2$ -generators for $\mathcal{P}_5$ in degree 8

Based on a result in [53], we have $\mathcal{B}_5(8) = \bigcup_{1 \leq j \leq 3} \mathcal{B}_5(\tilde{\omega}_{(j)})$ and $|\mathcal{B}_5(8)| = 174$. Here $\tilde{\omega}_{(1)} = (2, 1, 1)$, $\tilde{\omega}_{(2)} = (2, 3)$, $\tilde{\omega}_{(3)} = (4, 2)$. By a simple computation, we get

$$|\overline{\Phi}(\mathcal{B}_4(\tilde{\omega}_{(1)})) \cup \mathcal{B}(5, \tilde{\omega}_{(1)})| = 105, \quad |\overline{\Phi}(\mathcal{B}_4(\tilde{\omega}_{(2)})) \cup \mathcal{B}(5, \tilde{\omega}_{(2)})| = 24,$$

$$|\overline{\Phi}(\mathcal{B}_4(\tilde{\omega}_{(3)})) \cup \mathcal{B}(5, \tilde{\omega}_{(3)})| = 45,$$

where the set $\mathcal{B}(5, \tilde{\omega}_{(j)})$ is given in Notation 3.4.10. Furthermore,

$$\mathcal{B}_5(\tilde{\omega}_{(j)}) = \overline{\Phi}(\mathcal{B}_4(\tilde{\omega}_{(j)})) \cup \mathcal{B}(5, \tilde{\omega}_{(j)}), \quad j = 1, 2, 3.$$

This implies that Conjecture 3.3.1 is true for $d = 5$ and in degree 8.

Now, we have $\mathcal{B}_5(8) = \{\mathcal{Y}_{8,t} : 1 \leq t \leq 174\}$, where the monomials $\mathcal{Y}_{8,t} : 1 \leq t \leq 174$, are determined as below:

$$\begin{aligned}
\mathcal{V}_{8,1} &= x_4 x_5^7, & \mathcal{V}_{8,2} &= x_4^7 x_5, & \mathcal{V}_{8,3} &= x_3 x_5^7, & \mathcal{V}_{8,4} &= x_3 x_4^7, \\
\mathcal{V}_{8,5} &= x_3^7 x_5, & \mathcal{V}_{8,6} &= x_3^7 x_4, & \mathcal{V}_{8,7} &= x_2 x_5^7, & \mathcal{V}_{8,8} &= x_2 x_4^7, \\
\mathcal{V}_{8,9} &= x_2 x_3^7, & \mathcal{V}_{8,10} &= x_2^7 x_5, & \mathcal{V}_{8,11} &= x_2^7 x_4, & \mathcal{V}_{8,12} &= x_2^7 x_3, \\
\mathcal{V}_{8,13} &= x_1 x_5^7, & \mathcal{V}_{8,14} &= x_1 x_4^7, & \mathcal{V}_{8,15} &= x_1 x_3^7, & \mathcal{V}_{8,16} &= x_1 x_2^7, \\
\mathcal{V}_{8,17} &= x_1^7 x_2, & \mathcal{V}_{8,18} &= x_1^7 x_3, & \mathcal{V}_{8,19} &= x_1^7 x_4, & \mathcal{V}_{8,20} &= x_1^7 x_5, \\
\mathcal{V}_{8,21} &= x_4^3 x_5^5, & \mathcal{V}_{8,22} &= x_3^3 x_5^5, & \mathcal{V}_{8,23} &= x_3^3 x_4^5, & \mathcal{V}_{8,24} &= x_2^3 x_5^5, \\
\mathcal{V}_{8,25} &= x_2^3 x_4^5, & \mathcal{V}_{8,26} &= x_2^3 x_3^5, & \mathcal{V}_{8,27} &= x_1^3 x_5^5, & \mathcal{V}_{8,28} &= x_1^3 x_4^5, \\
\mathcal{V}_{8,29} &= x_1^3 x_3^5, & \mathcal{V}_{8,30} &= x_1^3 x_2^5, & \mathcal{V}_{8,31} &= x_3 x_4 x_5^6, & \mathcal{V}_{8,32} &= x_3 x_4^6 x_5, \\
\mathcal{V}_{8,33} &= x_2 x_4 x_5^6, & \mathcal{V}_{8,34} &= x_2 x_4^6 x_5, & \mathcal{V}_{8,35} &= x_2 x_3 x_5^6, & \mathcal{V}_{8,36} &= x_2 x_3 x_4^6, \\
\mathcal{V}_{8,37} &= x_2 x_3^6 x_5, & \mathcal{V}_{8,38} &= x_2 x_3^6 x_4, & \mathcal{V}_{8,39} &= x_1 x_4 x_5^6, & \mathcal{V}_{8,40} &= x_1 x_3 x_5^6, \\
\mathcal{V}_{8,41} &= x_1 x_3 x_4^6, & \mathcal{V}_{8,42} &= x_1 x_2 x_5^6, & \mathcal{V}_{8,43} &= x_1 x_2 x_4^6, & \mathcal{V}_{8,44} &= x_1 x_2 x_3^6, \\
\mathcal{V}_{8,45} &= x_1 x_4^6 x_5, & \mathcal{V}_{8,46} &= x_1 x_3^6 x_5, & \mathcal{V}_{8,47} &= x_1 x_3^6 x_4, & \mathcal{V}_{8,48} &= x_1 x_2^6 x_5, \\
\mathcal{V}_{8,49} &= x_1 x_2^6 x_4, & \mathcal{V}_{8,50} &= x_1 x_2^6 x_3, & \mathcal{V}_{8,51} &= x_3^3 x_4 x_5^4, & \mathcal{V}_{8,52} &= x_3^3 x_4^4 x_5, \\
\mathcal{V}_{8,53} &= x_2^3 x_4 x_5^4, & \mathcal{V}_{8,54} &= x_2^3 x_3 x_5^4, & \mathcal{V}_{8,55} &= x_2^3 x_3 x_4^4, & \mathcal{V}_{8,56} &= x_2^3 x_4^4 x_5, \\
\mathcal{V}_{8,57} &= x_2^3 x_3^4 x_5, & \mathcal{V}_{8,58} &= x_2^3 x_3^4 x_4, & \mathcal{V}_{8,59} &= x_1^3 x_4 x_5^4, & \mathcal{V}_{8,60} &= x_1^3 x_3 x_5^4, \\
\mathcal{V}_{8,61} &= x_1^3 x_3 x_4^4, & \mathcal{V}_{8,62} &= x_1^3 x_4^4 x_5, & \mathcal{V}_{8,63} &= x_1^3 x_3^4 x_5, & \mathcal{V}_{8,64} &= x_1^3 x_3^4 x_4, \\
\mathcal{V}_{8,65} &= x_1^3 x_2 x_5^4, & \mathcal{V}_{8,66} &= x_1^3 x_2 x_4^4, & \mathcal{V}_{8,67} &= x_1^3 x_2^4 x_5, & \mathcal{V}_{8,68} &= x_1^3 x_2^4 x_4, \\
\mathcal{V}_{8,69} &= x_1^3 x_2^4 x_3, & \mathcal{V}_{8,70} &= x_1^3 x_2^4 x_2, & \mathcal{V}_{8,71} &= x_3 x_4^2 x_5^5, & \mathcal{V}_{8,72} &= x_2 x_4^2 x_5^5, \\
\mathcal{V}_{8,73} &= x_2 x_3^2 x_5^5, & \mathcal{V}_{8,74} &= x_2 x_3^2 x_4^5, & \mathcal{V}_{8,75} &= x_1 x_4^2 x_5^5, & \mathcal{V}_{8,76} &= x_1 x_3^2 x_5^5, \\
\mathcal{V}_{8,77} &= x_1 x_3^2 x_4^5, & \mathcal{V}_{8,78} &= x_1 x_2^2 x_5^5, & \mathcal{V}_{8,79} &= x_1 x_2^2 x_4^5, & \mathcal{V}_{8,80} &= x_1 x_2^2 x_3^5, \\
\mathcal{V}_{8,81} &= x_3 x_4^3 x_5^4, & \mathcal{V}_{8,82} &= x_2 x_3^3 x_5^4, & \mathcal{V}_{8,83} &= x_2 x_3^3 x_4^5, & \mathcal{V}_{8,84} &= x_2 x_3^3 x_4^4, \\
\mathcal{V}_{8,85} &= x_1 x_4^3 x_5^4, & \mathcal{V}_{8,86} &= x_1 x_3^3 x_5^4, & \mathcal{V}_{8,87} &= x_1 x_3^3 x_4^4, & \mathcal{V}_{8,88} &= x_1 x_2^3 x_5^4, \\
\mathcal{V}_{8,89} &= x_1 x_2^3 x_4^4, & \mathcal{V}_{8,90} &= x_1 x_2^3 x_3^4, & \mathcal{V}_{8,91} &= x_2 x_3 x_4^2 x_5^4, & \mathcal{V}_{8,92} &= x_2 x_3^2 x_4 x_5^4, \\
\mathcal{V}_{8,93} &= x_2 x_3^2 x_4 x_5, & \mathcal{V}_{8,94} &= x_1 x_3 x_4^2 x_5^4, & \mathcal{V}_{8,95} &= x_1 x_2 x_4^2 x_5^4, & \mathcal{V}_{8,96} &= x_1 x_2 x_3^2 x_5^4, \\
\mathcal{V}_{8,97} &= x_1 x_2 x_3^2 x_4^4, & \mathcal{V}_{8,98} &= x_1 x_3^2 x_4 x_5^4, & \mathcal{V}_{8,99} &= x_1 x_3^2 x_4 x_5, & \mathcal{V}_{8,100} &= x_1 x_2^2 x_4 x_5^4, \\
\mathcal{V}_{8,101} &= x_1 x_2^2 x_4 x_5, & \mathcal{V}_{8,102} &= x_1 x_2^2 x_3 x_5^4, & \mathcal{V}_{8,103} &= x_1 x_2^2 x_3 x_4^4, & \mathcal{V}_{8,104} &= x_1 x_2^2 x_3^4 x_5, \\
\mathcal{V}_{8,105} &= x_1 x_2^2 x_3^4 x_4, & \mathcal{V}_{8,106} &= x_2 x_3^2 x_4 x_5^3, & \mathcal{V}_{8,107} &= x_2 x_3^2 x_4 x_5^2, & \mathcal{V}_{8,108} &= x_2 x_3^2 x_4^2 x_5^2, \\
\mathcal{V}_{8,109} &= x_2^3 x_3 x_4^2 x_5^2, & \mathcal{V}_{8,110} &= x_1 x_3^2 x_4 x_5^3, & \mathcal{V}_{8,111} &= x_1 x_3^2 x_4 x_5^2, & \mathcal{V}_{8,112} &= x_1 x_3^2 x_4^2 x_5^2, \\
\mathcal{V}_{8,113} &= x_1 x_2^2 x_4^2 x_5^3, & \mathcal{V}_{8,114} &= x_1 x_2^2 x_3^3 x_5^2, & \mathcal{V}_{8,115} &= x_1 x_2^2 x_3^3 x_4^2, & \mathcal{V}_{8,116} &= x_1 x_2^2 x_3^3 x_5^3, \\
\mathcal{V}_{8,117} &= x_1 x_2^2 x_3^3 x_4^2, & \mathcal{V}_{8,118} &= x_1 x_2^2 x_3^2 x_5^2, & \mathcal{V}_{8,119} &= x_1 x_2^2 x_3^2 x_4^3, & \mathcal{V}_{8,120} &= x_1 x_2^2 x_3^2 x_4^2, \\
\mathcal{V}_{8,121} &= x_1 x_2^2 x_3^2 x_4^2, & \mathcal{V}_{8,122} &= x_1^3 x_3 x_4^2 x_5^2, & \mathcal{V}_{8,123} &= x_1^3 x_2 x_4^2 x_5^2, & \mathcal{V}_{8,124} &= x_1^3 x_2 x_3^2 x_5^2, \\
\mathcal{V}_{8,125} &= x_1^3 x_2 x_3^2 x_4^2, & \mathcal{V}_{8,126} &= x_1 x_2 x_3^2 x_4^2 x_5^2, & \mathcal{V}_{8,127} &= x_1 x_2^2 x_3 x_4^2 x_5^2, & \mathcal{V}_{8,128} &= x_1 x_2^2 x_3 x_4 x_5^2, \\
\mathcal{V}_{8,129} &= x_1 x_2^2 x_3^2 x_4 x_5, & \mathcal{V}_{8,130} &= x_2 x_3 x_4^3 x_5^3, & \mathcal{V}_{8,131} &= x_2 x_3^3 x_4 x_5^3, & \mathcal{V}_{8,132} &= x_2 x_3^3 x_4 x_5, \\
\mathcal{V}_{8,133} &= x_2^3 x_3 x_4^3 x_5, & \mathcal{V}_{8,134} &= x_2^3 x_3 x_4 x_5^3, & \mathcal{V}_{8,135} &= x_2^3 x_3^3 x_4 x_5, & \mathcal{V}_{8,136} &= x_1 x_3 x_3^3 x_5^3, \\
\mathcal{V}_{8,137} &= x_1 x_3^3 x_4 x_5^3, & \mathcal{V}_{8,138} &= x_1 x_3^3 x_3 x_5^3, & \mathcal{V}_{8,139} &= x_1 x_2 x_3^3 x_5^3, & \mathcal{V}_{8,140} &= x_1 x_2 x_3^3 x_5^3, \\
\mathcal{V}_{8,141} &= x_1 x_2 x_3^3 x_4^3, & \mathcal{V}_{8,142} &= x_1 x_2^3 x_4 x_5^3, & \mathcal{V}_{8,143} &= x_1 x_2^3 x_3 x_5^3, & \mathcal{V}_{8,144} &= x_1 x_2^3 x_3 x_5^3, \\
\mathcal{V}_{8,145} &= x_1 x_2^3 x_3 x_4^3, & \mathcal{V}_{8,146} &= x_1 x_2^3 x_3^3 x_5, & \mathcal{V}_{8,147} &= x_1 x_2^3 x_3^3 x_4, & \mathcal{V}_{8,148} &= x_1^3 x_3 x_4 x_5^3, \\
\mathcal{V}_{8,149} &= x_1^3 x_3 x_3^3 x_5, & \mathcal{V}_{8,150} &= x_1^3 x_3^3 x_4 x_5, & \mathcal{V}_{8,151} &= x_1^3 x_2 x_4 x_5^3, & \mathcal{V}_{8,152} &= x_1^3 x_2 x_3^3 x_5, \\
\mathcal{V}_{8,153} &= x_1^3 x_2 x_3 x_5^3, & \mathcal{V}_{8,154} &= x_1^3 x_2 x_3 x_4^3, & \mathcal{V}_{8,155} &= x_1^3 x_2 x_3^3 x_5, & \mathcal{V}_{8,156} &= x_1^3 x_2 x_3^3 x_4, \\
\mathcal{V}_{8,157} &= x_1^3 x_2^3 x_4 x_5, & \mathcal{V}_{8,158} &= x_1^3 x_2^3 x_3 x_5, & \mathcal{V}_{8,159} &= x_1^3 x_2^3 x_3 x_4, & \mathcal{V}_{8,160} &= x_1 x_2 x_3 x_4^2 x_5^3, \\
\mathcal{V}_{8,161} &= x_1 x_2 x_3 x_4^2 x_5^3, & \mathcal{V}_{8,162} &= x_1 x_2 x_3^2 x_4 x_5^3, & \mathcal{V}_{8,163} &= x_1 x_2 x_3^2 x_3 x_5^3, & \mathcal{V}_{8,164} &= x_1 x_2 x_3^2 x_4 x_5^3, \\
\mathcal{V}_{8,165} &= x_1 x_2 x_3^2 x_4 x_5, & \mathcal{V}_{8,166} &= x_1 x_2^2 x_3 x_4 x_5^3, & \mathcal{V}_{8,167} &= x_1 x_2^2 x_3 x_4 x_5, & \mathcal{V}_{8,168} &= x_1 x_2^2 x_3^2 x_4 x_5, \\
\mathcal{V}_{8,169} &= x_1 x_2^2 x_3 x_4 x_5^3, & \mathcal{V}_{8,170} &= x_1 x_2^2 x_3 x_4 x_5, & \mathcal{V}_{8,171} &= x_1 x_2^2 x_3^3 x_4 x_5, & \mathcal{V}_{8,172} &= x_1^3 x_2 x_3 x_4 x_5^2, \\
\mathcal{V}_{8,173} &= x_1^3 x_2 x_3 x_4^2 x_5, & \mathcal{V}_{8,174} &= x_1^3 x_2 x_3^2 x_4 x_5.
\end{aligned}$$

6.2. $\mathcal{A}_2$ -generators for $\mathcal{P}_5^0$ in degree 21

Using the results in Sect. 3.4.1, we have $\mathcal{B}_5^0(21) = \mathcal{B}_5^0(3, 3, 1, 1) \cup \mathcal{B}_5^0(3, 3, 3)$, where $\mathcal{B}_5^0(3, 3, 1, 1)$ is the set of 340 admissible monomials: $\mathcal{V}_{21,t}$, $1 \leq t \leq 340$

$\mathcal{V}_{21,1} = x_3^3 x_4^3 x_5^{15}$	$\mathcal{V}_{21,2} = x_3^3 x_4^{15} x_5^3$	$\mathcal{V}_{21,3} = x_3^{15} x_4^3 x_5^3$	$\mathcal{V}_{21,4} = x_3^3 x_4^3 x_5^{15}$
$\mathcal{V}_{21,5} = x_2^3 x_4^{15} x_5^3$	$\mathcal{V}_{21,6} = x_2^3 x_3^3 x_5^{15}$	$\mathcal{V}_{21,7} = x_2^3 x_3^3 x_4^{15}$	$\mathcal{V}_{21,8} = x_2^3 x_3^{15} x_5^3$
$\mathcal{V}_{21,9} = x_2^3 x_3^{15} x_4^3$	$\mathcal{V}_{21,10} = x_2^{15} x_4^3 x_5^3$	$\mathcal{V}_{21,11} = x_2^{15} x_3^3 x_5^3$	$\mathcal{V}_{21,12} = x_2^{15} x_3^3 x_4^3$
$\mathcal{V}_{21,13} = x_1^3 x_4^3 x_5^{15}$	$\mathcal{V}_{21,14} = x_1^3 x_4^{15} x_5^3$	$\mathcal{V}_{21,15} = x_1^3 x_3^3 x_5^{15}$	$\mathcal{V}_{21,16} = x_1^3 x_3^3 x_4^{15}$
$\mathcal{V}_{21,17} = x_1^3 x_3^{15} x_5^3$	$\mathcal{V}_{21,18} = x_1^3 x_3^{15} x_4^3$	$\mathcal{V}_{21,19} = x_1^3 x_2^3 x_5^{15}$	$\mathcal{V}_{21,20} = x_1^3 x_2^3 x_4^{15}$
$\mathcal{V}_{21,21} = x_1^3 x_2^3 x_3^{15}$	$\mathcal{V}_{21,22} = x_1^3 x_2^{15} x_5^3$	$\mathcal{V}_{21,23} = x_1^3 x_2^{15} x_4^3$	$\mathcal{V}_{21,24} = x_1^3 x_2^{15} x_3^3$
$\mathcal{V}_{21,25} = x_1^{15} x_4^3 x_5^3$	$\mathcal{V}_{21,26} = x_1^{15} x_3^3 x_5^3$	$\mathcal{V}_{21,27} = x_1^{15} x_3^3 x_4^3$	$\mathcal{V}_{21,28} = x_1^{15} x_2^3 x_5^3$
$\mathcal{V}_{21,29} = x_1^{15} x_2^3 x_4^3$	$\mathcal{V}_{21,30} = x_1^{15} x_2^3 x_3^3$	$\mathcal{V}_{21,31} = x_3^3 x_4^7 x_5^{11}$	$\mathcal{V}_{21,32} = x_3^3 x_4^3 x_5^{11}$
$\mathcal{V}_{21,33} = x_3^7 x_4^{11} x_5^3$	$\mathcal{V}_{21,34} = x_2^3 x_4^7 x_5^{11}$	$\mathcal{V}_{21,35} = x_2^3 x_3^7 x_5^{11}$	$\mathcal{V}_{21,36} = x_2^3 x_3^7 x_4^{11}$
$\mathcal{V}_{21,37} = x_2^7 x_4^{11} x_5^3$	$\mathcal{V}_{21,38} = x_2^7 x_4^{11} x_3^3$	$\mathcal{V}_{21,39} = x_2^7 x_3^3 x_5^{11}$	$\mathcal{V}_{21,40} = x_2^7 x_3^3 x_4^{11}$
$\mathcal{V}_{21,41} = x_2^7 x_3^{11} x_5^3$	$\mathcal{V}_{21,42} = x_2^7 x_3^{11} x_4^3$	$\mathcal{V}_{21,43} = x_1^3 x_4^7 x_5^{11}$	$\mathcal{V}_{21,44} = x_1^3 x_3^7 x_5^{11}$
$\mathcal{V}_{21,45} = x_1^3 x_3^7 x_4^{11}$	$\mathcal{V}_{21,46} = x_1^3 x_2^7 x_5^{11}$	$\mathcal{V}_{21,47} = x_1^3 x_2^7 x_4^{11}$	$\mathcal{V}_{21,48} = x_1^3 x_2^7 x_3^{11}$
$\mathcal{V}_{21,49} = x_1^7 x_4^3 x_5^{11}$	$\mathcal{V}_{21,50} = x_1^7 x_4^{11} x_5^3$	$\mathcal{V}_{21,51} = x_1^7 x_3^3 x_5^{11}$	$\mathcal{V}_{21,52} = x_1^7 x_3^3 x_4^{11}$
$\mathcal{V}_{21,53} = x_1^7 x_3^{11} x_5^3$	$\mathcal{V}_{21,54} = x_1^7 x_3^{11} x_4^3$	$\mathcal{V}_{21,55} = x_1^7 x_2^3 x_5^{11}$	$\mathcal{V}_{21,56} = x_1^7 x_2^3 x_4^{11}$
$\mathcal{V}_{21,57} = x_1^7 x_2^3 x_3^{11}$	$\mathcal{V}_{21,58} = x_1^7 x_2^{11} x_5^3$	$\mathcal{V}_{21,59} = x_1^7 x_2^{11} x_4^3$	$\mathcal{V}_{21,60} = x_1^7 x_2^{11} x_3^3$
$\mathcal{V}_{21,61} = x_2 x_3^2 x_4^{15} x_5^2$	$\mathcal{V}_{21,62} = x_2 x_3^2 x_4^{15} x_5^3$	$\mathcal{V}_{21,63} = x_2 x_3^3 x_4^2 x_5^{15}$	$\mathcal{V}_{21,64} = x_2 x_3^3 x_4^{15} x_5^2$
$\mathcal{V}_{21,65} = x_2 x_3^{15} x_4^2 x_5^3$	$\mathcal{V}_{21,66} = x_2 x_3^{15} x_4^2 x_5^2$	$\mathcal{V}_{21,67} = x_2^3 x_3^3 x_4^2 x_5^{15}$	$\mathcal{V}_{21,68} = x_2^3 x_3^3 x_4^{15} x_5^2$
$\mathcal{V}_{21,69} = x_2^3 x_3^{15} x_4^2 x_5^2$	$\mathcal{V}_{21,70} = x_2^{15} x_3^2 x_4^2 x_5^3$	$\mathcal{V}_{21,71} = x_2^{15} x_3^2 x_4^2 x_5^2$	$\mathcal{V}_{21,72} = x_2^{15} x_3^3 x_4^2 x_5^2$
$\mathcal{V}_{21,73} = x_1 x_3^2 x_4^{15} x_5^2$	$\mathcal{V}_{21,74} = x_1 x_3^2 x_4^{15} x_5^3$	$\mathcal{V}_{21,75} = x_1 x_3^3 x_4^2 x_5^{15}$	$\mathcal{V}_{21,76} = x_1 x_3^3 x_4^2 x_5^2$
$\mathcal{V}_{21,77} = x_1 x_3^{15} x_4^2 x_5^3$	$\mathcal{V}_{21,78} = x_1 x_3^{15} x_4^2 x_5^2$	$\mathcal{V}_{21,79} = x_1 x_2^2 x_4^2 x_5^{15}$	$\mathcal{V}_{21,80} = x_1 x_2^2 x_4^{15} x_5^3$
$\mathcal{V}_{21,81} = x_1 x_2^2 x_3^3 x_5^{15}$	$\mathcal{V}_{21,82} = x_1 x_2^2 x_3^3 x_4^{15}$	$\mathcal{V}_{21,83} = x_1 x_2^2 x_3^{15} x_5^3$	$\mathcal{V}_{21,84} = x_1 x_2^2 x_3^{15} x_4^3$
$\mathcal{V}_{21,85} = x_1 x_2^3 x_4^{15} x_5^2$	$\mathcal{V}_{21,86} = x_1 x_2^3 x_4^{15} x_5^3$	$\mathcal{V}_{21,87} = x_1 x_2^3 x_3^2 x_5^{15}$	$\mathcal{V}_{21,88} = x_1 x_2^3 x_3^2 x_4^{15}$
$\mathcal{V}_{21,89} = x_1 x_2^3 x_3^{15} x_5^2$	$\mathcal{V}_{21,90} = x_1 x_2^3 x_3^{15} x_4^2$	$\mathcal{V}_{21,91} = x_1 x_2^{15} x_4^2 x_5^3$	$\mathcal{V}_{21,92} = x_1 x_2^{15} x_4^2 x_5^2$
$\mathcal{V}_{21,93} = x_1 x_2^{15} x_3^2 x_5^3$	$\mathcal{V}_{21,94} = x_1 x_2^{15} x_3^2 x_4^3$	$\mathcal{V}_{21,95} = x_1 x_2^{15} x_3^3 x_5^2$	$\mathcal{V}_{21,96} = x_1 x_2^{15} x_3^3 x_4^2$
$\mathcal{V}_{21,97} = x_1^3 x_3^2 x_4^{15} x_5^2$	$\mathcal{V}_{21,98} = x_1^3 x_3^2 x_4^{15} x_5^3$	$\mathcal{V}_{21,99} = x_1^3 x_3^{15} x_4^2 x_5^2$	$\mathcal{V}_{21,100} = x_1^3 x_2^2 x_4^2 x_5^{15}$
$\mathcal{V}_{21,101} = x_3^3 x_2 x_4^{15} x_5^2$	$\mathcal{V}_{21,102} = x_3^3 x_2 x_3^2 x_5^{15}$	$\mathcal{V}_{21,103} =$	

6.3. $\mathcal{A}_2$ -generators for $\mathcal{P}_5^+$ in degree 21

Note that Kameko's squaring operation $\widetilde{Sq}_*^0(5,21) : (Q\mathcal{P}_5)_{21} \rightarrow (Q\mathcal{P}_5)_8$ is an epimorphism of $\mathbb{Z}/2GL_5$ -modules. Hence, we have

$$\mathcal{B}_5(21) = \mathcal{B}_5^0(21) \cup \varphi(\mathcal{B}_5(8)) \cup (\mathcal{B}_5^+(21) \cap \text{Ker}((\widetilde{Sq}_*^0)_{(5,21)})),$$

where $\mathcal{B}_5^0(21) = \overline{\Phi}^0(\mathcal{B}_4(21))$, $|\mathcal{B}_5^0(21)| = 400$, $|\varphi(\mathcal{B}_5(8))| = 174$ with

$$\varphi : \mathcal{P}_5 \rightarrow \mathcal{P}_5, \varphi(Z) = X_{(0,5)}Z^2, \forall Z \in \mathcal{P}_5.$$

Based on the results in Sect.3.4.1, we see that

$$\mathcal{B}_5^+(21) \cap \text{Ker}((\widetilde{Sq}_*^0)_{(5,21)}) = \mathcal{B}_5^+(3,3,1,1) \cup \mathcal{B}_5^+(3,3,3),$$

where $\mathcal{B}_5^+(3,3,1,1)$ is the set of 196 admissible monomials: $\mathcal{V}_{21,t}$, $401 \leq t \leq 596$

$$\begin{aligned} \mathcal{V}_{21,401} &= x_1x_2^2x_3x_4^2x_5^{15}, & \mathcal{V}_{21,402} &= x_1x_2^2x_3x_4^{15}x_5^2, & \mathcal{V}_{21,403} &= x_1x_2^2x_3^{15}x_4x_5^2, & \mathcal{V}_{21,404} &= x_1x_2x_3^2x_4^2x_5^{15}, \\ \mathcal{V}_{21,405} &= x_1x_2x_3^2x_4^{15}x_5^2, & \mathcal{V}_{21,406} &= x_1x_2^{15}x_3^2x_4x_5^2, & \mathcal{V}_{21,407} &= x_1x_2x_3^{15}x_4^2x_5^2, & \mathcal{V}_{21,408} &= x_1x_2^{15}x_3x_4^2x_5^2, \\ \mathcal{V}_{21,409} &= x_1^{15}x_2x_3^2x_4x_5^2, & \mathcal{V}_{21,410} &= x_1^{15}x_2x_3x_4^2x_5^2, & \mathcal{V}_{21,411} &= x_1x_2x_3^6x_4^3x_5^{10}, & \mathcal{V}_{21,412} &= x_1x_2x_3^6x_4^{10}x_5^3, \\ \mathcal{V}_{21,413} &= x_1x_2x_3^3x_4^6x_5^{10}, & \mathcal{V}_{21,414} &= x_1x_2^2x_3^3x_4^3x_5^{12}, & \mathcal{V}_{21,415} &= x_1x_2^2x_3^3x_4^{12}x_5^3, & \mathcal{V}_{21,416} &= x_1x_2^2x_3^{12}x_4^3x_5^3, \\ \mathcal{V}_{21,417} &= x_1^3x_2^3x_3^4x_4^8x_5^8, & \mathcal{V}_{21,418} &= x_1^3x_2^3x_3^4x_4^8x_5^3, & \mathcal{V}_{21,419} &= x_1^3x_2^3x_3^3x_4^8x_5^8, & \mathcal{V}_{21,420} &= x_1x_2^2x_3^3x_4^3x_5^{14}, \\ \mathcal{V}_{21,421} &= x_1x_2^2x_3x_4^{14}x_5^3, & \mathcal{V}_{21,422} &= x_1x_2^2x_3^3x_4^{14}x_5^3, & \mathcal{V}_{21,423} &= x_1x_2^{14}x_3x_4^2x_5^3, & \mathcal{V}_{21,424} &= x_1x_2^{14}x_3x_4^3x_5^2, \\ \mathcal{V}_{21,425} &= x_1x_2^{14}x_3^3x_4^2x_5^2, & \mathcal{V}_{21,426} &= x_1x_2x_3^2x_4^{14}x_5^3, & \mathcal{V}_{21,427} &= x_1x_2x_3^2x_4^{14}x_5^3, & \mathcal{V}_{21,428} &= x_1x_2x_3^{14}x_4^2x_5^3, \\ \mathcal{V}_{21,429} &= x_1x_2x_3^{14}x_4^2x_5^2, & \mathcal{V}_{21,430} &= x_1x_2^3x_3^2x_4^{14}x_5^3, & \mathcal{V}_{21,431} &= x_1x_2^3x_3^{14}x_4x_5^2, & \mathcal{V}_{21,432} &= x_1x_2^3x_3^{14}x_4^2x_5^{14}, \\ \mathcal{V}_{21,433} &= x_1x_2x_3^3x_4^{14}x_5^2, & \mathcal{V}_{21,434} &= x_1x_2^3x_3x_4^{14}x_5^2, & \mathcal{V}_{21,435} &= x_1x_2^3x_3x_4^{14}x_5^2, & \mathcal{V}_{21,436} &= x_1x_2^2x_3^6x_4^6x_5^{11}, \\ \mathcal{V}_{21,437} &= x_1x_2^6x_3x_4^2x_5^{11}, & \mathcal{V}_{21,438} &= x_1x_2^6x_3x_4^{11}x_5^2, & \mathcal{V}_{21,439} &= x_1x_2^6x_3^{11}x_4x_5^2, & \mathcal{V}_{21,440} &= x_1x_2x_3^6x_4^6x_5^{11}, \\ \mathcal{V}_{21,441} &= x_1x_2x_3^6x_4^2x_5^{11}, & \mathcal{V}_{21,442} &= x_1x_2x_3^6x_4^{11}x_5^2, & \mathcal{V}_{21,443} &= x_1x_2^2x_3^7x_4^{10}x_5^2, & \mathcal{V}_{21,444} &= x_1x_2^2x_3^7x_4^5x_5^{10}, \\ \mathcal{V}_{21,445} &= x_1x_2x_3^2x_4^7x_5^{10}, & \mathcal{V}_{21,446} &= x_1x_2^7x_3^{10}x_4x_5^2, & \mathcal{V}_{21,447} &= x_1x_2^7x_3^2x_4^{10}x_5^2, & \mathcal{V}_{21,448} &= x_1x_2x_3^7x_4^{10}x_5^2, \\ \mathcal{V}_{21,449} &= x_1x_2x_3^7x_4^2x_5^{10}, & \mathcal{V}_{21,450} &= x_1x_2^7x_3x_4^{10}x_5^2, & \mathcal{V}_{21,451} &= x_1x_2^7x_3x_4^2x_5^{10}, & \mathcal{V}_{21,452} &= x_1x_2^2x_3^3x_4^{13}x_5^2, \\ \mathcal{V}_{21,453} &= x_1x_2^2x_3^{13}x_4^2x_5^3, & \mathcal{V}_{21,454} &= x_1x_2^2x_3^{13}x_4^2x_5^2, & \mathcal{V}_{21,455} &= x_1x_2^2x_3^2x_4^{13}x_5^2, & \mathcal{V}_{21,456} &= x_1x_2^3x_3^{13}x_4^2x_5^2, \\ \mathcal{V}_{21,457} &= x_1x_2^2x_3^5x_4^2x_5^{11}, & \mathcal{V}_{21,458} &= x_1x_2^2x_3^5x_4^{11}x_5^2, & \mathcal{V}_{21,459} &= x_1x_2^2x_3^7x_4^9x_5^2, & \mathcal{V}_{21,460} &= x_1x_2^7x_3^2x_4^9x_5^2, \\ \mathcal{V}_{21,461} &= x_1x_2^7x_3^9x_4^2x_5^2, & \mathcal{V}_{21,462} &= x_1x_2^6x_3^3x_4^{10}x_5^3, & \mathcal{V}_{21,463} &= x_1x_2^6x_3^3x_4^{10}x_5^3, & \mathcal{V}_{21,464} &= x_1x_2^6x_3^3x_4^2x_5^{10}, \\ \mathcal{V}_{21,465} &= x_1x_2^3x_3^6x_4^{10}x_5^3, & \mathcal{V}_{21,466} &= x_1x_2^3x_3^6x_4^{10}x_5^3, & \mathcal{V}_{21,467} &= x_1x_2^3x_3^{12}x_4^2x_5^3, & \mathcal{V}_{21,468} &= x_1x_2^3x_3^{12}x_4^3x_5^2, \\ \mathcal{V}_{21,469} &= x_1x_2^3x_3^2x_4^{12}x_5^3, & \mathcal{V}_{21,470} &= x_1x_2^3x_3^2x_4^{12}x_5^3, & \mathcal{V}_{21,471} &= x_1x_2^3x_3^3x_4^{12}x_5^2, & \mathcal{V}_{21,472} &= x_1x_2^3x_3^3x_4^2x_5^{12}, \\ \mathcal{V}_{21,473} &= x_1x_2^2x_3^3x_4^4x_5^{11}, & \mathcal{V}_{21,474} &= x_1x_2^3x_3^4x_4^2x_5^{11}, & \mathcal{V}_{21,475} &= x_1x_2^3x_3^4x_4^{11}x_5^2, & \mathcal{V}_{21,476} &= x_1x_2^3x_3^2x_4^4x_5^{11}, \\ \mathcal{V}_{21,477} &= x_1x_2^2x_3^4x_4^3x_5^{11}, & \mathcal{V}_{21,478} &= x_1x_2^2x_3^4x_4^{11}x_5^3, & \mathcal{V}_{21,479} &= x_1x_2^2x_3^3x_4^5x_5^{10}, & \mathcal{V}_{21,480} &= x_1x_2^2x_3^3x_4^3x_5^{10}, \\ \mathcal{V}_{21,481} &= x_1x_2^2x_3^5x_4^{10}x_5^3, & \mathcal{V}_{21,482} &= x_1x_2^3x_3^2x_4^5x_5^{10}, & \mathcal{V}_{21,483} &= x_1x_2^3x_3^5x_4^{10}x_5^2, & \mathcal{V}_{21,484} &= x_1x_2^3x_3^5x_4^2x_5^{10}, \\ \mathcal{V}_{21,485} &= x_1x_2^6x_3^3x_4^9x_5^2, & \mathcal{V}_{21,486} &= x_1x_2^6x_3^9x_4^2x_5^3, & \mathcal{V}_{21,487} &= x_1x_2^6x_3^9x_4^2x_5^2, & \mathcal{V}_{21,488} &= x_1x_2^3x_3^6x_4^9x_5^2, \\ \mathcal{V}_{21,489} &= x_1x_2^5x_3^3x_4^8x_5^8, & \mathcal{V}_{21,490} &= x_1x_2^5x_3^7x_4^3x_5^8, & \mathcal{V}_{21,491} &= x_1x_2^5x_3^7x_4^3x_5^8, & \mathcal{V}_{21,492} &= x_1x_2^3x_3^2x_4^7x_5^8, \\ \mathcal{V}_{21,493} &= x_1x_2^7x_3^8x_4^2x_5^3, & \mathcal{V}_{21,494} &= x_1x_2^7x_3^8x_4^2x_5^3, & \mathcal{V}_{21,495} &= x_1x_2^7x_3^2x_4^8x_5^8, & \mathcal{V}_{21,496} &= x_1x_2^7x_3^2x_4^8x_5^3, \\ \mathcal{V}_{21,497} &= x_1x_2^3x_3^7x_4^8x_5^2, & \mathcal{V}_{21,498} &= x_1x_2^3x_3^7x_4^2x_5^8, & \mathcal{V}_{21,499} &= x_1x_2^7x_3^3x_4^2x_5^2, & \mathcal{V}_{21,500} &= x_1x_2^7x_3^3x_4^2x_5^8, \\ \mathcal{V}_{21,501} &= x_1x_2^3x_3^4x_4^3x_5^{10}, & \mathcal{V}_{21,502} &= x_1x_2^3x_3^4x_4^{10}x_5^3, & \mathcal{V}_{21,503} &= x_1x_2^3x_3^3x_4^4x_5^{10}, & \mathcal{V}_{21,504} &= x_1x_2^6x_3^3x_4^3x_5^8, \\ \mathcal{V}_{21,505} &= x_1x_2^6x_3^3x_4^8x_5^3, & \mathcal{V}_{21,506} &= x_1x_2^3x_3^6x_4^3x_5^8, & \mathcal{V}_{21,507} &= x_1x_2^3x_3^6x_4^3x_5^3, & \mathcal{V}_{21,508} &= x_1x_2^2x_3^3x_4^6x_5^8, \\ \mathcal{V}_{21,509} &= x_1^3x_2x_3^3x_4^2x_5^{14}, & \mathcal{V}_{21,510} &= x_1^3x_2x_3^{14}x_4^2x_5^2, & \mathcal{V}_{21,511} &= x_1^3x_2x_3^2x_4^2x_5^{14}, & \mathcal{V}_{21,512} &= x_1^3x_2x_3^2x_4^2x_5^2, \\ \mathcal{V}_{21,513} &= x_1^3x_2x_3^{13}x_4^2x_5^2, & \mathcal{V}_{21,514} &= x_1^3x_2x_3^{13}x_4^2x_5^2, & \mathcal{V}_{21,515} &= x_1^3x_2^{13}x_3x_4^2x_5^2, & \mathcal{V}_{21,516} &= x_1^3x_2^{13}x_3x_4^2x_5^2, \\ \mathcal{V}_{21,517} &= x_1^3x_2x_3^6x_4^2x_5^{10}, & \mathcal{V}_{21,518} &= x_1^3x_2x_3^6x_4^2x_5^{10}, & \mathcal{V}_{21,519} &= x_1^3x_2x_3^{12}x_4^2x_5^3, & \mathcal{V}_{21,520} &= x_1^3x_2x_3^{12}x_4^2x_5^2, \\ \mathcal{V}_{21,521} &= x_1^3x_2x_3^3x_4^{12}x_5^2, & \mathcal{V}_{21,522} &= x_1^3x_2x_3^3x_4^{12}x_5^2, & \mathcal{V}_{21,523} &= x_1^3x_2x_3^3x_4^{12}x_5^2, & \mathcal{V}_{21,524} &= x_1^3x_2x_3^3x_4^{12}x_5^2, \\ \mathcal{V}_{21,525} &= x_1^3x_2^3x_3x_4^{12}x_5^2, & \mathcal{V}_{21,526} &= x_1^3x_2^3x_3x_4^{12}x_5^2, & \mathcal{V}_{21,527} &= x_1^3x_2^3x_3^{12}x_4x_5^2, & \mathcal{V}_{21,528} &= x_1^3x_2^3x_3^{12}x_4x_5^2, \\ \mathcal{V}_{21,529} &= x_1^3x_2^{12}x_3x_4^2x_5^2, & \mathcal{V}_{21,530} &= x_1^3x_2^{12}x_3x_4^2x_5^2, & \mathcal{V}_{21,531} &= x_1^3x_2x_3^4x_4^2x_5^{11}, & \mathcal{V}_{21,532} &= x_1^3x_2x_3^4x_4^{11}x_5^2, \end{aligned}$$

$$\begin{aligned}
\mathcal{V}_{21,533} &= x_1^3 x_2 x_3^2 x_4^{11} x_5^{11}, & \mathcal{V}_{21,534} &= x_1^3 x_2^4 x_3 x_4^2 x_5^{11}, & \mathcal{V}_{21,535} &= x_1^3 x_2^4 x_3 x_4^{11} x_5^2, & \mathcal{V}_{21,536} &= x_1^3 x_2^4 x_3^{11} x_4 x_5^2, \\
\mathcal{V}_{21,537} &= x_1^3 x_2 x_3^2 x_4^5 x_5^{10}, & \mathcal{V}_{21,538} &= x_1^3 x_2 x_3^5 x_4^{10} x_5^2, & \mathcal{V}_{21,539} &= x_1^3 x_2 x_3^5 x_4^2 x_5^{10}, & \mathcal{V}_{21,540} &= x_1^3 x_2^5 x_3 x_4^{10} x_5^2, \\
\mathcal{V}_{21,541} &= x_1^3 x_2^5 x_3 x_4^2 x_5^{10}, & \mathcal{V}_{21,542} &= x_1^3 x_2^5 x_3^2 x_4 x_5^{10}, & \mathcal{V}_{21,543} &= x_1^3 x_2^5 x_3^{10} x_4 x_5^2, & \mathcal{V}_{21,544} &= x_1^3 x_2 x_3^6 x_4^9 x_5^2, \\
\mathcal{V}_{21,545} &= x_1^3 x_2 x_3^2 x_4^7 x_5^8, & \mathcal{V}_{21,546} &= x_1^3 x_2 x_3^7 x_4^8 x_5^2, & \mathcal{V}_{21,547} &= x_1^3 x_2 x_3^7 x_4^2 x_5^8, & \mathcal{V}_{21,548} &= x_1^3 x_2^7 x_3 x_4^8 x_5^2, \\
\mathcal{V}_{21,549} &= x_1^3 x_2 x_3 x_4^2 x_5^8, & \mathcal{V}_{21,550} &= x_1^3 x_2 x_3^8 x_4 x_5^2, & \mathcal{V}_{21,551} &= x_1^3 x_2 x_3^4 x_4^3 x_5^{10}, & \mathcal{V}_{21,552} &= x_1^3 x_2 x_3^4 x_4^{10} x_5^3, \\
\mathcal{V}_{21,553} &= x_1^3 x_2 x_3^3 x_4^4 x_5^{10}, & \mathcal{V}_{21,554} &= x_1^3 x_2 x_3^4 x_4 x_5^{10}, & \mathcal{V}_{21,555} &= x_1^3 x_2^3 x_3^4 x_4 x_5^{10}, & \mathcal{V}_{21,556} &= x_1^3 x_2^4 x_3 x_4^3 x_5^{10}, \\
\mathcal{V}_{21,557} &= x_1^3 x_2^4 x_3 x_4^{10} x_5^3, & \mathcal{V}_{21,558} &= x_1^3 x_2^4 x_3^3 x_4 x_5^{10}, & \mathcal{V}_{21,559} &= x_1^3 x_2 x_3^6 x_4^3 x_5^8, & \mathcal{V}_{21,560} &= x_1^3 x_2 x_3^6 x_4^8 x_5^3, \\
\mathcal{V}_{21,561} &= x_1^3 x_2 x_3^3 x_4^6 x_5^8, & \mathcal{V}_{21,562} &= x_1^3 x_2^3 x_3 x_4^6 x_5^8, & \mathcal{V}_{21,563} &= x_1^3 x_2^5 x_3^2 x_4^9 x_5^2, & \mathcal{V}_{21,564} &= x_1^3 x_2^5 x_3^2 x_4^2 x_5^2, \\
\mathcal{V}_{21,565} &= x_1^3 x_2^3 x_3^4 x_4^9 x_5^2, & \mathcal{V}_{21,566} &= x_1^3 x_2^4 x_3^3 x_4^9 x_5^2, & \mathcal{V}_{21,567} &= x_1^3 x_2^4 x_3^9 x_4^2 x_5^3, & \mathcal{V}_{21,568} &= x_1^3 x_2^4 x_3^9 x_4^3 x_5^2, \\
\mathcal{V}_{21,569} &= x_1^3 x_2^5 x_3^8 x_4 x_5^3, & \mathcal{V}_{21,570} &= x_1^3 x_2^5 x_3^8 x_4^2 x_5^2, & \mathcal{V}_{21,571} &= x_1^3 x_2^5 x_3^2 x_4^3 x_5^8, & \mathcal{V}_{21,572} &= x_1^3 x_2^5 x_3^2 x_4^8 x_5^3, \\
\mathcal{V}_{21,573} &= x_1^3 x_2^5 x_3^3 x_4^8 x_5^2, & \mathcal{V}_{21,574} &= x_1^3 x_2^5 x_3^3 x_4^2 x_5^8, & \mathcal{V}_{21,575} &= x_1^3 x_2^3 x_3^5 x_4^8 x_5^2, & \mathcal{V}_{21,576} &= x_1^3 x_2^3 x_3^5 x_4^2 x_5^8, \\
\mathcal{V}_{21,577} &= x_1^7 x_2 x_3^{10} x_4 x_5^2, & \mathcal{V}_{21,578} &= x_1^7 x_2 x_3^2 x_4 x_5^{10}, & \mathcal{V}_{21,579} &= x_1^7 x_2 x_3 x_4^{10} x_5^2, & \mathcal{V}_{21,580} &= x_1^7 x_2 x_3 x_4^2 x_5^{10}, \\
\mathcal{V}_{21,581} &= x_1^7 x_2 x_3^2 x_4^2 x_5^2, & \mathcal{V}_{21,582} &= x_1^7 x_2 x_3^9 x_4^2 x_5^2, & \mathcal{V}_{21,583} &= x_1^7 x_2^9 x_3 x_4^2 x_5^2, & \mathcal{V}_{21,584} &= x_1^7 x_2^9 x_3^2 x_4 x_5^2, \\
\mathcal{V}_{21,585} &= x_1^7 x_2 x_3^8 x_4^3 x_5^3, & \mathcal{V}_{21,586} &= x_1^7 x_2 x_3^8 x_4^2 x_5^2, & \mathcal{V}_{21,587} &= x_1^7 x_2 x_3^2 x_4^3 x_5^8, & \mathcal{V}_{21,588} &= x_1^7 x_2 x_3^2 x_4^8 x_5^3, \\
\mathcal{V}_{21,589} &= x_1^7 x_2 x_3^3 x_4^8 x_5^2, & \mathcal{V}_{21,590} &= x_1^7 x_2 x_3^3 x_4^2 x_5^8, & \mathcal{V}_{21,591} &= x_1^7 x_2^3 x_3 x_4^8 x_5^2, & \mathcal{V}_{21,592} &= x_1^7 x_2^3 x_3^2 x_4^8 x_5^2, \\
\mathcal{V}_{21,593} &= x_1^7 x_2^3 x_3^8 x_4 x_5^2, & \mathcal{V}_{21,594} &= x_1^7 x_2^3 x_3^8 x_4^2 x_5^3, & \mathcal{V}_{21,595} &= x_1^7 x_2^8 x_3 x_4^3 x_5^2, & \mathcal{V}_{21,596} &= x_1^7 x_2^8 x_3^2 x_4 x_5^2,
\end{aligned}$$

$\mathcal{B}_5^+(3, 3, 3)$ is the set of 70 admissible monomials: $\mathcal{V}_{21,t}$, $597 \leq t \leq 666$

$$\begin{aligned}
\mathcal{V}_{21,597} &= x_1 x_2^2 x_3^4 x_4^7 x_5^7, & \mathcal{V}_{21,598} &= x_1 x_2^2 x_3^7 x_4^4 x_5^7, & \mathcal{V}_{21,599} &= x_1 x_2^2 x_3^7 x_4^7 x_5^4, & \mathcal{V}_{21,600} &= x_1 x_2^7 x_3^2 x_4^4 x_5^7, \\
\mathcal{V}_{21,601} &= x_1 x_2^7 x_3^2 x_4^4 x_5^4, & \mathcal{V}_{21,602} &= x_1 x_2^7 x_3^2 x_4^4 x_5^4, & \mathcal{V}_{21,603} &= x_1^7 x_2 x_3^2 x_4^4 x_5^7, & \mathcal{V}_{21,604} &= x_1^7 x_2 x_3^2 x_4^4 x_5^4, \\
\mathcal{V}_{21,605} &= x_1^7 x_2 x_3^7 x_4^2 x_5^4, & \mathcal{V}_{21,606} &= x_1^7 x_2 x_3^7 x_4^2 x_5^4, & \mathcal{V}_{21,607} &= x_1 x_2^6 x_3 x_4^6 x_5^7, & \mathcal{V}_{21,608} &= x_1 x_2^6 x_3 x_4^7 x_5^6, \\
\mathcal{V}_{21,609} &= x_1 x_2^6 x_3 x_4^6 x_5^6, & \mathcal{V}_{21,610} &= x_1 x_2^6 x_3 x_4^6 x_5^7, & \mathcal{V}_{21,611} &= x_1 x_2^6 x_3 x_4^7 x_5^6, & \mathcal{V}_{21,612} &= x_1 x_2^7 x_3^6 x_4 x_5^6, \\
\mathcal{V}_{21,613} &= x_1 x_2^6 x_3^6 x_4^6 x_5^6, & \mathcal{V}_{21,614} &= x_1 x_2^7 x_3^6 x_4^6 x_5^6, & \mathcal{V}_{21,615} &= x_1 x_2^7 x_3^6 x_4^6 x_5^5, & \mathcal{V}_{21,616} &= x_1 x_2^7 x_3^6 x_4^6 x_5^6, \\
\mathcal{V}_{21,617} &= x_1 x_2^5 x_3^5 x_4^7 x_5^7, & \mathcal{V}_{21,618} &= x_1 x_2^5 x_3^5 x_4^7 x_5^5, & \mathcal{V}_{21,619} &= x_1 x_2^5 x_3^7 x_4^5 x_5^6, & \mathcal{V}_{21,620} &= x_1 x_2^7 x_3^5 x_4^5 x_5^6, \\
\mathcal{V}_{21,621} &= x_1^7 x_2 x_3^2 x_4^5 x_5^6, & \mathcal{V}_{21,622} &= x_1 x_2^3 x_4^6 x_5^7, & \mathcal{V}_{21,623} &= x_1 x_2^3 x_4^7 x_5^6, & \mathcal{V}_{21,624} &= x_1 x_2^3 x_4^7 x_5^6, \\
\mathcal{V}_{21,625} &= x_1 x_2^7 x_3^3 x_4^6 x_5^6, & \mathcal{V}_{21,626} &= x_1^3 x_2 x_3^4 x_6^7, & \mathcal{V}_{21,627} &= x_1^3 x_2 x_3^4 x_7^6, & \mathcal{V}_{21,628} &= x_1^3 x_2 x_3^7 x_4^4 x_5^6, \\
\mathcal{V}_{21,629} &= x_1^3 x_2^7 x_3 x_4^6 x_5^6, & \mathcal{V}_{21,630} &= x_1^7 x_2 x_3^3 x_4^6 x_5^6, & \mathcal{V}_{21,631} &= x_1^7 x_2^3 x_3 x_4^6 x_5^6, & \mathcal{V}_{21,632} &= x_1 x_2^3 x_6^6 x_4^4 x_5^7, \\
\mathcal{V}_{21,633} &= x_1 x_2^3 x_6^6 x_4^4 x_5^7, & \mathcal{V}_{21,634} &= x_1 x_2^3 x_7^6 x_4^4 x_5^7, & \mathcal{V}_{21,635} &= x_1 x_2^7 x_3^3 x_4^6 x_5^4, & \mathcal{V}_{21,636} &= x_1^3 x_2 x_3^6 x_4^4 x_5^7, \\
\mathcal{V}_{21,637} &= x_1^3 x_2 x_3^6 x_7^4 x_5^4, & \mathcal{V}_{21,638} &= x_1^3 x_2 x_3^7 x_4^6 x_5^4, & \mathcal{V}_{21,639} &= x_1^3 x_2^7 x_3 x_4^6 x_5^4, & \mathcal{V}_{21,640} &= x_1^7 x_2 x_3^3 x_4^4 x_5^4, \\
\mathcal{V}_{21,641} &= x_1^7 x_2^3 x_3 x_4^4 x_5^4, & \mathcal{V}_{21,642} &= x_1^3 x_2^5 x_3^2 x_4^7 x_5^4, & \mathcal{V}_{21,643} &= x_1^3 x_2^5 x_3^2 x_4^7 x_5^4, & \mathcal{V}_{21,644} &= x_1^3 x_2^5 x_3^2 x_4^7 x_5^4, \\
\mathcal{V}_{21,645} &= x_1^3 x_2^7 x_3^5 x_4^2 x_5^4, & \mathcal{V}_{21,646} &= x_1^7 x_2^3 x_3^5 x_4^2 x_5^4, & \mathcal{V}_{21,647} &= x_1^3 x_2^3 x_3^4 x_4^4 x_5^7, & \mathcal{V}_{21,648} &= x_1^3 x_2^3 x_3^4 x_4^7 x_5^4, \\
\mathcal{V}_{21,649} &= x_1^3 x_2^3 x_3^7 x_4^4 x_5^4, & \mathcal{V}_{21,650} &= x_1^3 x_2^7 x_3^3 x_4^4 x_5^4, & \mathcal{V}_{21,651} &= x_1^7 x_2^3 x_3^3 x_4^4 x_5^4, & \mathcal{V}_{21,652} &= x_1 x_2^6 x_3^3 x_4^5 x_5^6, \\
\mathcal{V}_{21,653} &= x_1 x_2^3 x_3^6 x_5^6 x_6^6, & \mathcal{V}_{21,654} &= x_1 x_2^3 x_3^5 x_6^6 x_5^6, & \mathcal{V}_{21,655} &= x_1^3 x_2 x_3^6 x_5^6 x_6^6, & \mathcal{V}_{21,656} &= x_1^3 x_2 x_3^5 x_6^6 x_5^6, \\
\mathcal{V}_{21,657} &= x_1^3 x_2^5 x_3 x_4^6 x_5^6, & \mathcal{V}_{21,658} &= x_1^3 x_2^5 x_3^6 x_4 x_5^6, & \mathcal{V}_{21,659} &= x_1^3 x_2^5 x_3^2 x_4^5 x_5^6, & \mathcal{V}_{21,660} &= x_1^3 x_2^5 x_3^3 x_4^4 x_5^6, \\
\mathcal{V}_{21,661} &= x_1^3 x_2^3 x_3^5 x_4^6 x_5^6, & \mathcal{V}_{21,662} &= x_1^3 x_2^3 x_3^4 x_4^5 x_5^6, & \mathcal{V}_{21,663} &= x_1^3 x_2^3 x_3^5 x_4^6 x_5^4, & \mathcal{V}_{21,664} &= x_1^3 x_2^4 x_3^3 x_4^5 x_5^6, \\
\mathcal{V}_{21,665} &= x_1^3 x_2^5 x_3^3 x_4^5 x_5^4, & \mathcal{V}_{21,666} &= x_1^3 x_2^5 x_3^3 x_4^5 x_5^4.
\end{aligned}$$

6.4. $\mathcal{A}_2$ -generators for $\mathcal{P}_5^0$ in degree 22

According to the results in Sect. 3.4.2, we have

$$\mathcal{B}_5^0(22) = \overline{\Phi}^0(\mathcal{B}_4(\omega_{(1)})) \cup \overline{\Phi}^0(\mathcal{B}_4(\omega_{(4)})) \cup \overline{\Phi}^0(\mathcal{B}_4(\omega_{(5)})) = \{\mathcal{V}_{22,t} : 1 \leq t \leq 460\},$$

where $\omega_{(1)} = (2, 2, 2, 1)$, $\omega_{(4)} = (4, 3, 1, 1)$, $\omega_{(5)} = (4, 3, 3)$, and the monomials $\mathcal{V}_{22,t} : 1 \leq t \leq 460$, are determined as follows:

47

48

$$\begin{aligned}
\mathcal{V}_{22,393} &= x_2^3 x_3^7 x_4^9 x_5^3, & \mathcal{V}_{22,394} &= x_2^7 x_3^3 x_4^3 x_5^9, & \mathcal{V}_{22,395} &= x_2^7 x_3^3 x_4^9 x_5^3, & \mathcal{V}_{22,396} &= x_2^7 x_3^9 x_4^3 x_5^3, \\
\mathcal{V}_{22,397} &= x_1^3 x_3^3 x_4^7 x_5^9, & \mathcal{V}_{22,398} &= x_1^3 x_3^7 x_4^3 x_5^9, & \mathcal{V}_{22,399} &= x_1^3 x_3^7 x_4^9 x_5^3, & \mathcal{V}_{22,400} &= x_1^3 x_3^3 x_4^7 x_5^9, \\
\mathcal{V}_{22,401} &= x_1^3 x_3^3 x_4^7 x_5^9, & \mathcal{V}_{22,402} &= x_1^3 x_3^3 x_4^7 x_5^9, & \mathcal{V}_{22,403} &= x_1^3 x_2^7 x_3^3 x_5^9, & \mathcal{V}_{22,404} &= x_1^3 x_2^7 x_3^9 x_5^3, \\
\mathcal{V}_{22,405} &= x_1^3 x_2^7 x_3^3 x_5^9, & \mathcal{V}_{22,406} &= x_1^3 x_2^7 x_3^3 x_5^9, & \mathcal{V}_{22,407} &= x_1^3 x_2^7 x_3^3 x_5^9, & \mathcal{V}_{22,408} &= x_1^3 x_2^7 x_3^9 x_5^3, \\
\mathcal{V}_{22,409} &= x_1^7 x_3^3 x_4^3 x_5^9, & \mathcal{V}_{22,410} &= x_1^7 x_3^3 x_4^3 x_5^9, & \mathcal{V}_{22,411} &= x_1^7 x_3^3 x_4^3 x_5^9, & \mathcal{V}_{22,412} &= x_1^7 x_2^3 x_3^3 x_5^9, \\
\mathcal{V}_{22,413} &= x_1^7 x_2^3 x_3^3 x_5^9, & \mathcal{V}_{22,414} &= x_1^7 x_2^3 x_3^3 x_5^9, & \mathcal{V}_{22,415} &= x_1^7 x_2^3 x_3^3 x_5^9, & \mathcal{V}_{22,416} &= x_1^7 x_2^3 x_3^3 x_5^9, \\
\mathcal{V}_{22,417} &= x_1^7 x_2^3 x_3^9 x_4^3, & \mathcal{V}_{22,418} &= x_1^7 x_2^3 x_3^9 x_4^3, & \mathcal{V}_{22,419} &= x_1^7 x_2^3 x_3^9 x_4^3, & \mathcal{V}_{22,420} &= x_1^7 x_2^3 x_3^9 x_4^3, \\
\mathcal{V}_{22,421} &= x_2^3 x_3^5 x_4^{10} x_5^4, & \mathcal{V}_{22,422} &= x_1^3 x_3^5 x_4^{10} x_5^4, & \mathcal{V}_{22,423} &= x_1^3 x_3^5 x_4^{10} x_5^4, & \mathcal{V}_{22,424} &= x_1^3 x_3^5 x_4^{10} x_5^4, \\
\mathcal{V}_{22,425} &= x_1^3 x_3^5 x_4^{10} x_5^4, & \mathcal{V}_{22,426} &= x_2^3 x_3^5 x_4^6 x_5^8, & \mathcal{V}_{22,427} &= x_1^3 x_3^5 x_4^6 x_5^8, & \mathcal{V}_{22,428} &= x_1^3 x_3^5 x_4^6 x_5^8, \\
\mathcal{V}_{22,429} &= x_1^3 x_2^5 x_3^6 x_5^8, & \mathcal{V}_{22,430} &= x_1^3 x_2^5 x_3^6 x_5^8, & \mathcal{V}_{22,431} &= x_2^3 x_3^5 x_4^7 x_5^7, & \mathcal{V}_{22,432} &= x_2^3 x_3^5 x_4^7 x_5^7, \\
\mathcal{V}_{22,433} &= x_2^3 x_3^7 x_4^7 x_5^5, & \mathcal{V}_{22,434} &= x_2^3 x_3^7 x_4^7 x_5^5, & \mathcal{V}_{22,435} &= x_2^3 x_3^7 x_4^7 x_5^5, & \mathcal{V}_{22,436} &= x_2^3 x_3^7 x_4^7 x_5^5, \\
\mathcal{V}_{22,437} &= x_1^3 x_3^5 x_4^7 x_5^7, & \mathcal{V}_{22,438} &= x_1^3 x_3^5 x_4^7 x_5^7, & \mathcal{V}_{22,439} &= x_1^3 x_3^5 x_4^7 x_5^7, & \mathcal{V}_{22,440} &= x_1^3 x_2^5 x_4^7 x_5^7, \\
\mathcal{V}_{22,441} &= x_1^3 x_2^5 x_3^7 x_5^7, & \mathcal{V}_{22,442} &= x_1^3 x_2^5 x_3^7 x_5^7, & \mathcal{V}_{22,443} &= x_1^3 x_2^5 x_3^7 x_5^7, & \mathcal{V}_{22,444} &= x_1^3 x_2^5 x_3^7 x_5^7, \\
\mathcal{V}_{22,445} &= x_1^3 x_2^7 x_3^5 x_5^7, & \mathcal{V}_{22,446} &= x_1^3 x_2^7 x_3^5 x_5^7, & \mathcal{V}_{22,447} &= x_1^3 x_2^7 x_3^5 x_5^7, & \mathcal{V}_{22,448} &= x_1^3 x_2^7 x_3^5 x_5^7, \\
\mathcal{V}_{22,449} &= x_1^3 x_3^3 x_4^5 x_5^7, & \mathcal{V}_{22,450} &= x_1^3 x_3^3 x_4^5 x_5^7, & \mathcal{V}_{22,451} &= x_1^3 x_3^3 x_4^5 x_5^7, & \mathcal{V}_{22,452} &= x_1^3 x_3^3 x_4^5 x_5^7, \\
\mathcal{V}_{22,453} &= x_1^3 x_2^3 x_4^5 x_5^7, & \mathcal{V}_{22,454} &= x_1^3 x_2^3 x_4^5 x_5^7, & \mathcal{V}_{22,455} &= x_1^3 x_2^3 x_4^5 x_5^7, & \mathcal{V}_{22,456} &= x_1^3 x_2^3 x_4^5 x_5^7, \\
\mathcal{V}_{22,457} &= x_1^7 x_2^3 x_3^7 x_5^5, & \mathcal{V}_{22,458} &= x_1^7 x_2^3 x_3^7 x_5^5, & \mathcal{V}_{22,459} &= x_1^7 x_2^3 x_3^7 x_5^5, & \mathcal{V}_{22,460} &= x_1^7 x_2^3 x_3^7 x_5^5.
\end{aligned}$$

6.5. $\mathcal{A}_2$ -generators for $\mathcal{P}_5^+$ in degree 22

By Propositions 3.4.11 and 3.4.15, we have $\mathcal{B}_5^+(22) = \bigcup_{1 \leq j \leq 5} \mathcal{B}_5^+(\omega_{(j)})$. Here $\omega_{(1)} = (2, 2, 2, 1)$, $\omega_{(2)} = (2, 4, 1, 1)$, $\omega_{(3)} = (2, 4, 3)$, $\omega_{(4)} = (4, 3, 1, 1)$, $\omega_{(5)} = (4, 3, 3)$.

$\mathcal{B}_5^+(\omega_{(1)}) = \overline{\Phi}^+(\mathcal{B}_4((\omega_{(1)}))) \cup \mathcal{B}^+(5, \omega_{(1)}) \cup \mathcal{D} = \{\mathcal{V}_{22,t} : 461 \leq t \leq 510\}$, where the monomials $\mathcal{V}_{22,t} : 461 \leq t \leq 510$, are listed as follows:

$$\begin{aligned}
\mathcal{V}_{22,461} &= x_1 x_2 x_3^2 x_4^4 x_5^{14}, & \mathcal{V}_{22,462} &= x_1 x_2 x_3^2 x_4^4 x_5^{14}, & \mathcal{V}_{22,463} &= x_1 x_2 x_3^{14} x_4^2 x_5^4, & \mathcal{V}_{22,464} &= x_1 x_2^2 x_3 x_4^4 x_5^{14}, \\
\mathcal{V}_{22,465} &= x_1 x_2^2 x_3 x_4^{14} x_5^4, & \mathcal{V}_{22,466} &= x_1 x_2^2 x_3 x_4^{14} x_5^4, & \mathcal{V}_{22,467} &= x_1 x_2^{14} x_3 x_4^2 x_5^4, & \mathcal{V}_{22,468} &= x_1 x_2^2 x_3^2 x_4^4 x_5^{12}, \\
\mathcal{V}_{22,469} &= x_1 x_2 x_3^2 x_4^{12} x_5^6, & \mathcal{V}_{22,470} &= x_1 x_2 x_3^6 x_4^2 x_5^{12}, & \mathcal{V}_{22,471} &= x_1 x_2^2 x_3 x_4^6 x_5^{12}, & \mathcal{V}_{22,472} &= x_1 x_2^2 x_3 x_4^{12} x_5^6, \\
\mathcal{V}_{22,473} &= x_1 x_2^2 x_3^{12} x_4 x_5^6, & \mathcal{V}_{22,474} &= x_1 x_2^6 x_3 x_4^2 x_5^{12}, & \mathcal{V}_{22,475} &= x_1 x_2^2 x_3^{13} x_4^2 x_5^4, & \mathcal{V}_{22,476} &= x_1 x_2^2 x_3^5 x_4^2 x_5^{12}, \\
\mathcal{V}_{22,477} &= x_1 x_3^3 x_4^2 x_5^{12}, & \mathcal{V}_{22,478} &= x_1 x_2^3 x_3^{12} x_4^2 x_5^4, & \mathcal{V}_{22,479} &= x_1^3 x_2 x_3^4 x_4^2 x_5^{12}, & \mathcal{V}_{22,480} &= x_1^3 x_2 x_3^{12} x_4^2 x_5^4, \\
\mathcal{V}_{22,481} &= x_1 x_2^2 x_3^5 x_4^{10} x_5^4, & \mathcal{V}_{22,482} &= x_1 x_2^2 x_3^4 x_4^7 x_5^8, & \mathcal{V}_{22,483} &= x_1 x_2^2 x_3^4 x_4^8 x_5^7, & \mathcal{V}_{22,484} &= x_1 x_2^2 x_3^7 x_4^4 x_5^8, \\
\mathcal{V}_{22,485} &= x_1 x_2^2 x_3^4 x_4^8 x_5^8, & \mathcal{V}_{22,486} &= x_1 x_2^2 x_3^4 x_4^8 x_5^8, & \mathcal{V}_{22,487} &= x_1 x_2^2 x_3^5 x_4^6 x_5^8, & \mathcal{V}_{22,488} &= x_1 x_3^3 x_4^2 x_4^{10} x_5^4, \\
\mathcal{V}_{22,489} &= x_1^3 x_2 x_3^4 x_4^{10} x_5^4, & \mathcal{V}_{22,490} &= x_1 x_2^2 x_3^4 x_4^6 x_5^8, & \mathcal{V}_{22,491} &= x_1^3 x_2 x_3^4 x_4^6 x_5^8, & \mathcal{V}_{22,492} &= x_1 x_2^2 x_3^6 x_4^6 x_5^8, \\
\mathcal{V}_{22,493} &= x_1 x_2 x_3^2 x_4^{10} x_5^4, & \mathcal{V}_{22,494} &= x_1 x_2^2 x_3^3 x_4^4 x_5^{12}, & \mathcal{V}_{22,495} &= x_1 x_2^2 x_3^3 x_4^{12} x_5^4, & \mathcal{V}_{22,496} &= x_1 x_2^2 x_3^4 x_4^9 x_5^6, \\
\mathcal{V}_{22,497} &= x_1 x_2^2 x_3^5 x_4^8 x_5^6, & \mathcal{V}_{22,498} &= x_1 x_2^3 x_3^2 x_4^4 x_5^{12}, & \mathcal{V}_{22,499} &= x_1 x_2^3 x_3^2 x_4^{12} x_5^4, & \mathcal{V}_{22,500} &= x_1 x_3^3 x_4^2 x_4^8 x_5^6, \\
\mathcal{V}_{22,501} &= x_1 x_2^3 x_3^6 x_4^8 x_5^8, & \mathcal{V}_{22,502} &= x_1 x_2^3 x_3^6 x_4^8 x_5^8, & \mathcal{V}_{22,503} &= x_1^3 x_2 x_3^2 x_4^4 x_5^{12}, & \mathcal{V}_{22,504} &= x_1^3 x_2 x_3^2 x_4^{12} x_5^4, \\
\mathcal{V}_{22,505} &= x_1^3 x_2 x_3^4 x_4^8 x_5^6, & \mathcal{V}_{22,506} &= x_1^3 x_2 x_3^6 x_4^4 x_5^8, & \mathcal{V}_{22,507} &= x_1^3 x_2 x_3^6 x_4^8 x_5^4, & \mathcal{V}_{22,508} &= x_1^3 x_2^5 x_3^2 x_4^8 x_5^8, \\
\mathcal{V}_{22,509} &= x_1^3 x_2^5 x_3^2 x_4^8 x_5^8, & \mathcal{V}_{22,510} &= x_1^3 x_2^5 x_3^2 x_4^8 x_5^8.
\end{aligned}$$

$\mathcal{B}_5^+(\omega_{(2)}) = \mathcal{B}^+(5, \omega_{(2)}) \cup \mathcal{E} = \{\mathcal{V}_{22,t} : 511 \leq t \leq 535\}$, where the monomials $\mathcal{V}_{22,t} : 511 \leq t \leq 535$, are listed as follows:

$$\begin{aligned}
\mathcal{V}_{22,511} &= x_1 x_2^2 x_3^2 x_4^2 x_5^{15}, & \mathcal{V}_{22,512} &= x_1 x_2^2 x_3^2 x_4^2 x_5^{15}, & \mathcal{V}_{22,513} &= x_1 x_2^2 x_3^{15} x_4^2 x_5^2, & \mathcal{V}_{22,514} &= x_1 x_2^{15} x_3^2 x_4^2 x_5^2, \\
\mathcal{V}_{22,515} &= x_1^{15} x_2 x_3^2 x_4^2 x_5^2, & \mathcal{V}_{22,516} &= x_1 x_2^2 x_3^2 x_4^3 x_5^{14}, & \mathcal{V}_{22,517} &= x_1 x_2^2 x_3^3 x_4^2 x_5^{14}, & \mathcal{V}_{22,518} &= x_1 x_2^2 x_3^3 x_4^{14} x_5^2, \\
\mathcal{V}_{22,519} &= x_1 x_2^3 x_3^2 x_4^2 x_5^{14}, & \mathcal{V}_{22,520} &= x_1 x_2^3 x_3^2 x_4^{14} x_5^2, & \mathcal{V}_{22,521} &= x_1 x_2^3 x_3^{14} x_4^2 x_5^2, & \mathcal{V}_{22,522} &= x_1^3 x_2 x_3^2 x_4^2 x_5^{14}, \\
\mathcal{V}_{22,523} &= x_1^3 x_2 x_3^2 x_4^{14} x_5^2, & \mathcal{V}_{22,524} &= x_1^3 x_2 x_3^{14} x_4^2 x_5^2, & \mathcal{V}_{22,525} &= x_1 x_2^2 x_3^3 x_4^6 x_5^{10}, & \mathcal{V}_{22,526} &= x_1 x_2^2 x_3^2 x_4^6 x_5^{10}, \\
\mathcal{V}_{22,527} &= x_1 x_2^3 x_3^2 x_4^{10} x_5^{10}, & \mathcal{V}_{22,528} &= x_1 x_2^3 x_3^2 x_4^{10} x_5^{10}, & \mathcal{V}_{22,529} &= x_1^3 x_2 x_3^2 x_4^6 x_5^{10}, & \mathcal{V}_{22,530} &= x_1^3 x_2 x_3^2 x_4^6 x_5^{10}, \\
\mathcal{V}_{22,531} &= x_1^3 x_2 x_3^2 x_4^{10} x_5^{10}, & \mathcal{V}_{22,532} &= x_1^3 x_2^{13} x_3^2 x_4^2 x_5^2, & \mathcal{V}_{22,533} &= x_1^3 x_2^5 x_3^2 x_4^2 x_5^{10}, & \mathcal{V}_{22,534} &= x_1^3 x_2^5 x_3^2 x_4^2 x_5^{10}, \\
\mathcal{V}_{22,535} &= x_1^3 x_2^5 x_3^2 x_4^2 x_5^{10}.
\end{aligned}$$

$\mathcal{B}_5^+(\omega_{(3)}) = \{\mathcal{V}_{22,t} : 536 \leq t \leq 540\}$, where the monomials $\mathcal{V}_{22,t} : 536 \leq t \leq 540$, are determined as follows:

$$\mathcal{V}_{536} = x_1 x_2^3 x_3^6 x_4^6 x_5^6, \mathcal{V}_{537} = x_1^3 x_2 x_3^6 x_4^6 x_5^6, \mathcal{V}_{538} = x_1^3 x_2^5 x_3^2 x_4^6 x_5^6,$$

$$\mathcal{V}_{539} = x_1^3 x_2^5 x_3^6 x_4^2 x_5^6, \mathcal{V}_{540} = x_1^3 x_2^5 x_3^6 x_4^6 x_5^2,$$

$\mathcal{B}_5^+(\omega_{(4)}) = \overline{\Phi}^+(\mathcal{B}_4(\omega_{(4)})) \cup \mathcal{B}^+(5, \omega_{(4)}) = \{\mathcal{V}_{22,t} : 541 \leq t \leq 840\}$, where the monomials $\mathcal{V}_{22,t} : 541 \leq t \leq 840$, are listed as follows:

$$\begin{aligned} \mathcal{V}_{22,541} &= x_1 x_2 x_3^2 x_4^3 x_5^{15}, & \mathcal{V}_{22,542} &= x_1 x_2 x_3^2 x_4^{15} x_5^3, & \mathcal{V}_{22,543} &= x_1 x_2 x_3^3 x_4^2 x_5^{15}, & \mathcal{V}_{22,544} &= x_1 x_2 x_3^3 x_4^{15} x_5^2, \\ \mathcal{V}_{22,545} &= x_1 x_2 x_3^{15} x_4^2 x_5^3, & \mathcal{V}_{22,546} &= x_1 x_2 x_3^{15} x_4^3 x_5^2, & \mathcal{V}_{22,547} &= x_1 x_2^2 x_3 x_4^3 x_5^{15}, & \mathcal{V}_{22,548} &= x_1 x_2^2 x_3 x_4^{15} x_5^3, \\ \mathcal{V}_{22,549} &= x_1 x_2^2 x_3^3 x_4 x_5^{15}, & \mathcal{V}_{22,550} &= x_1 x_2^2 x_3^3 x_4^{15} x_5, & \mathcal{V}_{22,551} &= x_1 x_2^2 x_3^{15} x_4 x_5^3, & \mathcal{V}_{22,552} &= x_1 x_2^2 x_3^{15} x_4^3 x_5, \\ \mathcal{V}_{22,553} &= x_1 x_2^3 x_3 x_4^2 x_5^{15}, & \mathcal{V}_{22,554} &= x_1 x_2^3 x_3 x_4^{15} x_5^2, & \mathcal{V}_{22,555} &= x_1 x_2^3 x_3^2 x_4 x_5^{15}, & \mathcal{V}_{22,556} &= x_1 x_2^3 x_3^2 x_4^{15} x_5, \\ \mathcal{V}_{22,557} &= x_1 x_2^3 x_3^{15} x_4 x_5^2, & \mathcal{V}_{22,558} &= x_1 x_2^3 x_3^{15} x_4^2 x_5, & \mathcal{V}_{22,559} &= x_1 x_2^{15} x_3 x_4^2 x_5^3, & \mathcal{V}_{22,560} &= x_1 x_2^{15} x_3 x_4^3 x_5^2, \\ \mathcal{V}_{22,561} &= x_1 x_2^{15} x_3^2 x_4 x_5^3, & \mathcal{V}_{22,562} &= x_1 x_2^{15} x_3^2 x_4^2 x_5, & \mathcal{V}_{22,563} &= x_1 x_2^{15} x_3^3 x_4 x_5^2, & \mathcal{V}_{22,564} &= x_1 x_2^{15} x_3^3 x_4^2 x_5, \\ \mathcal{V}_{22,565} &= x_1^3 x_2 x_3 x_4^2 x_5^{15}, & \mathcal{V}_{22,566} &= x_1^3 x_2 x_3 x_4^{15} x_5^2, & \mathcal{V}_{22,567} &= x_1^3 x_2 x_3^2 x_4 x_5^{15}, & \mathcal{V}_{22,568} &= x_1^3 x_2 x_3^2 x_4^{15} x_5, \\ \mathcal{V}_{22,569} &= x_1^3 x_2 x_3^{15} x_4 x_5^2, & \mathcal{V}_{22,570} &= x_1^3 x_2 x_3^{15} x_4^2 x_5, & \mathcal{V}_{22,571} &= x_1^3 x_2^{15} x_3 x_4 x_5^2, & \mathcal{V}_{22,572} &= x_1^3 x_2^{15} x_3 x_4^2 x_5, \\ \mathcal{V}_{22,573} &= x_1^{15} x_2 x_3 x_4^2 x_5^3, & \mathcal{V}_{22,574} &= x_1^{15} x_2 x_3 x_4^3 x_5^2, & \mathcal{V}_{22,575} &= x_1^{15} x_2 x_3^2 x_4 x_5^3, & \mathcal{V}_{22,576} &= x_1^{15} x_2 x_3^2 x_4^2 x_5, \\ \mathcal{V}_{22,577} &= x_1^{15} x_2 x_3^3 x_4 x_5^2, & \mathcal{V}_{22,578} &= x_1^{15} x_2 x_3^3 x_4^2 x_5, & \mathcal{V}_{22,579} &= x_1^{15} x_2^2 x_3 x_4 x_5^2, & \mathcal{V}_{22,580} &= x_1^{15} x_2^2 x_3 x_4^2 x_5, \\ \mathcal{V}_{22,581} &= x_1 x_2 x_3^2 x_4^7 x_5^{11}, & \mathcal{V}_{22,582} &= x_1 x_2 x_3^2 x_4^{11} x_5^7, & \mathcal{V}_{22,583} &= x_1 x_2 x_3^3 x_4^7 x_5^{11}, & \mathcal{V}_{22,584} &= x_1 x_2 x_3^3 x_4^{11} x_5^7, \\ \mathcal{V}_{22,585} &= x_1 x_2^2 x_3^2 x_4^7 x_5^{11}, & \mathcal{V}_{22,586} &= x_1 x_2^2 x_3^2 x_4^{11} x_5^7, & \mathcal{V}_{22,587} &= x_1 x_2^2 x_3^3 x_4^7 x_5^{11}, & \mathcal{V}_{22,588} &= x_1 x_2^2 x_3^3 x_4^{11} x_5^7, \\ \mathcal{V}_{22,589} &= x_1 x_2^2 x_3^3 x_4^7 x_5^{11}, & \mathcal{V}_{22,590} &= x_1 x_2^2 x_3^3 x_4^{11} x_5^7, & \mathcal{V}_{22,591} &= x_1 x_2^2 x_3^{11} x_4 x_5^7, & \mathcal{V}_{22,592} &= x_1 x_2^2 x_3^{11} x_4^2 x_5, \\ \mathcal{V}_{22,593} &= x_1^7 x_2 x_3 x_4^2 x_5^{11}, & \mathcal{V}_{22,594} &= x_1^7 x_2 x_3 x_4^{11} x_5^2, & \mathcal{V}_{22,595} &= x_1^7 x_2 x_3^2 x_4 x_5^{11}, & \mathcal{V}_{22,596} &= x_1^7 x_2 x_3^2 x_4^{11} x_5, \\ \mathcal{V}_{22,597} &= x_1^7 x_2 x_3^{11} x_4 x_5^2, & \mathcal{V}_{22,598} &= x_1^7 x_2 x_3^{11} x_4^2 x_5, & \mathcal{V}_{22,599} &= x_1^7 x_2^{11} x_3 x_4 x_5^2, & \mathcal{V}_{22,600} &= x_1^7 x_2^{11} x_3 x_4^2 x_5, \\ \mathcal{V}_{22,601} &= x_1 x_2 x_3^3 x_4^{14} x_5^3, & \mathcal{V}_{22,602} &= x_1 x_2 x_3^3 x_4^{14} x_5^3, & \mathcal{V}_{22,603} &= x_1 x_2 x_3^{14} x_4^3 x_5^3, & \mathcal{V}_{22,604} &= x_1 x_3^3 x_3 x_4^3 x_5^{14}, \\ \mathcal{V}_{22,605} &= x_1 x_3^3 x_3 x_4^{14} x_5^3, & \mathcal{V}_{22,606} &= x_1 x_3^3 x_3^2 x_4 x_5^{14}, & \mathcal{V}_{22,607} &= x_1 x_2^2 x_3^3 x_4^{14} x_5, & \mathcal{V}_{22,608} &= x_1 x_2^2 x_3^3 x_4^3 x_5^{14}, \\ \mathcal{V}_{22,609} &= x_1 x_2^2 x_3^{14} x_3 x_5^3, & \mathcal{V}_{22,610} &= x_1 x_2^{14} x_3 x_3^2 x_5^3, & \mathcal{V}_{22,611} &= x_1 x_2^{14} x_3^3 x_4 x_5^3, & \mathcal{V}_{22,612} &= x_1 x_2^{14} x_3^3 x_3 x_5^3, \\ \mathcal{V}_{22,613} &= x_1^3 x_2 x_3 x_3^3 x_4^{14} x_5, & \mathcal{V}_{22,614} &= x_1^3 x_2 x_3 x_3^3 x_4^{14} x_5, & \mathcal{V}_{22,615} &= x_1^3 x_2 x_3^3 x_3 x_4^{14} x_5, & \mathcal{V}_{22,616} &= x_1^3 x_2 x_3^3 x_3^2 x_4^{14} x_5, \\ \mathcal{V}_{22,617} &= x_1^3 x_2 x_3^{14} x_3 x_4 x_5^3, & \mathcal{V}_{22,618} &= x_1^3 x_2 x_3^{14} x_3^2 x_4 x_5, & \mathcal{V}_{22,619} &= x_1^3 x_2^3 x_3 x_3 x_4^{14} x_5, & \mathcal{V}_{22,620} &= x_1^3 x_2^3 x_3 x_3^2 x_4^{14} x_5, \\ \mathcal{V}_{22,621} &= x_1 x_2 x_3^3 x_3^6 x_4 x_5^{11}, & \mathcal{V}_{22,622} &= x_1 x_2 x_3^3 x_3^6 x_4^{11} x_5, & \mathcal{V}_{22,623} &= x_1 x_2 x_3^6 x_3^{11} x_4 x_5^3, & \mathcal{V}_{22,624} &= x_1 x_2^6 x_3 x_3^6 x_4 x_5^{11}, \\ \mathcal{V}_{22,625} &= x_1 x_2^6 x_3^6 x_3 x_4 x_5^{11}, & \mathcal{V}_{22,626} &= x_1 x_2^6 x_3^6 x_3^{11} x_4 x_5, & \mathcal{V}_{22,627} &= x_1 x_2^6 x_3^3 x_3^4 x_4 x_5^{11}, & \mathcal{V}_{22,628} &= x_1 x_2^6 x_3^3 x_3^4 x_4^{11} x_5^3, \\ \mathcal{V}_{22,629} &= x_1 x_2^6 x_3^3 x_3^4 x_4 x_5^{11}, & \mathcal{V}_{22,630} &= x_1 x_2^6 x_3^3 x_3^{11} x_4 x_5, & \mathcal{V}_{22,631} &= x_1 x_2^6 x_3^{11} x_3 x_4 x_5^3, & \mathcal{V}_{22,632} &= x_1 x_2^6 x_3^{11} x_3^2 x_4 x_5^3, \\ \mathcal{V}_{22,633} &= x_1^3 x_2 x_3 x_3^6 x_4 x_5^{11}, & \mathcal{V}_{22,634} &= x_1^3 x_2 x_3^6 x_3 x_4 x_5^{11}, & \mathcal{V}_{22,635} &= x_1^3 x_2 x_3^6 x_3^2 x_4 x_5^{11}, & \mathcal{V}_{22,636} &= x_1 x_2 x_3^6 x_3^2 x_4^2 x_5^{10}, \\ \mathcal{V}_{22,637} &= x_1 x_2 x_3^7 x_3^3 x_4 x_5^{10}, & \mathcal{V}_{22,638} &= x_1 x_2 x_3^7 x_3^{10} x_4 x_5^3, & \mathcal{V}_{22,639} &= x_1 x_2^3 x_3^3 x_3^4 x_4 x_5^{10}, & \mathcal{V}_{22,640} &= x_1 x_2^3 x_3^3 x_3^4 x_4 x_5^{10}, \\ \mathcal{V}_{22,641} &= x_1 x_2^3 x_3^7 x_3^{10} x_4 x_5, & \mathcal{V}_{22,642} &= x_1 x_2^7 x_3 x_3^3 x_4 x_5^{10}, & \mathcal{V}_{22,643} &= x_1 x_2^7 x_3 x_3^{10} x_4 x_5^3, & \mathcal{V}_{22,644} &= x_1 x_2^7 x_3^2 x_3 x_4 x_5^{10}, \\ \mathcal{V}_{22,645} &= x_1 x_2^7 x_3^3 x_3^{10} x_4 x_5, & \mathcal{V}_{22,646} &= x_1 x_2^7 x_3^{10} x_3 x_4 x_5^3, & \mathcal{V}_{22,647} &= x_1 x_2^7 x_3^{10} x_3^2 x_4 x_5, & \mathcal{V}_{22,648} &= x_1^3 x_2 x_3 x_3^7 x_4 x_5^{10}, \\ \mathcal{V}_{22,649} &= x_1^3 x_2 x_3^7 x_3 x_4 x_5^{10}, & \mathcal{V}_{22,650} &= x_1^3 x_2 x_3^7 x_3^{10} x_4 x_5, & \mathcal{V}_{22,651} &= x_1^3 x_2^2 x_3 x_3 x_4 x_5^{10}, & \mathcal{V}_{22,652} &= x_1^3 x_2^2 x_3 x_3^2 x_4 x_5^{10}, \\ \mathcal{V}_{22,653} &= x_1^7 x_2 x_3 x_3^3 x_4 x_5^{10}, & \mathcal{V}_{22,654} &= x_1^7 x_2 x_3 x_3^{10} x_4 x_5^3, & \mathcal{V}_{22,655} &= x_1^7 x_2 x_3^3 x_3 x_4 x_5^{10}, & \mathcal{V}_{22,656} &= x_1^7 x_2 x_3^3 x_3^2 x_4 x_5^{10}, \\ \mathcal{V}_{22,657} &= x_1^7 x_2 x_3^{10} x_3 x_4 x_5^3, & \mathcal{V}_{22,658} &= x_1^7 x_2 x_3^{10} x_3^2 x_4 x_5, & \mathcal{V}_{22,659} &= x_1^7 x_2^2 x_3 x_3 x_4 x_5^{10}, & \mathcal{V}_{22,660} &= x_1^7 x_2^2 x_3^2 x_3 x_4 x_5^{10}, \\ \mathcal{V}_{22,661} &= x_1 x_2^2 x_3^3 x_3^4 x_5^{13}, & \mathcal{V}_{22,662} &= x_1 x_2^2 x_3^3 x_3^{13} x_4 x_5^3, & \mathcal{V}_{22,663} &= x_1 x_2^2 x_3^{13} x_3^2 x_4 x_5^3, & \mathcal{V}_{22,664} &= x_1 x_2^2 x_3^{13} x_3^2 x_4^2 x_5^3, \\ \mathcal{V}_{22,665} &= x_1 x_2^2 x_3^3 x_3^{13} x_4 x_5^3, & \mathcal{V}_{22,666} &= x_1 x_2^2 x_3^3 x_3^{13} x_4^2 x_5^3, & \mathcal{V}_{22,667} &= x_1 x_2^2 x_3^3 x_3^{13} x_4^2 x_5^2, & \mathcal{V}_{22,668} &= x_1 x_2^2 x_3^3 x_3^{13} x_4^2 x_5^2, \\ \mathcal{V}_{22,669} &= x_1 x_2^2 x_3^{13} x_3^2 x_4 x_5^3, & \mathcal{V}_{22,670} &= x_1^3 x_2 x_3^2 x_3^3 x_4 x_5^{13}, & \mathcal{V}_{22,671} &= x_1^3 x_2 x_3^2 x_3^{13} x_4 x_5^3, & \mathcal{V}_{22,672} &= x_1^3 x_2 x_3^2 x_3^{13} x_4^2 x_5^3, \\ \mathcal{V}_{22,673} &= x_1^3 x_2 x_3^3 x_3^{13} x_4 x_5^2, & \mathcal{V}_{22,674} &= x_1^3 x_2 x_3^{13} x_3^2 x_4 x_5^3, & \mathcal{V}_{22,675} &= x_1^3 x_2 x_3^{13} x_3^2 x_4 x_5^2, & \mathcal{V}_{22,676} &= x_1^3 x_2 x_3^3 x_3^2 x_4 x_5^{13}, \\ \mathcal{V}_{22,677} &= x_1^3 x_2^2 x_3 x_3^{13} x_4 x_5^2, & \mathcal{V}_{22,678} &= x_1^3 x_2^2 x_3^{13} x_3 x_4 x_5^2, & \mathcal{V}_{22,679} &= x_1^3 x_2^2 x_3^{13} x_3^2 x_4 x_5^2, & \mathcal{V}_{22,680} &= x_1^3 x_2^2 x_3^{13} x_3^2 x_4 x_5^2, \\ \mathcal{V}_{22,681} &= x_1^3 x_2^2 x_3^{13} x_3^2 x_4 x_5^2, & \mathcal{V}_{22,682} &= x_1^3 x_2^2 x_3^{13} x_3^2 x_4 x_5^3, & \mathcal{V}_{22,683} &= x_1^3 x_2^{13} x_3^2 x_3^2 x_4 x_5, & \mathcal{V}_{22,684} &= x_1^3 x_2^{13} x_3^2 x_3^2 x_4 x_5^2, \\ \mathcal{V}_{22,685} &= x_1^3 x_2^{13} x_3^2 x_3^2 x_4 x_5, & \mathcal{V}_{22,686} &= x_1 x_2^2 x_3^3 x_3^5 x_4 x_5^{11}, & \mathcal{V}_{22,687} &= x_1 x_2^2 x_3^3 x_3^5 x_4^{11} x_5^2, & \mathcal{V}_{22,688} &= x_1 x_2^2 x_3^3 x_3^5 x_4^{11} x_5^2, \\ \mathcal{V}_{22,689} &= x_1 x_2^2 x_3^3 x_3^5 x_4^{11} x_5^2, & \mathcal{V}_{22,690} &= x_1 x_2^2 x_3^3 x_3^5 x_4^{11} x_5^2, & \mathcal{V}_{22,691} &= x_1 x_2^2 x_3^3 x_3^5 x_4^{11} x_5^2, & \mathcal{V}_{22,692} &= x_1 x_2^2 x_3^3 x_3^5 x_4^{11} x_5^2, \\ \mathcal{V}_{22,693} &= x_1^3 x_2 x_3^5 x_3^2 x_4 x_5^{11}, & \mathcal{V}_{22,694} &= x_1^3 x_2 x_3^5 x_3^2 x_4^{11} x_5^2, & \mathcal{V}_{22,695} &= x_1^3 x_2^5 x_3 x_3^2 x_4 x_5^{11}, & \mathcal{V}_{22,696} &= x_1^3 x_2^5 x_3 x_3^2 x_4^{11} x_5^2, \end{aligned}$$

$\mathcal{B}_5^+(\omega_{(5)}) = \overline{\Phi}^+(\mathcal{B}_4(\omega_{(5)})) \cup \mathcal{B}^+(5, \omega_{(5)}) = \{\mathcal{Y}_{22,t} : 841 \leq t \leq 965\}$, where the monomials $\mathcal{Y}_{22,t} : 841 \leq t \leq 965$, are listed as follows:

51

$$\begin{aligned}
\mathcal{Y}_{22,873} &= x_1 x_2^3 x_3^7 x_4^7 x_5^4, & \mathcal{Y}_{22,874} &= x_1 x_2^7 x_3^3 x_4^4 x_5^7, & \mathcal{Y}_{22,875} &= x_1 x_2^7 x_3^3 x_4^7 x_5^4, & \mathcal{Y}_{22,876} &= x_1 x_2^7 x_3^7 x_4^3 x_5^4, \\
\mathcal{Y}_{22,877} &= x_1^3 x_2 x_3^4 x_4^7 x_5^7, & \mathcal{Y}_{22,878} &= x_1^3 x_2 x_3^7 x_4^4 x_5^7, & \mathcal{Y}_{22,879} &= x_1^3 x_2 x_3^7 x_4^7 x_5^4, & \mathcal{Y}_{22,880} &= x_1^3 x_2^4 x_3 x_4^7 x_5^7, \\
\mathcal{Y}_{22,881} &= x_1^3 x_2^4 x_3^7 x_4^7 x_5^5, & \mathcal{Y}_{22,882} &= x_1^3 x_2^4 x_3^7 x_4^7 x_5^5, & \mathcal{Y}_{22,883} &= x_1^3 x_2^7 x_3 x_4^4 x_5^7, & \mathcal{Y}_{22,884} &= x_1^3 x_2^7 x_3 x_4^7 x_5^4, \\
\mathcal{Y}_{22,885} &= x_1^3 x_2^7 x_3^4 x_4^7 x_5^7, & \mathcal{Y}_{22,886} &= x_1^3 x_2^7 x_3^4 x_4^7 x_5^5, & \mathcal{Y}_{22,887} &= x_1^3 x_2^7 x_3^7 x_4^4 x_5^4, & \mathcal{Y}_{22,888} &= x_1^3 x_2^7 x_3^7 x_4^4 x_5^5, \\
\mathcal{Y}_{22,889} &= x_1^7 x_2 x_3^3 x_4^7 x_5^7, & \mathcal{Y}_{22,890} &= x_1^7 x_2 x_3^3 x_4^7 x_5^4, & \mathcal{Y}_{22,891} &= x_1^7 x_2 x_3^7 x_4^3 x_5^4, & \mathcal{Y}_{22,892} &= x_1^7 x_2^3 x_3 x_4^7 x_5^7, \\
\mathcal{Y}_{22,893} &= x_1^7 x_2^3 x_3 x_4^7 x_5^4, & \mathcal{Y}_{22,894} &= x_1^7 x_2^3 x_3^4 x_4^7 x_5^7, & \mathcal{Y}_{22,895} &= x_1^7 x_2^3 x_3^4 x_4^7 x_5^4, & \mathcal{Y}_{22,896} &= x_1^7 x_2^3 x_3^4 x_4^7 x_5^4, \\
\mathcal{Y}_{22,897} &= x_1^7 x_2^3 x_3^7 x_4^4 x_5^5, & \mathcal{Y}_{22,898} &= x_1^7 x_2^7 x_3 x_4^3 x_5^4, & \mathcal{Y}_{22,899} &= x_1^7 x_2^7 x_3^3 x_4^4 x_5^4, & \mathcal{Y}_{22,900} &= x_1^7 x_2^7 x_3^3 x_4^4 x_5^5, \\
\mathcal{Y}_{22,901} &= x_1 x_2^3 x_3^5 x_4^6 x_5^7, & \mathcal{Y}_{22,902} &= x_1 x_2^3 x_3^5 x_4^7 x_5^6, & \mathcal{Y}_{22,903} &= x_1 x_2^3 x_3^6 x_4^5 x_5^7, & \mathcal{Y}_{22,904} &= x_1 x_2^3 x_3^6 x_4^7 x_5^5, \\
\mathcal{Y}_{22,905} &= x_1 x_2^3 x_3^7 x_4^5 x_5^6, & \mathcal{Y}_{22,906} &= x_1 x_2^3 x_3^7 x_4^6 x_5^5, & \mathcal{Y}_{22,907} &= x_1 x_2^6 x_3^3 x_4^5 x_5^7, & \mathcal{Y}_{22,908} &= x_1 x_2^6 x_3^3 x_4^7 x_5^5, \\
\mathcal{Y}_{22,909} &= x_1 x_2^6 x_3^7 x_4^3 x_5^5, & \mathcal{Y}_{22,910} &= x_1 x_2^7 x_3^3 x_4^5 x_5^6, & \mathcal{Y}_{22,911} &= x_1 x_2^7 x_3^3 x_4^6 x_5^5, & \mathcal{Y}_{22,912} &= x_1 x_2^7 x_3^6 x_4^3 x_5^5, \\
\mathcal{Y}_{22,913} &= x_1^3 x_2 x_3^5 x_4^6 x_5^7, & \mathcal{Y}_{22,914} &= x_1^3 x_2 x_3^5 x_4^6 x_5^6, & \mathcal{Y}_{22,915} &= x_1^3 x_2 x_3^6 x_4^5 x_5^7, & \mathcal{Y}_{22,916} &= x_1^3 x_2 x_3^6 x_4^7 x_5^5, \\
\mathcal{Y}_{22,917} &= x_1^3 x_2 x_3^7 x_4^5 x_5^6, & \mathcal{Y}_{22,918} &= x_1^3 x_2 x_3^7 x_4^6 x_5^5, & \mathcal{Y}_{22,919} &= x_1^3 x_2^5 x_3 x_4^6 x_5^7, & \mathcal{Y}_{22,920} &= x_1^3 x_2^5 x_3 x_4^7 x_5^6, \\
\mathcal{Y}_{22,921} &= x_1^3 x_2^5 x_3^6 x_4^7 x_5^7, & \mathcal{Y}_{22,922} &= x_1^3 x_2^5 x_3^6 x_4^7 x_5^5, & \mathcal{Y}_{22,923} &= x_1^3 x_2^5 x_3^7 x_4^6 x_5^6, & \mathcal{Y}_{22,924} &= x_1^3 x_2^5 x_3^7 x_4^6 x_5^5, \\
\mathcal{Y}_{22,925} &= x_1^3 x_2^7 x_3 x_4^5 x_5^6, & \mathcal{Y}_{22,926} &= x_1^3 x_2^7 x_3 x_4^6 x_5^5, & \mathcal{Y}_{22,927} &= x_1^3 x_2^7 x_3^5 x_4^6 x_5^6, & \mathcal{Y}_{22,928} &= x_1^3 x_2^7 x_3^5 x_4^6 x_5^5, \\
\mathcal{Y}_{22,929} &= x_1^7 x_2 x_3^3 x_4^5 x_5^6, & \mathcal{Y}_{22,930} &= x_1^7 x_2 x_3^3 x_4^6 x_5^5, & \mathcal{Y}_{22,931} &= x_1^7 x_2 x_3^6 x_4^3 x_5^5, & \mathcal{Y}_{22,932} &= x_1^7 x_2^3 x_3 x_4^5 x_5^6, \\
\mathcal{Y}_{22,933} &= x_1^7 x_2^3 x_3 x_4^6 x_5^5, & \mathcal{Y}_{22,934} &= x_1^7 x_2^3 x_3^5 x_4^6 x_5^6, & \mathcal{Y}_{22,935} &= x_1^7 x_2^3 x_3^5 x_4^6 x_5^5, & \mathcal{Y}_{22,936} &= x_1^3 x_2^5 x_3^2 x_4^5 x_5^7, \\
\mathcal{Y}_{22,937} &= x_1^3 x_2^5 x_3^2 x_4^7 x_5^5, & \mathcal{Y}_{22,938} &= x_1^3 x_2^5 x_3^7 x_4^2 x_5^5, & \mathcal{Y}_{22,939} &= x_1^3 x_2^7 x_3^5 x_4^2 x_5^5, & \mathcal{Y}_{22,940} &= x_1^7 x_2^3 x_3^5 x_4^2 x_5^5, \\
\mathcal{Y}_{22,941} &= x_1^3 x_2^3 x_3^4 x_5^7 x_5^7, & \mathcal{Y}_{22,942} &= x_1^3 x_2^3 x_3^4 x_5^7 x_5^5, & \mathcal{Y}_{22,943} &= x_1^3 x_2^3 x_3^5 x_4^7 x_5^7, & \mathcal{Y}_{22,944} &= x_1^3 x_2^3 x_3^5 x_4^7 x_5^4, \\
\mathcal{Y}_{22,945} &= x_1^3 x_2^3 x_3^7 x_4^4 x_5^5, & \mathcal{Y}_{22,946} &= x_1^3 x_2^3 x_3^7 x_4^4 x_5^4, & \mathcal{Y}_{22,947} &= x_1^3 x_2^4 x_3^3 x_5^7 x_5^7, & \mathcal{Y}_{22,948} &= x_1^3 x_2^4 x_3^3 x_5^7 x_5^5, \\
\mathcal{Y}_{22,949} &= x_1^3 x_2^4 x_3^3 x_5^7 x_5^5, & \mathcal{Y}_{22,950} &= x_1^3 x_2^5 x_3^3 x_4^7 x_5^7, & \mathcal{Y}_{22,951} &= x_1^3 x_2^5 x_3^3 x_4^7 x_5^4, & \mathcal{Y}_{22,952} &= x_1^3 x_2^5 x_3^3 x_4^7 x_5^4, \\
\mathcal{Y}_{22,953} &= x_1^3 x_2^7 x_3^3 x_4^4 x_5^5, & \mathcal{Y}_{22,954} &= x_1^3 x_2^7 x_3^3 x_4^5 x_5^4, & \mathcal{Y}_{22,955} &= x_1^3 x_2^7 x_3^4 x_4^3 x_5^5, & \mathcal{Y}_{22,956} &= x_1^3 x_2^7 x_3^5 x_4^3 x_5^4, \\
\mathcal{Y}_{22,957} &= x_1^7 x_2^3 x_3^3 x_4^4 x_5^5, & \mathcal{Y}_{22,958} &= x_1^7 x_2^3 x_3^3 x_4^5 x_5^4, & \mathcal{Y}_{22,959} &= x_1^7 x_2^3 x_3^4 x_4^3 x_5^5, & \mathcal{Y}_{22,960} &= x_1^7 x_2^3 x_3^5 x_4^3 x_5^4, \\
\mathcal{Y}_{22,961} &= x_1^3 x_2^3 x_3^5 x_4^6 x_5^6, & \mathcal{Y}_{22,962} &= x_1^3 x_2^3 x_3^5 x_4^6 x_5^5, & \mathcal{Y}_{22,963} &= x_1^3 x_2^5 x_3^3 x_4^5 x_5^6, & \mathcal{Y}_{22,964} &= x_1^3 x_2^5 x_3^3 x_4^6 x_5^5, \\
\mathcal{Y}_{22,965} &= x_1^3 x_2^5 x_3^6 x_4^3 x_5^5.
\end{aligned}$$

From the above data, we see that $\overline{\Phi}(\mathcal{B}_4(\omega_{(j)})) \subset \mathcal{B}_5(\omega_{(j)})$ for $j = 1, 4, 5$. Hence, Conjecture 3.3.1 is true for $d = 5$ and the degree 22.

6.6. $\mathcal{A}_2$ -generators for $\mathcal{P}_5^0$ in degree 47

Using the results in Sect. 3.4.2, we have

$$\mathcal{B}_5^0(47) = \overline{\Phi}(\mathcal{B}_4(47)) = \overline{\Phi}(\mathcal{B}_4(\overline{\omega}_{(1)})) = \{\mathcal{Y}_{47,t}; 1 \leq t \leq 560\},$$

where $\overline{\omega}_{(1)} = (3, 2, 2, 2, 1)$, and the monomials $\mathcal{Y}_{47,t} : 1 \leq t \leq 560$, are determined as follows:

$$\begin{aligned}
\mathcal{Y}_{47,1} &= x_3 x_4^{15} x_5^{31}, & \mathcal{Y}_{47,2} &= x_3 x_4^{31} x_5^{15}, & \mathcal{Y}_{47,3} &= x_3^{15} x_4 x_5^{31}, & \mathcal{Y}_{47,4} &= x_3^{15} x_4^{31} x_5, \\
\mathcal{Y}_{47,5} &= x_3^{31} x_4 x_5^{15}, & \mathcal{Y}_{47,6} &= x_3^{31} x_4^{15} x_5, & \mathcal{Y}_{47,7} &= x_2 x_4^{15} x_5^{31}, & \mathcal{Y}_{47,8} &= x_2 x_4^{31} x_5^{15}, \\
\mathcal{Y}_{47,9} &= x_2 x_3^{15} x_5^{31}, & \mathcal{Y}_{47,10} &= x_2 x_3^{31} x_4^{15}, & \mathcal{Y}_{47,11} &= x_2 x_3^{31} x_5^{15}, & \mathcal{Y}_{47,12} &= x_2 x_3^{31} x_4^{15}, \\
\mathcal{Y}_{47,13} &= x_2^{15} x_4 x_5^{31}, & \mathcal{Y}_{47,14} &= x_2^{15} x_4^{31} x_5, & \mathcal{Y}_{47,15} &= x_2^{15} x_3 x_5^{31}, & \mathcal{Y}_{47,16} &= x_2^{15} x_3 x_4^{31}, \\
\mathcal{Y}_{47,17} &= x_2^{15} x_3^{31} x_5, & \mathcal{Y}_{47,18} &= x_2^{15} x_3^{31} x_4, & \mathcal{Y}_{47,19} &= x_2^{31} x_4 x_5^{15}, & \mathcal{Y}_{47,20} &= x_2^{31} x_4^{15} x_5, \\
\mathcal{Y}_{47,21} &= x_2^{31} x_3 x_5^{15}, & \mathcal{Y}_{47,22} &= x_2^{31} x_3 x_4^{15}, & \mathcal{Y}_{47,23} &= x_2^{31} x_3^{15} x_5, & \mathcal{Y}_{47,24} &= x_2^{31} x_3^{15} x_4, \\
\mathcal{Y}_{47,25} &= x_1 x_4^{15} x_5^{31}, & \mathcal{Y}_{47,26} &= x_1 x_4^{31} x_5^{15}, & \mathcal{Y}_{47,27} &= x_1 x_3^{15} x_5^{31}, & \mathcal{Y}_{47,28} &= x_1 x_3^{15} x_4^{31}, \\
\mathcal{Y}_{47,29} &= x_1 x_3^{31} x_5^{15}, & \mathcal{Y}_{47,30} &= x_1 x_3^{31} x_4^{15}, & \mathcal{Y}_{47,31} &= x_1 x_2^{15} x_5^{31}, & \mathcal{Y}_{47,32} &= x_1 x_2^{15} x_4^{31}, \\
\mathcal{Y}_{47,33} &= x_1 x_2^{15} x_3^{31}, & \mathcal{Y}_{47,34} &= x_1 x_2^{31} x_5^{15}, & \mathcal{Y}_{47,35} &= x_1 x_2^{31} x_4^{15}, & \mathcal{Y}_{47,36} &= x_1 x_2^{31} x_3^{15}, \\
\mathcal{Y}_{47,37} &= x_1^{15} x_4 x_5^{31}, & \mathcal{Y}_{47,38} &= x_1^{15} x_4^{31} x_5, & \mathcal{Y}_{47,39} &= x_1^{15} x_3 x_5^{31}, & \mathcal{Y}_{47,40} &= x_1^{15} x_3 x_4^{31}, \\
\mathcal{Y}_{47,41} &= x_1^{15} x_3^{31} x_5, & \mathcal{Y}_{47,42} &= x_1^{15} x_3^{31} x_4, & \mathcal{Y}_{47,43} &= x_1^{15} x_2 x_5^{31}, & \mathcal{Y}_{47,44} &= x_1^{15} x_2 x_4^{31}, \\
\mathcal{Y}_{47,45} &= x_1^{15} x_2 x_3^{31}, & \mathcal{Y}_{47,46} &= x_1^{15} x_2^{31} x_5, & \mathcal{Y}_{47,47} &= x_1^{15} x_2^{31} x_4, & \mathcal{Y}_{47,48} &= x_1^{15} x_2^{31} x_3, \\
\mathcal{Y}_{47,49} &= x_1^{31} x_4 x_5^{15}, & \mathcal{Y}_{47,50} &= x_1^{31} x_4^{15} x_5, & \mathcal{Y}_{47,51} &= x_1^{31} x_3 x_5^{15}, & \mathcal{Y}_{47,52} &= x_1^{31} x_3 x_4^{15}, \\
\mathcal{Y}_{47,53} &= x_1^{31} x_3^{15} x_5, & \mathcal{Y}_{47,54} &= x_1^{31} x_3^{15} x_4, & \mathcal{Y}_{47,55} &= x_1^{31} x_2 x_5^{15}, & \mathcal{Y}_{47,56} &= x_1^{31} x_2 x_4^{15},
\end{aligned}$$

$\mathcal{V}_{47,57} = x_1^{31}x_2x_3^{15}$	$\mathcal{V}_{47,58} = x_1^{31}x_2^{15}x_5$	$\mathcal{V}_{47,59} = x_1^{31}x_2^{15}x_4$	$\mathcal{V}_{47,60} = x_1^{31}x_2^{15}x_3$
$\mathcal{V}_{47,61} = x_3^3x_4^{13}x_5^{31}$	$\mathcal{V}_{47,62} = x_3^3x_4^{31}x_5^{13}$	$\mathcal{V}_{47,63} = x_3^{31}x_4x_5^{13}$	$\mathcal{V}_{47,64} = x_2^3x_4^{13}x_5^{31}$
$\mathcal{V}_{47,65} = x_2^3x_4^{31}x_5^{13}$	$\mathcal{V}_{47,66} = x_2^3x_3^{13}x_5^{31}$	$\mathcal{V}_{47,67} = x_2^3x_3^{13}x_4^{13}$	$\mathcal{V}_{47,68} = x_2^3x_3^{31}x_5^{13}$
$\mathcal{V}_{47,69} = x_2^3x_3^{31}x_4^{13}$	$\mathcal{V}_{47,70} = x_2^3x_3^{31}x_5^{13}$	$\mathcal{V}_{47,71} = x_2^3x_3^{13}x_5^{31}$	$\mathcal{V}_{47,72} = x_2^3x_3^{13}x_4^{13}$
$\mathcal{V}_{47,73} = x_1^3x_4^{13}x_5^{31}$	$\mathcal{V}_{47,74} = x_1^3x_4^{31}x_5^{13}$	$\mathcal{V}_{47,75} = x_1^3x_3^{13}x_5^{31}$	$\mathcal{V}_{47,76} = x_1^3x_3^{13}x_4^{13}$
$\mathcal{V}_{47,77} = x_1^3x_3^{31}x_5^{13}$	$\mathcal{V}_{47,78} = x_1^3x_3^{31}x_4^{13}$	$\mathcal{V}_{47,79} = x_1^3x_2^{13}x_5^{31}$	$\mathcal{V}_{47,80} = x_1^3x_2^{13}x_4^{13}$
$\mathcal{V}_{47,81} = x_1^3x_2^{13}x_3^{31}$	$\mathcal{V}_{47,82} = x_1^3x_2^{31}x_5^{13}$	$\mathcal{V}_{47,83} = x_1^3x_2^{31}x_4^{13}$	$\mathcal{V}_{47,84} = x_1^3x_2^{31}x_3^{13}$
$\mathcal{V}_{47,85} = x_1^{31}x_4x_5^{13}$	$\mathcal{V}_{47,86} = x_1^{31}x_3x_5^{13}$	$\mathcal{V}_{47,87} = x_1^{31}x_3x_4^{13}$	$\mathcal{V}_{47,88} = x_1^{31}x_2x_5^{13}$
$\mathcal{V}_{47,89} = x_1^{31}x_2x_4^{13}$	$\mathcal{V}_{47,90} = x_1^{31}x_2x_3^{13}$	$\mathcal{V}_{47,91} = x_3^3x_4^{15}x_5^{29}$	$\mathcal{V}_{47,92} = x_3^3x_4^{29}x_5^{15}$
$\mathcal{V}_{47,93} = x_3^{15}x_4x_5^{29}$	$\mathcal{V}_{47,94} = x_2^3x_4^{15}x_5^{29}$	$\mathcal{V}_{47,95} = x_2^3x_4^{29}x_5^{15}$	$\mathcal{V}_{47,96} = x_2^3x_3^{15}x_5^{29}$
$\mathcal{V}_{47,97} = x_2^3x_3^{15}x_4^{29}$	$\mathcal{V}_{47,98} = x_2^3x_3^{29}x_5^{15}$	$\mathcal{V}_{47,99} = x_2^3x_3^{29}x_4^{15}$	$\mathcal{V}_{47,100} = x_2^{15}x_3x_5^{29}$
$\mathcal{V}_{47,101} = x_2^{15}x_3x_5^{29}$	$\mathcal{V}_{47,102} = x_2^{15}x_3x_4^{29}$	$\mathcal{V}_{47,103} = x_3^3x_4^{15}x_5^{29}$	$\mathcal{V}_{47,104} = x_3^3x_4^{29}x_5^{15}$
$\mathcal{V}_{47,105} = x_1^3x_3^{15}x_5^{29}$	$\mathcal{V}_{47,106} = x_1^3x_3^{15}x_4^{29}$	$\mathcal{V}_{47,107} = x_1^3x_3^{29}x_5^{15}$	$\mathcal{V}_{47,108} = x_1^3x_3^{29}x_4^{15}$
$\mathcal{V}_{47,109} = x_1^3x_2^{15}x_5^{29}$	$\mathcal{V}_{47,110} = x_1^3x_2^{15}x_4^{29}$	$\mathcal{V}_{47,111} = x_1^3x_2^{15}x_3^{29}$	$\mathcal{V}_{47,112} = x_1^3x_2^{29}x_5^{15}$
$\mathcal{V}_{47,113} = x_1^3x_2^{29}x_4^{15}$	$\mathcal{V}_{47,114} = x_1^3x_2^{29}x_3^{15}$	$\mathcal{V}_{47,115} = x_1^{15}x_3x_5^{29}$	$\mathcal{V}_{47,116} = x_1^{15}x_3x_4^{29}$
$\mathcal{V}_{47,117} = x_1^{15}x_3x_4^{29}$	$\mathcal{V}_{47,118} = x_1^{15}x_2x_5^{29}$	$\mathcal{V}_{47,119} = x_1^{15}x_2x_4^{29}$	$\mathcal{V}_{47,120} = x_1^{15}x_2x_3^{29}$
$\mathcal{V}_{47,121} = x_7^3x_4^{11}x_5^{29}$	$\mathcal{V}_{47,122} = x_7^3x_4^{11}x_5^{29}$	$\mathcal{V}_{47,123} = x_7^3x_3^{11}x_5^{29}$	$\mathcal{V}_{47,124} = x_7^3x_3^{11}x_4^{29}$
$\mathcal{V}_{47,125} = x_7^3x_4^{11}x_5^{29}$	$\mathcal{V}_{47,126} = x_7^3x_3^{11}x_5^{29}$	$\mathcal{V}_{47,127} = x_7^3x_3^{11}x_4^{29}$	$\mathcal{V}_{47,128} = x_7^3x_2^{11}x_5^{29}$
$\mathcal{V}_{47,129} = x_7^3x_2^{11}x_4^{29}$	$\mathcal{V}_{47,130} = x_7^3x_2^{11}x_3^{29}$	$\mathcal{V}_{47,131} = x_7^3x_4^{27}x_5^{13}$	$\mathcal{V}_{47,132} = x_7^3x_4^{27}x_5^{13}$
$\mathcal{V}_{47,133} = x_7^3x_3^{27}x_5^{13}$	$\mathcal{V}_{47,134} = x_7^3x_3^{27}x_4^{13}$	$\mathcal{V}_{47,135} = x_7^3x_4^{27}x_5^{13}$	$\mathcal{V}_{47,136} = x_7^3x_3^{27}x_5^{13}$
$\mathcal{V}_{47,137} = x_7^3x_3^{27}x_4^{13}$	$\mathcal{V}_{47,138} = x_7^3x_2^{27}x_5^{13}$	$\mathcal{V}_{47,139} = x_7^3x_2^{27}x_4^{13}$	$\mathcal{V}_{47,140} = x_7^3x_2^{27}x_3^{13}$
$\mathcal{V}_{47,141} = x_2x_3x_4^{14}x_5^{31}$	$\mathcal{V}_{47,142} = x_2x_3x_4^{31}x_5^{14}$	$\mathcal{V}_{47,143} = x_2x_3^{14}x_4x_5^{31}$	$\mathcal{V}_{47,144} = x_2x_3^{14}x_4^{31}x_5$
$\mathcal{V}_{47,145} = x_2x_3^{31}x_4x_5^{14}$	$\mathcal{V}_{47,146} = x_2x_3^{31}x_4^{14}x_5$	$\mathcal{V}_{47,147} = x_2^{31}x_3x_4x_5^{14}$	$\mathcal{V}_{47,148} = x_2^{31}x_3x_4^{14}x_5$
$\mathcal{V}_{47,149} = x_1x_3x_4^{14}x_5^{31}$	$\mathcal{V}_{47,150} = x_1x_3x_4^{31}x_5^{14}$	$\mathcal{V}_{47,151} = x_1x_3^{14}x_4x_5^{31}$	$\mathcal{V}_{47,152} = x_1x_3^{14}x_4^{31}x_5$
$\mathcal{V}_{47,153} = x_1x_3^{31}x_4x_5^{14}$	$\mathcal{V}_{47,154} = x_1x_3^{31}x_4^{14}x_5$	$\mathcal{V}_{47,155} = x_1x_2x_4^{14}x_5^{31}$	$\mathcal{V}_{47,156} = x_1x_2x_3^{14}x_5^{14}$
$\mathcal{V}_{47,157} = x_1x_2x_3^{14}x_5^{31}$	$\mathcal{V}_{47,$		

54

$$\begin{aligned}
\mathcal{V}_{47,457} &= x_1^7 x_2 x_3^{27} x_4^{12}, & \mathcal{V}_{47,458} &= x_1^7 x_2^{27} x_4 x_5^{12}, & \mathcal{V}_{47,459} &= x_1^7 x_2^{27} x_3 x_5^{12}, & \mathcal{V}_{47,460} &= x_1^7 x_2^{27} x_3 x_4^{12}, \\
\mathcal{V}_{47,461} &= x_2 x_3^7 x_4^{26} x_5^{13}, & \mathcal{V}_{47,462} &= x_2^7 x_3 x_4^{26} x_5^{13}, & \mathcal{V}_{47,463} &= x_1 x_3^7 x_4^{26} x_5^{13}, & \mathcal{V}_{47,464} &= x_1 x_2^7 x_4^{26} x_5^{13}, \\
\mathcal{V}_{47,465} &= x_1 x_2^7 x_3^{26} x_5^{13}, & \mathcal{V}_{47,466} &= x_1 x_2^7 x_3^{26} x_4^{13}, & \mathcal{V}_{47,467} &= x_1^7 x_3 x_4^{26} x_5^{13}, & \mathcal{V}_{47,468} &= x_1^7 x_2 x_4^{26} x_5^{13}, \\
\mathcal{V}_{47,469} &= x_1^7 x_2 x_3^{26} x_5^{13}, & \mathcal{V}_{47,470} &= x_1^7 x_2 x_3^{26} x_4^{13}, & \mathcal{V}_{47,471} &= x_2^3 x_3^{13} x_4^2 x_5^{29}, & \mathcal{V}_{47,472} &= x_2^3 x_3^{29} x_4^2 x_5^{13}, \\
\mathcal{V}_{47,473} &= x_1^3 x_3^{13} x_4^2 x_5^{29}, & \mathcal{V}_{47,474} &= x_1^3 x_3^{29} x_4^2 x_5^{13}, & \mathcal{V}_{47,475} &= x_1^3 x_2^{13} x_4^2 x_5^{29}, & \mathcal{V}_{47,476} &= x_1^3 x_2^{13} x_3^2 x_5^{29}, \\
\mathcal{V}_{47,477} &= x_1^3 x_2^{13} x_3^2 x_4^{29}, & \mathcal{V}_{47,478} &= x_1^3 x_2^{29} x_3^2 x_4^{13}, & \mathcal{V}_{47,479} &= x_1^3 x_2^{29} x_3^2 x_5^{13}, & \mathcal{V}_{47,480} &= x_1^3 x_2^{29} x_3^2 x_4^{13}, \\
\mathcal{V}_{47,481} &= x_2^3 x_3^{12} x_4^{29}, & \mathcal{V}_{47,482} &= x_2^3 x_3^{29} x_4^{12}, & \mathcal{V}_{47,483} &= x_2^3 x_3^{29} x_4^2 x_5^{12}, & \mathcal{V}_{47,484} &= x_1^3 x_3^{12} x_4^{29}, \\
\mathcal{V}_{47,485} &= x_1^3 x_3^{29} x_4^{12}, & \mathcal{V}_{47,486} &= x_1^3 x_3^{29} x_4^2 x_5^{12}, & \mathcal{V}_{47,487} &= x_1^3 x_2^3 x_4^{12} x_5^{29}, & \mathcal{V}_{47,488} &= x_1^3 x_2^3 x_4^{29} x_5^{12}, \\
\mathcal{V}_{47,489} &= x_1^3 x_2^3 x_3^{12} x_5^{29}, & \mathcal{V}_{47,490} &= x_1^3 x_2^3 x_3^{12} x_4^{29}, & \mathcal{V}_{47,491} &= x_1^3 x_2^3 x_3^{29} x_5^{12}, & \mathcal{V}_{47,492} &= x_1^3 x_2^3 x_3^{29} x_4^{12}, \\
\mathcal{V}_{47,493} &= x_1^3 x_2^{29} x_3^3 x_5^{12}, & \mathcal{V}_{47,494} &= x_1^3 x_2^{29} x_3^3 x_4^{12}, & \mathcal{V}_{47,495} &= x_1^3 x_2^{29} x_3^3 x_5^{12}, & \mathcal{V}_{47,496} &= x_2^3 x_3^3 x_4^{13} x_5^{28}, \\
\mathcal{V}_{47,497} &= x_2^3 x_3^3 x_4^{28} x_5^{13}, & \mathcal{V}_{47,498} &= x_2^3 x_3^{13} x_4^3 x_5^{28}, & \mathcal{V}_{47,499} &= x_1^3 x_3^3 x_4^{13} x_5^{28}, & \mathcal{V}_{47,500} &= x_1^3 x_3^3 x_4^{28} x_5^{13}, \\
\mathcal{V}_{47,501} &= x_1^3 x_3^{13} x_4^3 x_5^{28}, & \mathcal{V}_{47,502} &= x_1^3 x_2^3 x_4^{13} x_5^{28}, & \mathcal{V}_{47,503} &= x_1^3 x_2^3 x_4^{28} x_5^{13}, & \mathcal{V}_{47,504} &= x_1^3 x_2^3 x_3^{13} x_5^{28}, \\
\mathcal{V}_{47,505} &= x_1^3 x_2^3 x_3^{13} x_4^{28}, & \mathcal{V}_{47,506} &= x_1^3 x_2^3 x_3^{28} x_4^{13}, & \mathcal{V}_{47,507} &= x_1^3 x_2^3 x_3^{28} x_5^{13}, & \mathcal{V}_{47,508} &= x_1^3 x_2^{13} x_4^3 x_5^{28}, \\
\mathcal{V}_{47,509} &= x_1^3 x_2^{13} x_3^3 x_5^{28}, & \mathcal{V}_{47,510} &= x_1^3 x_2^{13} x_3^3 x_4^{28}, & \mathcal{V}_{47,511} &= x_2^3 x_3^4 x_4^{11} x_5^{29}, & \mathcal{V}_{47,512} &= x_1^3 x_3^4 x_4^{11} x_5^{29}, \\
\mathcal{V}_{47,513} &= x_1^3 x_2^4 x_4^{11} x_5^{29}, & \mathcal{V}_{47,514} &= x_1^3 x_2^4 x_3^{11} x_5^{29}, & \mathcal{V}_{47,515} &= x_1^3 x_2^4 x_3^{11} x_4^{29}, & \mathcal{V}_{47,516} &= x_2^3 x_3^4 x_4^{27} x_5^{13}, \\
\mathcal{V}_{47,517} &= x_1^3 x_2^4 x_3^{11} x_5^{13}, & \mathcal{V}_{47,518} &= x_1^3 x_2^4 x_3^{11} x_4^{13}, & \mathcal{V}_{47,519} &= x_1^3 x_2^4 x_3^{27} x_5^{13}, & \mathcal{V}_{47,520} &= x_1^3 x_2^4 x_3^{27} x_4^{13}, \\
\mathcal{V}_{47,521} &= x_2^3 x_3^5 x_4^{10} x_5^{29}, & \mathcal{V}_{47,522} &= x_1^3 x_3^5 x_4^{10} x_5^{29}, & \mathcal{V}_{47,523} &= x_1^3 x_2^5 x_4^{10} x_5^{29}, & \mathcal{V}_{47,524} &= x_1^3 x_2^5 x_3^{10} x_5^{29}, \\
\mathcal{V}_{47,525} &= x_1^3 x_2^5 x_3^{10} x_4^{29}, & \mathcal{V}_{47,526} &= x_2^3 x_3^5 x_4^{11} x_5^{28}, & \mathcal{V}_{47,527} &= x_1^3 x_3^5 x_4^{11} x_5^{28}, & \mathcal{V}_{47,528} &= x_1^3 x_2^5 x_4^{11} x_5^{28}, \\
\mathcal{V}_{47,529} &= x_1^3 x_2^5 x_3^{11} x_4^{28}, & \mathcal{V}_{47,530} &= x_1^3 x_2^5 x_3^{11} x_5^{28}, & \mathcal{V}_{47,531} &= x_2^3 x_3^5 x_4^{27} x_5^{12}, & \mathcal{V}_{47,532} &= x_1^3 x_3^5 x_4^{27} x_5^{12}, \\
\mathcal{V}_{47,533} &= x_1^3 x_2^5 x_3^{27} x_4^{12}, & \mathcal{V}_{47,534} &= x_1^3 x_2^5 x_3^{27} x_5^{12}, & \mathcal{V}_{47,535} &= x_1^3 x_2^5 x_3^{27} x_4^{12}, & \mathcal{V}_{47,536} &= x_2^3 x_3^5 x_4^{26} x_5^{13}, \\
\mathcal{V}_{47,537} &= x_1^3 x_2^5 x_3^{26} x_4^{13}, & \mathcal{V}_{47,538} &= x_1^3 x_2^5 x_3^{26} x_5^{13}, & \mathcal{V}_{47,539} &= x_1^3 x_2^5 x_3^{26} x_4^{13}, & \mathcal{V}_{47,540} &= x_1^3 x_2^5 x_3^{26} x_5^{13}, \\
\mathcal{V}_{47,541} &= x_2^3 x_3^7 x_4^9 x_5^{28}, & \mathcal{V}_{47,542} &= x_2^7 x_3^3 x_4^9 x_5^{28}, & \mathcal{V}_{47,543} &= x_1^3 x_3^7 x_4^9 x_5^{28}, & \mathcal{V}_{47,544} &= x_1^3 x_2^7 x_4^9 x_5^{28}, \\
\mathcal{V}_{47,545} &= x_1^3 x_2^7 x_3^9 x_5^{28}, & \mathcal{V}_{47,546} &= x_1^3 x_2^7 x_3^9 x_4^{28}, & \mathcal{V}_{47,547} &= x_1^7 x_3^3 x_4^9 x_5^{28}, & \mathcal{V}_{47,548} &= x_1^7 x_2^3 x_4^9 x_5^{28}, \\
\mathcal{V}_{47,549} &= x_1^7 x_2^3 x_3^9 x_5^{28}, & \mathcal{V}_{47,550} &= x_1^7 x_2^3 x_3^9 x_4^{28}, & \mathcal{V}_{47,551} &= x_2^3 x_3^7 x_4^{25} x_5^{12}, & \mathcal{V}_{47,552} &= x_2^3 x_3^7 x_4^{25} x_5^{12}, \\
\mathcal{V}_{47,553} &= x_1^3 x_3^7 x_4^{25} x_5^{12}, & \mathcal{V}_{47,554} &= x_1^3 x_2^7 x_4^{25} x_5^{12}, & \mathcal{V}_{47,555} &= x_1^3 x_2^7 x_3^{25} x_5^{12}, & \mathcal{V}_{47,556} &= x_1^3 x_2^7 x_3^{25} x_4^{12}, \\
\mathcal{V}_{47,557} &= x_1^7 x_3^3 x_4^{25} x_5^{12}, & \mathcal{V}_{47,558} &= x_1^7 x_2^3 x_4^{25} x_5^{12}, & \mathcal{V}_{47,559} &= x_1^7 x_2^3 x_3^{25} x_5^{12}, & \mathcal{V}_{47,560} &= x_1^7 x_2^3 x_3^{25} x_4^{12}.
\end{aligned}$$

6.7. $\mathcal{A}_2$ -generators for $\mathcal{P}_5^+$ in degree 47

Recall that Kameko's squaring operation $\widetilde{Sq}_*^0(5,47) : (Q\mathcal{P}_5)_{47} \rightarrow (Q\mathcal{P}_5)_{21}$ is an epimorphism of $\mathbb{Z}/2GL_5$ -modules. Therefore, we get

$$\mathcal{B}_5(47) = \mathcal{B}_5^0(47) \cup \varphi(\mathcal{B}_5(21)) \cup \left(\mathcal{B}_5^+(47) \cap \text{Ker}((\widetilde{Sq}_*^0(5,47)) \right),$$

where $|\mathcal{B}_5^0(47)| = 560$, $|\varphi(\mathcal{B}_5(21))| = 840$ with $\varphi : \mathcal{P}_5 \rightarrow \mathcal{P}_5$, $\varphi(u) = X_{(0,5)} u^2$, $\forall u \in \mathcal{P}_5$. From the results in Sect.3.4.2, we have

$$\mathcal{B}_5^+(47) \cap \text{Ker}((\widetilde{Sq}_*^0(5,47)) = \mathcal{B}_5^+(\overline{w}_{(1)}) \cup \mathcal{B}_5^+(\overline{w}_{(4)}) \cup \mathcal{B}_5^+(\overline{w}_{(5)}),$$

where $\overline{w}_{(1)} = (3, 2, 2, 2, 1)$, $\overline{w}_{(4)} = (3, 4, 3, 1, 1)$, and $\overline{w}_{(5)} = (3, 4, 3, 3)$.

$\mathcal{B}_5^+(\overline{w}_{(1)})$ is the set of 196 admissible monomials: $\mathcal{V}_{47,t}$, $1 \leq t \leq 370$

$$\begin{aligned}
\mathcal{V}_{47,1} &= x_1 x_2 x_3 x_4^{14} x_5^{30}, & \mathcal{V}_{47,2} &= x_1 x_2 x_3 x_4^{30} x_5^{14}, & \mathcal{V}_{47,3} &= x_1 x_2 x_3^{14} x_4 x_5^{30}, \\
\mathcal{V}_{47,4} &= x_1 x_2 x_3^{14} x_4^{30} x_5, & \mathcal{V}_{47,5} &= x_1 x_2 x_3^{30} x_4 x_5^{14}, & \mathcal{V}_{47,6} &= x_1 x_2 x_3^{30} x_4^{14} x_5, \\
\mathcal{V}_{47,7} &= x_1 x_2^{14} x_3 x_4 x_5^{30}, & \mathcal{V}_{47,8} &= x_1 x_2^{14} x_3 x_4^{30} x_5, & \mathcal{V}_{47,9} &= x_1 x_2^{30} x_3 x_4 x_5^{14}, \\
\mathcal{V}_{47,10} &= x_1 x_2^{30} x_3 x_4^{14} x_5, & \mathcal{V}_{47,11} &= x_1 x_2 x_3^2 x_4^{12} x_5^{31}, & \mathcal{V}_{47,12} &= x_1 x_2 x_3^2 x_4^{31} x_5^{12}, \\
\mathcal{V}_{47,13} &= x_1 x_2 x_3^{31} x_4^2 x_5^{12}, & \mathcal{V}_{47,14} &= x_1 x_2^2 x_3 x_4^{12} x_5^{31}, & \mathcal{V}_{47,15} &= x_1 x_2^2 x_3 x_4^{31} x_5^{12}, \\
\mathcal{V}_{47,16} &= x_1 x_2^2 x_3^{31} x_4 x_5^{12}, & \mathcal{V}_{47,17} &= x_1 x_2^{31} x_3 x_4^2 x_5^{12}, & \mathcal{V}_{47,18} &= x_1 x_2^{31} x_3^2 x_4 x_5^{12}, \\
\mathcal{V}_{47,19} &= x_1^{31} x_2 x_3 x_4^2 x_5^{12}, & \mathcal{V}_{47,20} &= x_1^{31} x_2 x_3^2 x_4 x_5^{12}, & \mathcal{V}_{47,21} &= x_1 x_2^2 x_3^{12} x_4 x_5^{31}, \\
\mathcal{V}_{47,22} &= x_1 x_2^2 x_3^{12} x_4^{31} x_5, & \mathcal{V}_{47,23} &= x_1 x_2^2 x_3^{31} x_4^{12} x_5, & \mathcal{V}_{47,24} &= x_1 x_2^{31} x_3^2 x_4^{12} x_5, \\
\mathcal{V}_{47,25} &= x_1^{31} x_2 x_3^2 x_4^{12} x_5, & \mathcal{V}_{47,26} &= x_1 x_2 x_3^2 x_4^{13} x_5^{30}, & \mathcal{V}_{47,27} &= x_1 x_2 x_3^2 x_4^{30} x_5^{13},
\end{aligned}$$

$$\begin{aligned}
\mathcal{V}_{47,28} &= x_1 x_2 x_3^{30} x_4^2 x_5^{13}, & \mathcal{V}_{47,29} &= x_1 x_2^2 x_3 x_4^{13} x_5^{30}, & \mathcal{V}_{47,30} &= x_1 x_2^2 x_3 x_4^{30} x_5^{13}, \\
\mathcal{V}_{47,31} &= x_1 x_2^2 x_3^{13} x_4 x_5^{30}, & \mathcal{V}_{47,32} &= x_1 x_2^2 x_3^{13} x_4^{30} x_5, & \mathcal{V}_{47,33} &= x_1 x_2^{30} x_3 x_4^2 x_5^{13}, \\
\mathcal{V}_{47,34} &= x_1 x_2 x_3^2 x_4^{14} x_5^{29}, & \mathcal{V}_{47,35} &= x_1 x_2 x_3^2 x_4^{29} x_5^{14}, & \mathcal{V}_{47,36} &= x_1 x_2 x_3^{14} x_4^2 x_5^{29}, \\
\mathcal{V}_{47,37} &= x_1 x_2^2 x_3 x_4^{14} x_5^{29}, & \mathcal{V}_{47,38} &= x_1 x_2^2 x_3 x_4^{29} x_5^{14}, & \mathcal{V}_{47,39} &= x_1 x_2^2 x_3^2 x_4 x_5^{14}, \\
\mathcal{V}_{47,40} &= x_1 x_2^2 x_3^2 x_4^{14} x_5, & \mathcal{V}_{47,41} &= x_1 x_2^{14} x_3 x_4^2 x_5^{29}, & \mathcal{V}_{47,42} &= x_1 x_2 x_3^2 x_4^{15} x_5^{28}, \\
\mathcal{V}_{47,43} &= x_1 x_2 x_3^2 x_4^{28} x_5^{15}, & \mathcal{V}_{47,44} &= x_1 x_2 x_3^{15} x_4^2 x_5^{28}, & \mathcal{V}_{47,45} &= x_1 x_2^2 x_3 x_4^{15} x_5^{28}, \\
\mathcal{V}_{47,46} &= x_1 x_2^2 x_3 x_4^{28} x_5^{15}, & \mathcal{V}_{47,47} &= x_1 x_2^2 x_3^{15} x_4 x_5^{28}, & \mathcal{V}_{47,48} &= x_1 x_2^{15} x_3 x_4^2 x_5^{28}, \\
\mathcal{V}_{47,49} &= x_1 x_2^{15} x_3^2 x_4 x_5^{28}, & \mathcal{V}_{47,50} &= x_1^{15} x_2 x_3 x_4^2 x_5^{28}, & \mathcal{V}_{47,51} &= x_1^{15} x_2^2 x_3 x_4 x_5^{28}, \\
\mathcal{V}_{47,52} &= x_1 x_2^2 x_3^{15} x_4^{28} x_5, & \mathcal{V}_{47,53} &= x_1 x_2^2 x_3^{28} x_4 x_5^{15}, & \mathcal{V}_{47,54} &= x_1 x_2^2 x_3^{28} x_4^{15} x_5, \\
\mathcal{V}_{47,55} &= x_1 x_2^{15} x_3^2 x_4^{28} x_5, & \mathcal{V}_{47,56} &= x_1^{15} x_2 x_3^2 x_4^{28} x_5, & \mathcal{V}_{47,57} &= x_1 x_2 x_3^3 x_4^{12} x_5^{30}, \\
\mathcal{V}_{47,58} &= x_1 x_2 x_3^3 x_4^{30} x_5^{12}, & \mathcal{V}_{47,59} &= x_1 x_2 x_3^{30} x_4^3 x_5^{12}, & \mathcal{V}_{47,60} &= x_1 x_2^3 x_3 x_4^{12} x_5^{30}, \\
\mathcal{V}_{47,61} &= x_1 x_2^2 x_3 x_4^{30} x_5^{12}, & \mathcal{V}_{47,62} &= x_1 x_2^3 x_3^{12} x_4 x_5^{30}, & \mathcal{V}_{47,63} &= x_1 x_2^3 x_3^{12} x_4^3 x_5^{30}, \\
\mathcal{V}_{47,64} &= x_1 x_2^2 x_3^3 x_4 x_5^{12}, & \mathcal{V}_{47,65} &= x_1 x_2^{30} x_3 x_4^3 x_5^{12}, & \mathcal{V}_{47,66} &= x_1 x_2^2 x_3^3 x_4 x_5^{12}, \\
\mathcal{V}_{47,67} &= x_1^3 x_2 x_3 x_4^{12} x_5^{30}, & \mathcal{V}_{47,68} &= x_1^3 x_2 x_3 x_4^{30} x_5^{12}, & \mathcal{V}_{47,69} &= x_1^3 x_2 x_3^{12} x_4 x_5^{30}, \\
\mathcal{V}_{47,70} &= x_1^3 x_2 x_3^{12} x_4^{30} x_5, & \mathcal{V}_{47,71} &= x_1^3 x_2 x_3^{30} x_4 x_5^{12}, & \mathcal{V}_{47,72} &= x_1 x_2^3 x_3^{30} x_4^{12} x_5, \\
\mathcal{V}_{47,73} &= x_1^3 x_2 x_3^{30} x_4^{12} x_5, & \mathcal{V}_{47,74} &= x_1 x_2 x_3^3 x_4^{14} x_5^{28}, & \mathcal{V}_{47,75} &= x_1 x_2 x_3^3 x_4^{28} x_5^{14}, \\
\mathcal{V}_{47,76} &= x_1 x_2 x_3^{14} x_4^3 x_5^{28}, & \mathcal{V}_{47,77} &= x_1 x_2^3 x_3 x_4^{14} x_5^{28}, & \mathcal{V}_{47,78} &= x_1 x_2^3 x_3 x_4^{28} x_5^{14}, \\
\mathcal{V}_{47,79} &= x_1 x_2^3 x_3^{14} x_4 x_5^{28}, & \mathcal{V}_{47,80} &= x_1 x_2^3 x_3^{28} x_4 x_5^{14}, & \mathcal{V}_{47,81} &= x_1 x_2^3 x_3^{28} x_4^{14} x_5, \\
\mathcal{V}_{47,82} &= x_1 x_2^{14} x_3 x_4^3 x_5^{28}, & \mathcal{V}_{47,83} &= x_1 x_2^{14} x_3^3 x_4 x_5^{28}, & \mathcal{V}_{47,84} &= x_1^3 x_2 x_3 x_4^{14} x_5^{28}, \\
\mathcal{V}_{47,85} &= x_1^3 x_2 x_3 x_4^{28} x_5^{14}, & \mathcal{V}_{47,86} &= x_1^3 x_2 x_3^{14} x_4 x_5^{28}, & \mathcal{V}_{47,87} &= x_1^3 x_2 x_3^{28} x_4 x_5^{14}, \\
\mathcal{V}_{47,88} &= x_1^3 x_2 x_3^{28} x_4^{14} x_5, & \mathcal{V}_{47,89} &= x_1 x_2^3 x_3^{14} x_4^{28} x_5, & \mathcal{V}_{47,90} &= x_1^3 x_2 x_3^{14} x_4^{28} x_5, \\
\mathcal{V}_{47,91} &= x_1 x_2 x_3^6 x_4^{10} x_5^{29}, & \mathcal{V}_{47,92} &= x_1 x_2^6 x_3 x_4^{10} x_5^{29}, & \mathcal{V}_{47,93} &= x_1 x_2 x_3^6 x_4^{11} x_5^{28}, \\
\mathcal{V}_{47,94} &= x_1 x_2^6 x_3 x_4^{11} x_5^{28}, & \mathcal{V}_{47,95} &= x_1 x_2^6 x_3^{11} x_4 x_5^{28}, & \mathcal{V}_{47,96} &= x_1 x_2 x_3^6 x_4^{27} x_5^{12}, \\
\mathcal{V}_{47,97} &= x_1 x_2^6 x_3 x_4^{27} x_5^{12}, & \mathcal{V}_{47,98} &= x_1 x_2^6 x_3^{27} x_4 x_5^{12}, & \mathcal{V}_{47,99} &= x_1 x_2 x_3^6 x_4^{26} x_5^{13}, \\
\mathcal{V}_{47,100} &= x_1 x_2^6 x_3 x_4^{26} x_5^{13}, & \mathcal{V}_{47,101} &= x_1 x_2 x_3^7 x_4^{10} x_5^{28}, & \mathcal{V}_{47,102} &= x_1 x_2^7 x_3 x_4^{10} x_5^{28}, \\
\mathcal{V}_{47,103} &= x_1 x_2^7 x_3^{10} x_4 x_5^{28}, & \mathcal{V}_{47,104} &= x_1^7 x_2 x_3 x_4^{10} x_5^{28}, & \mathcal{V}_{47,105} &= x_1^7 x_2 x_3^{10} x_4 x_5^{28}, \\
\mathcal{V}_{47,106} &= x_1 x_2 x_3^7 x_4^{26} x_5^{12}, & \mathcal{V}_{47,107} &= x_1 x_2^7 x_3 x_4^{26} x_5^{12}, & \mathcal{V}_{47,108} &= x_1 x_2^7 x_3^{26} x_4 x_5^{12}, \\
\mathcal{V}_{47,109} &= x_1^7 x_2 x_3 x_4^{26} x_5^{12}, & \mathcal{V}_{47,110} &= x_1^7 x_2 x_3^{26} x_4 x_5^{12}, & \mathcal{V}_{47,111} &= x_1 x_2^2 x_3^{13} x_4^2 x_5^{29}, \\
\mathcal{V}_{47,112} &= x_1 x_2^2 x_3^{29} x_4^2 x_5^{13}, & \mathcal{V}_{47,113} &= x_1 x_2^2 x_3^{12} x_4^{29} x_5^{13}, & \mathcal{V}_{47,114} &= x_1 x_2^2 x_3^{29} x_4^2 x_5^{12}, \\
\mathcal{V}_{47,115} &= x_1 x_2^2 x_3^{29} x_4^3 x_5^{12}, & \mathcal{V}_{47,116} &= x_1 x_2^3 x_3^{12} x_4^{29} x_5^{13}, & \mathcal{V}_{47,117} &= x_1 x_2^3 x_3^{29} x_4^2 x_5^{12}, \\
\mathcal{V}_{47,118} &= x_1 x_2^3 x_3^{12} x_4^2 x_5^{29}, & \mathcal{V}_{47,119} &= x_1 x_2^3 x_3^{29} x_4^2 x_5^{12}, & \mathcal{V}_{47,120} &= x_1^3 x_2 x_3^2 x_4^{12} x_5^{29}, \\
\mathcal{V}_{47,121} &= x_1^3 x_2 x_3^{12} x_4^{29} x_5^{12}, & \mathcal{V}_{47,122} &= x_1^3 x_2 x_3^{12} x_4^2 x_5^{29}, & \mathcal{V}_{47,123} &= x_1^3 x_2 x_3^{29} x_4^2 x_5^{12}, \\
\mathcal{V}_{47,124} &= x_1^3 x_2^{29} x_3 x_4^2 x_5^{12}, & \mathcal{V}_{47,125} &= x_1^3 x_2^{29} x_3^2 x_4 x_5^{12}, & \mathcal{V}_{47,126} &= x_1^3 x_2^{29} x_3^2 x_4^{12} x_5, \\
\mathcal{V}_{47,127} &= x_1 x_2^2 x_3^{12} x_4^3 x_5^{29}, & \mathcal{V}_{47,128} &= x_1 x_2^2 x_3^{13} x_4^3 x_5^{28}, & \mathcal{V}_{47,129} &= x_1 x_2^2 x_3^3 x_4^{28} x_5^{13}, \\
\mathcal{V}_{47,130} &= x_1 x_2^2 x_3^{13} x_4^3 x_5^{28}, & \mathcal{V}_{47,131} &= x_1 x_2^3 x_3^{12} x_4^{13} x_5^{28}, & \mathcal{V}_{47,132} &= x_1 x_2^3 x_3^2 x_4^{28} x_5^{13}, \\
\mathcal{V}_{47,133} &= x_1 x_2^3 x_3^{13} x_4^2 x_5^{28}, & \mathcal{V}_{47,134} &= x_1 x_2^3 x_3^{28} x_4^2 x_5^{13}, & \mathcal{V}_{47,135} &= x_1^3 x_2 x_3^2 x_4^{13} x_5^{28}, \\
\mathcal{V}_{47,136} &= x_1^3 x_2 x_3^2 x_4^{28} x_5^{13}, & \mathcal{V}_{47,137} &= x_1^3 x_2 x_3^{13} x_4^2 x_5^{28}, & \mathcal{V}_{47,138} &= x_1^3 x_2 x_3^{28} x_4^2 x_5^{13}, \\
\mathcal{V}_{47,139} &= x_1^3 x_2^{13} x_3 x_4^2 x_5^{28}, & \mathcal{V}_{47,140} &= x_1^3 x_2^{13} x_3^2 x_4 x_5^{28}, & \mathcal{V}_{47,141} &= x_1^3 x_2^{13} x_3^2 x_4^{28} x_5, \\
\mathcal{V}_{47,142} &= x_1 x_2^2 x_3^{28} x_4^3 x_5^{13}, & \mathcal{V}_{47,143} &= x_1 x_2^2 x_3^4 x_4^9 x_5^{31}, & \mathcal{V}_{47,144} &= x_1 x_2^2 x_3^4 x_4^{31} x_5^9, \\
\mathcal{V}_{47,145} &= x_1 x_2^2 x_3^{31} x_4^4 x_5^9, & \mathcal{V}_{47,146} &= x_1 x_2^{31} x_3^2 x_4^4 x_5^9, & \mathcal{V}_{47,147} &= x_1^{31} x_2 x_3^2 x_4^4 x_5^9, \\
\mathcal{V}_{47,148} &= x_1 x_2^2 x_3^4 x_4^{11} x_5^{29}, & \mathcal{V}_{47,149} &= x_1 x_2^2 x_3^4 x_4^{27} x_5^{13}, & \mathcal{V}_{47,150} &= x_1 x_2^{15} x_3^4 x_4^{15} x_5^{25}, \\
\mathcal{V}_{47,151} &= x_1 x_2^2 x_3^4 x_4^{25} x_5^{15}, & \mathcal{V}_{47,152} &= x_1 x_2^2 x_3^{15} x_4^4 x_5^{25}, & \mathcal{V}_{47,153} &= x_1 x_2^{15} x_3^2 x_4^4 x_5^{25}, \\
\mathcal{V}_{47,154} &= x_1^{15} x_2 x_3^2 x_4^4 x_5^{25}, & \mathcal{V}_{47,155} &= x_1 x_2^2 x_3^5 x_4^8 x_5^{31}, & \mathcal{V}_{47,156} &= x_1 x_2^2 x_3^5 x_4^{31} x_5^8, \\
\mathcal{V}_{47,157} &= x_1 x_2^2 x_3^{31} x_4^5 x_5^8, & \mathcal{V}_{47,158} &= x_1 x_2^{31} x_3^2 x_4^5 x_5^8, & \mathcal{V}_{47,159} &= x_1^{31} x_2 x_3^2 x_4^5 x_5^8, \\
\mathcal{V}_{47,160} &= x_1 x_2^2 x_3^5 x_4^{30} x_5^9, & \mathcal{V}_{47,161} &= x_1 x_2^2 x_3^5 x_4^9 x_5^{30}, & \mathcal{V}_{47,162} &= x_1 x_2^2 x_3^5 x_4^{10} x_5^{29}, \\
\mathcal{V}_{47,163} &= x_1 x_2^2 x_3^5 x_4^{11} x_5^{28}, & \mathcal{V}_{47,164} &= x_1 x_2^2 x_3^5 x_4^{27} x_5^{12}, & \mathcal{V}_{47,165} &= x_1 x_2^2 x_3^5 x_4^{26} x_5^{13}, \\
\mathcal{V}_{47,166} &= x_1 x_2^2 x_3^5 x_4^{14} x_5^{25}, & \mathcal{V}_{47,167} &= x_1 x_2^2 x_3^5 x_4^{25} x_5^{14}, & \mathcal{V}_{47,168} &= x_1 x_2^2 x_3^5 x_4^{15} x_5^{24}, \\
\mathcal{V}_{47,169} &= x_1 x_2^2 x_3^5 x_4^{24} x_5^{15}, & \mathcal{V}_{47,170} &= x_1 x_2^2 x_3^{15} x_4^5 x_5^{24}, & \mathcal{V}_{47,171} &= x_1 x_2^{15} x_3^2 x_4^5 x_5^{24}, \\
\mathcal{V}_{47,172} &= x_1^{15} x_2 x_3^2 x_4^5 x_5^{24}, & \mathcal{V}_{47,173} &= x_1 x_2^2 x_3^7 x_4^8 x_5^{29}, & \mathcal{V}_{47,174} &= x_1 x_2^2 x_3^7 x_4^{29} x_5^8, \\
\mathcal{V}_{47,175} &= x_1 x_2^7 x_3^2 x_4^8 x_5^{29}, & \mathcal{V}_{47,176} &= x_1 x_2^7 x_3^2 x_4^{29} x_5^8, & \mathcal{V}_{47,177} &= x_1^7 x_2 x_3^2 x_4^8 x_5^{29},
\end{aligned}$$

$$\begin{aligned}
\mathcal{V}_{47,178} &= x_1^7 x_2 x_3^2 x_4^{29} x_5^8, & \mathcal{V}_{47,179} &= x_1 x_2^2 x_3^7 x_4^9 x_5^{28}, & \mathcal{V}_{47,180} &= x_1 x_2^7 x_3^2 x_4^9 x_5^{28}, \\
\mathcal{V}_{47,181} &= x_1^7 x_2 x_3^2 x_4^9 x_5^{28}, & \mathcal{V}_{47,182} &= x_1 x_2^2 x_3^7 x_4^{25} x_5^{12}, & \mathcal{V}_{47,183} &= x_1 x_2^7 x_3^2 x_4^{25} x_5^{12}, \\
\mathcal{V}_{47,184} &= x_1^7 x_2 x_3^2 x_4^{25} x_5^{12}, & \mathcal{V}_{47,185} &= x_1 x_2^2 x_3^7 x_4^{24} x_5^{13}, & \mathcal{V}_{47,186} &= x_1 x_2^7 x_3^2 x_4^{13} x_5^{24}, \\
\mathcal{V}_{47,187} &= x_1 x_2^7 x_3^2 x_4^{24} x_5^{13}, & \mathcal{V}_{47,188} &= x_1 x_2^7 x_3^2 x_4^{13} x_5^{24}, & \mathcal{V}_{47,189} &= x_1^7 x_2 x_3^2 x_4^{24} x_5^{13}, \\
\mathcal{V}_{47,190} &= x_1^7 x_2 x_3^2 x_4^{13} x_5^{24}, & \mathcal{V}_{47,191} &= x_1 x_2^3 x_3^3 x_4^{12} x_5^{28}, & \mathcal{V}_{47,192} &= x_1 x_2^3 x_3^3 x_4^{28} x_5^{12}, \\
\mathcal{V}_{47,193} &= x_1 x_2^3 x_3^3 x_4^{12} x_5^{28}, & \mathcal{V}_{47,194} &= x_1 x_2^3 x_3^3 x_4^{12} x_5^{28}, & \mathcal{V}_{47,195} &= x_1^3 x_2 x_3^3 x_4^{12} x_5^{28}, \\
\mathcal{V}_{47,196} &= x_1^3 x_2 x_3^3 x_4^{28} x_5^{12}, & \mathcal{V}_{47,197} &= x_1^3 x_2 x_3^{12} x_4^3 x_5^{28}, & \mathcal{V}_{47,198} &= x_1^3 x_2 x_3^{28} x_4^3 x_5^{12}, \\
\mathcal{V}_{47,199} &= x_1^3 x_2^3 x_3 x_4^{12} x_5^{28}, & \mathcal{V}_{47,200} &= x_1^3 x_2^3 x_3 x_4^{28} x_5^{12}, & \mathcal{V}_{47,201} &= x_1^3 x_2^3 x_3^{12} x_4 x_5^{28}, \\
\mathcal{V}_{47,202} &= x_1^3 x_2^3 x_3^{28} x_4 x_5^{12}, & \mathcal{V}_{47,203} &= x_1 x_2^3 x_3^4 x_4^{31} x_5^8, & \mathcal{V}_{47,204} &= x_1 x_2^3 x_3^4 x_4^{31} x_5^8, \\
\mathcal{V}_{47,205} &= x_1 x_2^3 x_3^{31} x_4^4 x_5^8, & \mathcal{V}_{47,206} &= x_1 x_2^{31} x_3^3 x_4^4 x_5^8, & \mathcal{V}_{47,207} &= x_1^3 x_2 x_3^4 x_4^8 x_5^{31}, \\
\mathcal{V}_{47,208} &= x_1^3 x_2 x_3^4 x_4^{31} x_5^8, & \mathcal{V}_{47,209} &= x_1^3 x_2 x_3^{31} x_4^4 x_5^8, & \mathcal{V}_{47,210} &= x_1^3 x_2^{31} x_3 x_4^4 x_5^8, \\
\mathcal{V}_{47,211} &= x_1^{31} x_2 x_3^3 x_4^4 x_5^8, & \mathcal{V}_{47,212} &= x_1^{31} x_2 x_3 x_4^4 x_5^8, & \mathcal{V}_{47,213} &= x_1 x_2^3 x_3^4 x_4^{30} x_5^9, \\
\mathcal{V}_{47,214} &= x_1 x_2^3 x_3^{30} x_4^4 x_5^9, & \mathcal{V}_{47,215} &= x_1 x_2^3 x_3^4 x_4^9 x_5^{30}, & \mathcal{V}_{47,216} &= x_1^3 x_2 x_3^4 x_4^{30} x_5^9, \\
\mathcal{V}_{47,217} &= x_1^3 x_2 x_3^{30} x_4^4 x_5^9, & \mathcal{V}_{47,218} &= x_1^3 x_2 x_3^4 x_4^9 x_5^{30}, & \mathcal{V}_{47,219} &= x_1 x_2^3 x_3^4 x_4^{10} x_5^{29}, \\
\mathcal{V}_{47,220} &= x_1^3 x_2 x_3^4 x_4^{10} x_5^{29}, & \mathcal{V}_{47,221} &= x_1^3 x_2^4 x_3 x_4^{10} x_5^{29}, & \mathcal{V}_{47,222} &= x_1 x_2^3 x_3^4 x_4^{11} x_5^{28}, \\
\mathcal{V}_{47,223} &= x_1^3 x_2 x_3^4 x_4^{11} x_5^{28}, & \mathcal{V}_{47,224} &= x_1^3 x_2^4 x_3 x_4^{11} x_5^{28}, & \mathcal{V}_{47,225} &= x_1^3 x_2^4 x_3^{11} x_4 x_5^{28}, \\
\mathcal{V}_{47,226} &= x_1 x_2^3 x_3^4 x_4^{27} x_5^{12}, & \mathcal{V}_{47,227} &= x_1^3 x_2 x_3^4 x_4^{27} x_5^{12}, & \mathcal{V}_{47,228} &= x_1^3 x_2^4 x_3 x_4^{27} x_5^{12}, \\
\mathcal{V}_{47,229} &= x_1^3 x_2^4 x_3^{27} x_4 x_5^{12}, & \mathcal{V}_{47,230} &= x_1 x_2^3 x_3^4 x_4^{26} x_5^{13}, & \mathcal{V}_{47,231} &= x_1^3 x_2 x_3^4 x_4^{26} x_5^{13}, \\
\mathcal{V}_{47,232} &= x_1^3 x_2^4 x_3 x_4^{26} x_5^{13}, & \mathcal{V}_{47,233} &= x_1 x_2^3 x_3^4 x_4^{14} x_5^{25}, & \mathcal{V}_{47,234} &= x_1 x_2^3 x_3^{14} x_4^4 x_5^{25}, \\
\mathcal{V}_{47,235} &= x_1 x_2^3 x_3^4 x_4^{25} x_5^{14}, & \mathcal{V}_{47,236} &= x_1^3 x_2 x_3^4 x_4^{14} x_5^{25}, & \mathcal{V}_{47,237} &= x_1^3 x_2 x_3^{14} x_4^4 x_5^{25}, \\
\mathcal{V}_{47,238} &= x_1^3 x_2 x_3^4 x_4^{25} x_5^{14}, & \mathcal{V}_{47,239} &= x_1 x_2^3 x_3^4 x_4^{15} x_5^{24}, & \mathcal{V}_{47,240} &= x_1 x_2^3 x_3^4 x_4^{24} x_5^{15}, \\
\mathcal{V}_{47,241} &= x_1 x_2^3 x_3^{15} x_4^4 x_5^{24}, & \mathcal{V}_{47,242} &= x_1 x_2^{15} x_3^3 x_4^4 x_5^{24}, & \mathcal{V}_{47,243} &= x_1^3 x_2 x_3^4 x_4^{15} x_5^{24}, \\
\mathcal{V}_{47,244} &= x_1^3 x_2 x_3^4 x_4^{24} x_5^{15}, & \mathcal{V}_{47,245} &= x_1^3 x_2 x_3^{15} x_4^4 x_5^{24}, & \mathcal{V}_{47,246} &= x_1^3 x_2^{15} x_3 x_4^4 x_5^{24}, \\
\mathcal{V}_{47,247} &= x_1^{15} x_2 x_3^3 x_4^4 x_5^{24}, & \mathcal{V}_{47,248} &= x_1^{15} x_2 x_3 x_4^4 x_5^{24}, & \mathcal{V}_{47,249} &= x_1 x_2^3 x_3^{30} x_4^5 x_5^8, \\
\mathcal{V}_{47,250} &= x_1 x_2^3 x_3^5 x_4^8 x_5^{30}, & \mathcal{V}_{47,251} &= x_1 x_2^3 x_3^5 x_4^{30} x_5^8, & \mathcal{V}_{47,252} &= x_1^3 x_2 x_3^{30} x_4^5 x_5^8, \\
\mathcal{V}_{47,253} &= x_1^3 x_2 x_3^5 x_4^8 x_5^{30}, & \mathcal{V}_{47,254} &= x_1^3 x_2 x_3^5 x_4^{30} x_5^8, & \mathcal{V}_{47,255} &= x_1 x_2^3 x_3^5 x_4^{10} x_5^{28}, \\
\mathcal{V}_{47,256} &= x_1^3 x_2 x_3^5 x_4^{10} x_5^{28}, & \mathcal{V}_{47,257} &= x_1^3 x_2^5 x_3 x_4^{10} x_5^{28}, & \mathcal{V}_{47,258} &= x_1^3 x_2^5 x_3^{10} x_4 x_5^{28}, \\
\mathcal{V}_{47,259} &= x_1^3 x_2^5 x_3^{10} x_4 x_5^{28}, & \mathcal{V}_{47,260} &= x_1 x_2^3 x_3^5 x_4^{26} x_5^{12}, & \mathcal{V}_{47,261} &= x_1^3 x_2 x_3^5 x_4^{26} x_5^{12}, \\
\mathcal{V}_{47,262} &= x_1^3 x_2^5 x_3 x_4^{26} x_5^{12}, & \mathcal{V}_{47,263} &= x_1^3 x_2^5 x_3^{26} x_4 x_5^{12}, & \mathcal{V}_{47,264} &= x_1^3 x_2^5 x_3^{26} x_4^{12} x_5^5, \\
\mathcal{V}_{47,265} &= x_1 x_2^3 x_3^{14} x_4^5 x_5^{24}, & \mathcal{V}_{47,266} &= x_1 x_2^3 x_3^{14} x_4^{24} x_5^{24}, & \mathcal{V}_{47,267} &= x_1 x_2^3 x_3^5 x_4^{24} x_5^{14}, \\
\mathcal{V}_{47,268} &= x_1^3 x_2 x_3^{14} x_4^5 x_5^{24}, & \mathcal{V}_{47,269} &= x_1^3 x_2 x_3^5 x_4^{14} x_5^{24}, & \mathcal{V}_{47,270} &= x_1^3 x_2 x_3^5 x_4^{24} x_5^{14}, \\
\mathcal{V}_{47,271} &= x_1 x_2^3 x_3^6 x_4^8 x_5^{29}, & \mathcal{V}_{47,272} &= x_1 x_2^3 x_3^6 x_4^{29} x_5^8, & \mathcal{V}_{47,273} &= x_1^3 x_2 x_3^6 x_4^8 x_5^{29}, \\
\mathcal{V}_{47,274} &= x_1^3 x_2 x_3^6 x_4^{29} x_5^8, & \mathcal{V}_{47,275} &= x_1 x_2^3 x_3^6 x_4^9 x_5^{28}, & \mathcal{V}_{47,276} &= x_1 x_2^3 x_3^6 x_4^9 x_5^{28}, \\
\mathcal{V}_{47,277} &= x_1^3 x_2 x_3^6 x_4^9 x_5^{28}, & \mathcal{V}_{47,278} &= x_1 x_2^3 x_3^6 x_4^{28} x_5^9, & \mathcal{V}_{47,279} &= x_1^3 x_2 x_3^6 x_4^{28} x_5^9, \\
\mathcal{V}_{47,280} &= x_1 x_2^3 x_3^6 x_4^{25} x_5^{12}, & \mathcal{V}_{47,281} &= x_1 x_2^6 x_3^3 x_4^{25} x_5^{12}, & \mathcal{V}_{47,282} &= x_1^3 x_2 x_3^6 x_4^{25} x_5^{12}, \\
\mathcal{V}_{47,283} &= x_1 x_2^3 x_3^6 x_4^{12} x_5^{25}, & \mathcal{V}_{47,284} &= x_1^3 x_2 x_3^6 x_4^{12} x_5^{25}, & \mathcal{V}_{47,285} &= x_1 x_2^3 x_3^6 x_4^{24} x_5^{13}, \\
\mathcal{V}_{47,286} &= x_1 x_2^3 x_3^6 x_4^{13} x_5^{24}, & \mathcal{V}_{47,287} &= x_1^3 x_2 x_3^6 x_4^{24} x_5^{13}, & \mathcal{V}_{47,288} &= x_1^3 x_2 x_3^6 x_4^{13} x_5^{24}, \\
\mathcal{V}_{47,289} &= x_1 x_2^3 x_3^7 x_4^8 x_5^{28}, & \mathcal{V}_{47,290} &= x_1 x_2^7 x_3^3 x_4^8 x_5^{28}, & \mathcal{V}_{47,291} &= x_1^3 x_2 x_3^7 x_4^8 x_5^{28}, \\
\mathcal{V}_{47,292} &= x_1^3 x_2^7 x_3 x_4^8 x_5^{28}, & \mathcal{V}_{47,293} &= x_1^7 x_2 x_3^3 x_4^8 x_5^{28}, & \mathcal{V}_{47,294} &= x_1^7 x_2^3 x_3 x_4^8 x_5^{28}, \\
\mathcal{V}_{47,295} &= x_1 x_2^7 x_3^3 x_4^8 x_5^{28}, & \mathcal{V}_{47,296} &= x_1 x_2^7 x_3^3 x_4^8 x_5^{28}, & \mathcal{V}_{47,297} &= x_1^7 x_2 x_3^3 x_4^8 x_5^{28}, \\
\mathcal{V}_{47,298} &= x_1^3 x_2^7 x_3 x_4^{28} x_5^8, & \mathcal{V}_{47,299} &= x_1^7 x_2 x_3^3 x_4^{28} x_5^8, & \mathcal{V}_{47,300} &= x_1^7 x_2^3 x_3 x_4^{28} x_5^8, \\
\mathcal{V}_{47,301} &= x_1 x_2^7 x_3^3 x_4^{24} x_5^{12}, & \mathcal{V}_{47,302} &= x_1 x_2^7 x_3^3 x_4^{24} x_5^{12}, & \mathcal{V}_{47,303} &= x_1^3 x_2 x_3^7 x_4^{24} x_5^{12}, \\
\mathcal{V}_{47,304} &= x_1^3 x_2^7 x_3 x_4^{24} x_5^{12}, & \mathcal{V}_{47,305} &= x_1^7 x_2 x_3^3 x_4^{24} x_5^{12}, & \mathcal{V}_{47,306} &= x_1^7 x_2^3 x_3 x_4^{24} x_5^{12}, \\
\mathcal{V}_{47,307} &= x_1 x_2^3 x_3^7 x_4^{12} x_5^{24}, & \mathcal{V}_{47,308} &= x_1 x_2^7 x_3^3 x_4^{12} x_5^{24}, & \mathcal{V}_{47,309} &= x_1^3 x_2 x_3^7 x_4^{12} x_5^{24}, \\
\mathcal{V}_{47,310} &= x_1^3 x_2^7 x_3 x_4^{12} x_5^{24}, & \mathcal{V}_{47,311} &= x_1^7 x_2 x_3^3 x_4^{12} x_5^{24}, & \mathcal{V}_{47,312} &= x_1^7 x_2^3 x_3 x_4^{12} x_5^{24}, \\
\mathcal{V}_{47,313} &= x_1 x_2^7 x_3^{27} x_4^4 x_5^8, & \mathcal{V}_{47,314} &= x_1^7 x_2 x_3^{27} x_4^4 x_5^8, & \mathcal{V}_{47,315} &= x_1^7 x_2^{27} x_3 x_4^4 x_5^8, \\
\mathcal{V}_{47,316} &= x_1 x_2^7 x_3^{11} x_4^4 x_5^{24}, & \mathcal{V}_{47,317} &= x_1^7 x_2 x_3^{11} x_4^4 x_5^{24}, & \mathcal{V}_{47,318} &= x_1^7 x_2^{11} x_3 x_4^4 x_5^{24}, \\
\mathcal{V}_{47,319} &= x_1^3 x_2^{29} x_3^2 x_4^9 x_5^5, & \mathcal{V}_{47,320} &= x_1^3 x_2^{13} x_3^2 x_4^9 x_5^{25}, & \mathcal{V}_{47,321} &= x_1^3 x_2^5 x_3^2 x_4^9 x_5^{29},
\end{aligned}$$

$$\begin{aligned}
\mathcal{V}_{47,322} &= x_1^3 x_2^5 x_3^2 x_4^{29} x_5^8, & \mathcal{V}_{47,323} &= x_1^3 x_2^{29} x_3^2 x_4^5 x_5^8, & \mathcal{V}_{47,324} &= x_1^3 x_2^5 x_3^2 x_4^9 x_5^{28}, \\
\mathcal{V}_{47,325} &= x_1^3 x_2^5 x_3^2 x_4^{28} x_5^9, & \mathcal{V}_{47,326} &= x_1^3 x_2^5 x_3^2 x_4^{25} x_5^{12}, & \mathcal{V}_{47,327} &= x_1^3 x_2^5 x_3^2 x_4^{12} x_5^{25}, \\
\mathcal{V}_{47,328} &= x_1^3 x_2^5 x_3^2 x_4^{24} x_5^{13}, & \mathcal{V}_{47,329} &= x_1^3 x_2^5 x_3^2 x_4^{13} x_5^{24}, & \mathcal{V}_{47,330} &= x_1^3 x_2^{13} x_3^2 x_4^5 x_5^{24}, \\
\mathcal{V}_{47,331} &= x_1^3 x_2^3 x_3^4 x_4^8 x_5^{29}, & \mathcal{V}_{47,332} &= x_1^3 x_2^3 x_3^4 x_4^{29} x_5^8, & \mathcal{V}_{47,333} &= x_1^3 x_2^3 x_3^2 x_4^9 x_5^{28}, \\
\mathcal{V}_{47,334} &= x_1^3 x_2^{29} x_3^3 x_4^5 x_5^{12}, & \mathcal{V}_{47,335} &= x_1^3 x_2^3 x_3^4 x_4^9 x_5^{28}, & \mathcal{V}_{47,336} &= x_1^3 x_2^3 x_3^2 x_4^9 x_5^{28}, \\
\mathcal{V}_{47,337} &= x_1^3 x_2^3 x_3^4 x_4^{25} x_5^{12}, & \mathcal{V}_{47,338} &= x_1^3 x_2^3 x_3^2 x_4^{25} x_5^{12}, & \mathcal{V}_{47,339} &= x_1^3 x_2^3 x_3^4 x_4^{24} x_5^{13}, \\
\mathcal{V}_{47,340} &= x_1^3 x_2^3 x_3^4 x_4^{13} x_5^{24}, & \mathcal{V}_{47,341} &= x_1^3 x_2^3 x_3^{13} x_4^5 x_5^{24}, & \mathcal{V}_{47,342} &= x_1^3 x_2^{13} x_3^3 x_4^5 x_5^{24}, \\
\mathcal{V}_{47,343} &= x_1^3 x_2^3 x_3^5 x_4^8 x_5^{28}, & \mathcal{V}_{47,344} &= x_1^3 x_2^5 x_3^3 x_4^8 x_5^{28}, & \mathcal{V}_{47,345} &= x_1^3 x_2^3 x_3^5 x_4^8 x_5^{28}, \\
\mathcal{V}_{47,346} &= x_1^3 x_2^5 x_3^3 x_4^{28} x_5^8, & \mathcal{V}_{47,347} &= x_1^3 x_2^3 x_3^5 x_4^{24} x_5^{12}, & \mathcal{V}_{47,348} &= x_1^3 x_2^5 x_3^3 x_4^{24} x_5^{12}, \\
\mathcal{V}_{47,349} &= x_1^3 x_2^3 x_3^5 x_4^{12} x_5^{24}, & \mathcal{V}_{47,350} &= x_1^3 x_2^5 x_3^3 x_4^{12} x_5^{24}, & \mathcal{V}_{47,351} &= x_1^3 x_2^5 x_3^{27} x_4^5 x_5^8, \\
\mathcal{V}_{47,352} &= x_1^3 x_2^5 x_3^{26} x_4^9 x_5^9, & \mathcal{V}_{47,353} &= x_1^3 x_2^5 x_3^{10} x_4^5 x_5^{25}, & \mathcal{V}_{47,354} &= x_1^3 x_2^5 x_3^{11} x_4^5 x_5^{24}, \\
\mathcal{V}_{47,355} &= x_1^3 x_2^7 x_3^{25} x_4^8 x_5^{24}, & \mathcal{V}_{47,356} &= x_1^7 x_2^3 x_3^{25} x_4^8 x_5^{24}, & \mathcal{V}_{47,357} &= x_1^3 x_2^7 x_3^9 x_4^5 x_5^{24}, \\
\mathcal{V}_{47,358} &= x_1^7 x_2^3 x_3^9 x_4^5 x_5^{24}, & \mathcal{V}_{47,359} &= x_1^3 x_2^5 x_3^{26} x_4^5 x_5^8, & \mathcal{V}_{47,360} &= x_1^3 x_2^5 x_3^{10} x_4^5 x_5^{24}, \\
\mathcal{V}_{47,361} &= x_1^3 x_2^5 x_3^{10} x_4^{21} x_5^8, & \mathcal{V}_{47,362} &= x_1^3 x_2^5 x_3^{11} x_4^{20} x_5^8, & \mathcal{V}_{47,363} &= x_1^3 x_2^5 x_3^{10} x_4^{20} x_5^9, \\
\mathcal{V}_{47,364} &= x_1^3 x_2^5 x_3^{10} x_4^{12} x_5^{17}, & \mathcal{V}_{47,365} &= x_1^3 x_2^5 x_3^{10} x_4^{13} x_5^{16}, & \mathcal{V}_{47,366} &= x_1^3 x_2^5 x_3^{11} x_4^{12} x_5^{16}, \\
\mathcal{V}_{47,367} &= x_1^3 x_2^7 x_3^9 x_4^{20} x_5^8, & \mathcal{V}_{47,368} &= x_1^7 x_2^3 x_3^9 x_4^{20} x_5^8, & \mathcal{V}_{47,369} &= x_1^3 x_2^7 x_3^9 x_4^{12} x_5^{16}, \\
\mathcal{V}_{47,370} &= x_1^7 x_2^3 x_3^9 x_4^{12} x_5^{16}.
\end{aligned}$$

$\mathcal{B}_5^+(\overline{\omega}_{(4)})$ is the set of 110 admissible monomials: $\mathcal{V}_{47,t}$, $371 \leq t \leq 480$

$$\begin{aligned}
\mathcal{V}_{47,371} &= x_1 x_2^3 x_3^6 x_4^6 x_5^{31}, & \mathcal{V}_{47,372} &= x_1 x_2^3 x_3^6 x_4^{31} x_5^6, & \mathcal{V}_{47,373} &= x_1 x_2^3 x_3^{31} x_4^6 x_5^6, \\
\mathcal{V}_{47,374} &= x_1 x_2^{31} x_3^3 x_4^6 x_5^6, & \mathcal{V}_{47,375} &= x_1^3 x_2 x_3^6 x_4^6 x_5^{31}, & \mathcal{V}_{47,376} &= x_1^3 x_2 x_3^6 x_4^{31} x_5^6, \\
\mathcal{V}_{47,377} &= x_1^3 x_2 x_3^{31} x_4^6 x_5^6, & \mathcal{V}_{47,378} &= x_1^3 x_2^{31} x_3 x_4^6 x_5^6, & \mathcal{V}_{47,379} &= x_1^{31} x_2 x_3^3 x_4^6 x_5^6, \\
\mathcal{V}_{47,380} &= x_1^{31} x_2^3 x_3 x_4^6 x_5^6, & \mathcal{V}_{47,381} &= x_1 x_2^3 x_3^6 x_4^7 x_5^{30}, & \mathcal{V}_{47,382} &= x_1 x_2^3 x_3^7 x_4^6 x_5^{30}, \\
\mathcal{V}_{47,383} &= x_1 x_2^3 x_3^7 x_4^{30} x_5^6, & \mathcal{V}_{47,384} &= x_1 x_2^7 x_3^3 x_4^6 x_5^{30}, & \mathcal{V}_{47,385} &= x_1 x_2^7 x_3^3 x_4^{30} x_5^6, \\
\mathcal{V}_{47,386} &= x_1^3 x_2 x_3^6 x_4^7 x_5^{30}, & \mathcal{V}_{47,387} &= x_1^3 x_2 x_3^7 x_4^6 x_5^{30}, & \mathcal{V}_{47,388} &= x_1^3 x_2 x_3^7 x_4^{30} x_5^6, \\
\mathcal{V}_{47,389} &= x_1^3 x_2^7 x_3 x_4^6 x_5^{30}, & \mathcal{V}_{47,390} &= x_1^3 x_2^7 x_3 x_4^{30} x_5^6, & \mathcal{V}_{47,391} &= x_1^7 x_2 x_3^3 x_4^6 x_5^{30}, \\
\mathcal{V}_{47,392} &= x_1^7 x_2 x_3^3 x_4^{30} x_5^6, & \mathcal{V}_{47,393} &= x_1^7 x_2^3 x_3 x_4^6 x_5^{30}, & \mathcal{V}_{47,394} &= x_1^7 x_2^3 x_3 x_4^{30} x_5^6, \\
\mathcal{V}_{47,395} &= x_1 x_2^3 x_3^7 x_4^{14} x_5^{22}, & \mathcal{V}_{47,396} &= x_1 x_2^7 x_3^3 x_4^{14} x_5^{22}, & \mathcal{V}_{47,397} &= x_1^3 x_2^3 x_3^7 x_4^{14} x_5^{22}, \\
\mathcal{V}_{47,398} &= x_1^3 x_2^7 x_3 x_4^{14} x_5^{22}, & \mathcal{V}_{47,399} &= x_1^7 x_2 x_3^3 x_4^{14} x_5^{22}, & \mathcal{V}_{47,400} &= x_1^7 x_2^3 x_3 x_4^{14} x_5^{22}, \\
\mathcal{V}_{47,401} &= x_1 x_2^7 x_3^{27} x_4^6 x_5^6, & \mathcal{V}_{47,402} &= x_1^7 x_2 x_3^{27} x_4^6 x_5^6, & \mathcal{V}_{47,403} &= x_1^7 x_2^7 x_3 x_4^6 x_5^6, \\
\mathcal{V}_{47,404} &= x_1 x_2^7 x_3^{11} x_4^6 x_5^{22}, & \mathcal{V}_{47,405} &= x_1 x_2^7 x_3^{11} x_4^{22} x_5^6, & \mathcal{V}_{47,406} &= x_1^7 x_2 x_3^{11} x_4^6 x_5^{22}, \\
\mathcal{V}_{47,407} &= x_1^7 x_2 x_3^{11} x_4^{22} x_5^6, & \mathcal{V}_{47,408} &= x_1^7 x_2^{11} x_3 x_4^6 x_5^{22}, & \mathcal{V}_{47,409} &= x_1^7 x_2^{11} x_3 x_4^{22} x_5^6, \\
\mathcal{V}_{47,410} &= x_1^3 x_2^5 x_3^2 x_4^6 x_5^{31}, & \mathcal{V}_{47,411} &= x_1^3 x_2^5 x_3^2 x_4^{31} x_5^6, & \mathcal{V}_{47,412} &= x_1^3 x_2^5 x_3^6 x_4^2 x_5^{31}, \\
\mathcal{V}_{47,413} &= x_1^3 x_2^5 x_3^6 x_4^2 x_5^{31}, & \mathcal{V}_{47,414} &= x_1^3 x_2^5 x_3^{31} x_4^2 x_5^6, & \mathcal{V}_{47,415} &= x_1^3 x_2^5 x_3^{31} x_4^6 x_5^2, \\
\mathcal{V}_{47,416} &= x_1^3 x_2^{31} x_3^5 x_4^6 x_5^2, & \mathcal{V}_{47,417} &= x_1^3 x_2^{31} x_3^5 x_4^2 x_5^6, & \mathcal{V}_{47,418} &= x_1^{31} x_2^3 x_3^5 x_4^6 x_5^2, \\
\mathcal{V}_{47,419} &= x_1^{31} x_2^3 x_3^5 x_4^2 x_5^6, & \mathcal{V}_{47,420} &= x_1^3 x_2^5 x_3^2 x_4^7 x_5^{30}, & \mathcal{V}_{47,421} &= x_1^3 x_2^5 x_3^7 x_4^2 x_5^{30}, \\
\mathcal{V}_{47,422} &= x_1^3 x_2^5 x_3^7 x_4^2 x_5^{30}, & \mathcal{V}_{47,423} &= x_1^3 x_2^7 x_3^5 x_4^2 x_5^{30}, & \mathcal{V}_{47,424} &= x_1^3 x_2^7 x_3^5 x_4^{30} x_5^2, \\
\mathcal{V}_{47,425} &= x_1^7 x_2^3 x_3^5 x_4^{30} x_5^2, & \mathcal{V}_{47,426} &= x_1^7 x_2^3 x_3^5 x_4^{30} x_5^2, & \mathcal{V}_{47,427} &= x_1^3 x_2^7 x_3^{29} x_4^2 x_5^6, \\
\mathcal{V}_{47,428} &= x_1^3 x_2^7 x_3^{29} x_4^2 x_5^6, & \mathcal{V}_{47,429} &= x_1^7 x_2^3 x_3^{29} x_4^2 x_5^6, & \mathcal{V}_{47,430} &= x_1^7 x_2^3 x_3^{29} x_4^6 x_5^2, \\
\mathcal{V}_{47,431} &= x_1^3 x_2^7 x_3^{13} x_4^2 x_5^{22}, & \mathcal{V}_{47,432} &= x_1^3 x_2^7 x_3^{13} x_4^{22} x_5^2, & \mathcal{V}_{47,433} &= x_1^7 x_2^3 x_3^{13} x_4^2 x_5^{22}, \\
\mathcal{V}_{47,434} &= x_1^7 x_2^3 x_3^{13} x_4^{22} x_5^2, & \mathcal{V}_{47,435} &= x_1^7 x_2^{27} x_3^5 x_4^2 x_5^6, & \mathcal{V}_{47,436} &= x_1^7 x_2^{27} x_3^5 x_4^6 x_5^2, \\
\mathcal{V}_{47,437} &= x_1^7 x_2^{11} x_3^5 x_4^2 x_5^{22}, & \mathcal{V}_{47,438} &= x_1^7 x_2^{11} x_3^5 x_4^{22} x_5^2, & \mathcal{V}_{47,439} &= x_1^7 x_2^{11} x_3^3 x_4^2 x_5^6, \\
\mathcal{V}_{47,440} &= x_1^7 x_2^{11} x_3^3 x_4^2 x_5^6, & \mathcal{V}_{47,441} &= x_1^3 x_2^5 x_3^6 x_4^{30} x_5^3, & \mathcal{V}_{47,442} &= x_1^3 x_2^5 x_3^6 x_4^3 x_5^{30}, \\
\mathcal{V}_{47,443} &= x_1^3 x_2^5 x_3^6 x_4^3 x_5^{30}, & \mathcal{V}_{47,444} &= x_1^3 x_2^3 x_3^5 x_4^{30} x_5^6, & \mathcal{V}_{47,445} &= x_1^3 x_2^5 x_3^3 x_4^6 x_5^{30}, \\
\mathcal{V}_{47,446} &= x_1^3 x_2^5 x_3^3 x_4^{30} x_5^6, & \mathcal{V}_{47,447} &= x_1^3 x_2^3 x_3^5 x_4^{14} x_5^{22}, & \mathcal{V}_{47,448} &= x_1^3 x_2^5 x_3^3 x_4^{14} x_5^{22}, \\
\mathcal{V}_{47,449} &= x_1^3 x_2^3 x_3^{29} x_4^6 x_5^6, & \mathcal{V}_{47,450} &= x_1^3 x_2^{29} x_3^3 x_4^6 x_5^6, & \mathcal{V}_{47,451} &= x_1^3 x_2^3 x_3^{13} x_4^6 x_5^{22}, \\
\mathcal{V}_{47,452} &= x_1^3 x_2^3 x_3^{13} x_4^{22} x_5^6, & \mathcal{V}_{47,453} &= x_1^3 x_2^5 x_3^6 x_4^{27} x_5^6, & \mathcal{V}_{47,454} &= x_1^3 x_2^5 x_3^{27} x_4^6 x_5^6,
\end{aligned}$$

$$\begin{aligned}
\mathcal{V}_{47,455} &= x_1^3 x_2^5 x_3^6 x_4^7 x_5^{26}, & \mathcal{V}_{47,456} &= x_1^3 x_2^5 x_3^7 x_4^6 x_5^{26}, & \mathcal{V}_{47,457} &= x_1^3 x_2^5 x_3^7 x_4^{26} x_5^6, \\
\mathcal{V}_{47,458} &= x_1^3 x_2^7 x_3^5 x_4^6 x_5^{26}, & \mathcal{V}_{47,459} &= x_1^3 x_2^7 x_3^5 x_4^{26} x_5^6, & \mathcal{V}_{47,460} &= x_1^7 x_2^3 x_3^5 x_4^6 x_5^{26}, \\
\mathcal{V}_{47,461} &= x_1^7 x_2^3 x_3^5 x_4^{26} x_5^6, & \mathcal{V}_{47,462} &= x_1^3 x_2^5 x_3^{11} x_4^6 x_5^{22}, & \mathcal{V}_{47,463} &= x_1^3 x_2^5 x_3^{11} x_4^{22} x_5^6, \\
\mathcal{V}_{47,464} &= x_1^3 x_2^5 x_3^6 x_4^{14} x_5^{19}, & \mathcal{V}_{47,465} &= x_1^3 x_2^5 x_3^7 x_4^{10} x_5^{22}, & \mathcal{V}_{47,466} &= x_1^3 x_2^7 x_3^5 x_4^{10} x_5^{22}, \\
\mathcal{V}_{47,467} &= x_1^7 x_2^3 x_3^5 x_4^{10} x_5^{22}, & \mathcal{V}_{47,468} &= x_1^3 x_2^5 x_3^7 x_4^{14} x_5^{18}, & \mathcal{V}_{47,469} &= x_1^3 x_2^7 x_3^5 x_4^{14} x_5^{18}, \\
\mathcal{V}_{47,470} &= x_1^7 x_2^3 x_3^5 x_4^{14} x_5^{18}, & \mathcal{V}_{47,471} &= x_1^3 x_2^7 x_3^{25} x_4^6 x_5^6, & \mathcal{V}_{47,472} &= x_1^7 x_2^3 x_3^{25} x_4^6 x_5^6, \\
\mathcal{V}_{47,473} &= x_1^3 x_2^7 x_3^9 x_4^6 x_5^{22}, & \mathcal{V}_{47,474} &= x_1^3 x_2^7 x_3^9 x_4^{22} x_5^6, & \mathcal{V}_{47,475} &= x_1^7 x_2^3 x_3^9 x_4^6 x_5^{22}, \\
\mathcal{V}_{47,476} &= x_1^7 x_2^3 x_3^9 x_4^{22} x_5^6, & \mathcal{V}_{47,477} &= x_1^3 x_2^7 x_3^{13} x_4^6 x_5^{18}, & \mathcal{V}_{47,478} &= x_1^3 x_2^7 x_3^{13} x_4^{18} x_5^6, \\
\mathcal{V}_{47,479} &= x_1^7 x_2^3 x_3^{13} x_4^6 x_5^{18}, & \mathcal{V}_{47,480} &= x_1^7 x_2^3 x_3^{13} x_4^{18} x_5^6.
\end{aligned}$$

$\mathcal{B}_5^+(\overline{\omega}_{(5)})$ is the set of 14 admissible monomials: $\mathcal{V}_{47,t}$, $481 \leq t \leq 494$

$$\begin{aligned}
\mathcal{V}_{47,481} &= x_1 x_2^7 x_3^{11} x_4^{14} x_5^{14}, & \mathcal{V}_{47,482} &= x_1^7 x_2 x_3^{11} x_4^{14} x_5^{14}, & \mathcal{V}_{47,483} &= x_1^7 x_2^{11} x_3 x_4^{14} x_5^{14}, \\
\mathcal{V}_{47,484} &= x_1^7 x_2^{11} x_3^{13} x_4^2 x_5^{14}, & \mathcal{V}_{47,485} &= x_1^7 x_2^{11} x_3^{13} x_4^{14} x_5^2, & \mathcal{V}_{47,486} &= x_1^3 x_2^7 x_3^9 x_4^{14} x_5^{14}, \\
\mathcal{V}_{47,487} &= x_1^7 x_2^3 x_3^9 x_4^{14} x_5^{14}, & \mathcal{V}_{47,488} &= x_1^3 x_2^7 x_3^{13} x_4^{10} x_5^{14}, & \mathcal{V}_{47,489} &= x_1^3 x_2^7 x_3^{13} x_4^{14} x_5^{10}, \\
\mathcal{V}_{47,490} &= x_1^7 x_2^3 x_3^{13} x_4^{10} x_5^{14}, & \mathcal{V}_{47,491} &= x_1^7 x_2^3 x_3^{13} x_4^{14} x_5^{10}, & \mathcal{V}_{47,492} &= x_1^7 x_2^{11} x_3^5 x_4^{10} x_5^{14}, \\
\mathcal{V}_{47,493} &= x_1^7 x_2^{11} x_3^5 x_4^{10} x_5^{14}, & \mathcal{V}_{47,494} &= x_1^7 x_2^{11} x_3^{13} x_4^6 x_5^{10}.
\end{aligned}$$

References

- [1] J.F. Adams, *On the structure and applications of the Steenrod algebra*, Comment. Math. Helv. 32 (1958), 180-214, <https://doi.org/10.1007/BF02564578>.
- [2] M.A. Alghamdi, M.C. Crabb and J.R. Hubbuck, *Representations of the homology of BV and the Steenrod algebra I*, in Adams Memorial Symposium on Algebraic Topology 2, N. Ray and G. Walker (ed.), London Math. Soc. Lecture Note Ser. 176 (1992), 217-234, available at <http://citeseerx.ist.psu.edu/viewdoc/download?doi=10.1.1.22.7436&rep=rep1&type=pdf>.
- [3] S.V. Ault and W. Singer, *On the homology of elementary Abelian groups as modules over the Steenrod algebra*, J. Pure Appl. Algebra 215 (2011), 2847-2852, <https://doi.org/10.1016/j.jpaa.2011.04.004>.
- [4] J.M. Boardman, *Modular representations on the homology of power of real projective space*, in Algebraic Topology: Oaxtepec 1991, ed. M. C. Tangor; in Contemp. Math. 146 (1993), 49-70, <http://dx.doi.org/10.1090/conm/146>.
- [5] R.R. Bruner, L.M. Hà and N.H.V. Húng, *On behavior of the algebraic transfer*, Trans. Amer. Math. Soc. 357 (2005), 437-487, <https://doi.org/10.1090/S0002-9947-04-03661-X>.
- [6] T.W. Chen, *Determination of $\text{Ext}_{\mathcal{A}}^{5,*}(\mathbb{Z}/2, \mathbb{Z}/2)$* , Topol. Appl. 158 (2011), 660-689, <https://doi.org/10.1016/j.topol.2011.01.002>.
- [7] P.H. Chơn and L.M. Hà, *Lambda algebra and the Singer transfer*, C. R. Math. Acad. Sci. Paris 349 (2011), 21-23, <https://doi.org/10.1016/j.crma.2010.11.008>.
- [8] M.C. Crabb and J.R. Hubbuck, *Representations of the homology of BV and the Steenrod algebra II*, in Algebra Topology: New trend in localization and periodicity; in Progr. Math. 136 (1996), 143-154, https://doi.org/10.1007/978-3-0348-9018-2_9.
- [9] M.D. Crossley, *Monomial bases for $H^*(\mathbb{C}P^\infty \times \mathbb{C}P^\infty)$ over $\mathcal{A}(p)$* , Trans. Amer. Math. Soc. 351(1) (1999), 171-192, <https://doi.org/10.1090/S0002-9947-99-02060-7>.
- [10] M.D. Crossley, *$\mathcal{A}(p)$ generators for $H^*(V)$ and Singer's homological transfer*, Math. Z. 230(3) (1999), 401-411, <https://doi.org/10.1007/PL00004698>.
- [11] N.H.V. Húng and F.P. Peterson, *A-generators for the Dickson algebra*, Trans. Am. Math. Soc. 347(12) (1995), 4687-4728, <https://doi.org/10.1090/S0002-9947-1995-1316852-X>.
- [12] N.H.V. Húng and T.N. Nam, *The hit problem for the Dickson algebra*, Trans. Am. Math. Soc. 353(12) (2001), 5029-5040, <https://doi.org/10.1090/S0002-9947-01-02705-2>.
- [13] N.H.V. Húng, *The cohomology of the Steenrod algebra and representations of the general linear groups*, Trans. Amer. Math. Soc. 357 (2005), 4065-4089, <https://doi.org/10.1090/S0002-9947-05-03889-4>.
- [14] A.S. Janfada and R.M.W. Wood, *The hit problem for symmetric polynomials over the Steenrod algebra*, Math. Proc. Cambridge Philos. Soc. 133(2) (2002), 295-303, <https://doi.org/10.1017/S0305004102006059>.
- [15] M. Kameko, *Products of projective spaces as Steenrod modules*, PhD. thesis, The Johns Hopkins University, ProQuest LLC, Ann Arbor, MI, 1990, 29 pages.
- [16] M.H. Lê, *Sub-Hopf algebras of the Steenrod algebra and the Singer transfer*, Geom. Topol. Publ. 11 (2007), 101-124, <https://doi.org/10.2140/gtm.2007.11.81>.
- [17] W.H. Lin, *$\text{Ext}_{\mathcal{A}}^{4,*}(\mathbb{Z}/2, \mathbb{Z}/2)$ and $\text{Ext}_{\mathcal{A}}^{5,*}(\mathbb{Z}/2, \mathbb{Z}/2)$* , Topol. Appl. 155 (2008), 459-496, <https://doi.org/10.1016/j.topol.2007.11.003>.
- [18] A. Liulevicius, *The factorization of cyclic reduced powers by secondary cohomology operations*, Mem. Amer. Math. Soc. 42 (1962), 112, <http://dx.doi.org/10.1090/memo/0042>.
- [19] Ib H. Madsen, *On the action of the Dyer-Lashof algebra in $H_*(G)$* , Pac. J. Math. 60(1) (1975), 235-275, <https://doi.org/10.2140/pjm.1975.60.235>.

- [20] N. Minami, *The iterated transfer analogue of the new doomsday conjecture*, Trans. Amer. Math. Soc. 351 (1990), 2325-2351, <https://doi.org/10.1090/S0002-9947-99-02037-1>.
- [21] M.F. Mothebe, *Generators of the polynomial algebra $\mathbb{F}_2[x_1, x_2, \dots, x_n]$ as a module over the Steenrod algebra*, PhD. thesis, The University of Manchester, 1997.
- [22] M.F. Mothebe, *Dimension result for the polynomial algebra $\mathbb{F}_2[x_1, x_2, \dots, x_n]$ as a module over the Steenrod algebra*, Int. J. Math. Math. Sci. 2013, Article ID 150704, 6 pages, <https://doi.org/10.1155/2013/150704>.
- [23] M.F. Mothebe, *Products of admissible monomials in the polynomial algebra as a module over the Steenrod algebra*, Journal of Mathematics Research 8 (2016), 112-116, <https://doi.org/10.5539/jmr.v8n3p112>.
- [24] M.F. Mothebe, P. Kaelo and O. Ramatebele, *Dimension formula for the polynomial algebra as a module over the Steenrod algebra in degrees less than or equal to 12*, Journal of Mathematics Research 8 (2016), 92-100, <https://doi.org/10.5539/jmr.v8n5p92>.
- [25] T.N. Nam, *$\mathcal{A}$ -générateurs génériques pour l'algèbre polynomiale*, Adv. Math. 186 (2004) 334-362, <https://doi.org/10.1016/j.aim.2003.08.004>.
- [26] T.N. Nam, *Transfert algébrique et action du groupe linéaire sur les puissances divisées modulo 2*, Ann. Inst. Fourier (Grenoble) 58 (2008), 1785-1837, <https://doi.org/10.5802/aif.2399>.
- [27] D.J. Pengelley and F. Williams, *Global structure of the mod two symmetric algebra, $H^*(BO, \mathbb{F}_2)$, over the Steenrod algebra*, Algebr. Geom. Topol. 3 (2003), 1119-1138, <https://doi.org/10.2140/agt.2003.3.1119>.
- [28] D.J. Pengelley and F. Williams, *Beyond the hit problem: Minimal presentations of odd-primary Steenrod modules, with application to $CP(\infty)$ and BU* , Homology Homotopy Appl. 9(2) (2007), 363-395, <https://doi.org/10.4310/HHA.2007.v9.n2.a13>.
- [29] D. Pengelley and F. Williams, *A new action of the Kudo-Araki-May algebra on the dual of the symmetric algebras, with applications to the hit problem*, Algebr. Geom. Topol. 11(3), (2011), 1767-1780, <https://doi.org/10.2140/agt.2011.11.1767>.
- [30] D.J. Pengelley and F. Williams, *The hit problem for $H^*(BU(2); \mathbb{F}_p)$* , Algebr. Geom. Topol. 13 (2013), 2061-2085, <https://doi.org/10.2140/agt.2013.13.2061>.
- [31] D. Pengelley and F. Williams, *Sparseness for the symmetric hit problem at all primes*, Math. Proc. Cambridge Philos. Soc. 158(2), (2015), 269-274, <https://doi.org/10.1017/S0305004114000668>.
- [32] F.P. Peterson, *Generators of $H^*(\mathbb{R}P^\infty \times \mathbb{R}P^\infty)$ as a module over the Steenrod algebra*, Abstracts Amer. Math. Soc. 833 (1987).
- [33] F.P. Peterson, *$\mathcal{A}$ -generators for certain polynomial algebras*, Math. Proc. Cambridge Philos. Soc. 105 (1989), 311-312, <https://doi.org/10.1017/S0305004100067803>.
- [34] Đ.V. Phúc and N. Sum, *On the generators of the polynomial algebra as a module over the Steenrod algebra*, C.R.Math. Acad. Sci. Paris 353 (2015), 1035-1040, <https://doi.org/10.1016/j.crma.2015.09.002>.
- [35] D.V. Phuc and N. Sum, *On a minimal set of generators for the polynomial algebra of five variables as a module over the Steenrod algebra*, Acta Math. Vietnam. 42 (2017), 149-162, <https://doi.org/10.1007/s40306-016-0190-z>.
- [36] D.V. Phuc, *The hit problem for the polynomial algebra of five variables in degree seventeen and its application*, East-West J. Math. 18 (2016), 27-46, available at eastwestmath.org/index.php/ewm/article/view/35.
- [37] Đ.V. Phúc, *The "hit" problem of five variables in the generic degree and its application*, Topol. Appl. (2020) 107321, <https://doi.org/10.1016/j.topol.2020.107321>, in press.
- [38] Đ.V. Phúc, *$\mathcal{A}$ -generators for the polynomial algebra of five variables in degree $5(2^t - 1) + 6 \cdot 2^t$* , Commun. Korean Math. Soc. 35 (2020), 371-399, <https://doi.org/10.4134/CKMS.c190076>.
- [39] S. Priddy, *On characterizing summands in the classifying space of a group, I*, Amer. Jour. Math. 112 (1990), 737-748, <https://doi.org/10.2307/2374805>.
- [40] J. Repka and P. Selick, *On the subalgebra of $H_*(\mathbb{R}P^\infty)^n; \mathbb{F}_2$ annihilated by Steenrod operations*, J. Pure Appl. Algebra 127 (1998), 273-288, [https://doi.org/10.1016/S0022-4049\(96\)00177-6](https://doi.org/10.1016/S0022-4049(96)00177-6).
- [41] J.H. Silverman and W.M. Singer, *On the action of Steenrod squares on polynomial algebras II*, J. Pure Appl. Algebra 98 (1995), 95-103, [https://doi.org/10.1016/0022-4049\(95\)90027-6](https://doi.org/10.1016/0022-4049(95)90027-6).
- [42] W.M. Singer, *The transfer in homological algebra*, Math. Z. 202 (1989), 493-523, <https://doi.org/10.1007/BF01221587>.
- [43] W.M. Singer, *On the action of the Steenrod squares on polynomial algebras*, Proc. Amer. Math. Soc. 111 (1991), 577-583, <https://doi.org/10.1090/S0002-9939-1991-1045150-9>.
- [44] W.M. Singer, *Rings of symmetric functions as modules over the Steenrod algebra*, Algebr. Geom. Topol. 8 (2008), 541-562, <https://doi.org/10.2140/agt.2008.8.541>.
- [45] N.E. Steenrod and D.B.A. Epstein, *Cohomology operations*, Annals of Mathematics Studies 50, Princeton University Press, Princeton N.J., 1962.
- [46] N. Sum, *The negative answer to Kameko's conjecture on the hit problem*, Adv. Math. 225 (2010), 2365-2390, <https://doi.org/10.1016/j.aim.2010.04.026>.
- [47] N. Sum, *On the Peterson hit problem*, Adv. Math. 274 (2015), 432-489, <https://doi.org/10.1016/j.aim.2015.01.010>.
- [48] N. Sum, *On the determination of the Singer transfer*, Vietnam Journal of Science, Technology and Engineering, 60(1) (2018), 3-16, [https://doi.org/10.31276/VJSTE.60\(1\).03](https://doi.org/10.31276/VJSTE.60(1).03), available at <http://vietnamscience.vjst.vn/index.php/VJSTE/article/view/100>.
- [49] N. Sum, *On a construction for the generators of the polynomial algebra as a module over the Steenrod algebra*, in Singh M., Song Y., Wu J. (eds), Algebraic Topology and Related Topics. Trends in Mathematics. Birkhäuser/Springer, Singapore (2019), 265-286, https://doi.org/10.1007/978-981-13-5742-8_14.
- [50] N. Sum, *The kernel of Kameko's homomorphism and the Peterson hit problem*, Preprint 2019, 31 pages, VIASM.
- [51] N. Sum, *The squaring operation and the Singer algebraic transfer*, Vietnam J. Math. (2020),

- <https://doi.org/10.1007/s10013-020-00423-1>, in press.
- [52] M.C. Tangora, *On the cohomology of the Steenrod algebra*, Math. Z. 116 (1970), 18-64, <https://doi.org/10.1007/BF01110185>.
 - [53] N.K. Tin, *The admissible monomial basis for the polynomial algebra of five variables in degree eight*, Journal of Mathematical Sciences and Applications 2 (2014), 21-24, <https://doi.org/10.12691/jmsa-2-2-2>, available at <http://pubs.sciepub.com/jmsa/2/2/2/>.
 - [54] N.K. Tin, *The admissible monomial basis for the polynomial algebra of five variables in degree $2^{s+1} + 2^s - 5$* , East-West J. Math. 16 (2014), 34-46, available at eastwestmath.org/index.php/ewm/article/view/68.
 - [55] N.K. Tin, *On Singer's conjecture for the fifth algebraic transfer*, Preprint 2016, 25 pages; available online at <http://arxiv.org/abs/1609.02250>.
 - [56] N.K. Tin, *The admissible monomial basis for the polynomial algebra as a module over Steenrod algebra in some degrees*, JP J. Algebra Number Theory Appl. 46(1) (2020), 55-68, <http://dx.doi.org/10.17654/NT046010055>.
 - [57] N.K. Tin, *The hit problem for the polynomial algebra as a module over Steenrod algebra, I*, JP J. Algebra Number Theory Appl. 47(1) (2020), 67-86, <http://dx.doi.org/10.17654/NT047010067>.
 - [58] G. Walker and R.M.W. Wood, *Young tableaux and the Steenrod algebra*, in Proceedings of the School and Conference in Algebraic Topology, Hanoi 2004; in Geom. Topol. Monogr. 11 (2007), 379-397, <http://doi.org/10.2140/gtm.2007.11.379>.
 - [59] G. Walker and R.M.W. Wood, *Weyl modules and the mod 2 Steenrod algebra*, J. Algebra 311 (2007), 840-858, <http://doi.org/10.1016/j.jalgebra.2007.01.021>.
 - [60] G. Walker and R.M.W. Wood, *Polynomials and the mod 2 Steenrod Algebra: Volume 1, The Peterson hit problem*, in London Math. Soc. Lecture Note Ser., Cambridge Univ. Press, January 11, 2018.
 - [61] R.M.W. Wood, *Steenrod squares of polynomials and the Peterson conjecture*, Math. Proc. Cambriges Phil. Soc. 105 (1989), 307-309, <https://doi.org/10.1017/S0305004100067797>.
 - [62] R.M.W. Wood, *Problems in the Steenrod algebra*, Bull. London Math. Soc. 30 (1998), 449-517, <https://doi.org/10.1112/S002460939800486X>.
 - [63] H. Zare, *On the relation between Dyer-Lashof algebra and the hit problems*, Preprint 2016, 15 pages.