

On Peterson's open problem and representations of the general linear groups

Đặng Võ Phúc

*Faculty of Education Studies, University of Khanh Hoa,
01 Nguyen Chanh, Nha Trang, Khanh Hoa, Vietnam*

Abstract

Fix $\mathbb{Z}/2$ is the prime field of two elements and write $\mathcal{A}_2$ for the mod 2 Steenrod algebra. Denote by $GL_d := GL(d, \mathbb{Z}/2)$ the general linear group of rank d over $\mathbb{Z}/2$ and by $\mathcal{P}_d$ the polynomial algebra $\mathbb{Z}/2[x_1, x_2, \dots, x_d]$ as an unstable left $\mathcal{A}_2$ -module on d generators of degree one. We study the *Peterson hit problem* of finding a minimal set of generators for $\mathcal{A}_2$ -module $\mathcal{P}_d$. Equivalently, we need to determine a basis for the $\mathbb{Z}/2$ -vector space

$$Q\mathcal{P}_d := \mathbb{Z}/2 \otimes_{\mathcal{A}_2} \mathcal{P}_d \cong \mathcal{P}_d / \mathcal{A}_2^+ \mathcal{P}_d$$

in each degree $n \geq 1$. Note that this space is a representation of GL_d over $\mathbb{Z}/2$. The problem is still open for $d \geq 5$.

In this paper, we explicitly compute the hit problem in the 5-variable case and in generic degree $n = r(2^t - 1) + 2^t s$ with $r = d = 5$, $s = 8$ and t an arbitrary positive integer. This result is used to prove that the fifth algebraic transfer Tr_5 is an isomorphism in the bidegree $(5, 26)$. As a consequence, Singer's conjecture for Tr_5 is true in the internal degree $13 \cdot 2^t - 5$ with $t = 1$. This transfer was defined by himself [29] as an algebraic version of the geometrical transfer $tr_d : \pi_*^{\mathbb{S}}(B(\mathbb{Z}/2)_+^{\times d}) \rightarrow \pi_*^{\mathbb{S}}(\mathbb{S}^0)$.

Keywords:

Adams spectral sequences; Steenrod algebra; Peterson hit problem; Algebraic transfer

2010 MSC: Primary 55Q45, 55S10; Secondary 55S05, 55T15.

Contents

1	Introduction and main results	2
2	Preliminaries	5
2.1	Kameko's squaring operation	5
2.2	The weight vector of a monomial and the admissible monomial	6
2.3	Singer's criterion on the $\mathcal{A}_2$ -decomposable	8
3	Generators of $\mathcal{A}_2$-module P_5 in generic degree $13 \cdot 2^t - 5$	9
3.1	Some homomorphisms and Sum's conjecture	9
3.2	Proof of Theorem 1.1	11
3.2.1	The case $t = 1$	12
3.2.2	The case $t = 2$	15

Email address: dangphuc150497@gmail.com (Đặng Võ Phúc)

4	An application of Theorem 1.1	31
4.1	Computation of $(Q\mathcal{P}_5)_8^{GL_5}$	32
4.2	Computation of $(\text{Ker}(Sq_*^0)_{(5,21)})^{S_5}$	34
4.3	Proof of Theorem 1.2	37
5	Proof of Theorem 1.3	37
6	Appendix	40
6.1	$\mathcal{A}_2$ -generators for $\mathcal{P}_5$ in degree 8	40
6.2	$\mathcal{A}_2$ -generators for $\mathcal{P}_5^0$ in degree 21	41
6.3	$\mathcal{A}_2$ -generators for $\mathcal{P}_5^+$ in degree 21	43
6.4	$\mathcal{A}_2$ -generators for $\mathcal{P}_5^0$ in degree 22	45
6.5	$\mathcal{A}_2$ -generators for $\mathcal{P}_5^+$ in degree 22	47
6.6	$\mathcal{A}_2$ -generators for $\mathcal{P}_5^0$ in degree 47	51
6.7	$\mathcal{A}_2$ -generators for $\mathcal{P}_5^+$ in degree 47	54

1. Introduction and main results

Let $Sq^k : H^*(\mathbb{X}) \rightarrow H^{k+*}(\mathbb{X})$ be the stable cohomology operation of degree $k \geq 0$, which is introduced by Steenrod in 1947 (see [32]). Here $H^*(\mathbb{X})$ is the singular cohomology group of the topological space $\mathbb{X}$ with coefficients in the field $\mathbb{Z}/2$ of two elements. The $\mathbb{Z}/2$ -graded algebra $\mathcal{A}_2$ generated by the operations Sq^k is called *the mod 2 Steenrod algebra* and acts in a natural way on the cohomology of any space $\mathbb{X}$.

For d a natural number, we denote by $B(\mathbb{Z}/2)^{\times d}$ the classifying space of elementary abelian 2-group $(\mathbb{Z}/2)^{\times d}$ of rank d and by $\mathcal{P}_d = \mathbb{Z}/2[x_1, x_2, \dots, x_d]$ the polynomial algebra on d variables of degree 1. This algebra is isomorphic to the cohomology with $\mathbb{Z}/2$ -coefficients of $B(\mathbb{Z}/2)^{\times d}$, hence $\mathcal{P}_d$ has an unstable left $\mathcal{A}_2$ -module structure. The left action of $\mathcal{A}_2$ on $\mathcal{P}_d$ is determined by the elementary properties of the Steenrod operations Sq^k , in grading $k \geq 0$ and the Cartan formula (see [32]).

We are interested in Peterson's "hit" problem of determining a minimal set of generators for left $\mathcal{A}_2$ -module $\mathcal{P}_d$. Equivalently, we need to find a basis for the $\mathbb{Z}/2$ -graded vector space

$$Q\mathcal{P}_d := \mathbb{Z}/2 \otimes_{\mathcal{A}_2} \mathcal{P}_d \cong \mathcal{P}_d / \mathcal{A}_2^+ \mathcal{P}_d,$$

where $\mathcal{A}_2^+$ denotes the *augmentation ideal* of $\mathcal{A}_2$ and $\mathbb{Z}/2$ is viewed as a right $\mathcal{A}_2$ -module concentrated in grading 0.

For a non-negative integer n , let $(\mathcal{P}_d)_n$ be the subspace of $\mathcal{P}_d$ consisting of all the homogeneous polynomials of degree n in $\mathcal{P}_d$. Denote by $(Q\mathcal{P}_d)_n$ the subspace of $Q\mathcal{P}_d$ consisting of all the classes represented by the homogeneous polynomials in $(\mathcal{P}_d)_n$. As well known, the general linear group $GL_d = GL(d, \mathbb{Z}/2)$ acts regularly on $\mathcal{P}_d$ by matrix substitution. Further, the two actions of $\mathcal{A}_2$ and GL_d upon $\mathcal{P}_d$ commute with each other; hence there is an inherited action of GL_d on $Q\mathcal{P}_d$. From this event, the hit problem becomes one of the main tools for the studying of representations of the general linear groups over $\mathbb{Z}/2$. Moreover, according to Peterson [20], Priddy [26], Walker and his collaborator [43, 46], it has a deep connection with classical problems in the homotopy theory. The problem was then studied by many authors (see Ault and Singer [2], Crabb and Hubbuck [7], Kameko [9], Mothebe and his collaborators [13]-[16], Nam [17], Silverman and Singer [28], Singer [30, 31], Peterson [20], the present author and Sum [21]-[25], Sum [33]-[37], T  n [38]-[41], T  n and Sum [42], Walker and Wood [43, 44], Wood [45, 46] and others).

Let $P_{\mathcal{A}_2} H_*(B(\mathbb{Z}/2)^{\times d})$ be the subspace of $H_*(B(\mathbb{Z}/2)^{\times d})$ consisting of all elements that are $\mathcal{A}_2^+$ -annihilated. With the idea of studying the cohomology of the Steenrod algebra by means of

modular representations of the general linear groups, William Singer [29] constructed a transfer homomorphism from GL_d -coinvariants of the $\mathcal{A}_2^+$ -annihilated elements of the dual of $\mathcal{P}_d$ to the cohomology of the Steenrod algebra:

$$Tr_d : \mathbb{Z}/2 \otimes_{GL_d} P_{\mathcal{A}_2} H_*(B(\mathbb{Z}/2)^{\times d}) \rightarrow \text{Ext}_{\mathcal{A}_2}^{d,d+*}(\mathbb{Z}/2, \mathbb{Z}/2)$$

which is related to the geometrical transfer $tr_d : \pi_*^{\mathbb{S}}(B(\mathbb{Z}/2)_+^{\times d}) \rightarrow \pi_*^{\mathbb{S}}(\mathbb{S}^0)$ of the stable homotopy of spheres. More explicitly, tr_d induces Tr_d at the E_2 -page of the Adams spectral sequence [1]. The Singer transfer was studied by many authors (see Boardman [3], Bruner and his collaborators [4], Chơn and Hà [6], Hà [8], Minami [12], Nam [18], the present author [23, 24], Sum [35]-[37], Tín [41], Tín-Sum [42] and others).

Singer [29] has indicated the importance of this transfer by proving that Tr_d is an isomorphism with $d \leq 2$ and at some degrees with $d = 3, 4$. Latter, Boardman [3] confirmed this again by using a basis consisting of the all the classes represented by certain polynomials in $\mathcal{P}_3$ to show that Tr_3 is also an isomorphism. Therefore, the Singer transfer expected to be a useful tool in describing the cohomology groups of the Steenrod algebra, $\text{Ext}_{\mathcal{A}_2}^{d,*}(\mathbb{Z}/2, \mathbb{Z}/2)$.

In [29], Singer proved that Tr_5 is not an isomorphism in bidegree $(5, 14)$ and conjectured that Tr_d is a monomorphism for any positive integer d . By Boardman [3] and Singer [29], the conjecture is true for $d \leq 3$. However, for $d > 3$, it is still open. For $d = 4$, it can be verified by using the results in [33, 34].

One of the extremely useful tools for computing the hit problem and studying Singer's transfer is the Kameko squaring operation [9]

$$\widetilde{Sq}_*^0 = (\widetilde{Sq}_*^0)_{(d, 2n+d)} : (Q\mathcal{P}_d)_{2n+d} \rightarrow (Q\mathcal{P}_d)_n.$$

This operation is an $\mathbb{Z}/2GL_d$ -homomorphism induced by the $\mathbb{Z}/2$ -linear map

$$\delta : (\mathcal{P}_d)_{2n+d} \longrightarrow (\mathcal{P}_d)_n,$$

given by $\delta(x = x_1^{t_1} x_2^{t_2} \dots x_d^{t_d}) = x_1^{\frac{t_1-1}{2}} x_2^{\frac{t_2-1}{2}} \dots x_d^{\frac{t_d-1}{2}}$, if $t_1, t_2, \dots, t_d$ odd while $\delta(x) = 0$, for any monomial $x \in (\mathcal{P}_d)_{2n+d}$. Note that the map δ is not an $\mathcal{A}_2$ -homomorphism.

The hit problem is reduced to the case of the degree n of the form

$$(1.1) \quad n = r(2^t - 1) + 2^t s,$$

with r, t, s non-negative integers such that $1 \leq r \leq d$ and $\mu(s) < r$ (see Sect.2). Here, $\mu(s)$ is the smallest number u such that $\alpha(n + u) \leq u$, where $\alpha(k)$ denotes the number of 1's in dyadic expansion of a positive integer k . It was solved explicitly for the cases $d \leq 4$ (see [9, 19, 33, 34]).

For $r = d - 1$, the problem was partially studied by Crabb-Hubbuck [7], Nam [17], Repka-Selick [27], and Sum [33, 34] with $s > 0$, by Mothebe [13] and by us [21, 22] with $s = 0$. For $r = d - 2 = 3$, the problem was explicitly solved by the present author [24] with $s = 1$. For $r = d = 5$, it was explicitly determined by Tín [38, 39, 41], Tín-Sum [42] with $s = 1, 2, 3$, by Sum [35, 37] with $s = 5, 10$ and by the present author [25] with $s = 6$. unknown in general.

In the present paper, we study the hit problem of five variables and by using techniques of this problem, we survey Singer's algebraic transfer for the case $d = 5$ and the degree of the form (1.1). More precisely, we explicitly determine a basis of $Q\mathcal{P}_d$ for $d = 5$ and the degree n of the form (1.1) with $r = d = 5$, $s = 8$ and t an arbitrary positive integer. Using this result, we show that the fifth Singer transfer is an isomorphism in bidegree $(5, 13 \cdot 2^t)$ for $t = 1$. This gives an efficient method to study the Singer transfer and it is different from the ones of Boardman [3], Singer [29].

The following is one of our main results.

Theorem 1.1. *Let $n = 13 \cdot 2^t - 5$ with t a positive integer. Then, $\dim(Q\mathcal{P}_5)_n = 840$ for $t = 1$, and $\dim(Q\mathcal{P}_5)_n = 1897$ for $t \geq 2$.*

The proof of the theorem is long and very technical. It is proved by determining the admissible monomials of degree n in $\mathcal{A}_2$ -module $\mathcal{P}_5$ (see Sect.2). The ideas of proofs are from Kameko's squaring operation [9] combining with the results in [21], Mothebe [14, 16], Singer [30], Sum [34], T  n [39, 41], T  n and Sum [42].

Recently, in [37], Sum has proved some properties on the $\mathcal{A}_2$ -generators for $\mathcal{P}_d$. Then, he made a conjecture on the relation between the admissible monomials for the polynomial algebras (see Sect.3). The conjecture helps us to reduce remarkably in computing the hit problem. From the results of Peterson [19], Kameko [9] and Sum [34], this conjecture is true for $d \leq 4$. Sum proved in [37] that the conjecture is true for $d = 5$ and the degree n of the form (1.1) with $(r; s) = (5; 10)$ and $t \geq 0$. Based on the proof of Theorem 1.1 (see Sect.3) and our results in [21]-[24], the conjecture is also true for $d = 5$ and the degrees of the form (1.1) with $(r; s) = (3; 1)$, $(4; 0)$, $(5; 6)$, $(5; 8)$ and t an arbitrary positive integer.

For $t = 1$, we have $n = 13 \cdot 2^t - 5 = 21$. By applying Theorem 1.1, we have the following.

Theorem 1.2. *There exists uniquely a non-zero class in $(Q\mathcal{P}_5)_{21}$ invariant under the action of the group GL_5 .*

Note that the Kameko operation $(\widetilde{Sq}_*^0)_{(5,21)} : (Q\mathcal{P}_5)_{21} \rightarrow (Q\mathcal{P}_5)_8$ is an epimorphism of $\mathbb{Z}/2GL_5$ -modules. So, in order to prove Theorem 1.2, we describe the $\mathbb{Z}/2$ -vector space structure of $(Q\mathcal{P}_5)_8^{GL_5}$. Combining this with a basis of $(QP_5)_{21}$ in Theorem 1.1, we explicitly compute all GL_5 -invariants of $Q\mathcal{P}_5$ in degree 21. Using this result, we obtain the following theorem.

Theorem 1.3. *The fifth transfer homomorphism*

$$Tr_5 : \mathbb{Z}/2 \otimes_{GL_5} P_{\mathcal{A}_2} H_{21}(B(\mathbb{Z}/2)^{\times 5}) \rightarrow \text{Ext}_{\mathcal{A}_2}^{5,26}(\mathbb{Z}/2, \mathbb{Z}/2)$$

is an isomorphism.

This theorem shows that *Singer's conjecture is true in the 5-variable case and the internal degree $13 \cdot 2^t - 5$ for $t = 1$.*

We know that there is squaring operations Sq^0 (see [11]) acting on the cohomology groups of the Steenrod algebra; however it is not the identity. A family $\{u_t : t \geq 0\}$ of elements in $\text{Ext}_{\mathcal{A}_2}^{d,d+*}(\mathbb{Z}/2, \mathbb{Z}/2)$ is called a Sq^0 -family if $u_t = (Sq^0)^t(u_0)$ for every $t \geq 0$. According to Lin [10] and Chen [5], we have

$$\text{Ext}_{\mathcal{A}_2}^{5,13 \cdot 2^t}(\mathbb{Z}/2, \mathbb{Z}/2) = \langle h_t g_t \rangle, \quad h_t g_t = h_{t+1} f_{t-1} \neq 0,$$

where h_t is the Adams element in $\text{Ext}_{\mathcal{A}_2}^{1,2^t}(\mathbb{Z}/2, \mathbb{Z}/2)$, and $g_t = (Sq^0)^{t-1}(g_1)$, $f_{t-1} = (Sq^0)^{t-1}(f_0)$ are non-zero in $\text{Ext}_{\mathcal{A}_2}^{4,12 \cdot 2^t}(\mathbb{Z}/2, \mathbb{Z}/2)$ and $\text{Ext}_{\mathcal{A}_2}^{4,11 \cdot 2^t}(\mathbb{Z}/2, \mathbb{Z}/2)$, respectively, for any $t \geq 1$. On the other hand, Singer showed in [29] that the transfer Tr_1 detects the family $\{h_t | t \geq 1\}$. In [18], Nam proved that Tr_4 detects the family $\{f_{t-1} | t \geq 1\}$. Note that the family $\{g_t | t \geq 1\}$ was not detected by Tr_4 (see Bruner-H  -Hung [4]). Combining these data with the face that $Tr = \bigoplus_{d \geq 0} Tr_d$ is the homomorphism of algebras, we conclude that $h_t g_t$ is in the image of Tr_5 . In Sect.5, we prove this result in another way for the case $t = 1$. More specifically, it is proved by a direct computation using Theorem 1.2 and a representation in the E_1 -page of the Adams spectral sequence of Tr_5 .

We end this introduction with a short discussion on Singer's conjecture for Tr_5 in the bidegree $(5, 13.2^t)$ with $t \geq 2$. Since $\mu(13.2^t - 5) = 5$ for all $t > 2$, the iterated squaring operation $((\widetilde{Sq}_*^0)_{(5, 13.2^t - 5)})^{t-2} : (Q\mathcal{P}_5)_{13.2^t - 5} \rightarrow (Q\mathcal{P}_5)_{47=13.2^2 - 5}$ is an isomorphism of GL_5 -modules for every $t \geq 2$. So, to verify Singer's conjecture for Tr_5 in the above bidegree, we need only to determine all GL_5 -invariants of $(Q\mathcal{P}_5)_{13.2^t - 5}$ for $t = 2$.

Recall that Kameko's operation $(\widetilde{Sq}_*^0)_{(5, 47)} : (Q\mathcal{P}_5)_{47} \rightarrow (Q\mathcal{P}_5)_{21}$ is an epimorphism of GL_5 -modules and the element $h_2g_2 = h_3f_1 \in \text{Im } Tr_5$. So, by Theorem 1.2, we get

$$1 \leq \dim \left(\mathbb{Z}/2 \otimes_{GL_5} P_{\mathcal{A}_2} H_{47}(B(\mathbb{Z}/2)^{\times 5}) \right) \leq \dim(\text{Ker}(\widetilde{Sq}_*^0)_{(5, 47)})^{GL_5} + 1.$$

Furthermore, and all elements of $\mathbb{Z}/2 \otimes_{GL_5} P_{\mathcal{A}_2} H_{47}(B(\mathbb{Z}/2)^{\times 5})$ are of the form $(\gamma[\varphi(u_0)] + [v])^*$, where $\gamma \in \mathbb{Z}/2$; $\varphi : \mathcal{P}_5 \rightarrow \mathcal{P}_5$, which is determined by $\varphi(u) = x_1x_2x_3x_4x_5u^2$ for any $u \in \mathcal{P}_5$; $f \in (\mathcal{P}_5)_{47}$ such that $[f] \in \text{Ker}(\widetilde{Sq}_*^0)_{(5, 47)}$, and $u_0 \in (\mathcal{P}_5)_{21}$. By Theorem 1.2, $[u_0]$ is the only non-zero element in $(Q\mathcal{P}_5)_{21}^{GL_5}$. Direct calculating the elements $(\gamma[\varphi(u_0)] + [v])^*$ is very difficult. By using techniques of the hit problem of five variables, we will describe explicitly all these elements in the near future.

From these events and $h_tg_t \in \text{Im } Tr_5$ for any $t \geq 1$, we see that if $(\text{Ker}(\widetilde{Sq}_*^0)_{(5, 47)})^{GL_5} = 0$, then Singer's conjecture for Tr_5 is also true at the bidegree $(5, 13.2^t)$ for $t \geq 2$. However, it will be much more interesting if $(\text{Ker}(\widetilde{Sq}_*^0)_{(5, 47)})^{GL_5} \neq 0$ and $\dim(Q\mathcal{P}_5)_{47}^{GL_5} \neq 1$.

The paper is organized as follows. In Sect.2, we recall some results related to Kameko's squaring operation, the admissible monomials in $\mathcal{P}_d$, and Singer's criterion on the $\mathcal{A}_2$ -decomposable. In Sect.3, generators of $\mathcal{A}_2$ -module $\mathcal{P}_5$ in degree $13.2^t - 5$ are described explicitly by proving Theorem 1.1. The proof of Theorem 1.2 is presented in Sect.4 by using the results in Sect.3. In Sect.5, we prove Theorem 1.3. Finally, the Appendix describes all the admissible monomials of degrees 8, 21, 22, 47 in $\mathcal{A}_2$ -module $\mathcal{P}_5$.

Acknowledgment. I would like to give my deepest sincere thanks to Professor Nguyen Sum, my thesis advisor, for introducing me to the wonderful subject of the hit problem, for encouraging me to think about the relationship between the hit problem and the algebraic transfer, and for many wonderful discussions.

I would also like to thank Robert R. Bruner, Phan H. Chon, Mbakiso F. Mothebe for helpful conversations related to this work, and Le M. Ha for pointing out an error in the original manuscript.

2. Preliminaries

This section starts with a recollection of the Kameko homomorphism and some results related to the Peterson hit problem.

2.1. Kameko's squaring operation

Denote by $GL_d := GL(d, \mathbb{Z}/2)$ the general linear group of rank d over the field $\mathbb{Z}/2$. This GL_d left acts on $\mathcal{P}_d$ according to the following standard action

$$(w(f))(x_1, x_2, \dots, x_d) = f(w(x_1), w(x_2), \dots, w(x_d)),$$

where $w = (w_{ij}) \in GL_d$ and $w(x_j) = \sum_{i=1}^d w_{ij}x_i$, $1 \leq j \leq d$. Then, $\mathcal{P}_d$ has a left GL_d -module structure. Since the two actions of $\mathcal{A}_2$ and GL_d upon $\mathcal{P}_d$ commute with each other, there is an inherited action of GL_d on $Q\mathcal{P}_d$.

For n a positive integer, let $(Q\mathcal{P}_d)_n^{GL_d}$ be the subspace of $Q\mathcal{P}_d$ generated all GL_d -invariants of degree n . Then, we have the Kameko squaring operation

$$\widetilde{Sq}^0 : \mathbb{Z}/2 \otimes_{GL_d} P_{\mathcal{A}_2} H_n(B(\mathbb{Z}/2)^{\times d}) \rightarrow \mathbb{Z}/2 \otimes_{GL_d} P_{\mathcal{A}_2} H_{d+2n}(B(\mathbb{Z}/2)^{\times d}).$$

It was recognized by Boardman [3] for $d = 3$ and Minami [12] for general d that the operation $\widetilde{Sq}^0$ commutes with the classical squaring operation

$$Sq^0 : \text{Ext}_{\mathcal{A}_2}^{d,d+n}(\mathbb{Z}/2, \mathbb{Z}/2) \rightarrow \text{Ext}_{\mathcal{A}_2}^{d,2d+2n}(\mathbb{Z}/2, \mathbb{Z}/2)$$

on the cohomology of the Steenrod algebra through the Singer transfer Tr_d [29]. This means that the following diagram is commutative:

$$\begin{array}{ccc} \mathbb{Z}/2 \otimes_{GL_d} P_{\mathcal{A}_2} H_n(B(\mathbb{Z}/2)^{\times d}) & \xrightarrow{Tr_d} & \text{Ext}_{\mathcal{A}_2}^{d,d+n}(\mathbb{Z}/2, \mathbb{Z}/2) \\ \downarrow \widetilde{Sq}^0 & & \downarrow Sq^0 \\ \mathbb{Z}/2 \otimes_{GL_d} P_{\mathcal{A}_2} H_{d+2n}(B(\mathbb{Z}/2)^{\times d}) & \xrightarrow{Tr_d} & \text{Ext}_{\mathcal{A}_2}^{d,2d+2n}(\mathbb{Z}/2, \mathbb{Z}/2). \end{array}$$

Note that the homomorphism $\widetilde{Sq}_*^0 : (Q\mathcal{P}_d)_{d+2n}^{GL_d} \rightarrow (Q\mathcal{P}_d)_n^{GL_d}$ is the dual of $\widetilde{Sq}^0$, and it is induced the homomorphism $(\widetilde{Sq}_*^0)_{(d,d+2n)} : (Q\mathcal{P}_d)_{d+2n} \rightarrow (Q\mathcal{P}_d)_n$. The later homomorphism is given by the $\mathbb{Z}/2$ -linear map $\delta : (\mathcal{P}_d)_{d+2n} \rightarrow (\mathcal{P}_d)_n$, determined by

$$\delta(x) = \begin{cases} u & \text{if } x = x_1 x_2 \dots x_d u^2, \\ 0 & \text{otherwise,} \end{cases}$$

for any monomial $x \in (\mathcal{P}_d)_{d+2n}$. The map δ is not a homomorphism of the $\mathcal{A}_2$ -modules. However, $\delta Sq^{2k} = Sq^k \delta$, and $\delta Sq^{2k+1} = 0$ for any non-negative integer k .

For a positive integer n , denote by $\alpha(n)$ the number of 1's in dyadic expansion of n . We set

$$\mu(n) = \min\{u \in \mathbb{N} : \alpha(n+u) \leq u\}.$$

The following lemma can easily be proved by using a result in [35].

Lemma 2.1.1 (see [34]). *$\mu(n) = r \leq d$ if and only if there exist uniquely positive integers t, s such that $1 \leq r - 2 \leq \mu(s) < r$, $\alpha(n + \mu(s)) = \mu(s)$ and $n = r(2^t - 1) + 2^t s$.*

Theorem 2.1.2 (Kameko [9]). *If $\mu(d + 2n) = d$, then the homomorphism*

$$(\widetilde{Sq}_*^0)_{(d,d+2n)} : (Q\mathcal{P}_d)_{d+2n} \rightarrow (Q\mathcal{P}_d)_n$$

is an isomorphism of $\mathbb{Z}/2GL_d$ -modules.

From Lemma 2.1.1, Theorem 2.1.2 and a result in Wood [45], we need to compute the hit problem in degree n of the form (1.1), which is mentioned in Sect.1.

2.2. The weight vector of a monomial and the admissible monomial

For a natural number n , denote by $\alpha_t(n)$ the t -th coefficients in dyadic expansion of n . This means $\alpha(n) = \sum_{t \geq 0} \alpha_t(n)$.

Let $x = x_1^{u_1} x_2^{u_2} \dots x_d^{u_d}$ be a monomial $\mathcal{P}_d$. We define two sequences associated with x by $\omega(x) = (\omega_1(x), \omega_2(x), \dots, \omega_t(x), \dots)$ and $\xi(x) = (u_1, u_2, \dots, u_d)$, where $\omega_t(x) = \sum_{1 \leq j \leq d} \alpha_{t-1}(u_j)$. The sequence $\omega(x)$ is called the *weight vector* of the monomial x and $\xi(x)$ called the *exponent vector* of the monomial x .

Let $\omega = (\omega_1, \omega_2, \dots, \omega_t, \dots)$ be a sequence of non-negative integers. The sequence ω are called the weight vector if $\omega_t = 0$ for $t \gg 0$. The sets of all the weight vectors and the exponent vectors are given the left lexicographical order.

For a weight vector ω , we define $\deg \omega = \sum_{t \geq 1} 2^{t-1} \omega_t$. Denote by $\mathcal{P}_d(\omega)$ the subspace of $\mathcal{P}_d$ spanned by all monomials $x \in \mathcal{P}_d$ such that $\deg x = \deg \omega$, $\omega(x) \leq \omega$, and by $\mathcal{P}_d^-(\omega)$ the subspace of $\mathcal{P}_d$ spanned by all monomials $x \in \mathcal{P}_d$ such that $\omega(x) < \omega$.

Recall that a homogeneous element f of degree n in $\mathcal{P}_d$ is $\mathcal{A}_2$ -decomposable if there is a finite sum $f = \sum_{k > 0} Sq^k(f_k)$, where the homogeneous elements $f_k \in \mathcal{P}_d$ have grading strictly less than n . This means that f belong to $\mathcal{A}_2^+ \mathcal{P}_d$.

Definition 2.2.1. Let ω be a weight vector and f, g two homogeneous polynomials of the same degree in $\mathcal{P}_d$.

- (i) $f \equiv g$ if and only if $f = g$ modulo $(\mathcal{A}_2^+ \mathcal{P}_d)$. If $f \equiv 0$ then f is $\mathcal{A}_2$ -decomposable.
- (ii) $f \equiv_\omega g$ if and only if $f = g$ modulo $(\mathcal{A}_2^+ \mathcal{P}_d + \mathcal{P}_d^-(\omega))$.

Obviously, the relations $\equiv$ and $\equiv_\omega$ are equivalence ones. Denote by $Q\mathcal{P}_d(\omega)$ the quotient of $\mathcal{P}_d(\omega)$ by the equivalence relation $\equiv_\omega$. Then, we have

$$Q\mathcal{P}_d(\omega) = \mathcal{P}_d(\omega) / ((\mathcal{A}_2^+ \mathcal{P}_d \cap \mathcal{P}_d(\omega)) + \mathcal{P}_d^-(\omega)),$$

By Sum [35], $Q\mathcal{P}_d(\omega)$ is also an GL_d -module. Obviously, $(Q\mathcal{P}_d)_n \cong \bigoplus_{\deg \omega = n} Q\mathcal{P}_d(\omega)$.

Definition 2.2.2 (Kameko [9]). Let u, v be monomials of the same degree in $\mathcal{P}_d$. We say that $u < v$ if and only if one of the following holds:

- (i) $\omega(u) < \omega(v)$;
- (ii) $\omega(u) = \omega(v)$ and $\xi(u) < \xi(v)$.

Definition 2.2.3. A monomial u is said to be *inadmissible* if there exist monomials $x_1, x_2, \dots, x_k$ such that $x_j < u$ for $1 \leq j \leq k$ and $u \equiv \sum_{1 \leq i \leq k} x_i$. Then, u is said to be *admissible* if it is not inadmissible.

Definition 2.2.4. A monomial u is said to be *strictly inadmissible* if and only if there exist monomials $x_1, x_2, \dots, x_k$ such that $x_j < u$ for $1 \leq j \leq k$ and $u = \sum_{1 \leq j \leq k} x_j + \sum_{1 \leq t \leq 2^s - 1} Sq^t(y_t)$ with $s = \max\{i \in \mathbb{Z} : \omega_i(u) > 0\}$ and suitable polynomials $y_t \in \mathcal{P}_d$.

Obviously, if X is strictly inadmissible, then it is inadmissible. In general, the opposite is not true. For instance, the monomial $X = x_1 x_2^2 x_3^2 x_4^2 x_5^2 x_6 \in \mathcal{P}_6$ is inadmissible, but it is not strictly inadmissible.

Theorem 2.2.5 (Kameko [9], Sum [33]). Let x, y and u be monomials in $\mathcal{P}_d$ such that $\omega_i(x) = 0$ for $i > r > 0$, $\omega_t(u) \neq 0$ and $\omega_i(u) = 0$ for $i > t > 0$. Then, if u is inadmissible, then xu^{2^r} is also inadmissible. Furthermore, if u is strictly inadmissible, then uy^{2^t} is also strictly inadmissible.

For a natural number d , we now denote

$$\begin{aligned} \mathcal{P}_d^0 &= \langle x_1^{t_1} x_2^{t_2} \dots x_d^{t_d} \in \mathcal{P}_d \mid t_1 t_2 \dots t_d = 0 \rangle, \\ \mathcal{P}_d^+ &= \langle x_1^{t_1} x_2^{t_2} \dots x_d^{t_d} \in \mathcal{P}_d \mid t_1 t_2 \dots t_d > 0 \rangle. \end{aligned}$$

Then $\mathcal{P}_d^0$ and $\mathcal{P}_d^+$ are the $\mathcal{A}_2$ -submodules of $\mathcal{P}_d$. Furthermore, we have a direct summand decomposition of the $\mathbb{Z}/2$ -vector spaces

$$Q\mathcal{P}_d = Q\mathcal{P}_d^0 \oplus Q\mathcal{P}_d^+,$$

where $Q\mathcal{P}_d^0 := \mathbb{Z}/2 \otimes_{\mathcal{A}_2} \mathcal{P}_d^0$ and $Q\mathcal{P}_d^+ := \mathbb{Z}/2 \otimes_{\mathcal{A}_2} \mathcal{P}_d^+$.

We end this section by setting up a formula on the dimension of $Q\mathcal{P}_d^0$ in degree n , which will be used in the next sections. Note that this formula is similar to the one of Mothebe-Kaelo-Ramatebele [16].

Let $\mathcal{I} = (j_1, j_2, \dots, j_r)$, with $1 \leq j_1 < \dots < j_r \leq d$, $1 \leq r < d$, and let $r := \ell(\mathcal{I})$ be the length of $\mathcal{I}$. We denote $\mathcal{P}_{\mathcal{I}} = \langle x_{i_1}^{t_1} x_{i_2}^{t_2} \dots x_{i_r}^{t_r} \mid t_j \in \mathbb{N}, 1 \leq j \leq r \rangle \subset \mathcal{P}_d$. Then, $\mathcal{P}_{\mathcal{I}}$ is a $\mathcal{A}_2$ -submodule of $\mathcal{P}_d$. Further, it is isomorphic to $\mathcal{P}_r$. By a simple computation, one gets

$$Q\mathcal{P}_d^0 = \bigoplus_{1 \leq r \leq d-1} \bigoplus_{\ell(\mathcal{I})=r} Q\mathcal{P}_{\mathcal{I}}^+,$$

where $\mathcal{P}_{\mathcal{I}}^+ = \langle \{x_{i_1}^{t_1} x_{i_2}^{t_2} \dots x_{i_r}^{t_r} \in \mathcal{P}_{\mathcal{I}} \mid t_1 t_2 \dots t_r > 0\} \rangle$ for $1 \leq r \leq d-1$. It can be seen that $\dim(Q\mathcal{P}_{\mathcal{I}}^+)_n = \dim(Q\mathcal{P}_r^+)_n$ for all n , and that $\binom{d}{r}$ is the number of the sequences $\mathcal{I}$ of length r . Therefore, we get

$$\dim(Q\mathcal{P}_d^0)_n = \sum_{1 \leq r \leq d-1} \binom{d}{r} \dim(Q\mathcal{P}_r^+)_n.$$

2.3. Singer's criterion on the $\mathcal{A}_2$ -decomposable

Definition 2.3.1. A monomial $\mathcal{Z} = x_1^{t_1} x_2^{t_2} \dots x_d^{t_d}$ in $\mathcal{P}_d$ is called a *spike* if $t_j = 2^{a_j} - 1$ for a_j a non-negative integer and $1 \leq j \leq d$. If $\mathcal{Z}$ is a spike with $a_1 > a_2 > \dots > a_{r-1} \geq a_r > 0$ and $a_j = 0$ for $j > r$, then it is called a *minimal spike*.

The following result is mentioned in [21].

Lemma 2.3.2. *All the spikes in $\mathcal{P}_d$ are admissible and their weight vectors are weakly decreasing. Furthermore, if a weight vector $\omega = (\omega_1, \omega_2, \dots)$ is weakly decreasing and $\omega_1 \leq d$, then there is a spike $\mathcal{Z}$ in $\mathcal{P}_d$ such that $\omega(\mathcal{Z}) = \omega$.*

For convenience, we prove this lemma in detail.

Proof. Let $\mathcal{Z} = \prod_{1 \leq i \leq d} x_i^{2^{s_i}-1}$ be a spike monomial in $\mathcal{P}_d$. Suppose that $\mathcal{Z}$ is inadmissible. Then, there exist monomials $y_1, y_2, \dots, y_m \in \mathcal{P}_d$ such that $y_t < \mathcal{Z}$ for all t and $\mathcal{Z} = \sum_{1 \leq t \leq m} y_t + \sum_{j>0} Sq^j(w_j)$, with suitable monomial $w_j \in \mathcal{P}_d$. This implies that there is a monomial $u \in \mathcal{P}_d$ with $\deg u < \deg \mathcal{Z}$ such that $Sq^n(u) = \mathcal{Z} + \sum_i X_i$, where $n > 0$ and $X_i \in \mathcal{P}_d$, $\deg X_i = \deg \mathcal{Z}$. Suppose $u = \prod_{1 \leq \ell \leq k} x_{\ell}^{a_{\ell}}$. Since $\deg u < \deg \mathcal{Z}$, there is a number t , $1 \leq t \leq d$ such that $a_t < 2^{s_t} - 1$. According to Cartan's formula, we have

$$\mathcal{Z} + f = Sq^n(u) = \sum_{i \geq 0} Sq^i(x_t^{a_t}) Sq^{n-i}(x_1^{a_1} \dots x_{t-1}^{a_{t-1}} x_{t+1}^{a_{t+1}} \dots x_d^{a_d}).$$

Since $2^{s_t} - 1 - a_t > 0$ and $\mathcal{Z}$ is a term of $\sum_{i \geq 0} Sq^i(x_t^{a_t}) Sq^{n-i}(x_1^{a_1} \dots x_{t-1}^{a_{t-1}} x_{t+1}^{a_{t+1}} \dots x_d^{a_d})$, hence $\mathcal{Z}$ is a term of $Sq^{2^{s_t}-1-a_t}(x_t^{a_t}) Sq^{n-(2^{s_t}-1-a_t)}(x_1^{a_1} \dots x_{t-1}^{a_{t-1}} x_{t+1}^{a_{t+1}} \dots x_d^{a_d})$. On the other hand, $2^{s_t} - 1 - a_t \leq 2^{s_t} - 1$, so there exist a number $\ell \geq 0$ such that $\alpha_{\ell}(2^{s_t} - 1 - a_t) = 1$. This implies $\alpha_{\ell}(a_t) = 0$. Since $\binom{a_t}{2^{s_t}-1-a_t}$ is even, $Sq^{2^{s_t}-1-a_t}(x_t^{a_t}) = 0$. So, one gets $\mathcal{Z} = 0$. This contradicts the fact that $\mathcal{Z} = \prod_{1 \leq i \leq d} x_i^{2^{s_i}-1} \neq 0$. Obviously, if $\mathcal{Z}$ is a spike, then its weight vector is weakly decreasing.

Now, suppose that $\omega = (\omega_1, \omega_2, \dots)$ is weakly decreasing and $\omega_1 \leq d$. For $j \geq d$, we denote by a_j the non-negative integer such that

$$\alpha_i(a_j) = \begin{cases} 1, & \text{if } \omega_{i+1} \geq j, \\ 0, & \text{if } \omega_{i+1} < j. \end{cases}$$

Denote by s_j the greatest integer such that $\omega_{s_j} \geq j$. Since ω is weakly decreasing, $\alpha_i(a_j) = 1$ for $1 \leq i < s_j$ and $\alpha_i(a_j) = 0$ with $i \geq s_j$. So, $a_j = 2^{s_j} - 1$.

On the other hand, we have $\omega_{i+1} \leq \omega_1 \leq d$ for all $i \geq 0$. Hence, if $j > d$, then $a_j = 0$. Therefore, $\mathcal{Z} = x_1^{a_1} x_2^{a_2} \dots x_d^{a_d}$ is a spike in $\mathcal{P}_d$ and $\omega(\mathcal{Z}) = \omega$. The lemma is proved. $\square$

In [30], Singer showed that if $\mu(n) \leq d$, then there exists uniquely a minimal spike of degree n in $\mathcal{P}_d$. Further, we have the following.

Theorem 2.3.3 (Singer [30]). *Suppose that $X \in \mathcal{P}_d$ is a monomial of degree n , where $\mu(X) \leq d$. Let $\mathcal{Z}$ be the minimal spike of degree n in $\mathcal{P}_d$. If $\omega(X) < \omega(\mathcal{Z})$, then X is $\mathcal{A}_2$ -decomposable.*

3. Generators of $\mathcal{A}_2$ -module P_5 in generic degree $13.2^t - 5$

In this section, we prove Theorem 1.1. More precisely, we explicitly compute the admissible monomials of degree $13.2^t - 5$ in $\mathcal{A}_2$ -module $\mathcal{P}_5$. We first describe some homomorphisms in [34, 37].

3.1. Some homomorphisms and Sum's conjecture

For $1 \leq k \leq d$, define the homomorphism $\rho_{(k,d)} : \mathcal{P}_{d-1} \rightarrow \mathcal{P}_d$ of algebras by substituting

$$\rho_{(k,d)}(x_j) = \begin{cases} x_j & \text{if } 1 \leq j < k, \\ x_{j+1} & \text{if } k \leq j < d. \end{cases}$$

We now denote

$$\mathcal{N}_d := \{(k; \mathcal{K}) \mid \mathcal{K} = (k_1, k_2, \dots, k_r), 1 \leq k < k_1 < k_2 < \dots < k_r \leq d, 0 \leq r < d\}.$$

Here, by convention, $\mathcal{K} = \emptyset$, if $r = 0$. Denote by $r = \ell(\mathcal{K})$ the length of $\mathcal{K}$.

For any $(k; \mathcal{K}) \in \mathcal{N}_d$, we define the homomorphism $\pi_{(k; \mathcal{K})} : \mathcal{P}_d \rightarrow \mathcal{P}_{d-1}$ of algebras by substituting

$$\pi_{(k; \mathcal{K})}(x_j) = \begin{cases} x_j & \text{if } 1 \leq j < k, \\ \sum_{p \in \mathcal{K}} x_{p-1} & \text{if } j = k, \\ x_{j-1} & \text{if } k < j \leq d. \end{cases}$$

Note that $\rho_{(k,d)}$ and $\pi_{(k; \mathcal{K})}$ are also the homomorphisms of the $\mathcal{A}_2$ -modules. In particular, we have $\pi_{(k; \emptyset)}(x_k) = 0$ for $1 \leq k \leq d$ and $\pi_{(k; \mathcal{K})}(\rho_{(k,d)}(u)) = u$ for any $u \in \mathcal{P}_{d-1}$.

Proposition 3.1.1 (see [21]). *If x is a monomial in $\mathcal{P}_d$, then $\pi_{(k; \mathcal{K})}(x) \in \mathcal{P}_{d-1}(\omega(x))$.*

This result implies that if ω is a weight vector and $x \in \mathcal{P}_d(\omega(x))$, then $\pi_{(k; \mathcal{K})}(x) \in \mathcal{P}_{d-1}(\omega)$. Furthermore, $\pi_{(k; \mathcal{K})}$ passes to a homomorphism from $Q\mathcal{P}_d(\omega)$ to $Q\mathcal{P}_{d-1}(\omega)$.

Definition 3.1.2. Let $(k; \mathcal{K}) \in \mathcal{N}_d$, $r = \ell(\mathcal{K})$, and let u be an integer with $1 \leq u \leq r$. A monomial $x_1^{t_1} x_2^{t_2} \dots x_{d-1}^{t_{d-1}} \in \mathcal{P}_{d-1}$ is said to be u -compatible with $(k; \mathcal{K})$ if all of the following hold:

- (i) $t_{k_1-1} = t_{k_2-1} = \dots = t_{k_{u-1}-1} = 2^r - 1$,
- (ii) $t_{k_u-1} > 2^r - 1$,
- (iii) $\alpha_{r-m}(t_{k_u-1}) = 1$, $\forall m$, $1 \leq m \leq u$,
- (iv) $\alpha_{r-m}(t_{k_m-1}) = 1$, $\forall m$, $u < m \leq r$.

Clearly, a monomial x can be u -compatible with a given $(k; \mathcal{K}) \in \mathcal{N}_d$ for at most one value of u . By convention, x is 1-compatible with $(k; \emptyset)$.

Definition 3.1.3. Let $(k; \mathcal{K}) \in \mathcal{N}_d, 0 < r < d$. Denote

$$x_{(\mathcal{K}, u)} = x_{k_u}^{2^{r-1}+2^{r-2}+\dots+2^{r-u}} \prod_{u < m \leq r} x_{k_m}^{2^{r-m}}$$

for $1 \leq u \leq r$, $x_{(\emptyset, 1)} = 1$. For a monomial $x \in \mathcal{P}_{d-1}$, we define the monomial $\phi_{(k; \mathcal{K})}(x)$ in $\mathcal{P}_d$ by setting

$$\phi_{(k; \mathcal{K})}(x) = \begin{cases} \frac{x_k^{2^{r-1}} \rho_{(k, d)}(x)}{x_{(\mathcal{K}, u)}} & \text{if there exist } u \text{ such that } x \text{ is } u\text{-compatible with } (k; \mathcal{K}), \\ 0 & \text{otherwise.} \end{cases}$$

Then we have an $\mathbb{Z}/2$ -linear map $\phi_{(k; \mathcal{K})} : \mathcal{P}_{d-1} \rightarrow \mathcal{P}_d$. In particular, $\phi_{(k; \emptyset)} = \rho_k$. Note that $\phi_{(k; \mathcal{K})}$ is not an $\mathcal{A}_2$ -homomorphism. Moreover, we see that if $\phi_{(k; \mathcal{K})}(x) \neq 0$, then $\omega(\phi_{(k; \mathcal{K})}(x)) = \omega(x)$.

For a subset $\mathcal{U} \subset \mathcal{P}_{d-1}$, denote

$$\begin{aligned} \overline{\Phi}^0(\mathcal{U}) &= \bigcup_{1 \leq k \leq d} \phi_{(k; \emptyset)}(\mathcal{U}) = \bigcup_{1 \leq k \leq d} \rho_{(k, d)}(\mathcal{U}), \\ \overline{\Phi}^+(\mathcal{U}) &= \bigcup_{(k; \mathcal{K}) \in \mathcal{N}_d, 0 < \ell(\mathcal{K}) < d} \phi_{(k; \mathcal{K})}(\mathcal{U}) \setminus \mathcal{P}_d^0, \\ \overline{\Phi}(\mathcal{U}) &= \overline{\Phi}^0(\mathcal{U}) \cup \overline{\Phi}^+(\mathcal{U}). \end{aligned}$$

Since $\rho_{(k, d)}$ is a homomorphism of the $\mathcal{A}_2$ -modules, if $\mathcal{U}$ is a minimal set of generators for $\mathcal{A}_2$ -module $\mathcal{P}_{d-1}$ in degree n , then $\overline{\Phi}^0(\mathcal{U})$ is also a minimal set of generators for $\mathcal{A}_2$ -module $\mathcal{P}_d^0$ in degree n .

Now, for a polynomial $f \in \mathcal{P}_d$, we denote by $[f]$ the classes in $Q\mathcal{P}_d$ represented by f . If ω is a weight vector and $f \in \mathcal{P}_d(\omega)$, then denote by $[f]_\omega$ the classes in $Q\mathcal{P}_d(\omega)$ represented by f .

Denote by $\mathcal{B}_d(n)$ the set of all admissible monomials of degree n in $\mathcal{P}_d$. Set

$$\mathcal{B}_d^0(n) := \mathcal{B}_d(n) \cap (\mathcal{P}_d^0)_n, \quad \mathcal{B}_d^+(n) := \mathcal{B}_d(n) \cap (\mathcal{P}_d^+)_n.$$

If ω is a weight vector of degree n , we set

$$\mathcal{B}_d(\omega) := \mathcal{B}_d(n) \cap \mathcal{P}_d(\omega), \quad \mathcal{B}_d^0(\omega) := \mathcal{B}_d(\omega) \cap (\mathcal{P}_d^0)_n, \quad \mathcal{B}_d^+(\omega) := \mathcal{B}_d(\omega) \cap (\mathcal{P}_d^+)_n.$$

Then, $[\mathcal{B}_d(\omega)]_\omega$, $[\mathcal{B}_d^0(\omega)]_\omega$ and $[\mathcal{B}_d^+(\omega)]_\omega$ are respectively the bases of the $\mathbb{Z}/2$ -vector spaces

$$Q\mathcal{P}_d(\omega), \quad Q\mathcal{P}_d^0(\omega) := Q\mathcal{P}_d(\omega) \cap (Q\mathcal{P}_d^0)_n$$

and $Q\mathcal{P}_d^+(\omega) := Q\mathcal{P}_d(\omega) \cap (Q\mathcal{P}_d^+)_n$.

From now on, we use the following notations: For a natural number d , we set

$$\begin{aligned} \Gamma_d &= \{1, 2, \dots, d\}, \\ X_{(S, d)} &= X_{(\{s_1, s_2, \dots, s_r\}, d)} = \prod_{s \in \Gamma_d \setminus S} x_s, \quad S = \{s_1, s_2, \dots, s_r\} \subseteq \Gamma_d. \end{aligned}$$

In particular,

$$\begin{aligned} X_{\Gamma_d} &= 1, \quad X_{(\emptyset, d)} = x_1 x_2 \dots x_d, \\ X_{(\{s\}, d)} &= x_1 \dots \hat{x}_s \dots x_d, \quad 1 \leq s \leq d. \end{aligned}$$

Let $X = x_1^{t_1} x_2^{t_2} \dots x_d^{t_d} \in \mathcal{P}_d$. Set $S_j(X) = \{s \in \Gamma_d : \alpha_j(t_s) = 0\}$, for $j \geq 0$. Then we have

$$X = \prod_{j \geq 0} X_{(S_j(X), d)}^{2^j}.$$

For a subset $\mathcal{B} \subset \mathcal{P}_d$, we denote $[\mathcal{B}] = \{[f] : f \in \mathcal{B}\}$. If $\mathcal{B} \subset \mathcal{P}_d(\omega)$, then we set $[\mathcal{B}]_\omega = \{[f]_\omega : f \in \mathcal{B}\}$. Throughout this paper, we use Theorem 2.3.3, Proposition 3.1.1 and a result of Sum [34] to prove a certain subset of $Q\mathcal{P}_d$ is linearly independent.

More precisely, let $\mathcal{B}$ be a finite subset of $\mathcal{P}_d$ consisting of some monomials of degree n . Denote by $|\mathcal{B}|$ the cardinal of $\mathcal{B}$. To prove the set $[\mathcal{B}]$ is linearly independent in $(Q\mathcal{P}_d)_n$, we denote the elements of $\mathcal{B}$ by $\mathcal{Y}_{n,i}$, $1 \leq i \leq m = |\mathcal{B}|$ and assume that there is a linear relation

$$\mathcal{S} = \sum_{1 \leq i \leq m} \gamma_i \mathcal{Y}_{n,i} = 0 \text{ modulo } (\mathcal{A}_2^+ \mathcal{P}_d + \mathcal{P}_d^-(\omega)),$$

with $\gamma_i \in \mathbb{Z}/2$ for all i , $1 \leq i \leq m$. For $(k; \mathcal{K}) \in \mathcal{N}_d$, we explicitly compute $\pi_{(k; \mathcal{K})}(\mathcal{S})$ in terms of the admissible monomials in $\mathcal{P}_{d-1}$ (modulo $(\mathcal{A}_2^+ \mathcal{P}_{d-1} + \mathcal{P}_{d-1}^-(\omega))$). Computing from some relations $\pi_{(k; \mathcal{K})}(\mathcal{S}) = 0$ modulo $(\mathcal{A}_2^+ \mathcal{P}_{d-1} + \mathcal{P}_{d-1}^-(\omega))$ with $(k; \mathcal{K}) \in \mathcal{N}_d$, we obtain $\gamma_i = 0$ for all i .

We end this section by recalling the Sum conjecture [37] on the relation between the admissible monomials for the polynomial algebra. In [37], Sum made the following conjecture.

Conjecture 3.1.4 (Sum [37]). *If ω is a weight vector, then $\overline{\Phi}(\mathcal{B}_{d-1}(\omega)) \subseteq \mathcal{B}_d(\omega)$.*

We see that if this conjecture is true, then $\overline{\Phi}(\mathcal{B}_{d-1}(n)) \subseteq \mathcal{B}_d(n)$ for any positive integer n . In other words, if $x \in \mathcal{B}_{d-1}(n)$, then $\phi_{(k; \mathcal{K})}(x) \in \mathcal{B}_d(n)$. Based on the results of Peterson [19], Kameko [9] and Sum [34], the conjecture is true for $d \leq 4$. In particular, we have the following remark.

Remark 3.1.5. Consider the spike monomial $X = X_{\{d\}, d} = x_1 x_2 \dots x_{d-1} \in \mathcal{P}_{d-1}$. Let m be a positive integer such that $m > r = \ell(\mathcal{K})$. Then, we have

$$\phi_{(k; \mathcal{K})}(X^{2^m-1}) = x_k^{2^r-1} \prod_{1 \leq t \leq r} x_{i_t}^{2^m-2^{r-t}-1} X_{(\{k, k_1, \dots, k_r\}, d)}^{2^m-1}.$$

Indeed, since $m > r$, for each $(k; \mathcal{K}) \in \mathcal{N}_d$, $\mathcal{K} = (k_1; k_2; \dots; k_r)$, we have X^{2^m-1} is 1-compatible with $(k; \mathcal{K})$. Then, we get

$$\begin{aligned} \phi_{(k; \mathcal{K})}(X^{2^m-1}) &= \frac{x_k^{2^r-1} x_2^{2^m-1} \dots x_{k_1}^{2^m-1} \dots x_{k_r}^{2^m-1} \dots x_d^{2^m-1}}{x_{k_1}^{2^r-1} x_{k_2}^{2^r-2} \dots x_{k_r}^{2^r-r}} \\ &= x_k^{2^r-1} x_{k_1}^{2^m-2^{r-1}-1} \dots x_{k_2}^{2^m-2^{r-2}-1} \dots x_{k_r}^{2^m-2^{r-r}-1} X_{(\{k, k_1, \dots, k_r\}, d)}^{2^m-1} \\ &= x_k^{2^r-1} \prod_{1 \leq t \leq r} x_{k_t}^{2^m-2^{r-t}-1} X_{(\{k, k_1, \dots, k_r\}, d)}^{2^m-1}. \end{aligned}$$

Now, suppose $1 \leq r = q-1 < d$. Then $2 \leq q \leq d$ and $m \geq q$. Consider $(1; \mathcal{K})$, with $\mathcal{K} = (k_1; k_2; \dots; k_r) = (2; 3; \dots; q)$. Then, we have

$$\begin{aligned} \phi_{(1; \mathcal{K})}(X^{2^m-1}) &= x_1^{2^{q-1}-1} \prod_{1 \leq k \leq q-1} x_k^{2^m-2^{q-1-k}-1} X_{(\{1, 2, \dots, q\}, d)}^{2^m-1} \\ &= x_1^{2^{q-1}-1} \dots x_k^{2^m-2^{q-k}-1} \dots x_q^{2^m-2} x_{q+1}^{2^m-1} \dots x_d^{2^m-1}. \end{aligned}$$

Since X^{2^m-1} is a spike monomial, by Proposition 3.1.1, X^{2^m-1} is admissible. On the other hand, using a result in Mothebe [14], we conclude that $\phi_{(1; \mathcal{K})}(X^{2^m-1})$ is admissible. Thus, Sum's conjecture is also true in this case, for any d .

In [37], Sum showed that Conjecture 3.1.4 is true for $d = 5$ and any weight vector ω of the degree $15 \cdot 2^t - 5$ for t an arbitrary non-negative integer. In the next section, we will show that this conjecture is also true for $d = 5$ and in degree $13 \cdot 2^t - 5$ for any positive integer t .

3.2. Proof of Theorem 1.1

First, we see that $n = 13 \cdot 2^t - 5 = 2^{t+3} + 2^{t+2} + 2^t + 2^{t-1} + 2^{t-1} - 5$. This implies $\mu(n) = 5$ for any $t > 2$. So, by Theorem 2.1.2, the iterated Kameko's squaring operation

$$(\widetilde{Sq}_*^0)_{(5, n)}^{t-2} : (Q\mathcal{P}_5)_n \rightarrow (Q\mathcal{P}_5)_{13 \cdot 2^t - 5}$$

is an isomorphism of $\mathbb{Z}/2GL_5$ -modules for every $t \geq 2$. Therefore, we need only to compute $(Q\mathcal{P}_5)_n$ for $t = 1, 2$.

3.2.1. The case $t = 1$

Now, for $t = 1$, we have $n = 13 \cdot 2^t - 5 = 21$. Consider Kameko's squaring operation

$$(\widetilde{Sq_*^0})_{(5,21)} : (Q\mathcal{P}_5)_{21} \longrightarrow (Q\mathcal{P}_5)_8.$$

Since $(\widetilde{Sq_*^0})_{(5,21)}$ is an epimorphism of $\mathbb{Z}/2$ -vector spaces, one gets

$$(Q\mathcal{P}_5)_{21} \cong \text{Ker}(\widetilde{Sq_*^0})_{(5,21)} \bigoplus (Q\mathcal{P}_5)_8 \cong (Q\mathcal{P}_5^0)_{21} \bigoplus (\text{Ker}(\widetilde{Sq_*^0})_{(5,21)} \cap (Q\mathcal{P}_5^+)_{21}) \bigoplus (Q\mathcal{P}_5)_8.$$

By T  n [39], $\dim(Q\mathcal{P}_5)_8 = 174$. According to the results in [19], [9] and [34], we have

$$|\mathcal{B}_1^+(21)| = |\mathcal{B}_2^+(21)| = 0, \quad |\mathcal{B}_3^+(21)| = 7, \quad |\mathcal{B}_4^+(21)| = 66.$$

Note that $(Q\mathcal{P}_3)_{21} \cong (Q\mathcal{P}_3)_3$ and $\mathcal{B}_3(21) = \mathcal{B}_3^+(21) = \tilde{\varphi}^2(\mathcal{B}_3(3))$ with the $\mathbb{Z}/2$ -linear map $\tilde{\varphi} : \mathcal{P}_5 \rightarrow \mathcal{P}_5$, given by $\tilde{\varphi}(u) = X_{(\emptyset,5)}u^2$, $\forall u \in \mathcal{P}_5$. Since $(Q\mathcal{P}_5^0)_{21} = \bigoplus_{1 \leq r \leq 4} \bigoplus_{1 \leq j \leq \binom{5}{r}} (Q\mathcal{P}_r^+)_{21}$, one gets $\dim(Q\mathcal{P}_5^0)_{21} = \binom{5}{3} \cdot 7 + \binom{5}{4} \cdot 66 = 400$, and $\mathcal{B}_5^0(21) = \overline{\Phi}^0(\mathcal{B}_4(21)) = \{\mathcal{Y}_{21,i} : 1 \leq i \leq 400\}$, where the monomials $\mathcal{Y}_{21,i}$, $1 \leq i \leq 400$, are listed in Sect.6.2.

Proposition 3.2.1. *The set $\{[\mathcal{Y}_{21,i}] : 401 \leq i \leq 666\}$ is the basis of the $\mathbb{Z}/2$ -vector space $\text{Ker}(\widetilde{Sq_*^0})_{(5,21)} \cap (Q\mathcal{P}_5^+)_{21}$. Here the monomials $\mathcal{Y}_i := \mathcal{Y}_{21,i}$, $401 \leq i \leq 666$, which are determined in Sect.6.3.*

Combining Proposition 3.2.1 and the above arguments, we see that $\dim(Q\mathcal{P}_5)_{21} = 840$. We prepare some lemmas for the proof of the proposition.

Lemma 3.2.2. *If u is an admissible monomial of degree 21 in $\mathcal{P}_5$ and $[u]$ belong to $\text{Ker}(\widetilde{Sq_*^0})_{(5,21)}$, then we have either $\omega(u) = (3, 3, 1, 1)$ or $\omega(u) = (3, 3, 3)$.*

Proof. Observe $\mathcal{Z} = x_1^{15}x_2^3x_3^3$ is the minimal spike of degree 21 in $\mathcal{P}_5$ and $\omega(\mathcal{Z}) = (3, 3, 1, 1)$. Since $[u] \neq [0]$, by Theorem 2.3.3, we get either $\omega_1(u) = 3$ or $\omega_1(u) = 5$. If $\omega_1(u) = 5$ then $u = X_{(\emptyset,5)}y^2$ with y a monomial of degree 8 in $\mathcal{P}_5$. Since u is admissible, by Theorem 2.2.5, one gets $y \in \mathcal{B}_5(8)$. So $(\widetilde{Sq_*^0})_{(5,21)}([u]) = [y] \neq [0]$. This contradicts the fact that $[u] \in \text{Ker}(\widetilde{Sq_*^0})_{(5,21)}$. Hence, $\omega_1(u) = 3$. Then, we have $u = X_{\{i,j\},5}y_1^2$ with $1 \leq i < j \leq 5$ and $y_1 \in \mathcal{B}_5(9)$. Since y_1 is admissible, according to a result in [41], we have either $\omega(y_1) = (3, 1, 1)$ or $\omega(y_1) = (3, 3)$. The lemma is proved. $\square$

Based on Lemma 3.2.2, we conclude

$$\text{Ker}((\widetilde{Sq_*^0})_{(5,21)}) \cap (Q\mathcal{P}_5^+)_{21} = Q\mathcal{P}_5^+(3, 3, 1, 1) \bigoplus Q\mathcal{P}_5^+(3, 3, 3).$$

The following lemma is an immediate corollary from a result in [34].

Lemma 3.2.3. *The following monomials are strictly inadmissible:*

- (i) $x_i^2x_jx_k^3x_\ell^7$, $x_i^6x_jx_k^3x_\ell^3$, $x_i^2x_j^5x_k^3x_\ell^3$, $x_i^2x_jx_k^3x_\ell^3$, $1 \leq i < j \leq 5$, $1 \leq k, \ell \leq 5$, $k \neq \ell$, $k, \ell \neq i, j$;
- (ii) $x_i^3x_j^4x_k^3x_\ell^3$, $1 \leq i < j < k < \ell \leq 5$;
- (iii) $\rho_{(k,5)}(X)$, $1 \leq k \leq 5$, where X is one of the following monomials:

$$\begin{array}{cccc} x_1^3x_2^4x_3^7x_4^7, & x_1^3x_2^7x_3^4x_4^7, & x_1^3x_2^7x_3^7x_4^4, & x_1^7x_2^3x_3^4x_4^7, \\ x_1^7x_2^3x_3^7x_4^4, & x_1^7x_2^7x_3^3x_4^4, & x_1^7x_2^8x_3^3x_4^3. \end{array}$$

Lemma 3.2.4. *The following monomials are strictly inadmissible:*

- (i) $x_i^2x_j^2x_kx_\ell x_m^7$, $x_i^2x_j^6x_kx_\ell x_m^3$, $x_i^6x_j^2x_kx_\ell x_m^3$, $x_i^2x_j^2x_k^5x_\ell x_m^3$, $x_i^2x_j^4x_kx_\ell^3x_m^3$, $x_i^4x_j^2x_kx_\ell^3x_m^3$, $x_i^2x_jx_k^4x_\ell^3x_m^3$, where (i, j, k, ℓ, m) is a permutation of $(1, 2, 3, 4, 5)$;
- (ii) $x_1^3x_2^2x_3x_4x_5^6$, $x_1^3x_2^6x_3x_4x_5^2$, $x_1^3x_2^2x_3^5x_4x_5^2$, $x_1^3x_2^2x_3x_4^5x_5^2$,
 $x_1^3x_2^5x_3^2x_4x_5^2$, $x_1^3x_2^5x_3^2x_4^2x_5^2$, $x_1^3x_2^2x_3^2x_4x_5^2$, $x_1^3x_2^2x_3x_4^2x_5^2$.

Proof. Consider the monomials $X = x_i^2 x_j^2 x_k x_\ell x_m^7$ and $Y = x_1^3 x_2^2 x_3 x_4 x_5^6$. We prove that these monomials are strictly inadmissible. The others can be proved by a similar computation. Obviously, $\omega(X) = \omega(Y) = (3, 3, 1)$. By a direct computation using the Cartan formula, we obtain

$$\begin{aligned} X &= Sq^2(x_i x_j x_k x_\ell x_m^7) + Sq^4(x_i x_j x_k x_\ell x_m^5) + x_i x_j x_k^2 x_\ell^2 x_m^7 \\ &\quad + x_i x_j^2 x_k x_\ell^2 x_m^7 + x_i x_j^2 x_k^2 x_\ell x_m^7 \text{ modulo } (\mathcal{P}_5^-(3, 3, 1)); \\ Y &= Sq^1(x_1^3 x_2 x_3 x_4 x_5^6) + x_1^3 x_2 x_3^2 x_4 x_5^6 + x_1^3 x_2 x_3 x_4^2 x_5^6 \text{ modulo } (\mathcal{P}_5^-(3, 3, 1)). \end{aligned}$$

These equalities show that X and Y are strictly inadmissible. The lemma follows. $\square$

Lemma 3.2.5. *The following monomials are strictly inadmissible:*

$$\begin{aligned} X_1 &= x_1 x_2^6 x_3^8 x_4^3 x_5^3, & X_2 &= x_1 x_2^3 x_3^6 x_4^6 x_5^5, & X_3 &= x_1^3 x_2^5 x_3^2 x_4^6 x_5^5, & X_4 &= x_1 x_2^6 x_3^7 x_4^6 x_5^5, \\ X_5 &= x_1 x_2^7 x_3^6 x_4^6 x_5^5, & X_6 &= x_1^7 x_2 x_3^6 x_4^6 x_5^5, & X_7 &= x_1 x_2^2 x_3^6 x_4^5 x_5^7, & X_8 &= x_1 x_2^2 x_3^6 x_4^7 x_5^5, \\ X_9 &= x_1 x_2^6 x_3^5 x_4^7 x_5^5, & X_{10} &= x_1 x_2^6 x_3^7 x_4^5 x_5^5, & X_{11} &= x_1 x_2^2 x_3^7 x_4^6 x_5^5, & X_{12} &= x_1 x_2^6 x_3^7 x_4^2 x_5^5, \\ X_{13} &= x_1 x_2^7 x_3^2 x_4^6 x_5^5, & X_{14} &= x_1 x_2^7 x_3^6 x_4^2 x_5^5, & X_{15} &= x_1^7 x_2 x_3^2 x_4^6 x_5^5, & X_{16} &= x_1^7 x_2 x_3^6 x_4^2 x_5^5, \\ X_{17} &= x_1 x_2^6 x_3^3 x_4^7 x_5^5, & X_{18} &= x_1 x_2^6 x_3^3 x_4^7 x_5^5, & X_{19} &= x_1 x_2^6 x_3^7 x_4^3 x_5^5, & X_{20} &= x_1 x_2^7 x_3^6 x_4^3 x_5^5, \\ X_{21} &= x_1^3 x_2^4 x_3^6 x_4^7 x_5^5, & X_{22} &= x_1^3 x_2^4 x_3^7 x_4^6 x_5^5, & X_{23} &= x_1^3 x_2^4 x_3^7 x_4^6 x_5^5, & X_{24} &= x_1^3 x_2^7 x_4^3 x_4^6 x_5^5, \\ X_{25} &= x_1^7 x_2 x_3^6 x_4^3 x_5^5, & X_{26} &= x_1^7 x_2^3 x_4^3 x_4^6 x_5^5, & X_{27} &= x_1 x_2^6 x_3^6 x_4^3 x_5^5, & X_{28} &= x_1 x_2^6 x_3^3 x_4^6 x_5^5, \\ X_{29} &= x_1 x_2^6 x_3^6 x_4^7 x_5^5, & X_{30} &= x_1^3 x_2 x_3^6 x_4^6 x_5^5, & X_{31} &= x_1^3 x_2^5 x_3^6 x_4^6 x_5^5, & X_{32} &= x_1^3 x_2^4 x_3^5 x_4^2 x_5^5, \\ X_{33} &= x_1^3 x_2^4 x_3^5 x_4^7 x_5^5, & X_{34} &= x_1^3 x_2^4 x_3^7 x_4^5 x_5^5, & X_{35} &= x_1^3 x_2^5 x_3^4 x_4^2 x_5^5, & X_{36} &= x_1^3 x_2^5 x_3^4 x_4^7 x_5^5, \\ X_{37} &= x_1^3 x_2^7 x_3^4 x_4^5 x_5^5, & X_{38} &= x_1^3 x_2^5 x_3^7 x_4^4 x_5^5, & X_{39} &= x_1^3 x_2^5 x_3^4 x_4^5 x_5^5, & X_{40} &= x_1^7 x_2^3 x_3^4 x_4^5 x_5^5, \\ X_{41} &= x_1^3 x_2^5 x_3^4 x_4^5 x_5^5, & X_{42} &= x_1 x_2^6 x_3^6 x_4^7 x_5^5, & X_{43} &= x_1^3 x_2^5 x_3^6 x_4^2 x_5^5, & X_{44} &= x_1^3 x_2^5 x_3^6 x_4^2 x_5^5, \\ X_{45} &= x_1^3 x_2^5 x_3^2 x_4^6 x_5^5, & X_{46} &= x_1^3 x_2^5 x_3^5 x_4^6 x_5^5, & X_{47} &= x_1^3 x_2^5 x_3^4 x_4^3 x_5^5, & X_{48} &= x_1^3 x_2^4 x_3^4 x_4^7 x_5^5, \\ X_{49} &= x_1^3 x_2^4 x_3^3 x_4^4 x_5^5, & X_{50} &= x_1^3 x_2^4 x_3^7 x_4^4 x_5^5, & X_{51} &= x_1^3 x_2^4 x_3^3 x_4^7 x_5^5, & X_{52} &= x_1^3 x_2^4 x_3^7 x_4^3 x_5^5, \\ X_{53} &= x_1^3 x_2^7 x_3^4 x_4^3 x_5^5, & X_{54} &= x_1^3 x_2^7 x_3^4 x_3^3 x_5^5, & X_{55} &= x_1^7 x_2^3 x_3^4 x_4^3 x_5^5, & X_{56} &= x_1^7 x_2^3 x_3^4 x_3^3 x_5^5, \\ X_{57} &= x_1^3 x_2^4 x_3^5 x_4^6 x_5^5, & X_{58} &= x_1^3 x_2^4 x_3^5 x_4^3 x_5^5, & X_{59} &= x_1^3 x_2^5 x_3^4 x_4^6 x_5^5, & X_{60} &= x_1^3 x_2^5 x_3^6 x_4^3 x_5^5, \\ X_{61} &= x_1^3 x_2^5 x_3^4 x_3^3 x_5^5. \end{aligned}$$

Proof. It is easy to see that $\omega(X_1) = (3, 3, 1, 1)$ and $\omega(X_j) = (3, 3, 3)$ for $j = 2, 3, \dots, 61$. We prove the lemma for the monomials $X_1 = x_1 x_2^6 x_3^8 x_4^3 x_5^3$, $X_2 = x_1^3 x_2^5 x_3^2 x_4^6 x_5^5$ and $X_3 = x_1 x_2^6 x_3^7 x_4^6 x_5^5$. The others can be proven by a similar computation. By a direct computation, we have

$$\begin{aligned} X_1 &= x_1 x_2^3 x_3^3 x_4^6 x_5^8 + x_1 x_2^3 x_3^3 x_4^8 x_5^6 + x_1 x_2^3 x_3^6 x_4^3 x_5^8 + x_1 x_2^3 x_3^6 x_4^8 x_5^3 + x_1 x_2^3 x_3^8 x_4^3 x_5^6 \\ &\quad + x_1 x_2^3 x_3^8 x_4^6 x_5^3 + x_1 x_2^4 x_3^3 x_4^3 x_5^{10} + x_1 x_2^4 x_3^3 x_4^{10} x_5^3 + x_1 x_2^4 x_3^3 x_4^3 x_5^3 \\ &\quad + x_1 x_2^6 x_3^3 x_4^8 x_5^3 + x_1 x_2^6 x_3^3 x_4^8 x_5^3 + Sq^1(A_1) + Sq^2(A_2) + Sq^4(A_4) \text{ modulo } (\mathcal{P}_5^-(3, 3, 1, 1)), \end{aligned}$$

where

$$\begin{aligned} A_1 &= x_1^2 x_2^3 x_3^5 x_4^5 x_5^5 + x_1^2 x_2^5 x_3^3 x_4^5 x_5^5 + x_1^2 x_2^5 x_3^5 x_4^3 x_5^5 + x_1^2 x_2^5 x_3^5 x_4^5 x_5^3, \\ A_2 &= x_1 x_2^3 x_3^3 x_4^6 x_5^5 + x_1 x_2^3 x_3^5 x_4^5 x_5^5 + x_1 x_2^3 x_3^6 x_4^3 x_5^5 + x_1 x_2^3 x_3^6 x_4^5 x_5^3 + x_1 x_2^5 x_3^3 x_4^5 x_5^5 \\ &\quad + x_1 x_2^5 x_3^3 x_4^5 x_5^3 + x_1 x_2^5 x_3^5 x_4^5 x_5^3 + x_1 x_2^6 x_3^3 x_4^3 x_5^6 + x_1 x_2^6 x_3^3 x_4^6 x_5^3 + x_1 x_2^6 x_3^3 x_4^3 x_5^3, \\ A_4 &= x_1 x_2^4 x_3^3 x_4^3 x_5^6 + x_1 x_2^4 x_3^3 x_4^6 x_5^3 + x_1 x_2^4 x_3^6 x_4^3 x_5^3. \end{aligned}$$

This relation implies that X_1 is strictly inadmissible. By a similar technique, we obtain

$$X_2 = Sq^1(B_1) + Sq^2(B_2) + Sq^8(B_8) + x_1 x_2^3 x_3^5 x_4^6 x_5^6 + x_1 x_2^3 x_3^6 x_4^5 x_5^6 \text{ modulo } (\mathcal{P}_5^-(3, 3, 3)),$$

where

$$\begin{aligned} B_1 &= x_1^2 x_2^5 x_3^3 x_4^5 x_5^5 + x_1^2 x_2^5 x_3^5 x_4^3 x_5^5 + x_1^2 x_2^5 x_3^5 x_4^5 x_5^3, \\ B_2 &= x_1 x_2^3 x_3^3 x_4^6 x_5^6 + x_1 x_2^3 x_3^5 x_4^5 x_5^5 + x_1 x_2^3 x_3^6 x_4^3 x_5^5 + x_1 x_2^6 x_3^3 x_4^3 x_5^6 + x_1 x_2^6 x_3^3 x_4^6 x_5^3 + x_1 x_2^6 x_3^6 x_4^3 x_5^3, \\ B_8 &= x_1 x_2^3 x_3^3 x_4^3 x_5^3, \end{aligned}$$

$$X_3 = Sq^1(C_1) + Sq^2(C_2) + Sq^8(C_8) + x_1^3 x_2^5 x_3^6 x_4^6 x_5^6 + x_1^3 x_2^5 x_3^2 x_4^5 x_5^6 \text{ modulo } (\mathcal{P}_5^-(3, 3, 3)),$$

where

$$\begin{aligned} C_1 &= x_1^3 x_2^5 x_3 x_4^5 x_5^6 + x_1^3 x_2^5 x_3 x_4^6 x_5^5 + x_1^5 x_2^3 x_3^2 x_4^5 x_5^5 + x_1^5 x_2^5 x_3^2 x_4^3 x_5^5 + x_1^5 x_2^5 x_3^2 x_4^5 x_5^3, \\ C_2 &= x_1^3 x_2^3 x_3 x_4^6 x_5^6 + x_1^3 x_2^5 x_3 x_4^5 x_5^5 + x_1^3 x_2^6 x_3 x_4^3 x_5^6 + x_1^3 x_2^6 x_3 x_4^6 x_5^3 + x_1^6 x_2^3 x_3 x_4^6 x_5^3, \\ C_8 &= x_1^3 x_2^3 x_3 x_4^3 x_5^3. \end{aligned}$$

The lemma is proved. $\square$

Now we denote by $\mathcal{C}$ the set of the following monomials:

$$\begin{array}{cccc} x_1^3 x_2^{12} x_3 x_4^2 x_5^3 & x_1^3 x_2^{12} x_3 x_4^3 x_5^2 & x_1^3 x_2^{12} x_3^2 x_4 x_5^2 & x_1^3 x_2^4 x_3 x_4^2 x_5^{11}, \\ x_1^3 x_2^4 x_3 x_4^{11} x_5^2 & x_1^3 x_2^4 x_3^{11} x_4 x_5^2 & x_1^7 x_2^8 x_3 x_4^2 x_5^3 & x_1^7 x_2^8 x_3 x_4^3 x_5^2, \\ x_1^7 x_2^8 x_3^2 x_4 x_5^2 & x_1^3 x_2^4 x_3 x_4^3 x_5^{10} & x_1^3 x_2^4 x_3 x_4^{10} x_5^3 & x_1^3 x_2^4 x_3^2 x_4 x_5^{10}, \\ x_1^3 x_2^4 x_3^3 x_4 x_5^2 & x_1^3 x_2^4 x_3^9 x_4 x_5^2 & x_1^3 x_2^4 x_3^9 x_4^2 x_5^2 & \end{array}$$

A direct computation shows that $\overline{\Phi}^+(\mathcal{B}_4(3, 3, 1, 1)) \cup \mathcal{C}$ is the set of 196 monomials: $\mathcal{Y}_j := \mathcal{Y}_{21, j}$, $401 \leq j \leq 596$ (see Sect.6.3.)

Proposition 3.2.6. *Under the above notations, the $\mathbb{Z}/2$ -vector space $Q\mathcal{P}_5^+(3, 3, 1, 1)$ is spanned by the set*

$$[\overline{\Phi}^+(\mathcal{B}_4(3, 3, 1, 1)) \cup \mathcal{C}].$$

Proof. Let X be an admissible monomial in $\mathcal{P}_5$ such that $\omega(X) = (3, 3, 1, 1)$. Then $X = X_{(\{k, \ell\}, 5)} Y^2$ with $1 \leq k < \ell \leq 5$ and Y a monomial of degree 9 in $\mathcal{P}_5$. Since X is admissible, according to Theorem 2.2.5, $Y \in \mathcal{B}_5(3, 1, 1)$.

A direct computation shows that if $z \in \mathcal{B}_5^+(3, 1, 1)$, $1 \leq k < \ell \leq 5$, and $X_{(\{k, \ell\}, 5)} z^2 \neq \mathcal{Y}_j$, $\forall j$, $401 \leq j \leq 596$ then there exists a monomial w which is given in one of Lemmas 3.2.3-3.2.5 such that $X_{(\{k, \ell\}, 5)} z^2 = w z_1^{2^a}$ with a monomial $z_1 \in \mathcal{P}_5$ and $a = \max\{m \in \mathbb{Z} : \omega_m(w) > 0\}$. By Theorem 2.2.5, $X_{(\{k, \ell\}, 5)} z^2$ is inadmissible. Since $X = X_{(\{k, \ell\}, 5)} Y^2$ with $Y \in \mathcal{B}_5(3, 1, 1)$ and X is admissible, one can see that $X = \mathcal{Y}_j$, $401 \leq j \leq 596$. The lemma follows. $\square$

By a direct computation, we see that

$$\overline{\Phi}^+(\mathcal{B}_4(3, 3, 3)) \cup \{x_1^3 x_2^4 x_3 x_4^5 x_5^6, x_1^3 x_2^5 x_3 x_4^6 x_5^4, x_1^3 x_2^5 x_3^6 x_4^3 x_5^4\}$$

is the set consisting of 70 monomials: $\mathcal{Y}_t := \mathcal{Y}_{21, t}$, $597 \leq t \leq 666$ (see Sect.6.3.)

Proposition 3.2.7. *The $\mathbb{Z}/2$ -vector space $Q\mathcal{P}_5^+(3, 3, 3)$ is spanned by the set $\{[\mathcal{Y}_t] : 597 \leq t \leq 666\}$.*

Proof. Let u be an admissible monomial in $\mathcal{P}_5$ such that $\omega(u) = (3, 3, 3)$. Then $u = x_i x_j x_\ell y^2$ with $1 \leq i < j < \ell \leq 5$ and $y \in \mathcal{B}_5(3, 3)$.

By a direct computation, we can verify that for any $X \in \mathcal{B}_5(3, 3)$, $1 \leq i < j < \ell \leq 5$, such that $x_i x_j x_\ell X^2 \neq \mathcal{Y}_t$, $\forall t$, $597 \leq t \leq 666$, there is a monomial z which is given in one of Lemmas 3.2.3-3.2.5 such that $x_i x_j x_\ell X^2 = z w^{2^b}$ with suitable monomial $w \in \mathcal{P}_5$ and $b = \max\{r \in \mathbb{Z} : \omega_r(z) > 0\}$. Then, according to Theorem 2.2.5, $x_i x_j x_\ell X^2$ is inadmissible. Since $u = x_i x_j x_\ell y^2$ is admissible and $y \in \mathcal{B}_5(3, 3)$, one gets $u = \mathcal{Y}_t$, for some t . This proves the proposition. $\square$

Proof of Proposition 3.2.1. From Propositions 3.2.6 and 3.2.7, the space $\text{Ker}((\widetilde{Sq}_*^0)_{(5, 21)}) \cap (Q\mathcal{P}_5^+)_{21}$ is spanned by the set $\{[\mathcal{Y}_i] := \mathcal{Y}_{21, i} : 401 \leq i \leq 666\}$. Furthermore, this set is linearly independent in $(Q\mathcal{P}_5)_{21}$. Indeed, suppose there is a linear relation

$$\mathcal{S} = \sum_{401 \leq i \leq 666} \gamma_i \mathcal{Y}_i = 0 \text{ modulo } (\mathcal{A}_2^+ \mathcal{P}_5),$$

where $\gamma_i \in \mathbb{Z}/2$. Let $\mathcal{H}$ be a sequence of non-negative integers and $\gamma_h \in \mathbb{Z}/2$ for $h \in \mathcal{H}$. Denote $\gamma_{\mathcal{H}} = \sum_{h \in \mathcal{H}} \gamma_h \in \mathbb{Z}/2$. Based on Theorem 2.3.3 and Proposition 3.1.1, for $(k; \mathcal{H}) \in \mathcal{N}_5$, we explicitly compute $\pi_{(k; \mathcal{H})}(\mathcal{S})$ in terms of a given minimal set of $\mathcal{A}_2$ -generators in $\mathcal{P}_4$ (modulo $(\mathcal{A}_2^+ \mathcal{P}_4)$).

Computing directly from the relations $\pi_{(k;p)}(\mathcal{S}) \equiv 0$, $1 \leq k < p \leq 5$, one gets $\gamma_h = 0$, $h \in \mathbb{H}$, where the set $\mathbb{H} = \{401, 402, \dots, 425, 434, 435, 439, 441, 442, 443, 446, 451, 453, 454, 457, 458, 460, 461, 462, 463, 465, 467, 470, 471, \dots, 477, 479, 483, 487, 489, 497, 498, 500, 501, 502, 504, 505, 506, 507, 510, 513, 515, 517, 519, 520, 523, 527, 532, 533, 540, 542, 543, 560, 561, 562, 565, 566, 567, 568, 570, 573, 575, 577, 578, 579, 580, 581, 585, 597, 598, 599, 600, 601, 602, 604, 606, 607, 609, 610, 613, 615, 616, 617, 620, 622, 629, 630, 632, 633, \dots, 647, 650, 651, 652, 653, 655, 656, 657, 658, 659, 660, 662, 663\}$, and

$$\left\{ \begin{array}{l} \gamma_{430} = \gamma_{593}, \\ \gamma_{431} = \gamma_{594}, \gamma_{536} = \gamma_{541}, \\ \gamma_{603} = \gamma_{664}, \gamma_h = \gamma_{426}, h \in \mathbb{I}_1, \\ \gamma_{608} = \gamma_{627}, \gamma_h = \gamma_{427}, h \in \mathbb{I}_2, \\ \gamma_{619} = \gamma_{626}, \gamma_h = \gamma_{432}, h = 468, 491, 518, 595, \\ \gamma_{621} = \gamma_{665}, \gamma_h = \gamma_{433}, h = 596, 623, 666, \\ \gamma_h = \gamma_{436}, h = 437, 438, 469, 492, 495, 496, 499, 534, 539, 545, 614, 654, \\ \gamma_{\{450, 595, 661\}} = \gamma_{\{522, 648, 661\}} = \gamma_{\{531, 648, 661\}} = \gamma_{\{549, 654, 661\}} = 0, \\ \gamma_{\{555, 595, 654\}} = \gamma_{\{618, 626, 628\}} = 0, \\ \gamma_{\{582, 654, 661\}} = \gamma_{\{583, 654, 661\}} = \gamma_{\{584, 654, 661\}} = 0, \\ \gamma_{\{429, 648, 595, 661\}} = \gamma_{\{455, 595, 654, 661\}} = \gamma_{\{456, 654, 661, 666\}} = \gamma_{\{486, 648, 594, 654\}} = 0, \\ \gamma_{\{493, 664, 665, 594\}} = \gamma_{\{503, 654, 552, 666\}} = \gamma_{\{512, 648, 595, 661\}} = \gamma_{\{529, 664, 665, 594\}} = 0, \\ \gamma_{\{544, 654, 661, 595\}} = \gamma_{\{547, 654, 661, 595\}} = \gamma_{\{548, 654, 661, 666\}} = \gamma_{\{550, 654, 661, 595\}} = 0, \\ \gamma_{\{551, 648, 594, 654\}} = \gamma_{\{557, 654, 661, 666\}} = \gamma_{\{558, 654, 661, 648\}} = \gamma_{\{592, 648, 595, 661\}} = 0, \\ \gamma_{\{481, 648, 664, 665, 594\}} = \gamma_{\{526, 648, 664, 665, 594\}} = \gamma_{\{554, 654, 628, 595, 626\}} = 0, \\ \gamma_{\{624, 626, 628, 627, 664\}} = \gamma_{\{459, 628, 648, 595, 626, 666\}} = 0, \\ \gamma_{\{488, 628, 595, 626, 648, 594\}} = \gamma_{\{535, 627, 654, 595, 661, 626\}} = 0, \\ \gamma_{\{546, 661, 666, 648, 594, 552\}} = \gamma_{\{612, 628, 627, 665, 666, 664\}} = 0, \\ \gamma_{\{485, 627, 654, 648, 595, 661, 626\}} = \gamma_{\{490, 626, 628, 627, 665, 666, 664\}} = 0, \\ \gamma_{\{509, 666, 552, 654, 628, 595, 626\}} = \gamma_{\{524, 666, 552, 654, 628, 595, 626\}} = 0, \\ \gamma_{\{538, 627, 654, 648, 595, 661, 626\}} = \gamma_{\{556, 654, 628, 648, 595, 661, 626\}} = 0, \\ \gamma_{\{559, 648, 626, 628, 627, 666, 661\}} = \gamma_{\{625, 626, 628, 627, 665, 666, 664\}} = 0, \\ \gamma_{\{521, 654, 666, 648, 594, 552, 628, 664, 665\}} = \gamma_{\{553, 654, 648, 594, 626, 628, 627, 666, 664\}} = 0. \end{array} \right.$$

Here $\mathbb{I}_1 = \{440, 447, 448, 464, 466, 478, 484, 508, 514, 516, 586, 605, , 648\}$,

$\mathbb{I}_2 = \{428, 444, 445, 449, 452, 480, 482, 494, 511, 525, 528, 530, , 537, 563,$

$564, 569, 571, 572, 574, 576, 587, 588, 589, 590, 591, 611, 631, 649, 661\}$,

Combining the above computations and the relations $\pi_{(1,(2;j))}(\mathcal{S}) \equiv 0$, $j = 3, 4, 5$, $\pi_{(1,(3;4))}(\mathcal{S}) \equiv 0$, we obtain $\gamma_i = 0$, $\forall i$, $401 \leq i \leq 666$. This finishes the proof. $\square$

3.2.2. The case $t = 2$

For $t = 2$, we have $13 \cdot 2^t - 5 = 47$ and $\mu(47) = 3 < 5$. Since Kameko's operation

$$(\widetilde{Sq}_*^0)_{(5,47)} : (Q\mathcal{P}_5)_{47} \rightarrow (Q\mathcal{P}_5)_{21}$$

is an epimorphism of the $\mathbb{Z}/2GL_5$ -modules, hence $(Q\mathcal{P}_5)_{47} = \text{Ker}((\widetilde{Sq}_*^0)_{(5,47)}) \oplus (Q\mathcal{P}_5)_{21}$. Hence, we need to compute $\text{Ker}((\widetilde{Sq}_*^0)_{(5,47)})$.

Remark 3.2.8. If $Y \in \mathcal{B}_5(47)$ and $[Y] \in \text{Ker}((\widetilde{Sq}_*^0)_{(5,47)})$, then $\omega_1(Y) = 3$.

Indeed, we see that $\mathcal{Z} = x_1^{31}x_2^{15}x_3$ is the minimal spike of degree 47 in $\mathcal{P}_5$. By Lemma 2.3.2, $\mathcal{Z} \in \mathcal{B}_5(47)$. Since $[Y] \neq 0$, by Theorem 2.3.3, either $\omega_1(Y) = 3$ or $\omega_1(Y) = 5$. If $\omega_1(Y) = 5$, then $Y = X_{(\emptyset, 5)}Z^2$ with Z a monomial of degree 21 in $\mathcal{P}_5$. Since Y is admissible, by Theorem 2.2.5, $Z \in \mathcal{B}_5(21)$. So, we have $(\widetilde{Sq}_*^0)_{(5, 47)}([Y]) = [Z] \neq [0]$. This contradicts the fact that $[Y] \in \text{Ker}((\widetilde{Sq}_*^0)_{(5, 47)})$; hence we get $\omega_1(Y) = 3$.

From Remark 3.2.8, we have $Y = x_k x_\ell x_m g^2$ with $1 \leq k < \ell < m \leq 5$ and $g \in \mathcal{B}_5(22)$. Thus, to determine $\text{Ker}((\widetilde{Sq}_*^0)_{(5, 47)})$, we need to compute all the admissible monomials of degree 22 in $\mathcal{A}_2$ -module $\mathcal{P}_5$.

Computation of $(Q\mathcal{P}_5)_{22}$

We consider the weight vectors

$$\omega_{(1)} = (2, 2, 2, 1), \omega_{(2)} = (2, 4, 1, 1), \omega_{(3)} = (2, 4, 3), \omega_{(4)} = (4, 3, 1, 1), \omega_{(5)} = (4, 3, 3).$$

We have $\deg \omega_{(i)} = 22$, $1 \leq i \leq 5$. It is easy to see that $\mathcal{Z} = x_1^{15}x_2^7$ is the minimal spike of degree 22 in $\mathcal{P}_5$ and $\omega(z) = \omega_{(1)}$. Let u be an admissible monomial of degree 22 in $\mathcal{P}_5$. Then $[u] \neq [0]$ and by Theorem 2.3.3, either $\omega_1(u) = 2$ or $\omega_1(u) = 4$. Since $u \in \mathcal{B}_5(22)$, by Theorem 2.2.5, if $\omega_1(u) = 2$, then $u = X_{(\{i, j, k\}, 5)}y^2$ with $y \in \mathcal{B}_5(10)$ and $1 \leq i < j < k \leq 5$. According to T  n [40], $\omega(y)$ is one of the following sequences:

$$(2, 2, 1), (4, 1, 1), (4, 3).$$

If $\omega_1(u) = 4$, then $u = X_{(\{i\}, 5)}y_1^2$ with y_1 a monomial of degree 9 in $\mathcal{P}_5$ and $1 \leq i \leq 5$. By T  n [40], either $\omega(y_1) = (3, 1, 1)$ or $\omega(y_1) = (3, 3)$. Hence, one gets the following.

Remark 3.2.9. If $u \in \mathcal{B}_5(22)$, then $\omega(u)$ is one of the sequences $\omega_{(t)}$, $1 \leq t \leq 5$.

As it is known,

$$(Q\mathcal{P}_5)_{22} = (Q\mathcal{P}_5^0)_{22} \bigoplus (Q\mathcal{P}_5^+)_{22}.$$

By Sum [34], we have $|\mathcal{B}_4^+(22)| = 72$. Then, combining the results in [19] and [9] with fact that $(Q\mathcal{P}_5^0)_{22} = \bigoplus_{1 \leq s \leq 4} \bigoplus_{1 \leq u \leq \binom{5}{u}} (Q\mathcal{P}_s^+)_{22}$, we get

$$\dim(Q\mathcal{P}_5^0)_{22} = \binom{5}{2} \cdot 2 + \binom{5}{3} \cdot 8 + \binom{5}{4} \cdot 72 = 460,$$

and $\mathcal{B}_5^0(22) = \overline{\Phi}^0(\mathcal{B}_4(22)) = \{\mathcal{Y}_{22, t} : 1 \leq t \leq 460\}$, where the monomials $\mathcal{Y}_{22, t}$, $1 \leq t \leq 460$, are determined in Sect.6.4.

Next, we determine $(Q\mathcal{P}_5^+)_{22}$. From now on, we denote

$$\mathcal{U}(5, n) := \{r \in \mathbb{N} : \alpha(n - (2^r - 1) + 4) \leq 4\}$$

with n a natural number. For $r \in \mathcal{U}(5, n)$, let $X = x_1^{a_1} \dots x_4^{a_4}$ be an admissible monomial of degree $m = n - (2^r - 1)$ in $\mathcal{A}_2$ -module $\mathcal{P}_4$. We set

$$\mathcal{B}(5, n) := \bigcup_{r \in \mathcal{U}(5, 22)} \left\langle \{x_1^{a_1} \dots x_{j-1}^{a_{j-1}} x_j^{2^r-1} x_{j+1}^{a_{j+1}} \dots x_5^{a_4} \in (\mathcal{P}_5)_{n=m+2^r-1} : 1 \leq j \leq 5\} \right\rangle.$$

Then, according to Mothebe [14], $\mathcal{B}(5, n) \subseteq \mathcal{B}_5(n)$. If ω is a weight vector of degree n , then we write

$$\mathcal{B}(5, \omega) := \mathcal{B}(5, n) \cap \mathcal{P}_5(\omega), \quad \mathcal{B}^+(5, \omega) := \mathcal{B}(5, \omega) \cap (\mathcal{P}_5^+)_n.$$

By a simple computation, we see that $\overline{\Phi}^+(\mathcal{B}_4(\omega_{(1)})) \cup \mathcal{B}^+(5, \omega_{(1)})$ is the set of 31 admissible monomials: $\mathcal{Y}_{22, i}$, $461 \leq i \leq 491$ (see Sect.6.5.)

Denote by $\mathcal{D}$ is the set of the following monomials:

$$\begin{aligned} \mathcal{Y}_{22,492} &= x_1 x_2 x_3^6 x_4^6 x_5^8, & \mathcal{Y}_{22,493} &= x_1 x_2 x_3^6 x_4^{10} x_5^4, & \mathcal{Y}_{22,494} &= x_1 x_2^2 x_3^3 x_4^4 x_5^{12}, \\ \mathcal{Y}_{22,495} &= x_1 x_2^2 x_3^3 x_4^{12} x_5^4, & \mathcal{Y}_{22,496} &= x_1 x_2^2 x_3^4 x_4^9 x_5^6, & \mathcal{Y}_{22,497} &= x_1 x_2^2 x_3^5 x_4^8 x_5^6, \\ \mathcal{Y}_{22,498} &= x_1 x_2^3 x_3^2 x_4^4 x_5^{12}, & \mathcal{Y}_{22,499} &= x_1 x_2^3 x_3^2 x_4^{12} x_5^4, & \mathcal{Y}_{22,500} &= x_1 x_2^3 x_3^4 x_4^8 x_5^6, \\ \mathcal{Y}_{22,501} &= x_1 x_2^3 x_3^6 x_4^4 x_5^8, & \mathcal{Y}_{22,502} &= x_1 x_2^3 x_3^6 x_4^8 x_5^4, & \mathcal{Y}_{22,503} &= x_1^3 x_2 x_3^2 x_4^4 x_5^{12}, \\ \mathcal{Y}_{22,504} &= x_1^3 x_2 x_3^2 x_4^{12} x_5^4, & \mathcal{Y}_{22,505} &= x_1^3 x_2 x_3^4 x_4^8 x_5^6, & \mathcal{Y}_{22,506} &= x_1^3 x_2 x_3^6 x_4^4 x_5^8, \\ \mathcal{Y}_{22,507} &= x_1^3 x_2 x_3^6 x_4^8 x_5^4, & \mathcal{Y}_{22,508} &= x_1^3 x_2^5 x_3^2 x_4^4 x_5^8, & \mathcal{Y}_{22,509} &= x_1^3 x_2^5 x_3^2 x_4^8 x_5^4, \\ \mathcal{Y}_{22,510} &= x_1^3 x_2^8 x_3^2 x_4^4 x_5^4. \end{aligned}$$

Proposition 3.2.10. $\mathcal{B}_5^+(\omega_{(1)}) = \overline{\Phi}^+(\mathcal{B}_4(\omega_{(1)})) \cup \mathcal{B}^+(5, \omega_{(1)}) \cup \mathcal{D}$.

In order to prove the proposition, we need some lemmas.

Lemma 3.2.11. *The following monomials are strictly inadmissible:*

(i) $x_1^2 x_j x_k^2 x_l^3 x_m^6$, $x_1^6 x_j x_k x_l^2 x_m^4$, $l < m$, $x_1^2 x_j x_k x_l^4 x_m^6$.

Here (j, k, l, m) is a permutation of $(2, 3, 4, 5)$;

(ii) $x_1 x_2^2 x_3^4 x_4^6 x_5$, $x_1 x_2^2 x_3^6 x_4^4 x_5^4$, $x_1 x_2^2 x_3^6 x_4^4 x_5$, $x_1 x_2^6 x_3^2 x_4^4 x_5^4$,
 $x_1 x_2^6 x_3^2 x_4^4 x_5$, $x_1^3 x_2^3 x_3^4 x_4^2 x_5^2$, $x_1^3 x_2^4 x_3^2 x_4^2 x_5^3$, $x_1^3 x_2^4 x_3^2 x_4^3 x_5^2$,
 $x_1^3 x_2^4 x_3^2 x_4^4 x_5$, $x_1^3 x_2^2 x_3^2 x_4^6 x_5^2$, $x_1^3 x_2^2 x_3^2 x_4^6 x_5$, $x_1^3 x_2^2 x_3^2 x_4^6 x_5^2$,
 $x_1^3 x_2^2 x_3^2 x_4^6 x_5$, $x_1^3 x_2^2 x_3^6 x_4^2 x_5^2$, $x_1^3 x_2^2 x_3^6 x_4^2 x_5$, $x_1^3 x_2^2 x_3^6 x_4^2 x_5^2$,
 $x_1^3 x_2^6 x_3^2 x_4^2 x_5^2$, $x_1^3 x_2^6 x_3^2 x_4^2 x_5$, $x_1^2 x_2^3 x_3^3 x_4^3 x_5^3$, $x_1^2 x_2^3 x_3^3 x_4^3 x_5^2$,
 $x_1^3 x_2^3 x_3^3 x_4^3 x_5^3$, $x_1^3 x_2^3 x_3^3 x_4^3 x_5^2$.

Proof. We prove the lemma for the monomials $u = x_1^2 x_j x_k^2 x_l^3 x_m^6$, and $v = x_1 x_2^2 x_3^4 x_4^6 x_5$. The others can be proved by a similar computation. We have $\omega(u) = (2, 4, 1)$ and $\omega(v) = (2, 2, 2)$. By a simple computation, one gets

$$\begin{aligned} u &= x_1 x_j^2 x_k^2 x_l^3 x_m^6 + Sq^1(x_1 x_j x_k^2 x_l^3 x_m^6) \text{ modulo } (\mathcal{P}_5^-(2, 4, 1)), \\ v &= x_1 x_2 x_3^4 x_4^6 x_5^2 + x_1 x_2^2 x_3^4 x_4^5 x_5^2 + Sq^1(f_1) + Sq^2(f_2) \text{ modulo } (\mathcal{P}_5^-(2, 2, 2)), \end{aligned}$$

where $f_1 = x_1^2 x_2 x_3^4 x_4^5 x_5$ and $f_2 = x_1 x_2 x_3^4 x_4^5 x_5$. Hence, u and v are strictly inadmissible. The lemma follows. $\square$

The following lemma can easily be proved by a direct computation.

Lemma 3.2.12. *If (i, j, k, l, m) is a permutation of $(1, 2, 3, 4, 5)$, then the following monomials are strictly inadmissible:*

(i) $x_i^6 x_j x_k^7$, $x_i^2 x_j^5 x_k^7$, $x_i^3 x_j^4 x_k^7$, $x_i^2 x_j x_k^2 x_l^2 x_m^7$,
 $x_i^2 x_j^5 x_k^2 x_l^3 x_m^3$, $x_i^2 x_j^4 x_k^2 x_l^3 x_m^3$, $x_i^2 x_j x_k^2 x_l^2 x_m^3$, $i < j$;
(ii) $x_i x_j^6 x_k^3 x_l^2 x_m^2$, $j < k$, $x_i^3 x_j^6 x_k^5$, $x_i^6 x_j^3 x_k^5$, $x_i^2 x_j x_k^3 x_l^3 x_m^3$, $x_i^2 x_j x_k x_l^2$, $i < j < k$;
(iii) $x_i^2 x_j x_k^4 x_l^3 x_m^4$, $x_i^4 x_j x_k x_l^3 x_m^4$, $i < j < k$, $l < m$.

Lemma 3.2.13. *The following monomials are strictly inadmissible:*

$$\begin{aligned} x_1 x_2^2 x_3^2 x_4^7 x_5^{10}, & x_1 x_2^2 x_3^4 x_4^3 x_5^{12}, & x_1 x_2^2 x_3^4 x_4^{11} x_5^4, & x_1 x_2^2 x_3^7 x_4^2 x_5^{10}, \\ x_1 x_2^2 x_3^7 x_4^8 x_5^4, & x_1 x_2^2 x_3^7 x_4^{10} x_5^2, & x_1 x_2^2 x_3^{12} x_4^3 x_5^4, & x_1 x_2^6 x_3^3 x_4^6 x_5^8, \\ x_1 x_2^6 x_3^3 x_4^{10} x_5^4, & x_1 x_2^6 x_3^6 x_4^6 x_5^6, & x_1 x_2^6 x_3^6 x_4^3 x_5^6, & x_1 x_2^6 x_3^6 x_4^6 x_5^3, \\ x_1 x_2^6 x_3^9 x_4^2 x_5^4, & x_1 x_2^7 x_3^2 x_4^2 x_5^{10}, & x_1 x_2^7 x_3^2 x_4^8 x_5^4, & x_1 x_2^7 x_3^2 x_4^{10} x_5^2, \\ x_1 x_2^7 x_3^8 x_4^2 x_5^4, & x_1 x_2^7 x_3^{10} x_4^2 x_5^2, & x_1^3 x_2^3 x_3^4 x_4^4 x_5^8, & x_1^3 x_2^3 x_3^4 x_4^8 x_5^4, \\ x_1^3 x_2^4 x_3^3 x_4^2 x_5^{12}, & x_1^3 x_2^4 x_3^3 x_4^{10} x_5^4, & x_1^3 x_2^4 x_3^4 x_4^4 x_5^7, & x_1^3 x_2^4 x_3^4 x_4^5 x_5^6, \\ x_1^3 x_2^4 x_3^4 x_4^4 x_5^4, & x_1^3 x_2^4 x_3^5 x_4^4 x_5^6, & x_1^3 x_2^4 x_3^5 x_4^6 x_5^4, & x_1^3 x_2^4 x_3^7 x_4^4 x_5^4, \\ x_1^3 x_2^4 x_3^9 x_4^2 x_5^4, & x_1^3 x_2^5 x_3^4 x_4^4 x_5^6, & x_1^3 x_2^5 x_3^4 x_4^6 x_5^4, & x_1^3 x_2^5 x_3^6 x_4^4 x_5^4, \\ x_1^3 x_2^7 x_3^4 x_4^4 x_5^4, & x_1^3 x_2^{12} x_3 x_4^2 x_5^4, & x_1^7 x_2 x_3^2 x_4^2 x_5^{10}, & x_1^7 x_2 x_3^2 x_4^8 x_5^4, \\ x_1^7 x_2 x_3^2 x_4^{10} x_5^2, & x_1^7 x_2 x_3^8 x_4^2 x_5^4, & x_1^7 x_2 x_3^{10} x_4^2 x_5^2, & x_1^7 x_2 x_3^4 x_4^4 x_5^4, \\ x_1^7 x_2 x_3^8 x_4^2 x_5^4, & x_1^7 x_2 x_3^2 x_4^2 x_5^2. & & \end{aligned}$$

Proof. We prove the lemma for the monomials $x = x_1x_2^2x_3^4x_4^3x_5^{12}$, and $y = x_1^3x_2^3x_3^4x_4^5x_5^8$. The others can be proved by a similar computation. By a direct computation using the Cartan formula, we have

$$x = x_1x_2^2x_3^4x_4^3x_5^{12} + x_1x_2^2x_3^4x_4^2x_5^{13} + x_1x_2^2x_3^2x_4^4x_5^{13} + x_1x_2x_3^6x_4^6x_5^8 + x_1x_2x_3^6x_4^4x_5^{10} \\ + x_1x_2x_3^4x_4^6x_5^{10} + Sq^1(f_1) + Sq^2(f_2) + Sq^4(f_4) \text{ modulo } (\mathcal{P}_5^-(\omega_{(1)})),$$

where

$$f_1 = x_1x_2^4x_3^3x_4^4x_5^9 + x_1x_2^4x_3^3x_4^5x_5^8 + x_1x_2^4x_3^4x_4^3x_5^9 + x_1x_2^4x_3^4x_4^5x_5^7 \\ + x_1x_2^4x_3^5x_4^3x_5^8 + x_1x_2^4x_3^5x_4^4x_5^7 + x_1^2x_2x_3^4x_4^5x_5^9 + x_1^2x_2x_3^5x_4^4x_5^8 \\ + x_1^2x_2x_3^5x_4^5x_5^7 + x_1^4x_2^2x_3^3x_4^3x_5^7, \\ f_2 = x_1x_2x_3^4x_4^5x_5^9 + x_1x_2x_3^5x_4^4x_5^9 + x_1x_2x_3^5x_4^5x_5^8 + x_1x_2^2x_3^3x_4^4x_5^{10} \\ + x_1x_2^2x_3^3x_4^6x_5^8 + x_1x_2^2x_3^3x_4^4x_5^{10} + x_1x_2^2x_3^4x_4^6x_5^7 + x_1x_2^2x_3^6x_4^3x_5^8 \\ + x_1x_2^2x_3^6x_4^4x_5^7 + x_1x_2^4x_3^2x_4^2x_5^{11} + x_1^2x_2^4x_3^3x_4^3x_5^8 + x_1^2x_2^4x_3^3x_4^4x_5^7 \\ + x_1^2x_2^4x_3^4x_4^3x_5^7, \\ f_4 = x_1x_2^2x_3^2x_4^2x_5^{11} + x_1x_2^2x_3^4x_4^4x_5^7.$$

The above equalities show that x is strictly inadmissible. By a similar computation, we obtain

$$y = x_1^2x_2x_3^2x_4^{13}x_5^4 + x_1^2x_2x_3^3x_4^{12}x_5^4 + x_1^2x_2x_3^4x_4^7x_5^8 + x_1^2x_2x_3^4x_4^{13}x_5^2 + x_1^2x_2x_3^5x_4^6x_5^8 \\ + x_1^2x_2x_3^6x_4^9x_5^4 + x_1^2x_2x_3^8x_4^7x_5^4 + x_1^2x_2x_3^{12}x_4^5x_5^2 + x_1^2x_2^3x_3^5x_4^8x_5^4 + x_1^2x_2^3x_3^8x_5^4x_5^4 \\ + x_1^2x_2^5x_3^2x_4^5x_5^8 + x_1^2x_2^5x_3^2x_4^9x_5^4 + x_1^2x_2^5x_3^3x_4^8x_5^4 + x_1^2x_2^5x_4^3x_3^8x_5^8 + x_1^2x_2^5x_4^4x_3^9x_5^2 \\ + x_1^2x_2^5x_8^3x_5^4x_5^2 + x_1^3x_2x_3^2x_4^{12}x_5^4 + x_1^3x_2x_3^4x_4^6x_5^8 + x_1^3x_2x_3^4x_4^{10}x_5^4 + x_1^3x_2x_3^4x_4^{12}x_5^2 \\ + x_1^3x_2x_3^4x_4^{12}x_5^2 + x_1^3x_2^2x_3^4x_4^9x_5^4 + x_1^3x_2^2x_3^8x_5^4x_5^4 \\ + Sq^1(g_1) + Sq^2(g_2) + Sq^4(g_4) + Sq^8(x_1^3x_2^3x_3^2x_4^4x_5^2) \text{ modulo } (\mathcal{P}_5^-(\omega_{(1)})), \text{ where}$$

$$g_1 = x_1^3x_2^3x_3^2x_4^9x_5^4 + x_1^3x_2^3x_3^4x_4^9x_5^2 + x_1^3x_2^3x_3^2x_4^5x_5^8 + x_1^3x_2^3x_3^8x_4^5x_5^2 + x_1^5x_2x_3^2x_4^9x_5^4 \\ + x_1^5x_2x_3^4x_4^9x_5^2 + x_1^5x_2x_3^4x_4^9x_5^2 + x_1^5x_2x_3^8x_4^5x_5^2 + x_1^5x_2x_3^8x_4^5x_5^2 + x_1^5x_2x_3^6x_4^5x_5^4 \\ + x_1^5x_2^4x_3^5x_4^5x_5^4 + x_1^5x_2x_3^3x_4^6x_5^4 + x_1^5x_2x_3^3x_4^8x_5^4 + x_1^5x_2x_3^3x_4^8x_5^4 + x_1^5x_2x_3^5x_4^5x_5^8 \\ + x_1^5x_2x_3^4x_4^3x_5^8 + x_1^3x_2^3x_3^4x_4^8x_5^5, \\ g_2 = x_1^5x_2^3x_3^2x_4^8x_5^2 + x_1^5x_2^3x_3^2x_4^6x_5^4 + x_1^5x_2^3x_3^4x_4^6x_5^2 + x_1^2x_3^3x_3^2x_4^9x_5^4 + x_1^3x_2^2x_3^2x_4^9x_5^4 \\ + x_1^3x_2x_3^2x_4^{10}x_5^4 + x_1^6x_2x_3^2x_4^7x_5^4 + x_1^2x_3^2x_3^4x_4^9x_5^2 + x_1^3x_2^2x_3^4x_4^9x_5^2 + x_1^3x_2x_3^2x_4^{10}x_5^2 \\ + x_1^6x_2x_3^4x_4^7x_5^2 + x_1^2x_3^3x_3^8x_4^5x_5^2 + x_1^2x_3^2x_3^2x_4^5x_5^8 + x_1^3x_2^2x_3^8x_4^5x_5^2 + x_1^3x_2^2x_3^2x_4^5x_5^8 \\ + x_1^3x_2x_3^8x_4^6x_5^2 + x_1^6x_2x_3^6x_4^5x_5^2 + x_1^3x_2^2x_3^6x_4^5x_5^4 + x_1^3x_2^2x_3^5x_4^6x_5^4 + x_1^3x_2^2x_3^3x_4^8x_4^5 \\ + x_1^2x_2^3x_3^8x_4^4x_5^4 + x_1^6x_2x_3^2x_4^3x_5^8 + x_1^3x_2x_3^2x_4^6x_5^8 + x_1^3x_2^2x_4^3x_3^8x_5^8 + x_1^6x_2x_3^3x_4^6x_4^5 \\ + x_1^2x_2^3x_3^4x_4^8x_5^2 + x_1^2x_2x_3^{10}x_4^5x_5^2 + x_1^2x_2x_3^2x_4^7x_5^8 + x_1^2x_2x_3^2x_4^{11}x_5^4 + x_1^2x_2x_3^3x_4^6x_5^8 \\ + x_1^2x_2x_3^3x_4^{10}x_5^4 + x_1^2x_2x_3^4x_4^{11}x_5^2 + x_1^2x_2x_3^6x_4^9x_5^2 + x_1^2x_2x_3^8x_4^7x_5^2, \\ g_4 = x_1^3x_2^3x_3^2x_4^6x_5^4 + x_1^3x_2^3x_3^4x_4^6x_5^2 + x_1^4x_2x_3^2x_4^7x_5^4 \\ + x_1^4x_2x_3^4x_4^7x_5^2 + x_1^4x_2x_3^6x_4^5x_5^2 + x_1^4x_2x_3^3x_4^6x_4^5.$$

The above relations imply that y is also strictly inadmissible. The lemma is proved. $\square$

Proof of Proposition 3.2.10. We denote by $\mathcal{Y}_t := \mathcal{Y}_{22,t}$, $461 \leq t \leq 510$ the admissible monomials in $\mathcal{B}_5^+(\omega_{(1)})$ (see Sect.6.5.) For $x \in \mathcal{B}_5^+(\omega_{(1)})$, we have $x = X_{\{i,j,k\}}y^2$ with y a monomial of degree 10 in $\mathcal{P}_5$, and $1 \leq i < j < k \leq 5$. Since x is admissible, by Theorem 2.2.5, $y \in \mathcal{B}_5(2,2,1)$.

Let $z \in \mathcal{B}_5(2,2,1)$ such that $X_{\{i,j,k\},5}y_1^2 \in \mathcal{P}_5^+$. By a direct computation using a result in [41], we see that if $X_{\{i,j,k\},5}y_1^2 \neq \mathcal{Y}_t$, for all t , $461 \leq t \leq 510$, then there is a monomial w which is given in one of Lemmas 3.2.12 - 3.2.13 such that $X_{\{i,j,k\},5}y_1^2 = wz^{2^u}$ with suitable monomial $z \in \mathcal{P}_5$ and $u = \max\{j \in \mathbb{Z} : \omega_j(w) > 0\}$. By Theorem 2.2.5, $X_{\{i,j,k\},5}y_1^2$ is inadmissible. Since $x = X_{\{i,j,k\},5}y^2$ and x is admissible, one gets $x = \mathcal{Y}_t$. This implies $Q\mathcal{P}_5^+(\omega_{(1)})$ is spanned by the set $\{[\mathcal{Y}_t := \mathcal{Y}_{22,t}]_{\omega_{(1)}} : 461 \leq t \leq 510\}$.

We now prove the set $\{[\mathcal{Y}_t]_{\omega_{(1)}} : 461 \leq t \leq 510\}$ is linearly independent in $Q\mathcal{P}_5(\omega_{(1)})$. Suppose there is a linear relation

$$(3.2) \quad \mathcal{S} = \sum_{461 \leq t \leq 510} \gamma_t \mathcal{Y}_t \equiv_{\omega_{(1)}} 0,$$

where $\gamma_t \in \mathbb{Z}/2$. From a result in [34], $\dim Q\mathcal{P}_4^+(\omega_{(1)}) = 26$, with the basis $\{[u_j]_{\omega_{(1)}} : 1 \leq j \leq 26\}$, where

$$\begin{aligned} u_1. & x_1x_2x_3^6x_4^{14}, & u_2. & x_1x_2x_3^{14}x_4^6, & u_3. & x_1x_2^2x_3^4x_4^{15}, & u_4. & x_1x_2^2x_3^5x_4^{14}, & u_5. & x_1x_2^2x_3^7x_4^{12}, \\ u_6. & x_1x_2^2x_3^{12}x_4^7, & u_7. & x_1x_2^2x_3^{13}x_4^6, & u_8. & x_1x_2^2x_3^{15}x_4^4, & u_9. & x_1x_2^3x_3^4x_4^{14}, & u_{10}. & x_1x_2^3x_3^6x_4^{12}, \\ u_{11}. & x_1x_2^3x_3^{12}x_4^6, & u_{12}. & x_1x_2^3x_3^{14}x_4^4, & u_{13}. & x_1x_2^6x_3x_4^{14}, & u_{14}. & x_1x_2^7x_3^2x_4^{12}, & u_{15}. & x_1x_2^{14}x_3x_4^6, \\ u_{16}. & x_1x_2^{15}x_3^2x_4^4, & u_{17}. & x_1^3x_2x_3^4x_4^{14}, & u_{18}. & x_1^3x_2x_3^6x_4^{12}, & u_{19}. & x_1^3x_2x_3^{12}x_4^6, & u_{20}. & x_1^3x_2x_3^{14}x_4^4, \\ u_{21}. & x_1^3x_2^5x_3^2x_4^{12}, & u_{22}. & x_1^3x_2^5x_3^6x_4^8, & u_{23}. & x_1^3x_2^5x_3^{10}x_4^4, & u_{24}. & x_1^3x_2^{13}x_3^2x_4^4, & u_{25}. & x_1^7x_2x_3^2x_4^{12}, \\ u_{26}. & x_1^{15}x_2x_3^2x_4^4. \end{aligned}$$

Consider the homomorphism $\pi_{(1;2)} : \mathcal{P}_5 \rightarrow \mathcal{P}_4$. By a direct computation using Theorem 2.3.3 and Proposition 3.1.1, we have

$$\begin{aligned} \pi_{(1;2)}(\mathcal{S}) & \equiv_{\omega_{(1)}} (\gamma_{473} + \gamma_{475} + \gamma_{481} + \gamma_{483})u_1 + \gamma_{477}u_6 + \gamma_{478}u_7 \\ & + (\gamma_{476} + \gamma_{477} + \gamma_{478})u_4 + \gamma_{483}(u_9 + u_{14}) + \gamma_{481}(u_{10} + u_{11}) \\ & + (\gamma_{475} + \gamma_{476} + \gamma_{477} + \gamma_{478})u_{13} + (\gamma_{470} + \gamma_{481} + \gamma_{484})u_{18} \\ & + (\gamma_{471} + \gamma_{478} + \gamma_{481} + \gamma_{484})u_{19} + \gamma_{472}u_{20} + (\gamma_{479} + \gamma_{484})u_{21} \\ & + (\gamma_{480} + \gamma_{481} + \gamma_{484})u_{22} + \gamma_{482}u_{23} + \gamma_{485}u_{24} + \gamma_{495}u_{25} + \gamma_{497}u_{26} \\ & + (\gamma_{474} + \gamma_{478} + \gamma_{481} + \gamma_{484})u_2 + \gamma_{476}u_5 + (\gamma_{469} + \gamma_{475})u_{17} + \gamma_{484}u_{15} \equiv_{\omega_{(1)}} 0. \end{aligned}$$

This relation implies

$$\begin{aligned} (3.3) \quad \gamma_{469} & = \gamma_{470} = \gamma_{471} = \gamma_{472} = \gamma_{473} = \gamma_{474} = \gamma_{475} = \gamma_{476} \\ & = \gamma_{477} = \gamma_{478} = \gamma_{479} = \gamma_{480} = \gamma_{481} = \gamma_{482} = \gamma_{483} \\ & = \gamma_{484} = \gamma_{485} = \gamma_{495} = \gamma_{497} = 0. \end{aligned}$$

Substituting (3.3) into the relation (3.2), we have

$$(3.4) \quad \sum_{461 \leq t \leq 468} \gamma_t \mathcal{Y}_t + \sum_{486 \leq t \leq 494} \gamma_t \mathcal{Y}_t + \gamma_{496} \mathcal{Y}_{36} + \sum_{498 \leq t \leq 510} \gamma_t \mathcal{Y}_t \equiv_{\omega_{(1)}} 0.$$

Applying the homomorphisms $\pi_{(1;3)}, \pi_{(1;4)} : \mathcal{P}_5 \rightarrow \mathcal{P}_4$ to (3.4), we get

$$(3.5) \quad \begin{cases} \gamma_t = 0, \quad t \in \mathbb{J}, \\ \gamma_{464} = \gamma_{487} = \gamma_{499} = \gamma_{508}, \gamma_{468} = \gamma_{494} = \gamma_{506} = \gamma_{509}, \gamma_{498} = \gamma_{502}, \\ \gamma_{467} + \gamma_{468} + \gamma_{503} = \gamma_{467} + \gamma_{468} + \gamma_{505} = 0, \\ \gamma_{462} + \gamma_{468} + \gamma_{502} + \gamma_{508} = \gamma_{463} + \gamma_{468} + \gamma_{502} + \gamma_{508} = 0, \\ \gamma_{466} + \gamma_{501} + \gamma_{502} + \gamma_{508} = \gamma_{466} + \gamma_{468} + \gamma_{504} + \gamma_{507} + \gamma_{508} = 0. \end{cases}$$

Here $\mathbb{J} = \{461, 465, 486, 488, 489, 490, 491, 492, 493, 496, 500\}$. Then, combining (3.3), (3.5), and the relation $\pi_{(1;5)}(\mathcal{S}) \equiv_{\omega_{(1)}} 0$, we obtain $\gamma_t = 0$ for $461 \leq t \leq 510$. The proposition is proved. $\square$

Using a similar technique as mentioned in the proof of Proposition 3.2.10, we obtain

Proposition 3.2.14.

(I) $\mathcal{B}_5^+(\omega_{(2)}) = \mathcal{B}^+(5, \omega_{(2)}) \cup \mathcal{E}$, where $\mathcal{E}$ is the set of the following monomials:

$$\begin{aligned} & x_1x_2^2x_3^2x_4^3x_5^{14}, \quad x_1x_2^2x_3^3x_4^2x_5^{14}, \quad x_1x_2^2x_3^3x_4^6x_5^{10}, \quad x_1x_2^2x_3^3x_4^{14}x_5^2, \quad x_1x_2^3x_3^2x_4^2x_5^{14}, \\ & x_1x_2^3x_3^2x_4^6x_5^{10}, \quad x_1x_2^3x_3^2x_4^{14}x_5^2, \quad x_1x_2^3x_3^6x_4^2x_5^{10}, \quad x_1x_2^3x_3^6x_4^{10}x_5^2, \quad x_1x_2^3x_3^{14}x_4^2x_5^2, \\ & x_1^3x_2x_3^2x_4^2x_5^{14}, \quad x_1^3x_2x_3^2x_4^6x_5^{10}, \quad x_1^3x_2x_3^2x_4^{14}x_5^2, \quad x_1^3x_2x_3^6x_4^2x_5^{10}, \quad x_1^3x_2x_3^6x_4^{10}x_5^2, \\ & x_1^3x_2x_3^{14}x_4^2x_5^2, \quad x_1^3x_2x_3^5x_4^2x_5^{10}, \quad x_1^3x_2x_3^5x_4^{10}x_5^2, \quad x_1^3x_2x_3^{10}x_4^2x_5^2, \quad x_1^3x_2x_3^{13}x_4^2x_5^2, \end{aligned}$$

(II) $\mathcal{B}_5^+(\omega_{(3)}) = \{x_1x_2^3x_3^6x_4^6x_5^6, x_1^3x_2x_3^6x_4^6x_5^6, x_1^3x_2^5x_3^2x_4^6x_5^6, x_1^3x_2^5x_3^6x_4^2x_5^6, x_1^3x_2^5x_3^6x_4^6x_5^2\}$,

(III) $\mathcal{B}_5^+(\omega_{(4)}) = \overline{\Phi}^+(\mathcal{B}_4(\omega_{(4)}) \cup \mathcal{B}^+(5, \omega_{(4)}))$,

(IV) $\mathcal{B}_5^+(\omega_{(5)}) = \overline{\Phi}^+(\mathcal{B}_4(\omega_{(5)}) \cup \mathcal{B}^+(5, \omega_{(5)}))$.

A direct computation shows: $|\mathcal{B}_5^+(\omega_{(2)})| = 25$, $|\mathcal{B}_5^+(\omega_{(4)})| = 300$ and $|\mathcal{B}_5^+(\omega_{(5)})| = 125$ (see Sect.6.5.) On the other hand, by Remark 3.2.9, we have $(Q\mathcal{P}_5)_{22}^+ = \bigoplus_{1 \leq j \leq 5} Q\mathcal{P}_5^+(\omega_{(j)})$. Combining this with the above results, we obtain

Corollary 3.2.15. $(Q\mathcal{P}_5^+)_{22}$ is the $\mathbb{Z}/2$ -vector space of dimension 505 with a basis consisting of all the classes represented by the monomials $\mathcal{Y}_{22,t}$, $1 \leq t \leq 505$, which are determined in Sect.6.5.

Computation of $\text{Ker}((\widetilde{Sq}_*)_{(5,47)})$

We consider the weight vectors

$$\overline{\omega}_{(1)} = (3, 2, 2, 2, 1), \quad \overline{\omega}_{(2)} = (3, 2, 4, 1, 1), \quad \overline{\omega}_{(3)} = (3, 2, 4, 3),$$

$$\overline{\omega}_{(4)} = (3, 4, 3, 1, 1), \quad \overline{\omega}_{(5)} = (3, 4, 3, 3).$$

It is easy to see that $\deg \overline{\omega}_{(1)} = \deg \overline{\omega}_{(2)} = \dots = \deg \overline{\omega}_{(5)} = 47$. Then, from Remarks 3.2.8 and 3.2.9, one gets the following.

Lemma 3.2.16. *If X is an admissible monomial of degree 47 in $\mathcal{P}_5$ and $[X] \in \text{Ker}((\widetilde{Sq}_*)_{(5,47)})$, then $\omega(X)$ is one of the sequences $\overline{\omega}_{(k)}$, $1 \leq k \leq 5$.*

From this lemma, we have

$$\begin{aligned} \text{Ker}((\widetilde{Sq}_*)_{(5,47)}) &= (Q\mathcal{P}_5^0)_{47} \oplus (\text{Ker}((\widetilde{Sq}_*)_{(5,47)}) \cap (Q\mathcal{P}_5^+)_{47}) \\ &= (Q\mathcal{P}_5^0)_{47} \oplus (\oplus_{1 \leq i \leq 5} Q\mathcal{P}_5^+(\overline{\omega}_{(i)})). \end{aligned}$$

Recall that $(Q\mathcal{P}_5^0)_{47} = \bigoplus_{1 \leq t \leq 4} \bigoplus_{1 \leq \ell \leq \binom{5}{t}} (Q\mathcal{P}_t^+)_{47}$. Then, by a direct computation using a result in

[19], [9] and [34], we conclude

$$\dim(Q\mathcal{P}_5^0)_{47} = \binom{5}{3} \cdot 14 + \binom{5}{4} \cdot 84 = 560.$$

The set $\mathcal{B}_5^0(47) = \mathcal{B}_5^0(\overline{\omega}_{(1)}) = \overline{\Phi}^0(\mathcal{B}_4(47)) = \{\mathcal{Y}_{47,i} : 1 \leq i \leq 560\}$ is described in Sect.6.6. We now determine $Q\mathcal{P}_5^+(\overline{\omega}_{(j)})$ for $1 \leq j \leq 5$.

Lemma 3.2.17. *The following monomials are strictly inadmissible:*

- I) $X_1 = x_1^3 x_2^{12} x_k x_\ell^3 x_m^{12}$, $X_2 = x_1^3 x_2^4 x_k^3 x_\ell^8 x_m^{13}$, $X_3 = x_1 x_2^{14} x_k^2 x_\ell x_m^{13}$,
 $X_5 = x_1^3 x_2^{14} x_k^{12} x_\ell x_m$, $X_6 = x_1^7 x_2^{10} x_k^{12} x_\ell x_m$, $X_7 = x_1^3 x_2^2 x_k^{12} x_\ell x_m^{13}$,
 $X_8 = x_1^3 x_2^{12} x_k^2 x_\ell x_m^{13}$, $X_9 = x_1^{15} x_2^2 x_k x_\ell^4 x_m^9$, $X_{10} = x_1^{15} x_2^2 x_k x_\ell^5 x_m^8$,
 $X_{11} = x_1^7 x_2^2 x_k x_\ell^8 x_m^{13}$, $X_{12} = x_1^7 x_2^2 x_k x_\ell^9 x_m^{12}$, $X_{13} = x_1^{15} x_2^2 x_k^{12} x_\ell x_m$,
 $X_{14} = x_1^3 x_2^4 x_k x_\ell^8 x_m^{15}$, $X_{15} = x_1^3 x_2^4 x_k x_\ell^9 x_m^{14}$, $X_{16} = x_1^3 x_2^{14} x_k x_\ell^4 x_m^9$,
 $X_{17} = x_1^3 x_2^5 x_k x_\ell^8 x_m^{14}$, $X_{18} = x_1^3 x_2^6 x_k x_\ell^8 x_m^{13}$, $X_{19} = x_1^3 x_2^6 x_k x_\ell^9 x_m^{12}$,
 $X_{20} = x_1^7 x_2^{10} x_k x_\ell^4 x_m^9$, $X_{21} = x_1^7 x_2^{10} x_k x_\ell^5 x_m^8$, $X_{22} = x_1^3 x_2^2 x_k^5 x_\ell^8 x_m^{13}$,
 $X_{23} = x_1^3 x_2^2 x_k^5 x_\ell^9 x_m^{12}$, $X_{24} = x_1^7 x_2^2 x_k^5 x_\ell^8 x_m^9$, $X_{25} = x_1 x_2^{14} x_k^2 x_\ell x_m^{13}$,
 $X_{26} = x_1^3 x_2^6 x_k^5 x_\ell^8 x_m^9$, $X_{27} = x_1^3 x_2^{12} x_k^{14} x_\ell x_m$. Here (k, ℓ, m) is a permutation of $(3, 4, 5)$;
II) $X_{27} = x_i^3 x_j^2 x_k^{13} x_\ell^4 x_m^9$, $j < k$, $X_{28} = x_i x_j^2 x_k^6 x_\ell^9 x_m^{13}$, $X_{29} = x_i^3 x_j^4 x_k^6 x_\ell^9 x_m^9$, where (i, j, k, ℓ, m) is a permutation of $(1, 2, 3, 4, 5)$.

Proof. It is easy to see that $\omega(X_t) = \omega^* := (3, 2, 2, 2)$ for $1 \leq t \leq 29$. We prove the lemma for the monomials $X_1 = x_1^3 x_2^{12} x_k x_\ell^3 x_m^{12}$ and $X_2 = x_1^3 x_2^4 x_k^3 x_\ell^8 x_m^{13}$, where (k, ℓ, m) is a permutation of $(3, 4, 5)$. The others can be proved by a similar technique. We have

$$\begin{aligned} X_1 &= x_1^2 x_2^{11} x_k x_\ell^5 x_m^{12} + x_1^2 x_2^{11} x_k^4 x_\ell^5 x_m^9 + x_1^2 x_2^{11} x_k^8 x_\ell^5 x_m^5 + x_1^2 x_2^{13} x_k x_\ell^3 x_m^{12} \\ &\quad + x_1^2 x_2^{13} x_k^4 x_\ell^3 x_m^9 + x_1^2 x_2^{13} x_k^8 x_\ell^3 x_m^5 + x_1^3 x_2^3 x_k^8 x_\ell^5 x_m^{12} + x_1^3 x_2^5 x_k^8 x_\ell^5 x_m^{10} \\ &\quad + x_1^3 x_2^5 x_k^8 x_\ell^6 x_m^9 + x_1^3 x_2^7 x_k^4 x_\ell^8 x_m^9 + x_1^3 x_2^7 x_k^8 x_\ell^5 x_m^8 + x_1^3 x_2^7 x_k^8 x_\ell^8 x_m^5 \\ &\quad + x_1^3 x_2^9 x_k^2 x_\ell^5 x_m^{12} + x_1^3 x_2^9 x_k^4 x_\ell^3 x_m^{12} + x_1^3 x_2^9 x_k^4 x_\ell^5 x_m^{10} + x_1^3 x_2^9 x_k^4 x_\ell^6 x_m^9 \\ &\quad + x_1^3 x_2^9 x_k^8 x_\ell^5 x_m^6 + x_1^3 x_2^9 x_k^8 x_\ell^6 x_m^5 + x_1^3 x_2^{11} x_k x_\ell^4 x_m^{12} \\ &\quad + Sq^1(g_1) + Sq^2(g_2) + Sq^4(g_4) + Sq^8(x_1^6 x_2^5 x_k^4 x_\ell^3 x_m^5) \text{ modulo } (\mathcal{P}_5^-(\omega^*)), \text{ where} \\ g_1 &= x_1^3 x_2^7 x_k x_\ell^3 x_m^{16} + x_1^3 x_2^{11} x_k x_\ell^3 x_m^{12} + x_1^5 x_2^3 x_k^8 x_\ell^5 x_m^9 + x_1^5 x_2^7 x_k^4 x_\ell^5 x_m^9 + x_1^5 x_2^7 x_k^8 x_\ell^5 x_m^5, \\ g_2 &= x_1^2 x_2^{11} x_k x_\ell^3 x_m^{12} + x_1^2 x_2^{11} x_k^4 x_\ell^3 x_m^9 + x_1^2 x_2^{11} x_k^8 x_\ell^3 x_m^5 + x_1^3 x_2^3 x_k^8 x_\ell^5 x_m^{10} + x_1^3 x_2^3 x_k^8 x_\ell^6 x_m^9 \\ &\quad + x_1^3 x_2^7 x_k^4 x_\ell^5 x_m^{12} + x_1^3 x_2^7 x_k^4 x_\ell^6 x_m^9 + x_1^3 x_2^7 x_k^8 x_\ell^5 x_m^6 + x_1^3 x_2^7 x_k^8 x_\ell^6 x_m^5 + x_1^5 x_2^7 x_k^2 x_\ell^3 x_m^{12} \\ &\quad + x_1^6 x_2^3 x_k^8 x_\ell^3 x_m^9 + x_1^6 x_2^7 x_k^4 x_\ell^3 x_m^9 + x_1^6 x_2^7 x_k^8 x_\ell^3 x_m^5, \\ g_4 &= x_1^3 x_2^7 x_k^2 x_\ell^3 x_m^{12} + x_1^4 x_2^7 x_k^4 x_\ell^3 x_m^9 + x_1^4 x_2^7 x_k^8 x_\ell^3 x_m^5 + x_1^{10} x_2^5 x_k^4 x_\ell^3 x_m^5. \end{aligned}$$

This equality implies that X_1 is strictly inadmissible.

Next, we show that X_2 is also strictly inadmissible. Indeed, using Cartan's formula, we obtain

$$\begin{aligned} X_2 = & x_1^2 x_2 x_k^3 x_\ell^{12} x_m^{13} + x_1^2 x_2 x_k^5 x_\ell^9 x_m^{14} + x_1^2 x_2 x_k^5 x_\ell^{10} x_m^{13} + x_1^2 x_2 x_k^6 x_\ell^9 x_m^{13} + x_1^2 x_2 x_k^{10} x_\ell^5 x_m^{13} \\ & + x_1^2 x_2 x_k^{12} x_\ell^5 x_m^{11} + x_1^2 x_2 x_k^9 x_\ell^{14} x_m^{13} + x_1^2 x_2 x_k^8 x_\ell^{13} x_m^{13} + x_1^2 x_2 x_k^8 x_\ell^5 x_m^{11} \\ & + x_1^3 x_2 x_k^4 x_\ell^9 x_m^{14} + x_1^3 x_2 x_k^4 x_\ell^{10} x_m^{13} + x_1^3 x_2 x_k^6 x_\ell^8 x_m^{13} + x_1^3 x_2 x_k^8 x_\ell^6 x_m^{13} + x_1^3 x_2 x_k^4 x_\ell^9 x_m^{13} \\ & + x_1^3 x_2 x_k^5 x_\ell^8 x_m^{13} + x_1^3 x_2 x_k^8 x_\ell^4 x_m^{13} + x_1^3 x_2 x_k^8 x_\ell^5 x_m^{12} + x_1^3 x_2 x_k^9 x_\ell^{14} x_m^{13} + x_1^3 x_2 x_k^{10} x_\ell^{13} x_m^{13} \\ & + x_1^3 x_2 x_k^2 x_\ell^9 x_m^{13} \\ & + Sq^1(Z_1) + Sq^2(Z_2) + Sq^4(Z_4) + Sq^8(x_1^3 x_2^4 x_k^4 x_\ell^5 x_m^7) \text{ modulo } (\mathcal{P}_5^-(\omega^*)), \end{aligned}$$

where

$$\begin{aligned} Z_1 = & x_1^3 x_2 x_k^3 x_\ell^5 x_m^{18} + x_1^3 x_2 x_k^3 x_\ell^6 x_m^{17} + x_1^3 x_2 x_k^3 x_\ell^9 x_m^{14} + x_1^3 x_2 x_k^3 x_\ell^{10} x_m^{13} \\ & + x_1^3 x_2 x_k^6 x_\ell^9 x_m^{11} + x_1^3 x_2 x_k^{10} x_\ell^5 x_m^{11} + x_1^3 x_2 x_k^6 x_\ell^{17} x_m^{13} + x_1^3 x_2 x_k^{10} x_\ell^{13} x_m^{13} \\ & + x_1^3 x_2 x_k^{12} x_\ell^{11} x_m^{11} + x_1^3 x_2 x_k^{16} x_\ell^7 x_m^{11} + x_1^3 x_2 x_k^4 x_\ell^{17} x_m^{13} + x_1^3 x_2 x_k^8 x_\ell^{13} x_m^{13} \\ & + x_1^3 x_2 x_k^8 x_\ell^5 x_m^{11} + x_1^3 x_2 x_k^9 x_\ell^7 x_m^{11} + x_1^3 x_2 x_k^5 x_\ell^{13} x_m^{13} + x_1^3 x_2 x_k^5 x_\ell^{14} x_m^{13}, \\ Z_2 = & x_1^2 x_2 x_k^3 x_\ell^9 x_m^{14} + x_1^2 x_2 x_k^3 x_\ell^{10} x_m^{13} + x_1^2 x_2 x_k^6 x_\ell^9 x_m^{11} + x_1^2 x_2 x_k^{10} x_\ell^5 x_m^{11} \\ & + x_1^2 x_2 x_k^8 x_\ell^{14} x_m^{13} + x_1^2 x_2 x_k^{10} x_\ell^{12} x_m^{11} + x_1^2 x_2 x_k^8 x_\ell^{13} x_m^{13} \\ & + x_1^2 x_2 x_k^8 x_\ell^5 x_m^{11} + x_1^2 x_2 x_k^9 x_\ell^7 x_m^{11} + x_1^2 x_2 x_k^5 x_\ell^{14} x_m^{13} + x_1^2 x_2 x_k^5 x_\ell^{14} x_m^{13} \\ & + x_1^2 x_2 x_k^6 x_\ell^{14} x_m^{13} + x_1^2 x_2 x_k^6 x_\ell^{11} x_m^{11} + x_1^2 x_2 x_k^2 x_\ell^9 x_m^{11} + x_1^2 x_2 x_k^3 x_\ell^5 x_m^{14} \\ & + x_1^5 x_2^2 x_k^3 x_\ell^6 x_m^{13} + x_1^5 x_2^2 x_k^6 x_\ell^5 x_m^{11} + x_1^5 x_2^2 x_k^8 x_\ell^3 x_m^{11} + x_1^5 x_2^2 x_k^6 x_\ell^{14} x_m^{13} \\ & + x_1^5 x_2^3 x_k^2 x_\ell^6 x_m^{13} + x_1^5 x_2^3 x_k^2 x_\ell^8 x_m^{11} + x_1^5 x_2^3 x_k^2 x_\ell^{10} x_m^9 + x_1^5 x_2^3 x_k^2 x_\ell^{12} x_m^7 \\ & + x_1^5 x_2^3 x_k^4 x_\ell^3 x_m^{14} + x_1^5 x_2^3 x_k^4 x_\ell^6 x_m^{11} + x_1^5 x_2^3 x_k^4 x_\ell^{10} x_m^7 + x_1^6 x_2^3 x_k x_\ell^5 x_m^{14}, \\ Z_4 = & x_1^3 x_2 x_k^6 x_\ell^5 x_m^{12} + x_1^3 x_2 x_k^6 x_\ell^6 x_m^{11} + x_1^3 x_2 x_k^2 x_\ell^9 x_m^{11} + x_1^3 x_2 x_k^3 x_\ell^6 x_m^{13} + x_1^3 x_2 x_k^6 x_\ell^5 x_m^{11} \\ & + x_1^3 x_2 x_k^8 x_\ell^3 x_m^{11} + x_1^3 x_2 x_k^6 x_\ell^{14} x_m^{13} + x_1^3 x_2 x_k^2 x_\ell^6 x_m^{13} + x_1^3 x_2 x_k^2 x_\ell^8 x_m^{11} + x_1^3 x_2 x_k^2 x_\ell^{10} x_m^9 \\ & + x_1^3 x_2 x_k^2 x_\ell^{12} x_m^7 + x_1^3 x_2 x_k^4 x_\ell^3 x_m^{14} + x_1^3 x_2 x_k^4 x_\ell^6 x_m^{11} + x_1^3 x_2 x_k^4 x_\ell^{10} x_m^7 + x_1^3 x_2 x_k^4 x_\ell^5 x_m^{14} \\ & + x_1^3 x_2 x_k^8 x_\ell^4 x_\ell^5 x_m^7 + x_1^4 x_2^3 x_k x_\ell^5 x_m^{14}. \end{aligned}$$

The above relations imply that X_2 is strictly inadmissible. The lemma follows. $\square$

Lemma 3.2.18. *The following monomials are strictly inadmissible:*

$$\begin{aligned} Y_1 &= x_1^3 x_2^4 x_3^8 x_4^9, & Y_2 &= x_1^3 x_2^7 x_3^8 x_4^5 x_5^8, & Y_3 &= x_1 x_2^6 x_3^{10} x_4 x_5^{13}, & Y_4 &= x_1 x_2^6 x_3^{10} x_4^{13} x_5, \\ Y_5 &= x_1 x_2^6 x_3^{11} x_4^{12} x_5, & Y_6 &= x_1 x_2^7 x_3^{10} x_4^{12} x_5, & Y_7 &= x_1^7 x_2 x_3^{10} x_4^{12} x_5, & Y_8 &= x_1 x_2^2 x_3^{13} x_4^{13} x_5^{13}, \\ Y_9 &= x_1 x_2^2 x_3^{14} x_4^5 x_5^9, & Y_{10} &= x_1 x_2^{14} x_3^5 x_4^5 x_5^9, & Y_{11} &= x_1^3 x_2^3 x_3^{12} x_4^{12} x_5, & Y_{12} &= x_1^3 x_2^{15} x_3^4 x_4^8 x_5, \\ Y_{13} &= x_1^3 x_2^{15} x_3^4 x_4^8 x_5, & Y_{14} &= x_1^{15} x_2^3 x_3^4 x_4^8 x_5, & Y_{15} &= x_1^{15} x_2^3 x_3^4 x_4^8 x_5, & Y_{16} &= x_1 x_2^{14} x_3^3 x_4^4 x_5^9, \\ Y_{17} &= x_1^3 x_2^4 x_3^{10} x_4^{13} x_5, & Y_{18} &= x_1^3 x_2^4 x_3^{10} x_4^{13} x_5, & Y_{19} &= x_1^3 x_2^4 x_3^{11} x_4^{12} x_5, & Y_{20} &= x_1 x_2^{14} x_3^3 x_4^5 x_5^8, \\ Y_{21} &= x_1^3 x_2^{14} x_3^5 x_4^8 x_5, & Y_{22} &= x_1^3 x_2^{14} x_3^5 x_4^8 x_5, & Y_{23} &= x_1^3 x_2^{14} x_3^5 x_4^8 x_5, & Y_{24} &= x_1 x_2^6 x_3^3 x_4^{13} x_5^8, \\ Y_{25} &= x_1 x_2^{14} x_3^{14} x_4 x_5, & Y_{26} &= x_1^3 x_2^7 x_3^8 x_4^{12} x_5, & Y_{27} &= x_1^3 x_2^7 x_3^{12} x_4^8 x_5, & Y_{28} &= x_1^3 x_2^7 x_3^{12} x_4^8 x_5, \\ Y_{29} &= x_1^7 x_2^3 x_3^8 x_4^{12} x_5, & Y_{30} &= x_1^7 x_2^3 x_3^{12} x_4^8 x_5, & Y_{31} &= x_1^7 x_2^3 x_3^{12} x_4^8 x_5, & Y_{32} &= x_1 x_2^6 x_3^{11} x_4^4 x_5^9, \\ Y_{33} &= x_1^7 x_2^{11} x_3^4 x_4^8 x_5, & Y_{34} &= x_1^7 x_2^{11} x_3^4 x_4^8 x_5, & Y_{35} &= x_1 x_2^6 x_3^{11} x_4^5 x_5^8, & Y_{36} &= x_1 x_2^6 x_3^{10} x_4^5 x_5^9, \\ Y_{37} &= x_1^3 x_2^5 x_3^8 x_4^{13} x_5, & Y_{38} &= x_1^3 x_2^{12} x_3^5 x_4^5 x_5^9, & Y_{39} &= x_1^3 x_2^5 x_3^9 x_4^2 x_5^{12}, & Y_{40} &= x_1^7 x_2^2 x_3^4 x_4^9 x_5^9, \\ Y_{41} &= x_1^7 x_2^2 x_3^9 x_4^9 x_5, & Y_{42} &= x_1^3 x_2^4 x_3^3 x_4^{12} x_5^9, & Y_{43} &= x_1^3 x_2^{12} x_3^3 x_4^4 x_5^9, & Y_{44} &= x_1^3 x_2^4 x_3^9 x_4^3 x_5^{12}, \\ Y_{45} &= x_1^3 x_2^{12} x_3^3 x_4^5 x_5^8, & Y_{46} &= x_1^3 x_2^5 x_3^3 x_4^{12} x_5, & Y_{47} &= x_1^3 x_2^4 x_3^{11} x_4^4 x_5^9, & Y_{48} &= x_1^3 x_2^4 x_3^{11} x_4^5 x_5^8, \\ Y_{49} &= x_1^3 x_2^4 x_3^{10} x_4^5 x_5^9, & Y_{50} &= x_1 x_2^6 x_3^3 x_4^{12} x_5^9, & Y_{51} &= x_1^3 x_2^4 x_3^7 x_4^9 x_5^8, & Y_{52} &= x_1^3 x_2^7 x_3^8 x_4^4 x_5^9, \\ Y_{53} &= x_1^7 x_2^3 x_3^8 x_4^4 x_5^9, & Y_{54} &= x_1 x_2^{14} x_3^3 x_4^{12} x_5, & Y_{55} &= x_1^7 x_2^3 x_3^8 x_4^5 x_5^8. \end{aligned}$$

Proof. Note that $\omega(Y_j) = \omega^*$, $j = 1, 2, \dots, 55$. We prove this lemma for the monomials $Y_1 = x_1^3 x_2^4 x_3^8 x_4^9$, and $Y_2 = x_1^3 x_2^7 x_3^8 x_4^5 x_5^8$. The others can be proved by a similar computation. A direct computation shows:

$$\begin{aligned} Y_1 = & x_1^2 x_2 x_3^7 x_4^8 x_5^{13} + x_1^2 x_2 x_3^{13} x_4 x_5^{14} + x_1^2 x_2 x_3^{13} x_4^8 x_5^7 + x_1^2 x_2 x_3^9 x_4^4 x_5^{13} + x_1^2 x_2 x_3^{12} x_4 x_5^{13} \\ & + x_1^2 x_2^5 x_3^9 x_4^4 x_5^{11} + x_1^2 x_2^5 x_3^9 x_4^8 x_5^7 + x_1^2 x_2^5 x_3^{12} x_4 x_5^{11} + x_1^3 x_2 x_3^5 x_4^8 x_5^{14} + x_1^3 x_2 x_3^7 x_4^8 x_5^{12} \\ & + x_1^3 x_2 x_3^{12} x_4 x_5^{14} + x_1^3 x_2 x_3^{12} x_4^4 x_5^{11} + x_1^3 x_2^2 x_3^9 x_4^4 x_5^{13} + x_1^3 x_2^2 x_3^9 x_4^4 x_5^{12} + x_1^3 x_2^2 x_3^{12} x_4 x_5^{12} \\ & + x_1^3 x_2^2 x_3^{12} x_4^9 x_5^9 + Sq^1(u_1) + Sq^2(u_2) + Sq^4(u_4) + Sq^8(u_8) \text{ modulo } (\mathcal{P}_5^-(\omega^*)), \text{ where} \end{aligned}$$

$$u_1 = x_1^3 x_2 x_3^5 x_4^8 x_5^{13} + x_1^3 x_2 x_3^7 x_4 x_5^{18} + x_1^3 x_2 x_3^9 x_4^4 x_5^{13} + x_1^3 x_2 x_3^{11} x_4 x_5^{14} \\ + x_1^3 x_2^3 x_3^9 x_4^4 x_5^{11} + x_1^3 x_2^3 x_3^9 x_4^8 x_5^7 + x_1^3 x_2^3 x_3^{12} x_4 x_5^{11} + x_1^3 x_2^3 x_3^{16} x_4 x_5^7 \\ + x_1^3 x_2^4 x_3^9 x_4^4 x_5^{13} + x_1^5 x_2 x_3^5 x_4^8 x_5^{11} + x_1^5 x_2 x_3^7 x_4^8 x_5^9,$$

$$u_2 = x_1^2 x_2 x_3^7 x_4^8 x_5^{11} + x_1^2 x_2 x_3^{11} x_4 x_5^{14} + x_1^2 x_2 x_3^{11} x_4^8 x_5^7 + x_1^2 x_2^3 x_3^9 x_4^4 x_5^{11} \\ + x_1^2 x_2^3 x_3^9 x_4^8 x_5^7 + x_1^2 x_2^3 x_3^{12} x_4 x_5^{11} + x_1^3 x_2 x_3^6 x_4^8 x_5^{11} + x_1^3 x_2 x_3^7 x_4^8 x_5^{10} \\ + x_1^3 x_2 x_3^{10} x_4^4 x_5^{11} + x_1^3 x_2^2 x_3^5 x_4^8 x_5^{11} + x_1^3 x_2^2 x_3^7 x_4^8 x_5^9 + x_1^3 x_2^2 x_3^9 x_4^4 x_5^{11} \\ + x_1^5 x_2 x_3^7 x_4^2 x_5^{14} + x_1^5 x_2^2 x_3^7 x_4 x_5^{14} + x_1^5 x_2^2 x_3^9 x_4^2 x_5^{11} + x_1^5 x_2^3 x_3^{10} x_4^2 x_5^9 \\ + x_1^5 x_2^3 x_3^{10} x_4^4 x_5^7 + x_1^5 x_2^3 x_3^{12} x_4^2 x_5^7 + x_1^6 x_2 x_3^7 x_4^8 x_5^7,$$

$$u_4 = x_1^3 x_2 x_3^7 x_4^2 x_5^{14} + x_1^3 x_2^2 x_3^7 x_4 x_5^{14} + x_1^3 x_2^2 x_3^9 x_4^2 x_5^{11} + x_1^3 x_2^2 x_3^{10} x_4 x_5^{11} \\ + x_1^3 x_2^3 x_3^{10} x_4^2 x_5^9 + x_1^3 x_2^3 x_3^{10} x_4^4 x_5^7 + x_1^3 x_2^3 x_3^{12} x_4^2 x_5^7 + x_1^3 x_2^8 x_3^5 x_4^4 x_5^7 \\ + x_1^4 x_2 x_3^7 x_4^8 x_5^7 + x_1^{10} x_2 x_3^5 x_4^4 x_5^7,$$

$$u_8 = x_1^3 x_2^4 x_3^5 x_4^4 x_5^7 + x_1^6 x_2 x_3^5 x_4^4 x_5^7.$$

By a similar computation, we have

$$Y_2 = x_1^2 x_2^7 x_3^8 x_4^5 x_5^9 + x_1^2 x_2^{11} x_3 x_4^5 x_5^{12} + x_1^2 x_2^{11} x_3^8 x_4^5 x_5^5 + x_1^2 x_2^{13} x_3 x_4^5 x_5^{10} + x_1^2 x_2^{13} x_3 x_4^9 x_5^6 \\ + x_1^2 x_2^{13} x_3^8 x_4^5 x_5^5 + x_1^3 x_2^3 x_3 x_4^{12} x_5^{12} + x_1^3 x_2^3 x_3^4 x_4^9 x_5^{12} + x_1^3 x_2^3 x_3^8 x_4^5 x_5^{12} + x_1^3 x_2^5 x_3^4 x_4^9 x_5^{10} \\ + x_1^3 x_2^5 x_3^4 x_4^{10} x_5^9 + x_1^3 x_2^5 x_3^8 x_4^5 x_5^{10} + x_1^3 x_2^5 x_3^8 x_4^6 x_5^9 + x_1^3 x_2^5 x_3^8 x_4^9 x_5^6 + x_1^3 x_2^5 x_3^8 x_4^{10} x_5^5 \\ + x_1^3 x_2^7 x_3 x_4^8 x_5^{12} + x_1^3 x_2^7 x_3 x_4^{12} x_5^8 + x_1^3 x_2^7 x_3^4 x_4^9 x_5^8 + x_1^3 x_2^7 x_3^8 x_4^4 x_5^9 \\ + S q^1(v_1) + S q^2(v_2) + S q^4(v_4) + S q^8(v_8) \text{ modulo } (\mathcal{P}_5^-(\omega^*)), \text{ where}$$

$$v_1 = x_1^5 x_2^3 x_3 x_4^9 x_5^{12} + x_1^5 x_2^3 x_3^8 x_4^5 x_5^9 + x_1^5 x_2^7 x_3 x_4^5 x_5^{12} \\ + x_1^5 x_2^7 x_3 x_4^9 x_5^8 + x_1^5 x_2^7 x_3^4 x_4^5 x_5^9 + x_1^5 x_2^7 x_3^8 x_4^5 x_5^5,$$

$$v_2 = x_1^2 x_2^7 x_3^8 x_4^3 x_5^9 + x_1^2 x_2^{11} x_3 x_4^5 x_5^{10} + x_1^2 x_2^{11} x_3 x_4^9 x_5^6 + x_1^2 x_2^{11} x_3^8 x_4^3 x_5^5 + x_1^3 x_2^3 x_3 x_4^{10} x_5^{12} \\ + x_1^3 x_2^3 x_3^2 x_4^9 x_5^{12} + x_1^3 x_2^3 x_3^8 x_4^5 x_5^{10} + x_1^3 x_2^3 x_3^8 x_4^6 x_5^9 + x_1^3 x_2^7 x_3 x_4^6 x_5^{12} + x_1^3 x_2^7 x_3 x_4^{10} x_5^8 \\ + x_1^3 x_2^7 x_3^2 x_4^5 x_5^{12} + x_1^3 x_2^7 x_3^2 x_4^9 x_5^8 + x_1^3 x_2^7 x_3^4 x_4^5 x_5^{10} + x_1^3 x_2^7 x_3^4 x_4^6 x_5^9 + x_1^3 x_2^7 x_3^8 x_4^5 x_5^6 \\ + x_1^3 x_2^7 x_3^8 x_4^6 x_5^5 + x_1^6 x_2^2 x_3 x_4^9 x_5^{10} + x_1^6 x_2^3 x_3^8 x_4^3 x_5^9 + x_1^6 x_2^7 x_3 x_4^5 x_5^{10} + x_1^6 x_2^7 x_3 x_4^9 x_5^6 \\ + x_1^6 x_2^7 x_3^4 x_4^3 x_5^9 + x_1^6 x_2^7 x_3^8 x_4^3 x_5^5,$$

$$v_4 = x_1^3 x_2^5 x_3 x_4^6 x_5^{12} + x_1^3 x_2^5 x_3^2 x_4^5 x_5^{12} + x_1^3 x_2^5 x_3^4 x_4^5 x_5^{10} + x_1^3 x_2^5 x_3^4 x_4^6 x_5^9 + x_1^3 x_2^5 x_3^8 x_4^5 x_5^6 \\ + x_1^3 x_2^5 x_3^8 x_4^6 x_5^5 + x_1^3 x_2^{11} x_3^4 x_4^4 x_5^5 + x_1^4 x_2^5 x_3 x_4^5 x_5^{10} + x_1^4 x_2^7 x_3 x_4^9 x_5^6 + x_1^4 x_2^7 x_3^8 x_4^5 x_5^6 \\ + x_1^{10} x_2^5 x_3 x_4^5 x_5^6 + x_1^{10} x_2^5 x_3^4 x_4^3 x_5^5,$$

$$v_8 = x_1^3 x_2^7 x_3^4 x_4^4 x_5^5 + x_1^6 x_2^5 x_3^1 x_4^5 x_5^6 + x_1^6 x_2^5 x_3^4 x_4^3 x_5^5.$$

By Definition 2.2.4, we see that Y_1 and Y_2 are strictly inadmissible. The lemma is proved. $\square$

Lemma 3.2.19. *We have $Q\mathcal{P}_5^+(\bar{\omega}_{(1)}) = \langle [\bar{\Phi}^+(\mathcal{B}_4(\bar{\omega}_{(1)})) \cup \mathcal{B}^+(5, \bar{\omega}_{(1)}) \cup \mathcal{F}]_{\bar{\omega}_{(1)}} \rangle$, where $\bar{\Phi}^+(\mathcal{B}_4(\bar{\omega}_{(1)})) \cup \mathcal{B}^+(5, \bar{\omega}_{(1)}) \cup \mathcal{F} = \{\mathcal{Y}_{47,j} : 1 \leq j \leq 370\}$ with the monomials $\mathcal{Y}_{47,j} : 1 \leq j \leq 370$, are listed in Sect. 6.7.*

The proof of the lemma is based on Lemmas 3.2.17, 3.2.18 and some below results. The following lemma is an immediate consequence of a result in Sum [34].

Lemma 3.2.20. *The following monomials are strictly inadmissible:*

- I) $x_i^{15} x_j^2 x_k x_\ell^{13}, x_i^3 x_j^{12} x_k x_\ell^{15}, x_i^3 x_j^{14} x_k x_\ell^{13}, x_i^7 x_j^{10} x_k x_\ell^{13}, x_i^3 x_j^{13} x_k^8 x_\ell^7, x_i^3 x_j^{12} x_k^9 x_\ell^7, 1 \leq j < k \leq 5, 1 \leq i, \ell \leq 5, i \neq \ell, i, \ell \neq j, k;$
- II) $x_i^{15} x_j^{14} x_k x_\ell, x_i^3 x_j^2 x_k^{13} x_\ell^{13}, x_i^{15} x_j^2 x_k^5 x_\ell^9, x_i^{13} x_j^2 x_k^7 x_\ell^9, x_i^{13} x_j^7 x_k^2 x_\ell^9, x_i^{15} x_j^3 x_k^4 x_\ell^9, x_i^{15} x_j^3 x_k^5 x_\ell^8, x_i^{14} x_j^3 x_k^5 x_\ell^9, x_i^{13} x_j^3 x_k^6 x_\ell^9, x_i^{13} x_j^6 x_k^3 x_\ell^9, 1 \leq j < k < \ell \leq 5, 1 \leq i \leq 5, i \neq j, k, \ell;$
- III) $\rho_{(1,5)}(x_1^2 x_2^5 x_3^{11} x_4^{13}), \rho_{(1,5)}(x_1^6 x_2^{11} x_3^5 x_4^9), \rho_{(1,5)}(x_1^6 x_2 x_3^{11} x_4^{13}), \rho_{(1,5)}(x_1^6 x_2^{11} x_3 x_4^{13}), \rho_{(1,5)}(x_1^6 x_2^{11} x_3^4 x_4^{13});$

Lemma 3.2.21. *The following monomials are strictly inadmissible:*

$$\begin{aligned}
Z_1 &= x_1^3 x_2^3 x_3^{12} x_4^5 x_5^{24}, & Z_2 &= x_1^7 x_2 x_3^{11} x_4^{20} x_5^8, & Z_3 &= x_1^7 x_2 x_3^2 x_4^{28} x_5^9, & Z_4 &= x_1 x_2^2 x_3^7 x_4^{12} x_5^{25}, \\
Z_5 &= x_1 x_2^7 x_3^2 x_4^{12} x_5^{25}, & Z_6 &= x_1^7 x_2 x_3^2 x_4^{12} x_5^{25}, & Z_7 &= x_1 x_2^6 x_3^3 x_4^8 x_5^{29}, & Z_8 &= x_1 x_2^6 x_3^3 x_4^{24} x_5^{13}, \\
Z_9 &= x_1^3 x_2^7 x_3^8 x_4^5 x_5^{28}, & Z_{10} &= x_1^7 x_2^3 x_3^8 x_4^5 x_5^{28}, & Z_{11} &= x_1^3 x_2^7 x_3^{24} x_4^5 x_5^{12}, & Z_{12} &= x_1^7 x_2^3 x_3^{24} x_4^5 x_5^{12}, \\
Z_{13} &= x_1 x_2^7 x_3^2 x_4^4 x_5^9, & Z_{14} &= x_1^7 x_2 x_3^2 x_4^4 x_5^9, & Z_{15} &= x_1 x_2^7 x_3^{10} x_4^4 x_5^{25}, & Z_{16} &= x_1^7 x_2 x_3^{10} x_4^4 x_5^{25}, \\
Z_{17} &= x_1 x_2^7 x_3^2 x_4^5 x_5^8, & Z_{18} &= x_1^7 x_2 x_3^2 x_4^5 x_5^8, & Z_{19} &= x_1 x_2^7 x_3^{10} x_4^5 x_5^{24}, & Z_{20} &= x_1^7 x_2 x_3^{10} x_4^5 x_5^{24}, \\
Z_{21} &= x_1 x_2^6 x_3^{11} x_4^{13} x_5^{16}, & Z_{22} &= x_1 x_2^7 x_3^{10} x_4^{21} x_5^8, & Z_{23} &= x_1^7 x_2 x_3^{10} x_4^{21} x_5^8, & Z_{24} &= x_1 x_2^7 x_3^{11} x_4^{20} x_5^8, \\
Z_{25} &= x_1 x_2^7 x_3^2 x_4^4 x_5^9, & Z_{26} &= x_1^7 x_2^{11} x_3 x_4^{20} x_5^8, & Z_{27} &= x_1 x_2^7 x_3^{10} x_4^{20} x_5^9, & Z_{28} &= x_1^7 x_2 x_3^{10} x_4^{20} x_5^9, \\
Z_{29} &= x_1 x_2^7 x_3^{10} x_4^{13} x_5^{16}, & Z_{30} &= x_1^7 x_2 x_3^{10} x_4^{13} x_5^{16}, & Z_{31} &= x_1 x_2^7 x_3^{11} x_4^{12} x_5^{16}, & Z_{32} &= x_1^7 x_2 x_3^{11} x_4^{12} x_5^{16}, \\
Z_{33} &= x_1^7 x_2^{11} x_3 x_4^{12} x_5^{16}, & Z_{34} &= x_1^3 x_2^3 x_3^4 x_4^{28} x_5^9, & Z_{35} &= x_1^3 x_2^3 x_3^2 x_4^8 x_5^9, & Z_{36} &= x_1^3 x_2^3 x_3^4 x_4^{12} x_5^{25}, \\
Z_{37} &= x_1^3 x_2^3 x_3^{12} x_4^5 x_5^{25}, & Z_{38} &= x_1^3 x_2^3 x_3^{28} x_4^5 x_5^8, & Z_{39} &= x_1 x_2^7 x_3^2 x_4^{28} x_5^9, & Z_{40} &= x_1^3 x_2^3 x_3^{12} x_4^{21} x_5^8.
\end{aligned}$$

Proof. We have $\omega(Z_u) = \bar{\omega}_{(1)}, \forall u, 1 \leq u \leq 40$. Consider the monomials $Z_1 = x_1^3 x_2^3 x_3^{12} x_4^5 x_5^{24}$ and $Z_2 = x_1^7 x_2 x_3^{11} x_4^{20} x_5^8$. We prove that these monomials are strictly inadmissible. The others can be proved by a similar technique. Computing the monomials Z_1, Z_2 is long and technical. Indeed, by using Cartan's formula, we get

$$Z_1 = \sum X + Sq^1(\sum \sigma_1) + Sq^2(\sum \sigma_2) + Sq^4(\sum \sigma_4) + Sq^8(\sum \sigma_8) \text{ modulo } (\mathcal{P}_5^-(\bar{\omega}_{(1)})), \text{ where}$$

$$\begin{aligned}
\sum X &= x_1^2 x_2 x_3^3 x_4^{12} x_5^{29} + x_1^2 x_2 x_3^3 x_4^{13} x_5^{28} + x_1^2 x_2 x_3^5 x_4^{12} x_5^{27} + x_1^2 x_2 x_3^5 x_4^{13} x_5^{26} + x_1^2 x_2 x_3^5 x_4^{24} x_5^{15} \\
&+ x_1^2 x_2 x_3^5 x_4^{25} x_5^{14} + x_1^2 x_2 x_3^7 x_4^5 x_5^{28} + x_1^2 x_2 x_3^7 x_4^9 x_5^{24} + x_1^2 x_2 x_3^{12} x_4^5 x_5^{29} + x_1^2 x_2 x_3^{12} x_4^9 x_5^{27} \\
&+ x_1^2 x_2 x_3^{13} x_4^5 x_5^{24} + x_1^2 x_2 x_3^{13} x_4^9 x_5^{22} + x_1^2 x_2 x_3^{13} x_4^{16} x_5^{15} + x_1^2 x_2 x_3^{13} x_4^{17} x_5^{14} + x_1^2 x_2 x_3^{16} x_4^{13} x_5^{15} \\
&+ x_1^2 x_2 x_3^{24} x_4^5 x_5^{15} + x_1^2 x_2^3 x_3^5 x_4^9 x_5^{28} + x_1^2 x_2^3 x_3^5 x_4^{12} x_5^{25} + x_1^2 x_2^3 x_3^9 x_4^5 x_5^{28} + x_1^2 x_2^3 x_3^{12} x_4^5 x_5^{25} \\
&+ x_1^2 x_2^5 x_3^3 x_4^9 x_5^{28} + x_1^2 x_2^5 x_3^3 x_4^{12} x_5^{25} + x_1^2 x_2^5 x_3^9 x_4^5 x_5^{26} + x_1^2 x_2^5 x_3^{12} x_4^5 x_5^{25} + x_1^3 x_2 x_3^3 x_4^{12} x_5^{28} \\
&+ x_1^3 x_2 x_3^4 x_4^{12} x_5^{27} + x_1^3 x_2 x_3^4 x_4^{13} x_5^{26} + x_1^3 x_2 x_3^4 x_4^{24} x_5^{15} + x_1^3 x_2 x_3^4 x_4^{25} x_5^{14} + x_1^3 x_2 x_3^7 x_4^{12} x_5^{24} \\
&+ x_1^3 x_2 x_3^7 x_4^{12} x_5^{24} + x_1^3 x_2 x_3^9 x_4^{12} x_5^{22} + x_1^3 x_2 x_3^{12} x_4^5 x_5^{28} + x_1^3 x_2 x_3^{12} x_4^9 x_5^{27} + x_1^3 x_2 x_3^{12} x_4^{17} x_5^{24} \\
&+ x_1^3 x_2 x_3^{12} x_4^{18} x_5^{22} + x_1^3 x_2 x_3^{12} x_4^{16} x_5^{15} + x_1^3 x_2 x_3^{12} x_4^{17} x_5^{14} + x_1^3 x_2 x_3^{16} x_4^{12} x_5^{15} + x_1^3 x_2 x_3^{24} x_4^5 x_5^{15} \\
&+ x_1^3 x_2^2 x_3^3 x_4^9 x_5^{28} + x_1^3 x_2^2 x_3^3 x_4^{12} x_5^{25} + x_1^3 x_2^2 x_3^9 x_4^{12} x_5^{21} + x_1^3 x_2^2 x_3^{12} x_4^5 x_5^{25} + x_1^3 x_2^2 x_3^{12} x_4^9 x_5^{21} \\
&+ x_1^3 x_2^2 x_3^4 x_4^9 x_5^{28} + x_1^3 x_2^2 x_3^4 x_4^{12} x_5^{25} + x_1^3 x_2^2 x_3^5 x_4^8 x_5^{28} + x_1^3 x_2^2 x_3^5 x_4^{12} x_5^{24}
\end{aligned}$$

$$\begin{aligned}
\sum \sigma_1 &= x_1^3 x_2 x_3^3 x_4^{12} x_5^{27} + x_1^3 x_2 x_3^3 x_4^{13} x_5^{26} + x_1^3 x_2 x_3^3 x_4^{16} x_5^{23} + x_1^3 x_2 x_3^3 x_4^{17} x_5^{22} + x_1^3 x_2 x_3^3 x_4^{20} x_5^{19} \\
&+ x_1^3 x_2 x_3^3 x_4^{21} x_5^{18} + x_1^3 x_2 x_3^3 x_4^{24} x_5^{15} + x_1^3 x_2 x_3^3 x_4^{25} x_5^{14} + x_1^3 x_2 x_3^7 x_4^9 x_5^{26} + x_1^3 x_2 x_3^7 x_4^{11} x_5^{24} \\
&+ x_1^3 x_2 x_3^7 x_4^{16} x_5^{19} + x_1^3 x_2 x_3^7 x_4^{17} x_5^{18} + x_1^3 x_2 x_3^9 x_4^{13} x_5^{20} + x_1^3 x_2 x_3^{11} x_4^5 x_5^{24} + x_1^3 x_2 x_3^{11} x_4^9 x_5^{22} \\
&+ x_1^3 x_2 x_3^{11} x_4^{16} x_5^{15} + x_1^3 x_2 x_3^{11} x_4^{17} x_5^{14} + x_1^3 x_2 x_3^{12} x_4^5 x_5^{27} + x_1^3 x_2 x_3^{16} x_4^5 x_5^{23} + x_1^3 x_2 x_3^{16} x_4^7 x_5^{19} \\
&+ x_1^3 x_2 x_3^{16} x_4^{11} x_5^{15} + x_1^3 x_2 x_3^{20} x_4^5 x_5^{19} + x_1^3 x_2 x_3^{24} x_4^5 x_5^{15} + x_1^3 x_2^3 x_3^3 x_4^5 x_5^{32} + x_1^3 x_2^3 x_3^3 x_4^9 x_5^{28} \\
&+ x_1^3 x_2^3 x_3^3 x_4^{12} x_5^{25} + x_1^3 x_2^3 x_3^3 x_4^{16} x_5^{21} + x_1^3 x_2^3 x_3^9 x_4^5 x_5^{26} + x_1^3 x_2^3 x_3^9 x_4^9 x_5^{22} + x_1^3 x_2^3 x_3^{12} x_4^5 x_5^{25} \\
&+ x_1^3 x_2^3 x_3^{16} x_4^5 x_5^{21} + x_1^3 x_2^3 x_3^9 x_4^9 x_5^{21},
\end{aligned}$$

$$\begin{aligned}
\sum \sigma_2 &= x_1^2 x_2 x_3^3 x_4^{12} x_5^{27} + x_1^2 x_2 x_3^3 x_4^{13} x_5^{26} + x_1^2 x_2 x_3^3 x_4^{24} x_5^{15} + x_1^2 x_2 x_3^3 x_4^{25} x_5^{14} + x_1^2 x_2 x_3^7 x_4^9 x_5^{26} \\
&+ x_1^2 x_2 x_3^7 x_4^{11} x_5^{24} + x_1^2 x_2 x_3^{11} x_4^5 x_5^{22} + x_1^2 x_2 x_3^{11} x_4^{16} x_5^{15} + x_1^2 x_2 x_3^{11} x_4^{17} x_5^{14} \\
&+ x_1^2 x_2 x_3^{12} x_4^5 x_5^{27} + x_1^2 x_2 x_3^{16} x_4^{11} x_5^{15} + x_1^2 x_2 x_3^{24} x_4^5 x_5^{15} + x_1^2 x_2^3 x_3^3 x_4^9 x_5^{28} + x_1^2 x_2^3 x_3^3 x_4^{12} x_5^{25} \\
&+ x_1^2 x_2^3 x_3^9 x_4^5 x_5^{26} + x_1^2 x_2^3 x_3^9 x_4^9 x_5^{22} + x_1^2 x_2^3 x_3^{12} x_4^5 x_5^{25} + x_1^5 x_2 x_3^3 x_4^{14} x_5^{22} + x_1^5 x_2 x_3^3 x_4^{22} x_5^{14} \\
&+ x_1^5 x_2 x_3^7 x_4^{10} x_5^{22} + x_1^5 x_2 x_3^7 x_4^{18} x_5^{14} + x_1^5 x_2^2 x_3^3 x_4^9 x_5^{26} + x_1^5 x_2^2 x_3^3 x_4^{11} x_5^{24} + x_1^5 x_2^2 x_3^3 x_4^{12} x_5^{23} \\
&+ x_1^5 x_2^2 x_3^3 x_4^{13} x_5^{22} + x_1^5 x_2^2 x_3^3 x_4^{20} x_5^{15} + x_1^5 x_2^2 x_3^3 x_4^{21} x_5^{14} + x_1^5 x_2^2 x_3^7 x_4^5 x_5^{24} + x_1^5 x_2^2 x_3^7 x_4^9 x_5^{22} \\
&+ x_1^5 x_2^2 x_3^7 x_4^{16} x_5^{15} + x_1^5 x_2^2 x_3^7 x_4^{17} x_5^{14} + x_1^5 x_2^2 x_3^9 x_4^5 x_5^{19} + x_1^5 x_2^2 x_3^9 x_4^{11} x_5^{18} + x_1^5 x_2^2 x_3^{10} x_4^9 x_5^{19} \\
&+ x_1^5 x_2^2 x_3^{12} x_4^5 x_5^{23} + x_1^5 x_2^2 x_3^{16} x_4^5 x_5^{15} + x_1^5 x_2^2 x_3^{20} x_4^5 x_5^{15} + x_1^5 x_2^2 x_3^3 x_4^6 x_5^{28} + x_1^5 x_2^2 x_3^3 x_4^8 x_5^{26} \\
&+ x_1^5 x_2^2 x_3^3 x_4^{10} x_5^{24} + x_1^5 x_2^2 x_3^3 x_4^{12} x_5^{22} + x_1^5 x_2^2 x_3^5 x_4^6 x_5^{26} + x_1^5 x_2^2 x_3^5 x_4^{10} x_5^{22} + x_1^5 x_2^2 x_3^{10} x_4^3 x_5^{24} \\
&+ x_1^5 x_2^2 x_3^{10} x_4^5 x_5^{22} + x_1^5 x_2^2 x_3^{12} x_4^5 x_5^{22},
\end{aligned}$$

$$\sum \sigma_8 = x_1^3 x_2^4 x_3^5 x_4^5 x_5^{22} + x_1^3 x_2^4 x_3^5 x_4^7 x_5^{20} + x_1^3 x_2^4 x_3^5 x_4^{12} x_5^{15} + x_1^3 x_2^4 x_3^5 x_4^{13} x_5^{14} + x_1^3 x_2^4 x_3^{12} x_4^5 x_5^{15},$$

Let $X \in \mathcal{B}_5^+(\omega_{(1)})$ such that $x_t x_k x_m X^2 \in (P_5^+)_{47}$. By a direct computation using Proposition 3.2.10, we see that if $x_t x_k x_m X^2 \neq \mathcal{Y}_{47,j}$, $1 \leq j \leq 370$, then there is a monomial b which is given in one of Lemmas 3.2.17 - 3.2.22 such that $x_t x_k x_m X^2 = bY^{2^r}$ with suitable monomial $Y \in \mathcal{P}_5$, and $r = \max\{s \in \mathbb{Z} : \omega_s(b) > 0\}$. By Theorem 2.2.5, $x_t x_k x_m X^2$ is inadmissible. On the other hand, we have $a = x_t x_k x_m u^2$ and a is admissible, hence $a = \mathcal{Y}_j$ for some $j = 1, 2, \dots, 370$. This completes the proof of the lemma. $\square$

Remark 3.2.23. For $1 \leq j \leq 370$, we have $[\mathcal{Y}_j := \mathcal{Y}_{47,j}]_{\overline{\omega}_{(1)}} \neq [0]$. Indeed, suppose that there is a linear relation

$$\mathcal{S} = \sum_{1 \leq j \leq 170} \gamma_j \mathcal{Y}_j \equiv_{\overline{\omega}_{(1)}} 0,$$

where $\gamma_j \in \mathbb{Z}/2$, for all j . Based on Theorem 2.3.3 and Proposition 3.1.1, for $(k; \mathcal{K}) \in \mathcal{N}_5$, we explicitly compute $\pi_{(k; \mathcal{K})}(\mathcal{S})$ in terms of a given minimal set of $\mathcal{A}_2$ -generators in $\mathcal{P}_4$ (modulo $(\mathcal{A}_2^+ \mathcal{P}_4)$). By computing from the relations $\pi_{(k; \mathcal{K})}(\mathcal{S}) \equiv_{\overline{\omega}_{(1)}} 0$, $\ell(\mathcal{K}) = 1$, one gets $\gamma_j = 0$, $\forall j$, $1 \leq j \leq 370$. Note that these computations are similar to the proof of Propositions 3.2.1 and 3.2.10. Combining this and Proposition 3.2.24, we obtain

Proposition 3.2.24. *The set $[\overline{\Phi}^+(\mathcal{B}_4(\overline{\omega}_{(1)})) \cup \mathcal{B}^+(5, \overline{\omega}_{(1)}) \cup \mathcal{F}]_{\overline{\omega}_{(1)}}$ is a basis of $\mathbb{Z}/2$ -vector space $Q\mathcal{P}_5^+(\overline{\omega}_{(1)})$. Consequently, $\dim(Q\mathcal{P}_5^+(\overline{\omega}_{(1)})) = 370$.*

Remark 3.2.25. Consider the weight vector $\omega^{**} = (3, 2, 4)$ with $\deg \omega^{**} = 23$. By using a result in [41], we see that the following monomials are strictly inadmissible:

$$\begin{aligned} e_1 &= x_1^7 x_2^2 x_3^4 x_4^5 x_5^5, & e_2 &= x_1^7 x_2^2 x_3^5 x_4^4 x_5^5, & e_3 &= x_1^7 x_2^2 x_3^5 x_4^5 x_5^4, & e_4 &= x_1^3 x_2^7 x_3^4 x_4^4 x_5^5, \\ e_5 &= x_1^3 x_2^7 x_3^4 x_4^5 x_5^4, & e_6 &= x_1^3 x_2^7 x_3^5 x_4^4 x_5^4, & e_7 &= x_1^7 x_2^3 x_3^4 x_4^4 x_5^5, & e_8 &= x_1^7 x_2^3 x_3^4 x_4^5 x_5^4, \\ e_9 &= x_1^7 x_2^3 x_3^5 x_4^4 x_5^4, & e_{10} &= x_1^3 x_2^4 x_3^4 x_4^5 x_5^7, & e_{11} &= x_1^3 x_2^4 x_3^4 x_4^7 x_5^5, & e_{12} &= x_1^3 x_2^4 x_3^5 x_4^4 x_5^7, \\ e_{13} &= x_1^3 x_2^4 x_3^4 x_4^5 x_5^5, & e_{14} &= x_1^3 x_2^4 x_3^5 x_4^4 x_5^4, & e_{15} &= x_1^3 x_2^4 x_3^7 x_4^5 x_5^4, & e_{16} &= x_1^3 x_2^5 x_3^4 x_4^4 x_5^7, \\ e_{17} &= x_1^3 x_2^5 x_3^4 x_4^5 x_5^4, & e_{18} &= x_1^3 x_2^5 x_3^7 x_4^4 x_5^4, & e_{19} &= x_1^3 x_2^6 x_3^4 x_4^5 x_5^5, & e_{20} &= x_1^3 x_2^6 x_3^5 x_4^4 x_5^5, \\ e_{21} &= x_1^3 x_2^6 x_3^4 x_4^5 x_5^4, & e_{22} &= x_1^3 x_2^4 x_3^6 x_4^5 x_5^5, & e_{23} &= x_1^3 x_2^4 x_3^5 x_4^6 x_5^5, & e_{24} &= x_1^3 x_2^4 x_3^5 x_4^5 x_5^6, \\ e_{25} &= x_1^3 x_2^5 x_3^4 x_4^6 x_5^5, & e_{26} &= x_1^3 x_2^5 x_3^6 x_4^4 x_5^5, & e_{27} &= x_1^3 x_2^5 x_3^4 x_4^5 x_5^6, & e_{28} &= x_1^3 x_2^5 x_3^6 x_4^5 x_5^4, \\ e_{29} &= x_1^3 x_2^5 x_3^5 x_4^4 x_5^6, & e_{30} &= x_1^3 x_2^5 x_3^5 x_4^6 x_5^4, \end{aligned}$$

Note that $\omega(e_k) = \omega^{**}$, $k = 1, 2, \dots, 30$. Let $X \in \mathcal{B}_5^+(47)$ such that either $\omega(X) = \overline{\omega}_{(2)}$ or $\omega(X) = \overline{\omega}_{(3)}$. Then $X = X_{(\{\ell, m\}, 5)} y^2$ with $1 \leq \ell < m \leq 5$, and $y \in (\mathcal{P}_5^+)_{22}$. By a direct computation using Theorem 2.2.5, and Proposition 3.2.14(I), (II), we see that either $y \in \mathcal{B}_5^+(\omega_{(2)})$ or $y \in \mathcal{B}_5^+(\omega_{(3)})$ and there is a monomial $Z = e_i$ for some i , $1 \leq i \leq 30$ such that $X = X_{(\{\ell, m\}, 5)} y^2 = Zu^{2^t}$, $1 \leq \ell < m \leq 5$, with suitable monomial $u \in \mathcal{P}_5$, and $t = \max\{a \in \mathbb{Z} : \omega_a(Z) > 0\}$. By Theorem 2.2.5, we get either $[X] = [0]_{\overline{\omega}_{(2)}}$ or $[X] = [0]_{\overline{\omega}_{(3)}}$. As a consequence, we get the following proposition.

Proposition 3.2.26. $Q\mathcal{P}_5^+(\overline{\omega}_{(2)}) = Q\mathcal{B}_5^+(\overline{\omega}_{(3)}) = 0$.

Lemma 3.2.27. *The set $[\{\mathcal{Y}_{47,j} : 371 \leq j \leq 482\}]_{\overline{\omega}_{(4)}}$ is a $\mathbb{Z}/2$ -generator of $Q\mathcal{P}_5^+(\overline{\omega}_{(4)})$, where the monomials $\mathcal{Y}_{47,j} : 371 \leq j \leq 482$, are determined in Sect. 6.7.*

In order to prove the lemma, we need to use some results. As an immediate consequence of a result in [34], we obtain the following.

Lemma 3.2.28. *The following monomials are strictly inadmissible:*

$$\begin{aligned} I) & x_a^7 x_b^2 x_c^{15} x_d^7, x_a^7 x_b^{15} x_c^2 x_d^7, x_a^7 x_b^6 x_c^3 x_d^{15}, x_a^7 x_b^3 x_c^6 x_d^{15}, x_a^7 x_b^3 x_c^{14} x_d^7, \\ & x_a^7 x_b^{14} x_c^3 x_d^7, x_a^7 x_b^6 x_c^{11} x_d^7, x_a^7 x_b^{11} x_c^6 x_d^7, \quad 1 \leq b < c \leq 5, \quad 1 \leq a, d \leq 5, \quad a \neq d, \quad a, d \neq b, c; \end{aligned}$$

II) $\rho_{(k, 5)}(v)$, $1 \leq k \leq 5$, where v is one of the following monomials:

$$x_1^7 x_2^7 x_3^7 x_4^{10}, x_1^7 x_2^7 x_3^{10} x_4^7, x_1^7 x_2^{10} x_3^7 x_4^7.$$

We now consider the weight vector $\omega^{***} = (3, 4, 3, 1)$ with $\deg \omega^{***} = 31$. The following lemma can be easily proved by a direct computation.

Lemma 3.2.29. *All permutations of the following monomials are strictly inadmissible:*

$$\begin{aligned} & x_1 x_2 x_3^6 x_4^7 x_5^{15}, x_1 x_2 x_3^4 x_4^7 x_5^{15}, x_1 x_2 x_3^6 x_4^7 x_5^{10}, x_1 x_2 x_3^2 x_4^5 x_5^{15}, x_1 x_2 x_3^2 x_4^7 x_5^{13}, x_1 x_2 x_3^4 x_4^7 x_5^{15}, \\ & x_1^3 x_2^2 x_3^7 x_4^5^{12}, x_1^3 x_2^2 x_3^4 x_4^5^{11}, x_1^3 x_2^2 x_3^5 x_4^5^{10}, x_1^3 x_2^2 x_3^6 x_4^5^9, x_1^3 x_2^2 x_3^7 x_4^5^8, x_1^3 x_2^2 x_3^4 x_4^5^{15}, \\ & x_1^3 x_2^3 x_3^4 x_4^7 x_5^{14}, x_1^3 x_2^3 x_3^6 x_4^7 x_5^{12}, x_1^3 x_2^3 x_3^4 x_4^5^{11}, x_1^3 x_2^3 x_3^6 x_4^5^{10}, x_1^3 x_2^3 x_3^7 x_4^5^8. \end{aligned}$$

Lemma 3.2.30. *If (m, n, p, q, r) is a permutation of $(1, 2, 3, 4, 5)$, then the following monomials are strictly inadmissible:*

$$\begin{aligned} x_m^3 x_n^{13} x_p^6 x_q^6 x_r^6 & \neq X = x_1^3 x_2^3 x_3^{13} x_4^6 x_5^6, x_1^3 x_2^{13} x_3^3 x_4^6 x_5^6, \\ x_m^7 x_n^2 x_p^5 x_q^6 x_r^{11} & \neq Y = x_1^7 x_2^{11} x_3^5 x_4^6 x_5^2, x_1^7 x_2^{11} x_3^6 x_4^5 x_5^2, \\ x_m^3 x_n^5 x_p^6 x_q^6 x_r^{11} & \neq Z = x_1^3 x_2^5 x_3^6 x_4^6 x_5^{11}, x_1^3 x_2^5 x_3^{11} x_4^6 x_5^6, \\ x_m^3 x_n^7 x_p^9 x_q^6 x_r^6 & \neq G = x_1^3 x_2^7 x_3^9 x_4^6 x_5^6, x_1^3 x_2^9 x_3^7 x_4^6 x_5^6, x_1^3 x_2^9 x_3^6 x_4^6 x_5^6, \\ x_m^3 x_n^2 x_p^6 x_q^7 x_r^{13} & \neq H = x_1^3 x_2^7 x_3^{13} x_4^2 x_5^6, x_1^3 x_2^7 x_3^{13} x_4^6 x_5^2, x_1^3 x_2^7 x_3^{13} x_4^2 x_5^6, x_1^3 x_2^7 x_3^{13} x_4^6 x_5^2, \\ x_m^3 x_n^5 x_p^6 x_q^7 x_r^{10} & \neq F = x_1^3 x_2^5 x_3^6 x_4^7 x_5^{10}, x_1^3 x_2^5 x_3^7 x_4^6 x_5^{10}, x_1^3 x_2^5 x_3^7 x_4^{10} x_5^6, x_1^3 x_2^5 x_3^5 x_4^6 x_5^{10}, \\ & x_1^3 x_2^7 x_3^5 x_4^{10} x_5^6, x_1^3 x_2^7 x_3^5 x_4^6 x_5^{10}, x_1^3 x_2^7 x_3^5 x_4^{10} x_5^6, \\ x_m^3 x_n^5 x_p^{14} x_q^3 x_r^6 & \neq T = x_1^3 x_2^5 x_3^5 x_4^{14} x_5^6, x_1^3 x_2^5 x_3^5 x_4^{14} x_5^6, x_1^3 x_2^5 x_3^5 x_4^{14} x_5^6, \\ & x_1^3 x_2^5 x_3^3 x_4^{14} x_5^6, x_1^3 x_2^5 x_3^6 x_4^{14} x_5^3. \end{aligned}$$

Proof. It is easy to see that the weight vector of these monomials is ω^{***} . Note that the monomials X, Y, Z, G, H, F, T are admissible. We now prove the lemma for the monomials $f = x_m^3 x_n^{13} x_p^6 x_q^6 x_r^6$ and $g = x_m^3 x_n^5 x_p^{14} x_q^3 x_r^6$. The others can be proved by an argument similar to the proofs of Lemmas 3.2.17 and 3.2.18. Applying the Cartan formula, we get

$$f = x_m^3 x_n^{13} x_p^6 x_q^6 x_r^6 + Sq^1(\sum \bar{A}_1) + Sq^2(\sum \bar{B}_1) + Sq^4(\sum \bar{C}_1) \text{ modulo } (\mathcal{P}_5^-(\omega^{***})), \text{ where}$$

$$\begin{aligned} \sum \bar{A}_1 &= x_m^3 x_n^7 x_p^5 x_q^5 x_r^{10} + x_m^3 x_n^7 x_p^5 x_q^6 x_r^9 + x_m^3 x_n^7 x_p^6 x_q^5 x_r^9 + x_m^3 x_n^7 x_p^6 x_q^9 x_r^5 + x_m^3 x_n^7 x_p^9 x_q^6 x_r^5 \\ &\quad + x_m^3 x_n^{11} x_p^5 x_q^6 x_r^5 + x_m^3 x_n^{14} x_p^3 x_q^5 x_r^5 + x_m^6 x_n^{11} x_p^5 x_q^3 x_r^5, \\ \sum \bar{B}_1 &= x_m^3 x_n^{11} x_p^5 x_q^5 x_r^5 + x_m^3 x_n^{11} x_p^6 x_q^3 x_r^6 + x_m^3 x_n^{13} x_p^5 x_q^3 x_r^5 + x_m^5 x_n^7 x_p^3 x_q^5 x_r^9 + x_m^5 x_n^7 x_p^3 x_q^9 x_r^5 \\ &\quad + x_m^5 x_n^7 x_p^3 x_q^9 x_r^5 + x_m^5 x_n^9 x_p^3 x_q^3 x_r^5 + x_m^5 x_n^{11} x_p^3 x_q^3 x_r^5, \\ \sum \bar{C}_1 &= x_m^3 x_n^7 x_p^3 x_q^5 x_r^9 + x_m^3 x_n^7 x_p^3 x_q^9 x_r^5 + x_m^3 x_n^7 x_p^5 x_q^3 x_r^9 + x_m^3 x_n^7 x_p^9 x_q^3 x_r^5 + x_m^3 x_n^9 x_p^3 x_q^3 x_r^9 \\ &\quad + x_m^3 x_n^9 x_p^5 x_q^3 x_r^5 + x_m^3 x_n^{11} x_p^3 x_q^3 x_r^5 + x_m^3 x_n^{11} x_p^5 x_q^3 x_r^5, \end{aligned}$$

Since $x_m^3 x_n^{13} x_p^6 x_q^6 x_r^6 < f$, the monomial f is strictly inadmissible. Next, by a direct computation, we have

$$g = \sum \mathcal{L} + Sq^1(\sum \bar{A}_2) + Sq^2(\sum \bar{B}_2) + Sq^4(\sum \bar{C}_2) \text{ modulo } (\mathcal{P}_5^-(\omega^{***})), \text{ where}$$

$$\begin{aligned} \sum \mathcal{L} &= x_m^3 x_n^3 x_p^{14} x_q^5 x_r^6 + x_m^3 x_n^5 x_p^7 x_q^6 x_r^{10} + x_m^3 x_n^5 x_p^{11} x_q^6 x_r^6, \\ \sum \bar{A}_2 &= x_m^3 x_n^3 x_p^{13} x_q^5 x_r^6 + x_m^3 x_n^3 x_p^{14} x_q^5 x_r^5 + x_m^3 x_n^5 x_p^{11} x_q^5 x_r^6 + x_m^3 x_n^5 x_p^{13} x_q^3 x_r^6 + x_m^3 x_n^6 x_p^7 x_q^9 x_r^5 \\ &\quad + x_m^3 x_n^6 x_p^{11} x_q^5 x_r^5 + x_m^3 x_n^{10} x_p^7 x_q^5 x_r^5 + x_m^6 x_n^5 x_p^{11} x_q^3 x_r^5, \\ \sum \bar{B}_2 &= x_m^3 x_n^5 x_p^{11} x_q^5 x_r^5 + x_m^3 x_n^5 x_p^{13} x_q^3 x_r^5 + x_m^3 x_n^6 x_p^{11} x_q^3 x_r^6 + x_m^5 x_n^3 x_p^7 x_q^5 x_r^9 + x_m^5 x_n^3 x_p^7 x_q^9 x_r^5 \\ &\quad + x_m^5 x_n^3 x_p^9 x_q^3 x_r^9 + x_m^5 x_n^3 x_p^{11} x_q^3 x_r^5 + x_m^5 x_n^9 x_p^3 x_q^3 x_r^5 + x_m^5 x_n^9 x_p^7 x_q^3 x_r^5, \\ \sum \bar{C}_2 &= x_m^3 x_n^3 x_p^7 x_q^5 x_r^9 + x_m^3 x_n^3 x_p^7 x_q^9 x_r^5 + x_m^3 x_n^3 x_p^9 x_q^3 x_r^9 + x_m^3 x_n^3 x_p^{11} x_q^5 x_r^5 + x_m^3 x_n^5 x_p^7 x_q^3 x_r^9 \\ &\quad + x_m^3 x_n^5 x_p^7 x_q^6 x_r^6 + x_m^3 x_n^5 x_p^9 x_q^3 x_r^5 + x_m^3 x_n^5 x_p^{11} x_q^3 x_r^5 + x_m^3 x_n^9 x_p^7 x_q^3 x_r^5. \end{aligned}$$

These relations show that g is strictly inadmissible. The lemmas follows. $\square$

The proof of the following lemmas is analogous to the proofs of Lemmas 3.2.17, 3.2.18, 3.2.29 and 3.2.30.

Lemma 3.2.31. *If (p, q, r) is a permutation of $(3, 4, 5)$, then the following monomials are strictly inadmissible:*

$$\begin{aligned} u_1 &= x_1 x_2^6 x_p^3 x_q^{15}, & u_2 &= x_1^3 x_2^6 x_p^6 x_q^{15}, & u_3 &= x_1^{16} x_2^6 x_p^6 x_q^3, & u_4 &= x_1 x_2^6 x_p^3 x_q^{14}, \\ u_5 &= x_1 x_2^{14} x_p^3 x_q^7, & u_6 &= x_1^3 x_2^6 x_p^6 x_q^{14}, & u_7 &= x_1^3 x_2^{14} x_p^6 x_q^7, & u_8 &= x_1^7 x_2^6 x_p^3 x_q^{14}, \\ u_9 &= x_1^7 x_2^{14} x_p^3 x_q^6, & u_{10} &= x_1 x_2^6 x_p^6 x_q^{11}, & u_{11} &= x_1^7 x_2^6 x_p^6 x_q^{11}, & u_{12} &= x_1^3 x_2^5 x_p^5 x_q^{15}, \\ u_{13} &= x_1^3 x_2^6 x_p^2 x_q^{15}, & u_{14} &= x_1^{15} x_2^2 x_p^3 x_q^6, & u_{15} &= x_1^{15} x_2^6 x_p^2 x_q^5, & u_{16} &= x_1^3 x_2^5 x_p^5 x_q^{14}, \\ u_{17} &= x_1^3 x_2^{14} x_p^2 x_q^7, & u_{18} &= x_1^7 x_2^2 x_p^{14} x_q^5, & u_{19} &= x_1^7 x_2^{14} x_p^2 x_q^5. \end{aligned}$$

Lemma 3.2.32. *The following monomials are strictly inadmissible:*

$$\begin{aligned} I) \quad & u_{20} = x_1 x_2^{15} x_3^6 x_q^6, & u_{21} &= x_1^3 x_2^{15} x_3^6 x_q^6, & u_{22} &= x_1^{15} x_2 x_3^6 x_q^6, & u_{23} &= x_1^{15} x_2^3 x_3^6 x_q^6, \\ & u_{24} = x_1^7 x_2^3 x_3^6 x_q^{14}, & u_{25} &= x_1^7 x_2 x_3^{14} x_q^6, & u_{26} &= x_1^7 x_2^3 x_3^6 x_q^{14}, & u_{27} &= x_1^7 x_2^3 x_3^{14} x_q^6, \\ & u_{28} = x_1 x_2^3 x_3^{14} x_q^7, & u_{29} &= x_1 x_2^7 x_3^6 x_q^{14}, & u_{30} &= x_1 x_2^7 x_3^{14} x_q^6, & u_{31} &= x_1^3 x_2^3 x_3^{14} x_q^7, \\ & u_{32} = x_1^3 x_2^7 x_3^6 x_q^{14}, & u_{33} &= x_1^3 x_2^7 x_3^{14} x_q^6, & u_{34} &= x_1 x_2^7 x_3^6 x_q^{11}, & u_{35} &= x_1^7 x_2^3 x_3^6 x_q^{11}, \\ & u_{36} = x_1^7 x_2^{11} x_3^6 x_q^6, & u_{37} &= x_1^3 x_2^{15} x_3^2 x_q^6, & u_{38} &= x_1^3 x_2^{15} x_3^6 x_q^5, & u_{39} &= x_1^{15} x_2^3 x_3^2 x_q^6, \\ & u_{40} = x_1^{15} x_2^3 x_3^6 x_q^5, & u_{41} &= x_1^3 x_2^5 x_3^{14} x_q^7, & u_{42} &= x_1^3 x_2^5 x_3^{14} x_q^7, & u_{43} &= x_1^3 x_2^7 x_3^2 x_q^{14}, \\ & u_{44} = x_1^3 x_2^7 x_3^{14} x_q^5, & u_{45} &= x_1^7 x_2^3 x_3^2 x_q^{14}, & u_{46} &= x_1^7 x_2^3 x_3^{14} x_q^5, & & \text{where } q, r = 4, 5, \\ & & & & & & q \neq r; \end{aligned}$$

$$II) \quad u_{47} = x_1 x_2^3 x_3^6 x_4^{14} x_5^7, \quad u_{48} = x_1^3 x_2 x_3^6 x_4^{14} x_5^7, \quad u_{49} = x_1^3 x_2^5 x_3^2 x_4^{14} x_5^7.$$

Note that $\omega(u_t) = \omega^{***}$ for $t = 1, 2, \dots, 49$. Next we consider the following lemma.

Lemma 3.2.33. *The following monomials are strictly inadmissible:*

$$\begin{aligned} A_1 &= x_1 x_2^3 x_3^6 x_4^{15} x_5^{22}, & A_2 &= x_1 x_2^3 x_3^{15} x_4^6 x_5^{22}, & A_3 &= x_1 x_2^3 x_3^{15} x_4^{22} x_5^6, & A_4 &= x_1 x_2^{15} x_3^6 x_4^6 x_5^{22}, \\ A_5 &= x_1 x_2^{15} x_3^3 x_4^{22} x_5^6, & A_6 &= x_1 x_2^{15} x_3^{19} x_4^6 x_5^6, & A_7 &= x_1^3 x_2 x_3^6 x_4^{15} x_5^{22}, & A_8 &= x_1^3 x_2^5 x_3^3 x_4^{15} x_5^{22}, \\ A_9 &= x_1^3 x_2^5 x_3^6 x_4^{15} x_5^{18}, & A_{10} &= x_1^3 x_2 x_3^{15} x_4^6 x_5^{22}, & A_{11} &= x_1^3 x_2 x_3^{15} x_4^{22} x_5^6, & A_{12} &= x_1^3 x_2^5 x_3^{15} x_4^2 x_5^{22}, \\ A_{13} &= x_1^3 x_2^5 x_3^{15} x_4^6 x_5^{18}, & A_{14} &= x_1^3 x_2^5 x_3^{15} x_4^{18} x_5^6, & A_{15} &= x_1^3 x_2^5 x_3^{15} x_4^{22} x_5^2, & A_{16} &= x_1^3 x_2^{13} x_3^6 x_4^6 x_5^{22}, \\ A_{17} &= x_1^3 x_2^{13} x_3^3 x_4^{22} x_5^6, & A_{18} &= x_1^3 x_2^{13} x_3^{19} x_4^6 x_5^6, & A_{19} &= x_1^3 x_2^{15} x_3^6 x_4^6 x_5^{22}, & A_{20} &= x_1^3 x_2^{15} x_3^3 x_4^{22} x_5^6, \\ A_{21} &= x_1^3 x_2^{15} x_3^5 x_4^2 x_5^{22}, & A_{22} &= x_1^3 x_2^{15} x_3^5 x_4^{18} x_5^6, & A_{23} &= x_1^3 x_2^{15} x_3^5 x_4^{18} x_5^6, & A_{24} &= x_1^3 x_2^{15} x_3^5 x_4^{22} x_5^2, \\ A_{25} &= x_1^3 x_2^{15} x_3^{17} x_4^6 x_5^6, & A_{26} &= x_1^3 x_2^{15} x_3^{21} x_4^6 x_5^6, & A_{27} &= x_1^3 x_2^{15} x_3^{21} x_4^6 x_5^2, & A_{28} &= x_1^7 x_2^9 x_3^3 x_4^6 x_5^{22}, \\ A_{29} &= x_1^7 x_2^9 x_3^3 x_4^6 x_5^6, & A_{30} &= x_1^7 x_2^9 x_3^{19} x_4^6 x_5^6, & A_{31} &= x_1^7 x_2^{11} x_3^{17} x_4^6 x_5^6, & A_{32} &= x_1^7 x_2^{25} x_3^3 x_4^6 x_5^6, \\ A_{33} &= x_1^{15} x_2 x_3^3 x_4^6 x_5^{22}, & A_{34} &= x_1^{15} x_2 x_3^3 x_4^{22} x_5^6, & A_{35} &= x_1^{15} x_2 x_3^{19} x_4^6 x_5^6, & A_{36} &= x_1^{15} x_2^3 x_3^6 x_4^6 x_5^{22}, \\ A_{37} &= x_1^{15} x_2^3 x_3^3 x_4^{22} x_5^6, & A_{38} &= x_1^{15} x_2^3 x_3^5 x_4^{22} x_5^2, & A_{39} &= x_1^{15} x_2^3 x_3^5 x_4^{18} x_5^6, & A_{40} &= x_1^{15} x_2^3 x_3^5 x_4^{18} x_5^6, \\ A_{41} &= x_1^{15} x_2^3 x_3^5 x_4^{22} x_5^2, & A_{42} &= x_1^{15} x_2^3 x_3^{17} x_4^6 x_5^6, & A_{43} &= x_1^{15} x_2^3 x_3^{21} x_4^6 x_5^6, & A_{44} &= x_1^{15} x_2^3 x_3^{21} x_4^6 x_5^2, \\ A_{45} &= x_1^{15} x_2^{17} x_3^6 x_4^6 x_5^6, & A_{46} &= x_1^{15} x_2^{19} x_3^6 x_4^6 x_5^6, & A_{47} &= x_1^{15} x_2^{19} x_3^5 x_4^6 x_5^6, & A_{48} &= x_1^{15} x_2^{19} x_3^5 x_4^6 x_5^2. \end{aligned}$$

Proof. We have $\omega(A_i) = \overline{\omega}_{(4)}$, $1 \leq i \leq 48$. We prove the lemma for the monomials $A_{23} = x_1^3 x_2^{15} x_3^5 x_4^{18} x_5^6$ and $A_{29} = x_1^7 x_2^9 x_3^3 x_4^{22} x_5^6$. The others can be proved by using a similar technique as in Lemmas 3.2.21 and 3.2.22. Direct computing from Cartan's formula, we get

$$\begin{aligned} A_{23} &= \sum_{1 \leq i \leq 3} b_i + Sq^1(\sum f) + Sq^2(\sum g) + Sq^4(\sum h) + Sq^8(\sum p) \text{ modulo } (\mathcal{P}_5^-(\overline{\omega}_{(4)})), \text{ where} \\ b_1 &= x_1^3 x_2^{11} x_3^5 x_4^{22} x_5^6, \quad b_2 = x_1^3 x_2^{13} x_3^3 x_4^{22} x_5^6, \quad b_3 = x_1^3 x_2^{13} x_3^6 x_4^{19} x_5^6, \\ \sum f &= x_1^3 x_2^7 x_3^6 x_4^{21} x_5^9 + x_1^3 x_2^7 x_3^6 x_4^{25} x_5^5 + x_1^3 x_2^{11} x_3^6 x_4^{21} x_5^5 + x_1^3 x_2^{14} x_3^3 x_4^{21} x_5^5 + x_1^3 x_2^{15} x_3^5 x_4^{14} x_5^9 \\ &\quad + x_1^3 x_2^{15} x_3^5 x_4^{17} x_5^6 + x_1^3 x_2^{15} x_3^9 x_4^{14} x_5^5 + x_1^3 x_2^{19} x_3^5 x_4^{14} x_5^5 + x_1^3 x_2^{22} x_3^5 x_4^{11} x_5^5, \\ \sum g &= x_1^5 x_2^7 x_3^3 x_4^{21} x_5^9 + x_1^5 x_2^7 x_3^3 x_4^{25} x_5^5 + x_1^5 x_2^9 x_3^9 x_4^{19} x_5^3 + x_1^5 x_2^{11} x_3^3 x_4^{17} x_5^9 + x_1^5 x_2^{15} x_3^5 x_4^{11} x_5^9 \\ &\quad + x_1^5 x_2^{15} x_3^9 x_4^{11} x_5^5 + x_1^5 x_2^{17} x_3^3 x_4^{17} x_5^3 + x_1^5 x_2^{17} x_3^9 x_4^{11} x_5^5 + x_1^5 x_2^{19} x_3^5 x_4^{11} x_5^5, \\ \sum h &= x_1^3 x_2^7 x_3^3 x_4^{21} x_5^9 + x_1^3 x_2^7 x_3^3 x_4^{25} x_5^5 + x_1^3 x_2^9 x_3^5 x_4^{21} x_5^5 + x_1^3 x_2^9 x_3^9 x_4^{19} x_5^3 + x_1^3 x_2^{11} x_3^3 x_4^{17} x_5^9 \\ &\quad + x_1^3 x_2^{11} x_3^3 x_4^{21} x_5^5 + x_1^3 x_2^{13} x_3^9 x_4^{13} x_5^5 + x_1^3 x_2^{15} x_3^5 x_4^{11} x_5^9 + x_1^3 x_2^{15} x_3^6 x_4^{13} x_5^6 + x_1^3 x_2^{15} x_3^9 x_4^{11} x_5^5 \\ &\quad + x_1^3 x_2^{17} x_3^3 x_4^{17} x_5^3 + x_1^3 x_2^{17} x_3^9 x_4^{11} x_5^3 + x_1^3 x_2^{19} x_3^5 x_4^{11} x_5^5, \\ \sum p &= x_1^3 x_2^{13} x_3^5 x_4^{13} x_5^5 + x_1^3 x_2^{13} x_3^6 x_4^{11} x_5^6. \end{aligned}$$

Since $b_i < A_1$, $1 \leq i \leq 3$, by Definition 2.2.4, A_1 is strictly inadmissible. By a similar computation, we obtain

$$\begin{aligned}
A_{29} &= \sum Z + Sq^1(\sum \bar{f}) + Sq^2(\sum \bar{g}) + Sq^4(\sum \bar{h}) + Sq^8(x_1^7 x_2^5 x_3^6 x_4^{15} x_5^6) \text{ modulo } (\mathcal{P}_5^-(\bar{\omega}_{(4)})), \text{ where,} \\
\sum Z &= x_1^5 x_2^3 x_3^6 x_4^{23} x_5^{10} + x_1^5 x_2^3 x_3^6 x_4^{27} x_5^6 + x_1^5 x_2^3 x_3^{10} x_4^{23} x_5^6 + x_1^5 x_2^7 x_3^3 x_4^{22} x_5^{10} + x_1^5 x_2^7 x_3^3 x_4^{26} x_5^6 \\
&\quad + x_1^5 x_2^7 x_3^6 x_4^{19} x_5^{10} + x_1^5 x_2^7 x_3^{10} x_4^{19} x_5^6 + x_1^5 x_2^{11} x_3^3 x_4^{22} x_5^6 + x_1^5 x_2^{11} x_3^6 x_4^{19} x_5^6 + x_1^7 x_2^3 x_3^5 x_4^{22} x_5^{10} \\
&\quad + x_1^7 x_2^3 x_3^6 x_4^{23} x_5^8 + x_1^7 x_2^3 x_3^6 x_4^{25} x_5^6 + x_1^7 x_2^3 x_3^8 x_4^{23} x_5^6 + x_1^7 x_2^3 x_3^9 x_4^{22} x_5^6 + x_1^7 x_2^7 x_3^3 x_4^{22} x_5^8 \\
&\quad + x_1^7 x_2^7 x_3^3 x_4^{24} x_5^6, \\
\sum \bar{f} &= x_1^7 x_2^3 x_3^5 x_4^{19} x_5^{12} + x_1^7 x_2^3 x_3^5 x_4^{21} x_5^{10} + x_1^7 x_2^3 x_3^9 x_4^{21} x_5^6 + x_1^7 x_2^3 x_3^{12} x_4^{19} x_5^5 + x_1^7 x_2^6 x_3^9 x_4^{15} x_5^9 \\
&\quad + x_1^7 x_2^7 x_3^3 x_4^{19} x_5^8 + x_1^7 x_2^7 x_3^5 x_4^{21} x_5^6 + x_1^7 x_2^7 x_3^8 x_4^{19} x_5^5 + x_1^7 x_2^{10} x_3^5 x_4^{15} x_5^9 \\
&\quad + x_1^7 x_2^{10} x_3^9 x_4^{15} x_5^5, \\
\sum \bar{g} &= x_1^7 x_2^3 x_3^6 x_4^{19} x_5^{10} + x_1^7 x_2^3 x_3^6 x_4^{23} x_5^6 + x_1^7 x_2^3 x_3^9 x_4^{17} x_5^9 + x_1^7 x_2^3 x_3^{10} x_4^{19} x_5^6 + x_1^7 x_2^5 x_3^9 x_4^{15} x_5^9 \\
&\quad + x_1^7 x_2^7 x_3^3 x_4^{22} x_5^6 + x_1^7 x_2^7 x_3^6 x_4^{19} x_5^6 + x_1^7 x_2^9 x_3^3 x_4^{17} x_5^9 + x_1^7 x_2^9 x_3^5 x_4^{15} x_5^9 + x_1^7 x_2^9 x_3^9 x_4^{15} x_5^5 \\
&\quad + x_1^7 x_2^9 x_3^{17} x_4^{15} x_5^3 + x_1^9 x_2^3 x_3^9 x_4^{15} x_5^9 + x_1^9 x_2^3 x_3^{15} x_4^{15} x_5^9 + x_1^9 x_2^9 x_3^9 x_4^{15} x_5^3, \\
\sum \bar{h} &= x_1^5 x_2^3 x_3^6 x_4^{23} x_5^6 + x_1^5 x_2^7 x_3^3 x_4^{22} x_5^6 + x_1^5 x_2^7 x_3^6 x_4^{19} x_5^6.
\end{aligned}$$

The above equalities imply that A_2 is strictly inadmissible. The lemma is proved. $\square$

Proof of Lemma 3.2.27. Let b be an admissible monomial in $(\mathcal{P}_5^+)_{47}$ such that $\omega(b) = \bar{\omega}_{(4)}$. Then $\omega_1(b) = 3$ and $b = X_{(\{t,k\},5)} Y^2$ with $1 \leq t < k \leq 5$ and Y a monomial of degree 22 in $\mathcal{P}_5$. Since b is admissible, according to Theorem 2.2.5, $Y \in \mathcal{B}_5(\omega_{(4)})$.

Using Proposition 3.2.14(III) and a simple computation shows that if $Z \in \mathcal{B}_5(\omega_{(4)})$, $1 \leq t < k \leq 5$, and $X_{(\{t,k\},5)} Z^2 \neq \mathcal{Y}_j := \mathcal{Y}_{47,j}$, $\forall j$, $371 \leq j \leq 482$, then there is a monomial u which is given in one of Lemmas 3.2.28 - 3.2.33 such that $X_{(\{t,k\},5)} Z^2 = u g^{2^s}$ with a monomial $g \in \mathcal{P}_5$, and $s = \max\{\ell \in \mathbb{Z} : \omega_\ell(u) > 0\}$. Then, By Theorem 2.2.5, $X_{(\{t,k\},5)} Z^2$ is inadmissible. Finally, we see that $b = X_{(\{t,k\},5)} Y^2$ is admissible with $Y \in \mathcal{B}_5(\omega_{(4)})$; hence $b = \mathcal{Y}_j$ for some $j = 371, \dots, 482$. This implies $\mathcal{B}_5^+(\bar{\omega}_{(4)}) \subseteq \{\mathcal{Y}_{47,j} : 371 \leq j \leq 482\}$. $\square$

Proposition 3.2.34. $Q\mathcal{P}_5^+(\bar{\omega}_{(4)})$ is the $\mathbb{Z}/2$ -vector space of dimension 112 with a basis consisting of all the classes represented by the monomials $Y_{47,j}$, $371 \leq j \leq 482$.

Proof. First, we show that the set $[V := \{Y_{47,j} : 371 \leq j \leq 482\}]_{\bar{\omega}_{(4)}}$ is linearly independent in the space $Q\mathcal{P}_5^+(\bar{\omega}_{(4)})$. Indeed, suppose there is a linear relation $\mathcal{S} = \sum_{371 \leq j \leq 482} \gamma_j \mathcal{Y}_j \equiv_{\bar{\omega}_{(4)}} 0$ with $\gamma_j \in \mathbb{Z}/2$ and $\mathcal{Y}_j \in V$. By using Theorem 2.3.3 and Proposition 3.1.1, we determine explicitly $\pi_{(k;\mathcal{K})}(\mathcal{S})$ in terms of the admissible monomials in $(\mathcal{P}_4^+)_{47}$. From the relations $\pi_{(k;\mathcal{K})}(\mathcal{S}) \equiv_{\bar{\omega}_{(4)}} 0$ with $\ell(\mathcal{K}) > 0$, one gets $\gamma_j = 0$ for all $371 \leq j \leq 482$.

Now, by Lemma 3.2.27, to prove $[V]_{\bar{\omega}_{(4)}}$ is a basis of $Q\mathcal{P}_5^+(\bar{\omega}_{(4)})$ we need to show that $[\mathcal{Y}_j]_{\bar{\omega}_{(4)}} \neq [0]$ for all $\mathcal{Y}_j \in V$. By a similar argument as given in the proof of Propositions 3.2.1 and 3.2.10, we can prove that the set $[\mathcal{B}_5^+(\bar{\omega}_{(1)}) \cup V]$ is linearly independent in $(Q\mathcal{P}_5^+)_{47}$. This fact shows that $[\mathcal{Y}_j]_{\bar{\omega}_{(4)}} \neq [0]$ for all $\mathcal{Y}_j$. The proposition is proved. $\square$

Proposition 3.2.35. There exist exactly 15 admissible monomials in $(\mathcal{P}_5^+)_{47}$ such that their weight vectors are $\bar{\omega}_{(5)}$. Consequently $\dim(Q\mathcal{P}_5^+(\bar{\omega}_{(5)})) = 15$.

We prove the proposition by showing that

$$\mathcal{B}_5^+(\bar{\omega}_{(5)}) = \{\mathcal{Y}_{47,m} : 483 \leq m \leq 497\},$$

where the monomials $\mathcal{Y}_{47,m} : 283 \leq m \leq 297$, are described in Sect.6.7. We need some lemmas.

The following result is proved by using a result in [41].

Lemma 3.2.36. *The following monomials are strictly inadmissible:*

- a) $x_u x_v^2 x_m^6 x_n^7 x_p^7$, $x_u^7 x_v^2 x_m^2 x_n^5 x_p^7$, $x_u^3 x_v^2 x_m^4 x_n^7 x_p^7$, $x_u^3 x_v^4 x_m^6 x_n^3 x_p^7$, $x_u^3 x_v^6 x_m^6 x_n^3 x_p^5$,
where (u, v, m, n, p) is a permutation of $(1, 2, 3, 4, 5)$;
- b) $x_1 x_2^6 x_q^6 x_r^3 x_t^7$, $x_1^3 x_2^6 x_q^6 x_r x_t^7$, $x_1^7 x_2^6 x_q^6 x_r x_t^3$, $x_1^3 x_2^2 x_q^6 x_r^5 x_t^7$, $x_1^3 x_2^6 x_q^2 x_r^5 x_t^7$, $x_1^7 x_2^2 x_q^6 x_r^3 x_t^5$,
 $x_1^7 x_2^6 x_q^2 x_r^3 x_t^5$, where (q, r, t) is a permutation of $(3, 4, 5)$;
- c) $x_1 x_2^7 x_3^6 x_4^6 x_5^3$, $x_1 x_2^7 x_3^6 x_4^3 x_5^6$, $x_1^3 x_2^7 x_3^6 x_4^6 x_5$, $x_1^3 x_2^7 x_3^6 x_4 x_5^6$,
 $x_1^7 x_2^6 x_3^6 x_4^6 x_5^3$, $x_1^7 x_2^6 x_3^6 x_4^3 x_5^6$, $x_1^7 x_2^6 x_3^6 x_4^6 x_5$, $x_1^7 x_2^6 x_3^6 x_4 x_5^6$,
 $x_1^3 x_2^7 x_3^6 x_4^6 x_5^3$, $x_1^3 x_2^7 x_3^6 x_4^3 x_5^6$, $x_1^3 x_2^7 x_3^6 x_4^6 x_5$, $x_1^3 x_2^7 x_3^6 x_4 x_5^6$,
 $x_1^7 x_2^6 x_3^6 x_4^6 x_5^3$, $x_1^7 x_2^6 x_3^6 x_4^3 x_5^6$, $x_1^7 x_2^6 x_3^6 x_4^6 x_5$, $x_1^7 x_2^6 x_3^6 x_4 x_5^6$;
- d) $\rho_{(k,5)}(Z)$, $1 \leq k \leq 5$, where Z is one of the following monomials:
 $x_2^2 x_7^7 x_3^7 x_4^7$, $x_1^7 x_2^7 x_3^7 x_4^7$, $x_1^7 x_2^7 x_3^2 x_4^7$, $x_1^7 x_2^7 x_3^7 x_4^2$, $x_1^6 x_2^3 x_3^7 x_4^7$, $x_1^6 x_2^7 x_3^3 x_4^7$,
 $x_1^6 x_2^7 x_3^7 x_4^3$, $x_1^3 x_2^6 x_3^7 x_4^7$, $x_1^7 x_2^6 x_3^3 x_4^7$, $x_1^7 x_2^6 x_3^7 x_4^3$, $x_1^3 x_2^7 x_3^6 x_4^7$, $x_1^7 x_2^3 x_3^6 x_4^7$,
 $x_1^7 x_2^7 x_3^6 x_4^3$, $x_1^3 x_2^7 x_3^6 x_4^7$, $x_1^7 x_2^3 x_3^6 x_4^7$, $x_1^7 x_2^7 x_3^6 x_4^3$.

Lemma 3.2.37. *The following monomials are strictly inadmissible:*

$$\begin{aligned}
T_1 &= x_1 x_2^3 x_3^{14} x_4^{14} x_5^{15}, & T_2 &= x_1 x_2^3 x_3^{14} x_4^{15} x_5^{14}, & T_3 &= x_1 x_2^3 x_3^{15} x_4^{14} x_5^{14}, & T_4 &= x_1 x_2^{15} x_3^3 x_4^{14} x_5^{14}, \\
T_5 &= x_1^3 x_2 x_3^{14} x_4^{14} x_5^{15}, & T_6 &= x_1^3 x_2 x_3^{14} x_4^{15} x_5^{14}, & T_7 &= x_1^3 x_2 x_3^{15} x_4^{14} x_5^{14}, & T_8 &= x_1^3 x_2^{15} x_3 x_4^{14} x_5^{14}, \\
T_9 &= x_1^{15} x_2 x_3^3 x_4^{14} x_5^{14}, & T_{10} &= x_1^{15} x_2^3 x_3 x_4^{14} x_5^{14}, & T_{11} &= x_1^3 x_2^{13} x_3^2 x_4^{14} x_5^{15}, & T_{12} &= x_1^3 x_2^{13} x_3^2 x_4^{15} x_5^{14}, \\
T_{13} &= x_1^3 x_2^{13} x_3^{14} x_4^2 x_5^{15}, & T_{14} &= x_1^3 x_2^{13} x_3^{14} x_4^{15} x_5^2, & T_{15} &= x_1^3 x_2^{13} x_3^{15} x_4^2 x_5^{14}, & T_{16} &= x_1^3 x_2^{13} x_3^{15} x_4^{14} x_5^2, \\
T_{17} &= x_1^3 x_2^{15} x_3^{13} x_4^2 x_5^{14}, & T_{18} &= x_1^3 x_2^{15} x_3^{13} x_4^4 x_5^2, & T_{19} &= x_1^{15} x_2^3 x_3^{13} x_4^2 x_5^{14}, & T_{20} &= x_1^{15} x_2^3 x_3^{13} x_4^{14} x_5^2, \\
T_{21} &= x_1^3 x_2^3 x_3^{13} x_4^4 x_5^{14}, & T_{22} &= x_1^3 x_2^{13} x_3^3 x_4^{14} x_5^{14}, & T_{23} &= x_1^3 x_2^{13} x_3^{14} x_4^3 x_5^{14}, & T_{24} &= x_1^3 x_2^{13} x_3^{14} x_4^{14} x_5^3, \\
T_{25} &= x_1^3 x_2^3 x_3^{10} x_4^{14} x_5^{15}, & T_{26} &= x_1^3 x_2^3 x_3^{10} x_4^{15} x_5^{14}, & T_{27} &= x_1^3 x_2^5 x_3^{14} x_4^{10} x_5^{15}, & T_{28} &= x_1^3 x_2^5 x_3^{14} x_4^{15} x_5^{10}, \\
T_{29} &= x_1^3 x_2^5 x_3^{15} x_4^{10} x_5^{14}, & T_{30} &= x_1^3 x_2^5 x_3^{15} x_4^{14} x_5^{10}, & T_{31} &= x_1^3 x_2^{15} x_3^5 x_4^{10} x_5^{14}, & T_{32} &= x_1^3 x_2^{15} x_3^5 x_4^{14} x_5^{10}, \\
T_{33} &= x_1^{15} x_2^3 x_3^5 x_4^{10} x_5^{14}, & T_{34} &= x_1^{15} x_2^3 x_3^5 x_4^{14} x_5^{10}, & T_{35} &= x_1^3 x_2^5 x_3^{14} x_4^{11} x_5^{14}, & T_{36} &= x_1^3 x_2^5 x_3^{14} x_4^{14} x_5^{11}, \\
T_{37} &= x_1^3 x_2^{13} x_3^6 x_4^{10} x_5^{15}, & T_{38} &= x_1^3 x_2^{13} x_3^6 x_4^{15} x_5^{10}, & T_{39} &= x_1^3 x_2^{13} x_3^{15} x_4^6 x_5^{10}, & T_{40} &= x_1^3 x_2^{15} x_3^{13} x_4^6 x_5^{10}, \\
T_{41} &= x_1^{15} x_2^3 x_3^{13} x_4^6 x_5^{10}, & T_{42} &= x_1^3 x_2^{13} x_3^6 x_4^{11} x_5^{14}, & T_{43} &= x_1^3 x_2^{13} x_3^6 x_4^{14} x_5^{11}, & T_{44} &= x_1^3 x_2^{13} x_3^{14} x_4^6 x_5^{11}, \\
T_{45} &= x_1^7 x_2^9 x_3^3 x_4^{14} x_5^{14}, & T_{46} &= x_1^3 x_2^{13} x_3^7 x_4^{10} x_5^{14}, & T_{47} &= x_1^3 x_2^{13} x_3^7 x_4^{14} x_5^{10}, & T_{48} &= x_1^3 x_2^{13} x_3^{14} x_4^7 x_5^{10}.
\end{aligned}$$

Proof. It is easy to see that $\omega(T_s) = \overline{\omega}_{(5)}$ for $s = 1, 2, \dots, 48$. We prove the lemma for the monomials $T_{35} = x_1^3 x_2^5 x_3^{14} x_4^{11} x_5^{14}$ and $T_{42} = x_1^3 x_2^{13} x_3^6 x_4^{11} x_5^{14}$. The others can be proved by using a similar technique as in Lemmas 3.2.21, 3.2.22 and 3.2.33. Computing from the Cartan formula, we get

$$\begin{aligned}
T_{35} &= Sq^1 \left(x_1^3 x_2^6 x_3^{11} x_4^{13} x_5^{13} + x_1^3 x_2^6 x_3^{13} x_4^{11} x_5^{13} \right) + Sq^2 \left(x_1^5 x_2^3 x_3^7 x_4^{13} x_5^{17} + x_1^5 x_2^3 x_3^7 x_4^{17} x_5^{13} \right. \\
&\quad + x_1^5 x_2^3 x_3^9 x_4^{11} x_5^{17} + x_1^5 x_2^3 x_3^9 x_4^{17} x_5^{11} + x_1^5 x_2^3 x_3^{11} x_4^9 x_5^{17} + x_1^5 x_2^3 x_3^{11} x_4^{13} x_5^{13} \\
&\quad \left. + x_1^5 x_2^3 x_3^{13} x_4^7 x_5^{17} + x_1^5 x_2^3 x_3^{13} x_4^{11} x_5^{13} + x_1^5 x_2^3 x_3^{17} x_4^7 x_5^{13} + x_1^5 x_2^3 x_3^{17} x_4^9 x_5^{11} \right) \\
&\quad + Sq^4 \left(x_1^3 x_2^3 x_3^7 x_4^{13} x_5^{17} + x_1^3 x_2^3 x_3^7 x_4^{17} x_5^{13} + x_1^3 x_2^3 x_3^9 x_4^{11} x_5^{17} + x_1^3 x_2^3 x_3^9 x_4^{17} x_5^{11} \right. \\
&\quad + x_1^3 x_2^3 x_3^{11} x_4^9 x_5^{17} + x_1^3 x_2^3 x_3^{11} x_4^{13} x_5^{13} + x_1^3 x_2^3 x_3^{13} x_4^7 x_5^{17} + x_1^3 x_2^3 x_3^{13} x_4^{11} x_5^{13} \\
&\quad \left. + x_1^3 x_2^3 x_3^{17} x_4^7 x_5^{13} + x_1^3 x_2^3 x_3^{17} x_4^9 x_5^{11} + x_1^3 x_2^5 x_3^{11} x_4^{14} x_5^{14} \right) \text{ modulo } (\mathcal{P}_5^-(\overline{\omega}_{(5)})).
\end{aligned}$$

We have $x_1^3 x_2^5 x_3^{11} x_4^{14} x_5^{14} < T_{35}$. This implies that T_{35} is strictly inadmissible. By a similar computation, we obtain

$$T_{42} = (p_1 + p_2) \text{ modulo } (\mathcal{A}_2^+ \mathcal{P}_5 + \mathcal{P}_5^-(\overline{\omega}_{(5)})),$$

where $p_1 = x_1^3 x_2^7 x_3^{10} x_4^{13} x_5^{14}$, $p_2 = x_1^3 x_2^{13} x_3^3 x_4^{14} x_5^{14}$. Since $p_i < T_{42}$, $i = 1, 2$, the monomial T_{42} is strictly inadmissible. The lemma follows. $\square$

Proof of Proposition 3.2.35. Suppose that X is a admissible monomial in $\mathcal{P}_5^+$ and $\omega(X) = \bar{\omega}_{(5)}$. Then $\omega_1(X) = 3$ and $X = x_a x_b x_c u^2$ with $1 \leq a < b < c \leq 5$ and $u \in (P_5^+)_{22}$. Then, by Theorem 2.2.5, u is admissible and $u \in \mathcal{B}_5^+(\omega_{(5)})$.

Let $Z \in B_5^+(\omega_{(5)})$ such that $x_a x_b x_c Z^2 \in (P_5^+)_{47}$ with $1 \leq a < b < c \leq 5$. Denote by $\bar{V}$ the set of all the monomials as given in Proposition 3.2.35. By a direct computation using Proposition 3.2.14(IV), we see that if $x_a x_b x_c Z^2 \notin \bar{V}$ then there is a monomial X_1 which is given in Lemmas 3.2.36 and 3.2.37 such that $x_a x_b x_c Z^2 = X_1 Z_1^{2^t}$ with suitable monomial $Z_1 \in \mathcal{P}_5$ and $t = \max\{q \in \mathbb{Z} : \omega_q(X_1) > 0\}$. Based on Theorem 2.2.5, we conclude that $x_a x_b x_c Z^2$ is inadmissible. Combining this with the above data, one gets $X \in \bar{V}$. This means $\mathcal{B}_5^+(\bar{\omega}_{(5)}) \subseteq \bar{V}$.

Next, we show that the set $[\bar{V}]_{\bar{\omega}_{(5)}}$ is linearly independent in the space $Q\mathcal{P}_5^+(\bar{\omega}_{(5)})$. Indeed, suppose there is a linear relation $\mathcal{S} = \sum_{483 \leq m \leq 497} \gamma_m \mathcal{Y}_{47,m} \equiv_{\bar{\omega}_{(5)}} 0$, where with $\gamma_m \in \mathbb{Z}/2$, $m = 483, \dots, 497$ and $\mathcal{Y}_{47,m} \in \bar{V}$. By combining Theorem 2.3.3 and Proposition 3.1.1, we explicitly calculate $\pi_{(k;\mathcal{K})}(\mathcal{S})$ in terms of a given minimal set of $\mathcal{A}_2$ -generators in $\mathcal{P}_4$ (modulo $(\mathcal{A}_2^+ \mathcal{P}_4)$). From the relations $\pi_{(k;\mathcal{K})}(\mathcal{S}) \equiv_{\bar{\omega}_{(5)}} 0$ with $\ell(\mathcal{K}) \leq 2$, we get $\gamma_m = 0$ for all m .

To prove $[\bar{V}]_{\bar{\omega}_{(5)}}$ is a basis of $Q\mathcal{P}_5^+(\bar{\omega}_{(5)})$ we need to show that $[\mathcal{Y}_{47,m}]_{\bar{\omega}_{(5)}} \neq [0]$ for all $\mathcal{Y}_m \in \bar{V}$. Denote by $\bar{V}$ the set of all the monomials as given in Proposition 3.2.35. By a similar argument as given in the proof of Propositions 3.2.1 and 3.2.10, we can prove that the set $[\mathcal{B}_5^+(\bar{\omega}_{(1)}) \cup V \cup \bar{V}]$ is linearly independent in $(Q\mathcal{P}_5^+)_{47}$, where V the set of all the admissible monomials as given in Proposition 3.2.34. This implies $[\mathcal{Y}_{47,m}]_{\bar{\omega}_{(5)}} \neq [0]$ for all $\mathcal{Y}_{47,m}$. The proposition is proved. $\square$

Now, since $\dim(Q\mathcal{P}_5)_{47} = 560$ and $\dim(Q\mathcal{P}_5^0)_{21} = 460$, by Corollary 3.2.15 and Propositions 3.2.24, 3.2.26, 3.2.34, 3.2.35, we obtain $\dim(Q\mathcal{P}_5)_{47} = 1897$. The proof of Theorem 1.1 is completed.

Final remark. Since the homomorphism $(Sq_*^0)_{(5,13,2^t)} : (Q\mathcal{P}_5)_{13,2^t-5} \longrightarrow (Q\mathcal{P}_5)_{13,2^t-1-5}$ is an epimorphism of $\mathbb{Z}/2GL_5$ -modules, $(Q\mathcal{P}_5)_{13,2^t-5} \cong \text{Ker}(\widehat{Sq_*^0})_{(5,13,2^t-5)} \oplus (Q\mathcal{P}_5)_{13,2^t-1-5}$.

According to Lemma 3.2.2, we have

$$\text{Ker}(\widehat{Sq_*^0})_{(5,13,2^1-5=21)} = Q\mathcal{P}_5(3, 3, 1, 1) \bigoplus Q\mathcal{P}_5(3, 3, 3).$$

By a simple computation, we conclude $\mathcal{B}_5^0(21) = \bar{\Phi}(\mathcal{B}_4(21)) = \mathcal{B}_5^0(3, 3, 1, 1) \cup \mathcal{B}_5^0(3, 3, 3)$. Combining this with Propositions 3.2.6 and 3.2.7, we get

$$\bar{\Phi}(\mathcal{B}_4(3, 3, 1, 1)) \subset \mathcal{B}_5(3, 3, 1, 1), \quad \bar{\Phi}(\mathcal{B}_4(3, 3, 3)) \subset \mathcal{B}_5(3, 3, 3).$$

On the other hand, $(Q\mathcal{P}_5)_{13,2^t-5} \cong (Q\mathcal{P}_5)_{13,2^2-5=47}$ for any $t > 2$. By Lemma 3.2.16, $\text{Ker}(\widehat{Sq_*^0})_{(5,47)} = \bigoplus_{1 \leq j \leq 5} Q\mathcal{P}_5(\bar{\omega}_{(j)})$. From the above computations, we get $\mathcal{B}_5^0(\bar{\omega}_{(1)}) = \mathcal{B}_5^0(47) = \bar{\Phi}(\mathcal{B}_4(47)) = \bar{\Phi}(\mathcal{B}_4(\bar{\omega}_{(1)}))$. Then, by Propositions 3.2.24, 3.2.26, 3.2.34 and 3.2.35, one gets

$$\bar{\Phi}(\mathcal{B}_4(\bar{\omega}_{(1)})) \subset \mathcal{B}_5(\bar{\omega}_{(1)}).$$

If ω is a weight vector of degree 47 and $\omega \neq \bar{\omega}_{(1)}$, then $\mathcal{B}_4(\omega) = \emptyset$. Furthermore, if $t > 2$, then $\mathcal{B}_4(13, 2^t - 5) = \emptyset$.

From these remarks, we see that Conjecture 3.1.4 is true in case of five variables and the generic degree $n = 13, 2^t - 5$ with t an arbitrary positive integer.

4. An application of Theorem 1.1

The goal of this section is to prove Theorem 1.2. More precisely, by using our results in Sect.3 and a result in [38], we describe the $\mathbb{Z}/2GL_d$ -modules structure of $(Q\mathcal{P}_d)_n$ for $d = 5$ and $n = 8, 21$. Then, we explicitly determine all GL_5 -invariants of these spaces.

Before coming to the proof of the theorem, we introduce some notations and homomorphisms. We note that $(\mathbb{Z}/2)^{\times d}$ regarded as a $\mathbb{Z}/2$ -vector space of dimension d and

$$(\mathbb{Z}/2)^{\times d} \cong \langle x_1, \dots, x_d \rangle \subset \mathcal{P}_d.$$

For $1 \leq t \leq d$, define the $\mathbb{Z}/2$ -linear map $\tau_t : (\mathbb{Z}/2)^{\times d} \rightarrow (\mathbb{Z}/2)^{\times d}$, which is determined by $\tau_t(x_t) = x_{t+1}$, $\tau_t(x_{t+1}) = x_t$, $\tau_t(x_m) = x_m$ for $m \neq t, t+1$, $1 \leq t \leq d-1$ and $\tau_d(x_1) = x_1 + x_2$, $\tau_d(x_m) = x_m$ for $m > 1$.

Denote by S_d the symmetric group of degree d . Then, S_d is generated by τ_t , $1 \leq t \leq d-1$. For each permutation in S_d , consider corresponding permutation matrix; these form a group of matrices isomorphic to S_d . So, the general linear group $GL_d = GL(d, \mathbb{Z}/2) \cong GL((\mathbb{Z}/2)^{\times d})$ is generated by S_d and τ_d . Let $X = x_1^{a_1} x_2^{a_2} \dots x_d^{a_d}$ be a monomial in $\mathcal{P}_d$. Then, the weight vector $\omega(X)$ is invariant under the permutation of the generators x_j , $j = 1, 2, \dots, d$; hence $Q\mathcal{P}_d(\omega)$ also has a S_d -module structure. The homomorphism τ_t induces a homomorphism of $\mathcal{A}_2$ -algebras which is also denoted by $\tau_t : \mathcal{P}_d \rightarrow \mathcal{P}_d$. Hence, a class $[u]_\omega \in Q\mathcal{P}_d(\omega)$ is an GL_d -invariant if and only if $\tau_t(u) + u = 0$ modulo $(\mathcal{A}_2^+ \mathcal{P}_5 + \mathcal{P}_5^-(\omega))$ for $1 \leq t \leq d$. If $\tau_t(u) + u = 0$ modulo $(\mathcal{A}_2^+ \mathcal{P}_5 + \mathcal{P}_5^-(\omega))$ for $1 \leq t \leq d-1$, then $[u]_\omega$ is an S_d -invariant.

Let ω be a weight vector of degree n and the monomials $\mathcal{Y}_1, \mathcal{Y}_2, \dots, \mathcal{Y}_s \in \mathcal{P}_d(\omega)$, with $s \geq 1$. For a subgroup $L \subseteq GL_d$, we denote

$$\begin{aligned} L(\mathcal{Y}_1, \mathcal{Y}_2, \dots, \mathcal{Y}_s) &= \{\sigma(\mathcal{Y}_j) : \sigma \in L, 1 \leq j \leq s\} \subset \mathcal{P}_d(\omega), \\ [\mathcal{B}(\mathcal{Y}_1, \mathcal{Y}_2, \dots, \mathcal{Y}_s)]_\omega &= [\mathcal{B}_d(\omega)]_\omega \cap \langle [L(\mathcal{Y}_1, \mathcal{Y}_2, \dots, \mathcal{Y}_s)]_\omega \rangle, \\ \theta(\mathcal{Y}_j) &= \sum_{x \in \mathcal{B}_d(n) \cap L(\mathcal{Y}_j)} x, \end{aligned}$$

We note that $\langle [L(\mathcal{Y}_1, \mathcal{Y}_2, \dots, \mathcal{Y}_s)]_\omega \rangle$ is the L -submodule of $Q\mathcal{P}_d(\omega)$ generated by the set

$$\{[\mathcal{Y}_1]_\omega, [\mathcal{Y}_2]_\omega, \dots, [\mathcal{Y}_s]_\omega\}.$$

Now, recall that the squaring operation $(\widetilde{Sq}_*^0)_{(5,21)} : (Q\mathcal{P}_5)_{21} \rightarrow (Q\mathcal{P}_5)_8$ is an epimorphism of GL_5 -modules. So, to prove Theorem 1.2, we need to compute all GL_5 -invariants of $(Q\mathcal{P}_5)_8$.

4.1. Computation of $(Q\mathcal{P}_5)_8^{GL_5}$

According to T  n [38], $\dim(Q\mathcal{P}_5)_8 = 174$ with the basis $\{[\mathcal{Y}_{8,i}] : 1 \leq i \leq 174\}$, where the monomials $\mathcal{Y}_i := \mathcal{Y}_{8,i}$, $1 \leq i \leq 174$, as given in Sect.6.1.

Proposition 4.1.1. $(Q\mathcal{P}_5)_8^{GL_5} = 0$.

We prepare some lemmas for the proof of the proposition. We have

$$(Q\mathcal{P}_5)_8 = Q\mathcal{P}_5^0(\tilde{\omega}_{(1)}) \oplus Q\mathcal{P}_5(\tilde{\omega}_{(2)}) \oplus Q\mathcal{P}_5(\tilde{\omega}_{(3)}),$$

where $\tilde{\omega}_{(1)} = (2, 1, 1)$, $\tilde{\omega}_{(2)} = (2, 3)$, $\tilde{\omega}_{(3)} = (4, 2)$. We see that $\dim Q\mathcal{P}_5^0(\tilde{\omega}_{(1)}) = 105$ with the basis $\bigcup_{1 \leq i \leq 6} [\mathcal{B}(\mathcal{Y}_i)]_{\tilde{\omega}_{(1)}}$, where

$$\begin{aligned} \mathcal{Y}_1 &= x_4 x_5^7, \quad \mathcal{Y}_2 = x_4^3 x_5^5, \quad \mathcal{Y}_3 = x_3 x_4 x_5^6, \\ \mathcal{Y}_4 &= x_3 x_4^2 x_5^5, \quad \mathcal{Y}_5 = x_3 x_4^3 x_5^4, \quad \mathcal{Y}_6 = x_2 x_3 x_4^2 x_5^4. \end{aligned}$$

Observe that $\tilde{\omega}_{(1)}$ is the weight vector of the minimal spike $x_1^7 x_2$, so $[X]_{\tilde{\omega}_{(1)}} = [X]$ for all $X \in \mathcal{P}_5$. A direct computation shows that

$$\begin{aligned} [S_5(\mathcal{Y}_1)] &= \langle [\mathcal{Y}_i] : 1 \leq i \leq 20 \rangle, \\ [S_5(\mathcal{Y}_2)] &= \langle [\mathcal{Y}_i] : 21 \leq i \leq 30 \rangle, \\ [S_5(\mathcal{Y}_3, \mathcal{Y}_4, \mathcal{Y}_5)] &= \langle [\mathcal{Y}_i] : 31 \leq i \leq 90 \rangle, \\ [S_5(\mathcal{Y}_6)] &= \langle [\mathcal{Y}_i] : 91 \leq i \leq 105 \rangle \end{aligned}$$

are S_5 -submodules of $Q\mathcal{P}_5^0(\tilde{\omega}_{(1)})$. Hence, we have a direct summand decomposition of S_5 -modules: $QP_5^0(\tilde{\omega}_{(1)}) = [S_5(\mathcal{Y}_1)] \oplus [S_5(\mathcal{Y}_2)] \oplus [S_5(\mathcal{Y}_3, \mathcal{Y}_4, \mathcal{Y}_5)] \oplus [S_5(\mathcal{Y}_6)]$.

Lemma 4.1.2. $\dim(Q\mathcal{P}_5^0(\tilde{\omega}_{(1)}))^{S_5} = 4$.

Proof. We prove the following:

$$\begin{aligned} [S_5(\mathcal{Y}_j)]^{S_5} &= \langle [\theta(\mathcal{Y}_j)] \rangle, \quad j = 1, 2, 6 \\ [S_5(\mathcal{Y}_3, \mathcal{Y}_4, \mathcal{Y}_5)]^{S_5} &= \langle [q := \mathcal{Y}_{31} + \mathcal{Y}_{32} + \cdots + \mathcal{Y}_{70}] \rangle. \end{aligned}$$

Indeed, we compute $[S_5(\mathcal{Y}_j)]^{S_5}$ for $j = 2$ and $[S_5(\mathcal{Y}_3, \mathcal{Y}_4, \mathcal{Y}_5)]^{S_5}$. The others can be proved by a similar technique. Note that $\dim[S_5(\mathcal{Y}_2)] = 10$ with a basis consisting of all the classes represented by the monomials $\mathcal{Y}_i : 21 \leq i \leq 30$. Suppose that $a = \sum_{21 \leq i \leq 30} \gamma_i \mathcal{Y}_i$ with $\gamma_i \in \mathbb{Z}/2$ and $[a] \in [S_5(\mathcal{Y}_2)]^{S_5}$. By a direct computation using Theorem 2.3.3, we have

$$\begin{aligned} \tau_1(a) + a &= \sum X \text{ modulo } (\mathcal{A}_2^+ \mathcal{P}_5) = 0 \text{ modulo } (\mathcal{A}_2^+ \mathcal{P}_5), \\ \tau_2(a) + a &= \sum Y \text{ modulo } (\mathcal{A}_2^+ \mathcal{P}_5) = 0 \text{ modulo } (\mathcal{A}_2^+ \mathcal{P}_5), \\ \tau_3(a) + a &= \sum Z \text{ modulo } (\mathcal{A}_2^+ \mathcal{P}_5) = 0 \text{ modulo } (\mathcal{A}_2^+ \mathcal{P}_5), \\ \tau_4(a) + a &= \sum W \text{ modulo } (\mathcal{A}_2^+ \mathcal{P}_5) = 0 \text{ modulo } (\mathcal{A}_2^+ \mathcal{P}_5), \end{aligned}$$

where

$$\begin{aligned} \sum X &= (\gamma_{24} + \gamma_{27})(\mathcal{Y}_{24} + \mathcal{Y}_{27}) + (\gamma_{25} + \gamma_{28})(\mathcal{Y}_{25} + \mathcal{Y}_{28}) + (\gamma_{26} + \gamma_{29})(\mathcal{Y}_{26} + \mathcal{Y}_{29}), \\ \sum Y &= (\gamma_{22} + \gamma_{24})(\mathcal{Y}_{22} + \mathcal{Y}_{24}) + (\gamma_{23} + \gamma_{25})(\mathcal{Y}_{23} + \mathcal{Y}_{25}) + (\gamma_{29} + \gamma_{30})(\mathcal{Y}_{29} + \mathcal{Y}_{30}), \\ \sum Z &= (\gamma_{21} + \gamma_{22})(\mathcal{Y}_{21} + \mathcal{Y}_{22}) + (\gamma_{25} + \gamma_{26})(\mathcal{Y}_{25} + \mathcal{Y}_{26}) + (\gamma_{28} + \gamma_{29})(\mathcal{Y}_{28} + \mathcal{Y}_{29}), \\ \sum W &= (\gamma_{22} + \gamma_{23})(\mathcal{Y}_{22} + \mathcal{Y}_{23}) + (\gamma_{24} + \gamma_{25})(\mathcal{Y}_{24} + \mathcal{Y}_{25}) + (\gamma_{27} + \gamma_{28})(\mathcal{Y}_{27} + \mathcal{Y}_{28}). \end{aligned}$$

These relations imply that $\gamma_i = \gamma_{21}$ for $i = 22, \dots, 30$. Hence, we get $a = \theta(\mathcal{Y}_2) \text{ modulo } (\mathcal{A}_2^+ \mathcal{P}_5)$ with $\theta(\mathcal{Y}_2) = \sum_{21 \leq j \leq 30} \mathcal{Y}_j$.

Now, we have the set $\{[\mathcal{Y}_i] : 31 \leq i \leq 90\}$ is a basis of $[S_5(\mathcal{Y}_3, \mathcal{Y}_4, \mathcal{Y}_5)]$. Suppose that $b = \sum_{31 \leq i \leq 90} \gamma_i \mathcal{Y}_i$ with $\gamma_i \in \mathbb{Z}/2$ and $[b] \in [S_5(\mathcal{Y}_3, \mathcal{Y}_4, \mathcal{Y}_5)]^{S_5}$. By a similar computation from the relations $\tau_t(b) + b = 0 \text{ modulo } (\mathcal{A}_2^+ \mathcal{P}_5)$, $t = 1, 2, 3, 4$, one gets $\gamma_i = 0$ for $71 \leq i \leq 90$ and $\gamma_i = \gamma_{31}$, $i = 32, 33, \dots, 70$. This means $b = q \text{ modulo } (\mathcal{A}_2^+ \mathcal{P}_5)$ with $q = \sum_{31 \leq i \leq 70} \mathcal{Y}_i$. The lemma is proved. $\square$

Lemma 4.1.3. $(Q\mathcal{P}_5(\tilde{\omega}_{(3)}))^{GL_5} = 0$.

Proof. Using a result in [38], we see that $Q\mathcal{P}_5(\tilde{\omega}_{(3)})$ is a $\mathbb{Z}/2$ -vector space of dimension 45 with the basis

$$[\mathcal{B}(\mathcal{Y}_{130} = x_2 x_3 x_4^3 x_5^3)]_{\tilde{\omega}_{(3)}} \cup [\mathcal{B}(\mathcal{Y}_{160} = x_1 x_2 x_3 x_4^2 x_5^3)]_{\tilde{\omega}_{(3)}}.$$

Futhermore, $[S_5(\mathcal{Y}_{130})]$ and $[S_5(\mathcal{Y}_{160})]$ are S_5 -submodules of $Q\mathcal{P}_5^0(\tilde{\omega}_{(3)})$, where $[S_5(\mathcal{Y}_{130})]_{\tilde{\omega}_{(3)}} = \langle [\mathcal{Y}_i : 130 \leq i \leq 159]_{\tilde{\omega}_{(3)}} \rangle$ and $[S_5(\mathcal{Y}_{160})]_{\tilde{\omega}_{(3)}} = \langle [\mathcal{Y}_i : 160 \leq i \leq 174]_{\tilde{\omega}_{(3)}} \rangle$. So, we have a direct summand decomposition of S_5 -modules:

$$Q\mathcal{P}_5(\tilde{\omega}_{(3)}) = [S_5(\mathcal{Y}_{130})]_{\tilde{\omega}_{(3)}} \oplus [S_5(\mathcal{Y}_{160})]_{\tilde{\omega}_{(3)}}.$$

The set $[B(\mathcal{Y}_{130})]_{\tilde{\omega}_{(3)}}$ is a basis of $[S_5(\mathcal{Y}_{130})]_{\tilde{\omega}_{(3)}}$. The action of S_5 on $Q\mathcal{P}_5$ induces the one of it on $[B(\mathcal{Y}_{130})]_{\tilde{\omega}_{(3)}}$. On the other hand, this action is transitive, hence if $a = \sum_{130 \leq i \leq 159} \gamma_i \mathcal{Y}_i$ with $\gamma_i \in \mathbb{Z}/2$ and $[a] \in [S_5(\mathcal{Y}_{130})]^{S_5}$, then from the relations $\tau_t(a) + a = 0 \text{ modulo } (\mathcal{A}_2^+ \mathcal{P}_5 + \mathcal{P}_5^-(\tilde{\omega}_{(3)}))$, $1 \leq t \leq 4$, we get $\gamma_i = \gamma_{130}$, $\forall i$, $131 \leq i \leq 159$. In other words, $[S_5(\mathcal{Y}_j)]_{\tilde{\omega}_{(3)}}^{S_5} = \langle [\theta(\mathcal{Y}_{130})]_{\tilde{\omega}_{(3)}} \rangle$ with $\theta(\mathcal{Y}_{130}) = \sum_{130 \leq i \leq 159} \mathcal{Y}_i$.

Next, we have $\dim[S_5(\mathcal{Y}_{160})]_{\tilde{\omega}_{(3)}} = 15$ with the basis $[B(\mathcal{Y}_{160})]_{\tilde{\omega}_{(3)}}$. Suppose $b = \sum_{160 \leq i \leq 174} \gamma_i \mathcal{Y}_i$ with $\gamma_j \in \mathbb{Z}/2$ and $[b] \in [S_5(\mathcal{Y}_{160})]^{S_5}$. A direct computation shows:

$$\begin{aligned}
\tau_1(b) + b &\equiv_{\tilde{\omega}_{(3)}} \gamma_{169}(\mathcal{Y}_{160} + \mathcal{Y}_{162}) + \gamma_{170}(\mathcal{Y}_{161} + \mathcal{Y}_{163}) + \gamma_{171}(\mathcal{Y}_{164} + \mathcal{Y}_{165}) \\
&\quad + (\gamma_{166} + \gamma_{172})(\mathcal{Y}_{166} + \mathcal{Y}_{172}) + (\gamma_{167} + \gamma_{173})(\mathcal{Y}_{167} + \mathcal{Y}_{173}) \\
&\quad + (\gamma_{168} + \gamma_{174})(\mathcal{Y}_{168} + \mathcal{Y}_{174}) \\
\tau_2(b) + b &\equiv_{\tilde{\omega}_{(3)}} (\gamma_{162} + \gamma_{169})(\mathcal{Y}_{162} + \mathcal{Y}_{169}) + (\gamma_{163} + \gamma_{170})(\mathcal{Y}_{163} + \mathcal{Y}_{170}) \\
&\quad + (\gamma_{164} + \gamma_{166})(\mathcal{Y}_{164} + \mathcal{Y}_{166}) + (\gamma_{165} + \gamma_{167})(\mathcal{Y}_{165} + \mathcal{Y}_{167}) \\
&\quad + (\gamma_{168} + \gamma_{171})(\mathcal{Y}_{168} + \mathcal{Y}_{171}) + (\gamma_{165} + \gamma_{167})(\mathcal{Y}_{165} + \mathcal{Y}_{167}) \\
&\quad + \gamma_{174}(\mathcal{Y}_{172} + \mathcal{Y}_{173}), \\
\tau_3(b) + b &\equiv_{\tilde{\omega}_{(3)}} (\gamma_{160} + \gamma_{162})(\mathcal{Y}_{160} + \mathcal{Y}_{162}) + (\gamma_{161} + \gamma_{164})(\mathcal{Y}_{161} + \mathcal{Y}_{164}) \\
&\quad + (\gamma_{163} + \gamma_{165})(\mathcal{Y}_{163} + \mathcal{Y}_{165}) + (\gamma_{167} + \gamma_{168})(\mathcal{Y}_{167} + \mathcal{Y}_{168}) \\
&\quad + (\gamma_{170} + \gamma_{171})(\mathcal{Y}_{170} + \mathcal{Y}_{171}) + (\gamma_{173} + \gamma_{174})(\mathcal{Y}_{173} + \mathcal{Y}_{174}), \\
\tau_4(b) + b &\equiv_{\tilde{\omega}_{(3)}} (\gamma_{160} + \gamma_{161})(\mathcal{Y}_{160} + \mathcal{Y}_{161}) + (\gamma_{162} + \gamma_{163})(\mathcal{Y}_{162} + \mathcal{Y}_{163}) \\
&\quad + (\gamma_{164} + \gamma_{165})(\mathcal{Y}_{164} + \mathcal{Y}_{165}) + (\gamma_{166} + \gamma_{167})(\mathcal{Y}_{166} + \mathcal{Y}_{167}) \\
&\quad + (\gamma_{169} + \gamma_{170})(\mathcal{Y}_{169} + \mathcal{Y}_{170}) + (\gamma_{172} + \gamma_{173})(\mathcal{Y}_{172} + \mathcal{Y}_{173}).
\end{aligned}$$

Then, from the relations $\tau_t(b) + b = 0$ modulo $(\mathcal{A}_2^+ \mathcal{P}_5 + \mathcal{P}_5^-(\tilde{\omega}_{(3)}))$, we obtain $\gamma_j = 0, \forall j$.

Now, let $[X]_{\tilde{\omega}_{(3)}} \in (Q\mathcal{P}_5(\tilde{\omega}_{(3)}))^{GL_5}$ with $X \in \mathcal{P}_5(\tilde{\omega}_{(3)})$, then $[X]_{\tilde{\omega}_{(3)}} \in (Q\mathcal{P}_5(\tilde{\omega}_{(3)}))^{S_5}$. So, we have $X = \gamma\theta(\mathcal{Y}_{130})$ modulo $(\mathcal{A}_2^+ \mathcal{P}_5 + \mathcal{P}_5^-(\tilde{\omega}_{(3)}))$ with $\gamma \in \mathbb{Z}/2$. By a direct computation, we have

$$\tau_5(X) + X = \gamma\mathcal{Y}_{130} + \text{other terms modulo } (\mathcal{A}_2^+ \mathcal{P}_5 + \mathcal{P}_5^-(\tilde{\omega}_{(3)})) = 0 \text{ modulo } (\mathcal{A}_2^+ \mathcal{P}_5 + \mathcal{P}_5^-(\tilde{\omega}_{(3)})).$$

This implies $\gamma = 0$. The proposition follows. $\square$

By a simple computation using technique as given in the proof of Lemmas 4.1.2 and 4.1.3, we get the following.

Lemma 4.1.4.

i) We have a direct summand decomposition of the S_5 -modules:

$$Q\mathcal{P}_5(\tilde{\omega}_{(2)}) = [S_5(\mathcal{Y}_{106})]_{\tilde{\omega}_{(2)}} \bigoplus [S_5(\mathcal{Y}_{126})]_{\tilde{\omega}_{(2)}}.$$

ii) $(Q\mathcal{P}_5(\tilde{\omega}_{(2)}))^{GL_5} = 0$.

Proof of Proposition 4.1.1. Let $[X] \in (Q\mathcal{P}_5)^{GL_5}$. Then, from Lemmas 4.1.2 - 4.1.4, we have

$$X = \beta_1\theta(\mathcal{Y}_1) + \beta_2\theta(\mathcal{Y}_2) + \beta_3q + \beta_4\theta(\mathcal{Y}_3) \text{ modulo } (\mathcal{A}_2^+ \mathcal{P}_5),$$

with $\beta_t \in \mathbb{Z}/2, 1 \leq t \leq 4$. By using Theorem 2.3.3 and computing $\tau_5(X) + X$ in terms of the admissible monomials (modulo $(\mathcal{A}_2^+ \mathcal{P}_5)$), we conclude

$$\tau_5(X) + X \equiv \beta_1\mathcal{Y}_7 + (\beta_1 + \beta_2)\mathcal{Y}_{16} + \beta_3\mathcal{Y}_{33} + \beta_4(\mathcal{Y}_{91} + \mathcal{Y}_{92} + \mathcal{Y}_{93}) + \text{other terms} \equiv 0.$$

This relation shows $\beta_1 = \beta_2 = \beta_3 = \beta_4 = 0$. The proposition is proved. $\square$

4.2. Computation of $(\text{Ker}(\widetilde{Sq_*^0})_{(5,21)})^{S_5}$

From the results in Sect.3.2.1, we see that $\dim \text{Ker}(\widetilde{Sq_*^0})_{(5,21)} = 666$ with the basis $\{[\mathcal{Y}_{21,t}] : 1 \leq t \leq 666\}$. Here, the admissible monomials $\mathcal{Y}_t := \mathcal{Y}_{21,t}, 1 \leq t \leq 666$, as given in Sect.6.6 and Sect.6.7. Recall that $(\widetilde{Sq_*^0})_{(5,21)}$ is an epimorphism of GL_5 -modules. Combining this and the results

in Subsection 4.1, we get $(Q\mathcal{P}_5)_{21}^{GL_5} \subseteq (\text{Ker}(\widetilde{Sq_*^0})_{(5,21)})^{GL_5}$. By Lemma 3.2.2, we have a direct summand decomposition of the S_5 -modules:

$$\text{Ker}(\widetilde{Sq_*^0})_{(5,21)} = Q\mathcal{P}_5(3, 3, 1, 1) \oplus Q\mathcal{P}_5(3, 3, 3).$$

For $\omega = (3, 3, 1, 1)$, according to the results in Sect.3.2.1, we get $Q\mathcal{P}_5(\omega) = Q\mathcal{P}_5^0(\omega) \oplus Q\mathcal{P}_5^+(\omega)$ with $\dim Q\mathcal{P}_5^0(\omega) = 340$ and $\dim Q\mathcal{P}_5^+(\omega) = 196$. Note that $\mathcal{Z} = x_1^{15}x_2^3x_3^3$ is the minimal spike monomial in $(\mathcal{P}_5)_{21}$ and $\omega(\mathcal{Z}) = \omega$. So, $[x]_\omega = [x]$ for any $x \in (\mathcal{P}_5)_{21}$. By using the results in Sect.3.2.1, we see that there is a direct summand decomposition of the S_5 -modules:

$$Q\mathcal{P}_5^0(\omega) = \langle [S_5(\mathcal{Y}_1)] \rangle \oplus \langle [S_5(\mathcal{Y}_{31})] \rangle \oplus \langle [S_5(\mathcal{Y}_{61})] \rangle \oplus \langle [S_5(\mathcal{Y}_{121})] \rangle \oplus \langle \mathbb{V}_1 \rangle,$$

where

$$\begin{aligned} \mathcal{B}(\mathcal{Y}_1) &= \{\mathcal{Y}_t : 1 \leq t \leq 30\}, & \mathcal{B}(\mathcal{Y}_{31}) &= \{\mathcal{Y}_t : 31 \leq t \leq 60\}, \\ \mathcal{B}(\mathcal{Y}_{61}) &= \{\mathcal{Y}_t : 61 \leq t \leq 120\}, & \mathcal{B}(\mathcal{Y}_{121}) &= \{\mathcal{Y}_t : 121 \leq t \leq 150\}, \\ \mathbb{V}_1 &= \mathcal{B}(\mathcal{Y}_{151}, \mathcal{Y}_{181}, \mathcal{Y}_{201}, \mathcal{Y}_{241}, \mathcal{Y}_{256}, \mathcal{Y}_{266}, \mathcal{Y}_{286}, \mathcal{Y}_{296}, \mathcal{Y}_{301}, \mathcal{Y}_{316}) \\ &= \{\mathcal{Y}_t : 151 \leq t \leq 340\}, \end{aligned}$$

Lemma 4.2.1.

- a) $\langle [S_5(\mathcal{Y}_j)] \rangle^{S_5} = \langle [\theta(\mathcal{Y}_j)] \rangle$ for $j = 1, 31, 61, 121$.
- b) $\langle [S_5(\mathbb{V}_1)] \rangle^{S_5} = 0$.

Outline of the proof. For $j = 1$, let $[f_1] \in \langle [S_5(\mathcal{Y}_1)] \rangle^{S_5}$. Then, we have

$$\tau_m(f_1) = \sum_{X \in \mathcal{B}(\mathcal{Y}_1)} \beta_X \cdot X \text{ modulo } (\mathcal{A}_2^+ \mathcal{P}_5), \quad 0 < m < 5$$

with $f_1 = \sum_{X \in \mathcal{B}(\mathcal{Y}_1)} \beta_X \cdot X$ and $\beta_X \in \mathbb{Z}/2$. By a direct computation, we can see that the action of the symmetric group S_5 on $Q\mathcal{P}_5$ induces the one of it on the set $[\mathcal{B}(\mathcal{Y}_j)]$ and this action is transitive. So, we get $\beta_X = \beta_{X'} = \beta \in \mathbb{Z}/2$ for all $X, X' \in \mathcal{B}(\mathcal{Y}_j)$. This means $f_1 = \theta(\mathcal{Y}_1) \text{ modulo } (\mathcal{A}_2^+ \mathcal{P}_5)$. For $j = 31, 61, 121$, we determine $\tau_m(f_j) + f_j$ in terms of $\mathcal{Y}_j$. Then, by a simple computation using the relations $\tau_m(f_j) = f_j \text{ modulo } (\mathcal{A}_2^+ \mathcal{P}_5)$, we conclude $f_j = \theta(\mathcal{Y}_j) \text{ modulo } (\mathcal{A}_2^+ \mathcal{P}_5)$.

Next, we have $\dim \langle \mathbb{V}_1 \rangle = 190$ with the basis $\{\mathcal{Y}_t : 151 \leq t \leq 340\}$. Assume that $g = \sum_{u \in \mathbb{V}_1} \gamma_u \cdot u$ with $\gamma_u \in \mathbb{Z}/2$ and $[g] \in \langle [\mathbb{V}_1] \rangle^{S_5}$. By using Theorem 2.3.3 and a similar computation as given in the proof of Lemmas 4.1.2 and 4.1.3, we obtain $\gamma_u = 0$ for all $u \in \mathbb{V}_1$. This implies that g is $\mathcal{A}_2$ -decomposable. The lemma follows. $\square$

Lemma 4.2.2. $Q\mathcal{P}_5^+(\omega)^{S_5} = \langle [p := \mathcal{Y}_{411} + \mathcal{Y}_{412} + \cdots + \mathcal{Y}_{419}] \rangle$.

Proof. From Proposition 3.2.6, we see that the sets $[S_5(\mathcal{Y}_{401})] = \langle [\mathcal{Y}_t] : 401 \leq t \leq 410 \rangle$ and $\mathbb{V}_2 = \langle [\mathcal{Y}_t] : 411 \leq t \leq 596 \rangle$, are S_5 -submodules of $Q\mathcal{P}_5^+(\omega)$. Hence, we have a direct summand decomposition of the S_5 -modules:

$$Q\mathcal{P}_5^+(\omega) = [S_5(\mathcal{Y}_{401})] \oplus \mathbb{V}_2.$$

The set $[\mathcal{B}(\mathcal{Y}_{401})]$ is a basis of $[S_5(\mathcal{Y}_{401})]$. Assume that Z is a polynomial such that $[Z] \in [S_5(\mathcal{Y}_{401})]^{S_5}$ and $Z = \sum_{401 \leq t \leq 410} \ell_t \mathcal{Y}_t$ with $\ell_t \in \mathbb{Z}/2$. For $1 \leq j \leq 4$, we explicitly compute $\tau_j(Z) + Z$ in the terms of the admissible monomials $\mathcal{Y}_t$, $401 \leq t \leq 410$.

By a direct computation using Theorem 2.3.3, we get

$$\begin{aligned}
\tau_1(Z) + Z &\equiv \ell_{401}\mathcal{Y}_{404} + \ell_{402}\mathcal{Y}_{405} + \ell_{403}\mathcal{Y}_{407} + (\ell_{406} + \ell_{409})(\mathcal{Y}_{406} + \mathcal{Y}_{409}) \\
&\quad + (\ell_{408} + \ell_{410})(\mathcal{Y}_{408} + \mathcal{Y}_{410}), \\
\tau_2(Z) + Z &\equiv (\ell_{401} + \ell_{404})(\mathcal{Y}_{401} + \mathcal{Y}_{404}) + (\ell_{402} + \ell_{405})(\mathcal{Y}_{402} + \mathcal{Y}_{405}) \\
&\quad + (\ell_{403} + \ell_{406})(\mathcal{Y}_{403} + \mathcal{Y}_{406}) + (\ell_{407} + \ell_{408})(\mathcal{Y}_{407} + \mathcal{Y}_{408}) + \ell_{409}\mathcal{Y}_{410}, \\
\tau_3(Z) + Z &\equiv \ell_{401}\mathcal{Y}_{404} + (\ell_{402} + \ell_{403})(\mathcal{Y}_{402} + \mathcal{Y}_{403}) + (\ell_{405} + \ell_{407})(\mathcal{Y}_{405} + \mathcal{Y}_{407}), \\
&\quad + (\ell_{406} + \ell_{408})(\mathcal{Y}_{406} + \mathcal{Y}_{408}) + (\ell_{409} + \ell_{410})(\mathcal{Y}_{409} + \mathcal{Y}_{410}), \\
\tau_4(Z) + Z &\equiv (\ell_{401} + \ell_{402})(\mathcal{Y}_{401} + \mathcal{Y}_{402}) + \ell_{403}\mathcal{Y}_{407} + (\ell_{404} + \ell_{405})(\mathcal{Y}_{404} + \mathcal{Y}_{405}) \\
&\quad + \ell_{406}\mathcal{Y}_{408} + \ell_{409}\mathcal{Y}_{410}.
\end{aligned}$$

Then, by the relations $\tau_j(Z) = Z$ modulo $(\mathcal{A}_2^+ \mathcal{P}_5)$, $j = 1, 2, 3, 4$, one gets $\ell_t = 0$, $\forall t$. This implies that Z is $\mathcal{A}_2$ -decomposable.

Note that $\dim \mathbb{V}_2 = 186$ with the basis $\{[\mathcal{Y}_j] : 411 \leq j \leq 596\}$. Then if the polynomial $g = \sum_{u \in \mathbb{V}_2} \sigma_u u$, $\sigma_u \in \mathbb{Z}/2$ such that $[g] \in \mathbb{V}_2^{S_5}$, then by a similar argument as given above, we obtain $g = p$ modulo $(\mathcal{A}_2^+ \mathcal{P}_5)$. This completes the proof of the lemma. $\square$

Proposition 4.2.3. *For $\overline{\omega} = (3, 3, 3)$, we have $Q\mathcal{P}_5(\overline{\omega})^{GL_5} = \langle [q_2]_{\overline{\omega}} \rangle$, where*

$$\begin{aligned}
q_2 &= \mathcal{Y}_{652} + \mathcal{Y}_{653} + \mathcal{Y}_{654} + \mathcal{Y}_{656} + \mathcal{Y}_{657} + \mathcal{Y}_{658} + \sum_{661 \leq t \leq 666} \mathcal{Y}_t \\
&= x_1 x_2^6 x_3^3 x_4^5 x_5^6 + x_1 x_2^3 x_3^6 x_4^5 x_5^6 + x_1 x_2^3 x_3^5 x_4^6 x_5^6 + x_1^3 x_2 x_3^5 x_4^6 x_5^6 \\
&\quad + x_1^3 x_2^5 x_3 x_4^6 x_5^6 + x_1^3 x_2^5 x_3^6 x_4 x_5^6 + x_1^3 x_2^3 x_3^5 x_4^4 x_5^6 + x_1^3 x_2^3 x_3^4 x_4^5 x_5^6 \\
&\quad + x_1^3 x_2^3 x_3^5 x_4^4 x_5^6 + x_1^3 x_2^4 x_3^3 x_4^5 x_5^6 + x_1^3 x_2^5 x_3^3 x_4^6 x_5^4 + x_1^3 x_2^5 x_3^6 x_4^3 x_5^4.
\end{aligned}$$

Based on the results in Subsection 3.2, we have

$$\dim Q\mathcal{P}_5(\overline{\omega}) = \dim Q\mathcal{P}_5^0(\overline{\omega}) + \dim Q\mathcal{P}_5^+(\overline{\omega}) = 60 + 70 = 130.$$

Consider the following monomials:

$$\begin{aligned}
\mathcal{Y}_{341} &= x_3^7 x_4^7 x_5^7, & \mathcal{Y}_{351} &= x_2 x_3^6 x_4^7 x_5^7, & \mathcal{Y}_{381} &= x_2^3 x_3^5 x_4^6 x_5^7, \\
\mathcal{Y}_{597} &= x_1 x_2^2 x_3^4 x_4^7 x_5^7, & \mathcal{Y}_{607} &= x_1 x_2^6 x_3 x_4^6 x_5^7, & \mathcal{Y}_{617} &= x_1 x_2^2 x_3^5 x_4^6 x_5^7, \\
\mathcal{Y}_{622} &= x_1 x_2^3 x_3^4 x_4^6 x_5^7, & \mathcal{Y}_{642} &= x_1^3 x_2^5 x_3^2 x_4^4 x_5^7, & \mathcal{Y}_{647} &= x_1^3 x_2^3 x_3^4 x_4^4 x_5^7, \\
\mathcal{Y}_{652} &= x_1 x_2^6 x_3^3 x_4^5 x_5^6, & \mathcal{Y}_{659} &= x_1^3 x_2^5 x_3^2 x_4^5 x_5^6, & \mathcal{Y}_{660} &= x_1^3 x_2^5 x_3^3 x_4^4 x_5^6.
\end{aligned}$$

The following lemma can be easily proved by a direct computation.

Lemma 4.2.4.

i) The following subspaces are S_5 -submodules of $Q\mathcal{P}_5(\overline{\omega})$:

$$\begin{aligned}
\langle [S_5(\mathcal{Y}_a)]_{\overline{\omega}} \rangle, & \quad a = 341, 351, 381, 597, \mathbb{V}_3 := \langle [S_5(\mathcal{Y}_{607}, \mathcal{Y}_{617}, \mathcal{Y}_{622}, \mathcal{Y}_{642}, \mathcal{Y}_{647})]_{\overline{\omega}} \rangle, \\
& \quad \mathbb{V}_4 := \langle [S_5(\mathcal{Y}_{652}, \mathcal{Y}_{659}, \mathcal{Y}_{660})]_{\overline{\omega}} \rangle.
\end{aligned}$$

ii) We have a direct summand decomposition of the S_5 -modules:

$$\begin{aligned}
Q\mathcal{P}_5(\overline{\omega}) &= \langle [S_5(\mathcal{Y}_{341})]_{\overline{\omega}} \rangle \oplus \langle [S_5(\mathcal{Y}_{351})]_{\overline{\omega}} \rangle \oplus \langle [S_5(\mathcal{Y}_{381})]_{\overline{\omega}} \rangle \\
&\quad \oplus \langle [S_5(\mathcal{Y}_{597})]_{\overline{\omega}} \rangle \oplus \mathbb{V}_3 \oplus \mathbb{V}_4.
\end{aligned}$$

Lemma 4.2.5.

- i) $\langle [S_5(\mathcal{Y}_a)]_{\overline{\omega}} \rangle^{S_5} = \langle [\theta(\mathcal{Y}_a)]_{\overline{\omega}} \rangle$, for $a = 341, 351, 381, 597$.
- ii) $\langle [S_5(\mathbb{V}_3)]_{\overline{\omega}} \rangle^{S_5} = \langle [q_1 := \mathcal{Y}_{610} + \mathcal{Y}_{611} + \mathcal{Y}_{613} + \mathcal{Y}_{614} + \mathcal{Y}_{616} + \sum_{647 \leq t \leq 651} \mathcal{Y}_t]_{\overline{\omega}} \rangle$.
- iii) $\langle [S_5(\mathbb{V}_4)]_{\overline{\omega}} \rangle^{S_5} = \langle [q_2 := \mathcal{Y}_{652} + \mathcal{Y}_{653} + \mathcal{Y}_{654} + \mathcal{Y}_{656} + \mathcal{Y}_{657} + \mathcal{Y}_{658} + \sum_{661 \leq t \leq 666} \mathcal{Y}_t]_{\overline{\omega}} \rangle$,

The proof of the lemma is straightforward.

Proof of Proposition 4.2.3. By Lemmas 4.2.4 and 4.2.5, we get

$$Q\mathcal{P}_5(\overline{\omega})^{S_5} = \langle [\theta(\mathcal{Y}_{341})]_{\overline{\omega}}, [\theta(\mathcal{Y}_{351})]_{\overline{\omega}}, [\theta(\mathcal{Y}_{381})]_{\overline{\omega}}, [\theta(\mathcal{Y}_{597})]_{\overline{\omega}}, [q_1]_{\overline{\omega}}, [q_2]_{\overline{\omega}} \rangle.$$

Let X be a polynomial in $\mathcal{P}_5(\overline{\omega})$ such that $[X]_{\overline{\omega}} \in Q\mathcal{P}_5(\overline{\omega})^{GL_5}$. Then, we have

$$X = v_1\theta(\mathcal{Y}_{341}) + v_2\theta(\mathcal{Y}_{351}) + v_3\theta(\mathcal{Y}_{381}) + v_4\theta(\mathcal{Y}_{597}) + v_5q_1 + v_6q_2 \text{ modulo } (\mathcal{A}_2^+ \mathcal{P}_5 + \mathcal{P}_5^-(\overline{\omega})),$$

with $v_j \in \mathbb{Z}/2$ for $1 \leq j \leq 6$. We explicitly compute $\tau_5(X)$ in terms of the admissible monomials $\mathcal{Y}_t$ with $t = 341, 342, \dots, 400, 597, 598, \dots, 666$. By a direct computation, one gets

$$\begin{aligned} \tau_5(X) + X &= (v_1 + v_2)\mathcal{Y}_{342} + (v_2 + v_4)\mathcal{Y}_{351} + v_2\mathcal{Y}_{354} + v_3\mathcal{Y}_{383} \\ &\quad + v_5\mathcal{Y}_{650} + \text{other terms modulo } (\mathcal{A}_2^+ \mathcal{P}_5 + \mathcal{P}_5^-(\overline{\omega})). \end{aligned}$$

Since $[X]_{\overline{\omega}} \in Q\mathcal{P}_5(\overline{\omega})^{GL_5}$, $v_j = 0$, $1 \leq j \leq 5$. The proposition is proved. $\square$

Combining Lemmas 4.2.1, 4.2.2 and 4.2.5, one gets the following.

Corollary 4.2.6. $\dim(\text{Ker}(\widetilde{Sq_*^0}_{(5,21)})^{S_5}) = 11$.

4.3. Proof of Theorem 1.2

Suppose that $[T] \in (Q\mathcal{P}_5)_{21}^{GL_5}$ with T is a polynomial in $(\mathcal{P}_5)_{21}$. From Proposition 4.2.3, we have $T = T^* + \zeta_6 q_2 \text{ modulo } (\mathcal{A}_2^+ \mathcal{P}_5)$ with $T^* \in \mathcal{P}_5^-(\overline{\omega})$ and $\zeta_6 \in \mathbb{Z}/2$. By a simple computation, we see that $[q_2] \in (Q\mathcal{P}_5)_{21}^{S_5}$. This implies that $[T^*]$ is an S_5 -invariant. On the other hand, $[\mathcal{P}_5^-(\overline{\omega})] = Q\mathcal{P}_5(\omega)$. Hence, by Lemmas 4.2.1 and 4.2.2, we obtain

$$T^* = \zeta_1\theta(\mathcal{Y}_1) + \zeta_2\theta(\mathcal{Y}_{31}) + \zeta_3\theta(\mathcal{Y}_{61}) + \zeta_4\theta(\mathcal{Y}_{121}) + \zeta_5 p \text{ modulo } (\mathcal{A}_2^+ \mathcal{P}_5),$$

where $\zeta_i \in \mathbb{Z}/2$. Using Theorem 1.1, we computing $\tau_5(T) + T$ in terms of the admissible monomials $\mathcal{Y}_t$, $1 \leq t \leq 666$, we conclude

$$\tau_5(T) + T \equiv (\zeta_1 + \zeta_3)\mathcal{Y}_4 + (\zeta_2 + \zeta_4)\mathcal{Y}_{35} + \zeta_3\mathcal{Y}_{61} + (\zeta_3 + \zeta_4)\mathcal{Y}_{107} + \zeta_5\mathcal{Y}_{153} + \text{other terms}.$$

By the relation $\tau_5(T) = T \text{ modulo } (\mathcal{A}_2^+ \mathcal{P}_5)$, one gets $\zeta_i = 0$, $1 \leq i \leq 5$. This shows $T = \zeta_6 q_2 \text{ modulo } (\mathcal{A}_2^+ \mathcal{P}_5)$. The proof of the theorem is completed.

5. Proof of Theorem 1.3

First of all, we recall the definition of the Singer transfer. Let $\widetilde{\mathcal{P}}_1$ be the submdule of $\mathbb{Z}/2[x_1, x_1^{-1}]$ spanned by all powers x_1^i with $i \geq -1$. The usual $\mathcal{A}_2$ -action on $\mathcal{P}_1 = \mathbb{Z}/2[x_1]$ is canonically extended to an $\mathcal{A}_2$ -action on $\mathbb{Z}/2[x_1, x_1^{-1}]$. Then, $\widetilde{\mathcal{P}}_1$ is an $\mathcal{A}_2$ -submodule of $\mathbb{Z}/2[x_1, x_1^{-1}]$. The inclusion $\mathcal{P}_1 \subset \widetilde{\mathcal{P}}_1$ gives rise to a short exact sequence of $\mathcal{A}_2$ -module:

$$0 \longrightarrow \mathcal{P}_1 \longrightarrow \widetilde{\mathcal{P}}_1 \longrightarrow \Sigma^{-1} \mathbb{Z}/2 \longrightarrow 0.$$

Let e_1 be the corresponding element in $\text{Ext}_{\mathcal{A}_2}^1(\Sigma^{-1} \mathbb{Z}/2, \mathcal{P}_1)$. Singer set

$$e_d := e_1 \otimes \dots \otimes e_1 \in \text{Ext}_{\mathcal{A}_2}^d(\sum_{i=1}^d \mathbb{Z}/2, \mathcal{P}_d).$$

Then, he defined

$$Tr_d^* : \text{Tor}_d^{\mathcal{A}_2}(\mathbb{Z}/2, \Sigma^{-d} \mathbb{Z}/2) \longrightarrow \text{Tor}_0^{\mathcal{A}_2}(\mathbb{Z}/2, \mathcal{P}_d) = Q\mathcal{P}_d$$

by $Tr_d^*(z) = e_d \cap z$. Its image is a submodule of $(Q\mathcal{P}_d)^{GL_d}$. The d -th Singer transfer is defined to be the dual of Tr_d^* .

The algebra $\text{Ext}_{\mathcal{A}_2}^{*,*}(\mathbb{Z}/2, \mathbb{Z}/2)$ is described in term of the mod 2 lambda algebra Λ (see Lin [10]). Recall that Λ is a bigraded differential algebra over $\mathbb{Z}/2$ generated by $\lambda_j \in \Lambda^{1,j}$, $j \geq 0$, with the Adem relations

$$\lambda_j \lambda_{2j+1+m} = \sum_{\nu \geq 0} \binom{m - \nu - 1}{\nu} \lambda_{j+m-\nu} \lambda_{2j+1+\nu},$$

for $m \geq 0$ and the differential

$$\partial(\lambda_s) = \sum_{\nu \geq 0} \binom{s - \nu - 1}{\nu + 1} \lambda_{s-\nu-1} \lambda_\nu, \text{ for } s > 0$$

Further, $H^{s,t}(\Lambda, \partial) \cong \text{Ext}_{\mathcal{A}_2}^{s,t}(\mathbb{Z}/2, \mathbb{Z}/2)$. So, the algebra Λ is regarded as the E_1 -page of the Adams spectral sequence.

As well known, the dual of $\mathcal{P}_d = \mathbb{Z}/2[x_1, \dots, x_d]$ is isomorphic to $\Gamma(a_1, \dots, a_d)$, the divided power algebra generated by $a_1, \dots, a_d$, where a_t is dual to $x_t \in \mathcal{P}_d$ with respect to the basis of $\mathcal{P}_d$ consisting of all monomials in $x_1, \dots, x_d$. So, $H_*(B(\mathbb{Z}/2)^{\times d}) \cong \Gamma(a_1, \dots, a_d)$. The algebra $H_*(B(\mathbb{Z}/2)^{\times d})$ has a right $\mathcal{A}_2$ -module structure. The right action of $\mathcal{A}_2$ on $H_*(B(\mathbb{Z}/2)^{\times d})$ is given by $(a_t^{(s)})Sq^n = \binom{s-n}{n} a_t^{(s-n)}$ and Cartan's formula. In [6], Chơn and Hà have constructed the $\mathbb{Z}/2$ -linear map $\psi_d : H_*(B(\mathbb{Z}/2)^{\times d}) \longrightarrow \Lambda_d$, which is determined by the following inductive formula:

$$\psi_d(a^{(I,s)}) = \begin{cases} \lambda_s & \text{if } d-1 = \ell(I) = 0, \\ \sum_{t \geq 1} \psi_{d-1}(a^I Sq^{t-s}) \lambda_t & \text{if } d-1 = \ell(I) > 0, \end{cases}$$

for any $a^{(I,s)} = a_1^{(n_1)} \dots a_{d-1}^{(n_{d-1})} a_d^{(s)} \in H_*(B(\mathbb{Z}/2)^{\times d})$ and $I = (n_1, n_2, \dots, n_{d-1})$. This map is a presentation in the $\mathbb{Z}/2$ -lambda algebra of the transfer homomorphism Tr_d . Note that ψ_d is not an $\mathcal{A}_2$ -homomorphism. Then, we have the homomorphism of algebras

$$\psi = \oplus_{d \geq 1} \psi_d : \oplus_{d \geq 1} H_*(B(\mathbb{Z}/2)^{\times d}) \longrightarrow \oplus_{d \geq 1} \Lambda_d = \Lambda,$$

which induces the Singer transfer.

Proposition 5.1 (Chơn and Hà [6]). *If $Z \in P_{\mathcal{A}_2} H_*(B(\mathbb{Z}/2)^{\times d})$, then $\psi_d(Z)$ is a cycle in Λ and $\text{Tr}_d([Z]) = [\psi_d(Z)]$.*

We now are ready to prove Theorem 1.3.

We see that $\lambda_3 \in \Lambda^{1,3}$, and $\bar{f}_0 = \lambda_3 \lambda_5 \lambda_6 \lambda_4 + \lambda_3^2 \lambda_7 \lambda_5 + \lambda_7 \lambda_5 \lambda_2 \lambda_3^2 + \lambda_7 \lambda_5 \lambda_4 \lambda_2 \in \Lambda^{4,18}$ are the cycles in the lambda algebra Λ . By Lin [10], we have $\text{Ext}_{\mathcal{A}_2}^{5,26}(\mathbb{Z}/2, \mathbb{Z}/2) = \langle h_2 f_0 \rangle$, with $h_2 = [\lambda_3] \in \text{Ext}_{\mathcal{A}_2}^{1,4}(\mathbb{Z}/2, \mathbb{Z}/2)$ and $f_0 = [\bar{f}_0] \in \text{Ext}_{\mathcal{A}_2}^{4,22}(\mathbb{Z}/2, \mathbb{Z}/2)$. Note that $h_2 f_0 = h_1 g_1$ with $h_1 \in \text{Ext}_{\mathcal{A}_2}^{1,2}(\mathbb{Z}/2, \mathbb{Z}/2)$ and $g_1 \in \text{Ext}_{\mathcal{A}_2}^{4,24}(\mathbb{Z}/2, \mathbb{Z}/2)$.

Consider the element $Z \in H_{21}(B(\mathbb{Z}/2)^{\times 5})$, where

$$\begin{aligned}
Z = & a_1^{(3)} a_2^{(3)} a_3^{(5)} a_4^{(1)} a_5^{(9)} + a_1^{(3)} a_2^{(3)} a_3^{(5)} a_4^{(2)} a_5^{(8)} + a_1^{(3)} a_2^{(3)} a_3^{(6)} a_4^{(1)} a_5^{(8)} + a_1^{(3)} a_2^{(3)} a_3^{(6)} a_4^{(2)} a_5^{(7)} \\
& + a_1^{(3)} a_2^{(3)} a_3^{(5)} a_4^{(4)} a_5^{(6)} + a_1^{(3)} a_2^{(3)} a_3^{(6)} a_4^{(3)} a_5^{(6)} + a_1^{(3)} a_2^{(5)} a_3^{(6)} a_4^{(1)} a_5^{(6)} + a_1^{(3)} a_2^{(3)} a_3^{(5)} a_4^{(5)} a_5^{(5)} \\
& + a_1^{(3)} a_2^{(3)} a_3^{(6)} a_4^{(4)} a_5^{(5)} + a_1^{(3)} a_2^{(5)} a_3^{(6)} a_4^{(2)} a_5^{(5)} + a_1^{(3)} a_2^{(3)} a_3^{(9)} a_4^{(1)} a_5^{(5)} + a_1^{(3)} a_2^{(5)} a_3^{(7)} a_4^{(1)} a_5^{(5)} \\
& + a_1^{(3)} a_2^{(3)} a_3^{(9)} a_4^{(2)} a_5^{(4)} + a_1^{(3)} a_2^{(5)} a_3^{(7)} a_4^{(2)} a_5^{(4)} + a_1^{(3)} a_2^{(3)} a_3^{(10)} a_4^{(1)} a_5^{(4)} + a_1^{(3)} a_2^{(6)} a_3^{(7)} a_4^{(1)} a_5^{(4)} \\
& + a_1^{(3)} a_2^{(5)} a_3^{(6)} a_4^{(4)} a_5^{(3)} + a_1^{(3)} a_2^{(6)} a_3^{(7)} a_4^{(2)} a_5^{(3)} + a_1^{(3)} a_2^{(3)} a_3^{(10)} a_4^{(2)} a_5^{(3)} + a_1^{(3)} a_2^{(3)} a_3^{(11)} a_4^{(2)} a_5^{(2)} \\
& + a_1^{(3)} a_2^{(5)} a_3^{(9)} a_4^{(2)} a_5^{(2)} + a_1^{(3)} a_2^{(6)} a_3^{(10)} a_4^{(1)} a_5^{(1)} + a_1^{(3)} a_2^{(3)} a_3^{(5)} a_4^{(9)} a_5^{(1)} + a_1^{(3)} a_2^{(3)} a_3^{(5)} a_4^{(8)} a_5^{(2)} \\
& + a_1^{(3)} a_2^{(3)} a_3^{(6)} a_4^{(8)} a_5^{(1)} + a_1^{(3)} a_2^{(3)} a_3^{(6)} a_4^{(7)} a_5^{(2)} + a_1^{(3)} a_2^{(3)} a_3^{(5)} a_4^{(6)} a_5^{(4)} + a_1^{(3)} a_2^{(3)} a_3^{(6)} a_4^{(6)} a_5^{(3)} \\
& + a_1^{(3)} a_2^{(5)} a_3^{(6)} a_4^{(6)} a_5^{(1)} + a_1^{(3)} a_2^{(3)} a_3^{(5)} a_4^{(5)} a_5^{(5)} + a_1^{(3)} a_2^{(3)} a_3^{(6)} a_4^{(5)} a_5^{(4)} + a_1^{(3)} a_2^{(5)} a_3^{(6)} a_4^{(5)} a_5^{(2)} \\
& + a_1^{(3)} a_2^{(3)} a_3^{(9)} a_4^{(5)} a_5^{(1)} + a_1^{(3)} a_2^{(5)} a_3^{(7)} a_4^{(5)} a_5^{(1)} + a_1^{(3)} a_2^{(3)} a_3^{(9)} a_4^{(4)} a_5^{(2)} + a_1^{(3)} a_2^{(5)} a_3^{(7)} a_4^{(4)} a_5^{(2)} \\
& + a_1^{(3)} a_2^{(3)} a_3^{(10)} a_4^{(4)} a_5^{(1)} + a_1^{(3)} a_2^{(6)} a_3^{(7)} a_4^{(4)} a_5^{(1)} + a_1^{(3)} a_2^{(5)} a_3^{(6)} a_4^{(3)} a_5^{(4)} + a_1^{(3)} a_2^{(6)} a_3^{(7)} a_4^{(3)} a_5^{(2)} \\
& + a_1^{(3)} a_2^{(3)} a_3^{(10)} a_4^{(3)} a_5^{(2)} + a_1^{(3)} a_2^{(3)} a_3^{(11)} a_4^{(2)} a_5^{(2)} + a_1^{(3)} a_2^{(5)} a_3^{(9)} a_4^{(2)} a_5^{(2)} + a_1^{(3)} a_2^{(6)} a_3^{(10)} a_4^{(1)} a_5^{(1)} \\
& + a_1^{(3)} a_2^{(3)} a_3^{(12)} a_4^{(1)} a_5^{(2)} + a_1^{(3)} a_2^{(7)} a_3^{(8)} a_4^{(1)} a_5^{(2)} + a_1^{(3)} a_2^{(11)} a_3^{(4)} a_4^{(1)} a_5^{(2)} + a_1^{(3)} a_2^{(13)} a_3^{(2)} a_4^{(1)} a_5^{(2)} \\
& + a_1^{(3)} a_2^{(14)} a_3^{(1)} a_4^{(1)} a_5^{(2)} + a_1^{(3)} a_2^{(12)} a_3^{(3)} a_4^{(1)} a_5^{(2)} + a_1^{(3)} a_2^{(8)} a_3^{(7)} a_4^{(1)} a_5^{(2)} + a_1^{(3)} a_2^{(4)} a_3^{(11)} a_4^{(1)} a_5^{(2)} \\
& + a_1^{(3)} a_2^{(2)} a_3^{(13)} a_4^{(1)} a_5^{(2)} + a_1^{(3)} a_2^{(1)} a_3^{(14)} a_4^{(1)} a_5^{(2)} + a_1^{(3)} a_2^{(6)} a_3^{(6)} a_4^{(3)} a_5^{(3)} + a_1^{(3)} a_2^{(5)} a_3^{(5)} a_4^{(5)} a_5^{(3)} \\
& + a_1^{(3)} a_2^{(3)} a_3^{(3)} a_4^{(9)} a_5^{(3)} + a_1^{(3)} a_2^{(5)} a_3^{(3)} a_4^{(7)} a_5^{(3)} + a_1^{(3)} a_2^{(7)} a_3^{(7)} a_4^{(2)} a_5^{(2)} + a_1^{(3)} a_2^{(6)} a_3^{(9)} a_4^{(1)} a_5^{(2)} \\
& + a_1^{(3)} a_2^{(9)} a_3^{(6)} a_4^{(1)} a_5^{(2)} + a_1^{(3)} a_2^{(10)} a_3^{(5)} a_4^{(1)} a_5^{(2)} + a_1^{(3)} a_2^{(5)} a_3^{(10)} a_4^{(2)} a_5^{(1)} + a_1^{(3)} a_2^{(13)} a_3^{(3)} a_4^{(1)} a_5^{(1)} \\
& + a_1^{(3)} a_2^{(5)} a_3^{(11)} a_4^{(1)} a_5^{(1)} + a_1^{(3)} a_2^{(9)} a_3^{(7)} a_4^{(1)} a_5^{(1)}.
\end{aligned}$$

From the proof of Theorem 1.2 (see Sect.4.3), $\{[q_2]\}$ is a basis of $(Q\mathcal{P}_5)_{21}^{GL_5}$. Obviously, $\langle Z, q_2 \rangle = 1$. Then, it is easily verified that $Z \in P_{\mathcal{A}_2} H_{21}(B(\mathbb{Z}/2)^{\times 5})$. This implies $[Z] = ([q_2])^*$. By a direct computation, we have the representation in the $\mathbb{Z}/2$ -lambda algebra of Tr_5 :

$$a^{(n_1, \dots, n_5)} \longmapsto \sum_{i, j, h, m, k_1+k_2+k_3+k_4=i-n_1, \ell_1+\ell_2+\ell_3=j-n_2+k_4, u_1+u_2=h-n_3+k_3+\ell_3} \mathcal{T}_1 \cdot \mathcal{T}_2 \cdot \mathcal{T}_3 \cdot \mathcal{T}_4,$$

where

$$\begin{aligned}
\mathcal{T}_1 &= \binom{n_5-k_1}{k_1} \binom{n_4-k_2}{k_2} \binom{n_3-k_3}{k_3} \binom{n_2-k_4}{k_4}; \\
\mathcal{T}_2 &= \binom{n_5-k_1-\ell_1}{\ell_1} \binom{n_4-k_2-\ell_2}{\ell_2} \binom{n_3-k_3-\ell_3}{\ell_3}; \\
\mathcal{T}_3 &= \binom{n_5-k_1-\ell_1-u_1}{u_1} \binom{n_4-k_2-\ell_2-u_2}{u_2}; \\
\mathcal{T}_4 &= \binom{n_1+n_2+n_3+n_4+n_5-i-j-h-m}{m-n_4+k_2+\ell_2+u_2} \lambda_{i, j, h, m, n_1+n_2+n_3+n_4+n_5-i-j-h-m},
\end{aligned}$$

with the binomial factors (modulo 2).

Using the above presentation, we obtain

$$Z \longmapsto \lambda_3 \bar{f}_0 + \lambda_3^2 \lambda_5 \lambda_3 \lambda_7 = \lambda_3 \bar{f}_0 + \partial(\lambda_3^2 \lambda_5 \lambda_{11}).$$

Combining this and Proposition 5.1, we see that $h_2 f_0 = h_1 g_1$ is in the image of Tr_5 . Now, the theorem follows from Theorem 1.2.

6. Appendix

In this section, we describe all the admissible monomials in $\mathcal{B}_5(n)$ for $n = 8, 21, 22, 47$. These monomials are mentioned in Sect.3 and Sect.4. .

6.1. $\mathcal{A}_2$ -generators for $\mathcal{P}_5$ in degree 8

Based on a result in [38], we have $\mathcal{B}_5(8) = \bigcup_{1 \leq j \leq 3} \mathcal{B}_5(\tilde{\omega}_{(j)})$ and $|\mathcal{B}_5(8)| = 174$. Here $\tilde{\omega}_{(1)} = (2, 1, 1)$, $\tilde{\omega}_{(2)} = (2, 3)$, $\tilde{\omega}_{(3)} = (4, 2)$. By a simple computation, we get

$$|\overline{\Phi}(\mathcal{B}_4(\tilde{\omega}_{(1)})) \cup \mathcal{B}(5, \tilde{\omega}_{(1)})| = 105, \quad |\overline{\Phi}(\mathcal{B}_4(\tilde{\omega}_{(2)})) \cup \mathcal{B}(5, \tilde{\omega}_{(2)})| = 24,$$

$$|\overline{\Phi}(\mathcal{B}_4(\tilde{\omega}_{(3)})) \cup \mathcal{B}(5, \tilde{\omega}_{(3)})| = 45,$$

where the set $\mathcal{B}(5, \tilde{\omega}_{(j)})$ is given in Sect.3.2.2. Furthermore,

$$\mathcal{B}_5(\tilde{\omega}_{(j)}) = \overline{\Phi}(\mathcal{B}_4(\tilde{\omega}_{(j)})) \cup \mathcal{B}(5, \tilde{\omega}_{(j)}), \quad j = 1, 2, 3.$$

This implies that Conjecture 3.1.4 is true for $d = 5$ and in degree 8.

Now, we have $\mathcal{B}_5(8) = \{\mathcal{Y}_{8,t} : 1 \leq t \leq 174\}$, where the monomials $\mathcal{Y}_{8,t} : 1 \leq t \leq 174$, are determined as follows:

$$\begin{array}{llll} \mathcal{Y}_{8,1} = x_4 x_5^7, & \mathcal{Y}_{8,2} = x_4^7 x_5, & \mathcal{Y}_{8,3} = x_3 x_5^7, & \mathcal{Y}_{8,4} = x_3 x_4^7, \\ \mathcal{Y}_{8,5} = x_3^7 x_5, & \mathcal{Y}_{8,6} = x_3^7 x_4, & \mathcal{Y}_{8,7} = x_2 x_5^7, & \mathcal{Y}_{8,8} = x_2 x_4^7, \\ \mathcal{Y}_{8,9} = x_2 x_3^7, & \mathcal{Y}_{8,10} = x_2^7 x_5, & \mathcal{Y}_{8,11} = x_2^7 x_4, & \mathcal{Y}_{8,12} = x_2^7 x_3, \\ \mathcal{Y}_{8,13} = x_1 x_5^7, & \mathcal{Y}_{8,14} = x_1 x_4^7, & \mathcal{Y}_{8,15} = x_1 x_3^7, & \mathcal{Y}_{8,16} = x_1 x_2^7, \\ \mathcal{Y}_{8,17} = x_1^7 x_2, & \mathcal{Y}_{8,18} = x_1^7 x_3, & \mathcal{Y}_{8,19} = x_1^7 x_4, & \mathcal{Y}_{8,20} = x_1^7 x_5, \\ \mathcal{Y}_{8,21} = x_4^3 x_5^5, & \mathcal{Y}_{8,22} = x_3^3 x_5^5, & \mathcal{Y}_{8,23} = x_3^3 x_4^5, & \mathcal{Y}_{8,24} = x_2^3 x_5^5, \\ \mathcal{Y}_{8,25} = x_2^3 x_4^5, & \mathcal{Y}_{8,26} = x_2^3 x_3^5, & \mathcal{Y}_{8,27} = x_1^3 x_5^5, & \mathcal{Y}_{8,28} = x_1^3 x_4^5, \\ \mathcal{Y}_{8,29} = x_1^3 x_3^5, & \mathcal{Y}_{8,30} = x_1^3 x_2^5, & \mathcal{Y}_{8,31} = x_3 x_4 x_5^6, & \mathcal{Y}_{8,32} = x_3 x_4^6 x_5, \\ \mathcal{Y}_{8,33} = x_2 x_4 x_5^6, & \mathcal{Y}_{8,34} = x_2 x_3 x_5^6, & \mathcal{Y}_{8,35} = x_2 x_3 x_4^6, & \mathcal{Y}_{8,36} = x_2 x_3 x_3^6, \\ \mathcal{Y}_{8,37} = x_2 x_3^6 x_5, & \mathcal{Y}_{8,38} = x_2 x_3^6 x_4, & \mathcal{Y}_{8,39} = x_1 x_4 x_5^6, & \mathcal{Y}_{8,40} = x_1 x_3 x_5^6, \\ \mathcal{Y}_{8,41} = x_1 x_3 x_4^6, & \mathcal{Y}_{8,42} = x_1 x_2 x_5^6, & \mathcal{Y}_{8,43} = x_1 x_2 x_4^6, & \mathcal{Y}_{8,44} = x_1 x_2 x_3^6, \\ \mathcal{Y}_{8,45} = x_1 x_4^6 x_5, & \mathcal{Y}_{8,46} = x_1 x_3^6 x_5, & \mathcal{Y}_{8,47} = x_1 x_3^6 x_4, & \mathcal{Y}_{8,48} = x_1 x_2^6 x_5, \\ \mathcal{Y}_{8,49} = x_1 x_2^6 x_4, & \mathcal{Y}_{8,50} = x_1 x_2^6 x_3, & \mathcal{Y}_{8,51} = x_3^3 x_4 x_5^4, & \mathcal{Y}_{8,52} = x_3^3 x_4^4 x_5, \\ \mathcal{Y}_{8,53} = x_2^3 x_4 x_5^4, & \mathcal{Y}_{8,54} = x_2^3 x_3 x_5^4, & \mathcal{Y}_{8,55} = x_2^3 x_3 x_4^4, & \mathcal{Y}_{8,56} = x_2^3 x_4^4 x_5, \\ \mathcal{Y}_{8,57} = x_2^3 x_3^4 x_5, & \mathcal{Y}_{8,58} = x_2^3 x_3^4 x_4, & \mathcal{Y}_{8,59} = x_1^3 x_4 x_5^4, & \mathcal{Y}_{8,60} = x_1^3 x_3 x_5^4, \\ \mathcal{Y}_{8,61} = x_1^3 x_3 x_4^4, & \mathcal{Y}_{8,62} = x_1^3 x_4^4 x_5, & \mathcal{Y}_{8,63} = x_1^3 x_3^4 x_5, & \mathcal{Y}_{8,64} = x_1^3 x_3^4 x_4, \\ \mathcal{Y}_{8,65} = x_1^3 x_2 x_5^4, & \mathcal{Y}_{8,66} = x_1^3 x_2 x_4^4, & \mathcal{Y}_{8,67} = x_1^3 x_2^4 x_5, & \mathcal{Y}_{8,68} = x_1^3 x_2^4 x_4, \\ \mathcal{Y}_{8,69} = x_1^3 x_2 x_3^4, & \mathcal{Y}_{8,70} = x_1^3 x_2^4 x_3, & \mathcal{Y}_{8,71} = x_3 x_4^2 x_5^5, & \mathcal{Y}_{8,72} = x_2 x_4^2 x_5^5, \\ \mathcal{Y}_{8,73} = x_2 x_3^2 x_5^5, & \mathcal{Y}_{8,74} = x_2 x_3^2 x_4^5, & \mathcal{Y}_{8,75} = x_1 x_4^2 x_5^5, & \mathcal{Y}_{8,76} = x_1 x_3^2 x_5^5, \\ \mathcal{Y}_{8,77} = x_1 x_3^2 x_4^5, & \mathcal{Y}_{8,78} = x_1 x_2^2 x_5^5, & \mathcal{Y}_{8,79} = x_1 x_2^2 x_4^5, & \mathcal{Y}_{8,80} = x_1 x_2^2 x_3^5, \\ \mathcal{Y}_{8,81} = x_3 x_4^3 x_5^4, & \mathcal{Y}_{8,82} = x_2 x_4^3 x_5^4, & \mathcal{Y}_{8,83} = x_2 x_3^3 x_5^4, & \mathcal{Y}_{8,84} = x_2 x_3^3 x_4^4, \\ \mathcal{Y}_{8,85} = x_1 x_4^3 x_5^4, & \mathcal{Y}_{8,86} = x_1 x_3^3 x_5^4, & \mathcal{Y}_{8,87} = x_1 x_3^3 x_4^4, & \mathcal{Y}_{8,88} = x_1 x_2^3 x_5^4, \\ \mathcal{Y}_{8,89} = x_1 x_2^3 x_4^4, & \mathcal{Y}_{8,90} = x_1 x_2^3 x_3^4, & \mathcal{Y}_{8,91} = x_2 x_3 x_4^2 x_5^4, & \mathcal{Y}_{8,92} = x_2 x_3^2 x_4 x_5^4, \\ \mathcal{Y}_{8,93} = x_2 x_3^2 x_4^4 x_5, & \mathcal{Y}_{8,94} = x_1 x_3 x_4^2 x_5^4, & \mathcal{Y}_{8,95} = x_1 x_2 x_4^2 x_5^4, & \mathcal{Y}_{8,96} = x_1 x_2 x_3^2 x_5^4, \\ \mathcal{Y}_{8,97} = x_1 x_2 x_3^2 x_4^4, & \mathcal{Y}_{8,98} = x_1 x_2^2 x_4 x_5^4, & \mathcal{Y}_{8,99} = x_1 x_2^2 x_4^4 x_5, & \mathcal{Y}_{8,100} = x_1 x_2^2 x_4^4 x_5, \\ \mathcal{Y}_{8,101} = x_1 x_2^2 x_4^4 x_5, & \mathcal{Y}_{8,102} = x_1 x_2^2 x_3 x_5^4, & \mathcal{Y}_{8,103} = x_1 x_2^2 x_3 x_4^4, & \mathcal{Y}_{8,104} = x_1 x_2^2 x_3^4 x_5, \\ \mathcal{Y}_{8,105} = x_1 x_2^2 x_3^4 x_4, & \mathcal{Y}_{8,106} = x_2 x_3^2 x_4^2 x_5^3, & \mathcal{Y}_{8,107} = x_2 x_3^2 x_4^2 x_5^2, & \mathcal{Y}_{8,108} = x_2 x_3^3 x_4^2 x_5^2, \\ \mathcal{Y}_{8,109} = x_2^3 x_3 x_4^2 x_5^2, & \mathcal{Y}_{8,110} = x_1 x_3^2 x_4^2 x_5^3, & \mathcal{Y}_{8,111} = x_1 x_3^2 x_4^2 x_5^2, & \mathcal{Y}_{8,112} = x_1 x_3^3 x_4^2 x_5^2, \\ \mathcal{Y}_{8,113} = x_1 x_2^2 x_4^2 x_5^3, & \mathcal{Y}_{8,114} = x_1 x_2^2 x_4^2 x_5^2, & \mathcal{Y}_{8,115} = x_1 x_2^2 x_4^2 x_5^2, & \mathcal{Y}_{8,116} = x_1 x_2^2 x_3^2 x_5^3, \\ \mathcal{Y}_{8,117} = x_1 x_2^2 x_3^2 x_5^2, & \mathcal{Y}_{8,118} = x_1 x_2^2 x_3^2 x_5^2, & \mathcal{Y}_{8,119} = x_1 x_2^2 x_3^2 x_4^3, & \mathcal{Y}_{8,120} = x_1 x_2^2 x_3^2 x_4^2, \end{array}$$

$$\begin{aligned}
\mathcal{V}_{8,121} &= x_1 x_2^3 x_3^2 x_4^2, & \mathcal{V}_{8,122} &= x_1^3 x_3 x_4^2 x_5^2, & \mathcal{V}_{8,123} &= x_1^3 x_2 x_4^2 x_5^2, & \mathcal{V}_{8,124} &= x_1^3 x_2 x_3^2 x_5^2, \\
\mathcal{V}_{8,125} &= x_1^3 x_2 x_3^2 x_4^2, & \mathcal{V}_{8,126} &= x_1 x_2 x_3^2 x_4^2 x_5^2, & \mathcal{V}_{8,127} &= x_1 x_2^2 x_3 x_4^2 x_5^2, & \mathcal{V}_{8,128} &= x_1 x_2^2 x_3^2 x_4 x_5^2, \\
\mathcal{V}_{8,129} &= x_1 x_2^2 x_3^2 x_4^2 x_5, & \mathcal{V}_{8,130} &= x_2 x_3 x_4^3 x_5^3, & \mathcal{V}_{8,131} &= x_2 x_3^3 x_4 x_5^3, & \mathcal{V}_{8,132} &= x_2 x_3^3 x_4^3 x_5, \\
\mathcal{V}_{8,133} &= x_2^3 x_3 x_4 x_5, & \mathcal{V}_{8,134} &= x_2^3 x_3 x_4 x_5^3, & \mathcal{V}_{8,135} &= x_2^3 x_3^3 x_4 x_5, & \mathcal{V}_{8,136} &= x_1 x_3 x_4^3 x_5^3, \\
\mathcal{V}_{8,137} &= x_1 x_3^3 x_4 x_5^3, & \mathcal{V}_{8,138} &= x_1 x_3^3 x_4^3 x_5, & \mathcal{V}_{8,139} &= x_1 x_2 x_4^3 x_5^3, & \mathcal{V}_{8,140} &= x_1 x_2 x_3^3 x_5^3, \\
\mathcal{V}_{8,141} &= x_1 x_2 x_3^3 x_4^3, & \mathcal{V}_{8,142} &= x_1 x_2^3 x_4 x_5^3, & \mathcal{V}_{8,143} &= x_1 x_2^3 x_4^3 x_5, & \mathcal{V}_{8,144} &= x_1 x_2^3 x_3 x_5^3, \\
\mathcal{V}_{8,145} &= x_1 x_2^3 x_3^3 x_4^3, & \mathcal{V}_{8,146} &= x_1 x_2^3 x_3^3 x_5, & \mathcal{V}_{8,147} &= x_1 x_2^3 x_3^3 x_4, & \mathcal{V}_{8,148} &= x_1^3 x_3 x_4 x_5^3, \\
\mathcal{V}_{8,149} &= x_1^3 x_3 x_4^3 x_5, & \mathcal{V}_{8,150} &= x_1^3 x_3^3 x_4 x_5, & \mathcal{V}_{8,151} &= x_1^3 x_2 x_4 x_5^3, & \mathcal{V}_{8,152} &= x_1^3 x_2 x_4^3 x_5, \\
\mathcal{V}_{8,153} &= x_1^3 x_2 x_3 x_5^3, & \mathcal{V}_{8,154} &= x_1^3 x_2 x_3 x_4^3, & \mathcal{V}_{8,155} &= x_1^3 x_2 x_3^3 x_5, & \mathcal{V}_{8,156} &= x_1^3 x_2 x_3^3 x_4, \\
\mathcal{V}_{8,157} &= x_1^3 x_2^3 x_4 x_5, & \mathcal{V}_{8,158} &= x_1^3 x_2^3 x_3 x_5, & \mathcal{V}_{8,159} &= x_1^3 x_2^3 x_3 x_4, & \mathcal{V}_{8,160} &= x_1 x_2 x_3 x_4^2 x_5^3, \\
\mathcal{V}_{8,161} &= x_1 x_2 x_3 x_4^3 x_5^2, & \mathcal{V}_{8,162} &= x_1 x_2 x_3^2 x_4 x_5^3, & \mathcal{V}_{8,163} &= x_1 x_2 x_3^2 x_4^3 x_5, & \mathcal{V}_{8,164} &= x_1 x_2 x_3^3 x_4 x_5^2, \\
\mathcal{V}_{8,165} &= x_1 x_2 x_3^2 x_4^3 x_5, & \mathcal{V}_{8,166} &= x_1 x_2^3 x_3 x_4 x_5^2, & \mathcal{V}_{8,167} &= x_1 x_2^3 x_3 x_4^2 x_5, & \mathcal{V}_{8,168} &= x_1 x_2^3 x_3^2 x_4 x_5, \\
\mathcal{V}_{8,169} &= x_1 x_2^2 x_3 x_4 x_5^3, & \mathcal{V}_{8,170} &= x_1 x_2^2 x_3 x_4^3 x_5, & \mathcal{V}_{8,171} &= x_1 x_2^2 x_3^3 x_4 x_5, & \mathcal{V}_{8,172} &= x_1^3 x_2 x_3 x_4 x_5^2, \\
\mathcal{V}_{8,173} &= x_1^3 x_2 x_3 x_4^2 x_5, & \mathcal{V}_{8,174} &= x_1^3 x_2 x_3^2 x_4 x_5.
\end{aligned}$$

6.2. $\mathcal{A}_2$ -generators for $\mathcal{P}_5^0$ in degree 21

Using the results in Sect.3.2.1, we have $\mathcal{B}_5^0(21) = \mathcal{B}_5^0(3, 3, 1, 1) \cup \mathcal{B}_5^0(3, 3, 3)$, where $\mathcal{B}_5^0(3, 3, 1, 1)$ is the set of 340 admissible monomials: $\mathcal{V}_{21,t}$, $1 \leq t \leq 340$

$$\begin{aligned}
\mathcal{V}_{21,1} &= x_3^3 x_4^3 x_5^{15}, & \mathcal{V}_{21,2} &= x_3^3 x_4^{15} x_5^3, & \mathcal{V}_{21,3} &= x_3^{15} x_4^3 x_5^3, & \mathcal{V}_{21,4} &= x_3^3 x_4^3 x_5^{15}, \\
\mathcal{V}_{21,5} &= x_2^3 x_4^{15} x_5^3, & \mathcal{V}_{21,6} &= x_2^3 x_3^3 x_5^{15}, & \mathcal{V}_{21,7} &= x_2^3 x_3^3 x_4^{15}, & \mathcal{V}_{21,8} &= x_2^3 x_3^{15} x_5^3, \\
\mathcal{V}_{21,9} &= x_2^3 x_3^{15} x_4^3, & \mathcal{V}_{21,10} &= x_2^{15} x_3^3 x_5^3, & \mathcal{V}_{21,11} &= x_2^{15} x_3^3 x_5^3, & \mathcal{V}_{21,12} &= x_2^{15} x_3^3 x_4^3, \\
\mathcal{V}_{21,13} &= x_3^3 x_4^3 x_5^{15}, & \mathcal{V}_{21,14} &= x_3^3 x_4^{15} x_5^3, & \mathcal{V}_{21,15} &= x_1^3 x_3^3 x_5^{15}, & \mathcal{V}_{21,16} &= x_1^3 x_3^3 x_4^{15}, \\
\mathcal{V}_{21,17} &= x_1^3 x_3^{15} x_5^3, & \mathcal{V}_{21,18} &= x_1^3 x_3^{15} x_4^3, & \mathcal{V}_{21,19} &= x_1^3 x_2^3 x_5^{15}, & \mathcal{V}_{21,20} &= x_1^3 x_2^3 x_4^{15}, \\
\mathcal{V}_{21,21} &= x_1^3 x_2^3 x_3^{15}, & \mathcal{V}_{21,22} &= x_1^3 x_2^{15} x_5^3, & \mathcal{V}_{21,23} &= x_1^3 x_2^{15} x_4^3, & \mathcal{V}_{21,24} &= x_1^3 x_2^{15} x_3^3, \\
\mathcal{V}_{21,25} &= x_1^{15} x_4^3 x_5^3, & \mathcal{V}_{21,26} &= x_1^{15} x_3^3 x_5^3, & \mathcal{V}_{21,27} &= x_1^{15} x_3^3 x_4^3, & \mathcal{V}_{21,28} &= x_1^{15} x_2^3 x_5^3, \\
\mathcal{V}_{21,29} &= x_1^{15} x_2^3 x_4^3, & \mathcal{V}_{21,30} &= x_1^{15} x_2^3 x_3^3, & \mathcal{V}_{21,31} &= x_3^3 x_4^7 x_5^{11}, & \mathcal{V}_{21,32} &= x_3^3 x_4^7 x_5^{11}, \\
\mathcal{V}_{21,33} &= x_3^7 x_4^{11} x_5^3, & \mathcal{V}_{21,34} &= x_2^3 x_4^7 x_5^{11}, & \mathcal{V}_{21,35} &= x_2^3 x_3^7 x_5^{11}, & \mathcal{V}_{21,36} &= x_2^3 x_3^7 x_4^{11}, \\
\mathcal{V}_{21,37} &= x_2^7 x_4^3 x_5^{11}, & \mathcal{V}_{21,38} &= x_2^7 x_4^{11} x_5^3, & \mathcal{V}_{21,39} &= x_2^7 x_3^3 x_5^{11}, & \mathcal{V}_{21,40} &= x_2^7 x_3^3 x_4^{11}, \\
\mathcal{V}_{21,41} &= x_2^7 x_3^{11} x_5^3, & \mathcal{V}_{21,42} &= x_2^7 x_3^{11} x_4^3, & \mathcal{V}_{21,43} &= x_1^3 x_4^7 x_5^{11}, & \mathcal{V}_{21,44} &= x_1^3 x_3^7 x_5^{11}, \\
\mathcal{V}_{21,45} &= x_1^3 x_3^7 x_4^{11}, & \mathcal{V}_{21,46} &= x_1^3 x_2^7 x_5^{11}, & \mathcal{V}_{21,47} &= x_1^3 x_2^7 x_4^{11}, & \mathcal{V}_{21,48} &= x_1^3 x_2^7 x_3^{11}, \\
\mathcal{V}_{21,49} &= x_1^7 x_4^3 x_5^{11}, & \mathcal{V}_{21,50} &= x_1^7 x_4^{11} x_5^3, & \mathcal{V}_{21,51} &= x_1^7 x_3^3 x_5^{11}, & \mathcal{V}_{21,52} &= x_1^7 x_3^3 x_4^{11}, \\
\mathcal{V}_{21,53} &= x_1^7 x_3^{11} x_5^3, & \mathcal{V}_{21,54} &= x_1^7 x_3^{11} x_4^3, & \mathcal{V}_{21,55} &= x_1^7 x_2^3 x_5^{11}, & \mathcal{V}_{21,56} &= x_1^7 x_2^3 x_4^{11}, \\
\mathcal{V}_{21,57} &= x_1^7 x_2^3 x_3^{11}, & \mathcal{V}_{21,58} &= x_1^7 x_2^{11} x_5^3, & \mathcal{V}_{21,59} &= x_1^7 x_2^{11} x_4^3, & \mathcal{V}_{21,60} &= x_1^7 x_2^{11} x_3^3, \\
\mathcal{V}_{21,61} &= x_2 x_3^2 x_4^3 x_5^{15}, & \mathcal{V}_{21,62} &= x_2 x_3^2 x_4^{15} x_5^3, & \mathcal{V}_{21,63} &= x_2 x_3^3 x_4^2 x_5^{15}, & \mathcal{V}_{21,64} &= x_2 x_3^3 x_4^{15} x_5^2, \\
\mathcal{V}_{21,65} &= x_2 x_3^{15} x_4^2 x_5^3, & \mathcal{V}_{21,66} &= x_2 x_3^{15} x_4^3 x_5^2, & \mathcal{V}_{21,67} &= x_2^3 x_3 x_4^2 x_5^{15}, & \mathcal{V}_{21,68} &= x_2^3 x_3 x_4^{15} x_5^2, \\
\mathcal{V}_{21,69} &= x_2^3 x_3^{15} x_4 x_5^2, & \mathcal{V}_{21,70} &= x_2^{15} x_3 x_4^2 x_5^3, & \mathcal{V}_{21,71} &= x_2^{15} x_3 x_4^3 x_5^2, & \mathcal{V}_{21,72} &= x_2^{15} x_3^3 x_4 x_5^2, \\
\mathcal{V}_{21,73} &= x_1 x_3^2 x_4^3 x_5^{15}, & \mathcal{V}_{21,74} &= x_1 x_3^2 x_4^{15} x_5^3, & \mathcal{V}_{21,75} &= x_1 x_3^3 x_4^2 x_5^{15}, & \mathcal{V}_{21,76} &= x_1 x_3^3 x_4^{15} x_5^2, \\
\mathcal{V}_{21,77} &= x_1 x_3^{15} x_4^2 x_5^3, & \mathcal{V}_{21,78} &= x_1 x_3^{15} x_4^3 x_5^2, & \mathcal{V}_{21,79} &= x_1 x_2^2 x_4^3 x_5^{15}, & \mathcal{V}_{21,80} &= x_1 x_2^2 x_4^{15} x_5^3, \\
\mathcal{V}_{21,81} &= x_1 x_2^2 x_3^3 x_5^{15}, & \mathcal{V}_{21,82} &= x_1 x_2^2 x_3^3 x_4^{15}, & \mathcal{V}_{21,83} &= x_1 x_2^2 x_3^{15} x_5^3, & \mathcal{V}_{21,84} &= x_1 x_2^2 x_3^{15} x_4^3, \\
\mathcal{V}_{21,85} &= x_1 x_2^3 x_4^2 x_5^{15}, & \mathcal{V}_{21,86} &= x_1 x_2^3 x_4^{15} x_5^2, & \mathcal{V}_{21,87} &= x_1 x_2^3 x_3^2 x_5^{15}, & \mathcal{V}_{21,88} &= x_1 x_2^3 x_3^2 x_4^{15}, \\
\mathcal{V}_{21,89} &= x_1 x_2^3 x_3^{15} x_5^2, & \mathcal{V}_{21,90} &= x_1 x_2^3 x_3^{15} x_4^2, & \mathcal{V}_{21,91} &= x_1 x_2^{15} x_4^2 x_5^3, & \mathcal{V}_{21,92} &= x_1 x_2^{15} x_3^2 x_5^2, \\
\mathcal{V}_{21,93} &= x_1 x_2^{15} x_3^2 x_5^3, & \mathcal{V}_{21,94} &= x_1 x_2^{15} x_3^2 x_4^3, & \mathcal{V}_{21,95} &= x_1 x_2^{15} x_3^3 x_5^2, & \mathcal{V}_{21,96} &= x_1 x_2^{15} x_3^3 x_4^2, \\
\mathcal{V}_{21,97} &= x_1^3 x_3 x_4^2 x_5^{15}, & \mathcal{V}_{21,98} &= x_1^3 x_3 x_4^{15} x_5^2, & \mathcal{V}_{21,99} &= x_1^3 x_3^{15} x_4 x_5^2, & \mathcal{V}_{21,100} &= x_1^3 x_2 x_4^2 x_5^{15}, \\
\mathcal{V}_{21,101} &= x_1^3 x_2 x_4^{15} x_5^2, & \mathcal{V}_{21,102} &= x_1^3 x_2 x_3^2 x_5^{15}, & \mathcal{V}_{21,103} &= x_1^3 x_2 x_3^2 x_4^{15}, & \mathcal{V}_{21,104} &= x_1^3 x_2 x_3^{15} x_5^2, \\
\mathcal{V}_{21,105} &= x_1^3 x_2^3 x_3^{15} x_5^2, & \mathcal{V}_{21,106} &= x_1^3 x_2^3 x_3^{15} x_4^2, & \mathcal{V}_{21,107} &= x_1^3 x_2^{15} x_3 x_5^2, & \mathcal{V}_{21,108} &= x_1^3 x_2^{15} x_3 x_4^2, \\
\mathcal{V}_{21,109} &= x_1^{15} x_3 x_4^2 x_5^3, & \mathcal{V}_{21,110} &= x_1^{15} x_3 x_4^2 x_5^2, & \mathcal{V}_{21,111} &= x_1^{15} x_3^3 x_4 x_5^2, & \mathcal{V}_{21,112} &= x_1^{15} x_2 x_4^2 x_5^3, \\
\mathcal{V}_{21,113} &= x_1^{15} x_2 x_4^2 x_5^2, & \mathcal{V}_{21,114} &= x_1^{15} x_2 x_3^2 x_5^3, & \mathcal{V}_{21,115} &= x_1^{15} x_2 x_3^2 x_4^3, & \mathcal{V}_{21,116} &= x_1^{15} x_2 x_3^2 x_5^2, \\
\mathcal{V}_{21,117} &= x_1^{15} x_2 x_3^2 x_4^2, & \mathcal{V}_{21,118} &= x_1^{15} x_2^3 x_4 x_5^2, & \mathcal{V}_{21,119} &= x_1^{15} x_2^3 x_3 x_5^2, & \mathcal{V}_{21,120} &= x_1^{15} x_2^3 x_3 x_4^2,
\end{aligned}$$

42

$$\begin{aligned}
\mathcal{V}_{21,321} &= x_1^3 x_3^3 x_4^7 x_5^8, & \mathcal{V}_{21,322} &= x_1^3 x_3^7 x_4^3 x_5^8, & \mathcal{V}_{21,323} &= x_1^3 x_3^7 x_4^8 x_5^3, & \mathcal{V}_{21,324} &= x_1^3 x_2^3 x_4^7 x_5^8, \\
\mathcal{V}_{21,325} &= x_1^3 x_2^3 x_4^7 x_5^8, & \mathcal{V}_{21,326} &= x_1^3 x_2^3 x_3^7 x_4^8, & \mathcal{V}_{21,327} &= x_1^3 x_2^7 x_3^3 x_4^8, & \mathcal{V}_{21,328} &= x_1^3 x_2^7 x_4^8 x_5^3, \\
\mathcal{V}_{21,329} &= x_1^3 x_2^7 x_3^3 x_4^8, & \mathcal{V}_{21,330} &= x_1^3 x_2^7 x_3^3 x_4^8, & \mathcal{V}_{21,331} &= x_1^3 x_2^7 x_3^8 x_4^3, & \mathcal{V}_{21,332} &= x_1^3 x_2^7 x_3^8 x_4^3, \\
\mathcal{V}_{21,333} &= x_1^7 x_3^3 x_4^7 x_5^8, & \mathcal{V}_{21,334} &= x_1^7 x_3^3 x_4^7 x_5^8, & \mathcal{V}_{21,335} &= x_1^7 x_2^3 x_3^3 x_4^8, & \mathcal{V}_{21,336} &= x_1^7 x_2^3 x_4^8 x_5^3, \\
\mathcal{V}_{21,337} &= x_1^7 x_2^3 x_3^3 x_4^8, & \mathcal{V}_{21,338} &= x_1^7 x_2^3 x_3^3 x_4^8, & \mathcal{V}_{21,339} &= x_1^7 x_2^3 x_3^8 x_4^3, & \mathcal{V}_{21,340} &= x_1^7 x_2^3 x_3^8 x_4^3,
\end{aligned}$$

$\mathcal{B}_5^0(3, 3, 3)$ is the set of 60 admissible monomials: $\mathcal{V}_{21,t}$, $341 \leq t \leq 400$

$$\begin{aligned}
\mathcal{V}_{21,341} &= x_3^7 x_4^7 x_5^7, & \mathcal{V}_{21,342} &= x_2^7 x_4^7 x_5^7, & \mathcal{V}_{21,343} &= x_2^7 x_3^7 x_4^7, & \mathcal{V}_{21,344} &= x_2^7 x_3^7 x_4^7, \\
\mathcal{V}_{21,345} &= x_1^7 x_4^7 x_5^7, & \mathcal{V}_{21,346} &= x_1^7 x_3^7 x_4^7, & \mathcal{V}_{21,347} &= x_1^7 x_3^7 x_4^7, & \mathcal{V}_{21,348} &= x_1^7 x_2^7 x_4^7, \\
\mathcal{V}_{21,349} &= x_1^7 x_2^7 x_4^7, & \mathcal{V}_{21,350} &= x_1^7 x_2^7 x_3^7, & \mathcal{V}_{21,351} &= x_2^6 x_3^6 x_4^7 x_5^7, & \mathcal{V}_{21,352} &= x_2^6 x_3^6 x_4^7 x_5^7, \\
\mathcal{V}_{21,353} &= x_2^6 x_3^6 x_4^7 x_5^7, & \mathcal{V}_{21,354} &= x_2^6 x_3^6 x_4^7 x_5^7, & \mathcal{V}_{21,355} &= x_2^6 x_3^6 x_4^7 x_5^7, & \mathcal{V}_{21,356} &= x_2^6 x_3^6 x_4^7 x_5^7, \\
\mathcal{V}_{21,357} &= x_1^6 x_3^6 x_4^7 x_5^7, & \mathcal{V}_{21,358} &= x_1^6 x_3^6 x_4^7 x_5^7, & \mathcal{V}_{21,359} &= x_1^6 x_3^6 x_4^7 x_5^7, & \mathcal{V}_{21,360} &= x_1^6 x_2^6 x_4^7 x_5^7, \\
\mathcal{V}_{21,361} &= x_1^6 x_2^6 x_3^7 x_4^7, & \mathcal{V}_{21,362} &= x_1^6 x_2^6 x_3^7 x_4^7, & \mathcal{V}_{21,363} &= x_1^6 x_2^6 x_3^7 x_4^7, & \mathcal{V}_{21,364} &= x_1^6 x_2^6 x_3^7 x_4^7, \\
\mathcal{V}_{21,365} &= x_1^6 x_2^6 x_3^7 x_4^7, & \mathcal{V}_{21,366} &= x_1^6 x_2^6 x_3^7 x_4^7, & \mathcal{V}_{21,367} &= x_1^6 x_2^6 x_3^7 x_4^7, & \mathcal{V}_{21,368} &= x_1^6 x_2^6 x_3^7 x_4^7, \\
\mathcal{V}_{21,369} &= x_1^6 x_2^6 x_3^7 x_4^7, & \mathcal{V}_{21,370} &= x_1^6 x_2^6 x_3^7 x_4^7, & \mathcal{V}_{21,371} &= x_1^6 x_2^6 x_3^7 x_4^7, & \mathcal{V}_{21,372} &= x_1^6 x_2^6 x_3^7 x_4^7, \\
\mathcal{V}_{21,373} &= x_1^6 x_2^6 x_3^7 x_4^7, & \mathcal{V}_{21,374} &= x_1^6 x_2^6 x_3^7 x_4^7, & \mathcal{V}_{21,375} &= x_1^6 x_2^6 x_3^7 x_4^7, & \mathcal{V}_{21,376} &= x_1^6 x_2^6 x_3^7 x_4^7, \\
\mathcal{V}_{21,377} &= x_1^6 x_2^6 x_3^7 x_4^7, & \mathcal{V}_{21,378} &= x_1^6 x_2^6 x_3^7 x_4^7, & \mathcal{V}_{21,379} &= x_1^6 x_2^6 x_3^7 x_4^7, & \mathcal{V}_{21,380} &= x_1^6 x_2^6 x_3^7 x_4^7, \\
\mathcal{V}_{21,381} &= x_2^5 x_3^5 x_4^6 x_5^6, & \mathcal{V}_{21,382} &= x_2^5 x_3^5 x_4^6 x_5^6, & \mathcal{V}_{21,383} &= x_2^5 x_3^5 x_4^6 x_5^6, & \mathcal{V}_{21,384} &= x_2^5 x_3^5 x_4^6 x_5^6, \\
\mathcal{V}_{21,385} &= x_2^5 x_3^5 x_4^6 x_5^6, & \mathcal{V}_{21,386} &= x_2^5 x_3^5 x_4^6 x_5^6, & \mathcal{V}_{21,387} &= x_2^5 x_3^5 x_4^6 x_5^6, & \mathcal{V}_{21,388} &= x_2^5 x_3^5 x_4^6 x_5^6, \\
\mathcal{V}_{21,389} &= x_2^5 x_3^5 x_4^6 x_5^6, & \mathcal{V}_{21,390} &= x_2^5 x_3^5 x_4^6 x_5^6, & \mathcal{V}_{21,391} &= x_2^5 x_3^5 x_4^6 x_5^6, & \mathcal{V}_{21,392} &= x_2^5 x_3^5 x_4^6 x_5^6, \\
\mathcal{V}_{21,393} &= x_2^5 x_3^5 x_4^6 x_5^6, & \mathcal{V}_{21,394} &= x_2^5 x_3^5 x_4^6 x_5^6, & \mathcal{V}_{21,395} &= x_2^5 x_3^5 x_4^6 x_5^6, & \mathcal{V}_{21,396} &= x_2^5 x_3^5 x_4^6 x_5^6, \\
\mathcal{V}_{21,397} &= x_2^5 x_3^5 x_4^6 x_5^6, & \mathcal{V}_{21,398} &= x_2^5 x_3^5 x_4^6 x_5^6, & \mathcal{V}_{21,399} &= x_2^5 x_3^5 x_4^6 x_5^6, & \mathcal{V}_{21,400} &= x_2^5 x_3^5 x_4^6 x_5^6.
\end{aligned}$$

6.3. $\mathcal{A}_2$ -generators for $\mathcal{P}_5^+$ in degree 21

Note that Kameko's squaring operation $\widetilde{Sq}_*^0(5,21) : (Q\mathcal{P}_5)_{21} \rightarrow (Q\mathcal{P}_5)_8$ is an epimorphism of $\mathbb{Z}/2GL_5$ -modules. Hence, we have

$$\mathcal{B}_5(21) = \mathcal{B}_5^0(21) \cup \varphi(\mathcal{B}_5(8)) \cup (\mathcal{B}_5^+(21) \cap \text{Ker}((\widetilde{Sq}_*^0(5,21))),$$

where $\mathcal{B}_5^0(21) = \overline{\Phi}^0(\mathcal{B}_4(21))$, $|\mathcal{B}_5^0(21)| = 400$, $|\varphi(\mathcal{B}_5(8))| = 174$ with

$$\varphi : \mathcal{P}_5 \rightarrow \mathcal{P}_5, \quad \varphi(Z) = X_{(\emptyset,5)} Z^2, \quad \forall Z \in \mathcal{P}_5.$$

Based on the results in Sect.3.2.1, we see that

$$\mathcal{B}_5^+(21) \cap \text{Ker}((\widetilde{Sq}_*^0(5,21))) = \mathcal{B}_5^+(3, 3, 1, 1) \cup \mathcal{B}_5^+(3, 3, 3),$$

where $\mathcal{B}_5^+(3, 3, 1, 1)$ is the set of 196 admissible monomials: $\mathcal{V}_{21,t}$, $401 \leq t \leq 596$

$$\begin{aligned}
\mathcal{V}_{21,401} &= x_1 x_2^2 x_3 x_4^2 x_5^{15}, & \mathcal{V}_{21,402} &= x_1 x_2^2 x_3 x_4^2 x_5^{15}, & \mathcal{V}_{21,403} &= x_1 x_2^2 x_3^{15} x_4^2 x_5^2, & \mathcal{V}_{21,404} &= x_1 x_2 x_3^2 x_4^2 x_5^{15}, \\
\mathcal{V}_{21,405} &= x_1 x_2 x_3^2 x_4^2 x_5^{15}, & \mathcal{V}_{21,406} &= x_1 x_2^{15} x_3^2 x_4^2 x_5^2, & \mathcal{V}_{21,407} &= x_1 x_2 x_3^{15} x_4^2 x_5^2, & \mathcal{V}_{21,408} &= x_1 x_2^{15} x_3 x_4^2 x_5^2, \\
\mathcal{V}_{21,409} &= x_1^{15} x_2 x_3^2 x_4^2 x_5^2, & \mathcal{V}_{21,410} &= x_1^{15} x_2 x_3^2 x_4^2 x_5^2, & \mathcal{V}_{21,411} &= x_1 x_2 x_3^6 x_4^3 x_5^{10}, & \mathcal{V}_{21,412} &= x_1 x_2 x_3^6 x_4^3 x_5^{10}, \\
\mathcal{V}_{21,413} &= x_1 x_2 x_3^6 x_4^3 x_5^{10}, & \mathcal{V}_{21,414} &= x_1 x_2^2 x_3^3 x_4^3 x_5^{12}, & \mathcal{V}_{21,415} &= x_1 x_2^2 x_3^3 x_4^3 x_5^{12}, & \mathcal{V}_{21,416} &= x_1 x_2^2 x_3^3 x_4^3 x_5^{12}, \\
\mathcal{V}_{21,417} &= x_1^3 x_2^3 x_3^4 x_4^3 x_5^8, & \mathcal{V}_{21,418} &= x_1^3 x_2^3 x_3^4 x_4^3 x_5^8, & \mathcal{V}_{21,419} &= x_1^3 x_2^3 x_3^4 x_4^3 x_5^8, & \mathcal{V}_{21,420} &= x_1 x_2^2 x_3^3 x_4^3 x_5^{14}, \\
\mathcal{V}_{21,421} &= x_1 x_2^2 x_3^3 x_4^3 x_5^{14}, & \mathcal{V}_{21,422} &= x_1 x_2^2 x_3^3 x_4^3 x_5^{14}, & \mathcal{V}_{21,423} &= x_1 x_2^{14} x_3 x_4^3 x_5^3, & \mathcal{V}_{21,424} &= x_1 x_2^{14} x_3 x_4^3 x_5^3, \\
\mathcal{V}_{21,425} &= x_1 x_2^{14} x_3 x_4^3 x_5^3, & \mathcal{V}_{21,426} &= x_1 x_2 x_3^2 x_4^3 x_5^{14}, & \mathcal{V}_{21,427} &= x_1 x_2 x_3^2 x_4^3 x_5^{14}, & \mathcal{V}_{21,428} &= x_1 x_2 x_3^2 x_4^3 x_5^{14}, \\
\mathcal{V}_{21,429} &= x_1 x_2 x_3^2 x_4^3 x_5^{14}, & \mathcal{V}_{21,430} &= x_1 x_2^3 x_3^2 x_4^3 x_5^{14}, & \mathcal{V}_{21,431} &= x_1 x_2^3 x_3^2 x_4^3 x_5^{14}, & \mathcal{V}_{21,432} &= x_1 x_2^3 x_3^2 x_4^3 x_5^{14}, \\
\mathcal{V}_{21,433} &= x_1 x_2 x_3^3 x_4^3 x_5^{14}, & \mathcal{V}_{21,434} &= x_1 x_2^3 x_3 x_4^3 x_5^{14}, & \mathcal{V}_{21,435} &= x_1 x_2^3 x_3 x_4^3 x_5^{14}, & \mathcal{V}_{21,436} &= x_1 x_2^3 x_3 x_4^3 x_5^{14}, \\
\mathcal{V}_{21,437} &= x_1 x_2^6 x_3 x_4^3 x_5^{11}, & \mathcal{V}_{21,438} &= x_1 x_2^6 x_3 x_4^3 x_5^{11}, & \mathcal{V}_{21,439} &= x_1 x_2^6 x_3^{11} x_4^3 x_5^2, & \mathcal{V}_{21,440} &= x_1 x_2 x_3^2 x_4^3 x_5^{11}, \\
\mathcal{V}_{21,441} &= x_1 x_2 x_3^6 x_4^3 x_5^{11}, & \mathcal{V}_{21,442} &= x_1 x_2 x_3^6 x_4^3 x_5^{11}, & \mathcal{V}_{21,443} &= x_1 x_2^2 x_3^3 x_4^3 x_5^{10}, & \mathcal{V}_{21,444} &= x_1 x_2^2 x_3^3 x_4^3 x_5^{10}, \\
\mathcal{V}_{21,445} &= x_1 x_2 x_3^3 x_4^3 x_5^{10}, & \mathcal{V}_{21,446} &= x_1 x_2^7 x_3^{10} x_4^3 x_5^2, & \mathcal{V}_{21,447} &= x_1 x_2^7 x_3^{10} x_4^3 x_5^2, & \mathcal{V}_{21,448} &= x_1 x_2^7 x_3^{10} x_4^3 x_5^2, \\
\mathcal{V}_{21,449} &= x_1 x_2 x_3^3 x_4^3 x_5^{10}, & \mathcal{V}_{21,450} &= x_1 x_2^7 x_3^3 x_4^3 x_5^{10}, & \mathcal{V}_{21,451} &= x_1 x_2^7 x_3^3 x_4^3 x_5^{10}, & \mathcal{V}_{21,452} &= x_1 x_2^7 x_3^3 x_4^3 x_5^{10}, \\
\mathcal{V}_{21,453} &= x_1 x_2^7 x_3^{13} x_4^3 x_5^2, & \mathcal{V}_{21,454} &= x_1 x_2^7 x_3^{13} x_4^3 x_5^2, & \mathcal{V}_{21,455} &= x_1 x_2^7 x_3^3 x_4^3 x_5^{10}, & \mathcal{V}_{21,456} &= x_1 x_2^7 x_3^3 x_4^3 x_5^{10},
\end{aligned}$$

$$\begin{aligned}
\mathcal{V}_{21,457} &= x_1 x_2^2 x_3^5 x_4^2 x_5^{11}, & \mathcal{V}_{21,458} &= x_1 x_2^2 x_3^5 x_4^{11} x_5^2, & \mathcal{V}_{21,459} &= x_1 x_2^2 x_3^7 x_4^9 x_5^2, & \mathcal{V}_{21,460} &= x_1 x_2^7 x_3^2 x_4^9 x_5^2, \\
\mathcal{V}_{21,461} &= x_1 x_2^7 x_3^9 x_4^2 x_5^2, & \mathcal{V}_{21,462} &= x_1 x_2^6 x_3 x_4^3 x_5^{10}, & \mathcal{V}_{21,463} &= x_1 x_2^6 x_3 x_4^{10} x_5^3, & \mathcal{V}_{21,464} &= x_1 x_2^6 x_3^2 x_4^{10} x_5^2, \\
\mathcal{V}_{21,465} &= x_1 x_2^3 x_3^6 x_4^2 x_5^{10}, & \mathcal{V}_{21,466} &= x_1 x_2^3 x_3^6 x_4^{10} x_5^2, & \mathcal{V}_{21,467} &= x_1 x_2^3 x_3^{12} x_4^2 x_5^3, & \mathcal{V}_{21,468} &= x_1 x_2^3 x_3^{12} x_4^3 x_5^2, \\
\mathcal{V}_{21,469} &= x_1 x_2^3 x_3^2 x_4^3 x_5^{12}, & \mathcal{V}_{21,470} &= x_1 x_2^3 x_3^2 x_4^{12} x_5^3, & \mathcal{V}_{21,471} &= x_1 x_2^3 x_3^3 x_4^{12} x_5^2, & \mathcal{V}_{21,472} &= x_1 x_2^3 x_3^3 x_4^2 x_5^{12}, \\
\mathcal{V}_{21,473} &= x_1 x_2^3 x_3^3 x_4^4 x_5^{11}, & \mathcal{V}_{21,474} &= x_1 x_2^3 x_3^4 x_4^2 x_5^{11}, & \mathcal{V}_{21,475} &= x_1 x_2^3 x_3^4 x_4^{11} x_5^2, & \mathcal{V}_{21,476} &= x_1 x_2^3 x_3^3 x_4^4 x_5^{11}, \\
\mathcal{V}_{21,477} &= x_1 x_2^2 x_3^4 x_4^3 x_5^{11}, & \mathcal{V}_{21,478} &= x_1 x_2^2 x_3^4 x_4^{11} x_5^3, & \mathcal{V}_{21,479} &= x_1 x_2^2 x_3^3 x_4^5 x_5^{10}, & \mathcal{V}_{21,480} &= x_1 x_2^2 x_3^5 x_4^3 x_5^{10}, \\
\mathcal{V}_{21,481} &= x_1 x_2^2 x_3^5 x_4^{10} x_5^3, & \mathcal{V}_{21,482} &= x_1 x_2^2 x_3^5 x_4^5 x_5^{10}, & \mathcal{V}_{21,483} &= x_1 x_2^2 x_3^5 x_4^{10} x_5^2, & \mathcal{V}_{21,484} &= x_1 x_2^2 x_3^5 x_4^2 x_5^{10}, \\
\mathcal{V}_{21,485} &= x_1 x_2^6 x_3^3 x_4^9 x_5^2, & \mathcal{V}_{21,486} &= x_1 x_2^6 x_3^9 x_4^2 x_5^3, & \mathcal{V}_{21,487} &= x_1 x_2^6 x_3^9 x_4^3 x_5^2, & \mathcal{V}_{21,488} &= x_1 x_2^6 x_3^6 x_4^9 x_5^2, \\
\mathcal{V}_{21,489} &= x_1 x_2^2 x_3^3 x_4^7 x_5^8, & \mathcal{V}_{21,490} &= x_1 x_2^2 x_3^7 x_4^3 x_5^8, & \mathcal{V}_{21,491} &= x_1 x_2^2 x_3^7 x_4^8 x_5^3, & \mathcal{V}_{21,492} &= x_1 x_2^3 x_3^2 x_4^7 x_5^8, \\
\mathcal{V}_{21,493} &= x_1 x_2^7 x_3^8 x_4^2 x_5^3, & \mathcal{V}_{21,494} &= x_1 x_2^7 x_3^8 x_4^3 x_5^2, & \mathcal{V}_{21,495} &= x_1 x_2^7 x_3^2 x_4^3 x_5^8, & \mathcal{V}_{21,496} &= x_1 x_2^7 x_3^2 x_4^8 x_5^3, \\
\mathcal{V}_{21,497} &= x_1 x_2^3 x_3^8 x_4^2 x_5^2, & \mathcal{V}_{21,498} &= x_1 x_2^3 x_3^7 x_4^2 x_5^8, & \mathcal{V}_{21,499} &= x_1 x_2^7 x_3^3 x_4^8 x_5^2, & \mathcal{V}_{21,500} &= x_1 x_2^7 x_3^3 x_4^2 x_5^8, \\
\mathcal{V}_{21,501} &= x_1 x_2^3 x_3^3 x_4^4 x_5^{10}, & \mathcal{V}_{21,502} &= x_1 x_2^3 x_3^4 x_4^{10} x_5^3, & \mathcal{V}_{21,503} &= x_1 x_2^3 x_3^3 x_4^4 x_5^{10}, & \mathcal{V}_{21,504} &= x_1 x_2^6 x_3^3 x_4^3 x_5^8, \\
\mathcal{V}_{21,505} &= x_1 x_2^6 x_3^3 x_4^3 x_5^3, & \mathcal{V}_{21,506} &= x_1 x_2^3 x_3^6 x_4^3 x_5^8, & \mathcal{V}_{21,507} &= x_1 x_2^3 x_3^3 x_4^8 x_5^3, & \mathcal{V}_{21,508} &= x_1 x_2^3 x_3^3 x_4^4 x_5^8, \\
\mathcal{V}_{21,509} &= x_1^3 x_2^2 x_3^5 x_4 x_5^{14}, & \mathcal{V}_{21,510} &= x_1^3 x_2 x_3^{14} x_4 x_5^2, & \mathcal{V}_{21,511} &= x_1^3 x_2 x_3 x_4^2 x_5^{14}, & \mathcal{V}_{21,512} &= x_1^3 x_2 x_3 x_4^{14} x_5^2, \\
\mathcal{V}_{21,513} &= x_1^3 x_2 x_3^2 x_4^{13} x_5^2, & \mathcal{V}_{21,514} &= x_1^3 x_2 x_3^{13} x_4^2 x_5^2, & \mathcal{V}_{21,515} &= x_1^3 x_2^{13} x_3 x_4^2 x_5^2, & \mathcal{V}_{21,516} &= x_1^3 x_2^{13} x_3^2 x_4 x_5^2, \\
\mathcal{V}_{21,517} &= x_1^3 x_2 x_3^6 x_4 x_5^{10}, & \mathcal{V}_{21,518} &= x_1^3 x_2 x_3^6 x_4^{10} x_5^2, & \mathcal{V}_{21,519} &= x_1^3 x_2 x_3^{12} x_4^2 x_5^3, & \mathcal{V}_{21,520} &= x_1^3 x_2 x_3^{12} x_4^3 x_5^2, \\
\mathcal{V}_{21,521} &= x_1^3 x_2 x_3^2 x_4^3 x_5^{12}, & \mathcal{V}_{21,522} &= x_1^3 x_2 x_3^2 x_4^{12} x_5^3, & \mathcal{V}_{21,523} &= x_1^3 x_2 x_3^3 x_4^{12} x_5^2, & \mathcal{V}_{21,524} &= x_1^3 x_2 x_3^3 x_4^2 x_5^{12}, \\
\mathcal{V}_{21,525} &= x_1^3 x_2^3 x_3^2 x_4^{12} x_5^2, & \mathcal{V}_{21,526} &= x_1^3 x_2^3 x_3^2 x_4^2 x_5^{12}, & \mathcal{V}_{21,527} &= x_1^3 x_2^3 x_3^{12} x_4 x_5^2, & \mathcal{V}_{21,528} &= x_1^3 x_2^{12} x_3 x_4^2 x_5^3, \\
\mathcal{V}_{21,529} &= x_1^3 x_2^{12} x_3 x_4^2 x_5^{11}, & \mathcal{V}_{21,530} &= x_1^3 x_2^{12} x_3^3 x_4^2 x_5^2, & \mathcal{V}_{21,531} &= x_1^3 x_2 x_3^4 x_4^2 x_5^{11}, & \mathcal{V}_{21,532} &= x_1^3 x_2 x_3^4 x_4^{11} x_5^2, \\
\mathcal{V}_{21,533} &= x_1^3 x_2^2 x_3^3 x_4^4 x_5^{11}, & \mathcal{V}_{21,534} &= x_1^3 x_2^2 x_3^3 x_4^4 x_5^{11}, & \mathcal{V}_{21,535} &= x_1^3 x_2^4 x_3 x_4^{11} x_5^2, & \mathcal{V}_{21,536} &= x_1^3 x_2^4 x_3^3 x_4 x_5^2, \\
\mathcal{V}_{21,537} &= x_1^3 x_2^2 x_3^5 x_4^2 x_5^{10}, & \mathcal{V}_{21,538} &= x_1^3 x_2^2 x_3^5 x_4^2 x_5^{10}, & \mathcal{V}_{21,539} &= x_1^3 x_2^2 x_3^5 x_4^2 x_5^{10}, & \mathcal{V}_{21,540} &= x_1^3 x_2^5 x_3 x_4^{10} x_5^2, \\
\mathcal{V}_{21,541} &= x_1^3 x_2^5 x_3 x_4^2 x_5^{10}, & \mathcal{V}_{21,542} &= x_1^3 x_2^5 x_3^2 x_4 x_5^{10}, & \mathcal{V}_{21,543} &= x_1^3 x_2^5 x_3^{10} x_4 x_5^2, & \mathcal{V}_{21,544} &= x_1^3 x_2^6 x_3^9 x_4 x_5^2, \\
\mathcal{V}_{21,545} &= x_1^3 x_2^2 x_3^7 x_4^2 x_5^8, & \mathcal{V}_{21,546} &= x_1^3 x_2^2 x_3^7 x_4^8 x_5^2, & \mathcal{V}_{21,547} &= x_1^3 x_2^7 x_3^2 x_4^8 x_5^2, & \mathcal{V}_{21,548} &= x_1^3 x_2^7 x_3^2 x_4^8 x_5^2, \\
\mathcal{V}_{21,549} &= x_1^3 x_2^7 x_3^2 x_4^8 x_5^2, & \mathcal{V}_{21,550} &= x_1^3 x_2^7 x_3^8 x_4^2 x_5^2, & \mathcal{V}_{21,551} &= x_1^3 x_2^2 x_3^4 x_4^3 x_5^{10}, & \mathcal{V}_{21,552} &= x_1^3 x_2^2 x_3^4 x_4^{10} x_5^3, \\
\mathcal{V}_{21,553} &= x_1^3 x_2^3 x_3^3 x_4^4 x_5^{10}, & \mathcal{V}_{21,554} &= x_1^3 x_2^3 x_3^3 x_4^4 x_5^{10}, & \mathcal{V}_{21,555} &= x_1^3 x_2^3 x_3^4 x_4 x_5^{10}, & \mathcal{V}_{21,556} &= x_1^3 x_2^4 x_3 x_4^3 x_5^{10}, \\
\mathcal{V}_{21,557} &= x_1^3 x_2^2 x_3^4 x_4^{10} x_5^3, & \mathcal{V}_{21,558} &= x_1^3 x_2^4 x_3^3 x_4 x_5^{10}, & \mathcal{V}_{21,559} &= x_1^3 x_2^2 x_3^6 x_4^3 x_5^8, & \mathcal{V}_{21,560} &= x_1^3 x_2^2 x_3^6 x_4^3 x_5^8, \\
\mathcal{V}_{21,561} &= x_1^3 x_2^2 x_3^3 x_4^6 x_5^8, & \mathcal{V}_{21,562} &= x_1^3 x_2^3 x_3^6 x_4^6 x_5^8, & \mathcal{V}_{21,563} &= x_1^3 x_2^5 x_3^2 x_4^9 x_5^2, & \mathcal{V}_{21,564} &= x_1^3 x_2^5 x_3^2 x_4^9 x_5^2, \\
\mathcal{V}_{21,565} &= x_1^3 x_2^3 x_3^4 x_4^9 x_5^2, & \mathcal{V}_{21,566} &= x_1^3 x_2^4 x_3^3 x_4^9 x_5^2, & \mathcal{V}_{21,567} &= x_1^3 x_2^4 x_3^9 x_4^2 x_5^3, & \mathcal{V}_{21,568} &= x_1^3 x_2^4 x_3^9 x_4^2 x_5^3, \\
\mathcal{V}_{21,569} &= x_1^3 x_2^5 x_3^8 x_4^2 x_5^3, & \mathcal{V}_{21,570} &= x_1^3 x_2^5 x_3^8 x_4^2 x_5^3, & \mathcal{V}_{21,571} &= x_1^3 x_2^5 x_3^2 x_4^3 x_5^8, & \mathcal{V}_{21,572} &= x_1^3 x_2^5 x_3^2 x_4^3 x_5^8, \\
\mathcal{V}_{21,573} &= x_1^3 x_2^5 x_3^3 x_4^8 x_5^2, & \mathcal{V}_{21,574} &= x_1^3 x_2^5 x_3^3 x_4^8 x_5^2, & \mathcal{V}_{21,575} &= x_1^3 x_2^3 x_3^5 x_4^8 x_5^2, & \mathcal{V}_{21,576} &= x_1^3 x_2^3 x_3^5 x_4^8 x_5^2, \\
\mathcal{V}_{21,577} &= x_1^7 x_2 x_3^{10} x_4 x_5^2, & \mathcal{V}_{21,578} &= x_1^7 x_2 x_3^2 x_4 x_5^{10}, & \mathcal{V}_{21,579} &= x_1^7 x_2 x_3 x_4^{10} x_5^2, & \mathcal{V}_{21,580} &= x_1^7 x_2 x_3 x_4^{10} x_5^2, \\
\mathcal{V}_{21,581} &= x_1^7 x_2 x_3^2 x_4^9 x_5^2, & \mathcal{V}_{21,582} &= x_1^7 x_2 x_3^9 x_4^2 x_5^2, & \mathcal{V}_{21,583} &= x_1^7 x_2^9 x_3 x_4^2 x_5^2, & \mathcal{V}_{21,584} &= x_1^7 x_2^9 x_3^2 x_4 x_5^2, \\
\mathcal{V}_{21,585} &= x_1^7 x_2 x_3^8 x_4^3 x_5^2, & \mathcal{V}_{21,586} &= x_1^7 x_2 x_3^8 x_4^3 x_5^2, & \mathcal{V}_{21,587} &= x_1^7 x_2 x_3^2 x_4^3 x_5^8, & \mathcal{V}_{21,588} &= x_1^7 x_2 x_3^2 x_4^3 x_5^8, \\
\mathcal{V}_{21,589} &= x_1^7 x_2 x_3^3 x_4^8 x_5^2, & \mathcal{V}_{21,590} &= x_1^7 x_2 x_3^3 x_4^8 x_5^2, & \mathcal{V}_{21,591} &= x_1^7 x_2^3 x_3 x_4^8 x_5^2, & \mathcal{V}_{21,592} &= x_1^7 x_2^3 x_3 x_4^8 x_5^2, \\
\mathcal{V}_{21,593} &= x_1^7 x_2^3 x_3^8 x_4 x_5^2, & \mathcal{V}_{21,594} &= x_1^7 x_2^3 x_3^8 x_4 x_5^2, & \mathcal{V}_{21,595} &= x_1^7 x_2^8 x_3 x_4^3 x_5^2, & \mathcal{V}_{21,596} &= x_1^7 x_2^8 x_3 x_4^3 x_5^2,
\end{aligned}$$

$\mathcal{B}_5^+(3, 3, 3)$ is the set of 70 admissible monomials: $\mathcal{V}_{21,t}$, $597 \leq t \leq 666$

$$\begin{aligned}
\mathcal{V}_{21,597} &= x_1 x_2^2 x_3^4 x_4^7 x_5^7, & \mathcal{V}_{21,598} &= x_1 x_2^2 x_3^7 x_4^4 x_5^7, & \mathcal{V}_{21,599} &= x_1 x_2^2 x_3^7 x_4^4 x_5^7, & \mathcal{V}_{21,600} &= x_1 x_2^7 x_3^2 x_4^4 x_5^7, \\
\mathcal{V}_{21,601} &= x_1 x_2^7 x_3^2 x_4^4 x_5^7, & \mathcal{V}_{21,602} &= x_1 x_2^7 x_3^2 x_4^4 x_5^7, & \mathcal{V}_{21,603} &= x_1^7 x_2 x_3^2 x_4^4 x_5^7, & \mathcal{V}_{21,604} &= x_1^7 x_2 x_3^2 x_4^4 x_5^7, \\
\mathcal{V}_{21,605} &= x_1^7 x_2 x_3^7 x_4^4 x_5^7, & \mathcal{V}_{21,606} &= x_1^7 x_2 x_3^7 x_4^4 x_5^7, & \mathcal{V}_{21,607} &= x_1 x_2^6 x_3 x_4^6 x_5^7, & \mathcal{V}_{21,608} &= x_1 x_2^6 x_3 x_4^6 x_5^7, \\
\mathcal{V}_{21,609} &= x_1 x_2^6 x_3^6 x_4^6 x_5^7, & \mathcal{V}_{21,610} &= x_1 x_2^6 x_3^6 x_4^6 x_5^7, & \mathcal{V}_{21,611} &= x_1 x_2^6 x_3^6 x_4^6 x_5^7, & \mathcal{V}_{21,612} &= x_1 x_2^6 x_3^6 x_4^6 x_5^7, \\
\mathcal{V}_{21,613} &= x_1 x_2^6 x_3^7 x_4^6 x_5^7, & \mathcal{V}_{21,614} &= x_1 x_2^7 x_3 x_4^6 x_5^7, & \mathcal{V}_{21,615} &= x_1^7 x_2 x_3^6 x_4^6 x_5^7, & \mathcal{V}_{21,616} &= x_1^7 x_2 x_3^6 x_4^6 x_5^7, \\
\mathcal{V}_{21,617} &= x_1 x_2^2 x_3^5 x_4^6 x_5^7, & \mathcal{V}_{21,618} &= x_1 x_2^2 x_3^5 x_4^6 x_5^7, & \mathcal{V}_{21,619} &= x_1 x_2^2 x_3^7 x_4^5 x_5^6, & \mathcal{V}_{21,620} &= x_1 x_2^7 x_3^2 x_4^5 x_5^6, \\
\mathcal{V}_{21,621} &= x_1^7 x_2 x_3^2 x_4^5 x_5^6, & \mathcal{V}_{21,622} &= x_1 x_2^3 x_4^6 x_4^7 x_5^6, & \mathcal{V}_{21,623} &= x_1 x_2^3 x_4^6 x_4^7 x_5^6, & \mathcal{V}_{21,624} &= x_1 x_2^3 x_4^6 x_4^7 x_5^6, \\
\mathcal{V}_{21,625} &= x_1 x_2^7 x_3^3 x_4^6 x_5^6, & \mathcal{V}_{21,626} &= x_1^3 x_2 x_3^4 x_4^6 x_5^7, & \mathcal{V}_{21,627} &= x_1^3 x_2 x_3^4 x_4^6 x_5^7, & \mathcal{V}_{21,628} &= x_1^3 x_2 x_3^7 x_4^4 x_5^6, \\
\mathcal{V}_{21,629} &= x_1^3 x_2 x_3^7 x_4^4 x_5^6, & \mathcal{V}_{21,630} &= x_1^7 x_2 x_3^3 x_4^6 x_5^6, & \mathcal{V}_{21,631} &= x_1^7 x_2 x_3^3 x_4^6 x_5^6, & \mathcal{V}_{21,632} &= x_1 x_2^3 x_3^6 x_4^4 x_5^7, \\
\mathcal{V}_{21,633} &= x_1 x_2^3 x_3^6 x_4^4 x_5^7, & \mathcal{V}_{21,634} &= x_1 x_2^3 x_3^6 x_4^4 x_5^7, & \mathcal{V}_{21,635} &= x_1 x_2^3 x_3^6 x_4^4 x_5^7, & \mathcal{V}_{21,636} &= x_1^3 x_2 x_3^6 x_4^4 x_5^7, \\
\mathcal{V}_{21,637} &= x_1^3 x_2 x_3^6 x_4^4 x_5^7, & \mathcal{V}_{21,638} &= x_1^3 x_2 x_3^6 x_4^4 x_5^7, & \mathcal{V}_{21,639} &= x_1^3 x_2 x_3^6 x_4^4 x_5^7, & \mathcal{V}_{21,640} &= x_1^7 x_2 x_3^3 x_4^4 x_5^4, \\
\mathcal{V}_{21,641} &= x_1^7 x_2^3 x_3 x_4^6 x_5^4, & \mathcal{V}_{21,642} &= x_1^3 x_2^5 x_3^2 x_4^4 x_5^7, & \mathcal{V}_{21,643} &= x_1^3 x_2^5 x_3^2 x_4^4 x_5^7, & \mathcal{V}_{21,644} &= x_1^3 x_2^5 x_3^2 x_4^4 x_5^7,
\end{aligned}$$

$$\begin{aligned}
\mathcal{Y}_{21,645} &= x_1^3 x_2^7 x_3^5 x_4^2 x_5^4, & \mathcal{Y}_{21,646} &= x_1^7 x_2^3 x_3^5 x_4^2 x_5^4, & \mathcal{Y}_{21,647} &= x_1^3 x_2^3 x_3^4 x_4^4 x_5^7, & \mathcal{Y}_{21,648} &= x_1^3 x_2^3 x_3^4 x_4^7 x_5^4, \\
\mathcal{Y}_{21,649} &= x_1^3 x_2^3 x_3^7 x_4^4 x_5^4, & \mathcal{Y}_{21,650} &= x_1^3 x_2^7 x_3^3 x_4^4 x_5^4, & \mathcal{Y}_{21,651} &= x_1^7 x_2^3 x_3^3 x_4^4 x_5^4, & \mathcal{Y}_{21,652} &= x_1 x_2^6 x_3^3 x_4^5 x_5^6, \\
\mathcal{Y}_{21,653} &= x_1 x_2^3 x_3^6 x_4^5 x_5^6, & \mathcal{Y}_{21,654} &= x_1 x_2^3 x_3^5 x_4^6 x_5^6, & \mathcal{Y}_{21,655} &= x_1^3 x_2 x_3^6 x_4^5 x_5^6, & \mathcal{Y}_{21,656} &= x_1^3 x_2 x_3^5 x_4^6 x_5^6, \\
\mathcal{Y}_{21,657} &= x_1^3 x_2 x_3^5 x_4^6 x_5^6, & \mathcal{Y}_{21,658} &= x_1^3 x_2 x_3^6 x_4^5 x_5^6, & \mathcal{Y}_{21,659} &= x_1^3 x_2 x_3^5 x_4^5 x_5^6, & \mathcal{Y}_{21,660} &= x_1^3 x_2 x_3^3 x_4^4 x_5^6, \\
\mathcal{Y}_{21,661} &= x_1^3 x_2^3 x_3^4 x_4^5 x_5^6, & \mathcal{Y}_{21,662} &= x_1^3 x_2^3 x_3^4 x_4^5 x_5^6, & \mathcal{Y}_{21,663} &= x_1^3 x_2^3 x_3^5 x_4^5 x_5^6, & \mathcal{Y}_{21,664} &= x_1^3 x_2^3 x_3^3 x_4^5 x_5^6, \\
\mathcal{Y}_{21,665} &= x_1^3 x_2^3 x_3^6 x_4^5 x_5^6, & \mathcal{Y}_{21,666} &= x_1^3 x_2^3 x_3^6 x_4^5 x_5^6.
\end{aligned}$$

6.4. $\mathcal{A}_2$ -generators for $\mathcal{P}_5^0$ in degree 22

According to the results in Sect.3.2.2, we have

$$\mathcal{B}_5^0(22) = \overline{\Phi}^0(\mathcal{B}_4(\omega_{(1)})) \cup \overline{\Phi}^0(\mathcal{B}_4(\omega_{(4)})) \cup \overline{\Phi}^0(\mathcal{B}_4(\omega_{(5)})) = \{\mathcal{Y}_{22,t} : 1 \leq t \leq 460\},$$

where $\omega_{(1)} = (2, 2, 2, 1)$, $\omega_{(4)} = (4, 3, 1, 1)$, $\omega_{(5)} = (4, 3, 3)$, and the monomials $\mathcal{Y}_{22,t} : 1 \leq t \leq 460$, are determined as follows:

$$\begin{aligned}
\mathcal{Y}_{22,1} &= x_4^7 x_5^{15}, & \mathcal{Y}_{22,2} &= x_4^{15} x_5^7, & \mathcal{Y}_{22,3} &= x_3^7 x_5^{15}, & \mathcal{Y}_{22,4} &= x_3^7 x_4^{15}, \\
\mathcal{Y}_{22,5} &= x_3^{15} x_5^7, & \mathcal{Y}_{22,6} &= x_3^{15} x_4^7, & \mathcal{Y}_{22,7} &= x_2^7 x_5^{15}, & \mathcal{Y}_{22,8} &= x_2^7 x_4^{15}, \\
\mathcal{Y}_{22,9} &= x_2^7 x_3^{15}, & \mathcal{Y}_{22,10} &= x_2^{15} x_5^7, & \mathcal{Y}_{22,11} &= x_2^{15} x_4^7, & \mathcal{Y}_{22,12} &= x_2^{15} x_3^7, \\
\mathcal{Y}_{22,13} &= x_1^7 x_5^{15}, & \mathcal{Y}_{22,14} &= x_1^7 x_4^{15}, & \mathcal{Y}_{22,15} &= x_1^7 x_3^{15}, & \mathcal{Y}_{22,16} &= x_1^7 x_2^{15}, \\
\mathcal{Y}_{22,17} &= x_1^{15} x_5^7, & \mathcal{Y}_{22,18} &= x_1^{15} x_4^7, & \mathcal{Y}_{22,19} &= x_1^{15} x_3^7, & \mathcal{Y}_{22,20} &= x_1^{15} x_2^7, \\
\mathcal{Y}_{22,21} &= x_3 x_4^6 x_5^{15}, & \mathcal{Y}_{22,22} &= x_3 x_4^{15} x_5^6, & \mathcal{Y}_{22,23} &= x_3^{15} x_4 x_5^6, & \mathcal{Y}_{22,24} &= x_2 x_4^6 x_5^{15}, \\
\mathcal{Y}_{22,25} &= x_2 x_4^{15} x_5^6, & \mathcal{Y}_{22,26} &= x_2 x_3^6 x_5^{15}, & \mathcal{Y}_{22,27} &= x_2 x_3^{15} x_4^6, & \mathcal{Y}_{22,28} &= x_2 x_3^{15} x_5^6, \\
\mathcal{Y}_{22,29} &= x_2 x_3^{15} x_4^6, & \mathcal{Y}_{22,30} &= x_2^{15} x_4 x_5^6, & \mathcal{Y}_{22,31} &= x_2^{15} x_3 x_5^6, & \mathcal{Y}_{22,32} &= x_2^{15} x_3 x_4^6, \\
\mathcal{Y}_{22,33} &= x_1 x_4^6 x_5^{15}, & \mathcal{Y}_{22,34} &= x_1 x_4^{15} x_5^6, & \mathcal{Y}_{22,35} &= x_1 x_3^6 x_5^{15}, & \mathcal{Y}_{22,36} &= x_1 x_3^6 x_4^{15}, \\
\mathcal{Y}_{22,37} &= x_1 x_3^{15} x_5^6, & \mathcal{Y}_{22,38} &= x_1 x_3^{15} x_4^6, & \mathcal{Y}_{22,39} &= x_1 x_2^6 x_5^{15}, & \mathcal{Y}_{22,40} &= x_1 x_2^6 x_4^{15}, \\
\mathcal{Y}_{22,41} &= x_1 x_2^6 x_3^{15}, & \mathcal{Y}_{22,42} &= x_1 x_2^{15} x_5^6, & \mathcal{Y}_{22,43} &= x_1 x_2^{15} x_4^6, & \mathcal{Y}_{22,44} &= x_1 x_2^{15} x_3^6, \\
\mathcal{Y}_{22,45} &= x_1^{15} x_4 x_5^6, & \mathcal{Y}_{22,46} &= x_1^{15} x_3 x_5^6, & \mathcal{Y}_{22,47} &= x_1^{15} x_3 x_4^6, & \mathcal{Y}_{22,48} &= x_1^{15} x_2 x_5^6, \\
\mathcal{Y}_{22,49} &= x_1^{15} x_2 x_4^6, & \mathcal{Y}_{22,50} &= x_1^{15} x_2 x_3^6, & \mathcal{Y}_{22,51} &= x_3 x_4^{14} x_5^7, & \mathcal{Y}_{22,52} &= x_3 x_4^{14} x_5^7, \\
\mathcal{Y}_{22,53} &= x_3 x_4 x_5^{14}, & \mathcal{Y}_{22,54} &= x_2 x_4^{14} x_5^7, & \mathcal{Y}_{22,55} &= x_2 x_4^{14} x_5^7, & \mathcal{Y}_{22,56} &= x_2 x_3^7 x_5^{14}, \\
\mathcal{Y}_{22,57} &= x_2 x_3^7 x_4^{14}, & \mathcal{Y}_{22,58} &= x_2 x_3^{14} x_5^7, & \mathcal{Y}_{22,59} &= x_2 x_3^{14} x_4^7, & \mathcal{Y}_{22,60} &= x_2^7 x_4 x_5^{14}, \\
\mathcal{Y}_{22,61} &= x_2^7 x_3 x_5^{14}, & \mathcal{Y}_{22,62} &= x_2^7 x_3 x_4^{14}, & \mathcal{Y}_{22,63} &= x_1 x_4^{14} x_5^7, & \mathcal{Y}_{22,64} &= x_1 x_4^{14} x_5^7, \\
\mathcal{Y}_{22,65} &= x_1 x_3^7 x_5^{14}, & \mathcal{Y}_{22,66} &= x_1 x_3^7 x_4^{14}, & \mathcal{Y}_{22,67} &= x_1 x_3^{14} x_5^7, & \mathcal{Y}_{22,68} &= x_1 x_3^{14} x_4^7, \\
\mathcal{Y}_{22,69} &= x_1 x_2^7 x_5^{14}, & \mathcal{Y}_{22,70} &= x_1 x_2^7 x_4^{14}, & \mathcal{Y}_{22,71} &= x_1 x_2^{14} x_3^7, & \mathcal{Y}_{22,72} &= x_1 x_2^{14} x_5^7, \\
\mathcal{Y}_{22,73} &= x_1 x_2^{14} x_4^7, & \mathcal{Y}_{22,74} &= x_1 x_2^{14} x_3^7, & \mathcal{Y}_{22,75} &= x_1^7 x_4 x_5^{14}, & \mathcal{Y}_{22,76} &= x_1^7 x_3 x_5^{14}, \\
\mathcal{Y}_{22,77} &= x_1^7 x_3 x_4^{14}, & \mathcal{Y}_{22,78} &= x_1^7 x_2 x_5^{14}, & \mathcal{Y}_{22,79} &= x_1^7 x_2 x_4^{14}, & \mathcal{Y}_{22,80} &= x_1^7 x_2 x_3^{14}, \\
\mathcal{Y}_{22,81} &= x_3^3 x_4^5 x_5^{14}, & \mathcal{Y}_{22,82} &= x_3^3 x_4^5 x_5^{14}, & \mathcal{Y}_{22,83} &= x_3^3 x_3^5 x_5^{14}, & \mathcal{Y}_{22,84} &= x_3^3 x_3^5 x_4^{14}, \\
\mathcal{Y}_{22,85} &= x_3^3 x_4^5 x_5^{14}, & \mathcal{Y}_{22,86} &= x_3^3 x_3^5 x_5^{14}, & \mathcal{Y}_{22,87} &= x_3^3 x_3^5 x_4^{14}, & \mathcal{Y}_{22,88} &= x_3^3 x_2^5 x_5^{14}, \\
\mathcal{Y}_{22,89} &= x_3^3 x_3^5 x_4^{14}, & \mathcal{Y}_{22,90} &= x_3^3 x_2^5 x_4^{14}, & \mathcal{Y}_{22,91} &= x_3^3 x_4^{13} x_5^6, & \mathcal{Y}_{22,92} &= x_3^3 x_4^{13} x_5^6, \\
\mathcal{Y}_{22,93} &= x_3^3 x_3^{13} x_5^6, & \mathcal{Y}_{22,94} &= x_3^3 x_3^{13} x_4^6, & \mathcal{Y}_{22,95} &= x_3^3 x_4^{13} x_5^6, & \mathcal{Y}_{22,96} &= x_3^3 x_3^{13} x_5^6, \\
\mathcal{Y}_{22,97} &= x_3^3 x_3^{13} x_4^6, & \mathcal{Y}_{22,98} &= x_3^3 x_2^{13} x_5^6, & \mathcal{Y}_{22,99} &= x_3^3 x_2^{13} x_4^6, & \mathcal{Y}_{22,100} &= x_3^3 x_2^{13} x_3^6, \\
\mathcal{Y}_{22,101} &= x_2 x_3 x_4^{14} x_5^6, & \mathcal{Y}_{22,102} &= x_2 x_3 x_4^{14} x_5^6, & \mathcal{Y}_{22,103} &= x_2 x_3^6 x_4 x_5^{14}, & \mathcal{Y}_{22,104} &= x_2 x_3^{14} x_4 x_5^6, \\
\mathcal{Y}_{22,105} &= x_1 x_3 x_4^{14} x_5^6, & \mathcal{Y}_{22,106} &= x_1 x_3 x_4^{14} x_5^6, & \mathcal{Y}_{22,107} &= x_1 x_3^6 x_4 x_5^{14}, & \mathcal{Y}_{22,108} &= x_1 x_3^{14} x_4 x_5^6, \\
\mathcal{Y}_{22,109} &= x_1 x_2 x_4^{14} x_5^6, & \mathcal{Y}_{22,110} &= x_1 x_2 x_4^{14} x_5^6, & \mathcal{Y}_{22,111} &= x_1 x_2 x_3^6 x_5^{14}, & \mathcal{Y}_{22,112} &= x_1 x_2 x_3^6 x_4^{14}, \\
\mathcal{Y}_{22,113} &= x_1 x_2 x_3^{14} x_5^6, & \mathcal{Y}_{22,114} &= x_1 x_2 x_3^{14} x_4^6, & \mathcal{Y}_{22,115} &= x_1 x_2^6 x_4 x_5^{14}, & \mathcal{Y}_{22,116} &= x_1 x_2^6 x_3 x_5^{14}, \\
\mathcal{Y}_{22,117} &= x_1 x_2^6 x_3 x_4^{14}, & \mathcal{Y}_{22,118} &= x_1 x_2^{14} x_4 x_5^6, & \mathcal{Y}_{22,119} &= x_1 x_2^{14} x_3 x_5^6, & \mathcal{Y}_{22,120} &= x_1 x_2^{14} x_3 x_4^6, \\
\mathcal{Y}_{22,121} &= x_2 x_3^2 x_4 x_5^{15}, & \mathcal{Y}_{22,122} &= x_2 x_3^2 x_4^{15} x_5^4, & \mathcal{Y}_{22,123} &= x_2 x_3^{15} x_4^2 x_5^4, & \mathcal{Y}_{22,124} &= x_2^{15} x_3 x_4^2 x_5^4, \\
\mathcal{Y}_{22,125} &= x_1 x_3^2 x_4 x_5^{15}, & \mathcal{Y}_{22,126} &= x_1 x_3^2 x_4^{15} x_5^4, & \mathcal{Y}_{22,127} &= x_1 x_3^{15} x_4^2 x_5^4, & \mathcal{Y}_{22,128} &= x_1 x_2^2 x_4 x_5^{15}, \\
\mathcal{Y}_{22,129} &= x_1 x_2^2 x_4^{15} x_5^4, & \mathcal{Y}_{22,130} &= x_1 x_2^2 x_3^4 x_5^{15}, & \mathcal{Y}_{22,131} &= x_1 x_2^2 x_3^4 x_5^{15}, & \mathcal{Y}_{22,132} &= x_1 x_2^2 x_3^5 x_4^5,
\end{aligned}$$

46

$$\begin{aligned}
\mathcal{V}_{22,325} &= x_1^7 x_2^{11} x_4 x_5^3, & \mathcal{V}_{22,326} &= x_1^7 x_2^{11} x_4^3 x_5, & \mathcal{V}_{22,327} &= x_1^7 x_2^{11} x_3 x_5^3, & \mathcal{V}_{22,328} &= x_1^7 x_2^{11} x_3 x_4^3, \\
\mathcal{V}_{22,329} &= x_1^7 x_2^{11} x_3^3 x_5, & \mathcal{V}_{22,330} &= x_1^7 x_2^{11} x_3^3 x_4, & \mathcal{V}_{22,331} &= x_2 x_3^7 x_4^7 x_5^7, & \mathcal{V}_{22,332} &= x_2^7 x_3^7 x_4^7 x_5^7, \\
\mathcal{V}_{22,333} &= x_2^7 x_3^7 x_4 x_5^7, & \mathcal{V}_{22,334} &= x_2^7 x_3^7 x_4^3 x_5, & \mathcal{V}_{22,335} &= x_1 x_3^7 x_4^7 x_5^7, & \mathcal{V}_{22,336} &= x_1 x_2^7 x_4^7 x_5^7, \\
\mathcal{V}_{22,337} &= x_1 x_2^7 x_3^7 x_5^7, & \mathcal{V}_{22,338} &= x_1 x_2^7 x_3^7 x_4^7, & \mathcal{V}_{22,339} &= x_1^7 x_3 x_4^7 x_5^7, & \mathcal{V}_{22,340} &= x_1^7 x_3^7 x_4 x_5^7, \\
\mathcal{V}_{22,341} &= x_1^7 x_3^7 x_4 x_5^7, & \mathcal{V}_{22,342} &= x_1^7 x_2 x_4^7 x_5^7, & \mathcal{V}_{22,343} &= x_1^7 x_2 x_3^7 x_5^7, & \mathcal{V}_{22,344} &= x_1^7 x_2 x_3^7 x_4^7, \\
\mathcal{V}_{22,345} &= x_1^7 x_2^7 x_4 x_5^7, & \mathcal{V}_{22,346} &= x_1^7 x_2^7 x_4^3 x_5, & \mathcal{V}_{22,347} &= x_1^7 x_2^7 x_3 x_5^7, & \mathcal{V}_{22,348} &= x_1^7 x_2^7 x_3 x_4^7, \\
\mathcal{V}_{22,349} &= x_1^7 x_2^7 x_3^7 x_5, & \mathcal{V}_{22,350} &= x_1^7 x_2^7 x_3^7 x_4, & \mathcal{V}_{22,351} &= x_2^3 x_3^{13} x_4^2 x_5^4, & \mathcal{V}_{22,352} &= x_1^3 x_2^{13} x_4^2 x_5^4, \\
\mathcal{V}_{22,353} &= x_1^3 x_2^{13} x_4^2 x_5^4, & \mathcal{V}_{22,354} &= x_1^3 x_2^{13} x_3^2 x_5^4, & \mathcal{V}_{22,355} &= x_1^3 x_2^{13} x_3^2 x_4^4, & \mathcal{V}_{22,356} &= x_2^3 x_3^5 x_4^2 x_5^{12}, \\
\mathcal{V}_{22,357} &= x_1^3 x_3^5 x_4^2 x_5^{12}, & \mathcal{V}_{22,358} &= x_1^3 x_2^5 x_4^2 x_5^{12}, & \mathcal{V}_{22,359} &= x_1^3 x_2^5 x_3^2 x_5^{12}, & \mathcal{V}_{22,360} &= x_1^3 x_2^5 x_3^2 x_4^{12}, \\
\mathcal{V}_{22,361} &= x_2^3 x_3^3 x_4^3 x_5^{13}, & \mathcal{V}_{22,362} &= x_2^3 x_3^3 x_4^3 x_5^3, & \mathcal{V}_{22,363} &= x_2^3 x_3^{13} x_4^3 x_5^3, & \mathcal{V}_{22,364} &= x_1^3 x_3^3 x_4^3 x_5^{13}, \\
\mathcal{V}_{22,365} &= x_1^3 x_3^3 x_4^3 x_5^3, & \mathcal{V}_{22,366} &= x_1^3 x_3^{13} x_4^3 x_5^3, & \mathcal{V}_{22,367} &= x_1^3 x_2^3 x_4^3 x_5^{13}, & \mathcal{V}_{22,368} &= x_1^3 x_2^3 x_4^3 x_5^3, \\
\mathcal{V}_{22,369} &= x_1^3 x_2^3 x_3^3 x_5^{13}, & \mathcal{V}_{22,370} &= x_1^3 x_2^3 x_3^3 x_4^{13}, & \mathcal{V}_{22,371} &= x_1^3 x_2^3 x_3^{13} x_5^3, & \mathcal{V}_{22,372} &= x_1^3 x_2^3 x_3^{13} x_4^3, \\
\mathcal{V}_{22,373} &= x_1^3 x_2^{13} x_4^3 x_5^3, & \mathcal{V}_{22,374} &= x_1^3 x_2^{13} x_3^3 x_5^3, & \mathcal{V}_{22,375} &= x_1^3 x_2^{13} x_3^3 x_4^3, & \mathcal{V}_{22,376} &= x_2^3 x_3^3 x_4^3 x_5^{11}, \\
\mathcal{V}_{22,377} &= x_2^3 x_3^5 x_4^3 x_5^{11}, & \mathcal{V}_{22,378} &= x_2^3 x_3^5 x_4^{11} x_5^3, & \mathcal{V}_{22,379} &= x_1^3 x_3^3 x_4^5 x_5^{11}, & \mathcal{V}_{22,380} &= x_1^3 x_3^3 x_4^5 x_5^{11}, \\
\mathcal{V}_{22,381} &= x_1^3 x_3^5 x_4^{11} x_5^3, & \mathcal{V}_{22,382} &= x_1^3 x_3^3 x_4^5 x_5^{11}, & \mathcal{V}_{22,383} &= x_1^3 x_2^3 x_3^5 x_5^{11}, & \mathcal{V}_{22,384} &= x_1^3 x_2^3 x_3^5 x_4^{11}, \\
\mathcal{V}_{22,385} &= x_1^3 x_2^5 x_4^3 x_5^{11}, & \mathcal{V}_{22,386} &= x_1^3 x_2^5 x_4^{11} x_5^3, & \mathcal{V}_{22,387} &= x_1^3 x_2^5 x_3^3 x_5^{11}, & \mathcal{V}_{22,388} &= x_1^3 x_2^5 x_3^3 x_4^{11}, \\
\mathcal{V}_{22,389} &= x_1^3 x_2^5 x_3^{11} x_5^3, & \mathcal{V}_{22,390} &= x_1^3 x_2^5 x_3^{11} x_4^3, & \mathcal{V}_{22,391} &= x_2^3 x_3^3 x_4^7 x_5^9, & \mathcal{V}_{22,392} &= x_2^3 x_3^3 x_4^7 x_5^9, \\
\mathcal{V}_{22,393} &= x_2^3 x_3^7 x_4^3 x_5^9, & \mathcal{V}_{22,394} &= x_2^3 x_3^3 x_4^3 x_5^9, & \mathcal{V}_{22,395} &= x_2^7 x_3^3 x_4^3 x_5^9, & \mathcal{V}_{22,396} &= x_2^7 x_3^3 x_4^3 x_5^9, \\
\mathcal{V}_{22,397} &= x_1^3 x_3^3 x_4^3 x_5^9, & \mathcal{V}_{22,398} &= x_1^3 x_3^3 x_4^3 x_5^9, & \mathcal{V}_{22,399} &= x_1^3 x_3^7 x_4^3 x_5^9, & \mathcal{V}_{22,400} &= x_1^3 x_3^7 x_4^3 x_5^9, \\
\mathcal{V}_{22,401} &= x_1^3 x_2^3 x_3^7 x_5^9, & \mathcal{V}_{22,402} &= x_1^3 x_2^3 x_3^7 x_4^9, & \mathcal{V}_{22,403} &= x_1^3 x_2^7 x_4^3 x_5^9, & \mathcal{V}_{22,404} &= x_1^3 x_2^7 x_4^3 x_5^9, \\
\mathcal{V}_{22,405} &= x_1^3 x_2^7 x_3^3 x_5^9, & \mathcal{V}_{22,406} &= x_1^3 x_2^7 x_3^3 x_4^9, & \mathcal{V}_{22,407} &= x_1^3 x_2^7 x_3^9 x_5^3, & \mathcal{V}_{22,408} &= x_1^3 x_2^7 x_3^9 x_4^3, \\
\mathcal{V}_{22,409} &= x_1^7 x_3^3 x_4^3 x_5^9, & \mathcal{V}_{22,410} &= x_1^7 x_3^3 x_4^3 x_5^9, & \mathcal{V}_{22,411} &= x_1^7 x_3^9 x_4^3 x_5^3, & \mathcal{V}_{22,412} &= x_1^7 x_3^9 x_4^3 x_5^3, \\
\mathcal{V}_{22,413} &= x_1^7 x_2^3 x_4^3 x_5^9, & \mathcal{V}_{22,414} &= x_1^7 x_2^3 x_3^3 x_5^9, & \mathcal{V}_{22,415} &= x_1^7 x_2^3 x_3^3 x_4^9, & \mathcal{V}_{22,416} &= x_1^7 x_2^3 x_3^3 x_4^9, \\
\mathcal{V}_{22,417} &= x_1^7 x_2^3 x_3^9 x_4^3, & \mathcal{V}_{22,418} &= x_1^7 x_2^9 x_3^3 x_5^3, & \mathcal{V}_{22,419} &= x_1^7 x_2^9 x_3^3 x_4^3, & \mathcal{V}_{22,420} &= x_1^7 x_2^9 x_3^3 x_4^3, \\
\mathcal{V}_{22,421} &= x_2^3 x_3^5 x_4^{10} x_5^4, & \mathcal{V}_{22,422} &= x_1^3 x_3^5 x_4^{10} x_5^4, & \mathcal{V}_{22,423} &= x_1^3 x_2^5 x_4^{10} x_5^4, & \mathcal{V}_{22,424} &= x_1^3 x_2^5 x_4^{10} x_5^4, \\
\mathcal{V}_{22,425} &= x_1^3 x_2^5 x_3^{10} x_4^4, & \mathcal{V}_{22,426} &= x_2^3 x_3^5 x_4^6 x_5^8, & \mathcal{V}_{22,427} &= x_1^3 x_3^5 x_4^6 x_5^8, & \mathcal{V}_{22,428} &= x_1^3 x_2^5 x_4^6 x_5^8, \\
\mathcal{V}_{22,429} &= x_1^3 x_2^5 x_3^6 x_5^8, & \mathcal{V}_{22,430} &= x_1^3 x_2^5 x_3^6 x_4^8, & \mathcal{V}_{22,431} &= x_2^3 x_3^5 x_4^7 x_5^7, & \mathcal{V}_{22,432} &= x_2^3 x_3^5 x_4^7 x_5^7, \\
\mathcal{V}_{22,433} &= x_2^3 x_3^7 x_4^7 x_5^5, & \mathcal{V}_{22,434} &= x_2^7 x_3^3 x_5^7 x_4^7, & \mathcal{V}_{22,435} &= x_2^7 x_3^3 x_4^7 x_5^5, & \mathcal{V}_{22,436} &= x_2^7 x_3^3 x_4^7 x_5^5, \\
\mathcal{V}_{22,437} &= x_1^3 x_3^5 x_4^7 x_5^5, & \mathcal{V}_{22,438} &= x_1^3 x_3^7 x_4^5 x_5^7, & \mathcal{V}_{22,439} &= x_1^3 x_3^7 x_4^5 x_5^7, & \mathcal{V}_{22,440} &= x_1^3 x_3^7 x_4^5 x_5^7, \\
\mathcal{V}_{22,441} &= x_1^3 x_2^5 x_3^7 x_5^7, & \mathcal{V}_{22,442} &= x_1^3 x_2^5 x_3^7 x_4^7, & \mathcal{V}_{22,443} &= x_1^3 x_2^7 x_4^5 x_5^7, & \mathcal{V}_{22,444} &= x_1^3 x_2^7 x_4^5 x_5^7, \\
\mathcal{V}_{22,445} &= x_1^3 x_2^7 x_3^5 x_5^7, & \mathcal{V}_{22,446} &= x_1^3 x_2^7 x_3^5 x_4^7, & \mathcal{V}_{22,447} &= x_1^3 x_2^7 x_3^7 x_5^5, & \mathcal{V}_{22,448} &= x_1^3 x_2^7 x_3^7 x_5^5, \\
\mathcal{V}_{22,449} &= x_1^7 x_3^3 x_4^3 x_5^5, & \mathcal{V}_{22,450} &= x_1^7 x_3^3 x_4^3 x_5^5, & \mathcal{V}_{22,451} &= x_1^7 x_3^3 x_4^3 x_5^5, & \mathcal{V}_{22,452} &= x_1^7 x_3^3 x_4^3 x_5^5, \\
\mathcal{V}_{22,453} &= x_1^7 x_2^3 x_4^3 x_5^5, & \mathcal{V}_{22,454} &= x_1^7 x_2^3 x_3^5 x_5^7, & \mathcal{V}_{22,455} &= x_1^7 x_2^3 x_3^5 x_4^7, & \mathcal{V}_{22,456} &= x_1^7 x_2^3 x_3^5 x_4^7, \\
\mathcal{V}_{22,457} &= x_1^7 x_2^3 x_3^5 x_4^7, & \mathcal{V}_{22,458} &= x_1^7 x_2^3 x_4^3 x_5^5, & \mathcal{V}_{22,459} &= x_1^7 x_2^3 x_3^5 x_5^5, & \mathcal{V}_{22,460} &= x_1^7 x_2^3 x_3^5 x_4^5.
\end{aligned}$$

6.5. $\mathcal{A}_2$ -generators for $\mathcal{P}_5^+$ in degree 22

By Propositions 3.2.10 and 3.2.14, we have $\mathcal{B}_5^+(22) = \bigcup_{1 \leq j \leq 5} \mathcal{B}_5^+(\omega_j)$. Here $\omega_{(1)} = (2, 2, 2, 1)$, $\omega_{(2)} = (2, 4, 1, 1)$, $\omega_{(3)} = (2, 4, 3)$, $\omega_{(4)} = (4, 3, 1, 1)$, $\omega_{(5)} = (4, 3, 3)$.

$\mathcal{B}_5^+(\omega_{(1)}) = \overline{\Phi}^+(\mathcal{B}_4((\omega_{(1)}))) \cup \mathcal{B}^+(5, \omega_{(1)}) \cup \mathcal{D} = \{\mathcal{V}_{22,t} : 461 \leq t \leq 510\}$, where the monomials $\mathcal{V}_{22,t} : 461 \leq t \leq 510$, are listed as follows:

$$\begin{aligned}
\mathcal{V}_{22,461} &= x_1 x_2 x_3^2 x_4^4 x_5^{14}, & \mathcal{V}_{22,462} &= x_1 x_2 x_3^2 x_4^4 x_5^{14}, & \mathcal{V}_{22,463} &= x_1 x_2 x_3^{14} x_4^2 x_5^4, & \mathcal{V}_{22,464} &= x_1 x_2^2 x_3 x_4^4 x_5^{14}, \\
\mathcal{V}_{22,465} &= x_1 x_2^2 x_3 x_4^4 x_5^{14}, & \mathcal{V}_{22,466} &= x_1 x_2^2 x_3^4 x_4^4 x_5^{14}, & \mathcal{V}_{22,467} &= x_1 x_2^{14} x_3 x_4^2 x_5^4, & \mathcal{V}_{22,468} &= x_1 x_2^2 x_3^6 x_4^2 x_5^{12}, \\
\mathcal{V}_{22,469} &= x_1 x_2^2 x_3^2 x_4^{12} x_5^6, & \mathcal{V}_{22,470} &= x_1 x_2^2 x_3^6 x_4^2 x_5^{12}, & \mathcal{V}_{22,471} &= x_1 x_2^2 x_3^6 x_4^2 x_5^{12}, & \mathcal{V}_{22,472} &= x_1 x_2^2 x_3^2 x_4^{12} x_5^6, \\
\mathcal{V}_{22,473} &= x_1 x_2^2 x_3^{12} x_4^2 x_5^6, & \mathcal{V}_{22,474} &= x_1 x_2^2 x_3^2 x_4^{12} x_5^6, & \mathcal{V}_{22,475} &= x_1 x_2^2 x_3^{13} x_4^2 x_5^4, & \mathcal{V}_{22,476} &= x_1 x_2^2 x_3^5 x_4^2 x_5^{12}, \\
\mathcal{V}_{22,477} &= x_1 x_2^3 x_3^4 x_4^2 x_5^{12}, & \mathcal{V}_{22,478} &= x_1 x_2^3 x_3^{12} x_4^2 x_5^4, & \mathcal{V}_{22,479} &= x_1^3 x_2 x_3^4 x_4^2 x_5^{12}, & \mathcal{V}_{22,480} &= x_1^3 x_2 x_3^{12} x_4^2 x_5^4, \\
\mathcal{V}_{22,481} &= x_1 x_2^2 x_3^5 x_4^{10} x_5^4, & \mathcal{V}_{22,482} &= x_1 x_2^2 x_3^4 x_4^8 x_5^8, & \mathcal{V}_{22,483} &= x_1 x_2^2 x_3^4 x_4^8 x_5^8, & \mathcal{V}_{22,484} &= x_1 x_2^2 x_3^7 x_4^4 x_5^8,
\end{aligned}$$

$$\begin{aligned}
\mathcal{V}_{22,485} &= x_1 x_2^7 x_3^2 x_4^4 x_5^8, & \mathcal{V}_{22,486} &= x_1^7 x_2 x_3^2 x_4^4 x_5^8, & \mathcal{V}_{22,487} &= x_1 x_2^2 x_3^5 x_4^6 x_5^8, & \mathcal{V}_{22,488} &= x_1 x_2^3 x_3^4 x_4^{10} x_5^4, \\
\mathcal{V}_{22,489} &= x_1^3 x_2 x_3^4 x_4^{10} x_5^4, & \mathcal{V}_{22,490} &= x_1 x_2^3 x_3^4 x_4^6 x_5^8, & \mathcal{V}_{22,491} &= x_1^3 x_2 x_3^4 x_4^6 x_5^8, & \mathcal{V}_{22,492} &= x_1 x_2 x_3^6 x_4^6 x_5^8, \\
\mathcal{V}_{22,493} &= x_1 x_2 x_3^6 x_4^{10} x_5^4, & \mathcal{V}_{22,494} &= x_1 x_2^2 x_3^3 x_4^4 x_5^{12}, & \mathcal{V}_{22,495} &= x_1 x_2^2 x_3^3 x_4^{12} x_5^4, & \mathcal{V}_{22,496} &= x_1 x_2^2 x_3^4 x_4^9 x_5^6, \\
\mathcal{V}_{22,497} &= x_1 x_2^2 x_3^5 x_4^8 x_5^6, & \mathcal{V}_{22,498} &= x_1 x_2^3 x_3^2 x_4^4 x_5^{12}, & \mathcal{V}_{22,499} &= x_1 x_2^3 x_3^2 x_4^{12} x_5^4, & \mathcal{V}_{22,500} &= x_1 x_2^3 x_3^3 x_4^8 x_5^6, \\
\mathcal{V}_{22,501} &= x_1 x_2^3 x_3^6 x_4^8 x_5^6, & \mathcal{V}_{22,502} &= x_1 x_2^3 x_3^6 x_4^4 x_5^8, & \mathcal{V}_{22,503} &= x_1^3 x_2 x_3^2 x_4^4 x_5^{12}, & \mathcal{V}_{22,504} &= x_1^3 x_2 x_3^2 x_4^{12} x_5^4, \\
\mathcal{V}_{22,505} &= x_1^3 x_2 x_3^4 x_4^8 x_5^6, & \mathcal{V}_{22,506} &= x_1^3 x_2 x_3^6 x_4^4 x_5^8, & \mathcal{V}_{22,507} &= x_1^3 x_2 x_3^6 x_4^8 x_5^4, & \mathcal{V}_{22,508} &= x_1^3 x_2^5 x_3^2 x_4^8 x_5^8, \\
\mathcal{V}_{22,509} &= x_1^3 x_2^5 x_3^2 x_4^8 x_5^4, & \mathcal{V}_{22,510} &= x_1^3 x_2^5 x_3^8 x_4^4 x_5^4.
\end{aligned}$$

$\mathcal{B}_5^+(\omega_{(2)}) = \mathcal{B}^+(5, \omega_{(2)}) \cup \mathcal{E} = \{\mathcal{V}_{22,t} : 511 \leq t \leq 535\}$, where the monomials $\mathcal{V}_{22,t} : 511 \leq t \leq 535$, are listed as follows:

$$\begin{aligned}
\mathcal{V}_{22,511} &= x_1 x_2^2 x_3^2 x_4^2 x_5^{15}, & \mathcal{V}_{22,512} &= x_1 x_2^2 x_3^2 x_4^{15} x_5^2, & \mathcal{V}_{22,513} &= x_1 x_2^2 x_3^{15} x_4^2 x_5^2, & \mathcal{V}_{22,514} &= x_1 x_2^{15} x_3^2 x_4^2 x_5^2, \\
\mathcal{V}_{22,515} &= x_1^{15} x_2 x_3^2 x_4^2 x_5^2, & \mathcal{V}_{22,516} &= x_1 x_2^2 x_3^2 x_4^3 x_5^{14}, & \mathcal{V}_{22,517} &= x_1 x_2^2 x_3^3 x_4^2 x_5^{14}, & \mathcal{V}_{22,518} &= x_1 x_2^2 x_3^3 x_4^{14} x_5^2, \\
\mathcal{V}_{22,519} &= x_1 x_2^3 x_3^2 x_4^{14} x_5^2, & \mathcal{V}_{22,520} &= x_1 x_2^3 x_3^2 x_4^{14} x_5^2, & \mathcal{V}_{22,521} &= x_1 x_2^3 x_3^{14} x_4^2 x_5^2, & \mathcal{V}_{22,522} &= x_1^3 x_2 x_3^2 x_4^2 x_5^{14}, \\
\mathcal{V}_{22,523} &= x_1^3 x_2 x_3^2 x_4^{14} x_5^2, & \mathcal{V}_{22,524} &= x_1^3 x_2 x_3^{14} x_4^2 x_5^2, & \mathcal{V}_{22,525} &= x_1 x_2^2 x_3^3 x_4^6 x_5^{10}, & \mathcal{V}_{22,526} &= x_1 x_2^3 x_3^2 x_4^6 x_5^{10}, \\
\mathcal{V}_{22,527} &= x_1 x_2^3 x_3^6 x_4^2 x_5^{10}, & \mathcal{V}_{22,528} &= x_1 x_2^3 x_3^6 x_4^{10} x_5^2, & \mathcal{V}_{22,529} &= x_1^3 x_2 x_3^2 x_4^6 x_5^{10}, & \mathcal{V}_{22,530} &= x_1^3 x_2 x_3^6 x_4^2 x_5^{10}, \\
\mathcal{V}_{22,531} &= x_1^3 x_2 x_3^6 x_4^{10} x_5^2, & \mathcal{V}_{22,532} &= x_1^3 x_2^{13} x_3^2 x_4^2 x_5^2, & \mathcal{V}_{22,533} &= x_1^3 x_2^5 x_3^2 x_4^2 x_5^{10}, & \mathcal{V}_{22,534} &= x_1^3 x_2^5 x_3^2 x_4^{10} x_5^2, \\
\mathcal{V}_{22,535} &= x_1^3 x_2^5 x_3^{10} x_4^2 x_5^2.
\end{aligned}$$

$\mathcal{B}_5^+(\omega_{(3)}) = \{\mathcal{V}_{22,t} : 536 \leq t \leq 540\}$, where the monomials $\mathcal{V}_{22,t} : 536 \leq t \leq 540$, are determined as follows:

$$\begin{aligned}
\mathcal{V}_{536} &= x_1 x_2^3 x_3^6 x_4^6 x_5^6, & \mathcal{V}_{537} &= x_1^3 x_2 x_3^6 x_4^6 x_5^6, & \mathcal{V}_{538} &= x_1^3 x_2^5 x_3^2 x_4^6 x_5^6, \\
\mathcal{V}_{539} &= x_1^3 x_2^5 x_3^6 x_4^2 x_5^6, & \mathcal{V}_{540} &= x_1^3 x_2^5 x_3^6 x_4^6 x_5^2,
\end{aligned}$$

$\mathcal{B}_5^+(\omega_{(4)}) = \overline{\Phi}^+(\mathcal{B}_4(\omega_{(4)})) \cup \mathcal{B}^+(5, \omega_{(4)}) = \{\mathcal{V}_{22,t} : 541 \leq t \leq 840\}$, where the monomials $\mathcal{V}_{22,t} : 541 \leq t \leq 840$, are listed as follows:

$$\begin{aligned}
\mathcal{V}_{22,541} &= x_1 x_2 x_3^2 x_4^3 x_5^{15}, & \mathcal{V}_{22,542} &= x_1 x_2 x_3^2 x_4^{15} x_5^3, & \mathcal{V}_{22,543} &= x_1 x_2 x_3^3 x_4^2 x_5^{15}, & \mathcal{V}_{22,544} &= x_1 x_2 x_3^3 x_4^{15} x_5^2, \\
\mathcal{V}_{22,545} &= x_1 x_2 x_3^{15} x_4^2 x_5^3, & \mathcal{V}_{22,546} &= x_1 x_2 x_3^{15} x_4^3 x_5^2, & \mathcal{V}_{22,547} &= x_1 x_2^2 x_3^3 x_4^3 x_5^{15}, & \mathcal{V}_{22,548} &= x_1 x_2^2 x_3^3 x_4^{15} x_5^3, \\
\mathcal{V}_{22,549} &= x_1 x_2^2 x_3^3 x_4^3 x_5^{15}, & \mathcal{V}_{22,550} &= x_1 x_2^2 x_3^3 x_4^{15} x_5^3, & \mathcal{V}_{22,551} &= x_1 x_2^2 x_3^{15} x_4^3 x_5^3, & \mathcal{V}_{22,552} &= x_1 x_2^2 x_3^{15} x_4^3 x_5^3, \\
\mathcal{V}_{22,553} &= x_1 x_2^3 x_3^2 x_4^3 x_5^{15}, & \mathcal{V}_{22,554} &= x_1 x_2^3 x_3^2 x_4^{15} x_5^3, & \mathcal{V}_{22,555} &= x_1 x_2^3 x_3^2 x_4^3 x_5^{15}, & \mathcal{V}_{22,556} &= x_1 x_2^3 x_3^2 x_4^{15} x_5^3, \\
\mathcal{V}_{22,557} &= x_1 x_2^3 x_3^{15} x_4^2 x_5^3, & \mathcal{V}_{22,558} &= x_1 x_2^3 x_3^{15} x_4^2 x_5^3, & \mathcal{V}_{22,559} &= x_1 x_2^{15} x_3^2 x_4^3 x_5^3, & \mathcal{V}_{22,560} &= x_1 x_2^{15} x_3^2 x_4^3 x_5^3, \\
\mathcal{V}_{22,561} &= x_1 x_2^{15} x_3^2 x_4^3 x_5^3, & \mathcal{V}_{22,562} &= x_1 x_2^{15} x_3^2 x_4^3 x_5^3, & \mathcal{V}_{22,563} &= x_1 x_2^{15} x_3^3 x_4^2 x_5^2, & \mathcal{V}_{22,564} &= x_1 x_2^{15} x_3^3 x_4^2 x_5^2, \\
\mathcal{V}_{22,565} &= x_1^3 x_2 x_3^2 x_4^3 x_5^{15}, & \mathcal{V}_{22,566} &= x_1^3 x_2 x_3^2 x_4^{15} x_5^3, & \mathcal{V}_{22,567} &= x_1^3 x_2 x_3^2 x_4^3 x_5^{15}, & \mathcal{V}_{22,568} &= x_1^3 x_2 x_3^2 x_4^{15} x_5^3, \\
\mathcal{V}_{22,569} &= x_1^3 x_2 x_3^{15} x_4^2 x_5^3, & \mathcal{V}_{22,570} &= x_1^3 x_2 x_3^{15} x_4^2 x_5^3, & \mathcal{V}_{22,571} &= x_1^3 x_2^{15} x_3^2 x_4^3 x_5^2, & \mathcal{V}_{22,572} &= x_1^3 x_2^{15} x_3^2 x_4^3 x_5^2, \\
\mathcal{V}_{22,573} &= x_1^{15} x_2 x_3^2 x_4^3 x_5^3, & \mathcal{V}_{22,574} &= x_1^{15} x_2 x_3^2 x_4^3 x_5^3, & \mathcal{V}_{22,575} &= x_1^{15} x_2 x_3^2 x_4^3 x_5^3, & \mathcal{V}_{22,576} &= x_1^{15} x_2 x_3^2 x_4^3 x_5^3, \\
\mathcal{V}_{22,577} &= x_1^{15} x_2 x_3^3 x_4^2 x_5^2, & \mathcal{V}_{22,578} &= x_1^{15} x_2 x_3^3 x_4^2 x_5^2, & \mathcal{V}_{22,579} &= x_1^{15} x_2^2 x_3^2 x_4^3 x_5^3, & \mathcal{V}_{22,580} &= x_1^{15} x_2^2 x_3^2 x_4^3 x_5^3, \\
\mathcal{V}_{22,581} &= x_1 x_2^2 x_3^2 x_4^7 x_5^{11}, & \mathcal{V}_{22,582} &= x_1 x_2^2 x_3^2 x_4^7 x_5^{11}, & \mathcal{V}_{22,583} &= x_1 x_2^2 x_3^2 x_4^{11} x_5^2, & \mathcal{V}_{22,584} &= x_1 x_2^2 x_3^2 x_4^{11} x_5^2, \\
\mathcal{V}_{22,585} &= x_1 x_2^2 x_3^2 x_4^7 x_5^{11}, & \mathcal{V}_{22,586} &= x_1 x_2^2 x_3^2 x_4^{11} x_5^2, & \mathcal{V}_{22,587} &= x_1 x_2^2 x_3^2 x_4^{11} x_5^2, & \mathcal{V}_{22,588} &= x_1 x_2^2 x_3^2 x_4^{11} x_5^2, \\
\mathcal{V}_{22,589} &= x_1 x_2^2 x_3^2 x_4^{11} x_5^2, & \mathcal{V}_{22,590} &= x_1 x_2^2 x_3^2 x_4^{11} x_5^2, & \mathcal{V}_{22,591} &= x_1 x_2^2 x_3^2 x_4^{11} x_5^2, & \mathcal{V}_{22,592} &= x_1 x_2^2 x_3^2 x_4^{11} x_5^2, \\
\mathcal{V}_{22,593} &= x_1^7 x_2 x_3^2 x_4^3 x_5^{11}, & \mathcal{V}_{22,594} &= x_1^7 x_2 x_3^2 x_4^3 x_5^{11}, & \mathcal{V}_{22,595} &= x_1^7 x_2 x_3^2 x_4^3 x_5^{11}, & \mathcal{V}_{22,596} &= x_1^7 x_2 x_3^2 x_4^3 x_5^{11}, \\
\mathcal{V}_{22,597} &= x_1^7 x_2 x_3^{11} x_4^2 x_5^2, & \mathcal{V}_{22,598} &= x_1^7 x_2 x_3^{11} x_4^2 x_5^2, & \mathcal{V}_{22,599} &= x_1^7 x_2^{11} x_3^2 x_4^3 x_5^2, & \mathcal{V}_{22,600} &= x_1^7 x_2^{11} x_3^2 x_4^3 x_5^2, \\
\mathcal{V}_{22,601} &= x_1 x_2 x_3^3 x_4^3 x_5^{14}, & \mathcal{V}_{22,602} &= x_1 x_2 x_3^3 x_4^3 x_5^{14}, & \mathcal{V}_{22,603} &= x_1 x_2 x_3^{14} x_4^3 x_5^3, & \mathcal{V}_{22,604} &= x_1 x_2 x_3^{14} x_4^3 x_5^3, \\
\mathcal{V}_{22,605} &= x_1 x_2^3 x_3^3 x_4^3 x_5^3, & \mathcal{V}_{22,606} &= x_1 x_2^3 x_3^3 x_4^3 x_5^3, & \mathcal{V}_{22,607} &= x_1 x_2^3 x_3^3 x_4^3 x_5^3, & \mathcal{V}_{22,608} &= x_1 x_2^3 x_3^3 x_4^3 x_5^3, \\
\mathcal{V}_{22,609} &= x_1 x_2^3 x_3^{14} x_4^3 x_5^3, & \mathcal{V}_{22,610} &= x_1 x_2^{14} x_3^3 x_4^3 x_5^3, & \mathcal{V}_{22,611} &= x_1 x_2^{14} x_3^3 x_4^3 x_5^3, & \mathcal{V}_{22,612} &= x_1 x_2^{14} x_3^3 x_4^3 x_5^3, \\
\mathcal{V}_{22,613} &= x_1^3 x_2 x_3^3 x_4^3 x_5^{14}, & \mathcal{V}_{22,614} &= x_1^3 x_2 x_3^3 x_4^3 x_5^{14}, & \mathcal{V}_{22,615} &= x_1^3 x_2 x_3^3 x_4^3 x_5^{14}, & \mathcal{V}_{22,616} &= x_1^3 x_2 x_3^3 x_4^3 x_5^{14},
\end{aligned}$$

6.6. $\mathcal{A}_2$ -generators for $\mathcal{P}_5^0$ in degree 47

Using the results in Sect.3.2.2, we have

$$\mathcal{B}_5^0(47) = \overline{\Phi}(\mathcal{B}_4(47)) = \overline{\Phi}(\mathcal{B}_4(\overline{\omega}_{(1)})) = \{\mathcal{Y}_{47,t}; 1 \leq t \leq 560\},$$

where $\overline{\omega}_{(1)} = (3, 2, 2, 2, 1)$, and the monomials $\mathcal{Y}_{47,t} : 1 \leq t \leq 560$, are determined as follows:

$$\begin{aligned} \mathcal{Y}_{47,1} &= x_3 x_4^{15} x_5^{31}, & \mathcal{Y}_{47,2} &= x_3 x_4^{31} x_5^{15}, & \mathcal{Y}_{47,3} &= x_3^{15} x_4 x_5^{31}, & \mathcal{Y}_{47,4} &= x_3^{15} x_4^{31} x_5, \\ \mathcal{Y}_{47,5} &= x_3^{31} x_4 x_5^{15}, & \mathcal{Y}_{47,6} &= x_3^{31} x_4^{15} x_5, & \mathcal{Y}_{47,7} &= x_2 x_4^{15} x_5^{31}, & \mathcal{Y}_{47,8} &= x_2 x_4^{31} x_5^{15}, \\ \mathcal{Y}_{47,9} &= x_2 x_3^{15} x_5^{31}, & \mathcal{Y}_{47,10} &= x_2 x_3^{15} x_4^{31}, & \mathcal{Y}_{47,11} &= x_2 x_3^{31} x_5^{15}, & \mathcal{Y}_{47,12} &= x_2 x_3^{31} x_4^{15}, \\ \mathcal{Y}_{47,13} &= x_2^{15} x_4 x_5^{31}, & \mathcal{Y}_{47,14} &= x_2^{15} x_4^{31} x_5, & \mathcal{Y}_{47,15} &= x_2^{15} x_3 x_5^{31}, & \mathcal{Y}_{47,16} &= x_2^{15} x_3 x_4^{31}, \\ \mathcal{Y}_{47,17} &= x_2^{15} x_3^{31} x_5, & \mathcal{Y}_{47,18} &= x_2^{15} x_3^{31} x_4, & \mathcal{Y}_{47,19} &= x_2^{31} x_4 x_5^{15}, & \mathcal{Y}_{47,20} &= x_2^{31} x_4^{15} x_5, \\ \mathcal{Y}_{47,21} &= x_2^{31} x_3 x_5^{15}, & \mathcal{Y}_{47,22} &= x_2^{31} x_3 x_4^{15}, & \mathcal{Y}_{47,23} &= x_2^{31} x_3^{15} x_5, & \mathcal{Y}_{47,24} &= x_2^{31} x_3^{15} x_4, \\ \mathcal{Y}_{47,25} &= x_1 x_4^{15} x_5^{31}, & \mathcal{Y}_{47,26} &= x_1 x_4^{31} x_5^{15}, & \mathcal{Y}_{47,27} &= x_1 x_3^{15} x_5^{31}, & \mathcal{Y}_{47,28} &= x_1 x_3^{15} x_4^{31}, \\ \mathcal{Y}_{47,29} &= x_1 x_3^{31} x_5^{15}, & \mathcal{Y}_{47,30} &= x_1 x_3^{31} x_4^{15}, & \mathcal{Y}_{47,31} &= x_1 x_2^{15} x_5^{31}, & \mathcal{Y}_{47,32} &= x_1 x_2^{15} x_4^{31}, \\ \mathcal{Y}_{47,33} &= x_1 x_2^{15} x_3^{31}, & \mathcal{Y}_{47,34} &= x_1 x_2^{31} x_5^{15}, & \mathcal{Y}_{47,35} &= x_1 x_2^{31} x_4^{15}, & \mathcal{Y}_{47,36} &= x_1 x_2^{31} x_3^{15}, \\ \mathcal{Y}_{47,37} &= x_1^{15} x_4 x_5^{31}, & \mathcal{Y}_{47,38} &= x_1^{15} x_4^{31} x_5, & \mathcal{Y}_{47,39} &= x_1^{15} x_3 x_5^{31}, & \mathcal{Y}_{47,40} &= x_1^{15} x_3 x_4^{31}, \\ \mathcal{Y}_{47,41} &= x_1^{15} x_3^{31} x_5, & \mathcal{Y}_{47,42} &= x_1^{15} x_3^{31} x_4, & \mathcal{Y}_{47,43} &= x_1^{15} x_2 x_5^{31}, & \mathcal{Y}_{47,44} &= x_1^{15} x_2 x_4^{31}, \\ \mathcal{Y}_{47,45} &= x_1^{15} x_2 x_3^{31}, & \mathcal{Y}_{47,46} &= x_1^{15} x_2^{31} x_5, & \mathcal{Y}_{47,47} &= x_1^{15} x_2^{31} x_4, & \mathcal{Y}_{47,48} &= x_1^{15} x_2^{31} x_3, \\ \mathcal{Y}_{47,49} &= x_1^{31} x_4 x_5^{15}, & \mathcal{Y}_{47,50} &= x_1^{31} x_4^{15} x_5, & \mathcal{Y}_{47,51} &= x_1^{31} x_3 x_5^{15}, & \mathcal{Y}_{47,52} &= x_1^{31} x_3 x_4^{15}, \\ \mathcal{Y}_{47,53} &= x_1^{31} x_3^{15} x_5, & \mathcal{Y}_{47,54} &= x_1^{31} x_3^{15} x_4, & \mathcal{Y}_{47,55} &= x_1^{31} x_2 x_5^{15}, & \mathcal{Y}_{47,56} &= x_1^{31} x_2 x_4^{15}, \\ \mathcal{Y}_{47,57} &= x_1^{31} x_2 x_3^{15}, & \mathcal{Y}_{47,58} &= x_1^{31} x_2^{15} x_5, & \mathcal{Y}_{47,59} &= x_1^{31} x_2^{15} x_4, & \mathcal{Y}_{47,60} &= x_1^{31} x_2^{15} x_3, \\ \mathcal{Y}_{47,61} &= x_3^3 x_4^{13} x_5^{31}, & \mathcal{Y}_{47,62} &= x_3^3 x_4^{31} x_5^{13}, & \mathcal{Y}_{47,63} &= x_3^{31} x_4 x_5^{13}, & \mathcal{Y}_{47,64} &= x_3^{31} x_4^{13} x_5^{31}, \\ \mathcal{Y}_{47,65} &= x_3^3 x_4^{31} x_5^{13}, & \mathcal{Y}_{47,66} &= x_3^3 x_4^{13} x_5^{31}, & \mathcal{Y}_{47,67} &= x_3^3 x_3^{13} x_4^{31}, & \mathcal{Y}_{47,68} &= x_3^3 x_3^{31} x_4^{13}, \\ \mathcal{Y}_{47,69} &= x_3^3 x_3^{31} x_4^{13}, & \mathcal{Y}_{47,70} &= x_3^3 x_3^{13} x_4^{31}, & \mathcal{Y}_{47,71} &= x_3^3 x_3^{13} x_5^{31}, & \mathcal{Y}_{47,72} &= x_3^3 x_3^{31} x_4^{13}, \\ \mathcal{Y}_{47,73} &= x_3^3 x_4^{13} x_5^{31}, & \mathcal{Y}_{47,74} &= x_3^3 x_4^{31} x_5^{13}, & \mathcal{Y}_{47,75} &= x_3^3 x_3^{13} x_5^{31}, & \mathcal{Y}_{47,76} &= x_3^3 x_3^{13} x_4^{31}, \\ \mathcal{Y}_{47,77} &= x_3^3 x_3^{31} x_5^{13}, & \mathcal{Y}_{47,78} &= x_3^3 x_3^{13} x_4^{31}, & \mathcal{Y}_{47,79} &= x_3^3 x_2^{13} x_5^{31}, & \mathcal{Y}_{47,80} &= x_3^3 x_2^{13} x_4^{31}, \\ \mathcal{Y}_{47,81} &= x_3^3 x_2^{13} x_3^{31}, & \mathcal{Y}_{47,82} &= x_3^3 x_2^{31} x_5^{13}, & \mathcal{Y}_{47,83} &= x_3^3 x_2^{31} x_4^{13}, & \mathcal{Y}_{47,84} &= x_3^3 x_2^{31} x_3^{13}, \\ \mathcal{Y}_{47,85} &= x_3^3 x_4^{13} x_5^{31}, & \mathcal{Y}_{47,86} &= x_3^3 x_3^{13} x_5^{31}, & \mathcal{Y}_{47,87} &= x_3^3 x_3^{13} x_4^{31}, & \mathcal{Y}_{47,88} &= x_3^3 x_3^{13} x_5^{13}, \\ \mathcal{Y}_{47,89} &= x_3^3 x_2^{13} x_4^{31}, & \mathcal{Y}_{47,90} &= x_3^3 x_3^{13} x_4^{31}, & \mathcal{Y}_{47,91} &= x_3^3 x_4^{15} x_5^{29}, & \mathcal{Y}_{47,92} &= x_3^3 x_4^{29} x_5^{15}, \\ \mathcal{Y}_{47,93} &= x_3^{15} x_4 x_5^{29}, & \mathcal{Y}_{47,94} &= x_3^{15} x_4^{29} x_5, & \mathcal{Y}_{47,95} &= x_3^3 x_4^{29} x_5^{15}, & \mathcal{Y}_{47,96} &= x_3^3 x_3^{15} x_5^{29}, \\ \mathcal{Y}_{47,97} &= x_3^3 x_3^{15} x_4^{29}, & \mathcal{Y}_{47,98} &= x_3^3 x_3^{29} x_5^{15}, & \mathcal{Y}_{47,99} &= x_3^3 x_3^{29} x_4^{15}, & \mathcal{Y}_{47,100} &= x_2^{15} x_3 x_4^{29}, \\ \mathcal{Y}_{47,101} &= x_2^{15} x_3 x_5^{29}, & \mathcal{Y}_{47,102} &= x_2^{15} x_3 x_4^{29}, & \mathcal{Y}_{47,103} &= x_3^3 x_4^{15} x_5^{29}, & \mathcal{Y}_{47,104} &= x_3^3 x_4^{29} x_5^{15}, \\ \mathcal{Y}_{47,105} &= x_3^3 x_3^{15} x_5^{29}, & \mathcal{Y}_{47,106} &= x_3^3 x_3^{15} x_4^{29}, & \mathcal{Y}_{47,107} &= x_3^3 x_3^{29} x_5^{15}, & \mathcal{Y}_{47,108} &= x_3^3 x_3^{29} x_4^{15}, \\ \mathcal{Y}_{47,109} &= x_3^3 x_2^{15} x_5^{29}, & \mathcal{Y}_{47,110} &= x_3^3 x_2^{15} x_4^{29}, & \mathcal{Y}_{47,111} &= x_3^3 x_2^{15} x_3^{29}, & \mathcal{Y}_{47,112} &= x_3^3 x_2^{29} x_5^{15}, \\ \mathcal{Y}_{47,113} &= x_3^3 x_2^{29} x_4^{15}, & \mathcal{Y}_{47,114} &= x_3^3 x_2^{29} x_3^{15}, & \mathcal{Y}_{47,115} &= x_3^3 x_4^{15} x_5^{29}, & \mathcal{Y}_{47,116} &= x_3^3 x_3^{15} x_5^{29}, \\ \mathcal{Y}_{47,117} &= x_3^3 x_3^{15} x_4^{29}, & \mathcal{Y}_{47,118} &= x_3^3 x_2^{15} x_3^{29}, & \mathcal{Y}_{47,119} &= x_3^3 x_2^{15} x_4^{29}, & \mathcal{Y}_{47,120} &= x_3^3 x_2^{15} x_3^{29}, \\ \mathcal{Y}_{47,121} &= x_3^7 x_4^{11} x_5^{29}, & \mathcal{Y}_{47,122} &= x_3^7 x_4^{11} x_5^{29}, & \mathcal{Y}_{47,123} &= x_3^7 x_3^{11} x_5^{29}, & \mathcal{Y}_{47,124} &= x_3^7 x_3^{11} x_4^{29}, \\ \mathcal{Y}_{47,125} &= x_3^7 x_4^{11} x_5^{29}, & \mathcal{Y}_{47,126} &= x_3^7 x_3^{11} x_5^{29}, & \mathcal{Y}_{47,127} &= x_3^7 x_3^{11} x_4^{29}, & \mathcal{Y}_{47,128} &= x_3^7 x_2^{11} x_5^{29}, \\ \mathcal{Y}_{47,129} &= x_3^7 x_2^{11} x_4^{29}, & \mathcal{Y}_{47,130} &= x_3^7 x_2^{11} x_3^{29}, & \mathcal{Y}_{47,131} &= x_3^7 x_4^{27} x_5^{13}, & \mathcal{Y}_{47,132} &= x_3^7 x_4^{27} x_5^{13}, \\ \mathcal{Y}_{47,133} &= x_2^7 x_3^{27} x_5^{13}, & \mathcal{Y}_{47,134} &= x_2^7 x_3^{27} x_4^{13}, & \mathcal{Y}_{47,135} &= x_2^7 x_4^{27} x_5^{13}, & \mathcal{Y}_{47,136} &= x_2^7 x_3^{27} x_5^{13}, \\ \mathcal{Y}_{47,137} &= x_2^7 x_3^{27} x_4^{13}, & \mathcal{Y}_{47,138} &= x_2^7 x_2^{27} x_5^{13}, & \mathcal{Y}_{47,139} &= x_2^7 x_2^{27} x_4^{13}, & \mathcal{Y}_{47,140} &= x_2^7 x_2^{27} x_3^{13}, \\ \mathcal{Y}_{47,141} &= x_2 x_3 x_4^{14} x_5^{31}, & \mathcal{Y}_{47,142} &= x_2 x_3 x_4^{31} x_5^{14}, & \mathcal{Y}_{47,143} &= x_2 x_3^{14} x_4 x_5^{31}, & \mathcal{Y}_{47,144} &= x_2 x_3^{14} x_4^{31} x_5, \\ \mathcal{Y}_{47,145} &= x_2 x_3^{31} x_4 x_5^{14}, & \mathcal{Y}_{47,146} &= x_2 x_3^{31} x_4^{14} x_5, & \mathcal{Y}_{47,147} &= x_2^{31} x_3 x_4 x_5^{14}, & \mathcal{Y}_{47,148} &= x_2^{31} x_3 x_4^{14} x_5, \\ \mathcal{Y}_{47,149} &= x_1 x_3 x_4^{14} x_5^{31}, & \mathcal{Y}_{47,150} &= x_1 x_3 x_4^{31} x_5^{14}, & \mathcal{Y}_{47,151} &= x_1 x_3^{14} x_4 x_5^{31}, & \mathcal{Y}_{47,152} &= x_1 x_3^{14} x_4^{31} x_5, \\ \mathcal{Y}_{47,153} &= x_1 x_3^{31} x_4 x_5^{14}, & \mathcal{Y}_{47,154} &= x_1 x_3^{31} x_4^{14} x_5, & \mathcal{Y}_{47,155} &= x_1 x_2 x_4^{14} x_5^{31}, & \mathcal{Y}_{47,156} &= x_1 x_2 x_4^{31} x_5^{14}, \\ \mathcal{Y}_{47,157} &= x_1 x_2 x_3^{14} x_5^{31}, & \mathcal{Y}_{47,158} &= x_1 x_2 x_3^{14} x_4^{31}, & \mathcal{Y}_{47,159} &= x_1 x_2 x_3^{31} x_5^{14}, & \mathcal{Y}_{47,160} &= x_1 x_2 x_3^{31} x_4^{14}, \\ \mathcal{Y}_{47,161} &= x_1 x_2^{14} x_4 x_5^{31}, & \mathcal{Y}_{47,162} &= x_1 x_2^{14} x_4^{31} x_5, & \mathcal{Y}_{47,163} &= x_1 x_2^{14} x_3 x_5^{31}, & \mathcal{Y}_{47,164} &= x_1 x_2^{14} x_3 x_4^{31}, \end{aligned}$$

52

53

$$\begin{aligned}
\mathcal{Y}_{47,541} &= x_2^3 x_3^7 x_4^9 x_5^{28}, & \mathcal{Y}_{47,542} &= x_2^7 x_3^3 x_4^9 x_5^{28}, & \mathcal{Y}_{47,543} &= x_1^3 x_3^7 x_4^9 x_5^{28}, & \mathcal{Y}_{47,544} &= x_1^3 x_2^7 x_4^9 x_5^{28}, \\
\mathcal{Y}_{47,545} &= x_1^3 x_2^7 x_3^9 x_5^{28}, & \mathcal{Y}_{47,546} &= x_1^3 x_2^7 x_3^9 x_4^{28}, & \mathcal{Y}_{47,547} &= x_1^7 x_3^3 x_4^9 x_5^{28}, & \mathcal{Y}_{47,548} &= x_1^7 x_2^3 x_4^9 x_5^{28}, \\
\mathcal{Y}_{47,549} &= x_1^7 x_2^3 x_3^9 x_5^{28}, & \mathcal{Y}_{47,550} &= x_1^7 x_2^3 x_3^9 x_4^{28}, & \mathcal{Y}_{47,551} &= x_2^3 x_3^7 x_4^{25} x_5^{12}, & \mathcal{Y}_{47,552} &= x_2^7 x_3^3 x_4^{25} x_5^{12}, \\
\mathcal{Y}_{47,553} &= x_1^3 x_2^7 x_4^{25} x_5^{12}, & \mathcal{Y}_{47,554} &= x_1^3 x_2^7 x_4^{25} x_5^{12}, & \mathcal{Y}_{47,555} &= x_1^3 x_2^7 x_3^{25} x_5^{12}, & \mathcal{Y}_{47,556} &= x_1^3 x_2^7 x_3^{25} x_4^{12}, \\
\mathcal{Y}_{47,557} &= x_1^7 x_3^3 x_4^{25} x_5^{12}, & \mathcal{Y}_{47,558} &= x_1^7 x_2^3 x_4^{25} x_5^{12}, & \mathcal{Y}_{47,559} &= x_1^7 x_2^3 x_3^{25} x_5^{12}, & \mathcal{Y}_{47,560} &= x_1^7 x_2^3 x_3^{25} x_4^{12}.
\end{aligned}$$

6.7. $\mathcal{A}_2$ -generators for $\mathcal{P}_5^+$ in degree 47

Recall that Kameko's squaring operation $\widetilde{Sq}_*^0(5,47) : (Q\mathcal{P}_5)_{47} \rightarrow (Q\mathcal{P}_5)_{21}$ is an epimorphism of $\mathbb{Z}/2GL_5$ -modules. Therefore, we get

$$\mathcal{B}_5(47) = \mathcal{B}_5^0(47) \cup \varphi(\mathcal{B}_5(21)) \cup (\mathcal{B}_5^+(47) \cap \text{Ker}((\widetilde{Sq}_*^0(5,47))),$$

where $|\mathcal{B}_5^0(47)| = 560$, $|\varphi(\mathcal{B}_5(21))| = 840$ with $\varphi : \mathcal{P}_5 \rightarrow \mathcal{P}_5$, $\varphi(u) = X_{(0,5)}u^2$, $\forall u \in \mathcal{P}_5$. From the results in Sect.3.2.2, we have

$$\mathcal{B}_5^+(47) \cap \text{Ker}((\widetilde{Sq}_*^0(5,47)) = \mathcal{B}_5^+(\overline{\omega}_{(1)}) \cup \mathcal{B}_5^+(\overline{\omega}_{(4)}) \cup \mathcal{B}_5^+(\overline{\omega}_{(5)}),$$

where $\overline{\omega}_{(1)} = (3, 2, 2, 2, 1)$, $\overline{\omega}_{(4)} = (3, 4, 3, 1, 1)$, and $\overline{\omega}_{(5)} = (3, 4, 3, 3)$.

$\mathcal{B}_5^+(\overline{\omega}_{(1)})$ is the set of 196 admissible monomials: $\mathcal{Y}_{47,t}$, $1 \leq t \leq 370$

$$\begin{aligned}
\mathcal{Y}_{47,1} &= x_1 x_2 x_3 x_4^{14} x_5^{30}, & \mathcal{Y}_{47,2} &= x_1 x_2 x_3 x_4^{30} x_5^{14}, & \mathcal{Y}_{47,3} &= x_1 x_2 x_3^{14} x_4 x_5^{30}, \\
\mathcal{Y}_{47,4} &= x_1 x_2 x_3^{14} x_4 x_5^{30}, & \mathcal{Y}_{47,5} &= x_1 x_2 x_3^{30} x_4 x_5^{14}, & \mathcal{Y}_{47,6} &= x_1 x_2 x_3^{30} x_4 x_5^{14}, \\
\mathcal{Y}_{47,7} &= x_1 x_2^{14} x_3 x_4 x_5^{30}, & \mathcal{Y}_{47,8} &= x_1 x_2^{14} x_3 x_4 x_5^{30}, & \mathcal{Y}_{47,9} &= x_1 x_2^{30} x_3 x_4 x_5^{14}, \\
\mathcal{Y}_{47,10} &= x_1 x_2^{30} x_3 x_4 x_5^{14}, & \mathcal{Y}_{47,11} &= x_1 x_2^2 x_3^2 x_4^{12} x_5^{31}, & \mathcal{Y}_{47,12} &= x_1 x_2^2 x_3^2 x_4^{12} x_5^{31}, \\
\mathcal{Y}_{47,13} &= x_1 x_2^2 x_3^2 x_4^{12} x_5^{31}, & \mathcal{Y}_{47,14} &= x_1 x_2^2 x_3^2 x_4^{12} x_5^{31}, & \mathcal{Y}_{47,15} &= x_1 x_2^2 x_3^2 x_4^{12} x_5^{31}, \\
\mathcal{Y}_{47,16} &= x_1 x_2^2 x_3^2 x_4^{12} x_5^{31}, & \mathcal{Y}_{47,17} &= x_1 x_2^2 x_3^2 x_4^{12} x_5^{31}, & \mathcal{Y}_{47,18} &= x_1 x_2^2 x_3^2 x_4^{12} x_5^{31}, \\
\mathcal{Y}_{47,19} &= x_1 x_2^2 x_3^2 x_4^{12} x_5^{31}, & \mathcal{Y}_{47,20} &= x_1 x_2^2 x_3^2 x_4^{12} x_5^{31}, & \mathcal{Y}_{47,21} &= x_1 x_2^2 x_3^2 x_4^{12} x_5^{31}, \\
\mathcal{Y}_{47,22} &= x_1 x_2^2 x_3^2 x_4^{12} x_5^{31}, & \mathcal{Y}_{47,23} &= x_1 x_2^2 x_3^2 x_4^{12} x_5^{31}, & \mathcal{Y}_{47,24} &= x_1 x_2^2 x_3^2 x_4^{12} x_5^{31}, \\
\mathcal{Y}_{47,25} &= x_1 x_2^2 x_3^2 x_4^{12} x_5^{31}, & \mathcal{Y}_{47,26} &= x_1 x_2^2 x_3^2 x_4^{12} x_5^{31}, & \mathcal{Y}_{47,27} &= x_1 x_2^2 x_3^2 x_4^{12} x_5^{31}, \\
\mathcal{Y}_{47,28} &= x_1 x_2^2 x_3^2 x_4^{12} x_5^{31}, & \mathcal{Y}_{47,29} &= x_1 x_2^2 x_3^2 x_4^{12} x_5^{31}, & \mathcal{Y}_{47,30} &= x_1 x_2^2 x_3^2 x_4^{12} x_5^{31}, \\
\mathcal{Y}_{47,31} &= x_1 x_2^2 x_3^2 x_4^{12} x_5^{31}, & \mathcal{Y}_{47,32} &= x_1 x_2^2 x_3^2 x_4^{12} x_5^{31}, & \mathcal{Y}_{47,33} &= x_1 x_2^2 x_3^2 x_4^{12} x_5^{31}, \\
\mathcal{Y}_{47,34} &= x_1 x_2^2 x_3^2 x_4^{12} x_5^{31}, & \mathcal{Y}_{47,35} &= x_1 x_2^2 x_3^2 x_4^{12} x_5^{31}, & \mathcal{Y}_{47,36} &= x_1 x_2^2 x_3^2 x_4^{12} x_5^{31}, \\
\mathcal{Y}_{47,37} &= x_1 x_2^2 x_3^2 x_4^{12} x_5^{31}, & \mathcal{Y}_{47,38} &= x_1 x_2^2 x_3^2 x_4^{12} x_5^{31}, & \mathcal{Y}_{47,39} &= x_1 x_2^2 x_3^2 x_4^{12} x_5^{31}, \\
\mathcal{Y}_{47,40} &= x_1 x_2^2 x_3^2 x_4^{12} x_5^{31}, & \mathcal{Y}_{47,41} &= x_1 x_2^2 x_3^2 x_4^{12} x_5^{31}, & \mathcal{Y}_{47,42} &= x_1 x_2^2 x_3^2 x_4^{12} x_5^{31}, \\
\mathcal{Y}_{47,43} &= x_1 x_2^2 x_3^2 x_4^{12} x_5^{31}, & \mathcal{Y}_{47,44} &= x_1 x_2^2 x_3^2 x_4^{12} x_5^{31}, & \mathcal{Y}_{47,45} &= x_1 x_2^2 x_3^2 x_4^{12} x_5^{31}, \\
\mathcal{Y}_{47,46} &= x_1 x_2^2 x_3^2 x_4^{12} x_5^{31}, & \mathcal{Y}_{47,47} &= x_1 x_2^2 x_3^2 x_4^{12} x_5^{31}, & \mathcal{Y}_{47,48} &= x_1 x_2^2 x_3^2 x_4^{12} x_5^{31}, \\
\mathcal{Y}_{47,49} &= x_1 x_2^2 x_3^2 x_4^{12} x_5^{31}, & \mathcal{Y}_{47,50} &= x_1 x_2^2 x_3^2 x_4^{12} x_5^{31}, & \mathcal{Y}_{47,51} &= x_1 x_2^2 x_3^2 x_4^{12} x_5^{31}, \\
\mathcal{Y}_{47,52} &= x_1 x_2^2 x_3^2 x_4^{12} x_5^{31}, & \mathcal{Y}_{47,53} &= x_1 x_2^2 x_3^2 x_4^{12} x_5^{31}, & \mathcal{Y}_{47,54} &= x_1 x_2^2 x_3^2 x_4^{12} x_5^{31}, \\
\mathcal{Y}_{47,55} &= x_1 x_2^2 x_3^2 x_4^{12} x_5^{31}, & \mathcal{Y}_{47,56} &= x_1 x_2^2 x_3^2 x_4^{12} x_5^{31}, & \mathcal{Y}_{47,57} &= x_1 x_2^2 x_3^2 x_4^{12} x_5^{31}, \\
\mathcal{Y}_{47,58} &= x_1 x_2^2 x_3^2 x_4^{12} x_5^{31}, & \mathcal{Y}_{47,59} &= x_1 x_2^2 x_3^2 x_4^{12} x_5^{31}, & \mathcal{Y}_{47,60} &= x_1 x_2^2 x_3^2 x_4^{12} x_5^{31}, \\
\mathcal{Y}_{47,61} &= x_1 x_2^2 x_3^2 x_4^{12} x_5^{31}, & \mathcal{Y}_{47,62} &= x_1 x_2^2 x_3^2 x_4^{12} x_5^{31}, & \mathcal{Y}_{47,63} &= x_1 x_2^2 x_3^2 x_4^{12} x_5^{31}, \\
\mathcal{Y}_{47,64} &= x_1 x_2^2 x_3^2 x_4^{12} x_5^{31}, & \mathcal{Y}_{47,65} &= x_1 x_2^2 x_3^2 x_4^{12} x_5^{31}, & \mathcal{Y}_{47,66} &= x_1 x_2^2 x_3^2 x_4^{12} x_5^{31}, \\
\mathcal{Y}_{47,67} &= x_1 x_2^2 x_3^2 x_4^{12} x_5^{31}, & \mathcal{Y}_{47,68} &= x_1 x_2^2 x_3^2 x_4^{12} x_5^{31}, & \mathcal{Y}_{47,69} &= x_1 x_2^2 x_3^2 x_4^{12} x_5^{31}, \\
\mathcal{Y}_{47,70} &= x_1 x_2^2 x_3^2 x_4^{12} x_5^{31}, & \mathcal{Y}_{47,71} &= x_1 x_2^2 x_3^2 x_4^{12} x_5^{31}, & \mathcal{Y}_{47,72} &= x_1 x_2^2 x_3^2 x_4^{12} x_5^{31}, \\
\mathcal{Y}_{47,73} &= x_1 x_2^2 x_3^2 x_4^{12} x_5^{31}, & \mathcal{Y}_{47,74} &= x_1 x_2^2 x_3^2 x_4^{12} x_5^{31}, & \mathcal{Y}_{47,75} &= x_1 x_2^2 x_3^2 x_4^{12} x_5^{31}, \\
\mathcal{Y}_{47,76} &= x_1 x_2^2 x_3^2 x_4^{12} x_5^{31}, & \mathcal{Y}_{47,77} &= x_1 x_2^2 x_3^2 x_4^{12} x_5^{31}, & \mathcal{Y}_{47,78} &= x_1 x_2^2 x_3^2 x_4^{12} x_5^{31}, \\
\mathcal{Y}_{47,79} &= x_1 x_2^2 x_3^2 x_4^{12} x_5^{31}, & \mathcal{Y}_{47,80} &= x_1 x_2^2 x_3^2 x_4^{12} x_5^{31}, & \mathcal{Y}_{47,81} &= x_1 x_2^2 x_3^2 x_4^{12} x_5^{31}, \\
\mathcal{Y}_{47,82} &= x_1 x_2^2 x_3^2 x_4^{12} x_5^{31}, & \mathcal{Y}_{47,83} &= x_1 x_2^2 x_3^2 x_4^{12} x_5^{31}, & \mathcal{Y}_{47,84} &= x_1 x_2^2 x_3^2 x_4^{12} x_5^{31}, \\
\mathcal{Y}_{47,85} &= x_1 x_2^2 x_3^2 x_4^{12} x_5^{31}, & \mathcal{Y}_{47,86} &= x_1 x_2^2 x_3^2 x_4^{12} x_5^{31}, & \mathcal{Y}_{47,87} &= x_1 x_2^2 x_3^2 x_4^{12} x_5^{31},
\end{aligned}$$

$$\begin{aligned}
\mathcal{V}_{47,88} &= x_1^3 x_2 x_3^{28} x_4^{14} x_5, & \mathcal{V}_{47,89} &= x_1 x_2^3 x_3^{14} x_4^{28} x_5, & \mathcal{V}_{47,90} &= x_1^3 x_2 x_3^{14} x_4^{28} x_5, \\
\mathcal{V}_{47,91} &= x_1 x_2 x_3^6 x_4^{10} x_5^{29}, & \mathcal{V}_{47,92} &= x_1 x_2^6 x_3 x_4^{10} x_5^{29}, & \mathcal{V}_{47,93} &= x_1 x_2 x_3^6 x_4^{11} x_5^{28}, \\
\mathcal{V}_{47,94} &= x_1 x_2^6 x_3 x_4^{11} x_5^{28}, & \mathcal{V}_{47,95} &= x_1 x_2^6 x_3^{11} x_4 x_5^{28}, & \mathcal{V}_{47,96} &= x_1 x_2 x_3^6 x_4^{27} x_5^{12}, \\
\mathcal{V}_{47,97} &= x_1 x_2^6 x_3 x_4^{27} x_5^{12}, & \mathcal{V}_{47,98} &= x_1 x_2^6 x_3^{27} x_4 x_5^{12}, & \mathcal{V}_{47,99} &= x_1 x_2 x_3^6 x_4^{26} x_5^{13}, \\
\mathcal{V}_{47,100} &= x_1 x_2^6 x_3 x_4^{26} x_5^{13}, & \mathcal{V}_{47,101} &= x_1 x_2 x_3^7 x_4^{10} x_5^{28}, & \mathcal{V}_{47,102} &= x_1 x_2^7 x_3 x_4^{10} x_5^{28}, \\
\mathcal{V}_{47,103} &= x_1 x_2^7 x_3 x_4^{10} x_5^{28}, & \mathcal{V}_{47,104} &= x_1 x_2^7 x_3 x_4^{10} x_5^{28}, & \mathcal{V}_{47,105} &= x_1 x_2^7 x_3^{10} x_4 x_5^{28}, \\
\mathcal{V}_{47,106} &= x_1 x_2 x_3^7 x_4^{26} x_5^{12}, & \mathcal{V}_{47,107} &= x_1 x_2^7 x_3 x_4^{26} x_5^{12}, & \mathcal{V}_{47,108} &= x_1 x_2^7 x_3^{26} x_4 x_5^{12}, \\
\mathcal{V}_{47,109} &= x_1 x_2^7 x_3 x_4^{26} x_5^{12}, & \mathcal{V}_{47,110} &= x_1 x_2^7 x_3^{26} x_4 x_5^{12}, & \mathcal{V}_{47,111} &= x_1 x_2^2 x_3^{13} x_4^2 x_5^{29}, \\
\mathcal{V}_{47,112} &= x_1 x_2^2 x_3^{29} x_4^2 x_5^{13}, & \mathcal{V}_{47,113} &= x_1 x_2^2 x_3^{12} x_4^2 x_5^{29}, & \mathcal{V}_{47,114} &= x_1 x_2^2 x_3^{29} x_4^2 x_5^{12}, \\
\mathcal{V}_{47,115} &= x_1 x_2^2 x_3^{29} x_4^2 x_5^{12}, & \mathcal{V}_{47,116} &= x_1 x_2^2 x_3^{12} x_4^2 x_5^{29}, & \mathcal{V}_{47,117} &= x_1 x_2^2 x_3^{29} x_4^2 x_5^{12}, \\
\mathcal{V}_{47,118} &= x_1 x_2^2 x_3^{12} x_4^2 x_5^{29}, & \mathcal{V}_{47,119} &= x_1 x_2^2 x_3^{29} x_4^2 x_5^{12}, & \mathcal{V}_{47,120} &= x_1^3 x_2 x_3^2 x_4^{12} x_5^{29}, \\
\mathcal{V}_{47,121} &= x_1^3 x_2 x_3^2 x_4^{12} x_5^{29}, & \mathcal{V}_{47,122} &= x_1^3 x_2 x_3^{12} x_4^2 x_5^{29}, & \mathcal{V}_{47,123} &= x_1^3 x_2 x_3^2 x_4^2 x_5^{12}, \\
\mathcal{V}_{47,124} &= x_1^3 x_2^2 x_3 x_4^{12} x_5^{12}, & \mathcal{V}_{47,125} &= x_1^3 x_2^2 x_3^2 x_4 x_5^{12}, & \mathcal{V}_{47,126} &= x_1^3 x_2^2 x_3^2 x_4^{12} x_5, \\
\mathcal{V}_{47,127} &= x_1 x_2^2 x_3^{12} x_4^3 x_5^{29}, & \mathcal{V}_{47,128} &= x_1 x_2^2 x_3^3 x_4^{13} x_5^{28}, & \mathcal{V}_{47,129} &= x_1 x_2^2 x_3^3 x_4^{28} x_5^{13}, \\
\mathcal{V}_{47,130} &= x_1 x_2^2 x_3^{13} x_4^3 x_5^{28}, & \mathcal{V}_{47,131} &= x_1 x_2^3 x_3^2 x_4^{13} x_5^{28}, & \mathcal{V}_{47,132} &= x_1 x_2^3 x_3^2 x_4^{28} x_5^{13}, \\
\mathcal{V}_{47,133} &= x_1 x_2^3 x_3^{13} x_4^2 x_5^{28}, & \mathcal{V}_{47,134} &= x_1 x_2^3 x_3^{28} x_4^2 x_5^{13}, & \mathcal{V}_{47,135} &= x_1^3 x_2 x_3^2 x_4^{13} x_5^{28}, \\
\mathcal{V}_{47,136} &= x_1^3 x_2 x_3^2 x_4^{28} x_5^{13}, & \mathcal{V}_{47,137} &= x_1^3 x_2 x_3^{13} x_4^2 x_5^{28}, & \mathcal{V}_{47,138} &= x_1^3 x_2 x_3^{28} x_4^2 x_5^{13}, \\
\mathcal{V}_{47,139} &= x_1^3 x_2^{13} x_3 x_4^2 x_5^{28}, & \mathcal{V}_{47,140} &= x_1^3 x_2^{13} x_3^2 x_4 x_5^{28}, & \mathcal{V}_{47,141} &= x_1^3 x_2^{13} x_3^2 x_4^{28} x_5, \\
\mathcal{V}_{47,142} &= x_1 x_2^2 x_3^{28} x_4^3 x_5^{13}, & \mathcal{V}_{47,143} &= x_1 x_2^2 x_3^4 x_4^9 x_5^{31}, & \mathcal{V}_{47,144} &= x_1 x_2^2 x_3^4 x_4^{31} x_5^9, \\
\mathcal{V}_{47,145} &= x_1 x_2^2 x_3^{31} x_4^4 x_5^9, & \mathcal{V}_{47,146} &= x_1 x_2^2 x_3^{31} x_4^4 x_5^9, & \mathcal{V}_{47,147} &= x_1^3 x_2 x_3^3 x_4^4 x_5^9, \\
\mathcal{V}_{47,148} &= x_1 x_2^2 x_3^4 x_4^{11} x_5^{29}, & \mathcal{V}_{47,149} &= x_1 x_2^2 x_3^4 x_4^{27} x_5^{13}, & \mathcal{V}_{47,150} &= x_1 x_2^2 x_3^4 x_4^{15} x_5^{25}, \\
\mathcal{V}_{47,151} &= x_1 x_2^2 x_3^4 x_4^{25} x_5^{15}, & \mathcal{V}_{47,152} &= x_1 x_2^2 x_3^{15} x_4^4 x_5^{25}, & \mathcal{V}_{47,153} &= x_1 x_2^{15} x_3^2 x_4^4 x_5^{25}, \\
\mathcal{V}_{47,154} &= x_1^{15} x_2 x_3^2 x_4^4 x_5^{25}, & \mathcal{V}_{47,155} &= x_1 x_2^2 x_3^5 x_4^8 x_5^{31}, & \mathcal{V}_{47,156} &= x_1 x_2^2 x_3^5 x_4^{31} x_5^8, \\
\mathcal{V}_{47,157} &= x_1 x_2^2 x_3^{31} x_4^5 x_5^8, & \mathcal{V}_{47,158} &= x_1 x_2^{31} x_3^2 x_4^5 x_5^8, & \mathcal{V}_{47,159} &= x_1^{31} x_2 x_3^2 x_4^5 x_5^8, \\
\mathcal{V}_{47,160} &= x_1 x_2^2 x_3^5 x_4^{30} x_5^9, & \mathcal{V}_{47,161} &= x_1 x_2^2 x_3^5 x_4^9 x_5^{30}, & \mathcal{V}_{47,162} &= x_1 x_2^2 x_3^5 x_4^{10} x_5^{29}, \\
\mathcal{V}_{47,163} &= x_1 x_2^2 x_3^5 x_4^{11} x_5^{28}, & \mathcal{V}_{47,164} &= x_1 x_2^2 x_3^5 x_4^{27} x_5^{12}, & \mathcal{V}_{47,165} &= x_1 x_2^2 x_3^5 x_4^{26} x_5^{13}, \\
\mathcal{V}_{47,166} &= x_1 x_2^2 x_3^5 x_4^{14} x_5^{25}, & \mathcal{V}_{47,167} &= x_1 x_2^2 x_3^5 x_4^{25} x_5^{14}, & \mathcal{V}_{47,168} &= x_1 x_2^2 x_3^5 x_4^{15} x_5^{24}, \\
\mathcal{V}_{47,169} &= x_1 x_2^2 x_3^5 x_4^{24} x_5^{15}, & \mathcal{V}_{47,170} &= x_1 x_2^2 x_3^{15} x_4^5 x_5^{24}, & \mathcal{V}_{47,171} &= x_1 x_2^{15} x_3^2 x_4^5 x_5^{24}, \\
\mathcal{V}_{47,172} &= x_1^{15} x_2 x_3^2 x_4^5 x_5^{24}, & \mathcal{V}_{47,173} &= x_1 x_2^2 x_3^7 x_4^8 x_5^{29}, & \mathcal{V}_{47,174} &= x_1 x_2^2 x_3^7 x_4^{29} x_5^8, \\
\mathcal{V}_{47,175} &= x_1 x_2^7 x_3^2 x_4^8 x_5^{29}, & \mathcal{V}_{47,176} &= x_1 x_2^7 x_3^2 x_4^{29} x_5^8, & \mathcal{V}_{47,177} &= x_1^7 x_2 x_3^2 x_4^8 x_5^{29}, \\
\mathcal{V}_{47,178} &= x_1^7 x_2 x_3^2 x_4^{29} x_5^8, & \mathcal{V}_{47,179} &= x_1 x_2^7 x_3^7 x_4^9 x_5^{28}, & \mathcal{V}_{47,180} &= x_1 x_2^7 x_3^7 x_4^9 x_5^{28}, \\
\mathcal{V}_{47,181} &= x_1^7 x_2 x_3^7 x_4^{25} x_5^{12}, & \mathcal{V}_{47,182} &= x_1 x_2^7 x_3^7 x_4^{25} x_5^{12}, & \mathcal{V}_{47,183} &= x_1 x_2^7 x_3^7 x_4^{25} x_5^{12}, \\
\mathcal{V}_{47,184} &= x_1^7 x_2 x_3^7 x_4^{25} x_5^{12}, & \mathcal{V}_{47,185} &= x_1 x_2^7 x_3^7 x_4^{24} x_5^{13}, & \mathcal{V}_{47,186} &= x_1 x_2^7 x_3^7 x_4^{13} x_5^{24}, \\
\mathcal{V}_{47,187} &= x_1 x_2^7 x_3^7 x_4^{24} x_5^{13}, & \mathcal{V}_{47,188} &= x_1 x_2^7 x_3^7 x_4^{13} x_5^{24}, & \mathcal{V}_{47,189} &= x_1^7 x_2 x_3^2 x_4^{13} x_5^{24}, \\
\mathcal{V}_{47,190} &= x_1^7 x_2 x_3^2 x_4^{13} x_5^{24}, & \mathcal{V}_{47,191} &= x_1 x_2^3 x_3^3 x_4^{12} x_5^{28}, & \mathcal{V}_{47,192} &= x_1 x_2^3 x_3^3 x_4^{28} x_5^{12}, \\
\mathcal{V}_{47,193} &= x_1 x_2^3 x_3^{12} x_4^3 x_5^{28}, & \mathcal{V}_{47,194} &= x_1 x_2^3 x_3^{28} x_4^3 x_5^{12}, & \mathcal{V}_{47,195} &= x_1^3 x_2 x_3^3 x_4^{12} x_5^{28}, \\
\mathcal{V}_{47,196} &= x_1^3 x_2 x_3^3 x_4^{28} x_5^{12}, & \mathcal{V}_{47,197} &= x_1^3 x_2 x_3^{12} x_4^3 x_5^{28}, & \mathcal{V}_{47,198} &= x_1^3 x_2 x_3^{28} x_4^3 x_5^{12}, \\
\mathcal{V}_{47,199} &= x_1^3 x_2^3 x_3 x_4^{12} x_5^{28}, & \mathcal{V}_{47,200} &= x_1^3 x_2^3 x_3 x_4^{28} x_5^{12}, & \mathcal{V}_{47,201} &= x_1^3 x_2^3 x_3^{12} x_4 x_5^{28}, \\
\mathcal{V}_{47,202} &= x_1^3 x_2^3 x_3^{28} x_4 x_5^{12}, & \mathcal{V}_{47,203} &= x_1 x_2^3 x_3^4 x_4^8 x_5^{31}, & \mathcal{V}_{47,204} &= x_1 x_2^3 x_3^4 x_4^{31} x_5^8, \\
\mathcal{V}_{47,205} &= x_1 x_2^3 x_3^{31} x_4^4 x_5^8, & \mathcal{V}_{47,206} &= x_1 x_2^3 x_3^{31} x_4^4 x_5^8, & \mathcal{V}_{47,207} &= x_1^3 x_2 x_3^4 x_4^8 x_5^{31}, \\
\mathcal{V}_{47,208} &= x_1^3 x_2 x_3^4 x_4^{31} x_5^8, & \mathcal{V}_{47,209} &= x_1^3 x_2 x_3^3 x_4^4 x_5^8, & \mathcal{V}_{47,210} &= x_1^3 x_2^{31} x_3 x_4^4 x_5^8, \\
\mathcal{V}_{47,211} &= x_1^{31} x_2 x_3^3 x_4^4 x_5^8, & \mathcal{V}_{47,212} &= x_1^{31} x_2 x_3^3 x_4^4 x_5^8, & \mathcal{V}_{47,213} &= x_1 x_2^3 x_3^4 x_4^{30} x_5^9, \\
\mathcal{V}_{47,214} &= x_1 x_2^3 x_3^{30} x_4^4 x_5^9, & \mathcal{V}_{47,215} &= x_1 x_2^3 x_3^4 x_4^9 x_5^{30}, & \mathcal{V}_{47,216} &= x_1^3 x_2 x_3^4 x_4^{30} x_5^9, \\
\mathcal{V}_{47,217} &= x_1^3 x_2 x_3^{30} x_4^4 x_5^9, & \mathcal{V}_{47,218} &= x_1^3 x_2 x_3^4 x_4^9 x_5^{30}, & \mathcal{V}_{47,219} &= x_1 x_2^3 x_3^4 x_4^{10} x_5^{29}, \\
\mathcal{V}_{47,220} &= x_1^3 x_2 x_3^4 x_4^{10} x_5^{29}, & \mathcal{V}_{47,221} &= x_1^3 x_2^4 x_3 x_4^{10} x_5^{29}, & \mathcal{V}_{47,222} &= x_1 x_2^3 x_3^4 x_4^{11} x_5^{28}, \\
\mathcal{V}_{47,223} &= x_1^3 x_2^4 x_3 x_4^{11} x_5^{28}, & \mathcal{V}_{47,224} &= x_1^3 x_2^4 x_3 x_4^{11} x_5^{28}, & \mathcal{V}_{47,225} &= x_1^3 x_2^4 x_3^{11} x_4 x_5^{28}, \\
\mathcal{V}_{47,226} &= x_1 x_2^3 x_3^4 x_4^{27} x_5^{12}, & \mathcal{V}_{47,227} &= x_1^3 x_2^4 x_3^4 x_4^{27} x_5^{12}, & \mathcal{V}_{47,228} &= x_1^3 x_2^4 x_3^4 x_4^{27} x_5^{12}, \\
\mathcal{V}_{47,229} &= x_1^3 x_2^4 x_3^{27} x_4 x_5^{12}, & \mathcal{V}_{47,230} &= x_1 x_2^3 x_3^4 x_4^{26} x_5^{13}, & \mathcal{V}_{47,231} &= x_1^3 x_2^4 x_3^4 x_4^{26} x_5^{13}, \\
\mathcal{V}_{47,232} &= x_1^3 x_2^4 x_3^4 x_4^{26} x_5^{13}, & \mathcal{V}_{47,233} &= x_1 x_2^3 x_3^4 x_4^{14} x_5^{25}, & \mathcal{V}_{47,234} &= x_1 x_2^3 x_3^{14} x_4^4 x_5^{25},
\end{aligned}$$

$$\begin{aligned}
\mathcal{V}_{47,235} &= x_1 x_2^3 x_3^4 x_4^{25} x_5^{14}, & \mathcal{V}_{47,236} &= x_1^3 x_2 x_3^4 x_4^{14} x_5^{25}, & \mathcal{V}_{47,237} &= x_1^3 x_2 x_3^{14} x_4^4 x_5^{25}, \\
\mathcal{V}_{47,238} &= x_1^3 x_2 x_3^4 x_4^{25} x_5^{14}, & \mathcal{V}_{47,239} &= x_1 x_2^3 x_3^4 x_4^{15} x_5^{24}, & \mathcal{V}_{47,240} &= x_1 x_2^3 x_3^4 x_4^{24} x_5^{15}, \\
\mathcal{V}_{47,241} &= x_1 x_2^3 x_3^{15} x_4^4 x_5^{24}, & \mathcal{V}_{47,242} &= x_1 x_2^{15} x_3^3 x_4^4 x_5^{24}, & \mathcal{V}_{47,243} &= x_1^3 x_2 x_3^4 x_4^{15} x_5^{24}, \\
\mathcal{V}_{47,244} &= x_1^3 x_2 x_3^4 x_4^{24} x_5^{15}, & \mathcal{V}_{47,245} &= x_1^3 x_2 x_3^{15} x_4^4 x_5^{24}, & \mathcal{V}_{47,246} &= x_1^3 x_2^{15} x_3 x_4^4 x_5^{24}, \\
\mathcal{V}_{47,247} &= x_1^{15} x_2 x_3^3 x_4^4 x_5^{24}, & \mathcal{V}_{47,248} &= x_1^{15} x_2^3 x_3 x_4^4 x_5^{24}, & \mathcal{V}_{47,249} &= x_1 x_2^3 x_3^{30} x_4^5 x_5^8, \\
\mathcal{V}_{47,250} &= x_1 x_2^3 x_3^5 x_4^8 x_5^{30}, & \mathcal{V}_{47,251} &= x_1 x_2^3 x_3^5 x_4^{30} x_5^8, & \mathcal{V}_{47,252} &= x_1^3 x_2 x_3^{30} x_4^5 x_5^8, \\
\mathcal{V}_{47,253} &= x_1^3 x_2 x_3^5 x_4^{30} x_5^8, & \mathcal{V}_{47,254} &= x_1^3 x_2 x_3^5 x_4^{30} x_5^8, & \mathcal{V}_{47,255} &= x_1 x_2^3 x_3^5 x_4^{10} x_5^{28}, \\
\mathcal{V}_{47,256} &= x_1^3 x_2 x_3^5 x_4^{10} x_5^{28}, & \mathcal{V}_{47,257} &= x_1^3 x_2^5 x_3 x_4^{10} x_5^{28}, & \mathcal{V}_{47,258} &= x_1^3 x_2^5 x_3^{10} x_4 x_5^{28}, \\
\mathcal{V}_{47,259} &= x_1^3 x_2^5 x_3^{10} x_4 x_5^{28}, & \mathcal{V}_{47,260} &= x_1 x_2^3 x_3^5 x_4^{26} x_5^{12}, & \mathcal{V}_{47,261} &= x_1^3 x_2 x_3^5 x_4^{26} x_5^{12}, \\
\mathcal{V}_{47,262} &= x_1^3 x_2^5 x_3 x_4^{26} x_5^{12}, & \mathcal{V}_{47,263} &= x_1^3 x_2^5 x_3^{26} x_4 x_5^{12}, & \mathcal{V}_{47,264} &= x_1^3 x_2^5 x_3^{26} x_4^{12} x_5, \\
\mathcal{V}_{47,265} &= x_1 x_2^3 x_3^{14} x_4^5 x_5^{24}, & \mathcal{V}_{47,266} &= x_1 x_2^3 x_3^5 x_4^{14} x_5^{24}, & \mathcal{V}_{47,267} &= x_1 x_2^3 x_3^5 x_4^{24} x_5^{14}, \\
\mathcal{V}_{47,268} &= x_1^3 x_2 x_3^{14} x_4^5 x_5^{24}, & \mathcal{V}_{47,269} &= x_1^3 x_2 x_3^5 x_4^{14} x_5^{24}, & \mathcal{V}_{47,270} &= x_1^3 x_2 x_3^5 x_4^{24} x_5^{14}, \\
\mathcal{V}_{47,271} &= x_1 x_2^3 x_3^6 x_4^8 x_5^{29}, & \mathcal{V}_{47,272} &= x_1 x_2^3 x_3^6 x_4^{29} x_5^8, & \mathcal{V}_{47,273} &= x_1^3 x_2 x_3^6 x_4^8 x_5^{29}, \\
\mathcal{V}_{47,274} &= x_1^3 x_2 x_3^6 x_4^{29} x_5^8, & \mathcal{V}_{47,275} &= x_1 x_2^3 x_3^6 x_4^9 x_5^{28}, & \mathcal{V}_{47,276} &= x_1 x_2^6 x_3^3 x_4^9 x_5^{28}, \\
\mathcal{V}_{47,277} &= x_1^3 x_2 x_3^6 x_4^9 x_5^{28}, & \mathcal{V}_{47,278} &= x_1 x_2^3 x_3^6 x_4^{28} x_5^9, & \mathcal{V}_{47,279} &= x_1^3 x_2 x_3^6 x_4^{28} x_5^9, \\
\mathcal{V}_{47,280} &= x_1 x_2^3 x_3^6 x_4^{25} x_5^{12}, & \mathcal{V}_{47,281} &= x_1 x_2^6 x_3^3 x_4^{25} x_5^{12}, & \mathcal{V}_{47,282} &= x_1^3 x_2 x_3^6 x_4^{25} x_5^{12}, \\
\mathcal{V}_{47,283} &= x_1 x_2^3 x_3^6 x_4^{12} x_5^{25}, & \mathcal{V}_{47,284} &= x_1^3 x_2 x_3^6 x_4^{12} x_5^{25}, & \mathcal{V}_{47,285} &= x_1 x_2^3 x_3^6 x_4^{24} x_5^{13}, \\
\mathcal{V}_{47,286} &= x_1 x_2^3 x_3^6 x_4^{13} x_5^{24}, & \mathcal{V}_{47,287} &= x_1^3 x_2 x_3^6 x_4^{24} x_5^{13}, & \mathcal{V}_{47,288} &= x_1^3 x_2 x_3^6 x_4^{13} x_5^{24}, \\
\mathcal{V}_{47,289} &= x_1 x_2^3 x_3^7 x_4^8 x_5^{28}, & \mathcal{V}_{47,290} &= x_1 x_2^7 x_3^3 x_4^8 x_5^{28}, & \mathcal{V}_{47,291} &= x_1^3 x_2 x_3^7 x_4^8 x_5^{28}, \\
\mathcal{V}_{47,292} &= x_1^3 x_2 x_3^7 x_4^8 x_5^{28}, & \mathcal{V}_{47,293} &= x_1^7 x_2 x_3^3 x_4^8 x_5^{28}, & \mathcal{V}_{47,294} &= x_1^7 x_2 x_3^3 x_4^8 x_5^{28}, \\
\mathcal{V}_{47,295} &= x_1 x_2^3 x_3^7 x_4^{28} x_5^8, & \mathcal{V}_{47,296} &= x_1 x_2^7 x_3^3 x_4^{28} x_5^8, & \mathcal{V}_{47,297} &= x_1^3 x_2 x_3^7 x_4^{28} x_5^8, \\
\mathcal{V}_{47,298} &= x_1^3 x_2 x_3^7 x_4^{28} x_5^8, & \mathcal{V}_{47,299} &= x_1^7 x_2 x_3^3 x_4^{28} x_5^8, & \mathcal{V}_{47,300} &= x_1^7 x_2^3 x_3 x_4^{28} x_5^8, \\
\mathcal{V}_{47,301} &= x_1 x_2^3 x_3^7 x_4^{24} x_5^{12}, & \mathcal{V}_{47,302} &= x_1 x_2^7 x_3^3 x_4^{24} x_5^{12}, & \mathcal{V}_{47,303} &= x_1^3 x_2 x_3^7 x_4^{24} x_5^{12}, \\
\mathcal{V}_{47,304} &= x_1^3 x_2 x_3^7 x_4^{24} x_5^{12}, & \mathcal{V}_{47,305} &= x_1^7 x_2 x_3^3 x_4^{24} x_5^{12}, & \mathcal{V}_{47,306} &= x_1^7 x_2^3 x_3 x_4^{24} x_5^{12}, \\
\mathcal{V}_{47,307} &= x_1 x_2^3 x_3^7 x_4^{12} x_5^{24}, & \mathcal{V}_{47,308} &= x_1 x_2^7 x_3^3 x_4^{12} x_5^{24}, & \mathcal{V}_{47,309} &= x_1^3 x_2 x_3^7 x_4^{12} x_5^{24}, \\
\mathcal{V}_{47,310} &= x_1^3 x_2 x_3^7 x_4^{12} x_5^{24}, & \mathcal{V}_{47,311} &= x_1^7 x_2 x_3^3 x_4^{12} x_5^{24}, & \mathcal{V}_{47,312} &= x_1^7 x_2^3 x_3 x_4^{12} x_5^{24}, \\
\mathcal{V}_{47,313} &= x_1 x_2^3 x_3^7 x_4^4 x_5^8, & \mathcal{V}_{47,314} &= x_1^7 x_2 x_3^3 x_4^4 x_5^8, & \mathcal{V}_{47,315} &= x_1^7 x_2^3 x_3 x_4^4 x_5^8, \\
\mathcal{V}_{47,316} &= x_1 x_2^7 x_3^{11} x_4^4 x_5^{24}, & \mathcal{V}_{47,317} &= x_1^7 x_2 x_3^{11} x_4^4 x_5^{24}, & \mathcal{V}_{47,318} &= x_1^7 x_2^{11} x_3 x_4^4 x_5^{24}, \\
\mathcal{V}_{47,319} &= x_1^3 x_2^{29} x_3^2 x_4^9 x_5, & \mathcal{V}_{47,320} &= x_1^3 x_2^{13} x_3^2 x_4^9 x_5^{25}, & \mathcal{V}_{47,321} &= x_1^3 x_2^5 x_3^2 x_4^8 x_5^{29}, \\
\mathcal{V}_{47,322} &= x_1^3 x_2^5 x_3^2 x_4^{29} x_5^8, & \mathcal{V}_{47,323} &= x_1^3 x_2^{29} x_3^2 x_4^8 x_5^8, & \mathcal{V}_{47,324} &= x_1^3 x_2^5 x_3^2 x_4^9 x_5^{28}, \\
\mathcal{V}_{47,325} &= x_1^3 x_2^5 x_3^2 x_4^{28} x_5^9, & \mathcal{V}_{47,326} &= x_1^3 x_2^5 x_3^2 x_4^{25} x_5^{12}, & \mathcal{V}_{47,327} &= x_1^3 x_2^5 x_3^2 x_4^{12} x_5^{25}, \\
\mathcal{V}_{47,328} &= x_1^3 x_2^5 x_3^2 x_4^{13} x_5^{24}, & \mathcal{V}_{47,329} &= x_1^3 x_2^5 x_3^2 x_4^{13} x_5^{24}, & \mathcal{V}_{47,330} &= x_1^3 x_2^{13} x_3^2 x_4^5 x_5^{24}, \\
\mathcal{V}_{47,331} &= x_1^3 x_2^3 x_3^4 x_4^{29} x_5^8, & \mathcal{V}_{47,332} &= x_1^3 x_2^3 x_3^4 x_4^{29} x_5^8, & \mathcal{V}_{47,333} &= x_1^3 x_2^3 x_3^{29} x_4^4 x_5^8, \\
\mathcal{V}_{47,334} &= x_1^3 x_2^{29} x_3^3 x_4^8 x_5^8, & \mathcal{V}_{47,335} &= x_1^3 x_2^3 x_3^4 x_4^{28} x_5^8, & \mathcal{V}_{47,336} &= x_1^3 x_2^4 x_3^3 x_4^9 x_5^{28}, \\
\mathcal{V}_{47,337} &= x_1^3 x_2^3 x_3^4 x_4^{25} x_5^{12}, & \mathcal{V}_{47,338} &= x_1^3 x_2^4 x_3^3 x_4^{25} x_5^{12}, & \mathcal{V}_{47,339} &= x_1^3 x_2^3 x_3^4 x_4^{24} x_5^{13}, \\
\mathcal{V}_{47,340} &= x_1^3 x_2^3 x_3^4 x_4^{13} x_5^{24}, & \mathcal{V}_{47,341} &= x_1^3 x_2^3 x_3^{13} x_4^4 x_5^{24}, & \mathcal{V}_{47,342} &= x_1^3 x_2^{13} x_3^3 x_4^4 x_5^{24}, \\
\mathcal{V}_{47,343} &= x_1^3 x_2^3 x_3^5 x_4^8 x_5^{28}, & \mathcal{V}_{47,344} &= x_1^3 x_2^5 x_3^3 x_4^8 x_5^{28}, & \mathcal{V}_{47,345} &= x_1^3 x_2^3 x_3^5 x_4^{28} x_5^8, \\
\mathcal{V}_{47,346} &= x_1^3 x_2^5 x_3^3 x_4^{28} x_5^8, & \mathcal{V}_{47,347} &= x_1^3 x_2^3 x_3^5 x_4^{24} x_5^{12}, & \mathcal{V}_{47,348} &= x_1^3 x_2^5 x_3^3 x_4^{24} x_5^{12}, \\
\mathcal{V}_{47,349} &= x_1^3 x_2^3 x_3^5 x_4^{12} x_5^{24}, & \mathcal{V}_{47,350} &= x_1^3 x_2^5 x_3^3 x_4^{12} x_5^{24}, & \mathcal{V}_{47,351} &= x_1^3 x_2^5 x_3^{27} x_4^4 x_5^8, \\
\mathcal{V}_{47,352} &= x_1^3 x_2^5 x_3^3 x_4^9 x_5^8, & \mathcal{V}_{47,353} &= x_1^3 x_2^5 x_3^{10} x_4^4 x_5^{25}, & \mathcal{V}_{47,354} &= x_1^3 x_2^5 x_3^{11} x_4^4 x_5^{24}, \\
\mathcal{V}_{47,355} &= x_1^3 x_2^5 x_3^3 x_4^8 x_5^8, & \mathcal{V}_{47,356} &= x_1^3 x_2^5 x_3^{25} x_4^4 x_5^8, & \mathcal{V}_{47,357} &= x_1^3 x_2^7 x_3^9 x_4^4 x_5^{24}, \\
\mathcal{V}_{47,358} &= x_1^3 x_2^3 x_3^9 x_4^{24} x_5^8, & \mathcal{V}_{47,359} &= x_1^3 x_2^5 x_3^{26} x_4^4 x_5^8, & \mathcal{V}_{47,360} &= x_1^3 x_2^5 x_3^{10} x_4^4 x_5^{24}, \\
\mathcal{V}_{47,361} &= x_1^3 x_2^5 x_3^{10} x_4^{21} x_5^8, & \mathcal{V}_{47,362} &= x_1^3 x_2^5 x_3^{11} x_4^{20} x_5^8, & \mathcal{V}_{47,363} &= x_1^3 x_2^5 x_3^{10} x_4^{20} x_5^9, \\
\mathcal{V}_{47,364} &= x_1^3 x_2^5 x_3^{10} x_4^{12} x_5^{17}, & \mathcal{V}_{47,365} &= x_1^3 x_2^5 x_3^{10} x_4^{13} x_5^{16}, & \mathcal{V}_{47,366} &= x_1^3 x_2^5 x_3^{11} x_4^{12} x_5^{16}, \\
\mathcal{V}_{47,367} &= x_1^3 x_2^7 x_3^9 x_4^{20} x_5^8, & \mathcal{V}_{47,368} &= x_1^3 x_2^3 x_3^9 x_4^{20} x_5^8, & \mathcal{V}_{47,369} &= x_1^3 x_2^7 x_3^9 x_4^{12} x_5^{16}, \\
\mathcal{V}_{47,370} &= x_1^7 x_2^3 x_3^9 x_4^{12} x_5^{16}.
\end{aligned}$$

$\mathcal{B}_5^+(\overline{\omega}_{(4)})$ is the set of 112 admissible monomials: $\mathcal{Y}_{47,t}$, $371 \leq t \leq 482$

$$\begin{aligned}
\mathcal{Y}_{47,371} &= x_1 x_2^3 x_3^6 x_4^6 x_5^{31}, & \mathcal{Y}_{47,372} &= x_1 x_2^3 x_3^6 x_4^{31} x_5^6, & \mathcal{Y}_{47,373} &= x_1 x_2^3 x_3^{31} x_4^6 x_5^6, \\
\mathcal{Y}_{47,374} &= x_1 x_2^{31} x_3^6 x_4^6 x_5^6, & \mathcal{Y}_{47,375} &= x_1^3 x_2^6 x_3^6 x_4^{31} x_5^6, & \mathcal{Y}_{47,376} &= x_1^3 x_2^6 x_3^6 x_4^6 x_5^6, \\
\mathcal{Y}_{47,377} &= x_1^3 x_2^6 x_3^{31} x_4^6 x_5^6, & \mathcal{Y}_{47,378} &= x_1^3 x_2^{31} x_3^6 x_4^6 x_5^6, & \mathcal{Y}_{47,379} &= x_1^{31} x_2^6 x_3^6 x_4^6 x_5^6, \\
\mathcal{Y}_{47,380} &= x_1^{31} x_2^6 x_3^6 x_4^6 x_5^6, & \mathcal{Y}_{47,381} &= x_1 x_2^3 x_3^6 x_4^7 x_5^{30}, & \mathcal{Y}_{47,382} &= x_1 x_2^3 x_3^7 x_4^6 x_5^{30}, \\
\mathcal{Y}_{47,383} &= x_1 x_2^3 x_3^7 x_4^6 x_5^{30}, & \mathcal{Y}_{47,384} &= x_1 x_2^7 x_3^6 x_4^6 x_5^{30}, & \mathcal{Y}_{47,385} &= x_1 x_2^7 x_3^7 x_4^6 x_5^{30}, \\
\mathcal{Y}_{47,386} &= x_1^3 x_2^6 x_3^6 x_4^7 x_5^{30}, & \mathcal{Y}_{47,387} &= x_1^3 x_2^6 x_3^7 x_4^6 x_5^{30}, & \mathcal{Y}_{47,388} &= x_1^3 x_2^6 x_3^7 x_4^6 x_5^{30}, \\
\mathcal{Y}_{47,389} &= x_1^3 x_2^7 x_3^6 x_4^6 x_5^{30}, & \mathcal{Y}_{47,390} &= x_1^3 x_2^7 x_3^6 x_4^6 x_5^{30}, & \mathcal{Y}_{47,391} &= x_1^7 x_2^6 x_3^6 x_4^6 x_5^{30}, \\
\mathcal{Y}_{47,392} &= x_1^7 x_2^6 x_3^6 x_4^6 x_5^{30}, & \mathcal{Y}_{47,393} &= x_1^7 x_2^6 x_3^6 x_4^6 x_5^{30}, & \mathcal{Y}_{47,394} &= x_1^7 x_2^6 x_3^6 x_4^6 x_5^{30}, \\
\mathcal{Y}_{47,395} &= x_1 x_2^3 x_3^7 x_4^{14} x_5^{22}, & \mathcal{Y}_{47,396} &= x_1 x_2^7 x_3^6 x_4^{14} x_5^{22}, & \mathcal{Y}_{47,397} &= x_1^3 x_2^7 x_3^6 x_4^{14} x_5^{22}, \\
\mathcal{Y}_{47,398} &= x_1^3 x_2^7 x_3^6 x_4^{14} x_5^{22}, & \mathcal{Y}_{47,399} &= x_1^7 x_2^6 x_3^6 x_4^{14} x_5^{22}, & \mathcal{Y}_{47,400} &= x_1^7 x_2^6 x_3^6 x_4^{14} x_5^{22}, \\
\mathcal{Y}_{47,401} &= x_1 x_2^7 x_3^{27} x_4^6 x_5^6, & \mathcal{Y}_{47,402} &= x_1^7 x_2^6 x_3^{27} x_4^6 x_5^6, & \mathcal{Y}_{47,403} &= x_1^7 x_2^{27} x_3^6 x_4^6 x_5^6, \\
\mathcal{Y}_{47,404} &= x_1 x_2^7 x_3^{11} x_4^6 x_5^{22}, & \mathcal{Y}_{47,405} &= x_1 x_2^7 x_3^{11} x_4^6 x_5^{22}, & \mathcal{Y}_{47,406} &= x_1^7 x_2^6 x_3^{11} x_4^6 x_5^{22}, \\
\mathcal{Y}_{47,407} &= x_1^7 x_2^6 x_3^{11} x_4^6 x_5^{22}, & \mathcal{Y}_{47,408} &= x_1^7 x_2^{11} x_3^6 x_4^6 x_5^{22}, & \mathcal{Y}_{47,409} &= x_1^7 x_2^{11} x_3^6 x_4^6 x_5^{22}, \\
\mathcal{Y}_{47,410} &= x_1^3 x_2^5 x_3^6 x_4^6 x_5^{31}, & \mathcal{Y}_{47,411} &= x_1^3 x_2^5 x_3^6 x_4^6 x_5^{31}, & \mathcal{Y}_{47,412} &= x_1^3 x_2^5 x_3^6 x_4^6 x_5^{31}, \\
\mathcal{Y}_{47,413} &= x_1^3 x_2^5 x_3^6 x_4^6 x_5^{31}, & \mathcal{Y}_{47,414} &= x_1^3 x_2^5 x_3^{31} x_4^6 x_5^2, & \mathcal{Y}_{47,415} &= x_1^3 x_2^5 x_3^{31} x_4^6 x_5^2, \\
\mathcal{Y}_{47,416} &= x_1^3 x_2^{31} x_3^5 x_4^6 x_5^2, & \mathcal{Y}_{47,417} &= x_1^3 x_2^{31} x_3^5 x_4^6 x_5^2, & \mathcal{Y}_{47,418} &= x_1^{31} x_2^5 x_3^5 x_4^6 x_5^2, \\
\mathcal{Y}_{47,419} &= x_1^{31} x_2^5 x_3^5 x_4^6 x_5^2, & \mathcal{Y}_{47,420} &= x_1^3 x_2^5 x_3^7 x_4^6 x_5^{30}, & \mathcal{Y}_{47,421} &= x_1^3 x_2^5 x_3^7 x_4^6 x_5^{30}, \\
\mathcal{Y}_{47,422} &= x_1^3 x_2^5 x_3^7 x_4^6 x_5^{30}, & \mathcal{Y}_{47,423} &= x_1^3 x_2^5 x_3^7 x_4^6 x_5^{30}, & \mathcal{Y}_{47,424} &= x_1^3 x_2^7 x_3^5 x_4^6 x_5^{30}, \\
\mathcal{Y}_{47,425} &= x_1^7 x_2^3 x_3^5 x_4^6 x_5^{30}, & \mathcal{Y}_{47,426} &= x_1^7 x_2^3 x_3^5 x_4^6 x_5^{30}, & \mathcal{Y}_{47,427} &= x_1^7 x_2^3 x_3^{29} x_4^6 x_5^2, \\
\mathcal{Y}_{47,428} &= x_1^7 x_2^3 x_3^{29} x_4^6 x_5^2, & \mathcal{Y}_{47,429} &= x_1^7 x_2^3 x_3^{29} x_4^6 x_5^2, & \mathcal{Y}_{47,430} &= x_1^7 x_2^3 x_3^{29} x_4^6 x_5^2, \\
\mathcal{Y}_{47,431} &= x_1^3 x_2^7 x_3^{13} x_4^6 x_5^{22}, & \mathcal{Y}_{47,432} &= x_1^3 x_2^7 x_3^{13} x_4^6 x_5^{22}, & \mathcal{Y}_{47,433} &= x_1^7 x_2^3 x_3^{13} x_4^6 x_5^{22}, \\
\mathcal{Y}_{47,434} &= x_1^7 x_2^3 x_3^{13} x_4^6 x_5^{22}, & \mathcal{Y}_{47,435} &= x_1^7 x_2^7 x_3^5 x_4^6 x_5^2, & \mathcal{Y}_{47,436} &= x_1^7 x_2^7 x_3^5 x_4^6 x_5^2, \\
\mathcal{Y}_{47,437} &= x_1^7 x_2^{11} x_3^5 x_4^6 x_5^{22}, & \mathcal{Y}_{47,438} &= x_1^7 x_2^{11} x_3^5 x_4^6 x_5^{22}, & \mathcal{Y}_{47,439} &= x_1^7 x_2^{11} x_3^{21} x_4^6 x_5^6, \\
\mathcal{Y}_{47,440} &= x_1^7 x_2^{11} x_3^{21} x_4^6 x_5^6, & \mathcal{Y}_{47,441} &= x_1^3 x_2^5 x_3^6 x_4^6 x_5^{30}, & \mathcal{Y}_{47,442} &= x_1^3 x_2^5 x_3^6 x_4^6 x_5^{30}, \\
\mathcal{Y}_{47,443} &= x_1^3 x_2^5 x_3^6 x_4^6 x_5^{30}, & \mathcal{Y}_{47,444} &= x_1^3 x_2^5 x_3^6 x_4^6 x_5^{30}, & \mathcal{Y}_{47,445} &= x_1^3 x_2^5 x_3^6 x_4^6 x_5^{30}, \\
\mathcal{Y}_{47,446} &= x_1^3 x_2^5 x_3^6 x_4^6 x_5^{30}, & \mathcal{Y}_{47,447} &= x_1^3 x_2^5 x_3^{14} x_4^6 x_5^{22}, & \mathcal{Y}_{47,448} &= x_1^3 x_2^5 x_3^{14} x_4^6 x_5^{22}, \\
\mathcal{Y}_{47,449} &= x_1^3 x_2^9 x_3^6 x_4^6 x_5^6, & \mathcal{Y}_{47,450} &= x_1^3 x_2^9 x_3^6 x_4^6 x_5^6, & \mathcal{Y}_{47,451} &= x_1^3 x_2^9 x_3^{13} x_4^6 x_5^{22}, \\
\mathcal{Y}_{47,452} &= x_1^3 x_2^9 x_3^{13} x_4^6 x_5^{22}, & \mathcal{Y}_{47,453} &= x_1^3 x_2^5 x_3^6 x_4^6 x_5^{27}, & \mathcal{Y}_{47,454} &= x_1^3 x_2^5 x_3^6 x_4^6 x_5^{27}, \\
\mathcal{Y}_{47,455} &= x_1^3 x_2^5 x_3^6 x_4^6 x_5^{27}, & \mathcal{Y}_{47,456} &= x_1^3 x_2^5 x_3^6 x_4^6 x_5^{27}, & \mathcal{Y}_{47,457} &= x_1^3 x_2^5 x_3^6 x_4^6 x_5^{27}, \\
\mathcal{Y}_{47,458} &= x_1^3 x_2^5 x_3^6 x_4^6 x_5^{27}, & \mathcal{Y}_{47,459} &= x_1^3 x_2^5 x_3^6 x_4^6 x_5^{27}, & \mathcal{Y}_{47,460} &= x_1^7 x_2^3 x_3^5 x_4^6 x_5^{26}, \\
\mathcal{Y}_{47,461} &= x_1^7 x_2^3 x_3^5 x_4^6 x_5^{26}, & \mathcal{Y}_{47,462} &= x_1^3 x_2^5 x_3^{11} x_4^6 x_5^{22}, & \mathcal{Y}_{47,463} &= x_1^3 x_2^5 x_3^{11} x_4^6 x_5^{22}, \\
\mathcal{Y}_{47,464} &= x_1^3 x_2^5 x_3^{14} x_4^6 x_5^{22}, & \mathcal{Y}_{47,465} &= x_1^3 x_2^5 x_3^{10} x_4^6 x_5^{22}, & \mathcal{Y}_{47,466} &= x_1^3 x_2^5 x_3^{10} x_4^6 x_5^{22}, \\
\mathcal{Y}_{47,467} &= x_1^3 x_2^5 x_3^{10} x_4^6 x_5^{22}, & \mathcal{Y}_{47,468} &= x_1^3 x_2^5 x_3^{14} x_4^6 x_5^{18}, & \mathcal{Y}_{47,469} &= x_1^3 x_2^5 x_3^{14} x_4^6 x_5^{18}, \\
\mathcal{Y}_{47,470} &= x_1^3 x_2^5 x_3^{14} x_4^6 x_5^{18}, & \mathcal{Y}_{47,471} &= x_1^3 x_2^5 x_3^{25} x_4^6 x_5^6, & \mathcal{Y}_{47,472} &= x_1^3 x_2^5 x_3^{25} x_4^6 x_5^6, \\
\mathcal{Y}_{47,473} &= x_1^3 x_2^7 x_3^9 x_4^6 x_5^{22}, & \mathcal{Y}_{47,474} &= x_1^3 x_2^7 x_3^9 x_4^6 x_5^{22}, & \mathcal{Y}_{47,475} &= x_1^7 x_2^3 x_3^9 x_4^6 x_5^{22}, \\
\mathcal{Y}_{47,476} &= x_1^7 x_2^3 x_3^9 x_4^6 x_5^{22}, & \mathcal{Y}_{47,477} &= x_1^3 x_2^7 x_3^{13} x_4^6 x_5^{18}, & \mathcal{Y}_{47,478} &= x_1^3 x_2^7 x_3^{13} x_4^6 x_5^{18}, \\
\mathcal{Y}_{47,479} &= x_1^7 x_2^3 x_3^{13} x_4^6 x_5^{18}, & \mathcal{Y}_{47,480} &= x_1^7 x_2^3 x_3^{13} x_4^6 x_5^{18}, & \mathcal{Y}_{47,481} &= x_1^7 x_2^{11} x_3^5 x_4^6 x_5^{18}, \\
\mathcal{Y}_{47,482} &= x_1^7 x_2^{11} x_3^5 x_4^6 x_5^{18}.
\end{aligned}$$

$\mathcal{B}_5^+(\overline{\omega}_{(5)})$ is the set of 15 admissible monomials: $\mathcal{Y}_{47,t}$, $483 \leq t \leq 497$

$$\begin{aligned}
\mathcal{Y}_{47,483} &= x_1 x_2^7 x_3^{11} x_4^{14} x_5^{14}, & \mathcal{Y}_{47,484} &= x_1^7 x_2 x_3^{11} x_4^{14} x_5^{14}, & \mathcal{Y}_{47,485} &= x_1^7 x_2^{11} x_3 x_4^{14} x_5^{14}, \\
\mathcal{Y}_{47,486} &= x_1^7 x_2^{11} x_3^{13} x_4^{14} x_5^2, & \mathcal{Y}_{47,487} &= x_1^7 x_2^{11} x_3^{13} x_4^{14} x_5^2, & \mathcal{Y}_{47,488} &= x_1^3 x_2^5 x_3^{11} x_4^{14} x_5^{14}, \\
\mathcal{Y}_{47,489} &= x_1^3 x_2^5 x_3^9 x_4^{14} x_5^{14}, & \mathcal{Y}_{47,490} &= x_1^3 x_2^5 x_3^9 x_4^{14} x_5^{14}, & \mathcal{Y}_{47,491} &= x_1^3 x_2^5 x_3^{13} x_4^{10} x_5^{14}, \\
\mathcal{Y}_{47,492} &= x_1^3 x_2^5 x_3^{13} x_4^{10} x_5^{14}, & \mathcal{Y}_{47,493} &= x_1^3 x_2^5 x_3^{13} x_4^{10} x_5^{14}, & \mathcal{Y}_{47,494} &= x_1^7 x_2^3 x_3^{13} x_4^{14} x_5^{10}, \\
\mathcal{Y}_{47,495} &= x_1^7 x_2^{11} x_3^5 x_4^{14} x_5^{10}, & \mathcal{Y}_{47,496} &= x_1^7 x_2^{11} x_3^5 x_4^{14} x_5^{10}, & \mathcal{Y}_{47,497} &= x_1^7 x_2^{11} x_3^{13} x_4^6 x_5^{10}.
\end{aligned}$$

References

- [1] J.F. Adams, *On the structure and applications of the Steenrod algebra*, Comment. Math. Helv. 32 (1958), 180-214.
- [2] S.V. Ault and W. Singer, *On the homology of elementary Abelian groups as modules over the Steenrod algebra*, J. Pure Appl. Algebra 215 (2011), 2847-2852.
- [3] J.M. Boardman, *Modular representations on the homology of power of real projective space*; In: M. C. Tangora (Ed.), Algebraic Topology, Oaxtepec, 1991; In: Contemp. Math. 146 (1993), 49-70.
- [4] R.R. Bruner, L.M. Hà and N.H.V. Hùng, *On behavior of the algebraic transfer*, Trans. Amer. Math. Soc. 357 (2005), 437-487.
- [5] T.W. Chen, *Determination of $\text{Ext}_{\mathcal{A}}^{5,*}(\mathbb{Z}/2, \mathbb{Z}/2)$* , Topology Appl. 158 (2011), 660-689.
- [6] P.H. Chơn and L.M. Hà, *Lambda algebra and the Singer transfer*, C. R. Math. Acad. Sci. Paris, Ser. I 349 (2011), 21-23.
- [7] M.C. Crabb and J.R. Hubbuck, *Representations of the homology of BV and the Steenrod algebra II*, Algebra Topology: new trend in localization and periodicity, Progr. Math. 136 (1996), 143-154.
- [8] L. M. Hà, *Sub-Hopf algebras of the Steenrod algebra and the Singer transfer*; in Proceedings of the School and Conference in Algebraic Topology, Hà Nội 2004; in Geom. Topol. Publ., Coventry, 11 (2007), 101-124.
- [9] M. Kameko, *Products of projective spaces as Steenrod modules*, PhD. thesis, The Johns Hopkins University, ProQuest LLC, Ann Arbor, MI, 1990, 29 pages.
- [10] W.H. Lin, *$\text{Ext}_{\mathcal{A}}^{4,*}(\mathbb{Z}/2, \mathbb{Z}/2)$ and $\text{Ext}_{\mathcal{A}}^{5,*}(\mathbb{Z}/2, \mathbb{Z}/2)$* , Topology Appl. 155 (2008), 459-496.
- [11] A. Liulevicius, *The factorization of cyclic reduced powers by secondary cohomology operations*, Mem. Amer. Math. Soc. 42 (1962).
- [12] N. Minami, *The iterated transfer analogue of the new doomsday conjecture*, Trans. Amer. Math. Soc. 351 (1990), 2325-2351.
- [13] M.F. Mothebe, *Generators of the polynomial algebra $\mathbb{F}_2[x_1, x_2, \dots, x_n]$ as a module over the Steenrod algebra*, PhD. thesis, The University of Manchester, 1997.
- [14] M. F. Mothebe, *Dimension result for the polynomial algebra $\mathbb{F}_2[x_1, x_2, \dots, x_n]$ as a module over the Steenrod algebra*, Int. J. Math. Math. Sci. 2013, Art. ID 150704, 6 pp.
- [15] M. F. Mothebe, *Products of admissible monomials in the polynomial algebra as a module over the Steenrod algebra*, J. Math. Research 8 (2016), 112-116.
- [16] M.F. Mothebe, P. Kaelo and O. Ramatebele, *Dimension formula for the polynomial algebra as a module over the Steenrod algebra in degrees less than or equal to 12*, J. Math. Research, 8 (2016), 92-100.
- [17] T.N. Nam, *$\mathcal{A}$ -générateurs génériques pour l'algèbre polynomiale*, Adv. Math. 186 (2004) 334-362.
- [18] T.N. Nam, *Transfert algébrique et action du groupe linéaire sur les puissances divisées modulo 2*, Ann. Inst. Fourier (Grenoble) 58 (2008), 1785-1837.
- [19] F.P. Peterson, *Generators of $H^*(\mathbb{R}P^\infty \times \mathbb{R}P^\infty)$ as a module over the Steenrod algebra*, Abstracts Amer. Math. Soc. No. 833, April 1987.
- [20] F.P. Peterson, *$\mathcal{A}$ -generators for certain polynomial algebras*, Math. Proc. Cambridge Philos. Soc. 105 (1989), 311-312.
- [21] Đ.V. Phúc and N. Sum, *On the generators of the polynomial algebra as a module over the Steenrod algebra*, C.R.Math. Acad. Sci. Paris, Ser. I 353 (2015), 1035-1040.
- [22] Đ.V. Phúc and N. Sum, *On a minimal set of generators for the polynomial algebra of five variables as a module over the Steenrod algebra*, Acta Math. Vietnam. 42 (2017), 149-162.
- [23] Đ.V. Phúc, *The hit problem for the polynomial algebra of five variables in degree seventeen and its application*, East-West J. Math. 18 (2016), 27-46.
- [24] Đ.V. Phúc, *$\mathcal{A}$ -generators for the polynomial algebra of five variables and the algebraic transfer*, Preprint 2018, 31 pages; available online at <http://arxiv.org/abs/1810.06061>.
- [25] Đ.V. Phúc, *$\mathcal{A}$ -generators for the polynomial algebra of five variables in degree $5(2^t - 1) + 6 \cdot 2^t$* , Commun. Korean Math. Soc. 21 pages (accepted 2019).
- [26] S. Priddy, *On characterizing summands in the classifying space of a group*, Amer. Jour. Math. 112 (1990), 737-748.
- [27] J. Repka and P. Selick, *On the subalgebra of $H_*((\mathbb{R}P^\infty)^n; \mathbb{F}_2)$ annihilated by Steenrod operations*, J. Pure Appl. Algebra 127 (1998), 273-288.
- [28] J.H. Silverman and W.M. Singer, *On the action of Steenrod squares on polynomial algebras II*, J. Pure Appl. Algebra 98 (1995), 95-103.
- [29] W.M. Singer, *The transfer in homological algebra*, Math. Zeit. 202 (1989), 493-523.
- [30] W.M. Singer, *On the action of the Steenrod squares on polynomial algebras*, Proc. Amer. Math. Soc. 111 (1991), 577-583.
- [31] W.M. Singer, *Rings of symmetric functions as modules over the Steenrod algebra*, Algebr. Geom. Topol. 8 (2008), 541-562.
- [32] N.E. Steenrod and D.B.A. Epstein, *Cohomology operations*, Annals of Mathematics Studies 50, Princeton University Press, Princeton N.J, 1962.
- [33] N. Sum, *The negative answer to Kameko's conjecture on the hit problem*, Adv. Math. 225 (2010), 2365-2390.
- [34] N. Sum, *On the Peterson hit problem*, Adv. Math. 274 (2015), 432-489.
- [35] N. Sum, *The squaring operation and the Singer algebraic transfer*, Preprint 2018, 38 pages; available online at <http://arxiv.org/abs/1609.03006>.
- [36] N. Sum, *On the determination of the Singer transfer*, Vietnam Journal of Science, Technology and Engineering, 60(1) (2018), 3-16.
- [37] N. Sum, *On a construction for the generators of the polynomial algebra as a module over the Steenrod algebra*; In: Singh M., Song Y., Wu J. (eds), Algebraic Topology and Related Topics. Trends in Mathematics. Birkhäuser/Springer, Singapore (2019), 265-286.

- [38] N.K. Tín, *The admissible monomial basis for the polynomial algebra of five variables in degree eleven*, Journal of Science, Quy Nhơn University, 6 (2012), 81-89.
- [39] N.K. Tín, *The admissible monomial basis for the polynomial algebra of five variables in degree eight*, Journal of Mathematical Sciences and Applications, Vol. 2, No. 2 (2014), 21-24.
- [40] N.K. Tín, *The admissible monomial basis for the polynomial algebra of five variables in degree $2^{s+1} + 2^s - 5$* , East-West J. Math. 16 (2014), 34-46.
- [41] N.K. Tín, *On Singer's conjecture for the fifth algebraic transfer*, Preprint 2016, 25 pages; available online at <http://arxiv.org/abs/1609.02250>.
- [42] N.K. Tín and N. Sum, *Kameko's homomorphism and the algebraic transfer*, C. R. Acad. Sci. Paris, Ser. I 354 (2016), 940-943.
- [43] G. Walker and R.M.W. Wood, *Weyl modules and the mod 2 Steenrod algebra*, J. Algebra 311 (2007), 840-858.
- [44] G. Walker and R.M.W. Wood, *Polynomials and the mod 2 Steenrod Algebra: Volume 1, The Peterson hit problem*; In: London Math. Soc. Lecture Note Ser., Cambridge Univ. Press, January 11, 2018.
- [45] R.M.W. Wood, *Steenrod squares of polynomials and the Peterson conjecture*, Math. Proc. Cambriges Phil. Soc. 105 (1989), 307-309.
- [46] R.M.W. Wood, *Problems in the Steenrod algebra*, Bull. London Math. Soc. 30 (1998), 449-517.