## ALGEBRAS WITH LAURENT POLYNOMIAL IDENTITY

By

# Claudenir Freire Rodrigues, Ramon Códamo B. da Costa

Universidade Federal do Amazonas - Departamento de Matemática Av. General Rodrigo Octavio Jordão Ramos, 1200 - Coroado I, Manaus - AM - Brasil, 69067-005 claudiomao1@gmail.com (corresponding author) ramoncodamo@gmail.com

## ALGEBRAS WITH LAURENT POLYNOMIAL IDENTITY

CLAUDENIR FREIRE, RAMON CODAMO

ABSTRACT. In this article we shows some results about algebra with the group of units having special polynomial identity.

**Keywords**: Laurent polynomial identity, unit group, group identity, nil ideal, Brian Hartley Conjecture.

#### 1. Introduction

A Laurent polynomial  $f = f(x_1, ..., x_l)$  in the noncommutative variables  $x_i$ , i = 1, ..., l is an element non-zero in the group algebra  $RF_l$  over a ring R with free group  $F_l = \langle x_1, ..., x_l \rangle$ . One says f in  $RF_l$  is a Laurent Polynomial Identity (LPI) for an R algebra A (respc. for U(A) the group of units in A) if  $f(a_1, ..., a_l) = 0$  for all sequence  $a_1, ..., a_l$  in A (respc. in U(A)). A word w in  $RF_l$  is group identity for all U(A) if  $w(a_1, ..., a_l) = 1$ , for  $a_1, ..., a_l$  in U(A).

Brian Hartleys Conjecture: Let G be a torsion group and R a field. If the unit group U(RG) of RG satisfies a group identity w = 1, then RG satisfies a polynomial identity.

In [2] A.Giambruno, E.Jespers and A.Valenti shows that the Brian Hartleys Conjecture is true for a group algebra RG over an infinite commutative domnain R and a torsion group G and G has no divisible order elements by p with characteristic of R is p. For this purpose they proved the following crucial result.

**Proposition 1.1.** Let A be an algebra over an infinite commutative domain R and suppose that U(A) satisfies a group identity. There exists a positive m such that if a, b, c, u in A and  $a^2 = bc = 0$ , then bacA is nil right ideal of bounded exponent less or equal than m.

The case identity group w = 1 in the conjecture says the unity group satisfy equivalently the Laurent polynomial F = 1 - w. We use this idea and study the case  $F = a_1 + a_2w_2 + ... + a_nw_n$ . Clearly the case F = 1 - w is a special case of our generalization.

In this paper we prove the following results.

**Proposition 1.2.** Let A be an algebra with the group of units U(A) admits a LPI over a ring R whith unit whose non-constants words has the sum of exponents non-zero at least one of the variables. Then there exists a polynomial  $f \in R[X]$  with the limited degree  $d \leq 4(-l+r)+3$ , where  $l = \min\{\sum expw_i\}$  and  $r = \max\{\sum exp w_i\}$ , determinated by the LPI such that for all a, b, c, u in A with  $a^2 = bc = 0$ , f(bacu) = 0. In particular, bacA is an algebraic ideal.

As a consequence of Proposition 1.2 we prove the following:

**Corollary 1.3.** Let A be an algebra over a commutative domain R with |R| > dwhose unit group U(A) has a LPI. If a, b,  $c \in A$  with  $a^2 = bc = 0$ , then bacA is a nil ideal with limited exponent.

Notice that the restriction on LPI is necessary because we will show that without this assumption Proposition 1.2 i not valid. In this article we always adopt LPI with this restriction. Also, by changin  $x_i$  by  $x^{-i}yx^i$  we may assume that LPI in two variables.

### **Proof of Proposition 1.2:**

Let  $P = a_1 + a_2 w_2 + \dots + a_r w_t$  the LPI with  $w_i$  in the form

 $w_i = x_1^{r_1} x_2^{s_2} \cdots x_1^{r_k} x_2^{s_k}, k \ge 1, r_i, s_i$ 

integers. If  $\sum exp w_i = \sum exp_{x_1} w_i + \sum exp_{x_2} w_i = 0$  then we substitute  $x_1 = x_1^k$  or  $x_2 = x_2^k$  with k > 1 big enough, We get a new *LPI* in the form  $P = a_1 + a_2w'_2 + \dots + a_rw'_r$ , where

$$\sum exp \, w'_i = \sum exp_{x_1} \, w'_i + \sum exp_{x_2} \, w'_i = k \sum exp_{x_1} \, w_i + \sum exp_{x_2} \, w_i$$

$$\sum_{i=1}^{n} exp \, w'_i = \sum_{i=1}^{n} exp_{x_1} \, w_i + k \sum_{i=1}^{n} exp_{x_2} \, w_i$$

In any case that is not zero.

In general,  $P(1, 1) = a_1 + a_2 + ... + a_r = 0$ , so  $a_2 + ... + a_r \neq 0$ , because  $a_1 \neq 0$ . Hence any collection of sums with distinct parcels involving all coefficients  $a_2, ..., a_r$  has a non-zero term.

From this

or

$$P(\alpha, \alpha) = a_1 + a_2 \alpha^{\sum expw_2} + \dots + a_k \alpha^{\sum expw_k} = 0, \ 1 < k \le r,$$

is a non-zero polynomial with integers exponents over R, for all  $\alpha \in U(A)$ . In the following we see that all exponents are not necessarily positive.

Let  $\alpha^{-l}$  be where  $l = \min \{\sum expw_i\}$ , so  $\alpha^{-l}P(\alpha, \alpha) = f_0(\alpha) = 0$ , where  $f_0 \neq 0$ is a polynomial with degree  $\leq -l + r$ , over R. Therefore U(A) is algebraic over R. Now, we are going to use this polynomial with a particular unit in U(A). First let b = c be, so  $a^2 = b^2 = 0$ . Then we have (1 + aua),  $(1 + bauab) \in U(A)$ , for all  $u \in A$ . Let  $Q_n$  be the set of all products in *aua* and *bauab* with at most n - 1 factors, and let W be the set of all products in coefficients  $a_i$ 's, *aua* and *bauab*.

Thus  $\alpha = (1 + aua) \cdot (1 + bauab) \in U(A)$ , and for this unit,

$$\alpha^n = 1 + aua + bauab + \sum_{t \in Q_n} t + (auabauab)^n$$

for all  $n \ge 1$ .

It follows that there exists a polynomial  $f_0 \neq 0$  over R such that

$$f_0(\alpha) = a_1 + a_2 + \dots + a_r + \sum_{t \in W} t + \sum_{i=2}^r a_i (auabauab)^{s_i}$$
$$= \sum_{t \in W} t + \sum_{i=2}^r a_i (auabauab)^{s_i} = 0.$$

As  $a_2 + ... + a_r \neq 0$ , after to organize the powers of *auabauab* which appears in the last sum, has a non-zero coefficients. This term has the form  $(a_{j_1} + ... + a_{j_k})(auabauab)^s$  where  $s \leq (-l+r)$ . So,  $abf_0(\alpha)au = 0$  is a non-zero polynomial g over R such that g(abau) = 0 with degree  $\leq 2(-l+r) + 1$ . This finishs the case b = c. In general, if  $a, \alpha, \beta \in A$  and  $a^2 = \alpha\beta = 0$ , then we have  $(\beta u\alpha)^2 = 0$ , for all  $u \in A$ . So  $\beta u\alpha a\beta u\alpha A$  is algebraic and there exists g over R which is not zero and  $g((\beta u\alpha a\beta u\alpha)a) = 0$ , from this there exists a non-zero polynomial  $f = (\alpha a)f_0(\beta u)$ 

with  $f(\alpha a\beta u) = 0$  such that  $d = degree f \leq 4(-l+r) + 3$ . Therefore  $\alpha a\beta A$  is algebraic over R.

#### Proof of Corollary 1.3 :

By Proposition 1.2, there exists  $f \in R[x] \setminus \{0\}$  with degree  $d \leq m = 4(-l+r)+3$ , such that  $f(bac\lambda u) = 0$  for all  $\lambda \in R$ ,  $u \in A$ . That is,  $\lambda p_1 + \lambda^2 p_2 + \ldots + \lambda^d p_d = 0$ where  $p_i$  is polynomial on *bacu* with  $p_d = b_d(bacu)^d$ . Now, by the Vandermond argument,  $p_1 = \ldots = p_d = 0$ , so  $(bacu)^d = 0$ .

The next results are LPI versions of Lema 3.2, Lema 3.3 of [6] and Corolary 3 of [2]. Their proofs are analogous to the group identity case changing GI by LPI.

**Corollary 1.4.** Let A be a semiprime algebra over an commutative domain R with |R| > d. If U(A) satisfies a LPI, then every idempotent element of  $R^{-1}A$  is central.

**Corollary 1.5.** Let A be an algebra over a field K with U(A) satisfying a LPI, and  $a, b \in A$  such that  $a^2 = b^2 = 0$ .

- (1) If |K| > 2d, then  $(ab)^{2d} = 0$ .
- (2) If ab is nilpotent, then  $(ab)^{2d} = 0$ .

**Corollary 1.6.** Let K be any field and  $n \ge 2$ . Then the following are equivalent

- (1)  $U(M_n(K))$  satisfies the LPI.
- (2) K is a finite field.
- (3)  $U(M_n(K))$  satisfies a group identity.

In addition, if one of these conditions is hold, then  $|K| \leq 2d$  and  $n \leq 2log_{|K|}2d + 2 \leq 2log_22d + 2$ .

In general, Proposition 1.2 is not hold without the restrictions about exponents. If it does, then *bacu* is algebraic for all  $u \in A$ , in particular, for c = b and u = a, *ba* is algebraic. It follows that by Vandermonde augument with R a infinite commutative domain *ba* is nilpotent. In general, *ba* is not a nilpotent element. For example, let  $A = M_n(R)$ , n > 1. By *Amitsur - Levitzki* theorem [3] A satisfies a *LPI* 

$$f_1(x_1, \dots, x_{2n}) = S_{2n} \cdot (x_1 \cdots x_{2n})^{-1}$$

where  $s_{2n}$  is the polynomial standard. However A with  $a = e_{21}$  and  $b = e_{12}$ ,  $ba = e_{11}$  is not nilpotent. Note that this algebra also satisfies

$$f_2(x_1, ..., x_{2n}) = f_1(x_1, ..., x_{2n}) + S_{2n}$$

(an LPI with some words having sum of exponents zero at every variable).

By the same arguments we can see that if  $A = M_n(R)$  be n > 1 and R a infinite commutative domain, there is not exist a non-zero polynomial  $g \in R[X]$  not zero such that g(ab) = 0 for all  $a, b \in A$  with  $a^2 = b^2 = 0$ . Note that the other side of Proposition 1.2 is true. In fact, suppose that R is finite, so A is finite. From this for all matrices  $x \in A$  there exist integers r > t > 0 such that  $x^r = x^t$ . So x satisfies the polynomial  $p_x(X) = X^r - X^t$ . Then defining  $g = \prod_{x \in A} p_x$  one has g(ab) = 0 for

all  $a, b \in A$  with  $a^2 = b^2 = 0$ , which is a contradiction.

**Lemma 1.7.** Let  $A = M_n(R)$  be n > 1 and R a commutative domain. Then R is infinite if and only if there is not any non-zero polynomial  $g \in R[X]$  such that g(ab) = 0 for all  $a, b \in A$  with  $a^2 = b^2 = 0$ .

**Proposition 1.8.** Let  $F = \frac{R[x, y]}{(x^2, y^2)}$  where R is a infinite commutative domain. Then F satisfies any polynomial identity f of  $M_n(R)$  where  $n \ge 2$ . Also, U(F) does not satisfy the standard polynomial  $S_2$  and  $S_3$ . In particular, F satisfies  $S_{2n}$  if only if  $n \ge 2$ .

*Proof.* Suppose by contradiction that F does not satisfy an identity f of  $M_n(R)$  where  $n \ge 2$ . Then there exist  $\overline{u_1} = \overline{g_1(x, y)}, ..., \overline{u_{2n}} = \overline{g_{2n}(x, y)} \in F$  with

$$f(\overline{u_1}, ..., \overline{u_{2n}}) \neq 0$$

So (multiplying by x or y on left or right if necessary) we obtain

$$g(xy) = f(u_1, ..., u_{2n}) \neq \sum_i f_i u^2 g_i$$

u = x, y. Then, for all  $a, b \in A = M_n(R)$  such that  $a^2 = b^2 = 0$ , we have g(ab) = 0it is a contradiction by Lemma 1.7. By evaluation of  $S_2$  and  $S_3$  on units  $\overline{1+x}, \overline{1+y}$ and  $\overline{(1+x)(1+y)}$ , we see that U(F) does not satisfy  $S_2$  and  $S_3$ .

**Corollary 1.9.** Let B be an algebra over a commutative domain R whose U(A) satisfies the standard polynomial  $S_3$ . Then there exists a polynomial  $g \in R[X]$  such that g(ab) = 0 for all  $a, b \in B$  with  $a^2 = b^2 = 0$ .

*Proof.* By applying  $S_3$  on X, Y and XY we obtain

$$S_3(X, Y, XY) = (YX)^2 - X^2Y^2 - YX^2Y + XY^2X.$$

So  $S_3(1+x, 1+y, (1+x)(1+y)) \neq 0$  in F, then multiplying by x on the left and y on the right we obtain a polynomial  $g(xy) = xS_3y$  such that g(ab) = 0.

**Corollary 1.10.** Let F be the algebra in Proposition 1.8. The F does not satisfy the group identity w = 1, with R a infinite field.

*Proof.* If F satisfies a group identity w = 1. By Lemma 3.1 [6] there exists a polynomial  $g(t) \in R[t]$  such that  $g(\overline{x}\overline{y}) = 0$ . Thus  $\overline{g(xy)} = 0$ , that is,  $g(xy) = \sum f_i u^2 g_i$ , u = x, y. Then, g(ab) = 0 for all  $a, b \in M_n(R)$  such that  $a^2 = b^2 = 0$  and this contradicts Lemma 1.1.

With some adaptations Lemma 3.1 [6] holds whenever R is an infinite commutative domain. It follows that Corollary 1.5 holds if R is an infinite commutative domain. Note that F is an algebra with LPI (having more than one word which not constant) with  $\sum expw_i = 0$  without group identity (LPI with only one nonconstant word).

#### References

- A. Giambruno, S.Sehgal, and A. Valenti, Group algebras whose units satisfy a group identity, Proc. Amer. Math. Soc. 125 (1997), 629-634. MR 97f:16056
- [2] A. Giambruno, E. Jespers, and A. Valenti, *Group identities on units of rings*, Arch. Mathj. (Basel) 63 (1994), no. 4, 291-296. MR 95:16044
- [3] D.S. Passman, The algebraic structure of group rings, Dover, New York 2011
- [4] J. Z. Gonalves, M.A. Dokuchaev Identities on units of algebraic algebras, Journal of Algebra 250, 638-646 (2002)
- [5] J.Z. Gonalves, Free subgroups of units in group rings, Canad. Math. Bull.27 (1984), no. 3, 309-312. MR 85k:20021
- [6] Liu, Chia-Hsin, Group Algebras With Units Satisfying a Group Identity, Proc. Amer. Math. Soc. 127, (1999f), 327-336.

DEPARTMENTO DE MATEMTICA, UNIVERSIDADE FEDERAL DO AMAZONAS, AMAZONAS, MAN-AUS, BRASIL

*E-mail address*: claudiomao1@gmail.com *E-mail address*: ramoncodamo@gmail.com