

A new result for the global existence (and boundedness), regularity and stabilization of a three-dimensional Keller-Segel-Navier-Stokes system modeling coral fertilization

Jiashan Zheng*

School of Mathematics and Statistics Science,
Ludong University, Yantai 264025, P.R.China

Abstract

This paper deals with the following quasilinear Keller-Segel-Navier-Stokes system modeling coral fertilization

$$\left\{ \begin{array}{l} n_t + u \cdot \nabla n = \Delta n - \nabla \cdot (nS(x, n, c)\nabla c) - nm, \quad x \in \Omega, t > 0, \\ c_t + u \cdot \nabla c = \Delta c - c + m, \quad x \in \Omega, t > 0, \\ m_t + u \cdot \nabla m = \Delta m - nm, \quad x \in \Omega, t > 0, \\ u_t + \kappa(u \cdot \nabla)u + \nabla P = \Delta u + (n + m)\nabla \phi, \quad x \in \Omega, t > 0, \\ \nabla \cdot u = 0, \quad x \in \Omega, t > 0 \end{array} \right. \quad (*)$$

under no-flux boundary conditions in a bounded domain $\Omega \subset \mathbb{R}^3$ with smooth boundary, where $\phi \in W^{2,\infty}(\Omega)$. Here $S(x, n, c)$ denotes the rotational effect which satisfies $|S(x, n, c)| \leq S_0(c)(1+n)^{-\alpha}$ with $\alpha \geq 0$ and some nonnegative nondecreasing function

*Corresponding author. E-mail address: zhengjiashan2008@163.com (J.Zheng)

S_0 . Based on a new weighted estimate and some carefully analysis, if $\alpha > 0$, then for any $\kappa \in \mathbb{R}$, system $(*)$ possesses a global weak solution for which there exists $T > 0$ such that (n, c, m, u) is smooth in $\Omega \times (T, \infty)$. Furthermore, for any $p > 1$, this solution is uniformly bounded in with respect to the norm in $L^p(\Omega) \times L^\infty(\Omega) \times L^\infty(\Omega) \times L^2(\Omega; \mathbb{R}^3)$. Building on this boundedness property and some other analysis, it can finally even be proved that in the large time limit, any such solution approaches the spatially homogeneous equilibrium $(\hat{n}, \hat{m}, \hat{m}, 0)$ in an appropriate sense, where $\hat{n} = \frac{1}{|\Omega|} \{\int_\Omega n_0 - \int_\Omega m_0\}_+$ and $\hat{m} = \frac{1}{|\Omega|} \{\int_\Omega m_0 - \int_\Omega n_0\}_+$.

Key words: Navier-Stokes system; Keller-Segel model; Global existence; Large time behavior; Tensor-valued sensitivity

2010 Mathematics Subject Classification: 35K55, 35Q92, 35Q35, 92C17

1 Introduction

This work is concerned with the following chemotaxis-fluid system modelling coral fertilization:

$$\left\{ \begin{array}{l} n_t + u \cdot \nabla n = \Delta n - \nabla \cdot (nS(x, n, c)\nabla c) - nm, \quad x \in \Omega, t > 0, \\ c_t + u \cdot \nabla c = \Delta c - c + m, \quad x \in \Omega, t > 0, \\ m_t + u \cdot \nabla m = \Delta m - nm, \quad x \in \Omega, t > 0, \\ u_t + \kappa(u \cdot \nabla)u + \nabla P = \Delta u + (n + m)\nabla \phi, \quad x \in \Omega, t > 0, \\ \nabla \cdot u = 0, \quad x \in \Omega, t > 0 \\ (\nabla n - nS(x, n, c)) \cdot \nu = \nabla c \cdot \nu = \nabla m \cdot \nu = 0, u = 0, \quad x \in \partial\Omega, t > 0, \\ n(x, 0) = n_0(x), c(x, 0) = c_0(x), m(x, 0) = m_0(x), u(x, 0) = u_0(x), \quad x \in \Omega, \end{array} \right. \quad (1.1)$$

where $\Omega \subset \mathbb{R}^3$ is a bounded domain with smooth boundary $\partial\Omega$ and the matrix-valued function $S(x, n, c)$ indicates the rotational effect which satisfies

$$S \in C^2(\bar{\Omega} \times [0, \infty)^2; \mathbb{R}^{3 \times 3}) \quad (1.2)$$

and

$$|S(x, n, c)| \leq (1+n)^{-\alpha} S(c) \text{ for all } (x, n, c) \in \Omega \times [0, \infty)^2 \text{ with } S(c) \text{ nondecreasing on } [0, \infty) \quad (1.3)$$

and $\alpha \geq 0$. As described in [19, 18, 9, 10], problems of this type arise in the modeling of the phenomenon of coral broadcast spawning, where the sperm n chemotactically moves toward the higher concentration of the chemical c released by the egg m , while the egg m is merely affected by random diffusion, fluid transport and degradation upon contact with the sperm (see also [22]). Here κ, u, P and ϕ denote, respectively, the strength of nonlinear fluid convection, the velocity field, the associated pressure of the fluid and the potential of the gravitational field. We further note that the sensitivity tensor $S(x, n, c)$ may take values that are matrices possibly containing nontrivial off-diagonal entries, which reflects that the chemotactic migration may not necessarily be oriented along the gradient of the chemical signal, but may rather involve rotational flux components (see [56, 55] for the detailed model derivation).

Chemotaxis is the directed movement of the cells as a response to gradients of the concentration of the chemical signal substance in their environment, where the chemical signal substance may be produced or consumed by cells themselves (see e.g. Hillen and Painter [12] and [2]). The classical chemotaxis system was introduced in 1970 by Keller and Segel ([17]), which is called Keller-Segel system. Since then, the Keller-Segel model has attracted more and more attention, and also has been constantly modified by various authors to characterize more biological phenomena (see Cieřlak and Stinner [5], Cieřlak and Winkler [7], Ishida et al. [15], Painter and Hillen [27], Hillen and Painter [12], Wang et al. [34, 35], Winkler et al. [7, 13, 54, 43, 45, 44, 47], Zheng [58] and references therein for detailed results). For related works in this direction, we mention that a corresponding quasilinear version (see e.g. [31, 54, 60, 58, 61]), the logistic damping or the signal consumed by the cells, has been deeply investigated by Cieřlak and Stinner [5, 6], Tao and Winkler [31, 42, 54], and Zheng et al. [58, 66, 60, 67].

In various situations, however, the interaction of chemotactic movement of the gametes and the surrounding fluid is not negligible (see Tuval et al. [33]). In 2005, Tuval et al. ([33]) proposed the following prototypical **signal consuming** model (with tensor-valued sensitivity):

$$\begin{cases} n_t + u \cdot \nabla n = \Delta n - \nabla \cdot (nS(x, n, c)\nabla c), & x \in \Omega, t > 0, \\ c_t + u \cdot \nabla c = \Delta c - nf(c), & x \in \Omega, t > 0, \\ u_t + \kappa(u \cdot \nabla)u + \nabla P = \Delta u + n\nabla\phi, & x \in \Omega, t > 0, \\ \nabla \cdot u = 0, & x \in \Omega, t > 0, \end{cases} \quad (1.4)$$

where $f(c)$ denotes the **consumption** rate of the oxygen by the cells. Here S is a tensor-valued function or a scalar function which is the same as (2.1). The model (1.4) describes the interaction of oxygen-taxis bacteria with a surrounding incompressible viscous fluid in which the oxygen is dissolved. After this, assume that the chemotactic sensitivity $S(x, n, c) := S(c)$ is a scalar function. This kind of models have been studied by many researchers by making use of energy-type functionals (see e.g. Chae et. al. [4], Duan et. al. [8], Liu and Lorz [25, 26], Tao and Winkler [32, 46, 48, 50], Zhang and Zheng [57] and references therein). In fact, if $S(x, n, c) := S(c)$, Winkler ([46] and [48]) proved that in two-dimensional space

(1.4) admits a unique global classical solution which stabilizes to the spatially homogeneous equilibrium $(\frac{1}{|\Omega|} \int_{\Omega} n_0, 0, 0)$ in the large time limit. While in three-dimensional setting, he (see [50]) also showed that there exists a globally defined weak solution to (1.4).

Experiment [56] show that the chemotactic movement could be not directly along the signal gradient, but with a rotation, so that, the the corresponding chemotaxis-fluid system with tensor-valued sensitivity loses entropy-like functional structure, which gives rise to considerable mathematical difficulties during the process of analysis. The global solvability of corresponding initial value problem for chemotaxis-fluid system with tensor-valued sensitivity have been deeply investigated by Cao, Lankeit [3], Ishida [14], Wang et al. [36, 39] and Winkler [49].

If $-nf(c)$ in the c -equation is replaced by $-c+n$, and the u -equation is a (Navier-)Stokes equation, then (1.4) becomes the following chemotaxis-(Navier-)Stokes system in the context of signal **produced** other than consumed by cells (see [52, 38, 39, 40, 64, 16])

$$\begin{cases} n_t + u \cdot \nabla n = \Delta n - \nabla \cdot (nS(x, n, c) \cdot \nabla c), & x \in \Omega, t > 0, \\ c_t + u \cdot \nabla c = \Delta c - c + n, & x \in \Omega, t > 0, \\ u_t + \kappa(u \cdot \nabla)u + \nabla P = \Delta u + n\nabla\phi, & x \in \Omega, t > 0, \\ \nabla \cdot u = 0, & x \in \Omega, t > 0. \end{cases} \quad (1.5)$$

Due to the presence of the tensor-valued sensitivity $S(x, n, c)$ as well as the strongly nonlinear term $(u \cdot \nabla)u$ and lower regularity for n , the analysis of (1.5) with tensor-valued sensitivity began to flourish (see [52, 38, 39, 40, 64, 16]). In fact, the global boundedness of classical solutions to the Stokes-version ($\kappa = 0$ in the third equation of system (1.5)) of system (1.5) with the tensor-valued S satisfying $|S(x, n, c)| \leq C_S(1+n)^{-\alpha}$ with some $C_S > 0$ and $\alpha > 0$ which implies that the effect of chemotaxis is weakened when the cell density increases has been proved for any $\alpha > 0$ in two dimensions (see Wang and Xiang [39]) and for $\alpha > \frac{1}{2}$ in three dimensions (see Wang and Xiang [40]). Then Wang-Winkler-Xiang ([38]) further shows that when $\alpha > 0$ and $\Omega \subset R^2$ is a bounded **convex** domain with smooth boundary, system (1.5) possesses a global-in-time classical and bounded solution. Recently, Zheng ([62]) extends the results of [38] to the general bounded domain by some new entropy-energy estimates. More recently, by using new entropy-energy estimates, Zheng and Ke ([16]) presented the

existence of global and weak solutions for the system (1.5) under the assumption that S satisfies (1.2) and

$$|S(x, n, c)| \leq (1 + n)^{-\alpha} \text{ for all } (x, n, c) \in \Omega \times [0, \infty)^2$$

with $\alpha > \frac{1}{3}$, which, in light of the known results for the fluid-free system (see Horstmann and Winkler [13] and Bellomo et al. [2]), is an optimal restriction on α . For more works about the chemotaxis-(Navier-)Stokes models (1.5), we mention that a corresponding quasilinear version or the logistic damping has been deeply investigated by Zheng [59], Wang and Liu [24], Tao and Winkler [32], Wang et. al. [39, 40].

Other variants of the model (1.5) has been used in the mathematical study of coral broadcast spawning. In fact, Kiselev and Ryzhik ([19] and [18]) introduced the following Keller-Segel type system to model coral fertilization:

$$\begin{cases} \rho_t + u \cdot \nabla \rho = \Delta \rho - \chi \nabla \cdot (\rho \nabla c) - \varepsilon \rho^q, \\ 0 = \Delta c + \rho, \end{cases} \quad (1.6)$$

where ρ, u, χ and $-\varepsilon \rho^q$, respectively, denote the density of egg (sperm) gametes, the smooth divergence free sea fluid velocity as well as the positive chemotactic sensitivity constant and the reaction (fertilization) phenomenon. In fact, under suitable conditions, the global-in-time existence of the solution to (1.6) is presented by Kiselev and Ryzhik in [19]. Moreover, they proved that the total mass $m_0(t) = \int_{R^2} \rho(x, t) dx$ approaches a positive constant whose lower bound is $C(\chi, \rho_0, u)$ as $t \rightarrow \infty$ when $q > 2$. In the critical case of $N = q = 2$, a corresponding weaker but yet relevant effect within finite time intervals is detected (see [18]).

In order to analyze a further refinement of the model (1.6) which explicitly distinguishes between sperms and eggs, Espejo and Winkler ([10]) have recently considered the Navier-

Stokes version of (2.1):

$$\left\{ \begin{array}{l} n_t + u \cdot \nabla n = \Delta n - \nabla \cdot (n \nabla c) - nm, \quad x \in \Omega, t > 0, \\ c_t + u \cdot \nabla c = \Delta c - c + m, \quad x \in \Omega, t > 0, \\ m_t + u \cdot \nabla m = \Delta m - nm, \quad x \in \Omega, t > 0, \\ u_t + \kappa(u \cdot \nabla)u + \nabla P = \Delta u + (n + m)\nabla \phi, \quad x \in \Omega, t > 0, \\ \nabla \cdot u = 0, \quad x \in \Omega, t > 0 \end{array} \right. \quad (1.7)$$

in a bounded domain $\Omega \subset \mathbb{R}^2$. If $N = 2$, Espejo and Winkler ([10]) established the global existence of classical solutions to the associated initial-boundary value problem (1.7), which tend towards a spatially homogeneous equilibrium in the large time limit. Furthermore, if $S(x, n, c)$ satisfying (1.2) and (1.3) with $\alpha \geq \frac{1}{3}$ or $\alpha \geq 0$ and the initial data satisfy a certain smallness condition, Li-Pang-Wang ([22]) proved the same result for the three-dimensional Stokes ($\kappa = 0$ in the fourth equation of (1.1)) version of system (1.1). From [22], we know that $\alpha \geq \frac{1}{3}$ is enough to warrant the boundedness of solutions to system (2.1) for any large data (see Li-Pang-Wang [22]). We should point that the core step of [22] is to establish the estimates of the functional

$$\|n(\cdot, t)\|_{L^2(\Omega)}^2 + \|\nabla c(\cdot, t)\|_{L^2(\Omega)}^2 + \|u(\cdot, t)\|_{W^{1,2}(\Omega)}^2,$$

which strongly relies on $\alpha \geq \frac{1}{3}$ and $\kappa = 0$ (see the proof of Lemma 3.1 of [22]). To the best of our knowledge, it is yet unclear whether for $\alpha < \frac{1}{3}$ or $\kappa \neq 0$, the solutions of (2.1) exist (or even bounded) or not. Recently, relying on the functional

$$\left\{ \begin{array}{l} \int_{\Omega} n_{\varepsilon}^{4\alpha + \frac{2}{3}} + \int_{\Omega} |\nabla c_{\varepsilon}|^2 + \int_{\Omega} |u_{\varepsilon}|^2 \quad \text{if } \alpha \neq \frac{1}{12}, \\ \int_{\Omega} n_{\varepsilon} \ln n_{\varepsilon} + \int_{\Omega} |\nabla c_{\varepsilon}|^2 + \int_{\Omega} |u_{\varepsilon}|^2 \quad \text{if } \alpha = \frac{1}{12}, \end{array} \right.$$

we ([63]) presented the existence of global **weak** solutions for the system (1.1) under the assumption that S satisfies (1.2) and (1.3) with $\alpha > 0$. However, the existence of global (stronger than the result of [63]) **weak** solutions is still open. In this paper, by using a new weighted estimate (see Lemma 3.2), we try to obtain enough regularity and compactness properties (see Lemmas 3.2, 3.3, and 3.5), then show that system (1.1) possesses a globally defined **weak** solution, which improves the result of [63]. Therefore, collecting the above results, it is meaningful to analyze the following question:

Whether or not the assumption of α is optimal? Can we further relax the restriction on α , say, to $\alpha > 0$? Moreover, can we consider the regularity of global solution for system (1.1)?

Inspired by the above works, the first result of paper is to prove the existence of global (and bounded) solution for any $\alpha > 0$. Moreover, we also show that the corresponding solutions converge to a spatially homogeneous equilibrium exponentially as $t \rightarrow \infty$ as well.

Throughout this paper, we assume that

$$\phi \in W^{2,\infty}(\Omega) \quad (1.8)$$

and the initial data (n_0, c_0, u_0) fulfills

$$\begin{cases} n_0 \in C(\bar{\Omega}) & \text{with } n_0 \geq 0 \text{ and } n_0 \not\equiv 0, \\ c_0 \in W^{1,\infty}(\Omega) & \text{with } c_0 \geq 0 \text{ in } \bar{\Omega}, \\ m_0 \in C(\bar{\Omega}) & \text{with } m_0 \geq 0 \text{ and } m_0 \not\equiv 0, \\ u_0 \in D(A_r^\gamma) & \text{for some } \gamma \in (\frac{3}{4}, 1) \text{ and any } r \in (1, \infty), \end{cases} \quad (1.9)$$

where A_r denotes the Stokes operator with domain $D(A_r) := W^{2,r}(\Omega) \cap W_0^{1,r}(\Omega) \cap L_\sigma^r(\Omega)$, and $L_\sigma^r(\Omega) := \{\varphi \in L^r(\Omega) | \nabla \cdot \varphi = 0\}$ for $r \in (1, \infty)$ ([29]).

In the context of these assumptions, the first of our main results can be read as follows.

Theorem 1.1. *Let $\Omega \subset \mathbb{R}^3$ be a bounded domain with smooth boundary. Suppose that the assumptions (1.2)–(1.3) and (1.8)–(1.9) hold. If*

$$\alpha > 0, \quad (1.10)$$

then for any $\kappa \in \mathbb{R}$, there exist

$$\begin{cases} n \in L_{loc}^\infty([0, \infty), L^p(\Omega)) \cap L_{loc}^2([0, \infty), W^{1,2}(\Omega)) & \text{for any } p > 1, \\ c \in L^\infty(\Omega \times (0, \infty)) \cap L_{loc}^2([0, \infty), W^{2,2}(\Omega)) \cap L_{loc}^4([0, \infty), W^{1,4}(\Omega)), \\ m \in L^\infty(\Omega \times (0, \infty)) \cap L_{loc}^2([0, \infty), W^{2,2}(\Omega)) \cap L_{loc}^4([0, \infty), W^{1,4}(\Omega)), \\ u \in L_{loc}^2([0, \infty), W_{0,\sigma}^{1,2}(\Omega)) \cap L_{loc}^\infty([0, \infty), L^2(\Omega)), \end{cases} \quad (1.11)$$

such that (n, c, m, u) is a global weak solution of the problem (2.1) in the natural sense as specified in [63]. Moreover, if $\kappa = 0$, the problem (2.1) possesses at least one global classical

solution (n, c, m, u, P) . Moreover, this solution is bounded in $\Omega \times (0, \infty)$ in the sense that

$$\|n(\cdot, t)\|_{L^\infty(\Omega)} + \|c(\cdot, t)\|_{W^{1,\infty}(\Omega)} + \|m(\cdot, t)\|_{W^{1,\infty}(\Omega)} + \|u(\cdot, t)\|_{L^\infty(\Omega)} \leq C \text{ for all } t > 0. \quad (1.12)$$

Remark 1.1. (i) Theorem 1.1 indicates that $\alpha > 0$ and $\kappa = 0$ is enough to ensure the global existence and uniform boundedness of solution of the three-dimensional Keller-Segel-Stokes system (1.1), which improves the result obtained in [22], therein $\alpha \geq \frac{1}{3}$ is required.

(ii) This result also improves the result of our recent paper ([63]), where the more **weak** solution than our result was obtained by using different method (see [63]).

We can secondly prove that in fact any such **weak** solution (n, c, m, u) becomes smooth ultimately, and that it approaches the unique spatially homogeneous steady state $(\hat{n}, \hat{m}, \hat{m}, 0)$, where $\hat{n} = \frac{1}{|\Omega|} \{\int_\Omega n_0 - \int_\Omega m_0\}_+$ and $\hat{m} = \frac{1}{|\Omega|} \{\int_\Omega m_0 - \int_\Omega n_0\}_+$.

Theorem 1.2. *Under the assumptions of Theorem 1.1, then there are $T > 0$ and $\iota \in (0, 1)$ such that the solution (n, c, m, u) given by Theorem 1.1 satisfies*

$$n, c, m \in C^{2+\iota, 1+\frac{\iota}{2}}(\bar{\Omega} \times [T, \infty)), \quad u \in C^{2+\iota, 1+\frac{\iota}{2}}(\bar{\Omega} \times [T, \infty); \mathbb{R}^3).$$

Moreover,

$$n(\cdot, t) \rightarrow \hat{n}, \quad c(\cdot, t) \rightarrow \hat{m} \text{ as well as } m(\cdot, t) \rightarrow \hat{m} \text{ and } u(\cdot, t) \rightarrow 0 \text{ in } L^\infty(\Omega),$$

$$\text{where } \hat{n} = \frac{1}{|\Omega|} \{\int_\Omega n_0 - \int_\Omega m_0\}_+ \text{ and } \hat{m} = \frac{1}{|\Omega|} \{\int_\Omega m_0 - \int_\Omega n_0\}_+.$$

Remark 1.2. Theorem 1.1 indicates that if $\alpha > 0$, then for arbitrarily large initial data and for any $\kappa \in \mathbb{R}$, this problem admits at least one global weak solution for which there exists $T > 0$ such that (n, c, m, u) is smooth in $\Omega \times (T, \infty)$. Moreover, it is asserted that such solutions are shown to approach a spatially homogeneous equilibrium in the large time limit, which improves the result obtained in [22], therein $\kappa = 0$ is required.

Mathematical challenges for the regularity and stabilization of the solution for system (1.1). System (1.1) incorporates fluid and rotational flux, which involves more complex cross-diffusion mechanisms and brings about many considerable mathematical difficulties. Firstly, even when posed without any external influence, that is, $n = c = m \equiv 0$,

the corresponding Navier-Stokes system (1.1) does not admit a satisfactory solution theory up to now (see Leray [21] and Sohr [29], Wiegner [41]). As far as we know that the question of global solvability in classes of suitably regular functions yet remains open except in cases when the initial data are appropriately small (see e.g. Wiegner [41]). Moreover, the tensor-valued sensitivity functions result in new mathematical difficulties, mainly linked to the fact that a chemotaxis system with such rotational fluxes thereby loses an energy-like structure (see e.g. [56]). In [22] and [10], relying on globally **bounded** for the solution, Espejo-Winkler ([10]) Li-Pang-Wang ([22]) proved that all these solutions of problem (1.1) are shown to approach a spatially homogeneous equilibrium in the large time limit when $N = 2$ or $N = 3$ and $\kappa = 0$, respectively. As already mentioned in the above, in the case $N = 3$, it is not only unknown whether the incompressible Navier-Stokes equations possess global smooth solutions for arbitrarily large smooth initial data (see e.g. Wiegner [41] and Sohr [29]). Therefore, when $\kappa \neq 0$ and $N = 3$, we can not use the idea of [22] and [10] to discuss the large time behavior to problem (1.1), since, the globally **bounded** for the solutions are needed in [22] and [10].

In order to derive these theorems, in Section 2, we introduce the regularized system of (1.1), establish some basic estimates of the solutions and recall a local existence result. In Section 3, a key step of the proof of our main results is to establish a bound for $n_\varepsilon(\cdot, t)$ in $L^p(\Omega)$ for any $p > 1$. The approach is based on the weighted estimate of $\int_\Omega n_\varepsilon^p g(c_\varepsilon)$ with some weight function $g(c_\varepsilon)$ which is uniformly bounded both from above and below by positive constants. Here n_ε and c_ε are components of the solutions to (2.1) below. On the basis of the previously established estimates and the compactness properties thereby implied, we shall pass to the limit along an adequate sequence of numbers $\varepsilon = \varepsilon_j \searrow 0$ and thereby verify Theorem 1.1. Using the basic relaxation properties expressed in (2.13) and (2.14), Section 4 is devoted to showing the large time behavior of global solutions to (1.1) obtained in the above section. To this end, thanks to the decay property of $m_\varepsilon(\cdot, t) - \hat{m} + n_\varepsilon(\cdot, t) - \hat{n}$ formulated in Lemmas 4.6 and 4.8, this actually entails a certain eventual regularity and decay of u_ε also in the present situation, where $\hat{n} = \frac{1}{|\Omega|} \{ \int_\Omega n_0 - \int_\Omega m_0 \}_+$ and $\hat{m} = \frac{1}{|\Omega|} \{ \int_\Omega m_0 - \int_\Omega n_0 \}_+$. Using these bounds (see Lemmas 4.6–4.10), based on maximal Sobolev regularity

in the Stokes evolution system as well as inhomogeneous linear heat equations and the standard Schauder theory, we then prove eventual Hölder regularity and smoothness of solution $(n_\varepsilon, c_\varepsilon, m_\varepsilon, u_\varepsilon)$ (see Lemmas 4.11–4.12). For convergence as $t \rightarrow \infty$, we draw upon uniform Hölder bounds and smoothness for solution (n, c, m, u) (see Lemmas 4.13–4.14). Finally, applying an Ehrling-type lemma, we can prove any such solution approaches the spatially homogeneous equilibrium by using the above convergence properties (see Lemma 4.15).

2 Preliminaries

As mentioned in the introduction, the chemotactic sensitivity S in the first equation in (1.1) and the nonlinear convective term $\kappa(u \cdot \nabla)u$ in the Navier-Stokes subsystem of (1.1) bring about a great challenge to the study of system (1.1). To deal with these difficulties, according to the ideas in [50] (see also [53, 16, 51]), we first consider the approximate problems given by

$$\left\{ \begin{array}{l} n_{\varepsilon t} + u_\varepsilon \cdot \nabla n_\varepsilon = \Delta n_\varepsilon - \nabla \cdot (n_\varepsilon S_\varepsilon(x, n_\varepsilon, c_\varepsilon) \nabla c_\varepsilon) - n_\varepsilon m_\varepsilon, \quad x \in \Omega, t > 0, \\ c_{\varepsilon t} + u_\varepsilon \cdot \nabla c_\varepsilon = \Delta c_\varepsilon - c_\varepsilon + m_\varepsilon, \quad x \in \Omega, t > 0, \\ m_{\varepsilon t} + u_\varepsilon \cdot \nabla m_\varepsilon = \Delta m_\varepsilon - n_\varepsilon m_\varepsilon, \quad x \in \Omega, t > 0, \\ u_{\varepsilon t} + \nabla P_\varepsilon = \Delta u_\varepsilon - \kappa(Y_\varepsilon u_\varepsilon \cdot \nabla)u_\varepsilon + (n_\varepsilon + m_\varepsilon) \nabla \phi, \quad x \in \Omega, t > 0, \\ \nabla \cdot u_\varepsilon = 0, \quad x \in \Omega, t > 0, \\ \nabla n_\varepsilon \cdot \nu = \nabla c_\varepsilon \cdot \nu = \nabla m_\varepsilon \cdot \nu = 0, u_\varepsilon = 0, \quad x \in \partial\Omega, t > 0, \\ n_\varepsilon(x, 0) = n_0(x), c_\varepsilon(x, 0) = c_0(x), m_\varepsilon(x, 0) = m_0(x), u_\varepsilon(x, 0) = u_0(x), \quad x \in \Omega, \end{array} \right. \quad (2.1)$$

where

$$S_\varepsilon(x, n, c) = \rho_\varepsilon(x) \chi_\varepsilon(n) S(x, n, c), \quad x \in \bar{\Omega}, \quad n \geq 0, \quad c \geq 0 \quad (2.2)$$

and

$$Y_\varepsilon w := (1 + \varepsilon A)^{-1} w \quad \text{for all } w \in L_\sigma^2(\Omega) \quad (2.3)$$

is the standard Yosida approximation. Here $(\rho_\varepsilon)_{\varepsilon \in (0,1)} \in C_0^\infty(\Omega)$ and $(\chi_\varepsilon)_{\varepsilon \in (0,1)} \in C_0^\infty([0, \infty))$ are a family of functions which satisfy

$$0 \leq \rho_\varepsilon \leq 1 \quad \text{in } \Omega, \quad \rho_\varepsilon \nearrow 1 \quad \text{in } \Omega \quad \text{as } \varepsilon \searrow 0$$

and

$$0 \leq \chi_\varepsilon \leq 1 \quad \text{in } [0, \infty), \quad \chi_\varepsilon \nearrow 1 \quad \text{in } [0, \infty) \quad \text{as } \varepsilon \searrow 0.$$

Without essential difficulty, the local existence of approximate solutions to (2.1) can be easily proved according to the corresponding procedure in Lemma 2.1 of [50] (see also [49] and Lemma 2.1 of [27]). Therefore, we give the following lemma without proof.

Lemma 2.1. *Assume that $\varepsilon \in (0, 1)$. Then there exist $T_{max, \varepsilon} \in (0, \infty]$ and a classical solution $(n_\varepsilon, c_\varepsilon, m_\varepsilon, u_\varepsilon, P_\varepsilon)$ of (2.1) in $\Omega \times (0, T_{max, \varepsilon})$ such that*

$$\begin{cases} n_\varepsilon \in C^0(\bar{\Omega} \times [0, T_{max, \varepsilon})) \cap C^{2,1}(\bar{\Omega} \times (0, T_{max, \varepsilon})), \\ c_\varepsilon \in C^0(\bar{\Omega} \times [0, T_{max, \varepsilon})) \cap C^{2,1}(\bar{\Omega} \times (0, T_{max, \varepsilon})), \\ m_\varepsilon \in C^0(\bar{\Omega} \times [0, T_{max, \varepsilon})) \cap C^{2,1}(\bar{\Omega} \times (0, T_{max, \varepsilon})), \\ u_\varepsilon \in C^0(\bar{\Omega} \times [0, T_{max, \varepsilon})) \cap C^{2,1}(\bar{\Omega} \times (0, T_{max, \varepsilon})), \\ P_\varepsilon \in C^{1,0}(\bar{\Omega} \times (0, T_{max, \varepsilon})), \end{cases} \quad (2.4)$$

classically solving (2.1) in $\Omega \times [0, T_{max, \varepsilon})$. Moreover, $n_\varepsilon, c_\varepsilon$ and m_ε are nonnegative in $\Omega \times (0, T_{max, \varepsilon})$, and

$$\|n_\varepsilon(\cdot, t)\|_{L^\infty(\Omega)} + \|c_\varepsilon(\cdot, t)\|_{W^{1,\infty}(\Omega)} + \|m_\varepsilon(\cdot, t)\|_{W^{1,\infty}(\Omega)} + \|A^\gamma u_\varepsilon(\cdot, t)\|_{L^2(\Omega)} \rightarrow \infty \quad \text{as } t \nearrow T_{max, \varepsilon}, \quad (2.5)$$

where γ is given by (1.9).

Lemma 2.2. ([13, 43, 65]) *The Stokes operator A denotes the realization of the Stokes operator under homogeneous Dirichlet boundary conditions in the solenoidal subspace $L_\sigma^2(\Omega)$ of $L^2(\Omega)$. Let $\mathcal{P} : L^p(\Omega) \rightarrow L_\sigma^p(\Omega)$ stand for the Helmholtz projection in $L^p(\Omega)$. Then there exist positive constants $\kappa_i (i = 1, \dots, 3)$ such that*

$$\|e^{-tA} \mathcal{P} \varphi\|_{L^p(\Omega)} \leq \kappa_1(\Omega) t^{-\gamma} \|\varphi\|_{L^q(\Omega)} \quad \text{for all } t > 0 \quad \text{and any } \varphi \in L^q(\Omega) \quad (2.6)$$

as well as

$$\|e^{-tA} \mathcal{P} \nabla \cdot \varphi\|_{L^p(\Omega)} \leq \kappa_2(\Omega) t^{-\frac{1}{2} - \frac{3}{2}(\frac{1}{q} - \frac{1}{p})} \|\varphi\|_{L^q(\Omega)} \quad \text{for all } t > 0 \quad \text{and any } \varphi \in L^q(\Omega) \quad (2.7)$$

and

$$\|e^{-tA} \mathcal{P} \varphi\|_{L^p(\Omega)} \leq \kappa_3(\Omega) \|\varphi\|_{L^p(\Omega)} \quad \text{for all } t > 0 \quad \text{and any } \varphi \in L^p(\Omega). \quad (2.8)$$

Invoking the divergence free of the fluid and the homogeneous Neumann boundary conditions on $n_\varepsilon, m_\varepsilon$ and c_ε , we can establish the following basic estimates by using the maximum principle to the second and third equations. The proof of this lemma is very similar to that of Lemmas 2.2 and 2.6 of [32] (see also Lemma 3.2 of [37]), so we omit its proof here

Lemma 2.3. *There exists $\lambda > 0$ such that the solution of (2.1) satisfies*

$$\frac{d}{dt} \int_{\Omega} c_\varepsilon(\cdot, t) \leq 0, \quad \text{and} \quad \frac{d}{dt} \int_{\Omega} m_\varepsilon(\cdot, t) \leq 0 \quad \text{for all } t \in (0, T_{\max, \varepsilon}), \quad (2.9)$$

$$\|c_\varepsilon(\cdot, t)\|_{L^\infty(\Omega)} + \|m_\varepsilon(\cdot, t)\|_{L^\infty(\Omega)} \leq \lambda \quad \text{for all } t \in (0, T_{\max, \varepsilon}), \quad (2.10)$$

$$\int_{\Omega} n_\varepsilon - \int_{\Omega} m_\varepsilon = \int_{\Omega} n_0 - \int_{\Omega} m_0 \quad \text{for all } t \in (0, T_{\max, \varepsilon}), \quad (2.11)$$

$$\|c_\varepsilon(\cdot, t)\|_{L^2(\Omega)}^2 + 2 \int_0^t \int_{\Omega} |\nabla c_\varepsilon|^2 \leq \lambda \quad \text{for all } t \in (0, T_{\max, \varepsilon}), \quad (2.12)$$

$$\int_0^t \int_{\Omega} n_\varepsilon m_\varepsilon dx ds \leq \lambda \quad \text{for all } t \in (0, T_{\max, \varepsilon}) \quad (2.13)$$

as well as

$$\frac{1}{2} \int_0^t \int_{\Omega} |\nabla m_\varepsilon|^2 dx ds \leq \frac{1}{2} \int_{\Omega} m_0^2 \quad \text{for all } t \in (0, T_{\max, \varepsilon}) \quad (2.14)$$

and

$$\int_0^t \int_{\Omega} |\nabla c_\varepsilon|^2 \leq \lambda \quad \text{for all } t \in (0, T_{\max, \varepsilon}). \quad (2.15)$$

For simplicity, here and hereafter, we take the notations

$$C_S := \sup_{0 \leq s \leq \|c_0\|_{L^\infty(\Omega)}} S(s) \quad (2.16)$$

by using (2.10) and (1.3).

3 A-priori estimates

In this section we want to ensure that the time-local solutions obtained in Lemma 2.1 are in fact global solutions. To this end, for any $p > 1$, we firstly obtain boundedness of n_ε in $L^p(\Omega)$ under the assumption that $\alpha > 0$. Inspired by the weighted estimate argument developed in [44] (see also [48, 51]), we shall invoke a weight function $g(c_\varepsilon)$ which is uniformly bounded

from above and below by positive constants. Before deriving the uniform of L^p norm of n_ε , let us first recalling the well-known facts for $g(c_\varepsilon)$.

Lemma 3.1. *Let*

$$g(s) = e^{\beta s^2}, \quad \text{for any } s \in (0, \|c_0\|_{L^\infty(\Omega)}],$$

where

$$\beta = \frac{1}{8\|c_0\|_{L^\infty(\Omega)}^2} \quad (3.1)$$

and C_S is given by (2.16). Then for any $s \in (0, \|c_0\|_{L^\infty(\Omega)}]$,

$$1 \leq g(s) \leq \mu_0 := e^{\frac{1}{8}} \quad (3.2)$$

and

$$g'(s) \leq \mu_1 := \frac{1}{4\|c_0\|_{L^\infty}} e^{\frac{1}{8}}. \quad (3.3)$$

Proof. Obviously, (3.2) holds. On the other hand, a direct computation shows

$$g'(s) = 2\beta s e^{\beta s^2}. \quad (3.4)$$

This combined with the fact that $\beta = \frac{1}{8\|c_0\|_{L^\infty(\Omega)}^2}$ implies (3.3). \square

Lemma 3.2. *Let $\alpha > 0$. Then for any $p > 1$, there exists $C > 0$ such that the solution of (2.1) satisfies*

$$\int_{\Omega} n_\varepsilon^p \leq C \quad \text{for all } t \in (0, T_{\max, \varepsilon}). \quad (3.5)$$

Proof. Firstly, we define a functional

$$L(n_\varepsilon, c_\varepsilon) = \frac{1}{p} \int_{\Omega} n_\varepsilon^p g(c_\varepsilon),$$

where $p > \max\{1, 2\alpha\}$, $g(c_\varepsilon) = e^{\beta c_\varepsilon^2}$ and β is the same as (3.1). Using the first two equations

in (2.1), we find:

$$\begin{aligned}
& \frac{d}{dt} L(n_\varepsilon, c_\varepsilon) \\
&= \int_{\Omega} n_\varepsilon^{p-1} n_{\varepsilon t} g(c_\varepsilon) + \frac{1}{p} \int_{\Omega} n_\varepsilon^p g'(c_\varepsilon) c_{\varepsilon t} \\
&= \int_{\Omega} n_\varepsilon^{p-1} g(c_\varepsilon) (\Delta n_\varepsilon - \nabla \cdot (n_\varepsilon S_\varepsilon(x, n_\varepsilon, c_\varepsilon) \nabla c_\varepsilon) - u_\varepsilon \cdot \nabla n_\varepsilon - n_\varepsilon m_\varepsilon) \\
&\quad + \frac{1}{p} \int_{\Omega} n_\varepsilon^p g'(c_\varepsilon) (\Delta c_\varepsilon - c_\varepsilon + m_\varepsilon - u \cdot \nabla c_\varepsilon) \\
&= -\frac{1}{p} \int_{\Omega} n_\varepsilon^p g'(c_\varepsilon) u_\varepsilon \cdot \nabla c_\varepsilon - \int_{\Omega} n_\varepsilon^{p-1} g(c_\varepsilon) u_\varepsilon \cdot \nabla n_\varepsilon \\
&\quad - \frac{1}{p} \int_{\Omega} n_\varepsilon^p g'(c_\varepsilon) c_\varepsilon - \int_{\Omega} n_\varepsilon^p g(c_\varepsilon) m_\varepsilon + \frac{1}{p} \int_{\Omega} n_\varepsilon^p g'(c_\varepsilon) m_\varepsilon \\
&\quad + \int_{\Omega} n_\varepsilon^{p-1} g(c_\varepsilon) \Delta n_\varepsilon - \int_{\Omega} n_\varepsilon^{p-1} g(c_\varepsilon) \nabla \cdot (n_\varepsilon S_\varepsilon(x, n_\varepsilon, c_\varepsilon) \nabla c_\varepsilon) \\
&\quad + \frac{1}{p} \int_{\Omega} n_\varepsilon^p g'(c_\varepsilon) \Delta c_\varepsilon \\
&=: \sum_{i=1}^8 I_i \quad \text{for all } t \in (0, T_{max, \varepsilon}).
\end{aligned} \tag{3.6}$$

In the following, we will estimate the right-hand sides of (3.6) one by one. To this end, firstly, applying the elementary calculus identity

$$\frac{1}{p} n_\varepsilon^p g'(c_\varepsilon) \nabla c_\varepsilon + n_\varepsilon^{p-1} g(c_\varepsilon) \nabla n_\varepsilon = \frac{1}{p} \nabla (n_\varepsilon^p g(c_\varepsilon))$$

and the fact that

$$\nabla \cdot u_\varepsilon = 0, \quad x \in \Omega, t > 0,$$

we once more integrate by parts to find that

$$I_1 + I_2 = -\frac{1}{p} \int_{\Omega} n_\varepsilon^p g'(c_\varepsilon) u_\varepsilon \cdot \nabla c_\varepsilon - \int_{\Omega} n_\varepsilon^{p-1} g(c_\varepsilon) u_\varepsilon \cdot \nabla n_\varepsilon = 0 \tag{3.7}$$

by using $u_\varepsilon = 0, x \in \partial\Omega, t > 0$. Next, we derive from the non-negativity of g', g (see (3.4) and (3.2)), $c_\varepsilon, m_\varepsilon$ and n_ε that

$$I_3 + I_4 = -\frac{1}{p} \int_{\Omega} n_\varepsilon^p g'(c_\varepsilon) c_\varepsilon - \int_{\Omega} n_\varepsilon^p g(c_\varepsilon) m_\varepsilon \leq 0, \tag{3.8}$$

which combined with (3.6) and (3.7) yields

$$\begin{aligned}
& \frac{d}{dt} L(n_\varepsilon, c_\varepsilon) \\
&\leq \int_{\Omega} n_\varepsilon^{p-1} g(c_\varepsilon) \Delta n_\varepsilon - \int_{\Omega} n_\varepsilon^{p-1} g(c_\varepsilon) \nabla \cdot (n_\varepsilon S_\varepsilon(x, n_\varepsilon, c_\varepsilon) \nabla c_\varepsilon) + \frac{1}{p} \int_{\Omega} n_\varepsilon^p g'(c_\varepsilon) \Delta c_\varepsilon \\
&\quad + \frac{1}{p} \int_{\Omega} n_\varepsilon^p g'(c_\varepsilon) m_\varepsilon.
\end{aligned} \tag{3.9}$$

Now we proceed to estimate the fourth term on the right-hand side herein by using (2.10) and (3.3) to find that

$$\frac{1}{p} \int_{\Omega} n_{\varepsilon}^p g'(c_{\varepsilon}) m_{\varepsilon} \leq \frac{1}{p} \mu_1 \lambda \int_{\Omega} n_{\varepsilon}^p, \quad (3.10)$$

where μ_1 is the same as (3.3).

Now we estimate the term $\int_{\Omega} n_{\varepsilon}^{p-1} g(c_{\varepsilon}) \Delta n_{\varepsilon}$ and $\frac{1}{p} \int_{\Omega} n_{\varepsilon}^p g'(c_{\varepsilon}) \Delta c_{\varepsilon}$ in the right hand side of (3.9). In fact, we once more integrate by parts to see that

$$\begin{aligned} & \int_{\Omega} n_{\varepsilon}^{p-1} g(c_{\varepsilon}) \Delta n_{\varepsilon} + \frac{1}{p} \int_{\Omega} n_{\varepsilon}^p g'(c_{\varepsilon}) \Delta c_{\varepsilon} \\ &= -(p-1) \int_{\Omega} n_{\varepsilon}^{p-2} g(c_{\varepsilon}) |\nabla n_{\varepsilon}|^2 - \int_{\Omega} n_{\varepsilon}^{p-1} g'(c_{\varepsilon}) \nabla n_{\varepsilon} \cdot \nabla c_{\varepsilon} \\ & \quad - \frac{1}{p} \int_{\Omega} n_{\varepsilon}^p g''(c_{\varepsilon}) |\nabla c_{\varepsilon}|^2 - \int_{\Omega} n_{\varepsilon}^{p-1} g'(c_{\varepsilon}) \nabla c_{\varepsilon} \cdot \nabla n_{\varepsilon} \\ &\leq -(p-1) \int_{\Omega} n_{\varepsilon}^{p-2} g(c_{\varepsilon}) |\nabla n_{\varepsilon}|^2 - \frac{1}{p} \int_{\Omega} n_{\varepsilon}^p g''(c_{\varepsilon}) |\nabla c_{\varepsilon}|^2 \\ & \quad + 2 \int_{\Omega} n_{\varepsilon}^{p-1} g'(c_{\varepsilon}) |\nabla n_{\varepsilon}| |\nabla c_{\varepsilon}| \\ &\leq -\frac{3(p-1)}{4} \int_{\Omega} n_{\varepsilon}^{p-2} g(c_{\varepsilon}) |\nabla n_{\varepsilon}|^2 - \frac{1}{p} \int_{\Omega} n_{\varepsilon}^p g''(c_{\varepsilon}) |\nabla c_{\varepsilon}|^2 \\ & \quad + \frac{4}{p-1} \int_{\Omega} n_{\varepsilon}^p \frac{g'(c_{\varepsilon})^2}{g(c_{\varepsilon})} |\nabla c_{\varepsilon}|^2 \end{aligned} \quad (3.11)$$

by using the Young inequality. Next, recall (1.3), we can estimate second term on the right-hand side of (3.9) as follows:

$$\begin{aligned} & - \int_{\Omega} n_{\varepsilon}^{p-1} g(c_{\varepsilon}) \nabla \cdot (n_{\varepsilon} S_{\varepsilon}(x, n_{\varepsilon}, c_{\varepsilon}) \nabla c_{\varepsilon}) \\ &= (p-1) \int_{\Omega} n_{\varepsilon}^{p-1} g(c_{\varepsilon}) S(x, n_{\varepsilon}, c_{\varepsilon}) \nabla c_{\varepsilon} \cdot \nabla n_{\varepsilon} + \int_{\Omega} n_{\varepsilon}^p g'(c_{\varepsilon}) S(x, n_{\varepsilon}, c_{\varepsilon}) \nabla c_{\varepsilon} \cdot \nabla c_{\varepsilon} \\ &\leq (p-1) C_S \int_{\Omega} n_{\varepsilon}^{p-1-\alpha} g(c_{\varepsilon}) |\nabla c_{\varepsilon}| |\nabla n_{\varepsilon}| + C_S \int_{\Omega} n_{\varepsilon}^{p-\alpha} g'(c_{\varepsilon}) |\nabla c_{\varepsilon}|^2 \\ &\leq \frac{p-1}{4} \int_{\Omega} n_{\varepsilon}^{p-2} g(c_{\varepsilon}) |\nabla n_{\varepsilon}|^2 + C_S^2 (p-1) \int_{\Omega} n_{\varepsilon}^{p-2\alpha} g(c_{\varepsilon}) |\nabla c_{\varepsilon}|^2 \\ & \quad + C_S \int_{\Omega} n_{\varepsilon}^{p-\alpha} g'(c_{\varepsilon}) |\nabla c_{\varepsilon}|^2, \end{aligned} \quad (3.12)$$

where in the last inequality, we have used the Young inequality. Now, collecting (3.9)–(3.12), we may have

$$\begin{aligned} & \frac{d}{dt} L(n_{\varepsilon}, c_{\varepsilon}) + \frac{p-1}{2} \int_{\Omega} n_{\varepsilon}^{p-2} g(c_{\varepsilon}) |\nabla n_{\varepsilon}|^2 \\ & + \int_{\Omega} n_{\varepsilon}^p \left[\frac{1}{p} g''(c_{\varepsilon}) - \frac{4}{p-1} \frac{g'(c_{\varepsilon})^2}{g(c_{\varepsilon})} - C_S^2 (p-1) n_{\varepsilon}^{-2\alpha} g(c_{\varepsilon}) - C_S g'(c_{\varepsilon}) n_{\varepsilon}^{-\alpha} \right] |\nabla c_{\varepsilon}|^2 \\ &\leq \frac{1}{p} \mu_1 \lambda \int_{\Omega} n_{\varepsilon}^p, \end{aligned} \quad (3.13)$$

whence returning to the definition of $g(c_\varepsilon)$ we conclude that

$$\begin{aligned}
& \frac{d}{dt}L(n_\varepsilon, c_\varepsilon) + \frac{p-1}{2} \int_{\Omega} n_\varepsilon^{p-2} g(c_\varepsilon) |\nabla n_\varepsilon|^2 \\
& \leq - \int_{\Omega} n_\varepsilon^p e^{\beta c_\varepsilon^2} \left[\frac{1}{p} (4\beta^2 c_\varepsilon^2 + 2\beta) - \frac{16}{p-1} \beta^2 c_\varepsilon^2 - C_S^2 (p-1) n_\varepsilon^{-2\alpha} - 2C_S \beta c_\varepsilon n_\varepsilon^{-\alpha} \right] |\nabla c_\varepsilon|^2 \\
& \quad + \frac{1}{p} \mu_1 \lambda \int_{\Omega} n_\varepsilon^p \\
& \leq - \int_{\Omega} n_\varepsilon^p e^{\beta c_\varepsilon^2} \left[\frac{2\beta}{p} - \frac{16}{p-1} \beta^2 c_\varepsilon^2 - C_S^2 (p-1) n_\varepsilon^{-2\alpha} - 2C_S \beta c_\varepsilon n_\varepsilon^{-\alpha} \right] |\nabla c_\varepsilon|^2 \\
& \quad + \frac{1}{p} \mu_1 \lambda \int_{\Omega} n_\varepsilon^p \\
& \leq - \int_{\Omega} n_\varepsilon^p e^{\beta c_\varepsilon^2} \left[\frac{2\beta}{p} - \frac{16}{p-1} \beta^2 \|c_0\|_{L^\infty(\Omega)}^2 - C_S^2 (p-1) n_\varepsilon^{-2\alpha} - 2C_S \beta \|c_0\|_{L^\infty(\Omega)} n_\varepsilon^{-\alpha} \right] |\nabla c_\varepsilon|^2 \\
& \quad + \frac{1}{p} \mu_1 \lambda \int_{\Omega} n_\varepsilon^p \\
& \leq - \int_{\Omega} n_\varepsilon^p e^{\beta c_\varepsilon^2} \left[\frac{2\beta}{p} - \frac{8}{p} \beta^2 \|c_0\|_{L^\infty(\Omega)}^2 - C_S^2 p n_\varepsilon^{-2\alpha} - 2C_S \beta \|c_0\|_{L^\infty(\Omega)} n_\varepsilon^{-\alpha} \right] |\nabla c_\varepsilon|^2 \\
& \quad + \frac{1}{p} \mu_1 \lambda \int_{\Omega} n_\varepsilon^p
\end{aligned} \tag{3.14}$$

by using $p > \max\{1, 2\alpha\}$. In view of $\beta = \frac{1}{8\|c_0\|_{L^\infty(\Omega)}^2}$, thus, (3.14) implies that

$$\begin{aligned}
& \frac{d}{dt}L(n_\varepsilon, c_\varepsilon) + \frac{p-1}{2} \int_{\Omega} n_\varepsilon^{p-2} g(c_\varepsilon) |\nabla n_\varepsilon|^2 \\
& \leq - \int_{\Omega} n_\varepsilon^p e^{\beta c_\varepsilon^2} \left[\frac{\beta}{p} - C_S^2 p n_\varepsilon^{-2\alpha} - 2C_S \beta \|c_0\|_{L^\infty(\Omega)} n_\varepsilon^{-\alpha} \right] |\nabla c_\varepsilon|^2 + \frac{1}{p} \mu_1 \lambda \int_{\Omega} n_\varepsilon^p.
\end{aligned} \tag{3.15}$$

On the other hand, due to $\alpha > 0$, we may have

$$\lim_{s \rightarrow +\infty} [C_S^2 p s^{-2\alpha} + 2C_S \beta \|c_0\|_{L^\infty(\Omega)} s^{-\alpha}] = 0,$$

so that, there exists $\eta_0 > 0$, such that for any $s > \eta_0$,

$$[C_S^2 p s^{-2\alpha} + 2C_S \beta \|c_0\|_{L^\infty(\Omega)} s^{-\alpha}] < \frac{\beta}{2p}.$$

Therefore, by some basic calculation, we derive from (2.10) that

$$\begin{aligned}
& \int_{\Omega} n_{\varepsilon}^p e^{\beta c_{\varepsilon}^2} [C_S^2 p n_{\varepsilon}^{-2\alpha} + 2C_S \beta \|c_0\|_{L^{\infty}(\Omega)} n_{\varepsilon}^{-\alpha}] |\nabla c_{\varepsilon}|^2 \\
& \leq \int_{n_{\varepsilon} > \eta_0} n_{\varepsilon}^p e^{\beta c_{\varepsilon}^2} [C_S^2 p n_{\varepsilon}^{-2\alpha} + 2C_S \beta \|c_0\|_{L^{\infty}(\Omega)} n_{\varepsilon}^{-\alpha}] |\nabla c_{\varepsilon}|^2 \\
& \quad + \int_{n_{\varepsilon} \leq \eta_0} n_{\varepsilon}^p e^{\beta c_{\varepsilon}^2} [C_S^2 p n_{\varepsilon}^{-2\alpha} + 2C_S \beta \|c_0\|_{L^{\infty}(\Omega)} n_{\varepsilon}^{-\alpha}] |\nabla c_{\varepsilon}|^2 \\
& \leq \int_{n_{\varepsilon} > \eta_0} n_{\varepsilon}^p e^{\beta c_{\varepsilon}^2} \frac{\beta}{2p} |\nabla c_{\varepsilon}|^2 + \int_{n_{\varepsilon} \leq \eta_0} n_{\varepsilon}^p e^{\beta c_{\varepsilon}^2} [C_S^2 p n_{\varepsilon}^{-2\alpha} + 2C_S \beta \|c_0\|_{L^{\infty}(\Omega)} n_{\varepsilon}^{-\alpha}] |\nabla c_{\varepsilon}|^2 \quad (3.16) \\
& \leq \int_{\Omega} n_{\varepsilon}^p e^{\beta c_{\varepsilon}^2} \frac{\beta}{2p} |\nabla c_{\varepsilon}|^2 + \int_{n_{\varepsilon} \leq \eta_0} e^{\beta c_{\varepsilon}^2} [C_S^2 p n_{\varepsilon}^{p-2\alpha} + 2C_S \beta \|c_0\|_{L^{\infty}(\Omega)} n_{\varepsilon}^{p-\alpha}] |\nabla c_{\varepsilon}|^2 \\
& \leq \int_{\Omega} n_{\varepsilon}^p e^{\beta c_{\varepsilon}^2} \frac{\beta}{2p} |\nabla c_{\varepsilon}|^2 + \gamma_0 \int_{n_{\varepsilon} \leq \eta_0} |\nabla c_{\varepsilon}|^2 \\
& \leq \int_{\Omega} n_{\varepsilon}^p e^{\beta c_{\varepsilon}^2} \frac{\beta}{2p} |\nabla c_{\varepsilon}|^2 + \gamma_0 \int_{\Omega} |\nabla c_{\varepsilon}|^2
\end{aligned}$$

with

$$\gamma_0 = e^{\beta \|c_0\|_{L^{\infty}(\Omega)}^2} [C_S^2 p \eta_0^{p-2\alpha} + 2C_S \beta \|c_0\|_{L^{\infty}(\Omega)} \eta_0^{p-\alpha}]$$

by using (2.10) and $p > \max\{1, 2\alpha\}$. Substituting (3.16) into (3.15), we have

$$\begin{aligned}
& \frac{d}{dt} L(n_{\varepsilon}, c_{\varepsilon}) + \frac{p-1}{2} \int_{\Omega} n_{\varepsilon}^{p-2} g(c_{\varepsilon}) |\nabla n_{\varepsilon}|^2 + \int_{\Omega} n_{\varepsilon}^p e^{\beta c_{\varepsilon}^2} \frac{\beta}{2p} |\nabla c_{\varepsilon}|^2 \\
& \leq \gamma_0 \int_{\Omega} |\nabla c_{\varepsilon}|^2 + \frac{1}{p} \mu_1 \lambda \int_{\Omega} n_{\varepsilon}^p. \quad (3.17)
\end{aligned}$$

Now, according to (2.10), we therefore obtain on using the Gagliardo-Nirenberg inequality

that

$$\begin{aligned}
\int_{\Omega} n_{\varepsilon}^{p+\frac{1}{3}} &= \|n_{\varepsilon}^{\frac{p}{2}}\|_{L^{\frac{2(3p+1)}{3p}}(\Omega)}^{\frac{2(3p+1)}{3p}} \\
&\leq C_1 [\|\nabla n_{\varepsilon}^{\frac{p}{2}}\|_{L^{\frac{2}{p}}(\Omega)}^{\frac{2(3p-2)}{3p-1}} \|n_{\varepsilon}^{\frac{p}{2}}\|_{L^{\frac{2}{p}}(\Omega)}^{\frac{2(3p+1)}{3p} - \frac{2(3p-2)}{3p-1}} + \|n_{\varepsilon}^{\frac{p}{2}}\|_{L^{\frac{2}{p}}(\Omega)}^{\frac{2(3p+1)}{3p}}] \\
&\leq \frac{(p-1)}{4} \frac{4}{p^2} \|n_{\varepsilon}^{\frac{p}{2}}\|_{L^2(\Omega)}^2 + C_2 \quad (3.18) \\
&= \frac{(p-1)}{4} \int_{\Omega} n_{\varepsilon}^{p-2} |\nabla n_{\varepsilon}|^2 + C_2 \\
&\leq \frac{(p-1)}{4} \int_{\Omega} n_{\varepsilon}^{p-2} g(c_{\varepsilon}) |\nabla n_{\varepsilon}|^2 + C_2
\end{aligned}$$

for some positive constants C_1 and C_2 , where in the last inequality, we have used (3.2).

Collecting (3.17) and (3.18), we have

$$\begin{aligned}
& \frac{d}{dt} L(n_{\varepsilon}, c_{\varepsilon}) + \frac{p-1}{4} \int_{\Omega} n_{\varepsilon}^{p-2} g(c_{\varepsilon}) |\nabla n_{\varepsilon}|^2 + \int_{\Omega} n_{\varepsilon}^p e^{\beta c_{\varepsilon}^2} \frac{\beta}{2p} |\nabla c_{\varepsilon}|^2 + \int_{\Omega} n_{\varepsilon}^{p+\frac{1}{3}} \\
& \leq \gamma_0 \int_{\Omega} |\nabla c_{\varepsilon}|^2 + \frac{1}{p} \mu_1 \lambda \int_{\Omega} n_{\varepsilon}^p + C_2 \quad (3.19) \\
& \leq \gamma_0 \int_{\Omega} |\nabla c_{\varepsilon}|^2 + \frac{1}{2} \int_{\Omega} n_{\varepsilon}^{p+\frac{1}{3}} + C_3
\end{aligned}$$

by using the Young inequality. Therefore,

$$\begin{aligned} & \frac{d}{dt}L(n_\varepsilon, c_\varepsilon) + \frac{p-1}{4} \int_{\Omega} n_\varepsilon^{p-2} g(c_\varepsilon) |\nabla n_\varepsilon|^2 + \int_{\Omega} n_\varepsilon^p e^{\beta c_\varepsilon^2} \frac{\beta}{2p} |\nabla c_\varepsilon|^2 + \frac{1}{2} \int_{\Omega} n_\varepsilon^{p+\frac{1}{3}} \\ & \leq \gamma_0 \int_{\Omega} |\nabla c_\varepsilon|^2 + C_3. \end{aligned} \quad (3.20)$$

To track the time evolution of c_ε , testing the second equation in (2.1) by c_ε and using $\nabla \cdot u_\varepsilon = 0$ and (2.10) yields that for some positive constant C_4 such that

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \|c_\varepsilon\|_{L^2(\Omega)}^2 + \int_{\Omega} |\nabla c_\varepsilon|^2 &= - \int_{\Omega} c_\varepsilon^2 + \int_{\Omega} m_\varepsilon c_\varepsilon \\ &\leq -\frac{1}{2} \int_{\Omega} c_\varepsilon^2 + C_4 \quad \text{for all } t \in (0, T_{\max, \varepsilon}), \end{aligned} \quad (3.21)$$

wereafter integrating the above inequality in time yields

$$\int_t^{t+1} \int_{\Omega} |\nabla c_\varepsilon|^2 \leq C_5 \quad \text{for all } t > 0 \quad (3.22)$$

for some $C_5 > 0$ by an integration. This yields to

$$\int_t^{t+\tau} \left[\gamma_0 \int_{\Omega} |\nabla c_\varepsilon|^2 + C_3 \right] \leq C_6 \quad \text{for all } t \in (0, T_{\max, \varepsilon} - \tau) \quad (3.23)$$

with

$$\tau := \min\left\{1, \frac{1}{6} T_{\max, \varepsilon}\right\}. \quad (3.24)$$

Finally, (3.23) in conjunction with Lemma 2.3 of [38] (see also [62]) and (3.20) establish (3.25).

□

In a straightforward manner, the estimates gained above can be seen to imply the following ε -independent estimates, which plays an important role in proving Theorem 1.1.

Lemma 3.3. *Let $\alpha > 0$. Then there exists $C > 0$ such that the solution of (2.1) satisfies*

$$\int_{\Omega} (|\nabla c_\varepsilon|^2 + |\nabla m_\varepsilon|^2) \leq C \quad \text{for all } t \in (0, T_{\max, \varepsilon}). \quad (3.25)$$

Moreover, for $t \in (0, T_{\max, \varepsilon} - \tau)$, it holds that

$$\int_t^{t+\tau} \int_{\Omega} \left[|\nabla c_\varepsilon|^4 + |\nabla m_\varepsilon|^4 + |\nabla u_\varepsilon|^2 + |u_\varepsilon|^{\frac{10}{3}} \right] \leq C, \quad (3.26)$$

where τ is the same as (3.24).

Proof. We multiply the second equation in (2.1) by $-\Delta c_\varepsilon$ and integrate by parts to see that

$$\begin{aligned}
& \frac{1}{2} \frac{d}{dt} \|\nabla c_\varepsilon\|_{L^2(\Omega)}^2 + \int_{\Omega} |\Delta c_\varepsilon|^2 + \int_{\Omega} |\nabla c_\varepsilon|^2 \\
&= - \int_{\Omega} m_\varepsilon \Delta c_\varepsilon + \int_{\Omega} (u_\varepsilon \cdot \nabla c_\varepsilon) \Delta c_\varepsilon \\
&= - \int_{\Omega} m_\varepsilon \Delta c_\varepsilon - \int_{\Omega} \nabla c_\varepsilon \nabla (u_\varepsilon \cdot \nabla c_\varepsilon) \\
&= - \int_{\Omega} m_\varepsilon \Delta c_\varepsilon - \int_{\Omega} \nabla c_\varepsilon \nabla (\nabla u_\varepsilon \cdot \nabla c_\varepsilon),
\end{aligned} \tag{3.27}$$

where we have used the fact that

$$\int_{\Omega} \nabla c_\varepsilon \cdot (D^2 c_\varepsilon \cdot u_\varepsilon) = \frac{1}{2} \int_{\Omega} u_\varepsilon \cdot \nabla |\nabla c_\varepsilon|^2 = 0 \quad \text{for all } t \in (0, T_{max, \varepsilon}).$$

On the other hand, by the Young inequality and (2.14),

$$\begin{aligned}
- \int_{\Omega} m_\varepsilon \Delta c_\varepsilon &\leq \int_{\Omega} m_\varepsilon^2 + \frac{1}{4} \int_{\Omega} |\Delta c_\varepsilon|^2 \\
&\leq |\Omega| \|m_0\|_{L^\infty(\Omega)}^2 + \frac{1}{4} \int_{\Omega} |\Delta c_\varepsilon|^2 \quad \text{for all } t \in (0, T_{max, \varepsilon}).
\end{aligned} \tag{3.28}$$

In the last summand in (3.27), we use the Cauchy-Schwarz inequality to obtain

$$- \int_{\Omega} \nabla c_\varepsilon \nabla (\nabla u_\varepsilon \cdot \nabla c_\varepsilon) \leq \|\nabla u_\varepsilon\|_{L^2(\Omega)} \|\nabla c_\varepsilon\|_{L^4(\Omega)}^2 \quad \text{for all } t \in (0, T_{max, \varepsilon}). \tag{3.29}$$

Now thanks to (2.10) and in view of the Gagliardo-Nirenberg inequality, we can find $C_1 > 0$ and $C_2 > 0$ fulfilling integrate by parts to find that

$$\begin{aligned}
\|\nabla c_\varepsilon\|_{L^4(\Omega)}^2 &\leq C_1 \|\Delta c_\varepsilon\|_{L^2(\Omega)} \|c_\varepsilon\|_{L^\infty(\Omega)} + C_1 \|c_\varepsilon\|_{L^\infty(\Omega)}^4 \\
&\leq C_2 \|\Delta c_\varepsilon\|_{L^2(\Omega)} + C_2 \quad \text{for all } t \in (0, T_{max, \varepsilon}).
\end{aligned} \tag{3.30}$$

This, together with the Young inequality, yields

$$\begin{aligned}
& - \int_{\Omega} \nabla c_\varepsilon \nabla (\nabla u_\varepsilon \cdot \nabla c_\varepsilon) \\
&\leq \|\nabla u_\varepsilon\|_{L^2(\Omega)} [C_2 \|\Delta c_\varepsilon\|_{L^2(\Omega)} + C_2] \\
&\leq C_2^2 \|\nabla u_\varepsilon\|_{L^2(\Omega)}^2 + \frac{1}{4} \|\Delta c_\varepsilon\|_{L^2(\Omega)}^2 + C_3 \quad \text{for all } t \in (0, T_{max, \varepsilon}).
\end{aligned} \tag{3.31}$$

Inserting (3.28) and (3.31) into (3.27), we have

$$\begin{aligned}
& \frac{1}{2} \frac{d}{dt} \|\nabla c_\varepsilon\|_{L^2(\Omega)}^2 + \frac{1}{2} \int_{\Omega} |\Delta c_\varepsilon|^2 + \int_{\Omega} |\nabla c_\varepsilon|^2 \\
&\leq C_2^2 \|\nabla u_\varepsilon\|_{L^2(\Omega)}^2 + C_4.
\end{aligned} \tag{3.32}$$

Now, multiplying the third equation of (2.1) by u_ε , integrating by parts and using $\nabla \cdot u_\varepsilon = 0$

$$\frac{1}{2} \frac{d}{dt} \int_{\Omega} |u_\varepsilon|^2 + \int_{\Omega} |\nabla u_\varepsilon|^2 = \int_{\Omega} (n_\varepsilon + m_\varepsilon) u_\varepsilon \cdot \nabla \phi \quad \text{for all } t \in (0, T_{max, \varepsilon}). \quad (3.33)$$

Here we use the Hölder inequality, the Young inequality and the continuity of the embedding $W^{1,2}(\Omega) \hookrightarrow L^6(\Omega)$ and to find C_5 and $C_6 > 0$ such that

$$\begin{aligned} \int_{\Omega} (n_\varepsilon + m_\varepsilon) u_\varepsilon \cdot \nabla \phi &\leq \|\nabla \phi\|_{L^\infty(\Omega)} \|n_\varepsilon\|_{L^{\frac{6}{5}}(\Omega)} \|u_\varepsilon\|_{L^6(\Omega)} + \|\nabla \phi\|_{L^\infty(\Omega)} \|m_\varepsilon\|_{L^{\frac{6}{5}}(\Omega)} \|u_\varepsilon\|_{L^6(\Omega)} \\ &\leq C_5 \|\nabla \phi\|_{L^\infty(\Omega)} (\|n_\varepsilon\|_{L^{\frac{6}{5}}(\Omega)} + \|m_\varepsilon\|_{L^{\frac{6}{5}}(\Omega)}) \|\nabla u_\varepsilon\|_{L^2(\Omega)} \\ &\leq \frac{1}{2} \|\nabla u_\varepsilon\|_{L^2(\Omega)}^2 + C_6 \quad \text{for all } t \in (0, T_{max, \varepsilon}) \end{aligned} \quad (3.34)$$

by using (3.25) and (2.10). Inserting (3.34) into (3.33) and integrating in time to see that

$$\int_t^{t+\tau} \int_{\Omega} |\nabla u_\varepsilon|^2 \leq C_7 \quad \text{for all } t \in (0, T_{max, \varepsilon} - \tau) \quad (3.35)$$

and

$$\int_{\Omega} u_\varepsilon^2 \leq C_7 \quad \text{for all } t \in (0, T_{max, \varepsilon}), \quad (3.36)$$

where we use that once more employing the Gagliardo-Nirenberg inequality, the Hölder inequality and the Young inequality we can find $C_8 > 0$ and $C_9 > 0$ satisfying

$$\begin{aligned} \int_t^{t+\tau} \int_{\Omega} |u_\varepsilon|^{\frac{10}{3}} &= \int_t^{t+\tau} \|u_\varepsilon\|_{L^{\frac{10}{3}}(\Omega)}^{\frac{10}{3}} \\ &\leq C_8 \int_t^{t+\tau} \left(\|\nabla u_\varepsilon\|_{L^2(\Omega)}^2 \|u_\varepsilon\|_{L^2(\Omega)}^{\frac{4}{3}} + \|u_\varepsilon\|_{L^2(\Omega)}^{\frac{10}{3}} \right) \\ &\leq C_9 \quad \text{for all } t \in (0, T_{max, \varepsilon} - \tau). \end{aligned} \quad (3.37)$$

Next, combining (3.32), (3.35) and rearranging shows that

$$\int_t^{t+\tau} \int_{\Omega} (|\Delta c_\varepsilon|^2 + |\nabla c_\varepsilon|^4) \leq C_{10} \quad \text{for all } t \in (0, T_{max, \varepsilon} - \tau) \quad (3.38)$$

and

$$\int_{\Omega} |\nabla c_\varepsilon|^2 \leq C_{10} \quad \text{for all } t \in (0, T_{max, \varepsilon}) \quad (3.39)$$

by using (3.30).

Testing the third equation in (2.1) by $-\Delta m_\varepsilon$ and integrating by parts and using (3.25) and (3.35), one can finally derive

$$\int_t^{t+\tau} \int_{\Omega} (|\Delta m_\varepsilon|^2 + |\nabla m_\varepsilon|^4) \leq C_{11} \quad \text{for all } t \in (0, T_{max, \varepsilon} - \tau) \quad (3.40)$$

and

$$\int_{\Omega} |\nabla m_{\varepsilon}|^2 \leq C_{11} \quad \text{for all } t \in (0, T_{max, \varepsilon}). \quad (3.41)$$

□

With Lemmas 2.3 and 3.2–3.3 at hand, we can proceed to show that our approximate solutions are actually global in time.

Lemma 3.4. *For any $\varepsilon > 0$, then one can find $C > 0$ such that the solutions of (2.1) fulfill*

Proof. Firstly, under the assumption that $T_{max, \varepsilon} < \infty$, for any $\varepsilon > 0$, Lemmas 2.3–3.3 would provide us with $C_1 > 0$ such that

$$\int_{\Omega} n_{\varepsilon}^p \leq C_1 \quad \text{for all } t \in (0, T_{max, \varepsilon}) \quad \text{for all } p > 4, \quad (3.42)$$

$$\|c_{\varepsilon}(\cdot, t)\|_{L^{\infty}(\Omega)} \leq C_1 \quad \text{for all } t \in (0, T_{max, \varepsilon}) \quad (3.43)$$

as well as

$$\|m_{\varepsilon}(\cdot, t)\|_{L^{\infty}(\Omega)} \leq C_1 \quad \text{for all } t \in (0, T_{max, \varepsilon}) \quad (3.44)$$

and

$$\int_t^{t+\tau} \int_{\Omega} [|\nabla c_{\varepsilon}|^4 + |\nabla m|^4] \leq C_1. \quad (3.45)$$

Then, aided by the L^2 -estimate for ∇u_{ε} (from a testing argument), we can obtain that

$$\|A^{\gamma} u_{\varepsilon}(\cdot, t)\|_{L^2(\Omega)} \leq C_2 \quad \text{for all } t \in (0, T_{max, \varepsilon}) \quad (3.46)$$

with some $C_2 = C_2(\varepsilon) > 0$. Thus, the continuous embedding $D(A^{\gamma}) \hookrightarrow L^{\infty}(\Omega)$ implies the L^{∞} -estimate for u_{ε} . Therefore, employing the same arguments as in the proof of Lemma 3.2 in [63] (see also [16, 61, 49, 50]), and taking advantage of (3.42)–(3.46), we conclude the estimates

$$\|n_{\varepsilon}(\cdot, t)\|_{L^{\infty}(\Omega)} \leq C_3 \quad \text{for all } t \in (0, \infty) \quad (3.47)$$

as well as

$$\|c_{\varepsilon}(\cdot, t)\|_{W^{1, \infty}(\Omega)} \leq C_3 \quad \text{for all } t \in (0, \infty) \quad (3.48)$$

and

$$\|m_{\varepsilon}(\cdot, t)\|_{W^{1, \infty}(\Omega)} \leq C_3 \quad \text{for all } t \in (0, \infty) \quad (3.49)$$

and some positive constant C_3 . In view of (3.46)–(3.49), we apply Lemma 2.1 to reach a contradiction. □

3.1 Further a-priori estimates

With the help of Lemma 3.2 and the Gagliardo–Nirenberg inequality, one can derive the following Lemma:

Lemma 3.5. *Let $\alpha > 0$. Then for each $T \in (0, T_{max,\varepsilon})$, there exists $C > 0$ independent of ε such that the solution of (2.1) satisfies*

$$\int_0^T \int_{\Omega} |\nabla n_{\varepsilon}|^2 \leq C(T+1). \quad (3.50)$$

Proof. Multiply the first equation in (2.1) by n_{ε} and using $\nabla \cdot u_{\varepsilon} = 0$, we derive

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \|n_{\varepsilon}\|_{L^2(\Omega)}^2 + \int_{\Omega} |\nabla n_{\varepsilon}|^2 \\ &= - \int_{\Omega} n_{\varepsilon} \nabla \cdot (n_{\varepsilon} S_{\varepsilon}(x, n_{\varepsilon}, c_{\varepsilon}) \cdot \nabla c_{\varepsilon}) - \int_{\Omega} n_{\varepsilon}^2 m_{\varepsilon} \\ &\leq \int_{\Omega} n_{\varepsilon} |S_{\varepsilon}(x, n_{\varepsilon}, c_{\varepsilon})| |\nabla n_{\varepsilon}| |\nabla c_{\varepsilon}| \quad \text{for all } t \in (0, T_{max,\varepsilon}). \end{aligned} \quad (3.51)$$

Recalling (1.3) and using $\alpha \geq 0$, from Young inequality again, we derive from Lemma 3.2 that

$$\begin{aligned} & \int_{\Omega} n_{\varepsilon} |S_{\varepsilon}(x, n_{\varepsilon}, c_{\varepsilon})| |\nabla n_{\varepsilon}| |\nabla c_{\varepsilon}| \\ &\leq C_S \int_{\Omega} n_{\varepsilon} |\nabla n_{\varepsilon}| |\nabla c_{\varepsilon}| \\ &\leq \frac{1}{2} \int_{\Omega} |\nabla n_{\varepsilon}|^2 + \frac{C_S^2}{2} \int_{\Omega} n_{\varepsilon}^2 |\nabla c_{\varepsilon}|^2 \\ &\leq \frac{1}{2} \int_{\Omega} |\nabla n_{\varepsilon}|^2 + \frac{1}{4} \int_{\Omega} n_{\varepsilon}^4 + \frac{C_S^4}{4} \int_{\Omega} |\nabla c_{\varepsilon}|^4 \\ &\leq \frac{1}{2} \int_{\Omega} |\nabla n_{\varepsilon}|^2 + \frac{C_S^4}{4} \int_{\Omega} |\nabla c_{\varepsilon}|^4 + C_1 \quad \text{for all } t \in (0, T_{max,\varepsilon}), \end{aligned} \quad (3.52)$$

which combined with (3.51) implies that

$$\frac{1}{2} \frac{d}{dt} \|n_{\varepsilon}\|_{L^2(\Omega)}^2 + \frac{1}{2} \int_{\Omega} |\nabla n_{\varepsilon}|^2 \leq \frac{C_S^4}{4} \int_{\Omega} |\nabla c_{\varepsilon}|^4 + C_1 \quad \text{for all } t \in (0, T_{max,\varepsilon}), \quad (3.53)$$

so that, gather (3.26) and (3.53), one can get (3.50). □

4 Passing to the limit: The proof of Theorem 1.1

With the help of a priori estimates, in this subsection, by means of a standard extraction procedure we can now derive the following lemma which actually contains our main existence result (Theorem 1.1) already.

Lemma 4.1. *Assume that $\alpha > 0$. Then for any $\kappa \in \mathbb{R}$, there exists $(\varepsilon_j)_{j \in \mathbb{N}} \subset (0, 1)$ such that $\varepsilon_j \rightarrow 0$ as $j \rightarrow \infty$ and that*

$$n_\varepsilon \rightarrow n \text{ a.e. in } \Omega \times (0, \infty) \text{ and in } L^2_{loc}(\bar{\Omega} \times [0, \infty)) \quad (4.1)$$

$$n_\varepsilon \rightharpoonup n \text{ weak star in } L^\infty_{loc}([0, \infty), L^p(\Omega)) \text{ for any } p > 1, \quad (4.2)$$

$$\nabla n_\varepsilon \rightharpoonup \nabla n \text{ in } L^2_{loc}(\bar{\Omega} \times [0, \infty)), \quad (4.3)$$

$$c_\varepsilon \rightarrow c \text{ in } L^2_{loc}(\bar{\Omega} \times [0, \infty)) \text{ and a.e. in } \Omega \times (0, \infty), \quad (4.4)$$

$$m_\varepsilon \rightarrow m \text{ in } L^2_{loc}(\bar{\Omega} \times [0, \infty)) \text{ and a.e. in } \Omega \times (0, \infty), \quad (4.5)$$

$$\nabla c_\varepsilon \rightharpoonup \nabla c \text{ in } L^4_{loc}(\bar{\Omega} \times [0, \infty)), \quad (4.6)$$

$$\nabla m_\varepsilon \rightharpoonup \nabla m \text{ in } L^4_{loc}(\bar{\Omega} \times [0, \infty)), \quad (4.7)$$

$$u_\varepsilon \rightarrow u \text{ in } L^2_{loc}(\bar{\Omega} \times [0, \infty)) \text{ and a.e. in } \Omega \times (0, \infty), \quad (4.8)$$

$$\nabla c_\varepsilon \rightharpoonup \nabla c \text{ in } L^2_{loc}(\bar{\Omega} \times [0, \infty)) \quad (4.9)$$

as well as

$$\nabla m_\varepsilon \rightharpoonup \nabla m \text{ in } L^2_{loc}(\bar{\Omega} \times [0, \infty)) \quad (4.10)$$

and

$$\nabla u_\varepsilon \rightharpoonup \nabla u \text{ in } L^2_{loc}(\bar{\Omega} \times [0, \infty); \mathbb{R}^3) \quad (4.11)$$

some quadruple (n, c, m, u) which is a global weak solution of (1.1) in the natural sense as specified in [63].

Proof. Firstly, applying the discussion in Section 3, under the assumptions of Theorem 1.1, for each $T > 0$, we can find ε -independent constant $C(T)$ such that

$$\|n_\varepsilon(\cdot, t)\|_{L^p(\Omega)} + \|c_\varepsilon(\cdot, t)\|_{W^{1,2}(\Omega)} + \|m_\varepsilon(\cdot, t)\|_{W^{1,2}(\Omega)} + \|u_\varepsilon(\cdot, t)\|_{L^2(\Omega)} \leq C(T) \text{ for all } t \in (0, T) \text{ and } p > 1 \quad (4.12)$$

as well as

$$\int_0^T \int_{\Omega} (|\nabla n_{\varepsilon}|^2 + |\nabla c_{\varepsilon}|^4 + |\nabla m_{\varepsilon}|^4 + |\Delta c_{\varepsilon}|^2 + |\Delta m_{\varepsilon}|^2) \leq C(T) \quad \text{for all } t \in (0, T) \quad (4.13)$$

and

$$\int_0^T \int_{\Omega} |\nabla u_{\varepsilon}|^2 \leq C(T) \quad \text{for all } t \in (0, T). \quad (4.14)$$

Now, choosing $\varphi \in W^{1,2}(\Omega)$ as a test function in the first equation in (2.1) and using (4.12), we have

$$\begin{aligned} & \left| \int_{\Omega} (n_{\varepsilon,t}) \varphi \right| \\ &= \left| \int_{\Omega} [\Delta n_{\varepsilon} - \nabla \cdot (n_{\varepsilon} S_{\varepsilon}(x, n_{\varepsilon}, c_{\varepsilon}) \nabla c_{\varepsilon}) - u_{\varepsilon} \cdot \nabla n_{\varepsilon} - n_{\varepsilon} m_{\varepsilon}] \varphi \right| \\ &= \left| \int_{\Omega} [-\nabla n_{\varepsilon} \cdot \nabla \varphi + n_{\varepsilon} S_{\varepsilon}(x, n_{\varepsilon}, c_{\varepsilon}) \nabla c_{\varepsilon} \cdot \nabla \varphi + n_{\varepsilon} u_{\varepsilon} \cdot \nabla \varphi - n_{\varepsilon} m_{\varepsilon} \varphi] \right| \\ &\leq \left\{ \|\nabla n_{\varepsilon}\|_{L^2(\Omega)} + \|n_{\varepsilon} S_{\varepsilon}(x, n_{\varepsilon}, c_{\varepsilon}) \nabla c_{\varepsilon}\|_{L^2(\Omega)} + \|n_{\varepsilon} u_{\varepsilon}\|_{L^2(\Omega)} + \|n_{\varepsilon} m_{\varepsilon}\|_{L^2(\Omega)} \right\} \\ &\quad \times \|\varphi\|_{W^{1,2}(\Omega)} \end{aligned} \quad (4.15)$$

for all $t > 0$. Along with (4.12) and (4.13), further implies that

$$\begin{aligned} & \int_0^T \|\partial_t n_{\varepsilon}(\cdot, t)\|_{(W^{1,2}(\Omega))^*}^2 dt \\ &\leq \int_0^T \left\{ \|\nabla n_{\varepsilon}\|_{L^2(\Omega)} + \|n_{\varepsilon} S_{\varepsilon}(x, n_{\varepsilon}, c_{\varepsilon}) \nabla c_{\varepsilon}\|_{L^2(\Omega)} + \|n_{\varepsilon} u_{\varepsilon}\|_{L^2(\Omega)} \right\}^2 dt \\ &\leq C_1 \int_0^T \left\{ \|\nabla n_{\varepsilon}\|_{L^2(\Omega)}^2 + \|n_{\varepsilon}\|_{L^4(\Omega)}^4 \|\nabla c_{\varepsilon}\|_{L^4(\Omega)}^4 \right\} dt \\ &\quad + C_1 \int_0^T \left\{ \|n_{\varepsilon} u_{\varepsilon}\|_{L^2(\Omega)}^2 + \|m_{\varepsilon}\|_{L^\infty(\Omega)}^2 \|n_{\varepsilon}\|_{L^2(\Omega)}^2 \right\} dt \\ &\leq C_2 \int_0^T \left\{ \|\nabla n_{\varepsilon}\|_{L^2(\Omega)}^2 + \|\nabla c_{\varepsilon}\|_{L^4(\Omega)}^4 \right\} dt \\ &\quad + C_2 \int_0^T \left\{ \|n_{\varepsilon} u_{\varepsilon}\|_{L^2(\Omega)}^2 + 1 \right\} dt, \end{aligned} \quad (4.16)$$

where C_1 and C_2 are positive constants independent of ε . Now, due to the Hölder inequality, we have

$$\begin{aligned} & \int_0^T \int_{\Omega} |n_{\varepsilon} u_{\varepsilon}|^2 \\ &\leq \int_0^T \|n_{\varepsilon}\|_{L^3(\Omega)}^2 \|u_{\varepsilon}\|_{L^6(\Omega)}^2 \\ &\leq C_3 \int_0^T \|\nabla u_{\varepsilon}\|_{L^2(\Omega)}^2 \\ &\leq C_4(T+1) \quad \text{for all } T > 0 \end{aligned} \quad (4.17)$$

by using (4.12) and (4.13).

Inserting (4.17) into (4.16) and applying (4.12) and (4.13), we can obtain for some positive constant $C_1(T)$ such that

$$\|n_{\varepsilon t}\|_{L^2(0,T;(W^{1,2}(\Omega))^*)} \leq C_1(T) \quad (4.18)$$

by using (4.13) and Lemma 3.4.

In a similar way, one can derive

$$\|c_{\varepsilon t}\|_{L^2(0,T;(W^{1,2}(\Omega))^*)} \leq C_2(T) \quad (4.19)$$

and

$$\|m_{\varepsilon t}\|_{L^2(0,T;(W^{1,2}(\Omega))^*)} \leq C_2(T) \quad (4.20)$$

with some $C_2(T) > 0$.

Next, for any given $\varphi \in C_{0,\sigma}^\infty(\Omega; \mathbb{R}^3)$, we infer from the fourth equation in (2.1) that

$$\left| \int_{\Omega} \partial_t u_{\varepsilon}(\cdot, t) \varphi \right| = \left| - \int_{\Omega} \nabla u_{\varepsilon} \cdot \nabla \varphi - \kappa \int_{\Omega} (Y_{\varepsilon} u_{\varepsilon} \otimes u_{\varepsilon}) \cdot \nabla \varphi + \int_{\Omega} (n_{\varepsilon} + m_{\varepsilon}) \nabla \phi \cdot \varphi \right| \quad \text{for all } t > 0.$$

Now, by virtue of (4.14), Lemma 3.2 and Lemma 2.3, we thus infer that there exist positive constants C_3, C_4 and C_5 such that

$$\begin{aligned} & \int_0^T \|\partial_t u_{\varepsilon}(\cdot, t)\|_{(W_{0,\sigma}^{1,2}(\Omega))^*}^2 dt \\ & \leq C_3 \left(\int_0^T \int_{\Omega} |\nabla u_{\varepsilon}|^2 + \int_0^T \int_{\Omega} |Y_{\varepsilon} u_{\varepsilon} \otimes u_{\varepsilon}|^2 + \int_0^T \int_{\Omega} n_{\varepsilon}^2 + \int_0^T \int_{\Omega} m_{\varepsilon}^2 \right) \\ & \leq C_4 \left(\int_0^T \int_{\Omega} |\nabla u_{\varepsilon}|^2 + \int_0^T \int_{\Omega} |u_{\varepsilon}|^2 + \int_0^T \int_{\Omega} n_{\varepsilon}^2 + T \right) \\ & \leq C_5(T+1) \quad \text{for all } T > 0. \end{aligned} \quad (4.21)$$

Here we have used the fact that

$$\|Y_{\varepsilon} v\|_{L^2(\Omega)} \leq \|v\|_{L^2(\Omega)} \quad \text{for all } v \in L_{\sigma}^2(\Omega).$$

Combining estimates (4.12)–(4.21), we conclude from Aubin-Lions lemma (see e.g. [28]) that $(u_{\varepsilon})_{\varepsilon \in (0,1)}$ is relatively compact in $L_{loc}^2(\Omega \times [0, \infty); \mathbb{R}^3)$ and $(n_{\varepsilon})_{\varepsilon \in (0,1)}$, $(c_{\varepsilon})_{\varepsilon \in (0,1)}$, $(m_{\varepsilon})_{\varepsilon \in (0,1)}$ are relatively compact in $L_{loc}^2(\Omega \times [0, \infty))$. Therefore, in conjunction with (4.12)–(4.14) and

standard compactness arguments, we can thus find a sequence $(\varepsilon_j)_{j \in \mathbb{N}} \subset (0, 1)$ such that $\varepsilon_j \searrow 0$ as $j \rightarrow \infty$, and such that

$$n_\varepsilon \rightarrow n \text{ in } L_{loc}^2(\bar{\Omega} \times [0, \infty)) \quad \text{and} \quad n_\varepsilon \rightarrow n \text{ a.e. in } \Omega \times (0, \infty), \quad (4.22)$$

$$n_\varepsilon \rightharpoonup n \text{ weak star in } L_{loc}^\infty([0, \infty), L^p(\Omega)) \quad \text{for any } p > 1, \quad (4.23)$$

$$\nabla n_\varepsilon \rightharpoonup \nabla n \text{ in } L_{loc}^2(\bar{\Omega} \times [0, \infty)), \quad (4.24)$$

$$c_\varepsilon \rightarrow c \text{ in } L_{loc}^2(\bar{\Omega} \times [0, \infty)) \quad \text{and} \quad c_\varepsilon \rightarrow c \text{ a.e. in } \Omega \times (0, \infty), \quad (4.25)$$

$$\nabla c_\varepsilon \rightharpoonup \nabla c \text{ in } L_{loc}^4(\bar{\Omega} \times [0, \infty)) \quad (4.26)$$

$$\Delta c_\varepsilon \rightharpoonup \Delta c \text{ in } L_{loc}^2(\bar{\Omega} \times [0, \infty)) \quad (4.27)$$

$$m_\varepsilon \rightarrow m \text{ in } L_{loc}^2(\bar{\Omega} \times [0, \infty)) \quad \text{and} \quad m_\varepsilon \rightarrow m \text{ a.e. in } \Omega \times (0, \infty), \quad (4.28)$$

$$\nabla m_\varepsilon \rightharpoonup \nabla m \text{ in } L_{loc}^4(\bar{\Omega} \times [0, \infty)) \quad (4.29)$$

$$\Delta m_\varepsilon \rightharpoonup \Delta m \text{ in } L_{loc}^2(\bar{\Omega} \times [0, \infty)) \quad (4.30)$$

as well as

$$u_\varepsilon \rightarrow u \text{ in } L_{loc}^2(\bar{\Omega} \times [0, \infty)) \quad \text{and} \quad u_\varepsilon \rightarrow u \text{ a.e. in } \Omega \times (0, \infty) \quad (4.31)$$

and

$$\nabla u_\varepsilon \rightharpoonup \nabla u \text{ in } L_{loc}^2(\bar{\Omega} \times [0, \infty)). \quad (4.32)$$

for some limit function (n, c, m, u) . On the other hand, according to the bounds provided by Lemma 2.3 and Lemmas 3.2–3.3, this readily yields that, for any $\varepsilon \in (0, 1)$,

$$m_\varepsilon - c_\varepsilon - u_\varepsilon \cdot \nabla c_\varepsilon \text{ is bounded in } L^{\frac{5}{3}}(\Omega \times (0, T)), \quad (4.33)$$

where we have used the fact that

$$\begin{aligned} & \int_0^T \int_\Omega \left[|u_\varepsilon \cdot \nabla c_\varepsilon|^{\frac{5}{3}} \right] \\ & \leq C_6 \left[\int_0^T \int_\Omega |u_\varepsilon|^{\frac{10}{3}} \right]^{\frac{1}{2}} \left[\int_0^T \int_\Omega |\nabla c_\varepsilon|^{\frac{10}{3}} \right]^{\frac{1}{2}} \\ & \leq C_7 \left[\int_0^T \int_\Omega |u_\varepsilon|^{\frac{10}{3}} \right]^{\frac{1}{2}} \left[\int_0^T \int_\Omega |\nabla c_\varepsilon|^4 \right]^{\frac{5}{12}} \\ & \leq C_8(T+1) \end{aligned}$$

by using Lemma 3.3. Therefore, in light of (4.33), regularity estimates for the second equation of (2.1) (see e.g. [20]) ensure that $(c_\varepsilon)_{\varepsilon \in (0,1)}$ is bounded in $L^{\frac{5}{3}}((0, T); W^{2, \frac{5}{3}}(\Omega))$. Hence, by virtue of (4.20), we derive from the Aubin–Lions lemma that $(c_\varepsilon)_{\varepsilon \in (0,1)}$ relatively compacts in $L^{\frac{5}{3}}((0, T); W^{1, \frac{5}{3}}(\Omega))$. Thus, we can choose an appropriate subsequence that is still written as $(\varepsilon_j)_{j \in \mathbb{N}}$ such that $\nabla c_{\varepsilon_j} \rightarrow z_1$ in $L^{\frac{5}{3}}(\Omega \times (0, T))$ for all $T \in (0, \infty)$ and some $z_1 \in L^{\frac{5}{3}}(\Omega \times (0, T))$ as $j \rightarrow \infty$. Therefore, by (4.26), we can also derive that $\nabla c_{\varepsilon_j} \rightarrow z_1$ a.e. in $\Omega \times (0, \infty)$ as $j \rightarrow \infty$. In view of (4.26) and the Egorov theorem, we conclude that $z_1 = \nabla c$ and hence

$$\nabla c_\varepsilon \rightarrow \nabla c \quad a.e. \quad \text{in} \quad \Omega \times (0, \infty). \quad (4.34)$$

This combined with (4.22), (4.23) as well as (4.26) and (1.2) implies that

$$n_\varepsilon S_\varepsilon(x, n_\varepsilon, c_\varepsilon) \nabla c_\varepsilon \rightharpoonup n S(x, n, c) \nabla c \text{ in } L^2(\Omega \times (0, T)) \text{ as } \varepsilon = \varepsilon_j \searrow 0 \text{ for each } T \in (0, \infty) \quad (4.35)$$

by using the Egorov theorem. Next we shall prove that (n, c, m, u) is a weak solution of problem (2.1). To this end, testing the first equation in (2.1) by $\varphi \in C_0^\infty(\Omega \times [0, \infty))$, we obtain

$$\begin{aligned} - \int_0^\infty \int_\Omega n_\varepsilon \varphi_t - \int_\Omega n_0 \varphi(\cdot, 0) = & - \int_0^\infty \int_\Omega \nabla n_\varepsilon \cdot \nabla \varphi + \int_0^\infty \int_\Omega n_\varepsilon S_\varepsilon(x, n_\varepsilon, c_\varepsilon) \nabla c_\varepsilon \cdot \nabla \varphi \\ & + \int_0^\infty \int_\Omega n_\varepsilon u_\varepsilon \cdot \nabla \varphi - \int_0^\infty \int_\Omega n_\varepsilon m_\varepsilon \varphi \end{aligned} \quad (4.36)$$

for all $\varepsilon \in (0, 1)$. Then (4.24)–(4.35) and the dominated convergence theorem enables us to conclude

$$\begin{aligned} - \int_0^\infty \int_\Omega n \varphi_t - \int_\Omega n_0 \varphi(\cdot, 0) = & - \int_0^\infty \int_\Omega \nabla n \cdot \nabla \varphi + \int_0^\infty \int_\Omega n S(x, n, c) \nabla c \cdot \nabla \varphi \\ & + \int_0^\infty \int_\Omega n u \cdot \nabla \varphi - \int_0^\infty \int_\Omega n m \varphi \end{aligned} \quad (4.37)$$

by a limit procedure. Next, multiplying the second equation and the third equation in (2.1) by $\varphi \in C_0^\infty(\Omega \times [0, \infty))$, we derive from a limit procedure that

$$\begin{aligned} - \int_0^\infty \int_\Omega c \varphi_t - \int_\Omega c_0 \varphi(\cdot, 0) = & - \int_0^\infty \int_\Omega \nabla c \cdot \nabla \varphi - \int_0^\infty \int_\Omega c \varphi + \int_0^\infty \int_\Omega m \varphi + \int_0^\infty \int_\Omega c u \cdot \nabla \varphi \end{aligned} \quad (4.38)$$

and

$$-\int_0^\infty \int_\Omega m \varphi_t - \int_\Omega m_0 \varphi(\cdot, 0) = -\int_0^\infty \int_\Omega \nabla m \cdot \nabla \varphi - \int_0^\infty \int_\Omega nm \varphi + \int_0^\infty \int_\Omega mu \cdot \nabla \varphi \quad (4.39)$$

in a completed similar manner (see [63] for details). Then testing the fourth equation of (2.1) by $\varphi \in C_0^\infty(\bar{\Omega} \times [0, T]; \mathbb{R}^3)$, we obtain

$$-\int_0^\infty \int_\Omega u \varphi_t - \int_\Omega u_0 \varphi(\cdot, 0) + \kappa \int_0^T \int_\Omega u \otimes u \cdot \nabla \varphi = -\int_0^\infty \int_\Omega \nabla u \cdot \nabla \varphi - \int_0^\infty \int_\Omega (n+m) \nabla \phi \cdot \varphi \quad (4.40)$$

by using Lemma 3.4 and a limit procedure (see [63] for details). This means that (n, c, m, u) is a weak solution of (2.1), in the natural sense as specified in [63].

□

Moreover, if in addition we assume that $\kappa \in \mathbb{R}$, then our solutions will actually be bounded and smooth and hence classical. In fact, by applying the standard parabolic regularity and the classical Schauder estimates for the Stokes evolution, we will show that it is sufficiently regular so as to be a classical solution.

Lemma 4.2. *Let (n, c, m, u) be a weak solution of (1.1). Assume that $\alpha > 0$ and $\kappa = 0$. Then (n, c, m, u) solves (1.1) in the classical sense in $\Omega \times (0, \infty)$. Moreover, this solution is bounded in $\Omega \times (0, \infty)$ in the sense that*

$$\|n(\cdot, t)\|_{L^\infty(\Omega)} + \|c(\cdot, t)\|_{W^{1,\infty}(\Omega)} + \|m(\cdot, t)\|_{W^{1,\infty}(\Omega)} + \|A^\gamma u(\cdot, t)\|_{L^2(\Omega)} \leq C \text{ for all } t > 0. \quad (4.41)$$

Proof. In what follows, let C, C_i denote some different constants, and if no special explanation, they depend at most on $\Omega, \phi, m, n_0, c_0$ and u_0 .

Step 1. The boundedness of $\|A^\gamma u(\cdot, t)\|_{L^2(\Omega)}$ and $\|u(\cdot, t)\|_{L^\infty(\Omega)}$ for all $t \in (0, T_{max, \varepsilon})$

On the basis of the variation-of-constants formula for the projected version of the third equation in (1.1), we derive that

$$u(\cdot, t) = e^{-tA} u_0 + \int_0^t e^{-(t-\tau)A} \mathcal{P}((n(\cdot, \tau) + m(\cdot, \tau)) \nabla \phi) d\tau \text{ for all } t \in (0, T_{max, \varepsilon}).$$

On the other hand, in view of Lemma 3.2 as well as (1.8) and (2.10),

$$\|h(\cdot, t)\|_{L^{4+2\alpha}(\Omega)} \leq C \text{ for all } t \in (0, T_{max, \varepsilon})$$

with $h := \mathcal{P}((n(\cdot, \tau) + m(\cdot, \tau)\nabla\phi)$. Therefore, according to standard smoothing properties of the Stokes semigroup we see that there exist $C_1, C_2 > 0$ and $\lambda_1 > 0$ such that

$$\begin{aligned} \|A^\gamma u(\cdot, t)\|_{L^2(\Omega)} &\leq \|A^\gamma e^{-tA} u_0\|_{L^2(\Omega)} + \int_0^t \|A^\gamma e^{-(t-\tau)A} h(\cdot, \tau) d\tau\|_{L^2(\Omega)} d\tau \\ &\leq \|A^\gamma u_0\|_{L^2(\Omega)} + C_1 \int_0^t (t-\tau)^{-\gamma-\frac{3}{2}(\frac{1}{4+2\alpha}-\frac{1}{2})} e^{-\lambda_1(t-\tau)} \|h(\cdot, \tau)\|_{L^{4+2\alpha}(\Omega)} d\tau \\ &\leq C_2 \quad \text{for all } t \in (0, T_{\max, \varepsilon}) \end{aligned} \tag{4.42}$$

with $\gamma \in (\frac{3}{4}, 1)$, where in the last inequality, we have used the fact that

$$\int_0^t (t-\tau)^{-\gamma-\frac{3}{2}(\frac{1}{4+2\alpha}-\frac{1}{2})} e^{-\lambda_1(t-\tau)} d\tau \leq \int_0^\infty \sigma^{-\gamma-\frac{3}{2}(\frac{1}{4+2\alpha}-\frac{1}{2})} e^{-\lambda_1\sigma} d\sigma < +\infty$$

by using $-\gamma - \frac{3}{2}(\frac{1}{4+2\alpha} - \frac{1}{2}) > -1$. (4.42) implies to

$$\|u(\cdot, t)\|_{L^\infty(\Omega)} \leq \sigma_0 \quad \text{for all } t \in (0, T_{\max, \varepsilon}) \tag{4.43}$$

by using the fact that $D(A^\gamma)$ is continuously embedded into $L^\infty(\Omega)$ (by $\gamma > \frac{3}{4}$).

Step 2. The boundedness of $\|\nabla c(\cdot, t)\|_{L^4(\Omega)}$ for all $t \in (0, T_{\max, \varepsilon})$

Now, multiply the second equation in (1.1) by $-\Delta c$, in view of (3.25), (2.10) and (4.43), we derive from (3.22) and the Young inequality that

$$\|\nabla c(\cdot, t)\|_{L^2(\Omega)} \leq \sigma_1 \quad \text{for all } t \in (0, T_{\max, \varepsilon}). \tag{4.44}$$

Considering the fact that $\nabla c \cdot \nabla \Delta c = \frac{1}{2} \Delta |\nabla c|^2 - |D^2 c|^2$, by a straightforward computation using the second equation in (1.1) and several integrations by parts, we find that

$$\begin{aligned} &\frac{1}{4} \frac{d}{dt} \|\nabla c\|_{L^4(\Omega)}^4 \\ &= \int_\Omega |\nabla c|^2 \nabla c \cdot \nabla (\Delta c - c + m - u \cdot \nabla c) \\ &= \frac{1}{2} \int_\Omega |\nabla c|^2 \Delta |\nabla c|^2 - \int_\Omega |\nabla c|^2 |D^2 c|^2 - \int_\Omega |\nabla c|^4 \\ &\quad + \int_\Omega m \nabla \cdot (|\nabla c|^2 \nabla c) + \int_\Omega (u \cdot \nabla c) \nabla \cdot (|\nabla c|^2 \nabla c) \\ &= -\frac{1}{2} \int_\Omega |\nabla |\nabla c|^2|^2 - \int_\Omega |\nabla c|^4 + \frac{1}{2} \int_{\partial\Omega} |\nabla c|^2 \frac{\partial |\nabla c|^2}{\partial \nu} \\ &\quad - \int_\Omega |\nabla c|^2 |D^2 c|^2 + \int_\Omega m |\nabla c|^2 \Delta c + \int_\Omega m \nabla c \cdot \nabla (|\nabla c|^2) \\ &\quad + \int_\Omega (u \cdot \nabla c) |\nabla c|^2 \Delta c + \int_\Omega (u \cdot \nabla c) \nabla c \cdot \nabla (|\nabla c|^2) \end{aligned} \tag{4.45}$$

for all $t \in (0, T_{max, \varepsilon})$. On the other hand, since Lemma 2.2 of [67], we derive from (2.10) and the Young inequality that

$$\begin{aligned} \|\nabla c\|_{L^4(\Omega)}^4 &\leq \kappa_0 \| |\nabla c| D^2 c \|_{L^2(\Omega)}^{\frac{2}{3}} \|c\|_{L^\infty(\Omega)}^{\frac{8}{3}} + \kappa_0 \|c\|_{L^\infty(\Omega)}^4 \\ &\leq \frac{1}{4(1+16\sigma_0^2)} \int_{\Omega} |\nabla c|^2 |D^2 c|^2 + \kappa_1, \end{aligned} \quad (4.46)$$

where σ_0 is the same as (4.43) and κ_0 and κ_1 are some positive constants. Thanks to the pointwise inequality $|\Delta c| \leq \sqrt{3}|D^2 c|$, along with (1.8) as well as (2.10) and (2.10) this implies that

$$\begin{aligned} &\int_{\Omega} m |\nabla c|^2 \Delta c \\ &\leq \sqrt{3} \int_{\Omega} m |\nabla c|^2 |D^2 c| \\ &\leq \frac{1}{8} \int_{\Omega} |\nabla c|^2 |D^2 c|^2 + 6 \|m\|_{L^\infty(\Omega)}^2 \int_{\Omega} |\nabla c|^2 \\ &\leq \frac{1}{8} \int_{\Omega} |\nabla c|^2 |D^2 c|^2 + 6\lambda^2 \sigma_1^2 \\ &\leq \frac{1}{8} \int_{\Omega} |\nabla c|^2 |D^2 c|^2 + C_3 \quad \text{for all } t \in (0, T_{max, \varepsilon}) \end{aligned} \quad (4.47)$$

and

$$\begin{aligned} &\int_{\Omega} (u \cdot \nabla c) |\nabla c|^2 \Delta c \\ &\leq \sqrt{3} \int_{\Omega} |u \cdot \nabla c| |\nabla c|^2 |D^2 c| \\ &\leq \frac{1}{16} \int_{\Omega} |\nabla c|^2 |D^2 c|^2 + 12 \int_{\Omega} |u \cdot \nabla c|^2 |\nabla c|^2 \\ &\leq \frac{1}{16} \int_{\Omega} |\nabla c|^2 |D^2 c|^2 + 12 \|u\|_{L^\infty(\Omega)}^2 \int_{\Omega} |\nabla c|^4 \quad \text{for all } t \in (0, T_{max, \varepsilon}) \end{aligned} \quad (4.48)$$

by using (3.25) and the Young inequality. Now, inserting (4.46) into (4.48), this shows that

$$\begin{aligned} &\int_{\Omega} (u \cdot \nabla c) |\nabla c|^2 \Delta c \\ &\leq \frac{1}{16} \int_{\Omega} |\nabla c|^2 |D^2 c|^2 + 12 \|u\|_{L^\infty(\Omega)}^2 \times \left[\frac{1}{4(1+16\sigma_0^2)} \int_{\Omega} |\nabla c|^2 |D^2 c|^2 + \kappa_1 \right] \\ &\leq \frac{1}{4} \int_{\Omega} |\nabla c|^2 |D^2 c|^2 + C_4 \quad \text{for all } t \in (0, T_{max, \varepsilon}) \end{aligned} \quad (4.49)$$

by using (4.43). Again, from the Young inequality, (1.8) as well as (2.10) and (3.25), we have

$$\begin{aligned} &\int_{\Omega} m \nabla c \cdot \nabla (|\nabla c|^2) \\ &\leq \frac{1}{8} \int_{\Omega} |\nabla |\nabla c|^2|^2 + 2\lambda^2 \sigma_1^2 \\ &\leq \frac{1}{8} \int_{\Omega} |\nabla |\nabla c|^2|^2 + C_5 \end{aligned} \quad (4.50)$$

and

$$\begin{aligned}
& \int_{\Omega} (u \cdot \nabla c) \nabla c \cdot \nabla (|\nabla c|^2) \\
& \leq \frac{1}{16} \int_{\Omega} |\nabla |\nabla c|^2|^2 + 4 \int_{\Omega} |u \cdot \nabla c|^2 |\nabla c|^2 \\
& \leq \frac{1}{16} \int_{\Omega} |\nabla |\nabla c|^2|^2 + 4 \|u\|_{L^\infty(\Omega)}^2 \times \left[\frac{1}{4(1+16\sigma_0^2)} \int_{\Omega} |\nabla c|^2 |D^2 c|^2 + \kappa_1 \right] \\
& \leq \frac{1}{16} \int_{\Omega} |\nabla |\nabla c|^2|^2 + \frac{1}{16} \int_{\Omega} |\nabla c|^2 |D^2 c|^2 + C_6.
\end{aligned} \tag{4.51}$$

Given the the boundedness of $\|\nabla c\|_{L^2(\Omega)}^2$ (see (4.44)), it is well-known that (cf. [15, 32, 59]) the boundary trace embedding implies that

$$\begin{aligned}
\int_{\partial\Omega} |\nabla c|^2 \frac{\partial}{\partial \nu} |\nabla c|^2 & \leq \frac{1}{16} \int_{\Omega} |\nabla |\nabla c|^2|^2 + C_7 \left(\int_{\Omega} |\nabla c|^2 \right)^2 \\
& \leq \frac{1}{16} \int_{\Omega} |\nabla |\nabla c|^2|^2 + C_8.
\end{aligned} \tag{4.52}$$

Now, together with (4.45), (4.47)–(4.52), we can derive that, for some positive constant C_9 ,

$$\frac{1}{4} \frac{d}{dt} \|\nabla c\|_{L^4(\Omega)}^4 + \frac{3}{4} \int_{\Omega} |\nabla |\nabla c|^2|^2 + \frac{1}{2} \int_{\Omega} |\nabla c|^2 |D^2 c|^2 \leq C_9 \quad \text{for all } t \in (0, T_{max,\varepsilon}), \tag{4.53}$$

which combined with (4.46) yields to

$$\frac{1}{4} \frac{d}{dt} \|\nabla c\|_{L^4(\Omega)}^4 + C_{10} \|\nabla c\|_{L^4(\Omega)}^4 \leq C_{11} \quad \text{for all } t \in (0, T_{max,\varepsilon}). \tag{4.54}$$

This implies

$$\|\nabla c(\cdot, t)\|_{L^4(\Omega)} \leq C_{12} \quad \text{for all } t \in (0, T_{max,\varepsilon}) \tag{4.55}$$

by integration.

Step 3. The boundedness of $\|\nabla m(\cdot, t)\|_{L^4(\Omega)}$ for all $t \in (0, T_{max,\varepsilon})$

An application of the variation of constants formula for c leads to

$$\begin{aligned}
& \|\nabla m(\cdot, t)\|_{L^4(\Omega)} \\
& \leq \|\nabla e^{t(\Delta-1)} m_0\|_{L^4(\Omega)} + \int_0^t \|\nabla e^{(t-s)(\Delta-1)} (m(s) - n(s)m(s))\|_{L^4(\Omega)} ds \\
& \quad + \int_0^t \|\nabla e^{(t-s)(\Delta-1)} \nabla \cdot (u(s)m(s))\|_{L^4(\Omega)} ds.
\end{aligned} \tag{4.56}$$

To estimate the terms on the right of (4.56), in light of (2.10) and (3.25), applying the L^p - L^q estimates associated heat semigroup, for some positive constant λ_1 such that

$$\|\nabla e^{t(\Delta-1)} m_0\|_{L^4(\Omega)} \leq C_{13} \quad \text{for all } t \in (0, T_{max,\varepsilon}) \tag{4.57}$$

as well as

$$\begin{aligned}
& \int_0^t \|\nabla e^{(t-s)(\Delta-1)}(m(s) - n(s)m(s))\|_{L^4(\Omega)} ds \\
& \leq C_{14} \int_0^t [1 + (t-s)^{-\frac{1}{2}-\frac{3}{2}(\frac{1}{4}-\frac{1}{4})}] e^{-\lambda_1(t-s)} (\|n(s)\|_{L^4(\Omega)} + \|m(s)\|_{L^\infty(\Omega)}) ds \\
& \leq C_{15} \quad \text{for all } t \in (0, T_{max,\varepsilon})
\end{aligned} \tag{4.58}$$

and

$$\begin{aligned}
& \int_0^t \|\nabla e^{(t-s)(\Delta-1)} \nabla \cdot (u(s)c(s))\|_{L^4(\Omega)} ds \\
& \leq C_{16} \int_0^t \|(-\Delta + 1)^\iota e^{(t-s)(\Delta-1)} \nabla \cdot (u(s)c(s))\|_{L^4(\Omega)} ds \\
& \leq C_{17} \int_0^t (t-s)^{-\iota-\frac{1}{2}-\tilde{\kappa}} e^{-\lambda_1(t-s)} \|u(s)c(s)\|_{L^4(\Omega)} ds \\
& \leq C_{18} \int_0^t (t-s)^{-\iota-\frac{1}{2}-\tilde{\kappa}} e^{-\lambda_1(t-s)} \|u(s)\|_{L^\infty(\Omega)} \|c(s)\|_{L^\infty(\Omega)} ds \\
& \leq C_{19} \quad \text{for all } t \in (0, T_{max,\varepsilon}),
\end{aligned} \tag{4.59}$$

where $\iota = \frac{13}{28}$, $\tilde{\kappa} = \frac{1}{56}$. Inserting (4.57)–(4.59) into (4.56), one has

$$\|\nabla m(\cdot, t)\|_{L^4(\Omega)} \leq \sigma_2 \quad \text{for all } t \in (0, T_{max,\varepsilon}). \tag{4.60}$$

Step 4. The boundedness of $\|c(\cdot, t)\|_{W^{1,\infty}(\Omega)}$ and $\|m(\cdot, t)\|_{W^{1,\infty}(\Omega)}$ for all $t \in (\tau, T_{max,\varepsilon})$ with $\tau \in (0, T_{max,\varepsilon})$

Choosing $\theta \in (\frac{1}{2} + \frac{3}{8}, 1)$, then the domain of the fractional power $D((-\Delta + 1)^\theta) \hookrightarrow W^{1,\infty}(\Omega)$ (see e.g. [13, 43]). Thus, in light of $\alpha > 0$, using the Hölder inequality and the L^p - L^q estimates associated heat semigroup,

$$\begin{aligned}
& \|c(\cdot, t)\|_{W^{1,\infty}(\Omega)} \\
& \leq C_{20} \|(-\Delta + 1)^\theta c(\cdot, t)\|_{L^4(\Omega)} \\
& \leq C_{21} t^{-\theta} e^{-\lambda_1 t} \|c_0\|_{L^4(\Omega)} + C_{21} \int_0^t (t-s)^{-\theta} e^{-\lambda_1(t-s)} \|(m - u \cdot \nabla c)(s)\|_{L^4(\Omega)} ds \\
& \leq C_{22} + C_{22} \int_0^t (t-s)^{-\theta} e^{-\mu(t-s)} [\|m(s)\|_{L^\infty(\Omega)} + \|c(s)\|_{L^\infty(\Omega)} + \|u(s)\|_{L^\infty(\Omega)} \|\nabla c(s)\|_{L^4(\Omega)}] ds \\
& \leq C_{23} \quad \text{for all } t \in (\tau, T_{max,\varepsilon})
\end{aligned} \tag{4.61}$$

and

$$\begin{aligned}
& \|m(\cdot, t)\|_{W^{1,\infty}(\Omega)} \\
& \leq C_{24} \|(-\Delta + 1)^\theta m(\cdot, t)\|_{L^4(\Omega)} \\
& \leq C_{25} t^{-\theta} e^{-\lambda_1 t} \|m_0\|_{L^4(\Omega)} + C_{25} \int_0^t (t-s)^{-\theta} e^{-\lambda_1(t-s)} \|(m - mn - u \cdot \nabla m)(s)\|_{L^4(\Omega)} ds \\
& \leq C_{26} + C_{26} \int_0^t (t-s)^{-\theta} e^{-\mu(t-s)} [\|m(s)\|_{L^\infty(\Omega)} + \|n(s)\|_{L^4(\Omega)} + \|u(s)\|_{L^\infty(\Omega)} \|\nabla m(s)\|_{L^4(\Omega)}] ds \\
& \leq C_{27} \text{ for all } t \in (\tau, T_{max,\varepsilon})
\end{aligned} \tag{4.62}$$

with $\tau \in (0, T_{max,\varepsilon})$, where we have used (2.10), (4.55), (4.43), (4.60) as well as the Hölder inequality and

$$\int_0^t (t-s)^{-\theta} e^{-\lambda_1(t-s)} \leq \int_0^\infty \sigma^{-\theta} e^{-\lambda_1 \sigma} d\sigma < +\infty.$$

Step 5. The boundedness of $\|c(\cdot, t)\|_{W^{1,\infty}(\Omega)}$ and $\|n(\cdot, t)\|_{L^\infty(\Omega)}$ for all $t \in (0, T_{max,\varepsilon})$

Recalling Lemma 2.1, (4.61) and (4.62), we infer that

$$\|\nabla c(\cdot, t)\|_{L^\infty(\Omega)} \leq \kappa_1 \text{ for all } t \in (0, T_{max,\varepsilon}) \tag{4.63}$$

and

$$\|\nabla m(\cdot, t)\|_{L^\infty(\Omega)} \leq \kappa_2 \text{ for all } t \in (0, T_{max,\varepsilon}). \tag{4.64}$$

Step 6. The boundedness of $\|n(\cdot, t)\|_{L^\infty(\Omega)}$ for all $t \in (\tau, T_{max,\varepsilon})$ with $\tau \in (0, T_{max,\varepsilon})$

Fix $T \in (0, T_{max,\varepsilon})$. Let $M(T) := \sup_{t \in (0, T)} \|n(\cdot, t)\|_{L^\infty(\Omega)}$ and $\tilde{h} := nS(x, n, c)\nabla c + u$.

Then by (3.25), (1.3) and (4.55), there exists $C_{28} > 0$ such that

$$\|\tilde{h}(\cdot, t)\|_{L^4(\Omega)} \leq C_{28} \text{ } t \in (0, T_{max,\varepsilon}), \tag{4.65}$$

where we have used (2.5) and the boundedness of $\|c(\cdot, t)\|_{W^{1,\infty}(\Omega)}$ for all $t \in (\tau, T_{max,\varepsilon})$ with $\tau \in (0, T_{max,\varepsilon})$. Hence, due to the fact that $\nabla \cdot u = 0$, again, by means of an associate variation-of-constants formula for n , we can derive

$$n(t) = e^{(t-t_0)\Delta} n(\cdot, t_0) - \int_{t_0}^t e^{(t-s)\Delta} \nabla \cdot (n(\cdot, s) \tilde{h}(\cdot, s)) ds - \int_{t_0}^t e^{(t-s)\Delta} (n(\cdot, s) m(\cdot, s)) ds, \text{ } t \in (t_0, T), \tag{4.66}$$

where $t_0 := (t - 1)_+$. As the last summand in (4.67) is non-positive by the maximum principle, we can thus estimate

$$\|n(t)\|_{L^\infty(\Omega)} \leq \|e^{(t-t_0)\Delta}n(\cdot, t_0)\|_{L^\infty(\Omega)} + \int_{t_0}^t \|e^{(t-s)\Delta}\nabla \cdot (n(\cdot, s)\tilde{h}(\cdot, s))\|_{L^\infty(\Omega)} ds, \quad t \in (t_0, T). \quad (4.67)$$

If $t \in (0, 1]$, by virtue of the maximum principle, we derive that

$$\|e^{(t-t_0)\Delta}n(\cdot, t_0)\|_{L^\infty(\Omega)} \leq \|n_0\|_{L^\infty(\Omega)}, \quad (4.68)$$

while if $t > 1$ then with the help of the L^p - L^q estimates for the Neumann heat semigroup and (2.10), we conclude that

$$\|e^{(t-t_0)\Delta}n(\cdot, t_0)\|_{L^\infty(\Omega)} \leq C_{29}(t - t_0)^{-\frac{3}{2}}\|n(\cdot, t_0)\|_{L^1(\Omega)} \leq C_{30}. \quad (4.69)$$

Finally, we fix an arbitrary $\frac{7}{2} \in (3, 4)$ and then once more invoke known smoothing properties of the Stokes semigroup and the Hölder inequality to find $C_4 > 0$ such that

$$\begin{aligned} & \int_{t_0}^t \|e^{(t-s)\Delta}\nabla \cdot (n(\cdot, s)\tilde{h}(\cdot, s))\|_{L^\infty(\Omega)} ds \\ & \leq C_{31} \int_{t_0}^t (t-s)^{-\frac{1}{2}-\frac{3}{7}} \|n(\cdot, s)\tilde{h}(\cdot, s)\|_{L^p(\Omega)} ds \\ & \leq C_{32} \int_{t_0}^t (t-s)^{-\frac{1}{2}-\frac{3}{7}} \|n(\cdot, s)\|_{L^{28}(\Omega)} \|\tilde{h}(\cdot, s)\|_{L^4(\Omega)} ds \\ & \leq C_{33} \int_{t_0}^t (t-s)^{-\frac{1}{2}-\frac{3}{7}} \|u(\cdot, s)\|_{L^\infty(\Omega)}^{\frac{27}{28}} \|u(\cdot, s)\|_{L^1(\Omega)}^{\frac{1}{28}} \|\tilde{h}(\cdot, s)\|_{L^4(\Omega)} ds \\ & \leq C_{34} M^b(T) \quad \text{for all } t \in (0, T), \end{aligned} \quad (4.70)$$

In combination with (4.67)–(4.70) and using the definition of $M(T)$ we obtain $C_{35} > 0$ such that

$$M(T) \leq C_{35} + C_{35} M^{\frac{27}{28}}(T) \quad \text{for all } T \in (0, T_{\max, \varepsilon}). \quad (4.71)$$

Hence, with some basic calculation, in light of $T \in (0, T_{\max, \varepsilon})$ was arbitrary, one can get

$$\|n(\cdot, t)\|_{L^\infty(\Omega)} \leq C_{36} \quad \text{for all } t \in (0, T_{\max, \varepsilon}). \quad (4.72)$$

Finally, by virtue of Lemma 2.1 and (4.42), (4.63)–(4.64), (4.72), the local solution can be extend to the global-in-time solutions.

Employing almost exactly the same arguments as in the proof of Lemma 3.1 in [23] (see also [62]), and taking advantage of (4.41), we conclude the regularity theories for the Stokes semigroup and the Hölder estimate for local solutions of parabolic equations, we can obtain weak solution (n, c, m, u) is a classical solution. \square

The most important consequence of Lemmas 4.1–4.2 is the following:

Proof of Theorem 1.1: The theorem 1.1 is part of the statement proven by Lemmas 4.1–4.2.

4.1 Eventual smoothness and asymptotics

Given the preliminary lemma collected in the above, in this subsection, we now establish the claimed asymptotic behavior of the solutions to (2.1) under $\alpha > 0$. Before going further, we list the following lemma, which will be used to derive the convergence properties of solution with respect to the norm in $L^2(\Omega)$.

Lemma 4.3. *(Lemma 4.6 of [10]) Let $\lambda > 0, C > 0$, and suppose that $y \in C^1([0, \infty))$ and $h \in C^0([0, \infty))$ are nonnegative functions satisfying $y'(t) + \lambda y(t) \leq h(t)$ for some $\lambda > 0$ and all $t > 0$. Then if $\int_0^\infty h(s)ds \leq C$, we have $\lim_{t \rightarrow \infty} y(t) = 0$.*

To begin with, let us collect some basic solution properties which essentially have already been used in [10].

Lemma 4.4. *The global solution of (2.1) satisfies*

$$\int_0^\infty \int_\Omega (n_\varepsilon m_\varepsilon + |\nabla m_\varepsilon|^2) < +\infty. \quad (4.73)$$

Proof. These properties are immediate consequences of (2.15) and (2.12). \square

As an immediate consequence, we obtain the following which will firstly serve as a fundament for our proof of stabilization in the first and third solution components.

Lemma 4.5. *Under the assumptions of Lemma 4.1, for any $\eta > 0$, there are $T > 0$ and $\varepsilon_0 > 0$ such that for any $t > T$ and such that for any $\varepsilon \in (0, \varepsilon_0)$*

$$0 \leq \frac{1}{|\Omega|} \int_\Omega m_\varepsilon(\cdot, t) - \hat{m} < \eta \quad \text{for any } t > T \quad (4.74)$$

and

$$0 \leq \frac{1}{|\Omega|} \int_{\Omega} n_{\varepsilon}(\cdot, t) - \hat{n} < \eta \quad \text{for any } t > T, \quad (4.75)$$

where

$$\hat{m} = \left\{ \frac{1}{|\Omega|} \int_{\Omega} m_0 - \frac{1}{|\Omega|} \int_{\Omega} n_0 \right\}_+ \quad (4.76)$$

and

$$\hat{n} = \left\{ \frac{1}{|\Omega|} \int_{\Omega} n_0 - \frac{1}{|\Omega|} \int_{\Omega} m_0 \right\}_+. \quad (4.77)$$

Proof. Pursuing a strategy demonstrated in lemma 4.2 of [48], we start by noting that as a first consequence of Lemma 4.4 we know that

$$\int_{t-1}^t \int_{\Omega} (n_{\varepsilon} m_{\varepsilon} + |\nabla m_{\varepsilon}|^2) \rightarrow 0 \quad \text{as } t \rightarrow \infty. \quad (4.78)$$

Next, in view of (3.25), by using the Hölder inequality and the Poincaré inequality, for some positive constant K ,

$$\begin{aligned} \int_{t-1}^t \int_{\Omega} n_{\varepsilon} m_{\varepsilon} &= \int_{t-1}^t \int_{\Omega} n_{\varepsilon} (m_{\varepsilon} - \bar{m}) + \int_{t-1}^t \bar{m} \int_{\Omega} n_{\varepsilon} \\ &\geq - \int_{t-1}^t \|n_{\varepsilon}\|_{L^2(\Omega)} \|m_{\varepsilon} - \bar{m}\|_{L^2(\Omega)} + \int_{t-1}^t \bar{m} \int_{\Omega} n_{\varepsilon}(x, s) dx ds \\ &\geq -K \int_{t-1}^t \|\nabla m_{\varepsilon}\|_{L^2(\Omega)} + \frac{1}{|\Omega|} \int_{t-1}^t \left[\int_{\Omega} m_{\varepsilon}(x, s) dx \int_{\Omega} n_{\varepsilon}(x, s) dx \right] ds \\ &\geq -K \left(\int_{t-1}^t \|\nabla m_{\varepsilon}\|_{L^2(\Omega)}^2 \right)^{\frac{1}{2}} + \frac{1}{|\Omega|} \int_{t-1}^t \left[\int_{\Omega} m_{\varepsilon}(x, s) dx \int_{\Omega} n_{\varepsilon}(x, s) dx \right] ds. \end{aligned} \quad (4.79)$$

Inserting (4.78) into (4.79), we obtain

$$\int_{t-1}^t \left[\int_{\Omega} m_{\varepsilon}(x, s) dx \int_{\Omega} n_{\varepsilon}(x, s) dx \right] ds \rightarrow 0 \quad \text{as } t \rightarrow \infty. \quad (4.80)$$

Now if $\int_{\Omega} n_0 - \int_{\Omega} m_0 \geq 0$, (2.11) warrants that $\int_{\Omega} n_{\varepsilon} - \int_{\Omega} m_{\varepsilon} \geq 0$, which along with (4.80) implies that

$$\int_{t-1}^t \left(\int_{\Omega} m_{\varepsilon}(x, s) dx \right)^2 ds \rightarrow 0 \quad \text{as } t \rightarrow \infty. \quad (4.81)$$

Noticing that $\int_{\Omega} m_{\varepsilon}(s) \geq \int_{\Omega} m_{\varepsilon}(t)$ for all $t \geq s$, we have

$$0 \leq \left(\int_{\Omega} m_{\varepsilon}(x, t) dx \right)^2 \leq \int_{t-1}^t \left(\int_{\Omega} m_{\varepsilon}(x, s) dx \right)^2 ds \rightarrow 0 \quad \text{as } t \rightarrow \infty,$$

where we invoke (2.11) to obtain

$$\int_{\Omega} n_{\varepsilon}(\cdot, t) \rightarrow \int_{\Omega} n_0 - \int_{\Omega} m_0 \quad \text{as } t \rightarrow \infty.$$

By very similar argument, one can see that $\int_{\Omega} n_{\varepsilon} \rightarrow 0$ and $\int_{\Omega} m_{\varepsilon} \rightarrow \int_{\Omega} m_0 - \int_{\Omega} n_0$ as $t \rightarrow \infty$ in the case of $\int_{\Omega} n_0 - \int_{\Omega} m_0 < 0$. This readily establishes (4.74) and (4.75). \square

Lemma 4.6. *Under the assumptions of Lemma 4.1, for any $\eta > 0$, there are $T > 0$ and $\varepsilon_0 > 0$ such that for any $t > T$ and such that for any $\varepsilon \in (0, \varepsilon_0)$*

$$\int_t^{t+1} \|m_{\varepsilon}(\cdot, t) - \hat{m}\|_{L^{\infty}(\Omega)} < \eta \quad (4.82)$$

and

$$\|m_{\varepsilon}(\cdot, t) - \hat{m}\|_{L^p(\Omega)} < \eta, \quad (4.83)$$

where \hat{m} is give by (4.76).

Proof. Firstly, since Lemma 3.4 asserts the existence of κ_1 such that

$$\int_t^{t+1} \|\nabla m_{\varepsilon}(\cdot, t)\|_{L^4(\Omega)}^4 \leq \kappa_1 \quad (4.84)$$

and since (2.11) implies that

$$\frac{1}{|\Omega|} \int_{\Omega} m_{\varepsilon}(\cdot, t) \geq \left\{ \frac{1}{|\Omega|} \int_{\Omega} m_0 - \frac{1}{|\Omega|} \int_{\Omega} n_0 \right\}_+, \quad (4.85)$$

thus, by (4.74) we infer from the interpolation inequality and the Hölder inequality that

$$\begin{aligned} & \int_t^{t+1} \|m_{\varepsilon} - \hat{m}\|_{L^{\infty}(\Omega)} ds \\ & \leq C \left(\int_t^{t+1} (\|\nabla m_{\varepsilon}\|_{L^4(\Omega)}^{\frac{12}{13}} \|m_{\varepsilon} - \hat{m}\|_{L^1(\Omega)}^{\frac{1}{13}} + \|m_{\varepsilon} - \hat{m}\|_{L^1(\Omega)}) ds \right) \\ & \leq C \int_t^{t+1} \|\nabla m_{\varepsilon}\|_{L^4(\Omega)}^4 ds^{\frac{3}{13}} \left(\int_t^{t+1} \|m_{\varepsilon} - \hat{m}\|_{L^1(\Omega)} ds \right)^{\frac{1}{13}} + C \int_t^{t+1} \|m_{\varepsilon} - \hat{m}\|_{L^1(\Omega)} ds \\ & \leq C \int_t^{t+1} \|\nabla m_{\varepsilon}\|_{L^4(\Omega)}^4 ds^{\frac{3}{13}} \left(\sup_{t>0} \|m_{\varepsilon}(\cdot, t) - \hat{m}\|_{L^1(\Omega)} \right)^{\frac{1}{13}} + C \sup_{t>0} \|m_{\varepsilon}(\cdot, t) - \hat{m}\|_{L^1(\Omega)} \\ & \rightarrow 0 \quad \text{as } t \rightarrow +\infty, \end{aligned} \quad (4.86)$$

which immediately implies (4.82). Here we have used the fact that

$$\|m_{\varepsilon} - \hat{m}\|_{L^1(\Omega)} = \int_{\Omega} [m_{\varepsilon}(\cdot, t) - \hat{m}] = |\Omega| \left[\frac{1}{|\Omega|} \int_{\Omega} m_{\varepsilon}(\cdot, t) - \hat{m} \right] \rightarrow 0 \quad \text{as } t \rightarrow +\infty$$

by using (4.74). Next, for any $p > 1$, in view of Lemma 2.3, we derive from the the interpolation and the Hölder inequality that

$$\begin{aligned} & \|m_\varepsilon - \hat{m}\|_{L^p(\Omega)} \\ & \leq \|m_\varepsilon - \hat{m}\|_{L^\infty(\Omega)}^{\frac{p-1}{p}} \|m_\varepsilon - \hat{m}\|_{L^1(\Omega)}^{\frac{1}{p}} \\ & \rightarrow 0 \quad \text{as } t \rightarrow +\infty, \end{aligned} \tag{4.87}$$

which yields (4.83) directly. \square

Lemma 4.7. *Under the assumptions of Lemma 4.1, for any $\eta > 0$, there are $T > 0$ and $\varepsilon_0 > 0$ such that for any $t > T$ and such that for any $\varepsilon \in (0, \varepsilon_0)$*

$$\|c_\varepsilon(\cdot, t) - \hat{m}\|_{L^2(\Omega)} < \eta \tag{4.88}$$

and

$$\int_t^{t+1} \int_\Omega |\nabla c_\varepsilon|^2 < \eta. \tag{4.89}$$

where \hat{m} is give by (4.76).

Proof. Firstly, by means of the testing procedure, we may derive from the Young inequality that

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \|c_\varepsilon - \hat{m}\|_{L^2(\Omega)}^2 \\ & = \int_\Omega (c_\varepsilon - \hat{m}) [\Delta c_\varepsilon - u_\varepsilon \cdot \nabla c_\varepsilon - (c_\varepsilon - \hat{m}) + (m_\varepsilon - \bar{m})] \\ & = \int_\Omega (c_\varepsilon - \hat{m}) (\Delta c_\varepsilon - u_\varepsilon \cdot \nabla c_\varepsilon) - \int_\Omega (c_\varepsilon - \hat{m})^2 + \int_\Omega (c_\varepsilon - \hat{m}) (m_\varepsilon - \hat{m}) \\ & \leq - \int_\Omega |\nabla c_\varepsilon|^2 - \int_\Omega (c_\varepsilon - \hat{m})^2 + \frac{1}{2} \int_\Omega (m_\varepsilon - \hat{m})^2 \\ & \leq - \int_\Omega (c_\varepsilon - \hat{m})^2 + \frac{1}{2} \int_\Omega (m_\varepsilon - \hat{m})^2 \quad \text{for all } t > 0, \end{aligned} \tag{4.90}$$

where we have used the fact that $\nabla \cdot u_\varepsilon = 0$ and $u_\varepsilon|_{\partial\Omega} = 0$. On the other hand, the bounds from 4.6 entails

$$\int_t^{t+1} \int_\Omega (m_\varepsilon - \hat{m})^2 ds \rightarrow 0 \quad \text{as } t \rightarrow \infty. \tag{4.91}$$

This together with (4.90) and Lemma 4.3 imply (4.88) and (4.89). \square

Lemma 4.8. *Under the assumptions of Lemma 4.1, for any $p > 1$ and $\eta > 0$, there are $T > 0$ and $\varepsilon_0 > 0$ such that for any $t > T$ and such that for any $\varepsilon \in (0, \varepsilon_0)$*

$$\|n_\varepsilon(\cdot, t) - \hat{n}\|_{L^p(\Omega)} < \eta, \tag{4.92}$$

where \hat{n} is given by (4.77).

Proof. Firstly, for any $p > 1$, by Lemma 3.2, there exist positive constants α_1 and $q > p$ such that

$$\int_{\Omega} n_{\varepsilon}^q(x, t) \leq \alpha_1 \quad \text{for all } t > 0. \quad (4.93)$$

By the interpolation and the Hölder inequality, we have

$$\begin{aligned} \|n_{\varepsilon} - \hat{n}\|_{L^p(\Omega)} &\leq \|n_{\varepsilon} - \hat{n}\|_{L^q(\Omega)}^{\frac{q(p-1)}{p(q-1)}} \|n_{\varepsilon} - \hat{n}\|_{L^1(\Omega)}^{\frac{q-p}{p(q-1)}} \\ &\rightarrow 0 \quad \text{as } t \rightarrow +\infty \end{aligned} \quad (4.94)$$

by using (4.75). From (4.94) we readily derive (4.92) and thereby completes the proof. \square

The stabilization property implied by Lemmas 4.6 and 4.8 can now be turned into a preliminary statement on decay of u_{ε} by making use of Lemma 4.3 and the standard testing procedures.

Lemma 4.9. *Under the assumptions of Lemma 4.1, for any $\eta > 0$, there are $T > 0$ and $\varepsilon_0 > 0$ such that for any $t > T$ and such that for any $\varepsilon \in (0, \varepsilon_0)$*

$$\|u_{\varepsilon}(\cdot, t)\|_{L^2(\Omega)} < \eta, \quad (4.95)$$

$$\int_t^{t+1} \|\nabla u_{\varepsilon}\|_{L^2(\Omega)}^2 dx < \eta \quad (4.96)$$

as well as

$$\int_t^{t+1} \|u_{\varepsilon}\|_{L^q(\Omega)}^2 dx < \eta \quad (4.97)$$

and

$$\int_t^{t+1} \|u_{\varepsilon}\|_{L^q(\Omega)} dx < \eta \quad (4.98)$$

for any $q \in [1, 6)$.

Proof. From the fourth equation in (2.1) we obtain the associated Navier-Stokes energy

inequality in the form

$$\begin{aligned}
& \frac{1}{2} \frac{d}{dt} \|u_\varepsilon\|_{L^2(\Omega)}^2 \\
&= - \int_{\Omega} |\nabla u_\varepsilon|^2 + \int_{\Omega} (n_\varepsilon + m_\varepsilon) \nabla \phi \cdot u_\varepsilon - \int_{\Omega} \nabla P_\varepsilon \cdot u_\varepsilon \\
&= - \int_{\Omega} |\nabla u_\varepsilon|^2 + \int_{\Omega} (n_\varepsilon - \hat{n} + m_\varepsilon - \hat{m}) \nabla \phi \cdot u_\varepsilon \\
&\leq - \int_{\Omega} |\nabla u_\varepsilon|^2 + K_1 \left(\int_{\Omega} (n_\varepsilon - \hat{n} + m_\varepsilon - \hat{m})^2 \right)^{\frac{1}{2}} \left(\int_{\Omega} |u_\varepsilon|^2 \right)^{\frac{1}{2}} \\
&\leq - \int_{\Omega} |\nabla u_\varepsilon|^2 + K_1 \left(\int_{\Omega} |n_\varepsilon - \hat{n}|^2 + \int_{\Omega} |m_\varepsilon - \hat{m}|^2 \right)^{\frac{1}{2}} \left(\int_{\Omega} |u_\varepsilon|^2 \right)^{\frac{1}{2}},
\end{aligned} \tag{4.99}$$

where we have used the fact that $\hat{n} = -\hat{m}$ as well as $\nabla \cdot u_\varepsilon = 0$ and $u_\varepsilon|_{\partial\Omega} = 0$. Due to the Poincaré inequality again, we have

$$\eta_0 \int_{\Omega} |u_\varepsilon|^2 \leq \int_{\Omega} |\nabla u_\varepsilon|^2,$$

therefore, collecting (4.92) and (4.82), we derive from (4.99) that

$$\lim_{t \rightarrow +\infty} \int_{\Omega} |u_\varepsilon(x, t)|^2 dx = 0 \tag{4.100}$$

and

$$\int_t^{t+1} \|\nabla u_\varepsilon\|_{L^2(\Omega)}^2 dx \rightarrow 0 \quad \text{as } t \rightarrow \infty \tag{4.101}$$

and thereby proves (4.95)–(4.96). Finally, we make use of the embedding $W^{1,2}(\Omega) \hookrightarrow L^q(\Omega)$ (for any $q \in [1, 6)$) and the Young inequality to find that (4.97) and (4.98) hold. \square

Using the decay property of $m_\varepsilon(\cdot, t) - \hat{m} + n_\varepsilon(\cdot, t) - \hat{n}$ (see Lemmas 4.6 and 4.8), by means of a contraction mapping argument, we may derive a certain eventual regularity and decay of u_ε in $L^p(\Omega)$ with some $p \geq 6$.

Lemma 4.10. *For any $p \in [6, \infty)$ and $\eta > 0$, there are $T > 0$ and $\varepsilon_0 > 0$ such that for any $t > T$ and such that for any $\varepsilon \in (0, \varepsilon_0)$*

$$\|u_\varepsilon(\cdot, s)\|_{L^p(\Omega)} < \eta \quad \text{for any } s \in [t, t+1]. \tag{4.102}$$

Proof. We let $p \geq 6$ and choose $q \in (3, 6)$ which is close to 6 (e.g. $q = \frac{6p}{p+2}$) such that

$$\frac{1}{2} - \frac{3}{2p} - 3\left(\frac{1}{q} - \frac{1}{p}\right) > 0. \tag{4.103}$$

Now, we define $\gamma_0 = \frac{3}{2}(\frac{1}{q} - \frac{1}{p})$. Next, in view of (4.103) and using $p \geq 6$, $q \in (3, 6)$, we have

$$-2\gamma_0 - \frac{1}{2} - \frac{3}{2p} > -1 \quad \text{and} \quad -\frac{1}{2} - \frac{3}{2p} > -1.$$

Therefore,

$$2\kappa_2 \int_0^3 s^{-2\gamma_0 - \frac{1}{2} - \frac{3}{2p}} ds < +\infty \quad \text{and} \quad 2\kappa_2 \int_0^3 s^{-\frac{1}{2} - \frac{3}{2p}} ds < +\infty, \quad (4.104)$$

where κ_2 is give by Lemma 2.2. Thus for any $\eta > 0$, we any choose $\eta_0 \in (0, \eta)$ small enough such that

$$\eta_0 < \min\left\{\frac{1}{2\kappa_2 \int_0^3 s^{-2\gamma_0 - \frac{1}{2} - \frac{3}{2p}} ds}, \frac{1}{2\kappa_2 \int_0^3 s^{-\frac{1}{2} - \frac{3}{2p}} ds}\right\}. \quad (4.105)$$

On the other hand, by (4.98), Lemmas 4.6–4.9, we then pick T_0 and $\varepsilon_0 > 0$ such that for any $t > T$ and such that for any $\varepsilon \in (0, \varepsilon_0)$

$$\int_t^{t+1} \|u_\varepsilon\|_{L^q(\Omega)} dx < \frac{\eta_0}{3\kappa_1} \quad \text{and} \quad \|m_\varepsilon(\cdot, t) - \hat{m} + n_\varepsilon(\cdot, t) - \hat{n}\|_{L^p(\Omega)} < \frac{\eta_0}{3^{\gamma_0+1}\kappa_3\|\nabla\phi\|_{L^\infty(\Omega)}}, \quad (4.106)$$

where κ_1 and κ_3 are given by Lemma 2.2. In view of (4.106), for any $t_1 > T_0$ and $\varepsilon \in (0, \varepsilon_{t_1})$, one can find $\tilde{t}_0 \in (t_1, t_1 + 1)$ such that

$$\|u_\varepsilon(\cdot, \tilde{t}_0)\|_{L^q(\Omega)} dx < \frac{\eta_0}{3\kappa_1}. \quad (4.107)$$

Now, we define

$$T = T_0 + 2.$$

In the following, we will prove that (4.102) holds for any $t > T$. To this end, for the above \tilde{t}_0, p, γ_0 and η_0 , we let

$$X = \{v : \Omega \times (\tilde{t}_0, \tilde{t}_0 + 3) \rightarrow \mathbb{R}; \sup_{s \in (0, 3)} s^{\gamma_0} \|v(\tilde{t}_0 + s)\|_{L^p(\Omega)} \leq \eta_0\}. \quad (4.108)$$

Then we consider the mapping $\varphi : X \rightarrow \mathbb{R}$ defined by

$$\varphi(v) = e^{-tA} u_\varepsilon(\tilde{t}_0) + \int_{\tilde{t}_0}^t e^{-(t-\tau)A} \mathcal{P} [-\kappa \nabla \cdot (Y_\varepsilon v \otimes v)(\tau) + (n_\varepsilon(\tau) + m_\varepsilon(\tau)) \nabla \phi] d\tau.$$

Now, we will show that φ is a contraction on X . In fact, in view of Lemma 2.2, for any

$s > 1$ and for any such v we may derive from the Hölder inequality that

$$\begin{aligned}
& \|\varphi(v)(\cdot, t)\|_{L^p(\Omega)} \\
\leq & \kappa_1(t - \tilde{t}_0)^{-\gamma_0} \|u_\varepsilon(\tilde{t}_0)\|_{L^q(\Omega)} + \kappa_2 \int_{\tilde{t}_0}^t (t - \tau)^{-\frac{1}{2} - \frac{3}{2}(\frac{2}{p} - \frac{1}{p})} \|v \oplus v\|_{L^{\frac{p}{2}}(\Omega)} d\tau \\
& + \kappa_3 \|\nabla \phi\|_{L^\infty(\Omega)} \int_{\tilde{t}_0}^t \|m_\varepsilon(\cdot, \tau) - \hat{m} + n_\varepsilon(\cdot, \tau) - \hat{n}\|_{L^p(\Omega)} d\tau \\
\leq & \kappa_1(t - \tilde{t}_0)^{-\gamma_0} \|u_\varepsilon(\tilde{t}_0)\|_{L^q(\Omega)} + \kappa_2 \int_{\tilde{t}_0}^t (t - \tau)^{-\frac{1}{2} - \frac{3}{2}(\frac{2}{p} - \frac{1}{p})} \|v\|_{L^p(\Omega)}^2 d\tau \\
& + \kappa_3 \|\nabla \phi\|_{L^\infty(\Omega)} \int_{\tilde{t}_0}^{\tilde{t}_0+3} \|m_\varepsilon(\cdot, \tau) - \hat{m} + n_\varepsilon(\cdot, \tau) - \hat{n}\|_{L^p(\Omega)} d\tau.
\end{aligned} \tag{4.109}$$

Therefore, in light of (4.104), (4.107) as well as (4.108) and (4.106), we see that for every $t \in (\tilde{t}_0, \tilde{t}_0 + 3)$ and every $v \in X$

$$\begin{aligned}
& (t - \tilde{t}_0)^{\gamma_0} \|\varphi(v)(\cdot, t)\|_{L^p(\Omega)} \\
\leq & \kappa_1 \|u_\varepsilon(\tilde{t}_0)\|_{L^q(\Omega)} + \kappa_2 (t - \tilde{t}_0)^{\gamma_0} \int_{\tilde{t}_0}^t (t - \tau)^{-\frac{1}{2} - \frac{3}{2}(\frac{2}{p} - \frac{1}{p})} \|v \oplus v\|_{L^{\frac{p}{2}}(\Omega)} d\tau \\
& + \kappa_3 (t - \tilde{t}_0)^{\gamma_0} \|\nabla \phi\|_{L^\infty(\Omega)} \int_{\tilde{t}_0}^{\tilde{t}_0+3} \|m_\varepsilon(\cdot, \tau) - \hat{m} + n_\varepsilon(\cdot, \tau) - \hat{n}\|_{L^p(\Omega)} d\tau \\
\leq & \kappa_1 \|u_\varepsilon(\tilde{t}_0)\|_{L^q(\Omega)} + \kappa_2 \eta_0^2 3^{\gamma_0} \int_0^3 (t - \tau)^{-\frac{1}{2} - \frac{3}{2p}} \tau^{-2\gamma_0} d\tau \\
& + \kappa_3 3^{\gamma_0} \|\nabla \phi\|_{L^\infty(\Omega)} \frac{\eta_0}{3^{\gamma_0+1} \kappa_3 \|\nabla \phi\|_{L^\infty(\Omega)}} \\
< & \eta_0.
\end{aligned} \tag{4.110}$$

from which it readily follows that $\varphi(X) \subset X$. Likewise, for $v \in X$ and $w \in X$ we can use Lemma 2.2 to find that

$$\begin{aligned}
& \|\varphi(v)(\cdot, t) - \varphi(w)(\cdot, t)\|_{L^p(\Omega)} \\
\leq & \kappa_2 \int_{\tilde{t}_0}^t (t - \tau)^{-\frac{1}{2} - \frac{3}{2p}} \|v \oplus v - w \oplus w\|_{L^{\frac{p}{2}}(\Omega)} d\tau \\
= & \kappa_2 \int_{\tilde{t}_0}^t (t - \tau)^{-\frac{1}{2} - \frac{3}{2p}} \|v \oplus (v - w) + (v - w) \oplus w\|_{L^{\frac{p}{2}}(\Omega)} d\tau \\
\leq & \kappa_2 \int_{\tilde{t}_0}^t (t - \tau)^{-\frac{1}{2} - \frac{3}{2p}} (\|v\|_{L^p} + \|w\|_{L^p}) \|v - w\|_{L^p(\Omega)} d\tau \\
\leq & 2\kappa_2 \eta_0 \int_0^3 \tau^{-\frac{1}{2} - \frac{3}{2p}} d\tau \|v - w\|_{L^\infty((0,3); L^p(\Omega))}.
\end{aligned} \tag{4.111}$$

On the other hand, (4.105) implies that

$$2\kappa_2 \eta_0 \int_0^3 \tau^{-\frac{1}{2} - \frac{3}{2p}} d\tau < 1,$$

whence (4.111) shows that φ acts as a contraction on X and hence possesses a unique fixed point on X , which, in view of the definition of φ , must coincide with the unique weak solution u_ε of fourth equation of (2.1) on $(\tilde{t}_0, \tilde{t}_0 + 3)$ (see e.g. Thm. V.2.5.1 of [29]). Now, by (4.110), we also derive that

$$\|u_\varepsilon(\cdot, t)\|_{L^p(\Omega)} dx < \eta_0, \quad \text{for all } t \in (\tilde{t}_0 + 1, \tilde{t}_0 + 3), \quad (4.112)$$

from $(\tilde{t}_0 + 1, \tilde{t}_0 + 3) \supset (t_1 + 2, t_1 + 3)$ we readily derive (4.102). □

In the following lemmas, we next plan to prove Hölder regularity of the components of a solution on intervals of the form $(T_0, T_0 + 1)$ for $T_0 > 0$ by using the maximal Sobolev regularity. To this end, we introduce the following cut-off functions, which will play a key role in deriving higher order regularity for solution of problem (2.1).

Definition 4.1. Let $\xi : \mathbb{R} \rightarrow [0, 1]$ be a smooth, monotone function, satisfying $\xi \equiv 0$ on $(-\infty, 0]$ and $\xi \equiv 1$ on $(1, \infty)$ and for any $t_0 \in \mathbb{R}$ we let $\xi_{t_0} := \xi(t - t_0)$.

Due to the above cut-off function, it follows from maximal Sobolev regularity that the solution $(n_\varepsilon, c_\varepsilon, m_\varepsilon, u_\varepsilon)$ even satisfies estimates in appropriate Hölder spaces:

Lemma 4.11. *Let $\alpha > 0$. Then one can find $\mu \in (0, 1)$ and $T, \varepsilon_0, C > 0$ such that for all $\varepsilon \in (0, \varepsilon_0)$*

$$\|u_\varepsilon(\cdot, t)\|_{C^{1+\mu, \frac{\mu}{2}}(\bar{\Omega} \times [t, t+1])} \leq C \quad \text{for all } t > T, \quad (4.113)$$

$$\|c_\varepsilon(\cdot, t)\|_{C^{1+\mu, \frac{\mu}{2}}(\bar{\Omega} \times [t, t+1])} \leq C \quad \text{for all } t > T \quad (4.114)$$

as well as

$$\|m_\varepsilon(\cdot, t)\|_{C^{1+\mu, \frac{\mu}{2}}(\bar{\Omega} \times [t, t+1])} \leq C \quad \text{for all } t > T \quad (4.115)$$

and

$$\|n_\varepsilon(\cdot, t)\|_{C^{1+\mu, \frac{\mu}{2}}(\bar{\Omega} \times [t, t+1])} \leq C \quad \text{for all } t > T. \quad (4.116)$$

Proof. Firstly, for any $T_0 > 0$ let $\xi := \xi_{T_0}$ and $v := \xi u_\varepsilon$.

$$\begin{cases} v_t - \Delta v = h & \text{in } \Omega \times (T_0, \infty), \\ \nabla \cdot v = 0, & \text{in } \Omega \times (T_0, \infty), \\ v(x, T_0) = 0, & x \in \Omega, \\ v = 0, & \text{on } \partial\Omega \times (T_0, \infty), \end{cases}$$

where

$$h = -\kappa(Y_\varepsilon u_\varepsilon \cdot \nabla)v + \nabla(\xi P_\varepsilon) + \xi(n_\varepsilon + m_\varepsilon)\nabla\phi + \xi' u_\varepsilon.$$

To estimate the inhomogeneity h herein, we first note that the known maximal Sobolev regularity estimate for the Stokes semigroup ([11]) yields a constant $k_1 > 0$ such that

$$\begin{aligned} & \int_{T_0}^{T_0+2} \|v_t\|_{L^s(\Omega)}^s + \int_{T_0}^{T_0+2} \|D^2 v\|_{L^s(\Omega)}^s \\ & \leq k_1 \int_{T_0}^{T_0+2} \|\mathcal{P}(\xi Y_\varepsilon u_\varepsilon \cdot) u_\varepsilon\|_{L^s(\Omega)}^s + k_1 \int_{T_0}^{T_0+2} (\|\mathcal{P}\xi(m_\varepsilon(\cdot, \tau) - \hat{m} + n_\varepsilon(\cdot, \tau) - \hat{n})\nabla\phi\|_{L^s(\Omega)}^s) \\ & \quad + k_1 \int_{T_0}^{T_0+2} \|\mathcal{P}\xi' u_\varepsilon\|_{L^s(\Omega)}^s. \end{aligned} \tag{4.117}$$

From the boundedness of the Helmholtz projection in L^s -spaces and the Hölder inequality we derive from Lemma 4.10 that there exist positive constant k_2 and k_3 such that for any $T_0 > T$

$$\begin{aligned} & k_1 \int_{T_0}^{T_0+2} \|\mathcal{P}(Y_\varepsilon u_\varepsilon \cdot) v\|_{L^s(\Omega)}^s \\ & \leq k_2 \int_{T_0}^{T_0+2} \left(\|Y_\varepsilon u_\varepsilon\|_{L^{l'}(\Omega)}^s \|\nabla v\|_{L^l(\Omega)}^s \right) \\ & \leq k_2 \int_{T_0}^{T_0+2} \left(\|u_\varepsilon\|_{L^{l'}(\Omega)}^s \|\nabla v\|_{L^l(\Omega)}^s \right) \\ & \leq k_3 \int_{T_0}^{T_0+2} \|\nabla v\|_{L^l(\Omega)}^s \quad \text{for any } \varepsilon \in (0, \varepsilon_{T_0}), \end{aligned} \tag{4.118}$$

where $l > 2s$ and $\frac{1}{l} + \frac{1}{l'} = 1$. Thanks to the Gagliardo-Nirenberg inequality, from Lemma

4.10 again, we can estimate the right of (4.118) by following:

$$\begin{aligned}
& k_3 \int_{T_0}^{T_0+2} \|\nabla v\|_{L^l(\Omega)}^s \\
& \leq k_4 \int_{T_0}^{T_0+2} \left(\|D^2 v\|_{L^s(\Omega)}^{as} \|v\|_{L^{r_0}(\Omega)}^{(1-a)s} \right) \\
& \leq k_4 \int_{T_0}^{T_0+2} \left(\|D^2 v\|_{L^s(\Omega)}^{as} \|u_\varepsilon\|_{L^{r_0}(\Omega)}^{(1-a)s} \right) \\
& \leq k_5 \int_{T_0}^{T_0+2} \|D^2 v\|_{L^s(\Omega)}^{as} \quad \text{for any } \varepsilon \in (0, \varepsilon_{T_0}),
\end{aligned} \tag{4.119}$$

where $a \in (0, 1)$ satisfies

$$\frac{1}{l} - \frac{1}{3} = a\left(\frac{1}{s} - \frac{2}{3}\right) + (1-a)\frac{1}{r_0}.$$

Therefore, inserting (4.119) into (4.118) and applying the Young inequality, we find $k_6 > 0$ such that for all $t_0 > T$

$$k_1 \int_{T_0}^{T_0+2} \|\mathcal{P}(Y_\varepsilon u_\varepsilon \cdot) v\|_{L^s(\Omega)}^s \leq \frac{1}{2} \int_{T_0}^{T_0+2} \|D^2 v\|_{L^s(\Omega)}^s + k_6 \quad \text{for any } \varepsilon \in (0, \varepsilon_{T_0}). \tag{4.120}$$

Moreover, we derive from Definition 4.1 and Lemmas 4.6, 4.9 and 4.10, there is $k_7 > 0$ such that

$$\begin{aligned}
& k_1 \int_{T_0}^{T_0+2} \|\mathcal{P}\xi(m_\varepsilon(\cdot, \tau) - \hat{m} + n_\varepsilon(\cdot, \tau) - \hat{n}) \nabla \phi\|_{L^s(\Omega)}^s + k_1 \int_{T_0}^{T_0+2} \|\mathcal{P}\xi' u_\varepsilon\|_{L^s(\Omega)}^s \\
& \leq k_7,
\end{aligned} \tag{4.121}$$

so that invoking (4.120) and (4.117) we can estimate

$$\int_{T_0}^{T_0+2} \|v_t\|_{L^s(\Omega)}^s + \int_{T_0}^{T_0+2} \|D^2 v\|_{L^s(\Omega)}^s \leq k_8. \tag{4.122}$$

Therefore, by the definition of ξ , for any $s > 1$, there exist positive constants C and T such that for any $t > T$ there is $\varepsilon_0 > 0$ satisfying that for any $\varepsilon \in (0, \varepsilon_0)$

$$\|u_\varepsilon(\cdot, t)\|_{L^s((t, t+1); W^{2,s}(\Omega))} + \|u_{\varepsilon t}(\cdot, t)\|_{L^s(\Omega \times (t, t+1))} \leq C \quad \text{for all } t \geq T, \tag{4.123}$$

which in view of a known embedding result ([1]) implies that for all $t_0 > 0$, we can find $\theta_1 \in (0, 1)$ and C_{10} such that

$$\|u_\varepsilon(\cdot, t)\|_{C^{1+\theta_1, \theta_1}(\bar{\Omega} \times [t, t+1])} \leq C_{10} \quad \text{for all } t > T. \tag{4.124}$$

Likewise, again using the maximal Sobolev regularity estimates and the Gagliardo-Nirenberg inequality, we can claim that (4.114)–(4.116) hold by applying Lemmas 2.3, 3.2 and 4.10. \square

Straightforward applications of standard Schauder estimates for the Stokes evolution equation and the heat equation, respectively, finally yield eventual smoothness of the solution $(n_\varepsilon, c_\varepsilon, m_\varepsilon, u_\varepsilon)$.

Lemma 4.12. *Let $\alpha > 0$. Then one can find $\mu \in (0, 1)$ and $T > 0$ such that for some $C > 0$*

$$\|u_\varepsilon(\cdot, t)\|_{C^{2+\mu, 1+\frac{\mu}{2}}(\bar{\Omega} \times [t, t+1]; \mathbb{R}^3)} \leq C \quad \text{for all } t > T, \quad (4.125)$$

$$\|c_\varepsilon(\cdot, t)\|_{C^{2+\mu, 1+\frac{\mu}{2}}(\bar{\Omega} \times [t, t+1])} \leq C \quad \text{for all } t > T \quad (4.126)$$

as well as

$$\|m_\varepsilon(\cdot, t)\|_{C^{2+\mu, 1+\frac{\mu}{2}}(\bar{\Omega} \times [t, t+1])} \leq C \quad \text{for all } t > T \quad (4.127)$$

and

$$\|n_\varepsilon(\cdot, t)\|_{C^{2+\mu, 1+\frac{\mu}{2}}(\bar{\Omega} \times [t, t+1])} \leq C \quad \text{for all } t > T. \quad (4.128)$$

Proof. We first combine Lemma 4.11 to infer the existence of $\alpha_1 \in (0, 1)$, $T_1 > 0$ and $C_1 > 0$ such that for all $t > T_1$,

$$\begin{aligned} & \|u_\varepsilon \cdot \nabla c_\varepsilon\|_{C^{\alpha_1, \frac{\alpha_1}{2}}(\bar{\Omega} \times [t, t+1])} + \|u_\varepsilon \cdot \nabla m_\varepsilon\|_{C^{\alpha_1, \frac{\alpha_1}{2}}(\bar{\Omega} \times [t, t+1])} \\ & + \|n_\varepsilon m_\varepsilon\|_{C^{\alpha_1, \frac{\alpha_1}{2}}(\bar{\Omega} \times [t, t+1])} + \|m_\varepsilon\|_{C^{\alpha_1, \frac{\alpha_1}{2}}(\bar{\Omega} \times [t, t+1])} \\ & \leq C_1. \end{aligned} \quad (4.129)$$

Standard parabolic Schauder estimates applied to the second and third equation in (2.1) ([20]) thus provide $C_2 > 0$ fulfilling

$$\|c_\varepsilon\|_{C^{2+\alpha_1, 1+\frac{\alpha_1}{2}}(\bar{\Omega} \times [t, t+1])} + \|m_\varepsilon\|_{C^{2+\alpha_1, 1+\frac{\alpha_1}{2}}(\bar{\Omega} \times [t, t+1])} \leq C_2 \quad \text{for all } t > T_1 + 1. \quad (4.130)$$

According to Lemma 4.11, it is possible to fix $\alpha_2 \in (0, 1)$, $T_2 > 0$ and $C_3 > 0$ such that

$$\|n_\varepsilon\|_{C^{1+\alpha_2, \frac{\alpha_2}{2}}(\bar{\Omega} \times [t, t+1])} + \|u_\varepsilon\|_{C^{1+\alpha_2, \frac{\alpha_2}{2}}(\bar{\Omega} \times [t, t+1]; \mathbb{R}^3)} \leq C_3 \quad \text{for all } t > T_2. \quad (4.131)$$

We next set $T := T_2 + 1$ and let $t_0 > T$ be given. Then with ξ_{t_0} taken from Definition 4.1, we again use that $v(\cdot, t) := \xi_{t_0} u_\varepsilon(\cdot, t)$, $(x \in \Omega, t > t_0 - 1)$, is a solution of

$$\begin{cases} v_t - \Delta v = h & x \in \Omega, t > t_0 - 1, \\ v(x, t_0 - 1) = 0, & x \in \Omega, \end{cases} \quad (4.132)$$

where

$$h_\varepsilon = -\kappa(Y_\varepsilon u_\varepsilon \cdot \nabla)v + \nabla(\xi P_\varepsilon) + \xi(n_\varepsilon + m_\varepsilon)\nabla\phi + \xi' u_\varepsilon.$$

Now from (4.131) and the smoothness of ξ we readily obtain $\alpha_3 \in (0, 1)$ and $C_4 > 0$ fulfilling

$$\|h_\varepsilon\|_{C^{\alpha_3, \frac{\alpha_3}{2}}(\bar{\Omega} \times [t_0-1, t_0+1]; \mathbb{R}^3)} \leq C_4, \quad (4.133)$$

so that regularity estimates from Schauder theory for the Stokes evolution equation ([30]) ensure that (4.132) possesses a classical solution $\bar{v} \in C^{2+\alpha_3, 1+\frac{\alpha_3}{2}}(\bar{\Omega} \times [t_0-1, t_0+1])$ satisfying

$$\|\bar{v}\|_{C^{2+\alpha_3, 1+\frac{\alpha_3}{2}}(\bar{\Omega} \times [t_0-1, t_0+1]; \mathbb{R}^3)} \leq C_5 \quad (4.134)$$

with some $C_5 > 0$ which is independent of t_0 . This combined with the uniqueness property of (4.132), one can prove

$$\|u_\varepsilon\|_{C^{2+\alpha_3, 1+\frac{\alpha_3}{2}}(\bar{\Omega} \times [t, t+1]; \mathbb{R}^3)} \leq C_6. \quad (4.135)$$

Again relying on Lemma 4.11, this in turn warrants that for some $\alpha_4 \in (0, 1)$, $T_4 > 0$ and $C_7 > 0$ such that for all $t > T_4$

$$\|\nabla \cdot (n_\varepsilon S_\varepsilon(x, n_\varepsilon, c_\varepsilon) \nabla c_\varepsilon)\|_{C^{\alpha_4, \frac{\alpha_4}{2}}(\bar{\Omega} \times [t, t+1])} + \|u_\varepsilon \cdot \nabla n_\varepsilon\|_{C^{\alpha_4, \frac{\alpha_4}{2}}(\bar{\Omega} \times [t, t+1])} + \|n_\varepsilon m_\varepsilon\|_{C^{\alpha_4, \frac{\alpha_4}{2}}(\bar{\Omega} \times [t, t+1])} \leq C_7, \quad (4.136)$$

which along with the Schauder theory says establishes

$$\|n_\varepsilon\|_{C^{2+\alpha_4, 1+\frac{\alpha_4}{2}}(\bar{\Omega} \times [t, t+1])} \leq C_8. \quad (4.137)$$

Finally, choose $T = \max\{T_1, T_1, T_2, T_3, T_4\}$ and $\mu = \min\{\alpha_1, \alpha_2, \alpha_3, \alpha_4\}$, then (4.130), (4.135), (4.137) imply (4.125)–(4.128). \square

Having found uniform Hölder bounds on $n_\varepsilon, c_\varepsilon, m_\varepsilon$ and u_ε for $\varepsilon > 0$ in the previous three lemmas (see Lemmas 4.11 and 4.12), also n, c, m and u share this regularity and these bounds.

Lemma 4.13. *Assume that $\alpha > 0$. There exist $\theta \in (0, 1)$ as well as $T_0 > 0$, $(\varepsilon_j)_{j \in \mathbb{N}} \subset (0, 1)$*

of the sequence from Lemma 4.1 such that for any $t > T_0$

$$\begin{cases} n \in C^{2+\theta, 1+\frac{\theta}{2}}(\bar{\Omega} \times [t, t+1]), \\ c \in C^{2+\theta, 1+\frac{\theta}{2}}(\bar{\Omega} \times [t, t+1]), \\ m \in C^{2+\theta, 1+\frac{\theta}{2}}(\bar{\Omega} \times [t, t+1]), \\ u \in C^{2+\theta, 1+\frac{\theta}{2}}(\bar{\Omega} \times [t, t+1]; \mathbb{R}^3), \end{cases} \quad (4.138)$$

that $\varepsilon_j \searrow 0$ as $j \rightarrow \infty$ and

$$\begin{cases} n_\varepsilon \rightarrow n & \in C^{1+\theta, \frac{\theta}{2}}(\bar{\Omega} \times [t, t+1]), \\ c_\varepsilon \rightarrow c & \in C^{1+\theta, \frac{\theta}{2}}(\bar{\Omega} \times [t, t+1]), \\ m_\varepsilon \rightarrow m & \in C^{1+\theta, \frac{\theta}{2}}(\bar{\Omega} \times [t, t+1]), \\ u_\varepsilon \rightarrow u & \in C^{1+\theta, \frac{\theta}{2}}(\bar{\Omega} \times [t, t+1]; \mathbb{R}^3) \end{cases} \quad \text{and} \quad (4.139)$$

as $\varepsilon = \varepsilon_j \searrow 0$. Moreover, there is $C > 0$ such that

$$\|c(\cdot, t)\|_{C^{2+\theta, 1+\frac{\theta}{2}}(\bar{\Omega} \times [t, t+1])} + \|m(\cdot, t)\|_{C^{2+\theta, 1+\frac{\theta}{2}}(\bar{\Omega} \times [t, t+1])} \leq C \quad \text{for all } t > T_0 \quad (4.140)$$

as well as

$$\|n(\cdot, t)\|_{C^{2+\theta, 1+\frac{\theta}{2}}(\bar{\Omega} \times [t, t+1])} + \|u(\cdot, t)\|_{C^{2+\theta, 1+\frac{\theta}{2}}(\bar{\Omega} \times [t, t+1]; \mathbb{R}^3)} \leq C \quad \text{for all } t > T_0. \quad (4.141)$$

Proof. In conjunction with Lemmas 4.12 and 4.1 and the standard compactness arguments (see [28]), we can thus find a sequence $(\varepsilon_j)_{j \in \mathbb{N}} \subset (0, 1)$ such that $\varepsilon_j \searrow 0$ as $j \rightarrow \infty$, and such that (4.138)–(4.141) hold. The proof of Lemma 4.13 is completed. \square

Lemma 4.14. *Let $\alpha > 0$. Then one can find $\theta \in (0, 1)$ and $T > 0$ such that*

$$\|c(\cdot, t)\|_{C^{2+\theta, 1+\frac{\theta}{2}}(\bar{\Omega} \times [T, \infty))} \leq C \quad (4.142)$$

as well as

$$\|u(\cdot, t)\|_{C^{2+\theta, 1+\frac{\theta}{2}}(\bar{\Omega} \times [T, \infty); \mathbb{R}^3)} \leq C \quad (4.143)$$

and

$$\|n(\cdot, t)\|_{C^{2+\theta, 1+\frac{\theta}{2}}(\bar{\Omega} \times [T, \infty))} + \|m(\cdot, t)\|_{C^{2+\theta, 1+\frac{\theta}{2}}(\bar{\Omega} \times [T, \infty))} \leq C. \quad (4.144)$$

Proof. Let $g := -\xi c + m\xi - \xi u \cdot \nabla c + c\xi'$, where $\xi := \xi_{T_0}$ is given by Definition 4.1 and T_0 is same as the previous lemmas. Then we consider the following problem

$$\begin{cases} \tilde{c}_t - \Delta \tilde{c} = g & x \in \Omega, t > T_0, \\ \tilde{c}(T_0) = 0, & x \in \Omega, \\ \frac{\partial \tilde{c}}{\partial \nu} = 0, & x \in \partial\Omega, \end{cases} \quad (4.145)$$

In view of Lemma 4.13 and Definition 4.1, we drive that

$$g \text{ is bounded in } C^\theta(\bar{\Omega} \times (T, \infty)),$$

so that, regularity estimates from Schauder theory for the parabolic equation (see e.g. III.5.1 of [20]) ensure that problem (4.145) admits a unique solution $\tilde{c} \in C^{2+\theta, 1+\frac{\theta}{2}}(\bar{\Omega} \times [T_0 + 1, \infty))$. This combined with the property of ξ implies that

$$c \in C^{2+\theta, 1+\frac{\theta}{2}}(\bar{\Omega} \times [T_0 + 1, \infty)). \quad (4.146)$$

Applying the same argument one can derive the third equation of (1.1) that

$$m \in C^{2+\theta, 1+\frac{\theta}{2}}(\bar{\Omega} \times [T_0 + 1, \infty)). \quad (4.147)$$

Finally, employing almost exactly the same arguments as in the proof of Lemma 4.12 (the minor necessary changes are left as an easy exercise to the reader), and taking advantage of (4.141), we conclude that

$$u \in C^{2+\theta, 1+\frac{\theta}{2}}(\bar{\Omega} \times [T_0 + 1, \infty); \mathbb{R}^3) \quad (4.148)$$

and

$$n \in C^{2+\theta, 1+\frac{\theta}{2}}(\bar{\Omega} \times [T_0 + 1, \infty)), \quad (4.149)$$

whence combining the result of (4.146) with (4.147) completes the proof. \square

On the basis of the eventual uniform continuity properties implied by the estimates in this section (see Lemma 4.14), by using the interpolation inequality, we can now turn the weak stabilization properties of n, c, m and u from Lemmas 4.6–4.9 into convergence with regard to the norm in $L^\infty(\Omega)$.

Lemma 4.15. *Let $\alpha > 0$. The solution (n, c, m, u) of (2.1) constructed in Lemma 4.1 satisfies*

$$n(\cdot, t) \rightarrow \hat{n}, m(\cdot, t) \rightarrow \hat{m} \text{ as well as } c(\cdot, t) \rightarrow \hat{m} \text{ and } u(\cdot, t) \rightarrow 0 \text{ in } L^\infty(\Omega),$$

where $\hat{n} = \frac{1}{|\Omega|} \{ \int_\Omega n_0 - \int_\Omega m_0 \}_+$ and $\hat{m} = \frac{1}{|\Omega|} \{ \int_\Omega m_0 - \int_\Omega n_0 \}_+$.

Proof. Firstly, due to Lemmas 4.6–4.9, we derive from Lemma 4.14 that

$$n(t) \rightarrow \hat{n}, c(t) \rightarrow \hat{m}, m(t) \rightarrow \hat{m} \text{ and } u(t) \rightarrow 0 \text{ in } L^2(\Omega), \quad (4.150)$$

where \hat{m} and \hat{n} are given by (4.76) and (4.77), respectively. Next, due to Lemma 4.13, one can obtain there exist positive constants κ_1 and T such that for all $t > T$

$$\|n(\cdot, t)\|_{C^{2+\theta}(\bar{\Omega})} + \|c(\cdot, t)\|_{C^{2+\theta}(\bar{\Omega})} + \|m(\cdot, t)\|_{C^{2+\theta}(\bar{\Omega})} + \|u(\cdot, t)\|_{C^{2+\theta}(\bar{\Omega})} \leq \kappa_1. \quad (4.151)$$

Therefore, for any $\eta > 0$, we may use the compactness of the first of the embeddings $C^{2+\theta}(\bar{\Omega}) \hookrightarrow L^\infty(\Omega) \hookrightarrow L^2(\Omega)$ to fix, through an associated Ehrling lemma, a constant $\kappa_2 > 0$ such that

$$\|n(\cdot, t) - \hat{n}\|_{L^\infty(\Omega)} \leq \frac{\eta}{2\kappa_1} \|n(\cdot, t) - \hat{n}\|_{C^{2+\theta}(\bar{\Omega})} + \kappa_2 \|n(\cdot, t) - \hat{n}\|_{L^2(\Omega)} \quad (4.152)$$

$$\|c(\cdot, t) - \hat{m}\|_{L^\infty(\Omega)} \leq \frac{\eta}{2\kappa_1} \|c(\cdot, t) - \hat{m}\|_{C^{2+\theta}(\bar{\Omega})} + \kappa_2 \|c(\cdot, t) - \hat{m}\|_{L^2(\Omega)} \quad (4.153)$$

as well as

$$\|m(\cdot, t) - \hat{m}\|_{L^\infty(\Omega)} \leq \frac{\eta}{2\kappa_1} \|m(\cdot, t) - \hat{m}\|_{C^{2+\theta}(\bar{\Omega})} + \kappa_2 \|m(\cdot, t) - \hat{m}\|_{L^2(\Omega)} \quad (4.154)$$

and

$$\|u(\cdot, t)\|_{L^\infty(\Omega)} \leq \frac{\eta}{2\kappa_1} \|u(\cdot, t)\|_{C^{2+\theta}(\bar{\Omega})} + \kappa_2 \|u(\cdot, t)\|_{L^2(\Omega)}. \quad (4.155)$$

Now due to (4.150), we may choose $t_0 > \max\{1, T\}$ large enough such that for all $t > t_0$,

$$\|n(\cdot, t) - \hat{n}\|_{L^2(\Omega)} + \|c(\cdot, t) - \hat{m}\|_{L^2(\Omega)} + \|m(\cdot, t) - \hat{m}\|_{L^2(\Omega)} + \|u(\cdot, t)\|_{L^2(\Omega)} < \frac{\eta}{2\kappa_2}. \quad (4.156)$$

Combined with (4.152)–(4.156), this shows that in fact

$$\begin{aligned} \|n(\cdot, t) - \hat{n}\|_{L^\infty(\Omega)} &\leq \frac{\eta}{2\kappa_1} \|n(\cdot, t) - \hat{n}\|_{C^{2+\theta}(\bar{\Omega})} + \kappa_2 \|n(\cdot, t) - \hat{n}\|_{L^2(\Omega)} \\ &< \frac{\eta}{2\kappa_1} \kappa_1 + \kappa_2 \frac{\eta}{2\kappa_2} \\ &= \eta \text{ for all } t > t_0, \end{aligned}$$

$$\begin{aligned}
\|c(\cdot, t) - \hat{m}\|_{L^\infty(\Omega)} &\leq \frac{\eta}{2\kappa_1} \|c(\cdot, t) - \hat{m}\|_{C^{2+\theta}(\bar{\Omega})} + \kappa_2 \|c(\cdot, t) - \hat{m}\|_{L^2(\Omega)} \\
&< \frac{\eta}{2\kappa_1} \kappa_1 + \kappa_2 \frac{\eta}{2\kappa_2} \\
&= \eta \quad \text{for all } t > t_0
\end{aligned}$$

as well as

$$\begin{aligned}
\|m(\cdot, t) - \hat{m}\|_{L^\infty(\Omega)} &\leq \frac{\eta}{2\kappa_1} \|m(\cdot, t) - \hat{m}\|_{C^{2+\theta}(\bar{\Omega})} + \kappa_2 \|m(\cdot, t) - \hat{m}\|_{L^2(\Omega)} \\
&< \frac{\eta}{2\kappa_1} \kappa_1 + \kappa_2 \frac{\eta}{2\kappa_2} \\
&= \eta \quad \text{for all } t > t_0
\end{aligned}$$

and

$$\begin{aligned}
\|u(\cdot, t)\|_{L^\infty(\Omega)} &\leq \frac{\eta}{2\kappa_1} \|u(\cdot, t)\|_{C^{2+\theta}(\bar{\Omega})} + \kappa_2 \|u(\cdot, t)\|_{L^2(\Omega)} \\
&< \frac{\eta}{2\kappa_1} \kappa_1 + \kappa_2 \frac{\eta}{2\kappa_2} \\
&= \eta \quad \text{for all } t > t_0,
\end{aligned}$$

which together with the fact that $\eta > 0$ was arbitrary implies the claimed estimates. □

In order to prove Theorem 1.2, we now only have to collect the results prepared during this section:

Proof of Theorem 1.2.

Proof. Combining Lemmas 4.13–4.15 this convergence statement results immediately. □

Acknowledgement: This work is partially supported by the National Natural Science Foundation of China (No. 11601215), Shandong Provincial Science Foundation for Outstanding Youth (No. ZR2018JL005) and Project funded by China Postdoctoral Science Foundation (No. 2019M650927, 2019T120168).

References

- [1] H. Amann, *Compact embeddings of vector-valued Sobolev and Besov spaces*, Glasnik Mat., 35(55)(2000), 161–177.

- [2] N. Bellomo, A. Belloquid, Y. Tao, M. Winkler, *Toward a mathematical theory of Keller–Segel models of pattern formation in biological tissues*, Math. Models Methods Appl. Sci., 25(9)(2015), 1663–1763.
- [3] X. Cao, J. Lankeit, *Global classical small-Data solutions for a three-dimensional Chemotaxis Navier-Stokes System involving Matrix-Valued Sensitivities*, Cal. Var. Partial Differential Equations, 55(4)(2016), 107-123.
- [4] M. Chae, K. Kang, J. Lee, *Global Existence and temporal decay in Keller–Segel models coupled to fluid equations*, Comm. Part. Diff. Eqns., 39(2014), 1205–1235.
- [5] T. Cieřlak, C. Stinner, *Finite-time blowup and global-in-time unbounded solutions to a parabolic–parabolic quasilinear Keller–Segel system in higher dimensions*, J. Diff. Eqns., 252(2012), 5832–5851.
- [6] T. Cieřlak, C. Stinner, *New critical exponents in a fully parabolic quasilinear Keller–Segel system and applications to volume filling models*, J. Diff. Eqns., 258(2015), 2080–2113.
- [7] T. Cieřlak, M. Winkler, *Finite-time blow-up in a quasilinear system of chemotaxis*, Nonlinearity, 21(2008), 1057–1076.
- [8] R. Duan, A. Lorz, P. A. Markowich, *Global solutions to the coupled chemotaxis-fluid equations*, Comm. Part. Diff. Eqns., 35 (2010), 1635–1673.
- [9] E. E. Espejo, T. Suzuki, *Reaction enhancement by chemotaxis*, Nonlinear Anal. Real World Appl., 35(2017), 102–131.
- [10] E. E. Espejo, M. Winkler, *Global classical solvability and stabilization in a two-dimensional chemotaxis-Navier-Stokes system modeling coral fertilization*, Nonlinearity, 31(2018), 1227-1259.
- [11] Y. Giga, H. Sohr, *Abstract L^p estimate for the Cauchy problem with applications to the Navier-Stokes equations in exterior domains*, J. Funct. Anal., 102(1991), 72–94.

- [12] T. Hillen, K. Painter, *A user's guide to PDE models for chemotaxis*, J. Math. Biol., 58(2009), 183–217.
- [13] D. Horstmann, M. Winkler, *Boundedness vs. blow-up in a chemotaxis system*, J. Diff. Eqns, 215(2005), 52–107.
- [14] S. Ishida, *Global existence and boundedness for chemotaxis–Navier–Stokes system with position-dependent sensitivity in 2d bounded domains*, Discrete Contin. Dyn. Syst. Ser. A, 32(2015), 3463–3482.
- [15] S. Ishida, K. Seki and T. Yokota, *Boundedness in quasilinear Keller–Segel systems of parabolic–parabolic type on non-convex bounded domains*, J. Diff. Eqns., 256 (2014), 2993–3010.
- [16] Y. Ke, J. Zheng, *An optimal result for global existence in a three-dimensional Keller–Segel–Navier–Stokes system involving tensor-valued sensitivity with saturation*, Calculus of Variations and Partial Diff. Eqns., <https://doi.org/10.1007/s00526-019-1568-2>.
- [17] E. Keller, L. Segel, *Model for chemotaxis*, J. Theor. Biol., 30(1970), 225–234.
- [18] A. Kiselev, L. Ryzhik, *Biomixing by chemotaxis and efficiency of biological reactions: the critical reaction case*, J. Math. Phys., 53(11):115609, 9, 2012.
- [19] A. Kiselev and L. Ryzhik, *Biomixing by chemotaxis and enhancement of biological reactions*, Comm. Partial Diff. Eqns., 37(2012), 298–318.
- [20] O. A. Ladyzenskaja, V. A. Solonnikov and N. N. Ural'ceva, *Linear and quasi-Linear Equations of Parabolic Type*, Amer. Math. Soc. Transl., Providence, RI, 1968.
- [21] J. Leray, *Sur le mouvement d'un liquide visqueux emplissant l'espace*, Acta Math., 63(1934), 193–248.
- [22] J. Li, P. Pang, Y. Wang, *Global boundedness and decay property of a three-dimensional Keller–Segel–Stokes system modeling coral fertilization*, Nonlinearity, In press.

- [23] T. Li, A. Suen, M. Winkler and C. Xue, *Global small-data solutions in a chemotaxis system with rotation*, Math. Mod. Meth. Appl. Sci., 25(2015), 721–747.
- [24] J. Liu, Y. Wang, *Boundedness and decay property in a three-dimensional Keller–Segel–Stokes system involving tensor-valued sensitivity with saturation*, J. Diff. Eqns., 261(2)(2016), 967–999.
- [25] J.-G. Liu, A. Lorz, *A coupled chemotaxis–fluid model: global existence*, Ann. Inst. H. Poincaré Anal. Non Linéaire, 28(5)(2011), 643–652.
- [26] A. Lorz, *Coupled chemotaxis fluid equations*, Math. Models Methods Appl. Sci., 20(2010), 987–1004.
- [27] K. Painter, T. Hillen, *Volume-filling and quorum-sensing in models for chemosensitive movement*, Can. Appl. Math. Q. 10(2002), 501–543.
- [28] J. Simon, *Compact sets in the space $L^p(O, T; B)$* , Annali di Matematica Pura ed Applicata, 146(1)(1986), 65–96.
- [29] H. Sohr, *The Navier–Stokes equations, An elementary functional analytic approach*, Birkhäuser Verlag, Basel (2001).
- [30] V. A. Solonnikov, *Schauder estimates for the evolutionary generalized Stokes problem, Non-linear equations and spectral theory*, Amer. Math. Soc. Transl. Ser. 2, vol. 220, Amer. Math. Soc., Providence, RI, 2007, pp. 165–200.
- [31] Y. Tao, M. Winkler, *Boundedness in a quasilinear parabolic–parabolic Keller–Segel system with subcritical sensitivity*, J. Diff. Eqns., 252(2012), 692–715.
- [32] Y. Tao, M. Winkler, *Boundedness and decay enforced by quadratic degradation in a three-dimensional chemotaxis–fluid system*, Z. Angew. Math. Phys., 66(2015), 2555–2573.
- [33] I. Tuval, L. Cisneros, C. Dombrowski, et al., *Bacterial swimming and oxygen transport near contact lines*, Proc. Natl. Acad. Sci. USA, 102(2005), 2277–2282.

- [34] L. Wang, C. Mu, P. Zheng, *On a quasilinear parabolic–elliptic chemotaxis system with logistic source*, J. Diff. Eqns., 256(2014), 1847–1872.
- [35] L. Wang, Y. Li, C. Mu, *Boundedness in a parabolic–parabolic quasilinear chemotaxis system with logistic source*, Discrete Contin. Dyn. Syst. Ser. A., 34(2014), 789–802.
- [36] Y. Wang, X. Cao, *Global classical solutions of a 3d chemotaxis–Stokes system with rotation*, Discrete Contin. Dyn. Syst. Ser. B, 20(2015), 3235–3254.
- [37] Y. Wang, *Global weak solutions in a three-dimensional Keller–Segel–Navier–Stokes system with subcritical sensitivity*, Math. Models Methods Appl. Sci., (27)(14)(2017), 2745–2780.
- [38] Y. Wang, M. Winkler, Z. Xiang, *Global classical solutions in a two-dimensional chemotaxis–Navier–Stokes system with subcritical sensitivity*, Annali della Scuola Normale Superiore di Pisa–Classe di Scienze. XVIII, (2018), 2036–2145.
- [39] Y. Wang, Z. Xiang, *Global existence and boundedness in a Keller–Segel–Stokes system involving a tensor-valued sensitivity with saturation*, J. Diff. Eqns., 259(2015), 7578–7609.
- [40] Y. Wang, Z. Xiang, *Global existence and boundedness in a Keller–Segel–Stokes system involving a tensor-valued sensitivity with saturation: the 3D case*, J. Differ. Eqns. 261(2016), 4944–4973.
- [41] M. Wiegner, *The Navier–Stokes equations a neverending challenge?* Jahresber. Deutsch. Math.-Verein., 101(1)(1999), 1–25.
- [42] M. Winkler, *Does a volume-filling effect always prevent chemotactic collapse*, Math. Methods Appl. Sci., 33(2010), 12–24.
- [43] M. Winkler, *Aggregation vs. global diffusive behavior in the higher-dimensional Keller–Segel model*, J. Diff. Eqns., 248(2010), 2889–2905.

- [44] M. Winkler, *Absence of collapse in a parabolic chemotaxis system with signal-dependent sensitivity*, Math. Nachr., 283(11)(2010), 1664–1673.
- [45] M. Winkler, *Boundedness in the higher-dimensional parabolic–parabolic chemotaxis system with logistic source*, Comm. Partial Diff. Eqns., 35(2010), 1516–1537.
- [46] M. Winkler, *Global large-data solutions in a chemotaxis–(Navier–)Stokes system modeling cellular swimming in fluid drops*, Comm. Partial Diff. Eqns., 37(2012), 319–351.
- [47] M. Winkler, *Finite-time blow-up in the higher-dimensional parabolic–parabolic Keller–Segel system*, J. Math. Pures Appl., 100(2013), 748–767.
- [48] M. Winkler, *Stabilization in a two-dimensional chemotaxis–Navier–Stokes system*, Arch. Ration. Mech. Anal., 211(2014), 455–487.
- [49] M. Winkler, *Boundedness and large time behavior in a three-dimensional chemotaxis–Stokes system with nonlinear diffusion and general sensitivity*, Calculus of Variations and Partial Diff. Eqns., (54)(2015), 3789–3828.
- [50] M. Winkler, *Global weak solutions in a three-dimensional chemotaxis–Navier–Stokes system*, Ann. Inst. H. Poincaré Anal. Non Linéaire, 33(5)(2016), 1329–1352.
- [51] M. Winkler, *How far do oxytaxis-driven forces influence regularity in the Navier–Stokes system?* Trans. Am. Math. Soc., 369(2017), 3067–125.
- [52] M. Winkler, *Does fluid interaction affect regularity in the three-dimensional Keller–Segel System with saturated sensitivity?*, J. Math. Fluid Mechanics, 20(4)(2018), 1889–1909.
- [53] M. Winkler, *A three-dimensional Keller–Segel–Navier–Stokes system with logistic source: Global weak solutions and asymptotic stabilization*, J. Func. Anal., 276(5)(2019), 1339–1401.
- [54] M. Winkler, K. C. Djie, *Boundedness and finite-time collapse in a chemotaxis system with volume-filling effect*, Nonlinear Anal. TMA., 72(2010), 1044–1064.

- [55] C. Xue, *Macroscopic equations for bacterial chemotaxis: integration of detailed biochemistry of cell signaling*, J. Math. Biol., 70(2015), 1–44.
- [56] C. Xue, H. G. Othmer, *Multiscale models of taxis-driven patterning in bacterial populations*, SIAM J. Appl. Math., 70(1)(2009), 133–167.
- [57] Q. Zhang, X. Zheng, *Global well-posedness for the two-dimensional incompressible chemotaxis–Navier–Stokes equations*, SIAM J. Math. Anal., 46(2014), 3078–3105.
- [58] J. Zheng, *Boundedness of solutions to a quasilinear parabolic–elliptic Keller–Segel system with logistic source*, J. Diff. Eqns., 259(1)(2015), 120–140.
- [59] J. Zheng, *Boundedness in a three-dimensional chemotaxis–fluid system involving tensor-valued sensitivity with saturation*, J. Math. Anal. Appl., 442(1)(2016), 353–375.
- [60] J. Zheng, *A note on boundedness of solutions to a higher-dimensional quasi-linear chemotaxis system with logistic source*, Zeitsc. Angew. Mathe. Mech., 97(4)(2017), 414–421.
- [61] J. Zheng, *Global weak solutions in a three-dimensional Keller–Segel–Navier–Stokes system with nonlinear diffusion*, J. Diff. Eqns., 263(5)(2017), 2606–2629.
- [62] J. Zheng, *An optimal result for global classical and bounded solutions in a two-dimensional Keller–Segel–Navier–Stokes system with sensitivity*, arXiv:1903.01033, 2019.
- [63] J. Zheng, *Global weak solutions in a three-dimensional Keller–Segel–Navier–Stokes system modeling coral fertilization*, arXiv: 1905.08647, 2019.
- [64] J. Zheng, *An optimal result for global existence and boundedness in a three-dimensional Keller–Segel–Stokes system with nonlinear diffusion*, J. Diff. Eqns., (4)(267)(2019), 2385–2415.
- [65] J. Zheng, Y. Ke, *Large time behavior of solutions to a fully parabolic chemotaxis–haptotaxis model in N dimensions*, J. Diff. Eqns., 266(2019), 1969–2018.

- [66] J. Zheng, Y. Wang, *Boundedness and decay behavior in a higher-dimensional quasilinear chemotaxis system with nonlinear logistic source*, *Compu. Math. Appl.*, 72(10)(2016), 2604–2619.
- [67] J. Zheng, Y. Wang, *A note on global existence to a higher-dimensional quasilinear chemotaxis system with consumption of chemoattractant*, *Discrete Contin. Dyn. Syst. Ser. B*, 22(2)(2017), 669–686.