

Policy Evaluation with Latent Confounders via Optimal Balance

Andrew Bennett*
Cornell University
awb222@cornell.edu

Nathan Kallus*
Cornell University
kallus@cornell.edu

Abstract

Evaluating novel contextual bandit policies using logged data is crucial in applications where exploration is costly, such as medicine. But it usually relies on the assumption of no unobserved confounders, which is bound to fail in practice. We study the question of policy evaluation when we instead have proxies for the latent confounders and develop an importance weighting method that avoids fitting a latent outcome regression model. We show that unlike the unconfounded case no single set of weights can give unbiased evaluation for all outcome models, yet we propose a new algorithm that can still provably guarantee consistency by instead minimizing an adversarial balance objective. We further develop tractable algorithms for optimizing this objective and demonstrate empirically the power of our method when confounders are latent.

1 Introduction

Personalized intervention policies are of increasing importance in education (Mandel et al., 2014), healthcare (Bertsimas et al., 2017), and public policy (Kube et al., 2019). In many of these domains exploration is costly or otherwise prohibitive, and so it is crucial to evaluate new policies using existing observational data. Usually, this relies on an assumption of no unobserved confounding (aka unconfoundedness or ignorability): that conditioned on observables, interventions are independent of idiosyncrasies that affect outcomes, so that counterfactuals can be reliably and correctly predicted. In particular, this enables the use of inverse propensity score (IPS) estimators of policy value (Beygelzimer & Langford, 2009; Li et al., 2011; Kallus & Zhou, 2018) that eschew the need to actually fit outcome prediction models and doubly robust estimators that work even if such models are misspecified (Dudík et al., 2011).

In practice, however, it may be unlikely that we observe confounders exactly. Nonetheless, if we observe very many features they may serve as good proxies

*Alphabetical order.

for the true confounders, which can enable an alternative route to identification (Louizos et al., 2017; Kallus et al., 2018). In particular, noisy observations of true confounders can serve as valid proxies. For example, if intelligence is latent but affects both selection and outcome, we can instead use many noisy observations of intelligence such as school grades, IQ test, etc. Similarly, many medical measurements taken together can serve as proxies for underlying healthfulness.

In this paper, we study the problem of policy evaluation from observational data where we observe proxies instead of true confounders and we develop new weighting estimators based on optimizing balance in the latent confounders. Unlike the unconfounded setting where IPS weights ensure balance regardless of outcome model, we show that in this new setting there cannot exist any single set of weights that ensure such unbiasedness regardless of outcome model. Instead, we develop an adversarial objective that bounds the conditional mean square error (CMSE) of any weighted estimator and, by appealing to game theoretic and empirical process arguments, we show that this objective can actually be driven to zero by a single set of weights. We therefore propose a novel policy evaluation method that minimizes this objective, thus provably ensuring consistent estimation in the face of latent confounders. We develop tractable algorithms for this optimization problem. Finally, we provide empirical evidence demonstrating our method’s consistent evaluation compared to standard evaluation methods and its improved performance compared to using fitted latent outcome models.

2 Problem

2.1 Setting and Assumptions

We consider a contextual decision making setting with m possible treatments (aka actions or interventions). Each unit is associated with a set of potential outcomes $Y(1), \dots, Y(m) \in \mathbb{R}$ corresponding to the reward/loss for each treatment, an observed treatment $T \in \{1, \dots, m\}$, an observed outcome $Y = Y(T)$, true but latent confounders $Z \in \mathcal{Z} \subseteq \mathbb{R}^p$, and observed covariates $X \in \mathcal{X} \subseteq \mathbb{R}^q$. Our data consists of iid observations X_i, T_i, Y_i of X, T, Y . Both the latent confounders and potential outcomes of unassigned treatments are unobserved. Note that $Y_i = Y_i(T_i)$ encapsulates the assumptions of consistency between observed and potential outcomes and non-interference between units.

A *policy* is a rule for assigning the probability of each treatment option given the observed covariates X . Given a policy π , we use the notation $\pi_t(x)$ to indicate the probability of assigning treatment t when observed covariates are x . We define the *value* of a policy, τ^π , as the expected outcome that would be obtained from following the policy in the population. Formally:

Definition 1 (Policy Value). $\tau^\pi = \mathbb{E}[\sum_{t=1}^m \pi_t(X)Y(t)]$.

We encapsulate the assumption that Z are sufficient for unconfoundedness and that X is a proxy for Z in the following assumption. Figure 1 provides a representation of this




Figure 1: DAG representation of problem.

setting using a causal DAG (Pearl, 2000). Note importantly that we do *not* assume ignorability given X .

Assumption 1 (Z are true confounders). *For every $t \in \{1, \dots, m\}$, $Y(t)$ is independant of (X, T) , given Z .*

We next define the average mean outcome given Z and its conditional expectations given observables:

$$\begin{aligned}\mu_t(z) &= \mathbb{E}[Y(t) \mid Z = z], \\ \nu_t(x, t') &= \mathbb{E}[\mu_t(Z) \mid X = x, T = t'] = \mathbb{E}[Y(t) \mid X = x, T = t'], \\ \rho_t(x) &= \mathbb{E}[\mu_t(Z) \mid X = x] = \mathbb{E}[Y(t) \mid X = x].\end{aligned}$$

We further define the propensity function and its conditional expectation given observables:

$$\begin{aligned}e_t(z) &= \mathbb{P}(T = t \mid Z = z), \\ \eta_t(x) &= P(T = t \mid X = x) = \mathbb{E}[e_t(Z) \mid X = x].\end{aligned}$$

Finally, we denote by $\varphi(z; x, t)$ the conditional density of Z given $X = x, T = t$. This density represents the latent variable model underlying the observables. For example, this can be a Gaussian mixture model, a PCA-type model as in (Kallus et al., 2018), or a deep variational autoencoder as in (Louizos et al., 2017). Because we focus on how one might use such a latent model rather than the estimation of this model, we just assume we have some oracle for calculating its values. (Note that for fair comparison, in experiments in Section 5, we similarly let the outcome regression methods use this oracle.)

We further make the following regularity assumptions:

Assumption 2 (Weak Overlap). $\mathbb{E}[e_t^{-2}(Z)] < \infty$

Remark 1. *Given Assumption 2 it trivially follows that for every $t \in \{1, \dots, m\}$, $x \in \mathcal{X}$, $z \in \mathcal{Z}$ that $e_t(z) > 0$ and $\eta_t(x) > 0$.*

Assumption 3 (Bounded Variance). *The conditional variance of our potential outcomes given X, T is bounded: $\mathbb{V}[Y(t) \mid X, T] \leq \sigma^2$.*

2.2 The Policy Evaluation Task

The problem we consider is to estimate the policy value τ^π given a policy π and data $X_{1:n}, T_{1:n}, Y_{1:n}$. One standard approach to this is the *direct method* (Qian & Murphy, 2011), which given an estimate $\hat{\rho}_t$ of ρ_t predicts the policy value as

$$\hat{\tau}_{\hat{\rho}}^\pi = \frac{1}{n} \sum_{i=1}^n \sum_{t=1}^m \pi_t(X_i) \hat{\rho}_t(X_i). \quad (1)$$

However this method is known to be biased and doesn't generalize well (Beygelzimer & Langford, 2009). Furthermore given Assumption 1 $\hat{\rho}$ is not

straightforward to estimate, since the mean value of Y observed in our logged data given $X = x$ and $T = t$ is $\nu_t(x, t)$ not $\rho_t(x)$, so fitting $\hat{\rho}$ would require controlling for the effects of the unobserved Z .

An alternative to this is to come up with weights $W_{1:n} = f_W(X_{1:n}, T_{1:n})$ according to some function f_W of the observed covariates and treatments, in order to re-weight the outcomes to look more like those that would be observed under π . Using these weights we can define the *weighted* estimator

$$\hat{\tau}_W^\pi = \frac{1}{n} \sum_{i=1}^n W_i Y_i. \quad (2)$$

This weighted estimator has the advantage that it does not require modeling the outcome distributions. Furthermore we could combine the weights $W_{1:n}$ with an outcome model $\hat{\rho}_t$ to calculate the *doubly robust* estimator (Dudík et al., 2011), which is defined as

$$\hat{\tau}_{W, \hat{\rho}}^\pi = \frac{1}{n} \sum_{i=1}^n \sum_{t=1}^m \pi_t(X_i) \hat{\rho}_t(X_i) + \frac{1}{n} \sum_{i=1}^n W_i (Y_i - \hat{\rho}_{T_i}(X_i)). \quad (3)$$

The doubly robust estimator is known to be consistent when either the weighted or direct estimator is consistent and can attain local efficiency (Chernozhukov et al., 2016; Robins, 1999; Robins et al., 1994; Scharfstein et al., 1999).

Various approaches exist for coming up with weights for either the weighted or doubly robust estimators, which we discuss below. However none of these methods are applicable given Assumption 1, and so we develop a theory for weighting using proxy variables in Section 3.

2.3 Related Work

One of the most standard approaches for policy evaluation is using the weighted or doubly robust estimator defined in Eqs. (2) and (3), using inverse propensity score (IPS) weights. These are given by $W_i = \pi_{T_i}(X_i)/e_{T_i}(Z_i)$ (Bottou et al., 2013), where e_t are known or estimated logging probabilities. Since these weights can be extreme, both normalization (Austin & Stuart, 2015; Lunceford & Davidian, 2004; Swaminathan & Joachims, 2015b) and clipping (Elliott, 2008; Ionides, 2008; Swaminathan & Joachims, 2015a) are often employed. In addition some other approaches include recursive partitioning (Kallus, 2017a). None of these methods are applicable to our setting however, since we do not know the true confounders $Z_{1:n}$.

An alternative to approaches based on fixed formulae for computing the importance weights is to compute weights that optimize an imbalance objective function (Athey et al., 2018; Kallus, 2017b, 2018b). For policy evaluation, Kallus (2018a) propose to choose weights that adversarially minimize the conditional mean squared error of policy evaluation in the worst case of possible mean outcome functions in some reproducing kernel Hilbert space (RKHS) class, by solving a linearly constraint quadratic program (LCQP). Our work follows a

very similar style to this, however instead of using the true confounders we only assume access to proxies, and we prove our theory for more general families of functions.

Finally there has been a long history of work in causal inference using proxies for true confounders (Wickens, 1972; Frost, 1979). As in our problem setup, much of this work is based on the model of using an identified latent variable model for the proxies (Wooldridge, 2009; Pearl, 2012; Kuroki & Pearl, 2014; Miao et al., 2016; Edwards et al., 2015). Some recent work on this problem involves using techniques such as matrix completion (Kallus, 2018a) or variational autoencoders (Louizos et al., 2017) to infer confounders from the proxies. In addition, there is a variety of work that studies sufficient conditions for the identifiability of latent confounder models (Cai & Kuroki, 2008; Pearl, 2012; Miao et al., 2016). Our work is complementary to this line of research in that we assume access to an identified latent confounder model, but do not study how to identify such models. Furthermore our work is novel in combining proxy variable models with optimal balancing and applying it to finding importance weights for policy evaluation.

3 Weight-Balancing Objectives

3.1 Infeasibility of IPS-Style Unbiased Weighting

If we had unconfoundedness given X (*i.e.*, $Y(t) \perp\!\!\!\perp T \mid X$), the IPS weights $\pi_T(X)/\eta_T(X)$ are immediately gotten as the solution to making every term in the weighted sum Eq. (2) unbiased:

$$\mathbb{E}[W(X, T)\delta_{T=t}Y(t)] = \mathbb{E}[\pi_t(X)Y(t)]. \quad (4)$$

Notably the IPS weights do not depend on the outcome function. However, without unconfoundedness given X and given only Assumptions 1 to 3, this approach fails.

Theorem 1. *If $W(x, t)$ satisfies Eq. (4) then for any $t \in \{1, \dots, m\}$*

$$W(X, t) = \pi_t(X) \frac{\sum_{t' \in \tau} \eta_{t'}(X) \nu_t(X, t') + \Omega_t(X)}{\eta_t(X) \nu_t(X, t)}, \quad (5)$$

for some $\Omega_t(x)$ such that $\mathbb{E}\Omega_t(X) = 0 \forall t$.

The proof of Theorem 1 is given in Appendix A.1.

Note that *if* we had unconfoundedness given X then $\nu_t(x, t') = \nu_t(x, t) = \rho_t(x)$ so that choosing $\Omega_t(X) = 0$ would recover the standard IPS weights. However, in our setting we generally have $\nu_t(x, t') \neq \nu_t(x, t)$, and so Theorem 1 tells us that we cannot do unbiased IPS-style weighted evaluation without knowing the mean outcome functions $\nu_t(x, t')$. In particular, there exists no single weight function that is simultaneously unbiased for all outcome functions.

On the other hand, Theorem 1 tells us that there do exist some weights that give unbiased and consistent policy evaluation via Eq. (2) or Eq. (3): we just

may not be able to calculate them. The existence of such weights motivates our subsequent approach, which seeks weights that mimic these weights for a wide class of possible outcome functions.

3.2 Adversarial Error Objective

Over all weights that are functions of $X_{1:n}, T_{1:n}$, the *optimal* choice of weights for estimating τ^π via Eq. (2) would minimize the (unknown) conditional MSE (CMSE):

$$\mathbb{E}[(\hat{\tau}_W^\pi - \tau^\pi)^2 \mid X_{1:n}, T_{1:n}]. \quad (6)$$

In particular, the weights in Eq. (5) achieve $O_p(1/n)$ control on this CMSE for many outcome functions, as long as the denominator is well behaved, which can be seen by applying concentration inequalities to Eq. (6). However, as discussed above the outcome function is unknown and these weights are therefore practically infeasible. Our aim is to find weights with similar near-optimal behavior but that do not depend on the particular unknown outcome function. To do this, we will find an upper bound for Eq. (6) that we can actually compute.

Let $f_{it} = W_i \delta_{T_{it}} - \pi_t(X_i)$ and

$$J(W, \mu) = \left(\frac{1}{n} \sum_{i=1}^n \sum_{t=1}^m f_{it} \nu_t(X_i, T_i) \right)^2 + \frac{2\sigma^2}{n^2} \|W\|_2^2,$$

where we embedded the dependence on μ inside $\nu_t(x, t') = \mathbb{E}[\mu_t(Z) \mid X = x, T = t']$.

Theorem 2. $\mathbb{E}[(\hat{\tau}_W^\pi - \tau^\pi)^2 \mid X_{1:n}, T_{1:n}] \leq 2J(W, \mu) + O_p(1/n)$.

We prove this in Appendix A.2.

Therefore, if we find weights that obtain $O_p(1/n)$ control on $J(W, \mu)$, we can ensure that we also have $O_p(1/n)$ control on $\mathbb{E}[(\hat{\tau}_W^\pi - \tau^\pi)^2 \mid X_{1:n}, T_{1:n}]$. Combined with the following result, which follows from (Kallus, 2016, Lemma 31), this would give root-n consistent estimation.

Lemma 1. *If $\mathbb{E}[(\hat{\tau}_W^\pi - \tau^\pi)^2 \mid X_{1:n}, T_{1:n}] = O_p(1/n)$ then $\hat{\tau}_W^\pi = \tau^\pi + O_p(1/\sqrt{n})$.*

It remains to find weights that control $J(W, \mu)$. The key obstacle for this is that μ is unknown. Instead, we show how we can obtain weights that control $J(W, \mu)$ over a whole class of given functions μ .

Suppose we are given a set \mathcal{F} of functions mapping \mathcal{Z} to \mathbb{R}^m , where each $\mu \in \mathcal{F}$ corresponds to a vector of mean outcome functions $\mu = (\mu_1, \dots, \mu_m)$. Then, motivated by Theorem 2 and Lemma 1, we define our adversarial optimization problem as

$$\mathcal{W}^* = \arg \min_{W \in \mathcal{W}} \sup_{\mu \in \mathcal{F}} J(W, \mu). \quad (7)$$

One question the reader might ask at this point is why not solve the above optimization problem by ignoring the hidden confounders and directly balancing the conditional mean outcome functions $\nu_t(x, t')$. The problem is that this would

be impossible to do over any kind of generic flexible function space, since we have no data corresponding to terms in the form $\nu_t(x, t')$ when $t \neq t'$, so this is akin to an overlap problem. Conversely, if we were to ignore the conditioning on t and balance against functions of the form $\nu_t(x) = \nu_t(x, t)$ this would be inadequate, as we couldn't hope for such a space to cover the true μ since we don't assume ignorability given Z .

In light of these limitations, we can view what we are doing in optimizing Eq. (7) using an identified model $\varphi(z; x, t)$ as implicitly balancing some controlled space of functions $\nu_t(x, t')$ that do not have this overlap issue between different t values. The following lemma makes this explicit, as it implies that that the terms $\nu_t(x, t')$ are all mutually bounded by each other for fixed x and t :

Lemma 2. *Assuming $\|\mu_t\|_\infty \leq b$, under Assumption 2, for all $x \in \mathcal{X}$, and $t, t', t'' \in \{1, \dots, m\}$ we have*

$$|\nu_t(x, t'')| \leq \frac{\eta_{t'}(x)}{\eta_{t''}(x)} \sqrt{8b \mathbb{E}[e_t^{-2}(Z) \mid X = x, T = t']} |\nu_t(x, t')|.$$

We prove this in Appendix A.3.

3.3 Consistency of Adversarially Optimal Estimator

Now we analyze the consistency of our weighted estimator based on Eq. (7). Given Lemma 1, all we need to justify to prove consistency is that $\mu \in \mathcal{F}$ and that $\inf_W \sup_{\mu \in \mathcal{F}} J(W, \mu) \in O_p(\frac{1}{n})$. Define \mathcal{F}_t as the space of all functions for treatment level t allowed by \mathcal{F} . That is $\mathcal{F}_t = \{\mu_t : \exists(\mu'_1, \dots, \mu'_m) \in \mathcal{F} \text{ with } \mu'_t = \mu_t\}$. We will use the following assumptions about \mathcal{F} to prove control of J :

Assumption 4 (Normed). *For each $t \in \{1, \dots, m\}$ there exists a norm $\|\cdot\|_t$ on $\text{span}(\mathcal{F}_t)$, and there exists a norm $\|\cdot\|$ on $\text{span}(\mathcal{F})$ which is defined given some \mathbb{R}^m norm as $\|\mu\| = \|(\|\mu_1\|_1, \dots, \|\mu_m\|_m)\|$.*

Assumption 5 (Absolutely Star Shaped). *For every $\mu \in \mathcal{F}$ and $|\lambda| \leq 1$, we have $\lambda\mu \in \mathcal{F}$.*

Assumption 6 (Convex Compact). *\mathcal{F} is convex and compact*

Assumption 7 (Square Integrable). *For each $t \in \{1, \dots, m\}$ the space \mathcal{F}_t is a subset of $\mathcal{L}^2(\mathcal{Z})$, and its norm dominates the \mathcal{L}^2 norm (i.e., $\inf_{\mu_t \in \mathcal{F}_t} \|\mu_t\| / \|\mu_t\|_{\mathcal{L}^2} > 0$).*

Assumption 8 (Nondegeneracy). *Define $\mathcal{B}(\gamma) = \{\mu \in \text{span}(\mathcal{F}) : \|\mu\| \leq \gamma\}$. Then we have $\mathcal{B}(\gamma) \subseteq \mathcal{F}$ for some $\gamma > 0$.*

Assumption 9 (Boundedness). $\sup_{\mu \in \mathcal{F}} \|\mu\|_\infty < \infty$.

Definition 2 (Rademacher Complexity). $\mathcal{R}_n(\mathcal{F}) = \mathbb{E}[\sup_{f \in \mathcal{F}} \frac{1}{n} \sum_{i=1}^n \epsilon_i f(Z_i)]$, where ϵ_i are iid Rademacher random variables.

Assumption 10 (Complexity). *For each $t \in \{1 \dots, m\}$ we have $\mathcal{R}_n(\mathcal{F}_t) = o(1)$.*

These assumptions are satisfied for many commonly-used families of functions, such as RKHS spaces and families of neural networks. We shall prove this claim for RKHS spaces in Section 4.

In order to justify that we can control J , first we will show that these assumptions allow us to reverse the order of minimization and maximization in our optimization problem. This means we can reduce problem to finding weights to control any particular μ rather than controlling all of \mathcal{F} .

Lemma 3. *Let $B(W, \mu) = \frac{1}{n} \sum_{i=1}^n \sum_{t=1}^m f_{it} \nu_t(X_i, T_i)$. Then under Assumptions 5 to 7 for every $M > 0$ we have the bound*

$$\min_W \sup_{\mu \in \mathcal{F}} J(W, \mu) \leq \sup_{\mu \in \mathcal{F}} \min_{\|W\|_2 \leq M} B(W, \mu)^2 + \frac{\sigma^2}{n^2} M^2.$$

We prove this in Appendix A.4.

Next, we note that Lemma 3 means that we can choose of weights given μ to set $B(W, \mu) = 0$, and therefore we have our desired control as long as we can justify that these weights have controlled euclidean norm. Using this strategy and optimizing for the weights of this kind with minimum euclidean norm, we are able to prove the following:

Lemma 4. *Under Assumptions 4 to 10 we have $\inf_W \sup_{\mu \in \mathcal{F}} J(W, \mu) = O_p(1/n)$.*

We prove this in Appendix A.5. This is the key lemma in proving our main consistency theorem:

Theorem 3. *Under Assumptions 4 to 10 and assuming that $\mu \in \mathcal{F}$ we have $\hat{\tau}_{W^*}^\pi = \tau^\pi + O_p(1/\sqrt{n})$.*

This theorem follows immediately from our previous results, since $\mu \in \mathcal{F}$ and Lemma 4 imply that $J(W^*, \mu) = O_p(1/n)$. This combined with Theorem 2 imply that $\mathbb{E}[(\hat{\tau}_{W^*}^\pi - \tau^\pi)^2 \mid X_{1:n}, T_{1:n}] = O_p(1/n)$, which in turn combined with Lemma 1 gives us our result.

4 Algorithms for Optimal Kernel Balancing

4.1 Kernel Function Class

We now provide an algorithm for optimal balancing when our function class consists of vectors of RKHS functions. Formally, given a kernel K and corresponding RKHS norm $\|\cdot\|_K$, we define the space \mathcal{F}^K as follows:

Definition 3 (Kernel Class). $\mathcal{F}^K = \{\mu : \|\mu\| \leq 1\}$, where $\|(\mu_1, \dots, \mu_m)\| = \sqrt{\sum_{t=1}^m \|\mu_t\|_K^2}$.

Theorem 4. *Assuming K is a Mercer kernel (Zhou, 2002) and is bounded, \mathcal{F}^K satisfies Assumptions 4 to 10.*

We prove this in Appendix A.6.

We can remark that the commonly used Gaussian kernel is both Mercer and bounded, so it satisfies the conditions of Theorem 4. Given this, and assuming that \mathcal{F}^K covers the real mean outcome function μ , we can apply Theorem 3 to see that solving Eq. (7) using \mathcal{F}^K gives consistent evaluation.

We can note that the \mathcal{F}^K having maximum norm 1 is without loss of generality, because if we wanted the maximum norm to instead be γ we could replace the Σ matrix by $\Gamma = \frac{1}{\gamma}\Sigma$ in our objective function, resulting in an equivalent re-scaled optimization problem. To make this explicit, we will replace the Σ matrix in the objective with Γ in this section, where it is assumed that Γ is freely chosen as a hyperparameter.

4.2 Kernel Balancing Algorithm

In order to optimize Eq. (7) over a class of kernel functions as defined by Definition 3, we can observe that the definition of $J(W, \mu)$ looks very similar to the adversarial objective of Kallus (2018a), except that we have $\nu_t(X_i, T_i)$ terms instead of $\mu_t(X_i)$ terms. This motivates the idea that, given our identified posterior model $\varphi(z; x, t)$, we may be able to employ a similar quadratic programming (QP)-based approach. The following theorem makes this explicit, by defining a QP objective for W that we can approximate by sampling from φ :

Theorem 5. *Define $Q_{ij} = \mathbb{E}[K(Z_i, Z'_j)]$, $G_{ij} = \frac{1}{n^2}(Q_{ij}\delta_{T_i T_j} + \Gamma_{ij})$, and $a_i = \frac{2}{n^2} \sum_{j=1}^n Q_{ij}\pi_{T_j}(X_i)$, where for each i Z_i and Z'_i are iid shadow variables. Then for some c that is constant in W we have the identity*

$$\sup_{\mu \in \mathcal{F}^K} J(W, \mu) = W^T G W - a^T W + c.$$

We prove this in Appendix A.7.

Given this our balancing algorithm is natural and straightforward, and is summarized by Algorithm 1. Note that we provide an optional weight space constraint \mathcal{W} in this algorithm, since standard weighted estimator approaches for policy evaluation regularize by forcing constraints such as $W \in n\Delta^n$. Under this kind of constraint our unconstrained QP becomes a LCQP. However that our theory does not support this constraint, and that we find it hurts performance in practice, especially when Γ is large, so we do not use this constraint in our experiments.

5 Experiments

5.1 Experimental Setup

We now present a brief set of experiments to explore our methodology. The aim of these experiments is to be a proof of concept of our theory. We seek to show that given an identified posterior model φ policy as discussed in Section 2.1, evaluation using the weights defined by Eq. (7) can give unbiased policy evaluation

Algorithm 1 Optimal Kernel Balancing

Input: Data $(X_{1:n}, T_{1:n})$, policy π , kernel function K , posterior density φ , regularization matrix Γ , number samples B , optional weight space \mathcal{W} (defaults to \mathbb{R}^n if not provided)

Output: Optimal balancing weights $W_{1:n}$

- 1: **for** $i \in \{1, \dots, n\}$ **do**
 - 2: **Sample Data.** Draw B data points Z_i^b from the posterior $\varphi(\cdot; X_i, T_i)$
 - 3: **end for**
 - 4: **Estimate Q.** Calculate $Q_{ij} = \frac{1}{B^2} \sum_{b=1}^B \sum_{c=1}^B K(Z_i^b, Z_j^c)$
 - 5: **Calculate QP Inputs.** Calculate $G_{ij} = Q_{ij} \delta_{T_i T_j} + \Gamma_{ij}$, and $a_i = 2 \sum_{j=1}^n Q_{ij} \pi_{T_j}(X_i)$,
 - 6: **Solve Quadratic Program.** Calculate $W = \arg \min_{W \in \mathcal{W}} W^T G W - a^T W$
-

even in the face of sufficiently strong confounding where standard benchmark approaches that rely on ignorability given X fail. We experiment with the following generalized linear model-style scenario:

$$\begin{array}{lll} Z \sim \mathcal{N}(0, 1) & X \sim \mathcal{N}(\alpha^T Z + \alpha_0, \sigma_X^2) & P_T = \beta^T Z + \beta_0 \\ T \sim \text{softmax}(P_T) & W(t) \sim \mathcal{N}(\zeta(t)^T Z + \zeta_0(t), \sigma_Y^2) & Y(t) = g(W(t)) \end{array}$$

In our experiments Z is 1-dimensional, X is 10-dimensional, and we have two possible treatment levels ($m = 2$). We experiment with a parametric policy π and multiple link functions g as follows:

$$\pi_t(X) = \frac{\exp(\psi_t^T X)}{\exp(\psi_1^T X) + \exp(\psi_2^T X)}$$

step: $g(w) = 3\mathbb{1}_{\{w \geq 0\}} - 6$ **exp:** $g(w) = \exp(w)$ **cubic:** $g(w) = w^3$ **linear:** $g(w) = w$

We experiment with the following methods in this evaluation:

1. **OptZ** Our method, using $\Gamma = \gamma \text{Identity}(n)$ for $\gamma \in \{0.001, 0.2, 1.0, 5.0\}$.
2. **IPS** IPS weights based on X using estimated $\hat{\eta}_t$.
3. **OptX** The optimal weighting method of Kallus (2018a) with same values of Γ as our method.
4. **DirX** Direct method by fitting $\hat{\rho}_t(x)$ incorrectly assuming ignorability given X .
5. **DirZ** Direct method by first fitting $\hat{\mu}_t$ using posterior samples from φ , then using the estimate $\hat{\rho}_t(x) = (1/D) \sum_{i=1}^D \hat{\mu}_t(z'_i)$, where z'_i are sampled from $\varphi(\cdot; x, t)$.
6. **D:W** Doubly robust estimation using direct estimator **D** and weighted estimator **W**.

Finally we detail all choices for scenario parameters in Appendix B.1, and provide implementation details of our methods in Appendix B.2.¹

5.2 Results

We display results for our experiments using the **step** link function in Tables 1 and 2. For each of $n \in \{200, 500, 1000, 2000\}$ we estimate the RMSE of policy evaluation using each method, as well as doubly robust evaluation using our best performing weights, by averaging over 64 runs. In addition, in Tables 3 and 4 we display the estimated bias from the evaluations. It is clear that the naive methods that assume ignorability given X all hit a performance ceiling, where bias converges to some non-zero value. In particular for IPS we separately ran it on up to one million data points and found that the bias converged to 0.418 ± 0.001 . On the other hand, for our method it appears like we have consistency. This is particularly evident when we look at Table 3, as bias seems to be approximately converging to zero with vanishing variance. We can also observe that doubly robust estimation using either direct method does not appear to improve performance.

It is noteworthy that the **DirZ** benchmark method fails horribly, despite being a correctly specified regression estimate. From our experience we observed that it is difficult to train the μ_t functions accurately if there is a high amount of overlap in the $\varphi(\cdot; x, t)$ posteriors for fixed t . Therefore we postulate that in highly confounded settings this benchmark is inherently difficult to train using a finite number of samples from $\varphi(\cdot; x, t)$, and the result seems to collapse to degenerate solutions.

Finally we note that we observed similar trends to this using our other link functions, and other doubly robust estimators. We present more extensive tables of results in Appendix C. In addition we present some results there on the negative impact on our method’s performance using the constraint $W \in n\Delta^n$, as mentioned in Section 4.2.

6 Conclusion

We presented theory for how to do optimal balancing for policy evaluation when we only have proxies for the true confounders, given an already identified model for the confounders, treatment, and proxies, but not for the outcomes. We provided an adversarial objective for selecting optimal weights given some class of mean outcome functions to be balanced, and proved that under mild conditions these optimal weights result in consistent policy evaluation. In addition, we presented a tractable algorithm for minimizing this objective when our function class is an RKHS, and we conducted a series of experiments to demonstrate that our method can achieve consistent evaluation even under sufficient levels of confounding where standard approaches fail.

¹Code available online at <https://github.com/CausalML/LatentConfounderBalancing>.

n	OptZ_{0.001}	OptZ_{0.2}	OptZ_{1.0}	OptZ_{5.0}	DirX:OptZ_{0.001}	DirZ:OptZ_{0.001}
200	.39 \pm .07	.24 \pm .02	.36 \pm .02	.81 \pm .02	.57 \pm .06	.41 \pm .07
500	.19 \pm .02	.18 \pm .02	.23 \pm .02	.49 \pm .02	.55 \pm .03	.20 \pm .02
1000	.11 \pm .01	.11 \pm .01	.13 \pm .01	.27 \pm .01	.49 \pm .02	.11 \pm .01
2000	.08 \pm .01	.08 \pm .01	.09 \pm .01	.17 \pm .01	.48 \pm .01	.08 \pm .01

Table 1: Convergence of RMSE for for policy evaluation using our weights.

n	IPS	OptX_{0.001}	OptX_{0.2}	OptX_{1.0}	OptX_{5.0}	DirX	DirZ
200	.47 \pm .03	2.0 \pm .03	2.1 \pm .03	2.3 \pm .02	2.5 \pm .02	.52 \pm .02	2.6 \pm .02
500	.48 \pm .03	2.0 \pm .02	2.1 \pm .02	2.3 \pm .02	2.6 \pm .02	.48 \pm .02	2.6 \pm .01
1000	.39 \pm .02	2.0 \pm .01	2.1 \pm .01	2.3 \pm .01	2.5 \pm .01	.48 \pm .02	2.6 \pm .01
2000	.40 \pm .01	2.0 \pm .01	2.1 \pm .01	2.3 \pm .01	2.5 \pm .01	.45 \pm .02	2.6 \pm .01

Table 2: Convergence of RMSE for benchmark methods.

n	OptZ_{0.001}	OptZ_{0.2}	OptZ_{1.0}	OptZ_{5.0}	DirX:OptZ_{0.001}	DirZ:OptZ_{0.001}
200	.03 \pm .39	.11 \pm .21	.29 \pm .21	.78 \pm .18	.43 \pm .38	.05 \pm .40
500	.09 \pm .17	.10 \pm .15	.17 \pm .16	.47 \pm .15	.51 \pm .19	.10 \pm .18
1000	.02 \pm .11	.05 \pm .09	.08 \pm .09	.25 \pm .09	.47 \pm .13	.04 \pm .11
2000	.03 \pm .07	.05 \pm .06	.07 \pm .07	.16 \pm .07	.47 \pm .09	.03 \pm .07

Table 3: Convergence of bias for policy evaluation using our weights.

n	IPS	OptX_{0.001}	OptX_{0.2}	OptX_{1.0}	OptX_{5.0}	DirX	DirZ
200	.40 \pm .25	1.9 \pm .21	2.1 \pm .20	2.3 \pm .19	2.5 \pm .18	.49 \pm .18	2.6 \pm .14
500	.43 \pm .21	2.0 \pm .16	2.1 \pm .15	2.3 \pm .14	2.6 \pm .13	.45 \pm .16	2.6 \pm .12
1000	.37 \pm .12	2.0 \pm .10	2.1 \pm .09	2.3 \pm .09	2.5 \pm .08	.46 \pm .15	2.6 \pm .11
2000	.39 \pm .10	2.0 \pm .08	2.1 \pm .07	2.3 \pm .07	2.5 \pm .07	.42 \pm .17	2.6 \pm .11

Table 4: Convergence of bias for benchmark methods.

For future work we note that the adversarial objective and theory presented here is fairly general, and could be used to develop new algorithms for balancing different function classes such as neural networks. An alternative direction would be to study how best to apply this methodology when an identified model is not already given.

References

- Athey, S., Imbens, G. W., and Wager, S. Approximate residual balancing: debiased inference of average treatment effects in high dimensions. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 80(4):597–623, 2018.
- Austin, P. C. and Stuart, E. A. Moving towards best practice when using inverse probability of treatment weighting (iptw) using the propensity score to estimate causal treatment effects in observational studies. *Statistics in medicine*, 34(28):3661–3679, 2015.
- Bertsimas, D., Kallus, N., Weinstein, A. M., and Zhuo, Y. D. Personalized diabetes management using electronic medical records. *Diabetes care*, 40(2):210–217, 2017.
- Beygelzimer, A. and Langford, J. The offset tree for learning with partial labels. In *Proceedings of the 15th ACM SIGKDD international conference on Knowledge discovery and data mining*, pp. 129–138. ACM, 2009.
- Bottou, L., Peters, J., Candela, J. Q., Charles, D. X., Chickering, M., Portugaly, E., Ray, D., Simard, P. Y., and Snelson, E. Counterfactual reasoning and learning systems: the example of computational advertising. *Journal of Machine Learning Research*, 14(1):3207–3260, 2013.
- Cai, Z. and Kuroki, M. On identifying total effects in the presence of latent variables and selection bias. In *Proceedings of the Twenty-Fourth Conference on Uncertainty in Artificial Intelligence*, pp. 62–69. AUAI Press, 2008.
- Carpenter, B., Gelman, A., Hoffman, M. D., Lee, D., Goodrich, B., Betancourt, M., Brubaker, M., Guo, J., Li, P., and Riddell, A. Stan: A probabilistic programming language. *Journal of statistical software*, 76(1), 2017.
- Chernozhukov, V., Chetverikov, D., Demirer, M., Duflo, E., and Hansen, C. Double machine learning for treatment and causal parameters. *arXiv preprint arXiv:1608.00060*, 2016.
- Dudík, M., Langford, J., and Li, L. Doubly robust policy evaluation and learning. In *Proceedings of the 28th International Conference on International Conference on Machine Learning*, pp. 1097–1104. Omnipress, 2011.

- Edwards, J. K., Cole, S. R., and Westreich, D. All your data are always missing: incorporating bias due to measurement error into the potential outcomes framework. *International journal of epidemiology*, 44(4):1452–1459, 2015.
- Elliott, M. R. Model averaging methods for weight trimming. *Journal of official statistics*, 24(4):517, 2008.
- Frost, P. A. Proxy variables and specification bias. *The review of economics and Statistics*, pp. 323–325, 1979.
- Ionides, E. L. Truncated importance sampling. *Journal of Computational and Graphical Statistics*, 17(2):295–311, 2008.
- Kallus, N. Generalized optimal matching methods for causal inference. *arXiv preprint arXiv:1612.08321*, 2016.
- Kallus, N. Recursive partitioning for personalization using observational data. In *International Conference on Machine Learning (ICML)*, pp. 1789–1798, 2017a.
- Kallus, N. A framework for optimal matching for causal inference. In *Artificial Intelligence and Statistics (AISTATS)*, pp. 372–381, 2017b.
- Kallus, N. Balanced policy evaluation and learning. In *Advances in Neural Information Processing Systems*, pp. 8895–8906, 2018a.
- Kallus, N. Optimal a priori balance in the design of controlled experiments. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 80(1):85–112, 2018b.
- Kallus, N. and Zhou, A. Policy evaluation and optimization with continuous treatments. In *International Conference on Artificial Intelligence and Statistics*, pp. 1243–1251, 2018.
- Kallus, N., Mao, X., and Udell, M. Causal inference with noisy and missing covariates via matrix factorization. In *Advances in Neural Information Processing Systems*, pp. 6921–6932, 2018.
- Kingma, D. P. and Ba, J. Adam: A method for stochastic optimization. *arXiv preprint arXiv:1412.6980*, 2014.
- Kube, A., Das, S., and Fowler, P. J. Allocating interventions based on predicted outcomes: A case study on homelessness services. In *Proceedings of the AAAI Conference on Artificial Intelligence*, 2019.
- Kuroki, M. and Pearl, J. Measurement bias and effect restoration in causal inference. *Biometrika*, 101(2):423–437, 2014.
- Ledoux, M. and Talagrand, M. *Probability in Banach Spaces: isoperimetry and processes*. Springer Science & Business Media, 2013.

- Li, L., Chu, W., Langford, J., and Wang, X. Unbiased offline evaluation of contextual-bandit-based news article recommendation algorithms. In *Proceedings of the fourth ACM international conference on Web search and data mining*, pp. 297–306. ACM, 2011.
- Louizos, C., Shalit, U., Mooij, J. M., Sontag, D., Zemel, R., and Welling, M. Causal effect inference with deep latent-variable models. In *Advances in Neural Information Processing Systems*, pp. 6446–6456, 2017.
- Lunceford, J. K. and Davidian, M. Stratification and weighting via the propensity score in estimation of causal treatment effects: a comparative study. *Statistics in medicine*, 23(19):2937–2960, 2004.
- Mandel, T., Liu, Y.-E., Levine, S., Brunskill, E., and Popovic, Z. Offline policy evaluation across representations with applications to educational games. In *Proceedings of the International Conference on Autonomous Agents and Multiagent Systems*, pp. 1077–1084. International Foundation for Autonomous Agents and Multiagent Systems, 2014.
- Mendelson, S. On the performance of kernel classes. *Journal of Machine Learning Research*, 4(Oct):759–771, 2003.
- Miao, W., Geng, Z., and Tchetgen, E. T. Identifying causal effects with proxy variables of an unmeasured confounder. *arXiv preprint arXiv:1609.08816*, 2016.
- Pearl, J. *Causality: models, reasoning and inference*. Cambridge University Press, 2000.
- Pearl, J. On measurement bias in causal inference. *arXiv preprint arXiv:1203.3504*, 2012.
- Qian, M. and Murphy, S. A. Performance guarantees for individualized treatment rules. *Annals of statistics*, 39(2):1180, 2011.
- Robins, J. M. Robust estimation in sequentially ignorable missing data and causal inference models. In *Proceedings of the American Statistical Association*, pp. 6–10, 1999.
- Robins, J. M., Rotnitzky, A., and Zhao, L. P. Estimation of regression coefficients when some regressors are not always observed. *Journal of the American statistical Association*, 89(427):846–866, 1994.
- Scharfstein, D. O., Rotnitzky, A., and Robins, J. M. Adjusting for nonignorable drop-out using semiparametric nonresponse models. *Journal of the American Statistical Association*, 94(448):1096–1120, 1999.
- Swaminathan, A. and Joachims, T. Counterfactual risk minimization: Learning from logged bandit feedback. In *ICML*, pp. 814–823, 2015a.

- Swaminathan, A. and Joachims, T. The self-normalized estimator for counterfactual learning. In *Advances in Neural Information Processing Systems*, pp. 3231–3239, 2015b.
- Wickens, M. R. A note on the use of proxy variables. *Econometrica: Journal of the Econometric Society*, pp. 759–761, 1972.
- Wooldridge, J. M. On estimating firm-level production functions using proxy variables to control for unobservables. *Economics Letters*, 104(3):112–114, 2009.
- Zhou, D.-X. The covering number in learning theory. *Journal of Complexity*, 18(3):739–767, 2002.

A Omitted Proofs

A.1 Proof of Theorem 1

First we note that $W(X, T)Y - \sum_t \pi_t(X)Y(t) = \sum_t (W(X, t)\delta_{Tt} - \pi_t(X))Y(t)$. Then analyzing each summand separately, we can obtain:

$$\begin{aligned}
& \mathbb{E}[W(X, t)\delta_{Tt}Y(t) - \pi_t(X)Y(t)] \\
&= \mathbb{E}[\mathbb{E}[W(X, t)\delta_{Tt}Y(t) - \pi_t(X)Y(t) \mid X]] \\
&= \mathbb{E}[W(X, t)\mathbb{E}[\delta_{Tt}Y(t) \mid X] - \pi_t(X)\mathbb{E}[Y(t) \mid X]] \\
&= \mathbb{E}[W(X, t)\mathbb{E}[\mathbb{E}[\delta_{Tt}Y(t) \mid X, T] \mid X] - \pi_t(X)\mathbb{E}[\mathbb{E}[Y(t) \mid X, Z] \mid X]] \\
&= \mathbb{E}[W(X, t)\mathbb{E}[\delta_{Tt}\mathbb{E}[Y(t) \mid X, T = t] \mid X] - \pi_t(X)\mathbb{E}[\mu_t(Z) \mid X]] \\
&= \mathbb{E}[W(X, t)\mathbb{E}[\delta_{Tt} \mid X]\mathbb{E}[\mu_t(Z) \mid X, T = t] - \pi_t(X)\mathbb{E}[\mu_t(Z) \mid X]] \\
&= \mathbb{E}[W(X, t)\eta_t(X)\mathbb{E}[\mu_t(Z) \mid X, T = t] - \pi_t(X)\mathbb{E}[\mu_t(Z) \mid X]] \\
&= \mathbb{E}\left[W(X, t)\eta_t(X)\nu_t(X, t) - \pi_t(X)\sum_{t'}\eta_{t'}(X)\nu_t(X, t')\right]
\end{aligned}$$

Therefore the solution to the equation $\mathbb{E}[W(X, t)\delta_{Tt}Y(t) - \pi_t(X)Y(t)] = 0$ is given by:

$$W(X, t)\eta_t(X)\nu_t(X, t) - \pi_t(X)\sum_{t'}\eta_{t'}(X)\nu_t(X, t') = \Omega_t(X)$$

where $\Omega_t(X)$ is any arbitrary function of X with mean zero. Solving this for W gives

$$W(X, t) = \frac{\pi_t(X)\sum_{t'}\eta_{t'}(X)\nu_t(X, t') + \Omega_t(X)}{\eta_t(X)\nu_t(X, t)},$$

and finally replacing X with x gives the required solution.

A.2 Proof of Theorem 2

Define

$$\tau_{SAPE}^\pi = \frac{1}{n} \sum_i \sum_t \pi_t(X_i) \mu_t(Z_i).$$

Using $(x + y)^2 \leq 2x^2 + 2y^2$, we have

$$\mathbb{E}[(\hat{\tau}_W^\pi - \tau^\pi)^2 \mid X_{1:n}, T_{1:n}] \leq 2\mathbb{E}[(\hat{\tau}_W^\pi - \tau_{SAPE}^\pi)^2 \mid X_{1:n}, T_{1:n}] + 2\mathbb{E}[(\tau_{SAPE}^\pi - \tau^\pi)^2 \mid X_{1:n}, T_{1:n}].$$

Noting that $\tau^\pi = \mathbb{E}[\tau_{SAPE}^\pi]$, Assumption 3 implies that $\mathbb{E}[(\tau_{SAPE}^\pi - \tau^\pi)^2] = O(1/n)$. Markov's inequality yields $\mathbb{E}[(\tau_{SAPE}^\pi - \tau^\pi)^2 \mid X_{1:n}, T_{1:n}] = O_p(1/n)$.

Let $CMSE(W, \mu) = \mathbb{E}[(\hat{\tau}_W^\pi - \tau_{SAPE}^\pi)^2 \mid X_{1:n}, T_{1:n}]$. We proceed to bound

CMSE. By iterating expectations we can obtain:

$$\begin{aligned}
CMSE(W, \mu) &= \mathbb{E}[(\hat{\tau}_W^\pi - \tau_{SAPE}^\pi)^2 \mid X, T] \\
&= \mathbb{E}[\mathbb{E}[(\hat{\tau}_W^\pi - \tau_{SAPE}^\pi)^2 \mid X, T, Z \mid X, T]] \\
&= \mathbb{E}[\mathbb{E}[\hat{\tau}_W^\pi - \tau_{SAPE}^\pi \mid X, T, Z]^2 \mid X, T] + \mathbb{E}[\mathbb{V}[\hat{\tau}_W^\pi - \tau_{SAPE}^\pi \mid X, T, Z] \mid X, T] \\
&= \mathbb{E}[(\frac{1}{n} \sum_{i,t} f_{it} \mu_t(Z_i))^2 \mid X, T] + \mathbb{E}[\mathbb{V}[\frac{1}{n} \sum_i W_i Y(T_i) \mid X, T, Z] \mid X, T] \\
&\leq \mathbb{E}[(\frac{1}{n} \sum_{i,t} f_{it} \mu_t(Z_i))^2 \mid X, T] + \frac{\sigma^2}{n^2} \|W\|_2^2
\end{aligned}$$

where σ is the bound defined in Assumption 3.

Next observe that for any (possibly correlated) random variables A_1, \dots, A_n and numbers $p_1, \dots, p_n \in \mathbb{R}^n$ such that $\sum_i p_i = 1$, we have $\mathbb{V}[\sum_i p_i A_i] \leq \max_i \mathbb{V}[A_i]$. Given this, we can simplify the first term above further, as follows:

$$\begin{aligned}
\mathbb{E}[(\frac{1}{n} \sum_{i,t} f_{it} \mu_t(Z_i))^2 \mid X, T] &= (\frac{1}{n} \sum_{i,t} f_{it} \mathbb{E}[\mu_t(Z_i) \mid X_i, T_i])^2 + \mathbb{V}[\frac{1}{n} \sum_{i,t} f_{it} \mu_t(Z_i) \mid X, T] \\
&= (\frac{1}{n} \sum_{i,t} f_{it} \nu_t(X_i, T_i))^2 + \frac{1}{n^2} \sum_i \mathbb{V}[\sum_t f_{it} \mu_t(Z_i) \mid X, T] \\
&\leq (\frac{1}{n} \sum_{i,t} f_{it} \nu_t(X_i, T_i))^2 + \frac{1}{n^2} \sum_i \max_t f_{it}^2 \mathbb{V}[\mu_t(Z_i) \mid X, T] \\
&\leq (\frac{1}{n} \sum_{i,t} f_{it} \nu_t(X_i, T_i))^2 + \frac{2\sigma^2}{n^2} \sum_i \max_t W_i^2 \delta_{T_i t} + \pi_t(X_i)^2 \\
&\leq (\frac{1}{n} \sum_{i,t} f_{it} \nu_t(X_i, T_i))^2 + \frac{2\sigma^2}{n^2} \sum_i (W_i^2 + 1) \\
&= (\frac{1}{n} \sum_{i,t} f_{it} \nu_t(X_i, T_i))^2 + \frac{2\sigma^2}{n^2} \|W\|_2^2 + \frac{2\sigma^2}{n}
\end{aligned}$$

where for the second inequality we used the fact that $(x+y)^2 \leq 2x^2 + 2y^2$. This gives us $CMSE(W, \mu) \leq J(W, \mu) + O_p(1/n)$, and combining this with the above gives $\mathbb{E}[(\hat{\tau}_W^\pi - \tau^\pi)^2 \mid X_{1:N}, T_{1:N}] \leq 2J(W, \mu) + O_p(1/n)$ as required.

A.3 Proof of Lemma 2

First, we will use the notation $f(z; x, t)$ for the conditional measure of Z given $X = x$ and $T = t$, and observe that according to Bayes rule we have:

$$\frac{f(z; x, t'')}{f(z; x, t')} = \frac{e_{t''}(z) \eta_{t'}(x)}{e_{t'}(z) \eta_{t''}(x)}$$

Define \mathbb{E}_{xt} and \mathbb{P}_{xt} as shorthand for expectation and probability given $X = x, T = t$ respectively. Then given the above, for any $M > 0$ we can bound

$$\begin{aligned}
\nu_t(x, t'') &= \frac{\eta_{t''}(x)}{\eta_{t'}(x)} \mathbb{E}[\mu_t(Z) \mid X = x, T = t''] \\
&= \int_{\mathcal{Z}} f(z; x, t'') \mu_t(z) dz \\
&= \frac{\eta_{t''}(x)}{\eta_{t'}(x)} \int_{\mathcal{Z}} \frac{e_{t''}(z)}{e_{t'}(z)} \frac{\eta_{t'}(x)}{\eta_{t''}(x)} f(z; x, t') \mu_t(z) dz \\
&= \frac{\eta_{t'}(x)}{\eta_{t''}(x)} \int_{\mathcal{Z}} \frac{e_{t''}(z)}{e_{t'}(z)} f(z; x, t') \mu_t(z) dz \\
&\leq \frac{\eta_{t'}(x)}{\eta_{t''}(x)} \left(M \mathbb{E}_{xt'}[\mathbb{1}\{\frac{e_{t''}(z)}{e_{t'}(z)} \leq M\} \mu_t(Z)] + \mathbb{E}_{xt'}[\mathbb{1}\{\frac{e_{t''}(z)}{e_{t'}(z)} > M\} \frac{e_{t''}(z)}{e_{t'}(z)} \mu_t(Z)] \right)
\end{aligned}$$

Now we can use the fact that μ_t is b -bounded to bound the first term by

$$\begin{aligned}
M \mathbb{E}_{xt'}[\mathbb{1}\{\frac{e_{t''}(z)}{e_{t'}(z)} \leq M\} \mu_t(Z)] &= M \nu_t(x, t') - M \mathbb{E}_{xt'}[\mathbb{1}\{\frac{e_{t''}(z)}{e_{t'}(z)} > M\} \mu_t(Z)] \\
&\leq M \nu_t(x, t') + Mb \mathbb{P}_{xt'}[\frac{e_{t''}(z)}{e_{t'}(z)} > M],
\end{aligned}$$

and in addition applying Cauchy Schwartz we can bound the second term by

$$\begin{aligned}
\mathbb{E}_{xt'}[\mathbb{1}\{\frac{e_{t''}(z)}{e_{t'}(z)} > M\} \frac{e_{t''}(z)}{e_{t'}(z)} \mu_t(Z)] &\leq \sqrt{\mathbb{E}_{xt'}[\mathbb{1}\{\frac{e_{t''}(z)}{e_{t'}(z)} > M\} \mu_t(Z)^2] \mathbb{E}_{xt'}[\left(\frac{e_{t''}(z)}{e_{t'}(z)}\right)^2]} \\
&\leq \sqrt{b^2 \mathbb{P}_{xt'}[\frac{e_{t''}(z)}{e_{t'}(z)} > M] \mathbb{E}_{xt'}[\left(\frac{e_{t''}(z)}{e_{t'}(z)}\right)^2]} \\
&\leq b \sqrt{\mathbb{P}_{xt'}[\frac{1}{e_{t'}(z)} > M] \mathbb{E}_{xt'}[\left(\frac{1}{e_{t'}(z)}\right)^2]}.
\end{aligned}$$

Now define $g(x, t) = \mathbb{E}_{xt}[\left(\frac{1}{e_t(z)}\right)^2]$. By Assumption 2 we know $g(x, t)$ is finite for every x and t . Also by Markov's inequality we know that $\mathbb{P}_{xt}[\frac{1}{e_t(z)} > M] \leq \frac{g(x, t)}{M^2}$. Therefore putting all of the above together we can obtain

$$\begin{aligned}
\nu_t(x, t'') &\leq \frac{\eta_{t'}(x)}{\eta_{t''}(x)} \left(M \nu_t(x, t') + \frac{2bg(x, t')}{M} \right) \\
&\leq \frac{\eta_{t'}(x)}{\eta_{t''}(x)} \left(M |\nu_t(x, t')| + \frac{2bg(x, t')}{M} \right)
\end{aligned}$$

This inequality is valid every M , so we can pick M to make it as tight as possible.

Choosing $M = \sqrt{\frac{2bg(x, t')}{\nu_t(x, t')}}$ gives us:

$$\nu_t(x, t'') \leq \frac{\eta_{t'}(x)}{\eta_{t''}(x)} \sqrt{8bg(x, t') |\nu_t(x, t')|}$$

Finally note that, since by symmetry $\mathbb{E}[\mu(Z) \mid X, T] = -\mathbb{E}[-\mu(Z) \mid X, T]$, we can strengthen this inequality to the following

$$|\nu_t(x, t'')| \leq \frac{\eta_{t'}(x)}{\eta_{t''}(x)} \sqrt{8bg(x, t')|\nu_t(x, t')|},$$

and noting that $g(x, t') = \mathbb{E}[e_t^{-2}(Z) \mid X = x, T = t']$ gives us our final result.

A.4 Proof of Lemma 3

First note that by Assumption 6 \mathcal{F} is compact. Also $J(W, \cdot)$ is continuous for every W , since by Assumption 7 we know that the norm on each \mathcal{F}_t dominates the norm on $\mathcal{L}^2(\mathcal{Z})$ and this continuity result would be trivial if $\mathcal{F}_t = \mathcal{L}^2(\mathcal{Z})$. This means that by the Extreme Value theorem we can replace the supremum over μ with a maximum over μ in the quantity we are bounding. Given this, we will proceed by bounding $\min_W \max_{\mu \in \mathcal{F}} B(W, \mu)$ using von Neumann's minimax theorem to swap the minimum and the maximum, and then use this to establish the overall bound for $J(W, \mu)$.

Next we can observe that $B(W, \mu)$ is linear, and therefore both convex and concave, for each of W and μ . Next, by Assumption 6 \mathcal{F} is convex and compact, and following the same argument as above $B(W, \cdot)$ is continuous for every W . In addition, $B(W, \mu)$ is also clearly continuous in W for fixed μ , and the set $\{W : \|W\|_2 \leq M\}$ is obviously compact and convex for any constant M . Thus by von Neumann's minimax theorem we have the following for every finite M :

$$\min_{\|W\|_2 \leq M} \max_{\mu \in \mathcal{F}} B(W, \mu) = \max_{\mu \in \mathcal{F}} \min_{\|W\|_2 \leq M} B(W, \mu) \quad (8)$$

Given this, we can bound $\min_W \max_{\mu \in \mathcal{F}} \hat{J}(W, \mu)$ as follows, which is valid for any M :

$$\begin{aligned} \min_W \max_{\mu \in \mathcal{F}} \hat{J}(W, \mu) &\leq \min_W \max_{\mu \in \mathcal{F}} B(W, \mu)^2 + \frac{1}{n^2} W^T \Sigma W \\ &\leq \min_W \max_{\mu \in \mathcal{F}} B(W, \mu)^2 + \frac{\sigma^2}{n^2} \|W\|_2^2 \\ &\leq \min_{\|W\|_2 \leq M} \max_{\mu \in \mathcal{F}} B(W, \mu)^2 + \frac{\sigma^2}{n^2} \|W\|_2^2 \\ &\leq \left(\min_{\|W\|_2 \leq M} \max_{\mu \in \mathcal{F}} B(W, \mu) \right)^2 + \frac{\sigma^2}{n^2} M^2 \\ &= \left(\max_{\mu \in \mathcal{F}} \min_{\|W\|_2 \leq M} B(W, \mu) \right)^2 + \frac{\sigma^2}{n^2} M^2 \\ &= \max_{\mu \in \mathcal{F}} \min_{\|W\|_2 \leq M} B(W, \mu)^2 + \frac{\sigma^2}{n^2} M^2 \end{aligned}$$

In these inequalities we use the fact that $\min_W \max_{\mu} B(W, \mu)^2 = (\min_W \max_{\mu} B(W, \mu))^2$ and $\max_{\mu} \min_W B(W, \mu)^2 = (\max_{\mu} \min_W B(W, \mu))^2$ due to the symmetry of μ in \mathcal{F} implied by Assumption 5.

A.5 Proof of Lemma 4

Let \prod denote Cartesian product. First we note that without loss of generality we can prove this lemma in the case that $\mathcal{F} = \prod_t \mathcal{F}_t$, since in general $F \subseteq \prod_t \mathcal{F}_t$ so $\sup_{\mu \in \mathcal{F}} \inf_W J(W, \mu) \leq \sup_{\mu \in \prod_t \mathcal{F}_t} \inf_W J(W, \mu)$, and it is easy to verify that all of our assumptions would still hold on the larger set $\prod_t \mathcal{F}_t$.

Now define the set $\mathcal{H}^0 = \{\mu \in \prod_t \mathcal{L}^2(\mathcal{Z}) : \mathbb{E}[\nu_T(X, T)^2] = 0\}$. Each coordinate \mathcal{H}_t^0 of \mathcal{H}^0 is a subspace of $\mathcal{L}^2(\mathcal{Z})$, so we can also define its orthogonal complement \mathcal{H}_t^+ . Also, we have separability since from Section 2.1 we know that $\mathcal{Z} \subseteq \mathbb{R}^q$, so any function $f \in \mathcal{L}^2(\mathcal{Z})$ can be uniquely represented as $f = f^0 + f^+$ where $f^0 \in \mathcal{H}_t^0$ and $f^+ \in \mathcal{H}_t^+$. This means that for each $\mu_t \in \mathcal{F}_t$, we can similarly uniquely represent $\mu_t = \mu_t^0 + \mu_t^+$, and we can easily extend this to a unique representation of the vector $\mu = \mu^0 + \mu^+$. Now in the case that $\mathbb{E}[\nu_T(X, T)^2] = 0$ we have $\nu_{T_i}(X_i, T_i) = 0$ almost surely for all i , and it follows from Lemma 2 that $\nu_{T_i}(X_i, t) = 0$ almost surely also for all i and t . Therefore any component of μ in \mathcal{H}^0 has no effect on the function $J(W, \mu)$ which we are bounding, so without loss of generality we can restrict our attention to the following space:

$$\mathcal{F}^+ = \prod_t (\mathcal{H}_t^+ \cap \mathcal{F}_t).$$

By construction the only function in \mathcal{F}^+ such that $\mathbb{E}[\nu_T(X, T)^2] = 0$ is the zero function, which we can also ignore in our bounds below, since when $\mu = 0$ we can easily obtain $J(W, \mu) = 0$ by choosing $W = 0$. Furthermore, by Assumption 7 we know that for each t the \mathcal{F}_t norm dominates the $\mathcal{L}^2(\mathcal{Z})$ norm, so it must be the case that each space \mathcal{F}_t^+ is closed, since \mathcal{H}_t^+ is a closed subspace of $\mathcal{L}^2(\mathcal{Z})$ due to it being an orthogonal complement. Thus it follows easily from Assumption 4 that \mathcal{F}^+ is closed, given that its norm is an \mathbb{R}^m norm on top of the corresponding \mathcal{F}_t^+ norms and m is finite.

Now, based on Lemma 3, it is sufficient to pick weights in response to μ that control for a single mean outcome function. Instead of actually constructing a particular set of weights, we take the approach of viewing this as a convex optimization problem. Specifically, given μ , we calculate the minimum euclidean norm of all weights that set the bias term $B(W, \mu)$ to zero exactly. This can be formulated as the following convex optimization program

$$\begin{aligned} \min_W \quad & \sum_i W_i^2 \\ \text{s.t.} \quad & \sum_i W_i \nu_{T_i}(X_i, T_i) = \sum_{i,t} \pi_t(X_i) \nu_t(X_i, T_i). \end{aligned}$$

Given the program only has linear constraints with equality, it satisfies Slater's condition, and therefore satisfies strong duality, which we will use to find the optimal value of this program. First we calculate the Lagrangian as

$$\mathcal{L}_n(W, \lambda) = \sum_i W_i^2 + \lambda \left(\sum_i W_i \nu_{T_i}(X_i, T_i) - \sum_{i,t} \pi_t(X_i) \nu_t(X_i, T_i) \right).$$

It can easily be verified by taking derivatives that for any $\lambda \in \mathbb{R}$ this function is minimized by setting $W_i = -\frac{\lambda}{2}\nu_{T_i}(X_i, T_i)$, and plugging this value in gives the dual formulation of the program as

$$\max_{\lambda} -\frac{\lambda^2}{4} \sum_i \nu_{T_i}(X_i, T_i)^2 - \lambda \sum_{i,t} \pi_t(X_i) \nu_t(X_i, T_i),$$

which is unconstrained. Again by taking derivatives we can maximize this function, and we find the maximum value is given by

$$\lambda = -2 \frac{\sum_i \pi_t(X_i) \nu_t(X_i, T_i)}{\sum_i \nu_{T_i}(X_i, T_i)^2},$$

and finally plugging this value into the dual objective function we see that the euclidean norm of the weights W^* solving the convex program above is given by

$$\|W^*\|_2^2 = \frac{(\sum_i \pi_t(X_i) \nu_t(X_i, T_i))^2}{\sum_i \nu_{T_i}(X_i, T_i)^2}.$$

Now define \mathbb{E}_n as the mean with respect to the empirical distribution of the logged data. Then this objective value can be reformulated as

$$\|W^*\|_2^2 = \frac{n \mathbb{E}_n[\sum_t \nu_t(X, T)]^2}{\mathbb{E}_n[\nu_T(X, T)^2]}.$$

Therefore choosing $M = \|W^*\|$, combining this result with Lemma 3 gives us

$$\begin{aligned} \min_W \sup_{\mu \in \mathcal{F}^+} J(W, \mu) &\leq \sup_{\mu \in \mathcal{F}^+} \frac{1}{n} \left(\frac{\sigma^2 \mathbb{E}_n[\sum_t \nu_t(X, T)]^2}{\mathbb{E}_n[\nu_T(X, T)^2]} \right) \\ &= \sup_{\mu \in \mathcal{F}^+} \frac{1}{n} \left(\frac{\sigma^2 \mathbb{E}_n[\sum_t \nu_t(X, T)]^2}{\mathbb{E}[\nu_T(X, T)^2] + (\mathbb{E}_n[\nu_T(X, T)^2] - \mathbb{E}[\nu_T(X, T)^2])} \right). \end{aligned}$$

Given this we will proceed by arguing that we can bound the denominator away from zero. We can note that μ appears in both the numerator and denominator on the same scale, so without loss of generality we can further restrict our attention to μ with fixed norm. By Assumption 8 we know that we can rescale every $\mu \in \mathcal{F}$ to have norm γ for some $\gamma > 0$. Given this we will restrict ourselves to the set $\mathcal{F}_\gamma^+ = \{\mu \in \mathcal{F}^+ : \|\mu\| = \gamma\}$. Since \mathcal{F}_γ^+ is the intersection of two closed sets it must be closed. Furthermore by Assumption 6 it is also compact, so it satisfies the conditions for the extreme value theorem. By construction $\mathbb{E}[\nu_T(X, T)^2] > 0$ for every $\mu \in \mathcal{F}_\gamma^+$, so putting the above together we have $\inf_{\mu \in \mathcal{F}_\gamma^+} \mathbb{E}[\nu_T(X, T)^2] > 0$. We will define this value to be α .

Now the numerator in the above bound is clearly bounded above by some $\beta > 0$ uniformly over $\mu \in \mathcal{F}$, since by Assumption 9 we know that every $\mu_t \in \mathcal{F}_t$ is uniformly bounded by some global constant, and therefore all ν terms are bounded by some constant b . Given this all that remains to be shown is that

$\sup_{\mu \in \mathcal{F}} |\mathbb{E}_n[\nu_T(X, T)^2] - \mathbb{E}[\nu_T(X, T)^2]|$ converges in probability to zero. In order to show this we will define the following terms:

$$\begin{aligned} D_n &= \mathbb{E}_n[\nu_T(X, T)^2] \\ E_n &= \sup_{\mu \in \mathcal{F}} |D_n - \mathbb{E}[D_n]| \end{aligned}$$

We need to show that E_n converges uniformly to zero. Define D'_n as an arbitrary recalculation of D_n replacing $(X_{1:n}, T_{1:n})$ with $(X'_{1:n}, T'_{1:n})$, which differ from the originals at most in a single coordinate i , and define $E'_n = \sup_{\mu \in \mathcal{F}} |D'_n - \mathbb{E}[D'_n]|$. Furthermore as argued above all ν terms are bounded above by some constant b , so each $\nu_{T_i}(X_i, T_i)^2$ is bounded by b^2 . Given this we can obtain

$$\begin{aligned} |E_n - E'_n| &= \left| \sup_{\mu \in \mathcal{F}} |D_n - \mathbb{E}[D_n]| - \sup_{\mu \in \mathcal{F}} |D'_n - \mathbb{E}[D'_n]| \right| \\ &\leq \sup_{\mu \in \mathcal{F}} |(D_n - \mathbb{E}[D_n]) - (D'_n - \mathbb{E}[D'_n])| \\ &= \sup_{\mu \in \mathcal{F}} |D_n - D'_n| \\ &= \frac{1}{n} \sup_{\mu \in \mathcal{F}} |\nu_{T_i}(X_i, T_i)^2 - \nu_{T'_i}(X'_i, T'_i)^2| \\ &\leq \frac{2b^2}{n} \end{aligned}$$

Given this we can apply McDiarmid's inequality to obtain the following bound:

$$P(|E_n - \mathbb{E}[E_n]| \leq \epsilon) \leq 2 \exp\left(-\frac{n\epsilon^2}{2b^4}\right)$$

This implies that $E_n - \mathbb{E}[E_n] = o_p(1)$. Next we show that $\mathbb{E}[E_n] = o_p(1)$ also. We do this using a symmetrization argument as follows, where D'_n is defined identically to D_n using iid shadow variables (X'_i, T'_i) in place of (X_i, T_i)

for each i , and ϵ_i are iid Rademacher random variables:

$$\begin{aligned}
\mathbb{E}[E_n] &= \mathbb{E} \left[\sup_{\mu \in \mathcal{F}} |D_n - \mathbb{E}[D_n]| \right] \\
&= \mathbb{E} \left[\sup_{\mu \in \mathcal{F}} \left| \frac{1}{n} \sum_i \nu_{T_i}(X_i, T_i) - \mathbb{E}[\nu_{T'_i}(X'_i, T'_i)] \right| \right] \\
&\leq 2 \mathbb{E} \left[\sup_{\mu \in \mathcal{F}} \left| \frac{1}{n} \sum_i \epsilon_i \nu_{T_i}(X_i, T_i) \right| \right] \\
&\leq 2 \sum_t \mathbb{E} \left[\sup_{\mu \in \mathcal{F}} \left| \frac{1}{n} \sum_i \epsilon_i \delta_{T_i t} \nu_t(X_i, T_i) \right| \right] \\
&\leq 4 \sum_t \mathbb{E} \left[\sup_{\mu \in \mathcal{F}} \left| \frac{1}{n} \sum_i \epsilon_i \nu_t(X_i, T_i) \right| \right] \\
&\leq 4 \sum_t \mathbb{E} \left[\sup_{\mu \in \mathcal{F}} \left| \frac{1}{n} \sum_i \epsilon_i \mu_t(Z_i) \right| \right] \\
&\leq 4 \sum_t \mathcal{R}_n(\mathcal{F}_t)
\end{aligned}$$

where in the third inequality we appeal to the Rademacher comparison lemma (Ledoux & Talagrand, 2013, Thm. 4.12). Thus since from Assumption 10 we know that the Rademacher complexity of each set $\mathcal{R}_n(\mathcal{F}_n)$ vanishes, it follows that $\mathbb{E}[E_n] = o_p(1)$. Putting everything from above together we get

$$\min_W \sup_{\mu \in \mathcal{F}} J(W, \mu) \leq \frac{1}{n} \frac{\beta}{\alpha + o_p(1)},$$

so we have $\min_W \sup_{\mu \in \mathcal{F}} J(W, \mu) \leq O_p(1/n)$ as required.

A.6 Proof of Theorem 4

First, Assumption 4 follows trivially from the definition of \mathcal{F}^K . Next, Assumption 5 and Assumption 8 follow from the fact that \mathcal{F}^K consists of all functions in $\text{span}(\mathcal{F}_t)$ with norm at most 1, as does the fact that it is a closed space. Given that K is a Mercer kernel, balls in the corresponding RKHS have finite covering number (Zhou, 2002), and it follows easily from this that \mathcal{F}^K has finite covering numbers, as its covering number must be bounded above by the sum of the covering numbers of the spaces \mathcal{F}_t^K . So \mathcal{F}^K is closed and totally bounded with respect to its norm, and therefore compact, which gives us Assumption 6. Clearly each \mathcal{F}_t is contained in $\mathcal{L}^2(\mathcal{Z})$ since RKHS spaces are square integrable, and the fact that the K norm dominates the \mathcal{L}^2 follows from Mercer's Theorem, which implies that $\|f\|_K^2 = \sum_{i=1}^{\infty} f_i^2 / \sigma_i$, where f_i is the i th eigenvalue of f for some orthonormal basis of $\mathcal{L}^2(\mathcal{Z})$, and $\sigma_i \geq 0$ converges to zero. This gives

us Assumption 7. Next, by construction each \mathcal{F}_t^K consists of all functions in the RKHS up to norm 1. Therefore assuming the kernel K is bounded, it is trivial to verify that function application must be globally bounded, since for any function $f \in \mathcal{F}_t^K$ we have $f(x) \leq \langle f, K_x \rangle \leq \|f\| \sqrt{K(x, x)} \leq \sqrt{K(x, x)}$, which gives us Assumption 9. Finally, given this characterization of \mathcal{F}_t^K as the 1-ball of the RKHS, it has vanishing Rademacher complexity (Mendelson, 2003, Thm. 2.1), so we have Assumption 10.

A.7 Proof of Theorem 5

First we will find a closed form expression for $\sup_{\mu \in \mathcal{F}^K} (\frac{1}{n} \sum_{i,t} f_{it} \nu_t(X_i, T_i))^2$. In this derivation we will use the shorthand φ_{ti} for the conditional density of μ_t given X_i and T_i , and T_K for the kernel integral operator defined according to $T_K f = \int_{\mathcal{Z}} K(\cdot, z) f(z) dz$. In this derivation we will make use of the fact that $\langle f, g \rangle_{\mathcal{L}^2} = \langle f, T_K g \rangle_K$ for any square integrable f and g . Given all this we can obtain:

$$\begin{aligned}
\sup_{\mu \in \mathcal{F}^K} \left(\frac{1}{n} \sum_{i,t} f_{it} \nu_t(X_i, T_i) \right)^2 &= \sum_t \sup_{\mu_t \in \mathcal{F}_t^K} \left(\frac{1}{n} \sum_i f_{it} \langle \mu_t, \varphi_i \rangle_{\mathcal{L}^2} \right)^2 \\
&= \sum_t \sup_{\mu_t \in \mathcal{F}_t^K} \left\langle \mu_t, \frac{1}{n} \sum_i f_{it} \varphi_i \right\rangle_{\mathcal{L}^2}^2 \\
&= \sum_t \sup_{\mu_t \in \mathcal{F}_t^K} \left\langle \mu_t, T_K \frac{1}{n} \sum_i f_{it} \varphi_i \right\rangle_K^2 \\
&= \sum_t \frac{\langle T_K \frac{1}{n} \sum_i f_{it} \varphi_i, T_K \frac{1}{n} \sum_i f_{it} \varphi_i \rangle_K^2}{\|T_K \frac{1}{n} \sum_i f_{it} \varphi_i\|_K^2} \\
&= \sum_t \left\langle T_K \frac{1}{n} \sum_i f_{it} \varphi_i, T_K \frac{1}{n} \sum_i f_{it} \varphi_i \right\rangle_K \\
&= \sum_t \left\langle \frac{1}{n} \sum_i f_{it} \varphi_i, T_K \frac{1}{n} \sum_i f_{it} \varphi_i \right\rangle_{\mathcal{L}^2} \\
&= \frac{1}{n^2} \sum_{i,j,t} f_{it} f_{jt} \langle \varphi_i, T_K \varphi_j \rangle_{\mathcal{L}^2} \\
&= \frac{1}{n^2} \sum_{i,j,t} f_{it} f_{jt} \int_{\mathcal{Z}} \varphi_i(z) \left(\int_{\mathcal{Z}'} K(z, z') \varphi_j(z') dz' \right) dz \\
&= \frac{1}{n^2} \sum_{i,j,t} f_{it} f_{jt} \int_{\mathcal{Z}} \int_{\mathcal{Z}'} \varphi_i(z) \varphi_j(z') K(z, z') dz' dz \\
&= \frac{1}{n^2} \sum_{i,j,t} f_{it} f_{jt} \mathbb{E}[K(Z_i, Z'_j)]
\end{aligned}$$

Given this, and recalling that $f_{it} = W_i \delta_{T_i t} - \pi_t(X_i)$ we can derive a closed

form for our minimization objective, as follows:

$$\begin{aligned}
\sup_{\mu \in \mathcal{F}^K} J(W, \mu) &= \frac{1}{n^2} \sum_{i,j,t} f_{it} f_{jt} Q_{ij} + \frac{1}{n^2} \sum_{i,j} W_i W_j \Gamma_{ij} \\
&= \frac{1}{n^2} \sum_{i,j,t} Q_{ij} (W_i W_j \delta_{T_i t} \delta_{T_j t} - 2W_j \delta_{T_j t} \pi_t(X_i) + \pi_t(X_i) \pi_t(X_j)) \\
&\quad + \frac{1}{n^2} \sum_{i,j} W_i W_j \Gamma_{ij} \\
&= \frac{1}{n^2} \sum_{i,j} W_i W_j (Q_{ij} \delta_{T_i T_j} + \Gamma_{ij}) - \frac{2}{n^2} \sum_j W_j (\sum_i Q_{ij} \pi_{T_j}(X_i)) \\
&\quad + \frac{1}{n^2} \sum_{i,j,t} Q_{ij} \pi_t(X_i) \pi_t(X_j)
\end{aligned}$$

Finally we can conclude by noting that this corresponds to the quadratic program formulation given in the question with $c = \frac{1}{n^2} \sum_{i,j,t} Q_{ij} \pi_t(X_i) \pi_t(X_j)$.

B Additional Experimentation Details

B.1 Experiment Scenario

All our experiments were conducted using the setup described in Section 5.1. We used the following parameter values for our data-generating distribution:

$$\begin{aligned}
\alpha &= [1.0, -2.0, -1.0, 2.0, 4.0, 0.0, -2.0, -1.0, -3.0, 1.0] \\
\alpha_0 &= 0 \\
\sigma_X^2 &= 4.0 \\
\beta &= [0.5, -0.5] \\
\beta_0 &= 0 \\
\zeta(0) &= 1.0 \\
\zeta(0)_0 &= 0 \\
\zeta(1) &= -0.5 \\
\zeta(1)_0 &= 0 \\
\sigma_Y^2 &= 0.01
\end{aligned}$$

In addiiton, the policy π we are evaluating takes the form as described in Section 5.1, and we used the following parameter values for this policy:

$$\begin{aligned}
\psi_0 &= [-0.1, 0.2, 0.2, -0.1, -0.1, -0.1, 0.1, 0.1, 0.1, -0.1] \\
\psi_1 &= -\psi_0
\end{aligned}$$

B.2 Method Implementation Details

In all methods where we sampled from the posterior $\varphi(\cdot; x, t)$, this sampling was done using STAN (Carpenter et al., 2017), solving QPs and LCQPs was done using the Python package `quadprog`,² all stochastic gradient descent (SGD) learning was performed using the Adam (Kingma & Ba, 2014) optimizer with a learning rate of 0.001.

OptZ We ran Algorithm 1 with $B = 50$.

IPS Since the propensity scores $\eta_t(x)$ are not tractable to compute analytically, we trained a neural network $\hat{\eta}$ to estimate this function. This was done by sampling batches of (Z, X) pairs from the data model, and training the network using SGD to predict the vector of probabilities $P_T = \beta^T Z + \beta_0$ from X , using cross-entropy loss. We used a neural network with two hidden layers of size 200 for $\hat{\eta}$, and trained for 2000 iterations with a batch size of 32. We found in practice this training was very stable and gave accurate results.

DirX For each t we trained a neural network $\hat{\rho}_t$ to predict $\nu_t(x, t)$ by taking the set of (X, T, Y) triplets in our training data where $T = t$, and training the network using SGD to predict Y from X using MSE loss. Based on pilot experiments we used a network architecture with a single hidden layer of size 100, and trained using a batch size of 128. We used 80% of our data for training, and used the remaining 20% for the purpose of early stopping. We trained for a maximum of 500 epochs, or until we made no progress on development data for 20 epochs.

DirZ For each t we trained a neural network $\hat{\mu}_t$ to predict μ_t . This was done by taking the set of (X, T, Y) triplets in our training data, and for each sampling 200 Z values from the posterior using our identified model given X and T . This gives us a set of (Z, T, Y) triplets 200 times as large as our original training set. We then trained each $\hat{\mu}_t$ network by taking the set of these triplets where $T = t$, and optimized the network using SGD on this data predicting Y from Z . We used the same settings for this optimization as with the *direct-naive* method, except we allowed up to 1000 epochs. Note that for both training and inference we limited ourselves to sampling 200 Z values per data point due to computational limitations.

C Additional Experiment Results

We present here our additional experiment results. In these results **Simplex-OptZ** refers to our method using the simplex constraints discussed in Section 4.2.

²<https://pypi.org/project/quadprog/>

n	OptZ _{0.001}	OptZ _{0.2}	OptZ _{1.0}	OptZ _{5.0}
200	.39 ± .07	.24 ± .02	.36 ± .02	.81 ± .02
500	.19 ± .02	.18 ± .02	.23 ± .02	.49 ± .02
1000	.11 ± .01	.11 ± .01	.13 ± .01	.27 ± .01
2000	.08 ± .01	.08 ± .01	.09 ± .01	.17 ± .01

Table 5: Convergence of RMSE for weighted estimator using our weights, with **step** link

n	DirX:OptZ _{0.001}	DirX:OptZ _{0.2}	DirX:OptZ _{1.0}	DirX:OptZ _{5.0}
200	.57 ± .06	.42 ± .03	.39 ± .03	.43 ± .03
500	.55 ± .02	.46 ± .02	.39 ± .02	.37 ± .02
1000	.49 ± .02	.45 ± .01	.39 ± .01	.32 ± .01
2000	.48 ± .01	.47 ± .01	.42 ± .01	.34 ± .01

Table 6: Convergence of RMSE for doubly robust estimator using our weights and **DirX**, with **step** link

n	DirZ:OptZ _{0.001}	DirZ:OptZ _{0.2}	DirZ:OptZ _{1.0}	DirZ:OptZ _{5.0}
200	.41 ± .07	.29 ± .02	.50 ± .02	1.1 ± .03
500	.20 ± .02	.21 ± .02	.31 ± .02	.70 ± .02
1000	.11 ± .01	.13 ± .01	.18 ± .01	.42 ± .01
2000	.08 ± .01	.09 ± .01	.13 ± .01	.26 ± .01

Table 7: Convergence of RMSE for doubly robust estimator using our weights and **DirZ**, with **step** link

n	SimplexOptZ _{0.001}	SimplexOptZ _{0.2}	SimplexOptZ _{1.0}	SimplexOptZ _{5.0}
200	.30 ± .02	.25 ± .02	.38 ± .02	.91 ± .02
500	.18 ± .02	.19 ± .02	.24 ± .02	.54 ± .02
1000	.12 ± .01	.11 ± .01	.13 ± .01	.29 ± .01
2000	.07 ± .01	.08 ± .01	.10 ± .01	.18 ± .01

Table 8: Convergence of RMSE for weighted estimator using our weights and constraining $W \in n\Delta^n$, with **step** link

n	IPS	OptX _{0.001}	OptX _{0.2}	OptX _{1.0}	OptX _{5.0}	DirX	DirZ
200	.47 ± .03	2.0 ± .03	2.1 ± .03	2.3 ± .02	2.5 ± .02	.52 ± .02	2.6 ± .02
500	.48 ± .03	2.0 ± .02	2.1 ± .02	2.3 ± .02	2.6 ± .02	.48 ± .02	2.6 ± .01
1000	.39 ± .02	2.0 ± .01	2.1 ± .01	2.3 ± .01	2.5 ± .01	.48 ± .02	2.6 ± .01
2000	.40 ± .01	2.0 ± .01	2.1 ± .01	2.3 ± .01	2.5 ± .01	.45 ± .02	2.6 ± .01

Table 9: Convergence of RMSE for benchmark methods, with **step** link

n	OptZ _{0.001}	OptZ _{0.2}	OptZ _{1.0}	OptZ _{5.0}
200	.03 ± .39	.11 ± .21	.29 ± .21	.78 ± .18
500	.09 ± .17	.10 ± .15	.17 ± .16	.47 ± .15
1000	.02 ± .11	.05 ± .09	.08 ± .09	.25 ± .09
2000	.03 ± .07	.05 ± .06	.07 ± .07	.16 ± .07

Table 10: Convergence of bias for weighted estimator using our weights, with **step** link

n	DirX:OptZ _{0.001}	DirX:OptZ _{0.2}	DirX:OptZ _{1.0}	DirX:OptZ _{5.0}
200	.43 ± .38	.35 ± .24	.31 ± .24	.37 ± .22
500	.51 ± .19	.42 ± .18	.35 ± .18	.33 ± .17
1000	.47 ± .13	.44 ± .11	.37 ± .10	.30 ± .11
2000	.47 ± .09	.46 ± .08	.41 ± .08	.33 ± .08

Table 11: Convergence of bias for doubly robust estimator using our weights and **DirX**, with **step** link

n	DirZ:OptZ _{0.001}	DirZ:OptZ _{0.2}	DirZ:OptZ _{1.0}	DirZ:OptZ _{5.0}
200	.05 ± .40	.19 ± .22	.45 ± .22	1.1 ± .21
500	.10 ± .18	.14 ± .16	.26 ± .16	.68 ± .17
1000	.04 ± .11	.09 ± .10	.15 ± .10	.41 ± .10
2000	.03 ± .07	.06 ± .07	.10 ± .07	.25 ± .07

Table 12: Convergence of bias for doubly robust estimator using our weights and **DirZ**, with **step** link

n	SimplexOptZ _{0.001}	SimplexOptZ _{0.2}	SimplexOptZ _{1.0}	SimplexOptZ _{5.0}
200	.04 ± .30	.12 ± .21	.31 ± .21	.89 ± .20
500	.08 ± .15	.10 ± .15	.18 ± .16	.51 ± .16
1000	.01 ± .12	.06 ± .09	.09 ± .09	.27 ± .10
2000	.03 ± .07	.05 ± .06	.07 ± .07	.17 ± .07

Table 13: Convergence of bias for weighted estimator using our weights and constraining $W \in n\Delta^n$, with **step** link

n	IPS	OptX _{0.001}	OptX _{0.2}	OptX _{1.0}	OptX _{5.0}	DirX	DirZ
200	.40 ± .25	1.9 ± .21	2.1 ± .20	2.3 ± .19	2.5 ± .18	.49 ± .18	2.6 ± .14
500	.43 ± .21	2.0 ± .16	2.1 ± .15	2.3 ± .14	2.6 ± .13	.45 ± .16	2.6 ± .12
1000	.37 ± .12	2.0 ± .10	2.1 ± .09	2.3 ± .09	2.5 ± .08	.46 ± .15	2.6 ± .11
2000	.39 ± .10	2.0 ± .08	2.1 ± .07	2.3 ± .07	2.5 ± .07	.42 ± .17	2.6 ± .11

Table 14: Convergence of bias for benchmark methods, with **step** link

n	OptZ _{0.001}	OptZ _{0.2}	OptZ _{1.0}	OptZ _{5.0}
200	.07 ± .01	.04 ± .00	.04 ± .00	.07 ± .00
500	.04 ± .00	.03 ± .00	.03 ± .00	.04 ± .00
1000	.02 ± .00	.02 ± .00	.02 ± .00	.02 ± .00
2000	.01 ± .00	.01 ± .00	.01 ± .00	.01 ± .00

Table 15: Convergence of RMSE for weighted estimator using our weights, with **exp** link

n	DirX:OptZ _{0.001}	DirX:OptZ _{0.2}	DirX:OptZ _{1.0}	DirX:OptZ _{5.0}
200	.13 ± .01	.10 ± .01	.12 ± .01	.11 ± .01
500	.10 ± .01	.09 ± .01	.10 ± .01	.12 ± .01
1000	.08 ± .00	.08 ± .00	.08 ± .00	.10 ± .00
2000	.07 ± .00	.07 ± .00	.08 ± .00	.09 ± .00

Table 16: Convergence of RMSE for doubly robust estimator using our weights and **DirX**, with **exp** link

n	DirZ:OptZ _{0.001}	DirZ:OptZ _{0.2}	DirZ:OptZ _{1.0}	DirZ:OptZ _{5.0}
200	.15 ± .02	.15 ± .01	.25 ± .01	.44 ± .01
500	.10 ± .01	.11 ± .01	.18 ± .01	.32 ± .01
1000	.07 ± .01	.08 ± .01	.12 ± .01	.23 ± .01
2000	.04 ± .00	.05 ± .00	.08 ± .00	.16 ± .00

Table 17: Convergence of RMSE for doubly robust estimator using our weights and **DirZ**, with **exp** link

n	SimplexOptZ _{0.001}	SimplexOptZ _{0.2}	SimplexOptZ _{1.0}	SimplexOptZ _{5.0}
200	.05 ± .00	.11 ± .01	.22 ± .01	.39 ± .02
500	.04 ± .00	.08 ± .01	.15 ± .01	.28 ± .01
1000	.02 ± .00	.05 ± .00	.09 ± .00	.19 ± .00
2000	.02 ± .00	.03 ± .00	.07 ± .00	.14 ± .00

Table 18: Convergence of RMSE for weighted estimator using our weights and constraining $W \in n\Delta^n$, with **exp** link

n	IPS	OptX _{0.001}	OptX _{0.2}	OptX _{1.0}	OptX _{5.0}	DirX	DirZ
200	.12 ± .01	.76 ± .02	.81 ± .02	.92 ± .02	1.0 ± .02	.10 ± .01	1.0 ± .01
500	.11 ± .00	.76 ± .01	.82 ± .01	.92 ± .01	1.0 ± .01	.10 ± .01	1.0 ± .01
1000	.10 ± .00	.74 ± .01	.79 ± .01	.90 ± .01	1.0 ± .01	.09 ± .01	1.1 ± .01
2000	.10 ± .00	.73 ± .00	.78 ± .00	.88 ± .00	.99 ± .01	.10 ± .01	1.0 ± .01

Table 19: Convergence of RMSE for benchmark methods, with **exp** link

n	OptZ _{0.001}	OptZ _{0.2}	OptZ _{1.0}	OptZ _{5.0}
200	.01 ± .06	.00 ± .04	−0.00 ± .04	−0.05 ± .04
500	.01 ± .04	.01 ± .03	.00 ± .03	−0.03 ± .03
1000	.00 ± .02	.01 ± .02	−0.00 ± .02	−0.01 ± .02
2000	.01 ± .01	.01 ± .01	.00 ± .01	.00 ± .01

Table 20: Convergence of bias for weighted estimator using our weights, with **exp** link

n	DirX:OptZ _{0.001}	DirX:OptZ _{0.2}	DirX:OptZ _{1.0}	DirX:OptZ _{5.0}
200	.07 ± .10	.08 ± .06	.10 ± .05	.11 ± .04
500	.07 ± .07	.07 ± .05	.09 ± .05	.11 ± .04
1000	.06 ± .05	.07 ± .03	.07 ± .03	.09 ± .02
2000	.06 ± .04	.06 ± .03	.07 ± .03	.09 ± .03

Table 21: Convergence of bias for doubly robust estimator using our weights and **DirX**, with **exp** link

n	DirZ:OptZ _{0.001}	DirZ:OptZ _{0.2}	DirZ:OptZ _{1.0}	DirZ:OptZ _{5.0}
200	.05 ± .14	.12 ± .10	.23 ± .09	.43 ± .09
500	.03 ± .10	.09 ± .07	.16 ± .07	.32 ± .07
1000	.02 ± .06	.06 ± .05	.10 ± .05	.23 ± .06
2000	.01 ± .04	.03 ± .03	.07 ± .03	.16 ± .04

Table 22: Convergence of bias for doubly robust estimator using our weights and **DirZ**, with **exp** link

n	SimplexOptZ _{0.001}	SimplexOptZ _{0.2}	SimplexOptZ _{1.0}	SimplexOptZ _{5.0}
200	.02 ± .05	.10 ± .06	.20 ± .08	.38 ± .11
500	.02 ± .04	.07 ± .04	.14 ± .05	.27 ± .06
1000	.01 ± .02	.04 ± .02	.08 ± .03	.19 ± .04
2000	.01 ± .01	.03 ± .02	.06 ± .02	.14 ± .02

Table 23: Convergence of bias for weighted estimator using our weights and constraining $W \in n\Delta^n$, with **exp** link

n	IPS	OptX _{0.001}	OptX _{0.2}	OptX _{1.0}	OptX _{5.0}	DirX	DirZ
200	.10 ± .06	.74 ± .13	.80 ± .13	.91 ± .15	1.0 ± .16	.07 ± .07	1.0 ± .08
500	.11 ± .03	.76 ± .07	.81 ± .08	.92 ± .09	1.0 ± .10	.07 ± .08	1.0 ± .09
1000	.10 ± .03	.74 ± .05	.79 ± .05	.90 ± .06	1.0 ± .06	.06 ± .07	1.0 ± .10
2000	.10 ± .02	.73 ± .03	.78 ± .03	.88 ± .04	.99 ± .04	.07 ± .07	1.0 ± .09

Table 24: Convergence of bias for benchmark methods, with **exp** link

n	OptZ _{0.001}	OptZ _{0.2}	OptZ _{1.0}	OptZ _{5.0}
200	.47 ± .04	.35 ± .02	.39 ± .02	.48 ± .02
500	.36 ± .05	.27 ± .02	.30 ± .02	.40 ± .02
1000	.25 ± .02	.22 ± .01	.25 ± .01	.37 ± .01
2000	.14 ± .01	.14 ± .01	.17 ± .01	.27 ± .01

Table 25: Convergence of RMSE for weighted estimator using our weights, with **cubic** link

n	DirX:OptZ _{0.001}	DirX:OptZ _{0.2}	DirX:OptZ _{1.0}	DirX:OptZ _{5.0}
200	.58 ± .07	.41 ± .03	.42 ± .02	.38 ± .02
500	.37 ± .04	.33 ± .02	.35 ± .02	.37 ± .02
1000	.31 ± .02	.31 ± .02	.33 ± .02	.39 ± .01
2000	.21 ± .02	.23 ± .02	.26 ± .01	.32 ± .01

Table 26: Convergence of RMSE for doubly robust estimator using our weights and **DirX**, with **cubic** link

n	DirZ:OptZ _{0.001}	DirZ:OptZ _{0.2}	DirZ:OptZ _{1.0}	DirZ:OptZ _{5.0}
200	.49 ± .04	.42 ± .03	.54 ± .03	.76 ± .02
500	.38 ± .05	.30 ± .02	.38 ± .02	.59 ± .02
1000	.27 ± .02	.25 ± .02	.32 ± .02	.52 ± .02
2000	.16 ± .01	.16 ± .01	.22 ± .01	.39 ± .01

Table 27: Convergence of RMSE for doubly robust estimator using our weights and **DirZ**, with **cubic** link

n	SimplexOptZ _{0.001}	SimplexOptZ _{0.2}	SimplexOptZ _{1.0}	SimplexOptZ _{5.0}
200	.45 ± .04	.41 ± .03	.52 ± .03	.70 ± .03
500	.37 ± .05	.30 ± .02	.37 ± .02	.55 ± .02
1000	.26 ± .02	.24 ± .02	.31 ± .02	.50 ± .02
2000	.14 ± .01	.15 ± .01	.20 ± .01	.35 ± .01

Table 28: Convergence of RMSE for weighted estimator using our weights and constraining $W \in n\Delta^n$, with **cubic** link

n	IPS	OptX _{0.001}	OptX _{0.2}	OptX _{1.0}	OptX _{5.0}	DirX	DirZ
200	.46 ± .04	1.1 ± .02	1.1 ± .03	1.3 ± .03	1.4 ± .03	.36 ± .02	1.4 ± .01
500	.38 ± .02	1.1 ± .01	1.2 ± .01	1.3 ± .01	1.4 ± .01	.34 ± .02	1.4 ± .01
1000	.39 ± .01	1.1 ± .01	1.2 ± .01	1.3 ± .01	1.4 ± .01	.35 ± .02	1.4 ± .01
2000	.35 ± .01	1.1 ± .01	1.2 ± .01	1.3 ± .01	1.4 ± .01	.39 ± .02	1.4 ± .01

Table 29: Convergence of RMSE for benchmark methods, with **cubic** link

n	OptZ _{0.001}	OptZ _{0.2}	OptZ _{1.0}	OptZ _{5.0}
200	.01 ± .47	.16 ± .31	.29 ± .27	.45 ± .16
500	−0.01 ± .36	.12 ± .24	.22 ± .21	.37 ± .15
1000	.03 ± .25	.12 ± .18	.20 ± .15	.35 ± .12
2000	.02 ± .14	.07 ± .12	.13 ± .11	.26 ± .08

Table 30: Convergence of bias for weighted estimator using our weights, with **cubic** link

n	DirX:OptZ _{0.001}	DirX:OptZ _{0.2}	DirX:OptZ _{1.0}	DirX:OptZ _{5.0}
200	.13 ± .57	.26 ± .32	.33 ± .27	.34 ± .17
500	.12 ± .35	.22 ± .25	.28 ± .21	.34 ± .15
1000	.18 ± .26	.24 ± .19	.29 ± .15	.37 ± .11
2000	.14 ± .16	.18 ± .14	.22 ± .12	.31 ± .09

Table 31: Convergence of bias for doubly robust estimator using our weights and **DirX**, with **cubic** link

n	DirZ:OptZ _{0.001}	DirZ:OptZ _{0.2}	DirZ:OptZ _{1.0}	DirZ:OptZ _{5.0}
200	.04 ± .49	.25 ± .34	.45 ± .29	.74 ± .20
500	.03 ± .38	.17 ± .25	.32 ± .21	.57 ± .16
1000	.04 ± .26	.16 ± .19	.28 ± .16	.50 ± .13
2000	.03 ± .16	.10 ± .13	.19 ± .12	.37 ± .10

Table 32: Convergence of bias for doubly robust estimator using our weights and **DirZ**, with **cubic** link

n	SimplexOptZ _{0.001}	SimplexOptZ _{0.2}	SimplexOptZ _{1.0}	SimplexOptZ _{5.0}
200	.02 ± .45	.22 ± .35	.40 ± .34	.65 ± .28
500	.01 ± .37	.15 ± .26	.29 ± .23	.52 ± .18
1000	.04 ± .25	.15 ± .19	.27 ± .17	.47 ± .14
2000	.02 ± .14	.08 ± .13	.17 ± .11	.34 ± .09

Table 33: Convergence of bias for weighted estimator using our weights and constraining $W \in n\Delta^n$, with **cubic** link

n	IPS	OptX _{0.001}	OptX _{0.2}	OptX _{1.0}	OptX _{5.0}	DirX	DirZ
200	.26 ± .38	1.1 ± .20	1.1 ± .20	1.2 ± .21	1.4 ± .22	.32 ± .16	1.4 ± .12
500	.31 ± .22	1.1 ± .10	1.2 ± .10	1.3 ± .10	1.4 ± .11	.29 ± .18	1.4 ± .10
1000	.37 ± .14	1.1 ± .07	1.2 ± .07	1.3 ± .08	1.4 ± .08	.32 ± .16	1.4 ± .10
2000	.34 ± .09	1.1 ± .05	1.2 ± .06	1.3 ± .06	1.4 ± .06	.34 ± .18	1.4 ± .11

Table 34: Convergence of bias for benchmark methods, with **cubic** link

n	OptZ _{0.001}	OptZ _{0.2}	OptZ _{1.0}	OptZ _{5.0}
200	.09 ± .01	.08 ± .01	.13 ± .01	.23 ± .01
500	.06 ± .01	.06 ± .00	.09 ± .00	.16 ± .00
1000	.04 ± .01	.04 ± .00	.06 ± .00	.12 ± .00
2000	.02 ± .00	.03 ± .00	.04 ± .00	.08 ± .00

Table 35: Convergence of RMSE for weighted estimator using our weights, with **linear** link

n	DirX:OptZ _{0.001}	DirX:OptZ _{0.2}	DirX:OptZ _{1.0}	DirX:OptZ _{5.0}
200	.15 ± .01	.13 ± .01	.13 ± .01	.13 ± .01
500	.14 ± .01	.13 ± .01	.13 ± .01	.13 ± .00
1000	.13 ± .01	.13 ± .00	.13 ± .00	.13 ± .00
2000	.12 ± .00	.12 ± .00	.12 ± .00	.12 ± .00

Table 36: Convergence of RMSE for doubly robust estimator using our weights and **DirX**, with **linear** link

n	DirZ:OptZ _{0.001}	DirZ:OptZ _{0.2}	DirZ:OptZ _{1.0}	DirZ:OptZ _{5.0}
200	.11 ± .01	.11 ± .01	.18 ± .01	.35 ± .01
500	.06 ± .01	.08 ± .01	.13 ± .01	.24 ± .01
1000	.05 ± .01	.06 ± .00	.09 ± .00	.18 ± .00
2000	.03 ± .00	.04 ± .00	.06 ± .00	.12 ± .00

Table 37: Convergence of RMSE for doubly robust estimator using our weights and **DirZ**, with **linear** link

n	SimplexOptZ _{0.001}	SimplexOptZ _{0.2}	SimplexOptZ _{1.0}	SimplexOptZ _{5.0}
200	.09 ± .01	.09 ± .01	.15 ± .01	.29 ± .01
500	.06 ± .01	.07 ± .01	.10 ± .01	.19 ± .01
1000	.04 ± .01	.04 ± .00	.07 ± .00	.14 ± .00
2000	.02 ± .00	.03 ± .00	.05 ± .00	.09 ± .00

Table 38: Convergence of RMSE for weighted estimator using our weights and constraining $W \in n\Delta^n$, with **linear** link

n	IPS	OptX _{0.001}	OptX _{0.2}	OptX _{1.0}	OptX _{5.0}	DirX	DirZ
200	.15 ± .01	.57 ± .01	.60 ± .01	.66 ± .01	.72 ± .01	.13 ± .00	.76 ± .00
500	.15 ± .01	.57 ± .00	.60 ± .00	.66 ± .00	.72 ± .00	.13 ± .00	.76 ± .00
1000	.14 ± .00	.57 ± .00	.60 ± .00	.66 ± .00	.72 ± .00	.13 ± .00	.76 ± .00
2000	.14 ± .00	.57 ± .00	.60 ± .00	.66 ± .00	.72 ± .00	.13 ± .00	.76 ± .00

Table 39: Convergence of RMSE for benchmark methods, with **linear** link

n	OptZ _{0.001}	OptZ _{0.2}	OptZ _{1.0}	OptZ _{5.0}
200	.03 ± .09	.06 ± .06	.11 ± .05	.23 ± .04
500	.02 ± .05	.04 ± .05	.08 ± .04	.15 ± .04
1000	.01 ± .04	.03 ± .03	.05 ± .03	.11 ± .03
2000	.01 ± .02	.02 ± .02	.04 ± .02	.08 ± .02

Table 40: Convergence of bias for weighted estimator using our weights, with **linear** link

n	DirX:OptZ _{0.001}	DirX:OptZ _{0.2}	DirX:OptZ _{1.0}	DirX:OptZ _{5.0}
200	.12 ± .09	.12 ± .06	.12 ± .05	.13 ± .04
500	.13 ± .06	.12 ± .05	.12 ± .04	.12 ± .04
1000	.12 ± .04	.12 ± .03	.12 ± .03	.12 ± .03
2000	.12 ± .03	.12 ± .03	.12 ± .02	.12 ± .02

Table 41: Convergence of bias for doubly robust estimator using our weights and **DirX**, with **linear** link

n	DirZ:OptZ _{0.001}	DirZ:OptZ _{0.2}	DirZ:OptZ _{1.0}	DirZ:OptZ _{5.0}
200	.04 ± .10	.09 ± .07	.17 ± .06	.34 ± .05
500	.02 ± .06	.06 ± .05	.12 ± .05	.24 ± .05
1000	.02 ± .04	.05 ± .04	.08 ± .04	.18 ± .04
2000	.01 ± .03	.03 ± .03	.06 ± .03	.12 ± .02

Table 42: Convergence of bias for doubly robust estimator using our weights and **DirZ**, with **linear** link

n	SimplexOptZ _{0.001}	SimplexOptZ _{0.2}	SimplexOptZ _{1.0}	SimplexOptZ _{5.0}
200	.02 ± .08	.07 ± .06	.14 ± .06	.29 ± .06
500	.02 ± .05	.05 ± .05	.09 ± .05	.19 ± .04
1000	.01 ± .04	.03 ± .03	.06 ± .03	.14 ± .03
2000	.01 ± .02	.02 ± .02	.04 ± .02	.09 ± .02

Table 43: Convergence of bias for weighted estimator using our weights and constraining $W \in n\Delta^n$, with **linear** link

n	IPS	OptX _{0.001}	OptX _{0.2}	OptX _{1.0}	OptX _{5.0}	DirX	DirZ
200	.13 ± .08	.57 ± .05	.60 ± .05	.66 ± .05	.72 ± .05	.13 ± .04	.76 ± .04
500	.14 ± .05	.57 ± .04	.60 ± .03	.66 ± .03	.72 ± .03	.12 ± .04	.76 ± .02
1000	.14 ± .04	.57 ± .02	.60 ± .02	.66 ± .02	.72 ± .02	.13 ± .03	.76 ± .03
2000	.13 ± .03	.57 ± .02	.60 ± .02	.66 ± .02	.72 ± .02	.13 ± .03	.76 ± .02

Table 44: Convergence of bias for benchmark methods, with **linear** link