

PROOFS FOR CERTAIN Π_q -CONJECTURES OF GOSPER

BING HE

ABSTRACT. In 2001 W. Gosper introduced a constant Π_q and conjectured without proofs many intriguing identities on this constant. In this paper we establish some modular equations of degrees 3 and 5. From these modular equations we confirm two groups of Π_q -identities in Gosper's list. One group involves $\Pi_q, \Pi_{q^2}, \Pi_{q^3}$ or Π_{q^6} while the other is related to $\Pi_q, \Pi_{q^2}, \Pi_{q^5}$ or $\Pi_{q^{10}}$.

1. INTRODUCTION

Throughout this paper we assume that $|q| < 1$. W. Gosper [5, p. 85] first introduced the q -constant Π_q :

$$(1.1) \quad \Pi_q = q^{1/4} \frac{(q^2; q^2)_\infty^2}{(q; q^2)_\infty^2},$$

where $(a; q)_\infty$ is defined by

$$(a; q)_\infty = \prod_{n=0}^{\infty} (1 - aq^n),$$

and then stated without proofs many identities involving Π_q [5, pp. 102–104] by employing empirical evidence based on a computer program called MACSYMA. In particular, he [5, pp. 103–104] conjectured the following interesting Π_q -identities:

$$(1.2) \quad \frac{\Pi_q^2}{\Pi_{q^2}\Pi_{q^4}} - \frac{\Pi_{q^2}^2}{\Pi_{q^4}^2} = 4,$$

$$(1.3) \quad \Pi_{q^3}^2 + 3\Pi_q\Pi_{q^9} = \sqrt{\Pi_q\Pi_{q^9}}(\Pi_q + 3\Pi_{q^9}),$$

$$(1.4) \quad \frac{\Pi_{q^2}\Pi_{q^3}^2}{\Pi_{q^6}\Pi_q^2} = \frac{\Pi_{q^2} - \Pi_{q^6}}{\Pi_{q^2} + 3\Pi_{q^6}},$$

$$(1.5) \quad \Pi_{q^2}\Pi_{q^3}^4 = \Pi_{q^6}(\Pi_{q^2} - \Pi_{q^6})^3(\Pi_{q^2} + 3\Pi_{q^6}),$$

$$(1.6) \quad \Pi_{q^6}\Pi_q^4 = \Pi_{q^2}(\Pi_{q^2} - \Pi_{q^6})(\Pi_{q^2} + 3\Pi_{q^6})^3.$$

The formula (1.2) was deduced by Gosper [5, p. 93]. The Π_q -identity (1.3) was confirmed by the author and H.-C. Zhai [6] by establishing an identity involving q -trigonometric functions and Π_q , which is equivalent to a theta function identity that can be proved by using an addition formula for Jacobi's theta functions of Liu [8, Theorem 1] (See [5] for the definitions of the q -trigonometric functions and see [7] for applications of Liu's addition formula and the definitions of Jacobi's

2000 *Mathematics Subject Classification.* 33D15, 11F03, 14H42.

Key words and phrases. Π_q -identity; modular equation.

theta functions). M. El Bachraoui [4, Theorem 2.2] just gave a partial proof of the identity (1.4). Namely, he proved that

$$\left(\frac{\Pi_{q^2} \Pi_{q^3}^2}{\Pi_{q^6} \Pi_q^2} \right)^2 = \left(\frac{\Pi_{q^2} - \Pi_{q^6}}{\Pi_{q^2} + 3\Pi_{q^6}} \right)^2.$$

In [4, Theorem 2.3] El Bachraoui only showed that (1.5) is equivalent to (1.6). See [1] and [4] for many other Π_q -identities not mentioned by Gosper [5].

In this paper we will consider the Π_q -identities (1.4), (1.5), (1.6) and many other Π_q -identities of Gosper and adopt the notations of [2, Chapters 5 and 6].

The definition of modular equations [2, (6.3.2)] is very important.

Let $0 < k, l < 1$ and let n be a positive integer. A relation between k and l induced by the formula

$$n \frac{{}_2F_1(1/2, 1/2; 1; 1 - k^2)}{{}_2F_1(1/2, 1/2; 1; k^2)} = \frac{{}_2F_1(1/2, 1/2; 1; 1 - l^2)}{{}_2F_1(1/2, 1/2; 1; l^2)}$$

is called a modular equation of degree n . Take $\alpha = k^2$, $\beta = l^2$, we say that β has degree n over α . The multiplier m is given by

$$m = \frac{z_1}{z_n},$$

where

$$z_n = \varphi^2(q^n)$$

and

$$\varphi(q) = \sum_{k=-\infty}^{\infty} q^{k^2}.$$

In Sections 2 and 3 we will consider many Π_q -identities conjectured by W. Gosper. These identities can be divided into two groups, one group of identities involving the q -constants $\Pi_q, \Pi_{q^2}, \Pi_{q^3}$ or Π_{q^6} and the other concerning $\Pi_q, \Pi_{q^2}, \Pi_{q^5}$ or $\Pi_{q^{10}}$. They are confirmed by establishing several modular equations of degrees 3 or 5.

2. IDENTITIES INVOLVING $\Pi_q, \Pi_{q^2}, \Pi_{q^3}$ OR Π_{q^6}

2.1. Statement of results. In [5, p. 103] W. Gosper conjectured the following interesting identities:

$$(2.1) \quad \sqrt{\Pi_{q^2} \Pi_{q^6}} (\Pi_q^2 - 3\Pi_{q^3}^2) = \sqrt{\Pi_q \Pi_{q^3}} (\Pi_{q^2}^2 + 3\Pi_{q^6}^2),$$

$$(2.2) \quad \Pi_{q^2}^2 (\Pi_q^4 + 18\Pi_q^2 \Pi_{q^3}^2 - 27\Pi_{q^3}^4) = \Pi_q \Pi_{q^3} (\Pi_q^4 + 16\Pi_{q^2}^4),$$

$$(2.3) \quad \Pi_{q^6}^2 (\Pi_q^4 - 6\Pi_q^2 \Pi_{q^3}^2 - 3\Pi_{q^3}^4) = \Pi_q \Pi_{q^3} (\Pi_{q^3}^4 + 16\Pi_{q^6}^4),$$

$$(2.4) \quad \Pi_q \Pi_{q^3} (\Pi_q^2 \pm 4\Pi_{q^2}^2)^2 = \Pi_{q^2}^2 (\Pi_q \mp \Pi_{q^3}) (\Pi_q \pm 3\Pi_{q^3})^3,$$

$$(2.5) \quad \Pi_q \Pi_{q^3} (\Pi_{q^3}^2 \pm 4\Pi_{q^6}^2)^2 = \Pi_{q^6}^2 (\Pi_q \mp \Pi_{q^3})^3 (\Pi_q \pm 3\Pi_{q^3}).$$

In this section we will confirm these identities by establishing the following theorem.

Theorem 2.1. *The identities (1.4)–(1.6) and (2.1)–(2.5) are true.*

Both of the identities (1.4) and (2.1) involve all of the four constants $\Pi_q, \Pi_{q^2}, \Pi_{q^3}$ and Π_{q^6} , however, each of the other identities in Theorem 2.1 contains only three of these four constants. We will show these Π_q -identities by establishing several modular equations of degree 3.

2.2. Auxiliary results. Some auxiliary results are required to prove Theorem 2.1.

Theorem 2.2. *Let β have degree 3 over α . Then*

$$(2.6) \quad m - \sqrt{\frac{\beta}{\alpha}} = 1 + \frac{3}{m} \sqrt{\frac{\beta}{\alpha}},$$

$$(2.7) \quad m^2\alpha + 3\beta = 2(\alpha\beta)^{1/8}(m^2\alpha^{1/2} - 3\beta^{1/2}),$$

$$(2.8) \quad \frac{16}{m} \sqrt{\frac{\beta}{\alpha}} = \alpha \left(1 - \frac{1}{m} \sqrt{\frac{\beta}{\alpha}} \right) \left(1 + \frac{3}{m} \sqrt{\frac{\beta}{\alpha}} \right)^3,$$

$$(2.9) \quad \frac{1}{m} \sqrt[4]{\frac{\beta}{\alpha}} (1 + \alpha) = \frac{\alpha^{1/2}}{4} \left(1 + \frac{18}{m^2} \sqrt{\frac{\beta}{\alpha}} - \frac{27}{m^4} \frac{\beta}{\alpha} \right),$$

$$(2.10) \quad m^4 \sqrt{\frac{\alpha}{\beta}} (1 + \beta) = \frac{\beta^{1/2}}{4} \left(\frac{m^4\alpha}{\beta} - 6m^2 \sqrt{\frac{\alpha}{\beta}} - 3 \right),$$

$$(2.11) \quad \frac{1}{m} \sqrt[4]{\frac{\beta}{\alpha}} \left(1 \pm \alpha^{1/2} \right)^2 = \frac{\alpha^{1/2}}{4} \left(1 \mp \frac{1}{m} \sqrt[4]{\frac{\beta}{\alpha}} \right) \left(1 \pm \frac{3}{m} \sqrt[4]{\frac{\beta}{\alpha}} \right)^3,$$

$$(2.12) \quad m^4 \sqrt{\frac{\alpha}{\beta}} (1 \pm \beta^{1/2})^2 = \frac{\beta^{1/2}}{4} \left(m^4 \sqrt{\frac{\alpha}{\beta}} \mp 1 \right)^3 \left(m^4 \sqrt{\frac{\alpha}{\beta}} \pm 3 \right).$$

Proof. The identity (2.6) follows easily from [2, (6.3.23)].

We now show (2.7). It follows from [2, (6.3.19) and (6.3.20)] that

$$\begin{aligned} & (\alpha\beta)^{1/8}(m^2\alpha^{1/2} - 3\beta^{1/2}) \\ &= \frac{(m-1)(3+m)}{8m} \left((3+m)m - 3(m-1) \right) \\ &= \frac{(m-1)(3+m)(m^2+3)}{8m} \end{aligned}$$

and

$$\begin{aligned} & m^2\alpha + 3\beta \\ &= \frac{(m-1)(3+m)}{16m} ((3+m)^2 + 3(m-1)^2) \\ &= \frac{(m-1)(3+m)(m^2+3)}{4m}. \end{aligned}$$

From these identities the formula (2.7) follows readily.

The identity (2.8) can be obtained by combining [2, (6.3.19)] and [2, (6.3.23)].

We now prove (2.9). It can be deduced from [2, (6.3.19) and (6.3.23)] that

$$\begin{aligned} \frac{1}{m} \sqrt[4]{\frac{\beta}{\alpha}} (1 + \alpha) &= \frac{1}{m} \sqrt{\frac{m(m-1)}{3+m}} \left(1 + \frac{(m-1)(3+m)^3}{16m^3} \right) \\ &= \frac{m^4 + 24m^3 + 18m^2 - 27}{16m^3} \sqrt{\frac{m-1}{m(3+m)}} \end{aligned}$$

and

$$\begin{aligned}
& \frac{\alpha^{1/2}}{4} \left(1 + \frac{18}{m^2} \sqrt{\frac{\beta}{\alpha}} - \frac{27}{m^4} \frac{\beta}{\alpha} \right) \\
&= \frac{3+m}{16m} \sqrt{\frac{(m-1)(3+m)}{m}} \left(1 + \frac{18(m-1)}{m(3+m)} - \frac{27}{m^2} \frac{(m-1)^2}{(3+m)^2} \right) \\
&= \frac{m^4 + 24m^3 + 18m^2 - 27}{16m^3} \sqrt{\frac{m-1}{m(3+m)}}.
\end{aligned}$$

From these two identities (2.9) follows quickly.

We then deduce (2.10). It follows from [2, (6.3.20) and (6.3.23)] that

$$\begin{aligned}
m^4 \sqrt[4]{\frac{\alpha}{\beta}} \left(1 + \beta \right) &= \left(\frac{m(3+m)}{m-1} \right)^{1/2} \left(1 + \frac{(m-1)^3(3+m)}{16m} \right) \\
&= \frac{m^4 - 6m^2 + 24m - 3}{16} \left(\frac{3+m}{m(m-1)} \right)^{1/2}
\end{aligned}$$

and

$$\begin{aligned}
& \frac{\beta^{1/2}}{4} \left(\frac{m^4 \alpha}{\beta} - 6m^2 \sqrt{\frac{\alpha}{\beta}} - 3 \right) \\
&= \frac{m-1}{16} \left(\frac{(m-1)(3+m)}{m} \right)^{1/2} \left(\frac{m^2(3+m)^2}{(m-1)^2} - \frac{6m(3+m)}{m-1} - 3 \right) \\
&= \frac{m^4 - 6m^2 + 24m - 3}{16} \left(\frac{3+m}{m(m-1)} \right)^{1/2},
\end{aligned}$$

from which (2.10) is obtained.

We now derive (2.11). Using [2, (6.3.19) and (6.3.23)] we get

$$\begin{aligned}
& \frac{1}{m} \sqrt[4]{\frac{\beta}{\alpha}} \left(1 \pm \alpha^{1/2} \right)^2 \\
&= \sqrt{\frac{m-1}{m(3+m)}} \left(1 \pm \frac{3+m}{4m} \sqrt{\frac{(m-1)(3+m)}{m}} \right)^2 \\
&= \frac{m^4 + 24m^3 + 18m^2 - 27}{16m^3} \sqrt{\frac{m-1}{m(3+m)}} \pm \frac{m^2 + 2m - 3}{2m^2}
\end{aligned}$$

and

$$\begin{aligned}
& \frac{\alpha^{1/2}}{4} \left(1 \mp \frac{1}{m} \sqrt[4]{\frac{\beta}{\alpha}} \right) \left(1 \pm \frac{3}{m} \sqrt[4]{\frac{\beta}{\alpha}} \right)^3 \\
&= \frac{3+m}{16m} \sqrt{\frac{(m-1)(3+m)}{m}} \left(1 \mp \sqrt{\frac{m-1}{m(3+m)}} \right) \left(1 \pm 3 \sqrt{\frac{m-1}{m(3+m)}} \right)^3 \\
&= \frac{m^4 + 24m^3 + 18m^2 - 27}{16m^3} \sqrt{\frac{m-1}{m(3+m)}} \pm \frac{m^2 + 2m - 3}{2m^2}.
\end{aligned}$$

From these two identities we obtain (2.11).

Finally, we show (2.12). We deduce from [2, (6.3.20) and (6.3.23)] that

$$\begin{aligned} m\sqrt[4]{\frac{\alpha}{\beta}}(1 \pm \beta^{1/2})^2 &= \sqrt{\frac{m(3+m)}{m-1}} \left(1 \pm \frac{m-1}{4} \sqrt{\frac{(m-1)(3+m)}{m}}\right)^2 \\ &= \frac{m^4 - 6m^2 + 24m - 3}{16m} \sqrt{\frac{m(3+m)}{m-1}} \pm \frac{m^2 + 2m - 3}{2} \end{aligned}$$

and

$$\begin{aligned} &\frac{\beta^{1/2}}{4} \left(m\sqrt[4]{\frac{\alpha}{\beta}} \mp 1\right)^3 \left(m\sqrt[4]{\frac{\alpha}{\beta}} \pm 3\right) \\ &= \frac{m-1}{16} \sqrt{\frac{(m-1)(3+m)}{m}} \left(\sqrt{\frac{m(3+m)}{m-1}} \mp 1\right)^3 \left(\sqrt{\frac{m(3+m)}{m-1}} \pm 3\right) \\ &= \frac{m^4 - 6m^2 + 24m - 3}{16m} \sqrt{\frac{m(3+m)}{m-1}} \pm \frac{m^2 + 2m - 3}{2}. \end{aligned}$$

From these we arrive at (2.12). This completes the proof of Theorem 2.2. \square

2.3. Proof of Theorem 2.1. In this subsection, we only consider modular equations of degree 3 and then we always assumed that $n = 3$ and

$$(2.13) \quad m = \frac{z_1}{z_3}.$$

We are now ready to show Theorem 2.1.

Proof of Theorem 2.1. According to [4, Theorem 2.3], we know that (1.5) is equivalent to (1.6), so we only need to prove one of them. In this section we shall show the identities (1.4), (1.6), (2.1)–(2.5). If the identities in Theorem 2.1 hold for $0 < q < 1$, then, by analytic continuation, these identities are also true for $|q| < 1$. So we can assume that $0 < q < 1$.

Let

$$\psi(q) = \sum_{n=0}^{\infty} q^{n(n+1)/2}.$$

It follows from (1.1) and [2, (1.3.14)] that

$$(2.14) \quad \Pi_q = q^{1/4} \psi^2(q).$$

Then the identities (1.4), (1.6), (2.1)–(2.5) are respectively equivalent to

$$(2.15) \quad \frac{\psi^2(q^2)\psi^4(q^3)}{\psi^2(q^6)\psi^4(q)} = \frac{\psi^2(q^2) - q\psi^2(q^6)}{\psi^2(q^2) + 3q\psi^2(q^6)},$$

$$(2.16) \quad \frac{\psi^2(q^6)}{\psi^2(q^2)} \frac{\psi^8(q)}{\psi^8(q^2)} = \left(1 - q \frac{\psi^2(q^6)}{\psi^2(q^2)}\right) \left(1 + 3q \frac{\psi^2(q^6)}{\psi^2(q^2)}\right)^3,$$

$$(2.17) \quad \psi(q^2)\psi(q^6)(\psi^4(q) - 3q\psi^4(q^3)) = \psi(q)\psi(q^3)(\psi^4(q^2) + 3q^2\psi^4(q^6)),$$

$$(2.18) \quad \frac{\psi^2(q^3)}{\psi^2(q)} \left(1 + 16q \frac{\psi^8(q^2)}{\psi^8(q)}\right) = \frac{\psi^4(q^2)}{\psi^4(q)} \left(1 + 18q \frac{\psi^4(q^3)}{\psi^4(q)} - 27q^2 \frac{\psi^8(q^3)}{\psi^8(q)}\right),$$

$$(2.19) \quad \frac{\psi^2(q)}{\psi^2(q^3)} \left(1 + 16q^3 \frac{\psi^8(q^6)}{\psi^8(q^3)}\right) = \frac{\psi^4(q^6)}{\psi^4(q^3)} \left(\frac{\psi^8(q)}{\psi^8(q^3)} - 6q \frac{\psi^4(q)}{\psi^4(q^3)} - 3q^2\right),$$

$$(2.20) \quad \frac{\psi^2(q^3)}{\psi^2(q)} \left(1 \pm 4q^{1/2} \frac{\psi^4(q^2)}{\psi^4(q)}\right)^2 = \frac{\psi^4(q^2)}{\psi^2(q)} \left(1 \mp q^{1/2} \frac{\psi^2(q^3)}{\psi^2(q)}\right) \left(1 \pm 3q^{1/2} \frac{\psi^2(q^3)}{\psi^2(q)}\right)^3,$$

$$(2.21) \quad \frac{\psi^2(q)}{\psi^2(q^3)} \left(1 \pm 4q^{3/2} \frac{\psi^4(q^6)}{\psi^4(q^3)}\right)^2 = \frac{\psi^4(q^6)}{\psi^4(q^3)} \left(\frac{\psi^2(q)}{\psi^2(q^3)} \mp q^{1/2}\right)^3 \left(\frac{\psi^2(q)}{\psi^2(q^3)} \pm 3q^{1/2}\right).$$

We first prove (2.15). Let β have degree 3 over α . Then, by [2, Theorem 5.4.2, (i) and (iii)],

$$(2.22) \quad \psi(q) = \sqrt{\frac{z_1}{2}} (\alpha/q)^{1/8},$$

$$(2.23) \quad \psi(q^2) = \frac{1}{2} \sqrt{z_1} (\alpha/q)^{1/4},$$

$$(2.24) \quad \psi(q^3) = \sqrt{\frac{z_3}{2}} (\beta/q^3)^{1/8},$$

$$(2.25) \quad \psi(q^6) = \frac{1}{2} \sqrt{z_3} (\beta/q^3)^{1/4},$$

and so

$$(2.26) \quad \frac{\psi(q^3)}{\psi(q)} = q^{-1/4} \sqrt{\frac{z_3}{z_1}} (\beta/\alpha)^{1/8},$$

$$(2.27) \quad \frac{\psi(q^6)}{\psi(q^2)} = q^{-1/2} \sqrt{\frac{z_3}{z_1}} (\beta/\alpha)^{1/4}.$$

Hence, the formula (2.15) follows by dividing both sides of (2.6) by $m(1 + \frac{3}{m} \sqrt{\frac{\beta}{\alpha}})$ and then using (2.26), (2.27) and (2.13) in the resulting identity.

We now prove (2.16). It follows from (2.22) and (2.23) that

$$(2.28) \quad \frac{\psi(q)}{\psi(q^2)} = \frac{\sqrt{2}}{(\alpha/q)^{1/8}}.$$

Then (2.16) can be obtained by dividing both sides of (2.8) by α and then employing (2.27), (2.28) and (2.13) in the resulting identity.

The identity (2.17) follows easily by multiplying both sides of (2.7) by $\frac{(\alpha\beta)^{1/8}\sqrt{z_1z_3/q^3z_3^2}}{32}$ and then using (2.13) in the resulting equation.

The formula (2.18) can be deduced by dividing both sides of (2.9) by \sqrt{q} and then using (2.26) and (2.13).

We then show (2.19). It follows from (2.28) that

$$\frac{\psi(q^3)}{\psi(q^6)} = \frac{\sqrt{2}}{(\beta/q^3)^{1/8}}$$

and so

$$(2.29) \quad \frac{\psi(q^6)}{\psi(q^3)} = \frac{(\beta/q^3)^{1/8}}{\sqrt{2}}.$$

We multiply both sides of (2.10) by $q^{1/2}$ and then apply (2.26), (2.29) and (2.13) in the resulting identity to obtain (2.19).

The identity (2.20) can be derived by dividing both sides of (2.11) by $q^{1/2}$ and then using (2.26), (2.28) and (2.13) in the resulting formula.

The identity (2.21) follows readily by multiplying both sides of (2.12) by $q^{1/2}$ and then employing (2.26), (2.29) and (2.13) in the resulting identity. This finishes the proof of Theorem 2.1. \square

3. IDENTITIES INVOLVING $\Pi_q, \Pi_{q^2}, \Pi_{q^5}$ OR $\Pi_{q^{10}}$

3.1. Statement of results. Gosper [5, pp. 103–104] conjectured the following Π_q -identities:

$$(3.1) \quad \Pi_{q^2}\Pi_{q^5}(16\Pi_{q^{10}}^4 - \Pi_{q^5}^4) = \Pi_{q^{10}}^3(5\Pi_{q^{10}} - \Pi_{q^2})(\Pi_{q^2} - \Pi_{q^{10}})^5,$$

$$(3.2) \quad \Pi_{q^{10}}\Pi_q^4(16\Pi_{q^2}^4 - \Pi_q^4) = \Pi_{q^2}^3(5\Pi_{q^{10}} - \Pi_{q^2})^5(\Pi_{q^2} - \Pi_{q^{10}}),$$

$$(3.3) \quad \Pi_q\Pi_{q^5}(16\Pi_{q^2}^4 - \Pi_q^4)^2 = \Pi_{q^2}^4(5\Pi_{q^5} - \Pi_q)^5(\Pi_{q^5} - \Pi_q),$$

$$(3.4) \quad \Pi_q\Pi_{q^5}(16\Pi_{q^{10}}^4 - \Pi_{q^5}^4)^2 = \Pi_{q^{10}}^4(5\Pi_{q^5} - \Pi_q)(\Pi_{q^5} - \Pi_q)^5,$$

$$(3.5) \quad (\Pi_q\Pi_{q^{10}} - \Pi_{q^2}\Pi_{q^5})^2 = \Pi_{q^2}\Pi_{q^{10}}(\Pi_{q^5} - \Pi_q)(5\Pi_{q^5} - \Pi_q).$$

In this section we will confirm these results.

Theorem 3.1. *The identities (3.1)–(3.5) are true.*

The identities (3.1)–(3.4) only contain three of the constants $\Pi_q, \Pi_{q^2}, \Pi_{q^5}$ and $\Pi_{q^{10}}$, but the formula (3.5) includes all of these four constants. These five identities have similar styles so that our proofs share the same pattern. We will show these identities by setting up some modular equations of degree 5.

3.2. One lemma. The value of the multiplier m depends on n , but throughout this subsection and the next subsection we only consider modular equations of degree 5, then it is always assumed that $n = 5$ and

$$(3.6) \quad m = \frac{z_1}{z_5}.$$

In order to prove Theorem 3.1 we need several auxiliary results.

Theorem 3.2. *If β has degree 5 over α , then*

(3.7)

$$256 \frac{z_1}{z_5} \left(\frac{\alpha}{\beta} \right)^{1/2} \frac{1}{\beta} \left(1 - \frac{1}{\beta} \right) = \left(5 - \frac{z_1}{z_5} \left(\frac{\alpha}{\beta} \right)^{1/2} \right) \left(\frac{z_1}{z_5} \left(\frac{\alpha}{\beta} \right)^{1/2} - 1 \right)^5,$$

(3.8)

$$256 \frac{z_5}{z_1} \left(\frac{\beta}{\alpha} \right)^{1/2} \frac{1}{\alpha} \left(1 - \frac{1}{\alpha} \right) = \left(5 \frac{z_5}{z_1} \left(\frac{\beta}{\alpha} \right)^{1/2} - 1 \right)^5 \left(1 - \frac{z_5}{z_1} \left(\frac{\beta}{\alpha} \right)^{1/2} \right),$$

$$(3.9) \quad \frac{z_5}{z_1} \left(\frac{\beta}{\alpha} \right)^{1/4} \left(\alpha - 1 \right)^2 = \frac{\alpha}{16} \left(5 \frac{z_5}{z_1} \left(\frac{\beta}{\alpha} \right)^{1/4} - 1 \right)^5 \left(\frac{z_5}{z_1} \left(\frac{\beta}{\alpha} \right)^{1/4} - 1 \right),$$

$$(3.10) \quad \frac{z_1}{z_5} \left(\frac{\alpha}{\beta} \right)^{1/4} \left(\beta - 1 \right)^2 = \frac{\beta}{16} \left(5 - \frac{z_1}{z_5} \left(\frac{\alpha}{\beta} \right)^{1/4} \right) \left(1 - \frac{z_1}{z_5} \left(\frac{\alpha}{\beta} \right)^{1/4} \right)^5,$$

(3.11)

$$\left(\frac{z_1}{z_5} \left(\frac{\alpha}{\beta} \right)^{1/4} - \frac{z_1}{z_5} \left(\frac{\alpha}{\beta} \right)^{1/2} \right)^2 = \frac{z_1}{z_5} \left(\frac{\alpha}{\beta} \right)^{1/2} \left(1 - \frac{z_1}{z_5} \left(\frac{\alpha}{\beta} \right)^{1/4} \right) \left(5 - \frac{z_1}{z_5} \left(\frac{\alpha}{\beta} \right)^{1/4} \right).$$

Proof. We first prove (3.7) and (3.10). According to [3, Chapter 19, (13.12)–(13.15)] we have

$$(3.12) \quad \left(\frac{\alpha}{\beta} \right)^{1/4} = \frac{2m + \rho}{m(m-1)},$$

$$(3.13) \quad \left(\frac{\beta}{\alpha} \right)^{1/4} = \frac{2m - \rho}{5 - m},$$

$$\left(\frac{1 - \beta}{1 - \alpha} \right)^{1/4} = \frac{2m + \rho}{5 - m},$$

$$(3.14) \quad (\alpha\beta)^{1/2} = \frac{4m^3 - 16m^2 + 20m + \rho(m^2 - 5)}{16m^2},$$

$$(3.15) \quad \{(1 - \alpha)(1 - \beta)\}^{1/2} = \frac{4m^3 - 16m^2 + 20m - \rho(m^2 - 5)}{16m^2},$$

where

$$\rho = (m^3 - 2m^2 + 5m)^{1/2}.$$

Then

$$(3.16) \quad \beta = \left(\frac{2m - \rho}{5 - m} \right)^2 \frac{4m^3 - 16m^2 + 20m + \rho(m^2 - 5)}{16m^2},$$

$$(3.17) \quad 1 - \beta = \left(\frac{2m + \rho}{5 - m} \right)^2 \frac{4m^3 - 16m^2 + 20m - \rho(m^2 - 5)}{16m^2}.$$

Substituting (3.6), (3.12), (3.16) and (3.17) into both sides of each of the identities (3.7) and (3.10), noticing that $\rho = (m^3 - 2m^2 + 5m)^{1/2}$ and then simplifying we find that both sides of each of the identities (3.7) and (3.10) are respectively equal to

$$\left(\frac{2}{m-1} \right)^2 A(m)$$

and

$$\frac{m-5}{256m(m-1)} B(m),$$

where

$$A(m) = 4m^9 - 24m^8 + m^8\rho + 64m^7 - 2m^7\rho - 200m^6 + 6m^6\rho - 40m^5 - 98m^5\rho \\ - 40m^4 + 80m^4\rho - 1280m^3 - 470m^3\rho - 504m^2 - 470m^2\rho - 28m - 70m\rho - \rho$$

and

$$B(m) = 2m^6 - 10m^5 + m^5\rho - 5m^4\rho + 4m^4 + 10m^3\rho \\ - 4m^3 - 102m^2 - 42m^2\rho - 18m - 27m\rho - \rho.$$

These prove (3.7) and (3.10).

We now show (3.8) and (3.9). According to [3, Chapter 19, (13.12)] we get

$$(3.18) \quad \left(\frac{1-\alpha}{1-\beta} \right)^{1/4} = \frac{2m-\rho}{m(m-1)}.$$

It follows from (3.12), (3.14), (3.15) and (3.18) that

$$(3.19) \quad \alpha = \left(\frac{2m+\rho}{m(m-1)} \right)^2 \frac{4m^3 - 16m^2 + 20m + \rho(m^2 - 5)}{16m^2},$$

$$(3.20) \quad 1 - \alpha = \left(\frac{2m-\rho}{m(m-1)} \right)^2 \frac{4m^3 - 16m^2 + 20m - \rho(m^2 - 5)}{16m^2}.$$

We substitute (3.6), (3.13), (3.19) and (3.20) into both sides of each of the identities (3.8) and (3.9), note that $\rho = (m^3 - 2m^2 + 5m)^{1/2}$ and then simplify to deduce that both sides of each of the identities (3.8) and (3.9) equal

$$-\frac{2^{12}m^2}{(m-5)^{12}}C(m)$$

and

$$\frac{1-m}{256m^6(m-5)}D(m)$$

respectively, where

$$C(m) = 28m^9 + 2520m^8 - m^8\rho + 32000m^7 - 350m^7\rho + 5000m^6 - 11750m^6\rho \\ + 25000m^5 - 58750m^5\rho + 625000m^4 + 50000m^4\rho - 1000000m^3 - 306250m^3\rho \\ + 1875000m^2 + 93750m^2\rho - 1562500m - 156250m\rho + 390625\rho,$$

and

$$D(m) = 18m^6 + 510m^5 - m^5\rho + 100m^4 - 135m^4\rho - 1050m^3\rho \\ - 500m^3 + 1250m^2\rho + 6250m^2 - 6250m - 3125m\rho + 3125\rho,$$

which prove (3.8) and (3.9).

We finally prove (3.11). We substitute (3.6) and (3.12) into both sides of (3.11) and then simplify using the identity $\rho = (m^3 - 2m^2 + 5m)^{1/2}$ to derive that both sides of (3.11) are equal to

$$\frac{(m-5)^2(m^3 - m^2 + 7m + 2m\rho + 1 + 2\rho)}{(m-1)^4},$$

from which (3.11) follows readily. This concludes the proof of Theorem 3.2. \square

3.3. Proof of Theorem 3.1. In this subsection we will prove Theorem 3.1.

Proof of Theorem 3.1. Using the equation (2.14) we see that the identities (3.1)–(3.5) are respectively equivalent to

(3.21)

$$\frac{\psi^2(q^2)}{\psi^2(q^{10})} \frac{\psi^8(q^5)}{\psi^8(q^{10})} \left(16q^5 - \frac{\psi^8(q^5)}{\psi^8(q^{10})} \right) = \left(5q^2 - \frac{\psi^2(q^2)}{\psi^2(q^{10})} \right) \left(\frac{\psi^2(q^2)}{\psi^2(q^{10})} - q^2 \right)^5,$$

(3.22)

$$\frac{\psi^2(q^{10})}{\psi^2(q^2)} \frac{\psi^8(q)}{\psi^8(q^2)} \left(16q - \frac{\psi^8(q)}{\psi^8(q^2)} \right) = \left(5q^2 \frac{\psi^2(q^{10})}{\psi^2(q^2)} - 1 \right)^5 \left(1 - q^2 \frac{\psi^2(q^{10})}{\psi^2(q^2)} \right),$$

$$(3.23) \quad \frac{\psi^2(q^5)}{\psi^2(q)} \left(16q \frac{\psi^8(q^2)}{\psi^8(q)} - 1 \right)^2 = \frac{\psi^8(q^2)}{\psi^8(q)} \left(5q \frac{\psi^2(q^5)}{\psi^2(q)} - 1 \right)^5 \left(q \frac{\psi^2(q^5)}{\psi^2(q)} - 1 \right),$$

$$(3.24) \quad \frac{\psi^2(q)}{\psi^2(q^5)} \left(16q^5 \frac{\psi^8(q^{10})}{\psi^8(q^5)} - 1 \right)^2 = \frac{\psi^8(q^{10})}{\psi^8(q^5)} \left(5q - \frac{\psi^2(q)}{\psi^2(q^5)} \right) \left(q - \frac{\psi^2(q)}{\psi^2(q^5)} \right)^5,$$

$$(3.25) \quad \left(q \frac{\psi^2(q)}{\psi^2(q^5)} - \frac{\psi^2(q^2)}{\psi^2(q^{10})} \right)^2 = \frac{\psi^2(q^2)}{\psi^2(q^{10})} \left(q - \frac{\psi^2(q)}{\psi^2(q^5)} \right) \left(5q - \frac{\psi^2(q)}{\psi^2(q^5)} \right).$$

We temporarily assume that $0 < q < 1$. Let β have 5 degree over α . According to [2, Theorem 5.4.2 (i) and (iii)] we have

$$(3.26) \quad \psi(q) = \sqrt{\frac{z_1}{2}} (\alpha/q)^{1/8},$$

$$(3.27) \quad \psi(q^2) = \frac{1}{2} \sqrt{z_1} (\alpha/q)^{1/4},$$

$$(3.28) \quad \psi(q^5) = \sqrt{\frac{z_5}{2}} (\beta/q^5)^{1/8},$$

$$(3.29) \quad \psi(q^{10}) = \frac{1}{2} \sqrt{z_5} (\beta/q^5)^{1/4}.$$

It follows from (3.27), (3.28) and (3.29) that

$$(3.30) \quad \frac{\psi(q^2)}{\psi(q^{10})} = \sqrt{\frac{z_1}{z_5}} \left(\frac{\alpha}{\beta} \right)^{1/4} q,$$

$$(3.31) \quad \frac{\psi(q^5)}{\psi(q^{10})} = \frac{\sqrt{2}}{(\beta/q^5)^{1/8}}.$$

Multiplying both sides of (3.7) by q^{12} and then using (3.30) and (3.31) in the resulting equation we can easily obtain the identity (3.21).

It is easily deduced from (3.26) and (3.27) that

$$(3.32) \quad \frac{\psi(q)}{\psi(q^2)} = \frac{\sqrt{2}}{(\alpha/q)^{1/8}}.$$

Then (3.22) follows by substituting (3.30) and (3.32) into (3.8).

It is easily seen from (3.26) and (3.28) that

$$(3.33) \quad \frac{\psi(q^5)}{\psi(q)} = \sqrt{\frac{z_5}{z_1}} \left(\frac{\beta}{\alpha} \right)^{1/8} / q^{1/2}.$$

Then (3.23) follows easily by dividing both sides of (3.9) by q and then using (3.32) and (3.33) in the resulting identity.

Multiplying both sides of (3.10) by q and then employing (3.31) and (3.33) in the resulting equation we can attain (3.24).

The identity (3.25) follows readily by multiplying both sides of (3.11) by q^4 and then using (3.30) and (3.33) in the resulting identity.

From these we see that (3.21)–(3.25) holds for $0 < q < 1$. By analytic continuation, these identities are also true for $|q| < 1$. This completes the proof of Theorem 3.1. \square

ACKNOWLEDGEMENT

This work was partially supported by the National Natural Science Foundation of China (Grant No. 11801451) and the Natural Science Foundation of Hunan Province (Grant No. 2020JJ5682).

REFERENCES

- [1] S. Abo Touk, Z. Al Houchan and M. El Bachraoui, Proofs for two q -trigonometric identities of Gosper. *J. Math. Anal. Appl.* 456(1)(2017), 662–670.
- [2] B.C. Berndt, *Number Theory in the Spirit of Ramanujan*, American Mathematical Society, Providence, RI, 2006.
- [3] B.C. Berndt, *Ramanujan's Notebooks, Part III*, Springer-Verlag, New York, 1991.
- [4] M. El Bachraoui, On the Gosper's q -constant Π_q . *Acta Mathematica Sinica, English Series.* 34(11)(2018), 1755–1764
- [5] R.W. Gosper, Experiments and discoveries in q -trigonometry, in: F.G. Garvan, M.E.H. Ismail (Eds.), *Symbolic Computation, Number Theory, Special Functions, Physics and Combinatorics*, Kluwer, Dordrecht, Netherlands, 2001, pp.79–105.
- [6] B. He and H.-C. Zhai, Proofs for certain q -trigonometric identities of Gosper, *Science China–Mathematics*, 2020, <http://engine.scichina.com/doi/10.1007/s11425-019-9555-1>.
- [7] Z.-G. Liu, Addition formulas for Jacobi theta functions, Dedekind's eta functions, and Ramanujan's congruences, *Pacific J. Math.* 240(1)(2009), 135–150.
- [8] Z.-G. Liu, An addition formula for the Jacobian theta function and its applications, *Adv. Math.* 212(1)(2007), 389–406.

SCHOOL OF MATHEMATICS AND STATISTICS, CENTRAL SOUTH UNIVERSITY, CHANGSHA 410083, HUNAN, PEOPLE'S REPUBLIC OF CHINA
Email address: yuhe001@foxmail.com; yuhelingyun@foxmail.com