

Generalized KS transformations, ND singular oscillator and generalized MICZ-Kepler system

A. Lavrenov

Belarusian State Pedagogical University, Minsk, Belarus, 18, Sovetskaya St., Minsk, Belarus, 220050

E-mail: lavrenov@bspu.by

May 2019

Abstract. The description of number of dual (quasy)-exactly solvable models with its hidden symmetry algebra has been given at different levels of analysis within the framework of generalized Kustaanheimo-Stiefel (KS)-transformations. It's shown that N -dimensional singular oscillator and $(n+1)$ -dimensional generalized MICZ-Kepler system are dual to each other and the duality transformation is the generalized version of the KS transformation. The solvability of the Schrödinger equation of these problems by the variables separation method is given in a double, spherical and parabolic coordinates. The quadratic Hahn algebra $QH(3)$ as a hidden symmetry remains unchanged with different decomposition of the original real space \mathbb{R}^N into components in the framework of addition rule for $SU(1,1)$ algebra. Also the hidden symmetry algebra as Higgs/Hahn algebras is clearly shown by the commutant approach in the sense of Howe duality. A dimensional reduction is carried out to a singular oscillator of two dimensions where every variable is the n -dimensional hyper-radius r . The dual connection with $N \equiv 2n$ -dimensional singular oscillator and the $(n+1)$ general MICZ-Kepler system in the class of quasi-exact (QE) problems also are considered. The certain generalization both of harmonic oscillator model by anisotropic and nonlinear inharmonic terms and its dual analog is shown and analyzed in the framework of generalized KS transformations. The exact analytical solutions of the Schrödinger equation for abovementioned problems for QE class are discussed and given for four series of dual quasi-exact solvable models. In particular, a comparison with similar results in lower dimensions and its generalization are given.

1. Introduction

There is the Kustaanheimo and Stiefel transformation (KS transformation) [1] that establishes a connection between the problem of an isotropic harmonic oscillator (HO) in a real four-dimensional space and the Kepler problem in ordinary three-dimensional space (see [2]-[4]). On the other hand, it's possible to establish the connection between the Coulomb problem and the harmonic oscillator in the dimensions of its spaces, realizing the Hopf fibrations through the generalized KS transformations. The Hopf maps are fibrations of spheres over spheres with the fiber-sphere [5]. The term "Hopf fibration

or map" is usually restricted to the fibrations between spheres $S^{2n-1}/S^{n-1} = S^n$. Among them, four Hopf maps stand out: $S^{2n-1}/S^{n-1} = S^n$, ($n = 1, 2, 4, 8$):

- $S^1 \rightarrow S^1$ with fiber S^0 , (zero Hopf map $n = 1$),
- $S^3 \rightarrow S^2$ with fiber S^1 (first Hopf map $n = 2$),
- $S^7 \rightarrow S^4$ with fiber S^3 (second Hopf map $n = 4$)
- $S^{15} \rightarrow S^8$ with fiber S^7 (third Hopf map $n = 8$)

Each such Hopf map is closely related with its division algebra: $n = 1$ reflects the existence of division algebra of real numbers, $n = 2$ - complex numbers, $n = 4$ quaternions and $n = 8$ - octonions. Zero Hopf map is related with anyons (or magnetic vortices) [6]. The Dirac and Yang monopoles are related with the first and the second ones respectively. The first three cases have been studied in detail for a long time. There are a lot of literature that is devoted to this topic, which has already developed academic logic and style of presentation of the materials for this topic (for example [7] - [22]). The authors [23] - [28] have thoroughly worked out many questions of the above last case of the Hopf bundle associated with 16-dimensional harmonic oscillator and 9-dimensional Coulomb problems recently. All these connections between the Coulomb problem and the harmonic oscillator are very helpful for solutions of many problems of atomic physics (see [29]-[31]).

However, in his [32], Cordani, using the methods of group theory, asserted that one can construct the required KS transformations for the Kepler problem with dimension $n > 6$, but didn't answer the question "how these transformations can be realized?". The given theory was built in the article [33]. The authors of this article gave a simple graphical algorithm for constructing the generalized KS transformations that realizes the connection between two problems above and proved the following proposition.

Proposition. *A connection between the Kepler problem in a real space of dimension $n + 1$ and the problem of an isotropic harmonic oscillator in a real space of dimension N exists and can be established by means of generalized KS transformations in the cases when $N = 2n$ and $n = 2^h$ ($h = 0, 1, 2, \dots$).*

The question naturally arises - is it possible to obtain results realized in small dimensions for the general case?

From our point of view, the answer is YES. Therefore, in the framework of generalized KS transformations that is described above, we aim to describe a series of dual exactly solvable models with their hidden symmetry algebra at different levels of analysis. The article is organized as follows. In Section II, we give a brief results of the works by the authors of [33], recall the addition rule for $SU(1, 1)$ algebra, the problem of hidden symmetry for the systems admitting such an approach in accordance to Ref. [34] and the well-known results on Heisenberg-Weyl, $SU(1, 1)$, $U(16)$, Higgs and Hahn algebras.

Our main result will be the object of Sections III - V where the dual connection with the general MICZ-Kepler system and N -dimensional singular oscillator will be

made; the exact analytical solutions of the Schrödinger equation for these problems will be successfully built in a double, spherical and parabolic coordinates; the hidden symmetry algebra as Higgs/Hahn algebras will be clearly shown by both addition rule for $SU(1, 1)$ algebra and the commutant approach in the sense of Howe duality. In Section VI, a dimensional reduction is carried out to a singular oscillator of two dimensions, each variable of which is the n -dimensional hyper-radius r . The dual connection with $N \equiv 2n$ -dimensional singular oscillator and the $(n + 1)$ general MICZ-Kepler system in the class of quasi-exact problems are considered in the last sections VII.

2. Generalized KS transformations, the famous algebras and the addition rule for $SU(1, 1)$ algebra

The problem of an isotropic harmonic oscillator (IHO) in a real N -dimensional space is described by the next Schrödinger equation:

$$(H_0 - 4Z) \psi(u) \equiv \left(-\frac{1}{2} \frac{\partial^2}{\partial u_s \partial u_s} + \frac{1}{2} \omega^2 u_s u_s - 4Z \right) \psi(u) = 0, \quad (1)$$

Here, $u_s (s = \overline{1, N})$ are the Cartesian coordinates of the point u in N -dimensional space; summation over repeated indices is understood; w is a real positive number; Z is the eigenvalue. A generalized KS transformation is determined by

$$\begin{aligned} x_\lambda &= (\Gamma_\lambda)_{st} u_s u_t, \quad (\overline{\lambda} = \overline{1, n+1}), \\ \chi_a &= \chi_a(u) \quad (a = \overline{n+2, N \equiv 2n}), \end{aligned} \quad (2)$$

where the real symmetric $N \times N$ matrices Γ_λ satisfy

$$\begin{aligned} Sp(\Gamma_\lambda) &= 0, \quad \Gamma_\lambda \Gamma_\mu + \Gamma_\mu \Gamma_\lambda = 2\delta_{\lambda\mu} I; \\ x_\lambda x_\lambda &= (u_s u_s)^2 = r^2 \quad (r = u_s u_s); \end{aligned} \quad (3)$$

I is the unit $N \times N$ matrix and $\chi_a(u)$ are any functions with nonzero Jacobian of the transformation χ from the variables u to the variables x . In according to [33] among the solutions of Eq. (1) it is possible to find functions Ψ that are dependent on the variables x only and satisfied the equation of the Kepler problem in a space of dimension $n + 1$.

$$(H'_0 - E) \Psi(r) \equiv \left(-\frac{1}{2} \frac{\partial^2}{\partial x_\lambda \partial x_\lambda} - \frac{Z}{r} + \frac{1}{8} \omega^2 \right) \Psi(r) = 0, \quad (4)$$

On the other hand, with the help of the pair of creation (birth) $a_s^\dagger = \sqrt{\frac{\omega}{2}} \left(-\frac{1}{\omega} \frac{\partial}{\partial u_s} + u_s \right)$ and annihilation (destruction) operators $a_s = \sqrt{\frac{\omega}{2}} \left(\frac{1}{\omega} \frac{\partial}{\partial u_s} + u_s \right)$, the number operators $N_s = a_s^\dagger a_s \equiv \left[-\frac{1}{2\omega} \frac{\partial^2}{\partial u_s^2} + \frac{\omega}{2} u_s^2 - \frac{1}{2} \right]$ the Hamiltonian of the IHO can be rewritten in the operator form:

$$H_0 = \omega \left(\sum_{s=1}^{2n} N_s + n \right) \quad (5)$$

Recall the definition of the Heisenberg-Weyl algebra $W(n)$ and following commutation relations between its operators:

$$[a_i; a_j^\dagger] = \delta_{ij}, \quad [N_i, a_j] = -a_i \delta_{ij}, \quad [N_i, a_j^\dagger] = a_i^\dagger \delta_{ij}, \quad i, j = \overline{1, n}$$

The Lie algebra $U(n)$ with generators E_{ij} , $i, j = \overline{1, n}$ admits the following realization in $W(n)$:

$$E_{ij} = a_i^\dagger a_j, \quad i, j = \overline{1, n}.$$

Hamiltonian H_0 commutes with both E_{ij} (i.e. $[H_0, a_i^\dagger a_j] = 0$) and with generators of infinitesimal rotations

$$L_{jk} = \frac{i}{2}(a_j a_k^\dagger - a_j^\dagger a_k) = -\frac{i}{2} \left(u_j \frac{\partial}{\partial u_k} - u_k \frac{\partial}{\partial u_j} \right)$$

which have well known commutation relations

$$[L_{jk}, L_{\ell m}] = \frac{i}{2}(L_{j\ell}\delta_{km} - L_{k\ell}\delta_{jm} + L_{km}\delta_{j\ell} - L_{jm}\delta_{k\ell}), \quad j, k, \ell, m = \overline{1, n}.$$

The metaplectic representation of $SU(1, 1)$ is defined by the following map in $W(1)$:

$$J_0^{(i)} = \frac{1}{2} \left(N_i + \frac{1}{2} \right), \quad J_+^{(i)} = \frac{1}{2} a_i^{\dagger 2}, \quad J_-^{(i)} = \frac{1}{2} a_i^2, \quad (6)$$

where the generators J_0, J_\pm obey the commutation relations

$$[J_0, J_\pm] = \pm J_\pm, \quad [J_+, J_-] = -2J_0.$$

Its Casimir element is given by

$$Q = J_0^2 - J_+ J_- - J_0.$$

Consider now the addition of two irreducible representations of $SU(1, 1)$ for which the initial Casimir operators take values $Q^{(i)} = \lambda_i(\lambda_i - 1)$, $i = \overline{1, 2}$, and let us write

$$J_0^{(12)} = J_0^{(1)} + J_0^{(2)}, \quad J_\pm^{(12)} = J_\pm^{(1)} + J_\pm^{(2)} \quad (7)$$

with the superindex denoting on which of the two factors in $SU(1, 1)^{\otimes 2}$ the operator is acting. It is obvious that Hamiltonian $H = J_0^{(12)}$ has dynamical symmetry as $SU(1, 1) \oplus SU(1, 1)$ and it's spectrum $\epsilon = n_1 + n_2 + \lambda_1 + \lambda_2$ is degenerated. This means that there are the integrals of motion commuting with H . A trivial example of such integrals for the possible combinations from initial operators $(J_0^{(1)}, J_\pm^{(1)}, J_0^{(2)}, J_\pm^{(2)})$ is as follows:

$$\begin{aligned} K_1 &= J_0^{(1)} - J_0^{(2)} \\ K_2 &= Q^{(12)} = Q^{(1)} + Q^{(2)} + 2J_0^{(1)}J_0^{(2)} - J_+^{(1)}J_-^{(2)} - J_-^{(1)}J_+^{(2)} \end{aligned} \quad (8)$$

The hidden symmetry of our Hamiltonian $H = J_0^{(12)}$ means the existence of some algebra constructed from the integrals K_1, K_2 . According to the work [34] such algebra is quadratic Hahn algebra $QH(3)$ under the commutation relations:

$$\begin{aligned} [K_1, K_2] &= K_3; \\ [K_2, K_3] &= -2(K_1K_2 + K_2K_1) + \delta_1; \\ [K_3, K_1] &= -2K_1^2 - 4K_2 + \delta_2 \end{aligned} \quad (9)$$

Recall that the Higgs algebra can be considered as a polynomial deformation of $SU(2)$ with three generators D, A_\pm satisfying the following commutation relations [35] - [36]:

$$[D, A_\pm] = \pm 4A_\pm; \quad [A_+, A_-] = -D^3 + \alpha_1 D + \alpha_2 \quad (10)$$

with α_1, α_2 central elements. According to [37], the isomorphism of the discrete version of the Hahn algebra and the Higgs algebra is readily seen by taking

$$\begin{aligned} K_1 &= \frac{1}{2}D \\ K_2 &= -\frac{1}{4}\left(A_+ + A_- + \frac{1}{2}D^2\right) + \frac{\alpha_1}{8} \end{aligned} \quad (11)$$

and observing that the commutation relations (9) then follow from (10) with

$$\begin{aligned} \delta_1 &\equiv -\frac{\alpha_2}{4} \equiv 4\left(J_0^{(1)} + J_0^{(2)}\right)\left(Q^{(1)} - Q^{(2)}\right) \\ \delta_2 &\equiv \frac{\alpha_1}{2} \equiv 2\left(J_0^{(1)} + J_0^{(2)}\right)^2 + 4\left(Q^{(1)} + Q^{(2)}\right). \end{aligned} \quad (12)$$

3. Dual connection between the general MICZ-Kepler system and N -dimensional singular oscillator

In this section, we propose to discuss the following questions: How does the structure of matrices for the generalized version of the KS transformation impose restrictions on the splitting of the initial problem for the IHO into its components? Which version of the generalization for the IHO components choose? What will be its dual partner for the chosen option of generalization? Are the proposed dual models exactly solvable?

So, a generalized KS transformation is based on the matrices Γ_λ . The graphical algorithm for their construction, proposed in [33], is characterized by repetition of the structure of old matrices of dimension k in the structure of a new matrix of double dimension $2k$ on the it's main diagonal. In particular, the choice in [33] was such for the matrix Γ_{n+1}

$$\Gamma_{n+1} = \begin{pmatrix} I & 0 \\ 0 & -I \end{pmatrix}$$

where 0 and I are the null and unit $n \times n$ matrices, i.e.

$$\begin{aligned} x_{n+1} &= (\Gamma_{n+1})_{st} u_s u_t \\ &= u_1^2 + u_2^2 + \dots + u_n^2 - u_{n+1}^2 - u_{n+2}^2 - \dots - u_{2n}^2 \end{aligned} \quad (13)$$

Then the remaining matrices $\Gamma_\lambda, (\lambda = \overline{1, n})$ necessarily have the structure

$$\Gamma_\lambda = \begin{pmatrix} 0 & \gamma_\lambda \\ \gamma_\lambda & 0 \end{pmatrix}$$

where γ_λ are anticommuting real $n \times n$ matrices each of whose square is equal to the unit matrix.

Facts that have been described above and the results obtained in small dimensions clearly indicate the need to consider the original problem of an IHO in a real $N \equiv 2n$ -dimensional space as the sum of the two ones of dimension n :

$$H_0 \psi(\mathbf{u}, \mathbf{v}) = \left[-\frac{1}{2} \left(\frac{\partial^2}{\partial u_i \partial u_i} + \frac{\partial^2}{\partial v_i \partial v_i} \right) + \frac{\omega^2}{2} (u_i u_i + v_i v_i) \right] \psi = 4Z\psi, \quad (14)$$

Here ω, Z have positive real values and are correspondingly the frequency and energy of the IHO, $v_i \equiv u_{n+i}$, ($i = \overline{1, n}$).

To preserve symmetry between two IHO of dimension n and also taking into account the results obtained in small dimensions, we choose the same perturbation for both of them, which will be additive and the simplest. Therefore we proposed next simple generalization of our initial problem to so-called a double singular oscillator:

$$H \psi(\mathbf{u}, \mathbf{v}) = \left[H_0 + \frac{c_1}{u_1^2 + \dots + u_n^2} + \frac{c_2}{v_1^2 + \dots + v_n^2} \right] \psi = Z\psi, \quad (15)$$

where H_0 is Hamiltonian of an IHO determined earlier in the explicit formula for (1).

Thus, there is a transparent transition for any generalizations with additive terms to the original Hamiltonians. For example, it takes place

$$\begin{aligned} x_{n+1} &= u_1^2 + u_2^2 + \dots + u_n^2 - v_1^2 - v_2^2 - \dots - v_n^2 \\ &\equiv u_i u_i - v_i v_i; \\ r &= \sqrt{x_\lambda x_\lambda} = u_i u_i + v_i v_i; \end{aligned} \quad (16)$$

or

$$\begin{aligned} \frac{c_1}{u_1^2 + \dots + u_n^2} + \frac{c_2}{v_1^2 + \dots + v_n^2} &= \frac{2c_1}{2u_i u_i} + \frac{2c_2}{2v_i v_i} = \\ &= \frac{2c_1}{r + x_{n+1}} + \frac{2c_2}{r - x_{n+1}} = \frac{4\lambda_1}{r + x_{n+1}} + \frac{4\lambda_2}{r - x_{n+1}} \end{aligned}$$

This fact means that the dual analogue of so-called a double singular oscillator is the generalized MICZ-Kepler system with non central terms:

$$H' \psi(\mathbf{r}) = \left[H'_0 + \frac{\lambda_1}{r(r + x_{n+1})} + \frac{\lambda_2}{r(r - x_{n+1})} \right] \psi = E\psi, \quad (17)$$

where $2\lambda_a = c_a$, $a = 1, 2$ are nonnegative constants; H'_0 is Hamiltonian of the Kepler problem determined earlier (4).

Their dual nature underlines the fact that the roles of $E = -\frac{\omega^2}{8}$ and Z are interchanged. The variables E and Z become a negative number that denotes the energy of bound states and a parameter defining the ‘charge’ value in the Coulomb potential respectively.

So, on the one hand, our proposed generalization in according to the above formulas is reduced to additive terms in the Hamiltonians, and on the other hand we get a rich symmetry picture of new problem, which leads to a series of exactly-solvable cases.

3.1. Variables separation in double coordinates

The Schrödinger equation for a $N \equiv 2n$ -dimensional double singular oscillator (15) represents the sum of two singular oscillators of dimensions $D = n$:

$$\begin{aligned} H &= H_1 + H_2 = \sum_{1 \leq a \leq 2} H_a = \\ &= \sum_{1 \leq a \leq 2} \left[-\frac{1}{2} \frac{\partial^2}{\partial x_i^a \partial x_i^a} + \frac{\omega^2}{2} (x_i^a x_i^a) + \frac{c_a}{x_i^a x_i^a} \right], \end{aligned} \quad (18)$$

where $x_i^1 = u_i$, $x_i^2 = v_i$.

In each nD real space \mathbb{R}^n where x_1, x_2, \dots, x_n are Cartesian coordinates, we introduce it's hyperspherical coordinates: $(\phi_1, \dots, \phi_{n-1})$ are the hyperspherical angles and r is the hyperradius by

$$\begin{aligned} x_n &= r \cos(\phi_{n-1}), \\ x_{n-1} &= r \sin(\phi_{n-1}) \cos(\phi_{n-2}), \\ &\dots \\ &\dots \\ x_2 &= r \sin(\phi_{n-1}) \sin(\phi_{n-2}) \cdots \sin(\phi_2) \cos(\phi_1), \\ x_1 &= r \sin(\phi_{n-1}) \sin(\phi_{n-2}) \cdots \sin(\phi_2) \sin(\phi_1), \end{aligned} \quad (19)$$

Using the ansatz $\Psi(\mathbf{r}) = \Psi(r, \phi_1, \dots, \phi_{n-1}) = R(r)\Omega(\phi)$ the Schrödinger equation for each H_a can be rewritten in terms of it's separated equations of hyperradius r and angular variables ϕ as follows:

$$\left[\frac{1}{r^{n-1}} \frac{\partial}{\partial r} (r^{n-1} \frac{\partial}{\partial r}) + (2E_a - \omega^2 r^2) - \frac{\Lambda^2 + 2c_a}{r^2} \right] R(r) = 0, \quad (20)$$

$$[\Lambda^2 - L(L + n - 2)] \Omega(\phi) = 0, \quad (21)$$

where E_a are the eigenvalues of H_a , but $E_1 + E_2 = 4Z$; $L(L + n - 2)$ is the separation constant and is also an eigenvalue of the operator Λ^2 (21).

For convenience, we set a new variable $x = \omega r^2$ and function $R(x) = x^a e^{-x/2} g(x)$ and substitute them into the Eq. (20). We obtain a confluent hypergeometric equation for $g(x)$, which will give a solution of Eq. (20) in terms of a special function

$$R_{NL}(r) = C_{NL} r^{L'} e^{-\frac{\omega r^2}{2}} {}_1F_1(-N; L' + \frac{n}{2}; \omega r^2) \quad (22)$$

where $L'(L' + n - 2) = L(L + n - 2) + 2c_a$.

Hence we have the discrete energy E_a of H_a from (22) as

$$E_a = 2\omega \left(N_a + \frac{L'_a}{2} + \frac{n}{4} \right), \quad (23)$$

Thus, the energy spectrum of the our N -dimensional double singular oscillators

$$4Z = E_1 + E_2 = 2\omega \left(N_1 + N_2 + \frac{L'_1 + L'_2}{2} + \frac{n}{2} \right), \quad (24)$$

3.2. Variables separation in spherical coordinates

The Schrödinger equation for the generalized MICZ-Kepler system with non central terms has the next form (17):

$$\begin{aligned} H' \Psi(\mathbf{r}) &= \left[H'_0 + \frac{\lambda_1}{r(r + x_{n+1})} + \frac{\lambda_2}{r(r - x_{n+1})} \right] \Psi = \\ &\equiv \left(-\frac{1}{2} \frac{\partial^2}{\partial x_\lambda \partial x_\lambda} - \frac{Z}{r} + \frac{\lambda_1}{r(r + x_{n+1})} + \frac{\lambda_2}{r(r - x_{n+1})} \right) \Psi = \\ &\equiv \left(-\frac{1}{2} \Delta - \frac{Z}{r} + \frac{\lambda_1}{r(r + x_{n+1})} + \frac{\lambda_2}{r(r - x_{n+1})} \right) \Psi = E\psi, \end{aligned} \quad (25)$$

In the $(n + 1)$ -dimensional spherical coordinates

$$\begin{aligned}
 x_{n+1} &= r \cos(\theta), \\
 x_n &= r \sin(\theta) \cos(\phi_{n-2}), \\
 &\dots \\
 &\dots \\
 x_2 &= r \sin(\theta) \sin(\phi_{n-2}) \cdots \sin(\phi_1) \cos(\phi_0), \\
 x_1 &= r \sin(\theta) \sin(\phi_{n-2}) \cdots \sin(\phi_1) \sin(\phi_0),
 \end{aligned} \tag{26}$$

using the ansatz $\Psi(\mathbf{u}, \mathbf{v}) = \Psi(r, \theta, \phi_{n-2}, \dots, \phi_0, \phi'_{n-2}, \dots, \phi'_0) = R(r)\Theta(\theta)\Phi(\phi, \phi')$. the Schrödinger equation of our generalized MICZ-Kepler system with non central terms rewritten in terms of separated equations of variables r, θ and angular variables ϕ, ϕ'

$$\left[\frac{1}{r^n} \frac{\partial}{\partial r} \left(r^n \frac{\partial}{\partial r} \right) + 2 \left(\frac{Z}{r} + E \right) - \frac{\Lambda}{r^2} \right] R(r) = 0, \tag{27}$$

$$\left[\frac{1}{\sin^{n-1} \theta} \frac{\partial}{\partial \theta} \left(\sin^{n-1} \theta \frac{\partial}{\partial \theta} \right) - \frac{\hat{L}^2 + 4\lambda_2}{4 \sin^2 \frac{\theta}{2}} - \frac{\hat{L}^2 + 4\lambda_1}{4 \cos^2 \frac{\theta}{2}} + \Lambda \right] \Theta(\theta) = 0, \tag{28}$$

$$\left[\hat{L}^2 - L(L + n - 2) \right] \Phi(\phi, \phi') = 0, \tag{29}$$

where $\Lambda = \lambda(\lambda + n - 1)$ is the separation constant and is also an eigenvalue of the operator (28). Solutions of (27) - (28) are as follows

$$\begin{aligned}
 R(r) &= C_{k\lambda} r^\lambda e^{-\sqrt{-2E}r} {}_1F_1(-k + \lambda - Q/2, n + 2\lambda, 2\sqrt{-2E}r), \\
 \Theta(\theta) &= C_{\lambda J L} (1 + \cos \theta)^{J'/2} (1 - \cos \theta)^{L'/2} P_{\lambda - (J' + L')/2}^{(J' + \frac{n-2}{2}, L' + \frac{n-2}{2})}(\cos \theta),
 \end{aligned}$$

where

$$\begin{aligned}
 J'(J' + n - 2) &= L(L + n - 2) + 4\lambda_1, \\
 L'(L' + n - 2) &= L(L + n - 2) + 4\lambda_2.
 \end{aligned}$$

Hence, the energy spectrum

$$E = -\frac{Z^2}{2(k + \frac{n+Q}{2})^2} = -\frac{Z^2}{2(n_r + n_\theta + \frac{n+J'+L'}{2})^2}. \tag{30}$$

coincides with [27] in appearance only. For full compliance, it is necessary to take into account in the definition of the principal quantum number $k = n_r + n_\theta + (J' + L' - Q)/2$ that $J', L' \rightarrow L$ when $\lambda_1 = \lambda_2 = 0$.

3.3. Variables separation in parabolic coordinates

The Schrödinger equation $H'\Psi = E\Psi$ (25) is same, but the Laplace-Beltrami operator Δ in the hyperparabolic coordinates has the next form:

$$\Delta = \frac{4}{u+v} \left\{ \frac{1}{u^{\frac{n-2}{2}}} \frac{\partial}{\partial u} \left(u^{\frac{n}{2}} \frac{\partial}{\partial u} \right) + \frac{1}{v^{\frac{n-2}{2}}} \frac{\partial}{\partial v} \left(v^{\frac{n}{2}} \frac{\partial}{\partial v} \right) \right\} - \frac{L^2}{uv}$$

The parabolic coordinates on the S_{n-1} sphere are defined according to works [27] - [28] by

$$\begin{aligned} x_{n+1} &= \frac{u-v}{2}, \\ x_n &= \sqrt{uv} \cos(\phi_{n-2}), \\ &\dots \\ &\dots \\ x_2 &= \sqrt{uv} \sin(\phi_{n-2}) \cdots \cos(\phi_0), \\ x_1 &= \sqrt{uv} \sin(\phi_{n-2}) \cdots \sin(\phi_0), \\ r &= \frac{u+v}{2}, \end{aligned} \tag{31}$$

where the $(n+1)$ x_λ are Cartesian coordinates in the hyperparabolic coordinates, $\{\phi_0, \dots, \phi_{n-2}\}$ are the hyperparabolic angles and the parabolic coordinates u, v range from 0 to ∞ . In order to separate variables, the wave function is now choiced as follows $\Psi(\mathbf{u}, \mathbf{v}) = \Psi(u, v, \phi_{n-2}, \dots, \phi_0, \phi'_{n-2}, \dots, \phi'_0) = U(u)V(v)\Phi(\phi, \phi')$.

Thus, the Schrödinger equation of our generalized MICZ-Kepler system with non central terms in the hyperparabolic coordinates can be rewritten in terms of separated equations of variables u, v and angular variables ϕ, ϕ'

$$\left[\frac{1}{u^{\frac{n-2}{2}}} \frac{\partial}{\partial u} \left(u^{\frac{n}{2}} \frac{\partial}{\partial u} \right) - \frac{L(L+n-2) + 4\lambda_1}{u} + \frac{Z + Eu}{2} - P \right] U(u) = 0, \tag{32}$$

$$\left[\frac{1}{v^{\frac{n-2}{2}}} \frac{\partial}{\partial v} \left(v^{\frac{n}{2}} \frac{\partial}{\partial v} \right) - \frac{L(L+n-2) + 4\lambda_2}{v} + \frac{Z + Ev}{2} + P \right] V(v) = 0, \tag{33}$$

$$\left[\hat{L}^2 - L(L+n-2) \right] \Phi(\phi, \phi') = 0, \tag{34}$$

where P is the separation constant.

Solutions of (32) and (33) are given by the confluent hypergeometric polynomials,

$$\begin{aligned} U(u) &= C_{n_1} J u^{J'/2} e^{-\frac{\sqrt{-2Eu}}{2}} {}_1F_1(-n_1, \frac{n}{2} + J', \sqrt{-2Eu}), \\ V(v) &= C_{n_2} L v^{L'/2} e^{-\frac{\sqrt{-2Ev}}{2}} {}_1F_1(-n_2, \frac{n}{2} + L', \sqrt{-2Ev}), \end{aligned} \tag{35}$$

where

$$\begin{aligned} J'(J' + n - 2) &= L(L + n - 2) + 4\lambda_1; & -n_1 &= \frac{J'}{2} + \frac{n}{4} + \frac{P - Z/2}{\sqrt{-2E}}; \\ L'(L' + n - 2) &= L(L + n - 2) + 4\lambda_2; & -n_2 &= \frac{L'}{2} + \frac{n}{4} + \frac{-P - Z/2}{\sqrt{-2E}}. \end{aligned}$$

Set

$$n_1 + n_2 = \frac{Z}{\sqrt{-2E}} - \frac{J' + L' + n}{2}.$$

The energy spectrum is as follows

$$E = -\frac{Z^2}{2\{n_1 + n_2 + \frac{n+J'+L'}{2}\}^2}. \tag{36}$$

Making the identification $k = n_1 + n_2 + \frac{J'+L'-Q}{2}$, the energy spectrum becomes (30).

4. Hidden symmetry algebra and overlap coefficients

4.1. ($N \equiv 2n$)-dimensional oscillator

In according to the previous section 2, in order to have the quadratic Hahn algebra $QH(3)$ as a hidden symmetry algebra, two conditions must be satisfied. First of all, the resulting Hamiltonian must be the sum of the two original ones. Secondly, each of the starting Hamiltonians must have a symmetry $SU(1, 1)$.

In subsection 3.1 we have shown next two facts as detailed as possible:

- the ($N \equiv 2n$)-dimensional model of harmonic isotropic oscillator may be considered as the sum of the two independent ones of dimension n ;

$$\begin{aligned}\hat{H}_0\psi(\mathbf{u}, \mathbf{v}) &= Z\psi \equiv (H_1 + H_2)\psi = (Z_1 + Z_2)\psi \\ &\equiv \left[-\frac{1}{2} \left(\frac{\partial^2}{\partial u_i \partial u_i} + \frac{\partial^2}{\partial v_i \partial v_i} \right) + V_{sho}(u_i u_i) + V_{sho}(v_i v_i) \right] \psi = \\ &\equiv \sum_{1 \leq a \leq 2} H_a \psi = \sum_{1 \leq a \leq 2} \left[-\frac{1}{2} \frac{\partial^2}{\partial x_i \partial x_i} + V_{sho}(x_i x_i) \right] \psi\end{aligned}\quad (37)$$

where u_i, v_i ($i = 1, \dots, n$) are Cartesian coordinates of the space \mathbb{R}^n ; Z_a are the eigenvalues of H_a , but $Z_1 + Z_2 = Z$ and the potential of a singular oscillator V_{sho} is

$$V_{sho}(x_i x_i) = V_{ho}(x_i x_i) + \frac{c_a}{x_i x_i} = \frac{\omega^2 x_i x_i}{2} + \frac{c_a}{x_i x_i}, \quad (38)$$

- the nD model of singular oscillator is an exactly solvable model in the hyperspherical coordinates (19) with the following solution of Eq. (22).

Fixing quantum numbers (for the angular part in our case) allows us to reduce the problem to one-dimensional. This in turn means that the symmetry of the problem will now be determined by the first equation (20), which is the radial part of the n -dimensional non-relativistic Hamiltonian harmonic oscillator H_{ho} . Carrying out the replacement $R(r) = r^{\frac{1-n}{2}} u(r)$ and with some reassignment of the constants we have a convenient form of this equation to compare with the results of [34], namely:

$$H_{ho} = \left[-\frac{1}{4\omega} \frac{\partial^2}{\partial r^2} + \frac{\omega}{4} r^2 + \frac{g}{2\omega r^2} \right] \quad (39)$$

where $g = \frac{4L(L+n-2)+8c_a+(n-3)(n-1)}{8}$.

The following realization of the $SU(1, 1)$ operators solves our problem:

$$\begin{aligned}J_0 &\equiv H_{ho}; & J_2 &= \frac{i}{2} \left[r \frac{\partial}{\partial r} + \frac{1}{2} \right]; \\ J_1 &= \frac{\omega r^2}{2} - J_0; & J_{\pm} &= J_1 \pm iJ_2;\end{aligned}\quad (40)$$

Thus, the described realization allows to represent the original Hamiltonian as a sum of two items, each of them has the desired symmetry algebra $SU(1, 1)$ in a radial variable. The Casimir operator is $Q = \frac{8g-3}{16}$, and the angular part is fixed by its well-known quantum numbers.

It should be noted that in this analyzed case, the division of the original real space \mathbb{R}^N space into 2 equal nD parts, as well as the reduction to one-dimensional consideration is not significant. To show this fact, we will give one more realization of the $SU(1, 1)$ operators for any value of the dimension of the space $\mathbb{R}^{n'}$ (that is, the repeated index m is summed up from 1 to n'):

$$\begin{aligned} J_0 &= \left[-\frac{1}{4\omega} \frac{\partial^2}{\partial x_m \partial x_m} + \frac{\omega}{4} x_m x_m + \frac{g}{2\omega x_m x_m} \right]; \quad J_1 = \frac{\omega}{2} x_m x_m - J_0; \\ J_2 &= \frac{i}{2} \left[x_m \frac{\partial}{\partial x_m} + \frac{n'}{2} \right]; \quad J_{\pm} = J_1 \pm iJ_2; \quad Q = \frac{n(n-4) + 8g}{16} - \hat{L}^2. \end{aligned} \quad (41)$$

This result is interesting in several aspects. It turns out that the quadratic Hahn algebra $QH(3)$ as a hidden symmetry remains unchanged with different decomposition of the original real space \mathbb{R}^N into components and also when it's generalized to the model of a singular oscillator.

4.2. $(n+1)D$ related (MICZ-)Kepler-like systems

In according to (16), dual analog of so-called a double singular oscillator is the generalized MICZ-Kepler system with non central terms. In the hyperparabolic coordinates (31) it can be rewritten in terms of separated equations of variables u, v (32) - (33) and angular variables ϕ, ϕ' (34). It's solutions for (32) and (33) are given by the confluent hypergeometric polynomials (35) in subsection 3.3.

In other words, we have a similar situation as with a HO. The original Hamiltonian can be represent as a sum of two almost identical item in the parabolic coordinates u, v respectively (Eqs. (32) - (33)), i.e. if we fix the quantum numbers of the angular part we can reduce the problem to one-dimensional $u(v)$. The latter and will determine the symmetry of the problem now. The following realization of the $SU(1, 1)$ operators solves our problem in this case:

$$\begin{aligned} J_0 &= \frac{1}{2\gamma} \left[\frac{1}{x^{\frac{n}{2}-1}} \frac{\partial}{\partial x} (x^{\frac{n}{2}} \frac{\partial}{\partial x}) - \frac{c_a}{x} + \frac{E}{2} x \right]; \quad J_1 = \gamma x - J_0; \\ J_2 &= i \left[x \frac{\partial}{\partial x} + \frac{n}{4} \right]; \quad J_{\pm} = J_1 \pm iJ_2; \quad Q = \frac{n(n-4)}{16} + c_a. \end{aligned} \quad (42)$$

where are $c_a = L(L + n - 2) + 4\lambda_a$; $E = -2\gamma^2$.

Thus, the described realization allows to represent the original Hamiltonian of $(n+1)D$ related MICZ-Kepler systems as a sum of two items, each of them has the desired symmetry algebra $SU(1, 1)$. The angular part is fixed by its well-known quantum numbers.

4.3. The overlap coefficients

Above we have shown that the dual models to the generalized version of the KS-transformation models ($N \equiv 2n$)-dimensional singular oscillator and the $(n+1)$ -dimensional generalized MICZ-Kepler system) have the exact analytical solutions in

according to hidden symmetry as the quadratic Hahn algebra $QH(3)$. The solvability of the Schrödinger equation of these models by the variables separation method in spherical and parabolic (cylindrical) coordinates connects with $SU(1, 1) \oplus SU(1, 1)$ dynamical symmetry. So, in according to the work [34] the overlap coefficients between wavefunctions in these coordinates are Clebsch-Gordan coefficients (CGC) for $SU(1, 1)$ algebra. Thus, on the one hand, the diagonalization of the operator K_1 corresponds to choosing the unconnected basis $|n_1, \lambda_1\rangle \otimes |n_2, \lambda_2\rangle$ in the space of the direct sum $SU(1, 1) \oplus SU(1, 1)$ and, on the other hand, to separation of the variables in parabolic (cylindrical) coordinates. Similarly, on the one hand, the diagonalization of the operator K_2 corresponds to choosing the connected basis $|n_{12}, \lambda_{12}\rangle$ in the space of the direct sum $SU(1, 1) \oplus SU(1, 1)$ and, on the other hand, to separation of the variables in spherical coordinates. The overlaps between eigenstates of the operations K_1 and K_2 (or the wavefunctions in these coordinate systems) can be written in terms of CGC for $SU(1, 1)$. An explicit expression for CGC in terms of Hahn polynomials can be found in the work [34]:

$$C_{n, \lambda_1; N-n, \lambda_2}^{j, \lambda_{12}} = h_n w_p {}_3F_2 \left[\begin{matrix} -n, -p, 2\lambda_1 + \lambda_2 + p - 1 \\ -N, 2\lambda_1 \end{matrix} ; 1 \right] \quad (43)$$

5. The Higgs algebra as a commutant in $\mathcal{U}(U(2n))$

In this section we would like to explore hidden symmetry algebra as Higgs/Hahn algebras in the context of the new scientific direction, which has recently become particularly fashionable - to obtain a Howe duality setting for the interpretation of the Askey-Wilson (AW) and its degenerate algebras as commutants [37], [38]-[41]. Therefore, the goal of this section is to discuss the Howe duality of Higgs-Hahn algebra for $(N \equiv 2n)D$ harmonic oscillator.

By definition, an m -dimensional orthogonal group $O(m)$ is a collection of all linear transformations in m -dimensional linear space, which leave invariant the sum of squares of the components of any vector $x = (x_\alpha)$, $\alpha = 1, \dots, m$ from this space $x^2 = x_\alpha x_\alpha = x_1^2 + \dots + x_m^2$.

Choose a subalgebra $O(n) \oplus O(n)$ in $U(2n)$ generated by all rotations $L_a \equiv L_{ij}$, where $a \in (i, j = 1, \dots, n)$ and $L_b \equiv L_{ij}$, where $b \in (i, j = n+1, \dots, 2n)$ that leave the norm $x_a^2 = x_1^2 + \dots + x_n^2$ and $x_b^2 = x_{n+1}^2 + \dots + x_{2n}^2$ constant correspondingly; obviously $[L_a, L_b] = 0$. Define the next 3 operators

$$\begin{aligned} A_+ &= (a_1^{\dagger 2} + \dots + a_n^{\dagger 2})(a_{n+1}^2 + \dots + a_{2n}^2) \\ A_- &= (a_1^2 + \dots + a_n^2)(a_{n+1}^{\dagger 2} + \dots + a_{2n}^{\dagger 2}) \\ D &= \sum_{a=1}^{a=n} N_a - \sum_{b=n+1}^{b=2n} N_b. \end{aligned} \quad (44)$$

A_\pm and D are manifestly invariant under the rotations generated by L_a and L_b and they clearly commute with H [thus belonging to $U(2n)$]. All other elements of the

commutant are built from those. It is easy to get the following formulas or commutators:

$$[D, A_{\pm}] = \pm 4A_{\pm},$$

as well as the identities [37]:

$$\begin{aligned} a_i^{\dagger 2} a_i^2 &= (N_i - 1)N_i \\ a_i^2 a_i^{\dagger 2} &= (N_i + 1)(N_i + 2) \\ a_i^2 a_j^{\dagger 2} + a_i^{\dagger 2} a_j^2 &= 2N_i N_j + N_i + N_j - 4L_{ij}^2, \quad i, j = 1, \dots, 2n. \end{aligned} \quad (45)$$

Find the remaining commutator $[A_+, A_-]$:

$$[A_+, A_-] = 4 \left(\sum_{i=1}^{i=n} \sum_{j=1}^{j=n} a_i^{\dagger 2} a_j^2 \right) \left(\sum_{b=n+1}^{b=2n} N_b + n/2 \right) - 4 \left(\sum_{a=1}^{a=n} N_a + n/2 \right) \left(\sum_{i=n+1}^{i=2n} \sum_{j=n+1}^{j=2n} a_i^{\dagger 2} a_j^2 \right)$$

It with the help of (45) is readily converted to

$$\begin{aligned} [A_+, A_-] &= 4 \left(\left(\sum_{i=1}^{i=n} N_i \right)^2 + (n-2) \sum_{i=1}^{i=n} N_i - 4 \sum_{\substack{(i<j)=n \\ (i<j)=1}}^{(i<j)=n} L_{ij}^2 \right) \left(\sum_{b=n+1}^{b=2n} N_b + n/2 \right) \\ &\quad - 4 \left(\sum_{a=1}^{a=n} N_a + n/2 \right) \left(\left(\sum_{i=n+1}^{i=2n} N_i \right)^2 + (n-2) \sum_{i=n+1}^{i=2n} N_i - 4 \sum_{\substack{(i<j)=2n \\ (i<j)=n+1}}^{(i<j)=2n} L_{ij}^2 \right) \end{aligned}$$

Since

$$\sum_{a=1}^{a=n} N_a = \frac{1}{2} \left(\frac{H}{\omega} + D - n \right), \quad \sum_{b=n+1}^{b=2n} N_b = \frac{1}{2} \left(\frac{H}{\omega} - D - n \right),$$

upon substituting and after some transformations, one obtains

$$\begin{aligned} [A_+, A_-] &= \left[\left(\frac{H}{\omega} \right)^2 + 8 \left(\sum_{\substack{(i<j)=n \\ (i<j)=1}}^{(i<j)=n} L_{ij}^2 + \sum_{\substack{(i<j)=2n \\ (i<j)=n+1}}^{(i<j)=2n} L_{ij}^2 \right) + n(n-4) \right] D \\ &\quad - 8 \left(\sum_{\substack{(i<j)=n \\ (i<j)=1}}^{(i<j)=n} L_{ij}^2 - \sum_{\substack{(i<j)=2n \\ (i<j)=n+1}}^{(i<j)=2n} L_{ij}^2 \right) \frac{H}{\omega} - D^3 \end{aligned} \quad (46)$$

Thus, we have a Higgs algebra with structural constants of algebra, given as

$$\begin{aligned} \alpha_1 &= \left(\frac{H}{\omega} \right)^2 + 8 \left(\sum_{\substack{(i<j)=n \\ (i<j)=1}}^{(i<j)=n} L_{ij}^2 + \sum_{\substack{(i<j)=2n \\ (i<j)=n+1}}^{(i<j)=2n} L_{ij}^2 \right) + n(n-4) \\ \alpha_2 &= -8 \left(\sum_{\substack{(i<j)=n \\ (i<j)=1}}^{(i<j)=n} L_{ij}^2 - \sum_{\substack{(i<j)=2n \\ (i<j)=n+1}}^{(i<j)=2n} L_{ij}^2 \right) \frac{H}{\omega} \end{aligned} \quad (47)$$

Substituting (44) into (11) using the identities (45) and the resulting expression for the structure constant in (47) gives the following expressions for the operators

$$\begin{aligned} K_1 &= \frac{1}{2} \left[\sum_{a=1}^{a=n} N_a - \sum_{b=n+1}^{b=2n} N_b \right] \\ K_2 &= \sum_{\substack{(i<j)=2n \\ (i<j)=1}}^{(i<j)=2n} L_{ij}^2 + \frac{n(n-2)}{4} \\ K_3 &= [K_1, K_2], \end{aligned} \quad (48)$$

These operators obey the commutation relations of the Hahn algebra with structure constants given as

$$\begin{aligned}\delta_1 &= -\frac{\alpha_2}{4} = 2 \left(\sum_{(i<j)=1}^{(i<j)=n} L_{ij}^2 - \sum_{(i<j)=n+1}^{(i<j)=2n} L_{ij}^2 \right) \frac{H}{\omega} \\ \delta_2 &= \frac{\alpha_1}{2} = \frac{1}{2} \left(\frac{H}{\omega} \right)^2 + 4 \left(\sum_{(i<j)=1}^{(i<j)=n} L_{ij}^2 + \sum_{(i<j)=n+1}^{(i<j)=2n} L_{ij}^2 + \frac{n(n-4)}{8} \right)\end{aligned}\quad (49)$$

5.1. The Howe duality connection

Now we can consider the duality of Howe. It is known [40] - [41] that the state space of an $2n$ -dimensional harmonic oscillator is realized both in the representation $SU(1, 1)$ and $U(2n)$. Taking this fact into account, it can be shown that the embedding of a Hahn algebra into the double tensor product of one pair of $SU(1, 1)$ is in duality with the commutator $O(n) \oplus O(n)$ in the universal algebra of another algebra of the pair $U(2n)$.

For this purpose, we consider the addition of $2n$ metaplectic representations (6), grouped into two pairs. It means, that if we take

$$J^{(1-2n)} = J^{(1-n)} + J^{((n+1)-(2n))}$$

where

$$\begin{aligned}J_0^{(1-2n)} &= J_0^{(1-n)} + J_0^{((n+1)-(2n))} \equiv \frac{1}{2} \left[\left(\sum_{a=1}^{a=n} N_a + n/2 \right) + \left(\sum_{b=n+1}^{b=2n} N_b + n/2 \right) \right] \\ J_+^{(1-2n)} &= J_+^{(1-n)} + J_+^{((n+1)-(2n))} \equiv \frac{1}{2} \left[\sum_{a=1}^{a=n} a_a^{\dagger 2} + \sum_{b=n+1}^{b=2n} a_b^{\dagger 2} \right] \\ J_-^{(1-2n)} &= J_-^{(1-n)} + J_-^{((n+1)-(2n))} \equiv \frac{1}{2} \left[\sum_{a=1}^{a=n} a_a^2 + \sum_{b=n+1}^{b=2n} a_b^2 \right]\end{aligned}\quad (50)$$

then in accordance with the implementation of the operators of the Hahn algebra as in (8) we get

$$K_1 = J_0^{(1)} - J_0^{(2)} \equiv J_0^{(1-n)} - J_0^{((n+1)-(2n))} = \frac{1}{2} \left[\sum_{a=1}^{a=n} N_a - \sum_{b=n+1}^{b=2n} N_b \right]$$

which coincides with the result in the the commutant approach given in (48).

Similarly for K_2 we will have

$$\begin{aligned}K_2 = C^{(1-2n)} &= \left[J_0^{(1-n)} + J_0^{((n+1)-(2n))} \right]^2 - \left(J_0^{(1-n)} + J_0^{((n+1)-(2n))} \right) \\ &\quad - \left(J_+^{(1-n)} + J_+^{((n+1)-(2n))} \right) \left(J_-^{(1-n)} + J_-^{((n+1)-(2n))} \right)\end{aligned}$$

or taking into account (50)

$$K_2 = \frac{1}{4} \left(\frac{H}{\omega} \right)^2 - \frac{1}{2} \frac{H}{\omega} - \frac{1}{4} \left(\sum_{a=1}^{a=n} a_a^{\dagger 2} + \sum_{b=n+1}^{b=2n} a_b^{\dagger 2} \right) \left(\sum_{a=1}^{a=n} a_a^2 + \sum_{b=n+1}^{b=2n} a_b^2 \right)$$

Given the identities (45), we obtain

$$K_2 = \frac{1}{4}\left(\frac{H}{\omega}\right)^2 - \frac{1}{2}\frac{H}{\omega} - \frac{1}{4}\left[A_+ + A_- + \left(\sum_{a=1}^{a=n} N_a\right)^2 + \left(\sum_{b=n+1}^{b=2n} N_b\right)^2\right] - \\ - \frac{(n-2)}{4}\left(\sum_{a=1}^{a=n} N_a + \sum_{b=n+1}^{b=2n} N_b\right) + \sum_{(i<j)=1}^{(i<j)=n} L_{ij}^2 + \sum_{(i<j)=n+1}^{(i<j)=2n} L_{ij}^2$$

or

$$K_2 = -\frac{1}{8}\left[2(A_+ + A_-) + D^2 - \left(\frac{H}{\omega}\right)^2 - n(n-4)\right] + \sum_{(i<j)=1}^{(i<j)=n} L_{ij}^2 + \sum_{(i<j)=n+1}^{(i<j)=2n} L_{ij}^2$$

This expression is identical $K_2 = \sum_{(i<j)=1}^{(i<j)=2n} L_{ij}^2 + \frac{n(n-2)}{4}$ in formula (48), that was obtained earlier in the search for operators commuting during rotations L_{1-n} and $L_{((n+1)-(2n))}$. All of the above explicitly states that the operator K_2 , being as Casimir operator $SU(1,1)$, belongs to the commutant of L_{1-n} and $L_{((n+1)-(2n))}$ in $\mathcal{U}(U(2n))$.

A similar calculation shows that the $SU(1,1)$ Casimir operator for the representation $J^{(ij)}$ is given by the square of the corresponding rotation generator in $U(2n)$, namely

$$C^{(ij)} = L_{ij}^2 + \frac{n(n-4)}{16}. \quad (51)$$

It follows that the structural constants of the Hahn algebra become on the basis of (12):

$$\delta_1 = 4\left(J_0^{(1-n)} + J_0^{((n+1)-(2n))}\right)\left(C^{(1-n)} - C^{((n+1)-(2n))}\right) \\ \equiv 2\frac{H}{\omega}\left[\sum_{(i<j)=1}^{(i<j)=n} L_{ij}^2 - \sum_{(i<j)=n+1}^{(i<j)=2n} L_{ij}^2\right]; \\ \delta_2 = 2\left(J_0^{(1-n)} + J_0^{((n+1)-(2n))}\right)^2 + 4\left(C^{(1-8)} - C^{((n+1)-(2n))}\right) \\ = \frac{1}{2}\left(\frac{H}{\omega}\right)^2 + 4\left(\sum_{(i<j)=1}^{(i<j)=n} L_{ij}^2 + \sum_{(i<j)=n+1}^{(i<j)=2n} L_{ij}^2 + \frac{n(n-4)}{8}\right) \quad (52)$$

in perfect correspondance with (49).

Thus, it is established that the embedding of a Hahn algebra in $SU(1,1) \otimes SU(1,1)$ leads to its description as a commutant in $\mathcal{U}(U(2n))$ in according to the pairing of representations $SU(1,1)$ and $U(2n)$ in the framework of Howe's duality.

6. Dimensional reduction to the singular oscillator

To carry the dimensional reduction of the $2n$ -dimensional IHO in the approach with the embedding of the discrete version of the Hahn algebra in a double tensor product $SU(1,1) \otimes SU(1,1)$ it is necessary to pay attention to the structural constants δ_1 and δ_2 in (12). In according to formula (51), the Casimir operators are included in its definition are grand angular momentum operators \hat{L}_i^2 , $i = 1, 2$ in n -dimensional hyperspherical

coordinates $(\phi_1, \dots, \phi_{n-1})$ are the hyperspherical angles and r is the hyperradius). Therefore, eliminating the angular parts (hyperspherical angles) by separating the variables and performing the gauge transformation $\mathcal{O} \rightarrow \tilde{\mathcal{O}} = r_i^{(n-1)/2} \mathcal{O} r_i^{-(n-1)/2}$ for the radial part, we actually obtain the Hamiltonian of the singular oscillator in two dimensions:

$$\begin{aligned} \widetilde{H} &= 2\omega \left[\tilde{J}_0^{(1-n)} + \tilde{J}_0^{((n+1)-(2n))} \right] \\ &= \frac{1}{2} \left[- \left(\frac{\partial^2}{\partial r_1^2} + \frac{\partial^2}{\partial r_2^2} \right) + \omega^2 (r_1^2 + r_2^2) - \frac{a_1}{r_1^2} - \frac{a_2}{r_2^2} \right], \end{aligned} \quad (53)$$

where $a_i = \hat{L}_i^2 + \frac{(n-3)(n-1)}{4}$, $i = 1, 2$.

However, do not forget that each dimension i corresponds to its n -dimensional hyper-radius r_i . The motion constants are obvious and equal to the operators of the Hahn algebra K_i , which is here the algebra of hidden symmetry. They can be easily finding in according to the above material.

7. Our dual models in the class of quasi-exactly systems

In the previous sections we are considered the dual connection with N -dimensional singular oscillator and the general MICZ-Kepler system in the class of exactly solvable problems. On the other hand, recall, that there are quasi-exactly problems which occupy an intermediate place between exactly solvable problems and non-solvable ones. It is also well known that the theory of quasi-exact systems (CES) gives the following generalization or family of potentials, which includes HO, in the direction of degrees r less than 2, for a fixed N , by

$$V_{<2}(r) = V_{sho} + ar + \frac{b}{r} = \frac{\omega^2 r^2}{2} + \frac{c}{r^2} + ar + \frac{b}{r} \quad (54)$$

with the eigenfunctions

$$R(r) = p_{N-1}(r) r^{l'-c'} e^{-\frac{b'}{2} r^2 - a' r}, \quad (55)$$

where there are the following reassignment of constants from work [42] to our designation $\omega^2 = 2b'^2$; $a = 2a'b'$; $b = -a'(D - 2c')$; $c = c'(c' - D + 1)$; $d = a'^2 - b'(2N + D - 1 - 2c')$; $D = d' + 2l' - 1$; $p_{N-1}(r)$ are polynomial of the $(N - 1)$ -th degree.

This QES potential appears in a number of applications to the systems with two electrons ([43] - [44]).

Also at present we want to go to opposite direction and to consider the potential which includes HO and degrees of the variable more than 2. The QES theory gives the other generalization in this direction or the family of potentials for a fixed N , by see [42],

$$V_{>2}(r) = V_{sho} + br^4 + ar^6 = (\omega r)^2/2 + \frac{c}{r^2} + br^4 + ar^6, \quad (56)$$

with the eigenfunctions

$$R(r) = p_{N-1}(r^2) r^{l'-c'} e^{-\frac{a' r^4}{4} - \frac{b' r^2}{2}}, \quad (57)$$

where the following reassignment of constants from work [42] to our designation $\omega^2 = 2[b'^2 - (4N + D - 2c' - 1)a']$; $a = a'^2$; $b = 2a'b'$; $c = c'(c' - D + 1)$.

The one-dimensional Hamiltonian of the nonrelativistic quantum systems with this anharmonic potential (56) is well known as the crucial example that is stimulated the investigation of quasi-exactly solvable systems.

Thus, we can offer four different models of the $(N \equiv 2n)D$ anisotropic and nonlinear anharmonic oscillator in QES class. Each model is represented by a sum of two independent (this provides an anisotropic effect - $\omega_1 \neq \omega_2$) oscillators of dimensions $D = n$ with their various nonlinear anharmonic terms of the potential. In other words, we will further consider the following Hamiltonians of dimension $D = n$:

$$\begin{aligned} {}_{<2}H &= \left[-\frac{1}{2} \frac{\partial^2}{\partial x_s \partial x_s} + V_{<2}(x_s x_s) \right]; \\ V_{<2}(x_s x_s) &= V_{sho}(x_s x_s) + \frac{b}{\sqrt{x_s x_s}} + a\sqrt{x_s x_s} \\ &= \frac{\omega^2 x_s x_s}{2} + \frac{c}{x_s x_s} + \frac{b}{\sqrt{x_s x_s}} + a\sqrt{x_s x_s} \end{aligned} \quad (58)$$

and

$$\begin{aligned} {}_{>2}H &= \left[-\frac{1}{2} \frac{\partial^2}{\partial x_s \partial x_s} + V_{>2}(x_s x_s) \right]; \\ V_{>2}(x_s x_s) &= V_{sho}(x_s x_s) + b(x_s x_s)^2 + a(x_s x_s)^3 \\ &= \frac{\omega^2 x_s x_s}{2} + \frac{c}{x_s x_s} + b(x_s x_s)^2 + a(x_s x_s)^3. \end{aligned} \quad (59)$$

In hyperspherical coordinates, the potentials of these Hamiltonians are successively reduced to the potential (54) and (56). In any case, the final wave function $\psi(\mathbf{u}, \mathbf{v})$ of Eq. (37) will be represented by the product of the wave functions of each oscillator $\Psi(\mathbf{r}_a) \equiv R(r_a)\Omega(\phi_a)$ of Eq. (58) - (59). Recall that the hyperradius part $R(r_a)$ of the eigenfunction $\Psi(\mathbf{r}_a)$ has the form (55) and (57), respectively, depending on the potential (54) and (56).

Thus, now we are ready to determine the potential of the dual analog for our generalizations of the $(N \equiv 2n)D$ oscillator [4 models from the combination (58 – 59)] as $n + 1$ related MICZ-Kepler systems in different coordinates. However, the analysis performed in [45] showed the absence of new solvable models for the dual partner or the general MICZ-Kepler system in this generalization for the spherical coordinates. Therefore, it is not considered further and the solvability of the Schrödinger equation of the MICZ-Kepler problems by the variables separation method will be discussed only in QES class for parabolic coordinates. In according to its definition (31) and using the identities (16) it is not hard to receive:

$$\begin{aligned} r &= \sqrt{x_\lambda x_\lambda} = u_s u_s + v_s v_s; & x_{n+1} &= u_s u_s - v_s v_s \\ 2u_s u_s &= r + x_{n+1} \equiv u; & 2v_s v_s &= r - x_{n+1} \equiv v \end{aligned}$$

Therefore, we have additional additive terms, but each of them now is dependent on only one parabolic variable. Thus, the Schrödinger equation of our generalized MIC-

Kepler system with new terms in the hyperparabolic coordinates can be rewritten in terms of separated equations of variables u, v and angular variables ϕ, ϕ'

$$\left[\frac{1}{u^{\frac{n-2}{2}}} \frac{\partial}{\partial u} \left(u^{\frac{n}{2}} \frac{\partial}{\partial u} \right) - \frac{L(L+n-2) + 4\lambda_1}{u} - V_u + Z_1 - P \right] U(u) = 0, \quad (60)$$

$$\left[\frac{1}{v^{\frac{n-2}{2}}} \frac{\partial}{\partial v} \left(v^{\frac{n}{2}} \frac{\partial}{\partial v} \right) - \frac{L(L+n-2) + 4\lambda_2}{v} - V_v + Z_2 + P \right] V(v) = 0, \quad (61)$$

$$\left[\hat{L}^2 - L(L+n-2) \right] \Phi(\phi, \phi') = 0, \quad (62)$$

where P is the separation constant.

Let us specify the value of $V_u \equiv V(u_s u_s)$ and $V_v \equiv V(v_s v_s)$ for each model

- Model 1

$$\begin{aligned} H\psi_1(\mathbf{u}, \mathbf{v}) &= [{}_{<2}H_1 + {}_{<2}H_2] \psi_{<2}(\mathbf{u}) \psi_{<2}(\mathbf{v}) \equiv Z\psi_1(\mathbf{u}, \mathbf{v}) \\ V_u &= \frac{-uE_1}{2} + b_1 \sqrt{\frac{2}{u}} + a_1 \sqrt{\frac{u}{2}} \\ V_v &= \frac{-vE_2}{2} + b_2 \sqrt{\frac{2}{v}} + a_2 \sqrt{\frac{v}{2}} \end{aligned} \quad (63)$$

- Model 2

$$\begin{aligned} H\psi_2(\mathbf{u}, \mathbf{v}) &= [{}_{<2}H_1 + {}_{>2}H_2] \psi_{<2}(\mathbf{u}) \psi_{>2}(\mathbf{v}) \equiv Z\Psi_{<2}(\mathbf{r}_1) \Psi_{>2}(\mathbf{r}_2) \\ V_u &= \frac{-uE_1}{2} + b_1 \sqrt{\frac{2}{u}} + a_1 \sqrt{\frac{u}{2}} \\ V_v &= \frac{-uE_2}{2} + \frac{b_2 u^2}{4} + \frac{a_2 u^3}{8} \end{aligned} \quad (64)$$

- Model 3

$$\begin{aligned} H\psi_3(\mathbf{u}, \mathbf{v}) &= [{}_{>2}H_1 + {}_{<2}H_2] \psi_{>2}(\mathbf{u}) \psi_{<2}(\mathbf{v}) \equiv Z\Psi_{>2}(\mathbf{r}_1) \Psi_{<2}(\mathbf{r}_2) \\ V_u &= \frac{-uE_1}{2} + \frac{b_1 u^2}{4} + \frac{a_1 u^3}{8} \\ V_v &= \frac{-vE_2}{2} + b_2 \sqrt{\frac{2}{v}} + a_2 \sqrt{\frac{v}{2}} \end{aligned} \quad (65)$$

- Model 4

$$\begin{aligned} H\psi_4(\mathbf{u}, \mathbf{v}) &= [{}_{>2}H_1 + {}_{>2}H_2] \psi_{>2}(\mathbf{u}) \psi_{>2}(\mathbf{v}) \equiv Z\Psi_{>2}(\mathbf{r}_1) \Psi_{>2}(\mathbf{r}_2) \\ V_u &= \frac{-uE_1}{2} + \frac{b_1 u^2}{4} + \frac{a_1 u^3}{8} \\ V_v &= \frac{-uE_2}{2} + \frac{b_2 u^2}{4} + \frac{a_2 u^3}{8} \end{aligned} \quad (66)$$

At first glance, it seems that 4 models require solving 2 qualitatively different types of problems. In particular, for model 4 we will have the following equations after the gauge transformation $\mathcal{O} \rightarrow \tilde{\mathcal{O}} = x^{n/4} \mathcal{O} x^{-n/4}$ for (60) – (61):

$$\left[\frac{\partial^2}{\partial x^2} - \frac{d_i}{x^2} - \frac{a_i}{8} x^2 + \frac{E_i}{2} + \frac{Z_i \pm P}{x} - \frac{b_i}{4} x \right] X(x) = 0, \quad (67)$$

where $d_i = L(L+n-2) + 4\lambda_1 - \frac{n(n-4)}{16}$, $i = 1, 2$.

However, getting rid of irrationality by introducing a new variable $x \rightarrow z^2$ in one type of problem leads to the solution of a second type problem, which in turn coincides with the problem we considered earlier (54) with solution (55). In particular, for model 1 we will have the following equations after the gauge transformation $\mathcal{O} \rightarrow \tilde{\mathcal{O}} = z^{\frac{(n-1)}{2}} \mathcal{O} z^{-\frac{(n-1)}{2}}$ for (60) – (61):

$$\left[\frac{\partial^2}{\partial z^2} - \frac{d_i}{z^2} + 2E_i z^2 + 4(Z_i \pm P) - 4\frac{b_i}{z} - 4a_i z \right] Z(z) = 0, \quad (68)$$

where $d_i = 4L(L + n - 2) + 16\lambda_1 - \frac{(n-1)(n-3)}{4}$, $i = 1, 2$.

Thus, we have obtained that in parabolic coordinates all proposed 4 models (63) – (66) is the (quasi) exactly solvable models.

Now let us compare the results obtained with those available in the literature [46]. If we consider just a problem with potential $V_{ho}(x_i x_i) \equiv \frac{\omega^2 x_i x_i}{2}$ (38), but without a condition $\omega_1 = \omega_2$, then we obtain in the $(N \equiv 2n) - D$ the sum of two independent harmonic oscillators that will be dual to the $n + 1$ MICZ-problem with the potential $\cos \theta$:

$$\begin{aligned} \frac{\omega_1^2 u_s u_s}{8} + \frac{\omega_2^2 v_s v_s}{8} &\equiv -E_1 u_s u_s - E_2 v_s v_s \\ \rightarrow \frac{1}{r} \left[-\frac{E_1(r + x_{n+1})}{2} - \frac{E_2(r - x_{n+1})}{2} \right] &\equiv -\frac{E_1}{2} \left(1 + \frac{x_{n+1}}{r} \right) - \frac{E_2}{2} \left(1 - \frac{x_{n+1}}{r} \right) \\ = -\frac{E_1 + E_2}{2} - \frac{E_1 - E_2}{2} \cos \theta &\equiv -E_{MICZ} + \frac{\Delta w^2}{4} \cos \theta. \end{aligned}$$

The analog of the 4th order anisotropic potential term for oscillator system [46] is in our $(N \equiv 2n) - D$ case the sum of 2 harmonic oscillators with identical in value but different in sign potential coefficients for nonlinearity of the 4th order ($b_1 = -b_2$). In the $n + 1$ -dimensional space, this leads to the desired linear term:

$$\begin{aligned} b_1(u_s u_s)^2 + b_2(v_s v_s)^2 &= b \left[(u_s u_s)^2 - (v_s v_s)^2 \right] \\ \rightarrow \frac{b}{r} \left[\frac{(r + x_{n+1})^2}{4} - \frac{(r - x_{n+1})^2}{4} \right] &= \frac{b}{4r} [4 r x_{n+1}] \\ = b x_{n+1} &\equiv br \cos \theta. \end{aligned}$$

We note in particular that in our case there is still a new term with a higher degree of nonlinearity than the one considered above.

8. Conclusion

We have shown that N -dimensional singular oscillator and $(n + 1)$ -dimensional generalized MIC-Kepler system are dual to each other and the duality transformation is the generalized version of the Kustaanheimo-Stiefel transformation. The solvability of the Schrödinger equation of these problems by the variables separation method were given in different coordinates. We have successfully built the exact analytical

solutions of the Schrödinger equation for $N \equiv 2n$ -dimensional singular oscillator in double coordinates and for the $(n+1)$ -dimensional generalized MIC-Kepler system in spherical and parabolic coordinates. The overlap coefficients between wavefunctions in these coordinates were shown to coincide with Clebsch-Gordan coefficients for $SU(1,1)$ algebra. It turns out that the quadratic Hahn algebra $QH(3)$ as a hidden symmetry remains unchanged with different decomposition of the original space real space \mathbb{R}^N into components in the framework of addition rule for $SU(1,1)$ algebra. Also the hidden symmetry algebra as Higgs/Hahn algebras was clearly shown by the commutant approach in the sense of Howe duality. A dimensional reduction was carried out to a singular oscillator of two dimensions, each variable is the n -dimensional hyper-radius r . The dual connection with $N \equiv 2n$ -dimensional singular oscillator and the $(n+1)$ general MICZ-Kepler system in the class of quasi-exact problems was considered. In the above framework of generalized KS transformations, some generalization of HO by anisotropic and nonlinear inharmonic terms and its dual analog was shown and analyzed. The exact analytical solutions of the Schrödinger equation for abovementioned problems for QES class were discussed and given for four series of dual quasi-exact solvable models. In particular, a comparison with similar results in lower dimensions and its generalization were given.

References

- [1] Kustaanheimo, Stiefel E 1965 *J.Reine Angew.Math.*, **218**, 204 -219
- [2] Bergmann P, Frishman Y 1965 *J. Phys. A: Math. Theor.*, **6**, 1855 - 1856
- [3] Boiteux M 1973 *J Physica*, **65**, 381 - 395
- [4] Komarov L I, Romanova T S 1982 *Vesti AN BSSR. Seria phys.-math. nauk*, **2**, 98 -103
- [5] Hopf H, "Über die Abbildungen der dreidimensionalen Sphäre auf die Kugelfläche" 1931 *Mathematische Annalen (Berlin: Springer)*, **104**, 637665
- [6] Nersisyan A, Ter-Antonian V and Tsulaia M M 1996 *Mod. Phys. Lett. , A***11**, 1605
- [7] Bellucci S, Toppan J and Yeghikyan V 2010 *J. Phys. A: Math. Theor.*, **43**, 045205
- [8] Davtyan L S, Mardoyan L G, Pogosyan G S, Sissakian A N and Ter-Antonyan V M 1987 *J. Phys. A: Math. Theor.*, **20**, 6121
- [9] Iwai T and Uwano Y 1986 *J. Phys. A: Math. Theor.*, **27**, 1523
- [10] Mardoyan L G, Sissakian A N and Ter-Antonyan V M 1996 *JINR, Dubna, Preprint E2*, **96**, 24
- [11] Mardoyan L G, Sissakian A N and Ter-Antonyan V M 1998 *Phys. Atom. Nucl.*, **61**, 1746
- [12] Mardoyan L G, Sissakian A N and Ter-Antonyan V M 1999 *Mod. Phys. Lett. A*, **14**, 1303
- [13] Mardoyan L G, Sissakian A N and Ter-Antonyan V M 2000 *Theor. Math. Phys.*, **123**, 451
- [14] Mardoyan L G 2002 *Phys. Atomic Nuclei.*, **65**, 1096
- [15] Mardoyan L G 2003 *J. Math. Phys.*, **44**, 4981
- [16] Mardoyan L 2003 Dyon-oscillator duality. Hidden symmetry of the Yang-Coulomb monopole, [arXiv:quant-ph/0302162](https://arxiv.org/abs/quant-ph/0302162)
- [17] Mardoyan L G and Petrosyan M G 2007 *Phys. Atom. Nucl.*, **70**, 572
- [18] Marquette I 2012 *J. Math. Phys.*, **53**, 022103
- [19] Pletyukhov M V and Tolkachev E M 1999 *J. Math. Phys.*, **40**, 93
- [20] Pletyukhov M V and Tolkachev E A 1999 *Rep. Math. Phys. ,* **43**, 303
- [21] Pletyukhov M V and Tolkachev E A 1999 *J. Phys. A: Math. Theor.*, **32**, L249
- [22] Pletyukhov M V and Tolkachev E A 2000 *J. Math. Phys.*, **41**, 187
- [23] Le V.-H., Nguyen T.-S. and Phan N.-H. 2009 *J. Phys. A: Math. Theor.*, **42**, 175204

- [24] Le V.-H. and Nguyen T.-S. 2011 *J. Math. Phys.*, **52**, 032105
- [25] Le V.-H., Truong C.-T. and Phan T.-T. 2011 *J. Math. Phys.*, **52** (7), 072101
- [26] Phan N.-H. and Le V.-H. 2012 *Journal of Mathematical Physics*, **53** (8), 082103
- [27] Nguyen T.-S., Le D.-N., Thoi T.-Q. N. and Le V.-H. 2015 *J. Math. Phys.*, **56**, no. 5, 052103
- [28] Phan N.-H., Le D.-N., Thoi T.-Q. N. and Le V.-H. 2018 *J. Math. Phys.* **59**, no. 3, 032102
- [29] Komarov L I, Romanova T S, and Chan Za An 1987 *Vestsi Akad. Navuk B. SSR, Ser. Fiz. Mat. Navuk.*, No. 1, 90
- [30] Pris' I E and Tolkachev E A 1991 *Yad. Fiz.*, **54**, 962
- [31] Jaroszkiewicz G A and McHale G 1987 *Ann. Phys. (N. Y.)*, **175**, 267
- [32] Cordani B 1989 *J. Phys. A: Math. Theor.*, **22**, 2441
- [33] Le V.-H. and Komarov L I 1993 *Phys. Lett. A*, **177**, 121
- [34] Zhedanov A S 1993 *J. Phys. A: Math. Theor.*, **26**, 46334641
- [35] Higgs P W 1979 *J. Phys. A: Math. Theor.*, **12**, no. 4, 309-323
- [36] Leemon H I 1979 *J. Phys. A: Math. Theor.* **12**, no. 4, 489501
- [37] Frappat L, Gaboriaud J, Vinet L, Vinet S and Zhedanov A S 2018 The Higgs and Hahn algebras from a Howe duality perspective arXiv:1811.09359 [math-ph]
- [38] Gaboriaud J, Vinet L, Vinet S and Zhedanov A S 2018 The generalized Racah algebra as a commutant arXiv:1808.09518 [math-ph]
- [39] Gaboriaud J, Vinet L, Vinet S and Zhedanov A S 2018 The Racah algebra as a commutant and Howe duality arXiv:1808.05261 [math-ph]
- [40] Howe R 1989 *Transactions of the American Mathematical Society*, **313**(2), 539-570
- [41] Rowe D J, Carvalho M J and Repka J 2012 *Reviews of Modern Physics*, **84**(2), 711757
- [42] Turbiner A V 1988 *Funct. Anal. Appl.* **22**, 163
- [43] Turbiner A V 1994 *Phys. Rev. A*, **50**, 5335
- [44] Turbiner A V and Escobar-Ruiz M A 2013 *J. Phys. A: Math. Theor.*, **46**, 295204
- [45] Lavrenov A and Lavrenov I 2019 16D anisotropic inharmonic oscillator and 9D related (MICZ-) Kepler-like systems arXiv:1903.10847 [math-ph]
- [46] Nersessian A and Yeghikyan V 2008 *J. Phys. A: Math. Theor.*, **41**, 155203