Local realistic interpretation of entangled photon pairs in the Weyl-Wigner formalism

Emilio Santos Universidad de Cantabria. Santander. Spain. Email <emilio.santos@unican.es>.

February, 3, 2020

Abstract

A polarization correlation experiment with two maximally entangled photons created by spontaneous parametric down-conversion is studied in the Weyl-Wigner formalism, that reproduces the quantum predictions. An interpretation is proposed in terms of stochastic processes assuming that the quantum vacuum fields are real. This proves that local realism is compatible with the violation of Bell inequalities, thus rebutting the claim that local realism has been refuted by entangled photon experiments. Entanglement appears as a correlation between fluctuations of a signal field and vacuum fields.

1 The empirical refutation of Bell's local realism

In 2015 experiments were reported showing for the first time the loophole-free violation of a Bell inequality[1],[2]. The result has been interpreted as the "death by experiment for local realism", this being the hypothesis that "the world is made up of real stuff, existing in space and changing only through local interactions ... about the most intuitive scientific postulate imaginable" [3]. This statement, and many similar ones, emphasize both the relevance of *local realism* for our understanding of the physical world and the fact that it has been *refuted empirically*. Nevertheless it is worth studying the possibility of a loophole in the empirical refutation via a new definition of locality weaker than Bell's. In this article I search for such a weak locality, compatible with the said experiments [1],[2], that involved photon pairs entangled in polarization produced via spontaneous parametric down conversion. Thus I will analyze such experiments using the Weyl-Wigner formalism of quantum optics, rather than the more usual Hilbert-space formalism. Previously I revisit briefly the origin and meaning of the Bell inequalities[4].

Bell defined "local hidden variables" model, later named "local realistic", to be any model of an experiment where the results of all measurements may be interpreted according to the formulas

$$\langle A \rangle = \int \rho(\lambda) \, d\lambda M_A(\lambda, A) \,, \langle B \rangle = \int \rho(\lambda) \, d\lambda M_B(\lambda, B) \,,$$

$$\langle AB \rangle = \int \rho(\lambda) \, d\lambda M_A(\lambda, A) \, M_B(\lambda, B) \,,$$

$$(1)$$

where $\lambda \in \Lambda$ is one or several random ("hidden") variables, $\langle A \rangle$, $\langle B \rangle$ and $\langle AB \rangle$ being the expectation values of the results of measuring the observables A, Bor their product AB, respectively. Here we will consider that the observables correspond to detection, or not, of some signals (e. g. photons) by two parties named Alice and Bob, attaching the values 1 or 0 to these two possibilities. In this case $\langle A \rangle$, $\langle B \rangle$ correspond to the single and $\langle AB \rangle$ to the coincidence detection rates respectively. The following mathematical conditions are assumed

$$\rho(\lambda) \ge 0, \int \rho(\lambda) \, d\lambda = 1, M_A(\lambda, A) \in \{0, 1\}, M_B(\lambda, B) \in \{0, 1\}.$$
(2)

Eqs.(2) corresponds to a "deterministic model" where the statistical aspects derive from the probabilistic nature of the hidden random variables $\{\lambda\}$. More general models may be constructed where the whole interval [0, 1] is substituted for $\{0, 1\}$ in eq.(2). A constraint of locality is included, namely $M_A(\lambda, A)$ should be independent of the choice of the observable B, $M_B(\lambda, B)$ independent of Aand $\rho(\lambda)$ independent of both A and B[5]. From these conditions it is possible to derive empirically testable (Bell) inequalities[6], [7]. The tests are most relevant if the measurements performed by Alice and Bob are spacially separated in the sense of relativity theory.

For experiments measuring polarization correlation of photon pairs the Clauser-Horne inequality[6] may be written

$$\langle \theta_1 \rangle + \langle \phi_1 \rangle \ge \langle \theta_1 \phi_1 \rangle + \langle \theta_1 \phi_2 \rangle + \langle \theta_2 \phi_1 \rangle - \langle \theta_2 \phi_2 \rangle, \qquad (3)$$

where θ_j stands for the observable "detection of a photon with the Alice detector in front of a polarizer at angle θ_j ". Similarly ϕ_k for Bob detector. For simplicity in this paper I will study the case of maximally entangled photons, although in the mentioned experiments[1],[2] the photon pairs had partial entanglement, which made the experiments easier. I will present a local model that predicts the following single and coincidence rates by Alice and Bob

$$\langle \theta_j \rangle = \langle \phi_k \rangle = \frac{1}{2} K,$$

$$\langle \theta_j \phi_k \rangle = \frac{1}{2} K \cos^2 \left(\theta_j - \phi_k \right) = \frac{1}{4} K \left[1 + \cos \left(2\theta_j - 2\phi_k \right) \right],$$

$$(4)$$

where K is a constant that depends on the particular experimental setup. It is easy to check that the prediction eq.(4) violates the inequality eq.(3) for some choices of angles. For instance the choice

$$\theta_1 = \frac{\pi}{4}, \phi_1 = \frac{\pi}{8}, \theta_2 = 0, \phi_2 = \frac{3\pi}{8}, \theta_3 = 0, \phi_2 = \frac{3\pi}{8}, \theta_3 = 0, \phi_4 = \frac{3\pi}{8}, \phi_4$$

violates the inequality eq.(3) leading to

$$K \not\ge \frac{1}{2} \left(1 + \sqrt{2} \right) K \simeq 1.207 K$$

So far we have assumed ideal detectors, for real detectors the predicted single rates $\langle \theta_j \rangle$ and $\langle \phi_k \rangle$ should be multiplied times the detection efficiencies η_A and η_B , respectively, and the coincidence rate $\langle \theta_j \phi_k \rangle$ times $\eta_A \eta_B$, whence the empirical violation of the inequality eq.(3) would require high detection efficiencies, that is

$$\eta_A + \eta_B < (1 + \sqrt{2})\eta_A \eta_B \Rightarrow \eta > 0.828 \text{ if } \eta_A = \eta_B = \eta.$$

Experiments with some non-maximal entanglement need only $\eta > 2/3$ [7], that was the reason for using such entanglement in the actual experiments [1], [2].

In the following sections I shall shortly review the treatment within the Weyl-Wigner formalism of the polarization correlation measurement of two maximally entangled photons produced via spontaneous parametric down conversion (SPDC). Thus I continue a theoretical interpretation of SPDC experiments within the WW formalism in the Heisenberg picture, that was initiated in the nineties of the past century [8] - [18]. In many of those early studies the approach was heuristic and one of the purposes of this article is to provide a more formal foundation. The WW formalism suggests an intuitive picture for photon entanglement and the interpretation of SPDC experiments in terms of random variables and stochastic processes.

2 The Weyl-Wigner formalism in quantum optics

2.1 Definition

The WW formalism was developed for non-relativistic quantum mechanics, where the basic observables involved are positions, \hat{x}_j , and momenta, \hat{p}_j , of the particles[19], [20], [21], [22], [23], [24]. It may be trivially extended to quantum optics provided we interpret \hat{x}_j and \hat{p}_j to be the sum and the difference of the creation, \hat{a}_j^{\dagger} , and annihilation, \hat{a}_j , operators of the *j* normal mode of the radiation. That is

$$\hat{x}_{j} \equiv \frac{c}{\sqrt{2}\omega_{j}} \left(\hat{a}_{j} + \hat{a}_{j}^{\dagger} \right), \hat{p}_{j} \equiv \frac{i\hbar\omega_{j}}{\sqrt{2}c} \left(\hat{a}_{j} - \hat{a}_{j}^{\dagger} \right)
\Rightarrow \hat{a}_{j} = \frac{1}{\sqrt{2}} \left(\frac{\omega_{j}}{c} \hat{x}_{j} + \frac{ic}{\hbar\omega} \hat{p}_{j} \right), \hat{a}_{j}^{\dagger} = \frac{1}{\sqrt{2}} \left(\frac{\omega_{j}}{c} \hat{x}_{j} - \frac{ic}{\hbar\omega_{j}} \hat{p}_{j} \right).$$
(5)

Here \hbar is Planck constant, c the velocity of light and ω_j the frequency of the normal mode. In the following I will use units $\hbar = c = 1$. For the sake of clarity I shall represent the operators in a Hilbert space with a 'hat', e. g. $\hat{a}_j, \hat{a}_j^{\dagger}$ and the amplitudes in the WW formalism without 'hat', e. g. a_j, a_j^* .

The connection with the Hilbert-space formalism is made via the Weyl transform as follows. For any trace class operator \hat{M} of the former we define its Weyl transform to be a function of the field operators $\left\{\hat{a}_{j}, \hat{a}_{j}^{\dagger}\right\}$, that is

$$W_{\hat{M}} = \frac{1}{(2\pi^2)^n} \prod_{j=1}^n \int_{-\infty}^\infty d\lambda_j \int_{-\infty}^\infty d\mu_j \exp\left[-2i\lambda_j \operatorname{Re} a_j - 2i\mu_j \operatorname{Im} a_j\right] \\ \times Tr\left\{\hat{M} \exp\left[i\lambda_j \left(\hat{a}_j + \hat{a}_j^{\dagger}\right) + i\mu_j \left(\hat{a}_j - \hat{a}_j^{\dagger}\right)\right]\right\}.$$

The transform is *invertible* that is

$$\hat{M} = \frac{1}{(2\pi^2)^{2n}} \prod_{j=1}^n \int_{-\infty}^\infty d\lambda_j \int_{-\infty}^\infty d\mu_j \exp\left[i\lambda_j \left(\hat{a}_j + \hat{a}_j^\dagger\right) + i\mu_j \left(\hat{a}_j - \hat{a}_j^\dagger\right)\right] \\ \times \prod_{j=1}^n \int_{-\infty}^\infty d\operatorname{Re} a_j \int_{-\infty}^\infty d\operatorname{Im} a_j W_{\hat{M}}\left\{a_j, a_j^*\right\} \exp\left[-2i\lambda_j \operatorname{Re} a_j - 2i\mu_j \operatorname{Im} a_j\right]$$

The transform is *linear*, that is if f is the transform of \hat{f} and g the transform of \hat{g} , then the transform of $\hat{f} + \hat{g}$ is f + g.

It is standard wisdom that the WW formalism is unable to provide any intuitive picture of the quantum phenomena. The reason is that the Wigner function, that represents the quantum states, is not positive definite in general and therefore cannot be interpreted as a probability distribution (of positions and momenta in quantum mechanics, or field amplitudes in quantum optics). However we shall see that in quantum optics the formalism in the Heisenberg representation, where the evolution goes in the field amplitudes, allows the interpretation of the experiments using the Wigner function only for the vacuum state, that is positive definite.

The use of the WW formalism in quantum optics has the following features in comparison with the Hilbert-space formalism:

1. It is just quantum optics, therefore the predictions for experiments are the same.

2. The calculations using the WW formalism are generally no more involved than the corresponding ones in Hilbert space, and sometimes they are easier because no problem of non-commutativity arises.

3. The formalism suggests a physical picture in terms of random variables and stochastic processes. In particular the counterparts of creation and annihilation operators look like random amplitudes.

Here we shall use the formalism in the Heisenberg picture, where the evolution appears in the observables. On the other hand the concept of photon does not appear in the WW formalism.

2.2 Properties

All properties of the WW transform in particle systems may be translated to quantum optics via eqs.(5). The transform allows getting a function of (c-number) amplitudes for any trace-class operator (e. g. any function of the

creation and annihilation operators of 'photons'). In particular we may get the (Wigner) function corresponding to any quantum state. For instance the vacuum state, represented by the density matrix $|0\rangle\langle 0|$, is associated to the following Wigner function

$$W_0 = \prod_j \frac{2}{\pi} \exp\left(-2|a_j|^2\right).$$
 (6)

This function may be interpreted as a (positive) probability distribution. Hence the picture that emerges is that the quantum vacuum of the electromagnetic field (also named *zeropoint field*, *ZPF*) consists of stochastic fields with a probability distribution independent for every mode, having a Gaussian distribution with mean energy $\frac{1}{2}\hbar\omega$ per mode.

Similarly there are functions associated to the observables. For instance the following Weyl transforms are obtained

$$\hat{a}_{j} \quad \leftrightarrow \quad a_{j}, \hat{a}_{j}^{\dagger} \leftrightarrow a_{j}^{*}, \frac{1}{2} \left(\hat{a}_{j}^{\dagger} \hat{a}_{j} + \hat{a}_{j} \hat{a}_{j}^{\dagger} \right) \leftrightarrow a_{j} a_{j}^{*} = |a_{j}|^{2},$$

$$\hat{a}_{j}^{\dagger} \hat{a}_{j} \quad = \quad \frac{1}{2} \left(\hat{a}_{j}^{\dagger} \hat{a}_{j} + \hat{a}_{j} \hat{a}_{j}^{\dagger} \right) + \frac{1}{2} \left(\hat{a}_{j}^{\dagger} \hat{a}_{j} - \hat{a}_{j} \hat{a}_{j}^{\dagger} \right) \leftrightarrow |a_{j}|^{2} - \frac{1}{2},$$

$$\left(\hat{a}_{j}^{\dagger} + \hat{a}_{j} \right)^{n} \quad \leftrightarrow \quad \left(a_{j} + a_{j}^{*} \right)^{n}, \left(\hat{a}_{j}^{\dagger} - \hat{a}_{j} \right)^{n} \leftrightarrow \left(a_{j} - a_{j}^{*} \right)^{n}, n \text{ an integer.} \quad (7)$$

I stress that the quantities a_j and a_j^* are c-numbers and therefore they commute with each other. The former eqs.(7) mean that in expressions *linear in creation* and/or annihilation operator the Weyl transform just implies "removing the hats". However this is not the case in nonlinear expressions in general. In fact from the latter two eqs.(7) plus the linearity property it follows that for a product in the WW formalism the Hilbert space counterpart is

$$a_j^k a_j^{*^l} \leftrightarrow (\hat{a}_j^k \hat{a}_j^{\dagger l})_{sym}, \tag{8}$$

where the subindex sym means writing the product with all possible orderings and dividing for the number of terms. Hence the WW field amplitudes corresponding to products of field operators may be obtained putting the operators in symmetrical order via the commutation relations. Particular instances that will be useful latter are the following

$$\hat{a}_{j}^{\dagger}\hat{a}_{j} \rightarrow |a_{j}|^{2} - \frac{1}{2}, \hat{a}_{j}\hat{a}_{j}^{\dagger} \rightarrow |a_{j}|^{2} + \frac{1}{2}, \hat{a}_{j}^{2} \rightarrow a_{j}^{2}, \hat{a}_{j}^{\dagger 2} \rightarrow \hat{a}_{j}^{*2} \\
\hat{a}_{j}^{\dagger}\hat{a}_{j}\hat{a}_{j}^{\dagger}\hat{a}_{j} \rightarrow |a_{j}|^{4} - |a_{j}|^{2}, \hat{a}_{j}\hat{a}_{j}^{\dagger}\hat{a}_{j}\hat{a}_{j}^{\dagger} \rightarrow |a_{j}|^{4} + |a_{j}|^{2},$$

$$\hat{a}_{j}^{\dagger}\hat{a}_{j}^{\dagger}\hat{a}_{j}\hat{a}_{j} \rightarrow |a_{j}|^{4} - 2|a_{j}|^{2} + \frac{1}{2}, \hat{a}_{j}\hat{a}_{j}\hat{a}_{j}^{\dagger}\hat{a}_{j}^{\dagger} \rightarrow |a_{j}|^{4} + 2|a_{j}|^{2} + \frac{1}{2}.$$
(9)

Other properties may be easily obtained from well known results of the standard Weyl-Wigner formalism in particle quantum mechanics. I will present them omitting the proofs. Expectation values may be calculated in the WW formalism as follows. In the Hilbert-space formalism they read $Tr(\hat{\rho}\hat{M})$, or in particular $\langle \psi \mid \hat{M} \mid \psi \rangle$, whence the translation to the WW formalism is obtained taking into account that the trace of the product of two operators becomes

$$Tr(\hat{\rho}\hat{M}) = \int W_{\hat{\rho}}\left\{\hat{a}_{j}, \hat{a}_{j}^{\dagger}\right\} W_{\hat{M}}\left\{\hat{a}_{j}, \hat{a}_{j}^{\dagger}\right\} \prod_{j} d\operatorname{Re}a_{j}d\operatorname{Im}a_{j}.$$

That integral is the WW counterpart of the trace operation in the Hilbert-space formalism. Particular instances are the following expectations that will be of interest later on

$$\left\langle \left| a_{j} \right|^{2} \right\rangle \equiv \int d\Gamma W_{0} \left| a_{j} \right|^{2} = \frac{1}{2}, \left\langle a_{j}^{n} a_{j}^{*m} \right\rangle = 0 \text{ if } n \neq m.$$

$$\left\langle 0 \left| \hat{a}_{j}^{\dagger} \hat{a}_{j} \right| 0 \right\rangle = \int d\Gamma (a_{j}^{*} a_{j} - \frac{1}{2}) W_{0} = 0,$$

$$\left\langle 0 \left| \hat{a}_{j} \hat{a}_{j}^{\dagger} \right| 0 \right\rangle = \int d\Gamma (\left| a_{j} \right|^{2} + \frac{1}{2}) W_{0} = 2 \left\langle \left| a_{j} \right|^{2} \right\rangle = 1,$$

$$\left\langle \left| a_{j} \right|^{4} \right\rangle = 1/2, \left\langle \left| a_{j} \right|^{n} \left| a_{k} \right|^{m} \right\rangle = \left\langle \left| a_{j} \right|^{n} \right\rangle \left\langle \left| a_{k} \right|^{m} \right\rangle \text{ if } j \neq k.$$

$$(10)$$

where W_0 is the Wigner function of the vacuum, eq.(6). This means that in the WW formalism the field amplitude a_j (coming from the vacuum) behaves like a complex random variable with Gaussian distribution and mean square modulus $\langle |a_j|^2 \rangle = 1/2$. I point out that the integral for any mode not entering in the function $M(\{a_j, a_j^*\})$ gives unity in the integration due to the normalization of the Wigner function eq.(6). An important consequence of eq.(10) is that normal (antinormal) ordering of creation and annihilation operators in the Hilbert space formalism becomes subtraction (addition) of 1/2 in the WW formalism. The normal ordering rule is equivalent to subtracting the vacuum contribution.

2.3 Evolution

In the Heisenberg picture of the Hilbert-space formalism the density matrix is fixed and any observable, say \hat{M} , evolves according to

$$\frac{d}{dt}\hat{M} = i\left(\hat{H}\hat{M} - \hat{M}\hat{H}\right), \hat{M} = \hat{M}\left(t\right),$$

where \hat{H} is the Hamiltonian. Translated to the WW formalism this evolution of the observables is given by the Moyal equation with the sign changed. The standard Moyal equation applies to the evolution of the Wigner function, that represents a quantum state being the counterpart of the density matrix in the Schrödinger picture of the Hilbert space formalism. Thus in the WW formalism we have

$$\frac{\partial W_{\hat{M}}}{\partial t} = 2\left\{\sin\left[\frac{1}{4}\left(\frac{\partial}{\partial \operatorname{Re}a'_{j}}\frac{\partial}{\partial \operatorname{Im}a''_{j}} - \frac{\partial}{\partial \operatorname{Im}a'_{j}}\frac{\partial}{\partial \operatorname{Re}a''_{j}}\right)\right] \times W_{\hat{M}}\left\{a'_{j}, a^{*'}_{j}, t\right\} H\left(a''_{j}, a^{*''}_{j}\right)_{a_{j}},$$
(11)

where $\{\}_{a_j}$ means making $a'_j = a''_j = a_j$ and $a''_j = a''_j = a''_j$ after performing the derivatives.

A simple example is the free evolution of the field amplitude of a single mode. The Hamiltonian in the WW formalism may be trivially obtained translating the Hamiltonian of the Hilbert-space formalism, that is

$$\hat{H}_{free} = \omega_j \hat{a}_j^{\dagger} \hat{a}_j = \frac{1}{2} \omega_j (\hat{a}_j^{\dagger} \hat{a}_j + \hat{a}_j \hat{a}_j^{\dagger}) - \frac{1}{2} \omega_j$$

$$\rightarrow \quad H_{free} = \omega_j (|a_j|^2 - \frac{1}{2}) = \omega_j \left[(\operatorname{Re}a_j)^2 + (\operatorname{Im}a_j)^2 - \frac{1}{2} \right],$$

where we have taken the first eq.(10) into account. This leads to

$$\frac{d}{dt}a_j = \frac{1}{2}\omega_j \left[2(\operatorname{Im} a_j) - 2\left(\operatorname{Re} a_j\right)i\right] = -i\omega_j a_j \Rightarrow a_j\left(t\right) = a_j\left(0\right)\exp\left(-i\omega_j t\right)$$
(12)

Another example is the down-conversion process in a nonlinear crystal. Avoiding a detailed study of the physics inside the crystal[25], [26] we shall study a single mode problem with the model Hamiltonian[27]

$$\hat{H}_I = A\hat{a}_s^{\dagger}\hat{a}_i^{\dagger}\exp\left(-i\omega_P t\right) + A^*\hat{a}_s\hat{a}_i\exp\left(i\omega_P t\right), \qquad (13)$$

when the laser is treated as classically prescribed, undepleted and spatially uniform field of frequency ω_P . The parameter A is proportional to the pump amplitude and the nonlinear susceptibility. In the WW formalism this Hamiltonian becomes (see eqs.(7))

$$H_I = Aa_s^* a_i^* \exp\left(-i\omega_P t\right) + A^* a_s a_i \exp\left(i\omega_P t\right),$$

whence taking eqs.(11) and (12) into account we have

$$\frac{d}{dt}a_s = -i\omega_s a_s - iAa_i^* \exp\left(-i\omega_P t\right), \qquad (14)$$
$$\frac{d}{dt}a_i = -i\omega_i a_i - iAa_s^* \exp\left(-i\omega_P t\right).$$

We shall assume that the vacuum field a_s evolves as in eq.(12) before entering the crystal and then according to eqs.(14) inside the crystal, that is during the time T needed to cross it. In order to get the radiation intensity to second order in $AT \equiv C$ (see below section 2.4) we must solve these two coupled equations also to second order. After some algebra this leads to

$$a_{s}(t) = \left(1 + \frac{1}{2}|C|^{2}\right)a_{s}(0)\exp\left(-i\omega_{s}t\right) - iCa_{i}^{*}(0)\exp\left[i\left(\omega_{i} - \omega_{P}\right)t\right]$$
$$= \left[\left(1 + \frac{1}{2}|C|^{2}\right)a_{s}(0) - iCa_{i}^{*}(0)\right]\exp\left(-i\omega_{s}t\right),$$
(15)

and the latter equality takes the 'energy conservation' into account (that in the WW formalism looks like a condition of frequency matching, $\omega_P = \omega_s + \omega_i$, with no reference to photon energies).

Eq.(15) gives the time dependence of the relevant mode of signal after crossing the crystal, but we should take account of the field dependence on position including a factor $\exp(i\mathbf{k}_s \cdot \mathbf{r})$, that is a phase depending on the path length. Therefore the correct form of eq.(15) would be, modulo a global phase,

$$a_{s}(\mathbf{r},t) = \left[\left(1 + \frac{1}{2}|C|^{2}\right)a_{s}(0) - iCa_{i}^{*}(0)\right]\exp\left(i\mathbf{k}_{s}\cdot\mathbf{r} - i\omega_{s}t\right).$$
 (16)

A similar result is obtained for $a_i(t)$, that is

$$a_i(\mathbf{r},t) = \left[\left(1 + \frac{1}{2}\left|C\right|^2\right)a_i(0) - iCa_s^*(0)\right]\exp\left(i\mathbf{k}_i\cdot\mathbf{r} - i\omega_i t\right).$$
 (17)

Eq.(16) may be interpreted saying that the vacuum signal is modified by the addition of an amplification of the vacuum idler, but it travels in the same direction of the incoming vacuum signal, and therefore it has sense adding the initial vacuum signal plus the amplification of the idler. And similarly for $a_i \, \text{eq.}(17)$. We may perform a change from C to the new parameter $D = \left(1 + \frac{1}{2} |C|^2\right)^{-1} C$, whence eqs.(9) become, to order $O\left(|D|^2\right)$,

$$E_{s}^{+} = \left(1 + \frac{1}{2}|C|^{2}\right) \left[a_{s} + Da_{i}^{*}\right] \exp\left(i\mathbf{k}_{s}\cdot\mathbf{r} - i\omega_{s}t\right),$$

$$E_{i}^{+} = \left(1 + \frac{1}{2}|C|^{2}\right) \left[a_{i} + Da_{s}^{*}\right] \exp\left(i\mathbf{k}_{i}\cdot\mathbf{r} - i\omega_{i}t\right), |D| \ll 1, \quad (18)$$

and I will ignore the constant global factor $\left(1 + \frac{1}{2} |C|^2\right) \sim 1$ because we will be interested in calculating *relative* detection rates.

Still eqs.(16) and (17), although good enough for calculations are bad representations of the physics. In fact a physical beam corresponds to a superposition of the amplitudes, $a_{\mathbf{k}}^*$, of many modes with frequencies and wavevectors close to ω_s and \mathbf{k}_s , respectively. For instance we may represent the positive frequency part of the idler beam created in the crystal, at first order in D, as follows

$$E_{i}^{(+)}\left(\mathbf{r},t\right) = -iD\int f_{i}\left(\mathbf{k}\right)d^{3}\mathbf{k}a_{\mathbf{k}}^{*}\exp\left[i\left(\mathbf{k}-\mathbf{k}_{s}\right)\cdot\mathbf{r}-i\left(\omega-\omega_{s}\right)t\right] + E_{ZPF}^{(+)},$$
(19)

where $\omega = \omega(\mathbf{k})$ and $f_i(\mathbf{k})$ is an appropriate function, with domain in a region of \mathbf{k} around \mathbf{k}_s . The field $E_{ZPF}^{(+)}$ is the sum of amplitudes of all vacuum modes, including the one represented by a_s in eq.(16). (We have neglected a term of order $|C|^2$ so that $E_i^{(+)}$ is correct to order |C|). These vacuum modes have fluctuating amplitudes with a probability distribution given by the vacuum Wigner function eq.(6). It may appear that the amplitude a_s is lost 'as a needle in the haystack' within the background of many radiation modes, but it is relevant in photon correlation experiments. In fact the vacuum amplitude a_s in eqs.(15) or (16) is fluctuating and the same fluctuations appear also in the signal amplitude a_s^* of eq.(17). Therefore coincidence counts will be favoured when large positive fluctuations of the fields eqs.(15) and (17) arrive simultaneously to Alice and Bob detectors. In the Hilbert-space formamism this fact is named 'entanglement between a signal and the vacuum'. In the WW formalism of quantum optics entanglement appears as a correlation between fields in distant places, including the vacuum fields.

The mentioned properties of the WW formalism are sufficient for the interpretation of experiments involving pure radiation field interacting with macroscopic bodies, these defined by their bulk electric properties like the refraction index or the nonlinear electrical susceptibility. Within the WW formalism the interaction between the fields (either signals or vacuum fields) and macroscopic bodies may be treated as in classical electrodynamics. This is for instance the case for the action of a laser on a crystal with nonlinear susceptibility, studied elsewhere[25], [26].

3 Photon pairs entangled in polarization

In this section I will apply the WW formalism to the description of the polarization correlation of entangled photon pairs produced via spontaneous parametric down-conversion (SPDC). I will assume that the experimental set-up is made so that the fields arriving at the detectors correspond to photon pairs maximally entangled in polarization. These fields are obtained in the outgoing channels of a beam splitter after sending the signal and idler beams produced by SPDC to the incoming channels. The electromagnetic radiation is a vector field with two possible polarizations and I should take into account this fact including vectors in the description. Thus I will write the beams produced as follows

$$\mathbf{E}_{A}^{+} = (a_{s} + Da_{i}^{*}) \exp(-i\omega_{s}t) \mathbf{v} + i (a_{i} + Da_{s}^{*}) \exp(-i\omega_{i}t) \mathbf{h},
\mathbf{E}_{B}^{+} = -i (a_{s} + Da_{i}^{*}) \exp(-i\omega_{s}t) \mathbf{h} + (a_{i} + Da_{s}^{*}) \exp(-i\omega_{i}t) \mathbf{v}, \quad (20)$$

where **h** is a unit vector horizontal and **v** vertical. We have not written explicitly the dependence on position, that could be restored without difficulty, see eq.(19). Furthermore from now on I will ignore all spacetime dependence that usually contributes phase factors irrelevant for our argument. The complex conjugate of the above fields will be labelled as follows

$$(\mathbf{E}_A^+)^* \equiv \mathbf{E}_A^-, (\mathbf{E}_B^+)^* \equiv \mathbf{E}_B^-$$

Eqs.(20) represent "two photons entangled in polarization" as seen in the Weyl-Wigner formalism. The beams will arrive at the Alice and Bob polarization analyzers put at angles θ and ϕ with the vertical respectively. Hence the beams emerging from them will have field amplitudes

$$E_{A}^{+} = (a_{s} + Da_{i}^{*})\cos\theta + i(a_{i} + Da_{s}^{*})\sin\theta, E_{B}^{+} = -i(a_{s} + Da_{i}^{*})\sin\phi + (a_{i} + Da_{s}^{*})\cos\phi,$$
(21)

and polarizations at angles θ and ϕ with the vertical, respectively. For later convenience I define the partial fields

$$E_{A0}^{+} = a_s \cos\theta + ia_i \sin\theta, E_{B0}^{+} = -ia_s \sin\phi + a_i \cos\phi, \qquad (22)$$
$$E_{A1}^{+} = D\left[a_i^* \cos\theta + ia_s^* \sin\theta\right], E_{B1}^{+} = D\left[-ia_i^* \sin\phi + a_s^* \cos\phi\right].$$

Hence we may define intensities as follows

$$I_{A0} = E_{A0}^{+}E_{A0}^{-}, I_{A1} = E_{A0}^{+}E_{A1}^{-} + E_{A1}^{+}E_{A0}^{-} + E_{A1}^{+}E_{A1}^{-},$$

$$I_{B0} = E_{B0}^{+}E_{B0}^{-}, I_{B1} = E_{B0}^{+}E_{B1}^{-} + E_{B1}^{+}E_{B0}^{-} + E_{B1}^{+}E_{B1}^{-},$$

$$I_{A} = E_{A}^{+}E_{A}^{-} = I_{A0} + I_{A1}, I_{B} = E_{B}^{+}E_{B}^{-} = I_{B0} + I_{B1}.$$
(23)

The single, P_A , P_B , and coincidence, P_{AB} , detection rates in the WW formalism may be obtained by comparison with the rates calculated in the Hilbertspace formalism. Thus in the following we revisit briefly the Hilbert-space treatment of the entangled photon pairs. I will start with the quantum fields arriving at Alice and Bob respectively, that are the counterparts of the WW eqs.(21). It is trivial to get them either from eq.(13) or, taking eqs.(7) into account, that is "putting hats" in the WW eqs.(22). We get the field operators

$$\hat{E}_A^+ = \hat{E}_{A0}^+ + \hat{E}_{A1}^+, \hat{E}_B^+ = \hat{E}_{B0}^+ + \hat{E}_{B1}^+,$$

$$\hat{E}_{A0}^{+} = \hat{a}_{s}\cos\theta + i\hat{a}_{i}\sin\theta, \\
\hat{E}_{B0}^{+} = -i\hat{a}_{s}\sin\phi + \hat{a}_{i}\cos\phi, \\
\hat{E}_{A1}^{+} = D[\hat{a}_{i}^{\dagger}\cos\theta + i\hat{a}_{s}^{\dagger}\sin\theta], \\
\hat{E}_{B1}^{+} = D[-i\hat{a}_{i}^{\dagger}\sin\phi + \hat{a}_{s}^{\dagger}\cos\phi], \quad (24)$$

and similar for the Hermitean conjugates. Alice single detection rate is proportional to the following vacuum expectation (with $\hat{E}_A^- = \left(\hat{E}_A^+\right)^{\dagger}$)

$$P_{A} = \langle 0 | \hat{E}_{A}^{-} \hat{E}_{A}^{+} | 0 \rangle = |D|^{2} \langle 0 | \hat{E}_{A1}^{-} \hat{E}_{A1}^{+} | 0 \rangle$$

$$= |D|^{2} \langle 0 | (\hat{a}_{i} \cos \theta - i \hat{a}_{s} \sin \theta) (\hat{a}_{i}^{\dagger} \cos \theta + i \hat{a}_{s}^{\dagger} \sin \theta) | 0 \rangle$$

$$= |D|^{2} \langle 0 | \hat{a}_{i} \hat{a}_{i}^{\dagger} \cos^{2} \theta + \hat{a}_{s} \hat{a}_{s}^{\dagger} \sin^{2} \theta | 0 \rangle = |D|^{2}, \qquad (25)$$

where in the former equality I have neglected creation \hat{E}_{A0}^{-} (annihilation \hat{E}_{A0}^{+}) operators appearing on the left (right). A similar result may be obtained for the single detection rate of Bob, that is

$$P_B = \langle 0 \mid \hat{E}_B^- \hat{E}_B^+ \mid 0 \rangle = |D|^2, \\ \hat{E}_B^- = \left(\hat{E}_B^+\right)^{\dagger}.$$
(26)

We are assuming ideal detectors, but for real detectors P_A and P_B should be multiplied times the detection efficiencies η_A and η_B , and the coincidence rate P_{AB} times $\eta_A \eta_B$.

In order to get the detection rule for single rates in the WW formalism we should translate eq.(25) taking eqs.(10) into account. We get

$$P_A = \left[\left(\left\langle |a_i|^2 \right\rangle - \frac{1}{2} \right) \cos^2 \theta + \left(\left\langle |a_s|^2 \right\rangle - \frac{1}{2} \right) \sin^2 \theta \right] \\ + |D|^2 \left[\left(\left\langle |a_s|^2 \right\rangle + \frac{1}{2} \right) \cos^2 \theta + \left(\left\langle |a_i|^2 \right\rangle + \frac{1}{2} \right) \sin^2 \theta \right] \\ = |D|^2 \left[\cos^2 \theta + \sin^2 \theta \right] = |D|^2, \langle a_i a_s^* \rangle = \langle a_s a_i^* \rangle = 0, \quad (27)$$

that agrees with the result calculated in the Hilbert-space formalism, eq.(??), as it should. Now we compare eq.(27) with the average of the field intensity arriving at Alice (see eq.(21)), that is

$$I_{A} = |E_{A}^{+}|^{2}, \langle I_{A} \rangle = \left\langle |a_{i}|^{2} \right\rangle \cos^{2} \theta + \left\langle |a_{s}|^{2} \right\rangle \sin^{2} \theta + |D|^{2} \left[\left\langle |a_{s}|^{2} \right\rangle \cos^{2} \theta + \left\langle |a_{i}|^{2} \right\rangle \sin^{2} \theta \right] = \frac{1}{2} \left(1 + |D|^{2} \right).$$
(28)

We see that going from eq.(28) to eq.(27) the signal terms (those of order $|D|^2$) are multiplied times 2, whilst those coming from the vacuum (of order unity) are eliminated. This may be seen as a subtraction of the vacuum (ZPF) and multiplication of the signal times 2, which leads to the following rule for the single detection rate in the WW formalism:

$$P_A = 2 \langle I_A \rangle - 2 \langle I_{A0} \rangle, P_B = 2 \langle I_B \rangle - 2 \langle I_{B0} \rangle, \qquad (29)$$

the latter for Bob detection rate.

The Hilbert-space rule for the coincidence rate is the vacuum expectation value of the product of four field operators in normal order. Here we have two terms

$$P_{AB} = \frac{1}{2} \langle 0 \mid \hat{E}_{A}^{-} \hat{E}_{B}^{-} \hat{E}_{B}^{+} \hat{E}_{A}^{+} \mid 0 \rangle + \frac{1}{2} \langle 0 \mid \hat{E}_{B}^{-} \hat{E}_{A}^{-} \hat{E}_{A}^{+} \hat{E}_{B}^{+} \mid 0 \rangle, \qquad (30)$$

that would be equal if \hat{E}_A^+ and \hat{E}_B^+ commute. The former expectation may be evaluated to order $|D|^2$ as follows

$$\begin{aligned} \langle 0 & | \quad \hat{E}_{A}^{-} \hat{E}_{B}^{-} \hat{E}_{B}^{+} \hat{E}_{A}^{+} | 0 \rangle &= \langle 0 | \hat{E}_{A1}^{-} \hat{E}_{B}^{-} \hat{E}_{B}^{+} \hat{E}_{A1}^{+} | 0 \rangle &= \langle 0 | \hat{E}_{A1}^{-} \hat{E}_{B0}^{-} \hat{E}_{B0}^{+} \hat{E}_{A1}^{+} | 0 \rangle \\ &= \langle 0 | \hat{E}_{A1}^{-} \hat{E}_{B0}^{-} | 0 \rangle \langle 0 | \hat{E}_{B0}^{+} \hat{E}_{A1}^{+} | 0 \rangle &= \left| \langle 0 | \hat{E}_{B0}^{+} \hat{E}_{A1}^{+} | 0 \rangle \right|^{2} \\ &= \left| D \right|^{2} \left| \langle 0 | (-i \hat{a}_{s} \sin \phi + \hat{a}_{i} \cos \phi) \left(\hat{a}_{i}^{\dagger} \cos \theta + i \hat{a}_{s}^{\dagger} \sin \theta \right) | 0 \rangle \right|^{2}, \end{aligned}$$

where the former equality, similar to eq.(25), removes creation operators on the left and annihilation operators on the right, the second one removes terms of order $|D|^4$ and the rest is trivial. The latter term of eq.(30) gives a similar contribution so that we get

$$P_{AB} = \frac{1}{2} \left| \langle 0 \mid \hat{E}_{B0}^{+} \hat{E}_{A1}^{+} \mid 0 \rangle \right|^{2} + \frac{1}{2} \left| \langle 0 \mid \hat{E}_{A0}^{+} \hat{E}_{B1}^{+} \mid 0 \rangle \right|^{2} = |D|^{2} \cos^{2}(\theta - \phi).$$
(31)

Here the creation operators are placed to the right and those of annihilation to the left, so that no subtraction is required in passing to the WW formalism. It is enough to substitute c-number amplitudes multiplied times 2 for the field operators, in order to get the rule for the coincidence rate in the WW formalism. That is

$$\left\langle E_{A0}^{+} E_{B1}^{+} \right\rangle + \left\langle E_{A1}^{+} E_{B0}^{+} \right\rangle = \left\langle E_{A}^{+} E_{B}^{+} \right\rangle = D \cos(\theta - \phi)$$

$$\left| \left\langle E_{A}^{+} E_{B}^{+} \right\rangle \right|^{2} = |D|^{2} \cos^{2}(\theta - \phi),$$

$$(32)$$

where we have taken eqs.(22) and (10) into account. Here the vacuum subtraction is not explicit because the vacuum term would be zero, that is $\langle E_{A0}^+ E_{B0}^+ \rangle = 0$.

It is interesting to get the coincidence detection rate in terms of field intensities, rather than amplitudes. To do that we start calculating

$$\langle I_A I_B \rangle = \left\langle E_A^+ E_A^- E_B^+ E_B^- \right\rangle. \tag{33}$$

In the WW formalism the field amplitudes are c-numbers, therefore they commute, and the averages should be performed as in eq.(10). The expectation eq.(??) may be obtained taking into account that the fields have the mathematical properties of Gaussian random variables, see eq.(6) (although this section is devoted to calculations and for the moment I am not committed to any physical interpretation). Thus I apply a well known property of the average of the product of four Gaussian random variables, that is

$$\langle I_A I_B \rangle = \langle E_A^+ E_A^- \rangle \langle E_B^+ E_B^- \rangle + \langle E_A^+ E_B^- \rangle \langle E_A^- E_B^+ \rangle + \langle E_A^+ E_B^+ \rangle \langle E_A^- E_B^- \rangle = \langle I_A \rangle \langle I_B \rangle + |\langle E_A^+ E_B^- \rangle|^2 + |\langle E_A^+ E_B^+ \rangle|^2.$$
 (34)

A similar procedure but involving the vacuum intensities, gives

$$\langle I_{A0}I_{B0}\rangle = \langle I_{A0}\rangle \langle I_{B0}\rangle + \left| \left\langle E_{A0}^+ E_{B0}^- \right\rangle \right|^2 + \left| \left\langle E_{A0}^+ E_{B0}^+ \right\rangle \right|^2.$$
(35)

Here the third term does not contribute and the second one equals the second term of eq.(34) to order $|D|^2$. Hence we get the rule for the coincidence rate in the WW formalism that I write both in terms of fields and in terms of intensities as follows

$$P_{AB} = \left| \left\langle E_A^+ E_B^+ \right\rangle \right|^2 = \left\langle I_A I_B \right\rangle - \left\langle I_A \right\rangle \left\langle I_B \right\rangle - \left\langle I_{A0} I_{B0} \right\rangle + \left\langle I_{A0} \right\rangle \left\langle I_{B0} \right\rangle.$$
(36)

4 Locality in the experiments with entangled photon pairs

4.1 A realistic interpretation of the experiments

I emphasize again that the WW formalism provides an alternative formulation of quantum optics, physically equivalent to the more common the Hilbert-space formalism. But it suggests a picture of the optical phenomena quite different from the latter. Indeed the picture may provide a local realistic interpretation in terms of random variables and stochastic processes. In the following I present the main ideas of this stochastic interpretation. It rests upon several assumptions as follows.

The fundamental hypothesis is that the electromagnetic vacuum field is a real stochastic field (the zeropoint field, ZPF). If expanded in normal modes the ZPF has a (positive) probability distribution of the amplitudes given by eq.(6). Therefore we assume that all bodies are immersed in ZPF absorbing and emitting radiation continuously. In particular the ground state of an atom corresponds to a dynamical equilibrium with the ZPF. The hypothesis that vacuum fields are real stochastic fields leads to a general interpretation of quantum theory that has been sketched elsewhere[28].

According to that assumption any photodetector would be immersed in an extremely strong radiation, infinite if no cut-off existed. Thus how might we explain that detectors are not activated by the vacuum radiation? Firstly the strong vacuum field is effectively reduced to a weaker level if we assume that only radiation within some (small) frequency interval is able to activate a photodetector, that is the interval of sensibility (ω_1, ω_2) . However the problem is not vet solved because signals involved in experiments have typical intensities of order the vacuum radiation in the said frequency interval so that the detector would be unable to distinguish a signal from the ZPF noise. Our proposal is to assume that a detector may be activated only when the net Poynting vector (i. e. the directional energy flux) of the incoming radiation is different from zero, including both signal and vacuum fields. More specifically I will assume that the detector possesses an active area and the probability of a photocount is proportional to the net radiant energy flux crossing that area from the front side during the activation time, T, the probability being zero if the net flux crosing the area is in the reverse direction.

These assumptions allow to understand qualitatively why the signals, but not the vacuum fields, activate detectors. Indeed the ZPF arriving at any point (in particular the detector) would be isotropic on the average, whence its associated mean Poynting vector would be nil, therefore only the signal radiation should produce photocounts. A problem remains because the vacuum fields are fluctuating so that the net Poynting vector also fluctuates. These problems diminish due to the fact that *photocounts are not produced by an instantaneous interaction of the fields with the detectors but the activation requires some time interval*, a known fact in experiments. This would reduce the effect of the vacuum fluctuations.

Our aim is to achieve a realistic local interpretation of the experiments measuring polarization correlation of entangled photon pairs, that we studied with the WW formalism in the previous section. Thus I will consider two vacuum beams entering the nonlinear crystal, where they give rise to a "signal" and an "idler" beams. After crossing several appropriate devices they produce fields that will arrive at the Alice and Bob detectors. I do not attempt to present a detailed model that should involve many modes in order to represent the signals as (narrow) beams, see eq.(19).

In agreement with our previous hypotheses a photodetection should derive from the net energy flux crossing the active photocounter surface. Thus I will assume that the detection probabilities per time window, that is the single P_A, P_B and coincidence P_{AB} detection rates, are

$$P_{A} = \langle [M_{A}]_{+} \rangle, P_{B} = \langle [M_{A}]_{+} \rangle, P_{AB} = \langle [M_{A}]_{+} [M_{B}]_{+} \rangle,$$

$$M_{A} \equiv T^{-1} \int_{0}^{T} \vec{n}_{A} \cdot \vec{I}_{total}^{A} (\mathbf{r}_{A}, t) dt,$$

$$M_{B} \equiv T^{-1} \int_{0}^{T} \vec{n}_{B} \cdot \vec{I}_{total}^{B} (\mathbf{r}_{A}, t) dt,$$
(37)

where $[M]_+ = M$ if M > 0, $[M]_+ = 0$ otherwise, and \vec{n} is a unit vector in the direction of the incoming signal beams, that I assume perpendicular to the active area of the detector. I use units such that both the intensities and the detection rates are dimensionless (as explained above the rates are defined as probabilities of detection within a detection time window T). In eq.(37) the positivity constraint (i.e. putting $[M_A]_+$ rather than M_A) is needed because the detection probabilities must be non-negative for any particular run of an experiment whilst the quantities M_A and M_B are fluctuating and may be negative sometimes. Nevertheless the ensemble averages involved in eq.(37) are positive or zero and the fluctuations will not be too relevant due to the time integration that washes them out. Therefore I will ignore the positivity constraint substituting M_A for $[M_A]_+$ and M_B for $[M_B]_+$ in the following.

In order to have a realistic model of the experiments I will consider a simplified treatement involving just two vacuum radiation modes, with amplitudes a_s and a_i , as in the WW calculation of the previous section. After crossing several appropriate devices the fields will arrive at the Alice and Bob detectors. Each one of these two fields consists of two parts, one of order 0 and another of order |D| << 1. The former, given in eqs.(22), is what would arrive at the detectors if there was no pumping laser and therefore no signal. It is just a part of the ZPF, whilst the rest of the ZPF consists of radiation not appearing in the equations of the previous section because they were not needed in the calculations. The total ZPF should have the property of isotropy, therefore giving nil net flux in the detector (modulo fluctuations that we shall ignore at that moment). The terms of order |D|, given by eqs.(22) correspond to the signal produced in the nonlinear crystal after the modifications introduced by beam splitters and polarizers (and other devices like apertures, filters, lens systems, etc. whose action is not detailed in our simplified model). In summary the Poynting vectors of the radiation at the (center of the) active area of the detectors may be written

$$Alice : \vec{I}_{total}^{A}(t) = \vec{I}_{ZPF}^{A}(t) + \vec{I}_{A}(t), Bob : \vec{I}_{btal}^{B}(t) = \vec{I}_{ZPF}^{B}(t) + \vec{I}_{B}(t).$$
(38)

 \vec{I}_A , \vec{I}_B , are due to the fields E_A , E_B , eqs.(21), coming from the fields emerging from the non-linear crystal. The Poynting vectors $\vec{I}_A(t)$ and $\vec{I}_B(t)$ have the

direction of \vec{n}_A and \vec{n}_B respectively, see eqs.(37), and their moduli would be the field intensities. Furthermore these intensities are time independent (in our representation of the fields), so that we may write

$$M_A = I_{ZPF}^A + I_A, I_{ZPF}^A \equiv T^{-1} \int_0^T \vec{n}_A \cdot \vec{I}_{ZPF}^A \left(\mathbf{r}_A, t \right) dt$$
(39)

and similar for M_B . In order to get the Alice single detection rate we need the average of M_A , that we will evaluate by comparison with the case when there is no pumping on the nonlinear crystal. In this case I_A becomes the intensity I_{A0} and the Poynting vector of all vacuum fields arriving at the detector of Alice, i. e. $\vec{I}_{ZPF}^A(t) + \vec{I}_{A0}(t)$, should have nil average due to the isotropy of the ZPF. And similar for Bob. As a consequence the intensities I_{A0} and I_{B0} , eqs.(23), should fulfil the following equalities

$$\left\langle I_{ZPF}^{A} + I_{A0} \right\rangle = \left\langle I_{ZPF}^{B} + I_{B0} \right\rangle = 0.$$

$$\tag{40}$$

It would appear that this relation could not be true for all values of the angles θ, ϕ eqs.(22) because the ZPF Poynting vector \hat{I}_{ZPF}^A and \hat{I}_{ZPF}^B should not depend on our choice of angles whilst I_{A0} and I_{B0} do depend. However the positions of the polarizers may influence also the ZPZ arriving at the detectors and it is plausible that the total Poynting vector has always zero mean. From eqs.(39) and (40) we may derive the single rates of Alice and Bob, that is

$$P_A = \langle M_A \rangle = \langle I_A \rangle - \langle I_{A0} \rangle = \frac{1}{2} \left| D \right|^2, P_B = \langle M_B \rangle = \frac{1}{2} \left| D \right|^2.$$
(41)

The coincidence detection rate may be got taking eq.(37) into account, that is

$$P_{AB} = \langle M_A M_B \rangle = \langle \left[I_{ZPF}^A + I_A \right] \left[I_{ZPF}^B + I_B \right] \rangle.$$
(42)

For the sake of clarity the calculation of this average is made in some detail as follows. I will start deriving the joint probability distribution of M_A and M_B from the distributions of the amplitudes of normal modes, eq.(6), that is

$$\rho\left(M_A, M_B\right) = \int W_0 \prod_j da_j da_j^* \delta\left(M_A - I_{ZPF}^A - I_A\right) \delta\left(M_B - I_{ZPF}^B - I_B\right)$$

$$\tag{43}$$

where δ () is Dirac's delta and I_A, I_B are functions of the two modes involved, i. e. those with amplitudes a_s, a_i , see eqs.(21). Hence the expectation eq.(42) becomes

$$P_{AB} = \langle M_A M_B \rangle = \int M_A M_B \rho \left(M_A, M_B \right) dM_A dM_B$$

$$= \int W_0 \prod_j da_j da_j^* \left[I_{ZPF}^A + I_A \right] \left[I_{ZPF}^B + I_B \right]$$

$$\equiv \left\langle \left[I_{ZPF}^A + I_A \right] \left[I_{ZPF}^B + I_B \right] \right\rangle, \qquad (44)$$

where we have performed the M_A and M_B integrals in the former equality.

If there was no pumping laser on the nonlinear crystal the joint detection rate should be zero whence eq.(44) leads to

$$\left\langle \left[I_{ZPF}^{A} + I_{A0}\right] \left[I_{ZPF}^{B} + I_{B0}\right] \right\rangle = 0.$$

$$(45)$$

From eqs.(40), (44) and (45) it is possible to get P_{AB} . Firstly I point out that I_{ZPF}^A and I_{ZPF}^B could not depend on whether the pumping is switched on or out. More specifically the probability of the possible values of I_{ZPF} would be the same whether the pumping is on or out. Then subtracting eq.(45) from eq.(44) we get

$$P_{AB} = \langle (I_B - I_{B0}) I_{ZPF}^A \rangle + \langle (I_A - I_{A0}) I_{ZPF}^B \rangle + \langle I_A I_B \rangle - \langle I_{A0} I_{B0} \rangle$$

$$= \langle I_{B1} I_{ZPF}^A \rangle + \langle I_{A1} I_{ZPF}^B \rangle + \langle I_A I_B \rangle - \langle I_{A0} I_{B0} \rangle,$$

where we have taken into account the definition of I_{A1} and I_{B1} , eq.(23). The ZPF intensity I_{ZPF}^A should be independent of whether I_{B1} is zero (pumping out) or finite (pumping on) as commented above, therefore I_{ZPF}^A and I_{B1} are uncorrelated. Similarly for I_{ZPF}^B and I_{A1} . Hence we may write, taking eq.(40) into account,

$$P_{AB} = \langle I_{B1} \rangle \langle I_{ZPF}^{A} \rangle + \langle I_{A1} \rangle \langle I_{ZPF}^{B} \rangle + \langle I_{A}I_{B} \rangle - \langle I_{A0}I_{B0} \rangle$$

$$= \langle I_{A}I_{B} \rangle - \langle I_{A1} \rangle \langle I_{B0} \rangle - \langle I_{B1} \rangle \langle I_{A0} \rangle - \langle I_{A0}I_{B0} \rangle$$

$$= \langle I_{A0}I_{B1} \rangle + \langle I_{A1}I_{B0} \rangle - \langle I_{A1} \rangle \langle I_{B0} \rangle - \langle I_{B1} \rangle \langle I_{A0} \rangle.$$
(46)

Now we may write $\langle I_{A0}I_{B1}\rangle + \langle I_{A1}I_{B0}\rangle$ in terms of the fields taking into account the property of Gaussian random variables used in the evaluation of eq.(33), that is

$$\langle I_{A0}I_{B1} \rangle = \langle E_{A0}^{+}E_{A0}^{-} \rangle \langle E_{B1}^{+}E_{B1}^{-} \rangle + \langle E_{A0}^{+}E_{B1}^{-} \rangle \langle E_{A0}^{-}E_{B1}^{+} \rangle + \langle E_{A0}^{+}E_{B1}^{+} \rangle \langle E_{A0}^{-}E_{B1}^{-} \rangle$$

$$= \langle I_{A0} \rangle \langle I_{B1} \rangle + |\langle E_{A0}^{+}E_{B1}^{-} \rangle|^{2} + |\langle E_{A0}^{+}E_{B1}^{+} \rangle|^{2} ,$$

and similar for $\langle I_{A1}I_{B0}\rangle$. Then eq.(46) becomes

$$P_{AB} = \left| \left\langle E_{A0}^{+} E_{B1}^{+} \right\rangle \right|^{2} + \left| \left\langle E_{A1}^{+} E_{B0}^{+} \right\rangle \right|^{2} + \left| \left\langle E_{A0}^{+} E_{B1}^{-} \right\rangle \right|^{2} + \left| \left\langle E_{A1}^{+} E_{B0}^{-} \right\rangle \right|^{2}.$$

The two latter terms are nil and the former two give, taking eqs.(22) into account,

$$P_{AB} = \frac{1}{2} \left| D \right|^2 \cos^2(\theta - \phi).$$
(47)

As a conclusion the results of our realistic model, eqs.(41) and (47), agree with the quantum predictions eqs.(25), (26) and (31), modulo an scaling parameter 1/2.

Our model, resting upon the WW formalism of quantum optics, provides a picture quite different from the one suggested by the Hilbert-space in terms of photons. We do not assume that the photocount probability within a time window is the product of a *small* probability, of order $|D|^2$, that the pumping laser

produces in the crystal an "entangled photon pair" times a detection probability of order *unity* conditional to the photon production. The latter probability is defined as the detection efficiency (neglecting small losses). In our model the probabilities of photocounts do not factorize that way. Furthermore the concept of photon does not appear at all, but there are *continuous fluctuating fields including a real ZPF* arriving at the detectors that are activated when the fluctuations are big enough.

It is interesting to study more closely the "quantum correlation" qualified as strange from a classical point of view because it is a consequence of the phenomenon of entanglement. The origin is the correlation between the signal I_{B1} produced in the crystal and the part I_{A0} of the ZPF, that is essential for the large value of the coincidence rate eq.(46). It is remarkable that this correlation derives from the fact that the same normal mode appears in both radiation fields, E_{A0}^+ and E_{B1}^+ see eq.(32), received by Alice and Bob respectively. And similarly for E_{A1}^+ and E_{B0}^+ . Also with reference to eqs.(22) and (29) we see that if I_{A0} – $\langle I_{A0} \rangle$ is positive for some value of the random variable I_{A0} then I_{B1} is large, but if $I_{A0} - \langle I_{A0} \rangle$ is negative then I_{B1} is small. This argument explains intuitively why the product is positive on the average, that is proved quantitatively by eq.(46). Furthermore I stress that the ensemble average of $I_{A0} - \langle I_{A0} \rangle$ is zero, meaning that only the fluctuations are involved in the enhancement of detection probability by Bob due to the intensity I_{B1} . And similarly for the fluctuations of (Bob) term $I_{B0} - \langle I_{B0} \rangle$ that enhance the detection probability of Alice due to I_{A1} . This leads to an interesting interpretation of entanglement: it is a correlation between fluctuations involving the vacuum fields.

Finally I stress that the hypothesis that the quantum vacuum fields are real allows a concept of locality weaker than Bell's. Indeed the signal fields (accompanied by vacuum fields) travel causally from the source (the laser pumping beam and the nonlinear crystal folowed by a beam-splitter and other devices) to the detectors. Thus I claim that the model of this paper is local. On the other hand the results for the single and coincidence detection probabilities within a time window, eqs.(41) and (47), violate a Clauser-Horne inequality ([6]) as may be easily checked. Therefore it is possible a definition of locality, weaker than Bell's, able to explain the entangled photon experiments without the need of assuming the violation of relativistic causality.

References

- L. K. Shalm et al., A strong loophole-free test of local realism. *Phys. Rev. Lett.* 115, 250402 (2015).
- [2] M. Giustina et al., A significant loophole-free test of Bell's theorem with entangled photons. *Phys. Rev. Lett.* **115**, 250401 (2015).
- [3] H.Wiseman, Nature **526**, 687 (2015).

- [4] E. Santos, Mathematical and physical meaning of the Bell inequalities. *Eur. J. Phys.* 37, 055402 (2016).
- [5] J. S. Bell. On the Einstein, Podolsky and Rosen paradox. *Physics*, 1, 195 (1964).
- [6] J. F. Clauser, M. Horne. Experimental consequences of objective local theories. *Phys. Rev. D* 10, 526-535 (1974).
- [7] P. H. Eberhard. Background level and counter efficiencies required for a loophole-free Einstein-Podolsky-Rosen experiment. *Phys. Rev. A* 47, R747, R749 (1993).
- [8] A. Casado, T. W. Marshall, E. Santos. Parametric downconversion experiments in the Wigner representation. J. Opt. Soc. Amer. B 14, 494-502 (1997).
- [9] A. Casado, A. Fernández-Rueda, T. W. Marshall, R. Risco-Delgado, E. Santos. Fourth order interference in the Wigner representation for parametric downconversion experiments. *Phys. Rev. A* 55, 3879-3890 (1997).
- [10] A. Casado, A. Fernández-Rueda, T. W. Marshall, R. Risco-Delgado, E. Santos. Dispersion cancellation and quantum eraser experiments analyzed in the Wigner function formalism. *Phys. Rev. A* 56, 2477-2480 (1997).
- [11] A. Casado, T. W. Marshall, E. Santos. Type II parametric downconversion in the Wigner formalism. Entanglement and Bell's inequalities. J. Opt. Soc. Amer. B 15, 1572-1577 (1998).
- [12] A. Casado, T. W. Marshall, E.Santos. Parametric down and up conversion in the Wigner representation of quantum optics. J. Mod. Opt. 47, 1273-1287 (1999).
- [13] A. Casado, T. W. Marshall, R. Risco-Delgado, E. Santos. Spectrum of the parametric down converted radiation calculated in the Wigner function formalism. *Eur. Phys. J. D* 13, 109-119 (2001).
- [14] A. Casado, S. Guerra, J. Plácido. Wigner representation for experiments on quantum cryptography using two-photon polarization entanglement produced in parametric down-conversion. J. Phys. B: At. Mol. Opt. Phys. 41, 045501 (2008).
- [15] A. Casado, S. Guerra, J. Plácido. Partial Bell-state analysis with parametric down conversion in the Wigner function formalism. Adv. Math. Phys. 2010, 501521 (2010).
- [16] A. Casado, S. Guerra, J. Plácido. Wigner representation for polarization momentum hyperentanglement generated in parametric down-conversion, and its application to complete Bell-state measurement. *Eur. Phys. J. D* 68, 338-348 (2014).

- [17] A. Casado, S. Guerra, J. Plácido. Wigner representation for entanglement swapping using parametric down conversion: the role of vacuum fluctuations in teleportation. J. Mod. Opt. 62, 377-386 (2015).
- [18] A. Casado, S. Guerra, J. Plácido. Rome teleportation experiment analysed in the Wigner representation: the role of the zeropoint fluctuations in complete one-photon polarization-momentum Bell-state analysis. J. Mod. Opt. 65, 1960-1974 (2018).
- [19] H. Weyl, Z. Phys. 46, 1 (1927). H. Weyl. The Theory of Groups and Quantum Mechanics. Dover, New York, 1931. (German original, 1928).
- [20] E. P.Wigner. On the quantum correction for thermodynamic equilibrium. *Phys. Rev.* 40, 749 (1932).
- [21] H. J. Groenewold. On the principles of elementary quantum mechanics. *Physica* 12, 405-446 (1946).
- [22] J. E. Moyal. Quantum mechanics as a statistical theory. Proc. Cambridge Phil. Soc. 45, 99-124 (1949).
- [23] M. Hillery, R. F. O'Connell, M. O. Scully and E. P. Wigner. *Phys. Rep.* 106, 121-168 (1984).
- [24] C. K. Zachos, D. B. Fairlie, T. L. Curtright. Quantum Mechanics in Phase Space. World Scientic, Singapore, 2005.
- [25] K. Dechoum, T. W. Marshall, E. Santos. Parametric up and down conversion in the Wigner representation of quantum optics. J. Mod. Opt. 47, 1273-1287 (2000).
- [26] A. Casado, A. Fernández-Rueda, T. W. Marshall, J. Martínez, R. Risco-Delgado, E. Santos. Dependence on crystal parameters of the correlation time between signal and idler beams in parametric down conversion calculated in the Wigner representation. *Eur. Phys. J. D* 11, 465-472 (2000).
- [27] Z. Y. Ou, L. J. Wang, L. Mandel. Evidence for phase memory in two-photon down conversion through entanglement with the vacuum. *Phys. Rev. A* 41, 566 (1990).
- [28] E. Santos. Towards a realistic interpretation of quantum mechanics providing a model of the physical world. *Foundations of Science*, **20**, 357-386 (2015), arXiv: 1203.5688 (2012).