THE FIRST DETECTION OF ¹³C¹⁷O IN A PROTOPLANETARY DISK: A ROBUST TRACER OF DISK GAS MASS

Alice S. Booth,¹ Catherine Walsh,¹ John D. Ilee,¹ Shota Notsu,^{2,3} Chunhua Qi,⁴ Hideko Nomura,^{5,6} and Eiji Akiyama⁷

¹School of Physics and Astronomy, University of Leeds, Leeds LS2 9JT, UK

²Department of Astronomy, Graduate School of Science, Kyoto University, Kitashirakawa-Oiwake-cho, Sakyo-ku, Kyoto 606-8502, Japan

³ Leiden Observatory, Leiden University, P.O. Box 9513, NL-2300 RA, Leiden, the Netherlands

⁴Harvard-Smithsonian Center for Astrophysics, Cambridge, MA 02138, USA

⁵Department of Earth and Planetary Science, Tokyo Institute of Technology, 2-12-1 Ookayama, Meguro-ku, Tokyo 152-8551, Japan

⁶National Astronomical Observatory Japan (NAOJ), Osawa 2-21-1, Mitaka, Tokyo 181-8588, Japan

⁷ Institute for the Advancement of Higher Education, Hokkaido University, Kita 17, Nishi 8, Kita-ku, Sapporo, Hokkaido 060-0817, Japan

(Received June 14, 2019; Revised July 19, 2019; Accepted July 27, 2019)

ABSTRACT

Measurements of the gas mass are necessary to determine the planet formation potential of protoplanetary disks. Observations of rare CO isotopologues are typically used to determine disk gas masses; however, if the line emission is optically thick this will result in an underestimated disk mass. With ALMA we have detected the rarest stable CO isotopologue, ${}^{13}C^{17}O$, in a protoplanetary disk for the first time. We compare our observations with the existing detections of ${}^{12}CO$, ${}^{13}CO$, $C^{18}O$ and $C^{17}O$ in the HD 163296 disk. Radiative transfer modelling using a previously benchmarked model, and assuming interstellar isotopic abundances, significantly underestimates the integrated intensity of the ${}^{13}C^{17}O$ J=3-2 line. Reconciliation between the observations and the model requires a global increase in CO gas mass by a factor of 3.5. This is a factor of 2-6 larger than previous gas mass estimates using $C^{18}O$. We find that $C^{18}O$ emission is optically thick within the snow line, while the ${}^{13}C^{17}O$ emission is optically thick within the snow line, while the ${}^{13}C^{17}O$ emission is optically thick disk CO gas mass.

Keywords: astrochemistry—planets and satellites: formation—protoplanetary disks—submillimeter: planetary systems

1. INTRODUCTION

The mass of a disk sets a limit on the material available for forming a planetary system and can influence the mode of giant planet formation. Most disk gas masses rely on observations of CO that are extrapolated to a total gas mass by assuming a constant CO/H_2 abundance ratio in the disk. However, if the line emission is optically thick this will result in an underestimated disk mass (e.g., Bergin & Williams 2017). Spatially resolved ¹³C¹⁶O and ¹²C¹⁸O line emission are now commonly detected in protoplanetary disks and have been used to determine disk gas masses (e.g., Ansdell et al. 2016; Long et al. 2017). The second-most rarest CO isotopologue, ¹³C¹⁸O, has been detected in the TW Hya disk and this emission has been proposed to be optically thin and traces the disk midplane whereas the $C^{18}O$ emission is optically thick within the midplane snowline (Zhang et al. 2017). The under-estimation of disk gas mass due to optically thick emission will be more significant in more massive gas-rich disks, i.e., those around Herbig Ae/Be stars versus those around T Tauri stars.

Low CO gas masses, with respect to the dust mass, have been consistently measured in disks and complementary HD observations imply that this is because of the depletion of gas-phase CO in disks relative to that in the ISM (Bergin et al. 2013; McClure et al. 2016). CO can be depleted from the gas phase via freeze out onto the icy grains in the cold midplane, and subsequent conversion to CO_2 and more complex organic species e.g. CH_3OH (e.g. Bosman et al. 2018). Additionally, photodissociation via far-UV radiation destroys CO in the upper disk atmosphere, and isotope selective photodissociation can enhance the various isotopologue ratios relative to ${}^{12}C^{16}O$ in the atmosphere (e.g. Miotello et al. 2014). These chemical effects make the conversion from CO gas mass to total gas mass a non-trivial task.

The protoplanetary disk around HD 163296 has been well characterised with the Atacama Large Millimeter/submillimeter Array (ALMA). Band 6 and 7 observations show rings in both the continuum and the CO gas emission (e.g. Isella et al. 2016; Notsu et al. 2019, and see Figure 1a). There are four proposed ≈ 0.5 - $2 M_J$ planets in this disk inferred from the dust and gas rings, and deviations from Keplerian motion in the CO gas kinematics (Isella et al. 2016; Pinte et al. 2018; Teague et al. 2018; Liu et al. 2018). Recently, ≈ 5 au resolution observations of the continuum emission revealed an additional gap and ring in the inner disk as well as an azimuthal asymmetry in one of the previously detected rings (see Figure 1a, Isella et al. 2018). Hence, the proposed planet-induced structures in the HD 163296 disk make it an excellent observational laboratory to study planet formation.

We present the first detection of ${}^{13}C{}^{17}O$ in a protoplanetary disk providing a strong constraint on the CO gas mass in the HD 163296 disk.

2. OBSERVATIONS

HD 163296 was observed with ALMA in Band 7 during Cycle 3 on 2016 September 16 (2015.1.01259.S, PI: S. Notsu). See Notsu et al. (2019) for details on the calibration and self-calibration of the data. The spectral windows have a resolution of 1953.125 kHz. There are $14 {}^{13}C^{17}O$ J=3-2 hyperfine structure lines that lie between 321.851 and 321.852 GHz. All lines lie within a frequency range less than the spectral resolution of the data; hence, we are observing the blending of all of the lines. The ${}^{13}C^{17}O$ molecular data we use is from Klapper et al. (2003) and was accessed via the Cologne Database for Molecular Spectroscopy (CDMS, Müller et al. 2005). We detected ${}^{13}C^{17}O$ initially via a matched filter analysis¹ (Loomis et al. 2018) using a Keplerian mask assuming a disk position angle of 132° and an inclination of 42° (e.g. Isella et al. 2016). The resulting S/N ratio is ≈ 3.5 . The filter response is shown in Figure 1b with the black line marking the ${}^{13}C^{17}O$ J=3-2 transition after correction for the source velocity (5.8 km s⁻¹).

The line imaging was conducted using CLEAN with CASA version 4.6.0. The native spectral resolution of the data is 1.8 km s⁻¹; however, in order to optimise the S/N the final images were generated with a

3 km s⁻¹ channel width and a uv taper of 0.5 resulting in a synthesised beam of 0.87 x 0.51 (100°). Figures 1c and 1d present the ¹³C¹⁷O integrated intensity map and the intensity-weighted velocity map, respectively. The integrated intensity map was made using channels \pm 6 km s⁻¹ about the source velocity. The peak integrated intensity is 0.55 Jy beam⁻¹ km s⁻¹ with an rms noise level of 0.08 Jy beam⁻¹ km s⁻¹ (S/N = 7), that was extracted from the spatial region beyond the detected line emission. The intensity-weighted velocity map was made in the same manner but also with a 3 σ clip.

We also use archival data to benchmark our modelling, including the ${}^{12}C^{16}O$, ${}^{13}C^{16}O$, ${}^{12}C^{18}O$ J=2-1 transitions observed from Isella et al. (2016), and the ${}^{12}C^{16}O$ J=3-2 ALMA Science Verification data.² All integrated intensity maps were de-projected and azimuthally averaged and are shown in Figures 3a to 3e. The errors are the standard deviation of intensity of the pixels in each bin divided by the number of beams per annulus (e.g. Carney et al. 2018). In addition, Figure 3e shows the $^{12}C^{17}O$ J=3-2 total integrated intensity value with its associated errors (Qi et al. 2011). All data plotted assumes a source distance of 122 pc (van den Ancker et al. 1998). Although GAIA DR2 puts this source at 101.5 pc (Gaia Collaboration et al. 2018), in order to compare to previous analyses we use the previous value. We discuss the impact of the revised distance in Section 4.

3. ANALYSIS

Previous observations of the HD 163296 disk with the SMA and ALMA have detected multiple CO isotopologues: ${}^{12}C^{16}O$, ${}^{13}C^{16}O$, ${}^{12}C^{18}O$ and ${}^{12}C^{17}O$ (Qi et al. 2011; Isella et al. 2016). The models that were used to reproduce the line emission in Qi et al. (2011) recover the following global isotope ratios;

$$n({}^{12}C{}^{16}O)/n({}^{13}C{}^{16}O) = 67 \pm 8,$$

 $n({}^{12}C{}^{16}O)/n({}^{12}C{}^{18}O) = 444 \pm 88,$
 $n({}^{12}C{}^{18}O)/n({}^{12}C{}^{17}O) = 3.8 \pm 1.7,$

where $n({}^{X}C{}^{Y}O)$ is the number density of the molecule. This is consistent with the carbon and oxygen isotope ratios observed in the ISM (Wilson 1999).

We make a first estimate of the column density of gas traced by the ¹³C¹⁷O emission under the assumption of optically thin emission in local thermodynamic equilibrium. Following Carney et al. 2019 (their Equation 1, with molecular data obtained from CDMS, Müller et al.

 $^{^1}$ A python-based open-source implementation of VISIBLE is available at http://github.com/AstroChem/VISIBLE

² https://almascience.nrao.edu/alma-data/science-verification




Figure 1. a) The 1.25 mm continuum image from Isella et al. (2018). b) The matched filter response for the ¹³C¹⁷O J=3-2 detection where the black line marks the frequency of the hyper-fine transitions. c) The ¹³C¹⁷O J=3-2 integrated intensity map where the white dashed contours mark 3 and 5 σ . d) The ¹³C¹⁷O J=3-2 intensity-weighted velocity map.

2005), the average column density for the ${}^{13}C{}^{17}O$ within 50 au, assuming an excitation temperature of 50 K, is $7.1 \times 10^{15} \text{ cm}^{-2}$. This is equivalent to a $n_{\rm H}$ column density of $2.65 \times 10^{25} \text{ cm}^{-2}$ (44.4 g cm⁻²) at 50 au. In comparison, the corresponding value for the ${}^{12}C{}^{18}O$ is $1.7 \times 10^{16} \text{ cm}^{-2}$, resulting in a $n({}^{12}C{}^{18}O)/n({}^{13}C{}^{17}O)$ ratio of 2.5. Under the assumption that both the lines are optically thin (and taking the previously derived isotopic ratios), this value is a factor of 100 too small. Therefore, the ${}^{12}C{}^{18}O$ line emission is optically thick and the resulting gas mass derived from this tracer will be underestimated.

To quantify this more robustly, we utilise an existing disk model that has been shown to fit emission lines from multiple CO isotopologues (${}^{12}C^{16}O$, ${}^{13}C^{16}O$, ${}^{12}C^{18}O$ and ${}^{12}C^{17}O$) to model our ${}^{13}C^{17}O$ detection (Qi et al. 2011). The density (hydrogen nuclei density, $n_{\rm H}$) and temperature of the disk are shown in Figures 2a and 2b. The CO abundance distribution, shown in Figure 2c, was determined by setting n(CO) to a constant fractional abundance of 6.0×10^{-5} with respect to H₂ in the molecular layer following Qi et al. (2011). This assumes that 86% of volatile carbon inherited by the disk is in the form of CO (Graedel et al. 1982). This abundance was reduced by a factor of 10^{-4} in the midplane where T_{gas} ≤ 19 K and by a factor of 10^{-8} in the atmosphere where the vertically-integrated hydrogen column density, $\sigma(n_{\rm H})$, from the disk surface is $< 1.256 \times 10^{21}$ cm⁻². The depleted value in the midplane is consistent with the CO abundances derived from chemical models including non-thermal desorption (Walsh et al. 2010). The photodissociation and freeze out boundaries are shown in white contours overlaid on Figures 2a and 2b.

The first model that we test, Model 1, uses a constant $^{13}C^{17}O$ fractional abundance of 5.39 \times 10⁻¹⁰ relative to H_2 . This assumes isotope ratios that are consistent with the observations and modelling from Qi et al. (2011). Model 1 has a total disk mass of 0.089 M_{\odot} . Using the CDMS data for ¹³C¹⁷O we generated a LAMDA-like file in order to model the J=3-2 hyper-fine components in LIME³ (the Line Modeling Engine, Brinch & Hogerheijde 2010). Synthetic images cubes were computed assuming the appropriate position angle and inclination of the source, and the resulting images were smoothed with a Gaussian beam to the spatial resolution of the observations using the CASA task, IMSMOOTH. The generated integrated intensity map was then de-projected and azimuthally averaged. The radial profiles from Model 1 (orange) are shown alongside the observations in Figures 3a to 3e.

Model 1 under-predicts the ${}^{13}C{}^{17}O$ peak emission in the integrated intensity map by a factor of 2.5, yet provides a reasonable fit to the other lines (within a factor of two). The higher spatial resolution observations are effected by dust opacity within ≈ 50 au (see Isella et al. 2016); therefore we focus on reproducing the data beyond 50 au.

To better fit the ${}^{13}C{}^{17}O$ observations we globally increase the gas mass of Model 1. This was done by initially multiplying $n_{\rm H}$ by a factor of 1.5 and then increasing this factor in steps of 0.5 until the best by-eye fit of 3.5 was found. The results for Model 2 are shown in Figure 3 (purple). Model 2 provides a better fit to all of the lines. This model assumes a smooth radial gas density structure contrary to the most recent observations. However, our work is focused on reproducing the global disk mass rather than the underlying small scale gas surface density variations. Model 2 has a total disk mass of 0.31 M_{\odot} .

We note that a similar fit can be obtained using a different CO snowline location at 90 au as determined in Qi et al. (2015). This requires a corresponding increase in gas mass (\times 3.5) within the snowline, and we obtain a similar ¹²C¹⁸O column density profile as in Qi et al. (2015) beyond the snowline. Both of these models use the same underlying physical structure but have different CO snowline locations and levels of CO depletion beyond the snowline. The Qi et al. (2011) model has simpler assumptions regarding the freeze-out of CO, consistent with other work (e.g., Williams J. P. and Best

W. M. J. 2014), and was found to be a slightly better fit to the observations.

We also generated optical depth maps for a face on disk to recover the maximum value of τ for each transition. We then radially averaged these maps and plot the resulting optical depth of the ¹²C¹⁸O J=2-1 and ¹³C¹⁷O J=3-2 transitions for both models in Figure 2d. It can be seen that ¹²C¹⁸O is optically thick within the CO snowline (155 au) in both models, whereas the ¹³C¹⁷O remains optically thin across the full radial extent of the disk.

4. DISCUSSION

4.1. Comparison to other mass estimates

Using observations of ${}^{13}C{}^{17}O$ we derive a new gas mass for the HD 163296 disk of 0.31 M_{\odot}. The total disk mass depends on the gas to dust mass ratio (g/d), and using the dust mass from Isella et al. 2007 we find a g/d \approx 260. Here we compare our results to previous works.

The HD 163296 disk has been well studied and there are many mass measurements in the literature. In general, our estimate is the highest by a factor of 2 to 6 compared to previous studies using ${}^{12}C^{18}O$ (e.g. 0.17 M_{\odot}) and 0.048 ${\rm M}_{\odot}$ from Isella et al. 2007 and Williams & Mc-Partland 2016 respectively). There are a range of g/dvalues in the literature that span four orders of magnitude. Tilling et al. (2012) and Boneberg et al. (2016) models require a low g/d = 20. Isella et al. (2016) have a radially varying g/d covering a range from ≈ 30 to \approx 1100. Recent work from Powell et al. (2019) recover a total disk mass of 0.21 M_{\odot} with a high g/d $\sim 10^4$ in the outer disk. The one documented mass higher than our result is 0.58 M_{\odot} with a g/d = 350 (Woitke et al. 2019). The inconsistencies in these mass measurements and g/d from different models may be explained by trying to recover the gas density structure with optically thick lines. CO remains the best and most accessible tracer of mass that we have for disks (Molyarova et al. 2017), but robust lower limits to the gas mass can only be made by targeting the most optically thin isotopologues $({}^{12}C{}^{17}O, {}^{13}C{}^{18}O, \text{ and } {}^{13}C{}^{17}O).$

These masses have all been determined using a source distance of 122 pc. Considering the revised distance of 101.5 pc, the total disk gas mass from our work is thus 0.21 M_{\odot} (mass \propto flux \propto distance²).

4.2. The impact of CO chemistry on the disk mass

CO is susceptible to isotope-selective photodissociation which can reduce the abundance of the rarer isotopologues relative to ${}^{12}C{}^{16}O$ in the disk atmosphere. We find that the observations are well fit with interstel-

³ https://github.com/lime-rt/lime




Figure 2. The disk physical structure from Qi et al. (2011). a) The $n_{\rm H}$ density. b) The gas temperature. The white contours mark $\Sigma(n_{\rm H}) = 1.256 \times 10^{21}$ cm⁻² and $T_{\rm gas} = 19$ K, respectively. c) The $n({\rm CO})/n_{\rm H}$ distribution. d) The radially averaged optical depth (τ) of the ${}^{12}{\rm C}^{18}{\rm O}$ J=2-1 and ${}^{13}{\rm C}^{17}{\rm O}$ J=3-2 transitions from Model 1 (light purple and orange dashed lines) and Model 2 (dark purple and orange solid lines) assuming a face on disk. The vertical dashed line marks the location of the CO snowline in both models (155 au) and the horizontal dashed line marks where $\tau = 1$.

lar isotopic abundances. Because the ${}^{12}C^{16}O$, ${}^{13}C^{16}O$ and ${}^{12}C^{18}O$ line emission is optically thick, testing the significance of isotope-selective photodissociation in this disk requires higher sensitivity observations of the rarer isotopologues.

Observations have shown that CO is depleted with respect to H₂ in disks; however, without a better tracer of the H₂ column density, e.g., HD, the level of depletion is difficult to constrain. Carbon depletion effects are less significant in warmer disks around Herbig Ae stars compared to their T Tauri counterparts. Observations show moderate carbon depletion in the Herbig disk around HD 100546 with a model-derived [C]/[H] abundance ratio of 0.1 to 1.5×10^{-4} (Kama et al. 2016), and the value for CO adopted in our model is within this range. Consistent with this, models have also suggested that these disks have a close to canonical $n(\text{CO})/n(\text{H}_2)$ abundance (Bosman et al. 2018). These two chemical effects (isotope-selective photodissocation and carbon depletion) imply that our gas mass estimate is a lower limit.

4.3. Constraints on the location of the CO snowline

Locating the midplane CO snowline in disks is difficult due to the high optical depth of the more abundant CO isotopologues and the vertical temperature gradient of the disk. The location of the CO snowline can be determined directly by observing less abundant, optically




Figure 3. a-e) The de-projected and azimuthally averaged radial profiles of the observed and modelled CO lines. f) The value of integrated flux of the observed and modelled C¹⁷O J=3-2 line. The shaded regions are the errors as described in the text. Model 1 has a total disk mass of 0.089 M_{\odot} and Model 2 has a total disk mass of 0.31 M_{\odot} .

thin, CO isotopologues, or by detecting molecules that, due to chemistry, peak in abundance at a location related to the snowline. Qi et al. (2011) use observations CO and put the snowline at 155 au at 19 K: follow up work suggested that the snowline was instead at 90 au and 25 K (Qi et al. 2015). Out of these two scenarios our ¹³C¹⁷O observations fit best with the former option given the data in hand. The relationship between the cations, DCO⁺ (ring from 110 and 160 au) and N_2H^+ (inner edge of the ring at 90 au), and the location of the midplane CO snowline is not trivial, but both species have been detected in this disk (Mathews et al. 2013; Qi et al. 2015). Our analysis shows that the $C^{18}O$ emission in both models tested is optically thick, and thus cannot be used to easily locate the midplane CO snowline. However, the ¹³C¹⁷O emission is optically thin, so future observations at a higher spatial resolution and sensitivity could be used to directly constrain the radius of the midplane CO snowline. The new source distance from GAIA puts the proposed snowline locations at 75 au and 128 au. The former location is close one of the observed dust gaps in the disk and it may be the case that the drop in CO surface density detected here is due to gas depletion rather than the snowline. It is important to note that the snowline is not a simple sharp transition at the condensation temperature, but is instead determined by the balance of the rates of freeze out and thermal desorption, which should be considered in future disk models.

4.4. Is the disk gravitationally stable?

The potential exoplanet population currently probed with ALMA, via the ringed depletion of continuum emission, are gas giant planets on wide orbits. In the case of HD 163296 this would imply a multiple giant planet system and indeed, the presence of such a system has already been proposed (Isella et al. 2016; Pinte et al. 2018; Teague et al. 2018; Liu et al. 2018). The formation of massive planets on wide orbits can in some cases be achieved by core accretion, but a more economical route might involve the gravitational fragmentation of the outer regions of the disk (Boss 2011). Our new, higher disk mass estimate prompts us to investigate whether such processes may have occurred (or be occurring) in the HD 163296 disk.

The stability of a disk against fragmentation can be quantified via the Toomre Q parameter (Toomre 1964):

$$Q = \frac{c_s \kappa}{\pi G \Sigma}$$

where c_s is the sound speed of the gas, κ is the epicyclic frequency (equal to the angular velocity Ω in a Keplerian disk) and Σ is the surface density of the gas. Toomre Q




Figure 4. Toomre Q parameter for the new disk mass we derive in Model 2 and extrapolated back to 0.1 Myr. The shaded regions incorporate both the errors in stellar age and mass accretion rate. Vertical dashed lines mark the radial positions of gaps in the mm dust and kinematic perturbations which may be due to protoplanets.

values of 1 or less imply that the disk is susceptible to fragmentation, but simulations have shown that disks with $Q \lesssim 1.7$ begin to undergo instabilities in the form of non-axisymmetric spirals (Durisen et al. 2007). We calculate Q across the disk (Figure 4, orange) accounting for the lower mass due to the new source distance, assuming a g/d of 260, and the midplane temperature structure of our model (Figure 2b). We find that the minimum value of Q is ≈ 6 at ≈ 110 au, suggesting that the disk is currently gravitationally stable (in agreement with recent work from Powell et al. 2019).

The relatively large age of the HD 163296 system brings into question its stability earlier in its lifetime. The determination of previous disk masses is complicated by processes including episodic accretion (e.g. Mendigutía et al. 2013) and the decrease in accretion rate with time (e.g. Venuti et al. 2014). The magnitude of these effects are still under debate (see Hartmann et al. 2016, for a review), so we therefore assume that all of the accreted mass once resided in the disk, and that the accretion rate (\dot{M}) has been constant over disk lifetime. HD 163296 has an estimated age of $6.03^{+0.43}_{-0.14}$ Myr and log $\dot{M} = -6.81^{+0.16}_{-0.15} M_{\odot} \text{ yr}^{-1}$ (Wichittanakom et al., in prep.); thus under these assumptions, we estimate the disk mass at 0.1 Myr to be $1.13^{+0.51}_{-0.28} M_{\odot}$. The resulting minimum Toomre Q values for this stardisk configuration⁴ would be in the range of 1.3–0.7 (Figure 4, purple), placing regions of the disk from \sim 50– 220 au in the regime of instability. Such behaviour early in the disk lifetime has implications for the trapping and growth of dust (Rice et al. 2004) and the chemical composition of the disk (Evans et al. 2015). This previous unstable state could also be the source of the four massive planets currently proposed to reside in the disk around HD 163296.

5. CONCLUSIONS

We have presented the first detection of ${}^{13}C{}^{17}O$ in a protoplanetary disk showcasing the power of this optically thin isotopologue as a tracer of disk gas mass. This work provides robust evidence that disks are more massive than previously assumed (see also Zhu et al. 2019). Future observations of this tracer in more sources may help to address the discrepancy between the masses of disks and the observed exoplanet population (Manara et al. 2018).

We thank an anonymous referee for their constructive comments that improved the clarity of several sections of the paper. We thank Andrea Isella for the 12 CO, 13 CO and C¹⁸O J=2-1 data which was vital in our model comparison. We also thank Chumpon Wichittanakom for providing us with updated stellar parameters for HD 163296 pre-publication of their paper. AB acknowledges the studentship funded by the Science and Technology Facilities Council of the United Kingdom (STFC). CW acknowledges funds from the University of Leeds. JDI and CW acknowledge support from the STFC under ST/R000549/1. SN is grateful for support from JSPS (Japan Society for the Promotion of Science) Overseas Research Fellowships and the ALMA Japan Research Grant of NAOJ Chile Observatory, NAOJ-ALMA-211. This work is supported by MEXT/JSPS KAKENHI Grant Numbers 16J06887, 17K05399, 19K03910. Part of ALMA Data analysis was carried out on the Multi-wavelength Data Analysis System operated by the Astronomy Data Center (ADC), National Astronomical Observatory of Japan. This paper makes use of the following ALMA data: 2011.0.00010.SV, 2013.1.00601.S and 2015.1.01259.S. ALMA is a partnership of European Southern Observatory (ESO) (representing its member states), National Science Foundation (USA), and National Institutes of Natural Sciences (Japan), together with National Research Council (Canada), National Science Council and Academia Sinica Institute of Astronomy and Astrophysics (Taiwan), and Korea Astronomy and Space Science Institute (Korea), in cooperation with the Republic of Chile. The Joint ALMA Observatory is operated by ESO, Associated Universities, Inc/National Radio Astronomy Observatory (NRAO), and National Astronomical Observatory of Japan.

REFERENCES

- Ansdell, M., Williams, J. P., van der Marel, N., et al. 2016, ApJ, 828, 46
- Bergin, E. A., & Williams, J. P. 2017, in Astrophysics and Space Science Library, Vol. 445, Astrophysics and Space Science Library, ed. M. Pessah & O. Gressel, 1
- Bergin, E. A., Cleeves, L. I., Gorti, U., et al. 2013, Nature, 493, 644
- Boneberg, D. M., Panić, O., Haworth, T. J., Clarke, C. J., & Min, M. 2016, MNRAS, 461, 385
- Bosman, A. D., Walsh, C., & van Dishoeck, E. F. 2018, A&A, 618, A182
- Boss, A. P. 2011, ApJ, 731, 74
- Brinch, C., & Hogerheijde, M. R. 2010, A&A, 523, A25
- Carney, M. T., Fedele, D., Hogerheijde, M. R., et al. 2018, A&A, 614, A106

- Carney, M. T., Hogerheijde, M. R., Guzmán, V. V., et al. 2019, A&A, 623, A124
- Durisen, R. H., Boss, A. P., Mayer, L., et al. 2007, Protostars and Planets V, 607
- Evans, M. G., Ilee, J. D., Boley, A. C., et al. 2015, MNRAS, 453, 1147
- Gaia Collaboration, Brown, A. G. A., Vallenari, A., et al. 2018, A&A, 616, A1
- Graedel, T. E., Langer, W. D., & Frerking, M. A. 1982, ApJS, 48, 321
- Hartmann, L., Herczeg, G., & Calvet, N. 2016, ARA&A, 54, 135
- Isella, A., Testi, L., Natta, A., et al. 2007, A&A, 469, 213
- Isella, A., Guidi, G., Testi, L., et al. 2016, Physical Review Letters, 117, 251101
- Isella, A., Huang, J., Andrews, S. M., et al. 2018, ApJL, 869, L49
- Kama, M., Bruderer, S., van Dishoeck, E. F., et al. 2016, A&A, 592, A83

 $^{^4}$ We note this does not account for any change in the stellar mass over this time period, which would decrease Q further.

- Klapper, G., Surin, L., Lewen, F., et al. 2003, The Astrophysical Journal, 582, 262
- Liu, S.-F., Jin, S., Li, S., Isella, A., & Li, H. 2018, ApJ, 857, 87
- Long, F., Herczeg, G. J., Pascucci, I., et al. 2017, ApJ, 844, 99
- Loomis, R. A., Öberg, K. I., Andrews, S. M., et al. 2018, AJ, 155, 182
- Manara, C. F., Morbidelli, A., & Guillot, T. 2018, A&A, 618, L3
- Mathews, G. S., Klaassen, P. D., Juhász, A., et al. 2013, A&A, 557, A132
- McClure, M. K., Bergin, E. A., Cleeves, L. I., et al. 2016, ApJ, 831, 167
- Mendigutía, I., Brittain, S., Eiroa, C., et al. 2013, ApJ, 776, 44
- Miotello, A., Bruderer, S., & van Dishoeck, E. F. 2014, A&A, 572, A96
- Molyarova, T., Akimkin, V., Semenov, D., et al. 2017, ApJ, 849, 130
- Müller, H. S. P., Schlöder, F., Stutzki, J., & Winnewisser, G. 2005, JMoSt, 742, 215
- Notsu, S., Akiyama, E., Booth, A., et al. 2019, ApJ, 875, 96
- Pinte, C., Price, D. J., Ménard, F., et al. 2018, ApJL, 860, L13
- Powell, D., Murray-Clay, R., Pérez, L. M., Schlichting, H. E., & Rosenthal, M. 2019, ApJ, 878, 116

- Qi, C., D'Alessio, P., Öberg, K. I., et al. 2011, ApJ, 740, 84
- Qi, C., Öberg, K. I., Andrews, S. M., et al. 2015, ApJ, 813, 128
- Rice, W. K. M., Lodato, G., Pringle, J. E., Armitage, P. J.,
 & Bonnell, I. A. 2004, MNRAS, 355, 543
- Teague, R., Bae, J., Bergin, E. A., Birnstiel, T., & Foreman-Mackey, D. 2018, ApJL, 860, L12
- Tilling, I., Woitke, P., Meeus, G., et al. 2012, A&A, 538, A20
- Toomre, A. 1964, ApJ, 139, 1217
- van den Ancker, M. E., de Winter, D., & Tjin A Djie, H. R. E. 1998, A&A, 330, 145
- Venuti, L., Bouvier, J., Flaccomio, E., et al. 2014, A&A, 570, A82
- Walsh, C., Millar, T. J., & Nomura, H. 2010, ApJ, 722, 1607
- Williams, J. P., & McPartland, C. 2016, ApJ, 830, 32
- Williams J. P. and Best W. M. J. 2014, ApJ, 788, 59
- Wilson, T. L. 1999, Reports on Progress in Physics, 62, 143
- Woitke, P., Kamp, I., Antonellini, S., et al. 2019, PASP, 131, 064301
- Zhang, K., Bergin, E. A., Blake, G. A., Cleeves, L. I., & Schwarz, K. R. 2017, Nature Astronomy, 1, 0130
- Zhu, Z., Zhang, S., Jiang, Y.-F., et al. 2019, ApJL, 877, L18