

# Strong unitary uncertainty relations

Bing Yu

*School of Mathematics, South China University of Technology, Guangzhou 510640, China*

Naihuan Jing\*

*School of Mathematics, South China University of Technology, Guangzhou 510640, China and  
Department of Mathematics, North Carolina State University, Raleigh, NC 27695, USA*

Xianqing Li-Jost

*Max-Planck-Institute for Mathematics in the Sciences, 04103 Leipzig, Germany*

(Dated: August 13, 2019)

In this paper we provide a new set of uncertainty principles for unitary operators using a sequence of inequalities with the help of the geometric-arithmetic mean inequality. As these inequalities are “fine-grained” compared with the well-known Cauchy-Schwarz inequality, our framework naturally improves the results based on the latter. As such, the unitary uncertainty relations based on our method outperform the best known bound introduced in [Phys. Rev. Lett. 120, 230402 (2018)] to some extent. Explicit examples of unitary uncertainty relations are provided to back our claims.

## I. INTRODUCTION

At the foundation of quantum theory lies the Heisenberg uncertainty principle [1], which was first introduced in 1927. Traditionally, the textbook version of the uncertainty relation was established by Kennard [2] (see also the work of Weyl [3]) by means of variance in terms of position and momentum. The uncertainty principle lets us understand that if we were able to measure the momentum of a quantum system with certainty, then we would not gain the information of the measurement outcome of location with certainty. Robertson [4] generalized the uncertainty relation for position and momentum to any two bounded observables  $A$  and  $B$  as

$$\Delta A \Delta B \geq \frac{1}{2} |\langle \psi | [A, B] | \psi \rangle|, \quad (1)$$

where  $\Delta$  stands for the standard deviation of the observable relative to a fixed state  $|\psi\rangle$  and  $[A, B]$  represents the commutator of the observables  $A$  and  $B$ . Later Eq. (1) was improved by Schrödinger [5]. Recently, variance-based uncertainty relations have been intensely studied in [10–28].

Because of their relevance in quantum information theory, the entropies [29–45] have been employed to quantify the uncertainty relations between incompatible observables. The entropies are by no reason the best way to formulate joint uncertainties, and it is reasonable to consider all nonnegative Schur-concave functions as qualified uncertainty measures. This has lead to the well known universal uncertainty relations [46–49] expressed by majorization [50]. To this end, we shall remark that all these uncertainty relations play an important role in a wide range of applications such as entanglement detection [51, 52], quantum spin squeezing [53–57], quantum

metrology [58–62], quantum nonlocality [63, 64] and so on.

Now we turn to the variance-based uncertainty relations in the product form for unitary operators. Massar and Spiandel [10] have considered the uncertainty relation for two unitary operators that satisfy the commutation relation  $UV = e^{i\phi}VU$ . This uncertainty relation gives rise to the constraint for a quantum state to be simultaneously localized in two mutually unbiased bases related by a discrete Fourier transform (DFT). Other applications of Massar-Spiandel’s uncertainty relations include modular variables [7] and signal processing [8, 9]. Several further uncertainty relations for unitary operators related by DFT have been investigated in [11–14]. Later Bagchi and Pati [19] derived sum-form variance-based uncertainty relations for two general unitary operators, which have been tested experimentally with photonic qutrits [24]. The uncertainty relation for two general unitary operators is directly related to the preparation uncertainty principle that the amount of visibility for noncommuting unitary operators is nontrivially upper bounded. It is noted that a crucial technique underlying the variance-based uncertainty relations for two observables or unitary operators is the celebrated Cauchy-Schwarz inequality.

For multi-observables, the generalized uncertainty relation was first considered by Robertson using the positive semidefiniteness of a Hermitian matrix [6]. Recently, Bong et al used a similar method to derive a strong variance-based uncertainty relation for any  $n$  unitary operators [28]. The unitary uncertainty relation implies the famous Robertson-Schödinger uncertainty relation in the case of two Hermitian operators [5, 6]. However, the lower bound is implicitly given and sometimes hard to compute. This raises the question of explicitly extracting the uncertainty relation from the Gram determinant and also one wonders whether this strong uncertainty relation can be further improved.

The goal of this paper is to give new and improved un-

---

\* jing@ncsu.edu

certainty relations for general unitary operators. Following Xiao et al [21], a sequence of “fine-grained” inequalities compared with the Cauchy-Schwarz inequality are employed to derive uncertainty relations in connection with the Geometric-Arithmetic mean (AGM) inequality. We use this method to derive new variance-based unitary uncertainty relations in the product form for two and three operators in all quantum systems. The new uncertainty bounds for two unitary operators outperform those of Bong et al’s in the whole range. As the improvement is due to replacement of the Cauchy-Schwarz inequality underlying all previous uncertainty principles, our method provides fundamentally better bounds. We also generalize the uncertainty relation to the case of multiple unitary operators, and the new lower bounds are also shown to be tighter than that of Bong et al’s to some extent.

This paper is organized as follows. In Sec. II we introduce a fine-grained sequence of inequalities to generalize the Cauchy-Schwarz inequality, which was proved twice in this consideration. Our first main result (Thm. 1) of variance-based unitary uncertainty relations in the product-form is given in Sec. II A for two unitary operators. In Sec. II B, the bounds are strengthened by symmetry of permutations. In Sec. II C, examples are given to show our Theorem.1 provides tighter bounds than those of Bong et al’s. In Sec. III, we investigate product-form variance-based unitary uncertainty relations for three unitary operators. The uncertainty relations for multiple unitary operators are addressed in Sec. III A, and comparison is also provided with previous lower bounds for qutrit pure state, four-dimensional pure state and qutrit mixed state are studied in Sec. III B. Concluding remarks are given in IV. In the Appendix (Sec. V), we give some details of the proofs and calculations.

## II. UNCERTAINTY RELATIONS FOR TWO UNITARY OPERATORS

Let  $A$  and  $B$  be two unitary operators defined in a finite-dimensional Hilbert space with a fixed state  $|\psi\rangle$ . With respect to the mean value  $\langle A \rangle = \langle \psi | A | \psi \rangle$ , the variance of  $A$  over  $|\psi\rangle$  is defined by

$$\begin{aligned}\Delta A^2 &= \langle (A - \langle A \rangle)^\dagger (A - \langle A \rangle) \rangle \\ &= \langle \psi | \delta \hat{A}^\dagger \delta \hat{A} | \psi \rangle\end{aligned}\quad (2)$$

where  $\delta \hat{A} = A - \langle A \rangle$ . Note that the variance is bounded by  $0 \leq \Delta A^2 \leq 1$ .

Suppose  $\{|\psi_1\rangle, \dots, |\psi_n\rangle\}$  is a computational basis, then the state  $|f\rangle = \delta \hat{A} |\psi\rangle$  can be written as  $|f\rangle = \sum_{j=1}^n \alpha_j |\psi_j\rangle$  and similarly  $|g\rangle = \delta \hat{B} |\psi\rangle = \sum_{j=1}^n \beta_j |\psi_j\rangle$ . Thus the product of the variances obeys the unitary un-

certainty relation (UUR)

$$\begin{aligned}\Delta A^2 \Delta B^2 &= \langle f | f \rangle \langle g | g \rangle = \sum_{i,j} |\alpha_i|^2 |\beta_j|^2 \\ &\geq \left| \sum_{i=1}^n \alpha_i^* \beta_i \right|^2 = |\langle f | g \rangle|^2 \\ &= |\langle A^\dagger B \rangle - \langle A^\dagger \rangle \langle B \rangle|^2,\end{aligned}\quad (3)$$

where the inequality is due to the Cauchy-Schwarz inequality. Note that the last expression is independent from the choice of the computational basis.

Let  $\vec{X} = (x_1, x_2, \dots, x_n)$  and  $\vec{Y} = (y_1, y_2, \dots, y_n)$  be the (nonnegative) real vectors given by  $x_i = |\alpha_i|$ ,  $y_j = |\beta_j|$ , where  $(\alpha_1, \dots, \alpha_n)$  and  $(\beta_1, \dots, \beta_n)$  are the coordinate vectors of  $\delta \hat{A}$  and  $\delta \hat{B}$  respectively. Then the product of the variances can be rewritten as  $\Delta A^2 \Delta B^2 = |\vec{X}|^2 |\vec{Y}|^2 = \sum_{i,j} x_i^2 y_j^2$ . Note that the Cauchy-Schwarz inequality is in fact a consequence of  $n(n-1)/2$  AGM inequalities. Indeed,

$$\begin{aligned}\sum_{i,j} x_i^2 y_j^2 &= \sum_{i < j} (x_i^2 y_j^2 + x_j^2 y_i^2) + \sum_i x_i^2 y_i^2 \\ &\geq \sum_{i < j} 2x_i y_j x_j y_i + \sum_i x_i^2 y_i^2 \\ &= \left( \sum_{i=1}^n x_i y_i \right)^2\end{aligned}\quad (4)$$

with equality if and only if  $x_i y_j = x_j y_i$  for all  $i \neq j$ .

Now we refine the Cauchy-Schwarz inequality by introducing a sequence of partial ones. For each  $1 \leq k \leq n$ , define

$$\begin{aligned}I_k &= \sum_{1 \leq i \leq n} x_i^2 y_i^2 + \sum_{\substack{1 \leq i < j \leq n \\ k < j}} (x_i^2 y_j^2 + x_j^2 y_i^2) \\ &\quad + \sum_{1 \leq i < j \leq k} 2x_i y_j x_j y_i.\end{aligned}\quad (5)$$

In particular,  $I_1 = |\vec{X}|^2 |\vec{Y}|^2$  and  $I_n = (\sum_{i=1}^n x_i y_i)^2$ . The quantities  $I_k$  can be visualized by lattice dots within an  $n \times n$  square as follows. In Fig. 1 the black dot at  $i$ th column and  $j$ th row presents  $x_i^2 y_j^2$ , then  $I_k$  is the quantity  $(\sum_{i=1}^k x_i y_i)^2$  plus the dots outside of the  $k$ th principal square. It is easily seen that

$$I_{k+1} - I_k = -\left( \sum_{i=1}^k x_i y_{k+1} + y_i x_{k+1} \right)^2 \leq 0.$$

One therefore obtains the following descending sequence

$$I_1 \geq I_2 \geq \dots \geq I_{n-1} \geq I_n \quad (6)$$

and the Cauchy-Schwarz inequality also follows from the sequence:  $I_1 \geq I_n$ .

$x_i^2 y_j^2$	$x_1^2$	$x_2^2$	$x_3^2$	$x_4^2$	$\cdots$	$x_{n-2}^2$	$x_{n-1}^2$	$x_n^2$
$y_1^2$	•	•	•	•		•	•	•
$y_2^2$	•	•	•	•		•	•	•
$y_3^2$	•	•	•	•		•	•	•
$y_4^2$	•	•	•	•		•	•	•
$\cdots$					$\ddots$	$\vdots$	$\vdots$	$\vdots$
$y_{n-2}^2$	•	•	•	•	$\cdots$	•	•	•
$y_{n-1}^2$	•	•	•	•	$\cdots$	•	•	•
$y_n^2$	•	•	•	•	$\cdots$	•	•	•

FIG. 1. **Diagram for the  $I_k$**  ( $1 \leq k \leq n$ ). The black  $(i, j)$ -dot represents  $x_i^2 y_j^2$ . So  $I_k$  is  $(\sum_{i=1}^k x_i y_i)^2$  plus the dots outside of the  $k$ th principal square:  $I_k = (\sum_{i=1}^k x_i y_i)^2 + \sum_{1 \leq i < j \leq n} (x_i^2 y_j^2 + x_j^2 y_i^2) + \sum_{k+1 \leq i \leq n} x_i^2 y_i^2$ . The  $k$ th principal square shows the Cauchy-Schwarz inequality:  $\sum_{i,j=1}^k x_i^2 y_j^2 \geq (\sum_{i=1}^k x_i y_i)^2$ .

### A. Main Results

Let  $\rho$  be a mixed state on the Hilbert space. The variance of the unitary operator  $A$  with respect to  $\rho$  is defined as

$$(\Delta A)^2 = \text{Tr}(\rho \delta \hat{A}^\dagger \delta \hat{A}) \quad (7)$$

Let  $M = (m_{ij})_{l \times p}$  be a rectangular matrix, the vectorization  $|M\rangle$  (or  $\text{vec}(M)$ ) is the straightening vector  $(m_{11}, \dots, m_{1p}, \dots, m_{l1}, \dots, m_{lp}) \in \mathbb{C}^{lp}$ . As  $\rho$  is positive semi-definite, we will simply denote by  $|\sqrt{\rho}\rangle$  the pure state given by the vectorization  $\text{vec}(\sqrt{\rho})$  in the computational basis. Note that the vector  $|\sqrt{\rho}\rangle$  satisfies the following property [66]

$$|MT\rangle = (I \otimes M) |T\rangle \quad (8)$$

for two matrices  $M$  and  $T$  in suitable size. Thus

$$\begin{aligned} \Delta A^2 &= \text{Tr}(\sqrt{\rho} \delta \hat{A}^\dagger \delta \hat{A} \sqrt{\rho}) \\ &= \langle \sqrt{\rho} | (I \otimes \delta \hat{A}^\dagger \delta \hat{A}) | \sqrt{\rho} \rangle \\ &= |(I \otimes \delta \hat{A}) | \sqrt{\rho} \rangle|^2, \end{aligned} \quad (9)$$

where  $\sqrt{\rho}$  is the uniquely defined semi-definite positive matrix associated to  $\rho$ .

*Theorem 1.* Let  $A$  and  $B$  be two unitary operators on an  $n$ -dimensional Hilbert space  $H$  and  $\rho$  a quantum state on  $H$ . Suppose  $x_i$  and  $y_i$  are the probabilities of  $\delta \hat{A}$  and  $\delta \hat{B}$  with respect to a computational basis of  $H$ . Then the product of the variances of  $A$  and  $B$  satisfies the following uncertainty relations ( $k = 1, \dots, N$ )

$$\Delta A^2 \Delta B^2 \geq I_k, \quad (10)$$

where  $N = n$  (or  $n^2$ ) if  $\rho$  is pure (or mixed),  $I_k = \sum_{1 \leq i \leq N} x_i^2 y_i^2 + \sum_{1 \leq i < j \leq N} (x_i^2 y_j^2 + x_j^2 y_i^2) + \sum_{1 \leq i < j \leq k} 2x_i y_j x_j y_i$

and the equality holds if and only if  $x_i y_j = x_j y_i$  for all  $1 \leq i \neq j \leq k$ .

*Proof.* The uncertainty relations (10) for the case of pure state  $\rho$  were already shown in the last section. As for the mixed state  $\rho$ , we remarked that  $|\sqrt{\rho}\rangle$  is viewed as a pure state in an  $n^2$  dimensional Hilbert space [67], therefore the relations (10) also follow for all  $k = 1, \dots, n^2$ .

*Remark 1.* Note that  $|\vec{X}|^2 |\vec{Y}|^2 \geq I_k$  amounts to a partial Cauchy-Schwarz inequality [21] as it is obtained by applying the Cauchy-Schwarz inequality on the first  $k$  components. One can formulate an even more general inequality by selecting arbitrary  $x_i^2 y_j^2 + x_j^2 y_i^2$  instead of all the terms with  $1 \leq i < j \leq k$ .

Recently, Bong et al [28] derived a strong unitary uncertainty relation for any set of unitary operators based on the positive semi-definiteness of the Gram matrix. More precisely, let  $U_1, \dots, U_d$  be  $d$  unitary operators and  $U_0 = I$ . Their result says that the positive semidefiniteness of the Gram matrix  $G = G(\rho)$  of size  $d+1$  with  $G_{jk} = \langle U_j^\dagger U_k \rangle = \text{Tr}(\rho U_j^\dagger U_k)$  generalizes the UUR. In the case of two unitary operators  $A$  and  $B$ ,  $\det G(\rho) \geq 0$  turns out to be  $\Delta A^2 \Delta B^2 \geq |\langle A^\dagger B \rangle - \langle A^\dagger \rangle \langle B \rangle|^2$  [28], which is exactly the aforementioned (UUR) in Eq.(3).

We have seen that the lower bound of this UUR is weaker than our Theorem 1. In fact for any  $2n$  complex numbers  $\alpha_i, \beta_i$  [65]

$$|\sum_{i=1}^n \alpha_i^* \beta_i|^2 \leq (\sum_{i=1}^n |\alpha_i| |\beta_i|)^2 \leq \sum_{i,j} |\alpha_i|^2 |\beta_j|^2, \quad (11)$$

where the second inequality uses the Cauchy-Schwarz inequality. It follows from Eq.(6) that

$$\begin{aligned} \Delta A^2 \Delta B^2 &= I_1 \geq \dots \geq I_k \geq \dots \geq I_N = (\sum_{i=1}^N |\alpha_i| |\beta_i|)^2 \\ &\geq |\sum_{i=1}^N \alpha_i^* \beta_i|^2 = |\langle A^\dagger B \rangle - \langle A^\dagger \rangle \langle B \rangle|^2. \end{aligned} \quad (12)$$

This means that the UUR given in [28] for two unitary operators is the weakest bound in this sequence.

As the case of  $k = 1$  is trivial, we will include this in our statement of the result for simplicity.

## B. Improved UURs

The symmetric group  $S_N$ , which acts on the set  $\{1, 2, \dots, N\}$  naturally by permutation, can be used to strengthen the lower bounds of our UURs. For any two permutations  $\pi_1, \pi_2 \in S_N$ , the induced action of  $S_N \times S_N$  on  $I_k$  is given by

$$\begin{aligned} (\pi_1, \pi_2)I_k &= \sum_{1 \leq i \leq N} x_{\pi_1(i)}^2 y_{\pi_2(i)}^2 \\ &+ \sum_{\substack{1 \leq i < j \leq N \\ k < j}} (x_{\pi_1(i)}^2 y_{\pi_2(j)}^2 + x_{\pi_2(j)}^2 y_{\pi_1(i)}^2) \\ &+ \sum_{1 \leq i < j \leq k} 2x_{\pi_1(i)} y_{\pi_2(j)} x_{\pi_2(j)} y_{\pi_1(i)}. \end{aligned} \quad (13)$$

Clearly  $I_1$  is stable under the action of  $S_N \times S_N$ , subsequently

$$I_1 \geq (\pi_1, \pi_2)I_2 \geq \dots \geq (\pi_1, \pi_2)I_N. \quad (14)$$

Optimizing over the symmetric group  $S_N$ , we obtain the following stronger result.

*Theorem 2.* Let  $\rho$  be any quantum state on an  $n$ -dimensional Hilbert space  $H$ ,  $A$  and  $B$  two unitary operators on  $H$ . One has the following improved unitary uncertainty relations for the product of variances ( $k = 1, \dots, N$ )

$$\Delta A^2 \Delta B^2 \geq \max_{\pi_1, \pi_2 \in S_N} (\pi_1, \pi_2)I_k, \quad (15)$$

where  $N = n$  (or  $n^2$ ) if  $\rho$  is pure (or mixed),  $(\pi_1, \pi_2)I_k$  is defined in (13), and the equality holds if and only if  $x_{\pi_1(i)} y_{\pi_2(j)} = x_{\pi_2(j)} y_{\pi_1(i)}$  for all  $1 \leq i \neq j \leq k$ .

We remark that the lower bound in Theorem 2 is tighter than that of Theorem 1, since  $\max_{\pi_1, \pi_2 \in S_N} (\pi_1, \pi_2)I_k \geq I_k$  for any  $1 \leq k \leq N$ . An example is given to show strict strengthening of the bounds (see Example 1 and Fig. 3).

## C. Examples

*Example 1.* Let us consider the pure states  $|\psi\rangle = \cos\theta|0\rangle - \sin\theta|d-1\rangle$  on an  $d$ -dimensional Hilbert space [19], and  $A, B$  are the following unitary operators

$$\begin{aligned} A &= \sum_{j=-[\frac{d}{2}]}^{[\frac{d-1}{2}]} \omega^j |j\rangle \langle j| = \text{diag}(1, \omega, \omega^2, \dots, \omega^{d-1}), \\ B &= \sum_{j=-[\frac{d}{2}]}^{[\frac{d-1}{2}]} |j+1\rangle \langle j| = \begin{pmatrix} 0 & 1 \\ I_{d-1} & 0 \end{pmatrix}. \end{aligned} \quad (16)$$

where  $\omega = e^{i2\pi/d}$ . Note that  $AB = \omega BA$  [10].

*Case  $d = 2$ .* In this case

$$A = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad B = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}. \quad (17)$$

Both our UUR and Bong et al's are equal to  $\Delta A^2 \Delta B^2 = I_2$  (See Fig. 2). So we focus on  $d = 3, 4, 5$ , where the UURs are not saturated.

*Case  $d = 3$ .* The unitary operators are

$$A = \text{diag}(1, e^{\frac{2\pi i}{3}}, e^{\frac{4\pi i}{3}}), \quad B = \begin{pmatrix} 0 & 1 \\ I_2 & 0 \end{pmatrix}. \quad (18)$$

their associated real vectors  $\vec{X} = (x_1, x_2, x_3)$ ,  $\vec{Y} = (y_1, y_2, y_3)$  are given by

$$\begin{aligned} x_1 &= |(1 - e^{-\frac{2\pi i}{3}}) \sin^2 \theta \cos \theta|, \quad x_2 = 0, \\ x_3 &= |(1 - e^{-\frac{2\pi i}{3}}) \sin \theta \cos^2 \theta|, \end{aligned} \quad (19)$$

and

$$\begin{aligned} y_1 &= |-\sin^3 \theta|, \quad y_2 = |\cos \theta|, \\ y_3 &= |-\sin^2 \theta \cos \theta|, \end{aligned} \quad (20)$$

then  $I_2, I_3$  can be fixed and that  $\Delta A^2 \Delta B^2 \geq I_2 \geq I_3 = |\langle A^\dagger B \rangle - \langle A^\dagger \rangle \langle B \rangle|^2$ . Fig. 2 shows that our bounds are better than Bong et al's bound.

*Case  $d = 4, 5$ .* The vectors  $\vec{X}, \vec{Y}$  for  $d = 4, 5$  are respectively as follows.

$$\vec{X} = \begin{cases} |(1 - e^{\frac{-\pi i}{2}}) \frac{\sin 2\theta}{2}| (|\sin \theta|, 0, 0, |\cos \theta|), \\ |(1 - e^{\frac{-2\pi i}{5}}) \frac{\sin 2\theta}{2}| (|\sin \theta|, 0, 0, |\cos \theta|) \end{cases} \quad (21)$$

$$\vec{Y} = \begin{cases} (|-\sin^3 \theta|, |\cos \theta|, 0, |-\sin^2 \theta \cos \theta|) \\ (|-\sin^3 \theta|, |\cos \theta|, 0, 0, |-\sin^2 \theta \cos \theta|) \end{cases} \quad (22)$$

Then the lower bounds  $I_2, I_3, I_4$  (resp.  $I_2, I_3, I_4, I_5$ ) can be computed. It is readily seen that  $\Delta A^2 \Delta B^2 \geq I_2 = I_3 \geq I_4 = |\langle A^\dagger B \rangle - \langle A^\dagger \rangle \langle B \rangle|^2$  (resp.  $\Delta A^2 \Delta B^2 \geq I_2 = I_3 = I_4 \geq I_5 = |\langle A^\dagger B \rangle - \langle A^\dagger \rangle \langle B \rangle|^2$ ). Fig. 2 show that in all these cases, our bounds are better than that of Bong et al.

*Remark.* The bounds  $I_2, I_3, I_4$  can be further strengthened by Theorem 2. Consider the same qutrit state  $|\psi\rangle = \cos\theta|0\rangle - \sin\theta|2\rangle$ . Applying the symmetric group  $S_3$  as in Eq. (13) it follows that  $\Delta A^2 \Delta B^2 = \max_{\pi_1, \pi_2 \in S_3} (\pi_1, \pi_2)I_2 \geq$

$\max_{\pi_1, \pi_2 \in S_3} (\pi_1, \pi_2)I_3$ . Fig. 3 shows that the new bounds strictly outperform  $I_k$ .

*Example 2.* Consider the qubit mixed state  $\rho = \frac{1}{2}(I + \vec{r} \cdot \vec{\sigma})$  with  $\vec{r} = (\frac{1}{3}, \frac{2}{3} \cos \theta, \frac{2}{3} \sin \theta)$ , and  $\vec{\sigma} = (\sigma_x, \sigma_y, \sigma_z)$  where  $\sigma_x, \sigma_y, \sigma_z$  are the Pauli matrices.

Consider the unitary operators

$$A = e^{i\pi\sigma_y/8} = \begin{pmatrix} \cos \frac{\pi}{8} & \sin \frac{\pi}{8} \\ -\sin \frac{\pi}{8} & \cos \frac{\pi}{8} \end{pmatrix}, \quad (23)$$

$$B = e^{i\pi\sigma_z/8} = \begin{pmatrix} e^{i\frac{\pi}{8}} & 0 \\ 0 & e^{-i\frac{\pi}{8}} \end{pmatrix}, \quad (24)$$




FIG. 2. **Comparison of our bounds with Bong et al's bound for pure state.** The solid blue (upper) and green (lower) curves represent  $\Delta A^2 \Delta B^2$  and Bong et al's bound LB respectively. Our bounds  $I_2, I_3$  or  $I_4$  are tighter and shown in dashed yellow curves.




FIG. 3. Strengthened bounds vs. the bounds  $I_k$  for qutrit pure state. The solid blue curve represents  $\Delta A^2 \Delta B^2$  and  $\max_{\pi_1, \pi_2 \in S_3} (\pi_1, \pi_2) I_2$ . The dashed green curve represents  $\max_{\pi_1, \pi_2 \in S_3} (\pi_1, \pi_2) I_3$ . The dotted dashed and dotted curves represent  $I_2$  and LB (or  $I_3$ ) respectively.

which correspond to Bloch sphere rotations of  $-\pi/4$  about the  $y$  axis and  $z$  axis respectively.

It is seen that (cf. V. Appendix A)

$$|\sqrt{\rho}\rangle = \begin{pmatrix} \frac{\sqrt{3-\sqrt{5}}(\sqrt{5}-2\sin\theta) + \sqrt{3+\sqrt{5}}(\sqrt{5}+2\sin\theta)}{2\sqrt{30}} \\ -\frac{i(\sqrt{3-\sqrt{5}}-\sqrt{3+\sqrt{5}})(-i+2\cos\theta)}{2\sqrt{30}} \\ \frac{i(\sqrt{3-\sqrt{5}}-\sqrt{3+\sqrt{5}})(i+2\cos\theta)}{2\sqrt{30}} \\ \frac{\sqrt{3+\sqrt{5}}(\sqrt{5}-2\sin\theta) + \sqrt{3-\sqrt{5}}(\sqrt{5}+2\sin\theta)}{2\sqrt{30}} \end{pmatrix}. \quad (25)$$

Then the bounds  $I_2, I_3, I_4$  associated with  $\rho$  can be computed. We find that  $\Delta A^2 \Delta B^2 > I_2 > I_3 > I_4 \geq |\langle A^\dagger B \rangle - \langle A^\dagger \rangle \langle B \rangle|^2$ , which is the lower bound of Bong et al. Fig.4 shows that our bounds are almost always better than that of Bong et al. It seems that the bounds  $I_k$  are separated for mixed states.




FIG. 4. **Comparison of our bounds with that of Bong et al's for pure state.** The solid blue (upper) and purple (lower) curves represent  $\Delta A^2 \Delta B^2$  and Bong et al's bound LB respectively. Our bounds  $I_2, I_3$  or  $I_4$  are shown in dashed or dotted curves in yellow, green and red respectively.

### III. UNCERTAINTY RELATIONS FOR THREE UNITARY OPERATORS

We now study product-form variance-based unitary uncertainty relations for three unitary operators based upon our UUR for two unitary operators in terms of the quantities  $I_k$  in the preceding section.

#### A. Main results

Let  $A, B$  and  $C$  be three unitary operators defined on an  $n$ -dimensional Hilbert space. By Theorem 1 the UURs for the pairs  $\{A, B\}$ ,  $\{B, C\}$  and  $\{A, C\}$  over the quantum state  $\rho$  are written as  $\Delta A^2 \Delta B^2 \geq I_k$ ,  $\Delta B^2 \Delta C^2 \geq J_k$ , and  $\Delta A^2 \Delta C^2 \geq K_k$  where  $I_k, J_k, K_k$  are the quantities  $I_k$  defined above (6) for the pairs respectively. Taking the square root of the product, we have the following result.



*Corollary 1.* For a fixed quantum state  $\rho$  and three unitary operators  $A, B$  and  $C$  on an  $n$ -dimensional Hilbert space  $H$ , the product of the variances obeys the following inequalities ( $k = 2, \dots, N$ )

$$\Delta A^2 \Delta B^2 \Delta C^2 \geq (I_k J_k K_k)^{1/2} \quad (26)$$

where  $N = n$  (or  $n^2$ ) if  $\rho$  is pure (or mixed),  $I_k = I_k(A, B)$ ,  $J_k = I_k(A, C)$  and  $K_k = I_k(B, C)$ . Here  $I_k$  are defined in Sect. II A.

One can also strengthen the bound using the symmetry of  $S_N$ . Denote  $\max_{\pi_1, \pi_2 \in S_N} (\pi_1, \pi_2) I_i$  by  $\hat{I}_i$ , then the improved UURs are given in the following corollary.

*Corollary 2.* Let  $\rho, A, B, C$  as in Cor. 1. The strengthened UURs are given by

$$\Delta A^2 \Delta B^2 \Delta C^2 \geq (\hat{I}_k \hat{J}_k \hat{K}_k)^{1/2}, \quad (27)$$

where  $\hat{I}_k = \max_{\pi_1, \pi_2 \in S_N} (\pi_1, \pi_2) I_k$ ,  $\hat{J}_k = \max_{\pi_1, \pi_2 \in S_N} (\pi_1, \pi_2) J_k$  and  $\hat{K}_k = \max_{\pi_1, \pi_2 \in S_N} (\pi_1, \pi_2) K_k$ .

## B. Examples

For three unitary operators  $A, B, C$ , Bong et al's UUR is expressed as the positivity of the Gram matrix:

$$\det G(\rho) = \det \begin{pmatrix} 1 & \langle A \rangle & \langle B \rangle & \langle C \rangle \\ \langle A^\dagger \rangle & 1 & \langle A^\dagger B \rangle & \langle A^\dagger C \rangle \\ \langle B^\dagger \rangle & \langle B^\dagger A \rangle & 1 & \langle B^\dagger C \rangle \\ \langle C^\dagger \rangle & \langle C^\dagger A \rangle & \langle C^\dagger B \rangle & 1 \end{pmatrix} \geq 0 \quad (28)$$

which can be rewritten as

$$\begin{aligned} \Delta A^2 \Delta B^2 \Delta C^2 &\geq \Delta A^2 |\langle B^\dagger C \rangle - \langle B^\dagger \rangle \langle C \rangle|^2 \\ &+ \Delta B^2 |\langle A^\dagger C \rangle - \langle A^\dagger \rangle \langle C \rangle|^2 + \Delta C^2 |\langle A^\dagger B \rangle - \langle A^\dagger \rangle \langle B \rangle|^2 \\ &- 2 \operatorname{Re} \{ (\langle A^\dagger C \rangle - \langle A^\dagger \rangle \langle C \rangle) (\langle C^\dagger B \rangle - \langle C^\dagger \rangle \langle B \rangle) \\ &(\langle B^\dagger A \rangle - \langle B^\dagger \rangle \langle A \rangle) \}, \end{aligned} \quad (29)$$

where  $\operatorname{Re}$  denotes the real part. The right hand side (RHS) will be denoted by LB. This inequality is saturated for pure state when  $n = \dim H \leq 3$ , where the determinant of the Gram matrix vanishes.

Let us compare their result with our bounds in the cases of pure state ( $n \geq 4$ ) and mixed state separately.

*Example 3.* Let  $|\psi\rangle = \frac{1}{2} \cos \frac{\theta}{2} |0\rangle + \frac{\sqrt{3}}{2} \sin \frac{\theta}{2} |1\rangle + \frac{1}{2} \sin \frac{\theta}{2} |2\rangle + \frac{\sqrt{3}}{2} \cos \frac{\theta}{2} |3\rangle$  and we take three unitary operators:

$$\begin{aligned} A &= \operatorname{diag}(1, e^{i\frac{\pi}{2}}, e^{i\pi}, e^{i\frac{3\pi}{2}}), B = \begin{pmatrix} 0 & 1 \\ I_3 & 0 \end{pmatrix}, \\ C &= \begin{pmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix}. \end{aligned} \quad (30)$$

Using Corollary 1, the lower bounds  $(I_k J_k K_k)^{1/2}$  ( $2 \leq k \leq 4$ ) can be easily calculated and one sees that they are better than that of Bong et al's in significant regions. See Fig. 5 for the comparison.




FIG. 5. **Comparison of our bounds with Bong et al's for pure state.** The solid blue (upper) and purple (lower) curves are  $\Delta A^2 \Delta B^2 \Delta C^2$  and Bong et al's bound LB. Other three dotted dashed yellow, dashed green, dotted red lines (from top to bottom) represent our bounds  $(I_2 J_2 K_2)^{1/2}$ ,  $(I_3 J_3 K_3)^{1/2}$ ,  $(I_4 J_4 K_4)^{1/2}$  separately.

*Example 4.* Consider the mixed state analyzed in Example 2 and three unitary operators:

$$\begin{aligned} A &= e^{i\pi\sigma_y/8} = \begin{pmatrix} \cos \frac{\pi}{8} & \sin \frac{\pi}{8} \\ -\sin \frac{\pi}{8} & \cos \frac{\pi}{8} \end{pmatrix}, \\ B &= e^{i\pi\sigma_z/8} = \begin{pmatrix} e^{i\frac{\pi}{8}} & 0 \\ 0 & e^{-i\frac{\pi}{8}} \end{pmatrix}, \\ C &= e^{i\pi\sigma_x/8} = \begin{pmatrix} \cos \frac{\pi}{8} & i \sin \frac{\pi}{8} \\ i \sin \frac{\pi}{8} & \cos \frac{\pi}{8} \end{pmatrix}. \end{aligned} \quad (31)$$

The vectorized state  $|\sqrt{\rho}\rangle$  was given in Example 2, based on this the uncertainty bound  $(I_2 J_2 K_2)^{1/2}$  can be computed and is seen to be always tighter than the Bong et al's bound LB (cf. Fig. (6)). However,  $(I_3 J_3 K_3)^{1/2}$  and  $(I_4 J_4 K_4)^{1/2}$  are not as good as LB.




FIG. 6. **Comparison of our bounds with Bong et al's bound for mixed state.** The solid blue (upper) and purple (lower) curves represent  $\Delta A^2 \Delta B^2 \Delta C^2$  and Bong et al's bound LB. The other three dotted dashed yellow, dashed green, dotted red curves (from top to bottom) are our bounds  $(I_2 J_2 K_2)^{1/2}$ ,  $(I_3 J_3 K_3)^{1/2}$ ,  $(I_4 J_4 K_4)^{1/2}$  respectively.

*Example 5.* Consider the mixed qutrit state  $\rho = \frac{1}{3}(I + \sqrt{3}\vec{n} \cdot \vec{\lambda})$  [68] on  $\mathbb{C}^3$ , where  $\vec{\lambda}$  is the 8-dimensional vector of the Gell-Mann matrices of rank 3 and

$\vec{n} = (\frac{1}{\sqrt{3}}\cos\theta, 0, 0, 0, 0, \frac{1}{\sqrt{3}}\sin\theta, 0, 0)$ . As a matrix, the density operator  $\rho$  takes the following form

$$\rho = \frac{1}{3} \begin{pmatrix} 1 & \cos\theta & 0 \\ \cos\theta & 1 & \sin\theta \\ 0 & \sin\theta & 1 \end{pmatrix}. \quad (32)$$

The three unitary operators  $A, B, C$  are taken as the rotational operators  $R_{Z,\theta_z}, R_{Y,\theta_y}, R_{X,\theta_x}$  with the Euler angles  $\theta_z = \frac{\pi}{4}, \theta_y = -\frac{\pi}{4}, \theta_x = \frac{\pi}{3}$  around  $Z, Y$  and  $X$  axes respectively. i.e.

$$\begin{aligned} R_{Z,\theta_z} &= \begin{pmatrix} \cos\theta_z & \sin\theta_z & 0 \\ -\sin\theta_z & \cos\theta_z & 0 \\ 0 & 0 & 1 \end{pmatrix}, \\ R_{Y,\theta_y} &= \begin{pmatrix} \cos\theta_y & 0 & \sin\theta_y \\ 0 & 1 & 0 \\ -\sin\theta_y & 0 & \cos\theta_y \end{pmatrix}, \\ R_{X,\theta_x} &= \begin{pmatrix} 1 & 0 & 0 \\ 0 & \cos\theta_x & -\sin\theta_x \\ 0 & \sin\theta_x & \cos\theta_x \end{pmatrix}. \end{aligned} \quad (33)$$

The state  $|\sqrt{\rho}\rangle$  is seen as follows (cf. V. Appen. B)

$$\begin{aligned} |\sqrt{\rho}\rangle &= \left( \frac{\cos^2\theta}{\sqrt{6}} + \frac{\sin^2\theta}{\sqrt{3}}, \frac{\cos\theta}{\sqrt{6}}, \right. \\ &\quad \frac{(-2 + \sqrt{2})\sin 2\theta}{4\sqrt{3}}, \frac{\cos\theta}{\sqrt{6}}, \frac{1}{\sqrt{6}}, \frac{\sin\theta}{\sqrt{6}}, \\ &\quad \left. \frac{(-2 + \sqrt{2})\sin 2\theta}{4\sqrt{3}}, \frac{\sin\theta}{\sqrt{6}}, \frac{\cos^2\theta}{\sqrt{3}} + \frac{\sin^2\theta}{\sqrt{6}} \right) \end{aligned}$$

The lower bounds  $\{(I_k J_k K_k)^{1/2} | 2 \leq k \leq 8\}$  associated with  $\rho$  are then calculated and depicted in Fig. 7. The picture shows that our lower bounds  $\{(I_k J_k K_k)^{1/2} | 2 \leq k \leq 6\}$  are always tighter than  $LB$ , Bong et al's bound,  $(I_7 J_7 K_7)^{1/2}$  and  $(I_8 J_8 K_8)^{1/2}$  are better than  $LB$  in some region, and  $LB$  is better than  $(I_9 J_9 K_9)^{1/2}$ .

#### IV. CONCLUSION

In this paper, we have studied a stronger form of variance-based unitary uncertainty relations (UUR) for two and three operators relative to both pure and mixed quantum states. Our idea is to employ the partial Cauchy-Schwarz inequality to derive a sequence of effective lower bounds  $I_k$  for the product of the uncertainties. Moreover, our bounds  $I_k$  can be strengthened by permutation.

We have also shown that our new uncertainty bounds are tighter than the recently discovered UUR given by Bong et al using the positivity of the Gram matrix [28]




FIG. 7. **Comparison of our bounds with Bong et al's bound for qutrit state.** The solid blue (upper) and green (lower) curves represent  $\Delta A^2 \Delta B^2 \Delta C^2$  and Bong et al's bound  $LB$  respectively. The other eight dashed or dotted curves (from top to bottom) are the bounds  $(I_2 J_2 K_2)^{1/2}, \dots, (I_9 J_9 K_9)^{1/2}$ .

for two and multiple unitary operators. In one comparison with Bong et al's bound, two unitary operators related by the discrete Fourier transform are examined and it was found that our bounds outperform significantly their lower bounds, which could have potential implications for signal processing and modular variables. In another example of three unitary operators, most of our bounds demonstrated better effects than theirs for arbitrary quantum state and three unitary operators.

#### ACKNOWLEDGMENT

We are grateful to Yunlong Xiao for stimulating discussions and help in this work. The research is partially supported by National Natural Science Foundation of China grant no. 11531004, Simons Foundation grant no. 523868 and China Scholarship Council.

#### V. APPENDICES

##### Appendix A

The Hermitian matrix  $\rho$  is unitarily diagonalizable, so it can be expressed as  $\rho = U D U^\dagger$  for a unitary matrix  $U$  and a diagonal matrix  $D$ .

For the qubit mixed state  $\rho = \frac{1}{2}(I + \vec{r} \cdot \vec{\sigma})$  with  $\vec{r} = \{\frac{1}{3}, \frac{2}{3}\cos\theta, \frac{2}{3}\sin\theta\}$  and  $\vec{\sigma} = \{\sigma_x, \sigma_y, \sigma_z\}$ . The unitary matrix  $U = (\frac{v_1}{|v_1|}, \frac{v_2}{|v_2|})$ , where the orthogonal eigenvectors

$u_i, u_2$  are given by

$$v_1 = \left( -\frac{i(2\sin\theta + \sqrt{5})}{-i + 2\cos\theta}, 1 \right)^T,$$

$$v_2 = \left( \frac{i(\sqrt{5} - 2\sin\theta)}{-i + 2\cos\theta}, 1 \right)^T.$$

The diagonal matrix  $D$  is determined by the corresponding eigenvalues and

$$D^{\frac{1}{2}} = \begin{pmatrix} \sqrt{\frac{1}{6}(3+\sqrt{5})} & 0 \\ 0 & \sqrt{\frac{1}{6}(3-\sqrt{5})} \end{pmatrix}. \quad (34)$$

Therefore the unique positive semidefinite square root of the Hermitian matrix  $\rho$  is given by

$$\sqrt{\rho} = UD^{\frac{1}{2}}U^\dagger = \begin{pmatrix} \frac{\sqrt{3+\sqrt{5}}(2\sin\theta+\sqrt{5})+\sqrt{3-\sqrt{5}}(\sqrt{5}-2\sin\theta)}{2\sqrt{30}} & \frac{i(\sqrt{3-\sqrt{5}}-\sqrt{3+\sqrt{5}})(i+2\cos\theta)}{2\sqrt{30}} \\ -\frac{i(\sqrt{3-\sqrt{5}}-\sqrt{3+\sqrt{5}})(-i+2\cos\theta)}{2\sqrt{30}} & \frac{\sqrt{3-\sqrt{5}}(2\sin\theta+\sqrt{5})+\sqrt{3+\sqrt{5}}(\sqrt{5}-2\sin\theta)}{2\sqrt{30}} \end{pmatrix}. \quad (35)$$

Consequently, the vectorization  $|\sqrt{\rho}\rangle$  for the mixed state  $\rho$  is obtained as a 4-dimensional pure state.

### Appendix B

For the qutrit mixed state  $\rho = \frac{1}{3}(I + \sqrt{3}\vec{n} \cdot \vec{\lambda})$  with  $\vec{n} = (\frac{1}{\sqrt{3}}\cos\theta, 0, 0, 0, 0, \frac{1}{\sqrt{3}}\sin\theta, 0, 0)$  and  $\vec{\lambda} = (\lambda_1, \lambda_2, \dots, \lambda_8)$  is the vector of the Gell-Mann matrices. Using a similar procedure as Appendix A, we diagonalize the matrix  $\rho$  as

$$D = U^\dagger \rho U = \begin{pmatrix} \frac{2}{3} & 0 & 0 \\ 0 & \frac{1}{3} & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad (36)$$

where the unitary matrix  $U = (\frac{v_1}{|v_1|}, \frac{v_2}{|v_2|}, \frac{v_3}{|v_3|})$  is given by the eigenvectors

$$v_1 = (\cot\theta, \csc\theta, 1)^T,$$

$$v_2 = (-\tan\theta, 0, 1)^T,$$

$$v_3 = (\cot\theta, -\csc\theta, 1)^T.$$

Then the unique semidefinite square root of matrix  $\rho$  is

$$\sqrt{\rho} = UD^{\frac{1}{2}}U^\dagger = \begin{pmatrix} \frac{\cos^2\theta}{\sqrt{6}} + \frac{\sin^2\theta}{\sqrt{3}} & \frac{\cos\theta}{\sqrt{6}} & \frac{(-2+\sqrt{2})\sin 2\theta}{4\sqrt{3}} \\ \frac{\cos\theta}{\sqrt{6}} & \frac{1}{\sqrt{6}} & \frac{\sin\theta}{\sqrt{6}} \\ \frac{(-2+\sqrt{2})\sin 2\theta}{4\sqrt{3}} & \frac{\sin\theta}{\sqrt{6}} & \frac{\cos^2\theta}{\sqrt{3}} + \frac{\sin^2\theta}{\sqrt{6}} \end{pmatrix}. \quad (37)$$

By stacking columns of the matrix  $\sqrt{\rho}$  on top of one another, we have the pure state  $|\sqrt{\rho}\rangle$  on the 9-dimensional Hilbert space.

### Appendix C

To highlight our method, we further consider the strengthened UURs for four unitary operators.

Let  $A, B, C$  and  $D$  be four unitary operators on an  $n$ -dimensional Hilbert space, the product form of variance-based unitary uncertainty relations with two pairs of unitary operators  $\{A, B\}$  and  $\{C, D\}$  in quantum state  $|\psi\rangle$  can be written as  $\Delta A^2 \Delta B^2 \geq I_k$ ,  $\Delta C^2 \Delta D^2 \geq J_k$  respectively.

Therefore UURs for four unitary operators is then given as follows:

$$\Delta A^2 \Delta B^2 \Delta C^2 \Delta D^2 \geq I_k J_k, \quad (38)$$

with  $2 \leq k \leq N$ .

Though the above seems to be a trivial step beyond the case of two unitary operators, it still outperforms Bong et al's bound in many situations.

*Example 6.* Let us consider the pure state  $|\psi\rangle = \cos\theta|0\rangle + \frac{1}{2}\sin\theta|1\rangle + \frac{\sqrt{3}}{2}\sin\theta|4\rangle$  on 5-dimensional Hilbert space, and take four unitary operators  $A, B, C$  and  $D$  as follows.

$$A = \text{diag}(e^{-\frac{4\pi i}{5}}, e^{-\frac{2\pi i}{5}}, 1, e^{\frac{2\pi i}{5}}, e^{\frac{4\pi i}{5}}),$$

$$B = \text{diag}(e^{\frac{4\pi i}{5}}, e^{\frac{2\pi i}{5}}, 1, e^{-\frac{2\pi i}{5}}, e^{-\frac{4\pi i}{5}}), \quad (39)$$

$$C = \begin{pmatrix} 0 & 1 \\ I_4 & 0 \end{pmatrix}, D = i \begin{pmatrix} 0 & 1 \\ I_4 & 0 \end{pmatrix}.$$

It is not difficult to check that  $\Delta A^2 \Delta B^2 \Delta C^2 \Delta D^2 = I_k J_k$  with  $2 \leq k \leq 5$  in our UURs due to its saturated conditions.

For four unitary operators, Bong et al's UUR is

$$\det G(\rho) = \det \begin{pmatrix} 1 & \langle A \rangle & \langle B \rangle & \langle C \rangle & \langle D \rangle \\ \langle A^\dagger \rangle & 1 & \langle A^\dagger B \rangle & \langle A^\dagger C \rangle & \langle A^\dagger D \rangle \\ \langle B^\dagger \rangle & \langle B^\dagger A \rangle & 1 & \langle B^\dagger C \rangle & \langle B^\dagger D \rangle \\ \langle C^\dagger \rangle & \langle C^\dagger A \rangle & \langle C^\dagger B \rangle & 1 & \langle C^\dagger D \rangle \\ \langle D^\dagger \rangle & \langle D^\dagger A \rangle & \langle D^\dagger B \rangle & \langle D^\dagger C \rangle & 1 \end{pmatrix} \geq 0 \quad (40)$$

It is complicated and cumbersome to simplify the above into a form of  $\Delta A^2 \Delta B^2 \Delta C^2 \Delta D^2 \geq M$ , the uncertainty lower bound. So we simply sketch  $\det G(|\psi\rangle)$  in Fig.8.



We find that the determinant  $G(|\psi\rangle)$  vanishes only when  $\{\theta = n\pi | n \in \mathbb{Z}\}$ , i.e., when the uncertainty relation is saturated.

This means that our bound is tighter than Bong et al's bound in the whole range except at the points  $n\pi$ . Given the complexity of straightening out the product of the variances from  $\det G(\rho)$  as required from Bong et al's method, our procedure is simpler and provides direct lower bounds in this case.




FIG. 8. The solid blue curve represents  $\Delta A^2 \Delta B^2 \Delta C^2 \Delta D^2$ , the dotted orange line denotes  $\det G(|\psi\rangle)$ .

- 
- [1] W. Heisenberg, Z. Phys. **43**, 172 (1927).
  - [2] E. H. Kennard, Z. Phys. **44**, 326 (1927).
  - [3] H. Weyl, *Gruppentheorie und Quantenmechanik* (Hirzel, Leipzig, 1928) [English translation, H. P. Robertson, *The Theory of Groups and Quantum Mechanics* (Dover, New York, 1931)].
  - [4] H. P. Robertson, Phys. Rev. **34**, 163 (1929).
  - [5] E. Schrödinger, Ber. Kgl. Akad. Wiss. Berlin **24**, 296 (1930).
  - [6] H. P. Robertson, Phys. Rev. **35**, 667A (1930); **46**, 794 (1934).
  - [7] Y. Aharonov, H. Pendleton, and A. Petersen, Int. J. Theor. Phys. **2**, 213 (1969).
  - [8] T. Opatrný, V. Bužek, J. Bajer, and G. Drobný, Phys. Rev. A **52**, 2419 (1995).
  - [9] T. Opatrný, D.-G. Welsch, and V. Bužek, Phys. Rev. A **53**, 3822 (1996).
  - [10] S. Massar and P. Spindel, Phys. Rev. Lett. **100**, 190401 (2008).
  - [11] A. B. Klimov, C. Muñoz, and L. L. Sánchez-Soto, Phys. Rev. A **80**, 043836 (2009).
  - [12] M. A. Marchioli and M. Ruzzi, Ann. Phys. (NY) **327**, 1538 (2012).
  - [13] R. Namiki and Y. Tokunaga, Phys. Rev. Lett. **108**, 230503 (2012).
  - [14] M. A. Marchioli and P. E. M. F. Mendonça, Ann. Phys. (NY) **336**, 76 (2013).
  - [15] L. Maccone and A. K. Pati, Phys. Rev. Lett. **113**, 260401 (2014).
  - [16] J.-L. Li and C.-F. Qiao, Sci. Rep. **5**, 12708 (2015).
  - [17] A. K. Pati, U. Singh, and U. Sinha, Phys. Rev. A **92**, 052120 (2015).
  - [18] M. J. W. Hall, A. K. Pati, and J. Wu, Phys. Rev. A **93**, 052118 (2016).
  - [19] S. Bagchi and A. K. Pati, Phys. Rev. A **94**, 042104 (2016).
  - [20] Y. Xiao, N. Jing, X. Li-Jost, and Shao-Ming Fei, Sci. Rep. **6**, 23201 (2016).
  - [21] Y. Xiao, N. Jing, B. Yu, S.-M. Fei, and X. Li-Jost, arXiv:1610.01692 (2016).
  - [22] D. Mondal, S. Bagchi, and A. K. Pati, Phys. Rev. A **95**, 052117 (2017).
  - [23] Y. Xiao, C. Guo, F. Meng, N. Jing, and M.-H. Yung, arXiv:1706.05650 (2017).
  - [24] L. Xiao, K. Wang, X. Zhan, Z. Bian, J. Li, Y. Zhang, P. Xue, and A. K. Pati, Opt. Express **25**, 17904 (2017).
  - [25] C. Qian, J.-L. Li, and C.-F. Qiao, Quantum Inf. Process. **17**, 84 (2018).
  - [26] S. Sazim, S. Adhikari, A. K. Pati, and P. Agrawal, Phys. Rev. A **98**, 032123 (2018).
  - [27] G. Sharma, C. Mukhopadhyay, S. Sazim, and A. K. Pati, Phys. Rev. A **98**, 032106 (2018).
  - [28] K. Bong, N. Tischler, R. B. Patel, S. Wollmann, G. J. Pryde, and M. J. W. Hall, Phys. Rev. Lett. **120**, 230402 (2018).
  - [29] I. Białynicki-Birula and J. Mycielski, Commun. Math. Phys. **44**, 129 (1975).
  - [30] D. Deutsch, Phys. Rev. Lett. **50**, 631 (1983).
  - [31] M. H. Partovi, Phys. Rev. Lett. **50**, 1883 (1983).
  - [32] K. Kraus, Phys. Rev. D **35**, 3070 (1987).

- [33] H. Maassen and J. B. M. Uffink, Phys. Rev. Lett. **60**, 1103 (1988).
- [34] I. D. Ivanovic, J. Phys. A: Math. Gen. **25**, L363 (1992).
- [35] J. Sánchez, Phys. Lett. A **173**, 233 (1993).
- [36] M. A. Ballester and S. Wehner, Phys. Rev. A **75**, 022319 (2007).
- [37] S. Wu, S. Yu, and K. Mølmer, Phys. Rev. A **79**, 022104 (2009).
- [38] Y. Huang, Phys. Rev. A **83**, 052124 (2011).
- [39] M. Tomamichel and R. Renner, Phys. Rev. Lett. **106**, 110506 (2011).
- [40] P. J. Coles, R. Colbeck, L. Yu, and M. Zwolak, Phys. Rev. Lett. **108**, 210405 (2012).
- [41] P. J. Coles and M. Piani, Phys. Rev. A **89**, 022112 (2014).
- [42] Y. Xiao, N. Jing, S.-M. Fei, T. Li, X. Li-Jost, T. Ma, and Z.-X. Wang, Phys. Rev. A **93**, 042125 (2016).
- [43] Y. Xiao, N. Jing, S.-M. Fei, and X. Li-Jost, J. Phys. A **49**, 49LT01 (2016).
- [44] Y. Xiao, N. Jing, and X. Li-Jost, Quantum Inf. Process. **16**, 104 (2017).
- [45] P. J. Coles, M. Berta, M. Tomamichel, and S. Wehner, Rev. Mod. Phys. **89**, 015002 (2017).
- [46] S. Friedland, V. Gheorghiu, and G. Gour, Phys. Rev. Lett. **111**, 230401 (2013).
- [47] Z. Puchała, L. Rudnicki, and K. Życzkowski, J. Phys. A: Math. Theor. **46**, 272002 (2013).
- [48] L. Rudnicki, Z. Puchała, and K. Życzkowski, Phys. Rev. A **89**, 052115 (2014).
- [49] V. Narasimhachar, A. Poostindouz, and G. Gour, New J. Phys. **18**, 033019 (2016).
- [50] A. W. Marshall, I. Olkin, and B. C. Arnold, *Inequalities: Theory of majorization and its applications*, Springer, New York (2011).
- [51] H. F. Hofmann and S. Takeuchi, Phys. Rev. A **68**, 032103 (2003).
- [52] O. Gühne, Phys. Rev. Lett. **92**, 117903 (2004).
- [53] D. F. Walls and P. Zoller, Phys. Rev. Lett. **47**, 709 (1981).
- [54] K. Wódkiewicz and J. Eberly, J. Opt. Soc. Am. B **2**, 458 (1985).
- [55] D. J. Wineland, J. J. Bollinger, W. M. Itano, F. L. Moore, and D. J. Heinzen, Phys. Rev. A **46**, R6797 (1992).
- [56] M. Kitagawa and M. Ueda, Phys. Rev. A **47**, 5138 (1993).
- [57] J. Ma, X. G. Wang, C. P. Sun, and F. Nori, Phys. Rep. **509**, 89 (2011).
- [58] S. L. Braunstein and C. M. Caves, Phys. Rev. Lett. **72**, 3439 (1994).
- [59] S. L. Braunstein, C. M. Caves, and G. J. Milburn, Ann. Phys. (NY) **247**, 135 (1996).
- [60] V. Giovannetti, S. Lloyd, and L. Maccone, Science **306**, 1330 (2004).
- [61] V. Giovannetti, S. Lloyd, and L. Maccone, Phys. Rev. Lett. **96**, 010401 (2006).
- [62] V. Giovannetti, S. Lloyd, and L. Maccone, Nat. Photonics **5**, 222 (2011).
- [63] J. Oppenheim, and S. Wehner, Science **330**, 1072 (2010).
- [64] Y. Xiao, Y. Xiang, Y. Q. He, B. C. Sanders, arXiv:1807.07829 (2018).
- [65] G. H. Hardy, J. E. Littlewood, G. Polya, *Inequalities*, Cambridge Univ. Press, London (1952).
- [66] Phoebus J. Dhrymes, *Mathematics for econometrics*, Springer (1978).
- [67] J. Watrous, *Theory of quantum information*, Cambridge Univ. Press, Cambridge (2011).
- [68] S. K. Goyal, B. Neethi Simon, R. Singh, and S. Simon, J. Phys. A: Math. Theor. **49**, 165203 (2016).