

Tripartite realism-based quantum nonlocality

D. M. Fucci and R. M. Angelo

Department of Physics, Federal University of Paraná, P.O. Box 19044, 81531-980 Curitiba, Paraná, Brazil

From an operational criterion of physical reality, a quantifier of realism-based nonlocality was recently introduced for two-part quantum states. This measure has shown to capture aspects that are rather different from Bell nonlocality. Here we take a step further and introduce a tripartite realism-based nonlocality quantifier. We show that this measure reduces to genuine tripartite entanglement for a certain class of pure tripartite states and manifests itself in correlated mixed states even in the absence of quantum correlations. A case study for noisy GHZ and W states points out the existence of scenarios where the realism-based nonlocality is monogamous.

I. INTRODUCTION

One of the most astonishing aspects of nature is unveiled by Bell's theorem, which proves that the predictions of quantum mechanics are incompatible with physical theories relying upon the local causality hypothesis [1], which is mathematically formulated as $p(a, b|A, B) = \sum_{\lambda} p_{\lambda} p(a|A, \lambda) p(b|B, \lambda)$. This relation assumes that the joint probability $p(a, b|A, B)$ of finding the outcomes a and b in measurements of observables A and B , respectively, is factorable into local probability distributions, $p(a|A, \lambda)$ and $p(b|B, \lambda)$, by virtue of the specification of a hidden variable λ governed by a probability distribution p_{λ} . Today, it is a fact firmly backed by solid experimental evidence [2–8] that nature does not submit itself to such hypothesis. This suggests that “spooky actions” may take place at distance, a phenomenon known as Bell nonlocality [9].

Quantum mechanics provides accurate accounts for experimental data. On the one hand, it is known that separable states $\rho_s = \sum_{\lambda} p_{\lambda} \rho_{\lambda}^A \otimes \rho_{\lambda}^B$ on $\mathcal{H}_A \otimes \mathcal{H}_B$ produce distributions $p(a, b|A, B) = \text{Tr}(A_a \otimes B_b \rho_s)$ admitting the factorization prescribed by the local causality hypothesis, for projectors A_a on \mathcal{H}_A and B_b on \mathcal{H}_B . In this case, no Bell inequality is violated and the ρ_s is said Bell local. On the other hand, quantum mechanics predicts that all entangled pure states are Bell nonlocal [10, 11]. For mixed states, Bell nonlocality is known to demand entanglement, while the converse is not true [9], which implies that the class of Bell nonlocal states form a subset of the entangled states. The existence of maximally entangled states that are not maximally Bell nonlocal [12–15]—the so called anomaly—reveals how tricky the quantification of Bell nonlocality may be. In effect, unlike entanglement quantification [16], the task of quantifying Bell nonlocality still elicits debate. Some of the recent approaches make reference to maximal violations of Bell inequalities [17], performances in communication tasks [18–22], noise resistance [23, 24], and the volume of violation [25], the latter solving the anomaly problem for specific states.

Even though Bell's original argument makes no formal link with realism [26], one can argue that this concept—actually the absence of it—is mandatory for the manifestation of nonlocality. The example given in Ref. [27] illustrates this point. Suppose that two qubits share a singlet state, that is, a physical preparation that constraints the total spin as $s_z = s_z^A + s_z^B = 0$. The definiteness of the total spin allows us to say that it is an element of reality, even though the spins of the parts are not. This conservation law is preserved as the spins are sent

to far distant locations. Now, if s_z^A is measured and becomes an element of reality, the conservation law forces s_z^B to instantaneously become so. Since the emergence of reality in part B is then causally induced by a remote disturbance, some aspect of nonlocality is presumed to take place. Inspired by the lack of such a clear link between realism and violations of the local causality hypothesis, a notion of nonlocality has recently been put forward that makes explicit use of a criterion of realism [27]. This novel aspect of nonlocality counts with a nonanomalous quantifier, is remarkably more resilient to noise than other quantum resources, and reduces to bipartite entanglement for pure states [28, 29].

While on the Bell nonlocality side, a significant literature exists concerning multipartite settings [30–35], it is still not clear whether it is possible to extend the notion of realism-based nonlocality even to tripartite scenarios. This work aims at starting this research program. Our basic strategy consists of performing a slight adaptation of the current measure of realism-based nonlocality so as to make it applicable to all possible bipartitions of a tripartite state. Then, we construct a quantifier of genuine tripartite realism-based nonlocality, derive its properties, and prove that it reduces to genuine tripartite entanglement for a given class of pure states. In addition, we discuss whether the introduced measure is monogamous.

II. PRELIMINARY CONCEPTS

Here we present a brief review of the realism-based nonlocality. As seminally introduced by Einstein, Podolsky, and Rosen (EPR) [36], the concept of “elements of reality” was related with the idea of full predictability without disturbance. This criterion, however, fails to diagnose situations where the elements of reality cannot be predicted because of mere subjective ignorance. For instance, if Alice measures the z -spin of a particle but does not let Bob know the outcome, one would not say that the measured spin is not an element of reality just because Bob cannot predict the result before the next measurement on this particle. To overcome this difficulty, Bilobran and Angelo introduced an operational criterion to identify elements of reality for a generic preparation ρ on $\mathcal{H}_A \otimes \mathcal{H}_B$ [27]. Their key premise is that once a discrete-spectrum observable $A = \sum_a a A_a$ is measured, there must be an element of reality associated with A , even when the measurement outcome is not

revealed. The post-measurement state reads

$$\sum_a (A_a \otimes \mathbb{1}_B) \rho (A_a \otimes \mathbb{1}_B) = \sum_a p_a A_a \otimes \rho_{B|a} =: \Phi_A(\rho), \quad (1)$$

where $p_a = \text{Tr}(A_a \otimes \mathbb{1}_B \rho)$, $\rho_{B|a} = \text{Tr}_{\mathcal{A}}(A_a \otimes \mathbb{1}_B \rho) / p_a$, and $A_a A_{a'} = A_a \delta_{aa'}$. The unrevealed measurement of A is henceforth denoted by Φ_A , which is a completely positive trace-preserving unital map. Bilobran and Angelo then take $\Phi_A(\rho) = \rho$ as a criterion of realism with basis on the following rationale: if measuring A and not revealing the outcome do not effectively change the state of the system, then this state is just an epistemic description of the part \mathcal{A} , where A is already an element of reality. As expected, one has $\Phi_A \Phi_A(\rho) = \Phi_A(\rho)$, meaning that successive applications of an unrevealed measurement of A over an A -reality state does not create “irreality” for A . It is then proposed the measure

$$\mathfrak{I}_A(\rho) := S(\Phi_A(\rho)) - S(\rho) \quad (2)$$

of the irreality of A for a given preparation ρ , where S denotes the von Neumann entropy. It can be shown that irreality is nonnegative, vanishes if and only if $\rho = \Phi_A(\rho)$, and is nonincreasing under completely positive trace-preserving maps. By use of \mathfrak{I}_A , an information-reality complementarity relation was recently derived for generic unrevealed measurements [37], with some results having been experimentally verified through a photonics platform [38].

In the same work, Bilobran and Angelo propose to use measure (2) to quantify by how much the irreality of an observable A in the site \mathcal{A} changes due to unrevealed measurements of an observable B acting on part \mathcal{B} . The authors then introduced the contextual realism-based nonlocality

$$\eta_{A|B}(\rho) := \mathfrak{I}_A(\rho) - \mathfrak{I}_A(\Phi_B(\rho)), \quad (3)$$

where $B = \sum_b b B_b$ and $\Phi_B(\rho) = \sum_b (\mathbb{1}_{\mathcal{A}} \otimes B_b) \rho (\mathbb{1}_{\mathcal{A}} \otimes B_b)$, for projectors B_b . Here the context is defined by the pair $\{A, B\}$. It has been shown that $\eta_{A|B}(\rho) \geq 0$, with equality holding for product states, $\rho = \rho_{\mathcal{A}} \otimes \rho_{\mathcal{B}}$, and for states of reality, that is, $\rho = \Phi_A(\rho)$, $\rho = \Phi_B(\rho)$, or $\rho = \Phi_{A,B}(\rho) \equiv \Phi_A \Phi_B(\rho)$.

By maximizing the contextual realism-based nonlocality over all possible contexts, Gomes and Angelo then introduced the bipartite realism-based nonlocality [28]

$$\mathcal{N}_2(\rho) := \max_{\{A,B\}} \eta_{A|B}(\rho), \quad (4)$$

which diagnoses the nonlocality of the state ρ on $\mathcal{H}_{\mathcal{A}} \otimes \mathcal{H}_{\mathcal{B}}$. This quantity has shown to be nonanomalous, since for maximally entangled bipartite states, $|\psi\rangle = \sum_{i=1}^d |i\rangle |i\rangle / \sqrt{d}$, it reduces to the entanglement E of $|\psi\rangle$, that is, $\mathcal{N}_2(\varrho) = S(\varrho_{\mathcal{R}}) \equiv E(\varrho)$, where $\mathcal{R} \in \{\mathcal{A}, \mathcal{B}\}$, $\varrho = |\psi\rangle \langle \psi|$, and $\varrho_{\mathcal{A}(\mathcal{B})} = \text{Tr}_{\mathcal{B}(\mathcal{A})} \varrho$. A distinctive feature of \mathcal{N}_2 can be readily appreciated for the classical-classical state $\rho_{cc} = \sum_i p_i A'_i \otimes B'_i$, where $A' = \sum_i a'_i A'_i$ and $B' = \sum_i b'_i B'_i$. As shown in Ref. [29] this state has none of the well-established nonclassical features, namely, Bell nonlocality, EPR steering, entanglement, and quantum discord. Still, by choosing a context $\{A, B\}$ maximally incompatible with $\{A', B'\}$ we find $\mathcal{N}_2(\rho_{cc}) = \eta_{A|B}(\rho_{cc}) = H(\{p_i\}) > 0$, where

$H(\{p_i\})$ is the Shannon entropy of the probability distribution p_i . Since $\eta_{A'|B'}(\rho_{cc}) = 0$, one sees that \mathcal{N}_2 is able to capture aspects of incompatibility which suffice to produce realism changes at distance.

III. TRIPARTITE REALISM-BASED NONLOCALITY

With basis on Bilobran and Angelo’s approach to realism and nonlocality, we now construct the notion of genuine tripartite realism-based nonlocality. Hereafter, for the sake of notational simplicity, we reserve the term nonlocality for referring to the bi- and tripartite versions of the realism-based nonlocality, in distinction to Bell nonlocality. Consider a tripartite preparation $\rho \in \mathcal{H}_{\mathcal{A}} \otimes \mathcal{H}_{\mathcal{B}} \otimes \mathcal{H}_{\mathcal{C}}$ such that the parts of the system are sent to distinct laboratories, \mathcal{A} , \mathcal{B} , and \mathcal{C} , which are far apart from each other. Let $R \in \{A, B, C\}$ on $\mathcal{H}_{\mathcal{R}}$ be an observable accessible only in the laboratory $\mathcal{R} \in \{\mathcal{A}, \mathcal{B}, \mathcal{C}\}$. We then introduce the bipartite contextual nonlocality

$$\eta_{A|B,C}(\rho) := \mathfrak{I}_A(\rho) - \mathfrak{I}_A(\Phi_{B,C}(\rho)), \quad (5)$$

where $\Phi_{B,C} = \Phi_B \Phi_C$. This formula applies for any permutations of A , B , and C . Being a natural extension of definition (3), $\eta_{A|B,C}$ captures changes in the reality of A given that local unrevealed measurements Φ_B and Φ_C are conducted in the remote labs \mathcal{B} and \mathcal{C} . Because irreality never increases upon completely positive trace-preserving maps, it follows that $\eta_{A|B,C}(\rho) \geq 0$, the equality applying for states of reality $\rho = \Phi_A(\rho)$ and $\rho = \Phi_{B,C}(\rho)$, for the full realism state $\bigotimes_{\mathcal{R}=\mathcal{A},\mathcal{B},\mathcal{C}} \frac{\mathbb{1}_{\mathcal{R}}}{d_{\mathcal{R}}} \equiv \frac{\mathbb{1}}{d}$, where $d = d_{\mathcal{A}} d_{\mathcal{B}} d_{\mathcal{C}}$, and for uncorrelated states as $\rho = \rho_{\mathcal{A}} \otimes \rho_{\mathcal{BC}}$. From this we see that irreality and correlations are prerequisites for nonlocality. It is also noteworthy that $\eta_{A|B,C}(\rho_{\mathcal{AB}} \otimes \rho_{\mathcal{C}}) = \eta_{A|B}(\rho_{\mathcal{AB}})$, which is desirable since part \mathcal{C} is fully irrelevant in this case. The same applies if \mathcal{B} is uncorrelated.

By maximizing over all trios $\{A, B, C\}$ of local observables, we introduce the amount of nonlocality $\mathcal{N}_{\mathcal{A|BC}}$ associated with realism changes in part \mathcal{A} induced by local measurements in parts \mathcal{B} and \mathcal{C} for a given preparation ρ , that is,

$$\mathcal{N}_{\mathcal{A|BC}}(\rho) := \max_{\{A,B,C\}} \eta_{A|B,C}(\rho). \quad (6)$$

When $\mathcal{N}_{\mathcal{A|BC}}(\rho) > 0$ we are certain that there exists at least one setting $\{A, B, C\}$ through which one is able to spot a change in the reality of A when B and C are measured. It can be verified that $\mathcal{N}_{\mathcal{A|BC}}(\rho)$ vanishes when $\rho = \mathbb{1}/d$ or $\rho = \rho_{\mathcal{A}} \otimes \rho_{\mathcal{BC}}$. In addition, one has $\mathcal{N}_{\mathcal{A|BC}}(\rho_{\mathcal{AB}} \otimes \rho_{\mathcal{C}}) = \mathcal{N}_2(\rho_{\mathcal{AB}})$.

Now, to build our quantifier of genuine tripartite nonlocality, we invoke Bennett *et al.*’s proposal [39] for the identification of genuine multipartite correlations: “A state of n particles has genuine n -partite correlations if it is nonproduct in every bipartite cut”. As shown by Ma *et al.* [40], this approach is able to produce, for instance, a measure of genuine tripartite entanglement for pure states, namely,

$$E_3(\rho) := \min \{E_{\mathcal{A|BC}}(\rho), E_{B|\mathcal{AC}}(\rho), E_{C|\mathcal{AB}}(\rho)\}, \quad (7)$$

where $E_{\mathcal{A|BC}}(\rho) = S(\rho_{\mathcal{A}})$ is the entanglement entropy of the partition $\mathcal{A|BC}$, with a similar interpretation for the other

terms. Being different from quantum discord, $\mathcal{N}_{\mathcal{A}|\mathcal{B}\mathcal{C}}$ cannot be termed a strict measure of quantum correlations. Still, since it makes reference to nonlocal connections between the parts, the above rationale involving bipartite cuts is applicable. This allows us to introduce the quantifier of genuine tripartite realism-based nonlocality:

$$\mathcal{N}_3(\rho) := \min \{ \mathcal{N}_{\mathcal{A}|\mathcal{B}\mathcal{C}}(\rho), \mathcal{N}_{\mathcal{B}|\mathcal{A}\mathcal{C}}(\rho), \mathcal{N}_{\mathcal{C}|\mathcal{A}\mathcal{B}}(\rho) \}. \quad (8)$$

From what we have for measure (6), it follows that \mathcal{N}_3 vanishes for states like $\mathbb{1}/d$ and $\rho_{\mathcal{A}} \otimes \rho_{\mathcal{B}\mathcal{C}}$ (including states with permuted indexes).

We now prove an interesting result with respect to the special class of pure states $|\varphi\rangle$ admitting a Schmidt decomposition $|\varphi\rangle = \sum_i \sqrt{\xi_i} |\alpha_i\rangle |\beta_i\rangle |\gamma_i\rangle$ [41], for bases connected with the observables $\alpha = \sum_i \alpha_i |\alpha_i\rangle \langle \alpha_i|$, $\beta = \sum_i \beta_i |\beta_i\rangle \langle \beta_i|$, and $\gamma = \sum_i \gamma_i |\gamma_i\rangle \langle \gamma_i|$ acting on respective spaces $\mathcal{H}_{\mathcal{A},\mathcal{B},\mathcal{C}}$. We start by plugging definition (2) into the contextual nonlocality, (5), with $\varsigma = |\varphi\rangle \langle \varphi|$, to write

$$\eta_{\mathcal{A}|\mathcal{B},\mathcal{C}}(\varsigma) = S(\Phi_{\mathcal{A}}(\varsigma)) + S(\Phi_{\mathcal{B},\mathcal{C}}(\varsigma)) - S(\Phi_{\mathcal{A},\mathcal{B},\mathcal{C}}(\varsigma)) - S(\varsigma). \quad (9)$$

For pure states, $S(\varsigma) = 0$. Given the entropy monotonicity, $S(\Phi_R(\varsigma)) \geq S(\varsigma)$, which implies that $S(\Phi_{R,Q}(\varsigma)) \geq S(\Phi_R(\varsigma))$ and $S(\Phi_{\mathcal{A},\mathcal{B},\mathcal{C}}(\varsigma)) \geq S(\Phi_{R,Q}(\varsigma))$, with $R, Q \in \{\mathcal{A}, \mathcal{B}, \mathcal{C}\}$, we find

$$2S(\Phi_{\mathcal{A},\mathcal{B},\mathcal{C}}(\varsigma)) \geq S(\Phi_{\mathcal{A}}(\varsigma)) + S(\Phi_{\mathcal{B},\mathcal{C}}(\varsigma)), \quad (10)$$

which saturates when $\Phi_{\mathcal{A},\mathcal{B},\mathcal{C}}(\varsigma) = \Phi_{\mathcal{B},\mathcal{C}}(\varsigma) = \Phi_{\mathcal{A}}(\varsigma)$. Combining this inequality with Eq. (9) gives

$$\eta_{\mathcal{A}|\mathcal{B},\mathcal{C}}(\varsigma) \leq \frac{1}{2} [S(\Phi_{\mathcal{A}}(\varsigma)) + S(\Phi_{\mathcal{B},\mathcal{C}}(\varsigma))]. \quad (11)$$

Therefore, the maximization of $\eta_{\mathcal{A}|\mathcal{B},\mathcal{C}}(\varsigma)$ will come by the saturation of this inequality. This is accomplished by the choice $A = \alpha$, $B = \beta$, and $C = \gamma$, which can be shown to give $\Phi_{\mathcal{A}}(\varsigma) = \Phi_{\beta,\gamma}(\varsigma) = \Phi_{\alpha,\beta,\gamma}(\varsigma) = \sum_i \xi_i |\alpha_i\rangle \langle \alpha_i| \otimes |\beta_i\rangle \langle \beta_i| \otimes |\gamma_i\rangle \langle \gamma_i|$. Hence, $\mathcal{N}_{\mathcal{A}|\mathcal{B}\mathcal{C}}(\varsigma) = \eta_{\alpha|\beta,\gamma}(\varsigma) = S(\Phi_{\mathcal{A}}(\varsigma)) = H(\{\xi_i\})$. Given the symmetry of $|\varphi\rangle$, one does not expect different results for the other bipartitions, so that $\mathcal{N}_3(\varsigma) = H(\{\xi_i\})$. The entanglement of the cut $\mathcal{A}|\mathcal{B}\mathcal{C}$ is given by $E_{\mathcal{A}|\mathcal{B}\mathcal{C}}(\varsigma) = S(\text{Tr}_{\mathcal{A}}\varsigma) = H(\{\xi_i\})$. Using symmetry again and definition (7), we obtain $E_3(\varsigma) = H(\{\xi_i\})$, which finally gives

$$\mathcal{N}_3(\varsigma) = E_3(\varsigma) \quad (\varsigma = |\varphi\rangle \langle \varphi|). \quad (12)$$

This implies that, as far as the class $|\varphi\rangle$ of tripartite pure states is concerned, \mathcal{N}_3 is surely nonanomalous with respect to genuine tripartite entanglement as measured by the formula (7).

Consider now a classical-classical-classical state, defined as $\rho_{\text{ccc}} = \sum_i p_i A'_i \otimes B'_i \otimes C'_i$, with $A' = \sum_i a'_i A'_i$, $B' = \sum_i b'_i B'_i$, and $C' = \sum_i c'_i C'_i$. It is clear that this is a separable state, thus possessing no form of entanglement. Yet, it does present tripartite nonlocality. To see this, let the context $\{\mathcal{A}, \mathcal{B}, \mathcal{C}\}$ be maximally incompatible with $\{A', B', C'\}$. Since $\Phi_{\mathcal{A}}(A'_i) = \mathbb{1}_{\mathcal{A}}/d_{\mathcal{A}}$, with similar relations for the other parts, direct calculations give $\eta_{\mathcal{A}|\mathcal{B},\mathcal{C}}(\rho_{\text{ccc}}) = H(\{p_i\})$. This guarantees that $\mathcal{N}_{\mathcal{A}|\mathcal{B}\mathcal{C}}(\rho_{\text{ccc}}) > 0$ and, via symmetry, that $\mathcal{N}_3(\rho_{\text{ccc}}) > 0$. In other words, just as \mathcal{N}_2 , tripartite nonlocality may manifest itself even when no quantum correlation is present.

IV. APPLICATIONS

Here we calculate \mathcal{N}_3 for the noisy three-qubit states

$$\rho_n^\chi := n \frac{\mathbb{1}}{8} + (1-n) |\Psi_\chi\rangle \langle \Psi_\chi|, \quad (13)$$

where $n \in [0, 1]$ gives the noise (or impurity) added to the tripartite pure state $|\Psi_\chi\rangle$, which, with $\chi \in \{\text{GHZ}, \text{W}\}$, assumes

$$|\Psi_{\text{GHZ}}\rangle = \frac{1}{\sqrt{2}} (|000\rangle + |111\rangle), \quad (14a)$$

$$|\Psi_{\text{W}}\rangle = \frac{1}{\sqrt{3}} (|100\rangle + |010\rangle + |001\rangle). \quad (14b)$$

Even for these relatively simple states, the evaluation of the genuine tripartite nonlocality \mathcal{N}_3 is a hard computational problem due to the optimizations demanded by definitions (6) and (8). Incidentally, due to the subsystem-permutation symmetry, we have $\mathcal{N}_3 = \mathcal{N}_{\mathcal{A}|\mathcal{B}\mathcal{C}} = \mathcal{N}_{\mathcal{B}|\mathcal{A}\mathcal{C}} = \mathcal{N}_{\mathcal{C}|\mathcal{A}\mathcal{B}}$, which simplifies our task and makes it numerically feasible. To obtain $\mathcal{N}_{\mathcal{A}|\mathcal{B}\mathcal{C}}$ via maximization of $\eta_{\mathcal{A}|\mathcal{B},\mathcal{C}}$, we considered spin operators $A = \hat{a} \cdot \vec{\sigma}$ with $\hat{a} = (\sin \theta_a \cos \varphi_a, \sin \theta_a \sin \varphi_a, \cos \theta_a)$ and $\vec{\sigma} = (\sigma_x, \sigma_y, \sigma_z)$, for Pauli matrices $\sigma_{x,y,z}$. Similarly, we parametrized B and C with unit vectors \hat{b} and \hat{c} , respectively, thus getting the angles $\theta_{a,b,c} \in [0, \pi]$ and $\varphi_{a,b,c} \in [0, 2\pi]$, over which the optimization were performed. We then defined a grid by letting these angles vary in their domain with increments of $\pi/8$, which yielded a set of 2,985,984 distinct settings $\{\mathcal{A}, \mathcal{B}, \mathcal{C}\}$. We computed $\eta_{\mathcal{A}|\mathcal{B}\mathcal{C}}$ for each setting of the set and then picked the maximum. This process was repeated for each value n of noise, which ranged in the domain $[0, 1]$ with increments of 0.01, and for $\chi = \{\text{GHZ}, \text{W}\}$. Our results are presented in Fig. 1. Such numerical analysis was realized also for 10^6 randomly generated settings $\{\mathcal{A}, \mathcal{B}, \mathcal{C}\}$ and no appreciable difference was found, meaning that these results are statistically reliable. Throughout our numerical investigations, we found some noteworthy settings. For the GHZ state ($n = 0$), we found that $\eta_{\mathcal{A}|\mathcal{B},\mathcal{C}}$ reaches the maximum value $\ln 2$ when $A = \sigma_z$ and at least one of the observables B and C is equal to A . The contextual nonlocality is also maximal when (A, B, C) assumes $(\sigma_x, \sigma_x, \sigma_x)$, $(\sigma_y, \sigma_y, \sigma_x)$, $(\sigma_y, \sigma_x, \sigma_y)$, or $(\sigma_x, \sigma_y, \sigma_y)$, a set of measurement operators that when acting over a GHZ state is known to provide predictions that are in conflict with the local realism hypothesis [42]. For the W state, the maximum contextual nonlocality, 0.6364, was found for $A = B = C = \sigma_z$.

Notably, the genuine tripartite nonlocality \mathcal{N}_3 is a monotonically decreasing function of the noise n , strictly vanishing only in the scenario of no purity whatsoever ($n = 1$). This points out the stronger resilience of \mathcal{N}_3 in comparison with other measures of nonclassicality which abruptly vanish under high levels of noise. For instance, as far as ρ_n^{GHZ} is concerned, it is known that tripartite entanglement disappears for $n \geq 4/5$ [43, 44], Bell nonlocality for $n > 1/2$ (with two up to five measurements per site) [45], and steering for $n \geq 0.225$ [46]. For ρ_n^{W} , entanglement vanishes for $n \geq 0.8220$ [48], Bell nonlocality for $n > 0.3558$ (0.3952) with two (three) measurements per site [45], and steering for $n \geq 0.1634$ [46]. An equivalent noise resilience was verified for \mathcal{N}_2 in Ref. [28].




FIG. 1. Genuine tripartite nonlocality $N_3(\rho_n^\chi)$ for the noisy three-qubit states (13) as a function of the noise amount η . Blue circles correspond to the noisy GHZ state ($\chi = \text{GHZ}$) and red squares to the noisy W state ($\chi = \text{W}$). Tripartite nonlocality monotonically decreases with noise.

V. MONOGAMY OF NONLOCALITY

A monogamy inequality $Q_{ABC} \geq Q_{AB} + Q_{AC}$ for a given resource measure Q gives an upper bound to the shareability of the related resource among the parts of the system. We now assess whether the genuine tripartite nonlocality N_3 exhibits this property. In fact, we want to test the relation

$$N_3^\alpha(\rho_{ABC}) \geq N_2^\alpha(\rho_{AB}) + N_2^\alpha(\rho_{AC}), \quad (15)$$

where ρ_{AB} and ρ_{AC} are reduced states, and $\alpha \in \mathbb{R}_{>0}$ is a parameter intended to give to a further measure N_3^α , monotonically related with N_3 , an extra chance to satisfy monogamy. This strategy has proven successful in establishing monogamy for general measures of nonclassicality [47].

We start by proving that N_3^α is not monogamous in general. Consider the tripartite GHZ state $\rho_{ABC} = \rho_{n=0}^{\text{GHZ}}$, whose reduced states read $\rho_{AB} = \rho_{AC} = (|00\rangle\langle 00| + |11\rangle\langle 11|)/2 \equiv \rho_{cc}$. We then find $N_3(\rho_{ABC}) = N_2(\rho_{AB}) = N_2(\rho_{AC}) = \ln 2$, which, for all α , can never satisfy the monogamy relation (15). Interestingly, however, violations of monogamy for the state ρ_n^χ tend to be relatively rare in the parameter space. To show this, we present in Fig. 2 numerical results for the quantity

$$\delta N_3^\alpha(\rho_{ABC}) := N_3^\alpha(\rho_{ABC}) - [N_2^\alpha(\rho_{AB}) + N_2^\alpha(\rho_{AC})], \quad (16)$$

with $\rho_{ABC} = \rho_n^\chi$. This is a quantifier that, whenever nonnegative, witnesses monogamy for the measure N_3^α (see the colored region in Fig. 2). The results show that monogamy is prevented for small values of $\{\eta, \alpha\}$ and is saturated for large values. Also, they confirm that the pure GHZ state does not allow for monogamy and that for the pure W state monogamy holds for $\alpha \gtrsim 2.1641$, with $\delta N_3^\alpha(\rho_{n=0}^{\text{W}})$ peaking at $\alpha \cong 3.8372$.

VI. CONCLUDING REMARKS

With basis on the criterion of reality proposed in Ref. [27], we introduce a bipartite realism-based nonlocality measure that applies for three parts and, from that, a genuine tripartite realism-based nonlocality measure. This quantifier is shown to reduce, for a given class of pure states, to genuine tripartite entanglement and to manifest itself even in cases where no quantum correlations are present. In addition, genuine tripartite realism-based nonlocality turns out to be greatly resistant to noise and deformable into further monogamous quantities for some states. This work paves the way to incursions on the unexplored realm of n -partite realism-based nonlocality.

ACKNOWLEDGMENTS

This work was financed in part by the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior - Brasil (CAPES) - Finance Code 001. R.M.A acknowledges CNPq (Brazil) and the National Institute for Science and Technology of Quantum Information (INCT-IQ/CNPq).




FIG. 2. Contour plots for the normalized monogamy witness $\delta N_3^\alpha(\rho_n^\chi)/N_\chi$ (see color code), with the respective normalization factors $N_{\text{GHZ}} \cong 0.019997$ and $N_{\text{W}} \cong 0.073296$, for the state ρ_n^χ , as a function of the parameters α and η . Monogamy holds in the whole colored regions, where $\delta N_3^\alpha \geq 0$, and does not apply for the pure GHZ state (left panel with $\eta = 0$ and $\forall \alpha > 0$).

- [1] J. S. Bell, On the Einstein-Podolsky-Rosen paradox, *Physics* **1**, 195 (1964).
- [2] B. Hensen *et al.*, Loophole-free Bell inequality violation using electron spins separated by 1.3 kilometres, *Nature (London)* **526**, 682 (2015).
- [3] M. Giustina *et al.*, Significant Loophole-Free Test of Bells Theorem with Entangled Photons, *Phys. Rev. Lett.* **115**, 250401

(2015).

- [4] L. K. Shalm *et al.*, Strong Loophole-Free Test of Local Realism, *Phys. Rev. Lett.* **115**, 250402 (2015).
- [5] B. Hensen *et al.*, Loophole-free Bell test using electron spins in diamond: Second experiment and additional analysis, *Sci. Rep.* **6**, 30289 (2016).

- [6] W. Rosenfeld *et al.*, Event-Ready Bell Test Using Entangled Atoms Simultaneously Closing Detection and Locality Loopholes, *Phys. Rev. Lett.* **119**, 010402 (2017).
- [7] J. Handsteiner *et al.*, Cosmic Bell Test: Measurement Settings from Milky Way Stars, *Phys. Rev. Lett.* **118**, 060401 (2017).
- [8] D. Rauch *et al.*, Cosmic Bell Test using Random Measurement Settings from High-Redshift Quasars, *Phys. Rev. Lett.* **121**, 080403 (2018).
- [9] N. Brunner, D. Cavalcanti, S. Pironio, V. Scarani, and S. Wehner, Bell nonlocality, *Rev. Mod. Phys.* **86**, 419 (2014); **86**, 839 (2014).
- [10] N. Gisin, Bells inequality holds for all non-product states, *Phys. Lett. A* **154**, 201 (1991).
- [11] S. Yu, Q. Chen, C. Zhang, C. H. Lai, and C. H. Oh, All Entangled Pure States Violate a Single Bells Inequality, *Phys. Rev. Lett.* **109**, 120402(2012).
- [12] A. Acín, T. Durt, N. Gisin, and J. I. Latorre, Quantum nonlocality in two three-level systems, *Phys. Rev. A* **65**, 052325 (2002).
- [13] A. A. Méthot and V. Scarani, An anomaly of non-locality, *Quantum Inf. Comput.* **7**, **157** (2007).
- [14] T. Vidick and S. Wehner, More nonlocality with less entanglement, *Phys. Rev. A* **83**, 052310 (2011).
- [15] S. Camalet, Measure-independent anomaly of nonlocality, *Phys. Rev. A* **96**, 052332 (2017).
- [16] R. Horodecki, P. Horodecki, M. Horodecki, and K. Horodecki, Quantum entanglement, *Rev. Mod. Phys.* **81**, 865 (2009).
- [17] A. C. S. Costa, M. W. Beims, and R. M. Angelo, Generalized discord, entanglement, Einstein-Podolsky-Rosen steering, and Bell nonlocality in two-qubit systems under (non-)Markovian channels: Hierarchy of quantum resources and chronology of deaths and births, *Physica A* **461**, 469 (2016).
- [18] T. Maudlin, in *Proceedings of the 1992 Meeting of the Philosophy of Science Association*, edited by D. Hull, M. Forbes, and K. Okruhlik (Philosophy of Science Association, East Lansing, 1992), Vol. 1, pp. 404417.
- [19] G. Brassard, R. Cleve, and A. Tapp, Cost of Exactly Simulating Quantum Entanglement with Classical Communication, *Phys. Rev. Lett.* **83**, 1874 (1999).
- [20] M. Steiner, Towards quantifying non-local information transfer: Finite-bit non-locality, *Phys. Lett. A* **270**, 239 (2000).
- [21] D. Bacon and B. F. Toner, Bell Inequalities with Auxiliary Communication, *Phys. Rev. Lett.* **90**, 157904 (2003).
- [22] C. Branciard and N. Gisin, Quantifying the Nonlocality of Greenberger-Horne-Zeilinger Quantum Correlations by a Bounded Communication Simulation Protocol, *Phys. Rev. Lett.* **107**, 020401 (2011).
- [23] D. Kaszlikowski, P. Gnacinski, M. Zukowski, W. Miklaszewski, and A. Zeilinger, Violations of Local Realism by Two Entangled N-Dimensional Systems Are Stronger than for Two Qubits, *Phys. Rev. Lett.* **85**, 4418 (2000).
- [24] W. Laskowski, J. Ryu, and M. Zukowsky, Noise resistance of the violation of local causality for pure three-qutrit entangled states, *J. Phys. A* **47**, 424019 (2014).
- [25] E. A. Fonseca and F. Parisio, Measure of nonlocality which is maximal for maximally entangled qutrits, *Phys. Rev. A* **92**, 030101(R) (2015).
- [26] N. Gisin, Non-realism: Deep thought or a soft option?, *Found. Phys.* **42**, **80** (2012).
- [27] A. L. O. Bilobran and R. M. Angelo, A measure of physical reality, *Europhys. Lett.* **112**, 40005 (2015).
- [28] V. S. Gomes and R. M. Angelo, Nonanomalous measure of realism-based nonlocality, *Phys. Rev. A* **97**, 012123 (2018).
- [29] V. S. Gomes and R. M. Angelo, Resilience of realism-based nonlocality to local disturbance, *Phys. Rev. A* **99**, 012109 (2019).
- [30] G. Svetlichny, Distinguishing three-body from two-body nonseparability by a Bell-type inequality, *Phys. Rev. D* **35**, 3066 (1987).
- [31] J.-D. Bancal, C. Branciard, N. Gisin, and S. Pironio, Quantifying multipartite nonlocality, *Phys. Rev. Lett.* **103**, 090503 (2009).
- [32] J.-D. Bancal, N. Brunner, N. Gisin, Y.-C. Liang, Detecting genuine multipartite quantum nonlocality: A simple approach and generalization to arbitrary dimensions, *Phys. Rev. Lett.* **106**, 020405 (2011).
- [33] J.-D. Bancal, J. Barrett, N. Gisin, S. Pironio, Definitions of multipartite nonlocality, *Phys. Rev. A* **88**, 014102 (2013).
- [34] A. de Rosier, J. Gruca, F. Parisio, T. Vértesi, and W. Laskowski, Multipartite nonlocality and random measurements, *Phys. Rev. A* **96**, 012101 (2017).
- [35] R. Chaves, D. Cavalcanti, and L. Aolita, Causal hierarchy of multipartite Bell nonlocality, *Quantum* **1**, 23 (2017).
- [36] A. Einstein, B. Podolsky, and N. Rosen, Can Quantum-Mechanical Description of Physical Reality Be Considered Complete?, *Phys. Rev.* **47**, 777 (1935).
- [37] P. R. Dieguez and R. M. Angelo, Information-reality complementarity: The role of measurements and quantum reference frames, *Phys. Rev. A* **97**, 022107 (2018).
- [38] L. Mancino, M. Sbroscia, E. Roccia, I. Gianani, V. Cimini, M. Paternostro, and M. Barbieri, Information-reality complementarity in photonic weak measurements, *Phys. Rev. A* **97**, 062108 (2018).
- [39] C. H. Bennett, A. Grudka, M. Horodecki, P. Horodecki, and R. Horodecki, Postulates for measures of genuine multipartite correlations, *Phys. Rev. A* **83**, 012312 (2011).
- [40] Z.-H. Ma, Z.-H. Chen, J.-L. Chen, C. Spengler, A. Gabriel, and M. Huber, Measure of genuine multipartite entanglement with computable lower bounds, *Phys. Rev. A* **83**, 062325 (2011).
- [41] Arun K. Pati, Existence of the Schmidt decomposition for tripartite systems, *Phys. Lett. A* **278**, 118 (2000).
- [42] D. Bouwmeester, A. Ekert and A. Zeilinger, *The Physics of Quantum Information: Quantum Cryptography, Quantum Teleportation, Quantum Computation* (Springer, Berlin Heidelberg, 2000), pp. 198200.
- [43] R. Schack and C. M. Caves, Explicit product ensembles for separable quantum states, *J. Mod. Opt.* **47**, 387, (2000).
- [44] W. Dür and J. I. Cirac, Classification of multi-qubit mixed states: separability and distillability properties, *Phys. Rev. A* **61**, 042314 (2000).
- [45] J. Gruca, W. Laskowski, M. Zukowski, N. Kiesel, W. Wic-zorek, C. Schmid, H. Weinfurter, Non-classicality thresholds for multiqubit states - numerical analysis, *Phys. Rev. A* **82**, 012118 (2010).
- [46] A. C. S. Costa, R. Uola and O. Gühne, Entropic Steering Criteria: Applications to Bipartite and Tripartite Systems, *Entropy* **20**, 763 (2018).
- [47] Z.-X. Jin and S.-M. Fei, Monogamy relations of all quantum correlation measures for multipartite quantum systems, *Opt. Commun.* **446**, 39 (2019).
- [48] Z.-H. Chen, Z.-H. Ma, O. Gühne, S. Severini, Estimating entanglement monotones with a generalization of the Wootters formula, *Phys. Rev. Lett.* **109**, 200503 (2012).