

Beyond mean-field corrections to the quasiparticle spectrum of superfluid Fermi gases

Senne Van Loon,^{*} Jacques Tempere, and Hadrien Kurkjian
TQC, Universiteit Antwerpen, Universiteitsplein 1, B-2610 Antwerpen, België
 (Dated: December 13, 2019)

We investigate the fermionic quasiparticle branch of superfluid Fermi gases in the BCS-BEC crossover and calculate the quasiparticle lifetime and energy shift due to its coupling with the collective mode. The only close-to-resonance process that a quasiparticle can undergo at zero temperature is the emission of a bosonic excitation from the phononic branch. Close to the minimum of the branch we find that the quasiparticles remain undamped, allowing us to find the energy correction in a self-consistent way, that we express in experimentally relevant quantities such as the energy gap, location of the minimum, effective mass, and Landau critical velocity.

Introduction— The notion of quasiparticles is an essential tool for the study of interacting many-body systems. Within this formalism, the many-body problem can be considerably simplified by introducing weakly interacting elementary excitations above a known ground state. It is, however, generally known that fermionic quasiparticles in an interacting Fermi system are only well-defined in a small region around the Fermi level, and that elsewhere they acquire a finite lifetime, even at zero temperature [1]. In a superfluid, the fermionic branch, which consists of pair-breaking excitations, obtains a finite energy gap described by BCS theory [2], and the system exhibits a gapless bosonic mode (unlike in superconductors [3]) representing the collective motion of fermion pairs [4]. In this Letter, we identify the coupling of the fermionic quasiparticle to the bosonic collective branch as the mechanism responsible for its finite lifetime away from the Fermi level, and compute the corresponding damping rate and energy shift from first principles. This problem is quite similar to the Bose polaron (an impurity in a sea of bosons) [5–7], but a crucial difference is that the fermionic quasiparticle has a roton-like dispersion $\epsilon_k = \Delta + \hbar^2(k - k_0)^2/2m^*$, rather than an impurity-like dispersion $\epsilon_k = \hbar^2k^2/2m^*$.

The appearance of a finite quasiparticle lifetime away from the energy minimum is a many-body phenomenon that occurs in many quantum systems such as normal Fermi liquids [1, 8], superconductors [9], or rotonic systems (superfluid Helium [10] or dipolar gases [11]). Using ultracold fermionic atoms [12–18], it can be studied analytically from a first-principles microscopic approach, and compared to experimental observations. Measurements of the quasiparticle spectrum using rf-spectroscopy [19] and momentum-resolved rf-spectroscopy [20], in particular of the quasiparticle gap [21], are in fact already available. Moreover, using Feshbach resonances to tune the interaction strength [22], experiments can study this effect in the whole range between a weakly-interacting BCS-type superfluid and a Bose-Einstein Condensate (BEC) of tightly-bound dimers, including at unitarity where interactions are resonant.

Yet, this problem was somehow overlooked in recent theoretical studies, concentrating rather on computing

the equation-of-state [23–25], the order parameter [26–28], or the bosonic collective mode spectrum [4, 29–33]. Pioneering studies have looked at beyond mean-field corrections to the single-particle Green’s function [34–36], identifying the coupling to the collective mode as the most important effect [36], but could not extract analytically the corrected eigenenergy and damping rate; moreover, their numerical approaches suffered limitations, predicting in particular a finite lifetime of the quasiparticles at the dispersion minimum. Here, we aim to fill this gap by analytically studying the coupling of the fermionic branch with the bosonic collective mode, modifying its dispersion in the entire BCS-BEC crossover. As expected, we find the quasiparticles to be well-defined around the Fermi level, while their lifetime becomes finite away from the energy minimum. The correction is perturbative in both the BCS and the BEC limit, expressing a vanishing damping rate in these limits.

Quasiparticle Hamiltonian— We study a gas of neutral fermionic atoms of mass m in two different hyperfine states, interacting via an attractive short-range potential, fully characterized, at low energy, by its s -wave scattering length a . Instead of using the full microscopic Hamiltonian, we describe the weakly-excited state of the system in terms of its quasiparticles. Specifically, we use an effective Hamiltonian derived from first principles that describes the fermionic quasiparticles and their coupling to the Anderson-Bogoliubov bosonic collective modes [31]

$$\begin{aligned} \hat{H}_{\text{qp}} = & \sum_{\mathbf{k}, \sigma=\uparrow, \downarrow} \epsilon_{\mathbf{k}} \hat{\gamma}_{\mathbf{k}, \sigma}^{\dagger} \hat{\gamma}_{\mathbf{k}, \sigma} + \sum_{\mathbf{q}} \hbar \omega_{\mathbf{q}} \hat{b}_{\mathbf{q}}^{\dagger} \hat{b}_{\mathbf{q}} \\ & + \frac{1}{\sqrt{V}} \sum_{\mathbf{k}, \mathbf{q}, \sigma} \left(\mathcal{A}_{\mathbf{k}-\mathbf{q}, \mathbf{q}} \hat{b}_{\mathbf{q}}^{\dagger} + \mathcal{A}_{\mathbf{k}, -\mathbf{q}} \hat{b}_{-\mathbf{q}} \right) \hat{\gamma}_{\mathbf{k}-\mathbf{q}, \sigma}^{\dagger} \hat{\gamma}_{\mathbf{k}, \sigma}. \end{aligned} \quad (1)$$

Here, the first term describes the BCS quasiparticles with creation (annihilation) operators $\hat{\gamma}_{\mathbf{k}, \sigma}^{\dagger}$ ($\hat{\gamma}_{\mathbf{k}, \sigma}$) and energy $\epsilon_{\mathbf{k}} = \sqrt{\xi_{\mathbf{k}}^2 + \Delta^2}$, with $\xi_{\mathbf{k}} = \frac{\hbar^2 k^2}{2m} - \mu$ the free-fermion dispersion relation, Δ the mean-field gap and μ the chemical potential. The second term in Eq. (1) represents the free bosonic collective modes, calculated within the RPA or Gaussian pair fluctuations (GPF) approximation [3, 4, 24], with operators $\hat{b}_{\mathbf{q}}^{\dagger}$, $\hat{b}_{\mathbf{q}}$; their eigenenergy $\hbar \omega_{\mathbf{q}}$ is

the only positive real root of the gaussian fluctuation matrix $\det M(\omega_{\mathbf{q}}, \mathbf{q}) = 0$ [4], with

$$M_{\pm\pm}(\omega, \mathbf{q}) = \frac{1}{2V} \sum_{\mathbf{k}} \left[\frac{\epsilon_{\mathbf{k}_+} + \epsilon_{\mathbf{k}_-}}{\epsilon_{\mathbf{k}_+} \epsilon_{\mathbf{k}_-}} \frac{\epsilon_{\mathbf{k}_+} \epsilon_{\mathbf{k}_-} + \xi_{\mathbf{k}_+} \xi_{\mathbf{k}_-} \pm \Delta^2}{\hbar^2 \omega^2 - (\epsilon_{\mathbf{k}_+} + \epsilon_{\mathbf{k}_-})^2} + \frac{1}{\epsilon_{\mathbf{k}}} \right]$$

$$M_{+-}(\omega, \mathbf{q}) = \frac{\hbar \omega}{2V} \sum_{\mathbf{k}} \frac{1}{\epsilon_{\mathbf{k}_+} \epsilon_{\mathbf{k}_-}} \frac{\epsilon_{\mathbf{k}_+} \xi_{\mathbf{k}_-} + \xi_{\mathbf{k}_+} \epsilon_{\mathbf{k}_-}}{\hbar^2 \omega^2 - (\epsilon_{\mathbf{k}_+} + \epsilon_{\mathbf{k}_-})^2} \quad (2)$$

and $\mathbf{k}_{\pm} = \mathbf{k} \pm \mathbf{q}/2$. Finally, the second line of Eq. (1) describes the three-body coupling between the fermionic quasiparticles and the collective modes, for which the coupling amplitude is given by

$$\mathcal{A}_{\mathbf{k}, \mathbf{q}} = \frac{w_{\mathbf{k}\mathbf{q}}^+ \sqrt{M_{++}(\omega_{\mathbf{q}}, \mathbf{q})} + w_{\mathbf{k}\mathbf{q}}^- \sqrt{M_{--}(\omega_{\mathbf{q}}, \mathbf{q})}}{\sqrt{-\frac{2}{\hbar} \frac{\partial}{\partial \omega} \det M(\omega, \mathbf{q})|_{\omega=\omega_{\mathbf{q}}}}}, \quad (3)$$

with the weights $w_{\mathbf{k}\mathbf{q}}^{\pm} = U_{\mathbf{k}+\mathbf{q}} V_{\mathbf{k}} \pm U_{\mathbf{k}} V_{\mathbf{k}+\mathbf{q}}$, and $U_{\mathbf{k}} = \sqrt{(1 + \xi_{\mathbf{k}}/\epsilon_{\mathbf{k}})/2}$, $V_{\mathbf{k}} = \sqrt{(1 - \xi_{\mathbf{k}}/\epsilon_{\mathbf{k}})/2}$. In the effective Hamiltonian of Eq. (1), we only take into account the process of emission or absorption of Anderson-Bogoliubov bosons, and omit highly off-resonant processes such as four-fermion interactions (at $T = 0$: $\gamma^\dagger \gamma^\dagger \gamma^\dagger \gamma$ or $\gamma^\dagger \gamma^\dagger \gamma^\dagger \gamma^\dagger$) and simultaneous creation of two fermions and a boson ($\gamma^\dagger \gamma^\dagger b^\dagger$). The decay channels not taken into account in our description will lead to a nonzero contribution to the damping rate only when the quasiparticle energy is at least 3Δ . The process described by \hat{H}_{qp} is then the only relevant damping process at zero temperature and low energy.

Energy corrections— To study the fermionic branch we compute its Green's function, which contains all the information on the quasiparticle. At zero temperature, the collective modes are unoccupied, which forbids the absorption of a boson by the fermionic quasiparticle. In this way, the only second order diagram contributing to the Green's function is the self-energy where the fermion emits and reabsorbs a boson:

$$G_{\mathbf{k}}(z) = \left(z - \epsilon_{\mathbf{k}} - \frac{1}{V} \sum_{\mathbf{q}} \frac{|\mathcal{A}_{\mathbf{k}-\mathbf{q}, \mathbf{q}}|^2}{z - \epsilon_{\mathbf{k}-\mathbf{q}} - \hbar \omega_{\mathbf{q}}} \right)^{-1}. \quad (4)$$

As explained in the Appendix, one obtains the same quasiparticle Green's function within the formalism of Refs. [34, 36] provided one omits all far-off-shell processes. The poles of the Green's function $G_{\mathbf{k}}^{-1}(z_{\mathbf{k}}) = 0$ are the eigenenergies of fermionic quasiparticles dressed by their interactions with the boson bath. When the coupling amplitude \mathcal{A} is small, one can replace z by $\epsilon_{\mathbf{k}} + i0^+$ in the last term between brackets of Eq. (4), to obtain the energy correction $z_{\mathbf{k}}^{(2)} = E_{\mathbf{k}}^{(2)} - i\hbar\Gamma_{\mathbf{k}}/2$ to second order perturbation theory:

$$E_{\mathbf{k}}^{(2)} = \epsilon_{\mathbf{k}} + \frac{1}{V} \mathcal{P} \sum_{\mathbf{q}} \frac{|\mathcal{A}_{\mathbf{k}-\mathbf{q}, \mathbf{q}}|^2}{\epsilon_{\mathbf{k}} - \epsilon_{\mathbf{k}-\mathbf{q}} - \hbar \omega_{\mathbf{q}}}, \quad (5)$$

$$\hbar\Gamma_{\mathbf{k}} = \frac{2\pi}{V} \sum_{\mathbf{q}} |\mathcal{A}_{\mathbf{k}-\mathbf{q}, \mathbf{q}}|^2 \delta(\epsilon_{\mathbf{k}} - \epsilon_{\mathbf{k}-\mathbf{q}} - \hbar \omega_{\mathbf{q}}). \quad (6)$$




FIG. 1. (Color online) The imaginary part of the quasiparticle Green's function (4) is shown as a function of the wavenumber k and energy ϵ in units of Δ . It is nonzero only above the threshold energy ϵ_{th} (solid gray line). Black dotted line: mean field energy $\epsilon_{\mathbf{k}}$, blue dashed line: perturbative eigenfrequency $E_{\mathbf{k}}^{(2)}$, blue solid line: perturbative damping rate $\Gamma_{\mathbf{k}}$. The latter is nonzero only for $k > k_{\text{th}} \simeq 1.17\sqrt{2m\Delta}/\hbar$ (vertical dashed dotted line) with $|\partial\epsilon_{\mathbf{k}}/\partial k|_{k=k_{\text{th}}} = \hbar c$. Below the threshold energy, a self-consistent energy $z_{\mathbf{k}}$ can be found until $k \simeq 1.62\sqrt{2m\Delta}/\hbar > k_{\text{th}}$ (orange solid line). It remains relatively close to the perturbative energy.

The resonance condition $\epsilon_{\mathbf{k}} - \epsilon_{\mathbf{k}-\mathbf{q}} - \hbar \omega_{\mathbf{q}} = 0$ in (6) is satisfied provided $\epsilon_{\mathbf{k}}$ is inside the boson emission continuum $\{\epsilon_{\mathbf{k}-\mathbf{q}} + \hbar \omega_{\mathbf{q}}, \mathbf{q}\}$, that is strictly superior to the threshold energy $\epsilon_{\text{th}} = \min_{\mathbf{q}}[\epsilon_{\mathbf{k}-\mathbf{q}} + \hbar \omega_{\mathbf{q}}]$, which is also the lower edge of the branch cut of $G_{\mathbf{k}}(z)$ [37].

Close to the minimum k_0 of the unperturbed fermionic branch, the group velocity of the quasiparticle is smaller than the sound velocity c of the collective mode $|\partial\epsilon_{\mathbf{k}}/\partial k| < \hbar c$. For these values of k , the minimum $\min_u[\epsilon_{\mathbf{k}-\mathbf{q}} + \hbar \omega_{\mathbf{q}}]$ over the scattering angle $u = \mathbf{k} \cdot \mathbf{q}/kq$ is a strictly increasing function of q , starting from its lowest value $\epsilon_{\mathbf{k}}$ in $q = 0$, such that $\epsilon_{\text{th}} = \epsilon_{\mathbf{k}}$ and the decay by emission of collective modes is energetically forbidden [31, 36]. The perturbative damping rate is zero ($\Gamma_{\mathbf{k}} = 0$) and we find correspondingly a real pole of $G_{\mathbf{k}}$, indicating that the quasiparticles, despite their renormalisation by the bosonic bath, remain well-defined close to the minimum of their dispersion.

When the group velocity $|\partial\epsilon_{\mathbf{k}}/\partial k|$ becomes larger than $\hbar c$ (which can happen in both the increasing [$k > k_0$] and decreasing parts [$k < k_0$] of the BCS branch), the unperturbed energy $\epsilon_{\mathbf{k}}$ becomes greater than ϵ_{th} and the resonance condition of Eq. (6) can be satisfied. Although the perturbative damping rate $\Gamma_{\mathbf{k}}$ becomes nonzero, the self-consistent solution $z_{\mathbf{k}}$ below ϵ_{th} remains real for larger values of $|k - k_0|$. Eventually, $z_{\mathbf{k}}$ also enters the continuum and becomes imaginary, which results in a broadened peak in the spectral function $\epsilon \mapsto \text{Im}[G_{\mathbf{k}}(\epsilon + i0^+)]$.

In Fig. 1, we illustrate this by plotting $\epsilon \mapsto \text{Im}[G_{\mathbf{k}}(\epsilon + i0^+)]$ at unitarity ($1/k_{\text{F}}a = 0$, where k_{F} is the Fermi

wavenumber) as a function of ε and k . We superimpose the eigenenergy $z_{\mathbf{k}}$ obtained by self-consistently solving for the real pole of the Green's function below ε_{th} , and the perturbative result $E_{\mathbf{k}}^{(2)}$ of Eq. (5), which remains fairly close to $z_{\mathbf{k}}$ everywhere. Once the self-consistent solution hits the continuum, the exact resonance of $G_{\mathbf{k}}$ turns into a broadened peak at energies $\varepsilon > \varepsilon_{\text{th}}$. The perturbative damping $\Gamma_{\mathbf{k}}$ becomes nonzero at $k = k_{\text{th}}$ with $|\partial \varepsilon_{\mathbf{k}} / \partial k|_{k=k_{\text{th}}} = \hbar c$. It is then highly peaked when the energy $\varepsilon_{\mathbf{k}}$ is around 3Δ . This could suggest that the $1 \rightarrow 3$ fermionic processes that we excluded from our Hamiltonian \hat{H}_{qp} in Eq. (1) become important above 3Δ . At higher wavenumbers, $E_{\mathbf{k}}^{(2)}$ approaches the mean field result (as does the maximum of $\text{Im}[G_{\mathbf{k}}(\varepsilon + i0^+)]$). This is not a surprise since the coupling \mathcal{A} is comparatively small in the limit $k \rightarrow \infty$.

BCS and BEC limits— The perturbative treatment Eqs. (5)–(6) (already close to the self-consistent solution at unitarity) gives asymptotically the exact solution of $G_{\mathbf{k}}^{-1} = 0$ in the BCS and BEC limits ($\Delta/|\mu| \rightarrow 0$). In the BCS limit, the bosonic wavenumbers should be expressed in units of $\Delta/\hbar c$, such that the dispersion $\hbar\omega_{\mathbf{q}}/\Delta$ becomes a universal function of $\tilde{\mathbf{q}} = \hbar c \mathbf{q}/\Delta$ [38]. The energy correction $|z_{\mathbf{k}} - \varepsilon_{\mathbf{k}}|$ is of order Δ^2/μ^2 :

$$\frac{z_{\mathbf{k}}^{(2)} - \varepsilon_{\mathbf{k}}}{\Delta} = \frac{\Delta^2}{\mu^2} \frac{1}{V} \sum_{\tilde{\mathbf{q}}} \frac{(w_{\mathbf{k}\mathbf{q}}^-)^2}{K(\tilde{\mathbf{q}})} \frac{\Delta}{\varepsilon_{\mathbf{k}} - \varepsilon_{\mathbf{k}-\mathbf{q}} - \hbar\omega_{\mathbf{q}} + i0^+}, \quad (7)$$

with $K(\tilde{\mathbf{q}}) = -32\hbar\Delta/(3\sqrt{3}mk_{\text{F}})\partial M_{++}/\partial\omega|_{\omega=\omega_{\mathbf{q}}}$ a universal function of $\tilde{\mathbf{q}}$. In the BEC limit, the energy should be scaled to the chemical potential $|\mu|$ and analytic results are available for the collective mode dispersion [4] at arbitrary momentum q . Consequently, the energy $E_{\mathbf{k}}^{(2)}$ and damping rate $\Gamma_{\mathbf{k}}$ of Eqs. (5-6) can be computed analytically, resulting in

$$\frac{z_{\mathbf{k}}^{(2)} - \varepsilon_{\mathbf{k}}}{|\mu|} = \frac{\Delta^2}{|\mu|^2} \frac{2\tilde{k}^2 - 6 - 8i\tilde{k}}{\tilde{k}^4 + 10\tilde{k}^2 + 9}, \quad (8)$$

with $\tilde{k} = \hbar k/\sqrt{2m|\mu|}$. The quasiparticle lifetime, which is also the lifetime of the bosonic dimers of \uparrow/\downarrow fermions, thus diverges like μ^2/Δ^2 .

In Fig. 2 we examine the perturbative damping rate $\Gamma_{\mathbf{k}}$ of Eq. (6) in the BCS and BEC limits. In the BCS limit, the undamped region lies around $k_0 \approx k_{\text{F}}$ and its width in units of k_{F} tends to zero like $\Delta/\varepsilon_{\text{F}}$. Outside this region, a highly peaked behavior can be observed, which, as in the unitary case, occurs when the energy is close to 3Δ . In the BEC limit, the undamped region lies around $k_0 = 0$, while the threshold wavenumber vanishes as $\hbar k_{\text{th}}/\sqrt{2m|\mu|} = \Delta/4|\mu|$. As a function of k , the damping rate smoothens, and exhibits a $1/k^3$ tail at high k .

Quadratic dispersion near the minimum— To analyze the characteristics of the corrected fermionic energy




FIG. 2. (Color online) Perturbative damping rate $\Gamma_{\mathbf{k}}$ (6) for different interaction regimes. (a) The damping rate as a function of $\xi_k = \hbar^2 k^2/2m - \mu$ at $\mu/\Delta = 10$ (full red line) is rescaled to compare with the universal behavior in the BCS limit (dotted blue line), where the damping rate vanishes as Δ^2/μ^2 , according to Eq. (7). (b) Damping rate in the BEC regime as a function of $\hbar k/\sqrt{2m|\mu|}$ for $\mu/\Delta = -5$ (full red line), and $\mu/\Delta = -1$ (dashed orange line) and $\mu/\Delta = -\infty$ (blue line, see Eq. (8)).

branch, we fit a quadratic dispersion

$$\varepsilon_k^{\text{fit}} = \varepsilon^* + \frac{\hbar^2(k - k_m^*)^2}{2m^*} \quad (9)$$

to the minimum of the energy. This allows us to extract the most interesting features of the energy correction, in the region where the description in terms of quasiparticles is certainly accurate. Concretely, these parameters represent the effective energy gap ε^* , the location of the minimum of the branch k_m^* and the effective mass m^* . In Fig. 3 we present these fitting parameters in the BCS-BEC crossover for both the self-consistent energy solution and the perturbative energy correction, comparing with the mean-field version of Eq. (9) obtained by expanding the BCS energy $\varepsilon_{\mathbf{k}}$ around its minimum. All parameters tend to the mean-field result in the BCS and BEC limits, confirming that the energy correction is perturbative for $\Delta/|\mu| \rightarrow 0$. Additionally, the differences between the self-consistent and perturbative results are never substantial, although perturbation theory somewhat overestimates the correction. The interaction with the bosonic collective mode lowers the fermionic energy gap ε^* , which is to be expected as it is generally known that the mean-field theory exaggerates the gap. At unitarity, we find $\varepsilon^* \simeq 0.88\Delta \simeq 0.41\varepsilon_{\text{F}}$ (using the GPF equation of state [24]), close to the experimental result $\Delta = 0.44\varepsilon_{\text{F}}$ [21], and $k_m^* \simeq 1.01\sqrt{2m\Delta}/\hbar \simeq 0.69k_{\text{F}}$. We furthermore see that the location k_m^* of the energy minimum reaches 0 at some critical value $\mu/\Delta \simeq -0.26$, depicted by a vertical line on Fig. 3, corresponding to $1/k_{\text{F}}a \simeq 0.56$. The fact that this happens while the chemical potential is already negative has been theoretically predicted before [36].




FIG. 3. (Color online) Minimum of the corrected fermionic energy branch fitted to a quadratic dispersion. (a) Dimensionless inverse effective mass m/m^* , (b) location of the minimum $\hbar k_m^*/\sqrt{2m\Delta}$, (c) energy gap ϵ^*/Δ , shown in function of the inverse scattering length $1/k_F a$ (bottom x -axis) and μ/Δ (top x -axis). Full curves: fitting parameters to the self-consistent (SC) energy solution z_k . Dashed lines: perturbative (Pert) results. Dotted lines: mean-field (MF) results. The vertical black lines specify the critical values of the interaction after which the minimum of the energy is located at $k = 0$, in their respective line styles. The BCS and BEC asymptotes are drawn for the energy gap in inset.

Critical velocity— Another interesting characterization of the fermionic branch is its Landau critical ve-

locity, which determines the maximal velocity for frictionless flow at $T = 0$ in a superfluid. As there are two branches of elementary excitations in a superfluid Fermi gas, the critical velocity will be given by the smallest of the two velocities $v_c = \min[v_f, c]$, with $v_f = \min_{\mathbf{k}_1, \mathbf{k}_2} (\epsilon_{\mathbf{k}_1} + \epsilon_{\mathbf{k}_2})/\hbar|\mathbf{k}_1 + \mathbf{k}_2|$ the fermionic critical velocity [39]. In the BCS limit the critical velocity is reached for k close to k_F and we can use the effective quadratic dispersion (9) near the minimum to compute it. This yields $m^* v_f^{\text{quad}} = \sqrt{2m^* \epsilon^* + (\hbar k_m^*)^2} - \hbar k_m^*$, from which we can extract the first deviation to the mean-field velocity $\frac{v_f}{v_f^{\text{MF}}} = \frac{\epsilon^*}{\Delta} + O\left(\frac{\Delta}{\mu}\right)^3 \simeq 1 - 0.5\left(\frac{\Delta}{\mu}\right)^2 + O\left(\frac{\Delta}{\mu}\right)^3$.

Conclusion— We have corrected the fermionic quasiparticle branch by including its interaction with the bosonic collective mode. At low energy, this is the only relevant decay channel, thus giving the sole contribution to the damping of the single-particle excitations. We have computed this damping rate in the entire BCS-BEC crossover, and found real poles of the corrected Green's function close to the minimum of the branch, indicating well-defined quasiparticles.

The boson-emission process we have studied occurs in systems where rotons are present, such as superfluid Helium [10] and dipolar Bose gases [11], and should be responsible at $T = 0$ for both a shift of the roton gap and a finite lifetime away from the roton minimum. It occurs also in normal Fermi liquids [1, 8] when the quasiparticles (in this case gapless) are coupled to the zero sound branch. In superconductors, the Bogoliubov-Anderson branch is gapped due to Coulomb interactions, but the quasiparticle can emit crystal phonons [9], a process quite similar to the one we have studied.

This research was supported by the Bijzonder Onderzoeksfonds (BOF) of the University of Antwerp, the Fonds Wetenschappelijk Onderzoek Vlaanderen, project G.0429.15.N, and the European Union's Horizon 2020 research and innovation program under the Marie Skłodowska-Curie grant agreement number 665501.

Appendix: Quasiparticle Green's function

In this Appendix, we show that the quasiparticle Green's function introduced in Eq. (4) (which follows immediately [40] from the low-energy Hamiltonian (1)) can be rederived in a more general microscopic formalism [34, 36] whose starting point are the particle, hole and anomalous Green's functions. The lowest order self-energy correcting the mean-field particle-hole Green's function

$$\mathcal{G}_0(z, \mathbf{k}) = -\frac{1}{z + \epsilon_{\mathbf{k}}} \begin{pmatrix} V_{\mathbf{k}}^2 & U_{\mathbf{k}} V_{\mathbf{k}} \\ U_{\mathbf{k}} V_{\mathbf{k}} & U_{\mathbf{k}}^2 \end{pmatrix} + \frac{1}{z - \epsilon_{\mathbf{k}}} \begin{pmatrix} -U_{\mathbf{k}}^2 & U_{\mathbf{k}} V_{\mathbf{k}} \\ U_{\mathbf{k}} V_{\mathbf{k}} & -V_{\mathbf{k}}^2 \end{pmatrix} \quad (\text{A.10})$$

was derived in Ref. [34]: $\mathcal{G}^{-1} = \mathcal{G}_0^{-1} - \Sigma$, with

$$\Sigma_{\alpha\alpha'}(z, \mathbf{k}) = \frac{1}{V} \sum_{\mathbf{q}} \int_{-\infty}^{+\infty} \frac{dz_q}{2\pi i} \mathcal{G}_{0,\alpha'\alpha}(z - z_q, \mathbf{q} - \mathbf{k}) \tilde{M}_{\alpha\alpha'}^{-1}(z_q, \mathbf{q}). \quad (\text{A.11})$$

Here, $\alpha, \alpha' = 1$ or 2 and \tilde{M} is a unitary transform of M defined in Eq. (2)

$$\tilde{M} = -\frac{1}{2} \begin{pmatrix} M_{++} + M_{--} + 2M_{+-} & M_{++} - M_{--} \\ M_{++} - M_{--} & M_{++} + M_{--} - 2M_{+-} \end{pmatrix}. \quad (\text{A.12})$$

This self-energy contains corrections from four-fermion processes (the lowest order processes allowed for fermions), as can be seen by expanding \tilde{M}^{-1} in powers of the coupling constant $g_0 = -(\sum_{\mathbf{k}} 1/2\epsilon_{\mathbf{k}})^{-1}$. To correctly describe the bosonic spectrum, two of the intermediate fermions are propagated using the ladder-resummed pair propagator \tilde{M}^{-1} , see Fig. 7 in Ref. [36]. To obtain analytic results, and avoid spurious effects (such as the appearance of a nonzero imaginary part of Δ at $T = 0$), we have dropped the self-consistent treatment of Ref. [36] and used in Eq. (A.11) the BCS/RPA expressions $\mathcal{G}_0, \tilde{M}^{-1}$. At $T = 0$, \tilde{M}^{-1} has two real poles $\pm\hbar\omega_{\mathbf{q}}$, and two gapped branch cuts. The Matsubara frequency integral in Eq. (A.11) can then be performed by closing the contour on the real axis, (the most straightforward way is to avoid the poles of \mathcal{G}_0). Neglecting the far-off-shell contributions of the branch cuts of \tilde{M}^{-1} , and performing the orthogonal transform to the quasiparticle basis $\tilde{\mathcal{G}}^{-1} = P\mathcal{G}^{-1}P^\dagger$ with

$$P = \begin{pmatrix} U_{\mathbf{k}} & -V_{\mathbf{k}} \\ V_{\mathbf{k}} & U_{\mathbf{k}} \end{pmatrix} \quad (\text{A.13})$$

we obtain the corrected Green's function

$$\tilde{\mathcal{G}}^{-1}(z, \mathbf{k}) = \begin{pmatrix} -z + \epsilon_{\mathbf{k}} + \frac{1}{V} \sum_{\mathbf{q}} \left[\frac{\mathcal{A}_{\mathbf{k}-\mathbf{q}, \mathbf{q}}^2}{z - \epsilon_{\mathbf{k}-\mathbf{q}} - \hbar\omega_{\mathbf{q}}} - \frac{\mathcal{B}_{\mathbf{k}, \mathbf{q}}^2}{-z - \epsilon_{\mathbf{k}+\mathbf{q}} - \hbar\omega_{\mathbf{q}}} \right] & \frac{1}{V} \sum_{\mathbf{q}} \left[\frac{\mathcal{A}_{\mathbf{k}-\mathbf{q}, \mathbf{q}} \mathcal{B}_{\mathbf{k}-\mathbf{q}, \mathbf{q}}}{z - \epsilon_{\mathbf{k}-\mathbf{q}} - \hbar\omega_{\mathbf{q}}} + \frac{\mathcal{A}_{\mathbf{k}, \mathbf{q}} \mathcal{B}_{\mathbf{k}, \mathbf{q}}}{-z - \epsilon_{\mathbf{k}+\mathbf{q}} - \hbar\omega_{\mathbf{q}}} \right] \\ \frac{1}{V} \sum_{\mathbf{q}} \left[\frac{\mathcal{A}_{\mathbf{k}-\mathbf{q}, \mathbf{q}} \mathcal{B}_{\mathbf{k}-\mathbf{q}, \mathbf{q}}}{z - \epsilon_{\mathbf{k}-\mathbf{q}} - \hbar\omega_{\mathbf{q}}} + \frac{\mathcal{A}_{\mathbf{k}, \mathbf{q}} \mathcal{B}_{\mathbf{k}, \mathbf{q}}}{-z - \epsilon_{\mathbf{k}+\mathbf{q}} - \hbar\omega_{\mathbf{q}}} \right] & -z - \epsilon_{\mathbf{k}} + \frac{1}{V} \sum_{\mathbf{q}} \left[\frac{\mathcal{B}_{\mathbf{k}-\mathbf{q}, \mathbf{q}}^2}{z - \epsilon_{\mathbf{k}-\mathbf{q}} - \hbar\omega_{\mathbf{q}}} - \frac{\mathcal{A}_{\mathbf{k}, \mathbf{q}}^2}{-z - \epsilon_{\mathbf{k}+\mathbf{q}} - \hbar\omega_{\mathbf{q}}} \right] \end{pmatrix}. \quad (\text{A.14})$$

Introducing the weights $W_{\mathbf{k}\mathbf{q}}^\pm = U_{\mathbf{k}+\mathbf{q}}U_{\mathbf{k}} \pm V_{\mathbf{k}}V_{\mathbf{k}+\mathbf{q}}$, the amplitude \mathcal{B} is given by

$$\mathcal{B}_{\mathbf{k}, \mathbf{q}} = \frac{W_{\mathbf{k}\mathbf{q}}^+ \sqrt{M_{--}(\omega_{\mathbf{q}}, \mathbf{q})} - W_{\mathbf{k}\mathbf{q}}^- \sqrt{M_{++}(\omega_{\mathbf{q}}, \mathbf{q})}}{\sqrt{-\frac{2}{\hbar} \frac{\partial}{\partial \omega} \det M(\omega, \mathbf{q})|_{\omega=\omega_{\mathbf{q}}}}}, \quad (\text{A.15})$$

while $\mathcal{A}_{\mathbf{k}, \mathbf{q}}$ coincides with Eq. (3). The quasiparticle energy is given by the poles of $\tilde{\mathcal{G}}(z)$, hence the roots of $\det \tilde{\mathcal{G}}^{-1}(z)$. Note the property $\det \tilde{\mathcal{G}}^{-1}(-z) = \det \tilde{\mathcal{G}}^{-1}(z)$, which ensures that the quasiparticle and quasihole energies are simply opposite. To be consistent with the omission of the branch cut contributions, we drop in Eq. (A.14) the terms corresponding to the far-off-shell processes $\hat{\gamma}_{-\mathbf{k}-\mathbf{q}, \sigma}^\dagger \hat{\gamma}_{\mathbf{k}, \sigma}^\dagger \hat{b}_{\mathbf{q}}^\dagger$, and we restrict to first order in Σ (which amounts to setting $\mathcal{B} = 0$). Then $\det \tilde{\mathcal{G}}^{-1}(z)$ is simply the product of the inverse quasiparticle $\tilde{\mathcal{G}}_{11}(z) = -G_{\mathbf{k}}(z) + \mathcal{O}(\|\Sigma\|^2)$ and quasihole $\tilde{\mathcal{G}}_{22}(z) = G_{\mathbf{k}}(-z) + \mathcal{O}(\|\Sigma\|^2)$ Green's functions

$$\det \mathcal{G}^{-1} = -G_{\mathbf{k}}^{-1}(z)G_{\mathbf{k}}^{-1}(-z) + \mathcal{O}(\|\Sigma\|^2). \quad (\text{A.16})$$

This finally explains the use of Eq. (4) to study the perturbed quasiparticle spectrum.

* Senne.VanLoon@UAntwerpen.be

- [1] Philippe Nozières, *Theory of interacting Fermi systems* (W.A. Benjamin, New York, 1964).
- [2] J. Bardeen, L. N. Cooper, and J. R. Schrieffer, "Theory of Superconductivity," *Phys. Rev.* **108**, 1175–1204 (1957).
- [3] P.W. Anderson, "Random-Phase Approximation in the Theory of Superconductivity," *Phys. Rev.* **112**, 1900–1916 (1958).
- [4] R. Combescot, M. Yu. Kagan, and S. Stringari, "Collective mode of homogeneous superfluid Fermi gases in the BEC-BCS crossover," *Phys. Rev. A* **74**, 042717 (2006).
- [5] L.D. Landau, *Phys. Z. Sowjetunion* **3**, 664 (1933).
- [6] O. I. Utesov, M. I. Baglay, and S. V. Andreev, "Effective interactions in a quantum Bose-Bose mixture," *Phys.*

Rev. A **97**, 053617 (2018).

- [7] Senne Van Loon, Wim Casteels, and Jacques Tempere, "Ground-state properties of interacting Bose polarons," *Phys. Rev. A* **98**, 063631 (2018).
- [8] Henri Godfrin, Matthias Meschke, Hans-Jochen Lauter, Ahmad Sultan, Helga M. Böhm, Eckhard Krotscheck, and Martin Panholzer, "Observation of a roton collective mode in a two-dimensional Fermi liquid," *Nature* **483**, 576 EP – (2012).
- [9] S. B. Kaplan, C. C. Chi, D. N. Langenberg, J. J. Chang, S. Jafarey, and D. J. Scalapino, "Quasiparticle and phonon lifetimes in superconductors," *Phys. Rev. B* **14**, 4854–4873 (1976).
- [10] B. Fåk, T. Keller, M. E. Zhitomirsky, and A. L. Chernyshev, "Roton-Phonon Interactions in Superfluid ^4He ," *Phys. Rev. Lett.* **109**, 155305 (2012).
- [11] L. Chomaz, R. M. W. van Bijnen, D. Petter, G. Faraoni, S. Baier, J. H. Becher, M. J. Mark, F. Wächtler, L. San-

- tos, and F. Ferlaino, “Observation of roton mode population in a dipolar quantum gas,” *Nature Physics* **14**, 442–446 (2018).
- [12] Immanuel Bloch, Jean Dalibard, and Wilhelm Zwerger, “Many-body physics with ultracold gases,” *Rev. Mod. Phys.* **80**, 885–964 (2008).
- [13] Wilhelm Zwerger, *The BCS-BEC crossover and the unitary Fermi gas*, Vol. 836 (Springer Science & Business Media, 2011).
- [14] Markus Greiner, Cindy A. Regal, and Deborah S. Jin, “Emergence of a molecular Bose-Einstein condensate from a Fermi gas,” *Nature* **426**, 537–540 (2003).
- [15] J. Joseph, B. Clancy, L. Luo, J. Kinast, A. Turlapov, and J. E. Thomas, “Measurement of Sound Velocity in a Fermi Gas near a Feshbach Resonance,” *Phys. Rev. Lett.* **98**, 170401 (2007).
- [16] Wolfgang Ketterle and Martin W Zwiernlein, “Making, probing and understanding ultracold Fermi gases,” in *Proceedings of the International School of Physics Enrico Fermi, Course CLXIV*, edited by M Inguscio, W Ketterle, and C Salomon (IOS Press, Amsterdam, 2008).
- [17] Leonid A. Sidorenkov, Meng Khoon Tey, Rudolf Grimm, Yan-Hua Hou, Lev Pitaevskii, and Sandro Stringari, “Second sound and the superfluid fraction in a Fermi gas with resonant interactions,” *Nature* **498**, 78–81 (2013).
- [18] Sascha Hoinka, Paul Dyke, Marcus G. Lingham, Jami J. Kinnunen, Georg M. Bruun, and Chris J. Vale, “Goldstone mode and pair-breaking excitations in atomic Fermi superfluids,” *Nature Physics* **13**, 943–946 (2017).
- [19] S. Gupta, Z. Hadzibabic, M. W. Zwiernlein, C. A. Stan, K. Dieckmann, C. H. Schunck, E. G. M. van Kempen, B. J. Verhaar, and W. Ketterle, “Radio-frequency spectroscopy of ultracold fermions,” *Science* **300**, 1723–1726 (2003).
- [20] J. T. Stewart, J. P. Gaebler, and D. S. Jin, “Using photoemission spectroscopy to probe a strongly interacting Fermi gas,” *Nature* **454**, 744 (2008).
- [21] André Schirrotzek, Yong-il Shin, Christian H Schunck, and Wolfgang Ketterle, “Determination of the superfluid gap in atomic Fermi gases by quasiparticle spectroscopy,” *Physical Review Letters* **101**, 140403 (2008).
- [22] M. Bartenstein, A. Altmeyer, S. Riedl, S. Jochim, C. Chin, J. Hecker Denschlag, and R. Grimm, “Crossover from a Molecular Bose-Einstein Condensate to a Degenerate Fermi Gas,” *Phys. Rev. Lett.* **92**, 120401 (2004).
- [23] H. Hu, X.-J. Liu, and P. D. Drummond, “Equation of state of a superfluid Fermi gas in the BCS-BEC crossover,” *EPL* **74**, 574 (2006).
- [24] Roberto B. Diener, Rajdeep Sensarma, and Mohit Randeria, “Quantum fluctuations in the superfluid state of the BCS-BEC crossover,” *Phys. Rev. A* **77**, 023626 (2008).
- [25] G. E. Astrakharchik, J. Boronat, J. Casulleras, and S. Giorgini, “Equation of State of a Fermi Gas in the BEC-BCS Crossover: A Quantum Monte Carlo Study,” *Phys. Rev. Lett.* **93**, 200404 (2004).
- [26] L.P. Gor’kov and T.K. Melik-Barkhudarov, “Contribution to the theory of superfluidity in an imperfect Fermi gas,” *Sov. Phys. JETP* **13**, 1018 (1958 [*ZhETF*, **40**, 1452]).
- [27] P. Pieri, L. Pisani, and G. Calvanese Strinati, “BCS-BEC crossover at finite temperature in the broken-symmetry phase,” *Physical Review B* **70**, 094508 (2004).
- [28] L. Pisani, P. Pieri, and G. Calvanese Strinati, “Gap equation with pairing correlations beyond the mean-field approximation and its equivalence to a Hugenholtz-Pines condition for fermion pairs,” *Physical Review B* **98**, 104507 (2018).
- [29] Vladimir Andreevich Andrianov and Viktor Nikolaevich Popov, “Gidrodinamieskoe dejstvie i Boze-spektr sverhtekuih Fermi-sistem,” *Teoreticheskaya i Matematicheskaya Fizika* **28**, 341–352 (1976), [English translation: *Theoretical and Mathematical Physics*, 1976, 28:3, 829837].
- [30] H. Kurkjian, Y. Castin, and A. Sinatra, “Three-Phonon and Four-Phonon Interaction Processes in a Pair-Condensed Fermi Gas,” *Annalen der Physik* **529**, 1600352 (2017).
- [31] Hadrien Kurkjian and Jacques Tempere, “Absorption and emission of a collective excitation by a fermionic quasiparticle in a Fermi superfluid,” *New Journal of Physics* **19**, 113045 (2017).
- [32] H. Kurkjian, S. N. Klimin, J. Tempere, and Y. Castin, “Pair-Breaking Collective Branch in BCS Superconductors and Superfluid Fermi Gases,” *Phys. Rev. Lett.* **122**, 093403 (2019).
- [33] S. N. Klimin, H. Kurkjian, and J. Tempere, “Phononic collective excitations in superfluid Fermi gases at nonzero temperatures,” *arXiv:1811.07796* (2018).
- [34] R. Haussmann, “Crossover from BCS superconductivity to Bose-Einstein condensation: A self-consistent theory,” *Zeitschrift für Physik B Condensed Matter* **91**, 291–308 (1993).
- [35] Nils Lerch, Lorenz Bartosch, and Peter Kopietz, “Absence of Fermionic Quasiparticles in the Superfluid State of the Attractive Fermi Gas,” *Phys. Rev. Lett.* **100**, 050403 (2008).
- [36] Rudolf Haussmann, Matthias Punk, and Wilhelm Zwerger, “Spectral functions and rf response of ultracold fermionic atoms,” *Physical Review A* **80**, 063612 (2009).
- [37] The boson emission continuum may also have an upper bound in the regimes where the Anderson-Bogoliubov branch has a compact domain of existence [4].
- [38] H Kurkjian, Yvan Castin, and A Sinatra, “Concavity of the collective excitation branch of a Fermi gas in the BEC-BCS crossover,” *Physical Review A* **93**, 013623 (2016).
- [39] Yvan Castin, Igor Ferrier-Barbut, and Christophe Salomon, “La vitesse critique de Landau d’une particule dans un superfluide de fermions,” *Comptes Rendus Physique* **16**, 241–253 (2015).
- [40] C. Cohen-Tannoudji, J. Dupont-Roc, and G. Grynberg, “Processus d’interaction entre photons et atomes,” (InterEditions et Éditions du CNRS, Paris, 1988) Chap. III. Étude non perturbative des amplitudes de transition.