On the existence of t_r -norm and t_r -conorm not in convolution form^{*}

Xinxing Wu^{a,b,c,*}, Guanrong Chen^d, Lidong Wang^c

^aSchool of Mathematics and Statistics, Guizhou University of Finance and Economics, Guiyang 550025, China

^bInstitute for Artificial Intelligence, Southwest Petroleum University, Chengdu, Sichuan 610500, China ^cZhuhai College of Jilin University, Zhuhai, Guangdong 519041, China

^dDepartment of Electrical Engineering, City University of Hong Kong, Hong Kong SAR, China

Abstract

This paper constructs a t_r -norm and a t_r -conorm on the set of all normal and convex functions from [0, 1] to [0, 1], which are not obtained by using the following two formulas on binary operations λ and γ :

$$(f \land g)(x) = \sup \{f(y) \ast g(z) \mid y \land z = x\},$$

$$(f \land g)(x) = \sup \{f(y) \ast g(z) \mid y \lor z = x\},$$

where $f, g \in Map([0,1], [0,1])$, \triangle and \forall are respectively a *t*-norm and a *t*-conorm on [0,1], and * is a binary operation on [0,1]. This result answers affirmatively an open problem posed in [3]. Moreover, the duality between t_r -norms and t_r -conorms is obtained by the introduction of operations dual to binary operations on Map([0,1], [0,1]).

Keywords: Normal and convex function, t-norm, t-conorm, t_r -norm, t_r -conorm, type-2 fuzzy set.

1. Introduction

In 1975, Zadeh [15] introduced the notion of type-2 fuzzy sets (T2FSs), that is, a fuzzy set with fuzzy sets as truth values (simply, "fuzzy-fuzzy sets"), as an extension of type-1 fuzzy sets (FSs) and interval-valued fuzzy sets (IVFSs), which was then equivalently expressed in different forms by Mendel et al. [4, 5, 7, 8]. The definitions of triangular norms (briefly *t*-norms) and triangular conorms (briefly *t*-conorms) on the real unit interval were introduced by Schweizer and Sklar [10] in the framework of probabilistic metric spaces. These definitions exploits the main idea of Menger [9] that extends the classical triangle inequality in metric spaces to probabilistic metric spaces. In 2006, Walker and Walker [12] extended *t*-norm and *t*-conorm on *I* to the algebra of truth values on T2FSs and IVFSs. Then, Hernández et al. [3] modified Walker and Walker's definition and introduced the notions of a t_r -norm and a t_r -conorm by adding some "restrictive axioms" (see Definition 2.8 below). In particular, in [3] they proved that the binary operation λ (resp., γ) on the set **L** of all normal and convex functions is a t_r -norm (resp., a t_r -conorm). They also proposed the following two open problems on the binary operations λ and γ (see Definition 2.5 below).

Question 1. [3] Apart from the t-norm, does there exist other binary operation '*' on I such that ' λ ' and ' γ ' are, respectively, a t_r -norm and a t_r -conorm on **L**?

Question 2. [3] Determine other binary operations, which are not obtained using the formulas given for the operations ' λ ' and ' γ ', that are either a t_r -norm or a t_r -conorm on **L**.

Recently, in [13] we have answered negatively Question 1, proving that, if a binary operation * ensures \land be a t_r -norm on \mathbf{L} or Υ be a t_r -conorm on \mathbf{L} , then * is a t-norm. This paper is devoted to solving Question 2 by constructing a t_r -norm ' \bigstar ' (see Section 4) and a t_r -conorm ' \bigstar ' (see Section 6) on \mathbf{L} , which cannot be obtained by the formulas defining the operations ' \land ' and ' Υ '.

^{*}This work was supported by National Natural Science Foundation of China (No. 11601449), Yibin University Cultivating Foundation (No. 02180199), Science and Technology Innovation Team of Education Department of Sichuan for Dynamical System and its Applications (No. 18TD0013), Youth Science and Technology Innovation Team of Southwest Petroleum University for Nonlinear Systems (No. 2017CXTD02), and Key Natural Science Foundation of Universities in Guangdong Province (No. 2019KZDXM027).

^{*}Corresponding author

Email addresses: wuxinxing5201314@163.com (Xinxing Wu), gchen@ee.cityu.edu.hk (Guanrong Chen), wld0707@126.com (Lidong Wang)

2. Preliminaries and basic concepts

Throughout this paper, let I = [0, 1], Map(X, Y) be the set of all mappings from X to Y, and ' \leq ' denote the usual order relation in the lattice of real numbers, with $\mathbf{M} = Map(I, I)$. Let \vee and \wedge be the maximum and minimum operations, respectively, on lattice I.

Definition 2.1. [14] A type-1 fuzzy set A in space X is a mapping from X to I, i.e., $A \in Map(X, I)$.

Definition 2.2. [11] A type-2 fuzzy set A in space X is a mapping

$$A: X \to \mathbf{M},$$

i.e., $A \in Map(X, \mathbf{M})$.

Definition 2.3. [11] A function $f \in \mathbf{M}$ is

- (1) normal if $\sup\{f(x) \mid x \in I\} = 1;$
- (2) convex if, for any $0 \le x \le y \le z \le 1$, $f(y) \ge f(x) \land f(z)$.

Let N and L denote the set of all normal functions in M and the set of all normal and convex functions in M, respectively.

For any subset B of X, a special fuzzy set $\mathbf{1}_B$, called the *characteristic function* of B, is defined by

$$\mathbf{1}_B(x) = \begin{cases} 1, & x \in B, \\ 0, & x \in X \setminus B \end{cases}$$

Let $\mathbf{J} = {\mathbf{1}_{\{x\}} \mid x \in I}$ and $\mathbf{K} = {\mathbf{1}_{[a,b]} \mid 0 \le a \le b \le 1}.$

Definition 2.4. [6] A binary operation $*: I^2 \to I$ is a *t*-norm on I if it satisfies the following axioms:

- (T1) (commutativity) x * y = y * x for $x, y \in I$;
- (T2) (associativity) (x * y) * z = x * (y * z) for $x, y, z \in I$;
- (T3) (monotonicity) * is increasing in each argument;
- (T4) (neutral element) 1 * x = x * 1 = x for $x \in I$.

A binary operation $*: I^2 \to I$ is a *t*-conorm on I if it satisfies axioms (T1), (T2), and (T3) above; and axiom (T4'): 0 * x = x * 0 = x for $x \in I$.

Definition 2.5. [3] Let * be a binary operation on I, \triangle be a *t*-norm on I, and ∇ be a *t*-conorm on I. Define the binary operations \wedge and $\Upsilon : \mathbf{M}^2 \to \mathbf{M}$ as follows: for $f, g \in \mathbf{M}$,

$$(f \land g)(x) = \sup\left\{f(y) * g(z) \mid y \land z = x\right\},\tag{2.1}$$

and

$$(f \uparrow g)(x) = \sup \{ f(y) * g(z) \mid y \forall z = x \}.$$
(2.2)

Definition 2.6. [11] The operations of \sqcup (*union*), \sqcap (*intersection*), \neg (*complementation*) on **M** are defined as follows: for $f, g \in \mathbf{M}$,

$$(f \sqcup g)(x) = \sup\{f(y) \land g(z) \mid y \lor z = x\},\$$

$$(f \sqcap g)(x) = \sup\{f(y) \land g(z) \mid y \land z = x\},\$$

and

$$(\neg f)(x) = \sup\{f(y) \mid 1 - y = x\} = f(1 - x).$$

From [11], it follows that $\mathfrak{M} = (\mathbf{M}, \sqcup, \sqcap, \neg, \mathbf{1}_{\{0\}}, \mathbf{1}_{\{1\}})$ does not have a lattice structure, although \sqcup and \sqcap satisfy the De Morgan's laws with respect to the complementation \neg .

Walker and Walker [11] introduced the following partial orders \sqsubseteq and \preccurlyeq on M.

Definition 2.7. [11] $f \sqsubseteq g$ if $f \sqcap g = f$; $f \preccurlyeq g$ if $f \sqcup g = g$.

It follows from [11, Proposition 14] that both \sqsubseteq and \preccurlyeq are partial orders on **M**. Generally, the partial orders \sqsubseteq and \preccurlyeq do not coincide. In [1, 2, 11], it was proved that \sqsubseteq and \preccurlyeq coincide on **L**, and the subalgebra $\mathfrak{L} = (\mathbf{L}, \sqcup, \sqcap, \neg, \mathbf{1}_{\{0\}}, \mathbf{1}_{\{1\}})$ is a bounded complete lattice. In particular, $\mathbf{1}_{\{0\}}$ and $\mathbf{1}_{\{1\}}$ are the minimum and the maximum of \mathfrak{L} , respectively.

Definition 2.8. [3] A binary operation $\widetilde{T} : \mathbf{L}^2 \to \mathbf{L}$ is a t_r -norm (t-norm according to the restrictive axioms), if

- (O1) \widetilde{T} is commutative, i.e., $\widetilde{T}(f,g) = \widetilde{T}(g,f)$ for $f,g \in \mathbf{L}$;
- (O2) \widetilde{T} is associative, i.e., $\widetilde{T}(\widetilde{T}(f,g),h) = \widetilde{T}(f,\widetilde{T}(g,h))$ for $f,g,h \in \mathbf{L}$;
- (O3) $\widetilde{T}(f, \mathbf{1}_{\{1\}}) = f$ for $f \in \mathbf{L}$ (neutral element);
- (O4) \widetilde{T} is increasing, i.e., for $f, g, h \in \mathbf{L}$ such that $f \sqsubseteq g, \widetilde{T}(f, h) \sqsubseteq \widetilde{T}(g, h)$;
- (O5) $\widetilde{T}(\mathbf{1}_{[0,1]},\mathbf{1}_{[a,b]}) = \mathbf{1}_{[0,b]};$
- (O6) \widetilde{T} is closed on **J**;
- (O7) \widetilde{T} is closed on **K**.

A binary operation $\widetilde{S} : \mathbf{L}^2 \to \mathbf{L}$ is a t_r -conorm if it satisfies axioms (O1), (O2), (O4), (O6), and (O7) above; axiom (O3'): $\widetilde{S}(f, \mathbf{1}_{\{0\}}) = f$; and axiom (O5'): $\widetilde{S}(\mathbf{1}_{[0,1]}, \mathbf{1}_{[a,b]}) = \mathbf{1}_{[a,1]}$. Axioms (O1), (O2), (O3), (O3'), and (O4) are called "basic axioms", and an operation that complies with these axioms will be referred to as *t*-norm and *t*-conorm, respectively.

Definition 2.9. For $f \in \mathbf{M}$, define

$$f^{L}(x) = \sup\{f(y) \mid y \le x\},\$$

$$f^{L_{w}}(x) = \begin{cases} \sup\{f(y) \mid y < x\}, & x \in (0, 1], \\ f(0), & x = 0, \end{cases}$$

and

$$f^{R}(x) = \sup\{f(y) \mid y \ge x\},\$$

$$f^{R_{w}}(x) = \begin{cases} \sup\{f(y) \mid y > x\}, & x \in [0, 1), \\ f(1), & x = 1. \end{cases}$$

Clearly, (1) f^L , f^{L_w} and f^R , f^{R_w} are increasing and decreasing, respectively; (2) $f^L(x) \vee f^R(x) = f^L(x) \vee f^{R_w}(x) = \sup_{z \in I} \{f(z)\}$ and $f^R(x) \vee f^{L_w}(x) = \sup_{z \in I} \{f(z)\}$ for all $x \in I$. The following properties of f^L and f^R were obtained by Walker et al. [1, 2, 11].

Proposition 2.1. [11] For $f, g \in \mathbf{M}$,

- (1) $f \leq f^{L} \wedge f^{R};$ (2) $(f^{L})^{L} = f^{L}, (f^{R})^{R} = f^{R};$ (3) $(\neg f)^{L} = \neg (f^{R}), (\neg f)^{R} = \neg (f^{L});$
- (4) $(f^L)^R = (f^R)^L = \sup_{x \in I} \{f(x)\};$
- (5) $f \sqsubseteq g$ if and only if $f^R \land g \le f \le g^R$;
- (6) $f \preccurlyeq g$ if and only if $f \land g^L \leq g \leq f^L$;
- (7) f is convex if and only if $f = f^L \wedge f^R$.

Theorem 2.1. [1, 2] Let $f, g \in \mathbf{L}$. Then, $f \sqsubseteq g$ if and only if $f^L \ge g^L$ and $f^R \le g^R$.

The following result follows from the definitions of f^L and f^R .

Lemma 2.1. For $f \in \mathbf{M}$, $f^{L}(1) = f^{R}(0) = \sup_{x \in I} \{f(x)\}.$

Proposition 2.2. For $f \in \mathbf{M}$, $f^{L_{w}}(x) = \sup_{t \in [0,x)} \{f^{L}(t)\}$ for all $x \in (0,1]$.

Proof. Fix any $x \in (0,1]$, noting that $f(t) \leq f^{L}(t)$ for all $t \in [0,x)$, we have

$$f^{L_{\mathbf{w}}}(x) = \sup_{t \in [0,x)} \{f(t)\} \le \sup_{t \in [0,x)} \{f^{L}(t)\}.$$

Moreover, for any $t \in [0, x)$, it follows from $t < \frac{t+x}{2} < x$ that $f^L(t) \le f^{L_w}(\frac{t+x}{2}) \le f^{L_w}(x)$, implying that

$$\sup_{t \in [0,x)} \{ f^L(t) \} \le f^{L_{\mathbf{w}}}(x).$$

Thus,

$$f^{L_{\mathbf{w}}}(x) = \sup_{t \in [0,x)} \{f^{L}(t)\}.$$

Lemma 2.2. For $f \in \mathbf{N}$, $\inf\{x \in I \mid f^L(x) = 1\} \le \sup\{x \in I \mid f^R(x) = 1\}$.

Proof. From $f \in \mathbf{N}$ and Lemma 2.1, it follows that $f^L(1) = f^R(0) = \sup\{f(x) \mid x \in I\} = 1$, which means that both $\{x \in I \mid f^L(x) = 1\}$ and $\{x \in I \mid f^R(x) = 1\}$ are nonempty sets. Denote $\eta = \inf\{x \in I \mid f^L(x) = 1\}$ and $\xi = \sup\{x \in I \mid f^R(x) = 1\}$. If $\eta = 0$, this holds trivially. If $\eta > 0$, then for any $0 \le \alpha < \eta$, $f^L(\alpha) < 1$. This, together with $f^L(\alpha) \lor f^R(\alpha) = \sup_{x \in I}\{f(x)\} = 1$, implies that $f^R(\alpha) = 1$. Thus, $\alpha \le \xi$. Therefore,

$$\xi \ge \sup\{\alpha \mid 0 \le \alpha < \eta\} = \eta.$$

3. Basic properties of *

In this section, basic properties of * determined by the binary operations Δ , ∇ , λ , and Υ are obtained.

Proposition 3.1. (1) Let * be a t-norm on I. Then, x * y = 1 if and only if x = y = 1.

(2) Let * be a t-conorm on I. Then, x * y = 0 if and only if x = y = 0.

Lemma 3.1. (1) Let \triangle be a t-norm on I and \ast be a binary operation on I. Then,

$$(f \land g)(1) = f(1) * g(1)$$

(2) Let \forall be a t-conorm on I and * be a binary operation on I. Then,

$$(f \uparrow g)(0) = f(0) * g(0).$$

Proof. Since \triangle is a *t*-norm, from Proposition 3.1, we have

$$(f \land g)(1) = \sup\{f(y) * g(z) \mid y \land z = 1\} = f(1) * g(1).$$

Similarly, we have

$$(f \uparrow g)(0) = \sup\{f(y) * g(z) \mid y \forall z = 1\} = f(0) * g(0).$$

Proposition 3.2. Let \triangle be a t-norm on I and \ast be a binary operation on I. If λ is commutative on L, then \ast is commutative.

Proof. Suppose, on the contrary, that * is not commutative. Then, there exist $u, v \in I$ such that $u * v \neq v * u$. Choose two functions $f, g \in \mathbf{M}$, as follows

$$f(x) = (u-1)x + 1$$

and

$$g(x) = (v-1)x + 1,$$

for $x \in I$. It can be verified that $f, g \in \mathbf{L}$, as both f and g are decreasing. Since \wedge is commutative, Lemma 3.1 yields

$$u * v = f(1) * g(1) = (f \land g)(1)$$

= $(g \land f)(1) = g(1) * f(1) = v * u$

which is a contradiction. Therefore, * is commutative.

Proposition 3.3. Let \triangle be a t-norm on I and \ast be a binary operation on I. If \land is a t-norm on L, then $0 \ast 0 = 0 \ast 1 = 1 \ast 0 = 0$ and $1 \ast 1 = 1$.

Proof. Since $\mathbf{1}_{\{1\}}$ is the neutral element of λ , from Lemma 3.1 and Proposition 3.2, it follows that

$$0 = \mathbf{1}_{\{0\}}(1) = (\mathbf{1}_{\{1\}} \land \mathbf{1}_{\{0\}})(1)$$

= $\mathbf{1}_{\{1\}}(1) * \mathbf{1}_{\{0\}}(1)$
= $1 * 0 = 0 * 1;$
$$0 = \mathbf{1}_{\{0.5\}}(0) = (\mathbf{1}_{\{0.5\}} \land \mathbf{1}_{\{1\}})(0)$$

\geq $\mathbf{1}_{\{0.5\}}(1) * \mathbf{1}_{\{1\}}(0) \text{ (as } 1 \bigtriangleup 0 = 0)$
= $0 * 0;$

and

$$1 = \mathbf{1}_{\{1\}}(1) = (\mathbf{1}_{\{1\}} \land \mathbf{1}_{\{1\}})(1)$$

= $\mathbf{1}_{\{1\}}(1) * \mathbf{1}_{\{1\}}(1) = 1 * 1.$

Proposition 3.4. Let \forall be a t-conorm on I and * be a binary operation on I. If Υ is a t-conorm on L, then 0 * 0 = 0 * 1 = 1 * 0 = 0 and 1 * 1 = 1.

Proof. Since $\mathbf{1}_{\{0\}}$ is the neutral element of Υ , from Lemma 3.1, it follows that

$$0 = \mathbf{1}_{\{1\}}(0) = (\mathbf{1}_{\{1\}} \uparrow \mathbf{1}_{\{0\}})(0)$$

= $\mathbf{1}_{\{1\}}(0) * \mathbf{1}_{\{0\}}(0) = 0 * 1;$ (3.1)

$$0 = \mathbf{1}_{\{1\}}(0) = (\mathbf{1}_{\{0\}} \land \mathbf{1}_{\{1\}})(0)$$

= $\mathbf{1}_{\{0\}}(0) \ast \mathbf{1}_{\{1\}}(0) = 1 \ast 0;$ (3.2)

and

$$0 = \mathbf{1}_{\{0.5\}}(1) = (\mathbf{1}_{\{0.5\}} \lor \mathbf{1}_{\{0\}})(1)$$

$$\geq \mathbf{1}_{\{0.5\}}(0) * \mathbf{1}_{\{0\}}(1) \text{ (as } 0 \bigtriangledown 1 = 1)$$

$$= 0 * 0.$$
(3.3)

It follows from (3.1)–(3.3) that, for $y, z \in I$, one has $\mathbf{1}_{\{1\}}(y) * \mathbf{1}_{\{0\}}(z) \in \{0, 1 * 1\}$. This implies that

$$1 = \mathbf{1}_{\{1\}}(1) = (\mathbf{1}_{\{1\}} \lor \mathbf{1}_{\{0\}})(1)$$

= sup{0, $\mathbf{1}_{\{1\}}(1) * \mathbf{1}_{\{0\}}(0)$ } (as $1 \lor 0 = 1$)
= 1 * 1.

Proposition 3.5. Let \triangle be a t-norm on I and * be a binary operation on I. If \wedge is a t-norm on \mathbf{L} , then, for $x_1, x_2 \in I$, one has $\mathbf{1}_{\{x_1\}} \wedge \mathbf{1}_{\{x_2\}} = \mathbf{1}_{\{x_1 \triangle x_2\}}$.

Proof. Proposition 3.3 yields

- (a) for $y, z \in I$, $\mathbf{1}_{\{x_1\}}(y) * \mathbf{1}_{\{x_2\}}(z) \in \{0, 1\};$
- (b) $\mathbf{1}_{\{x_1\}}(y) * \mathbf{1}_{\{x_2\}}(z) = 1$ if and only if $y = x_1$ and $z = x_2$.

This, together with

$$(\mathbf{1}_{\{x_1\}} \land \mathbf{1}_{\{x_2\}})(x) = \sup\{\mathbf{1}_{\{x_1\}}(y) * \mathbf{1}_{\{x_2\}}(z) \mid y \vartriangle z = x\},\$$

implies that

$$\mathbf{1}_{\{x_1\}} \land \mathbf{1}_{\{x_2\}} = \mathbf{1}_{\{x_1 \land x_2\}}.$$

Proposition 3.6. Let ∇ be a t-conorm on I and * be a binary operation on I. If Υ is a t-conorm, then, for $x_1, x_2 \in I$, one has $\mathbf{1}_{\{x_1\}} \Upsilon \mathbf{1}_{\{x_2\}} = \mathbf{1}_{\{x_1 \nabla x_2\}}$.

Proof. Proposition 3.4 yields

- (a) for $y, z \in I$, $\mathbf{1}_{\{x_1\}}(y) * \mathbf{1}_{\{x_2\}}(z) \in \{0, 1\};$
- (b) $\mathbf{1}_{\{x_1\}}(y) * \mathbf{1}_{\{x_2\}}(z) = 1$ if and only if $y = x_1$ and $z = x_2$.

This, together with

$$(\mathbf{1}_{\{x_1\}} \Upsilon \mathbf{1}_{\{x_2\}})(x) = \sup\{\mathbf{1}_{\{x_1\}}(y) * \mathbf{1}_{\{x_2\}}(z) \mid y \triangledown z = x\},\$$

implies that

$$1_{\{x_1\}} \curlyvee 1_{\{x_2\}} = 1_{\{x_1 \triangledown x_2\}}.$$

4. Construction of a t_r -norm on L

For any $f, g \in \mathbf{L}$, let $\eta_{f,g} = \inf\{x \in I \mid f^L(x) = 1\} \land \inf\{x \in I \mid g^L(x) = 1\}$ and $\xi_{f,g} = \sup\{x \in I \mid f^R(x) = 1\} \land \sup\{x \in I \mid g^R(x) = 1\}$. By Lemma 2.2, we have $\eta_{f,g} \leq \xi_{f,g}$.

Definition 4.1. Define a binary operation $\mathbf{\bigstar} : \mathbf{L}^2 \to \mathbf{M}$ as follows: for $f, g \in \mathbf{L}$,

- (1) f = 1_{1}, f ★g = g ★f = g;
 (2) g = 1_{{1}}, f ★g = g ★f = f;
- (3) $f \neq \mathbf{1}_{\{1\}}$ and $g \neq \mathbf{1}_{\{1\}}$,

$$(f \bigstar g)(t) = \begin{cases} f^{L}(t) \lor g^{L}(t), & t \in [0, \eta_{f,g}), \\ 1, & t \in [\eta_{f,g}, \xi_{f,g}), \\ f^{R}(\xi) \land g^{R}(\xi), & t = \xi_{f,g}, \\ 0, & t \in (\xi_{f,g}, 1]. \end{cases}$$
(4.1)

Clearly, $f \bigstar g$ is increasing on $[0, \xi_{f,g})$.

Proposition 4.1. For $f, g \in \mathbf{L}$, $f \triangleq g$ is normal and convex, i.e., $f \triangleq g \in \mathbf{L}$.

Proof. Consider the following two cases:

- (1) if $f = \mathbf{1}_{\{1\}}$ or $g = \mathbf{1}_{\{1\}}$, it is clear that $f \bigstar g \in \mathbf{L}$;
- (2) if $f \neq \mathbf{1}_{\{1\}}$ and $g \neq \mathbf{1}_{\{1\}}$, applying (4.1), it is easy to see that $f \triangleq g$ is convex, since it is increasing on $[0, \xi_{f,g})$ and decreasing on $[\xi_{f,g}, 1]$. It remains to show that $f \triangleq g$ is normal.
 - (a) If $\eta_{f,g} < \xi_{f,g}$, then $(f \bigstar g)(t) = 1$ for all $t \in [\eta_{f,g}, \xi_{f,g})$;
 - (b) If $\eta_{f,g} = \xi_{f,g}$, consider the following two subcases:

(b.1) $\eta_{f,g} = 0$. It follows from (4.1) that

$$(f \bigstar g)(t) = \begin{cases} f^R(0) \land g^R(0), & t = 0, \\ 0, & t \in (0, 1] \end{cases}$$

Since f and g are normal, from Lemma 2.1, it is clear that

$$f^{R}(0) \wedge g^{R}(0) = \sup_{x \in I} \{f(x)\} \wedge \sup_{x \in I} \{g(x)\} = 1.$$

(b.2) $\eta_{f,g} > 0$. From Proposition 2.2, it follows that

$$(f \bigstar g)^{L_{w}}(\eta_{f,g})$$

$$= \sup_{t \in [0,\eta_{f,g})} \{(f \bigstar g)(t)\}$$

$$= \sup_{t \in [0,\eta_{f,g})} \{f^{L}(t)\} \lor \sup_{t \in [0,\eta_{f,g})} \{g^{L}(t)\}$$

$$= f^{L_{w}}(\eta_{f,g}) \lor g^{L_{w}}(\eta_{f,g}).$$

If $f^{L_{\mathbf{w}}}(\eta_{f,g}) \vee g^{L_{\mathbf{w}}}(\eta_{f,g}) = 1$, then clearly $f \not\approx g$ is normal. If $f^{L_{\mathbf{w}}}(\eta_{f,g}) \vee g^{L_{\mathbf{w}}}(\eta_{f,g}) < 1$, noting that $1 = \sup_{t \in I} \{f(t)\} = f^{L_{\mathbf{w}}}(\eta_{f,g}) \vee f^{R}(\eta_{f,g})$ and $1 = \sup_{t \in I} \{g(t)\} = g^{L_{\mathbf{w}}}(\eta_{f,g}) \vee g^{R}(\eta_{f,g})$, we have

$$f^{R}(\eta_{f,g}) = g^{R}(\eta_{f,g}) = 1,$$

which, together with $\eta_{f,g} = \xi_{f,g}$, implies that

$$(f \bigstar g)(\eta_{f,g}) = 1.$$

Thus, $f \bigotimes g$ is normal.

Remark 1. (i) Proposition 4.1 shows that the binary operation \bigstar is closed on L, i.e., $\bigstar(L^2) \subset L$.

(ii) From the proof of Proposition 4.1, it follows that, for $f, g \in \mathbf{L}$, if $\eta_{f,g} = \xi_{f,g}$, then $(f \bigstar g)^{L_w}(\xi_{f,g}) = 1$ or $(f \, \bigstar \, g)(\xi_{f,g}) = 1.$

Proposition 4.2. For $f, g \in \mathbf{L} \setminus \{\mathbf{1}_{\{1\}}\},\$

$$(f \bigstar g)^{L}(t) = \begin{cases} f^{L}(t) \lor g^{L}(t), & t \in [0, \eta_{f,g}), \\ 1, & t \in [\eta_{f,g}, 1], \end{cases}$$
(4.2)

$$(f \bigstar g)^{R}(t) = \begin{cases} 1, & t \in [0, \xi_{f,g}), \\ f^{R}(\xi_{f,g}) \wedge g^{R}(\xi_{f,g}), & t = \xi_{f,g}, \\ 0, & t \in (\xi_{f,g}, 1]. \end{cases}$$
(4.3)

Proof. (1) If $\eta_{f,g} < \xi_{f,g}$, since $f^L(t) \lor g^L(t)$ is increasing, (4.1) evidently implies (4.2) and (4.3).

(2) If $\eta_{f,g} = \xi_{f,g}$, the result follows from Remark 1 (ii) and (4.1).

Theorem 4.1. \Leftrightarrow satisfies (01).

Proof. For
$$f, g \in \mathbf{L}$$
,

(A.1) if $f = \mathbf{1}_{\{1\}}$ or $g = \mathbf{1}_{\{1\}}$, then clearly $f \not\approx g = g \not\approx f$;

(A.2) if $f \neq \mathbf{1}_{\{1\}}$ and $g \neq \mathbf{1}_{\{1\}}$, then

$$(f \bigstar g)(t) = \begin{cases} f^L(t) \lor g^L(t), & t \in [0, \eta_{f,g}), \\ 1, & t \in [\eta_{f,g}, \xi_{f,g}), \\ f^R(\xi_{f,g}) \land g^R(\xi_{f,g}), & t = \xi_{f,g}, \\ 0, & t \in (\xi_{f,g}, 1], \end{cases}$$

and the commutativity of \bigstar follows from the commutativity of \lor and \land .

Lemma 4.1. For $f, g \in \mathbf{L} \setminus \{\mathbf{1}_{\{1\}}\}, f \not \approx g \neq \mathbf{1}_{\{1\}}$.

Proof. Suppose on the contrary that $f \not\approx g = \mathbf{1}_{\{1\}}$. Then, $\eta_{f,g} = \xi_{f,g} = 1$ and $f^L(t) \lor g^L(t) = 0$ for $t \in [0, 1)$. Since $f^L \ge f$ and $g^L \ge g$, we have f(t) = g(t) = 0 for $t \in [0, 1)$. This, together with $f, g \in \mathbf{L}$, implies that

$$f = g = \mathbf{1}_{\{1\}},$$

which is a contradiction.

Theorem 4.2. \Leftrightarrow satisfies (O2).

Proof. For $f, g, h \in \mathbf{L}$,

(B.1) if one of f, g, and h is equal to $\mathbf{1}_{\{1\}}$, then it is easy to verify that $(f \not\approx g) \not\approx h = f \not\approx (g \not\approx h)$;

(B.2) if none of f, g, and h are equal to $\mathbf{1}_{\{1\}}$, then

$$(f \bigstar g)(t) = \begin{cases} f^L(t) \lor g^L(t), & t \in [0, \eta_{f,g}), \\ 1, & t \in [\eta_{f,g}, \xi_{f,g}), \\ f^R(\xi_{f,g}) \land g^R(\xi_{f,g}), & t = \xi_{f,g}, \\ 0, & t \in (\xi_{f,g}, 1], \end{cases}$$

and

$$(g \bigstar h)(t) = \begin{cases} g^{L}(t) \lor h^{L}(t), & t \in [0, \eta_{g,h}), \\ 1, & t \in [\eta_{g,h}, \xi_{g,h}), \\ g^{R}(\xi_{g,h}) \land h^{R}(\xi_{g,h}), & t = \xi_{g,h}, \\ 0, & t \in (\xi_{g,h}, 1]. \end{cases}$$

By Lemma 4.1, we have $f \triangleq g \neq \mathbf{1}_{\{1\}}$ and $g \triangleq h \neq \mathbf{1}_{\{1\}}$. Proposition 4.2 implies that

$$(f \bigstar g)^{L}(t) = \begin{cases} f^{L}(t) \lor g^{L}(t), & t \in [0, \eta_{f,g}), \\ 1, & t \in [\eta_{f,g}, 1], \end{cases}$$
$$(f \bigstar g)^{R}(t) = \begin{cases} 1, & t \in [0, \xi_{f,g}), \\ f^{R}(\xi_{f,g}) \land g^{R}(\xi_{f,g}), & t = \xi_{f,g}, \\ 0, & t \in (\xi_{f,g}, 1], \end{cases}$$

and

$$(g \bigstar h)^{L}(t) = \begin{cases} g^{L}(t) \lor h^{L}(t), & t \in [0, \eta_{g,h}), \\ 1, & t \in [\eta_{g,h}, 1], \end{cases}$$
$$(g \bigstar h)^{R}(t) = \begin{cases} 1, & t \in [0, \xi_{g,h}), \\ g^{R}(\xi_{g,h}) \land h^{R}(\xi_{g,h}), & t = \xi_{g,h}, \\ 0, & t \in (\xi_{g,h}, 1]. \end{cases}$$

Since $f \triangleq g, g \triangleq h, f, h \in \mathbf{L} \setminus \{\mathbf{1}_{\{1\}}\}$, we have

$$((f \bigstar g) \bigstar h)(t) = \begin{cases} (f \bigstar g)^{L}(t) \lor h^{L}(t), & t \in [0, \eta_{f \bigstar g, h}), \\ 1, & t \in [\eta_{f \bigstar g, h}, \xi_{f \bigstar g, h}), \\ (f \bigstar g)^{R}(\xi_{f \bigstar g, h}) \land h^{R}(\xi_{f \bigstar g, h}), & t = \xi_{f \bigstar g, h}, \\ 0, & t \in (\xi_{f \bigstar g, h}, 1], \end{cases}$$
(4.4)

and

$$(f \not \approx (g \not \approx h))(t) = \begin{cases} f^{L}(t) \lor (g \not \approx h)^{L}(t), & t \in [0, \eta_{f,g \not \approx h}), \\ 1, & t \in [\eta_{f,g \not \approx h}, \xi_{f,g \not \approx h}), \\ f^{R}(\xi_{f,g \not \approx h}) \land (g \not \approx h)^{R}(\xi_{f,g \not \approx h}), & t = \xi_{f,g \not \approx h}, \\ 0, & t \in (\xi_{f,g \not \approx h}, 1]. \end{cases}$$

$$(4.5)$$

Clearly,

$$\eta := \eta_{f \bigstar g,h} = \eta_{f,g \bigstar h}$$
$$= \inf\{x \in I \mid f^L(x) = 1\}$$
$$\land \inf\{x \in I \mid g^L(x) = 1\}$$
$$\land \inf\{x \in I \mid h^L(x) = 1\}$$
$$= \eta_{f,g} \land \eta_{g,h},$$

and

$$\xi := \xi_{f \bigstar g,h} = \xi_{f,g \bigstar h}$$
$$= \sup\{x \in I \mid f^R(x) = 1\}$$
$$\wedge \sup\{x \in I \mid g^R(x) = 1\}$$
$$\wedge \sup\{x \in I \mid h^R(x) = 1\}$$
$$= \xi_{f,g} \land \xi_{g,h}.$$

Thus, for $t \in [0, \eta)$,

$$((f \bigstar g) \bigstar h)(t) = (f^L(t) \lor g^L(t)) \lor h^L(t),$$

and

$$(f \bigstar (g \bigstar h))(t) = f^L(t) \lor (g^L(t) \lor h^L(t)),$$

and the associativity holds. Clearly, for $t \in [\eta, \xi) \cup (\xi, 1]$,

$$((f \bigotimes g) \bigotimes h)(t) = (f \bigotimes (g \bigotimes h))(t).$$

To finish the proof we have to show that $((f \bigstar g) \bigstar h)(\xi) = (f \bigstar (g \bigstar h))(\xi)$.

Consider the following three cases:

(B.2.1) If $\xi_{f,g} = \xi_{g,h}$, then $\xi = \xi_{f,g} = \xi_{g,h}$, implying that

$$(f \bigstar g)^R(\xi) = f^R(\xi) \land g^R(\xi)$$

and

$$(g \bigstar h)^R(\xi) = g^R(\xi) \wedge h^R(\xi)$$

Then, (4.4) and (4.5) yield

$$((f \bigstar g) \bigstar h)(\xi) = f^R(\xi) \land g^R(\xi) \land h^R(\xi) = (f \bigstar (g \bigstar h))(\xi).$$

(B.2.2) If $\xi_{f,g} < \xi_{g,h}$, then $\xi_{f,g} = \sup\{x \in I \mid f^R(x) = 1\} < \sup\{x \in I \mid g^R(x) = 1\}$ (as $\sup\{x \in I \mid f^R(x) = 1\} \ge \sup\{x \in I \mid g^R(x) = 1\}$ implies that $\xi_{f,g} = \sup\{x \in I \mid g^R(x) = 1\} \ge \sup\{x \in I \mid g^R(x) = 1\} \ge \sup\{x \in I \mid g^R(x) = 1\} = \xi_{g,h}$), which means that there exists $\hat{x} > \xi_{f,g}$ such that $g^R(\hat{x}) = 1$. Thus,

$$g^R(\xi_{f,g}) \ge g^R(\hat{x}) = 1$$

Therefore, since $\xi = \xi_{f,g} \wedge \xi_{g,h} = \xi_{f,g}$, we get

$$(f \bigstar g)^{R}(\xi) = (f \bigstar g)^{R}(\xi_{f,g}) = f^{R}(\xi_{f,g}) \wedge g^{R}(\xi_{f,g}) = f^{R}(\xi_{f,g}).$$
(4.6)

From $\xi_{f,g} < \xi_{g,h} \le \sup\{x \in I \mid h^R(x) = 1\}$, it follows that there exists $x' > \xi_{f,g}$ such that $h^R(x') = 1$, implying that

$$h^R(\xi) = h^R(\xi_{f,g}) \ge h^R(x') = 1.$$
 (4.7)

(4.7) together with (4.4) and (4.6) implies that

$$((f \bigstar g) \bigstar h)(\xi) = f^R(\xi).$$

Since $\xi = \xi_{f,g} < \xi_{g,h}$, then we have $(g \not\approx h)^R(\xi) = 1$, which together with (4.5) implies that

$$(f \bigstar (g \bigstar h))(\xi) = f^R(\xi) \land (g \bigstar h)^R(\xi) = f^R(\xi)$$

Therefore,

$$((f \bigstar g) \bigstar h)(\xi) = (f \bigstar (g \bigstar h))(\xi) = f^R(\xi).$$

(B.2.3) If $\xi_{f,g} > \xi_{g,h}$, then similarly as in the previous case we can show that

$$((f \bigstar g) \bigstar h)(\xi) = (f \bigstar (g \bigstar h))(\xi) = h^R(\xi).$$

Summing up (B.2.1)–(B.2.3), we have

$$((f \otimes g) \otimes h)(\xi) = (f \otimes (g \otimes h))(\xi)$$

Theorem 4.3. \Leftrightarrow satisfies (O3).

Proof. This follows directly from (1) and (2) of Definition 4.1.

Theorem 4.4. \Leftrightarrow satisfies (O4).

Proof. We have to show that, for $f, g, h \in \mathbf{L}$ with $f \sqsubseteq g, f \not\approx h \sqsubseteq g \not\approx h$. There are the following possible cases:

(D.1) if $h = \mathbf{1}_{\{1\}}$, then $f \not\approx h = f \sqsubseteq g = g \not\approx h$; (D.2) if $f = \mathbf{1}_{\{1\}}$, then $g = \mathbf{1}_{\{1\}}$ (as $f \sqsubseteq g$), implying that $f \not\approx h = h \sqsubseteq h = g \not\approx h$; (D.3) if $g = \mathbf{1}_{\{1\}}$ and $f \neq \mathbf{1}_{\{1\}} \neq h$, then

$$(f \bigstar h)(t) = \begin{cases} f^L(t) \lor h^L(t), & t \in [0, \eta_{f,h}), \\ 1, & t \in [\eta_{f,h}, \xi_{f,h}) \\ f^R(\xi_{f,h}) \land h^R(\xi_{f,h}), & t = \xi_{f,h}, \\ 0, & t \in (\xi_{f,h}, 1]. \end{cases}$$

By Proposition 4.2, one has

$$(f \bigstar h)^{L}(t) = \begin{cases} f^{L}(t) \lor h^{L}(t), & t \in [0, \eta_{f,h}), \\ 1, & t \in [\eta_{f,h}, 1], \end{cases}$$
(4.8)

and

$$(f \bigstar h)^{R}(t) = \begin{cases} 1, & t \in [0, \xi_{f,h}), \\ f^{R}(\xi_{f,h}) \wedge h^{R}(\xi_{f,h}), & t = \xi_{f,h}, \\ 0, & t \in (\xi_{f,h}, 1]. \end{cases}$$
(4.9)

Clearly,

and

$$(f \not\approx h)^R(\xi_{f,h}) \le h^R(\xi_{f,h}).$$
 (4.10)

Additionally, for $t < \xi_{f,h}$, it follows from $\xi_{f,h} \leq \sup\{x \in I \mid h^R(x) = 1\}$ that there exists $t < \hat{x} \leq \xi$ such that $h^R(\hat{x}) = 1$. Thus, $h^R(t) \geq h^R(\hat{x}) = 1$ since h^R is decreasing, implying that, for $t \in [0, \xi)$,

 $(f \bigstar h)^L \ge h^L,$

$$(f \not\approx h)^R(t) \le 1 = h^R(t).$$
 (4.11)

(4.11) together with (4.10) and (4.9) implies that

$$(f \bigstar h)^R \le h^R.$$

By Theorem 2.1 there is

$$f \, \bigstar h \sqsubseteq h = g \, \bigstar h$$

(D.4) If $f \neq \mathbf{1}_{\{1\}}$, $g \neq \mathbf{1}_{\{1\}}$, and $h \neq \mathbf{1}_{\{1\}}$, then from the definition of \mathfrak{A} , it follows that

$$(f \bigstar h)(t) = \begin{cases} f^L(t) \lor h^L(t), & t \in [0, \eta_{f,h}), \\ 1, & t \in [\eta_{f,h}, \eta_{f,h}), \\ f^R(\eta_{f,h}) \land h^R(\eta_{f,h}), & t = \eta_{f,h}, \\ 0, & t \in (\eta_{f,h}, 1], \end{cases}$$

and

$$(g \bigstar h)(t) = \begin{cases} g^{L}(t) \lor h^{L}(t), & t \in [0, \eta_{g,h}), \\ 1, & t \in [\eta_{g,h}, \eta_{g,h}), \\ g^{R}(\eta_{g,h}) \land h^{R}(\eta_{g,h}), & t = \eta_{g,h}, \\ 0, & t \in (\eta_{g,h}, 1]. \end{cases}$$

From $f \sqsubseteq g$ and Theorem 2.1, it follows that $f^L \ge g^L$ and $f^R \le g^R$. Therefore,

$$\{x \in I \mid g^{L}(x) = 1\} \subseteq \{x \in I \mid f^{L}(x) = 1\},\$$

and

$$\{x \in I \mid f^{R}(x) = 1\} \subseteq \{x \in I \mid g^{R}(x) = 1\},\$$

implying that

$$\inf\{x \in I \mid g^{L}(x) = 1\} \ge \inf\{x \in I \mid f^{L}(x) = 1\},\$$

and

$$\sup\{x \in I \mid f^{R}(x) = 1\} \le \sup\{x \in I \mid g^{R}(x) = 1\}$$

Thus,

$$\eta_{f,h} \leq \eta_{g,h}$$
 and $\eta_{f,h} \leq \eta_{g,h}$.

Further, by Proposition 4.2 there is

$$(f \bigstar h)^{L}(t) = \begin{cases} f^{L}(t) \lor h^{L}(t), & t \in [0, \eta_{f,h}), \\ 1, & t \in [\eta_{f,h}, 1], \end{cases}$$
$$(f \bigstar h)^{R}(t) = \begin{cases} 1, & t \in [0, \eta_{f,h}), \\ f^{R}(\eta_{f,h}) \land h^{R}(\eta_{f,h}), & t = \eta_{f,h}, \\ 0, & t \in (\eta_{f,h}, 1], \end{cases}$$

and

$$(g \bigstar h)^{L}(t) = \begin{cases} g^{L}(t) \lor h^{L}(t), & t \in [0, \eta_{g,h}), \\ 1, & t \in [\eta_{g,h}, 1], \end{cases}$$
$$(g \bigstar h)^{R}(t) = \begin{cases} 1, & t \in [0, \eta_{g,h}), \\ g^{R}(\eta_{g,h}) \land h^{R}(\eta_{g,h}), & t = \eta_{g,h}, \\ 0, & t \in (\eta_{g,h}, 1]. \end{cases}$$

From $f^R \leq g^R$, it follows that

$$(f \bigstar h)^R \le (g \bigstar h)^R. \tag{4.12}$$

From $f^L \ge g^L$, it follows that, for $t \in [0, \eta_{f,h})$,

$$(f \bigstar h)^L(t) \ge (g \bigstar h)^L(t).$$

 $(f \bigstar h)^L(t) = 1 \ge (g \bigstar h)^L(t).$

It is clear that, for $t \in [\eta_{f,h}, 1]$,

Thus,

$$(f \bigstar h)^L \ge (g \bigstar h)^L. \tag{4.13}$$

(4.13) together with (4.12) and Theorem 2.1 implies that

$$f \bigstar h \sqsubseteq g \bigstar h$$

Theorem 4.5. \Leftrightarrow satisfies (05).

Proof. Assume $0 \le a \le b \le 1$. Then we can distinguish the following cases:

(E.1) If
$$a = 1$$
, then $\mathbf{1}_{[0,1]} \bigstar \mathbf{1}_{[a,b]} = \mathbf{1}_{[0,1]} \bigstar \mathbf{1}_{\{1\}} = \mathbf{1}_{[0,1]};$
(E.2) If $a < 1$, then
 $\mathbf{1}_{[0,1]}^L \equiv 1,$

$$\mathbf{1}_{[0,1]}^{R} \equiv 1,$$
$$\mathbf{1}_{[a,b]}^{L}(x) = \begin{cases} 0, & x \in [0,a), \\ 1, & x \in [a,1], \end{cases}$$
$$\mathbf{1}_{R}^{R}(x) = \int 1, & x \in [0,b], \end{cases}$$

and

$$\mathbf{1}_{[a,b]}^{R}(x) = \begin{cases} 1, & x \in [0,b], \\ 0, & x \in (b,1], \end{cases}$$

which implies that $\inf\{x \in I \mid \mathbf{1}_{[0,1]}^L(x) = 1\} \land \inf\{x \in I \mid \mathbf{1}_{[a,b]}^L(x) = 1\} = 0$ and $\sup\{x \in I \mid \mathbf{1}_{[0,1]}^R(x) = 1\} \land \sup\{x \in I \mid \mathbf{1}_{[a,b]}^R(x) = 1\} = b.$

Consider the following two subcases:

(E.2.1) If b = 0, we have

$$(\mathbf{1}_{[0,1]} \bigstar \mathbf{1}_{[a,b]})(t) = \begin{cases} \mathbf{1}_{[0,1]}^{L}(t) \lor \mathbf{1}_{[a,b]}^{L}(t), & t \in [0,0), \\ 1, & t = 0, \\ 0, & t \in (0,1], \end{cases}$$

implying that $\mathbf{1}_{[0,1]} \bigstar \mathbf{1}_{[a,b]} = \mathbf{1}_{[0,0]} = \mathbf{1}_{[0,b]}$ as $[0,0) = \emptyset$.

(E.2.2) If b > 0, we have

$$(\mathbf{1}_{[0,1]} \bigstar \mathbf{1}_{[a,b]})(t) = \begin{cases} \mathbf{1}_{[0,1]}^{L}(t) \lor \mathbf{1}_{[a,b]}^{L}(t), & t \in [0,0), \\ 1, & t \in [0,b), \\ \mathbf{1}_{[0,1]}^{R}(b) \land \mathbf{1}_{[a,b]}^{R}(b), & t = b, \\ 0, & t \in (b,1]. \end{cases}$$
(4.14)

(4.14) together with $\mathbf{1}_{[0,1]}^R(b) \wedge \mathbf{1}_{[a,b]}^R(b) = 1$ implies that $\mathbf{1}_{[0,1]} \mathbf{\bigstar} \mathbf{1}_{[a,b]} = \mathbf{1}_{[0,b]}$.

Theorem 4.6. For $x_1, x_2 \in I$, $\mathbf{1}_{\{x_1\}} \bigstar \mathbf{1}_{\{x_2\}} = \mathbf{1}_{\{x_1 \land x_2\}}$. In particular, \bigstar satisfies (O6).

Proof. Clearly, $\mathbf{1}_{\{x_1\}} \not\approx \mathbf{1}_{\{x_2\}} = \mathbf{1}_{\{x_1\}}$ when $x_2 = 1$ by Theorem 4.3. Moreover, for $x \in I$,

$$\mathbf{1}^L_{\{x\}}(t) = \begin{cases} 0, & t \in [0, x), \\ 1, & t \in [x, 1], \end{cases}$$

and

$$\mathbf{1}^{R}_{\{x\}}(t) = \begin{cases} 1, & t \in [0, x], \\ 0, & t \in (x, 1]. \end{cases}$$

Then, for $0 \le x_1 \le x_2 < 1$, $\inf\{x \in I \mid \mathbf{1}_{\{x_1\}}^L(x) = 1\} \land \inf\{x \in I \mid \mathbf{1}_{\{x_2\}}^L(x) = 1\} = x_1$ and $\sup\{x \in I \mid \mathbf{1}_{\{x_1\}}^R(x) = 1\} \land \sup\{x \in I \mid \mathbf{1}_{\{x_2\}}^R(x) = 1\} = x_1$. Clearly, $(\mathbf{1}_{\{x_1\}} \bigstar \mathbf{1}_{\{x_2\}})^{L_w}(x_1) = 0$. From Remark 1 (ii), it follows that

$$(\mathbf{1}_{\{x_1\}} \bigstar \mathbf{1}_{\{x_2\}})(t) = \begin{cases} 0, & t \in [0, x_1), \\ 1, & t = x_1, \\ 0, & t \in (x_1, 1], \end{cases}$$

which, together with the commutativity of \bigstar , implies that

$$\mathbf{1}_{\{x_1\}} \bigstar \mathbf{1}_{\{x_2\}} = \mathbf{1}_{\{x_2\}} \bigstar \mathbf{1}_{\{x_1\}} = \mathbf{1}_{\{x_1\}} \in \mathbf{J}.$$

Theorem 4.7. For $[a_1, b_1]$, $[a_2, b_2] \subset I$, $\mathbf{1}_{[a_1, b_1]} \bigstar \mathbf{1}_{[a_2, b_2]} = \mathbf{1}_{[a_1 \land a_2, b_1 \land b_2]}$. In particular, \bigstar satisfies (O7). Proof. Clearly, $\mathbf{1}_{[a_1, b_1]} \bigstar \mathbf{1}_{[a_2, b_2]} = \mathbf{1}_{[a_2, b_2]} \bigstar \mathbf{1}_{[a_1, b_1]} \in \mathbf{K}$ when $[a_1, b_1] = \{1\}$ or $[a_2, b_2] = \{1\}$ by Theorem 4.3.

Moreover, for $0 \le a \le b \le 1$,

$$\mathbf{1}_{[a,b]}^{L}(t) = \begin{cases} 0, & t \in [0,a), \\ 1, & t \in [a,1], \end{cases}$$

and

$$\mathbf{1}_{[a,b]}^{R}(t) = \begin{cases} 1, & t \in [0,b], \\ 0, & t \in (b,1]. \end{cases}$$

Then, for $[a_1, b_1], [a_2, b_2] \subset I$ with $[a_1, b_1] \neq \{1\}$ and $[a_2, b_2] \neq \{1\}$, we have $\inf\{x \in I \mid \mathbf{1}_{[a_1, b_1]}^L(x) = 1\} \wedge \inf\{x \in I \mid \mathbf{1}_{[a_2, b_2]}^R(x) = 1\} = a_1 \wedge a_2$ and $\sup\{x \in I \mid \mathbf{1}_{[a_1, b_1]}^R(x) = 1\} \wedge \sup\{x \in I \mid \mathbf{1}_{[a_2, b_2]}^R(x) = 1\} = a_1 \wedge a_2$.

 $b_1 \wedge b_2$. From (4.1), it follows that

$$\begin{aligned} &(\mathbf{1}_{[a_1,b_1]} \bigstar \mathbf{1}_{[a_2,b_2]})(t) \\ &= \begin{cases} \mathbf{1}_{[a_1,b_1]}^L(t) \lor \mathbf{1}_{[a_2,b_2]}^L(t), & t \in [0,a_1 \land a_2), \\ 1, & t \in [a_1 \land a_2, b_1 \land b_2), \\ \mathbf{1}_{[a_1,b_1]}^R(t) \land \mathbf{1}_{[a_2,b_2]}^R(t), & t = b_1 \land b_2, \\ 0, & t \in (b_1 \land b_2, 1], \end{cases} \\ &= \begin{cases} 0, & t \in [0,a_1 \land a_2), \\ 1, & t \in [a_1 \land a_2, b_1 \land b_2], \\ 0, & t \in (b_1 \land b_2, 1], \end{cases} \end{aligned}$$

which, together with the commutativity of \bigstar , implies that

$$\mathbf{1}_{[a_1,b_1]} \bigstar \mathbf{1}_{[a_2,b_2]} = \mathbf{1}_{[a_2,b_2]} \bigstar \mathbf{1}_{[a_1,b_1]} = \mathbf{1}_{[a_1 \land a_2, b_1 \land b_2]} \in \mathbf{K}.$$

Theorems 4.1–4.7 imply the following result.

Theorem 4.8. The binary operation \Leftrightarrow is a t_r -norm on **L**.

5. \bigstar cannot be obtained by \land and \curlyvee

This section shows that the t_r -norm \Leftrightarrow constructed in Section 4 cannot be obtained by operations \land or \curlyvee .

The following theorem provides a sufficient condition ensuring that * is a *t*-norm on *I*.

Theorem 5.1. [13, Theorem 21] Let \triangle be a continuous t-norm on I and * be a binary operation on I. If λ is a t-norm on \mathbf{L} , then * is a t-norm.

Theorem 5.2. For any binary operation * on I and any t-norm \triangle on I, there exist $f, g \in \mathbf{L}$ such that $f \not \approx g \neq f \land g$, *i.e.*, $\not \approx$ cannot be obtained by \land .

Proof. Suppose, on the contrary, that there exist a binary operation * on I and a t-norm \triangle on I such that, for any $f, g \in \mathbf{L}$, one has $f \not\approx g = f \land g$.

First we will show that $\triangle = \land$.

For $x_1, x_2 \in I$, Theorem 4.6 gives

$$1_{\{x_1\}} lpha 1_{\{x_2\}} = 1_{\{x_1 \land x_2\}}.$$

Further, Theorem 4.8 and Proposition 3.5 yield

$$\mathbf{1}_{\{x_1\}} \bigstar \mathbf{1}_{\{x_2\}} = \mathbf{1}_{\{x_1\}} \land \mathbf{1}_{\{x_2\}} = \mathbf{1}_{\{x_1 \triangle x_2\}}.$$

Thus,

$$x_1 \wedge x_2 = x_1 \bigtriangleup x_2$$
 for all $x_1, x_2 \in I$, i.e., $\bigtriangleup = \land$.

Clearly, $\Delta = \wedge$ is a continuous *t*-norm on *I*. From Theorem 4.8 and Theorem 5.1, it follows that * is a *t*-norm on *I* and

$$(f \bigstar g)(x) = \sup\{f(y) \ast g(z) \mid y \land z = x\}.$$
(5.1)

Choose a function $\psi \in \mathbf{L}$ by

$$\psi(x) = \begin{cases} 1, & x \in [0, 0.75], \\ 0.5, & x \in (0.75, 1]. \end{cases}$$




Figure 1: The function ψ .

 $\psi^L(x) \equiv 1,$

Then,

$$\begin{split} \psi^R(x) &= \begin{cases} 1, & x \in [0, 0.75], \\ 0.5, & x \in (0.75, 1], \end{cases} \\ \mathbf{1}^L_{\{0.8\}}(x) &= \begin{cases} 0, & x \in [0, 0.8), \\ 1, & x \in [0.8, 1], \end{cases} \end{split}$$

and

$$\mathbf{1}_{\{0.8\}}^{R}(x) = \begin{cases} 1, & x \in [0, 0.8], \\ 0, & x \in (0.8, 1]. \end{cases}$$

From (4.1), we have

$$(\psi \bigstar \mathbf{1}_{\{0.8\}})(t) = \begin{cases} 1, & x \in [0, 0.75], \\ 0, & x \in (0.75, 1]. \end{cases}$$

In particular,

$$(\psi \bigstar \mathbf{1}_{\{0,8\}})(0.8) = 0. \tag{5.2}$$

(5.2) together with (5.1) and the fact that * is a *t*-norm implies that

$$\begin{split} 0 &= (\psi \bigstar \mathbf{1}_{\{0.8\}})(0.8) \\ &= \sup \left\{ \psi(y) \ast \mathbf{1}_{\{0.8\}}(z) \mid y \land z = 0.8 \right\} \\ &\geq \psi(0.8) \ast \mathbf{1}_{\{0.8\}}(0.8) = 0.5 \ast 1 = 0.5 \end{split}$$

which is a contradiction.

Theorem 5.3. Let * be a binary operation on I, \triangle be a t-norm on I, and ∇ be a t-conorm on I. Then, we have

- (1) the binary operation Υ defined by (2.2) is not a t-norm on L;
- (2) the binary operation \land defined by (2.1) is not a t-conorm on **L**.

Proof. (1) Suppose on the contrary that Υ is a *t*-norm on **L**. Then, we have $f \Upsilon \mathbf{1}_{\{1\}} = f$ for any $f \in \mathbf{L}$ as $\mathbf{1}_{\{1\}}$ is the neutral element of Υ . For $0 \leq \zeta \leq 1$, take $f_{\zeta} : I \to I$ as $f_{\zeta}(x) = (1-\zeta)x + \zeta$. Clearly, $f_{\zeta} \in \mathbf{L}$, which together with Lemma 3.1 implies that

$$\zeta = f_{\zeta}(0) = (f_{\zeta} \uparrow \mathbf{1}_{\{1\}})(0) = f_{\zeta}(0) \ast \mathbf{1}_{\{1\}}(0) = \zeta \ast 0.$$
(5.3)

Similarly, we have $0 * \zeta = \zeta$. Moreover, for $g \in \mathbf{L}$ with g(x) = 1 - x $(x \in I)$, we have

$$\begin{split} 0.5 &= g(0.5) = (g \curlyvee \mathbf{1}_{\{1\}})(0.5) \\ &= \sup\{g(y) * \mathbf{1}_{\{1\}}(z) \mid y \bigtriangledown z = 0.5\} \\ &\geq g(0) * \mathbf{1}_{\{1\}}(0.5) \text{ (as } 0 \bigtriangledown 0.5 = 0.5) \\ &= g(0) * 0 = 1 \text{ (by } (5.3)), \end{split}$$

which is a contradiction.

(2) Similarly as in the previous case we can prove that λ is not a *t*-conorm on **L**.

Theorems 4.8 and 5.3 imply the following result.

Corollary 5.1. The t_r -norm \bigstar cannot be obtained by \curlyvee .

Remark 2. Theorems 4.8 and 5.2, and Corollary 5.1 show that there exists a t_r -norm \bigstar on **L**, which cannot be obtained using the formulas defining the operations λ and γ . This gives a positive answer to Question 2.

6. A t_r -conorm that is not obtained by \land and \curlyvee

This section introduces the dual operation for every binary operation on **M** and proves the duality between t_r -norm and t_r -conorm, also between λ and γ . As an application, a t_r -conorm on **L**, which cannot be obtained by λ and γ , is obtained.

Definition 6.1. Let \Diamond be a binary operation on **M**. Define the *dual operation* \Diamond^{\complement} of \Diamond as follows: for $f, g \in \mathbf{M}$,

$$f \Diamond^{\mathsf{C}} g = \neg((\neg f) \Diamond(\neg g)).$$

Proposition 6.1. For a binary operation \Diamond on \mathbf{M} , $(\Diamond^{\complement})^{\complement} = \Diamond$.

Proof. For $f, g \in \mathbf{M}$, from Definition 6.1, it follows that $f(\Diamond^{\complement})^{\complement}g = \neg((\neg f)\Diamond^{\complement}(\neg g)) = \neg(\neg(f\Diamond g)) = f\Diamond g$.

Theorem 6.1. Let \Diamond be a binary operation on \mathbf{M} such that $\Diamond(\mathbf{L}^2) \subset \mathbf{L}$, *i.e.*, \Diamond is closed on \mathbf{L} . Then, $\Diamond|_{\mathbf{L}^2}$ is a t_r -norm (resp., t-norm) on \mathbf{L} if and only if $\Diamond^{\complement}|_{\mathbf{L}^2}$ is a t_r -conorm (resp., t-conorm) on \mathbf{L} .

Proof. Clearly, \Diamond^{\complement} is closed on **L**. By Proposition 6.1, it suffices to show that $\Diamond^{\complement}|_{\mathbf{L}^2}$ is a t_r -conorm provided that $\Diamond|_{\mathbf{L}^2}$ is a t_r -norm.

(i) \Diamond^{\complement} satisfies (O1).

For $f, g \in \mathbf{L}$, since \Diamond satisfies (O1), we have $f \Diamond^{\complement} g = \neg((\neg f) \Diamond(\neg g)) = \neg((\neg g) \Diamond(\neg f)) = g \Diamond^{\complement} f$.

(ii) \Diamond^{\complement} satisfies (O2).

For $f, g, h \in \mathbf{L}$, we get

$$(f \diamond^{\complement} g) \diamond^{\complement} h = \neg((\neg(f \diamond^{\complement} g)) \diamond(\neg h)) = \neg(((\neg f) \diamond(\neg g)) \diamond(\neg h)),$$

and

$$f\Diamond^{\complement}(g\Diamond^{\complement}h) = \neg((\neg f)\Diamond(\neg(g\Diamond^{\complement}h))) = \neg((\neg f)\Diamond((\neg g)\Diamond(\neg h))),$$

and the associativity of \Diamond^{\complement} follows from the associativity of \Diamond .

(iii) \Diamond^{\complement} satisfies (O3').

For $f \in \mathbf{L}$, since $\mathbf{1}_{\{1\}}$ is the neutral element of \Diamond , we have

$$f \diamond^{\complement} \mathbf{1}_{\{0\}} = \neg ((\neg f) \diamond (\neg \mathbf{1}_{\{0\}})) = \neg ((\neg f) \diamond \mathbf{1}_{\{1\}}) = \neg (\neg f) = f.$$

(iv) \Diamond^{\complement} satisfies (O4).

For $f, g, h \in \mathbf{L}$ with $f \sqsubseteq g$, Proposition 2.1 and Theorem 2.1 yield

$$(\neg f)^{L}(x) = (\neg (f^{R}))(x) = f^{R}(1-x) \le g^{R}(1-x) = (\neg (g^{R}))(x) = (\neg g)^{L}(x),$$

and

$$(\neg f)^{R}(x) = (\neg (f^{L}))(x) = f^{L}(1-x) \ge g^{L}(1-x) = (\neg (g^{L}))(x) = (\neg g)^{R}(x).$$

Applying again Theorem 2.1, we obtain $\neg g \sqsubseteq \neg f$. Since \Diamond satisfies (O4), we have

$$(\neg g)\Diamond(\neg h) \sqsubseteq (\neg f)\Diamond(\neg h).$$

Thus,

$$f \Diamond^{\mathsf{L}} h = \neg((\neg f) \Diamond(\neg h)) \sqsubseteq \neg((\neg g) \Diamond(\neg h)) = g \Diamond^{\mathsf{L}} h$$

(v) \Diamond^{\complement} satisfies (O5').

Since \Diamond satisfies (O5), it follows that

$$\begin{split} \mathbf{1}_{[0,1]} &\diamond^{\complement} \mathbf{1}_{[a,b]} \\ &= \neg ((\neg \mathbf{1}_{[0,1]}) \diamond (\neg \mathbf{1}_{[a,b]})) \\ &= \neg (\mathbf{1}_{[0,1]} \diamond \mathbf{1}_{[1-b,1-a]}) \\ &= \neg \mathbf{1}_{[0,1-a]} = \mathbf{1}_{[a,1]}. \end{split}$$

(vi) \Diamond^{\complement} satisfies (O6).

For $x_1, x_2 \in I$, since \diamond satisfies (O6), then there exists $x_3 \in I$ such that $\mathbf{1}_{\{1-x_1\}} \diamond \mathbf{1}_{\{1-x_2\}} = \mathbf{1}_{\{x_3\}}$, implying that

$$\begin{aligned} \mathbf{1}_{\{x_1\}} \Diamond^{\mathbb{C}} \mathbf{1}_{\{x_2\}} \\ &= \neg ((\neg \mathbf{1}_{\{x_1\}}) \Diamond (\neg \mathbf{1}_{\{x_2\}})) \\ &= \neg (\mathbf{1}_{\{1-x_1\}} \Diamond \mathbf{1}_{\{1-x_2\}}) \\ &= \neg \mathbf{1}_{\{x_3\}} = \mathbf{1}_{\{1-x_3\}} \in \mathbf{J}. \end{aligned}$$

(vii) \Diamond^{\complement} satisfies (O7).

For $[a_1, b_1]$, $[a_2, b_2] \subset I$, since \Diamond satisfies (O7), then there exist $[a_3, b_3] \subset I$ such that $\mathbf{1}_{[1-b_1, 1-a_1]} \Diamond \mathbf{1}_{[1-b_2, 1-a_2]} = \mathbf{1}_{[a_3, b_3]}$, implying that

Theorem 6.2. A binary operation on **M** is obtained by λ if and only if its dual operation is obtained by γ .

Proof. By Proposition 6.1, it suffices to prove the necessity. Assume that λ is a binary operation satisfying that there exist a binary operation * on I and a *t*-norm Δ on I such that, for $f, g \in \mathbf{M}$, $(f \land g)(x) = \sup\{f(y) * g(z) \mid y \Delta z = x\}$. Take $\nabla : I^2 \to I$ as $x \nabla y = 1 - (1 - x)\Delta(1 - y)$ for any $(x, y) \in I^2$. Clearly, ∇ is a *t*-conorm on I. For any $f, g \in \mathbf{M}$ and $x \in I$, we have

$$(f \wedge^{\mathsf{C}} g)(x) = (\neg((\neg f) \wedge (\neg g)))(x) = ((\neg f) \wedge (\neg g))(1 - x) = \sup\{(\neg f)(y) * (\neg g)(z) \mid y \land z = 1 - x\} = \sup\{f(1 - y) * g(1 - z) \mid y \land z = 1 - x\} = \sup\{f(y) * g(z) \mid y \nabla z = x\},$$

implying that $\mathcal{A}^{\complement}$ can be obtained by Υ .

Theorems 4.8, 6.1, and 6.2 imply the following result.

Corollary 6.1. Let $\star = \star^{\complement}$. Then,

- (1) \star is a t_r -conorm on **L**;
- (2) \bigstar cannot be obtained by \land and \curlyvee .

7. Conclusion

Employing the functions f^L and f^R , we have constructed in this paper two binary operations \bigstar and $\bigstar = \bigstar^{\mathbb{C}}$ on \mathbf{L} (see Definitions 4.1 and 6.1) and proved that \bigstar is a t_r -norm on \mathbf{L} and \bigstar is a t_r -conorm on \mathbf{L} . Both \bigstar and \bigstar cannot be obtained by using the formulas defining the operations \land and \curlyvee . These results give a positive answer to an open problem (see Question 2) in [3]. Combining this result with our main results in [13], the two open problems posed by Hernández et al. [3] are completely solved.

Acknowledgements

The authors express their sincere thanks to the editors and reviewers for their most valuable comments and suggestions in improving this paper greatly.

References

References

- [1] J. Harding, C. Walker, E. Walker, Convex normal functions revisited, Fuzzy Sets Syst., 161 (2010), 1343–1349.
- [2] J. Harding, C. Walker, E. Walker, Lattices of convex normal functions, Fuzzy Sets Syst., 159 (2008), 1061–1071.
- [3] P. Hernández, S. Cubillo, C. Torres-Blanc, On t-norms for type-2 fuzzy sets, IEEE Trans. Fuzzy Syst., 23 (2015), 1155–1163.
- [4] N. N. Karnik, J. M. Mendel, Operations on type-2 fuzzy sets, Fuzzy Sets Syst., 122 (2001), 327–348.
- [5] N. N. Karnik, J. M. Mendel, Centroid of a type-2 fuzzy set, Inform. Sci., 132 (2001), 195–220.
- [6] P. Klement, R. Mesiar, E. Pap, Triangular Norms. in *Trends in Logic: Studia Logica Library*, vol. 8. Dordrecht, The Netherlands: Kluwer, 2000.
- [7] J. M. Mendel, Uncertain Rule-Based Fuzzy Logic Systems: Introduction and New Directions. Prentice-Hall, Upper Saddle River, NJ, 2001.
- [8] J. M. Mendel, R. John, Type-2 fuzzy sets made simple, IEEE Trans. Fuzzy Syst., 10 (2002), 117–127.
- [9] K. Menger, Statistical metrics, Proc. Nat. Acad. Sci. USA, 37 (1942), 535-537.
- [10] B. Schweizer, A. Sklar, Associative functions and statistical triangle inequalities, Publ. Math., 8 (1961), 169–186.
- [11] C. Walker, E. Walker, The algebra of fuzzy truth values, Fuzzy Sets Syst., 149 (2005), 309–347.
- [12] C. Walker, E. Walker, T-norms for type-2 fuzzy sets. in Proc. IEEE Int Conf Fuzzy Syst, Vancouver, BC, Canada, 2006, pp. 1235–1239.
- [13] X. Wu, G. Chen, Answering an open problem on t-norms for type-2 fuzzy sets, Inform. Sci., 522 (2020), 124–133.
- [14] L. Zadeh, Fuzzy sets, Information and Control, 20 (1965), 301–312.
- [15] L. Zadeh, The concept of a linguistic variable and its application to approximate reasoning–I, Inform. Sci., 8 (1975), 199–249.