

CYCLOTOMIC QUIVER HECKE ALGEBRAS CORRESPONDING TO MINUSCULE REPRESENTATIONS

EUIYONG PARK

ABSTRACT. In the paper, we give an explicit basis of the cyclotomic quiver Hecke algebra corresponding to a minuscule representation of finite type.

INTRODUCTION

The quiver Hecke algebras (also known as Khovanov-Lauda-Rouquier algebras) were introduced by Khovanov and Lauda [15, 16] and independently Rouquier [18] to categorify the half of a quantum group. For a dominant integral weight Λ , the quiver Hecke algebras have special quotient algebras R^Λ which give a categorification of the irreducible highest weight module $V(\Lambda)$ [10]. In the viewpoint of categorification, these algebras have been the focus of many studies and various new features were discovered.

In the paper, we give an explicit basis of the cyclotomic quiver Hecke algebra corresponding to a *minuscule representation* in terms of crystals. A minuscule representation is an irreducible highest weight $U_q(\mathfrak{g})$ -module $V(\Lambda_i)$ such that the Weyl group acts transitively on its weight space. The possible types for minuscule representations are A_n , B_n , C_n , D_n , E_6 and E_7 , and the nontrivial highest weights of minuscule representations are marked by \circ in (1.2). Let $V(\Lambda_i)$ be a minuscule representation. Since every weight space $V(\Lambda_i)_\xi$ is extremal, the corresponding cyclotomic quiver Hecke algebra $R^{\Lambda_i}(\xi)$ is simple. Thus, since every irreducible R -module is absolutely irreducible, we obtain an explicit basis of $R^{\Lambda_i}(\xi)$ by constructing a simple $R^{\Lambda_i}(\xi)$ -module combinatorially in terms of the crystal $B(\Lambda_i)$. Our main tool is a *homogeneous representation*, which was introduced by Kleshchev and Ram [17]. In [17], they gave a combinatorial construction of homogeneous representations of simply-laced

2010 *Mathematics Subject Classification.* 16G10, 17B37, 05E10.

Key words and phrases. Categorification, Minuscule representations, Quantum groups, Quiver Hecke algebras.

The research was supported by the National Research Foundation of Korea (NRF) Grant funded by the Korean Government (NRF-2017R1A1A1A05001058).

type and investigated a connection with *fully commutative elements* including dominant minuscule elements. The homogeneous representations of type A_n and D_n are classified and enumerated in [5, 6] by studying fully commutative elements.

For simply-laced ADE type, essential parts for the main theorem come from the results of [17]. Let $b \in B(\Lambda_i)$, $\xi = \text{wt}(b)$ and $\mathcal{P}(b)$ the set of all paths from b_{Λ_i} to b on the crystal $B(\Lambda_i)$ (see (2.3)). We show that $\mathcal{P}(b)$ is in 1-1 correspondence to the set of reduced expressions of the minuscule element $w(b)$, which says that $\mathcal{P}(b)$ has a homogeneous $R^{\Lambda_i}(\xi)$ -module structure by the result of [17] (see Theorem 2.9). For type C_n , since a simple $R^{\Lambda_n}(\xi)$ -module is 1-dimensional for all weights ξ , we easily obtain that the algebra $R^{\Lambda_n}(\xi)$ is trivial (see Proposition 2.11). The main novelty of this paper lies in the case of type B_n . Note that, since B_n is not simply-laced, we cannot use the result of [17] directly. In type B_n , we show that there is a 1-1 correspondence between the set \mathfrak{P}_n of all strict partitions $\lambda = (\lambda_1, \lambda_2, \dots)$ with $\lambda_1 \leq n$ and the Weyl group orbit $W \cdot \Lambda_1$. We then give a homogeneous $R^{\Lambda_1}(\xi)$ -module structure on the set $\text{ST}(\lambda)$ of standard tableaux of shape λ using combinatorics of shifted Young tableaux. Thus, we obtain a \mathbf{k} -basis of $R^{\Lambda_1}(\xi)$ indexed by the set of all pairs of shifted Young tableaux (see Theorem 3.4). Remark 3.3 tells us that shifted Young tableaux also can be regarded as paths on the crystal $B(\Lambda_1)$. As a corollary, we show that there exist algebra isomorphisms $R_{B_n}^{\Lambda_1}(m) \simeq R_{D_{n+1}}^{\Lambda_1}(m) \simeq R_{D_{n+1}}^{\Lambda_2}(m)$, where $R_X^{\Lambda_i}(m)$ is the cyclotomic quiver Hecke algebra of type X (see Corollary 3.5).

This paper is organized as follows: In Section 1, we review briefly minuscule representations of quantum groups of finite types. In Section 2, we review cyclotomic quiver Hecke algebras and show an explicit basis of $R^{\Lambda_i}(\xi)$ corresponding to minuscule representations for finite types ADE and C . In Section 3, we construct simple R^{Λ_1} -module of type B_n using combinatorics of shifted Young tableaux and show an explicit basis of $R^{\Lambda_1}(\xi)$.

Acknowledgments. The author would like to thank Myungho Kim for fruitful discussions. The author also would like to thank the anonymous reviewers for their valuable comments and suggestions.

1. MINUSCULE REPRESENTATIONS

1.1. Quantum groups. Let $I = \{1, 2, \dots, n\}$. A quintuple $(A, P, \Pi, P^\vee, \Pi^\vee)$ is called a *Cartan datum* if it consists of

- (a) a generalized *Cartan matrix* $A = (a_{ij})_{i,j \in I}$,
- (b) a free abelian group P , called the *weight lattice*,
- (c) $\Pi = \{\alpha_i \mid i \in I\} \subset P$, called the set of *simple roots*,
- (d) $P^\vee = \text{Hom}_{\mathbb{Z}}(P, \mathbb{Z})$, called the *coweight lattice*,

(e) $\Pi^\vee = \{h_i \in \mathcal{P}^\vee \mid i \in I\}$, called the set of *simple coroots*

satisfying the following:

- (i) $\langle h_i, \alpha_j \rangle = a_{ij}$ for $i, j \in I$,
- (ii) Π is linearly independent over \mathbb{Q} ,
- (iii) for each $i \in I$, there exists $\Lambda_i \in \mathcal{P}$, called the *fundamental weight*, such that $\langle h_j, \Lambda_i \rangle = \delta_{j,i}$ for all $j \in I$.

There is a nondegenerate symmetric bilinear form (\cdot, \cdot) on \mathcal{P} satisfying

$$(\alpha_i, \alpha_j) = d_i a_{ij} \quad \text{and} \quad \langle h_i, \lambda \rangle = \frac{2(\alpha_i, \lambda)}{(\alpha_i, \alpha_i)},$$

where d_i 's are relatively prime integers such that $\text{diag}(d_i \mid i \in I) \cdot \mathbf{A}$ is symmetric. Set $\mathcal{Q} := \bigoplus_{i \in I} \mathbb{Z} \alpha_i$ and $\mathcal{Q}_+ := \sum_{i \in I} \mathbb{Z}_{\geq 0} \alpha_i$ and define $\text{ht}(\beta) = \sum_{i \in I} k_i$ for $\beta = \sum_{i \in I} k_i \alpha_i \in \mathcal{Q}_+$. We denote by $\mathcal{P}_+ := \{\lambda \in \mathcal{P} \mid \langle h_i, \lambda \rangle \geq 0 \text{ for all } i \in I\}$ the set of *dominant integral weights*.

Let $U_q(\mathfrak{g})$ be the *quantum group* associated with the Cartan datum $(\mathbf{A}, \mathcal{P}, \mathcal{P}^\vee, \Pi, \Pi^\vee)$, which is a $\mathbb{Q}(q)$ -algebra generated by f_i, e_i ($i \in I$) and q^h ($h \in \mathcal{P}^\vee$) with certain defining relations (see [8, Chater 3] for details). For $\lambda \in \mathcal{P}_+$, we set $V(\lambda)$ to be the irreducible highest weight $U_q(\mathfrak{g})$ -module with highest weight λ and denote by $B(\lambda)$ its crystal (see [11, 12], [8, Chapter 4] for the definition). Let $\mathbb{A} = \mathbb{Z}[q, q^{-1}]$. We denote by $U_{\mathbb{A}}^-(\mathfrak{g})$ and $V_{\mathbb{A}}(\lambda)$ the \mathbb{A} -lattices of $U_q^-(\mathfrak{g})$ and $V_q(\lambda)$ respectively.


1.2. Minuscule representations. We write Δ for the set of all *roots* associated with \mathbf{A} , and denote by W the *Weyl group*, which is the subgroup of $\text{Aut}(\mathcal{P})$ generated by $s_i(\lambda) := \lambda - \langle h_i, \lambda \rangle \alpha_i$ for $i \in I$. When $w \in W$ is written as $w = s_{i_1} \dots s_{i_\ell}$ (ℓ minimal), we call the expression *reduced* and write $\ell(w) = \ell$. An element w is *fully commutative* if for every pair of noncommuting generators s_i and s_j there is no reduced expression for w containing a subword of length m of the form $s_i s_j s_i s_j \dots$, where m denotes the order of $s_i s_j$ in W .


For $\lambda \in \mathcal{P}_+$, an element $w \in W$ is λ -*minuscule* if there exists a reduced expression $w = s_{i_1} \dots s_{i_\ell}$ such that


$$(1.1) \quad \langle h_{i_k}, s_{i_{k+1}} s_{i_{k+2}} \dots s_{i_\ell} \lambda \rangle = 1 \quad \text{for } k = 1, \dots, \ell.$$


A weight $\lambda \in \mathcal{P}$ is called *minuscule* if $\langle \alpha^\vee, \lambda \rangle \in \{0, \pm 1\}$ for all $\alpha \in \Delta$, where α^\vee is the coroot corresponding to α . Irreducible highest weight modules corresponding to the dominant minuscule weights are called *minuscule representations*. In the following Dynkin diagrams,


(1.2)


$A_n :$ 

 $C_n :$ 

 $E_6 :$ 

$B_n :$ 

 $D_n :$ 

 $E_7 :$ 

For a dominant minuscule weight Λ_i , the set of vertices of the crystal $B(\Lambda_i)$ can be identified with the Weyl group orbit $W \cdot \Lambda_i$. Set $\mathbb{B}(\Lambda_i) := W \cdot \Lambda_i$ and define

$$\begin{aligned}\varepsilon_i(\mu) &:= \max\{0, -\langle h_i, \mu \rangle\}, & \varphi_i(\mu) &:= \max\{0, \langle h_i, \mu \rangle\}, \\ \tilde{f}_i(\mu) &:= \begin{cases} \mu - \alpha_i & \text{if } \langle h_i, \mu \rangle = 1, \\ 0 & \text{otherwise,} \end{cases} & \tilde{e}_i(\mu) &:= \begin{cases} \mu + \alpha_i & \text{if } \langle h_i, \mu \rangle = -1, \\ 0 & \text{otherwise,} \end{cases}\end{aligned}$$

2. CYCLOTOMIC QUIVER HECKE ALGEBRAS AND MINUSCULE WEIGHTS

$$Q_{i,j}(u,v) = \delta_{i,j} \sum_{\substack{(p,q) \in \mathbb{Z}_{\geq 0}^2 \\ (\alpha_i, \alpha_i)p + (\alpha_j, \alpha_j)q = -2(\alpha_i, \alpha_j)}} t_{i,j;p,q} u^p v^q \in \mathbf{k}[u,v]$$

with $t_{i,j;p,q} \in \mathbf{k}$, $t_{i,j;p,q} = t_{j,i;q,p}$ and $t_{i,j;-a_{ij},0} \in \mathbf{k}^\times$. Note that $Q_{i,j}(u,v) = Q_{j,i}(v,u)$ for $i, j \in I$. Let $\mathfrak{S}_n = \langle s_1, \dots, s_{n-1} \rangle$ be the symmetric group on n letters with the action of \mathfrak{S}_n on I^n by place permutation. For $\beta \in \mathbf{Q}_+$ with $\text{ht}(\beta) = n$, we set

$$I^\beta = \{\nu = (\nu_1, \dots, \nu_n) \in I^n \mid \alpha_{\nu_1} + \dots + \alpha_{\nu_n} = \beta\}.$$

Definition 2.1. For $\beta \in \mathbf{Q}_+$ with $\text{ht}(\beta) = n$, the *quiver Hecke algebra* $R(\beta)$ associated with $\{Q_{i,j}\}_{i,j \in I}$ is the \mathbf{k} -algebra generated by $\{e(\nu)\}_{\nu \in I^\beta}$, $\{x_k\}_{1 \leq k \leq n}$, $\{\tau_m\}_{1 \leq m \leq n-1}$ satisfying the following defining relations:

$$e(\nu)e(\nu') = \delta_{\nu,\nu'}e(\nu), \quad \sum_{\nu \in I^\beta} e(\nu) = 1,$$

$$\begin{aligned}
x_k x_m &= x_m x_k, \quad x_k e(\nu) = e(\nu) x_k, \\
\tau_m e(\nu) &= e(s_m(\nu)) \tau_m, \quad \tau_k \tau_m = \tau_m \tau_k \quad \text{if } |k - m| > 1, \\
\tau_k^2 e(\nu) &= Q_{\nu_k, \nu_{k+1}}(x_k, x_{k+1}) e(\nu), \\
(\tau_k x_m - x_{s_k(m)} \tau_k) e(\nu) &= \begin{cases} -e(\nu) & \text{if } m = k, \nu_k = \nu_{k+1}, \\ e(\nu) & \text{if } m = k + 1, \nu_k = \nu_{k+1}, \\ 0 & \text{otherwise,} \end{cases} \\
(\tau_{k+1} \tau_k \tau_{k+1} - \tau_k \tau_{k+1} \tau_k) e(\nu) &= \begin{cases} \frac{Q_{\nu_k, \nu_{k+1}}(x_k, x_{k+1}) - Q_{\nu_k, \nu_{k+1}}(x_{k+2}, x_{k+1})}{x_k - x_{k+2}} e(\nu) & \text{if } \nu_k = \nu_{k+2}, \\ 0 & \text{otherwise.} \end{cases}
\end{aligned}$$

Note that $R(\beta)$ has the \mathbb{Z} -grading defined by

$$\deg e(\nu) = 0, \quad \deg x_k e(\nu) = (\alpha_{\nu_k}, \alpha_{\nu_k}), \quad \deg \tau_l e(\nu) = -(\alpha_{\nu_l}, \alpha_{\nu_{l+1}}).$$

For $n \in \mathbb{Z}_{\geq 0}$, we set $R(n) := \bigoplus_{\beta \in \mathbf{Q}_+, \text{ht}(\beta)=n} R(\beta)$. For $w \in \mathfrak{S}_n$, we fix a preferred reduced expression $w = s_{i_1} \cdots s_{i_l}$ and define $\tau_w := \tau_{s_{i_1}} \cdots \tau_{s_{i_l}}$. Note that τ_w depends on the choice of reduced expression w unless w is fully commutative.

Throughout the paper, we assume that

$$(2.1) \quad Q_{i,j}(u, v) = 1 \quad \text{for any } i, j \in I \text{ with } (\alpha_i, \alpha_j) = 0.$$

If every element of the base field \mathbf{k} has a square root, then one can show that any quiver Hecke algebra associated with $Q_{i,j}(u, v)$ is isomorphic to the quiver Hecke algebra associated with

$$Q'_{i,j}(u, v) = \begin{cases} 1 & \text{if } (\alpha_i, \alpha_j) = 0, \\ Q_{i,j}(u, v) & \text{otherwise,} \end{cases}$$

under the isomorphism given by $e(\nu) \mapsto e(\nu)$, $x_k e(\nu) \mapsto x_k e(\nu)$ and $\tau_k e(\nu) \mapsto c_{\nu_k, \nu_{k+1}} \tau_k e(\nu)$, where

$$c_{i,j} = \begin{cases} (Q_{i,j}(u, v))^{1/2} & \text{if } (\alpha_i, \alpha_j) = 0, \\ 1 & \text{otherwise.} \end{cases}$$

(See [2, Lemma 3.2] for the isomorphism).

For a \mathbb{Z} -graded algebra A , we denote by $A\text{-Mod}$ the category of graded left A -modules, and write $A\text{-proj}$ (resp. $A\text{-gmod}$) for the full subcategory of $A\text{-Mod}$ consisting of finitely generated projective (resp. finite-dimensional) graded A -modules. We set $R\text{-proj} := \bigoplus_{\beta \in \mathbf{Q}_+} R(\beta)\text{-proj}$ and $R\text{-gmod} := \bigoplus_{\beta \in \mathbf{Q}_+} R(\beta)\text{-gmod}$.

For $M \in R(\beta)\text{-Mod}$ and $N \in R(\gamma)\text{-Mod}$, we define

$$M \circ N = R(\beta + \gamma)e(\beta, \gamma) \otimes_{R(\beta) \otimes R(\gamma)} (M \boxtimes N),$$

where $e(\beta, \gamma) = \sum_{\nu_1 \in I^\beta, \nu_2 \in I^\gamma} e(\nu_1 * \nu_2)$. Here $\nu_1 * \nu_2$ is the concatenation of ν_1 and ν_2 . Then, the Grothendieck groups $[R\text{-proj}]$ and $[R\text{-gmod}]$ admit \mathbb{A} -algebra structures with the multiplication given by the functor \circ .

Theorem 2.2. ([15, 16, 18]) *There exist \mathbb{A} -algebra isomorphisms*

$$U_{\mathbb{A}}^-(\mathfrak{g}) \xrightarrow{\sim} [R\text{-proj}], \quad U_{\mathbb{A}}^-(\mathfrak{g})^\vee \xrightarrow{\sim} [R\text{-gmod}].$$

For $\Lambda \in \mathbf{P}_+$ and $\beta \in \mathbf{Q}_+$, let I_β^Λ be the two-sided ideal of $R(\beta)$ generated by the elements $\{x^{(h_{\nu_{\text{ht}(\beta)}, \Lambda})} e(\nu) \mid \nu \in I^\beta\}$. The *cyclotomic quiver Hecke algebra* $R^\Lambda(\Lambda - \beta)$ is defined to be the quotient algebra

$$R^\Lambda(\Lambda - \beta) := R(\beta)/I_\beta^\Lambda.$$

For $m \in \mathbb{Z}_{\geq 0}$, we set $R^\Lambda(m) := \bigoplus_{\beta \in \mathbf{Q}_+, \text{ht}(\beta)=m} R^\Lambda(\Lambda - \beta)$. When \mathbf{A} is of type X , we sometimes write $R_X(m)$ and $R_X^\Lambda(m)$ to emphasize the type X .

For $M \in R^\Lambda(\mu)\text{-Mod}$, we set $\text{wt}(M) = \mu$ and define functors F_i^Λ and E_i^Λ by

$$F_i^\Lambda M := R^\Lambda(\mu - \alpha_i)e(\alpha_i, \beta) \otimes_{R^\Lambda(\mu)} M \quad \text{and} \quad E_i^\Lambda M := e(\alpha_i, \beta - \alpha_i)M,$$

where $\beta = \Lambda - \mu$. We denote by Ind_m^{m+1} and Res_m^{m+1} the induction and restriction functors via the embedding $R^\Lambda(m) \hookrightarrow R^\Lambda(m+1)$. Note that $\text{Ind}_m^{m+1} = \sum_{i \in I} F_i^\Lambda$ and $\text{Res}_m^{m+1} = \sum_{i \in I} E_i^\Lambda$. The functors F_i^Λ and E_i^Λ give a $U_{\mathbb{A}}(\mathfrak{g})$ -module structure on $[R^\Lambda\text{-proj}]$ and $[R^\Lambda\text{-gmod}]$.

Theorem 2.3. ([10]) *There exist $U_{\mathbb{A}}(\mathfrak{g})$ -module isomorphisms*

$$[R^\Lambda\text{-proj}] \xrightarrow{\sim} V_{\mathbb{A}}(\Lambda), \quad [R^\Lambda\text{-gmod}] \xrightarrow{\sim} V_{\mathbb{A}}(\Lambda)^\vee.$$

If ξ is an extremal weight of $V(\Lambda)$, then we know the representation type of $R^\Lambda(\xi)$.

Proposition 2.4. ([1, Corollary 4.7], [14, Lemma 1.11]) *Let $\Lambda \in \mathbf{P}_+$ and $\xi = w\Lambda$ for $w \in \mathbf{W}$. Then the algebra $R^\Lambda(\xi)$ is simple.*

Proposition 2.5. ([15, Corollary 3.19]) *Every irreducible R -module is absolutely irreducible.*

We now assume that \mathbf{A} is of simply laced type. Then the construction for homogeneous representations was given in [17]. We denote by G_β the graph with the set of vertices I^β , and with $\nu, \mu \in I^\beta$ connected by an edge if and only if there exists k such that $\mu = s_k \nu$ and

$a_{\nu_k, \nu_{k+1}} = 0$. Let C be a connected component of G_β . We say that C is *homogeneous* if for each $\nu \in C$ the following condition holds:

$$(2.2) \quad \begin{aligned} &\text{if } \nu_r = \nu_s \text{ for some } r < s \text{ then there exist } t, u \text{ with } r < t < u < s \\ &\text{such that } a_{\nu_r, \nu_t} = -1 \text{ and } a_{\nu_u, \nu_s} = -1. \end{aligned}$$

It was proved in [17, Theorem 3.4] that C has a simple $R(\beta)$ -module structure.

2.2. Minuscul representations of symmetric type ADE . Let $V(\Lambda_i)$ be a minuscule representation and $B(\Lambda_i)$ its crystal. Let $b \in B(\Lambda_i)$, $\xi = \text{wt}(b)$ and $\ell = \text{ht}(\Lambda_i - \xi)$. We set

$$(2.3) \quad \mathcal{P}(b) := \{(i_1, \dots, i_\ell) \in I^\ell \mid \tilde{f}_{i_1} \cdots \tilde{f}_{i_\ell} b_{\Lambda_i} = b\},$$

where b_{Λ_i} is the highest weight vector of $B(\Lambda_i)$. From a viewpoint of the crystal graph $B(\Lambda_i)$, an element of $\mathcal{P}(b)$ can be thought as a sequence of arrows from b_{Λ_i} to b on the graph $B(\Lambda_i)$. In this sense, an element of $\mathcal{P}(b)$ is called a *path* from b_{Λ_i} to b on the crystal $B(\Lambda_i)$.

From now on, we suppose that $X \in \{A_n, D_n, E_6, E_7\}$ and $i \in \mathbf{I}_X$.

Lemma 2.6. *Let $(i_1, \dots, i_\ell) \in \mathcal{P}(b)$ and set $w = s_{i_1} s_{i_2} \cdots s_{i_\ell} \in \mathbf{W}$. Then $s_{i_1} s_{i_2} \cdots s_{i_\ell}$ is a reduced expression of w .*

Proof. As $(i_1, \dots, i_\ell) \in \mathcal{P}(b)$, we have $\text{wt}(b) = \Lambda_i - \sum_{k=1}^\ell \alpha_{i_k}$ by the definition of $\mathcal{P}(b)$. On the other hand, since $B(\Lambda_i)$ is the crystal of the minuscule representation $V(\Lambda_i)$, we have $\text{wt}(b) = w\Lambda_i$, which says that

$$\Lambda_i - w\Lambda_i = \sum_{k=1}^\ell \alpha_{i_k}.$$

Thus we have

$$(2.4) \quad \text{ht}(\Lambda_i - w\Lambda_i) = \ell.$$

We now set $m := \ell(w)$ and take a reduced expression $s_{j_1} \cdots s_{j_m}$ of w . For $k = 1, \dots, m$, define $w_k := s_{j_k} s_{j_{k+1}} \cdots s_{j_m}$ and $w_{m+1} := \text{id}$. Then we have

$$\begin{aligned} \Lambda_i - w\Lambda_i &= \sum_{k=1}^m (w_{k+1}\Lambda_i - w_k\Lambda_i) = \sum_{k=1}^m (w_{k+1}\Lambda_i - s_{j_k}(w_{k+1}\Lambda_i)) \\ &= \sum_{k=1}^m \langle h_{j_k}, w_{k+1}\Lambda_i \rangle \alpha_{j_k}. \end{aligned}$$

Since Λ_i is minuscule, we have $\langle h_{j_k}, w_{k+1}\Lambda_i \rangle = \langle w_{k+1}^{-1} h_{j_k}, \Lambda_i \rangle \in \{0, \pm 1\}$ for any k , which implies that

$$(2.5) \quad \text{ht}(\Lambda_i - w\Lambda_i) \leq \sum_{k=1}^m |\langle h_{j_k}, w_{k+1}\Lambda_i \rangle| \leq m = \ell(w).$$

Combining (2.5) with (2.4), we obtain

$$\ell \leq \ell(w),$$

which tells us that $s_{i_1}s_{i_2}\cdots s_{i_\ell}$ is a reduced expression of w . \square

Lemma 2.7. *For $(i_1, \dots, i_\ell), (j_1, \dots, j_\ell) \in \mathcal{P}(b)$, we have*

$$s_{i_1} \cdots s_{i_\ell} = s_{j_1} \cdots s_{j_\ell}.$$

Proof. Let $(i_1, \dots, i_\ell) \in \mathcal{P}(b)$ and set $w := s_{i_1} \cdots s_{i_\ell}$. Thanks to Lemma 2.6, $s_{i_1} \cdots s_{i_\ell}$ is a reduced expression of w . Since $B(\Lambda_i)$ is minuscule, $s_i b' = \tilde{f}_i b'$ for any $b' \in B(\Lambda_i)$ with $\varphi_i(b') > 0$. Thus, the reduced expression $w = s_{i_1} \cdots s_{i_\ell}$ satisfies (1.1), i.e. w is Λ_i -minuscule. It was shown in [19, Proposition 2.1] that every reduced expression of w satisfies (1.1). We take $j \in I$ with $j \neq i$.

We suppose that $\ell(ws_j) < \ell(w)$. Then one can write a reduced expression of $w = s_{t_1} \cdots s_{t_{\ell-1}} s_j$ for some $t_1, \dots, t_\ell \in I$. Since $\langle h_j, \Lambda_i \rangle = 0$, $w = s_{t_1} \cdots s_{t_{\ell-1}} s_j$ is not Λ_i -minuscule which is a contradiction. Thus, we have

$$(2.6) \quad \ell(ws_j) > \ell(w) \text{ for all } j \in I \text{ with } j \neq i.$$

We now consider an element $(j_1, \dots, j_\ell) \in \mathcal{P}(b)$. Set $u := s_{j_1} \cdots s_{j_\ell}$. By the same argument as above,

$$(2.7) \quad \ell(us_j) > \ell(u) \text{ for all } j \in I \text{ with } j \neq i$$

Considering the realization $\mathbb{B}(\Lambda_i)$ given in Section 1.2, we have $w \cdot \Lambda_i = u \cdot \Lambda_i = \text{wt}(b)$, which tells us that

$$u^{-1}w \cdot \Lambda_i = \Lambda_i.$$

Hence $u^{-1}w$ is contained in the parabolic subgroup W_J of W generated by s_j for $j \in J := I \setminus \{i\}$. We set $X_J := \{w \in W \mid \ell(ws_j) > \ell(w) \text{ for all } j \in J\}$. Note that $u, w \in X_J$ by (2.6) and (2.7). It is well-known that $\ell(xw) = \ell(x) + \ell(w)$ for $x \in X_J$ and $w \in W_J$ (see [7, Proposition 2.1.1] for example).

Suppose that $u^{-1}w \neq \text{id}$. Then there exists $j \in I \setminus \{i\}$ such that $\ell(u^{-1}ws_j) = \ell(u^{-1}w) - 1$. It follows from (2.7) that

$$\begin{aligned} \ell(ws_j) &= \ell(u(u^{-1}ws_j)) = \ell(u) + \ell((u^{-1}w)s_j) \\ &= \ell(u) + \ell(u^{-1}w) - 1 = \ell(u(u^{-1}w)) - 1 = \ell(w) - 1, \end{aligned}$$

which is a contradiction to (2.6). The second and forth identities follow from the fact that $u \in X_J$ and $u^{-1}w, (u^{-1}w)s_j \in W_J$. Therefore, we conclude that $u^{-1}w = \text{id}$. \square

We define

$$w(b) := s_{i_1} \cdots s_{i_\ell} \in \mathbf{W},$$

where (i_1, \dots, i_ℓ) is an element of $\mathcal{P}(b)$. Note that it is well-defined by Lemma 2.7.

Lemma 2.8. *For $b \in B(\Lambda_i)$, the set $\mathcal{P}(b)$ is in 1-1 correspondence to the set of all reduced expressions of $w(b)$ via the map $(i_1, \dots, i_\ell) \mapsto s_{i_1} \cdots s_{i_\ell}$.*

Proof. Let $E(b)$ be the set of all reduced expressions of $w(b)$. We consider the map

$$F : \mathcal{P}(b) \longrightarrow E(b), \quad (i_1, \dots, i_\ell) \mapsto s_{i_1} \cdots s_{i_\ell}.$$

Thanks to Lemma 2.6 and Lemma 2.7, the map F is well-defined.

Recall the crystal realization $\mathbb{B}(\Lambda_i)$ given in Section 1.2. Note that, for $\mu \in \mathbb{B}(\Lambda_i)$, $\tilde{f}_i(\mu) = s_i \mu$ if $\langle h_i, \mu \rangle = 1$. Since every reduced expression of $w(b)$ satisfies (1.1) by [19, Proposition 2.1], the map

$$G : E(b) \longrightarrow \mathcal{P}(b), \quad s_{i_1} \cdots s_{i_\ell} \mapsto (i_1, \dots, i_\ell),$$

is well-defined. Then, it is clear that $F \circ G = \text{id}$ and $G \circ F = \text{id}$. \square

Since $w(b)$ is fully commutative [19, Proposition 2.1], Lemma 2.8 says that for $\nu, \mu \in \mathcal{P}(b)$, there exists $w_{\nu, \mu} \in \mathfrak{S}_\ell$ such that $\nu = w_{\nu, \mu} \mu$. We set

$$(2.8) \quad c_{\nu, \mu} := e(\nu) \tau_{w_{\nu, \mu}} e(\mu).$$

Note that $c_{\nu, \mu}$ does not depend on the choice of the reduced expressions of $w_{\nu, \mu}$.

We now set

$$\mathcal{S}(b) := \bigoplus_{\nu \in \mathcal{P}(b)} \mathbf{k} \mathbf{p}(\nu)$$

and define an $R(\beta)$ -module structure on $\mathcal{S}(b)$ by

$$e(\mu) \mathbf{p}(\nu) = \delta_{\nu, \mu} \mathbf{p}(\nu), \quad x_i e(\mu) \mathbf{p}(\nu) = 0, \quad \tau_j e(\mu) \mathbf{p}(\nu) = \begin{cases} \mathbf{p}(s_j \nu), & \text{if } \nu = \mu \text{ and } s_j \nu \in \mathcal{P}(b), \\ 0 & \text{otherwise,} \end{cases}$$

for $\nu \in \mathcal{P}(b)$ and admissible i, j, μ .

Theorem 2.9. *Let $X \in \{A_n, D_n, E_6, E_7\}$ and $i \in \mathbf{I}_X$. Let $b \in B(\Lambda_i)$ and write $\xi = \text{wt}(b)$.*

- (1) $\mathcal{S}(b)$ is a homogeneous simple $R^{\Lambda_i}(\xi)$ -module.
- (2) $R^{\Lambda_i}(\xi)$ has a \mathbf{k} -basis

$$\mathcal{C}(b) := \{c_{\nu, \mu} \mid \nu, \mu \in \mathcal{P}(b)\},$$

where $c_{\nu, \mu}$ is given in (2.8).

Proof. (1) By Lemma 2.8, the set $\mathcal{P}(b)$ can be identify with the set of all reduced expression of $w(b)$. Then it follows from [19, Proposition 2.5] (c.f. [17, Proposition 3.8]) that $\mathcal{P}(b)$ satisfies the condition (2.2). Thus, $\mathcal{P}(b)$ has a homogeneous simple R -module structure which is same as the above actions on $\mathcal{S}(b)$. Moreover, since $\nu_\ell = i$ for any $(\nu_1, \dots, \nu_\ell) \in \mathcal{P}(b)$ and x_ℓ acts as zero, $\mathcal{S}(b)$ is an $R^{\Lambda_i}(\xi)$ -module.

(2) Since every element of $B(\Lambda_i)$ is extremal, $R^{\Lambda_i}(\xi)$ is simple by Proposition 2.4. Thus, by Proposition 2.5, $R^{\Lambda_i}(\xi)$ is isomorphic to the endomorphism ring of $\mathcal{S}(b)$, which implies that $\mathcal{C}(b)$ is a basis of $R^{\Lambda_i}(\xi)$. \square

Remark 2.10. For type A_n , the simple module $\mathcal{S}(b)$ can be realized as the set of standard tableaux of a usual Young diagram (see [4, Section 5]). Thus, the algebra $R^{\Lambda_i}(m)$ has a \mathbf{k} -basis indexed by all pairs of standard tableaux. In Section 3, we shall give a B_n -type analogue using shifted Young tableaux.

2.3. Minuscule representations of type C_n . In this subsection, we assume that $X = C_n$ and consider the dominant minuscule weight Λ_n . Then the crystal $B(\Lambda_n)$ is given as follows:

$$\boxed{\overline{n}} \xleftarrow{n} \dots \xleftarrow{3} \boxed{\overline{2}} \xleftarrow{2} \boxed{\overline{1}} \xleftarrow{1} \boxed{1} \xleftarrow{2} \boxed{2} \xleftarrow{3} \dots \xleftarrow{n} \boxed{n}$$

For $b \in B(\Lambda_n)$, if we write $b = \tilde{f}_{i_1} \dots \tilde{f}_{i_\ell} b_{\Lambda_n}$, then we set

$$\nu_b := (i_1, \dots, i_\ell).$$

We set $\mathcal{S}(b) := \mathbf{kp}(\nu_b)$ and define an R -module structure by

$$e(\mu)\mathbf{p}(\nu_b) = \delta_{\nu_b, \mu}\mathbf{p}(\nu_b), \quad x_i e(\mu)\mathbf{p}(\nu_b) = \tau_j e(\mu)\mathbf{p}(\nu_b) = 0,$$

for admissible i, j, μ . Then one can prove that $\mathcal{S}(b)$ is a 1-dimensional R^{Λ_n} -module. Note that this construction of $\mathcal{S}(b)$ is the same as the one for type ADE . Proposition 2.4 and Proposition 2.5 imply the following immediately.

Proposition 2.11. *Let $X = C_n$, $b \in B(\Lambda_n)$, and write $\xi = \text{wt}(b)$. Then $R^{\Lambda_n}(\xi)$ is isomorphic to \mathbf{k} .*

3. SIMPLE R^{Λ_1} -MODULES OF TYPE B_n

3.1. Shifted Young tableaux. For a *strict partition* $\lambda = (\lambda_1 > \lambda_2 > \dots > \lambda_\ell > 0)$ with $N = \sum_{i=1}^\ell \lambda_i$, we write $\lambda \vdash N$ and set $|\lambda| := N$. We identify strict partitions with shifted Young diagrams and depict shifted Young diagrams using English convention. We write $b = (i, j) \in \lambda$ if there exists a box in the i th row and the j th column. By an *addable* (resp. a *removable*) box $b \in \lambda$, we mean a box which can be added to (resp. removed from) λ to

obtain a valid shifted Young diagram $\lambda \swarrow b$ (resp. $\lambda \nearrow b$). For $n \in \mathbb{Z}_{>0}$, let \mathfrak{P}_n be the set of all strict partitions $(\lambda_1, \lambda_2, \dots)$ such that $\lambda_1 \leq n$. It is easy to check that $|\mathfrak{P}_0| = 1$ and $|\mathfrak{P}_n| = 2|\mathfrak{P}_{n-1}|$ for $n \geq 1$, which say that

$$(3.1) \quad |\mathfrak{P}_n| = 2^n \quad \text{for } n \geq 1.$$

A *standard tableau* T of shape $\lambda \vdash n$ is a filling of boxes of λ with $\{1, 2, \dots, n\}$ such that (i) each entry is used exactly once, (ii) the entries in rows and columns increase from left to right and top to bottom, respectively. Let $\text{ST}(\lambda)$ be the set of all standard tableaux of shape λ . For example, the following is a standard tableaux of shape $\lambda = (7, 4)$.

1	2	4	5	7	8	11
	3	6	9	10		

Let $\lambda \vdash n$ be a shifted Young diagram. We define the residue of $(i, j) \in \lambda$ as follows:

$$(3.2) \quad \text{res}(i, j) := j - i + 1.$$

For example, the residues for $\lambda = (9, 6, 3, 1)$ are as follows. In this example, $\text{res}(2, 6) = 5$.

1	2	3	4	5	6	7	8	9
	1	2	3	4	5	6		
		1	2	3				
			1					

Remark 3.1. The above residues can be viewed as the limit of the residues given in [1] as ℓ goes to infinity.

For $T \in \text{ST}(\lambda)$, we define the *residue sequence* of T by

$$\text{res}(T) = (\text{res}(i_n, j_n), \dots, \text{res}(i_2, j_2), \text{res}(i_1, j_1)) \in I^n,$$

where $(i_k, j_k) \in \lambda$ is the box filled with k in T .

3.2. Irreducible representations. For $\lambda \in \mathfrak{P}_n$, we set

$$\text{wt}(\lambda) := \Lambda_1 - \sum_{b \in \lambda} \alpha_{\text{res}(b)} \in \mathcal{P}.$$

Recall the crystal realization $\mathbb{B}(\Lambda_i)$ given in Section 1.2.

Lemma 3.2. *The map $\lambda \mapsto \text{wt}(\lambda)$ gives 1-1 correspondence between \mathfrak{P}_n and $\mathbb{B}(\Lambda_1)$.*

Proof. Let $f : \mathfrak{P}_n \rightarrow \mathbf{P}$ be the map defined by

$$f(\lambda) = \text{wt}(\lambda) \quad \text{for } \lambda \in \mathfrak{P}_n.$$

Let $\lambda \in \mathfrak{P}_n$. We first show that $f(\lambda) \in \mathbf{W} \cdot \Lambda_1$ by induction argument on $|\lambda|$. Since $\text{wt}(\emptyset) = \Lambda_1 \in \mathbf{W} \cdot \Lambda_1$, we assume that $|\lambda| > 0$. Thus we can write $\lambda = (\lambda_1 > \lambda_2 > \cdots > \lambda_\ell > 0) \in \mathfrak{P}_n$, and set

$$\beta_k := \sum_{t=1}^k \alpha_t \quad \text{for } k = 1, \dots, n.$$

Then we have $\text{wt}(\lambda) = \Lambda_1 - \sum_{k=1}^\ell \beta_{\lambda_k}$ and

$$\langle h_i, \beta_{\lambda_j} \rangle = \begin{cases} 2 & \text{if } i = \lambda_j = 1, \\ 1 & \text{if } i = \lambda_j > 1, \\ -1 & \text{if } i = \lambda_j + 1, \\ 0 & \text{otherwise.} \end{cases}$$

Let $b := (\ell, \lambda_\ell + \ell - 1) \in \lambda$ and $\mu := \lambda \nearrow b$. By the induction hypothesis, we have $\text{wt}(\mu) \in \mathbf{W} \cdot \Lambda_1$. Setting $i := \text{res}(b)$, we obtain

$$\langle h_i, \text{wt}(\mu) \rangle = \langle h_i, \Lambda_1 \rangle - \sum_{k=1}^\ell \langle h_i, \beta_{\lambda_k} \rangle + \langle h_i, \alpha_i \rangle = 1,$$

which tells us that

$$\text{wt}(\lambda) = \text{wt}(\mu) - \alpha_i = s_i(\text{wt}(\mu)) \in \mathbf{W} \cdot \Lambda_1.$$

Thus, we have that $\text{Im} f \subset \mathbf{W} \cdot \Lambda_1$.

We now show that f is injective. Let $\lambda = (\lambda_1, \lambda_2, \dots), \mu = (\mu_1, \mu_2, \dots) \in \mathfrak{P}_n$, and assume that $\lambda \neq \mu$, i.e., there exists k such that $\lambda_i = \mu_i$ for $i < k$ and $\lambda_k \neq \mu_k$. Then by the definition, we have

$$\max\{\lambda_k, \mu_k\} \in \text{supp}(\beta_{\lambda_k} - \beta_{\mu_k}) \quad \text{and} \quad \max\{\lambda_k, \mu_k\} \notin \text{supp}(\beta_{\lambda_t} - \beta_{\mu_t}) \text{ for } t > k,$$

where $\text{supp}(\beta) := \{i \in I \mid b_i \neq 0\}$ for $\beta = \sum_{i \in I} b_i \alpha_i \in \mathbf{Q}$. This tells us that

$$\text{wt}(\lambda) - \text{wt}(\mu) = \beta_{\lambda_k} - \beta_{\mu_k} + \sum_{t>k} (\beta_{\lambda_t} - \beta_{\mu_t}) \neq 0.$$

Hence f is injective.

Moreover, if we denote by P the subgroup of \mathbf{W} generated by s_2, \dots, s_n , then by (3.1), we have

$$|\mathfrak{P}_n| = 2^n = \frac{2^n n!}{n!} = \left| \frac{\mathbf{W}}{P} \right| = |\mathbf{W} \cdot \Lambda_1|.$$

Therefore, the map f gives a bijection between \mathfrak{P}_n and $\mathbf{W} \cdot \Lambda_1$. □

For $b \in B(\Lambda_1)$, we denote by λ_b the strict partition corresponding to b under the bijection given in Lemma 3.2. We set

$$\mathcal{S}(b) := \bigoplus_{T \in \text{ST}(\lambda_b)} \mathbf{k}T,$$

and define an $R(\text{wt}(b))$ -module structure on $\mathcal{S}(b)$ by

$$(3.3) \quad \begin{aligned} e(\nu)T &= \delta_{\nu, \text{res}(T)}T, \\ x_i e(\nu)T &= 0, \\ \tau_j e(\nu)T &= \begin{cases} s_j T & \text{if } \nu = \text{res}(T) \text{ and } s_j T \in \text{ST}(\lambda_b), \\ 0 & \text{otherwise,} \end{cases} \end{aligned}$$

for $T \in \text{ST}(\lambda_b)$ and admissible i, j, ν . Here, $s_j T$ is the tableau obtained from T by exchanging the entries j and $j+1$. For $k \in \mathbb{Z}_{\geq 0}$, we simply write $\mathcal{S}(k)$ for the simple module $\mathcal{S}(b)$ where b is the element of $B(\Lambda_1)$ such that $\lambda_b = (k)$.

Remark 3.3. One can describe the $U_q(B_n)$ -crystal structure on \mathfrak{P}_n directly via the 1-1 correspondence given in Lemma 3.2. In this description, the crystal operators \tilde{f}_i and \tilde{e}_i are defined as follows: for $i \in I$ and $\lambda \in \mathfrak{P}_n$,

$$\begin{aligned} \tilde{f}_i \lambda &= \begin{cases} \lambda \swarrow b & \text{if } b \text{ is an addable box of } \lambda \text{ with residue } i, \\ 0 & \text{otherwise,} \end{cases} \\ \tilde{e}_i \lambda &= \begin{cases} \lambda \nearrow b & \text{if } b \text{ is a removable box of } \lambda \text{ with residue } i, \\ 0 & \text{otherwise.} \end{cases} \end{aligned}$$

Thus, for $b \in B(\Lambda_1)$, we can identify $\text{ST}(\lambda_b)$ with the set $\mathcal{P}(b)$ of all paths from b_{Λ_1} to b on the crystal $B(\Lambda_1)$. This crystal realization also can be obtained from the B_∞ -Fock space given in [9] by restricting to type B_n . From the viewpoint of crystal, if the formulas (3.3) were written in terms of $\mathcal{P}(b)$, then it is same as the ones for type ADE and C .

For $S, T \in \text{ST}(\lambda_b)$, let $w_{T,S}$ be an element of $\mathfrak{S}_{|\lambda_b|}$ such that $T = w_{T,S}S$, and set

$$(3.4) \quad c_{T,S} := e(\text{res}(T))\tau_{w_{T,S}}e(\text{res}(S)).$$

Theorem 3.4. *Let $X = B_n$, $b \in B(\Lambda_1)$, and write $\xi = \text{wt}(b)$.*

- (1) $\mathcal{S}(b)$ is a homogeneous simple $R^{\Lambda_1}(\xi)$ -module.
- (2) If we write $\lambda_b = (\lambda_1, \dots, \lambda_\ell)$, then

$$\mathcal{S}(b) \simeq \text{hd}(\mathcal{S}(\lambda_\ell) \circ \dots \mathcal{S}(\lambda_2) \circ \mathcal{S}(\lambda_1)),$$

where $\text{hd}(M)$ denotes by the head of a module M .

(3) Let $m = \text{ht}(\Lambda_1 - \xi)$. Then

$$\text{Ind}_m^{m+1} \mathcal{S}(b) \simeq \bigoplus_{i \in I, \tilde{f}_i b \neq 0} \mathcal{S}(\tilde{f}_i b), \quad \text{Res}_{m-1}^m \mathcal{S}(b) \simeq \bigoplus_{i \in I, \tilde{e}_i b \neq 0} \mathcal{S}(\tilde{e}_i b),$$

where Ind_m^{m+1} and Res_{m-1}^m are the induction and restriction functors given in Section 2.1.

(4) For $T, S \in \text{ST}(\lambda_b)$, $c_{T,S}$ does not depend on the choice of the reduced expressions of $w_{T,S}$. Moreover, the algebra $R^{\Lambda_1}(\xi)$ has a \mathbf{k} -basis

$$\mathcal{C}(b) := \{c_{T,S} \mid T, S \in \text{ST}(\lambda_b)\}.$$

Proof. (1) For $i < j$, by the assumption (2.1), we can write

$$Q_{i,j}(u, v) = \begin{cases} \mathbf{a}_{i,j}u^2 + \mathbf{b}_{i,j}v & \text{if } i = 1, j = 2, \\ \mathbf{a}_{i,j}u + \mathbf{b}_{i,j}v & \text{if } j = i + 1 > 2, \\ 1 & \text{otherwise,} \end{cases}$$

for some nonzero $\mathbf{a}_{i,j}, \mathbf{b}_{i,j} \in \mathbf{k}^\times$. Note that, if every element of the base field \mathbf{k} has a square root, then the isomorphism given in [2, Lemma 3.2] allows us to assume that $\mathbf{a}_{i,j} = 1$ and $\mathbf{b}_{i,j} = -1$.

Let $T \in \text{ST}(\lambda_b)$, $N := |\lambda_b|$ and write $\text{res}(T) = (i_1, \dots, i_N)$. Considering the residue pattern (3.2) and combinatorics of standard tableaux, we have

- (i) $i_k \neq i_{k+1}$ for any $k = 1, \dots, N-1$,
- (ii) $s_k T$ is standard if and only if $a_{i_k, i_{k+1}} = 0$,
- (iii) if $i_k = i_{k+2}$, then $(i_k, i_{k+1}, i_{k+2}) = (1, 2, 1)$.

We shall only check that the defining relations for $(\tau_k x_m - x_{s_k(m)} \tau_k)$, τ_k^2 and $\tau_{k+1} \tau_k \tau_{k+1} - \tau_k \tau_{k+1} \tau_k$ since it is straightforward to check other relations.

By (i), it suffices to show that $(\tau_k x_m - x_{s_k(m)} \tau_k) e(\nu) = 0$, which always holds by the definition of $\mathcal{S}(b)$.

If $a_{i_k, i_{k+1}} \neq 0$, then $\deg Q_{i_k, i_{k+1}}(u, v) > 0$ and $s_k T \notin \text{ST}(\lambda_b)$ by (ii). Thus we have

$$\tau_k^2 e(\nu) T = 0 = Q_{i_k, i_{k+1}}(x_k, x_{k+1}) e(\nu) T.$$

If $a_{i_k, i_{k+1}} = 0$, then $Q_{i_k, i_{k+1}}(u, v) = 1$ and

$$\tau_k^2 e(\nu) T = \delta_{\nu, \text{res}(T)} T = Q_{i_k, i_{k+1}}(x_k, x_{k+1}) e(\nu) T$$

by (ii). Thus, the defining relation for $\tau_k^2 e(\nu)$ holds.

We now consider the last case. If $i_k \neq i_{k+2}$, then it is easy to check that $\tau_{k+1} \tau_k \tau_{k+1} e(\nu) T = \tau_k \tau_{k+1} \tau_k e(\nu) T$. Suppose that $i_k = i_{k+2}$. Then (iii) says that $(i_k, i_{k+1}, i_{k+2}) = (1, 2, 1)$. Thus,

by the definition of $\mathcal{S}(b)$, we have

$$(\tau_{k+1}\tau_k\tau_{k+1} - \tau_k\tau_{k+1}\tau_k)e(\nu)T = 0.$$

On the other hand, since $(i_k, i_{k+1}, i_{k+2}) = (1, 2, 1)$, we obtain

$$\frac{Q_{1,2}(x_k, x_{k+1}) - Q_{1,2}(x_{k+2}, x_{k+1})}{x_k - x_{k+2}} = \mathbf{a}_{1,2}(x_k + x_{k+2}),$$

which implies that

$$\frac{Q_{1,2}(x_k, x_{k+1}) - Q_{1,2}(x_{k+2}, x_{k+1})}{x_k - x_{k+2}}e(\nu)T = 0.$$

Therefore, the defining relations for $\tau_{k+1}\tau_k\tau_{k+1} - \tau_k\tau_{k+1}\tau_k$ holds, which completes the proof.

(2) We set $e := e(\nu_{\lambda_\ell} * \dots * \nu_{\lambda_2} * \nu_{\lambda_1})$ where $\nu_k = (k, \dots, 2, 1)$ for $k \in \mathbb{Z}_{\geq 0}$. Considering the combinatorics of shifted Young tableaux, it is easy to see that $e\mathcal{S}(b) = \mathbf{k}T_0$, where T_0 is the *initial tableau*, which is filled with $\{1, \dots, N\}$ from left to right and top to bottom in order. Thus, there exists a surjective homomorphism

$$\mathcal{S}(\lambda_\ell) \otimes \dots \otimes \mathcal{S}(\lambda_2) \otimes \mathcal{S}(\lambda_1) \twoheadrightarrow e\mathcal{S}(b),$$

which yields the surjective homomorphism

$$(3.5) \quad \mathcal{S}(\lambda_\ell) \circ \dots \circ \mathcal{S}(\lambda_2) \circ \mathcal{S}(\lambda_1) \twoheadrightarrow \mathcal{S}(b).$$

On the other hand, by the shuffle lemma [15, Lemma 2.20], we have

$$\dim e(\mathcal{S}(\lambda_\ell) \circ \dots \circ \mathcal{S}(\lambda_2) \circ \mathcal{S}(\lambda_1)) = 1.$$

For any quotient Q of $\mathcal{S}(\lambda_\ell) \circ \dots \circ \mathcal{S}(\lambda_2) \circ \mathcal{S}(\lambda_1)$, the natural surjection

$$\mathcal{S}(\lambda_\ell) \circ \dots \circ \mathcal{S}(\lambda_2) \circ \mathcal{S}(\lambda_1) \twoheadrightarrow Q$$

tells us that $\dim(eQ) = 1$. Thus, we conclude that $\mathcal{S}(\lambda_\ell) \circ \dots \circ \mathcal{S}(\lambda_2) \circ \mathcal{S}(\lambda_1)$ has a unique simple head. Hence, (3.5) implies the desired result.

(3) Note that every element of $B(\Lambda_1)$ is extremal. By Proposition 2.4, $R^{\Lambda_1}(\zeta)$ is simple for any weight ζ with $B(\Lambda_1)_\zeta \neq 0$, which says that $R^{\Lambda_1}(t)$ is semisimple for any $t > 0$. Since $\mathcal{S}(b)$ has a basis indexed by $\mathbf{ST}(\lambda_b)$, we have $E_i^{\Lambda_1}\mathcal{S}(b) \simeq \mathcal{S}(\tilde{e}_i b)$ by considering their characters. Here we set $\mathcal{S}(0) := 0$. Thus, we have $\text{Res}_{m-1}^m \mathcal{S}(b) \simeq \bigoplus_{i \in I, \tilde{e}_i b \neq 0} \mathcal{S}(\tilde{e}_i b)$. The isomorphism for Ind_m^{m+1} follows from the fact that F_i^Λ and E_i^Λ form an adjoint pair [13, Theorem 3.5].

(4) It follows from the proof of (1) that $\tau_{k+1}\tau_k\tau_{k+1} - \tau_k\tau_{k+1}\tau_k$ acts as 0 on $\mathcal{S}(b)$. Thus, for any $T \in \mathcal{S}(b)$, $\tau_w T$ does not depend on the choice of the reduced expressions of $w \in \mathfrak{S}_N$. By Proposition 2.4 and Proposition 2.5, $R^{\Lambda_1}(\xi)$ is isomorphic to the endomorphism ring of $\mathcal{S}(b)$. Therefore, for $T, S \in \mathbf{ST}(\lambda_b)$, $c_{T,S}$ does not depend on the choice of the reduced expressions of $w_{T,S}$ and $\mathcal{C}(b)$ is a basis of $R^{\Lambda_1}(\xi)$. \square

Corollary 3.5. *For $n \in \mathbb{Z}_{\geq 2}$ and $m \in \mathbb{Z}_{\geq 0}$, there exist algebra isomorphisms*

$$R_{B_n}^{\Lambda_1}(m) \simeq R_{D_{n+1}}^{\Lambda_1}(m) \simeq R_{D_{n+1}}^{\Lambda_2}(m).$$

In particular, $\dim R_{B_n}^{\Lambda_1}(m) = \dim R_{D_{n+1}}^{\Lambda_1}(m) = \dim R_{D_{n+1}}^{\Lambda_2}(m) = \sum_{\lambda \in \mathfrak{P}_n, \lambda \vdash m} |\mathrm{ST}(\lambda)|^2$

Proof. It is clear that $R_{D_{n+1}}^{\Lambda_1}(m) \simeq R_{D_{n+1}}^{\Lambda_2}(m)$ by the symmetry of the Dynkin diagram. We shall define a simple $R_{D_{n+1}}^{\Lambda_1}(m)$ -module using shifted Young tableaux. Let $\lambda \vdash m$ be a strict partition. We define

$$\mathrm{res}_D(i, j) := \begin{cases} 1 & \text{if } i = j \text{ and } i \text{ is odd,} \\ 2 & \text{if } i = j \text{ and } i \text{ is even,} \\ j - i + 2 & \text{otherwise,} \end{cases}$$

and $\mathrm{wt}_D(\lambda) := \Lambda_1 - \sum_{b \in \lambda} \alpha_{\mathrm{res}_D(b)}$. For $T \in \mathrm{ST}(\lambda)$, we set

$$\mathrm{res}_D(T) = (\mathrm{res}_D(i_m, j_m), \dots, \mathrm{res}_D(i_1, j_1)),$$

where $(i_k, j_k) \in \lambda$ is the box filled with k in T . For example, the residue are given as follows:

1	3	4	5	6	7	8	9	10
	2	3	4	5	6	7		
		1	3	4				
			2					

Then one can show that there exists $b \in B(\Lambda_1)$ with $\mathrm{wt}(b) = \mathrm{wt}_D(\lambda)$. We set

$$S(b) := \bigoplus_{T \in \mathrm{ST}(\lambda)} \mathbf{k}T,$$

and define an $R_{D_{n+1}}$ -module structure on $S(b)$ by

$$e(\nu)T = \delta_{\nu, \mathrm{res}_D(T)}T, \quad x_i e(\nu)T = 0, \quad \tau_j e(\nu)T = \begin{cases} s_j T & \text{if } \nu = \mu \text{ and } s_j T \in \mathrm{ST}(\lambda), \\ 0 & \text{otherwise,} \end{cases}$$

for $T \in \mathrm{ST}(\lambda)$ and admissible i, j, ν . Then it is straightforward to check that $S(b)$ is a homogeneous $R_{D_{n+1}}$ -module and it is isomorphic to $\mathcal{S}(b)$ given in Section 2.2.

Therefore, for each $\lambda \vdash m$, $R_{B_n}^{\Lambda_1}(\mathrm{wt}(\lambda))$ and $R_{D_{n+1}}^{\Lambda_1}(\mathrm{wt}_D(\lambda))$ are simple and their simple modules have the same dimension. Thus we conclude that $R_{B_n}^{\Lambda_1}(\mathrm{wt}(\lambda)) \simeq R_{D_{n+1}}^{\Lambda_1}(\mathrm{wt}_D(\lambda))$, which implies the desired result. \square

REFERENCES

1. S. Ariki and E. Park, *Representation type of finite quiver Hecke algebras of type $A_{2\ell}^{(2)}$* , J. Algebra **397** (2014), 457–488.
2. S. Ariki, K. Iijima, and E. Park, *Representation type of finite quiver Hecke algebras of type $A_\ell^{(1)}$ for arbitrary parameters* Int. Math. Res. Not. IMRN (2015), no. 15, 6070–6135.
3. N. Bourbaki, *Lie groups and Lie algebras Chapters 79*. Translated from the 1975 and 1982 French originals by Andrew Pressley. Elements of Mathematics (Berlin). Springer-Verlag, Berlin, 2005. xii+434 pp.
4. J. Brundan and A. Kleshchev, *Blocks of cyclotomic Hecke algebras and Khovanov-Lauda algebras*, Invent. Math. **178** (2009), 451–484.
5. G. Feinberg, K.-H. Lee, *Fully commutative elements of type D and homogeneous representations of KLR-algebras*, J. Comb. **6** (2015), no. 4, 535–557.
6. ———, *Homogeneous representations of type A KLR-algebras and Dyck paths*, Sémin. Lothar. Combin. 75 ([2015–2018]), Art. B75b, 19 pp.
7. M. Geck and G. Pfeiffer, *Characters of finite Coxeter groups and Iwahori-Hecke algebras*, London Mathematical Society Monographs. New Series, **21**. The Clarendon Press, Oxford University Press, New York, 2000.
8. J. Hong and S.-J. Kang, *Introduction to Quantum Groups and Crystal Bases*, Graduate Studies in Mathematics, **42**. American Mathematical Society, Providence, RI, 2002.
9. M. Jimbo and T. Miwa, *Solitons and infinite-dimensional Lie algebras*, Publ. Res. Inst. Math. Sci. **19** (1983), no. 3, 943–1001.
10. S.-J. Kang and M. Kashiwara, *Categorification of Highest Weight Modules via Khovanov-Lauda-Rouquier Algebras*, Invent. Math. **190** (2012), no. 3, 699–742.
11. M. Kashiwara, *Crystalizing the q -analogue of universal enveloping algebras*, Comm. Math. Phys. **133** (1990), no. 2, 249–260.
12. ———, *On crystal bases of the Q -analogue of universal enveloping algebras*, Duke. Math. J. **63** (1991), no. 2, 465–516.
13. ———, *Biadjointness in cyclotomic Khovanov-Lauda-Rouquier algebras*, Publ. Res. Inst. Math. Sci. **48** (2012), no. 3, 501–524.
14. M. Kashiwara, M. Kim, S.-j. Oh and E. Park, *Monoidal categories associated with strata of flag manifolds*, Adv. Math. **328** (2018), 959–1009.
15. M. Khovanov and A. D. Lauda, *A diagrammatic approach to categorification of quantum groups I*, Represent. Theory **13** (2009), 309–347.
16. ———, *A diagrammatic approach to categorification of quantum groups II*, Trans. Amer. Math. Soc. **363** (2011), no. 5, 2685–2700.
17. A. Kleshchev and A. Ram, *Homogeneous representations of Khovanov-Lauda algebras*, J. Eur. Math. Soc. (JEMS) **12** (2010), no. 5, 1293–1306.
18. R. Rouquier, *2 Kac-Moody algebras*, arXiv:0812.5023 (2008).
19. J. R. Stembridge, *Minuscule elements of Weyl groups*, J. Algebra **235** (2001), no. 2, 722–743.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF SEOUL, SEOUL 02504, KOREA

E-mail address: epark@uos.ac.kr